The timing and spectral properties of the 2022 outburst of SGR J1935+2154 observed with NICER

Fu Yu-Cong,^{1,2} Lin Lin,^{1,2} Ge Ming-Yu,³ Enoto Teruaki,⁴ Hu Chin-Ping,⁵ Younes George,^{6,7} Göğüş Ersin,⁸ and Malacaria Christian⁹

¹Institute for Frontiers in Astronomy and Astrophysics, Beijing Normal University, Beijing, 102206, China

²School of Physics and Astronomy, Beijing Normal University, Beijing, 100875, China

³Key Laboratory of Particle Astrophysics, Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, 100049, China

⁴Department of Physics, Graduate School of Science, Kyoto University, Kyoto, 606-8502, Japan

⁵Department of Physics, National Changhua University of Education, Changhua City, Taiwan

⁶Astrophysics Science Division, NASA Goddard Space Flight Center, 8800 Greenbelt Road, Greenbelt, MD, 20771, USA

⁷Center for Space Sciences and Technology, UMBC, Baltimore, MD, 21250, USA

⁸Sabancı University, Faculty of Engineering and Natural Sciences, İstanbul, 34956, Turkey

⁹ INAF-Osservatorio Astronomico di Roma, Via Frascati 33, Monteporzio Catone (RM), I-00078, Italy

ABSTRACT

The magnetar SGR J1935+2154 entered a new active episode on October 10, 2022, with X-ray bursts and enhanced persistent emission. At the tail of high burst rate interval, lasting several hours, radio bursts were detected, revealing the connection between the X-ray activities and radio emissions. We analyzed observations of SGR J1935+2154 for nearly three months, using data from Neutron Star Interior Composition Explorer (NICER). We report the timing and spectral results following the onset of this outburst. In general, the X-ray flux of the persistent emission decays exponentially. While a flare is evident on the light curve, a fast radio burst (FRB) was detected immediately following the peak of this flare. We found a phase jump of pulse profile, with a deviation of 0.16 ± 0.03 phase, which is related to the glitch. The spectra are well fit with the combination of a blackbody and a power law model. The decay of the outburst is dominated by the drop of the non-thermal component, which also leads to the increase of thermal proportion. The photon index of the power law is inversely correlated with both the unabsorbed flux and the burst rate. We find that unlike the large variety of the persistent emission around FRB 221014, the X-ray properties are very stable when FRBs 221021 and 221201 happened. These results manifest the connection between glitch, phase jump, X-ray burst, and radio burst, crucial for studying the mutation in twisted magnetic fields and constraining the trigger mechanism of radio bursts.

Keywords: Neutron stars (1108) — Magnetars (992) — X-ray transient sources (1852)

1. INTRODUCTION

Magnetars are a special type of isolated neutron stars (NSs), with distinctive phenomenons in X-ray and radio bands, which are mainly powered by their immense (up to $\sim 10^{15}$ G) magnetic fields (e.g., Kouveliotou et al. 1998; Olausen & Kaspi 2014; Kaspi & Beloborodov 2017). During the active period, the brightening of Xray emission from magnetars is often accompanied by bursts. The bursts can occur sporadically with a few bursts, or massive bursts are concentrated within a few hours (e.g., Israel et al. 2008; Lin et al. 2011; An et al. 2015; Cai et al. 2022). The persistent emission level also increases rapidly, accompanied by the variability of spectral and timing properties (e.g., Woods et al. 2004; Younes et al. 2017a; Borghese et al. 2022). Discovered for the first time in 2007 (Lorimer et al. 2007), Fast radio bursts (FRBs) are short pulses observed in the radio sky, and the explanation of their origin has become a central unresolved problem in astronomy (Cordes & Chatterjee 2019; Petroff et al. 2019). Although the physical mechanism of FRBs is not yet clear (Zhang 2020; Xiao et al. 2021), the connection between the magnetars and the origin of some FRBs has been established (Petroff et al.

Corresponding author: Lin Lin llin@bnu.edu.cn

2019; Bochenek et al. 2020; Lin et al. 2020c; Ge et al. 2023, 2024).

SGR J1935+2154 was discovered during its outburst in 2014 (Stamatikos et al. 2014). Then, the spin period of NS was calculated as about 3.24 s with a spin-down rate of $\dot{P} \sim 1.4 \times 10^{-11} \mathrm{s} \, \mathrm{s}^{-1}$, which implies a surface dipolar magnetic field strength of $B \sim 2.2 \times 10^{14} \text{ G}$ (Israel et al. 2016). Since then, as one of the most active magnetars, SGR J1935+2154 has shown outbursts in 2014, 2015, 2016, 2019, and 2020, which exhibit the different properties of persistent emission of the source (e.g., Younes et al. 2017b; Lin et al. 2020a,b; Younes et al. 2020; Borghese et al. 2022). The X-ray spectra below 10 keV are described well with a blackbody+powerlaw (BB+PL) or BB+BB model, and the cool BB temperature kT_{BB} is about 0.47 keV (Younes et al. 2017b). In 2020 outburst, with the decay in the luminosity, the contribution of the PL component was observed to decrease from $\sim 75\%$ to $\sim 45\%$ (Borghese et al. 2022; Younes et al. 2017b). X-ray burst and FRB 200428 are both observed in 2020 outburst, suggesting that they originate from the same source (CHIME/FRB Collaboration et al. 2020; Bochenek et al. 2020; Li et al. 2021; Ridnaia et al. 2021).

In 2022 October, SGR J1935+2154 entered a new active period, with several instruments triggered on the bursts (Palm 2022; Roberts et al. 2022; Li et al. 2022a; Enoto et al. 2022). During the peak of the active period, many bursts tend to cluster together, resulting in a high burst rate interval (Younes et al. 2022). After this interval, several FRB-like bursts were detected by GBT (Maan et al. 2022), CHIME (Dong & Chime/Frb Collaboration 2022; Pearlman & Chime/Frb Collaboration 2022), and Yunnan 40m radio telescope (Huang et al. 2022). The details of the FRBs are outlined as follows:

- FRB 221014, detected by CHIME/FRB on October 14, 2022, at 19:21:39.130 UTC (~ MJD 59866.82), with a fluence of 9.7±6.7 kJy ms (Dong & Chime/Frb Collaboration 2022; Giri et al. 2023), was accompanied by X-ray bursts captured by Konus-Wind (Frederiks et al. 2022) and GECAM (Wang et al. 2022).
- FRB 221021, detected by Yunnan 40m radio telescope on October 21, 2022, at 10:01:45.84215 UT (~ MJD 59873.42, Huang et al. 2022), was accompanied by an X-ray burst captured by *Insight*-HXMT (Li et al. 2022b,c).
- FRB 221201, was detected by CHIME/FRB on December 1, 2022, at 22:06:59.0762 UTC (\sim MJD 59914.92), with a fluence of 23.7 ± 18.0 kJy ms

(Pearlman & Chime/Frb Collaboration 2022; Giri et al. 2023).

During the 2022 outburst, the spectrum of the persistent emission observed with XMM-Newton and NuS-TAR is also well described by the BB+PL model below ~ 25 keV, and there are no significant changes in the BB temperature ($kT_{\rm BB} \sim 0.4$ keV) between the two epochs on October 15–18 and October 22, 2022 (Ibrahim et al. 2024). Hu et al. (2024) analyzed the data observed with NICER and NuSTAR during a month following the outburst, and discovered an unprecedented double glitch¹ associated with FRB 221014.

In this study, using NICER data of SGR J1935+2154 during its 2022 outburst, spanning nearly three months, we report the long-term evolution of the timing and spectral properties of this source. The observations and data reduction are presented in Section 2, the data analysis and results are described in Section 3, and, finally, the discussions and conclusions are given in Section 4.

2. OBSERVATIONS AND DATA REDUCTION

NICER is an X-ray timing and spectral instrument, which is successfully launched in 2017 and installed on the International Space Station (Gendreau & Arzoumanian 2017). NICER consists of 56 X-ray concentrator optics, 52 of which are currently operating. It covers the 0.2–12 keV energy range and provides a collecting area of 1900 cm² at 1.5 keV (Gendreau et al. 2016).

NICER started monitoring SGR J1935+2154 on October 12, 2022, at 17:32:40 UTC, during its 2022 active episode, with the detection of over 100 bursts and enhanced persistent emission (Younes et al. 2022). In this work, we analyzed the data from October 12 to December 21, 2022, with a total of 38 observations, as shown in Table 1. Each observation consists of several good time intervals (GTIs). For observations with high emission levels within the first three days, we divide them into smaller segments based on GTIs for further analysis.

We processed the NICER observations using NICER-DAS (v.10) as part of HEASOFT (v.6.31.1), along with the calibration database (CALDB) version 20221001. The standard calibration and filtering processes are performed by the task nicerl2² with the default parameters. The spectra, background, and associated responses in the NICER-recommended way³ are generated by the

- 1 The glitch means a sudden spin-up of the NS.
- ² https://heasarc.gsfc.nasa.gov/lheasoft/ftools/headas/nicerl2.html
 ³ https://heasarc.gsfc.nasa.gov/docs/nicer/analysis_threads/calrecommend/



Figure 1. NICER light curve in the 0.3–12 keV energy range. Top panel: the light curve of Obs. ID 5020560107 with a 4 ms resolution. The GTI between two blue dashed vertical lines is enlarged and displayed in the middle panel. Middle panel: the zoomed-in light curve of a GTI with a 4 ms resolution. Bottom panel: the zoomed-in light curve with a 3.2 s resolution is calculated after removing all the identified bursts.

task $nicerl3-spect^4$ with the SCORPEON background model. The light curves and associated background esti-

mation are generated using a pipeline task nicerl3-lc⁵ with the Space Weather background model.

To study the evolution of the persistent emission, we excluded all the identified bursts. We apply a Poisso-

⁴ https://heasarc.gsfc.nasa.gov/lheasoft/ftools/headas/nicerl3spect.html

 $^{^5}$ https://heasarc.gsfc.nasa.gov/docs/nicer/analysis_threads/nicerl3-lc/

nian procedure to identify the bursts (e.g., Gavriil et al. 2004) with a significance higher than $3-\sigma$, as described in Younes et al. (2020). To avoid interference from burst tails, we carefully remove the nearby bases of the bursts as much as possible, ensuring a clear analysis of the persistent emission. Figure 1 shows an example of burst subtraction, where the initial light curve of Obs. ID 5020560107 with a 4 ms resolution is displayed in the top panel, which is divided into 11 GTIs. The middle panel of Figure 1 presents a zoomed-in view of the GTI between two blue dashed vertical lines. After removing all the identified bursts, the light curve with 3.2 s resolution is shown in the bottom panel.

The events in the energy range 0.8–4 keV after the barycenter correction are used for timing analysis, because the effective area within this range is larger. We

use the epoch-folding technique (Leahy 1987) to search the initial period of the neutron star (e.g., Borghese et al. 2020; Tuo et al. 2020; Fu et al. 2023), and use the TEMPO2 (Edwards et al. 2006) version 2022.05.1 to update the more accurate ephemeris of the source. Times of arrival (ToAs) are acquired through Z_1^2 searching, and the minimum phase in each profile is used as the TOA for that specific observation (Ge et al. 2012, 2019; Younes et al. 2020). For the spectral analysis, we fit all the spectra using XSPEC (Arnaud 1996) version 12.13.0c. A photo-electric absorption model wabs (Anders & Ebihara 1982; Morrison & McCammon 1983) is used to calculate the interstellar absorption of SGR J1935+2154. The errors of the parameters are given at the level of 1- σ uncertainty.

Table 1. NICER observations of SGR J1935+2154 analyzed in this work.

Obs. ID-GTI ^a	Time ^b (MJD)	Exposure (s)	Burst ^c	Count Rate ^d	Index	Flux ^e	Part ^f	
5020560106-01	59864.7364	532.97	5	2.15 ± 0.08	2.42 ± 0.12	2.30 ± 0.12	I	
5020560106-02	59864.8011	655.96	2	2.27 ± 0.06	2.31 ± 0.13	1.75 ± 0.14	Ι	
5020560106-03	59864.8667	75.00	0	2.03 ± 0.16	2.40 ± 0.61	1.70 ± 0.44	Ι	
5020560107-01	59865.0068	212.98	2	2.48 ± 0.11	2.25 ± 0.27	1.62 ± 0.18	Ι	
5020560107-02	59865.0555	1320.92	11	2.71 ± 0.05	2.63 ± 0.08	2.10 ± 0.06	Ι	
5020560107-03	59865.1193	1391.92	6	2.68 ± 0.05	2.29 ± 0.08	2.05 ± 0.06	Ι	
5020560107-04	59865.1838	1515.91	9	2.62 ± 0.04	2.59 ± 0.07	2.03 ± 0.06	Ι	
5020560107-05	59865.2482	1574.90	7	2.29 ± 0.04	2.63 ± 0.09	1.77 ± 0.05	Ι	
5020560107-06	59865.3128	1581.90	7	2.37 ± 0.04	2.68 ± 0.08	1.87 ± 0.06	Ι	
5020560107-07	59865.3846	1027.93	7	2.29 ± 0.05	2.49 ± 0.10	1.75 ± 0.07	Ι	
5020560107-08	59865.4419	1631.90	6	2.23 ± 0.04	2.78 ± 0.08	1.68 ± 0.05	Ι	
5020560107-09	59865.5145	1764.86	2	1.71 ± 0.03	2.59 ± 0.09	1.31 ± 0.04	Ι	
5020560107-10	59865.5814	1569.87	2	1.86 ± 0.04	2.55 ± 0.10	1.30 ± 0.05	Ι	
5020560107-11	59865.6396	1310.91	3	1.98 ± 0.04	2.77 ± 0.10	1.56 ± 0.06	Ι	
5576010101-01	59865.7016	1951.87	2	2.00 ± 0.03	2.74 ± 0.09	1.17 ± 0.04	Ι	
5576010101-02	59865.7676	1160.93	2	1.97 ± 0.04	2.59 ± 0.13	1.14 ± 0.06	Ι	
5576010102-01	59866.1514	1892.87	10	2.41 ± 0.04	2.74 ± 0.09	1.17 ± 0.04	Ι	
5576010102-02	59866.2160	1889.87	5	2.67 ± 0.04	2.58 ± 0.08	1.79 ± 0.26	Ι	
5576010102-03	59866.4094	1289.92	6	5.14 ± 0.06	1.79 ± 0.10	3.06 ± 0.35	Ι	
5576010102-04	59866.4813	1303.90	10	5.54 ± 0.07	2.07 ± 0.06	4.35 ± 0.10	Ι	
5576010102-05	59866.6097	877.94	1	3.87 ± 0.07	2.15 ± 0.12	1.95 ± 0.08	Ι	
5576010102-06	59866.7331	1535.90	36	12.28 ± 0.09	1.31 ± 0.02	19.29 ± 0.18	II	
5576010102-07	59866.8676	807.95	5	6.02 ± 0.09	2.01 ± 0.06	4.97 ± 0.13	II	
5576010102-08	59866.9279	378.98	0	6.77 ± 0.13	1.96 ± 0.12	4.19 ± 0.20	II	
5576010103-01	59867.0008	867.93	1	4.83 ± 0.07	2.14 ± 0.11	3.73 ± 0.17	II	
5576010103-02	59867.0680	1311.89	0	5.97 ± 0.07	2.07 ± 0.10	3.59 ± 0.15	II	
5576010103-03	59867.1835	1973.87	0	4.12 ± 0.05	2.37 ± 0.08	2.81 ± 0.09	II	
5576010103-04	59867.3236	998.92	0	3.98 ± 0.06	2.24 ± 0.14	2.37 ± 0.14	II	
5576010103-05	59867.3880	999.92	0	3.24 ± 0.06	2.56 ± 0.14	2.20 ± 0.10	II	
5576010103-06	59867.5167	1019.92	0	2.40 ± 0.06	2.32 ± 0.14	2.03 ± 0.12	II	
Continued on next page								

Table 1 –Continued from previous page

Obs. ID-GTI ^a	$Time^{b}$ (MJD)	Exposure (s)	$\operatorname{Burst}^{\operatorname{c}}$	$Count \ Rate^d$	Index	$\operatorname{Flux}^{\operatorname{e}}$	Part $^{\rm f}$
5576010103-07	59867.7805	584.95	0	1.41 ± 0.06	2.49 ± 0.21	1.13 ± 0.15	II
5576010104	59868.0968	9993	2	1.99 ± 0.02	2.62 ± 0.03	1.80 ± 0.04	II
5576010105	59869.0539	2609	4	1.83 ± 0.03	2.61 ± 0.07	1.94 ± 0.05	II
5576010106	59870.1655	2474	0	1.28 ± 0.03	2.97 ± 0.06	1.43 ± 0.05	II
5576010108	59872.2922	1693	0	1.23 ± 0.04	3.22 ± 0.10	1.27 ± 0.07	II
5576010109	59873.0013	3021	0	0.98 ± 0.03	3.05 ± 0.20	0.92 ± 0.06	II
5576010110	59874.0871	2034	1	1.46 ± 0.04	2.88 ± 0.11	1.38 ± 0.05	III
5020560108	59875.9127	260	0	2.07 ± 0.22	2.39 ± 0.84	0.76 ± 0.18	III
5020560109	59877.2640	670	0	0.99 ± 0.07	3.04 ± 0.18	0.89 ± 0.06	III
5576010111	59878.3503	6596	0	1.04 ± 0.02	2.90 ± 0.06	1.17 ± 0.03	III
5576010112	59886.5579	753	0	1.02 ± 0.08	2.82 ± 0.28	1.24 ± 0.18	III
5020560110	59888.3810	1194	0	0.96 ± 0.04	3.55 ± 0.24	0.80 ± 0.06	III
5020560111	59889.0319	728	0	0.89 ± 0.05	2.81 ± 0.19	0.76 ± 0.05	III
5576010113	59889.3457	1094	0	0.90 ± 0.03	3.21 ± 0.17	0.77 ± 0.03	III
5020560112	59892.9453	1161	2	1.03 ± 0.04	2.68 ± 0.12	1.38 ± 0.05	III
5576010114	59893.0096	1134	0	1.27 ± 0.05	2.82 ± 0.13	1.19 ± 0.06	III
5020560113	59893.9125	1717	1	1.04 ± 0.03	3.35 ± 0.15	1.06 ± 0.04	III
5020560114	59894.1060	8347	1	1.41 ± 0.13	3.01 ± 0.05	1.38 ± 0.02	III
5020560115	59895.0090	10778	2	1.42 ± 0.07	2.72 ± 0.03	1.28 ± 0.03	III
5576010115	59895.7272	437	0	0.89 ± 0.05	3.03 ± 0.51	0.78 ± 0.12	III
5020560116	59896.0497	7068	0	0.92 ± 0.03	3.08 ± 0.05	1.24 ± 0.03	III
5020560117	59897.0086	10996	2	0.87 ± 0.02	3.02 ± 0.04	1.31 ± 0.02	III
5576010116	59898.4921	779	0	0.79 ± 0.06	3.11 ± 0.22	1.03 ± 0.14	III
5576010117	59901.3300	1209	0	0.77 ± 0.03	3.87 ± 0.25	0.93 ± 0.07	III
5020560118	59902.4340	375	0	0.85 ± 0.05	2.70 ± 0.42	1.01 ± 0.18	III
5576010121	59913.4562	909	0	0.84 ± 0.03	3.19 ± 0.27	0.62 ± 0.06	III
5020560122	59924.0454	760	0	0.91 ± 0.07	3.62 ± 0.26	0.93 ± 0.09	III
5576010122	59925.2061	2078	0	0.80 ± 0.03	2.95 ± 0.12	0.82 ± 0.06	III
5576010123	59929.9123	7	0	0.49 ± 0.23	2.89 ± 0.57	< 0.69	III
5576010124	59930.9444	37	0	0.53 ± 0.19	3.55 ± 0.46	< 0.57	III
5576010125	59931.0090	36	0	0.67 ± 0.14	3.78 ± 0.56	< 0.25	III
5576010126	59932.9440	42	0	0.74 ± 0.21	2.69 ± 0.49	< 0.38	III
5576010127	59933.0087	31	0	0.61 ± 0.16	2.50 ± 0.47	< 0.71	III
5576010128	59934.2338	64	0	0.81 ± 0.17	2.67 ± 0.50	< 0.83	III

NOTES-

The errors are calculated with 1- σ level uncertainties.

^a Obs. ID-NN, NN represents the serial number of the good time intervals (GTIs).

^b The start time of each observation or GTI.

 $^{\rm c}$ The number of the bursts with a significance higher than 3- $\sigma.$

^d The 0.3–12 keV net count rate after removing bursts.

^e The 0.8–12 keV unabsorbed flux in units of 10^{-11} erg s⁻¹ cm⁻².

^f The three temporal parts are defined from the pulse profiles analysis in Section 3.1.

3. RESULTS

3.1. Timing analysis

According to the study by Hu et al. (2024), there are double glitches around the high burst rate interval. The timing parameter \dot{f} between glitches jumps significantly, and the order of magnitude of \dot{f} before the first



6



Figure 2. The evolution of the pulse profiles. (a): The two-dimensional (2D) maps describe the evolution of the pulse profiles with time for NICER (0.8–4 keV). The colors representing the values of the pulse profile are normalized by Pulse/Average count rate; the red represents pulse-on phase, and the blue represents pulse-off phase. 20 bins within a phase are used to generate the pulse profiles, and the plot is smoothed through interpolation and Gaussian filtering for clarity. Phase zero is defined as the reference epoch MJD 59865 as shown in Table 2. The black dash-dotted line marks the position of the phase jump. The grey solid, dashed, and dotted lines mark the FRB 221014, FRB 221021, and FRB 221201, respectively. Two cycles are shown for clarity. (b): The average pulse profiles of SGR J1935+2154. As shown in Table 1, Part I represents the pulse profile from the start to the phase jump. Part II represents the pulse profile from the phase jump to FRB 221021. Part III represents the pulse profile from FRB 221021 to the end. The errors are calculated with 1- σ level uncertainties.

Table 2. Best fits spin parameters of SGR J1935+2154 from the 2022 October 12 to December 21 (excluding glitch time interval). The reference epoch is designated as the phase zero of the pulse profile.

Parameter	Value
R.A. (J2000)	19:34:55.68
Decl. (J2000)	21:53:48.2
Solar ephemeris	DE405
Start (MJD)	59864
Finish (MJD)	59934
Reference epoch (MJD)	59865
Spin frequency f (Hz)	0.30752804(6)
Spin-down rate \dot{f} (Hz s ⁻¹)	$-4.82(3) \times 10^{-12}$
Second derivative \ddot{f} (Hz s ⁻²)	$-4.9(7) \times 10^{-21}$
χ^2/dof	56/43

and after the second glitches is consistent. Therefore, we deducted the data between the two glitches (MJD 59866.63–59866.99) to reduce their impact on the longterm evolution of the timing solution. As shown in Table 2, from the best-fitting of long-term data over a period of MJD 59864–59934, the spin frequency and the spin-down rate are calculated as f = 0.30752804(6) Hz and $\dot{f} = -4.82(3) \times 10^{-12}$ Hz s⁻¹, respectively. This is slower than the spin measurement of 0.30789626(2) from the 2020 October outburst (Younes et al. 2023), with a difference of 0.00036822(6) Hz. The difference is consistent with a straightforward estimate; considering a two-year interval and only \dot{f} , the difference is calculated as ~ 0.0003 Hz, which is consistent with the above values in terms of magnitude.

The pulse profiles exhibit a single-peaked shape in the 0.8–4 keV energy range, consistent with the result in Lu et al. (2024). As shown in Figure 2(a), the colors representing the values of the pulse profile are normalized by Pulse/Average count rate, the red represents pulse-on, and the blue represents pulse-off. During the outburst, the shape of single-peaked profiles shows no significant change, the gradual broadening of the pulse in the green area in Figure 2(a) is caused by a decrease in signal-to-noise ratio (SNR), and it cannot indicate that the profile has evolved. However, the phase of the pulse profile changes significantly, the phase of the pulse of the pulse of the main peak remains at ~ 0.50 after MJD 59866.73. Between the MJD 59866.62 and MJD 59866.73, a phase



Figure 3. Spectra and residual of Part I (top figures), Part II (middle figures), and Part III (bottom figures). Left figures: the total model (orange line) is plotted together with powerlaw (blue line) and the background model (grey line). Right figures: the total model (orange line) is plotted together with powerlaw (sky-blue line), bbody (red line), and the background model (grey line). The spectra are re-binned to display clarity. The errors are calculated with $1-\sigma$ level uncertainties.

jump seems to occur, with a deviation of ~ 0.15 , which is consistent with Extended Data Figure 3 of Hu et al. (2024). For the convenience of representation, we use the midpoint of this time interval (MJD 59866.68) as the time when the phase jump occurs marked with the black dash-dotted line. The grey solid, dashed, and dotted lines mark the positions of the FRB 221014, FRB 221021, and FRB 221201, respectively. The phase jump and FRB 221021 divide the data into three parts: Part I, Part II, and Part III.

As shown in Figure 2(b), the average pulse profiles of there parts are displayed. Part I represents the pulse profile from the start (~ MJD 59864.73) to the phase jump (~ MJD 59866.68), Part II represents the pulse profile from the phase jump to FRB 221021 (~ MJD 59873.42), and Part III represents the pulse profile from FRB 221021 to the end (~ MJD 59934.24). The intensity of the pulse profiles are normalized by their average counts. Phase zero is defined as the reference epoch of ephemeris as shown in Table 2. The pulse profiles of Part I and Part II are similar in shape, but there is a significant deviation in phase, and the main peak phase of Part II shifts 0.16 ± 0.03 to the left. After ignoring statistical fluctuations, the pulse profiles of Part II and Part II are basically consistent.

The position of the radio bursts on the 0.8–4 keV Xray pulse profile of persistent emission is also marked in Figure 2(b). The phase of radio bursts is calculated using the timing solution obtained from X-ray. Consistent with the result from Hu et al. (2024), FRB 221014 appears at the position of pulse-off after the peak (part II), which seems to be consistent with the radio burst B1 in Figure 5 by Younes et al. (2020). FRB 221201 appears in the rising phase before the peak (part III), which seems to be consistent with the radio burst B2 in Figure 5 by Younes et al. (2020). It should be noted that the position of FRB 221201 is greatly influenced by the ephemeris \dot{f} or \ddot{f} , as it spans a large time interval from the reference epoch.

3.2. Spectral analysis

3.2.1. Phase-averaged spectral analysis

The selection of spectral models has been systematically studied during the previous outbursts; the spectra below 10 keV are described by the combination of bbody and powerlaw (Younes et al. 2017b, 2020; Borghese et al. 2022). Shao et al. (2024) also reported that most of the spectra (in the range of 1–5 keV) could be better described with an absorbed bbody plus powerlaw model during 2022 outburst. We attempted to fit the spectra in 0.8–12 keV energy range using different model combinations of **bbody** and **powerlaw**. Based on the selection criteria of residual showing no significant structure, combined with χ^2 /dof values less than 1.3 and the null hypothesis probability for the improvement from Ftest less than 0.05, we determined that wabs*powerlaw (Abbreviated as Model I) and wabs*(bbody+powerlaw) (Model II) were the best choices for fitting spectra. The hydrogen column density of wabs is fixed at $N_{\rm H} =$ $2.3 \times 10^{22} \,{\rm cm}^{-2}$ according to the previous studies (e.g. Younes et al. 2017b; Coti Zelati et al. 2018; Borghese et al. 2022).

In Figure 3, the average spectra of Part I, Part II, and Part III fitted with these two model combinations are displayed. The reason why the residuals are higher on the high energy side is probably because the hard power law above 10 keV is contaminating the data. The Chi-squared test indicates that there is no significant difference between the two combinations for Part I and Part II.

For Part I, the χ^2/dof of Model I and Model II are $1596/1418 \approx 1.126$ and $1591/1416 \approx 1.124$, respectively, as shown in Figure 3 (a1) and (b1). The Chi-squared statistics of Model I and Model II are almost equivalent, thus making it impossible to distinguish which model is the best fit based on this criterion. The F-test gives a null hypothesis probability for the improvement of the fit of 0.11. Considering the inclusion of bbody, it is not statistically necessary, but is needed in the fit. This is because bbody has been observed in previous outbursts (Younes et al. 2017b, 2020; Borghese et al. 2022) and at other temporal parts within this outburst, indicating its possible physical presence. However, due to spectral variability, its statistical significance may not be prominent. The flux of bbody (red line) is one order of magnitude smaller than the flux of powerlaw (sky-blue line), thus bbody does not have a significant impact on the residual of spectra

For Part II, the χ^2/dof of Model I and Model II are 1395/1418 ≈ 0.983 and 1363/1416 ≈ 0.960 , respectively, as shown in Figure 3 (a2) and (b2). The F-test gives a null hypothesis probability for the improvement of the fit of 7.32×10^{-8} , indicating that the inclusion of the bbody component is statistically significant and thus necessary in the model.

For Part III, the χ^2 /dof of Model I and Model II are 1270/1099 \approx 1.16 and 1127/1097 \approx 1.03, respectively. The obvious residual structure in Figure 3 (a3) and (b3) demonstrates the significance of the **bbody** component. Additionally, the flux of **bbody** accounts for $42.9 \pm 1.1\%$ of the total flux.

Table 3. Results of the spectral parameters presented in Section 3.2.

Model	Model I, wabs*powerlaw			$Model \ II, \ \texttt{wabs}^*(\texttt{bbody} + \texttt{powerlaw})$			
Epoch	Part I	Part II	Part III	Part I	Part II	Part III	
$kT_{\rm BB}~({\rm keV})$	-	-	-	0.43 ± 0.02	0.37 ± 0.02	0.35 ± 0.01	
R^{a}_{BB} (km)	-	-	-	1.9 ± 0.1	2.9 ± 0.2	4.2 ± 0.2	
Γ	2.30 ± 0.01	2.09 ± 0.01	2.95 ± 0.02	2.08 ± 0.06	1.81 ± 0.07	2.00 ± 0.14	
$Norm^b PL$	6.7 ± 0.1	6.1 ± 0.1	6.9 ± 0.1	4.5 ± 0.4	3.7 ± 0.3	1.6 ± 0.2	
Total $\operatorname{Flux}^{\operatorname{c}}$	2.12 ± 0.01	2.26 ± 0.01	1.32 ± 0.01	2.06 ± 0.01	2.23 ± 0.01	1.19 ± 0.02	
BB $\operatorname{Flux}^{\operatorname{c}}$	-	-	-	0.22 ± 0.02	0.28 ± 0.01	0.51 ± 0.01	
$PL \ Flux^{c}$	-	-	-	1.85 ± 0.02	1.94 ± 0.02	0.68 ± 0.02	
Ratio ^d (%)	-	-	-	10.7 ± 0.9	12.6 ± 0.5	42.9 ± 1.1	
$\chi^2/{ m dof}$	1596/1418	1395/1418	1270/1099	1591/1416	1363/1416	1127/1097	

Note—

The errors are calculated with 1- σ level uncertainties.

^a The source radius in km at the distance of 6.6 kpc (Zhou et al. 2020).

^b The normalization of the power law in units of $\times 10^{-3}$ photons $\cdot \text{ keV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$.

^c The 0.8–12 keV unabsorbed total, BB and PL flux in units of $\times 10^{-11}$ erg s⁻¹ cm⁻².

^d The ratio of the **bbody** flux to total flux.

As shown in Table 3, the decrease in the total flux of Part III relative to Part I and Part II is mainly contributed by the decrease in powerlaw, while accompanied by a slight increase in bbody. The temperature of bbody slightly decreases, while its radius grows from $\sim 2 \text{ km to} \sim 4 \text{ km}$. The proportion of bbody component increases from $\sim 10\%$ to $\sim 40\%$ (powerlaw component proportion decreases from $\sim 90\%$ to $\sim 60\%$), which is consistent with the results of 2020 outburst by Borghese et al. (2022), where the powerlaw component proportion decreased from $\sim 75\%$ to $\sim 45\%$.

3.2.2. Spectral evolution

We perform the spectral fitting of each observation, and the results are shown in Table 1. In Figure 4, the evolution of the flux, burst rate, photon index (Γ), and the ratio of the **bbody** flux to the total flux with time are displayed from top to bottom panels.

In Figure 4 (a), the unabsorbed total flux initially remains at ~ $(1-2) \times 10^{-11} \mathrm{erg \, s^{-1} \, cm^{-2}}$. Then there is a flare around the phase jump which is marked with the black dash-dotted line, and the flux increases more than ten times at the peak of the flare, reaching ~ $19 \times 10^{-11} \mathrm{erg \, s^{-1} \, cm^{-2}}$. The phase jump is close to the peak of the flare, and it can be discerned that it precedes the peak. The flare starts on ~ MJD 59866.4 and ends on ~ MJD 59867.2, lasting for less than a day. The blue line fits the exponential decay of the flux, with the fitting result being $Flux(10^{-11}) = 2.5(2)e^{-0.025(5)T(\mathrm{days})}$, based on data after MJD 59867.068. The characteristic

time of the decay is ~ 40 days. The pink dashed line represents the quiescent flux level as reported by Shao et al. (2024). It is worth noting that FRB 221014 (Maan et al. 2022; Dong & Chime/Frb Collaboration 2022) appears slightly behind the peak of the flare, marked with the grey solid line. The grey dashed and grey dotted lines also mark the FRB 221021 and FRB 221201, respectively.

In Figure 4 (b), the burst rate is defined as the number of bursts divided by the exposure time per observation or GTI. From the start of the outburst, the burst rate decreases until the beginning of the flare, during which the burst rate significantly increases, and there is a high burst rate interval at the peak of the flare. Higher energy bursts from Fermi/GBM are marked with orange arrows. The orange arrows gather more densely just behind the black dash-dotted line, which also indicates that there is a high burst rate interval. After the flare, the burst rate rapidly decreases to zero, and there are also almost no higher energy bursts. There is a positive correlation between the burst rate and the flux as shown in Figure 5 (b). The P-value of the Pearson correlation test is 2×10^{-14} , indicating a statistically significant positive correlation, with a correlation coefficient of 0.79.

In Figure 4 (c), the evolution of the photon index from Model I and II is displayed. The red data points come from Model I (PL). When the outburst decays normally, the photon index is ~ 2.5 and slowly increases, but decreases to ~ 1.3 during the flare. After the flare, the photon index gradually increases and remains at ~ 3 .



Figure 4. Evolution of the flux, burst rate, photon index (Γ), and the ratio of the bbody flux to the total flux with time. Panel (a): the evolution of the unabsorbed flux in units of 10^{-11} erg s⁻¹ cm⁻². The fluxes are computed in the energy range 0.8–12 keV. The blue line fits the exponential decay of the flux, with the result of $y = 2.5(2)e^{-0.025(5)x}$. The pink dashed line represents the quiescent flux level as reported by Shao et al. (2024). Panel (b): the evolution of the burst rate. Bursts are identified with a significance higher than 3- σ . Black hollow points represent zero. The orange arrows represent the identified bursts from Fermi/GBM. Panel (c): the evolution of the photon index of power law. The red data points come from Model I (PL), and the gray points come from Model II (BB+PL). Panel (d): the evolution of the temperature of BB. Panel (e): the evolution of the ratio of the bbody flux to the total flux. The errors are calculated with 1- σ level uncertainties. The black dash-dotted line marks the phase jump time of the pulse profile which is ~ MJD 59866.68, and grey solid, dashed, and dotted lines mark the FRB 221014, FRB 221021, and FRB 221201, respectively.

The gray points come from Model II (BB+PL). When the proportion of BB is small ($\leq 30\%$), it is consistent with the results of Model I. And when the proportion of BB increases ($\geq 30\%$), the photon index remains at ~ 2, which is smaller than the results of Model I. After MJD 59875, the decrease in data SNR leads to the dispersion of the photon index and larger errors. As shown in Figure 5 (a) and (c), the evolution of the photon index (Model II) is inversely proportional to the flux and the burst rate. The Pearson correlation test yields a Pvalue of 8×10^{-8} with a correlation coefficient of -0.61for the flux, and a P-value of 4×10^{-6} with a correlation coefficient of -0.54 for the burst rate, respectively.

In Figure 4 (d), the temperature of bbody varies between ~ 0.3 keV and ~ 0.5 keV, which is roughly consistent with the results from Shao et al. (2024). There is no significant evolution in temperature, with an average value of 0.38 keV.

In Figure 4 (e), the ratio of the bbody flux to the total flux increases from ~ 10% at the beginning to ~ 60% at the end, with a maximum value of $75 \pm 7\%$. A linear model is used to fit this growth trend, and the fitting result is Ratio(%) = 0.90(6)T(days) + 12(2), which means that the bbody ratio is growing at a rate of ~ 1% per day. Some missing data points are due to the poor data quality of these observations which is not sufficient to limit the complexity of Model II.

3.2.3. Phase-resolved spectral analysis

In order to analyze the possible transition of the spectral property before and after the phase jump, we study the phase-resolved spectral analysis on the average spectra of Part I, Part II, and Part III, respectively.

For Model I, the entire cycle is divided into ten phase intervals. In Figure 6, the photon index and the normalization of the power law of three parts are presented. respectively. For Part I, there is an inverse correlation between the photon index and the pulse profile, with a correlation coefficient of -0.53. The photon index varies between 2.34 and 2.55, with the minimum pulse (~ 0.3) corresponding to 2.52 ± 0.06 and the maximum pulse (~ 0.7) corresponding to 2.35 ± 0.04 . The normalization is proportional to the pulse profile, with a correlation coefficient of 0.87. For Part II, there is also an inverse correlation between the photon index and the pulse profile, with a correlation coefficient of -0.52. The photon index varies between 2.03 and 2.28, with the minimum pulse (~ 0.9) corresponding to 2.22 ± 0.05 and the maximum pulse (~ 0.6) corresponding to 2.05 ± 0.03 . The normalization is proportional to the pulse profile, with a correlation coefficient of 0.67. For Part III, the evolution is similar to Part II, and the correlation coefficient

between the photon index and the pulse is -0.47. The normalization is proportional to the pulse profile, with a correlation coefficient of 0.48.

In brief, when the phase jump occurs (Part I and Part II), the spectral parameters also correspondingly change, when the pulse phase remains constant (Part II and Part III), the spectral parameters also approximately remain constant. The evolution of spectral parameters with phase does not change.

For Model II, the entire cycle is divided into two phase segments, pulse on and pulse off. As shown in Figure 7, there is no significant change in $kT_{\rm BB}$ and Γ between pulse on and pulse off for all three parts, which is consistent with the result from Ibrahim et al. (2024). The source radius R_{BB} remains consistent across the different phase segments for Parts I and II, but for Part III, the R_{BB} of pulse off is larger than that of pulse on, increasing by ~15%. Meanwhile, for three Parts, from pulse on to pulse off, the PL normalization is decreasing.

4. DISCUSSION

We analyzed the NICER observations of SGR J1935+2154 during its 2022 active period and performed the timing and spectral analysis of the X-ray persistent emission. During the decay of the outburst, we found a phenomenon of the phase jump of the pulse profile, with a deviation of 0.16 ± 0.03 , as shown in Figure 2. Before and after the phase jump, the phase-resolved spectral properties also changed correspondingly, as shown in Figure 6. Around the phase jump, a flare appeared on the light curve accompanied by changes in spectral properties. The phase jump is slightly earlier than the flare, and FRB 221014 is closely following the flare, as shown in Figure 4. Furthermore, there are no significant changes in the properties of X-ray persistent emission near FRB 221021 and FRB 221201.

The increase of the persistent X-ray flux is quite common in magnetars, accompanied by spectral hardening, manifested as a decrease in the power law (PL) index or an increase in the blackbody (BB) temperature (e.g., Israel et al. 2010; Coti Zelati et al. 2018). This spectral evolution is also ubiquitous in the previous active period of SGR J1935+2154 (Israel et al. 2016; Younes et al. 2017b, 2020), as well as other sources such as 1E 1547– 5408 (Scholz & Kaspi 2011) and 1E 2259+586 (Kaspi et al. 2003). In this study, the spectra are well modeled by a single PL with a photon index $\Gamma \sim (1.3 - 3.0)$. The Chi-squared statistics and the insensitive fitting of the BB indicate that the thermal component is not important. These characteristics are slightly different from the 2020 outburst of SGR J1935+2154, where a thermal component is required to model the spectra (Younes



Figure 5. The correlation between the 0.8–12 keV flux, the photon index Γ , and the burst rate in Figure 4. In Panels (b) and (c), the black hollow square points represent the burst rate of zero. The errors are calculated with 1- σ level uncertainties.



Figure 6. Fitting results of the phase-resolved spectral analysis of Model I. Solid lines represent the pulse profiles as shown in Figure 2(b). The errors are calculated with 1- σ level uncertainties. The values in parentheses represent the correlation coefficient between the pulse and parameters, with positive numbers indicating positive correlation and negative numbers indicating negative correlation.

et al. 2020; Borghese et al. 2022). In the early observation of the 2020 outburst, the proportion of PL accounting for ~ 75% (Borghese et al. 2022) is less than ~ 90% of the 2022 outburst, which could explain the insensitivity of the BB component in the spectral fitting in this study. In Figure 4, the PL photon index fitting results of Model I and Model II show that when the proportion of BB is greater than ~ 30%, the thermal component will affect the fitting of the photon index. When the proportion of BB is less than ~ 30%, the influence of thermal components on the photon index is negligible. The proportion of BB seems to have a potential relationship with the burst rate. On the ~ MJD 59895, where the burst rate increases, the BB ratio decreases, with a slight increase in flux, indicating the possible existence of a small flare here.

The combination of a BB and a PL component is generally interpreted as the thermal emission from the surface of NS, and the scattering of seed photons by physical mechanisms taking place in the magnetosphere, such as the Resonant Cyclotron Scattering (Nobili et al. 2008) and the currents scattering in the twisted magnetosphere model (Thompson et al. 2002). The degree of scattering increases with the increasing magnetic twist, which could be localized to a restricted portion of the magnetosphere (e.g., Özel & Güver 2007; Beloborodov 2009). The persistent emission during the SGR J1935+2154 outburst dominated by the PL com-



Figure 7. Fitting results of the phase-resolved spectral analysis of Model II. Solid lines represent the pulse profiles as shown in Figure 2(b). The errors are calculated with 1- σ level uncertainties. The source radius in km at the distance of 6.6 kpc (Zhou et al. 2020). The normalization of the power law in units of 10^{-3} photons $\cdot \text{keV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$.

ponent (~ 90%) indicates that the twisted magnetic field plays an important role in the outburst. During the decay of the outburst, the temperature and flux of BB do not show a significant evolution, while the flux of PL shows a decreasing trend. This similar phenomenon also has been observed in 2014, 2015, and 2016 outbursts; the flux of BB remains at $\sim 1.5 \times 10^{-12} \mathrm{erg \, s^{-1} \, cm^{-2}}$, and the flux of PL decreases from $\sim (3-8) \times 10^{-12} \mathrm{erg \, s^{-1} \, cm^{-2}}$ to $\sim 1 \times 10^{-12} \text{erg s}^{-1} \text{ cm}^{-2}$ (Younes et al. 2017b). According to the spectral results obtained by Hu et al. (2024), during the early stage of outburst decay, the BB temperature first increases and then decreases with the flare, with this rapid decline terminating approximately 10 hours after the FRB. This rapid decrease is consistent with the result of the 2020 outburst, where BB temperature rapidly decreased within 3 days after the outburst, then entered a slow decrease stage lasting for several tens of days (Younes et al. 2020).

The pulse profile exhibits a single-peaked shape without a significant variability in the 0.8–4 keV energy band, consistent with the previous observations (Borghese et al. 2022; Younes et al. 2023). However, particularly, a phase jump was discovered around the peak of a flare during this outburst. Continuous observations indicate that the phase jump occurs between MJD 59866.62 and MJD 59866.73, which coincides with the first glitch time (MJD 59866.63) from NICER and NuSTAR data (Hu et al. 2024). Figure 2(b) shows that the overall left shift of the pulse profile is 0.16(3) phase, including the pulse-on and the pulse-off. The phase-resolved spectral analysis also supports the phase jump, before and after the phase jump, the fitting parameters of phase-resolved spectra always maintain a constant correlation with the pulse profile. In other words, the spectral properties also

shift 0.16(3) phases to the left after the phase jump. The consistency between phase jump and glitch time indicates that this is due to the glitch in the timing solution, which caused by magnetospheric wind (Younes et al. 2023; Hu et al. 2024). However, at the second glitch time (MJD 59866.99), there is no obvious phase jump here, which may be due to data quality blurring possible phase jumps, but it may also indicate that only a few special glitches are associated with the phase jumps, indicating that there may be other physical mechanisms that cause this phase jump.

Figure 4 shows that the long-term evolution of the unabsorbed flux, burst rate, and photon index of PL, with the strong correlations between each parameter. Around the phase jump, a brief high burst rate interval appears at the peak of the flare, and the phase jump occurs just before this interval, which seems to suggest a connection between the phase jump and the X-ray burst. The high burst rate interval accompanied by the enhancement of persistent emission has been observed in previous studies (e.g. Younes et al. 2020; Borghese et al. 2022), and the spectrum becomes harder and the photon index decreases (Mereghetti et al. 2009), which are consistent with the results in this study. The burst rate rapidly decreases to zero within a few hours after the flare, and almost no bursts are detected thereafter. The connections with the bursts and the hardness are related to the evolution of a twisted magnetic field (Thompson et al. 2002; Israel et al. 2010). After the flare, the photon index tends to recover to the pre-flare level, the recovery process is relatively slow and does not reach the previous level in a short period of days. This may indicate a certain degree of lag in the evolution of the system (Woods et al. 2007). The simultaneous occurrence of phase jump, glitch, high burst rate interval, and the spectral evolution may indicate more complex scenarios, which may be caused by the mutation in twisted magnetic fields.

We also noted the appearance of FRB 221014 just behind the flare as marked with the black dash-dotted line in Figure 4, and this temporal consistency indicated a potential physical connection between FRBs and the twisted magnetic fields. However, The triggering mechanism of FRBs is still unclear. It is generally suggested that FRBs are generated by the magnetospheric activity of the magnetars, including internal (e.g. Yang & Zhang 2018; Lu et al. 2020) and external (e.g. Zhang 2017; Dai 2020) triggering. From the study of FRB 200428, FRBs preferably occur after the most active episodes triggered by giant glitches, which points to an active magnetic field, as the magnetic field rearrangement is completed, magnetic activity becomes less frequent and FRBs become more difficult to generate (Ge et al. 2024). This supports the conjecture that a significant change in twisted magnetic fields leads to the phase jump, while the change in the magnetospheric structure results in the gradual appearance of the high burst rate interval and the FRB. Following FRB 200428, several radio bursts emerge at different phases without significant X-ray variability (Younes et al. 2020). Similarly,

subsequent to FRB 221014, there are also several radio bursts detected, with no significant changes observed in the timing or spectral properties of the persistent X-ray emission near FRB 221021 and FRB 221201 (Ge et al. 2024). The outburst decays by $\gtrsim 50\%$, and lower data quality may make small changes invisible, which also indicates the diversity and potential differences of FRBs.

This work has made use of data from the NICER mission, and data and/or software provided by the High Energy Astrophysics Science Archive Research Center (HEASARC), a service of the Astrophysics Science Division at NASA/GSFC. This work is supported by the National Key R&D Program of China (2021YFA0718500) from the Minister of Science and Technology of China (MOST). The authors thank supports from the National Natural Science Foundation of China under Grants U2038103, U2038101, U2038102 and 12373051. This work is also supported by International Partnership Program of Chinese Academy of Sciences (Grant No.113111KYSB20190020).

Software: ASTROPY (Astropy Collaboration et al. 2013, 2018), XSPEC (Arnaud 1996), and STINGRAY (Huppenkothen et al. 2019).

REFERENCES

- An, H., Archibald, R. F., Hascoët, R., et al. 2015, ApJ, 807, 93, doi: 10.1088/0004-637X/807/1/93
- Anders, E., & Ebihara, M. 1982, GeoCoA, 46, 2363, doi: 10.1016/0016-7037(82)90208-3
- Arnaud, K. A. 1996, in Astronomical Society of the Pacific Conference Series, Vol. 101, Astronomical Data Analysis Software and Systems V, ed. G. H. Jacoby & J. Barnes, 17
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, A&A, 558, A33, doi: 10.1051/0004.6261/201222068
 - doi: 10.1051/0004-6361/201322068
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, AJ, 156, 123, doi: 10.3847/1538-3881/aabc4f
- Beloborodov, A. M. 2009, ApJ, 703, 1044, doi: 10.1088/0004-637X/703/1/1044
- Bochenek, C. D., Ravi, V., Belov, K. V., et al. 2020, Nature, 587, 59, doi: 10.1038/s41586-020-2872-x
- Borghese, A., Coti Zelati, F., Rea, N., et al. 2020, ApJL, 902, L2, doi: 10.3847/2041-8213/aba82a
- Borghese, A., Coti Zelati, F., Israel, G. L., et al. 2022, MNRAS, 516, 602, doi: 10.1093/mnras/stac1314

- Cai, C., Xue, W.-C., Li, C.-K., et al. 2022, ApJS, 260, 24, doi: 10.3847/1538-4365/ac6172
- CHIME/FRB Collaboration, Andersen, B. C., Bandura, K. M., et al. 2020, Nature, 587, 54, doi: 10.1038/s41586-020-2863-y
- Cordes, J. M., & Chatterjee, S. 2019, ARA&A, 57, 417, doi: 10.1146/annurev-astro-091918-104501
- Coti Zelati, F., Rea, N., Pons, J. A., Campana, S., & Esposito, P. 2018, MNRAS, 474, 961, doi: 10.1093/mnras/stx2679
- Dai, Z. G. 2020, ApJL, 897, L40, doi: 10.3847/2041-8213/aba11b
- Dong, F. A., & Chime/Frb Collaboration. 2022, The Astronomer's Telegram, 15681, 1
- Edwards, R. T., Hobbs, G. B., & Manchester, R. N. 2006, MNRAS, 372, 1549,

doi: 10.1111/j.1365-2966.2006.10870.x

- Enoto, T., Hu, C.-P., Guver, T., et al. 2022, The Astronomer's Telegram, 15690, 1
- Frederiks, D., Ridnaia, A., Svinkin, D., et al. 2022, The Astronomer's Telegram, 15686, 1

- Fu, Y.-C., Song, L. M., Ding, G. Q., et al. 2023, MNRAS, 521, 893, doi: 10.1093/mnras/stad614
- Gavriil, F. P., Kaspi, V. M., & Woods, P. M. 2004, ApJ, 607, 959, doi: 10.1086/383564
- Ge, M. Y., Lu, F. J., Qu, J. L., et al. 2012, ApJS, 199, 32, doi: 10.1088/0067-0049/199/2/32
- Ge, M. Y., Lu, F. J., Yan, L. L., et al. 2019, Nature Astronomy, 3, 1122, doi: 10.1038/s41550-019-0853-5
- Ge, M. Y., Liu, C. Z., Zhang, S. N., et al. 2023, ApJ, 953, 67, doi: 10.3847/1538-4357/acda1d
- Ge, M.-Y., Yang, Y.-P., Lu, F.-J., et al. 2024, Research in Astronomy and Astrophysics, 24, 015016, doi: 10.1088/1674-4527/ad0f0c
- Gendreau, K., & Arzoumanian, Z. 2017, Nature Astronomy, 1, 895, doi: 10.1038/s41550-017-0301-3
- Gendreau, K. C., Arzoumanian, Z., Adkins, P. W., et al. 2016, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9905, Space Telescopes and Instrumentation 2016: Ultraviolet to Gamma Ray, ed. J.-W. A. den Herder, T. Takahashi, & M. Bautz, 99051H, doi: 10.1117/12.2231304
- Giri, U., Andersen, B. C., Chawla, P., et al. 2023, arXiv e-prints, arXiv:2310.16932, doi: 10.48550/arXiv.2310.16932
- Hu, C.-P., Narita, T., Enoto, T., et al. 2024, Nature, 626, 500, doi: 10.1038/s41586-023-07012-5
- Huang, Y. X., Xu, H., Xu, Y. H., et al. 2022, The Astronomer's Telegram, 15707, 1
- Huppenkothen, D., Bachetti, M., Stevens, A. L., et al. 2019, ApJ, 881, 39, doi: 10.3847/1538-4357/ab258d
- Ibrahim, A. Y., Borghese, A., Coti Zelati, F., et al. 2024, ApJ, 965, 87, doi: 10.3847/1538-4357/ad293b
- Israel, G. L., Romano, P., Mangano, V., et al. 2008, ApJ, 685, 1114, doi: 10.1086/590486
- Israel, G. L., Esposito, P., Rea, N., et al. 2010, MNRAS, 408, 1387, doi: 10.1111/j.1365-2966.2010.17001.x
- —. 2016, MNRAS, 457, 3448, doi: 10.1093/mnras/stw008
- Kaspi, V. M., & Beloborodov, A. M. 2017, ARA&A, 55, 261, doi: 10.1146/annurev-astro-081915-023329
- Kaspi, V. M., Gavriil, F. P., Woods, P. M., et al. 2003, ApJL, 588, L93, doi: 10.1086/375683
- Kouveliotou, C., Dieters, S., Strohmayer, T., et al. 1998, Nature, 393, 235, doi: 10.1038/30410
- Leahy, D. A. 1987, A&A, 180, 275
- Li, C. K., Cai, C., Xiong, S. L., et al. 2022a, The Astronomer's Telegram, 15698, 1
- Li, C. K., Lin, L., Xiong, S. L., et al. 2021, Nature Astronomy, 5, 378, doi: 10.1038/s41550-021-01302-6
- Li, X. B., Zhang, S. N., Xiong, S. L., et al. 2022b, The Astronomer's Telegram, 15708, 1

- —. 2022c, The Astronomer's Telegram, 15714, 1
- Lin, L., Göğüş, E., Roberts, O. J., et al. 2020a, ApJL, 902, L43, doi: 10.3847/2041-8213/abbefe
- ---. 2020b, ApJ, 893, 156, doi: 10.3847/1538-4357/ab818f
- Lin, L., Kouveliotou, C., Göğüş, E., et al. 2011, ApJL, 740, L16, doi: 10.1088/2041-8205/740/1/L16
- Lin, L., Zhang, C. F., Wang, P., et al. 2020c, Nature, 587, 63, doi: 10.1038/s41586-020-2839-y
- Lorimer, D. R., Bailes, M., McLaughlin, M. A., Narkevic, D. J., & Crawford, F. 2007, Science, 318, 777, doi: 10.1126/science.1147532
- Lu, W., Kumar, P., & Zhang, B. 2020, MNRAS, 498, 1397, doi: 10.1093/mnras/staa2450
- Lu, X.-F., Song, L.-M., Ge, M.-Y., et al. 2024, Research in Astronomy and Astrophysics, 24, 065018, doi: 10.1088/1674-4527/ad4599
- Maan, Y., Leeuwen, J. v., Straal, S., & Pastor-Marazuela, I. 2022, The Astronomer's Telegram, 15697, 1
- Mereghetti, S., Götz, D., Weidenspointner, G., et al. 2009, ApJL, 696, L74, doi: 10.1088/0004-637X/696/1/L74
- Morrison, R., & McCammon, D. 1983, ApJ, 270, 119, doi: 10.1086/161102
- Nobili, L., Turolla, R., & Zane, S. 2008, MNRAS, 386, 1527, doi: 10.1111/j.1365-2966.2008.13125.x
- Olausen, S. A., & Kaspi, V. M. 2014, ApJS, 212, 6, doi: 10.1088/0067-0049/212/1/6
- Özel, F., & Güver, T. 2007, ApJL, 659, L141, doi: 10.1086/517933
- Palm, D. M. 2022, The Astronomer's Telegram, 15667, 1
- Pearlman, A. B., & Chime/Frb Collaboration. 2022, The Astronomer's Telegram, 15792, 1
- Petroff, E., Hessels, J. W. T., & Lorimer, D. R. 2019, A&A Rv, 27, 4, doi: 10.1007/s00159-019-0116-6
- Ridnaia, A., Svinkin, D., Frederiks, D., et al. 2021, Nature Astronomy, 5, 372, doi: 10.1038/s41550-020-01265-0
- Roberts, O. J., Dalessi, S., & Malacaria, C. 2022, The Astronomer's Telegram, 15672, 1
- Scholz, P., & Kaspi, V. M. 2011, ApJ, 739, 94, doi: 10.1088/0004-637X/739/2/94
- Shao, Y.-X., Zhou, P., Li, X.-D., et al. 2024, arXiv e-prints, arXiv:2410.00635, doi: 10.48550/arXiv.2410.00635
- Stamatikos, M., Malesani, D., Page, K. L., & Sakamoto, T. 2014, GRB Coordinates Network, 16520, 1
- Thompson, C., Lyutikov, M., & Kulkarni, S. R. 2002, ApJ, 574, 332, doi: 10.1086/340586
- Tuo, Y. L., Ji, L., Tsygankov, S. S., et al. 2020, Journal of High Energy Astrophysics, 27, 38, doi: 10.1016/j.jheap.2020.03.003
- Wang, C. W., Xiong, S. L., Zhang, Y. Q., et al. 2022, The Astronomer's Telegram, 15682, 1

- 16
- Woods, P. M., Kouveliotou, C., Finger, M. H., et al. 2007, ApJ, 654, 470, doi: 10.1086/507459
- Woods, P. M., Kaspi, V. M., Thompson, C., et al. 2004, ApJ, 605, 378, doi: 10.1086/382233
- Xiao, D., Wang, F., & Dai, Z. 2021, Science China Physics, Mechanics, and Astronomy, 64, 249501, doi: 10.1007/s11433-020-1661-7
- Yang, Y.-P., & Zhang, B. 2018, ApJ, 868, 31, doi: 10.3847/1538-4357/aae685
- Younes, G., Baring, M. G., Kouveliotou, C., et al. 2017a, ApJ, 851, 17, doi: 10.3847/1538-4357/aa96fd

- Younes, G., Kouveliotou, C., Jaodand, A., et al. 2017b, ApJ, 847, 85, doi: 10.3847/1538-4357/aa899a
- Younes, G., Güver, T., Kouveliotou, C., et al. 2020, ApJL, 904, L21, doi: 10.3847/2041-8213/abc94c
- Younes, G., Enoto, T., Hu, C.-P., et al. 2022, The Astronomer's Telegram, 15674, 1
- Younes, G., Baring, M. G., Harding, A. K., et al. 2023, Nature Astronomy, 7, 339, doi: 10.1038/s41550-022-01865-y
- Zhang, B. 2017, ApJL, 836, L32, doi: 10.3847/2041-8213/aa5ded
- -. 2020, Nature, 587, 45, doi: 10.1038/s41586-020-2828-1
- Zhou, P., Zhou, X., Chen, Y., et al. 2020, ApJ, 905, 99, doi: 10.3847/1538-4357/abc34a