Dualities in random matrix theory

Peter J. Forrester

School of Mathematics and Statistics, The University of Melbourne, Victoria 3010, Australia. Email: pjforr@unimelb.edu.au;

Abstract

Duality identities in random matrix theory for products and powers of characteristic polynomials, and for moments, are reviewed. The structure of a typical duality identity for the average of a positive integer power k of the characteristic polynomial for particular ensemble of $N \times N$ matrices is that it is expressed as the average of the power N of the characteristic polynomial of some other ensemble of random matrices, now of size $k \times k$. With only a few exceptions, such dualities involve (the β generalised) classical Gaussian, Laguerre and Jacobi ensembles Hermitian ensembles, the circular Jacobi ensemble, or the various non-Hermitian ensembles relating to Ginibre random matrices. In the case of unitary symmetry in the Hermitian case, they can be studied using the determinantal structure. The β generalised case requires the use of Jack polynomial theory, and in particular Jack polynomial based hypergeometric functions. Applications to the computation of the scaling limit of various β ensemble correlation and distribution functions are also reviewed. The non-Hermitian case relies on the particular cases of Jack polynomials corresponding to zonal polynomials, and their integration properties when their arguments are eigenvalues of certain matrices. The main tool to study dualities for moments of the spectral density, and generalisations, is the loop equations.

[1 Introduction](#page-1-0) 2

Contents

1 Introduction

As a prelude to reviewing duality relations in random matrix theory, a few remarks about dualities more generally are in order. In the essay "Duality in mathematics and physics", Atiyah [\[5\]](#page-47-0) posits "Fundamentally, duality gives two different points of view of looking at the same object". In internet forums, an obvious key word search reveals the descriptions "In laymen terms, it's when you consider an opposite concept in such a way that some properties have 'flipped' analogous properties"; "The common idea is that there are two things which basically are just two sides of the same coin"; and "When applied to specific examples, there is usually a precise definition for just that context", among others.

Indeed dualities in random matrix theory involve 'flipping'. The flipping may be of two parameters (e.g. the order of a moment, and the size of the matrix), or of two ensembles (e.g. orthogonal symmetry, and symplectic symmetry), these being the primary classes of dualities to be discussed below. It is furthermore the case that these contexts offer a precise mathematical definition. Introductory examples can be given.

In relation to a duality relation involving flipping the order of a moment and the size of the matrix, one can consider the first moment (i.e. average value) of the characteristic polynomial $\det(x\mathbb{I}_N - H)$ for the matrix H an Hermitian Wigner matrix. The construction of the latter is according to $H=\frac{1}{2}$ $\frac{1}{2}(X+X^{\dagger})$, where the $N \times N$ random matrix X has all elements independently distributed with zero mean and variance σ^2 (for complex elements with zero mean, $\sigma^2 := \langle |x_{ij}|^2 \rangle$). With the ensemble of such matrices denoted by \mathcal{H}_N , one has [\[64,](#page-49-0) equivalent to Prop. 11]

$$
\langle \det(x\mathbb{I}_N - H) \rangle_{H \in \mathcal{H}_N} = i^{-N} \langle (ix - h)^N \rangle_{h \in \mathcal{N}(0, \sigma^2/2)},\tag{1.1}
$$

where $\mathcal{N}(0, \sigma^2/2)$ denotes the zero mean normal distribution with variance $\sigma^2/2$. Comparing both sides, one sees the interchange (flipping) of the pair (N, n) , where on the left hand side N is the matrix size, and $n = 1$ is the power of the characteristic polynomial, while on the right hand side the meaning of these parameters is reversed.

We again turn to moments for our introductory example illustrating a random matrix duality flipping two ensembles. The moments of interest here are those associated with the spectral density for finite N , which itself after normalisation can be considered as a probability density function. As an ensemble average, this is equivalent to considering the expected value of the power sum $\sum_{j=1}^{N} \lambda_j^k$ $(k = 1, 2, ...)$, where $\{\lambda_j\}$ are the eigenvalues. For a given k, we will do this for the two distinct ensembles of the Gaussian orthogonal ensemble (GOE) of real symmetric matrices, and the Gaussian symplectic ensemble (GSE) of quaternion self dual Hermitian matrices. Only the even moments are non-zero, with the $2k$ -th moment in fact a polynomial in N of degree $k+1$, which furthermore vanishes at $N=0$. Before listing some of these polynomials, we recall that matrices from the GOE and GSE are specific classes of Hermitian Wigner matrices, where the elements of the matrix X introduced above are zero mean Gaussians. For the GOE, these elements are all real. For the GSE, the elements are themselves 2×2 complex matrices representing quaternions (see e.g. $[54, \{1.3.2\}]$), and the eigenvalues here are two fold degenerate. The probability measure on the space of real symmetric matrices for the GOE, being proportional to $e^{-\text{Tr }H^2/2}$, is invariant under the similarity transformation, real symmetric matrices automorphism, $H \mapsto O^T H O$ for any real orthogonal matrix O of the same size as H. This is the reason for the 'orthogonal' in the naming of the ensemble. Similar considerations apply to the naming of the GSE, where the relevant similarity transformation, quaternion self dual Hermitian matrices automorphism, is $H \mapsto S^{\dagger} H S$, where S is unitary symplectic matrix.

A classical result in random matrix theory is that the joint eigenvalue probability density function (PDF) of the GOE and GSE is proportional to (see e.g. [\[54,](#page-49-1) Prop. 1.3.4])

$$
\prod_{l=1}^{N} e^{-\beta \lambda_l^2/2} \prod_{1 \le j < k \le N} |\lambda_k - \lambda_j|^\beta,\tag{1.2}
$$

for $\beta = 1$ (GOE) and $\beta = 4$ (GSE). The parameter β is referred to as the Dyson index. The most efficient computational scheme for the moments ${m_{2k}}$ is via the use of certain fourth order linear difference equations found by Ledoux [\[112\]](#page-51-0), which furthermore only require knowledge of the initial conditions $m_0 = N$ and $m_2^{\text{GOE}} = N^2 + N$, $m_2^{\text{GSE}} = N^2 - N/2$ for their unique solution. For the particular low order cases $k = 3, 4$, one then finds

$$
m_6^{\text{GOE}} = 5N^4 + 22N^3 + 52N^2 + 41N, \ m_8^{\text{GOE}} = 14N^5 + 93N^4 + 374N^3 + 690N^2 + 509N \tag{1.3}
$$

$$
8m_6^{\text{GSE}} = 40N^4 - 88N^3 + 104N^2 - 41N, \ 16m_8^{\text{GSE}} = 224N^5 - 744N^4 + 1496N^3 - 1380N^2 + 509N. \tag{1.4}
$$

Inspection reveals the functional (duality) relation

$$
m_{2k}^{\text{GOE}} = (-1)^{k+1} 2^{k+1} m_{2k}^{\text{GSE}} \Big|_{N \mapsto -N/2}, \quad (k = 3, 4)
$$
 (1.5)

which is in fact valid for each $k \in \mathbb{Z}_{\geq 0}$ ([\[123,](#page-52-0) Eq. (5.3)], special case of [\[40,](#page-48-0) Th. 2.8]). Hence there is a flipping of ensembles on different sides of this equation, as well as a particular algebraic

mapping of N (we say algebraic, as there is no matrix construction once N is so mapped). We remark that the GOE and GSE can be linked in other ways. For example, integrating over all the odd labelled eigenvalues in the GOE of size $2N + 1$ gives the GSE of size N; for a recent review on results in random matrix theory of this sought, see [\[61\]](#page-49-2).

The class of dualities involving characteristic polynomials will be considered first. As this class is large, the subject matter under this heading is to be further subdivided. This we do by having separate sections for when the dualities involve classical random matrix ensembles with $\beta = 2$ (Section [2\)](#page-3-0), then the general β case (Section [3\)](#page-10-0). Consequences for the large N limit are considered in Section [4.](#page-20-0) Dualities relating to characteristic polynomials for non-Hermitian ensembles are the topic of Section [5.](#page-31-0) Section [6](#page-43-0) gives an account of dualities relating to (mixed) moments for a large class of β ensembles.

Not considered in this review (due to the present author's lack of expertise) are random matrix group integral dualities coming from the field theory method of the colour-flavour transformation [\[147,](#page-53-0) [86\]](#page-50-0), or from the related mathematical theory of Howe pairs [\[32,](#page-48-1) [148\]](#page-53-1).

2 Characteristic polynomial dualities for classical random matrix ensembles with $\beta = 2$

2.1 Invariant ensemble definitions

Consider the generalisation of [\(1.2\)](#page-2-0), defined as the eigenvalue PDF on the real line proportional to

$$
\prod_{l=1}^{N} w(\lambda_l) \prod_{1 \le j < k \le N} |\lambda_k - \lambda_j|^\beta. \tag{2.1}
$$

Viewed as relating to an ensemble of Hermitian random matrices, (2.1) is denoted $ME_{\beta,N}[w]$. The function $w(x)$ is referred to as the weight. As is consistent with the situation for [\(1.2\)](#page-2-0), which corresponds to choosing $w(\lambda) = e^{-\beta \lambda^2/2}$, the cases $\beta = 1, 4$ relate to orthogonal (symplectic) invariant ensembles of real (quaternion) valued Hermitian random matrices. Similarly complex valued Hermitian random matrices permit the automorphism $H \mapsto U^{\dagger} H U$ for U a member of the unitary group. If the joint element PDF is invariant under this mapping, an eigenvalue PDF of the form [\(2.1\)](#page-3-2) results with $\beta = 2$ (specifically, the element PDF reduces to the functional form $\prod_{l=1}^{N} w(\lambda_l)$, while $\prod_{1 \leq j < k \leq N} |\lambda_k - \lambda_j|^\beta$ is identified as the eigenvalue factor of the Jacobian in the change of variables from the matrix elements to the eigenvalues and eigenvectors; see [\[54,](#page-49-1) Prop. 1.3.4]).

It turns out that the dualities of certain classes of Hermitian random matrices to be considered will also involve unitary random matrices. The analogue of (2.1) in this setting is

$$
\prod_{l=1}^{N} w(\theta_l) \prod_{1 \le j < k \le N} |e^{2\pi i \theta_k} - e^{2\pi i \theta_j}|^{\beta}, \quad -1/2 < \theta_l \le 1/2,\tag{2.2}
$$

to be denoted $CE_{\beta,N}[w]$ (the "C" here stands for circular). The naming circular β ensemble is used for the case $w = 1$, while the naming circular orthogonal, circular unitary, and circular symplectic ensemble (notation COE, CUE and CSE respectively) is used for the cases $\beta = 1, 2$ and 4 of the circular β ensemble.

A variation of [\(2.1\)](#page-3-2) is the eigenvalue PDF proportional to

$$
\prod_{l=1}^{N} w(\lambda_l) \prod_{1 \le j < k \le N} |\lambda_k^2 - \lambda_j^2|^{\beta}.\tag{2.3}
$$

Typically this results from matrices with the property that the eigenvalues come in \pm pairs, which is termed a chiral symmetry. Thus the naming $\text{chME}_{\beta,N}(w)$. An example of matrices with a chiral symmetry is the structure

$$
Y = \begin{bmatrix} 0_{n \times n} & Z \\ Z^{\dagger} & 0_{N \times N} \end{bmatrix}
$$
 (2.4)

for Z an $n \times N$ ($n \geq N$) matrix — in general there are N eigenvalues in \pm pairs, and $n - N$ zero eigenvalues. In the case that Z has independent standard complex entries, and thus being a so called (complex) rectangular Ginibre matrix [\[27\]](#page-48-2), the positive eigenvalues of Y have PDF chME_{2,N}(w), $w(x) = x^{2(n-N)+1}e^{-x^2}$; see e.g. [\[54,](#page-49-1) Prop. 3.1.3].

2.2 Products of characteristic polynomials and the GUE

In light of [\(1.1\)](#page-2-1), an immediate question to ask is if there is an analogous duality for

$$
\left\langle \prod_{l=1}^{p} \det(x_l \mathbb{I}_N - H) \right\rangle_{\mathcal{H}_N}
$$
\n(2.5)

for $p > 1$. Here we may consider the cases of \mathcal{H}_N a class of Wigner matrices, or an invariant ensemble characterised by the eigenvalue PDF (2.1) (note that choosing the matrix X — recall the text above (1.1) — to have independent standard Gaussian elements, which are real, complex or quaternion for $\beta = 1, 2$ and 4 respectively gives rise to random matrix ensembles, the GOE. GUE, GSE, which are both of the Wigner and invariant class). The simplest case to investigate is for \mathcal{H}_N an invariant ensemble with complex elements, and thus corresponding to $ME_{2,N}(w)$. Introduce the family of monic orthogonal polynomials $\{p_j(x)\}_{j=0,1,\ldots}$ by the requirement that $\int_I w(x)p_j(x)p_k(x) dx = h_j \delta_{j,k}$ (here I is the support of $w(x)$ and $h_j > 0$ is the normalisation). Then it is a standard result [\[21\]](#page-47-1), [\[90\]](#page-50-1), [\[7\]](#page-47-2), [\[54\]](#page-49-1) that

$$
\left\langle \prod_{l=1}^{p} \det(x_{l} \mathbb{I}_{N} - H) \right\rangle_{\text{ME}_{2,N}[w]} = \frac{1}{\prod_{1 \leq j < k \leq p} (x_{k} - x_{j})} \det[p_{N+j-1}(x_{k})]_{j,k=1,\dots,p}.\tag{2.6}
$$

In the case $p = 1$ [\(2.6\)](#page-4-1) gives

$$
\left\langle \det(x\mathbb{I}_N - H) \right\rangle_{\text{ME}_{2,N}[w]} = p_N(x),\tag{2.7}
$$

which in fact is a classical formula in orthogonal polynomial theory due to Heine [\[138\]](#page-52-1). In the case $w(x) = e^{-x^2}$ corresponding to the GUE, one has $p_N(x) = 2^{-N} H_N(x)$, where $H_N(x)$ denotes the Hermite polynomial of degree N . Use of the integral form of the Hermite polynomials

$$
H_N(x) = \frac{(2i)^N}{\sqrt{\pi}} e^{x^2} \int_{-\infty}^{\infty} e^{-t^2} t^N e^{-2ixt} dt
$$
 (2.8)

gives agreement with [\(1.1\)](#page-2-1).

We would like to identify the RHS of (2.6) with a random matrix average over an ensemble of $p \times p$ matrices for general $p \geq 1$. It turns out that this is possible for both the GUE and LUE, although in the latter case the ensemble that appears is hard to anticipate (see [§3.5\)](#page-14-1). For the GUE weight $w(x) = e^{-x^2}$, the RHS consists of another GUE average.

Proposition 2.1. ([\[22\]](#page-47-3)) Let $X = diag(x_1, \ldots, x_p)$. We have

$$
\left\langle \prod_{l=1}^{p} \det(x_{l} \mathbb{I}_{N} - H) \right\rangle_{\text{GUE}_{N}} = i^{-pN} \left\langle \det(iX - H)^{N} \right\rangle_{\text{GUE}_{p}}.
$$
\n(2.9)

In particular

$$
\left\langle \det(x\mathbb{I}_N - H)^p \right\rangle_{\text{GUE}_N} = i^{-p} \left\langle \det(ix\mathbb{I}_p - H)^N \right\rangle_{\text{GUE}_p}.\tag{2.10}
$$

Proof. We know that for the Gaussian weight, $p_k(x) = 2^{-k}H_k(x)$. The task is to show that the RHS of [\(2.9\)](#page-5-0) evaluates to the determinant formula in [\(2.6\)](#page-4-1) with this substitution. We begin by noting that, up to proportionality, the average on the RHS can be rewritten as

$$
\int_{H \in \mathcal{H}_p} (\det H)^N e^{-\text{Tr}\,(H + iX)^2} \,(dH),\tag{2.11}
$$

where \mathcal{H}_p is the space of $p \times p$ Hermitian matrices and (dH) denotes the product of all the independent differentials (real and imaginary parts) of H. Changing variables according to the diagonalisation formula $H = U\Lambda U^{\dagger}$, where $\Lambda = \text{diag}(\lambda_1, \ldots, \lambda_p)$, shows that (2.11) is proportional to

$$
e^{\sum_{j=1}^{p}x_j^2} \left\langle \prod_{l=1}^{p} \lambda_l^N \int e^{-2i \text{Tr } U^\dagger \Lambda U X} d^{\text{H}} U \right\rangle_{\Lambda \in \text{GUE}_p}, \tag{2.12}
$$

where $d^H U$ denotes the (normalised) Haar measure on the unitary group $U(p)$. This group integral is evaluated according to the well known HCIZ formula — see e.g. [\[54,](#page-49-1) Prop. 11.6.1] and [\[125\]](#page-52-2) for recent developments — showing that [\(2.12\)](#page-5-2) is proportional to

$$
e^{\sum_{j=1}^p x_j^2} \int_{-\infty}^{\infty} d\lambda_1 \cdots \int_{-\infty}^{\infty} d\lambda_p \prod_{l=1}^p \lambda_l^N e^{-\sum_{j=1}^p \lambda_j^2} \prod_{1 \le j < k \le p} (\lambda_k - \lambda_j) \det[e^{-2i\lambda_j x_k}]_{j,k=1,\dots,p}.
$$
 (2.13)

Application now of Andréief's integration formula [\[57\]](#page-49-3) reduces this to

$$
e^{\sum_{j=1}^{p}x_j^2} \det \left[\int_{-\infty}^{\infty} \lambda^{N+j-1} e^{-\lambda^2} e^{-2i\lambda x_k} d\lambda \right]_{j,k=1,\dots,p}.
$$
 (2.14)

Using now the integral form for the Hermite polynomials [\(2.8\)](#page-4-2) allows [\(2.14\)](#page-5-3) to be recognised as being proportional to the determinant formula in [\(2.6\)](#page-4-1) in the GUE case. The proportionality is fixed by noting that both sides of [\(2.9\)](#page-5-0) reduce to $\prod_{l=1}^{p} x_l^N$ is the limit that each $x_l \to \infty$.

The duality [\(2.10\)](#page-5-4) follows from [\(2.9\)](#page-5-0) by setting all the x_l equal. However, for reference in settings where an analogue of [\(2.9\)](#page-5-0) is not available, we outline a direct strategy beginning with [\(2.6\)](#page-4-1). This is to first set $x_1 = x$ and then subtract column 1 from column two, divide by the existing prefactor $(x_2 - x)$ and take the limit $x_2 \rightarrow x$. Next subtract columns 1 and $(x_3 - x)$ times column 2 from column 3, divide by the existing prefactor $(x_3 - x)^2$ and take the limit $x_3 \rightarrow x$. By repeating this procedure in succession, it is possible to subtract from column k the first $(k-1)$ terms in its Taylor series expansion about x, which after dividing by $(x_k - x)^{k-1}$ and taking the limit $x_k \to x$ gives $D_x^{k-1} p_{N+j-1}(x)/(k-1)!$ (here the D_x denotes derivative operation) as the entry in row j and column k. This shows

$$
\left\langle \det(x\mathbb{I}_N - H)^p \right\rangle_{\text{ME}_{2,N}[w]} = \frac{1}{\prod_{l=1}^p l!} \det[D_x^{k-1} p_{N+j-1}(x)]_{j,k=1,\dots,p}.
$$
 (2.15)

Repeated use of the Hermite polynomial identity $H'_n(x) = 2xH_n(x) - H_{n+1}(x)$ then allows us to deduce that in the Gaussian case the RHS of [\(2.15\)](#page-5-5) is proportional to the Hankel determinant

$$
\det[H_{N+j+k-2}(x)]_{j,k=1,\dots,p}.\tag{2.16}
$$

Next, a well known formula in random matrix theory [\[54,](#page-49-1) Eq. (5.75)], due to Heine in the context of the study of orthogonal polynomials [\[138\]](#page-52-1), gives that in general for $\alpha_j := \int_I a(t) t^j dt$, one has

$$
\det[\alpha_{j+k}]_{j,k=0,\dots,n-1} = \frac{1}{n!} \int_I dx_1 \cdots \int_I dx_n \prod_{l=1}^n a(x_l) \prod_{1 \le j < k \le n} (x_k - x_j)^2. \tag{2.17}
$$

Using the integral form for the Hermite polynomials (2.8) allows for (2.16) to be rewritten in the form of the RHS of [\(2.17\)](#page-6-2), which after minor manipulation can be recognised as the RHS of $(2.10).$ $(2.10).$ \Box

If we consider now [\(2.5\)](#page-4-3) for \mathcal{H}_N the class of Wigner matrices beyond the GUE, and restrict to the case $p = 2$, a duality formula analogous to (2.9) is known from [\[136,](#page-52-3) Eq. (2.16)) after scaling].

Proposition 2.2. Consider a complex Hermitian Wigner matrix $\tilde{H} = [\tilde{H}_{jk}]_{j,k=1,\dots,N}$. Require that the real and imaginary parts for $j < k$ are independently drawn from the same distribution, which has first and third moments zero, second moment $\frac{1}{2}$, fourth moment b and corresponding cumulant $\kappa_4 = b - \frac{1}{4}$ $\frac{1}{4}$. Let \tilde{H}_{jj} also be independently drawn from this distribution, together with the scaling of multiplying by $\sqrt{2}$. Further scale all the entries by considering $H = \frac{1}{\sqrt{2}}$ $\frac{1}{2}\tilde{H}$ to define the ensemble \mathcal{H}_N . With $X = \text{diag}(x_1, x_2)$ one has

$$
\left\langle \det(x_1 \mathbb{I}_N - H) \det(x_2 \mathbb{I}_N - H) \right\rangle_{\mathcal{H}_N}
$$

= $i^{-2N} \left\langle \left(\det(iX - Q) + t\epsilon(\kappa_4) \right)^N \right\rangle_{Q \in \text{GUE}_2, t \in \mathcal{N}(0, 1/(2|\kappa_4|))}, (2.18)$

where $\epsilon(x) = x$, $(x > 0)$ and $\epsilon(x) = -ix$, $(x < 0)$.

Proof. (Comment only.) The proof is based on Grassmann integration involving formal antisymmetric variables $\{\psi_j\}, \{\bar{\psi}_j\}$. Each determinant on the LHS can be rewritten according to the key formula

$$
\int \exp\Big(\sum_{j,k=1}^N A_{j,k}\bar{\psi}_j\psi_k\Big) \prod_{j=1}^N d\bar{\psi}_j d\psi_j = \det A;\tag{2.19}
$$

see e.g. [\[95\]](#page-51-1). An instructive preliminary exercise is to make use of this identity and the calculus of Grassmann integration to give a derivation of [\(1.1\)](#page-2-1). 口

2.3 Averages of reciprocals of characteristic polynomials for the GUE

A companion to the determinant identity [\(2.6\)](#page-4-1) applying to the average of a product of characteristic polynomials in a general $\beta = 2$ matrix ensemble ME_{2,N} [w] relates to a product of reciprocals of characteristic polynomials [\[90\]](#page-50-1), [\[7\]](#page-47-2), [\[54,](#page-49-1) Prop. 5.3.1],

$$
\left\langle \frac{1}{\prod_{l=1}^{q} \det(v_j - X)} \right\rangle_{\text{ME}_{2,N}[w]} = \frac{(-1)^{q(q-1)/2}}{\prod_{j=N-q}^{N-1} h_j} \det \left[\int_{-\infty}^{\infty} \frac{p_{N-q+k-1}(x)}{v_j - x} w(x) \, dx \right]_{j,k=1,\dots,q} . \tag{2.20}
$$

Set $q = 1$ and consider the case of the Gaussian weight, for which the $p_n(x)$ are proportional to the Hermite polynomials. Since the support of the Gaussian weight is the whole real line, it is required that the v_j have a nonzero imaginary part. Using the contour integral form (equivalent to Rodrigues formula)

$$
e^{-x^2}H_{N-1}(x) = \frac{(N-1)!}{2\pi i} \int_{I_+ + I_-} \frac{e^{-z^2}}{(x-z)^N} dz,
$$

cf. [\(2.8\)](#page-4-2), where for x real I_{+} (I_{-}) can be taken as the real line shifted up (down) by $i\epsilon$ ($\epsilon > 0$) and traversed in the negative (positive) sense. Substituting in the integral on the RHS of [\(2.20\)](#page-6-3), and taking the z integral outside of the x integral allows the former to be computed, with the result

$$
\left\langle \frac{1}{\det(v - X)} \right\rangle_{\text{ME}_{2, N}[e^{-x^{2}}]} = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \frac{e^{-z^{2}}}{(v - z)^{N}} dz = \left\langle \frac{1}{(v - z)^{N}} \right\rangle_{z \in \mathcal{N}(0, 1/\sqrt{2})},
$$
(2.21)

which itself is a companion duality to the GUE case of (1.1) .

Extending [\(2.21\)](#page-7-1), Proposition [2.1](#page-5-6) has a counterpart for averaged reciprocal characteristic polynomials with respect to the GUE.

Proposition 2.3. ([\[35\]](#page-48-3)) Let $X = \text{diag}(x_1, \ldots, x_p)$, where each x_i has a nonzero imaginary part. We have

$$
\left\langle \prod_{l=1}^{p} \frac{1}{\det(x_l \mathbb{I}_N - H)} \right\rangle_{\text{GUE}_N} = \left\langle \frac{1}{\det(X - H)^N} \right\rangle_{\text{GUE}_p}.
$$
\n(2.22)

In particular

$$
\left\langle \frac{1}{\det(x\mathbb{I}_N - H)^p} \right\rangle_{\text{GUE}_N} = \left\langle \frac{1}{\det(x\mathbb{I}_p - H)^N} \right\rangle_{\text{GUE}_p}.
$$
\n(2.23)

We will see in Section [3.5](#page-14-1) that these results are special cases of dualities applying to the GUE generalised to include a source matrix.

2.4 Powers of characteristic polynomials and the LUE and JUE

There are no known (literal) analogues of [\(2.9\)](#page-5-0) beyond the Gaussian case. However progress can be made if one seeks analogues of [\(2.10\)](#page-5-4) only. This is possible for the case of the unitary invariant ensemble with weight $w(x) = x^a e^{-x} \mathbb{1}_{x>0}$, referred to as the Laguerre unitary ensemble and denoted LUE (or $LUE_{N,a}$ when the size of the matrix N and the Laguerre parameter a are to be emphasised), or the weight $w(x) = x^{a_1}(1-x)^{a_2} \mathbb{1}_{x>0}$, referred to as the Jacobi unitary ensemble JUE (or JUE_{N,(a_{1,a2)}). Whereas the GUE case of the determinant in [\(2.15\)](#page-5-5) was shown} to give rise to a Hankel determinant, it turns out that both the LUE and JUE cases give rise to Toeplitz determinants.

Proposition 2.4. ([\[65,](#page-49-4) for the LUE case with $\alpha = 2$]) We have

$$
\det[D_x^{k-1} L_{N+j-1}^{(a)}(x)]_{j,k=1,\dots,p} = (-1)^{p(p-1)/2} \det[L_{N+j-k}^{(a+k-j)}(x)]_{j,k=1,\dots,p},
$$

\n
$$
\det[D_x^{k-1} P_{N+j-1}^{(a_1,a_2)}(x)]_{j,k=1,\dots,p} \propto \det[P_{N+j-k}^{(a_1+k-j,a_2+k-j)}(x)]_{j,k=1,\dots,p}.
$$
\n(2.24)

Proof. Consider for definiteness the LUE case. The relevant orthogonal polynomials (up to the normalisation condition of being monic) are the Laguerre polynomials $\{L_n^{(a)}(x)\}_{n=0,1,...}$. The

determinant in [\(2.15\)](#page-5-5) is to be manipulated by the elementary row operation of replacing the row $p - j$ by row $p - j$ minus row $p - j + 1$ for $j = 0, \ldots, p - 2$ in order, then simplifying using the Laguerre polynomial identity $L_n^{(a)}(x) = L_n^{(a+1)}(x) - L_{n-1}^{(a+1)}(x)$ $\binom{(a+1)}{n-1}(x)$. We repeat this same procedure, finishing one further row down from the top at each iteration, a total of $p-2$ further times to conclude

$$
\det[D_x^{k-1}L_{N+j-1}^{(a)}(x)]_{j,k=1,\dots,p} = \det[D_x^{k-1}L_{N+j-1}^{(a+1-j)}(x)]_{j,k=1,\dots,p}.
$$

Use of the Laguerre polynomial formula $D_x L_p^{(a)}(x) = -L_{p-1}^{(a+1)}$ $p-1 \choose p-1}$ (x) a total of j times in row j allows for a substitution of the matrix element on the RHS of this to give the RHS of [\(2.24\)](#page-7-2). \Box

The identities in Proposition [2.4](#page-7-3) provide duality formulas for the average in [\(2.15\)](#page-5-5) in the cases of the LUE and JUE. However, unlike the case of the GUE, these dualities no longer relate back to the same ensembles, but rather to a class of $\beta = 2$ circular ensembles [\(2.2\)](#page-3-3). The reason

for this is that with $\beta_j = \int_{-1/2}^{1/2}$ $\int_{-1/2}^{1/2} b(e^{2\pi i\theta})e^{2\pi i\theta(j-k)} d\theta$ one has the general identity [\[138\]](#page-52-1)

$$
\det[\beta_{j-k}]_{j,k=1,\dots,n} = \frac{1}{n!} \int_{-1/2}^{1/2} d\theta_1 \cdots \int_{-1/2}^{1/2} d\theta_n \prod_{l=1}^n b(e^{2\pi i \theta_l}) \prod_{1 \le j < k \le n} |e^{2\pi i \theta_k} - e^{2\pi i \theta_j}|^2; \tag{2.25}
$$

cf. [\(2.17\)](#page-6-2).

Corollary 2.1. We have [\[51\]](#page-49-5)

$$
\left\langle \det(x\mathbb{I}_N - H)^p \right\rangle_{\text{LUE}_{N,a}} \propto \left\langle \prod_{l=1}^p e^{-xe^{2\pi i\theta_l}} \right\rangle_{\text{CE}_{2,p}[e^{\pi i(a-N)\theta}|1+e^{2\pi i\theta}|^{a+N}]}
$$
(2.26)

and

$$
\left\langle \det(x\mathbb{I}_N - H)^p \right\rangle_{\text{JUE}_{N,(a_1,a_2)}} \propto (1-x)^{pN} \left\langle \prod_{l=1}^p \left(1 - \frac{x}{1-x} e^{2\pi i \theta_l} \right)^{N+a_2} \right\rangle_{\text{CE}_{2,p}[e^{\pi i (a_1 - N)\theta}]1 + e^{2\pi i \theta} | a_1 + N]}.
$$
 (2.27)

Proof. To apply [\(2.25\)](#page-8-0) it is necessary to be able to identify the corresponding generating functions $b(z)$ that are consistent with the Toeplitz determinants in [\(2.24\)](#page-7-2). For this we first note the hypergeometric polynomial forms of the Laguerre and Jacobi polynomials

$$
L_n^{(a)}(x) \propto {}_1F_1(-n, a+1; x), \quad P_n^{(a_1, a_2)}(1-2x) \propto {}_2F_1(-n, 1+a_1+a_2+n; a_1+1; x) \tag{2.28}
$$

(the relevant orthogonal polynomials for the Jacobi weight $w(x) = x^{\alpha_1}(1-x)^{\alpha_2}\mathbb{1}_{0 \le x \le 1}$ are the Jacobi polynomials $P_n^{(a_1,a_2)}(1-2x)$. For these hypergeometric functions $_1F_1$ and $_2F_1$ more generally we have the circular integral forms (see e.g. [\[54,](#page-49-1) Exercises 13.1 q.4(i) with $N = 1$ and Eq. (13.11) with $N = 1$

$$
{}_{1}F_{1}(-b;a+1;t) \propto \int_{-1/2}^{1/2} e^{\pi i \theta(a-b)} |1+e^{2\pi i \theta}|^{a+b} e^{-te^{2\pi i \theta}} d\theta
$$

$$
{}_{2}F_{1}(r,-b;a+1;t) \propto \int_{-1/2}^{1/2} e^{\pi i \theta(a-b)} |1+e^{2\pi i \theta}|^{a+b} (1+te^{2\pi i \theta})^{-r} d\theta,
$$
 (2.29)

valid for Re $(a + b) > -1$.

The LUE duality [\(2.26\)](#page-8-1) follows immediately by substituting for the Laguerre polynomial on the RHS of the first identity in (2.24) as is consistent with (2.28) and (2.29) . In the Jacobi case the particular ${}_2F_1$ in [\(2.24\)](#page-7-2) (after the replacement $x \mapsto 2x - 1$) does not allow the circular integral in [\(2.29\)](#page-8-3) to be identified in a form as required by [\(2.25\)](#page-8-0). However, upon use of the Euler transformation

$$
{}_2F_1(-n, 1+a_1+a_2+n; \alpha_1+1; x) = (1-x)^n {}_2F_1(-a_2-n, -n; a_1+1; -\frac{x}{1-x}) \qquad (2.30)
$$

and factoring $(1-x)^n$ out of the determinant, the form as required by (2.25) is obtained, and [\(2.27\)](#page-8-4) results. \Box

Remark 2.1.

1. The RHS of [\(2.26\)](#page-8-1) and [\(2.27\)](#page-8-4) equals unity for $x = 0$. Requiring the same on the LHS provides a way to fix the proportionality constants.

2. Later (see [\(3.15\)](#page-12-1) below with $\beta = 2$), as an alternative to [\(2.27\)](#page-8-4) we will obtain a result which implies that the RHS can be rewritten in the manifestly polynomial form [\[54,](#page-49-1) Exercises 13.2 q.6(ii) with $\beta = 2$]

$$
(1-x)^{pN} \left\langle \prod_{l=1}^p \left(1 - \frac{x}{1-x} e^{2\pi i \theta_l} \right)^N \right\rangle_{\text{CE}_{2,p}[e^{\pi i (a_1 - a_2 - N)\theta}]1 + e^{2\pi i \theta} | a_1 + a_2 + N]}.
$$
(2.31)

2.5 Characteristic polynomial dualities for unitary random matrices

The Wronskian type determinant formula [\(2.15\)](#page-5-5) remains valid in the setting of circular ensembles [\(2.2\)](#page-3-3) with $\beta = 2$. In saying this, it is assumed that the orthogonality relation now has the form $\int_1^{1/2}$ $-\frac{1}{2}w(\theta)p_j(e^{2\pi i\theta})p_k(e^{-2\pi i\theta}) d\theta = h_j\delta_{j,k}$, and thus involves the complex conjugate operation. For the choice of weight

$$
w(\theta) = e^{\pi i \theta (a_1 - a_2)} |1 + e^{2\pi i \theta}|^{a_1 + a_2}
$$
\n(2.32)

one has the hypergeometric polynomial expressions

$$
p_n(z) \propto {}_2F_1(-n, a_1+1; -a_2-n+1; -z) \propto {}_2F_1(-n, a_1+1; a_1+a_2+1; 1+z), \tag{2.33}
$$

where the second form follows upon use of a transformation formula satisfied by the ${}_2F_1$ function (for the first see [\[54,](#page-49-1) Exercises 5.5 q.3]). Upon use now of the classical integral representation of Euler type we can deduce the duality

$$
\left\langle \prod_{l=1}^{N} (z - e^{2\pi i \theta_l}) \right\rangle_{\text{ME}_{2,N}[w]} \propto \left\langle (1 - (1+z)x)^N \right\rangle_{x \in B[a_1+1, a_2]},\tag{2.34}
$$

where $B[b, c]$ denotes the beta distribution, supported on $(0, 1)$ with density proportional to $x^{b-1}(1-x)^{c-1}.$

For circular ensembles, of more interest than averages of characteristic polynomials is averages of their absolute value squared. Noting that for $|z|=1$

$$
\prod_{l=1}^{N} |z - e^{2\pi i \theta_l}|^2 = (-1)^N z^N \prod_{l=1}^{N} e^{-2\pi i \theta_l} (1 - \bar{z} e^{2\pi i \theta_l})^2,
$$
\n(2.35)

the equivalent problem in relation to averages of even powers of this quantity is to consider

$$
\left\langle \prod_{l=1}^{N} (1 - \bar{z}e^{2\pi i \theta_l})^{2q} \right\rangle_{\text{ME}_{2,N}[e^{-2\pi i q \theta}w]}
$$
\n(2.36)

for q a positive integer. With N replaced by p, 2q by N, and $a_1 + a_2$ replaced by $a_1 + a_2 + 2q$ we recognise this average from (2.31) , which we know satisfies a duality with the JUE average in [\(2.27\)](#page-8-4). Working along this line, together with its β extension, will be given in [§3.6.](#page-18-0)

3 Powers and products of characteristic polynomial dualities for β ensembles

3.1 Jack polynomial theory

Common to all β generalisations is the essential use of Jack polynomial theory [\[116,](#page-52-4) Ch. VI.6], [\[54,](#page-49-1) Ch. 12], [\[111,](#page-51-2) Ch. 7]. Jack polynomials, to be denoted $P_{\kappa}^{(\alpha)}(\mathbf{x})$, depend on a set of variables $x = \{x_i\}$ and a parameter $\alpha > 0$. When expanded in terms of the monomial symmetric polynomials $\{m_u(\mathbf{x})\}$ they have the triangular structure

$$
P_{\kappa}^{(\alpha)}(\mathbf{x}) = m_{\kappa}(\mathbf{x}) + \sum_{\mu < \kappa} c_{\kappa,\mu}^{(\alpha)} m_{\mu}(\mathbf{x}).\tag{3.1}
$$

Here the notation $\mu < \kappa$ denotes the partial order on partitions defined by the requirement that $\sum_{i=1}^{s} \mu_i \leq \sum_{i=1}^{s} \kappa_i$, for each $s = 1, \ldots, \ell(\kappa)$ ($\ell(\kappa)$) denotes the number of nonzero parts of κ). Once this structure is stipulated, the Jack polynomials can be uniquely determined by an orthogonality property. Within the theory, several orthogonalities hold. Here we make note the one which relates to a random matrix average

$$
\left\langle P_{\kappa}^{(2/\beta)}(\mathbf{z})P_{\mu}^{(2/\beta)}(\bar{\mathbf{z}})\right\rangle_{\mathrm{CE}_{\beta,N}[1]} \propto \delta_{\kappa,\mu},\tag{3.2}
$$

where $\mathbf{z} = \{e^{2\pi i \theta_l}\}_{l=1}^N$; see e.g. [\[57,](#page-49-3) Prop. 12.6.3] for the normalisation.

The Jack polynomials with $\alpha = 1$ are the familiar Schur polynomials [\[116\]](#page-52-4), denoted $s_{\kappa}(\mathbf{x})$, while for $\alpha = 2$ they up to choice of normalisation the zonal polynomials of mathematical statistics [\[122\]](#page-52-5). Both these cases permit an interpretation in terms of spherical functions, as does the case $\alpha = 1/2$ [\[116\]](#page-52-4). These three cases furthermore relate to certain random matrix integrals. In this regard, let us define the particular hypergeometric function in two sets of variables $\mathbf{x} = \{x_i\}_{i=1,\dots,N}$ and $\mathbf{y} = \{y_i\}_{i=1,\dots,N}$ as a series in Jack polynomials according to

$$
{}_{0}\mathcal{F}_{0}^{(\alpha)}(\mathbf{x};\mathbf{y}) = \sum_{\kappa} \frac{\alpha^{|\kappa|}}{h_{\kappa}'} \frac{P_{\kappa}^{(\alpha)}(\mathbf{x}) P_{\kappa}^{(\alpha)}(\mathbf{y})}{P_{\kappa}^{(\alpha)}((1)^{N})}.
$$
(3.3)

Here $(1)^N$ denotes the point $x_i = 1$ $(i = 1, \ldots, N)$ and

$$
h'_{\kappa} = \prod_{s \in \kappa} (\alpha(a(s) + 1) + l(s))
$$
\n(3.4)

(in some Jack polynomial literature, for example in [\[54,](#page-49-1) Ch. 12 and 13], this quantity is alteratively denoted d'_{κ}), where the quantities $a(s)$, $l(s)$ are the arm and leg lengths at position s in the diagram associated with κ [\[116\]](#page-52-4).

While the quantity [\(3.3\)](#page-10-2) is not restricted to the three special α , its matrix integral interpretation is. For the latter, let U be a complex unitary $(\beta = 2)$, real orthogonal $(\beta = 1)$, or symplectic ($\beta = 4$ unitary matrix, and let $d^H U$ denote the corresponding Haar measure. Then in terms of (3.3) we have (see e.g. [\[54,](#page-49-1) Eq. (13.146)])

$$
\int e^{\text{Tr}(UGU^{\dagger}X_0)} d^{\text{H}}U =: {}_0\mathcal{F}_0^{(2/\beta)}(\mathbf{x}; \mathbf{x}^{(0)}), \tag{3.5}
$$

where G, X_0 are Hermitian matrices with complex $(\beta = 2)$, real $(\beta = 1)$ and quaternion $(\beta = 4)$ elements with eigenvalues **x** and $\mathbf{x}^{(0)}$ respectively. Note the simplification in the case that X_0 is proportional to the identity (i.e. all eigenvalues take on the constant value c)

$$
{}_{0}\mathcal{F}_{0}^{(\alpha)}(\mathbf{x};\mathbf{y})\Big|_{\mathbf{y}=(c)^{N}}=e^{cx_{1}+\cdots+cx_{N}},\tag{3.6}
$$

which in fact holds for general $\alpha \geq 0$ (see e.g. [\[54,](#page-49-1) Eq. (13.3)]). General aspects of the group integral in [\(3.5\)](#page-11-1), along the lines initiated by Harish-Chandra [\[97\]](#page-51-3), can be found in [\[118\]](#page-52-6).

3.2 Gaussian β ensemble

There is a very clean generalisation of [\(2.10\)](#page-5-4) and [\(2.23\)](#page-7-4) to the case of the Gaussian β ensemble. Proposition 3.1. We have [\[10\]](#page-47-4), [\[35\]](#page-48-3), [\[54,](#page-49-1) Eq. (13.162)]

$$
\left\langle \prod_{j=1}^{N} (x - \sqrt{\alpha} \lambda_j)^p \right\rangle_{\text{ME}_{2/\alpha, N}[e^{-\lambda^2}]} = i^{-pN} \left\langle \prod_{j=1}^{p} (ix - \lambda_j)^N \right\rangle_{\text{ME}_{2\alpha, p}[e^{-\lambda^2}]} \tag{3.7}
$$

and

$$
\left\langle \frac{1}{\det(x\mathbb{I}_N - H)^{p\beta/2}} \right\rangle_{\text{ME}_{N,\beta}[e^{-\lambda^2}]} = \left\langle \frac{1}{\det(x\mathbb{I}_p - H)^{N\beta/2}} \right\rangle_{\text{ME}_{p,\beta}[e^{-\lambda^2}]},\tag{3.8}
$$

(note that in this latter duality, in distinction to [\(3.7\)](#page-11-2), the value of β in the ensemble is the same on both sides, and x must have a nonzero imaginary part).

In Section [3.5](#page-14-1) the Gaussian β ensemble will be generalised to involve a source. Specialising dualities that can be derived in that setting will be shown also to imply Proposition [3.1.](#page-11-3) Other derivations of [\(3.7\)](#page-11-2) are also known. The first [\[10\]](#page-47-4) begins by generalising the LHS from a function of one variable, to a function of p variables by the replacement

$$
\left\langle \det(x\mathbb{I}_N - H)^p \right\rangle \mapsto \left\langle \prod_{l=1}^p \det(x_l \mathbb{I}_N - H) \right\rangle. \tag{3.9}
$$

It is then verified that this generalised quantity satisfies a set of partial differential equations associated with the Calogero-Sutherland quantum many body problem with harmonic confinement. This leads to the conclusion that the generalised average is in fact the polynomial part of a particular class of eigenfunctions for that model system, known as the generalised Hermite polynomials $\{P_{\kappa}^{(H)}(\mathbf{x};\alpha)\}\$ (specifically, with $\kappa=(p)^N$). On the other hand, by developing the theory associated with the latter, an alternative integral representation can be deduced, which leads to the RHS of [\(3.9\)](#page-11-4). For this, introduce the measure

$$
d\mu^{\mathcal{G}}(\mathbf{y}) := \mathrm{ME}_{p,2/\alpha}[e^{-y^2}]d\mathbf{y}.
$$
\n(3.10)

Then we have from [\[10,](#page-47-4) Cor. 3.2]

$$
e^{-(x_1^2 + \cdots + x_p^2)} P_{\kappa}^{(\mathrm{H})}(\mathbf{x}; \alpha) \propto \int_{\mathbb{R}^p} {}_0\mathcal{F}_0^{(\alpha)}(2\mathbf{y}; -i\mathbf{x}) P_{\kappa}^{(\alpha)}(i\mathbf{y}) d\mu^{\mathrm{G}}(\mathbf{y}). \tag{3.11}
$$

Now note that for $\kappa = (p)^N$, $P_{\kappa}^{(\alpha)}(i\mathbf{y}) = \prod_{l=1}^p y_l^N$, then set all the entries of **x** to equal x so that the ${}_{0}\mathcal{F}_{0}$ function can be simplified according to [\(3.6\)](#page-11-5). Then the RHS of [\(3.9\)](#page-11-4) can be identified as the RHS of [\(3.7\)](#page-11-2).

Another distinct derivation of [\(3.7\)](#page-11-2) proceeds by first establishing a class of duality formulas of independent interest.

Proposition 3.2. ([\[39,](#page-48-4) Th. 8.5.3], [\[40,](#page-48-0) Lemma 2.6], [\[35,](#page-48-3) Prop. 4]) Let κ' denote the conjugate partition, defined by interchanging the roles of the rows and columns in the diagram of κ . For $\ell(\kappa) \leq N$ and $\ell(\kappa') \leq p$ we have

$$
(-2/\beta)^{|\kappa|/2} \left\langle \frac{P_{\kappa}^{(2/\beta)}(\mathbf{x})}{P_{\kappa}^{(2/\beta)}((1)^{N})} \right\rangle_{\text{ME}_{\beta,N}[e^{-\lambda^{2}}]} = \left\langle \frac{P_{\kappa'}^{(\beta/2)}(\mathbf{x})}{P_{\kappa'}^{(\beta/2)}((1)^{p})} \right\rangle_{\text{ME}_{4/\beta,p}[e^{-\lambda^{2}}]}.
$$
(3.12)

(Note in particular that under the stated conditions there is no dependence on N or p in this identity.)

To derive [\(3.7\)](#page-11-2) from knowledge of this, we require too knowledge of the Jack polynomial homogeniety property $P_{\kappa}^{(\alpha)}(c\mathbf{x})=c^{|\kappa|}P_{\kappa}^{(\alpha)}(\mathbf{x})$ for any scalar c, and also the dual Cauchy identity (see e.g. [\[54,](#page-49-1) Eq. (12.187)])

$$
\prod_{k=1}^{N} \prod_{l=1}^{p} (1 - x_k y_l) = \sum_{\kappa \subseteq (p)^N} (-1)^{|\kappa|} P_{\kappa}^{(\alpha)}(\mathbf{x}) P_{\kappa'}^{(1/\alpha)}(\mathbf{y}),\tag{3.13}
$$

where $(N)^p$ denotes the partition with p parts all equal to N. In [\(3.12\)](#page-12-2) we cross multiply the denominators, and multiply both sides by $(-1)^{|k|/2}c^{|k|}$. Summing both sides over $\kappa \subseteq (N)^p$ (which is equivalent to $\kappa' \subseteq (p)^N$), we can apply the dual Cauchy identity [\(3.13\)](#page-12-3) to obtain, after replacing c by $1/x$ and multiplying both sides by x^{pN} , [\(3.7\)](#page-11-2).

3.3 Laguerre and Jacobi β ensembles — positive powers of characteristic polynomials

The dualites of Corollary [2.1](#page-8-5) also allow for clean β generalisations [\[51\]](#page-49-5), [\[54,](#page-49-1) Exercises 13.2 q.6].

Proposition 3.3. We have

$$
\left\langle \det(x\mathbb{I}_N - H)^p \right\rangle_{\text{ME}_{\beta,N}[x^a e^{-\beta x/2}]} \propto \left\langle \prod_{l=1}^p e^{-x e^{2\pi i \theta_l}} \right\rangle_{\text{CE}_{4/\beta,p}[e^{2\pi i (a+1)\theta/\beta - \pi i \theta(N+1)}|1 + e^{2\pi i \theta}|^{2(a+1)/\beta+N-1}]}
$$
(3.14)

and

$$
\langle \det(x\mathbb{I}_{N} - H)^{p} \rangle_{\text{ME}_{\beta,N}[x^{a_{1}}(1-x)^{a_{2}}]} \times (1-x)^{pN} \langle \prod_{l=1}^{p} \left(1 - \frac{x}{1-x} e^{2\pi i \theta_{l}}\right)^{N} \rangle_{\text{CE}_{4/\beta,p}[e^{2\pi i (a_{1}-a_{2})\theta/\beta-\pi i \theta_{N}}]1+e^{2\pi i \theta}|^{2(a_{1}+a_{2}+2)/\beta+N-2}]}\n\times (1-x)^{pN} \langle \prod_{l=1}^{p} \left(1 - \frac{x}{1-x} e^{2\pi i \theta_{l}}\right)^{N-1+2(a_{2}+1)/\beta} \rangle_{\text{CE}_{4/\beta,p}[e^{2\pi i (a_{1}+1)\theta/\beta-\pi i \theta(N+1)}]1+e^{2\pi i \theta}|^{2(a_{1}+1)/\beta+N-1]}}(3.15)
$$

In providing a proof outline, use will be made of hypergeometric functions based on Jack polynomials which relate to the series implied by the LHS of [\(3.6\)](#page-11-5); an extended account is given in [\[54,](#page-49-1) §13.1]. To define these, introduce the generalised Pochhammer symbol

$$
[u]_{\kappa}^{(\alpha)} := \prod_{l=1}^{N} \frac{\Gamma(u - (j-1)/\alpha + \kappa_l)}{\Gamma(u - (j-1)/\alpha)};
$$
\n(3.16)

see e.g. [\[54,](#page-49-1) Eq. (12.46)]. The family of hypergeometric functions of interest are then specified by

$$
{}_{p}F_{q}^{(\alpha)}(a_{1},\ldots,a_{p};b_{1},\ldots,b_{q};\mathbf{x}) := \sum_{\kappa} \frac{\alpha^{|\kappa|}}{h_{\kappa}'} \frac{[a_{1}]_{\kappa}^{(\alpha)} \ldots [a_{p}]_{\kappa}^{(\alpha)}}{[b_{1}]_{\kappa}^{(\alpha)} \ldots [b_{q}]_{\kappa}^{(\alpha)}} P_{\kappa}^{(\alpha)}(\mathbf{x}).
$$
\n(3.17)

With $\mathbf{x} = \{x_i\}_{i=1}^m$, the sum is over all partitions $\kappa_1 \geq \kappa_2 \geq \cdots \geq \kappa_m \geq 0$, and conventionally the sum is performed in order of increasing $|\kappa|$.

Proof outline of Prop. [3.3.](#page-12-4) In both the Laguerre and Jacobi cases, a pathway to the dualities is to generalise the average on the LHS by introducing variables $\mathbf{x} = \{x_l\}_{l=1}^p$ according to [\(3.9\)](#page-11-4). This function of p variables can be uniquely characterised by a set of p partial differential equations which moreover admit the Jack polynomial based hypergeometric function series solutions [\[104\]](#page-51-4)

$$
{}_1F_1^{(\beta/2)}(-N;2(a+p)/\beta;\mathbf{x}), \quad {}_2F_1^{(\beta/2)}(-N,N-1+2(a_1+a_2+p+1)/\beta;2(a_1+p)/\beta;\mathbf{x}). \tag{3.18}
$$

One notes that the first of these can be obtained from the second by the scaling $\mathbf{x} \mapsto \frac{\beta}{2a_2} \mathbf{x}$ and then taking the limit $\alpha_2 \to \infty$, as is consistent with the relation between the Jacobi and Laguerre weights. Specialising then to the Jacobi case, it follows

$$
\left\langle \det(x\mathbb{I}_N - H)^p \right\rangle_{\text{ME}_{\beta,N}[x^{a_1}(1-x)^{a_2}]} \propto {}_2F_1^{(\beta/2)}(-N, N-1+2(a_1+a_2+p+1)/\beta; 2(a_1+p)/\beta; (x)^p). \tag{3.19}
$$

With $\mathbf{x} = \{x_l\}_{l=1}^p$, the method of partial differential equations can be used to establish the explicit functional form (a generalised binomial formula) [\[104\]](#page-51-4)

$$
{}_{1}F_{0}^{(\beta/2)}(r;\underline{\ };\mathbf{x})=\prod_{l=1}^{p}(1-x_{l})^{-r}.
$$
\n(3.20)

This together with the orthogonality [\(3.2\)](#page-10-3) can be used to deduce the integration formula [\[54,](#page-49-1) Eq. (12.142)]

$$
\left\langle P_{\kappa}^{(\alpha)}(-z) \right\rangle_{\mathrm{CE}_{2/\alpha,p}[e^{\pi i (a-b)\theta}]1+e^{2\pi i \theta}|a+b]} = P_{\kappa}^{(\alpha)}((1)^p) \frac{[-b]_{\kappa}^{(\alpha)}}{[1+a+(p-1)/\alpha]_{\kappa}^{(\alpha)}}.
$$
(3.21)

Note that for the LHS to be well defined, it is required that $\text{Re}(a+b) > -1$. Multiplying through by $(t\alpha)^{|\kappa|}[r]_{\kappa}^{(\alpha)}/h'_{\kappa}$ and summing over κ , we see that on the LHS we encounter ${}_{1}F_{0}^{(\alpha)}$ $\mathcal{L}_0^{(\alpha)}(r; \underline{\ }; -\underline{\mathbf{z}}),$ which we know can be summed according to (3.20) , while the RHS is an example of the generalised hypergeometric function ${}_2F_1^{(\alpha)}$ $\int_1^{(\alpha)}$ of p variables all equal to t. Explicitly, it follows from [\(2.27\)](#page-8-4) that [\[54,](#page-49-1) Eq. (13.11)]

$$
\left\langle \prod_{l=1}^{p} (1 + te^{2\pi i \theta_l})^{-r} \right\rangle_{\text{CE}_{2/\alpha, p}[e^{\pi i (a-b)\theta}]1 + e^{2\pi i \theta} | a+b]} = {}_{2}F_{1}^{(\alpha)}\left(r, -b; \frac{1}{\alpha}(p-1) + a + 1; (t)^{p}\right). \tag{3.22}
$$

As with [\(3.21\)](#page-13-1), for this to be well defined, it is required that $\text{Re}(a + b) > -1$. However, comparison with ${}_2F_1^{(\beta/2)}$ $i_{1}^{(p/2)}$ in [\(3.19\)](#page-13-2) shows that for this case of interest, the latter requirement is violated.

To overcome this circumstance, we can use the fact that ${}_2F_1^{(\alpha)}$ $\int_1^{(\alpha)}$ satisfies an analogue of the classical Euler transformation [\(2.30\)](#page-9-2) [\[146\]](#page-53-2)

$$
{}_{2}F_{1}^{(\alpha)}(a,b;c;t_{1},\ldots,t_{p}) = \prod_{j=1}^{p} (1-t_{j})^{-a} {}_{2}F_{1}^{(\alpha)}\Big(a,c-b;c;-\frac{t_{1}}{1-t_{1}},\ldots,-\frac{t_{p}}{1-t_{p}}\Big); \tag{3.23}
$$

cf. [\(2.30\)](#page-9-2). Applying [\(3.23\)](#page-14-2) in [\(3.19\)](#page-13-2) gives rise to a $_2F_1^{(\alpha)}$ $f_1^{(\alpha)}$ for which [\(3.22\)](#page-13-3) holds, and [\(3.15\)](#page-12-1) results (the two stated dualities result by utilising the fact that ${}_2F_1^{(\alpha)}$ $i_1^{(\alpha)}$ is symmetric in its first two arguments).

3.4 Laguerre and Jacobi β ensembles — negative powers of characteristic polynomials

As for the case of the positive powers of these ensembles, use will be made of the hypergeometric functions based on Jack polynomials. As they are analytic at the origin, the first step is to write (considering the Jacobi case for definiteness)

$$
\left\langle \det(x\mathbb{I}_N - H)^{-r} \right\rangle_{\text{ME}_{\beta,N}[x^{a_1}(1-x)^{a_2}]} = y^{rN} \left\langle \det(\mathbb{I}_N - yH)^{-r} \right\rangle_{\text{ME}_{\beta,N}[x^{a_1}(1-x)^{a_2}]}, \quad y = \frac{1}{x}, (3.24)
$$

so that the average on the RHS is analytic at the origin in the variable y . Further progress is possible for $r = \beta p/2$, when the later average permits the ${}_2F_1^{(\alpha)}$ $1^{(4)}$ evaluation [\[104\]](#page-51-4), [\[54,](#page-49-1) Eq. (13.10)]

$$
\left\langle \det(\mathbb{I}_N - yH)^{-\beta p/2} \right\rangle_{\text{ME}_{\beta,N}[x^{a_1}(1-x)^{a_2}]} = {}_2F_1^{(\alpha)} \left(\frac{\beta N}{2}, \frac{\beta}{2}(N-1) + a_1 + 1; \beta(N-1) + a_1 + a_2 + 2; (y)^p \right). \tag{3.25}
$$

With this as the starting point, duality formulas for negative powers of characteristic polynomials in the Laguerre and Jacobi β ensemble cases can now be deduced.

Proposition 3.4. Require that $y < 0$. We have

$$
\left\langle \det(\mathbb{I}_N - yH)^{-\beta p/2} \right\rangle_{\text{ME}_{\beta, N}[x^{a_1}(1-x)^{a_2}]} = \left\langle \det(\mathbb{I}_p - yH)^{-\beta N/2} \right\rangle_{\text{ME}_{\beta, p}[x^{a_1}(1-x)^{a_2}]} \tag{3.26}
$$

and

$$
\left\langle \det(\mathbb{I}_N - yH)^{-\beta p/2} \right\rangle_{\text{ME}_{\beta,N}[x^a e^{-x}]} = \left\langle \det(\mathbb{I}_p - yH)^{-\beta N/2} \right\rangle_{\text{ME}_{\beta,p}[x^a e^{-x}]}.\tag{3.27}
$$

Proof. The duality (3.27) follows from (3.26) by a suitable scaling and limit process. In relation to [\(3.26\)](#page-14-4), we read off from [\[54,](#page-49-1) Eq. (13.12)] a matrix average formula for the ${}_2F_1^{(\alpha)}$ $j_1^{(\alpha)}$ function in [\(3.25\)](#page-14-5) distinct from that given therein, which implies the stated result. \Box

3.5 Gaussian and Laguerre β ensembles with a source

A generalisation of the classical Gaussian ensembles, which can be traced back to Dyson [\[42\]](#page-48-5), is to introduce a parameter $\tau \geq 0$ by defining $G = |1 - e^{-2\tau}|^{1/2}H + e^{-\tau}H^{(0)}$. Here H is a member of a classical Gaussian ensemble and $H^{(0)}$ is a fixed matrix of the same class (e.g. real symmetric for $H \in GOE$). The joint element PDF is then proportional to

$$
e^{-(\beta/2)\text{Tr}\,(G-e^{-\tau}H^{(0)})^2/(1-e^{-2\tau})},\tag{3.28}
$$

showing that the model of independent Gaussian elements is maintained, but now the means are no longer zero, but rather determined by $H^{(0)}$ (referred to as an external source [\[23\]](#page-47-5)). Changing variables to the eigenvalues and eigenvectors, there is no longer a separation of the two classes of variables, since [\(3.28\)](#page-14-6) depends on both. Rather integration over the eigenvectors gives for the eigenvalue PDF, up to proportionality

$$
\prod_{1 \leq j < k \leq N} |\lambda_k - \lambda_j|^\beta e^{-\tilde{\beta} \sum_{j=1}^N \lambda_j^2 - \tilde{\beta} t^2 \sum_{j=1}^N \mu_j^2} \int e^{2\tilde{\beta} t \text{Tr}(U\Lambda U^\dagger L)} d^{\text{H}} U. \tag{3.29}
$$

Here $\{\lambda_j\}$ are the eigenvalues of G, $\{\mu_j\}$ are the eigenvalues of $H^{(0)}$, $\Lambda = \text{diag}(\lambda_1, \ldots, \lambda_N)$, $L = \text{diag}(\mu_1, \dots, \mu_N)$, $\tilde{\beta} = \beta/(2(1 - e^{-2\tau}))$ and $t = e^{-\tau}$. Recalling [\(3.5\)](#page-11-1) gives the rewrite of [\(3.29\)](#page-15-0)

$$
\prod_{1 \leq j < k \leq N} |\lambda_k - \lambda_j|^\beta e^{-\tilde{\beta} \sum_{j=1}^N \lambda_j^2 - \tilde{\beta} t^2 \sum_{j=1}^N \mu_j^2} \sigma_0^{\mathcal{L}(2/\beta)}(2\tilde{\beta} t \lambda; \mu). \tag{3.30}
$$

With $\tilde{\beta} = t = 1$ (these parameters are effectively scales associated with the eigenvalues), we will denote the PDF corresponding to [\(3.30\)](#page-15-1) by $ME_{\beta,N}[e^{-\lambda^2};\mu]$. The case $\mu = 0$ reduces back to $ME_{\beta,N}[e^{-\lambda^2}]$. In the case $\beta=2$, a generalisation of [\(2.9\)](#page-5-0) to involve this matrix ensemble with a source was given by Brézin and Hikami [\[22\]](#page-47-3), which in turn was generalised to all $\beta > 0$ by Desrosiers [\[35\]](#page-48-3).

Proposition 3.5. We have

$$
\left(\frac{1}{i}\sqrt{\frac{2}{\beta}}\right)^{pN/2} \left\langle \prod_{l=1}^{p} \det\left(i\sqrt{\frac{\beta}{2}}\nu_{l}\mathbb{I}_{N}-H\right)\right\rangle_{\mathrm{ME}_{\beta,\mathrm{N}}[e^{-\lambda^{2}};\mu]} = \left\langle \prod_{j=1}^{N} \det\left(i\sqrt{\frac{2}{\beta}}\mu_{j}\mathbb{I}_{p}-H\right)\right\rangle_{\mathrm{ME}_{\frac{4}{\beta},\mathrm{p}}[e^{-\lambda^{2}};\nu]},\tag{3.31}
$$

with the case $v = 0$, $\beta = 2$ being equivalent to [\(2.9\)](#page-5-0). Also

$$
\left\langle \prod_{l=1}^{p} \det \left(\nu_l \mathbb{I}_N - H \right)^{-\beta/2} \right\rangle_{\text{ME}_{\beta,\text{N}}[e^{-\lambda^2};\mu]} = \left\langle \prod_{j=1}^{N} \det \left(\mu_j \mathbb{I}_p - H \right)^{-\beta/2} \right\rangle_{\text{ME}_{\beta,p}[e^{-\lambda^2};\nu]},\tag{3.32}
$$

which with $\{\nu_l\}$ and $\{\mu_j\}$ real and distinct requires $\beta < 2$ to be well defined, and for consistent branches of the fractional power function to be chosen.

Proof. We will consider only (3.31) , as the working required to establish (3.32) is similar. Introduce the operator

$$
\Delta_{N,\mathbf{z}}^{(\alpha)} := \sum_{j=1}^{N} \frac{\partial^2}{\partial z_j^2} + \frac{2}{\alpha} \sum_{1 \le j < k \le N} \frac{1}{z_j - z_k} \Big(\frac{\partial}{\partial z_j} - \frac{\partial}{\partial z_k} \Big). \tag{3.33}
$$

A formula following from the development of the theory of generalised Hermite polynomials based on Jack polynomials [\[10,](#page-47-4) Eq. (3.22)] (see also [\[132\]](#page-52-7)) gives

$$
\langle f(i\boldsymbol{\lambda}) \rangle_{\text{ME}_{2/\alpha,N}[e^{-\lambda^2};\mathbf{z}]} = e^{-\frac{1}{4}\Delta_{N,\mathbf{z}}^{(\alpha)}} f(\mathbf{z}) \Big|_{\mathbf{z} \mapsto i\mathbf{z}}.
$$
 (3.34)

Choose

$$
f(\mathbf{z}) = f(\mathbf{z}; \nu) = \prod_{l=1}^{p} \prod_{k=1}^{N} (\nu_l + \sqrt{\alpha} z_k).
$$
 (3.35)

One observes $f(i\mathbf{z}; \boldsymbol{\nu}) = i^{pN} f(\mathbf{z}; -i\boldsymbol{\nu})$ and moreover, via a direct calculation, that

$$
\Delta_{N,\mu}^{(\alpha)} f(\mu; \nu) \Big|_{\mu \mapsto i\mu} = i^{pN} \Delta_{p,\nu}^{(1/\alpha)} f(\mu; -\nu) \Big|_{\nu \mapsto i\nu}.
$$
\n(3.36)

Consequently

$$
e^{-\frac{1}{4}\Delta_{N,\mu}^{(\alpha)}}f(\mu;\nu)\Big|_{\mu\mapsto i\mu} = i^{pN}e^{-\frac{1}{4}\Delta_{p,\nu}^{(1/\alpha)}}f(\mu;-\nu)\Big|_{\nu\mapsto i\nu}.\tag{3.37}
$$

Comparing [\(3.34\)](#page-15-4) and [\(3.37\)](#page-16-0) shows

$$
\langle f(i\boldsymbol{\lambda}, \boldsymbol{\nu}) \rangle_{\text{ME}_{1/\alpha, N}[e^{-\lambda^2}; \boldsymbol{\mu}]} = i^{pN} \langle f(\boldsymbol{\lambda}, -i\boldsymbol{\mu}) \rangle_{\text{ME}_{\alpha, p}[e^{-\lambda^2}; \boldsymbol{\nu}]},
$$
(3.38)

which is [\(3.31\)](#page-15-2).

 \ddotsc

Remark 3.1. In the case $p = 1$, $\beta = 2$, [\(3.31\)](#page-15-2) can be rewritten to read

$$
\langle \det(\mu \mathbb{I}_N - (H + H_0)) \rangle_{\text{GUE}_N} = i^{pN/2} \Big\langle \prod_{j=1}^N (\mu - \nu_j + ix) \Big\rangle_{x \in \mathcal{N}(0, 1/2)},\tag{3.39}
$$

where $\{\nu_i\}$ are the eigenvalues of H_0 . In fact this holds in the more general case when H is a Wigner matrix as specified in (1.1) [\[55\]](#page-49-6), with the latter corresponding to the case that H_0 is the zero matrix.

The chiral Gaussian random matrices based on the structure [\(2.4\)](#page-4-4) admit a parametric extension involving a source matrix by replacing Z therein by $Y = |1 - e^{-2\tau}|^{1/2} Z + e^{-\tau} Z^{(0)}$. Of interest are the squared non-zero singular values Y (i.e. the eigenvalues $\{y_j\}_{j=1}^N$ of $Y^{\dagger}Y$). We know from [\[54,](#page-49-1) Eq. (11.105)] that the PDF for these singular values is given by, up to proportionality

$$
\prod_{j=1}^{N} y_j^{\beta a} \prod_{1 \le j < k \le N} |y_k - y_j|^\beta e^{-\tilde{\beta} \sum_{j=1}^N y_j - \tilde{\beta} t^2 \sum_{j=1}^N \mu_j} \sigma_{1}^{\{2/\beta\}}(\beta(a+N-1)/2+1; \tilde{\beta} t \mathbf{y}; \boldsymbol{\mu}), \quad (3.40)
$$

where $a = n - N + 1 - 2/\beta$. Here, with the components of **z** given by $\{z_j\}_{j=1}^N$, we have denoted by z^2 the vector with components $\{z_j^2\}_{j=1}^N$, and $\{\mu_j\}_{j=1}^N$ are the eigenvalues of $(Z^{(0)})^{\dagger}Z^{(0)}$. Further, with $\Lambda(L)$ the $n \times N$ matrix with non-zero entries on the diagonal only, $\{y_1, \ldots, y_N\}$ $({\mu_1, ..., \mu_N})$, and with $\beta = 1, 2$ or 4 the function ${}_0\mathcal{F}_1^{(2/\beta)}$ $i_1^{(2/p)}$ is defined as the matrix integral relating to the singular value decomposition of Z

$$
\int d^{\rm H} U \int d^{\rm H} V \, e^{\tilde{\beta} t \text{Tr}(V \Lambda^{\dagger} U^{\dagger} L + L^{\dagger} U \Lambda V^{\dagger})}.
$$
\n(3.41)

Here the unitary matrices U, V are of size $N \times N$ and $n \times n$ respectively, from the orthogonal $(\beta = 1)$, unitary $(\beta = 2)$ and symplectic $(\beta = 4)$ matrix groups respectively. Important is that with $\tilde{\beta} = t = 1$ (this is just for convenience as done in relation to [\(3.28\)](#page-14-6)), and $\alpha = 2/\beta$, the latter admits the Jack polynomial expansion (see e.g. [\[55,](#page-49-6) §4])

$$
{}_{0}\mathcal{F}_{1}^{(\alpha)}(u; \mathbf{y}; \boldsymbol{\mu}) = \sum_{\kappa} \frac{\alpha^{|\kappa|}}{[u]_{\kappa}^{(\alpha)} h'_{\kappa}} \frac{P_{\kappa}^{(\alpha)}(\mathbf{y}) P_{\kappa}^{(\alpha)}(\boldsymbol{\mu})}{P_{\kappa}^{(\alpha)}((1)^{N})},
$$
(3.42)

 \Box

with $u = (a + N - 1)/\alpha + 1 = \beta n/2$, thus allowing for a β generalisation. This function of two sets of variables relates to the $p = 0$, $q = 1$ Jack polynomial based hypergeometric function defined in [\(3.17\)](#page-13-4) according to

$$
{}_{0}\mathcal{F}_{1}^{(\alpha)}(u; \mathbf{y}; (1)^{N}) = {}_{0}F_{1}^{(\alpha)}(u; \mathbf{y}).
$$
\n(3.43)

In the case $\mu = 0, \,{}_0^{\vphantom{1}}\mathcal{F}_1^{(\alpha)}$ $_{1}^{(\alpha)}$ equals unity and we recognise [\(3.40\)](#page-16-1) as $ME_{\beta,N}[y^{a}e^{-y}]$; the generalised PDF with source variables μ will be denoted $\text{ME}_{\beta,N}[y^a e^{-y};\mu]$. In both cases we have assumed $\beta = t = 1$ and it is regarded as implicit that the eigenvalues can only take on positive values. Desrosiers [\[35\]](#page-48-3) has deduced an analogue of the duality [\(3.31\)](#page-15-2) for this source generalised ensemble.

Proposition 3.6. We have

$$
\left\langle \prod_{l=1}^p \det \left(\nu_l \mathbb{I}_N + \frac{2}{\beta} H \right) \right\rangle_{\text{ME}_{\beta, N}[\lambda^a e^{-\lambda}; \mu]} = \left(\frac{2}{\beta} \right)^{pN} \left\langle \prod_{j=1}^N \det \left(\mu_j \mathbb{I}_p + \frac{\beta}{2} H \right) \right\rangle_{\text{ME}_{\frac{4}{\beta}, p}[\lambda^{a'} e^{-\lambda}; \nu]}, \tag{3.44}
$$

where $a' = \frac{2}{\beta}$ $\frac{2}{\beta}(a+1)-1$. Also

$$
\left\langle \prod_{l=1}^p \det \left(\nu_l \mathbb{I}_N + H \right)^{-\beta/2} \right\rangle_{\text{ME}_{\beta, N}[\lambda^a e^{-\lambda}; \mu]} = \left\langle \prod_{j=1}^N \det \left(\mu_j \mathbb{I}_p + H \right)^{-\beta/2} \right\rangle_{\text{ME}_{\beta, p}[\lambda^{a'} e^{-\lambda}; \nu]}.
$$
 (3.45)

Proof. (Sketch.) A strategy analogous to that used in the proof of Proposition [3.5](#page-15-5) applies. Taking the place of [\(3.33\)](#page-15-6) is the operator

$$
D_{N,a,\mathbf{z}}^{(\alpha)} := \sum_{j=1}^{N} z_j \frac{\partial^2}{\partial z_j^2} + \frac{2}{\alpha} \sum_{1 \le j < k \le N} \frac{1}{z_j - z_k} \left(z_j \frac{\partial}{\partial z_j} - z_k \frac{\partial}{\partial z_k} \right) + (a+1) \sum_{j=1}^{N} z_j \frac{\partial}{\partial z_j},\tag{3.46}
$$

while taking the place of (3.34) is the operator based integration formula [\[10,](#page-47-4) Eq. (3.22)]

$$
\langle f(-\boldsymbol{\lambda}) \rangle_{\mathrm{ME}_{2/\alpha,N}[\lambda^a e^{-\lambda};\mathbf{z}]} = e^{-D_{N,a,\mathbf{z}}^{(\alpha)}} f(\mathbf{z}). \tag{3.47}
$$

In relation to [\(3.44\)](#page-17-0) the crucial observation from here is that with the choice

$$
f(\mathbf{z}) = f(\mathbf{z}; \nu) = \prod_{l=1}^{p} \prod_{k=1}^{N} (\nu_l - \alpha z_k),
$$
\n(3.48)

one has

$$
D_{N,a,\mu}^{(\alpha)} f(\mu;\nu) = D_{p,a',\nu}^{(\alpha)} f(\mu;\nu).
$$
\n(3.49)

Remark 3.2.

1. Let M be an $n \times N$ $(n \geq N)$ standard Gaussian complex matrix. Let M_0 be of the same size as M and have singular values $\{\mu_l\}_{l=1,\dots,N}$. It follows from [\(3.44\)](#page-17-0) with $p=1, \beta=2$, that

$$
\langle \det(\mu \mathbb{I}_N - (M + M_0)^{\dagger} (M + M_0)) \rangle = \Big\langle \prod_{j=1}^N (\mu - |\mathbf{x} + \boldsymbol{\mu}|^2) \Big\rangle_{x_j \in \mathcal{N}(0, 1/2)},\tag{3.50}
$$

where $\mathbf{x} = (x_1, \ldots, x_N)$. In [\[55,](#page-49-6) Cor. 3], a result equivalent to this was shown to hold for all $n \times N$ random matrices M with independent entries of mean zero and averaged modulus squared equal to unity.

2. Setting $\nu_1 = \cdots = \nu_p = -\beta x/2$ and $\mu = 0$ in [\(3.44\)](#page-17-0) gives on the LHS, up to a simple factor, the same average as on the LHS of [\(3.14\)](#page-12-5). However the matrix averages on the RHS are different. 3. In [\[83\]](#page-50-2) the duality [\(3.45\)](#page-17-1) with $\beta = 2$ is considered from a different perspective. In the case that all the components of ν are equal, it is used therein the study the large N asymptotics.

3.6 Circular and circular Jacobi β ensembles

 \ddotsc

The original circular β ensemble of Dyson [\[41\]](#page-48-6) is the case $w = 1$ of [\(2.2\)](#page-3-3). The name given to [\(2.2\)](#page-3-3) with the more general weight [\(2.32\)](#page-9-3) is the (generalised) circular Jacobi β ensemble (the restriction to $a_1 = a_2$ was the original meaning to this ensemble [\[54,](#page-49-1) §3.9]). It turns out that there are special features of averaged (squared modulus) characteristic polynomials in the case $w = 1$ relative to (2.32) , so warranting a separate discussion.

First, in the case $w = 1$, for $|z| = 1$ we observe that $\langle \prod_{l=1}^{N} |z+e^{2\pi i \theta_l}|^{2q} \rangle_{\text{CE}_{\beta, N}[1]}$ is independent of z (we write $z + e^{2\pi i \theta_l}$ rather than $z - e^{2\pi i \theta_l}$ to account for the range being $-1/2 < \theta_l <$ 1/2 in [\(2.2\)](#page-3-3) rather than $0 < \theta_l < 1$ as is conventional for $CE_{\beta,N}[1]$.) This is a consequence of rotational invariance. A natural generalisation is to replace each factor $|z + e^{2\pi i \theta_l}|^{2q}$ by $|z_1 + e^{2\pi i \theta_l}|^{2q} |z_2 + e^{2\pi i \theta_l}|^{2\mu}$. Rotational invariance then gives

$$
\left\langle \prod_{l=1}^{N} |z_1 + e^{2\pi i \theta_l}|^{2q} |z_2 + e^{2\pi i \theta_l}|^{2\mu} \right\rangle_{\text{CE}_{\beta, N}[1]} \propto \left\langle \prod_{l=1}^{N} |z + e^{2\pi i \theta_l}|^{2q} \right\rangle_{\text{CE}_{\beta, N}[|1 + e^{2\pi i \theta}|^{2\mu}]}, \quad z := \frac{z_1}{z_2}.\tag{3.51}
$$

The working which lead to (2.36) shows that for q a positive integer, upon simple manipulation, the absolute value signs can be removed, giving that [\(3.51\)](#page-18-1) is proportional to

$$
z^{qN} \Big\langle \prod_{l=1}^{N} (1 + \bar{z}e^{2\pi i \theta_l})^{2q} \Big\rangle_{\text{CE}_{\beta, N}[e^{-2\pi i \theta q}|1 + e^{2\pi i \theta}|^{2\mu}]}.
$$
(3.52)

The average in [\(3.51\)](#page-18-1) in the case $2q = 2\mu = \beta$ relates to the two-point correlation function of Dyson's circular β ensemble. For β even an evaluation in terms of the Jack polynomial based hypergeometric function ${}_2F_1^{(\alpha)}$ was given in the 1992 work [\[48\]](#page-48-7), and a duality identity obtained two years later in [\[53\]](#page-49-7) — these findings are summarised in [\[54,](#page-49-1) §13.2.1]. The same considerations can be used to obtain a duality identity for the average in [\(3.51\)](#page-18-1).

Proposition 3.7. Let q be a positive integer, let $\mu \ge q$ be real, and let $z = z_1/z_2$ with $|z_1| =$ $|z_2|=1$. We have

$$
\left\langle \prod_{l=1}^{N} |z_1 + e^{2\pi i \theta_l}|^{2q} |z_2 + e^{2\pi i \theta_l}|^{2\mu} \right\rangle_{\text{CE}_{\beta, N}[1]} \propto z^{qN} \left\langle \prod_{l=1}^{2q} (1 + (1 - \bar{z})e^{2\pi i \theta_l})^N \right\rangle_{\text{CE}_{4/\beta, 2q}[e^{\pi i (a - b)\theta}]1 + e^{2\pi i \theta} |^{a+b}]}\n\right|_{a+b=-1+2(\mu-q+1)/\beta} \propto z^{qN} \left\langle \prod_{l=1}^{2q} (1 - (1 - \bar{z})x_l)^N \right\rangle_{\text{ME}_{4/\beta, 2q}[x^a(1-x)^a]}\n\right|_{a=-1+2(\mu-q+1)/\beta}.\n\tag{3.53}
$$

Proof. As noted, it suffices to consider the average in (3.52) . Reading off from [\[54,](#page-49-1) Eq. (13.6)] we have that this average is equal to

$$
{}_{2}F_{1}^{(\beta/2)}\Big(-N,\frac{2}{\beta}(\mu+q);-(N-1)-\frac{2}{\beta}(\mu-q+1);(\bar{z})^{2q}\Big). \tag{3.54}
$$

Next we would like the apply (3.22) to express this as an average over 2q variables. However, this is not immediately possible as the requirement that $\text{Re}(a + b) > -1$ therein is violated. To

overcome this, we make use of the transformation identity [\[54,](#page-49-1) Prop. 13.1.7]

$$
{}_{2}F_{1}^{(\beta/2)}(-N,b;c;t_{1},\ldots,t_{m}) \propto {}_{2}F_{1}^{(\beta/2)}\Big(-N,b;-N+b+1+2(m-1)/\beta-c;1-t_{1},\ldots,1-t_{m}\Big). \tag{3.55}
$$

This shows [\(3.54\)](#page-18-3) can be replaced by

$$
{}_{2}F_{1}^{(\beta/2)}\left(-N,\frac{2}{\beta}(\mu+q);\frac{4}{\beta}(\mu+q);(v)^{2q}\right)\Big|_{v=1-\bar{z}}.\tag{3.56}
$$

In the notation of [\(3.22\)](#page-13-3) we now have $a + b = -1 + 2(\mu - q + 1)/\beta$, which is greater than -1 under the assumption $\mu \geq q$, allowing us to deduce the first line on the RHS.

In relation to the second line of the RHS, we require knowledge of a companion to (3.22) , which expresses the ${}_2F_1^{(\alpha)}$ $\int_1^{(\alpha)}$ function of p variables all equal as an average over the Jacobi β ensemble [\[54,](#page-49-1) Eq. (13.12)]

$$
\left\langle \prod_{l=1}^{p} (1 - tx_l)^{-r} \right\rangle_{\text{ME}_{2/\alpha, p}[x^{a_1}(1-x)^{a_2}]} = {}_2F_1^{(\alpha)} \left(r, \frac{1}{\alpha}(p-1) + a_1 + 1; \frac{2}{\alpha}(p-1) + a_1 + a_2 + 2; (t)^p \right). \tag{3.57}
$$

Comparing with [\(3.56\)](#page-19-0) gives the stated result.

We next consider

$$
I_{N,2q}(z) := \left\langle \prod_{l=1}^{N} |z - e^{2\pi i \theta_l}|^{2q} \right\rangle_{\text{CE}_{\beta,N}[1]}
$$
(3.58)

 \Box

in the case that $|z| < 1$. We know from [\[68\]](#page-49-8) that in this circumstance we have the rewrite

$$
I_{N,2q}(z) = \left\langle \prod_{l=1}^{N} (1+|z|^2 e^{2\pi i \theta_l})^q \right\rangle_{\text{CE}_{\beta,N}[e^{-\pi i \theta q}|1+e^{2\pi i \theta} |^q]}
$$
(3.59)

(cf. [\(3.52\)](#page-18-2)), valid for $q > -1$. Using this as the starting point, with q a positive integer, a minor modification of the working of the proof of Proposition [3.7](#page-18-4) allows for the deduction of a duality formula.

Proposition 3.8. Let q be a positive integer, let $|z| < 1$, and define $I_{N,2g}(z)$ as in [\(3.58\)](#page-19-1). We have

$$
I_{N,2q}(z) \propto \left\langle \prod_{l=1}^{q} (1 + (1 - |z|^2) e^{2\pi i \theta_l})^N \right\rangle_{\text{CE}_{4/\beta,q}[e^{\pi i (a-b)\theta}]1 + e^{2\pi i \theta} |a+b]}
$$

$$
\propto \left\langle \prod_{l=1}^{q} (1 - (1 - |z|^2) x_l)^N \right\rangle_{\text{ME}_{4/\beta,q}[x^a(1-x)^a]} \Big|_{a=-1+2/\beta} \tag{3.60}
$$

Neither [\(3.51\)](#page-18-1) nor [\(3.52\)](#page-18-2) give rise to dualities for the circular Jacobi ensemble unless $|z| = 1$. Then

$$
\left\langle \prod_{l=1}^{N} |z + e^{2\pi i \theta_l}|^{2q} \right\rangle_{\text{CE}_{\beta, N}[e^{\pi i (a-b)}]1 + e^{2\pi i \theta} |a+b]} \propto z^{qN} \left\langle \prod_{l=1}^{N} (1 + \bar{z}e^{2\pi i \theta_l})^{2q} \right\rangle_{\text{CE}_{\beta, N}[e^{\pi i \theta (a-b-2q)}]1 + e^{2\pi i \theta} |a+b]},
$$
\n(3.61)

which as follows from the working of the proof of Proposition [3.7](#page-18-4) permits a duality identity (for the special case $2q = \beta$ see [\[70\]](#page-49-9)).

Proposition 3.9. Let q be a positive integer, let $|z| = 1$, and require that $a \geq q - 1$. We have

$$
\left\langle \prod_{l=1}^{N} |z + e^{2\pi i \theta_l}|^{2q} \right\rangle_{\text{CE}_{\beta, N}[e^{\pi i (a-b)}|1 + e^{2\pi i \theta}|^{a+b}]}
$$
\n
$$
\propto z^{qN} \left\langle \prod_{l=1}^{2q} (1 + (1 - \bar{z})e^{2\pi i \theta_l})^N \right\rangle_{\text{CE}_{4/\beta, q}[e^{\pi i (a'-b')\theta}|1 + e^{2\pi i \theta}|^{a'+b'}]} \left| \begin{array}{c} a'+b'=-1+2(a-q+1/\beta) \\ a'-b'=-1+2(2b+a+q+1)/\beta \\ a'-b'=-1+2(2b+a+q+1)/\beta \\ a' - b'=-1+2(b-q+1)/\beta \end{array} \right. \quad (3.62)
$$

(The final of these averages also requires $b \geq q-1$.)

4 Interpretations and scaled large N asymptotics

4.1 Wigner semi-circle law

With $\sigma^2 = 1/(2N)$, and making use of [\(2.8\)](#page-4-2), we see that [\(1.1\)](#page-2-1) can be written

$$
\langle \det(x\mathbb{I}_N - \tilde{H}) \rangle_{\tilde{H}\in\mathcal{H}_N} = 2^{-N} (\sqrt{2N})^N H_N(\sqrt{2N}x). \tag{4.1}
$$

The significance of this scaling is that the Wigner class \mathcal{H}_N then has eigenvalue support on the compact interval $(-1, 1)$. Thus in the setting that the normalised limiting eigenvalue density, $\rho_{\infty,(1)}(y)$ say, has support on a compact interval I, it is true that for a large class of $a(y)$ (slow enough decay at infinity), for N large

$$
\left\langle e^{\sum_{l=1}^{N} a(y_l)} \right\rangle_{\mathcal{H}_N} = e^{N \int_I a(\lambda)\rho_{\infty,(1)}(y) \, dy + \mathcal{O}(1)};
$$
\n(4.2)

see e.g. $[80, §3.1], [85]$ $[80, §3.1], [85]$, which can be viewed as a law of large numbers. Now the LHS of (4.1) is of the form of the LHS of [\(4.2\)](#page-20-3) with $a(y) = a(y; x) = \log(x - y)$. Substituting [\(4.2\)](#page-20-3) in [\(4.1\)](#page-20-2) and taking the logarithmic derivative with respect to x then gives that to leading order

$$
W_1(x) := \int_{-1}^1 \frac{\rho_{\infty,(1)}(y)}{x - y} dy = \frac{1}{N} \frac{1}{u(x)} \frac{d}{dx} u(x), \quad u(x) = H_N(\sqrt{2N}x), \quad x \notin (-1, 1).
$$
 (4.3)

Consideration of the second order linear differential equation satisfied by $u(x)$ leads to the conclusion that $W_1(x)$ satisfies the quadratic equation $W_1^2/4 - xW_1 + 1 = 0$. Choosing the root such that $W_1(x) \sim 1/x$ as $x \to \infty$, and applying the Sokhotski-Plemelj formula $\rho_{\infty, (1)}(y) =$ 1 $\frac{1}{\pi} \text{Im}_{\epsilon \to 0^+} W_1(y + i\epsilon)$, then gives

$$
\rho_{\infty,(1)}(y) = \rho^{\mathcal{W}}(y), \qquad \rho^{\mathcal{W}}(y) := \frac{2}{\pi} (1 - y^2)^{1/2} \mathbb{1}_{|y| < 1},\tag{4.4}
$$

which is the density function specifying the Wigner semi-circle law; see e.g. [\[126\]](#page-52-8).

4.2 Gaussian β ensemble

4.2.1 Global density

Generally for the functional form [\(2.1\)](#page-3-2) specifying $ME_{\beta,N} [w]$, the definition of the eigenvalue density $\rho_{N,(1)}(\lambda) = \rho_{N,(1)}(\lambda; w)$ as an integral over all but one of the λ_i gives

$$
\rho_{N,\beta,(1)}(\lambda;w) \propto w(\lambda) \Big\langle \prod_{l=1}^{N-1} |\lambda - x_l|^\beta \Big\rangle_{\text{ME}_{\beta,N}[w]}.
$$
\n(4.5)

In the case of the Gaussian β ensemble it therefore follows from [\(3.7\)](#page-11-2) that for β even

$$
\rho_{N,(1)}(\lambda; e^{-\beta \lambda^2/2}) \propto e^{-\beta \lambda^2/2} \Big\langle \prod_{j=1}^{N-1} (ix - \lambda_j)^{N-1} \Big\rangle_{\text{ME}_{4/\beta, N-1}[e^{-\lambda^2}]}.
$$
\n(4.6)

Starting with this duality identity, several scaled asymptotic limits have been analysed using saddle point analysis; for a detailed account of this method in the present context see [\[37\]](#page-48-8). One is the direct evaluation of the global density limit [\[10,](#page-47-4) §5.3],

$$
\lim_{N \to \infty} \sqrt{\frac{2}{N}} \rho_{N,(1)}(\lambda; e^{-\beta N \lambda^2}) = \rho^{\mathcal{W}}(\lambda). \tag{4.7}
$$

In [\[36\]](#page-48-9) this result was extended to an asymptotic expansion in which the first $\frac{1}{\beta}{2}$ oscillatory terms were specified. Here we make note of the first of these.

Proposition 4.1. With $P^{\mathcal{W}}(\lambda) := \int_{-1}^{\lambda} \rho^{\mathcal{W}}(x) dx$, for large N and $|\lambda| < 1$,

$$
\sqrt{\frac{2}{N}}\rho_{N,(1)}(\lambda; e^{-\beta N\lambda^2}) = \rho^{\mathcal{W}}(\lambda) - \frac{2}{\pi} \frac{\Gamma(1+2/\beta)}{(\pi \rho^{\mathcal{W}}(\lambda))^{6/\beta - 1}} \frac{1}{N^{2/\beta}} \cos\left(2\pi N P^{\mathcal{W}}(\lambda) + (1 - 2/\beta)\arcsin\lambda\right) + \cdots,
$$
\n(4.8)

where the next oscillatory term is at order $N^{-8/\beta}$, while the next non-oscillatory term is at order N^{-1} ($\beta \neq 2$) and order N^{-2} for $\beta = 2$.

Analysis analogous to that of [§4.1](#page-20-1) can be applied to the Gaussian β ensemble. This is because up to a scaling the formula [\(4.1\)](#page-20-2) remains valid,

$$
\left\langle \prod_{j=1}^{N} (x - \lambda_j) \right\rangle_{\text{ME}_{\beta, N}[e^{-\beta \lambda^2/2}]} = 2^{-N} H_N(x),\tag{4.9}
$$

as follows from [\(3.7\)](#page-11-2) with $p = 1$ and the integral form of the Hermite polynomials [\(2.8\)](#page-4-2). In particular the RHS is independent of β , and so the working of [§4.1](#page-20-1) with respect to the global limit can be repeated to reclaim [\(4.4\)](#page-20-4).

4.2.2 Edge density

With $w(\lambda) = e^{-\beta \lambda^2/2}$, to leading order the right spectrum edge is at $\lambda = \sqrt{2N}$. It has been known since [\[50\]](#page-49-10) that a well defined limiting state results from the use of the so-called soft edge scaling variables $\{x_i\}$, specified by the requirement that $\lambda_i = \sqrt{2N} + x_i/(\sqrt{2}N^{1/6})$ (i.e. the origin is shifted to the location of the right edge, and the distances are measured on the scale of $1/N^{1/6}$). In [\[36\]](#page-48-9) a β dimensional integral formula was obtained for the soft edge scaling limit of the eigenvalue density for β even, based on a coalescing saddle point analysis applied to (4.6) . In [\[78\]](#page-50-5) this analysis was extended to give the next two large terms in the large N asymptotic expansion. In fact some structure was uncovered in this latter study, with the first order correction (occurring at order $N^{-1/3}$) being shown to be related to the leading order asymptotic form by a derivative operation (see also [\[63\]](#page-49-11), [\[29,](#page-48-10) Eq. (1.35)] for the case $\beta = 1$). Equivalently, by redefining the soft edge scaling to include a particular constant shift of the scaling variables x_i , this correction term can be cancelled to obtain an optimal $N^{-2/3}$ correction term [\[77,](#page-50-6) §5].

Proposition 4.2. For large N and β even

$$
\frac{1}{\sqrt{2}N^{1/6}}\rho_{N,(1)}\left(\sqrt{2N} + 2^{-1/2}N^{-1/6}\left(x + 1/2 - 1/\beta\right); e^{-\beta\lambda^2/2}\right) = \rho_{\infty,(1)}^{\text{soft}}(x;\beta) + \mathcal{O}(N^{-2/3}),\tag{4.10}
$$

where with

$$
K_{n,\beta}(x) = -\frac{1}{(2\pi i)^n} \int_{-i\infty}^{i\infty} dv_1 \cdots \int_{-i\infty}^{i\infty} dv_n \prod_{j=1}^n e^{v_j^3/3 - xv_j} \prod_{1 \le k < l \le n} |v_k - v_l|^{4/\beta}, \tag{4.11}
$$

(a multidimensional extension of the integral form of the Airy function) the limiting soft edge density $\rho_{\infty,(1)}^{\text{soft}}(x)$ is specified by

$$
\rho_{\infty,(1)}^{\text{soft}}(x;\beta) = \frac{1}{2\pi} \left(\frac{4\pi}{\beta}\right)^{\beta/2} \Gamma(1+\beta/2) \prod_{j=1}^{\beta} \frac{\Gamma(1+2/\beta)}{\Gamma(1+2j/\beta)} K_{\beta,\beta}(x). \tag{4.12}
$$

Remark 4.1.

1. Define $N' = N + \frac{2-\beta}{2\beta}$. An alternative to shifting x as in [\(4.10\)](#page-22-0) is to consider

$$
\frac{1}{\sqrt{2}(N')^{1/6}} \rho_{N,\beta} \left(\sqrt{2N'} + 2^{-1/2} (N')^{-1/6} x; e^{-\beta \lambda^2/2} \right), \tag{4.13}
$$

in which the leading order mean and standard deviation are shifted. In the cases $\beta = 1, 2$ and 4, results of Bornemann [\[16\]](#page-47-6) imply that this quantity has an asymptotic expansion in powers of $(N')^{-2/3}.$

2. The multidimensional integral form [\(4.11\)](#page-22-1) is well suited to determining the $|x| \to \infty$ asymptotics of $\rho_{\infty,(1)}^{\text{soft}}(x;\beta)$ [\[36\]](#page-48-9), [\[54,](#page-49-1) Eq. (13.68) with factor $1/(2\pi)$ corrected to $1/\pi$ in the first line].

A generalisation of $K_{n,\beta}(x)$ [\(4.11\)](#page-22-1) is specified by [\[35,](#page-48-3) [37\]](#page-48-8)

$$
\mathrm{Ai}_{n}^{(\alpha)}(\mathbf{f}) = -\frac{1}{(2\pi i)^{n}} \int_{-i\infty}^{i\infty} dv_{1} \cdots \int_{-i\infty}^{i\infty} dv_{n} \prod_{j=1}^{n} e^{v_{j}^{3}/3} \sigma_{0}^{\alpha}(\mathbf{v}; \mathbf{f}) \prod_{1 \leq k < l \leq n} |v_{k} - v_{j}|^{2/\alpha}, \qquad (4.14)
$$

(a further generalisation to $\text{Ai}_{n,m}^{(\alpha)}(\mathbf{f},\mathbf{s})$, where **s** is a second set of variables (*m* in number) has also shown itself [\[38\]](#page-48-11).) We have from [\[35,](#page-48-3) Prop. 10] that this results from a soft edge scaling limit of [\(3.31\)](#page-15-2) with $\mu = 0$. Note that [\(4.11\)](#page-22-1) corresponds to the case that all the entries of f are equal. The soft edge scaling limit in the case $\beta = 2$ (i.e. [\(2.9\)](#page-5-0)) has relevance to the computation of a certain class of algebraic-geometric quantities known as intersection numbers [\[23,](#page-47-5) §7.2].

As already remarked, [\(4.11\)](#page-22-1) (and [\(4.14\)](#page-22-2)) can be considered as generalisations of the Airy function. We remark that the asymptotic study of the duality [\(3.31\)](#page-15-2) in the case that the source μ consists of of an even number of variables, half of which take on the value a, and the other half take on the value $-a$, there is a scaling limit which leads to multidimensional generalisations of the Pearcey integral [\[24\]](#page-47-7), [\[71\]](#page-50-7).

4.2.3 Moments of the characteristic polynomial

Consider next

$$
\left\langle \prod_{l=1}^{N} |\lambda - x_l|^p \right\rangle_{\text{ME}_{\beta, N}[e^{-\beta N x^2}]},\tag{4.15}
$$

which corresponds to a global scaling of the power of the characteristic polynomial in the Gaus-sian β ensemble. Motivation to study the large N form of [\(4.15\)](#page-23-1) in the case $\beta = 2$ corresponding to the GUE came from the application of random matrix ensembles with a unitary symmetry to the modelling of the Riemann zeta function of the critical line due to Keating and Snaith [\[107\]](#page-51-5). The required analysis was undertaken by Brézin and Hikami [\[21\]](#page-47-1) (see also [\[23,](#page-47-5) Theorem 5.4 with $\lambda \mapsto 2\lambda$, $2\rho(2\lambda) \mapsto \rho^{\mathbf{W}}(x)$], [\[91,](#page-50-8) Eq. (21) with $m = 1$, after minor correction], [\[108,](#page-51-6) Th. 1 with $m = 1$, [\[31,](#page-48-12) Eq. (1.13) with $m = 1$, [\[101,](#page-51-7) expanded in terms of λ]).

Proposition 4.3. For large N and p a positive integer

$$
\left\langle \prod_{l=1}^{N} |\lambda - x_l|^{2p} \right\rangle_{\text{ME}_{2,N}[e^{-2Nx^2}]} = (\pi N \rho^{\text{W}}(\lambda))^{p^2} e^{2Np(\lambda^2 - 1/2 - \log 2)} \prod_{l=0}^{p-1} \frac{l!}{(p+l)!} \left(1 + \text{O}\left(\frac{1}{N}\right)\right). \tag{4.16}
$$

After noting the identity $\prod_{l=0}^{p-1} \frac{l!}{(p+l)!} = (G(1+p))^2/G(1+2p)$, where $G(z)$ denotes the Barnes G-function, this asymptotic expression was proved in [\[108\]](#page-51-6) to be valid for general $\text{Re}(2p) > -1$ (now with error term $O(log N/N)$). In keeping with (4.2) we expect (4.16) to exhibit the leading large N asymptotic form

$$
\log \left\langle \prod_{l=1}^{N} |\lambda - x_l|^{2p} \right\rangle_{\text{ME}_{2,N}[e^{-2Nx^2}]} = 2pN \int_{-1}^{1} \log |\lambda - x| \rho^{\text{W}}(x) \, dx + \cdots, \tag{4.17}
$$

which from the integral evaluation $\int_{-1}^{1} \log |\lambda - x| \rho^{W}(x) dx = \lambda^{2} - \frac{1}{2} - \log 2$ is indeed true. One sees from [\(4.16\)](#page-23-2) that the next term in [\(4.17\)](#page-23-3) is $p^2 \log N$. In the case of the circular ensemble for $\beta = 2$, when averages relate to Toeplitz determinants according to [\(2.25\)](#page-8-0), the analogous expansion is a modification of the Szegő asymptotic formula known as Fisher-Hartwig asymptotics [\[34\]](#page-48-13). Another point to note is that with $p = 1$ [\(4.16\)](#page-23-2) is consistent with [\(4.5\)](#page-21-4) with $\beta = 2$ and [\(4.7\)](#page-21-5).

For a large class of GUE linear statistics $\sum_{l=1}^{N} a(\lambda_l)$, the asymptotic formula [\(4.2\)](#page-20-3) has the generalisation that for large N and (continuous) k small enough

$$
\log \left\langle e^{k\sum_{l=1}^N a(x_l)} \right\rangle = kN \int_I a(x)\rho_{\infty,(1)}(x) dx + \frac{k^2}{2} \sum_{n=1}^\infty n a_n^2 + \cdots, \ \ a_n = \frac{1}{\pi} \int_0^\pi a(\cos\theta) \cos n\theta d\theta. \tag{4.18}
$$

The term proportional to k^2 corresponds to the variance of the fluctuation of the linear statistic — for more on this see [\[60\]](#page-49-12). Using the expansion (see e.g. [\[54,](#page-49-1) Exercises 1.4 q.4]) $\log(2|\cos\theta - \sin\theta|)$ $\cos \phi| = -\sum_{n=1}^{\infty} \frac{2}{n}$ $\frac{2}{n}$ cos n θ cos n ϕ , one computes that for $a(x) = a(x; \lambda) = \log |x - \lambda|$, $a_n =$ $-\frac{2}{n}$ $\frac{2}{n}$ cos $n\phi$ where $\lambda = \cos \phi$. Simple manipulation then gives

$$
\sum_{n=1}^{\infty} n a_n^2 = \frac{1}{2} \left(\sum_{n=1}^{\infty} \frac{1}{n} - \sum_{n=1}^{\infty} \frac{\cos 2n\phi}{n} \right) = \frac{1}{2} \sum_{n=1}^{\infty} \frac{1}{n} + \frac{1}{2} \log |2 \sin \phi|; \tag{4.19}
$$

in relation to the second equality see e.g. $[54, Eq. (14.95)]$. Here the asterisk indicates that a regularisation of the otherwise divergent series is required. One sees that choosing the latter to be $log N$, [\(4.19\)](#page-23-4) precisely matches the term raised to the power of p^2 in [\(4.16\)](#page-23-2).

By making use of the duality [\(3.7\)](#page-11-2), the recent work [\[137\]](#page-52-9) considers the analogue of [\(4.16\)](#page-23-2) for the general Gaussian β ensemble. In fact the compatibility of [\(4.16\)](#page-23-2) with [\(4.18\)](#page-23-5) makes its structure easy to anticipate. Thus the β generalisation of the latter leaves the first term on the RHS unchanged. It gives rise too to a further term proportional to k but which is independent of N [\[100\]](#page-51-8)

$$
k\left(\frac{1}{\beta} - \frac{1}{2}\right) \int_{-\infty}^{\infty} \left(\delta(x - 1) + \delta(x + 1) - \frac{1}{\pi} \frac{1}{(1 - x^2)^{1/2}} 1_{|x| < 1}\right) a(x) \, dx,\tag{4.20}
$$

which comes from the first subleading term of the large N expansion of the smoothed density [\[145,](#page-53-3) $\S3.2$], while the only effect on the term proportional to k is the requirement of the factor 2 $\frac{2}{\beta}$; see e.g. [\[60\]](#page-49-12). Indeed as shown in [\[137\]](#page-52-9) a steepest descent strategy applied to the RHS of [\(3.7\)](#page-11-2) shows that this anticipated functional form holds true (for purpose of comparison one notes that for $a(x) = a(x; \lambda) = \log |x - \lambda|$ with $|\lambda| < 1$ the integral in [\(4.20\)](#page-24-0) evaluates to $\log |1 - \lambda^2| + \log 2$; see [\[54,](#page-49-1) §1.4.2]).

Proposition 4.4. Let p be a positive integer. For large N we have

$$
\left\langle \prod_{l=1}^{N} |\lambda - x_{l}|^{2p} \right\rangle_{\text{ME}_{\beta,N}[e^{-\beta Nx^{2}}]} = A_{\beta,p}^{G} (\pi \rho^{W}(\lambda))^{p(2-\beta)/\beta} (\pi N \rho^{W}(\lambda))^{2p^{2}/\beta} e^{2Np(\lambda^{2}-1/2-\log 2)} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right), \tag{4.21}
$$

where, with $G(z)$ denoting the Barnes G-function,

$$
A_{\beta,p}^{\mathcal{G}} = \binom{2p}{p} \prod_{j=1}^{p} \frac{\Gamma(1+2j/\beta)}{\Gamma(1+2(j+p)/\beta)} = \prod_{l=0}^{\beta/2} \frac{(G(2(p-l)/\beta+1))^2}{G(2(2p-l)/\beta+1)G(1-2l/\beta)},\tag{4.22}
$$

 \Box

where the equality requires β even.

Proof. (Sketch only) The details in the case $p = 2$ have been given in [\[36\]](#page-48-9). One first has to modify the integration domain corresponding to the RHS of [\(3.7\)](#page-11-2) by certain contours so that the product of differences factor in the integrand is an analytic function. Next, the exponent of the N dependent factors in the integrand is considered from the viewpoint of its saddle points, with the conclusion being that these occur at $u_{\pm} = \frac{1}{2}$ $\frac{1}{2}(i\lambda \pm \sqrt{1-\lambda^2})$. After this, half of the integration contours are deformed to pass through u_+ , and the other half through u_- . Expansion about these points gives rise to the Gaussian form of the Selberg integral, which has a known evaluation as a product of gamma functions [\[54,](#page-49-1) Eq. (4.140)], giving rise to [\(4.22\)](#page-24-1).

Remark 4.2.

1. Substituting $2p = \beta$ gives that the average [\(4.21\)](#page-24-2) is proportional to $\rho^{\rm W}(\lambda)$, which is consistent with [\(4.7\)](#page-21-5) and [\(4.5\)](#page-21-4).

2. The constant factor $A_{\beta,p}^G$ in [\(4.21\)](#page-24-2), in the second of its stated forms, was first identified in the context of the chemical potential of an impurity charge of a log-gas on the circle (circular β ensemble), and can be given an evaluation in terms of the Barnes G-function for all rational β [\[49\]](#page-49-13), [\[54,](#page-49-1) Prop. (14.5.3)].

3. Generally

$$
\lim_{n\to 0}\frac{1}{n}\bigg(\Big\langle\prod_{l=1}^N|x-x_l|^n\Big\rangle-1\bigg)=\bigg\langle\sum_{l=1}^N\log|x-x_l|\bigg\rangle,
$$

an identity which underlies the so-called replica trick [\[45\]](#page-48-14). The problem of taking the limit $p \to 0$ on the RHS of the duality (2.10) , under the assumption that N is large, is discussed in [\[103\]](#page-51-9) and [\[105,](#page-51-10) [106\]](#page-51-11).

In the context of application to the Riemann zeta function on the critical line, specifically to an effect known as singularity dominated strong fluctuations, the large N form of the reciprocal characteristic polynomial moments

$$
\left\langle \frac{1}{|\det((x+i\epsilon)\mathbb{I}_N - H)|^{2p}} \right\rangle_{\text{ME}_{2,N}(e^{-2Nx^2})} \tag{4.23}
$$

is of interest [\[12\]](#page-47-8). Due to the absolute value sign in [\(4.23\)](#page-25-2), the relevant duality identity is not [\(2.23\)](#page-7-4) but rather [\(2.22\)](#page-7-5) with $p \mapsto 2p$, $x_1 = \cdots = x_p = x+i\epsilon$, $x_{p+1} = \cdots = x_{2p} = x-i\epsilon$. It is found in [\[82,](#page-50-9) Eq. (4) with $\mu_i \mapsto 2\mu_i$] (see too [\[84\]](#page-50-10)) that after normalisation by $|\langle \det((x+i\epsilon)\mathbb{I}_N - H)^{-p} \rangle|^2$, the $N \to \infty$ limit of [\(4.23\)](#page-25-2) exhibits, for $\epsilon \to 0$, the leading order behaviour

$$
\left(\frac{\pi \rho^{\mathcal{W}}(\lambda)}{4\epsilon}\right)^{p^2};\tag{4.24}
$$

note in particular the exponent p^2 as in [\(4.16\)](#page-23-2).

4.3 Laguerre and Jacobi β ensemble

4.3.1 Global scaling

For the density of the Laguerre β ensemble to have compact limiting support there are two distinct possible scaled Laguerre weights,

$$
ME_{\beta,N}[x^a e^{-\beta Nx/2}], \quad ME_{\beta,N}[x^{\beta N\gamma/2} e^{-\beta Nx/2}], \quad (a > -1, \gamma \ge 0),
$$
 (4.25)

and so two distinct ensembles to consider. In both cases the normalised limiting eigenvalue density is independent of β , while for the first of these it is independent of a and given by the $\gamma = 0$ case of the second. The functional form is specified by the Marchenko-Pastur density [\[126\]](#page-52-8)

$$
\rho^{\rm MP}(x) = \frac{1}{2\pi x} \sqrt{(x - c_{-})(c_{+} - x)} \mathbb{1}_{x \in [c_{-}, c_{+}]}, \quad c_{\pm} := (\sqrt{\gamma + 1} \pm 1)^{1/2}.
$$
 (4.26)

In the case of the Jacobi weight, the interval of support is compact from its definition, so there is no need to scale the eigenvalues. Still, there are distinct cases depending on the scaling of the exponents, with the analogue of [\(4.25\)](#page-25-3) being

$$
ME_{\beta,N}[x^{a_1}(1-x)^{a_2}], \quad ME_{\beta,N}[x^{\beta N\gamma_1/2}(1-x)^{\beta N\gamma_2/2}]. \tag{4.27}
$$

In further analogy to [\(4.25\)](#page-25-3), in both cases the normalised limiting eigenvalue density is independent of β, while for the first of these it is independent of a_1, a_2 and is equal to the $\gamma_1 = \gamma_2 = 0$ case of the second. It is given by a functional form first identified by Wachter [\[141\]](#page-53-4),

$$
\rho^{J}(x) = (\gamma_1 + \gamma_2 + 2) \frac{\sqrt{(x - c^{J})(d^{J} - x)}}{2\pi x(1 - x)},
$$
\n(4.28)

supported on $(c^{\{J}}, d^{\{J\}})$ with these endpoints specified by

$$
\frac{1}{(\gamma_1 + \gamma_2 + 2)^2} \left(\sqrt{(\gamma_1 + 1)(\gamma_1 + \gamma_2 + 1)} \pm \sqrt{(\gamma_2 + 1)} \right)^2.
$$
 (4.29)

One expects that the analogue of Proposition [4.4](#page-24-3) for the ensembles [\(4.25\)](#page-25-3) and [\(4.27\)](#page-25-4), which for $p = \beta/2$ relates to the computation of the global density, can be obtained by use of the dualities of Proposition [3.3.](#page-12-4) In fact the required working is carried out in the recent paper [\[137\]](#page-52-9). We know from [§4.1](#page-20-1) that another pathway to the computation of the global density is via the computation of the averaged characteristic polynomial.

4.3.2 Soft edge scaling

The right hand spectrum edge of both versions of the Laguerre ensemble in [\(4.25\)](#page-25-3) are soft edges, as they do not border a region where the eigenvalue density is strictly zero. On the other hand, for the Jacabi ensemble, only the second version in (4.27) exhibits a soft edge at its right boundary of leading order support. It is further true that the left edge of the second Laguerre and Jacobi versions exhibit a soft edge at their left boundary of leading order support.

At the present time only an analysis of the even β density at the right hand soft edge in both the Laguerre ensembles is available in the literature. The first such study was in [\[36\]](#page-48-9), considering the case of a fixed Laguerre exponent, with the limiting density computed to reclaim $\rho_{\infty,(1)}^{\text{soft}}(x;\beta)$ as appears in [\(4.10\)](#page-22-0) for the Gaussian β ensemble. Some years later, in [\[78\]](#page-50-5), the analysis was extended to include the case that the Laguerre exponent is proportional to N , and moreover to the calculation of the first two corrections. The working in [\[36\]](#page-48-9) and [\[78\]](#page-50-5) was based on the duality [\(3.14\)](#page-12-5), with it being a common feature of the β ensembles defined by [\(2.1\)](#page-3-2) that for even β the density is given in terms of an even moment of the characteristic polynomial. For both the cases of fixed and growing with N Laguerre exponent it was found that the leading order correction was related to $\rho_{\infty,(1)}^{\text{soft}}$ by a derivative operation. Due to this, what was the first correction term in the large N expansion can be eliminated by tuning the scaling parameters (recall (4.10)) and [\(4.13\)](#page-22-3)). However, as presented in [\[78\]](#page-50-5) the required tuning is different in both cases.

In a recent development Bornemann [\[16\]](#page-47-6) has found a parametrisation of the mean and standard derivation associated with the soft edge scaling which allows both cases to be treated on an equal footing. This requires first considering the Laguerre β ensemble with the Laguerre exponent parametrised as is natural from the viewpoint of the underlying Wishart matrix (see [\[54,](#page-49-1) Eq. (3.16)]). This is specified by $ME_{\beta,N}[x^{\alpha}e^{-\beta x/2}]$ with $\alpha = \beta(n-N+1)/2-1$, where $n - N + 1 > 0$. In terms of N, n define

$$
\mu_{N,n} = (\sqrt{N} + \sqrt{n})^2, \quad \sigma_{N,n} = (\sqrt{N} + \sqrt{n})\left(\frac{1}{\sqrt{N}} + \frac{1}{\sqrt{n}}\right), \quad h_{N,n} = \frac{1}{4}\left(\frac{1}{\sqrt{N}} + \frac{1}{\sqrt{n}}\right)^{4/3} \tag{4.30}
$$

(note that $\mu_{N,n}/N \to c_+$, with c_+ defined as in [\(4.26\)](#page-25-5)). Covering both the circumstances of fixed and N dependent Laguerre exponent, these variables can be used to write the respective results of [\[78\]](#page-50-5) as a single statement.

Proposition 4.5. Define $N' = N + \frac{2-\beta}{2\beta}$, $n' = n + \frac{2-\beta}{2\beta}$. For β even and N large we have

$$
\sigma_{N',n'}\rho_{N,\beta}\left(\mu_{N',n'} + \sigma_{N',n'}x; \lambda^{\beta(n-N+1)/2-1}e^{-\beta\lambda/2}\right) = \rho_{\infty,(1)}^{\text{soft}}(x;\beta) + O(h_{N',n'}); \tag{4.31}
$$

cf. LHS with [\(4.13\)](#page-22-3). Moreover, for $\beta = 1, 2$ and 4, results from [\[16\]](#page-47-6) imply that higher order terms on the RHS are a power series in $h_{N',n'}$.

4.3.3 Hard edge scaling

In distinction to the Gaussian ensemble, both the Laguerre and Jacobi ensembles exhibit a hard edge. This occurs at the left edge in the neighbourhood of the origin for the first ensemble of (4.25) , and both at the left and right edges, in the neighbourhood of $x = 0$ and $x = 1$ for the first ensemble of [\(4.27\)](#page-25-4). Considering the left edge for definiteness, in the Laguerre case a well defined limiting statistical state (the hard edge state) results from the scalings $\lambda_i \mapsto \lambda_i/(4N)$ (the factor of 4 is just for convenience), while in the Jacobi case one requires $\lambda_i \mapsto \lambda_i/(2N^2)$. For even β Corollary [2.1](#page-8-5) provides access to study the limiting functional form (first carried out for the Laguerre case in [\[51\]](#page-49-5)). In fact, as for the soft edge, the leading correction was found to be related to the limiting density by a derivative operation (first observed in [\[92,](#page-51-12) [63\]](#page-49-11) in relation to the cases $\beta = 1, 2$ and 4), and so can be eliminated by tuning the definition of the hard edge scaling.

At the hard edge, a further observable can similarly be analysed. This is the probability of no eigenvalue in a neighbourhood of the origin, $E_{N,\beta}(0,(0,s);\lambda^a e^{-\beta\lambda/2})$ say. In the hard edge scaling limit for general $\beta > 0$ and with a a non-negative its limiting functional form was computed in [\[51\]](#page-49-5), both in terms of the generalised hypergeometric function $_0F_1^{(\alpha)}$ with a variables all equal, and as an a-dimensional integral. Using in an essential way the duality of Corollary [2.1,](#page-8-5) the first two correction terms in powers of $1/N$ were computed in [\[79\]](#page-50-11) (Laguerre case) [\[69,](#page-49-14) Appendix, arXiv version] (Jacobi case, first correction only) with this same analysis in the Jacobi case also carried out directly from the finite N generalised hypergeometric expression in [\[144\]](#page-53-5). As for the density, the first correction was found to be related to the limiting probability by a derivative operation (see also [\[44,](#page-48-15) [13,](#page-47-9) [127,](#page-52-10) [96,](#page-51-13) [121\]](#page-52-11) in relation to this effect specific to the case $\beta = 2$). For brevity of presentation we make note here only of the results in relation to the probability $E_{N,\beta}(0,(0,s);\cdot)$. The limit will be given in terms of a particular ${}_{0}F_{1}^{(\alpha)}$ $j_1^{(\alpha)}$ in the case of equal variables, when we have available the multidimensional integral form ([\[54,](#page-49-1) modification of Eq. (13.27)])

$$
{}_{0}F_{1}^{(\beta/2)}(c+2(p-1)/\beta; (u)^{p}) = \prod_{j=1}^{p} \frac{\Gamma(1+2/\beta)\Gamma(c+2(j-1)/\beta)}{\Gamma(1+2j/\beta)}
$$

$$
\times \int_{-1/2}^{1/2} d\theta_{1} \cdots \int_{-1/2}^{1/2} d\theta_{p} \prod_{j=1}^{p} e^{2\pi i (c-1)\theta_{l}} e^{ue^{2\pi i \theta_{l}} + e^{-2\pi i \theta_{l}}} \prod_{1 \leq j < k \leq p} |e^{2\pi i \theta_{k}} - e^{2\pi i \theta_{j}}|^{4/\beta}. \quad (4.32)
$$

With $z_l = e^{2\pi i \theta_l}$, each integration herein can be thought of as over the unit circle in the complex z_l -plane. In fact for [\(4.32\)](#page-27-0) to hold with this contour of integration it is required that c be a positive integer. Otherwise the contour has to run along the negative real axis (lower half plane side), before traversing the unit circle, and returning along the negative real axis (upper half plane side); see the discussion in [\[56,](#page-49-15) Proof of Prop. 2].

Proposition 4.6. Consider $ME_{\beta,N}[\lambda^a e^{-\beta\lambda/2}]$ and require that a be a non-negative integer. Define $N_L = N + a/\beta$. For large N we have

$$
E_{N,\beta}(0, (0, s/(4N_{\rm L}); \lambda^a e^{-\beta \lambda/2}) = e^{-\beta s/8} {}_0F_1^{(\beta/2)}(2a/\beta; (s/4)^a) + O(N_{\rm L}^{-2}). \tag{4.33}
$$

Consider $ME_{\beta,N}[\lambda^{a_1}(1-\lambda)^{a_2}e^{-\beta\lambda/2}]$ and require that a_1 be a non-negative integer. Define $N_J = N - 1/2 + (1 + a_1 + a_2)/\beta$. We have

$$
E_{N,\beta}(0, (0, s/(2(N_J)^2); \lambda^{a_1}(1-\lambda)^{a_2}) = e^{-\beta s/8} {}_0F_1^{(\beta/2)}(2a_1/\beta; (s/4)^a) + O(N_J^{-2}). \tag{4.34}
$$

Remark 4.3.

1. For finite N the probability of no eigenvalues in $(0, s)$ for the Laguerre and Jacobi ensembles

with a and a_1 respectively non-negative integers admit evaluation formulas in terms of Jack polynomial hypergeometric functions according to ([\[51\]](#page-49-5) Laguerre case, use of [\(3.19\)](#page-13-2) Jacobi case)

$$
E_{N,\beta}(0,(0,s);\lambda^{a}e^{-\beta\lambda/2}) = e^{-\beta N_{\rm L}s/2}e^{as/2}{}_1F_1^{(\beta/2)}(-N_{\rm L};2a/\beta;(-s)^a)
$$

\n
$$
E_{N,\beta}(0,(0,s);\lambda^{a_1}(1-\lambda)^{a_2}) = (1-s)^{(\beta/2)(N_3^2-\gamma^2)}{}_2F_1^{(\beta/2)}(-N_{\rm J}+\gamma,N_{\rm J}+\gamma;2a/\beta;(-s/(1-s))^{a_1}),
$$
\n(4.35)

where the notation N_L, N_J is as in Proposition [4.6](#page-27-1) and $\gamma := (a_1 + a_2 + 1)/\beta - 1/2$. From the symmetry ${}_2F_1^{(\beta/2)}$ $I_1^{(\beta/2)}(a,b;c;{\bf x})={}_2F_1^{(\beta/2)}$ $1^{(p/2)}(b, a; c; \mathbf{x})$, we see immediately that the second of these is even in N_J , telling us that the large N expansion with $s \mapsto s/(2N_J^2)$ is in powers of $1/N_J^2$. Also, the generalised Kummer transformation for ${}_{1}F_{1}^{(\alpha)}$ $1^{(1)}$ (see e.g. [\[54,](#page-49-1) Eq. (13.16)])

$$
{}_{1}F_{1}^{(\alpha)}(a;c;t_{1},\ldots,t_{m}) = \prod_{j=1}^{m} e^{t_{j}} {}_{1}F_{1}^{(\alpha)}(c-a;c;-t_{1},\cdots,-t_{m}),
$$
\n(4.36)

applied to the first formula in [\(4.35\)](#page-28-2) tells us that $E_{N,\beta}(0,(0,s/(4N_{\rm L}));\lambda^a e^{-\beta\lambda/2})$ is even in $N_{\rm L}$ and so has an expansion for large N in powers of $1/N_{\rm L}^2$; cf. Remark [4.1.](#page-22-4)1.

2. From the relation between [\(3.41\)](#page-16-2) and [\(3.42\)](#page-16-3) specialised to $n = N$ and $L = \mathbb{I}_N$ we have that for $\beta = 1, 2$ and 4 (after changing variables $V^{\dagger}U \mapsto U$)

$$
{}_{0}F_{1}^{(2/\beta)}(\beta N/2; \mathbf{y}) = \int e^{\text{Tr}(U^{\dagger}\Lambda + U\Lambda)} d^{H}U.
$$
\n(4.37)

Here the matrix group integral is over the orthogonal group ($\beta = 1$), unitary group ($\beta = 2$) and symplectic unitary group ($\beta = 4$). In the case that $\Lambda = y\mathbb{I}_N$, the same ${}_0F_1^{(2/\beta)}$ $_{1}^{(2/\rho)}$ appear in [\(4.33\)](#page-27-2) and [\(4.34\)](#page-27-3), provided $2a/\beta$ is a non-negative integer, thus giving an evaluation of those hard edge probabilities in terms of classical matrix group integrals [\[81,](#page-50-12) §5.2] (see also [\[72,](#page-50-13) §3.1] and [\[14,](#page-47-10) Th. 2.1]). On the other hand these same group integrals appear as generating functions for the longest increasing subsequence length of classes of random permutations [\[130\]](#page-52-12), [\[8\]](#page-47-11). This fact has been put to use as a strategy for the analysis of the asymptotics of the longest increasing subsequence length in these cases [\[17\]](#page-47-12), [\[15\]](#page-47-13), [\[14\]](#page-47-10).

4.4 Circular and circular Jacobi β ensembles

4.4.1 Bulk scaling for the circular ensemble two-point correlation

Generally the bulk of the spectrum is any portion that is away from the edge or a singularity. In the circular ensemble the statistical state is rotationally invariant, and there is no edge or singularity associated with the spectrum. Bulk scaling refers to the use of scaled eigenvalues so that the mean spacing between eigenvalues is a nonzero constant (which we take to be unity) independent of N. This is achieved by setting $\theta_i = x_i/N$.

The two-point correlation function — defined as the integral over all but two of the eigenvalues in the joint eigenvalue PDF, multiplied by $N(N-1)$ as a type of normalisation — is simply related to the average on the LHS of [\(3.53\)](#page-18-5) with $N \mapsto N-2$ and $2q = 2\mu = \beta$. Thus, following [\[53\]](#page-49-7), a β-dimensional integral evaluation for the limiting two-point correlation function, $\rho_{(2)}^{\text{bulk}}(x_1, x_2; \beta)$ say, can be obtained by the duality averages on the RHS of [\(3.53\)](#page-18-5) (for definiteness, the second of these has been chosen).

Proposition 4.7. Define

$$
S_N(\lambda_1, \lambda_2, \beta) = \prod_{j=0}^{N-1} \frac{\Gamma(\lambda_1 + 1 + j\beta/2)\Gamma(\lambda_2 + 1 + j\beta/2)\Gamma(1 + (j+1)\beta/2)}{\Gamma(\lambda_1 + \lambda_2 + 2 + (N+j-1)\beta/2)\Gamma(1 + \beta/2)},
$$
(4.38)

which is the normalisation associated with $ME_{\beta,N}[x^{\lambda_1}(1-x)^{\lambda_2})$ as evaluated by Selberg; see e.g. [\[54,](#page-49-1) §4.1]. For β even we have

$$
\rho_{(2)}^{\text{bulk}}(x_1, x_2; \beta) = (\beta/2)^{\beta} \frac{((\beta/2)!)^3}{\beta!(3\beta/2)!} \frac{e^{-\pi i \beta (x_1 - x_2)} (2\pi (x_1 - x_2))^{\beta}}{S_{\beta}(-1 + 2/\beta, -1 + 2/\beta, 4/\beta)} \int_{[0,1]^{\beta}} du_1 \cdots du_{\beta}
$$

$$
\times \prod_{j=1}^{\beta} e^{2\pi i (x_1 - x_2) u_j} u_j^{-1 + 2/\beta} (1 - u_j)^{-1 + 2/\beta} \prod_{1 \le j < k \le \beta} |u_k - u_j|^{4/\beta}. \tag{4.39}
$$

4.4.2 Moments of moments of the characteristic polynomial

The moments of moments of the characteristic polynomial for the circular β ensemble are defined as [\[6\]](#page-47-14)

$$
\text{MoM}^{(\beta)}(k;q) = \left\langle \left(\int_0^1 \prod_{l=1}^N |e^{2\pi i \phi} + e^{2\pi i \theta_l}|^{2q} d\phi \right)^k \right\rangle_{\text{ME}_{\beta,N}[1]} \n= \int_0^1 d\phi_1 \cdots \int_0^1 d\phi_k \left\langle \prod_{l=1}^N |e^{2\pi i \phi_1} + e^{2\pi i \theta_l}|^{2q} \cdots |e^{2\pi i \phi_k} + e^{2\pi i \theta_l}|^{2q} \right\rangle_{\text{ME}_{\beta,N}[1]}, \quad (4.40)
$$

where the second line applies for k a positive integer. In the case $\beta = 2$ this quantity is of interest for its application to the local maxima of the Riemann zeta function on the critical line, and more generally for its relevance to the study of Gaussian multiplicative chaos; see the review [\[9\]](#page-47-15).

In the case $k = 2$ and q a positive integer the duality [\(3.53\)](#page-18-5) can be applied to give a rewrite of the second average in [\(4.40\)](#page-29-1). Taking into consideration that the implied proportionality constant therein is

$$
\left\langle \prod_{l=1}^{N} |1 + e^{2\pi i \theta_l}|^{4q} \right\rangle_{\text{ME}_{\beta,N}[1]} = \prod_{l=1}^{2q} \frac{\Gamma(N + 2(2q + l)/\beta)\Gamma(2l/\beta)}{\Gamma(N + 2l/\beta)\Gamma(2(2q + l)/\beta)},\tag{4.41}
$$

where use has been made of the so-called Morris integral evaluation $[54, Eq. (4.4)],$ with a simplification particular to the assumption that q is integer, we can reclaim a result of Assiotis [\[6,](#page-47-14) Eq. (18) written in the form of the first displayed equation in §3.3 therein] obtained using different working.

Proposition 4.8. For $4q^2 > \beta$ we have

$$
\lim_{N \to \infty} \frac{1}{N^{8q^2/\beta - 1}} \text{MoM}^{(\beta)}(k; q)|_{k=2} = \frac{1}{(2q)!} \prod_{j=0}^{2q-1} \frac{\Gamma(2/\beta)}{(\Gamma(2(j+1)/\beta))^2} \times \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{iqs} \int_0^1 dx_1 \cdots \int_0^1 dx_{2q} \prod_{l=1}^{2q} e^{-isx_l} (x_l(1 - x_l))^{-1+2/\beta} \prod_{1 \le j < k \le 2q} |x_k - x_j|^{4/\beta}. \tag{4.42}
$$

Proof. On the RHS of [\(4.40\)](#page-29-1) we use the periodicity in ϕ to shift the integration range to $[-1/2, 1/2]$. As already remarked, for $k = 2$ and q a positive integer the duality [\(3.53\)](#page-18-5) can be applied to give a rewrite of the second average in [\(4.40\)](#page-29-1). Changing variables $\phi = s/(2\pi N)$ and use of the elementary limit $(1 + u/N)^N \to e^u$ shows that, up to the factor corresponding to the normalisation of the average and a factor of $1/N$, the second line of (4.42) results. However this procedure is only well defined if the resulting integral over s is absolutely integrable. Using a method to analyse the multidimensional integral over x_1, \ldots, x_{2q} for large s presented in [\[52\]](#page-49-16), which involves considering the contribution of half of the integration variables clustered near 0 , and the other half clustered near 1 using a simple scaling of variables, we see that this requires $4q^2 > \beta$, with the decay of the integrand then being of order $|s|^{-4q^2/\beta}$. The factors of N result from the large N form of ratios of N dependent gamma functions in (4.41) , and the factor of a power of $1/N$ from the change of variables. The factors on the first line of the RHS of (4.42) result from [\(4.41\)](#page-29-3) and the product of gamma function function form of the normalisation factor for the average in the duality as read off from [\(4.38\)](#page-29-4). \Box

4.5 Circular Jacobi density for even β

In [\(2.32\)](#page-9-3) we set $a_1 = (\beta p + iq)/2$, $a_2 = \bar{a}_1$ so that the weight reads

$$
w(\theta) = e^{-\pi q \theta} |1 + e^{2\pi i \theta}|^{\beta p}.
$$
\n(4.43)

The points $\theta = \pm 1/2$ are referred to as a spectrum singularity of Fisher-Hartwig type (the parameter p can be viewed as determining the degeneracy of a conditioned eigenvalue at $\theta =$ $\pm 1/2$, while q determines the amplitude of the discontinuity in the factor $e^{-\pi q\theta}$ about the points $\theta = \pm \pi$). In the case β even, the average on the LHS of [\(3.62\)](#page-20-5) is (upon appropriate identification of the parameters) simply related to the density in the Jacobi circular ensemble with weight [\(4.43\)](#page-30-1). Use of (either of) the dualities on the RHS of [\(3.62\)](#page-20-5) leads to a β -dimensional integral form of the limiting density in the neighbourhood of the singularity, which is obtained by the scalings $\theta \mapsto -1/2 + x/(2\pi N)$ ($x > 0$) and $\theta \mapsto 1/2 + x/(2\pi N)$ ($x < 0$) [\[70\]](#page-49-9). Alternatively, from the underlying theory as outlined in the proof of Proposition [3.7,](#page-18-4) the evaluation formula can be given in terms of a particular ${}_{1}F_{1}^{(\alpha)}$ $\int_1^{(\alpha)}$ [\[113\]](#page-51-14) in the case of β equal variables, using the integral representation (see e.g. $[54, \text{Exercises } 13.1 \text{ q.}4(i)]$)

$$
{}_{1}F_{1}^{(\alpha)}(-b;a+1+(N-1)/\alpha;(t)^{n})\n\times \int_{-1/2}^{1/2} dx_{1} \cdots \int_{-1/2}^{1/2} dx_{n} \prod_{l=1}^{n} e^{\pi i x_{l}(a-b)} |1+e^{2\pi i x_{l}}|^{a+b} e^{-te^{2\pi i x_{l}}} \prod_{j\n(4.44)
$$

(We remark that in the cases $\beta = 2$ and 4 linear differential equations of degree 3 and 5 for the limiting density have been given in [\[76\]](#page-50-14).)

Proposition 4.9. Consider the Jacobi circular weight (4.43) in (2.2) , and for convenience shift each θ_l by $\theta_l \mapsto -1/2 + \theta_l$. Denote by $\rho^{\text{cJ}}_{\infty,(1)}(x;\beta)$ the limiting scaled density about the origin, and define $N_{cJ} = N + p$. We have that for large N

$$
\frac{1}{N_{\rm cJ}} \rho_{N,(1)} \left(x / N_{\rm cJ}; e^{-\pi q (\theta - 1/2)} |1 - e^{2\pi i \theta}|^{\beta p} \right) = \rho_{\infty,(1)}^{\rm cJ}(x) + \mathcal{O}(N_{\rm cJ}^{-2}),\tag{4.45}
$$

where

$$
\rho_{\infty,(1)}^{\rm cJ}(x) \propto e^{q\pi \text{sgn}(x)} e^{i\beta x/2} |x|^{p\beta} {}_1F_1^{(\beta/2)}(p+1-2iq/\beta;2p+2;(-ix)^\beta)).\tag{4.46}
$$

5 Characteristic polynomial dualities for non-Hermitian random matrices

5.1 Introductory remarks

In view of the result [\(1.1\)](#page-2-1), a logical starting point to explore dualities in relation to characteristic polynomials for non-Hermitian matrices is to consider

$$
\langle \det(z_1 \mathbb{I}_N - Z) \det(z_2 \mathbb{I}_N - \bar{Z}) \rangle \tag{5.1}
$$

for Z an $N \times N$ random matrix with all elements independently and identically distributed, chosen from a distribution with mean zero and variance σ^2 (interpreted as $\sigma^2 = \langle |z_{ij}|^2 \rangle$ is the elements are complex, which relates to the requirement of a complex conjugate in the second factor of (5.1)). One notes that if the second determinant were absent, (5.1) would trivially evaluate to $(z_1)^N$. In the case of standard Gaussian real entries (a general variance can be inserted by scaling), use of Grassmann integration by Akemann, Phillips and Sommers [\[2\]](#page-46-0) gave the duality relation

$$
\langle \det(z_1 \mathbb{I}_N - Z) \det(z_2 \mathbb{I}_N - Z) \rangle = \langle (z_1 z_2 + |w|^2)^N \rangle_{w \in \mathcal{CN}(0, \sigma^2)},\tag{5.2}
$$

where $CN(0, \sigma^2)$ denotes the complex normal distribution with mean zero and variance σ^2 ; cf. [\(1.1\)](#page-2-1). The case $z_1 = z_2$ was considered earlier in [\[43\]](#page-48-16), where a duality per se was not identified but rather the average was evaluated as an incomplete gamma function, which is noted in [\[2\]](#page-46-0) is consistent with the computation of the RHS of [\(5.2\)](#page-31-3) as a series in (z_1z_2) .

A generalisation of [\(5.1\)](#page-31-2) is to consider

$$
\Big\langle \prod_{l=1}^{k} \det(z_{l} \mathbb{I}_{N} - Z) \prod_{l=1}^{k} \det(w_{l} \mathbb{I}_{N} - \bar{Z}) \Big\rangle. \tag{5.3}
$$

Note that if all the entries of Z are real, the second product is not necessary. An evaluation of [\(5.3\)](#page-31-4) in the case of the so-called normal matrix model (see e.g. [\[27,](#page-48-2) Ch. 5]), defined as having an eigenvalue PDF in the complex plane of the form proportional to

$$
\prod_{l=1}^{N} e^{-V(z_l, \bar{z}_l)} \prod_{1 \le j < k \le N} |z_j - z_k|^2,\tag{5.4}
$$

and moreoever with the requirement of rotational invariance $V(z_l, \bar{z}_l) = V(|z_l|)$, an evaluation formula analogous to [\(2.6\)](#page-4-1) has been obtained in [\[3\]](#page-46-1). Prominent in the class of rotationally invariant normal matrix models is the Ginibre unitary ensemble (GinUE) consisting of $N \times N$ matrices with independent standard complex Gaussian entries. One then has $V(z_l, \bar{z}_l) = |z|^2$; see [\[27,](#page-48-2) Eq. (1.7)]. A duality identity for multiple products of characteristic polynomials for a non-Hermitian ensemble was first given by Nishigaki and Kamenev [\[124\]](#page-52-13), who deduced

$$
\left\langle |\det(z_1 \mathbb{I}_N - Z)|^{2k} |\det(z_2 \mathbb{I}_N - Z)|^{2k} \right\rangle_{Z \in \text{GinUE}_N} = \left\langle \det \begin{bmatrix} D & -Y \\ Y^{\dagger} & D^{\dagger} \end{bmatrix}^N \right\rangle_{Y \in \text{GinUE}_{2k}},\tag{5.5}
$$

where D is the $2k \times 2k$ diagonal matrix with the first k diagonal entries equal to z_1 and the remaining k diagonal entries equal to z_2 . For the generalisation of [\(5.5\)](#page-31-5) applying to [\(5.3\)](#page-31-4), see Proposition [5.2](#page-34-2) below.

The β generalisation of [\(5.4\)](#page-31-6) is to replace the exponent 2 on the product of differences with a general $\beta > 0$, which gives the two-dimensional Coulomb gas analogue of [\(2.1\)](#page-3-2). While in the Hermitian case of [\(2.1\)](#page-3-2) we have seen that duality relations enjoy a rich β generalisation, this is no longer true of two-dimensional Coulomb gas non-Hermitian β generalisations. Rather the variety of cases comes from the distinction of real, complex and quaternion elements, distinguished choices of invariant measures related to the classical weights in the Hermitian case, and possible symmetries of the matrices.

5.2 Zonal polynomials

Relationships between the Jack polynomial based hypergeometric functions with parameter $\alpha = 2/\beta$ and matrix integrals over the orthogonal group ($\beta = 1$), unitary group ($\beta = 2$) and symplectic unitary group ($\beta = 4$) have been seen in [\(3.5\)](#page-11-1), [\(3.41\)](#page-16-2) and [\(3.42\)](#page-16-3), and [\(4.37\)](#page-28-3). With parameter $\alpha = 2/\beta$ and $\beta = 1, 2$ and 4 the Jack polynomials are known as zonal polynomials, or sometimes as zonal spherical polynomials, due to their appearance in the broader theory of spherical functions [\[116\]](#page-52-4). Due to their relation to group integrals, they have received attention long before the more general Jack polynomials, particularly in the case $\alpha = 2$ ($\beta = 1$) in the context of multivariate statistics [\[122,](#page-52-5) Ch. 7]. A comprehensive historical account of the early literature is given in the thesis [\[131\]](#page-52-14).

Underpinning the evaluation of the group integrals (3.5) and (4.37) are group integral identities applying directly to the zonal polynomials. Conventionally, the normalisation used for the zonal polynomials is no longer that implied by [\(3.1\)](#page-10-5), but rather that given by introducing a scaling factor and defining,

$$
C_{\kappa}^{(\alpha)}(\mathbf{x}) := \frac{\alpha^{|\kappa|} |\kappa|!}{h_{\kappa}'} P_{\kappa}^{(\alpha)}(\mathbf{x});\tag{5.6}
$$

recall [\(3.4\)](#page-10-6) in relation to h'_κ and cf. the definition of the Jack polynomial based hypergeometric function [\(3.17\)](#page-13-4).

Proposition 5.1. ([\[99,](#page-51-15) case $\beta = 1$], [\[116\]](#page-52-4), [\[129\]](#page-52-15)) Let U be from the matrix group of unitary matrices as specified above depending on the value of β . Let A and B be $N \times N$ matrices with real, complex and real quaternion entries for $\beta = 1, 2, 4$, respectively. We have

$$
\int C_{\kappa}^{(2/\beta)} (AU^{\dagger} BU) d^{\text{H}} U = \frac{C_{\kappa}^{(2/\beta)} (A) C_{\kappa}^{(2/\beta)} (B)}{C_{\kappa}^{(2/\beta)} ((1)^{N})},\tag{5.7}
$$

where $C_{\kappa}^{(2/\beta)}(AU^{\dagger}BU)$ is defined as $C_{\kappa}^{(2/\beta)}(\mathbf{y})$ with $\mathbf{y}=(y_1,\ldots,y_N)$ denoting the eigenvalues of $AU^{\dagger}BU$, and similarly the meaning of $C_{\kappa}^{(2/\beta)}(A)$, $C_{\kappa}^{(2/\beta)}(B)$. In the quaternion case it is required that all matrices have doubly degenerate eigenvalues, and only one copy is included in the arguments of $C_{\kappa}^{(1/2)}$.

Furthermore, with $s_{\kappa}(\mathbf{x}) = P_{\kappa}^{(\alpha)}(\mathbf{x})|_{\alpha=1}$ denoting the Schur polynomial, one has

$$
\langle s_{\lambda}(AO) \rangle_{O \in O(N)} = \begin{cases} \frac{C_{\kappa}^{(2)}(AA^T)}{C_{\kappa}^{(2)}((1)^N)}, & \lambda = 2\kappa, \\ 0 & \text{otherwise,} \end{cases}
$$
(5.8)

$$
\langle s_{\lambda}(AU)s_{\kappa}(U^{\dagger}A^{\dagger})\rangle_{U\in U(N)} = \delta_{\lambda,\kappa}\frac{C_{\kappa}^{(1)}(AA^{\dagger})}{C_{\kappa}^{(1)}((1)^{N})},\tag{5.9}
$$

$$
\langle s_{\lambda}(AS) \rangle_{S \in \text{Sp}(2N)} = \begin{cases} \frac{C_{\kappa}^{(1/2)}(AA^{\dagger})}{C_{\kappa}^{(1/2)}((1)^{N})}, & \lambda = \kappa^{2}, \\ 0 & \text{otherwise}, \end{cases}
$$
(5.10)

where in [\(5.8\)](#page-33-0) the partition 2κ is the partition obtained by doubling each part of κ , while in [\(5.10\)](#page-33-0), κ^2 is the partition obtained by repeating each part of κ twice. We note too that in [\(5.10\)](#page-33-0), the meaning of $C_{\kappa}^{(1/2)}(AA^{\dagger})$ is $C_{\kappa}^{(1/2)}(\mathbf{x})$ where \mathbf{x} is the N independent eigenvalues of AA^{\dagger} (for A quaternion, the eigenvalues of AA^{\dagger} are doubly degenerate).

Application to random non-Hermitian random matrices relies on consequences of these group integrals in the case that the non-Hermitian random matrices being averaged over are bi-unitary invariant [\[139\]](#page-53-6), [\[129\]](#page-52-15), [\[75\]](#page-50-15), [\[54,](#page-49-1) Exercises 13.4 q.1].

Corollary 5.1. For $\beta = 1, 2$ and 4, let X, A, B be $N \times N$ matrices with real, complex and quaternion entries respectively. Let X be random matrix chosen from a left and right unitary invariant distribution, with the unitary matrices drawn from the orthogonal group, unitary group and symplectic unitary group respectively. We have

$$
\langle C_{\kappa}^{(2/\beta)}(AX^{\dagger}BX) \rangle_{X} = \frac{C_{\kappa}^{(2/\beta)}(A)C_{\kappa}^{(2/\beta)}(B)}{(C_{\kappa}^{(2/\beta)}((1)^{N}))^{2}} \langle C_{\kappa}^{(2/\beta)}(XX^{\dagger}) \rangle_{X}.
$$
 (5.11)

As in Proposition [5.1,](#page-32-1) in the quaternion case it is required that all matrices have double degenerate eigenvalues, and only one copy is included in the arguments of $C_{\kappa}^{(1/2)}$. Also

$$
\langle s_{\lambda}(AX) \rangle_{X} = \begin{cases} \frac{C_{\kappa}^{(2)}(AA^{T})}{C_{\kappa}^{(2)}((1)^{N})} \langle C_{\kappa}^{(2)}(XX^{T}) \rangle_{X}, & \lambda = 2\kappa, \\ 0 & \text{otherwise,} \end{cases}
$$
(5.12)

$$
\langle s_{\lambda}(AX)s_{\kappa}(X^{\dagger}A^{\dagger})\rangle_{X} = \delta_{\lambda,\kappa} \frac{C_{\kappa}^{(1)}(AA^{\dagger})}{C_{\kappa}^{(1)}((1)^{N})} \langle C_{\kappa}^{(1)}(XX^{\dagger})\rangle_{X},
$$
\n(5.13)

$$
\langle s_{\lambda}(AX) \rangle_{X} = \begin{cases} \frac{C_{\kappa}^{(1/2)}(AA^{\dagger})}{C_{\kappa}^{(1/2)}((1)^{N})} \langle C_{\kappa}^{(1/2)}(XX^{\dagger}) \rangle_{X}, & \lambda = \kappa^{2}, \\ 0 & \text{otherwise}, \end{cases}
$$
(5.14)

where for the averages on the LHS the matrices are as required for beta = 1, 2 and 4 respectively. As in [\(5.10\)](#page-33-0), in [\(5.14\)](#page-33-1) the meaning of $C_{\kappa}^{(1/2)}(AA^{\dagger})$ is $C_{\kappa}^{(1/2)}(\mathbf{x})$ where \mathbf{x} is the N independent eigenvalues of AA† .

Proof. Let $d\mu(X)$ be a left and right unitary invariant measure, and let $f(X)$ be integrable with respect to this measure. Left unitary invariance tells us that

$$
\langle f(AXBX^{\dagger})\rangle_X = \langle \langle f(AUXBX^{\dagger}U^{\dagger})\rangle_U \rangle_X. \tag{5.15}
$$

Choosing $f = C_{\kappa}^{(\alpha)}$ and using [\(5.7\)](#page-32-2) to evaluate the average over U on the RHS shows

$$
\langle C_{\kappa}^{(\alpha)}(AXBX^{\dagger})\rangle_{X} = \frac{C_{\kappa}^{(\alpha)}(A)}{C_{\kappa}^{(\alpha)}((1)^{N})}\langle C_{\kappa}^{(\alpha)}(XBX^{\dagger})\rangle_{X}.
$$
 (5.16)

Regarding the RHS, right unitary invariance and further use of [\(5.7\)](#page-32-2) gives

$$
\langle C_{\kappa}^{(\alpha)}(XBX^{\dagger})\rangle_{X} = \langle \langle f(XUBU^{\dagger}X^{\dagger}U^{\dagger})\rangle_{U}\rangle_{X} = \frac{C_{\kappa}^{(\alpha)}(B)}{C_{\kappa}^{(\alpha)}((1)^{N})}\langle C_{\kappa}^{(\alpha)}(XX^{\dagger})\rangle_{X}
$$
(5.17)

and [\(5.11\)](#page-33-2) results.

The derivation of the remaining identities from their companion identities in Proposition [5.1](#page-32-1) is similar. \Box

5.3 Dualities for products and powers of characteristic polynomials

5.3.1 Complex entry matrices

A duality formula for the average of the product of random matrices in [\(5.3\)](#page-31-4), generalising [\(5.5\)](#page-31-5), has been given in the thesis of Serebryakov [\[133,](#page-52-16) second statement of Th. 4.1].

Proposition 5.2. We have

$$
\left\langle \prod_{l=1}^{k} \det(z_{l} \mathbb{I}_{N} - Z) \det(w_{l} \mathbb{I}_{N} - \bar{Z}) \right\rangle_{Z \in \text{GinUE}_{N}} = \left\langle \det \begin{bmatrix} D_{1} & -Y \\ Y^{\dagger} & D_{2} \end{bmatrix}^{N} \right\rangle_{Y \in \text{GinUE}_{k}},\tag{5.18}
$$

where D_1, D_2 are diagonal matrices with diagonal entries **z** and **w** respectively.

Proof. Simple manipulation and use of the dual Cauchy identity [\(3.13\)](#page-12-3) with $\alpha = 1$ gives that the LHS of [\(5.18\)](#page-34-3) is equal to

$$
\prod_{l=1}^k (w_l z_l)^N \sum_{\mu,\kappa \subseteq (k)^N} (-1)^{|\mu|+|\kappa|} s_{\mu'}(1/\mathbf{z}) s_{\kappa'}(1/\mathbf{w}) \langle s_\kappa(Z) s_\mu(\bar{Z}) \rangle_{Z \in \text{GinUE}_N},\tag{5.19}
$$

where the notation $1/\mathbf{z}$ (similarly $1/\mathbf{w}$) is used to denote the vector with entries the reciprocal of the entries of z . The average in this expression is a particular case of (5.13) , reducing the task to computing $\langle s_{\kappa}(ZZ^{\dagger})\rangle_{Z\in\text{GinUE}_N}$. This can be achieved by first changing variables $W = ZZ^{\dagger}$ (see [\[54,](#page-49-1) Prop. 3.2.7]), then changing variables to the eigenvalues and eigenvectors of W (see [\[54,](#page-49-1) Prop. 3.2.2]), and finally making use of the special case $\alpha = 1$, $a = 0$ of the known Jack polynomial average (see e.g. [\[54,](#page-49-1) Eq. (12.152)])

$$
\langle P_{\kappa}^{(\alpha)}(\mathbf{x}) \rangle_{\mathrm{ME}_{2/\alpha,N}[\lambda^{a}e^{-\lambda}]} = P_{\kappa}^{(\alpha)}((1)^{N})[a+1+(N-1)/\alpha]_{\kappa}^{(\alpha)}.
$$
\n(5.20)

Consequently the LHS of [\(5.18\)](#page-34-3) is reduced to the form

$$
\prod_{l=1}^{k} (w_l z_l)^N \sum_{\kappa \subseteq (k)^N} s_{\kappa'}(1/\mathbf{z}) s_{\kappa'}(1/\mathbf{w}) [N]_{\kappa}^{(1)} = \prod_{l=1}^{k} (w_l z_l)^N \sum_{\kappa \subseteq (N)^k} s_{\kappa}(1/\mathbf{z}) s_{\kappa}(1/\mathbf{w}) [N]_{\kappa'}^{(1)}.
$$
 (5.21)

On the other hand, the RHS of [\(5.21\)](#page-34-4) can be rewritten by substituting for $s_{\kappa}(1/\mathbf{z})s_{\kappa}(1/\mathbf{w})$ according to [\(5.11\)](#page-33-2) with $\alpha = 1$, $N \mapsto k$, $A = D_1^{-1}$, $B = D_2^{-1}$, simplified by making further use of [\(5.20\)](#page-34-5), now with $\alpha = 1$, $N = k$ and $a = 0$. This shows that [\(5.21\)](#page-34-4) is equal to

$$
\prod_{l=1}^{k} (w_l z_l)^N \sum_{\kappa \subseteq (N)^k} [N]_{\kappa'}^{(1)} \frac{s_{\kappa}((1)^k)}{[k]_{\kappa}^{(1)}} \langle s_{\kappa}(D_1^{-1} X^\dagger D_2^{-1} X) \rangle_{X \in \text{GinUE}_k}.
$$
\n(5.22)

But from [\[54,](#page-49-1) Eqns. (12.104), (12.105), (13.1) and the fact that $h_{\kappa}|_{\alpha=1} = h'_{\kappa}|_{\alpha=1}$ we have

$$
[N]_{\kappa'}^{(1)} \frac{s_{\kappa}((1)^k)}{[k]_{\kappa}^{(1)}} = s_{\kappa'}((1)^N). \tag{5.23}
$$

With this noted, the sum over $\kappa \subseteq (N)^k$ can be performed using the dual Cauchy identity [\(3.13\)](#page-12-3) with $\alpha = 1$ to give in place of [\(5.22\)](#page-35-0) the expression

$$
\prod_{l=1}^{k} (w_l z_l)^N \left\langle \prod_{l=1}^{k} (1 + \gamma_l)^N \right\rangle, \tag{5.24}
$$

where $\{\gamma_l\}$ are the eigenvalues of $Z_1^{-1}X^{\dagger}Z_2^{-1}X$. Use of the block determinant identity

$$
\det\begin{bmatrix} A & B \\ C & D \end{bmatrix} = \det AD \det(\mathbb{I}_k - A^{-1}BD^{-1}C), \tag{5.25}
$$

identifies [\(5.24\)](#page-35-1) with the RHS of [\(5.18\)](#page-34-3).

Remark 5.1.

1. The duality [\(5.18\)](#page-34-3) can be generalised so that Z and \overline{Z} are replaced by ΩZ and $\Sigma \overline{Z}$ on the LHS [\[133,](#page-52-16) first statement of Th. 4.1]. Such generalised dualities have found application to the study of the spectral density in the Ginibre ensemble subject to an additive or multiplicative low rank perturbation [\[114\]](#page-51-16).

2. An alternative to the use of Schur polynomials in the derivation of the just mentioned generalisation of [\(5.18\)](#page-34-3) is to make use of a certain two-sided matrix diffusion operator identity $\Delta_X Q(X, Y) \propto \Delta_Y Q(X, Y)$ for $Q(X, Y)$ a particular kernel function relating to the averages and Δ_Z a (generalised) Laplacian [\[94\]](#page-51-17), [\[115\]](#page-51-18).

3. For H_1, H_2 independent GUE matrices, the elliptic Ginibre ensemble is specified by matrices of the form $\sqrt{1+\tau}H_1 + \sqrt{1-\tau}iH_2$, $|\tau| \leq 1$; see e.g. [\[27,](#page-48-2) §2.3]. This ensemble thus interpolates between the GUE ($\tau = 1$) and the GUE ($\tau = 0$). Grassmann integration methods have been used to deduce the duality identity for products of characteristic polynomials [\[3\]](#page-46-1), [\[133,](#page-52-16) Eq. (5.3)]

$$
\left\langle \prod_{j=1}^{k} \det(J - z_j \mathbb{I}_N) \det(J^{\dagger} - w_j \mathbb{I}_N) \right\rangle_{J \in \text{GinUE}_N} \propto \left\langle \prod_{j=1}^{k} \det \begin{bmatrix} -iD_1 + \sqrt{2\tau}A & C \\ C^{\dagger} & -iD_2 + \sqrt{2\tau}B \end{bmatrix}^{N} \right\rangle_{\substack{A, B \in \text{GUE}_k \\ C \in \text{GinUE}_k}} , (5.26)
$$

where $D_1 = \text{diag}(z_1, \ldots, z_k)$ and $D_2 = \text{diag}(w_1, \ldots, w_k)$. In the GinUE case $\tau = 0$, this agrees with [\(5.18\)](#page-34-3). Also, taking the limit $w_1, \ldots, w_k \to \infty$ gives agreement with the GUE result [\(2.9\)](#page-5-0).

 \Box

Inspection of the proof of Proposition [5.2](#page-34-2) shows that a key role is played by the eigenvalue distribution of ZZ^{\dagger} belonging to a classical Hermitian ensemble (Laguerre ensemble), which moreover permit the structured formula [\(5.20\)](#page-34-5) for the Jack polynomial average. In addition, to identify the evaluation of the LHS of the duality identity in the form of [\(5.21\)](#page-34-4) with the RHS, it is crucial that the Jack polynomial average contains as a factor $P_{\kappa}^{(1)}((1)^N)[N]_{\kappa}^{(1)}$.

The first of these requirements is met by the ensemble of $N \times N$ non-Hermitian matrices formed by deleting K rows and K columns from an $(N+K) \times (N+K)$ unitary matrix chosen with Haar measure. We will denote this ensemble by $\text{TrUE}_{(N,K)}$. One has that the distribution of squared singular values are the case $a_1 = 0$ and $a_2 = K - N$ of the Jacobi ensemble $ME_{2,N}[x^{a_1}(1 (x)^{a_2}$ [\[89\]](#page-50-16). This requirement is met too by the so-called complex spherical ensemble of non-Hermitian matrices $G_1^{-1}G_2$, where G_1, G_2 are independent GinUE matrices, as it is by the generalisation of the complex spherical ensemble defined by the construction $(G^{\dagger}G)^{-1/2}X$, where G is an $(N + K) \times N$ complex standard Gaussian matrix, and X is a GinUE matrix (the spherical ensemble corresponds to the case $K = 0$). Denoting this latter ensemble by $\text{StUE}_{(N,K)}$, one has that the squared singular values are the case $b_1 = 0$, $b_2 = K$, of the so-called Jacobi prime ensemble [\[66,](#page-49-17) §2.4] (a particular linear fractional transformation of the Jacobi ensemble) $ME_{2,N}[x^{b_1}(1+x)^{-b_2-2N}]$, supported on $x > 0$ [\[109\]](#page-51-19), [\[54,](#page-49-1) Exercises 3.6 q.3].

It is also the case that the Jacobi and Jacobi prime ensembles admit structured formulas for averages of Jack polynomials. Thus we have [\[102\]](#page-51-20), [\[104\]](#page-51-4) (see [\[54,](#page-49-1) Eq. (12.143)])

$$
\langle P_{\kappa}^{(\alpha)}(\mathbf{x})\rangle_{\mathrm{ME}_{2/\alpha,N}[\lambda^{a_1}(1-\lambda)^{a_2}]} = P_{\kappa}^{(\alpha)}((1)^N) \frac{[a_1+1+(N-1)/\alpha]_{\kappa}^{(\alpha)}}{[a_1+a_2+2+2(N-1)/\alpha]_{\kappa}^{(\alpha)}},\tag{5.27}
$$

and [\[142\]](#page-53-7), [\[47\]](#page-48-17)

$$
\langle P_{\kappa}^{(\alpha)}(\mathbf{x}) \rangle_{\text{ME}_{2/\alpha,N}[\lambda^{b_1}(1+\lambda)^{-b_1-b_2-2-(N-1)/\alpha}]} = P_{\kappa}^{(\alpha)}((-1)^N) \frac{[b_1+1+(N-1)/\alpha]_{\kappa}^{(\alpha)}}{[-b_2]_{\kappa}^{(\alpha)}}. \tag{5.28}
$$

It fact seeking a duality identity analogous to (5.18) for TrUE $_{(N,K)}$ gives rise to an average involving $\text{StUE}_{(N',K')}$ for certain N', K' [\[134\]](#page-52-17). Similarly for the case of starting with the average of a product of characteristic polynomials over $\text{StUE}_{(N,K)}$.

Proposition 5.3. We have

$$
\left\langle \prod_{l=1}^{k} \det(z_{l} \mathbb{I}_{N} - Z) \det(w_{l} \mathbb{I}_{N} - \bar{Z}) \right\rangle_{Z \in \text{TrUE}_{(N,K)}} = \left\langle \det \begin{bmatrix} D_{1} & -Y \\ Y^{\dagger} & D_{2} \end{bmatrix}^{N} \right\rangle_{Y \in \text{StrUE}_{(k,N+K)}}, \quad (5.29)
$$

where D_1, D_2 are diagonal matrices with diagonal entries **z** and **w** respectively. Also

$$
\left\langle \prod_{l=1}^{k} \det(z_{l} \mathbb{I}_{N} - Z) \det(w_{l} \mathbb{I}_{N} - \bar{Z}) \right\rangle_{Z \in \text{StUE}_{(N,K)}} = \left\langle \det \begin{bmatrix} D_{1} & -Y \\ Y^{\dagger} & D_{2} \end{bmatrix}^{N} \right\rangle_{Y \in \text{TrUE}_{(k,K-k)}}, \quad (5.30)
$$

where it is required $K \geq k$.

Proof. Following the working which lead to (5.21) , with use of (5.27) replacing use of (5.20) , shows that the LHS is equal to

$$
\prod_{l=1}^{k} (w_l z_l)^N \sum_{\mu \subseteq (N)^k} s_{\mu} (1/\mathbf{z}) s_{\mu} (1/\mathbf{w}) \frac{[N]_{\mu'}^{(1)}}{[N+K]_{\mu'}^{(1)}}.
$$
\n(5.31)

In this we replace the sum over $\mu \subseteq (N)^k$ by the sum over $\mu \subseteq (k)^N$, which is valid provided we replace μ by μ' in the summand. We then return to the identity [\(5.21\)](#page-34-4), now with $N \mapsto k$ to substitute for $s_{\mu'}(1/\mathbf{z})s_{\mu'}(1/\mathbf{w})$ analogous to the strategy of the proof of Proposition [5.2](#page-34-2) (this requires too further use of [\(5.27\)](#page-36-0), now with $\alpha = 1, b_2 = K + N$). Further use of the dual Cauchy identity [\(3.13\)](#page-12-3) with $\alpha = 1$, and use too of [\(5.25\)](#page-35-2) identifies the resulting expression with [\(5.31\)](#page-36-1). This line of working provides too the derivation of [\(5.30\)](#page-36-2). \Box

Consider the special case of [\(5.18\)](#page-34-3) $D_1 = z\mathbb{I}_k, D_2 = \bar{z}\mathbb{I}_k$, with [\(5.25\)](#page-35-2) used to rewrite the determinant on the RHS, and then with the change of variables $W = YY^{\dagger}$. The duality now takes the form [\[75,](#page-50-15) Eq. (5.6) with $\Sigma = \mathbb{I}_N$]

$$
\langle |\det(z\mathbb{I}_N - Z)|^{2k} \rangle_{Z \in \text{GinUE}_N} = |z|^{2k} \langle \det \left(\mathbb{I}_k + \frac{1}{|z|^2} W \right)^N \rangle_{W \in \text{ME}_{2,k}[e^{-x}]},\tag{5.32}
$$

thus relating a non-Hermitian average to one over an Hermitian ensemble. With k chosen to be proportional to N , this identity has recently been used to study the large N form of the Coulomb gas partition function interpretation of the LHS [\[30\]](#page-48-18); it was used earlier in [\[33\]](#page-48-19) to determine the asymptotics with k fixed, with the result

$$
\langle |\det(z\mathbb{I}_N - N^{-1/2}Z)|^{2k} \rangle_{Z \in \text{GinUE}_N} = N^{k^2/2} e^{kN(|z|^2 - 1)} \frac{(2\pi)^{k/2}}{G(1+k)} \Big(1 + o(1) \Big), \quad |z| < 1,\tag{5.33}
$$

where $G(z)$ denotes the Barnes G-function. From [\[143\]](#page-53-8) this is known to be valid for continuous $k > -1$, while [\[33\]](#page-48-19) also gives an extension of [\(5.33\)](#page-37-0) to the case that $|1-z|\sqrt{N}$ is of order unity.

Identities analogous to (5.32) follow from (5.29) and (5.30) . In the case of TrUE_{N,K}, with $N/(N + K) =: \mu$, an application is the asymptotic formula analogous to [\(5.33\)](#page-37-0) [\[135\]](#page-52-18), [\[133\]](#page-52-16)

$$
\langle |\det(z\mathbb{I}_N - Z)|^{2k} \rangle_{Z \in \text{TrUE}_{N,K}} = N^{k^2/2} \mu^{Nk} \Big(\frac{1 - \mu}{1 - |z|^2} \Big)^{Nk(1 - 1/\mu)} \Big(\frac{\sqrt{1 - \mu}}{1 - |z|^2} \Big)^{k^2} \frac{(2\pi)^{k/2}}{G(1 + k)} \Big(1 + o(1) \Big), \quad |z| < |\mu|^{1/2}, \tag{5.34}
$$

(note that the constant factor is the same as in [\(5.33\)](#page-37-0)). Potential theoretic aspects of the Coulomb gas interpretation of the LHS of (5.34) in the case that k is proportional to N is the subject of the recent work [\[26\]](#page-48-20).

The duality formula [\[75,](#page-50-15) Eq. (5.6)] for $\Sigma = A^{\dagger} A \neq \mathbb{I}_N$ referred to in the above paragraph in relation to [\(5.32\)](#page-37-1) is distinct from [\(5.18\)](#page-34-3), and moreover permits analogues that are distinct from those of Proposition [5.3.](#page-36-4) There are two stages to such identities. The first is to express the averages in terms of Jack polynomial based hypergeometric functions, and the second is the alternative form of the latter as random matrix averages over ensembles of size r.

Proposition 5.4. We have

$$
\langle |\det(x\mathbb{I}_N - AX)|^{2r} \rangle_{X \in \text{GinUE}_N} = |x|^{2r} \cdot 2F_0^{(1)}(-r, -r; |x|^{-2}AA^{\dagger})
$$

$$
\langle |\det(x\mathbb{I}_N - AX)|^{2r} \rangle_{X \in \text{TrUE}_{N,K}} = |x|^{2r} \cdot 2F_1^{(1)}(-r, -r; N + K; |x|^{-2}AA^{\dagger})
$$

$$
\langle |\det(x\mathbb{I}_N - AX)|^{2r} \rangle_{X \in \text{SrUE}_{N,K}} = |x|^{2r} \cdot 2F_1^{(1)}(-r, -r; -K; -|x|^{-2}AA^{\dagger}), \quad (K > r).
$$
 (5.35)

Consequently

$$
\langle |\det(x\mathbb{I}_N - AX)|^{2r} \rangle_{X \in \text{GinUE}_N} = \langle \prod_{l=1}^r \det(|x|^2 \mathbb{I}_N + t_l A A^{\dagger}) \rangle_{\mathbf{t} \in \text{ME}_{2,r}[e^{-\lambda}]}
$$

$$
\langle |\det(x\mathbb{I}_N - AX)|^{2r} \rangle_{X \in \text{TrUE}_{N,K}} \propto \det(|x|^2 \mathbb{I}_N - AA^{\dagger})^r \langle \prod_{l=1}^r \det(t_l \mathbb{I}_N - \Lambda) \rangle_{\mathbf{t} \in \text{ME}_{2,r}[\lambda^K \mathbb{1}_{0 < \lambda < 1}]}
$$

$$
\langle |\det(x\mathbb{I}_N - AX)|^{2r} \rangle_{X \in \text{SrUE}_{N,K}} = \langle \prod_{l=1}^r \det(|x|^2 \mathbb{I}_N + t_l A A^{\dagger}) \rangle_{\mathbf{t} \in \text{ME}_{2,r}[(1-\lambda)^{K-2r} \mathbb{1}_{0 < \lambda < 1}]}, \quad (5.36)
$$

where $\Lambda = -AA^{\dagger}(|x|^2 - AA^{\dagger})^{-1}$, and in the final equation it is required that $K \geq 2r$.

Proof. The derivation of [\(5.35\)](#page-37-3) is similar in all three cases. For definiteness we will consider the first only. Since the parameter x can be inserted by scaling Σ , it suffices to consider the case $x = 1$. With this done, we write $|\det(xI_M - AX)|^{2r} = \det(I_M - AX)^r \det(I_M - A^{\dagger}X^{\dagger})^r$, and then expand each of these powers of characteristic polynomials according to

$$
\det(\mathbb{I}_N - Q)^r = \sum_{\kappa \subseteq (N)^r} \frac{[-r]_{\kappa}^{(1)}}{|\kappa|!} C_{\kappa}^{(1)}(Q),\tag{5.37}
$$

(as implied by [\(3.20\)](#page-13-0) with $\alpha = 1$) with parameters appropriately identified. The computation of the average is therefore reduced to computing $\langle C_{\kappa}^{(1)}(AX)C_{\kappa}^{(1)}(A^{\dagger}X^{\dagger})\rangle_{X\in\text{GinUE}_N}$, which we do using [\(5.13\)](#page-33-1) to further reduce the problem to computing the average $\langle C_{\kappa}^{(1)}(XX^{\dagger})\rangle_{X\in \text{GinUE}_N}$. This latter average has appeared in the proof of Proposition [5.2,](#page-34-2) with its method of computation using [\(5.20\)](#page-34-5) detailed therein. Use of [\(5.6\)](#page-32-3),

$$
\frac{[k]_{\kappa}^{(1)}}{C_{\kappa}^{(1)}((1)^k)} = \frac{1}{|\kappa|!} (h_{\kappa}'|_{\alpha=1})^2
$$
\n(5.38)

(which is equivalent to [\(5.23\)](#page-35-3)) and [\(3.17\)](#page-13-4) then allows the resulting expression for the LHS of the first average in [\(5.35\)](#page-37-3) can be identified with the Jack polynomial based hypergeometric function on the RHS.

The task of going from (5.35) to (5.36) is to identify the Jack polynomial based hypergeometric functions with averages over matrix ensembles of size r . In the case of the first appearing ${}_2F_1^{(1)}$ this is done by applying the transformation [\(3.23\)](#page-14-2) and then making use of the theory in 1 the first paragraph of the proof of Proposition [3.3](#page-12-4) with $N \mapsto r$, $p \mapsto N$ in [\(3.18\)](#page-13-5). For the second appearing ${}_2F_1^{(1)}$ $1^{(1)}$, we consider its series form [\(3.17\)](#page-13-4) with the sum over κ replaced by a sum over κ' . The identity

$$
[u]_{\kappa}^{(\alpha)} = (-\alpha)^{-|\kappa|} [-\alpha u]_{\kappa'}^{(1/\alpha)}, \tag{5.39}
$$

can then be used to rewrite the generalised Pochhammer symbols back in terms of κ . Also, we have that [\[75,](#page-50-15) displayed equation above Eq. (5.4) and Eq. (3.18)]

$$
h'_{\kappa'} = \frac{[r\alpha]_{\kappa}^{(1/\alpha)}}{P_{\kappa}^{(1/\alpha)}((1)^{r})}
$$
(5.40)

(note that this is independent of r for the number of parts of κ' less that or equal to r). Hence

$$
{}_{2}F_{1}^{(\alpha)}(-r, -b; -K; -AA^{\dagger}) = \sum_{\kappa' \subseteq (N)^{r}} \frac{[\alpha b]_{\kappa}^{(1/\alpha)}}{[\alpha K]_{\kappa}^{(1/\alpha)}} P_{\kappa}^{(1/\alpha)}((1)^{r}) P_{\kappa'}^{(\alpha)}(AA^{\dagger}). \tag{5.41}
$$

Except for the factor of $(-1)^{|\kappa|}$ we see from [\(5.27\)](#page-36-0) that the prefactors of $P_{\kappa}^{\alpha}(AA^{\dagger})$ in this expression can be replaced by the average over $ME_{2/\alpha,N}[\lambda^{a_1}(1-\lambda)^{a_2}]$ with $\alpha \mapsto 1/\alpha$, $N = r$, $a_1 = -1 + (b - r + 1)\alpha$ and $a_2 = -1 + (K - b - r + 1)\alpha$. After using this as a substitution, the sum over κ' can be carried out according to the dual Cauchy identity [\(3.13\)](#page-12-3) to give the ensemble average form

$$
{}_{2}F_{1}^{(\alpha)}(-r,-b;-K;AA^{\dagger}) = \left\langle \prod_{l=1}^{r} \det(\mathbb{I}_{N} - x_{l}AA^{\dagger}) \right\rangle_{\mathbf{x} \in \text{ME}_{2\alpha,r}[\lambda^{a_{1}}(1-\lambda)^{a_{2}}]} \Big|_{\substack{a_{1}=-1+(b-r+1)\alpha \\ a_{2}=-1+(K-b-r+1)\alpha}}, (5.42)
$$

valid provided $a_1, a_2 > -1$. Application of this gives the final duality in [\(5.36\)](#page-38-0).

In the case of the evaluation in terms of ${}_2F_0^{(1)}$ \int_0^{1} , which is the first identity in [\(5.35\)](#page-37-3), one has available the identity [\[18\]](#page-47-16)

$$
{}_{2}F_{0}^{(\alpha)}(-a, 1+b-a+(n-1)/\alpha; x_{1}^{-1}, \ldots, x_{n}^{-1}) \propto (x_{1} \cdots x_{n})^{-a} {}_{1}F_{1}^{(\alpha)}(-a;-b;-x_{1}, \ldots, -x_{n}), \ (5.43)
$$

valid for $a \in \mathbb{Z}^+$ and $b > a$. Scaling $x_j \mapsto \beta x_j/2a_2$ and then taking the limit $a_2 \to \infty$ we have that ${}_{1}F_{1}^{(\alpha)}$ with the first parameter a negative integer can be written as the average

$$
{}_{1}F_{1}^{(\alpha)}(-a;(a_{1}+n)/\alpha;x_{1},\ldots,x_{n}) \propto \left\langle \prod_{j=1}^{n} \prod_{l=1}^{a}(x_{j}-y_{l}) \right\rangle_{\mathbf{y} \in \mathrm{ME}_{2\alpha,a}[x^{a_{1}}e^{-\beta x/2}]}.
$$
(5.44)

 \Box

Use of this with $\alpha = 1$, $a_1 = 0$, $a = n = r$ gives the first average in [\(5.36\)](#page-38-0).

Remark 5.2.

1. In the case $K = 0$, by its definition TrUE_{N,K} becomes equal to $U(N)$ with Haar measure. For this latter ensemble, a duality identity of Fyodorov and Khoruzhenko [\[86\]](#page-50-0) states

$$
\langle |\det(x\mathbb{I}_N - AU)|^{2r} \rangle_{U \in U(N)} \propto \langle \prod_{l=1}^r \det(|x|^2 \mathbb{I}_N + t_l A A^{\dagger}) \rangle_{\mathbf{t} \in \text{ME}_{2,r}[(1+\lambda)^{-N-2r}]}.
$$
(5.45)

After a change of variables $t_l/(1 + t_l) \mapsto t_l$ on the RHS, this can be checked to agree with the case $K = 0$ of the second duality relation in [\(5.36\)](#page-38-0). When A is a multiplicative rank 1 perturbation of the identity, $A = diag(a, 1, \ldots, 1)$ with $|a| < 1$, the eigenvalues of AU relate to the point process for the zeros of the random power series $1/\mu - \sum_{j=1}^{N} c_j z^j$ where each c_j is an independent standard complex Gaussian [\[67\]](#page-49-18) (this requires setting $a = 1/(\mu\sqrt{N})$ and taking $N \to \infty$). This point process is further studied in the recent work [\[88\]](#page-50-17).

2. One can check that the third identity in [\(5.36\)](#page-38-0) with $A = \mathbb{I}_N$ is consistent with the case $z_1 = \cdots = z_k = z, w_1 = \cdots = w_k = \overline{z}$ of [\(5.30\)](#page-36-2). Also, as with the first two identities in [\(5.36\)](#page-38-0), setting $A = \mathbb{I}_N$ allows for a Coulomb gas interpretation. Aspects of this in the case r proportional to N have been the subject of the recent work [\[28\]](#page-48-21).

5.3.2 Real and quaternion entry matrices

Under this heading attention will first be focussed on powers of characteristic polynomial averages extending Proposition [5.4.](#page-37-4) As some preliminaries, we recall the notation GinOE_N (GinSE_N) for $N \times N$ standard real (quaternion) Gaussian random matrices. We denote by $TroE_{N,K}$ (TrSE_{N,K}) the ensemble of $N \times N$ non-Hermitian matrices formed by deleting K rows and K columns from an $(N+K) \times (N+K)$ unitary matrix with real (quaternion) elements chosen with Haar measure. Also, we use the notation $SroE_{N,K}$ ($SrSE_{N,K}$) for the ensemble of matrices of the form

 $(G^{\dagger}G)^{-1/2}X$, where G is an $(N+K) \times N$ real (quaternion) standard Gaussian matrix, and X is a GinOE (GinSE) matrix. A feature of the eigenvalues in both the real and quaternion cases is that they appear in complex conjugate pairs, and typically in the real case some eigenvalues will also be real [\[43\]](#page-48-16). Further, in the quaternion case, when we define the characteristic polynomial $\det(x\mathbb{I}_N - B)$ of a matrix B, the viewpoint is that entries are quaternions in both \mathbb{I}_N and B. Thus as a complex matrix the size of $x\mathbb{I}_N - B$ is $2N \times 2N$, and the characteristic polynomial is of degree 2N.

Proposition 5.5. ([\[75\]](#page-50-15) in relation to the first two equations and [\[134\]](#page-52-17) for the next two.) We have

$$
\langle \det(xI_N - AX)^r \rangle_{X \in \text{GinoE}_N} = x^{rN} {}_2F_0^{(2)}(-r/2, (-r+1)/2; 2|x|^{-2}AA^{\dagger})
$$

\n
$$
\langle \det(xI_N - AX)^r \rangle_{X \in \text{GinsE}_N} = x^{2rN} {}_2F_0^{(1/2)}(-r, -r-1; |x|^{-2}AA^{\dagger}/2)
$$

\n
$$
\langle \det(xI_N - AX)^r \rangle_{X \in \text{TrOE}_{N,K}} = x^{rN} {}_2F_1^{(2)}(-r/2, (-r+1)/2; (N+K)/2; |x|^{-2}AA^{\dagger})
$$

\n
$$
\langle \det(xI_N - AX)^r \rangle_{X \in \text{TrSE}_{N,K}} = x^{2rN} {}_2F_1^{(1/2)}(-r, -r-1; 2(N+K); |x|^{-2}AA^{\dagger})
$$

\n
$$
\langle \det(xI_N - AX)^r \rangle_{X \in \text{SrOE}_{N,K}} = x^{rN} {}_2F_1^{(2)}(-r/2, (-r+1)/2; -(K-1)/2; -|x|^{-2}AA^{\dagger})
$$

\n
$$
\langle \det(xI_N - AX)^r \rangle_{X \in \text{SrSE}_{N,K}} = x^{2rN} {}_2F_1^{(1/2)}(-r, -r-1; -2K-1; -|x|^{-2}AA^{\dagger}), \qquad (5.46)
$$

where in the last two identities we require $K - 1 > r$ and $2K + 1 > r$ respectively.

Proof. The derivations begin with the use of [\(5.37\)](#page-38-1), $(N, r) \mapsto (2N, r)$ in the quaternion case. Also, we replace $C_{\kappa}^{(1)}(\mathbf{x})$ by $|\kappa|!s_{\kappa}(\mathbf{x})/h_{\kappa}'|_{\alpha=1}$ as is consistent with [\(5.6\)](#page-32-3). The next step is to make use of (5.12) or (5.14) as appropriate, together with the key identities [\[17\]](#page-47-12), [\[75\]](#page-50-15)

$$
\frac{1}{|\kappa|!2^{|\kappa|}}h'_{2\kappa}|_{\alpha=1} = \frac{2^{|\kappa|}[N/2]_{\kappa}^{(2)}}{C_{\kappa}^{(2)}((1)^{N})}, \qquad \frac{1}{|\kappa|!2^{|\kappa|}}h'_{\kappa^2}|_{\alpha=1} = \frac{[2N]_{\kappa}^{(1/2)}}{2^{|\kappa|}C_{\kappa}^{(1/2)}((1)^{N})},
$$

\n
$$
[u]_{2\kappa}^{(1)} = 2^{2|\kappa|}[u/2]_{\kappa}^{(2)}[(u+1)/2]_{\kappa}^{(2)}, \qquad [u]_{\kappa^2}^{(1)} = [u]_{\kappa}^{(1/2)}[u-1]_{\kappa}^{(1/2)},
$$
\n(5.47)

inter-relating Jack polynomial quantities for $\alpha = 1$ with $\alpha = 2, 1/2$. This reduces the task to computing $\langle P_{\kappa}^{(2/\beta)}(XX^{\dagger})\rangle$ for the ensemble of interest, which in turn, after a change of variables, reduces the problem to computing $\langle P_{\kappa}^{(2/\beta)}(\mathbf{x})\rangle$, where the average now is over the corresponding squared singular values. For $\widehat{\text{GinOE}_N}$ ($\widehat{\text{GinSE}_N}$) the squared singular values belong to $ME_{\beta,N}[\lambda^{\beta/2-1}e^{-\beta\lambda/2}]$ with $\beta = 1$ ($\beta = 4$) [\[54,](#page-49-1) case $n-m$ of (3.16)]; for TrOE_N (TrSE_N) the squared singular values belong to $ME_{\beta,N}[\lambda^{\beta/2-1}(1-\lambda)^{(\beta/2)(K-N+1)-1}]$ with $\beta=1$ ($\beta=4$) [\[54,](#page-49-1) Eq. (3.113) with $N \mapsto N + K$, $n_1, n_2 \mapsto N$]; SrOE_N (SrSE_N) the squared singular values belong to $\text{ME}_{\beta,N}[\lambda^{\beta/2-1}(1+\lambda)^{-(\beta/2)(K+2N)}]$ with $\beta=1$ ($\beta=4$) [\[54,](#page-49-1) Eq. (3.81)]. Consequently the sought averages of $P_{\kappa}^{(2/\beta)}(\mathbf{x})$ over the squared singular values can be computed by making use of [\(5.20\)](#page-34-5) or [\(5.28\)](#page-36-5) as appropriate. Assembling the results, the series form of various Jack polynomial based hypergeometric functions with $\alpha = 2$ (real case) and $\alpha = 1/2$ (quaternion case) are recognised. \Box

We can now proceed to express the averages given in terms of Jack polynomial based hypergeometric functions in the above proposition, in terms of random matrix averages of sizes relating to the power according to the formulas identified in [\(5.42\)](#page-39-1), [\(5.43\)](#page-39-2) and [\(5.44\)](#page-39-3).

Proposition 5.6. ([\[75\]](#page-50-15) in relation to the Ginibre ensembles and [\[134\]](#page-52-17) in relation to the truncated ensembles.) Let Λ be as in Proposition [5.4.](#page-37-4) We have

$$
\langle \det(\mathbb{I}_N - AX)^r \rangle_{X \in \text{GinOE}_N} = \Big\langle \prod_{l=1}^{\lfloor r/2 \rfloor} \det(\mathbb{I}_N + t_l A A^{\dagger}) \Big\rangle_{\mathbf{t} \in \text{ME}_{4, \lfloor r/2 \rfloor}[\lambda^{\nu} e^{-\lambda}]}
$$

$$
\langle \det(\mathbb{I}_N - AX)^r \rangle_{X \in \text{TrOE}_{N,K}} \propto \det(\mathbb{I}_N - AA^{\dagger})^{\lfloor r/2 \rfloor} \Big\langle \prod_{l=1}^{\lfloor r/2 \rfloor} \det(t_l \mathbb{I}_N - \Lambda) \Big\rangle_{\mathbf{t} \in \text{ME}_{4, \lfloor r/2 \rfloor}[\lambda^K(1-\lambda)^{\nu}]}
$$

$$
\langle \det(\mathbb{I}_N - AX)^r \rangle_{X \in \mathrm{SroE}_{N,K}} = \langle \prod_{l=1}^{\lfloor t/2 \rfloor} \det(\mathbb{I}_N + t_l A A^{\dagger}) \rangle_{\mathbf{t} \in \mathrm{ME}_{4, \lfloor r/2 \rfloor}[\lambda^{\nu}(1-\lambda)^{K-2r+1})]},\tag{5.48}
$$

where $\nu = 0$ (r even) and $\nu = 2$ (r odd). Also

$$
\langle \det(\mathbb{I}_{N} - AX)^{r} \rangle_{X \in \text{GinsE}_{N}} = \left\langle \prod_{l=1}^{r} \det(\mathbb{I}_{N} + t_{l}AA^{\dagger}) \right\rangle_{\mathbf{t} \in \text{ME}_{1,r}[e^{-\lambda}]}
$$

$$
\langle \det(\mathbb{I}_{N} - AX)^{r} \rangle_{X \in \text{TrSE}_{N,K}} \propto \det(\mathbb{I}_{N} - AA^{\dagger})^{r} \left\langle \prod_{l=1}^{r} \det(t_{l} \mathbb{I}_{N} - \Lambda) \right\rangle_{\mathbf{t} \in \text{ME}_{1,r}[X^{K} \mathbb{I}_{0 < \lambda < 1}]}
$$

$$
\langle \det(\mathbb{I}_{N} - AX)^{r} \rangle_{X \in \text{SrSE}_{N,K}} = \left\langle \prod_{l=1}^{r} \det(\mathbb{I}_{N} + t_{l}AA^{\dagger}) \right\rangle_{\mathbf{t} \in \text{ME}_{1,r}[(1-\lambda)^{K-r-1/2}]}.
$$
(5.49)

Remark 5.3. We note that the first identity in (5.48) with $r = 2$ is consistent with (5.2) .

We next consider analogues of the dualities for products of non-Hermitian random matrix ensembles with complex entries [\(5.18\)](#page-34-3), [\(5.29\)](#page-36-3) and [\(5.30\)](#page-36-2), in the case of real entries or quaternion entries. In the case of GinUE in [\(5.18\)](#page-34-3) we saw that the RHS also considered of an average over a particular GinUE. In the case of GinOE (GinSE) it turns out that the duality gives rise to an average over complex anti-symmetric (complex symmetric) matrices [\[140\]](#page-53-9), [\[124\]](#page-52-13), [\[133\]](#page-52-16). Let the set of these latter matrices be denoted $\mathcal{A}_N(\mathbb{C})$ and $\mathcal{S}_N(\mathbb{C})$ respectively. Fundamental in their derivation are integration formulas over zonal polynomials analogous to [\(5.12\)](#page-33-1) and [\(5.14\)](#page-33-1) [\[47\]](#page-48-17) (see too [\[133,](#page-52-16) pg. 44] for a clear statement).

Proposition 5.7. Let A be an $N \times N$ matrix. For $X \in S_N(\mathbb{C})$ chosen from a measure unchanged by $X \mapsto UXU^T$, $U \in U(N)$ we have

$$
\left\langle C_{\kappa}^{(2)}(AXA^TX^\dagger)\right\rangle_X = \frac{s_{2\kappa}(A)}{s_{2\kappa}((1)^N)} \langle C_{\kappa}^{(2)}(XX^\dagger)\rangle_X. \tag{5.50}
$$

Also, for $X \in \mathcal{A}_N(\mathbb{C})$ chosen from a measure unchanged by $X \mapsto UXU^T$, $U \in U(N)$ we have

$$
\left\langle C_{\kappa}^{(1/2)}(AXA^TX^\dagger)\right\rangle_X = \frac{s_{\kappa^2}(A)}{s_{\kappa^2}((1)^N)} \langle C_{\kappa}^{(1/2)}(XX^\dagger)\rangle_X\tag{5.51}
$$

(note that here the matrices in $C_{\kappa}^{(1/2)}(\cdot)$ for N even are doubly degenerate and so are only to be included once, and similarly for N odd apart from a zero eigenvalue which also is not included).

Corollary 5.2. Denote by GA_N (GS_N) the set of $N \times N$ anti-symmetric complex (symmetric complex) matrices chosen with measure proportional to $e^{-\text{Tr}XX^{\dagger}/2}$ ($e^{-\text{Tr}XX^{\dagger}}$). We have

$$
\left\langle \prod_{j=1}^{2k} \det(G - z_j \mathbb{I}_N) \right\rangle_{G \in \text{GinOE}_N} = \left\langle \det \begin{bmatrix} X & Z \\ -Z & X^{\dagger} \end{bmatrix}^{N/2} \right\rangle_{X \in \text{G} \mathcal{A}_{2k}}
$$
(5.52)

and

$$
\left\langle \prod_{j=1}^{2k} \det(G - z_j \mathbb{I}_N) \right\rangle_{G \in \text{GinsE}_N} = \left\langle \det \begin{bmatrix} X & Z \\ -Z & X^{\dagger} \end{bmatrix}^N \right\rangle_{X \in \text{G} \mathcal{S}_{2k}}.
$$
 (5.53)

Proof. Following [\[133\]](#page-52-16), we adopt a strategy analogous to that used in the proof of Proposition [5.2,](#page-34-2) giving the details only of [\(5.52\)](#page-41-1). Consider the LHS. Making use of the dual Cauchy identity [\(3.13\)](#page-12-3) with $\alpha = 1$, then [\(5.12\)](#page-33-1) with $A = Z^{-1}$ shows that it can be written

$$
\prod_{l=1}^{2k} z_l^N \sum_{\kappa \subseteq (2k)^N} s_{\kappa'}(1/\mathbf{z}) \delta_{\kappa=2\mu} \frac{\langle C_{\mu}^{(2)}(GG^T) \rangle_{G \in \text{GinoE}_N}}{C_{\mu}^{(2)}((1)^N)} \n= \prod_{l=1}^{2k} z_l^N \sum_{\kappa \subseteq (2k)^N} s_{\kappa'}(1/\mathbf{z}) \delta_{\kappa=2\mu} 2^{|\mu|} [N/2]_{\mu}^{(2)} = \prod_{l=1}^{2k} z_l^N \sum_{\mu \subseteq (N)^k} s_{\mu^2}(1/\mathbf{z}) 2^{|\mu|} [N/2]_{\mu'}^{(2)}, \quad (5.54)
$$

where the first equality follows from the fact that the eigenvalues of GG^T belong to the matrix ensemble $\text{ME}_{1,N}[\lambda^{-1/2}e^{-\lambda/2}]$ (recall the text below [\(5.47\)](#page-40-0)) and use of [\(5.20\)](#page-34-5) with $\alpha = 2, \alpha =$ $-1/2$, while the second equality results upon replace μ by μ' in the summand.

The next step is to make use of (5.51) with $N = k$, for the purpose of substituting for $s_{\mu^2}(1/\mathbf{z})$. In this $\langle C_{\mu}^{(1/2)}(XX^{\dagger})\rangle_{X\in\text{G}\mathcal{A}_{2k}} = C_{\mu}^{(1/2)}((1)^k)[2k-1]_{\mu}^{(1/2)}$ as follows from [\(5.20\)](#page-34-5) and the fact that for $X \in \widehat{G\mathcal{A}}_{2k}(\mathbb{C})$, the eigenvalues of XX^{\dagger} are doubly degenerate, with independent eigenvalues distributed according to $ME_{4,k}[e^{-\lambda}]$ [\[54,](#page-49-1) Table 3.1]. Noting the formulas

$$
\frac{s_{\mu^2}((1)^k)}{C_{\mu}^{(1/2)}((1)^k)[2k-1]_{\mu}^{(1/2)}} = \frac{1}{|\mu|!}, \qquad P_{\mu'}^{(2)}((1)^N) = \frac{[N/2]_{\mu'}^{(2)}}{h'_{\mu}|_{\alpha=1/2}},
$$

(see [\[133,](#page-52-16) Eq. (2.62)] for the first, and [\[54,](#page-49-1) Eq. (13.8) together with $h_{\kappa'} = \alpha^{|\kappa|} h'_{\kappa}|_{\alpha \mapsto 1/\alpha}$ as noted three lines below] for the second), and making use too of (5.6) then shows that (5.54) is equal to

$$
\prod_{l=1}^{2k} z_l^N \sum_{\mu \subseteq (N)^k} P_{\mu'}^{(2)}((1)^N) \langle P_{\mu}^{(1/2)}(Z^{-1}XZ^{-1}X^{\dagger}) \rangle_{X \in \text{GA}_{2k}}.
$$
\n(5.55)

The sum herein is carried out using the dual Cauchy identity [\(3.13\)](#page-12-3) with $\alpha = 1/2$, to obtain an expression analogous to (5.24) . To obtain the RHS of (5.52) from this requires the use of (5.25) , and the fact that the eigenvalues of the matrix in the latter are doubly degenerate. \Box

Remark 5.4.

1. Let A, B be $N \times N$ complex symmetric (anti-symmetric) matrices. Let $X \in \mathcal{S}_N(\mathbb{C})$ $(\mathcal{A}_N(\mathbb{C}))$ be chosen from a measure unchanged by $X \mapsto UXU^T$ for $U \in U(N)$. Then in addition to [\(5.50\)](#page-41-3) and [\(5.51\)](#page-41-2) one has [\[47\]](#page-48-17)

$$
\left\langle C_{\kappa}^{(2)}(AX)C_{\mu}^{(2)}(BX^{\dagger})\right\rangle_{X\in\mathcal{S}_{N}(\mathbb{C})} = \delta_{\kappa,\mu}\frac{C_{\kappa}^{(2)}(AB)}{s_{2\kappa}((1)^{N})}\langle C_{\kappa}^{(2)}(XX^{\dagger})\rangle_{X\in\mathcal{S}_{N}(\mathbb{C})}
$$

$$
\left\langle C_{\kappa}^{(1/2)}(AX)C_{\mu}^{(1/2)}(BX^{\dagger})\right\rangle_{X\in\mathcal{A}_{N}(\mathbb{C})} = \delta_{\kappa,\mu}\frac{C_{\kappa}^{(1/2)}(AB)}{s_{\kappa}^{2}((1)^{N})}\langle C_{\kappa}^{(1/2)}(XX^{\dagger})\rangle_{X\in\mathcal{A}_{N}(\mathbb{C})}.
$$
(5.56)

2. A duality which reverses the role of GS_{2k} and $G_{1}S_{N}$ in [\(5.53\)](#page-42-1), and its companion reversing

the roles of the ensembles on either sides of [\(5.52\)](#page-41-1), but with the Gaussian complex anti-symmetric matrices replaced by Gaussian complex self dual matrices (the latter are obtained from even size anti-symmetric matrices by premultiplying by $\mathbb{I}_N \otimes \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$), has attracted recent interest in the works [\[1\]](#page-46-2), [\[110\]](#page-51-21), [\[62\]](#page-49-19).

Documented in the literature is the analogue of Corollary [5.2](#page-41-4) for the non-Hermitian ensembles $TroE_{N,K}$ and $TrSE_{N,K}$. For this one requires the spherical anti-symmetric (symmetric) ensemble $S\mathcal{A}_{N,K}$ ($S\mathcal{S}_{N,K}$), defined to have joint element PDF proportional to

$$
\det(\mathbb{I}_N + XX^\dagger)^{-N-K},\tag{5.57}
$$

where X is anti-symmetric (symmetric).

Proposition 5.8. ([\[135\]](#page-52-18), [\[133\]](#page-52-16)) We have

$$
\left\langle \prod_{j=1}^{2k} \det(G - z_j \mathbb{I}_N) \right\rangle_{G \in \text{TPOE}_{N,K}} = \left\langle \det \begin{bmatrix} X & Z \\ -Z & X^\dagger \end{bmatrix}^{N/2} \right\rangle_{X \in \text{SA}_{2k, (N+K)/2 - 1}} \tag{5.58}
$$

and

$$
\left\langle \prod_{j=1}^{2k} \det(G - z_j \mathbb{I}_N) \right\rangle_{G \in \text{TrSE}_{N,K}} = \left\langle \det \begin{bmatrix} X & Z \\ -Z & X^{\dagger} \end{bmatrix}^N \right\rangle_{X \in \text{SS}_{2k,N+K+1}}.
$$
 (5.59)

Proof. The general strategy of the proof of Corollary [5.2](#page-41-4) suffices, with changes of detail as implied by the use of (5.28) rather than (5.20) . \Box

Remark 5.5. By taking the limit $z_{2k} \to \infty$ in the identities of Corollary [5.2](#page-41-4) and Proposition [5.8,](#page-43-1) they remain formally unchanged except for the replacement of $2k$ by $2k-1$ (odd number of products). Intermediate working, such as the in-text identity five lines below [\(5.54\)](#page-42-0) require modification if one was to not distinguish the parity from the outset.

6 Dualities for moments

Consider the class of Hermitian β ensembles $ME_{\beta,N}[e^{-N\beta V(x)/2}]$, where $V(x)$ is such that in the large N limit the support of the eigenvalue density is a compact, single interval. Classical examples of this setting are $V(x)$ equal to

$$
x^{2}, \qquad (x - \gamma \log x) \mathbb{1}_{x>0}, \qquad (\gamma_{1} \log x + \gamma_{2} \log(1-x)) \mathbb{1}_{0
$$

as discussed in the context of global scaling in §4.2.1 and §4.3.1.

With $m_{\kappa}(\mathbf{x})$ denoting the monomial symmetric polynomial in the variables x indexed by the partition κ , the mixed moments are specified by $m(\kappa; N, \beta) := \langle m_{\kappa}(\mathbf{x}) \rangle$. For example, with $\kappa =$ $(k,(0)^{N-1})$ one has that this average corresponds to the power sum of the eigenvalues $\langle \sum_{l=1}^{N} x_l^k \rangle$, which in turn is equal to the k-th moment of the spectral density. When κ consists of two or more nonzero parts, one defines from the mixed moments $\mu(\kappa; N, \beta)$ the corresponding mixed cumulants, defined in the usual way according to the logarithm of the exponential generating function of the former. For the cumulants specified by partitions of no more than n parts, the Laurent series generating function is specified by

$$
W_n(y_1,\ldots,y_n;N,\beta) := \frac{1}{y_1\cdots y_n} \sum_{p_1,\ldots,p_n=0}^{\infty} \frac{\mu((p_1,\ldots,p_n);N,\beta)}{y_1^{p_1}\cdots y_n^{p_n}};
$$
(6.2)

here after suitable ordering (p_1, \ldots, p_n) corresponds to the partition labelling the cumulant. Introducing the averages

$$
\left\langle \prod_{l=1}^{q} (A_l - \langle A_l \rangle) \right\rangle, \quad A_i = \sum_{j=1}^{N} \frac{1}{y_i - x_j},\tag{6.3}
$$

the generating function [\(6.2\)](#page-43-2) can be written as a multinomial in such terms for $q = 2, \ldots, n$ which is homogeneous of degree n in the total number of factors of the form $(A_l - \langle A_l \rangle)$. In fact for $n = 2$ and 3, [\(6.2\)](#page-43-2) is precisely equal to [\(6.3\)](#page-44-0) with $q = 2$ and 3. We will write $\langle \prod_{l=1}^{q} A_l \rangle_c$ to denote the linear combination of products of averages relating to the cumulants (the symbol "c" denotes connected, or fully truncated; here we omit the $\langle A_l \rangle$ in $(A - \langle A_l \rangle)$ which can be absorbed into the linear combination).

In addition to the quantity W_n [\(6.2\)](#page-43-2), introduce the connected quantity

$$
P_{n+1}(y_1, \dots, y_n; y; \beta, N) = \left\langle \prod_{l=1}^n A_l \sum_{l=1}^N \frac{V'(y) - V'(y_l)}{y - y_l} \right\rangle_c.
$$
 (6.4)

Introduce too the notation $I = \{y_1, \ldots, y_{n-1}\}.$ Together with the W_n [\(6.2\)](#page-43-2) these quantities satisfy the coupled hierarchy of equations, referred to as loop equations,

$$
(\beta/2) \sum_{J \subseteq I} W_{|J|+1}(y, J) W_{n-|J|}(y, I \setminus J) + (\beta/2) W_{n+1}(y, y, I) + (\beta/2 - 1) \frac{\partial W_n(y, I)}{\partial y}
$$

=
$$
\frac{\beta N}{2} \Big(V'(y) W_n(y, I) - P_n(I; y) \Big) - \sum_{y_i \in I} \frac{\partial}{\partial y_i} \Big(\frac{W_{n-1}(y, I \setminus \{y_i\}) - W_{n-1}(I)}{y - y_i} \Big); \quad (6.5)
$$

see [\[19\]](#page-47-17), [\[25\]](#page-48-22), [\[145,](#page-53-3) Eq. (2.5)]. For ease of presentation, the dependence of the W_n and P_n on N, β has been suppressed. The remaining step is to introduce the scaled quantity

$$
\widetilde{W}_n(y_1, \dots, y_n) := \frac{1}{N} (N\beta/2)^{n-1} W_n(y_1, \dots, y_n), \tag{6.6}
$$

which for large N is expected to be independent of N at leading order (for justifications, see [\[20\]](#page-47-18)), and furthermore independent of β . One similarly introduces a scaling of P_n with an analogous large N property. This gives the rewrite of (6.5)

$$
(\beta N/2) \sum_{J \subseteq I} \widetilde{W}_{|J|+1}(y,J) \widetilde{W}_{n-|J|}(y,I \setminus J) + \frac{1}{N} \widetilde{W}_{n+1}(y,y,I) + (\beta/2 - 1) \frac{\partial \widetilde{W}_{n}(y,I)}{\partial y}
$$

=
$$
\frac{\beta N}{2} \Big(V'(y) \widetilde{W}_{n}(y,I) - \widetilde{P}_{n}(I;y) \Big) - \frac{\beta N}{2} \sum_{y_i \in I} \frac{\partial}{\partial y_i} \Big(\frac{\widetilde{W}_{n-1}(y,I \setminus \{y_i\}) - \widetilde{W}_{n-1}(I)}{y - y_i} \Big). \tag{6.7}
$$

The following duality relation is now evident.

Proposition 6.1. ([\[46\]](#page-48-23), [\[59\]](#page-49-20)) The quantity $\widetilde{W}_n(y_1, \ldots, y_n) = \widetilde{W}_n(y_1, \ldots, y_n; N, \beta)$ is invariant under the mappings $\beta \mapsto 4/\beta$, $N \mapsto -\beta N/2$.

Proof. Upon the stated mappings, the loop equations [\(6.5\)](#page-44-1) remain formally unchanged. \Box

Remark 6.1.

1. One recalls that the meaning of the mapping in N is with respect to a $1/N$ expansion. It follows from Proposition [6.1](#page-44-2) that such an expansion for $\beta = 2$ involves only powers of $1/N^2$.

2. A viewpoint of the duality linking $\beta = 1$ and $\beta = 4$ as first identified in [\[123\]](#page-52-0) within the theory of Lie groups and symmetric spaces is given in [\[120\]](#page-52-19); see also [\[11\]](#page-47-19).

Corollary 6.1. The scaled moments (independent of N and β to leading order in N) $\widetilde{m}_k =$ $\widetilde{m}_k(\beta, N)$, and smoothed density $\widetilde{\rho}_{N,(1)}(x;\beta)$ satisfy the duality relations

$$
\widetilde{m}_k(\beta, N) = \widetilde{m}_k(4/\beta, -\beta N/2), \qquad \widetilde{\rho}_{N,(1)}(x; \beta) = \widetilde{\rho}_{-\beta N/2,(1)}(x; 4/\beta). \tag{6.8}
$$

The duality for the scaled moments is easy to illustrate. Thus, with $\tau := \beta/2$, for the Gaussian β ensemble one has [\[40\]](#page-48-0), [\[119\]](#page-52-20), [\[145\]](#page-53-3)

$$
\widetilde{m}_2^{\text{G}}(\beta, N) = 1 + N^{-1}(-1 + \tau^{-1}) \n\widetilde{m}_4^{\text{G}}(\beta, N) = 2 + 5N^{-1}(-1 + \tau^{-1}) + N^{-2}(3 - 5\tau^{-1} + 3\tau^{-2}) \n\widetilde{m}_6^{\text{G}}(\beta, N) = 5 + 22N^{-1}(-1 + \tau^{-1}) + N^{-2}(32 - 54\tau^{-1} + 32\tau^{-2}) \n+ N^{-3}(-15 + 32\tau^{-1} - 32\tau^{-2} + 15\tau^{-3}),
$$
\n(6.9)

and for the Laguerre β ensemble [\[117\]](#page-52-21), [\[74,](#page-50-18) Prop. 3.11],

$$
\widetilde{m}_{2}^{L}(\beta, N) = (1 + \gamma) + N^{-1}(-1 + \tau^{-1}) \n\widetilde{m}_{4}^{L}(\beta, N) = (2 + 3\gamma + \gamma^{2}) + N^{-1}(4 + 3\gamma)(-1 + \tau^{-1}) + 2N^{-2}(1 - 2\tau^{-1} + \tau^{-2}) \n\widetilde{m}_{6}^{L}(\beta, N) = (5 + 10\gamma + 6\gamma^{2} + \gamma^{3}) + N^{-1}(16 + 21\gamma + 6\gamma^{2})(-1 + \tau^{-1}) \n+ N^{-2}((17 + 11\gamma)\tau^{-2} - (33 + 21\gamma)\tau^{-1} + (17 + 11\gamma)) + N^{-3}(-1 + \tau^{-1})(-6 + 11\tau^{-1} - 6\tau^{-2}).
$$
\n(6.10)

An observation from [\[145\]](#page-53-3) is that the polynomials in τ^{-1} in the 1/N expansion [\(6.9\)](#page-45-0) have all their zeros on the unit circle and also exhibit an interlacing property. Explicit functional forms are also available for low order moments in the Jacobi β ensemble [\[117\]](#page-52-21), [\[74\]](#page-50-18), which exhibit a rational function, rather than a polynomial, structure. A consequence of the density duality in [\(6.8\)](#page-45-1) is that the 5th order linear differential equations which can be derived in the cases of the GOE and GSE are related to each other by the implied mapping [\[145\]](#page-53-3). In [\[128\]](#page-52-22) the 7th order linear differential equation satisfied by the soft edge scaled density for the Gaussian β ensemble with $\beta = 6$ is transformed to 7th order linear differential equation for the soft edge scaled density of the Gaussian β ensemble with $\beta = 2/3$.

It is known how to map from the Jacobi β ensemble to the (generalised) circular Jacobi β ensemble [\[76,](#page-50-14) Eq. (1.8)]. Consequently, there is an analogue of Proposition [6.1](#page-44-2) for the latter ensemble; see [\[54,](#page-49-1) Remark 4.3]. In relation to a consequence, denote by $\rho_{(2),\infty}((x,0);\beta)$ the bulk scaled (unit density) two-point correlation for the circular β ensemble. Define the structure function as the Fourier transform

$$
S(k; \beta) := \int_{-\infty}^{\infty} \left(\rho_{(2),\infty}^T((x,0); \beta) + \delta(x) \right) e^{ikx} dx, \tag{6.11}
$$

where $\rho_{(2),\infty}^T((x,0);\beta) := \rho_{(2),\infty}((x,0);\beta) - 1$. It is known that $f(k;\beta) := \pi \beta S(k;\beta)/|k|$ extends to an analytic function in the range $|k| < \min(2\pi, \pi\beta)$. Most significantly from the viewpoint of the present theme is that f exhibits the particular duality [\[73\]](#page-50-19)

$$
f(k; \beta) = f\left(-\frac{2k}{\beta}; \frac{4}{\beta}\right),\tag{6.12}
$$

which as discussed in $[76]$ is in keeping with the more general (but finite N) duality exhibited by the circular Jacobi β ensemble. We remark too that as is consistent with [\(6.12\)](#page-45-2), the small k expansion of [\(6.11\)](#page-45-3) has for the coefficient of k^p polynomials in $u = 2/\beta$ of degree p, which are palindromic or anti-palindromic [\[73,](#page-50-19) [58\]](#page-49-21); see also [\[76\]](#page-50-14) for observations relating to the zeros.

An important detail with regards to the results of Proposition [6.1](#page-44-2) and Corollary [6.1](#page-45-4) is the factor $N\beta$ in the weight function $e^{-N\beta V(x)/2}$. If this factor is removed, and one considers instead the matrix ensemble $\text{ME}_{\beta,N}[e^{-\tilde{V}(x)}]$ for some \tilde{V} independent of N, β , a distinct statistical state can be obtained by taking the scaled high temperature limit $\beta^{-1} \to \infty$ with $N\beta/2 = \alpha$ fixed. As noted in [\[59\]](#page-49-20), in the classical cases of the Gaussian, Laguerre and Jacobi weights, a duality relation for the low temperature, $\beta \rightarrow 0$ with N fixed, limit applies.

Consider for definiteness the Gaussian case with $\tilde{V}(x) = x^2/2$ [\[4\]](#page-47-20). Let the moments of the resulting normalised density be denoted $\{m_{2k}^{G^*}(\alpha)\}_{k=1,2,...}$. With $\{\tilde{m}_{2k}^{G}(\beta,N)\}\$ as in Corollary [6.1,](#page-45-4) from the definitions $m_{2k}^{G^*}(\alpha) = \lim_{N \to \infty, N\beta/2 = \alpha} (\frac{N}{\beta} / 2)^k \tilde{m}_{2k}^G(\beta, N)$ and we read off from [\(6.9\)](#page-45-0) that

$$
\widetilde{m}_{2}^{\mathrm{G}^{*}}(\alpha) = \alpha + 1 \n\widetilde{m}_{4}^{\mathrm{G}^{*}}(\alpha) = 2\alpha^{2} + 5\alpha + 3 \n\widetilde{m}_{6}^{\mathrm{G}^{*}}(\alpha) = 5\alpha^{3} + 22\alpha^{2} + 32\alpha + 15.
$$
\n(6.13)

Insert now a factor β in front of \tilde{V} and consider $ME_{\beta,N}[e^{-\beta \tilde{V}(x)}]$. Denoting the corresponding moments by $\{m_{2k}^{\text{G}}(\beta, N)\}\$, we have $m_{2k}^{\text{G}}(\beta, N) = N^{k+1} \tilde{m}_{2k}^{\text{G}}(\beta, N)$. We see from [\(6.9\)](#page-45-0) that these moments have a well defined expansion in $1/\beta$ with leading terms

$$
\lim_{\beta \to \infty} m_2^G(\beta, N) = N^2 - N
$$

\n
$$
\lim_{\beta \to \infty} m_4^G(\beta, N) = 2N^3 - 5N^2 + 3N
$$

\n
$$
\lim_{\beta \to \infty} m_6^G(\beta, N) = 5N^4 - 22N^3 + 32N^2 - 15N,
$$
\n(6.14)

thus providing evidence for a duality formula linking low temperature moments to the scaled high temperature moments

$$
\lim_{\beta \to \infty} m_{2k}^{\mathcal{G}}(\beta, N) = (-1)^k N m_{2k}^{\mathcal{G}^*}(\alpha) \Big|_{\alpha = -N}.
$$
\n(6.15)

This has been established in [\[59\]](#page-49-20) using a loop equation analysis.

Acknowledgements

Feedback on an earlier draft from Sungsoo Byun and Bojian Shen is appreciated. This work was supported by the Australian Research Council Discovery Project grant DP250102552.

References

- [1] G. Akemann, N. Aygün, M. Kieburg, and P. Pässler, Complex symmetric, self-dual, and Ginibre random matrices: Analytical results for three classes of bulk and edge statistics, [arXiv:2410.21032.](http://arxiv.org/abs/2410.21032)
- [2] G. Akemann, M.J. Phillips and H.-J. Sommers, Characteristic polynomials in real Ginibre ensembles, J. Phys. A 42 (2009), 012001.
- [3] G. Akemann and G. Vernizzi, *Characteristic polynomials of complex matrix models*, Nucl. Phys. B, 660 (2003), 532–556.
- [4] R. Allez, J.P. Bouchard and A. Guionnet, Invariant beta ensembles and the Gauss-Wigner crossover, Phys. Rev. Lett. 109 (2012), 09412.
- [5] M.F. Atiyah, Duality in mathematics and physics, In Conferències FME. Vol. V, Curs Riemann, 2007-2008, Barcelona: Facultat de Matemàtiques i Estadística, pp. 69–91.
- [6] T. Assiotis, On the moments of the partition function of the CβE field, J. Stat. Phys. 187 (2022), 14.
- [7] J. Baik, P. Deift, and E. Strahov, Products and ratios of characteristic polynomials of random Hermitian matrices, J. Math. Phys. 44 (2003), 3657–3670.
- [8] J. Baik and E.M. Rains, Algebraic aspects of increasing subsequences, Duke Math. J., 109 (2001), $1-65.$
- [9] E.C. Bailey and J.P. Keating, Maxima of log-correlated fields: some recent developments, J. Phys. A 55 (2022), 053001.
- [10] T.H. Baker and P.J. Forrester, The Calogero-Sutherland model and generalized classical polynomials, Commun. Math. Phys. 188 (1997), 175–216.
- [11] C.W.J. Beenakker, GOE/GSE duality and Bott periodicity, (2019), mathoverflow.net/questions/258363/goe-gse-duality-and-bott-periodicity
- [12] M.V. Berry and J.P. Keating, Clusters of near degenerate levels dominate negative moments of spectral determinants, J. Phys. A 35 (2002), L1–L6.
- [13] F. Bornemann, A note on the expansion of the smallest eigenvalue distribution of the LUE at the hard edge, Ann. Appl. Prob. 26 (2016), 1942–1946.
- [14] F. Bornemann, Asymptotic expansions relating to the lengths of longest monotone subsequences of involutions, Experimental Math. (2024), 1–45. https://doi.org/10.1080/10586458.2024.2397334.
- [15] F. Bornemann, Asymptotic expansions relating to the distribution of the length of longest increasing subsequences, Forum Math. Sigma. 12:e36. (2024) doi:10.1017/fms.2024.13
- [16] F. Bornemann, Asymptotic expansions of the limit laws of Gaussian and Laguerre (Wishart) ensembles at the soft edge, [arXiv:2403.07628.](http://arxiv.org/abs/2403.07628)
- [17] A. Borodin and P.J. Forrester, Increasing subsequences and the hard-to-soft transition in matrix ensembles, J.Phys. A 36 (2003), 2963–2981.
- [18] A. Borodin and G. Olshanski Z-measures on partitions and their scaling limits, Electr. J. Combin. 26 (2005), 795–834.
- [19] G. Borot, B. Eynard, S.N. Majumdar, and C. Nadal, Large deviations of the maximal eigenvalue of random matrices, J. Stat. Mech. 2011 (2011), P11024.
- [20] G. Borot, A. Guionnet, Asymptotic expansion of β matrix models in the one-cut regime, Commun. Math. Phys. 317 (2013), 447–483.
- [21] E. Brézin and S. Hikami, Characteristic polynomials of random matrices, Commun. Math. Phys. 214 (2000), 111–135.
- [22] E. Brézin and S. Hikami, Characteristic polynomials of real symmetric random matrices, Commun. Math. Phys. 223 (2001), 363–382.
- [23] E. Brézin and S. Hikami, Random matrix theory with an external source, Springer briefs in mathematical physics vol. 19, 2016.
- [24] E. Brézin, S. Hikami, A. Zee, Universal correlations for deterministic plus random hamiltonians, Phys. Rev. E 51 (1995), 5442.
- [25] A. Brini, M. Mariño, and S. Stevan, The uses of the refined matrix model recursion, J. Math. Phys. 52 (2011), 35–51.
- [26] S.-S. Byun, C. Charlier, P. Moreillon and N. Simm, Planar equilibrium measure for the truncated ensembles with a point charge, manuscript in progress.
- [27] S.-S. Byun and P.J. Forrester, *Progress on the study of the Ginibre ensembles*, KIAS Springer Series in Mathematics 3, Springer, 2024.
- [28] S.-S. Byun, P.J. Forrester and S. Lahiry, Properties of the one-component Coulomb gas on a sphere with two macroscopic external charges, a xiv:2501.05061.
- [29] S.-S. Byun and Y.-W. Lee, Finite size corrections for real eigenvalues of the elliptic Ginibre matrices, Random Matrices: Th. Appl. 13 (2024), 2450005.
- [30] S.-S. Byun, S.-M. Seo, and M. Yang, Free energy expansions of a conditional GinUE and large deviations of the smallest eigenvalue of the LUE, [arXiv:2402.18983.](http://arxiv.org/abs/2402.18983)
- [31] C. Charlier, Asymptotics of Hankel determinants with a one-cut regular potential and Fisher-Hartwig singularities, Int. Math. Res. Not. 2019 (2019), 7515–7576.
- [32] J.B. Conrey, D.W. Farmer, and M.R. Zirnbauer, Howe pairs, supersymmetry, and ratios of random *characteristic polynomials for the unitary groups* $U(N)$ *, [arXiv:math-ph/0511024.](http://arxiv.org/abs/math-ph/0511024)*
- [33] A. Deaño and N. Simm, Characteristic polynomials of complex random matrices and Painlevé transcendents, Int. Math. Res. Not., 2022 (2022), 210–264.
- [34] P. Deift, A. Its and I. Krasovky, Asymptotics of Toeplitz, Hankel, and Toeplitz+Hankel determinants with Fisher-Hartwig singularities, Ann. Math. 174 (2011) 1243-1299.
- [35] P. Desrosiers, Duality in random matrix ensembles for all β, Nucl. Phys. B 817 (2009), 224–251.
- [36] P. Desrosiers and P.J. Forrester, Hermite and Laguerre β-ensembles: asymptotic corrections to the eigenvalue density, Nucl. Phys. B 743 (2006), 307-332.
- [37] P. Desrosiers and D.-Z. Liu, Asymptotics for products of characteristic polynomials in classical βensembles, Constr. Approx. 39 (2014), 273.
- [38] P. Desrosiers and D.-Z. Liu, Scaling limits of correlations of characteristic polynomials for the Gaussian β -ensemble with external source, Int. Math. Res. Not. 2015 (2015), 3751–3781.
- [39] I. Dumitriu, Eigenvalue statistics for beta-ensembles, Ph.D. Thesis, MIT, 2003.
- [40] I. Dumitriu and A. Edelman, Global spectrum fluctuations for the β-Hermite and β-Laguerre ensembles via matrix models, J. Math. Phys. 47 (2006), 063302.
- [41] F.J. Dyson, Statistical theory of energy levels of complex systems I, J. Math. Phys. 3 (1962), 140–156.
- [42] F.J. Dyson, A Brownian motion model for the eigenvalues of a random matrix, J. Math. Phys. 3 (1962), 1191–1198.
- [43] A. Edelman, The probability that a random real Gaussian matrix has k real eigenvalues, related distributions, and the circular law, J. Multivariate. Anal. **60** (1997), 203-232.
- [44] A. Edelman, A. Guionnet and S. Péché, Beyond universality in random matrix theory, Ann. Appl. Prob. 26, 1659–1697.
- [45] S.F. Edwards and P.W. Anderson, Theory of spin glasses, J. Phys. F 5 (1975), 965–974.
- [46] B. Eynard, T. Kimura and S. Ribault, Random matrices, [arXiv:1510.04430.](http://arxiv.org/abs/1510.04430)
- [47] Z.M. Feng and J.P. Song, Integrals over the circular ensembles relating to classical domains, J. Phys. A 42 (2009), 325204.
- [48] P.J. Forrester, Selberg correlation integrals and the $1/r^2$ quantum many body system, Nucl. Phys. B 388 (1992), 671–699.
- [49] P.J. Forrester, A constant term identity and its relationship to the log-gas and some quantum many body systems, Phys. Lett. A 163 (1992), 121–126.
- [50] P.J. Forrester, The spectrum edge of random matrix ensembles, Nucl. Phys. B 402 (1993), 709–728.
- [51] P.J. Forrester, Exact results and universal asymptotics in the Laguerre random matrix ensemble, J. Math. Phys. 35 (1993), 2539–2551.
- [52] P. J. Forrester, Exact integral formulas and asymptotics for the correlations in the $1/r^2$ quantum many body system, Phys. Lett. A 179 (1993), 127–130.
- [53] P.J. Forrester, Addendum to Selberg correlation integrals and the $1/r^2$ quantum many body system, Nucl. Phys. B 416 (1994), 377–385.
- [54] P.J. Forrester, Log-gases and random matrices, Princeton University Press, Princeton, NJ, 2010.
- [55] P.J. Forrester, The averaged characteristic polynomial for the Gaussian and chiral Gaussian ensemble with a source, J. Phys. A **46** (2013), 345204.
- [56] P.J. Forrester, Asymptotics of spacing distributions at the hard edge for β-ensembles, Random Matrices: Th. Appl. 2 (2013), 1350002.
- [57] P.J. Forrester, Meet Andréief, Bordeaux 1886, and Andreev, Kharkov 1882-1883, Random Matrices: Th. Appl. 8 (2019), 1930001.
- [58] P.J. Forrester, Differential identities for the structure function of some random matrix ensembles, J. Stat. Phys. 183 (2021), 33.
- [59] P.J. Forrester, High-low temperature dualities for the classical β-ensembles, Random Matrices: Th. Appl. 11, (2022) 2250035.
- [60] P.J. Forrester, A review of exact results for fluctuation formulas in random matrix theory, Probab. Surveys 20 (2023), 170–225.
- [61] P.J. Forrester, Ensemble inter-relations in random matrix theory, [arXiv:2409.00964.](http://arxiv.org/abs/2409.00964)
- [62] P.J. Forrester, Dualities for characteristic polynomial averages of complex symmetric and self dual non-Hermitian random matrices, [arXiv:2411.07356.](http://arxiv.org/abs/2411.07356)
- [63] P.J. Forrester, N.E. Frankel and T. Garoni. Asymptotic form of the density profile for Gaussian and Laguerre random matrix ensembles with orthogonal and symplectic symmetry. J. Math. Phys., 47 (2006), 023301.
- [64] P.J. Forrester and A. Gamburd, Counting formulas associated with some random matrix ensembles, J. Combin. Th. Series A 113 (2006), 934–951.
- [65] P.J. Forrester and T.D. Hughes, Complex Wishart matrices and conductance in mesoscopic systems: exact results, J. Math. Phys. 35 (1994), 6736–6747.
- [66] P.J. Forrester and J.R. Ipsen, Selberg integral theory and Muttalib-Borodin ensembles, Adv. Appl. Math. 95 (2018), 152–176.
- [67] P.J. Forrester and J.R. Ipsen, A generalisation of the relation between zeros of the complex Kac polynomial and eigenvalues of truncated unitary matrices, Prob. Theory Related Fields 175 (2019), 833–847.
- [68] P.J. Forrester and J.P. Keating, Singularity dominated strong fluctuations for some random matrix averages, Commun. Math. Phys. 250 (2004), 119–131.
- [69] P.J. Forrester and S.-H. Li, Rate of convergence at the hard edge for various Pólya ensembles of positive definite matrices, Integral Transforms Spec. Funct. 33 (2022), 466–484.
- [70] P.J. Forrester, S.-H. Li and A.K. Trinh, Asymptotic correlations with corrections for the circular Jacobi β-ensemble, J. Approx. Th. 271 (2021), 105633.
- [71] P.J. Forrester and D.-Z. Liu, Phase transitions for products of characteristic polynomials under Dyson Brownian motion, Acta. Math. Sin.-English Ser. 37 (2021), 509–524.
- [72] P.J. Forrester and A. Mays, Finite size corrections relating to distributions of the length of longest increasing subsequences, Adv. Applied Math. 145 (2023), 102482.
- [73] P.J. Forrester, B. Jancovici, and D.S. McAnally, Analytic properties of the structure function for the one-dimensional one-component log-gas, J. Stat. Phys. 102 (2000), 737–780.
- [74] P.J. Forrester, A.A. Rahman, and N.S. Witte, Large N expansions for the Laguerre and Jacobi β ensembles from the loop equations, J. Math. Phys. 58 (2017), 113303.
- [75] P.J. Forrester and E.M. Rains, Matrix averages relating to the Ginibre ensemble, J. Phys. A 42 (2009), 385205
- [76] P.J. Forrester and B.-J. Shen, Expanding the Fourier transform of the scaled circular Jacobi β ensemble density, J. Stat. Phys. 190 (2023), 160.
- [77] P.J. Forrester and A.K. Trinh. Functional form for the leading correction to the distribution of the largest eigenvalue in the GUE and LUE. J. Math. Phys., 59 (2018), 053302.
- [78] P.J. Forrester and A. K. Trinh, Optimal soft edge scaling variables for the Gaussian and Laguerre even β ensembles, Nucl. Phys. B 938 (2019), 621–639.
- [79] P.J. Forrester and A. K. Trinh, Finite size corrections at the hard edge for the Laguerre β ensemble, Stud. Appl. Math. 143 (2019), 315–336.
- [80] P.J. Forrester and D. Wang, Muttalib-Borodin ensembles in random matrix theory realisations and correlation functions, Elec. J. Probab. 22 (2017), 54.
- [81] P.J. Forrester and N.S. Witte. Application of the τ-function theory of Painlevé equations to random matrices: P_{VI} , the JUE, CyUE, cJUE and scaled limits. Nagoya Math. J., 174: 29-114, 2004.
- [82] Y.V. Fyodorov, Negative moments of characteristic polynomials of random matrices: Ingham-Siegel integral as an alternative to Hubbard-Stratonovich transformation, Nucl. Phys. B 621 (2002), 643– 674.
- [83] Y.V. Fyodorov, J. Grela, and E. Strahov, On characteristic polynomials for a generalized chiral random matrix ensemble with a source, J. Phys. A 51 (2018) 134003.
- [84] Y.V. Fyodorov and J.P. Keating, Negative moments of characteristic polynomials of GOE matrices and singularity- dominated strong fluctuations, J. Phys. A 36 (2003), 4035–4046.
- [85] Y.V. Fyodorov and B.A. Khoruzhenko, Averages of spectral determinants and 'Single ring theorem' of Feinberg and Zee, Acta Phys. Polonica B, 38 (2007), 4067–4077.
- [86] Y.V. Fyodorov and B.A. Khoruzhenko, On absolute moments of characteristic polynomials of a certain class of complex random matrices, Comm. Math. Phys. 273 (2007), 561–599.
- [87] Y. V. Fyodorov and B. A. Khoruzhenko, A few remarks on colour-flavour transformations, truncations of random unitary matrices, Berezin reproducing kernels and Selberg-type integrals, J. Phys. A 40 (2007), 669-699.
- [88] Y.V. Fyodorov, B.A. Khoruzhenko, T. Prellberg, Zeros of conditional Gaussian analytic functions, random sub-unitary matrices and q-series, [arXiv:2412.06086.](http://arxiv.org/abs/2412.06086)
- [89] Y.V. Fyodorov and H.-J. Sommers, Random matrices close to hermitian or unitary: overview of methods and results, J. Phys. A 36 (2003), 3303–3347.
- [90] Y.V. Fyodorov and E. Strahov, An exact formula for general spectral correlation function of random Hermitian matrices, J. Phys. A 36 (2003), 3203.
- [91] T.M. Garoni, On the asymptotics of some large Hankel determinants generated by Fisher-Hartwig symbols defined on the real line, J. Math. Phys. 46 (2005), 043516.
- [92] T.M. Garoni, P.J. Forrester, and N.E. Frankel, Asymptotic corrections to the eigenvalue density of the GUE and LUE, J. Math. Phys. 46 (2005), 103301.
- [93] I.P. Goulden and D.M. Jackson, Maps in locally orientable surfaces and integrals over real symmetric matrices, Canad. J. Math. 49 (1997), 865–882.
- [94] J. Grela, Diffusion method in random matrix theory, J. Phys. A 49 (2016), 015201.
- [95] F. Haake, Quantum signatures of chaos, 2nd ed., Springer, Berlin, 2000.
- [96] W. Hachem, A. Hardy and J. Najim, Large complex correlated Wishart matrices: the Pearcey kernel and expansion at the hard edge, Elec. J. Probab. 21 (2016), 1–36.
- [97] Harish-Chandra, Differential operators on a semisimple Lie algebra. Amer. J. Math. (1957) 87–120.
- [98] C. Itzykson and J.-B. Zuber, Matrix integration and combinatorics of modular groups, Commun. Math. Phys. 134 (1990), 197–207.
- [99] A.T. James, Distributions of matrix variate and latent roots derived from normal samples, Ann. Math. Statist. 35 (1964), 475–501.
- [100] K. Johansson, On fluctuation of eigenvalues of random Hermitian matrices, Duke Math. J. 91 (1998), 151–204.
- [101] B. Jonnadula, J. P. Keating and F. Mezzadri, On the moments of characteristic polynomials, Glasg. Math. J. 65(S1) (2023), S102–S122.
- [102] K.W.J. Kadell, The Selberg-Jack symmetric functions, Adv. Math. 130 (1997), 33–102.
- [103] A. Kamenev and M. Mézard, Wigner-Dyson statistics from the replica method, J. Phys. A 32 (1999), 4373.
- [104] J. Kaneko, Selberg integrals and hypergeometric functions associated with Jack polynomials, SIAM J. Math Anal. 24 (1993), 1086–1110.
- [105] E. Kanzieper, Replica field theories, Painlevé transcendents, and exact correlation functions, Phys. Rev. Lett., 89 (2002) 250201.
- [106] E. Kanzieper, Exact replica treament of non-Hermitean complex random matrices. In O. Kovras, editor, Frontiers in Field Theory, pages 23–51. Nova Science Publishers, 2005.
- [107] J.P. Keating and N.C. Snaith, Random matrix theory and $\zeta(1/2 + it)$, Commun. Math. Phys. 214 (2001), 57–89.
- [108] I. Krasovsky, Correlations of the characteristic polynomials in the Gaussian unitary ensemble or a singular Hankel determinant, Duke Math J. 139 (2007), 581–619.
- [109] M. Krishnapur, From random matrices to random analytic functions, Ann. Prob. 37 (2009), 314– 346.
- [110] A. Kulkarni, K. Kawabata and S. Ryu, Non-linear sigma models for non-Hermitian random matrices in symmetry classes $A\bar{I}^{\dagger}$ and AII^{\dagger} , arXiv: 2410.24043.
- [111] Y. Kuramoto and Y. Kato, Dynamics of one-dimensional quantum systems, CUP, Cambridge, 2009.
- [112] M. Ledoux, A recursion formula for the moments of the Gaussian orthogonal ensemble, Ann. Inst. Henri Poincaré Probab. Stat. 45 (2009), 754–769.
- [113] D.-Z. Liu, Limits for circular Jacobi beta-ensembles. J. Approx. Theory, 215 (2017), 40–67.
- [114] D.-Z. Liu and L. Zhang, Phase transition of eigenvalues in deformed Ginibre ensembles, [arXiv:2204.13171.](http://arxiv.org/abs/2204.13171)
- [115] D.-Z. Liu and L. Zhang, Duality in non-Hermitian random matrix theory, Nucl. Phys. B 1004 (2024), 116559.
- [116] I.G. Macdonald, Hall polynomials and symmetric functions, 2nd ed., Oxford University Press, Oxford, 1995.
- [117] F. Mezzadri, A.K. Reynolds and B. Winn, Moments of the eigenvalue densities and of the secular coefficients of β -ensembles, Nonlinearity 30 (2017), 1034.
- [118] C. McSwiggen, The Harish-Chandra integral: An introduction with examples, L'Enseignement Mathématique 67 (2021), 229–299.
- [119] A.D. Mironov, A.Yu. Morozov, A.V. Popolitov, and Sh.R. Shakirov, Resolvents and Seiberg-Witten representation for a Gaussian β -ensemble, Theor. Math. Phys., 171 (2012), 505–522.
- [120] R. L. Mkrtchyan and A. P. Veselov, On duality and negative dimensions in the theory of Lie groups and symmetric spaces, J. Math. Phys. 52 (2011), 083514.
- [121] L. Moreno-Pozas, D. Morales-Jimenez and M.R. McKay, Extreme eigenvalue distributions of Jacobi ensembles: new exact representations, asymptotics and finite size corrections, Nucl. Phys. B 947, (2019) 114724.
- [122] R.J. Muirhead, *Aspects of multivariate statistical theory*, Wiley, New York, 1982.
- [123] M. Mulase and A. Waldron, Duality of orthogonal and symplectic matrix integrals and quaternionic Feynman graphs, Commun. Math. Phys. 240 (2003), 553–586.
- [124] S. M. Nishigaki and A. Kamenev, Replica treatment of non-Hermitian disordered Hamiltonians, J. Phys. A 35 (2002), 4571–4590.
- [125] J. Novak, Hypergeometric functions of random matrices and quasimodular forms, [arXiv:2410.04243.](http://arxiv.org/abs/2410.04243)
- [126] L. Pastur and M. Shcherbina, Eigenvalue distribution of large random matrices, American Mathematical Society, Providence, RI, 2011.
- [127] A. Perret and G. Schehr, Finite N corrections to the limiting distribution of the smallest eigenvalue of Wishart complex matrices, Random Matrices: Theory and Applications, 5 (2016), 1650001.
- [128] A.A. Rahman and P.J. Forrester, Linear differential equations for the resolvents of the classical matrix ensembles, Random Matrices Theory Appl. 10 (2021), 2250003.
- [129] E.M. Rains, Attack of the zonal polynomials, unpublished manuscript, 1995.
- [130] E.M. Rains, Increasing subsequences and the classical groups, Elect. J. of Combinatorics 5 (1998), $\#\mathrm{R12}.$
- [131] L.A. Roberts, Towards the calculation of Jack polynomials, Ph.D. Thesis, Victoria University Wellington, 2001.
- [132] M. Rösler, Generalized Hermite polynomials and the heat equation for Dunkl operators, Commun. Math. Phys. 192 (1998) 519–542.
- [133] A. Serebryakov, *Multi-point correlators in non-Hermitian matrices and beyond*, Ph.D. Thesis, University of Sussex, 2023.
- [134] A. Serebryakov, N. Simm, and G. Dubach, Characteristic polynomials of random truncations: Moments, duality and asymptotics, Random Matrices: Theory and Applications, 12 (2023), 2250049.
- [135] A. Serebryakov and N. Simm, *Schur function expansion in non-Hermitian ensembles and averages* of characteristic polynomials, Ann. Henri Poincaré (2024). https://doi.org/10.1007/s00023-024-01483-6
- [136] T. Shcherbina, On the correlation function of the characteristic polynomials of the Hermitian Wigner ensemble, Commun. Math. Phys. 308 (2011), 1–21.
- [137] B.-J. Shen and P.J. Forrester, Moments of characteristic polynomials for classical β ensembles, in preparation.
- [138] G. Szegő, Orthogonal polynomials, 4th ed., American Mathematical Society, Providence R.I., 1975.
- [139] A. Takemura, Zonal polynomials, Institute of Mathematical Statistics, Hayward, CA, 1984.
- [140] R. Tribe and O. Zaboronski, The Ginibre evolution in the large N limit, J. Math. Phys. 55 (2014), 063304.
- [141] K.W. Wachter, The strong limits of random matrix spectra for sample matrices of independent elements, Annal. Prob. 6 (1978), 1–18.
- [142] S.O. Warnaar, q-Selberg integrals and Macdonald polynomials, Ramanujan J. 10 (2005), 237–268.
- [143] C. Webb and M. D. Wong, On the moments of the characteristic polynomial of a Ginibre random matrix, Proc. Lond. Math. Soc., 118 (2019), 1017–1056.
- [144] B. Winn, Extreme eigenvalues of random matrices from Jacobi ensembles, [arXiv:2302.12082.](http://arxiv.org/abs/2302.12082)
- [145] N.S. Witte and P.J. Forrester, Moments of the Gaussian β ensembles and the large N expansion of the densities, J. Math. Phys. 55 (2014), 083302.
- [146] Z. Yan, A class of generalized hypergeometric functions in several variables, Can J. Math. 44 (1992), 1317–1338.
- [147] M.R. Zirnbauer, Supersymmetry for systems with unitary disorder: Circular ensembles, J Phys A 29 (1996), 7113–7136.
- [148] M.R. Zirnbauer, Color-flavor transformation revisited, [arXiv:2109.10272.](http://arxiv.org/abs/2109.10272)