

# SEMICLASSICAL RESOLVENT ESTIMATES FOR THE MAGNETIC SCHRÖDINGER OPERATOR

GEORGI VODEV

**ABSTRACT.** We obtain semiclassical resolvent estimates for the Schrödinger operator  $(ih\nabla + b)^2 + V$  in  $\mathbb{R}^d$ ,  $d \geq 3$ , where  $0 < h \ll 1$  is a semiclassical parameter,  $V$  and  $b$  are real-valued electric and magnetic potentials independent of  $h$ . If  $V \in L^\infty(\mathbb{R}^d)$ ,  $b \in L^\infty(\mathbb{R}^d; \mathbb{R}^d)$ ,  $\operatorname{div} b \in L^\infty(\mathbb{R}^d)$  satisfy  $V(x) = \mathcal{O}(|x|^{-1-\epsilon})$ ,  $b(x) = \mathcal{O}(|x|^{-1-\epsilon})$ ,  $\operatorname{div} b(x) = \mathcal{O}(|x|^{-1-\epsilon})$ ,  $\epsilon > 0$ , for  $|x| \gg 1$ , we prove that the norm of the weighted resolvent is bounded by  $\exp(C h^{-2} \log(h^{-1}))$ ,  $C > 0$ . We get better resolvent bounds for electric potentials which are Hölder with respect to the radial variable and magnetic potentials which are Hölder with respect to the space variable. For long-range electric potentials which are Lipschitz with respect to the radial variable and long-range magnetic potentials which are Lipschitz with respect to the space variable we obtain a resolvent bound of the form  $\exp(C h^{-1})$ ,  $C > 0$ .

Key words: Schrödinger operator, magnetic potentials, resolvent estimates.

## 1. INTRODUCTION AND STATEMENT OF RESULTS

In this paper we are going to study the resolvent of the Schrödinger operator

$$P(h) = (ih\nabla + b(x))^2 + V(x) \quad \text{in } \mathbb{R}^d, d \geq 3,$$

where  $0 < h \ll 1$  is a semiclassical parameter,  $\nabla$  is the gradient,  $V \in L^\infty(\mathbb{R}^d; \mathbb{R})$  and  $b = (b_1, \dots, b_d) \in L^\infty(\mathbb{R}^d; \mathbb{R}^d)$  are electric and magnetic potentials, respectively. We are interested in bounding the quantity

$$g_s^\pm(h, \varepsilon) := \log \left\| (|x| + 1)^{-s} (P(h) - E \pm i\varepsilon)^{-1} (|x| + 1)^{-s} \right\|_{L^2 \rightarrow L^2}$$

from above by an explicit function of  $h$ , independent of  $\varepsilon$ . Here  $L^2 := L^2(\mathbb{R}^d)$ ,  $0 < \varepsilon < 1$ ,  $s > 1/2$  is independent of  $h$  and  $E > 0$  is a fixed energy level independent of  $h$ . Our goal is to obtain the best possible upper bounds for  $g_s^\pm(h, \varepsilon)$  for potentials which are Lipschitz, Hölder or just  $L^\infty$ . There have been recently many papers studying this problem when  $b \equiv 0$ . To our best knowledge, no such results exist for non-trivial magnetic potentials when  $d \geq 2$  and it seems that this paper is the first one where upper bounds for  $g_s^\pm$  are proved in this case. When  $d = 1$  and  $b$  not identically zero, semiclassical resolvent bounds have been recently proved in [9] for a very large class of electric and magnetic potentials. Note also that sharp high-frequency resolvent bounds for the operator  $P(1)$  with  $L^\infty$  potentials are proved in [11] when  $d \geq 2$  and in [16] when  $d \geq 3$ . In [7] exponential high-frequency resolvent bounds for the operator  $P(1)$  with smooth potentials on non-compact Riemannian manifolds have been recently proved, extending the results in [1].

We will be looking for bounding  $g_s^\pm$  for the largest possible class of electric and magnetic potentials that our method allows to cover. To describe it we introduce the polar coordinates  $r = |x|$ ,  $w = x/|x|$ . We suppose that  $V = V_L + V_S$ ,  $b = b^L + b^S$ , where  $V_L, b^L$  (resp.  $V_S, b^S$ ) are long-range (resp. short-range) electric and magnetic potentials satisfying the following conditions.

We suppose that  $V_L$  satisfies

$$(1.1) \quad V_L(rw) \leq p(r),$$

where  $p > 0$  is a decreasing function such that  $p(r) \rightarrow 0$  as  $r \rightarrow \infty$ . We also suppose that  $V_L(rw)$  is Lipschitz with respect to the radial variable  $r$  and its first derivative satisfies the bound

$$(1.2) \quad \partial_r V_L(rw) \leq C(r+1)^{-1-\rho}.$$

Hereafter  $C > 0$  and  $0 < \rho \ll 1$  will denote positive constants that may change from line to line. The short-range part  $V_S \in L^\infty(\mathbb{R}^d)$  is supposed to satisfy

$$(1.3) \quad |V_S(rw)| \leq C(r+1)^{-1-\rho}.$$

The long-range part  $b^L \in L^\infty(\mathbb{R}^d; \mathbb{R}^d)$  is supposed to be Lipschitz with respect to  $r$  and satisfies

$$(1.4) \quad |\partial_r^k b^L(rw)| \leq C(r+1)^{-k-\rho}, \quad k = 0, 1.$$

The short-range part  $b^S \in L^\infty(\mathbb{R}^d; \mathbb{R}^d)$  satisfies

$$(1.5) \quad |b^S(rw)| \leq C(r+1)^{-1-\rho}.$$

Finally, we suppose that  $\operatorname{div} b^S \in L^\infty(\mathbb{R}^d)$  exists and satisfies

$$(1.6) \quad |\operatorname{div} b^S(rw)| \leq C(r+1)^{-1-\rho}.$$

This condition is fulfilled if each  $b_j^S$  is Lipschitz with respect to the variable  $x_j$  and  $\partial_{x_j} b_j^S(x) = \mathcal{O}(|x|^{-1-\rho})$  for  $|x| \gg 1$ .

Throughout this paper, given two vectors  $a, b \in \mathbb{R}^d$ ,  $a \cdot b$  will denote the scalar product. Our first result is the following

**Theorem 1.1.** *In addition to the above conditions we assume either the condition*

$$(1.7) \quad |w \cdot b^L(rw)| \leq C(r+1)^{-1-\rho},$$

*or we suppose that the function  $x \cdot b^L(x)$  is Lipschitz in  $x$  and satisfies*

$$(1.8) \quad |\partial_x^\beta (x \cdot b^L(x))| \leq C, \quad |\beta| = 1.$$

*Then, there exist constants  $C > 0$ ,  $0 < h_0 \ll 1$ , independent of  $h$  and  $\varepsilon$ , such that the bound*

$$(1.9) \quad g_s^\pm(h, \varepsilon) \leq Ch^{-2} \log(h^{-1})$$

*holds for all  $0 < h \leq h_0$ . If  $V_S \equiv 0$  and  $b^S \equiv 0$ , under the condition (1.8), we have the better bound*

$$(1.10) \quad g_s^\pm(h, \varepsilon) \leq Ch^{-1}.$$

Note that the bound (1.10) is proved in [9] when  $d = 1$ , while for  $d \geq 3$  it seems to be new. Proving (1.10) when  $d = 2$  in the presence of a magnetic potential remains an open problem. When  $b \equiv 0$ , however, the bound (1.10) is well-known. Indeed, it was proved in [2] when  $d \geq 3$  for potentials satisfying (1.1) with  $p(r) = C(r+1)^{-\rho}$ , and extended to the general case in [5] and [18]. When  $d = 2$  the bound (1.10) is proved in [13] for potentials  $V \in C^1(\mathbb{R}^2)$  satisfying (1.1) with  $p(r) = C(r+1)^{-\rho}$  as well as the condition

$$(1.11) \quad |\nabla V(x)| \leq C(|x|+1)^{-1-\rho}.$$

The bound (1.10) has been recently proved in [12] for long-range Lipschitz potentials having singularities at the origin in all dimensions  $d \geq 2$ . When  $d = 1$  the bound (1.10) is proved in [4] for potentials  $V \in L^1(\mathbb{R})$  and in [9] for more general measure potentials. It turns out that (1.10) also holds for  $V \in L^\infty(\mathbb{R}^d)$ ,  $d \geq 2$ , provided  $V$  is compactly supported and depends only on the radial variable  $r$ . Indeed, this is proved in [3] when  $d \geq 2$ . A simpler proof of this result is given

in [21] when  $d \geq 3$ . Without a radial symmetry or some regularity of the potential, however, the bound (1.10) seems to fail even for compactly supported potentials. Indeed, for potentials  $V \in L^\infty$  satisfying

$$(1.12) \quad |V(x)| \leq C(|x| + 1)^{-\delta}$$

where  $\delta > 1$ , the following bounds are known to hold:

$$(1.13) \quad g_s^\pm(h, \varepsilon) \leq Ch^{-4/3} \log(h^{-1}) \quad \text{if } \delta > 2,$$

$$(1.14) \quad g_s^\pm(h, \varepsilon) \leq C_\epsilon h^{-\frac{2\delta}{2\delta-1}-\epsilon} \quad \text{if } 1 < \delta \leq 2,$$

for every  $0 < \epsilon \ll 1$ . When  $d \geq 3$  the bound (1.13) is proved in [6] and in [17] if  $\delta > 3$ . Previously, (1.13) was proved for compactly supported  $L^\infty$  potentials in [8] and [14]. The bound (1.13) is extended in [19] to more general Riemannian manifolds. The bounds (1.13) and (1.14) have been recently proved in [15] when  $d \geq 2$  for potentials  $V = V_L + V_S$  with  $V_L$  satisfying (1.1) and (1.2), and  $V_S$  satisfying (1.12). In fact, the conditions on the behavior of  $V_L$  and  $V_S$  at infinity in this paper are a little bit more general than this. Moreover, the potential in [15] is allowed to have singularities at the origin. It turns out that (1.13) and (1.14) can be improved for potentials  $V \in L^\infty$  depending only on  $r$ . Indeed, for radial potentials satisfying (1.12) it is shown in [20] and [21] that when  $d \geq 3$  we have the bounds

$$(1.15) \quad g_s^\pm(h, \varepsilon) \leq Ch^{-\frac{\delta}{\delta-1}} \quad \text{if } \delta > 4,$$

$$(1.16) \quad g_s^\pm(h, \varepsilon) \leq Ch^{-4/3} \quad \text{if } 2 < \delta \leq 4,$$

$$(1.17) \quad g_s^\pm(h, \varepsilon) \leq Ch^{-\frac{2\delta}{2\delta-1}} (\log(h^{-1}))^{\frac{\delta+1}{2\delta-1}} \quad \text{if } 1 < \delta \leq 2.$$

It turns out that the above bounds can be improved if some small regularity of the potential is assumed. To describe these results, given  $0 < \alpha \leq 1$  and  $k > 0$ , we introduce the space  $C_k^\alpha(\overline{\mathbb{R}^+})$  of all Hölder functions  $a$  such that

$$\sup_{r' \geq 0: 0 < |r-r'| \leq 1} \frac{|a(r) - a(r')|}{|r - r'|^\alpha} \leq C(r+1)^{-k}, \quad \forall r \in \overline{\mathbb{R}^+}.$$

We suppose that the function  $V(r, w) := V(rw)$  satisfies the condition

$$(1.18) \quad V(\cdot, w) \in C_{3+\rho}^\alpha(\overline{\mathbb{R}^+}), \quad 0 < \alpha < 1,$$

uniformly in  $w \in \mathbb{S}^{d-1}$ . For potentials  $V$  satisfying (1.1) and (1.18) and  $d \geq 3$  the following bound is known to hold:

$$(1.19) \quad g_s^\pm(h, \varepsilon) \leq Ch^{-4/(\alpha+3)} \log(h^{-1}).$$

The bound (1.19) is proved in [6] and in [18] if  $\rho = 1$ . It has been extended in [18] to the case  $d = 2$  as well as to exterior domains for potentials which are Hölder with respect to the space variable  $x$ . The bound (1.19) is improved in [20] for radial potentials satisfying (1.1) and (1.18) with  $\rho = 0$  to

$$(1.20) \quad g_s^\pm(h, \varepsilon) \leq Ch^{-4/(\alpha+3)}.$$

The proof of the above bounds is based on global Carleman estimates with suitable phase and weight functions. In order that the Carleman estimates work the phase and weight functions must satisfy some conditions. Therefore, the task consists of constructing explicitly these functions in such a way that we get the best possible bound for  $g_s^\pm$ . In fact, there are many ways to do so and the class the potential belongs to is determined by the way we do it. This explains why in the papers mentioned above the same bound is proved for different classes of potentials.

To prove Theorem 1.1 we follow the same strategy as in the case  $b \equiv 0$ . The Carleman estimates for the magnetic Schrödinger operator, however, are much more difficult to handle with. The main difference is that conjugating this operator with  $e^{\varphi/h}$ ,  $\varphi > 0$  being a phase function depending only on  $r$ , makes appear a complex-valued effective potential of the form (see Section 3)

$$-2i\varphi'(r)w \cdot b(rw),$$

where  $\varphi' > 0$  denotes the first derivative of  $\varphi$ . This term is not easy to handle with even in the case when  $b^S \equiv 0$ . One way to treat it is to consider it as a short-range potential and to add it to  $V_S$ . To do so, however, we need the condition (1.7). It allows us to make the Carleman estimates work. However, this approach does not lead to better bounds than what we have in (1.9). Therefore, we propose another way that requires the condition (1.8). This approach allows us to improve the bound (1.9) if  $b_S \equiv 0$  or  $b_S$  is Hölder. To be more precise, given  $0 < \alpha \leq 1$  and  $k > 0$ , we introduce the space  $C_k^\alpha(\mathbb{R}^d)$  of all Hölder functions  $a$  such that

$$\sup_{x' \in \mathbb{R}^d: 0 < |x-x'| \leq 1} \frac{|a(x) - a(x')|}{|x - x'|^\alpha} \leq C(|x| + 1)^{-k}, \quad \forall x \in \mathbb{R}^d.$$

Note that the case  $\alpha = 1$  corresponds to the Lipschitz functions. We suppose that

$$(1.21) \quad V_S(\cdot, w) \in C_{3+\rho}^{\alpha}(\overline{\mathbb{R}^+}), \quad 0 < \alpha \leq 1, \quad \rho > 0,$$

uniformly in  $w$ , and

$$(1.22) \quad b^S \in C_{3/2+\rho_1}^{\alpha'}(\mathbb{R}^d; \mathbb{R}^d), \quad 0 < \alpha' \leq 1, \quad \rho_1 > 0.$$

We have the following

**Theorem 1.2.** *Assume that  $V_L$  satisfies (1.1), (1.2),  $b^L$  satisfies (1.4), (1.8),  $V_S$  satisfies (1.21) and (1.1) with possibly a new decreasing function  $p$  tending to zero,  $b^S$  satisfies (1.6), (1.22). We also suppose that  $b^L$  and  $b^S$  satisfy*

$$(1.23) \quad |b^L(rw)| + |b^S(rw)| \leq C(r+1)^{-\rho_2},$$

with some  $\rho_2 > 0$  such that  $\rho_1 + \rho_2 > 3/2$ . Then we have the bound

$$(1.24) \quad g_s^\pm(h, \varepsilon) \leq Ch^{-n} \log(h^{-1}),$$

where

$$n = \max \left\{ \frac{4}{\alpha + 3}, \frac{2}{\alpha' + 1} \right\}.$$

When  $\alpha' \leq \frac{1}{2}$  the bound (1.24) also holds for  $L^\infty$  potentials  $V_S$ . More precisely, we have the following

**Theorem 1.3.** *Assume that  $V_L$  satisfies (1.1), (1.2),  $b^L$  satisfies (1.4), (1.8),  $b^S$  satisfies (1.6), (1.22) and (1.23),  $V_S$  satisfies*

$$(1.25) \quad |V_S(rw)| \leq C(r+1)^{-\delta},$$

with  $\delta > 1$ . Then we have the bounds

$$(1.26) \quad g_s^\pm(h, \varepsilon) \leq Ch^{-2/(\alpha'+1)} \log(h^{-1}) \quad \text{if } \delta > \frac{1}{1-\alpha'}, \quad \alpha' \leq \frac{1}{2},$$

$$(1.27) \quad g_s^\pm(h, \varepsilon) \leq C_\epsilon h^{-\frac{2\delta}{2\delta-1}-\epsilon} \quad \text{if } \alpha' \geq \frac{\delta-1}{\delta}, \quad 1 < \delta \leq 2,$$

for every  $0 < \epsilon \ll 1$  independent of  $h$ .

Most probably, the conditions in Theorems 1.1, 1.2 and 1.3 are not optimal, but they allow us to make a simple choice of the phase and weight functions, and hence to keep the proof of the Carleman estimates relatively short and simple. However, the bounds (1.9), (1.24) and (1.26) seem to be sharp even for compactly supported potentials, unless one assumes a radial symmetry of the potentials. Indeed, when  $b \equiv 0$  and  $V$  radial, it is shown in [20] that the logarithmic term in (1.9), (1.24), (1.26) and the  $h^{-\epsilon}$  term in (1.27) can be removed. Probably this is still the case for non-trivial radial magnetic and electric potentials.

The paper is organized as follows. In Section 2 we construct the phase and weight functions needed for the proof of the Carleman estimates. In Section 3 we prove Carleman estimates for the magnetic Schrödinger operator under the conditions of Theorem 1.1 needed for the proof of the bound (1.9). In Section 4 we improve the Carleman estimates when  $V_S \equiv 0$  and  $b^S \equiv 0$ , which leads to the bound (1.10). In Section 5 we adapt the Carleman estimates to Hölder potentials  $V_S$  and  $b^S$ , which leads to the bound (1.24). In Section 6 we adapt the Carleman estimates to  $L^\infty$  potentials  $V_S$  and Hölder potentials  $b^S$ , which leads to the bounds (1.26) and (1.27). Finally, in Section 7 we show that the Carleman estimates imply the resolvent ones.

## 2. CONSTRUCTION OF THE PHASE AND WEIGHT FUNCTIONS

We first construct the weight function  $\mu$  as follows:

$$\mu(r) = \begin{cases} r^2 + r & \text{for } 0 \leq r \leq 1, \\ r^2 + r^{2-\ell} & \text{for } 1 \leq r \leq a, \\ (a^2 + a^{2-\ell})(1 + (a+1)^{-2s+1} - (r+1)^{-2s+1}) & \text{for } r \geq a, \end{cases}$$

where  $a \gg 1$  is either independent of  $h$  or it is of the form  $a = h^{-m}$ ,  $m > 0$ . The parameters  $\ell$  and  $s$  are independent of  $h$  such that  $0 < \ell \ll 1$ ,  $1/2 < s < 1$ . Clearly, the first derivative of  $\mu$  is given by

$$\mu'(r) = \begin{cases} 2r + 1 & \text{for } 0 \leq r < 1, \\ 2r + (2 - \ell)r^{1-\ell} & \text{for } 1 < r < a, \\ (2s - 1)(a^2 + a^{2-\ell})(r + 1)^{-2s} & \text{for } r > a. \end{cases}$$

We have the following

**Lemma 2.1.** *We have the identity*

$$(2.1) \quad 2r^{-1}\mu(r) - \mu'(r) = \begin{cases} 1 & \text{for } 0 < r < 1, \\ \ell r^{1-\ell} & \text{for } 1 < r < a. \end{cases}$$

For all  $r > a$ , we have the lower bound

$$(2.2) \quad 2r^{-1}\mu(r) - \mu'(r) \geq a^2(r+1)^{-1}.$$

For all  $r \neq 1$ ,  $r \neq a$ , we have

$$(2.3) \quad \mu'(r) \geq (r+1)^{-2s},$$

$$(2.4) \quad \frac{\mu(r)^j}{\mu'(r)} \lesssim a^{2j-2}(r+1)^{2s},$$

for every  $j \geq 1$ .

*Proof.* The identity (2.1) is obvious, while (2.2) follows from the inequalities

$$2r^{-1}\mu(r) \geq 2(a^2 + a^{2-\ell})(r+1)^{-1}, \quad \mu'(r) \leq (2s-1)(a^2 + a^{2-\ell})(r+1)^{-1}.$$

It is also clear that the inequality (2.3) holds as long as  $a^2 \geq (2s-1)^{-1}$ . The bound (2.4) with  $j=1$  is obvious, while for  $j > 1$  it follows from the fact that  $\mu(r) \lesssim a^2$ .  $\square$

We will now construct a phase function  $\psi \in C^1([0, +\infty))$  such that  $\psi(0) = 0$  and  $\psi(r) > 0$  for  $r > 0$ . We define the first derivative of  $\psi$  by

$$\psi'(r) = \begin{cases} c_\nu & \text{for } 0 \leq r \leq 1, \\ r^{-1}(1+r^{-\ell})^{-1/2}(1-(r+1)^{-\nu}) & \text{for } 1 \leq r \leq a, \\ K_a(r+1)^{-1-\kappa} & \text{for } r \geq a, \end{cases}$$

where  $0 < \nu, \kappa \ll 1$  are independent of  $h$  to be fixed later on, and

$$c_\nu = 2^{-1/2}(1-2^{-\nu}), \quad K_a = a^{-1}(a+1)^{1+\kappa}(1+a^{-\ell})^{-1/2}(1-(a+1)^{-\nu}) = \mathcal{O}(a^\kappa),$$

are chosen in such a way that the function  $\psi'$  is Lipschitz. It is easy to check that the first derivative of  $\psi'$  satisfies

$$\psi''(r) = \begin{cases} 0 & \text{for } 0 \leq r < 1, \\ \mathcal{O}(r^{-2}) & \text{for } 1 < r < a, \\ \mathcal{O}(a^\kappa r^{-2-\kappa}) & \text{for } r > a. \end{cases}$$

**Lemma 2.2.** *For all  $r \geq 0$  we have the bound*

$$(2.5) \quad \psi(r) \lesssim \log a.$$

*Proof.* We have

$$\begin{aligned} \max \psi &= \int_0^1 \psi'(r) dr + \int_1^a \psi'(r) dr + \int_a^\infty \psi'(r) dr \\ &\leq c_\nu \int_0^1 dr + \int_1^a r^{-1} dr + K_a \int_a^\infty (r+1)^{-1-\kappa} dr = c_\nu + \log a + \kappa^{-1}(a+1)^{-\kappa} K_a. \end{aligned}$$

$\square$

The next lemma will play a crucial role in the proof of the Carleman estimates.

**Lemma 2.3.** *For  $0 < r < a$ ,  $r \neq 1$ , we have the inequality*

$$(2.6) \quad (\mu\psi'^2)'(r) \geq C_\nu(r+1)^{-1-\nu},$$

*with some constant  $C_\nu > 0$ . For all  $r > a$  we have the inequality*

$$(2.7) \quad (\mu\psi'^2)'(r) \geq -\tilde{C}_\kappa a^{-3+2s} \mu'(r),$$

*with some constant  $\tilde{C}_\kappa > 0$ .*

*Proof.* For  $0 < r < 1$  we have

$$(\mu\psi'^2)'(r) = c_\nu^2(2r+1),$$

which clearly implies (2.6) in this case. For  $1 < r < a$  we have

$$(\mu\psi'^2)'(r) = ((1-(r+1)^{-\nu})^2)' = 2\nu(1-(r+1)^{-\nu})(r+1)^{-1-\nu} \geq 2\nu(1-2^{-\nu})(r+1)^{-1-\nu}.$$

For  $r > a$ , in view of (2.4) with  $j=1$ , we have

$$(\mu\psi'^2)'(r) = \mu'\psi'^2 + 2\mu\psi'\psi'' \geq -2(r+1)^{2s}\mu'\psi'|\psi''|$$

$$\geq -Ca^{2\kappa}(r+1)^{-3-2\kappa+2s} \geq -Ca^{-3+2s},$$

as desired.  $\square$

### 3. CARLEMAN ESTIMATES IN THE GENERAL CASE

Let  $\mu$  and  $\psi$  be the functions introduced in the previous section and set  $\varphi = \tau\psi$ , where  $\tau = \tau_0 h^{-1}$ ,  $\tau_0 \gg 1$  being a parameter independent of  $h$  to be chosen later on. In what follows we set  $\mathcal{D}_r = ih\partial_r$ . In this section we will prove the following

**Theorem 3.1.** *Under the conditions of Theorem 1.1, there are constants  $C > 0$  and  $0 < h_0 \ll 1$  such that for all  $0 < h \leq h_0$ ,  $0 < \varepsilon \leq 1$ ,  $s > 1/2$ , and for all functions  $f \in H^2(\mathbb{R}^d)$  satisfying*

$$(|x| + 1)^s (P(h) - E \pm i\varepsilon)f \in L^2(\mathbb{R}^d)$$

we have the estimate

$$(3.1) \quad \begin{aligned} & \|(|x| + 1)^{-s} e^{\varphi/h} f\|_{L^2} \leq Cah^{-1} \|(|x| + 1)^s e^{\varphi/h} (P(h) - E \pm i\varepsilon)f\|_{L^2} \\ & + Ca (\tau h^{-1} \varepsilon)^{1/2} \|e^{\varphi/h} f\|_{L^2} + Ca (\tau^{-1} h^{-1} \varepsilon)^{1/2} \|e^{\varphi/h} \mathcal{D}_r f\|_{L^2}. \end{aligned}$$

*Proof.* Clearly, it suffices to prove (3.1) for  $0 < s - 1/2 \ll 1$  as this would imply (3.1) for all  $s > 1/2$ . So, in what follows  $s$  will be as in Section 2 with additional restrictions made later on. We write the operator  $P(h)$  in the form

$$P(h) = -h^2 \Delta + ihb^L \cdot \nabla + ih\nabla \cdot b^L + 2ihb^S \cdot \nabla + \tilde{V}_L + \tilde{V}_S,$$

where

$$\begin{aligned} \tilde{V}_L &= |b^L|^2 + V_L, \\ \tilde{V}_S &= ih \operatorname{div} b^S + 2b^S \cdot b^L + |b^S|^2 + V_S. \end{aligned}$$

Note that our conditions guarantee that

$$|\tilde{V}_S(x)| \lesssim (|x| + 1)^{-1-\rho}, \quad \partial_r \tilde{V}_L(x) \lesssim (|x| + 1)^{-1-\rho}.$$

Moreover,  $\tilde{V}_L$  satisfies (1.1) with a new decreasing function  $p$ .

We will write  $P(h)$  in the polar coordinates  $(r, w) \in \mathbb{R}^+ \times \mathbb{S}^{d-1}$ ,  $r = |x|$ ,  $w = x/|x|$ . Recall that  $L^2(\mathbb{R}^d) = L^2(\mathbb{R}^+ \times \mathbb{S}^{d-1}, r^{d-1} dr dw)$ . In what follows in this section we denote by  $\|\cdot\|$  and  $\langle \cdot, \cdot \rangle$  the norm and the scalar product in  $L^2(\mathbb{S}^{d-1})$ . We will make use of the identity

$$(3.2) \quad r^{(d-1)/2} \Delta r^{-(d-1)/2} = \partial_r^2 + r^{-2} \Delta_w - (d-1)(d-3)(2r)^{-2}$$

where  $\Delta_w$  denotes the negative Laplace-Beltrami operator on  $\mathbb{S}^{d-1}$ . We also have

$$(3.3) \quad r^{(d-1)/2} \partial_{x_j} r^{-(d-1)/2} = w_j \partial_r + r^{-1} q_j(w, \partial_w)$$

where  $q_j$  is a first-order differential operator on  $\mathbb{S}^{d-1}$  which is antisymmetric with respect to the scalar product in  $L^2(\mathbb{S}^{d-1})$ . Clearly, we have the estimates

$$(3.4) \quad \|q_j(w, \partial_w)v\| \lesssim \|(-\Delta_w)^{1/2}v\| + \|v\| \quad \text{for all } v \in H^1(\mathbb{S}^{d-1}).$$

Set  $\Lambda = -h^2 \Delta_w$ ,  $\mathcal{D}_r = ih\partial_r$  and  $Q_j = ihq_j(w, \partial_w)$ . Then the operator  $Q_j$  is symmetric with respect to the scalar product in  $L^2(\mathbb{S}^{d-1})$ . Moreover, (3.4) implies the estimates

$$(3.5) \quad \|Q_j v\| \lesssim \|\Lambda^{1/2}v\| + h\|v\| \quad \text{for all } v \in H^1(\mathbb{S}^{d-1}).$$

Set  $u(r, w) = r^{(d-1)/2} e^{\varphi/h} f(rw)$  and

$$\begin{aligned} \mathcal{P}(h) &= r^{(d-1)/2} (P(h) - E) r^{-(d-1)/2}, \\ \mathcal{P}_\varphi(h) &= e^{\varphi/h} \mathcal{P}(h) e^{-\varphi/h}. \end{aligned}$$

We now define the function  $\sigma$  as follows. We put  $\sigma = 0$  if (1.7) holds and we put

$$\sigma(r, w) = \sum_{j=1}^d w_j b_j^L(rw)$$

if (1.8) holds. Using (3.2) and (3.3) we can write the operator  $\mathcal{P}(h)$  in the coordinates  $(r, w)$  as follows

$$\begin{aligned} \mathcal{P}(h) &= \mathcal{D}_r^2 + r^{-2}\Lambda - E + h^2(d-1)(d-3)(2r)^{-2} + \tilde{V}_L + \tilde{V}_S \\ &+ r^{-1} \sum_{j=1}^d (b_j^L Q_j + Q_j b_j^L) + 2 \sum_{j=1}^d w_j b_j \mathcal{D}_r + 2r^{-1} \sum_{j=1}^d b_j^S Q_j + ih \sum_{j=1}^d w_j \partial_r (b_j^L), \end{aligned}$$

if (1.7) holds, and

$$\begin{aligned} \mathcal{P}(h) &= (\mathcal{D}_r + \sigma)^2 + r^{-2}\Lambda - E + h^2(d-1)(d-3)(2r)^{-2} + \tilde{V}_L + \tilde{V}_S - \sigma^2 - 2\sigma \sum_{j=1}^d w_j b_j^S \\ &+ r^{-1} \sum_{j=1}^d (b_j^L Q_j + Q_j b_j^L) + 2 \sum_{j=1}^d w_j b_j^S (\mathcal{D}_r + \sigma) + 2r^{-1} \sum_{j=1}^d b_j^S Q_j, \end{aligned}$$

if (1.8) holds. Since the function  $\varphi$  depends only on the variable  $r$ , in the first case we get

$$\begin{aligned} \mathcal{P}_\varphi(h) &= \mathcal{D}_r^2 + r^{-2}\Lambda - E + h^2(d-1)(d-3)(2r)^{-2} - 2i\varphi' \mathcal{D}_r + W_L + W_S \\ &+ 2 \sum_{j=1}^d w_j b_j \mathcal{D}_r + r^{-1} \sum_{j=1}^d (b_j^L Q_j + Q_j b_j^L) + 2r^{-1} \sum_{j=1}^d b_j^S Q_j, \end{aligned}$$

where

$$\begin{aligned} W_L &= \tilde{V}_L - \varphi'^2, \\ W_S &= \tilde{V}_S + h\varphi'' - 2i\varphi' \sum_{j=1}^d w_j b_j + ih \sum_{j=1}^d w_j \partial_r (b_j^L). \end{aligned}$$

In the second case we get

$$\begin{aligned} \mathcal{P}_\varphi(h) &= (\mathcal{D}_r + \sigma)^2 + r^{-2}\Lambda - E + h^2(d-1)(d-3)(2r)^{-2} - 2i\varphi' (\mathcal{D}_r + \sigma) + W_L + W_S \\ &+ 2 \sum_{j=1}^d w_j b_j^S (\mathcal{D}_r + \sigma) + r^{-1} \sum_{j=1}^d (b_j^L Q_j + Q_j b_j^L) + 2r^{-1} \sum_{j=1}^d b_j^S Q_j, \end{aligned}$$

where

$$\begin{aligned} W_L &= \tilde{V}_L - \sigma^2 - \varphi'^2, \\ W_S &= \tilde{V}_S + h\varphi'' - 2(i\varphi' + \sigma) \sum_{j=1}^d w_j b_j^S. \end{aligned}$$

For  $r > 0$ ,  $r \neq 1$ ,  $r \neq a$ , introduce the function

$$\begin{aligned} F(r) &= - \langle (r^{-2}\Lambda + h^2(d-1)(d-3)(2r)^{-2} - E + W_L)u(r, \cdot), u(r, \cdot) \rangle \\ &+ \|(\mathcal{D}_r + \sigma)u(r, \cdot)\|^2 - 2r^{-1} \sum_{j=1}^d \operatorname{Re} \langle b_j^L Q_j u(r, \cdot), u(r, \cdot) \rangle. \end{aligned}$$

Observe now that in the first case we have

$$\frac{h}{2} \frac{d}{dr} \|\mathcal{D}_r u\|^2 = \operatorname{Im} \langle \mathcal{D}_r^2 u, \mathcal{D}_r u \rangle = \operatorname{Im} \langle \mathcal{P}_\varphi(h)u, \mathcal{D}_r u \rangle$$



$$\begin{aligned}
& -\operatorname{Im} \langle (r^{-2}\Lambda + h^2(d-1)(d-3)(2r)^{-2} - E + W_L + W_S)u, \mathcal{D}_r u \rangle \\
& + 2\varphi' \|\mathcal{D}_r u\|^2 - 2 \sum_{j=1}^d \operatorname{Im} \langle w_j b_j \mathcal{D}_r u, \mathcal{D}_r u \rangle \\
& - \sum_{j=1}^d \operatorname{Im} \langle r^{-1}(b_j^L Q_j + Q_j b_j^L)u, \mathcal{D}_r u \rangle - 2 \sum_{j=1}^d \operatorname{Im} \langle r^{-1} b_j^S Q_j u, \mathcal{D}_r u \rangle,
\end{aligned}$$

while in the second case we have

$$\begin{aligned}
& \frac{h}{2} \frac{d}{dr} \|(\mathcal{D}_r + \sigma)u\|^2 = \operatorname{Im} \langle (\mathcal{D}_r + \sigma)^2 u, (\mathcal{D}_r + \sigma)u \rangle = \operatorname{Im} \langle \mathcal{P}_\varphi(h)u, (\mathcal{D}_r + \sigma)u \rangle \\
& - \operatorname{Im} \langle (r^{-2}\Lambda + h^2(d-1)(d-3)(2r)^{-2} - E + W_L + W_S)u, (\mathcal{D}_r + \sigma)u \rangle \\
& + 2\varphi' \|(\mathcal{D}_r + \sigma)u\|^2 - 2 \sum_{j=1}^d \operatorname{Im} \langle w_j b_j^S (\mathcal{D}_r + \sigma)u, (\mathcal{D}_r + \sigma)u \rangle \\
& - \sum_{j=1}^d \operatorname{Im} \langle r^{-1}(b_j^L Q_j + Q_j b_j^L)u, (\mathcal{D}_r + \sigma)u \rangle \\
& - 2 \sum_{j=1}^d \operatorname{Im} \langle r^{-1} b_j^S Q_j u, (\mathcal{D}_r + \sigma)u \rangle.
\end{aligned}$$

On the other hand, we have the identities

$$\begin{aligned}
& \operatorname{Im} \langle w_j b_j \mathcal{D}_r u, \mathcal{D}_r u \rangle = 0, \\
& \operatorname{Re} \frac{d}{dr} \langle b_j^L Q_j u, u \rangle = \operatorname{Re} \langle \partial_r (b_j^L) Q_j u, u \rangle \\
& - h^{-1} \operatorname{Im} \langle (b_j^L Q_j + Q_j b_j^L)u, \mathcal{D}_r u \rangle
\end{aligned}$$

in the first case, and

$$\begin{aligned}
& \operatorname{Im} \langle w_j b_j^S (\mathcal{D}_r + \sigma)u, (\mathcal{D}_r + \sigma)u \rangle = 0, \\
& \operatorname{Re} \frac{d}{dr} \langle b_j^L Q_j u, u \rangle = \operatorname{Re} \langle \partial_r (b_j^L) Q_j u, u \rangle \\
& - h^{-1} \operatorname{Im} \langle (b_j^L Q_j + Q_j b_j^L)u, (\mathcal{D}_r + \sigma)u \rangle - \operatorname{Re} \langle b_j^L u, [q_j, \sigma]u \rangle
\end{aligned}$$

in the second case. Using the above identities we find that in both cases the first derivative of  $F$  is given by

$$\begin{aligned}
F'(r) &= \frac{2}{r} \langle r^{-2}(\Lambda + (h/2)^2(d-1)(d-3))u, u \rangle - W'_L \|u\|^2 \\
& + 2r^{-2} \sum_{j=1}^d \operatorname{Re} \langle b_j^L Q_j u, u \rangle - 2r^{-1} \sum_{j=1}^d \operatorname{Re} \langle \partial_r (b_j^L) Q_j u, u \rangle + 2r^{-1} \sum_{j=1}^d \operatorname{Re} \langle b_j^L u, [q_j, \sigma]u \rangle \\
& + 2h^{-1} \operatorname{Im} \langle \mathcal{P}_\varphi(h)u, (\mathcal{D}_r + \sigma)u \rangle + 4h^{-1} \varphi' \|(\mathcal{D}_r + \sigma)u\|^2 - 2h^{-1} \operatorname{Im} \langle \widetilde{W}_S u, (\mathcal{D}_r + \sigma)u \rangle,
\end{aligned}$$

where

$$\widetilde{W}_S = W_S + 2r^{-1} \sum_{j=1}^d b_j^S Q_j.$$

Thus we obtain the identity

$$\begin{aligned}
(\mu F)' &= \mu' F + \mu F' \\
&= (2r^{-1}\mu - \mu') \langle r^{-2}(\Lambda + (h/2)^2(d-1)(d-3))u, u \rangle
\end{aligned}$$

$$\begin{aligned}
& +(E\mu' - (\mu W_L)')\|u\|^2 + (\mu' + 4h^{-1}\varphi'\mu)\|(\mathcal{D}_r + \sigma)u\|^2 \\
& - 2(r^{-1}\mu)' \sum_{j=1}^d \operatorname{Re} \langle b_j^L Q_j u, u \rangle - 2r^{-1}\mu \sum_{j=1}^d \operatorname{Re} \langle \partial_r(b_j^L) Q_j u, u \rangle \\
& + 2r^{-1}\mu \sum_{j=1}^d \operatorname{Re} \langle b_j^L u, [q_j, \sigma]u \rangle - 2h^{-1}\mu \operatorname{Im} \langle \widetilde{W}_S u, (\mathcal{D}_r + \sigma)u \rangle \\
& + 2h^{-1}\mu \operatorname{Im} \langle (\mathcal{P}_\varphi(h) \pm i\varepsilon)u, (\mathcal{D}_r + \sigma)u \rangle \mp 2\varepsilon h^{-1}\mu \operatorname{Re} \langle u, (\mathcal{D}_r + \sigma)u \rangle.
\end{aligned}$$

In view of (3.5), we have the estimates

$$(3.6) \quad \left\| \widetilde{W}_S u \right\|^2 \lesssim B_1 \|u\|^2 + B_2 \left\| r^{-1} \Lambda^{1/2} u \right\|^2,$$

where

$$\begin{aligned}
B_1(r) &= (r+1)^{-2-\rho} + (r+1)^{-2-\rho} \varphi'^2 + h^2 \varphi''^2 + h^2 r^{-2} (r+1)^{-2-\rho}, \\
B_2(r) &= (r+1)^{-2-\rho},
\end{aligned}$$

and

$$\begin{aligned}
(3.7) \quad & |\langle b_j^L Q_j u, u \rangle| \lesssim B_3 \|u\| \left\| \Lambda^{1/2} u \right\| + h B_3 \|u\|^2 \\
& \lesssim (\gamma_1^{-1} (r B_3)^2 + h B_3) \|u\|^2 + \gamma_1 \left\| r^{-1} \Lambda^{1/2} u \right\|^2,
\end{aligned}$$

$$(3.8) \quad |\langle \partial_r(b_j^L) Q_j u, u \rangle| \lesssim (\gamma_2^{-1} (r B_4)^2 + h B_4) \|u\|^2 + \gamma_2 \left\| r^{-1} \Lambda^{1/2} u \right\|^2,$$

where  $\gamma_1$  and  $\gamma_2$  are arbitrary positive functions of  $r$  to be fixed below and

$$B_3(r) = (r+1)^{-\rho}, \quad B_4(r) = (r+1)^{-1} B_3(r).$$

Write now (3.3) in the form

$$w_j \partial_r + r^{-1} q_j(w, \partial_w) = \partial_{x_j} - \frac{d-1}{4r^2} \frac{\partial r^2}{\partial x_j} = \partial_{x_j} - (d-1)(2r)^{-1} w_j.$$

Therefore we obtain the identity

$$[q_j, \sigma] = [r^{-1} q_j, r\sigma] = \partial_{x_j}(x \cdot b^L(x)) - w_j \partial_r(r\sigma).$$

Hence, in view of (1.4) and (1.8), we get the bound

$$|[q_j, \sigma]| \lesssim 1,$$

which implies

$$(3.9) \quad |\langle b_j^L u, [q_j, \sigma]u \rangle| \lesssim B_5 \|u\|^2,$$

where  $B_5(r) = B_3(r)$ . Since  $2r^{-1}\mu - \mu' > 0$  and  $d \geq 3$ , by (3.6), (3), (3.8) and (3.9), we get the inequality

$$\begin{aligned}
& (\mu F)' \geq (2r^{-1}\mu - \mu') \langle r^{-2} \Lambda u, u \rangle \\
& + (E\mu' - (\mu W_L)')\|u\|^2 + (\mu' + 4h^{-1}\varphi'\mu)\|(\mathcal{D}_r + \sigma)u\|^2 \\
& - 2|(r^{-1}\mu)'| \sum_{j=1}^d |\langle b_j^L Q_j u, u \rangle| - 2r^{-1}\mu \sum_{j=1}^d |\langle \partial_r(b_j^L) Q_j u, u \rangle| - 2r^{-1}\mu \sum_{j=1}^d |\langle b_j^L u, [q_j, \sigma]u \rangle| \\
& - 3h^{-2}\mu^2 (\mu' + 4h^{-1}\varphi'\mu)^{-1} \left\| \widetilde{W}_S u \right\|^2 - \frac{1}{3} (\mu' + 4h^{-1}\varphi'\mu) \|(\mathcal{D}_r + \sigma)u\|^2
\end{aligned}$$

$$\begin{aligned}
& -\frac{3h^{-2}\mu^2}{\mu'}\|(\mathcal{P}_\varphi(h) \pm i\varepsilon)u\|^2 - \frac{\mu'}{3}\|(\mathcal{D}_r + \sigma)u\|^2 - \varepsilon h^{-1}\mu\|u\|\|(\mathcal{D}_r + \sigma)u\| \\
& \geq A_1\|u\|^2 + A_2\|r^{-1}\Lambda^{1/2}u\|^2 - \frac{3h^{-2}\mu^2}{\mu'}\|(\mathcal{P}_\varphi(h) \pm i\varepsilon)u\|^2 - \varepsilon h^{-1}\mu\|u\|\|(\mathcal{D}_r + \sigma)u\|,
\end{aligned}$$

where

$$\begin{aligned}
A_1(r) &= (\varphi'^2\mu)' + (E - \tilde{V}_L)\mu' - C(r+1)^{-1-\rho}\mu - Ch^{-2}B_1\mu^2(\mu' + h^{-1}\varphi'\mu)^{-1} \\
&\quad - C|(r^{-1}\mu)'|(\gamma_1^{-1}(rB_3)^2 + hB_3) - Cr^{-1}\mu(\gamma_2^{-1}(rB_4)^2 + hB_4 + B_5), \\
A_2(r) &= 2r^{-1}\mu - \mu' - Ch^{-2}B_2\mu^2(\mu' + h^{-1}\varphi'\mu)^{-1} \\
&\quad - C\gamma_1|(r^{-1}\mu)'| - C\gamma_2r^{-1}\mu.
\end{aligned}$$

We now choose the functions  $\gamma_1$  and  $\gamma_2$  so that

$$C\gamma_1|(r^{-1}\mu)'| = C\gamma_2r^{-1}\mu = \frac{1}{4}(2r^{-1}\mu - \mu').$$

Thus we conclude that the above inequality holds with

$$\begin{aligned}
A_1(r) &= (\varphi'^2\mu)' + (E - \tilde{V}_L)\mu' - Cr^{-1}(r+1)^{-\rho}\mu - Ch^{-2}B_1\mu^2(\mu' + h^{-1}\varphi'\mu)^{-1} \\
&\quad - C(rB_3)^2M_1 - ChB_3M_2, \\
2A_2(r) &= 2r^{-1}\mu - \mu' - Ch^{-2}B_2\mu^2(\mu' + h^{-1}\varphi'\mu)^{-1},
\end{aligned}$$

with a new constant  $C > 0$ , where

$$\begin{aligned}
M_1(r) &= (2r^{-1}\mu - \mu')^{-1} (|(r^{-1}\mu)'|^2 + (r(r+1))^{-2}\mu^2), \\
M_2(r) &= |(r^{-1}\mu)'| + (r(r+1))^{-1}\mu.
\end{aligned}$$

Now we will use Lemmas 2.1 and 2.3 to bound the functions  $A_1$  and  $A_2$  from below. Observe first that  $M_1(r) = M_2(r) = 1$  for  $r < 1$ . For  $1 < r < a$  we have the bounds

$$M_1(r) \lesssim (r+1)^{-1+\ell}, \quad M_2(r) \lesssim 1,$$

while for  $r > a$  we have the bounds

$$M_1(r) \lesssim (r+1)^{-3+2s}\mu'(r), \quad M_2(r) \lesssim (r+1)^{-2+2s}\mu'(r).$$

Therefore, for  $0 < r < 1$  we have

$$\begin{aligned}
A_1(r) &\geq E\mu'(r) + C_\nu^\flat\tau^2 - Ch^{-2} - Ch^{-1}\tau, \quad C_\nu^\flat > 0, \\
A_2(r) &\geq 2^{-1} - Ch^{-1}\tau^{-1}.
\end{aligned}$$

We take  $\tau = \tau_0 h^{-1}$ , where  $\tau_0 \gg 1$  is independent of  $h$ . It is clear that if  $\tau_0$  is big enough we can arrange that

$$A_1(r) \geq E\mu'(r), \quad A_2(r) \geq 0, \quad \text{for } 0 < r < 1.$$

For  $1 < r < a$  we have  $r^{-1}\mu(r) < \mu'(r)$  and

$$(3.10) \quad (rB_3)^2M_1(r) \lesssim (r+1)^{\ell-2\rho}\mu'(r) \lesssim (r+1)^{-\rho}\mu'(r)$$

if  $\ell < \rho$ . Hence

$$\begin{aligned}
A_1(r) &\geq \tau^2 C_\nu(r+1)^{-1-\nu} + (E - \tilde{V}_L)\mu' - C(r+1)^{-\rho}\mu' \\
&\quad - Ch^{-1}\tau^{-1}(r+1)^{-\rho}\mu' - Ch\tau(r+1)^{-2}\mu' - Ch^{-1}\tau(r+1)^{-1-\rho}, \\
A_2(r) &\geq 2^{-1}\ell r^{1-\ell} - Ch^{-1}\tau^{-1}r^{1-\rho}.
\end{aligned}$$

Since  $\tilde{V}_L$  satisfies (1.1), there exists  $\lambda_0 \gg 1$ , independent of  $h$  and  $\tau_0$ , such that

$$(3.11) \quad \tilde{V}_L(rw) + C(r+1)^{-\rho} \leq E/4 \quad \text{for } r \geq \lambda_0.$$

On the other hand, for  $r < \lambda_0$ , we have

$$(3.12) \quad \tilde{V}_L \mu'(r) + C(r+1)^{-\rho} \mu'(r) \lesssim (r+1)^{-1-\nu}.$$

Similarly, there exists  $\lambda \gg 1$ , independent of  $h$  but depending on  $\tau_0$ , such that

$$C\tau_0(r+1)^{-2} \leq E/4 \quad \text{for } r \geq \lambda,$$

and

$$C\tau_0(r+1)^{-2} \mu'(r) \lesssim (r+1)^{-1-\nu} \quad \text{for } r < \lambda.$$

Therefore, if we choose  $\nu < \rho$ ,  $\ell < \rho$ , taking  $\tau_0$  big enough and  $h$  small enough we can arrange that

$$A_1(r) \geq 3^{-1} E \mu'(r), \quad A_2(r) \geq 0, \quad \text{for } 1 < r < a.$$

We now take  $1/2 < s < (1+\rho)/2$ ,  $0 < \kappa < 1+\rho-2s$ . In view of (1.1) with  $\tilde{V}_L$ , for  $r > a$  we have

$$\begin{aligned} A_1(r) &\geq 2^{-1} E \mu'(r) - C \mu'(r) (\tau^2 a^{-3+2s} + a^{-\rho+2s-1} + \tau h^{-1} a^{-4+2s}), \\ A_2(r) &\geq 2^{-1} a^2 (r+1)^{-1} (1 - C h^{-1} \tau^{-1} a^{-1}). \end{aligned}$$

Taking  $a = h^{-m}$  with  $m > 0$  big enough and  $h$  small enough we can arrange that

$$A_1(r) \geq 3^{-1} E \mu'(r), \quad A_2(r) \geq 0, \quad \text{for } r > a.$$

By the above inequalities we obtain

$$(3.13) \quad (\mu F)' \geq \frac{E}{3} \mu' \|u\|^2 - \frac{3h^{-2} \mu^2}{\mu'} \|(\mathcal{P}_\varphi(h) \pm i\varepsilon)u\|^2 - \varepsilon h^{-1} \mu \|u\| \|(\mathcal{D}_r + \sigma)u\|.$$

Integrating (3.13) with respect to  $r$  and using that

$$\int_0^\infty (\mu F)' dr = 0,$$

we obtain the estimate

$$(3.14) \quad \frac{E}{3} \int_0^\infty \mu' \|u\|^2 dr \leq 3h^{-2} \int_0^\infty \frac{\mu^2}{\mu'} \|(\mathcal{P}_\varphi(h) \pm i\varepsilon)u\|^2 dr + \varepsilon h^{-1} \int_0^\infty \mu \|u\| \|(\mathcal{D}_r + \sigma)u\| dr.$$

Using that  $\mu(r) \lesssim \zeta(r)a^2$ , where  $\zeta(r) = r(r+1)^{-1}$ , together with (2.3) and (2.4) we get from (3.14),

$$\begin{aligned} \int_0^\infty (r+1)^{-2s} \|u\|^2 dr &\leq C a^2 h^{-2} \int_0^\infty (r+1)^{2s} \|(\mathcal{P}_\varphi(h) \pm i\varepsilon)u\|^2 dr \\ &+ C \varepsilon h^{-1} a^2 \int_0^\infty (\tau \|u\|^2 + \tau^{-1} \zeta^2 \|\mathcal{D}_r u\|^2) dr \end{aligned} \quad (3.15)$$

with some constant  $C > 0$  independent of  $h$  and  $\varepsilon$ . Since

$$\mathcal{D}_r u = r^{(d-1)/2} e^{\varphi/h} \mathcal{D}_r f + (i\varphi' + (d-1)(2r)^{-1})u,$$

we have

$$(3.16) \quad \zeta^2 \|\mathcal{D}_r u\|^2 \lesssim \left\| r^{(d-1)/2} e^{\varphi/h} \mathcal{D}_r f \right\|^2 + \tau^2 \|u\|^2.$$

Clearly, the estimate (3.1) follows from (3) and (3.16).  $\square$

4. CARLEMAN ESTIMATES WHEN  $V_S \equiv 0$  AND  $b^S \equiv 0$ 

In this section we will prove the following

**Theorem 4.1.** *Under the conditions of Theorem 1.1 with (1.8), if  $V_S \equiv 0$  and  $b^S \equiv 0$ , the estimate (3.1) holds true with new parameters  $a, \tau \gg 1$  independent of  $h$ .*

*Proof.* We will make use of the Carleman estimates obtained in the previous section under the condition (1.8), which in our case take a much simpler form. Indeed,  $V_S \equiv 0$ ,  $b^S \equiv 0$  imply that  $\tilde{V}_S \equiv 0$  and  $\tilde{W}_S = h\varphi''$ . Hence  $B_1 = h^2\varphi''^2$  and  $B_2 = 0$ . The functions  $B_3$ ,  $B_4$  and  $B_5$  remain unchanged. Therefore the functions  $A_1$  and  $A_2$  take the form

$$\begin{aligned} A_1(r) &= (\varphi'^2\mu)' + (E - \tilde{V}_L)\mu' - Cr^{-1}(r+1)^{-\rho}\mu \\ &\quad - Ch\tau\mu(\psi'')^2(\psi')^{-1} - C(rB_3)^2M_1 - ChB_3M_2, \\ 2A_2(r) &= 2r^{-1}\mu - \mu' > 0. \end{aligned}$$

For  $0 < r < a$ ,  $r \neq 1$ , in view of (3.10), (3.11) and (3.12), we have

$$A_1(r) \geq \tau^2 C_\nu (r+1)^{-1-\nu} + (E - \tilde{V}_L)\mu' - C(r+1)^{-\rho}\mu' - Ch\tau(r+1)^{-2}\mu' \geq E\mu'/3,$$

if  $\tau$  is taken big enough, independent of  $h$ , and  $h$  is taken small enough. For  $r > a$  we have

$$A_1(r) \geq 2^{-1}E\mu' - C\mu'(\tau^2 a^{-3+2s} + a^{-\rho+2s-1}) \geq E\mu'/3,$$

if  $a$  is taken big enough, independent of  $h$ . Thus we conclude that the estimate (3) still holds in this case with parameters  $a, \tau \gg 1$  independent of  $h$ . Clearly, this implies the theorem.  $\square$

## 5. CARLEMAN ESTIMATES FOR HÖLDER POTENTIALS

In this section we will prove the following

**Theorem 5.1.** *Under the conditions of Theorem 1.2, the estimate (3.1) holds true with new parameters  $a = h^{-m}$  and  $\tau = \tau_0 h^{-n+1}$  with some positive constants  $m$  and  $\tau_0$ .*

*Proof.* Let  $\phi_0 \in C_0^\infty([0, 1])$ ,  $\phi_0 \geq 0$ , be a real-valued function independent of  $h$  such that  $\int_0^\infty \phi_0(t)dt = 1$ . If  $V_S$  satisfies (1.21), we approximate it by the function

$$V_{\theta_1}(r, w) = \theta_1^{-1} \int_0^\infty \phi_0((r' - r)/\theta_1) V_S(r', w) dr' = \int_0^\infty \phi_0(t) V_S(r + \theta_1 t, w) dt$$

where  $0 < \theta_1 < 1$  is a parameter depending on  $h$  to be fixed later on. We have

$$\begin{aligned} |V_S(r, w) - V_{\theta_1}(r, w)| &\leq \int_0^\infty \phi_0(t) |V_S(r + \theta_1 t, w) - V_S(r, w)| dt \\ (5.1) \quad &\lesssim \theta_1^\alpha (r+1)^{-3-\rho} \int_0^\infty t^\alpha \phi_0(t) dt \lesssim \theta_1^\alpha (r+1)^{-3-\rho}. \end{aligned}$$

Since  $V_S$  satisfies (1.1), the above bound implies

$$(5.2) \quad V_{\theta_1}(r, w) \leq p(r) + \mathcal{O}((r+1)^{-3-\rho}).$$

Clearly,  $V_{\theta_1}$  is  $C^1$  with respect to the variable  $r$  and its first derivative is given by

$$\begin{aligned} \partial_r V_{\theta_1}(r, w) &= -\theta_1^{-2} \int_0^\infty \phi_0'((r' - r)/\theta_1) V_S(r', w) dr' \\ &= -\theta_1^{-1} \int_0^\infty \phi_0'(t) V_S(r + \theta_1 t, w) dt = -\theta_1^{-1} \int_0^\infty \phi_0'(t) (V_S(r + \theta_1 t, w) - V_S(r, w)) dt \end{aligned}$$

where we have used that  $\int_0^\infty \phi'_0(t)dt = 0$ . Hence

$$(5.3) \quad |\partial_r V_{\theta_1}(r, w)| \lesssim \theta_1^{-1+\alpha}(r+1)^{-3-\rho} \int_0^\infty t^\alpha |\phi'_0(t)| dt \lesssim \theta_1^{-1+\alpha}(r+1)^{-3-\rho}.$$

Let  $\phi \in C_0^\infty(\mathbb{R}^d)$ ,  $\phi \geq 0$ , be a real-valued function independent of  $h$  such that  $\int_{\mathbb{R}^d} \phi(x)dx = 1$ . If  $b^S$  satisfies (1.22), we approximate it by the function

$$b_{\theta_2}(x) = \theta_2^{-d} \int_{\mathbb{R}^d} \phi((x' - x)/\theta_2) b^S(x') dx' = \int_{\mathbb{R}^d} \phi(y) b^S(x + \theta_2 y) dy$$

where  $0 < \theta_2 < 1$  is a parameter depending on  $h$  to be fixed later on. We have

$$(5.4) \quad \begin{aligned} |b^S(x) - b_{\theta_2}(x)| &\leq \int_{\mathbb{R}^d} \phi(y) |b^S(x + \theta_2 y) - b^S(x)| dy \\ &\lesssim \theta_2^{\alpha'} (r+1)^{-3/2-\rho_1} \int_{\mathbb{R}^d} |y|^{\alpha'} \phi(y) dy \lesssim \theta_2^{\alpha'} (r+1)^{-3/2-\rho_1}. \end{aligned}$$

By (1.23) and (5),

$$(5.5) \quad |b_{\theta_2}(rw)| \lesssim (r+1)^{-\rho_2} + (r+1)^{-3/2-\rho_1}.$$

If  $\beta$  is a multi-index such that  $|\beta| = 1$ , we have

$$\begin{aligned} \partial_x^\beta b_{\theta_2}(x) &= -\theta_2^{-d-1} \int_{\mathbb{R}^d} (\partial_x^\beta \phi)((x' - x)/\theta_2) b^S(x') dx' = -\theta_2^{-1} \int_{\mathbb{R}^d} (\partial_y^\beta \phi)(y) b^S(x + \theta_2 y) dy \\ &= -\theta_2^{-1} \int_{\mathbb{R}^d} (\partial_y^\beta \phi)(y) (b^S(x + \theta_2 y) - b^S(x)) dy. \end{aligned}$$

Hence

$$(5.6) \quad \begin{aligned} |\partial_x^\beta b_{\theta_2}(x)| &\leq \theta_2^{-1} \int_{\mathbb{R}^d} |(\partial_y^\beta \phi)(y)| |b^S(x + \theta_2 y) - b^S(x)| dy \\ &\lesssim \theta_2^{-1+\alpha'} (r+1)^{-3/2-\rho_1} \int_{\mathbb{R}^d} |y|^{\alpha'} |(\partial_y^\beta \phi)(y)| dy \lesssim \theta_2^{-1+\alpha'} (r+1)^{-3/2-\rho_1}. \end{aligned}$$

By (5.5) and (5) we deduce that

$$(5.7) \quad |\partial_r b_{\theta_2}(rw)| \lesssim \theta_2^{-1+\alpha'} (r+1)^{-3/2-\rho_1},$$

$$(5.8) \quad |\partial_x^\beta (x \cdot b_{\theta_2}(x))| \lesssim \theta_2^{-1+\alpha'} (r+1)^{-1/2-\rho_1} + (r+1)^{-\rho_2}.$$

It is also easy to see that (1.6) implies

$$(5.9) \quad |\operatorname{div}(b^S(x) - b_{\theta_2}(x))| \lesssim (r+1)^{-1-\rho}.$$

We will make use of the Carleman estimates obtained in Section 3 under the condition (1.8) with  $V_S$ ,  $V_L$ ,  $b^S$  and  $b^L$  replaced by  $V_S - V_{\theta_1}$ ,  $V_L + V_{\theta_1}$ ,  $b^S - b_{\theta_2}$  and  $b^L + b_{\theta_2}$ , respectively. Using the above estimates one can easily check that our assumptions guarantee that the new functions  $\tilde{V}_S$ ,  $\tilde{V}_L$  and  $\sigma$  satisfy the bounds

$$\begin{aligned} |\tilde{V}_S| &\lesssim \left( \theta_1^\alpha + \theta_2^{\alpha'} \right) (r+1)^{-3-\rho} + h(r+1)^{-1-\rho}, \\ \partial_r(\tilde{V}_L - \sigma^2) &\lesssim (r+1)^{-1-\rho} + \left( \theta_1^{-1+\alpha} + \theta_2^{-1+\alpha'} \right) (r+1)^{-3-\rho}, \end{aligned}$$

with some  $\rho > 0$ . Moreover,  $\tilde{V}_L$  satisfies (1.1) with a new decreasing function  $p$  independent of the parameters  $\theta_1$  and  $\theta_2$ . Then the functions  $B_j$  in this case become

$$B_1(r) = \left( \theta_1^{2\alpha} + \theta_2^{2\alpha'} \right) (r+1)^{-6-\rho} + h^2 (r+1)^{-2-\rho} + h^2 \theta_2^{2\alpha'} r^{-2} (r+1)^{-3-\rho}$$

$$\begin{aligned}
& +\theta_2^{2\alpha'}(r+1)^{-3-\rho}\varphi'^2 + h^2\varphi''^2, \\
B_2(r) &= \theta_2^{2\alpha'}(r+1)^{-3-\rho}, \quad B_3(r) = (r+1)^{-\rho}, \\
B_4(r) &= (r+1)^{-1}B_3(r) + \theta_2^{-1+\alpha'}(r+1)^{-3/2-\rho}, \\
B_5(r) &= B_3(r) + \theta_2^{-1+\alpha'}(r+1)^{-2-\rho}.
\end{aligned}$$

For the new functions  $A_1$  and  $A_2$  we obtain

$$\begin{aligned}
A_1(r) &= \tau^2(\psi'^2\mu)' + (E - \tilde{V}_L)\mu' - C\mu r^{-1}(r+1)^{-\rho} \\
& - C \left( M_2\theta_1^{-1+\alpha} + M_2\theta_2^{-1+\alpha'} + M_1\theta_2^{-2+2\alpha'} \right) (r+1)^{-1-\rho} \\
& - Ch^{-1}\tau^{-1} \left( \theta_1^{2\alpha} + \theta_2^{2\alpha'} \right) (r+1)^{-6-\rho}\mu(\psi')^{-1} - C\tau h^{-1}\theta_2^{2\alpha'}(r+1)^{-3-\rho}\mu\psi' \\
& - Ch\tau^{-1}(r+1)^{-2-\rho}\mu(\psi')^{-1} - Ch\tau\mu(\psi'')^2(\psi')^{-1} \\
& - C\theta_2^{2\alpha'}\mu^2(r\mu')^{-2}(r+1)^{-3-\rho}\mu' - C(rB_3)^2M_1 - ChB_3M_2, \\
2A_2(r) &= 2r^{-1}\mu - \mu' - Ch^{-1}\tau^{-1}\theta_2^{2\alpha'}(r+1)^{-3-\rho}\mu(\psi')^{-1},
\end{aligned}$$

where the functions  $M_1$  and  $M_2$  are the same as in Section 3. Note that for  $1 < r < a$  we have

$$\mu(\psi')^{-1} = \mathcal{O}(r^3), \quad \mu\psi' = \mathcal{O}(r), \quad \mu(\psi'')^2(\psi')^{-1} = \mathcal{O}(r^{-1}), \quad \mu^2(r\mu')^{-2} = \mathcal{O}(1).$$

Set

$$\chi_1 = \tau^{-2}\theta_1^{-1+\alpha}, \quad \eta_1 = h^{-1}\tau^{-3}\theta_1^{2\alpha}, \quad \chi_2 = \tau^{-2}\theta_2^{-2+2\alpha'}, \quad \eta_2 = h^{-1}\tau^{-1}\theta_2^{2\alpha'}.$$

We take  $\theta_1 = h^{2/(3+\alpha)}$ ,  $\theta_2 = h^{1/(1+\alpha')}$ ,  $\tau = h^{-n+1}\tau_0$ , where  $\tau_0 \gg 1$  is independent of  $h$ . It is easy to check that with this choice we have the inequalities

$$\chi_1 \leq \tau_0^{-2}, \quad \eta_1 \leq \tau_0^{-3}, \quad \chi_2 \leq \tau_0^{-2}, \quad \eta_2 \leq \tau_0^{-1}.$$

Therefore, we can make all these parameters small enough by taking  $\tau_0$  big enough. On the other hand, if we choose  $\nu < \rho$ ,  $\ell < \rho$  and the above parameters are small enough, it is easy to see that we have  $A_1 \geq E\mu'/3$  and  $A_2 \geq 0$ . Indeed, in view of (3.10), for  $r < a$ ,  $r \neq 1$ , we have

$$\begin{aligned}
A_1(r) &\geq \tau^2(C_\nu - C(\chi_1 + \eta_1 + \chi_2 + \eta_2))(r+1)^{-1-\nu} \\
& + (E - \tilde{V}_L)\mu' - C(r+1)^{-\rho}\mu' - C(h\tau + \theta_2^{2\alpha'})\mu' \\
& \geq 2^{-1}\tau^2C_\nu(r+1)^{-1-\nu} + (E - \tilde{V}_L)\mu' - C(r+1)^{-\rho}\mu' - C(h\tau + \theta_2^{2\alpha'})\mu', \\
A_2(r) &\geq C_\ell - C\eta_2 > 0.
\end{aligned}$$

Since  $n < 2$ , we can arrange  $C(h\tau + \theta_2^{2\alpha'}) \leq E/4$  by taking  $h$  small enough. Therefore, by (3.11) and (3.12) we obtain the desired lower bound for  $A_1$  in this case. For  $r > a$  the lower bounds for  $A_1$  and  $A_2$  are obtained in the same way as in Section 3 taking  $a = h^{-m}$  with  $m > 0$  big enough. Thus we conclude that the estimate (3) still holds in this case with the new parameters  $a, \tau$ , as desired.  $\square$

## 6. CARLEMAN ESTIMATES FOR $L^\infty$ POTENTIALS $V_S$ AND HÖLDER POTENTIALS $b^S$

In this section we will prove the following

**Theorem 6.1.** *Under the conditions of Theorem 1.3, the estimate (3.1) holds true with new parameters  $a = h^{-m}$ ,  $\tau = \tau_0 h^{-\frac{1-\alpha'}{1+\alpha'}}$  if  $\delta > (1 - \alpha')^{-1}$ ,  $\alpha' \leq \frac{1}{2}$ , and  $\tau = h^{-\frac{1}{2\delta-1}-\epsilon} \tau_0$ ,  $0 < \epsilon \ll 1$  being arbitrary, independent of  $h$ , if  $\alpha' \geq 1 - \delta^{-1}$ ,  $1 < \delta \leq 2$ , with some positive constants  $m$  and  $\tau_0$ .*

*Proof.* In this case we will make use of the Carleman estimates obtained in Section 3 under the condition (1.8) with  $b^S$  and  $b^L$  replaced by  $b^S - b_{\theta_2}$  and  $b^L + b_{\theta_2}$ , respectively. The functions  $V_S$  and  $V_L$  remain unchanged. Then the new functions  $\tilde{V}_S$ ,  $\tilde{V}_L$  and  $\sigma$  satisfy the bounds

$$\begin{aligned} |\tilde{V}_S| &\lesssim (r+1)^{-\delta} + \theta_2^{\alpha'} (r+1)^{-3-\rho} + h(r+1)^{-1-\rho}, \\ \partial_r(\tilde{V}_L - \sigma^2) &\lesssim (r+1)^{-1-\rho} + \theta_2^{-1+\alpha'} (r+1)^{-3-\rho}. \end{aligned}$$

Moreover,  $\tilde{V}_L$  satisfies (1.1) with a new decreasing function  $p$  independent of the parameter  $\theta_2$ . Then the functions  $B_j$  in this case become

$$\begin{aligned} B_1(r) &= (r+1)^{-2\delta} + \theta_2^{2\alpha'} (r+1)^{-6-\rho} + h^2 (r+1)^{-2-\rho} + h^2 \theta_2^{2\alpha'} r^{-2} (r+1)^{-3-\rho} \\ &\quad + \theta_2^{2\alpha'} (r+1)^{-3-\rho} \varphi'^2 + h^2 \varphi''^2, \\ B_2(r) &= \theta_2^{2\alpha'} (r+1)^{-3-\rho}, \quad B_3(r) = (r+1)^{-\rho}, \\ B_4(r) &= (r+1)^{-1} B_3(r) + \theta_2^{-1+\alpha'} (r+1)^{-3/2-\rho}, \\ B_5(r) &= B_3(r) + \theta_2^{-1+\alpha'} (r+1)^{-2-\rho}. \end{aligned}$$

For the new functions  $A_1$  and  $A_2$  we obtain

$$\begin{aligned} A_1(r) &= \tau^2 (\psi'^2 \mu)' + (E - \tilde{V}_L) \mu' - C \mu r^{-1} (r+1)^{-\rho} \\ &\quad - C \left( M_2 \theta_2^{-1+\alpha'} + M_1 \theta_2^{-2+2\alpha'} \right) (r+1)^{-1-\rho} \\ &\quad - C h^{-1} \tau^{-1} (r+1)^{-2\delta} \mu (\psi')^{-1} - C h^{-1} \tau^{-1} \theta_2^{2\alpha'} (r+1)^{-6-\rho} \mu (\psi')^{-1} \\ &\quad - C \tau h^{-1} \theta_2^{2\alpha'} (r+1)^{-3-\rho} \mu \psi' - C h \tau^{-1} (r+1)^{-2-\rho} \mu (\psi')^{-1} - C h \tau \mu (\psi'')^2 (\psi')^{-1} \\ &\quad - C \theta_2^{2\alpha'} \mu^2 (r \mu')^{-2} (r+1)^{-3-\rho} \mu' - C (r B_3)^2 M_1 - C h B_3 M_2, \\ 2A_2(r) &= 2r^{-1} \mu - \mu' - C h^{-1} \tau^{-1} \theta_2^{2\alpha'} (r+1)^{-3-\rho} \mu (\psi')^{-1}. \end{aligned}$$

We have to show that the functions  $A_1$  and  $A_2$  satisfy the same lower bounds as in the previous section. It suffices to do so for  $r < a$  since for  $r > a$  the lower bounds for  $A_1$  and  $A_2$  can be obtained in the same way as in Section 3 taking  $a = h^{-m}$  with  $m > 0$  big enough. Set

$$\chi = h^{-1} \tau^{-3}, \quad \chi_2 = \tau^{-2} \theta_2^{-2+2\alpha'}, \quad \eta_2 = h^{-1} \tau^{-1} \theta_2^{2\alpha'}.$$

We take  $\theta_2 = h^{\frac{1}{1+\alpha'}}$ ,  $\tau = h^{-\frac{1-\alpha'}{1+\alpha'}} \tau_0$ , where  $\tau_0 \gg 1$  is independent of  $h$ . Then we have

$$\chi_2 = \tau_0^{-2}, \quad \eta_2 = \tau_0^{-1}.$$

Consider first the case  $\alpha' = \frac{1}{2}$  and  $\delta > 2$ . Then

$$\chi = \tau_0^{-3}.$$

Choose  $\nu < \min\{\rho, 2(\delta - 2)\}$ . For  $r < a$ ,  $r \neq 1$ , we have

$$\begin{aligned} A_1(r) &\geq \tau^2 (C_\nu - C(\chi + \chi_2 + \eta_2)) (r+1)^{-1-\nu} \\ &\quad + (E - \tilde{V}_L) \mu' - C (r+1)^{-\rho} \mu' - C (h\tau + \theta_2^{2\alpha'}) \mu' \end{aligned}$$



$$\begin{aligned} &\geq 2^{-1}\tau^2 C_\nu(r+1)^{-1-\nu} + (E - \tilde{V}_L)\mu' - C(r+1)^{-\rho}\mu' - C(h\tau + \theta_2^{2\alpha'})\mu', \\ &A_2(r) \geq C_\ell - C\eta_2 > 0, \end{aligned}$$

if  $\tau_0$  is taken big enough. In the same way as in the previous section one can conclude that  $A_1 \geq E\mu'/3$ , which leads to the desired estimate in this case.

Let now  $\alpha' < \frac{1}{2}$  and  $(1 - \alpha')^{-1} < \delta \leq 2$ . Let  $1 < T < a$  be a parameter of the form

$$T = T_0(h\tau)^{-\frac{1}{2(\delta-1)}},$$

where  $T_0 \gg 1$  is independent of  $h$  and  $\tau_0$ . For  $T \leq r < a$  we have

$$Ch^{-1}\tau^{-1}(r+1)^{-2\delta}\mu(\psi')^{-1} \lesssim h^{-1}\tau^{-1}(r+1)^{-2\delta+2}\mu'(r) \lesssim T_0^{-2\delta+2}\mu'(r) \leq 4^{-1}E\mu'(r),$$

provided  $T_0$  is taken big enough. On the other hand, for  $r < T$ , we have

$$\begin{aligned} Ch^{-1}\tau^{-1}(r+1)^{-2\delta}\mu(\psi')^{-1} &\lesssim h^{-1}\tau^{-1}(r+1)^{-2\delta+3} \\ &\lesssim \chi\tau^2(r+1)^{-1-\nu}T^{4-2\delta+\nu} \lesssim \tilde{\chi}\tau^2(r+1)^{-1-\nu}, \end{aligned}$$

where

$$\tilde{\chi} = \chi(h\tau)^{-\frac{2-\delta+\nu/2}{\delta-1}} = h^{-\frac{1+\nu/2}{\delta-1}}\tau^{-\frac{2\delta-1+\nu/2}{\delta-1}} \leq \tau_0^{-\frac{2\delta-1+\nu/2}{\delta-1}},$$

provided  $\nu$  is chosen small enough. Hence we can make  $\tilde{\chi}$  small enough by taking  $\tau_0$  big enough, depending on  $T_0$ . Thus we conclude that the above lower bound for  $A_1$  holds with  $\chi$  replaced by  $\tilde{\chi}$  and  $E$  replaced by  $3E/4$ , which again leads to the inequality  $A_1 \geq E\mu'/3$  in this case.

Let now  $\alpha' \geq 1 - \delta^{-1}$ ,  $1 < \delta \leq 2$ . In this case we take  $\tau = h^{-\frac{1}{2\delta-1}-\epsilon}\tau_0$ , where  $0 < \epsilon \ll 1$  is arbitrary, independent of  $h$ , and  $\tau_0 \gg 1$  is independent of  $h$  and  $\epsilon$ . Let  $\theta_2$  be as above. Clearly, we have  $\tau \geq h^{-\frac{1-\alpha'}{1+\alpha'}}\tau_0$ , which implies the inequalities

$$\chi_2 \leq \tau_0^{-2}, \quad \eta_2 \leq \tau_0^{-1}.$$

Moreover, it is easy to see that the above bound of  $\tilde{\chi}$  still holds, provided  $\nu$  is chosen small enough depending on  $\epsilon$ . Therefore, the inequality  $A_1 \geq E\mu'/3$  holds in this case, too.  $\square$

## 7. RESOLVENT ESTIMATES

In this section we will show that the Carleman estimates (3.1), proved under various conditions, imply the resolvent bounds in Theorems 1.1, 1.2 and 1.3. These arguments are well-known when  $b \equiv 0$ , but the presence of a non-trivial magnetic potential does not change anything. The key points are the symmetry of the operator  $P(h)$  on the Hilbert space  $L^2(\mathbb{R}^d)$  and the fact that the potentials are  $L^\infty$ . In what follows we will sketch the proof for the sake of completeness. It follows from (3.1) and Lemma 2.2 that we have the estimate

$$(7.1) \quad \|(|x| + 1)^{-s}f\|_{L^2} \leq N_1\|(|x| + 1)^s(P(h) - E \pm i\varepsilon)f\|_{L^2} + N_2\varepsilon^{1/2}(\|f\|_{L^2} + \tau^{-1}\|h\nabla f\|_{L^2})$$

where

$$N_1 = ah^{-1}\exp(Ch^{-1}\tau \log a), \quad N_2 = (h\tau)^{1/2}N_1,$$

with a constant  $C > 0$  independent of  $h$  and  $\varepsilon$ . On the other hand, since the operator  $P(h)$  is symmetric, we have

$$\begin{aligned} \varepsilon\|f\|_{L^2}^2 &= \pm \text{Im} \langle (P(h) - E \pm i\varepsilon)f, f \rangle_{L^2} \\ &\leq \gamma^2 N_2^{-2} \|(|x| + 1)^{-s}f\|_{L^2}^2 + \gamma^{-2} N_2^2 \|(|x| + 1)^s(P(h) - E \pm i\varepsilon)f\|_{L^2}^2 \end{aligned}$$

for every  $\gamma > 0$ , which yields

$$(7.2) \quad N_2\varepsilon^{1/2}\|f\|_{L^2} \leq \gamma\|(|x| + 1)^{-s}f\|_{L^2} + \gamma^{-1}N_2^2\|(|x| + 1)^s(P(h) - E \pm i\varepsilon)f\|_{L^2}.$$

We also have

$$\begin{aligned} \operatorname{Re} \langle (P(h) - E \pm i\varepsilon)f, f \rangle_{L^2} &= \|(ih\nabla + b)f\|_{L^2}^2 + \operatorname{Re} \langle (V - E)f, f \rangle_{L^2} \\ &\geq \|h\nabla f\|_{L^2}^2 - C\|f\|_{L^2}^2. \end{aligned}$$

Thus we get

$$(7.3) \quad \|h\nabla f\|_{L^2} \lesssim \|f\|_{L^2} + \|(P(h) - E \pm i\varepsilon)f\|_{L^2}.$$

Taking  $\gamma$  small enough, independent of  $N_2$  and  $\varepsilon$ , by (7.1), (7.2) and (7.3) we obtain the estimate

$$(7.4) \quad \|(|x| + 1)^{-s}f\|_{L^2} \lesssim (N_1 + N_2^2)\|(|x| + 1)^s(P(h) - E \pm i\varepsilon)f\|_{L^2}.$$

It follows from (7.4) that the resolvent estimate

$$(7.5) \quad \|(|x| + 1)^{-s}(P(h) - E \pm i\varepsilon)^{-1}(|x| + 1)^s\|_{L^2 \rightarrow L^2} \lesssim N_1 + h\tau N_1^2 \lesssim N_1^2$$

holds for all  $0 < h \ll 1$ . Clearly, (7.5) implies the desired bounds for  $g_s^\pm$ .

## REFERENCES

- [1] F. CARDOSO AND G. VODEV, *Uniform estimates of the resolvent of the Laplace-Beltrami operator on infinite volume Riemannian manifolds*, Ann. Henri Poincaré **4** (2002), 673-691.
- [2] K. DATCHEV, *Quantitative limiting absorption principle in the semiclassical limit*, Geom. Funct. Anal. **24** (2014), 740-747.
- [3] K. DATCHEV, J. GALKOWSKI AND J. SHAPIRO, *Semiclassical resolvent bounds for compactly supported radial potentials*, J. Funct. Anal. **284** (2023), paper No. 109835, 28pp.
- [4] K. DATCHEV AND J. SHAPIRO, *Semiclassical estimates for scattering on the real line*, Commun. Math. Phys. **376** (2020), 2301-2308.
- [5] J. GALKOWSKI AND J. SHAPIRO, *Semiclassical resolvent bounds for long-range Lipschitz potentials*, IMRN **18** (2022), 14134-14150.
- [6] J. GALKOWSKI AND J. SHAPIRO, *Semiclassical resolvent bounds for weakly decaying potentials*, Math. Res. Lett. **29** (2022), 373-397.
- [7] V. GRASSELLI, *High frequency resolvent estimates for the magnetic Laplacian on non compact manifolds*, preprint 2024.
- [8] F. KLOPP AND M. VOGEL, *Semiclassical resolvent estimates for bounded potentials*, Pure Appl. Analysis **1** (2019), 1-25.
- [9] A. LARRAÍN-HUBACH AND J. SHAPIRO, *Semiclassical resolvent estimates for measure potentials on the real line*, J. Spectral Theory **14** (2024), 1033-1062.
- [10] A. LARRAÍN-HUBACH AND J. SHAPIRO, *Semiclassical estimates for the magnetic Schrödinger operator on the line*, preprint 2024.
- [11] C. J. MEROÑO, L. POTENCIANO-MACHADO AND M. SALO, *Resolvent estimates for the magnetic Schrödinger operator in dimensions  $\geq 2$* , Rev. Mat. Complut. **33** (2020), 619-641.
- [12] D. OBOVU, *Resolvent bounds for Lipschitz potentials in dimension two and higher with singularities at the origin*, J. Spectral Theory **14** (2024), 163-183.
- [13] J. SHAPIRO, *Semiclassical resolvent bounds in dimension two*, Proc. Amer. Math. Soc. **147** (2019), 1999-2008.
- [14] J. SHAPIRO, *Semiclassical resolvent bounds for compactly supported  $L^\infty$  potentials*, J. Spectral Theory **10** (2020), 651-672.
- [15] J. SHAPIRO, *Semiclassical resolvent bounds for short-range  $L^\infty$  potentials with singularities at the origin*, Asympt. Anal. **136** (2024), 157-180.
- [16] G. VODEV, *Resolvent estimates for the magnetic Schrödinger operator*, Anal. PDE **7** (2014), 1639-1648.
- [17] G. VODEV, *Semiclassical resolvent estimates for short-range  $L^\infty$  potentials*, Pure Appl. Analysis **1** (2019), 207-214.
- [18] G. VODEV, *Semiclassical resolvent estimates for Hölder potentials*, Pure Appl. Analysis **4** (2020), 841-860.
- [19] G. VODEV, *Semiclassical resolvent estimates for  $L^\infty$  potentials on Riemannian manifolds*, Ann. Henri Poincaré **21** (2020), 437-459.
- [20] G. VODEV, *Improved resolvent bounds for radial potentials*, Lett. Math. Phys. **111** (2021), 21pp.

- [21] G. VODEV, *Improved resolvent bounds for radial potentials.II*, Arch. Math. **119** (2022), 427-438.

UNIVERSITÉ DE NANTES, LABORATOIRE DE MATHÉMATIQUES JEAN LERAY, 2 RUE DE LA HOUSSINIÈRE, BP  
92208, 44322 NANTES CEDEX 03, FRANCE

*Email address:* [Georgi.Vodev@univ-nantes.fr](mailto:Georgi.Vodev@univ-nantes.fr)