

# Novel possible symmetries of $S$ -matrix generated by $\mathbb{Z}_2^n$ -graded Lie superalgebras

Ren Ito and Akio Nago

Department of Physics, Osaka Metropolitan University,  
Sugimoto Campus, Osaka 558-8585, Japan

January 17, 2025

arXiv:2501.07311v2 [hep-th] 16 Jan 2025

## Abstract

In this paper, we explore the  $\mathbb{Z}_2^n$ -graded Lie (super)algebras as novel possible generators of symmetries of  $S$ -matrix. As the results, we demonstrate that a  $\mathbb{Z}_2^n$ -graded extension of the supersymmetric algebra can be a symmetry of  $S$ -matrix. Furthermore, it turns out that a  $\mathbb{Z}_2^n$ -graded Lie algebra appears as internal symmetries. They are natural extensions of Coleman-Mandula theorem and Haag-Łopuszański-Sohnius theorem, which are the no-go theorems for generators of symmetries of  $S$ -matrix.

# 1 Introduction

Since the introduction of  $SU(4)$  symmetry in the classification of nuclei by Wigner in 1937, the properties and fundamental structure of particles have been described using group theory [1, 2]. The quark model introduced in the Standard Model is a model using  $SU(3)$  flavor symmetry proposed by Gell-Mann in 1964 [3]. It successfully describes the relationship among hadrons with the same spin. Furthermore, if we consider the  $SU(6)$  symmetry for quark flavor and spin, we can embed hadrons with different spins in the same representation.

However, such  $SU(6)$  symmetry that unify and describe different spins are approximate symmetries that exist in the non-relativistic limit, and despite some success, all attempts to generalize them to relativistic quantum theory have failed [4]. Indeed, it was shown by Coleman and Mandula in 1967 that such symmetries are forbidden in relativistic quantum field theory (QFT), known as the Coleman-Mandula theorem [5]. The theorem says that all possible symmetries of  $S$ -matrix generated by Lie algebra are only Poincaré symmetry and internal symmetry such that its generators commute with Poincaré algebra. This prevents us to unify Poincaré symmetry and internal symmetries into a single multiplet. The reason for the strong restriction of this theorem is that the symmetries allowed in the  $S$ -matrix of QFT were searched only in the framework of Lie algebras. If we want to avoid the Coleman-Mandula theorem, it is natural to explore symmetries generated by algebras beyond the Lie algebra.

In searching for symmetries that do not conflict with the Coleman-Mandula theorem, Haag, Lopsuzański and Sohnius extended the algebra to Lie superalgebra, so that the symmetry generators are closed not only for commutation relations but also for anti-commutation relations [7]. This showed that supersymmetry is the only possible symmetry as an extension of Poincaré symmetry, which is known as the Haag-Lopsuzański-Sohnius theorem. As its result, if we consider the Lie superalgebra as symmetry generators of the  $S$ -matrix, supersymmetry is allowed in QFT.

In 1978, further extensions of the Lie algebra were introduced by Rittenberg and Wyler [8, 9] (see also [10, 11]). This is called  $\mathbb{Z}_2^n$ -graded Lie (super)algebra, which is the extended algebra with the grading by abelian group  $\mathbb{Z}_2^n := \mathbb{Z}_2 \times \mathbb{Z}_2 \times \cdots$  (repeated  $n$  times). It reduces to an ordinary Lie algebra when  $n = 0$ , and to a Lie superalgebra when  $n = 1$ .

This type of algebra exhibits more complicated (anti-)commutation relations and possesses interesting properties in both physics and mathematics. In physics, symmetries generated by  $\mathbb{Z}_2^n$ -graded Lie superalgebras have been identified in various contexts, such as de Sitter SUGRA [12], nuclear quasi-spin [13], parastatistics [14], non-relativistic Dirac equation [15, 16], etc. Additionally, several  $\mathbb{Z}_2^n$ -graded extension of spacetime supersymmetries have also been proposed [17–22]. In particular,  $\mathbb{Z}_2^n$ -graded extension of super-Poincaré algebra introduced by Bruce [22] has some interesting properties and  $\mathbb{Z}_2^n$ -graded version of classical and quantum mechanics, sigma model, etc. have been discussed based on the extension [23–37].

This indicates that  $\mathbb{Z}_2^n$ -graded Lie (super)algebras are not uncommon in physics. Instead, they provide new insights into the analysis of physical systems based on symmetries, which is one of the key motivations for studying their structure and representations. The

structure and representation theories of  $\mathbb{Z}_2^n$ -graded Lie (super)algebras have been continuously explored since their introduction [38–59].

However, despite the extensive studies carried out so far, the application of  $\mathbb{Z}_2^n$ -graded Lie (super)algebras to  $(1+3)$ -dimensional relativistic quantum field theory has not been considered at all. To this end, one of the most important issues is the  $\mathbb{Z}_2^n$ -graded Lie (super)algebra can be symmetries of  $S$ -matrix. If this is the case, it provides an extension of the Haag-Łopuszański-Sohnius theorem. The purpose of this paper is to investigate the possible  $\mathbb{Z}_2^n$ -graded Lie (super)algebra symmetries of  $S$ -matrix in relativistic quantum field theory, following the approach of the Coleman-Mandula theorem and Haag-Łopuszański-Sohnius theorem. As the results, we demonstrate that a  $\mathbb{Z}_2^n$ -graded extension of the supersymmetric algebra can be a symmetry of  $S$ -matrix. Furthermore, it turns out that a  $\mathbb{Z}_2^n$ -graded Lie algebra appears as internal symmetries.

Our paper is organized as follows. In §2, we recall the definition of  $\mathbb{Z}_2^2$ -graded Lie superalgebra [8, 9], and briefly review Coleman-Mandula theorem and Haag-Łopuszański-Sohnius theorem [5, 7] in the case of massive particles. In §3, we consider the novel generators of symmetries of  $S$ -matrix in the case of massive particles. We first analyze the scenario with  $\mathbb{Z}_2^2$ -grading, following the frameworks of Coleman-Mandula theorem and Haag-Łopuszański-Sohnius theorem. Subsequently, using the same method as for the  $\mathbb{Z}_2^2$ -grading, we will consider the possible generators with  $\mathbb{Z}_2^n$ -gradings. Finally we will discuss the relation between the generators of internal symmetries and  $\mathbb{Z}_2^n$ -graded Lie algebra.

## 2 Preliminaries

### 2.1 $\mathbb{Z}_2^n$ -graded Lie superalgebra

We recall the definition of the  $\mathbb{Z}_2^n$ -graded LSA [8, 9].  $\mathbb{Z}_2^n$  refers to the  $n$ -fold direct product of the abelian group  $\mathbb{Z}_2 = \{0, 1\}$ , namely  $\mathbb{Z}_2^n = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \cdots$  (repeated  $n$  times).

**Definition 1.** Let  $\mathfrak{g}$  be a  $\mathbb{Z}_2^n$ -graded vector space (over  $\mathbb{R}$  or  $\mathbb{C}$ ), which is the direct sum of homogeneous vector subspaces labeled by an element of  $\mathbb{Z}_2^n$ :

$$\mathfrak{g} = \bigoplus_{\vec{a} \in \mathbb{Z}_2^n} \mathfrak{g}_{\vec{a}}. \quad (2.1)$$

An element of  $\mathfrak{g}_{\vec{a}}$  is said to have the  $\mathbb{Z}_2^n$ -grading  $\vec{a}$ .

**Definition 2.** We define the  $\mathbb{Z}_2^n$ -Lie bracket by

$$[[A, B]] = AB - (-1)^{\vec{a} \cdot \vec{b}} BA, \quad (2.2)$$

where  $A \in \mathfrak{g}_{\vec{a}}$ ,  $B \in \mathfrak{g}_{\vec{b}}$ , and  $\vec{a} \cdot \vec{b}$  is the standard scalar product of  $n$ -dimensional vectors. A  $\mathbb{Z}_2^n$ -graded vector space is called a  $\mathbb{Z}_2^n$ -graded Lie superalgebra if  $[[A, B]] \in \mathfrak{g}_{\vec{a}+\vec{b}}$  and the Jacobi identity is satisfied:

$$[[A, [[B, C]]]] = [[[A, B], C]] + (-1)^{\vec{a} \cdot \vec{b}} [[B, [[A, C]]]]. \quad (2.3)$$

The  $\mathbb{Z}_2^n$ -graded LSA can be regarded as an extended Lie algebra because the  $\mathbb{Z}_2^n$ -Lie bracket is a commutator (anti-commutator) for  $\vec{a} \cdot \vec{b}$  is even (odd). Actually when  $n = 0$ , a  $\mathbb{Z}_2^n$ -graded LSA is equivalent to an ordinary Lie algebras. Furthermore, when  $n = 1$ , it corresponds to the standard LSA, which consists of a bosonic sector (0-grading) and a fermionic sector (1-grading). It is known that for  $n \geq 2$ , a  $\mathbb{Z}_2^n$ -graded LSA also generates a symmetry.

## 2.2 Coleman-Mandula theorem

In this subsection, we briefly review the Coleman-Mandula theorem, which is known as the no-go theorem for symmetries in QFT [5, 6]. The theorem states that ‘‘The continuous symmetries of the  $S$ -matrix in QFT can only be generated by a Lie algebra consisting of the translation generator  $P_\mu$ , the homogeneous Lorentz transformation generator  $M_{\mu\nu}$ , and the internal symmetry generator  $B_l$ ’’. Namely they discussed possible symmetries of the  $S$ -matrix generated by  $\mathbb{Z}_2^n$ -graded LSA’s with  $n = 0$  and proved Poincaré symmetry and internal symmetry were only allowed in QFT. To consider the avoidance of this theorem, we look at the proof of this theorem [2, 6]. They used the following reasonable assumptions in their proof,

1. The  $S$ -matrix is defined in relativistic local field theory,
2. The finite number of particle types with masses lighter than an arbitrary mass  $M > 0$  is finite,
3. The amplitudes for elastic two-body scattering are analytic functions of the scattering angle at almost all energies and angles.

The symmetry generator here means a Hermitian operator on the Hilbert space which is commutative with the  $S$ -matrix and acts on multi particle states as well as single particle ones. Note that we work in the momentum space so that any matrix element of the group generator is a distribution in momentum space.

First, we consider all generators which commute with  $P_\mu$ . Any compact Lie algebra can be decomposed into a direct sum of a semi-simple part  $\mathfrak{g}^S$  and an Abelian part  $\mathfrak{g}^A$  (Levi-Decomposition). Since  $P_\mu$  commute with itself,  $P_\mu$  is in Abelian part:  $P_\mu \in \mathfrak{g}^A$ . Thus, we consider the generators  $B_\alpha \in \mathfrak{g}^S$ . We transform the commutation relation between  $P_\mu$  and  $B_\alpha$  by Lorentz transformation

$$[U(\Lambda)B_\alpha U^{-1}(\Lambda), \Lambda^\nu{}_\mu P_\nu] = 0 \quad (2.4)$$

where  $U(\Lambda)$  is the unitary operator on Hilbert space and represents the Lorentz transformation. Since  $\Lambda^\nu{}_\mu$  is regular,  $U(\Lambda)B_\alpha U^{-1}(\Lambda)$  must be a linear combination of  $B_\alpha$ :

$$U(\Lambda)B_\alpha U^{-1}(\Lambda) = D(\Lambda)^\beta{}_\alpha B_\beta, \quad (2.5)$$

where  $D(\Lambda)$  is the representation matrix of Lorentz group. Since the metric of semi-simple part  $\mathfrak{g}^S$  is positive definite,  $D(\Lambda)$  must be an identity matrix for  $B_\alpha$  to satisfy the same algebraic relation after Lorentz transformation. Noting that the Lorentz group is not

compact, the symmetry generated by  $B_\alpha \in \mathfrak{g}^S$  is independent of Lorentz symmetry. The  $B_\alpha$  is independent of the momentum and acts as an identity matrix with respect to the spin, which is the generator of internal symmetry  $B_l$ .

Next, we consider all symmetry generators which do not commute with  $P_\mu$ . All symmetry generators  $G_\alpha$  act on single particle states as follows

$$G_\alpha |n, p\rangle = \sum_{n'} \int d^4 p' (K_\alpha(p', p))_{n', n} |n', p'\rangle, \quad (2.6)$$

where  $n', n$  denote the spin and particle type. The kernel  $K_\alpha(p', p)$  will be zero if both  $p$  and  $p'$  are not on the mass hyperboloid. Furthermore, the kernel  $K(p', p)$  is zero for all  $p \neq p'$ :

$$K_\alpha(p', p) = K_\alpha(p) \delta^4(p - p'). \quad (2.7)$$

From the second assumption, since the mass of the particle takes only discrete values,  $G_\alpha$  commute with the mass square operator. Thus, the commutation relation of  $P_\mu$  and  $G_\alpha$  satisfies

$$[P^\mu, G_\alpha] = a_\alpha^{\mu\nu} P_\nu, \quad (2.8)$$

where  $a_\alpha^{\mu\nu}$  is anti-symmetric with respect to  $\mu, \nu$ . It follows that  $G_\alpha$ , which does not commute with  $P_\mu$ , is the Lorentz generator

$$[P_\mu, M_{\nu\rho}] = i(g_{\nu\mu} P_\rho - g_{\rho\mu} P_\nu). \quad (2.9)$$

Therefore, the symmetries of the  $S$ -matrix in QFT are only the Poincaré and internal symmetries.

### 2.3 Haag-Lopuszański-Sohnius theorem

In this subsection, we briefly review the Haag-Lopuszański-Sohnius theorem [7], which extends the Coleman-Mandula theorem. In (2.6),  $G_\alpha$  can be separated into two different types; bosonic type for same statistics  $n', n$ , or fermionic type for different statistics  $n', n$ . All bosonic type generators are fully analyzed in the Coleman-Mandula theorem. A complete basis of bosonic type generators are given by the  $P_\mu, M_{\mu\nu}$  and  $B^l$ . On the other hand, the fermionic type generator is not mentioned in the Coleman-Mandula theorem because of the assumption that the all generators consist of Lie algebra. They extended the Lie algebra to include anti-commutation relations in order to consider the fermionic type generators. This theorem says that, ‘‘Poincaré symmetry can only be extended to supersymmetry’’

Now, let us see how this supersymmetry avoids the Coleman-Mandula theorem. Note that  $G_\alpha$  in (2.9) can be bosonic or fermionic. However, the proof in the previous section concludes that the possible  $G_\alpha$  is only (2.9). Therefore, it is enough to consider fermionic generator  $Q$  commute with  $P_\mu$ . Let  $\mathfrak{c}$  denote the set of all generators commute with  $P_\mu$ . Since  $\mathfrak{c}$  is stable under the Lorentz transformations, it is a finite dimensional representation

space of Lorentz groups, and all generators in  $\mathfrak{c}$  can be decomposed into irreducible representation. The irreducible representation of the Lorentz group is specified by two spinors, which we write as  $(a, b)$ -rep. In general, because fermionic generators can have an odd number of spin indices, we denote the generator of  $(j, j')$ -rep and its conjugate  $(j', j)$ -rep characterizing two independent spins  $\alpha, \beta$  as  $Q_{\alpha_1 \dots \alpha_{2j} \dot{\beta}_1 \dots \dot{\beta}_{2j'}}$ ,  $\bar{Q}_{\beta_1 \dots \beta_{2j'} \dot{\alpha}_1 \dots \dot{\alpha}_{2j}}$ , where  $j, j'$  is an integer or half-integer, and  $2j$  undotted and  $2j'$  dotted indices are symmetric in each part. Now consider the anti-commutation relation between  $Q$  and  $\bar{Q}$ . Since it is produced from two fermionic generators, it must be bosonic and belong to  $(j + j', j + j')$ -rep. The bosonic generators are defined by the Coleman-Mandula theorem and are either  $P_\mu, M_{\mu\nu}, B_l$ . The irreducible representation of the Lorentz group of each generator is as follows

$$P_\mu \rightarrow \left(\frac{1}{2}, \frac{1}{2}\right)\text{-rep}, \quad M_{\mu\nu} \rightarrow (1, 0)\text{-rep} + (0, 1)\text{-rep}, \quad B_l \rightarrow (0, 0)\text{-rep}.$$

It follows that  $j + j' \leq \frac{1}{2}$ , which means  $Q \in \mathfrak{c}$  is only rank-1 spinor charge. Therefore, the anti-commutator of  $Q$  with  $\bar{Q}$  becomes  $P_\mu$ <sup>1</sup>

$$\{Q_\alpha, \bar{Q}_{\dot{\beta}}\} = \sigma_{\alpha\dot{\beta}}^\mu P_\mu. \quad (2.10)$$

$Q_\alpha, \bar{Q}_{\dot{\alpha}}$  are the supersymmetry generators. The other relations are obtained in the same way

$$[Q_\alpha, M_{\mu\nu}] = (\sigma_{\mu\nu})_\alpha^{\alpha_1} Q_{\alpha_1}, \quad \{Q_\alpha, Q_\beta\} = 0. \quad (2.11)$$

If we consider that there are several charges  $Q_{\alpha L}, \bar{Q}_{\dot{\alpha} M}$  ( $L, M = 1, 2, \dots$ ), anti-commutation relations of  $Q_{\alpha L}$  is non-zero:

$$\begin{aligned} \{Q_{\alpha K}, Q_{\beta L}\} &= \epsilon_{\alpha\beta} \sum_l a_{l, KL} B_l =: \epsilon_{\alpha\beta} Z_{KL}, \\ \{\bar{Q}_{\dot{\alpha} K}, \bar{Q}_{\dot{\beta} L}\} &= \varepsilon_{\dot{\alpha}\dot{\beta}} \sum_l \bar{a}_{l, KL} B_l =: \varepsilon_{\dot{\alpha}\dot{\beta}} \bar{Z}_{KL}, \end{aligned} \quad (2.12)$$

where  $a_{LM}^l$  is antisymmetric with respect to  $L, M$ . From the Jacobi identity, we can get

$$[Z_{KL}, Z_{MN}] = [Z_{KL}, \bar{Z}_{MN}] = 0. \quad (2.13)$$

Since  $Z_{KL}$  commutes with itself,  $Z_{KL}$  is in the Abelian part  $\mathfrak{g}^A$  of the Lie algebra and commutative with  $B_l \in \mathfrak{g}^S$ . Therefore,  $Z_{KL}$  commutes with all elements of the algebra. It is called the central charge. The LSA  $\langle P_\mu, M_{\mu\nu}, Z_{KL}, \bar{Z}_{KL}, Q_{\alpha L}, \bar{Q}_{\dot{\alpha} M} \rangle$  is called the super-Poincaré algebra, and the LSA  $\langle P_\mu, Z_{KL}, \bar{Z}_{KL}, Q_{\alpha L}, \bar{Q}_{\dot{\alpha} M} \rangle$  is called the supersymmetric algebra.

<sup>1</sup>Our notations are as follows. Let  $g_{\mu\nu} = \text{diag}(+, -, -, -)$  and  $(\sigma^\mu)_{\alpha\dot{\alpha}} = (\sigma^0, \sigma^1, \sigma^2, \sigma^3)_{\alpha\dot{\alpha}}$ , where

$$\sigma^0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Define the complex conjugate for any matrices as  $(M_{\alpha\beta})^* =: \bar{M}_{\dot{\alpha}\dot{\beta}}$  and the Hermitian conjugate as  $(M_{\alpha\beta})^\dagger =: \bar{M}_{\dot{\beta}\dot{\alpha}}$ . Then we obtain  $\{(\sigma^\mu)_{\alpha\dot{\beta}}\}^\dagger = (\bar{\sigma}^\mu)_{\dot{\beta}\alpha} = (\sigma^\mu)_{\beta\dot{\alpha}}$  and  $(\bar{\sigma}^\mu)^{\dot{\alpha}\alpha} := \varepsilon^{\dot{\alpha}\dot{\beta}} \varepsilon^{\alpha\beta} \sigma_{\beta\dot{\beta}}^\mu = (\sigma^0, -\sigma^i)$ , where  $\varepsilon_{\alpha\beta}$  is the antisymmetric tensor characterized by  $\varepsilon^{12} = \varepsilon_{21} = 1$  (the same for dotted indices).

### 3 Novel possible generators of the $S$ -matrix

#### 3.1 Possible $\mathbb{Z}_2^2$ -graded generators of symmetries

In this section, we search for novel possible generators of symmetries of  $S$ -matrix within the framework of the  $\mathbb{Z}_2^2$ -graded LSA. From (2.1)  $\mathbb{Z}_2^2$ -graded LSA  $\mathfrak{g}$  has four types of homogeneous vector subspaces:

$$\mathfrak{g} = \mathfrak{g}_{(0,0)} \oplus \mathfrak{g}_{(1,1)} \oplus \mathfrak{g}_{(0,1)} \oplus \mathfrak{g}_{(1,0)}, \quad (3.1)$$

and, from (2.2),  $\mathbb{Z}_2^2$ -Lie brackets between each subspace are given, in terms of (anti-)commutator as in Table 1.

	$\mathfrak{g}_{(0,0)}$	$\mathfrak{g}_{(1,1)}$	$\mathfrak{g}_{(0,1)}$	$\mathfrak{g}_{(1,0)}$
$\mathfrak{g}_{(0,0)}$	$[\, , \,]$	$[\, , \,]$	$[\, , \,]$	$[\, , \,]$
$\mathfrak{g}_{(1,1)}$	$[\, , \,]$	$[\, , \,]$	$\{ , \}$	$\{ , \}$
$\mathfrak{g}_{(0,1)}$	$[\, , \,]$	$\{ , \}$	$\{ , \}$	$[\, , \,]$
$\mathfrak{g}_{(1,0)}$	$[\, , \,]$	$\{ , \}$	$[\, , \,]$	$\{ , \}$

Table 1:  $\mathbb{Z}_2^2$ -Lie brackets between each subspace

Thus, the generator  $G_\alpha$  in (2.6) is categorized into two types of bosonic generators in  $\mathfrak{g}_{(0,0)}$  and  $\mathfrak{g}_{(1,1)}$ , and two types of fermionic generator in  $\mathfrak{g}_{(0,1)}$  and  $\mathfrak{g}_{(1,0)}$ . The  $(0,0)$ -graded generators are (ordinary) bosonic because the algebraic relations between them and all generators are given by the commutator. For the  $(0,1)$  and the  $(1,0)$ -graded generators, the relations among generators with the same grading are given by the anti-commutators, while those between generators with different gradings are given by commutators. They are called commuting fermionic generators. The  $(1,1)$ -graded generators are referred to as exotic bosonic ones because the relations among them are given by commutators, whereas their relations with fermionic generators are given by anti-commutators. Note that there are four types of symmetry generators of  $S$ -matrix. This means there are four types of particles in QFT within this framework.

We focus on the following three subalgebras:

$$\mathfrak{h}_1 = \mathfrak{g}_{00} \oplus \mathfrak{g}_{11}, \quad \mathfrak{h}_2 = \mathfrak{g}_{00} \oplus \mathfrak{g}_{01}, \quad \mathfrak{h}_3 = \mathfrak{g}_{00} \oplus \mathfrak{g}_{10}.$$

As indicated in Table 1,  $\mathfrak{h}_1$  is an ordinary Lie algebra, while  $\mathfrak{h}_2$  and  $\mathfrak{h}_3$  are nothing but LSA's. This shows that we can obtain generators by applying the Coleman-Mandula theorem to  $\mathfrak{h}_1$  and the Haag-Lopuszański-Sohnius theorem to  $\mathfrak{h}_2$  and  $\mathfrak{h}_3$ . Therefore, in  $\mathfrak{h}_1$ , there are only the translation generators  $P_\mu$ , Lorentz generator  $M_{\mu\nu}$  and two types of generators of internal symmetries:  $B_l^{00} \in \mathfrak{g}_{(0,0)}$ ,  $B_l^{11} \in \mathfrak{g}_{(1,1)}$ .  $P_\mu$  and  $M_{\mu\nu}$ , of course, commute with  $B_l^{00}$  and  $B_l^{11}$ . The algebraic relations between  $B$ 's are given by

$$\begin{aligned} [B_l^{00}, B_m^{00}] &= i \sum_k f_{lmk} B_k^{00}, & [B_l^{11}, B_m^{11}] &= i \sum_k g_{lmk} B_k^{00}, \\ [B_l^{00}, B_m^{11}] &= i \sum_k h_{lmk} B_k^{11}, \end{aligned} \quad (3.2)$$

where  $f_{lmk}$ ,  $g_{lmk}$  must be antisymmetric  $f_{lmk} = -f_{mlk}$ ,  $g_{lmk} = -g_{mlk}$ , but  $h_{lmk}$  has no conditions because  $B_l^{00}$  has the different grading from  $B_l^{11}$ .

On the other hand,  $\mathfrak{h}_2$  is identical to the LSA introduced in §2.3. Denoting the  $(0, 1)$ -graded supercharges by  $Q_{\alpha K}^{01}$ ,  $\bar{Q}_{\dot{\alpha} K}^{01} \in \mathfrak{g}_{(0,1)}$ , the algebraic relations between  $P_\mu$ ,  $B_l^{00}$  and  $Q^{01}$ 's are given by

$$\begin{aligned} \{Q_{\alpha K}^{01}, \bar{Q}_{\dot{\alpha} L}^{01}\} &= \sigma_{\alpha\dot{\alpha}}^\mu P_\mu \delta_{KL}, & [Q_{\alpha K}^{01}, B_l^{00}] &= \sum_L s_{l,KL} Q_{\alpha L}^{01}, \\ \{Q_{\alpha K}^{01}, Q_{\beta L}^{01}\} &= \varepsilon_{\alpha\beta} \sum_l a_{l,KL} B_l^{00}, & \{\bar{Q}_{\dot{\alpha} K}^{01}, \bar{Q}_{\dot{\beta} L}^{01}\} &= \varepsilon_{\dot{\alpha}\dot{\beta}} \sum_l \bar{a}_{l,KL} B_l^{00}, \end{aligned} \quad (3.3)$$

where  $a_{l,KL} = -a_{l,LK}$ . Introducing the generator  $Z_{KL}^{(1)} := \sum_l a_{l,KL} B_l^{00} \in \mathfrak{g}_{(0,0)}$ , it becomes the central charge in  $\mathfrak{h}_2$ , which means

$$\begin{aligned} \{Q_{\alpha K}^{01}, Q_{\beta L}^{01}\} &= \varepsilon_{\alpha\beta} Z_{KL}^{(1)}, & [Z_{KL}^{(1)}, \circ] &= 0, \\ \{\bar{Q}_{\dot{\alpha} K}^{01}, \bar{Q}_{\dot{\beta} L}^{01}\} &= \varepsilon_{\dot{\alpha}\dot{\beta}} \bar{Z}_{KL}^{(1)}, & [\bar{Z}_{KL}^{(1)}, \circ] &= 0. \end{aligned} \quad (3.4)$$

Similarly,  $\mathfrak{h}_3$  is also identical to the LSA introduced in §2.3. Denoting the  $(1, 0)$ -graded supercharges by  $Q_{\alpha K}^{10}$ ,  $\bar{Q}_{\dot{\alpha} K}^{10} \in \mathfrak{g}_{(1,0)}$  in  $\mathfrak{h}_3$ , the algebraic relations between  $P_\mu$  and  $Q^{10}$ 's are given by

$$\begin{aligned} \{Q_{\alpha K}^{10}, \bar{Q}_{\dot{\alpha} L}^{10}\} &= \sigma_{\alpha\dot{\alpha}}^\mu P_\mu \delta_{KL}, & [Q_{\alpha K}^{10}, B_l^{00}] &= \sum_L t_{l,KL} Q_{\alpha L}^{10}, \\ \{Q_{\alpha K}^{10}, Q_{\beta L}^{10}\} &= \varepsilon_{\alpha\beta} \sum_l b_{l,KL} B_l^{00}, & \{\bar{Q}_{\dot{\alpha} K}^{10}, \bar{Q}_{\dot{\beta} L}^{10}\} &= \varepsilon_{\dot{\alpha}\dot{\beta}} \sum_l \bar{b}_{l,KL} B_l^{00}, \end{aligned} \quad (3.5)$$

where  $b_{l,KL} = -b_{l,LK}$ . Introducing the generator  $Z_{KL}^{(2)} := \sum_l b_{l,KL} B_l^{00} \in \mathfrak{g}_{(0,0)}$ , it also becomes the central charge in  $\mathfrak{h}_3$ , which means

$$\begin{aligned} \{Q_{\alpha K}^{10}, Q_{\beta L}^{10}\} &= \varepsilon_{\alpha\beta} Z_{KL}^{(2)}, & [Z_{KL}^{(2)}, \circ] &= 0, \\ \{\bar{Q}_{\dot{\alpha} K}^{10}, \bar{Q}_{\dot{\beta} L}^{10}\} &= \varepsilon_{\dot{\alpha}\dot{\beta}} \bar{Z}_{KL}^{(2)}, & [\bar{Z}_{KL}^{(2)}, \circ] &= 0. \end{aligned} \quad (3.6)$$

We need to determine the other algebraic relations. Taking into account that  $Q_{\alpha L}^{01}$  and  $Q_{\alpha L}^{10}$  are a  $(\frac{1}{2}, 0)$ -rep,  $[Q_{\alpha K}^{01}, Q_{\beta L}^{10}] \in \mathfrak{g}_{(1,1)}$  will be a  $(1, 0)$ -rep or  $(0, 0)$ -rep, and  $[Q_{\alpha K}^{01}, \bar{Q}_{\dot{\alpha} L}^{10}] \in \mathfrak{g}_{(1,1)}$  will be a  $(\frac{1}{2}, \frac{1}{2})$ -rep. Moreover, because  $B_l^{11} \in \mathfrak{g}_{11}$  is  $(0, 0)$ -rep, we can determine the following relations:

$$\begin{aligned} [Q_{\alpha K}^{01}, Q_{\beta L}^{10}] &= \sum_l \varepsilon_{\alpha\beta} c_{l,KL} B_l^{11}, & [\bar{Q}_{\dot{\alpha} K}^{01}, \bar{Q}_{\dot{\beta} L}^{10}] &= -\sum_l \varepsilon_{\dot{\alpha}\dot{\beta}} \bar{c}_{l,KL} B_l^{11}, \\ \{Q_{\alpha K}^{01}, B_l^{11}\} &= \sum_L u_{l,KL} Q_{\alpha L}^{10}, & \{Q_{\alpha K}^{10}, B_l^{11}\} &= \sum_L v_{l,KL} Q_{\alpha L}^{01}, \\ [Q_{\alpha K}^{01}, \bar{Q}_{\dot{\alpha} L}^{10}] &= 0. \end{aligned} \quad (3.7)$$

We compute  $\mathbb{Z}_2^2$ -Jacobi identities (2.3) to consider the conditions for structure constants such that  $\mathfrak{g}$  becomes  $\mathbb{Z}_2^2$ -graded LSA. We obtain the following conditions from  $(A, B, C) = (Q, \bar{Q}, B)$ :

$$s_{l,LK} = \bar{s}_{l,KL}, \quad t_{l,LK} = \bar{t}_{l,KL}, \quad u_{l,LK} = \bar{v}_{l,KL}, \quad (3.8)$$



from  $(A, B, C) = (Q, Q, \bar{Q})$ :

$$\begin{aligned}\sum_l a_{l,KL} \bar{s}_{l,KL} &= 0, & \sum_l b_{l,KL} \bar{t}_{l,KL} &= 0, \\ \sum_l c_{l,KL} \bar{u}_{l,KL} &= 0, & \sum_l c_{l,KL} \bar{v}_{l,KL} &= 0,\end{aligned}\quad (3.9)$$

from  $(A, B, C) = (Q, Q, Q)$ :

$$\begin{aligned}\sum_l (a_{l,KL} s_{l,NM} + a_{l,NL} s_{l,KM}) &= 0, & \sum_l (b_{l,KL} t_{l,NM} + b_{l,NL} t_{l,KM}) &= 0, \\ \sum_l (c_{l,KL} u_{l,NM} + c_{l,NL} u_{l,KM}) &= 0, & \sum_l (c_{l,LK} v_{l,NM} + c_{l,LN} v_{l,KM}) &= 0, \\ \sum_l (a_{l,KL} t_{l,NM} + c_{l,LN} u_{l,KM}) &= 0, & \sum_l (b_{l,KL} s_{l,NM} - c_{l,NL} v_{l,KM}) &= 0,\end{aligned}\quad (3.10)$$

and from  $(A, B, C) = (Q, Q, B)$ :

$$\begin{aligned}\sum_M (s_{l,LM} a_{m,KM} - s_{l,KM} a_{m,LM}) &= i \sum_k f_{klm} a_{k,KL}, \\ \sum_M (u_{l,LM} c_{m,KM} - u_{l,KM} c_{m,LM}) &= i \sum_k h_{klm} a_{k,KL}, \\ \sum_M (t_{l,LM} b_{m,KM} - t_{l,KM} b_{m,LM}) &= i \sum_k f_{klm} b_{k,KL}, \\ \sum_M (v_{l,LM} c_{m,MK} - v_{l,KM} c_{m,ML}) &= -i \sum_k h_{klm} b_{k,KL}, \\ \sum_M (t_{l,LM} c_{m,KM} - s_{l,KM} c_{m,ML}) &= -i \sum_k g_{lkm} c_{k,KL}, \\ \sum_M (v_{l,LM} a_{m,KM} + u_{l,KM} b_{m,LM}) &= i \sum_k g_{klm} c_{k,KL}.\end{aligned}\quad (3.11)$$

From these conditions, introducing the generator  $Z_{KL}^{11} := \sum_l c_{l,ML} B_l^{11} \in \mathfrak{g}_{(1,1)}$ , it becomes the  $\mathbb{Z}_2^2$ -central charge, which means

$$\begin{aligned}[Q_{\alpha K}^{01}, Q_{\beta L}^{10}] &= \varepsilon_{\alpha\beta} Z_{KL}^{11}, & [[Z_{KL}^{11}, T]] &= 0, \\ [Q_{\dot{\alpha} K}^{01}, Q_{\dot{\beta} L}^{10}] &= \varepsilon_{\dot{\alpha}\dot{\beta}} \bar{Z}_{KL}^{11}, & [[\bar{Z}_{KL}^{11}, T]] &= 0,\end{aligned}\quad (3.12)$$

where  $T \in \{P_\mu, Z_{MN}^{(1)}, \bar{Z}_{MN}^{(1)}, Z_{MN}^{(2)}, \bar{Z}_{MN}^{(2)}, Z_{MN}^{11}, \bar{Z}_{MN}^{11}, Q_{\alpha K}^{01}, \bar{Q}_{\dot{\alpha} K}^{01}, Q_{\alpha K}^{10}, \bar{Q}_{\dot{\alpha} K}^{10}\}$ . From these observations, we can obtain the novel possible generators of symmetries of the  $S$ -matrix:

$$\begin{aligned}\mathfrak{g}_{(0,0)} &\ni P_\mu, M_{\mu\nu}, B_l^{00}, Z_{KL}^{(1)}, \bar{Z}_{KL}^{(1)}, Z_{KL}^{(2)}, \bar{Z}_{KL}^{(2)}, & \mathfrak{g}_{(1,1)} &\ni B_l^{11}, Z_{KL}^{11}, \bar{Z}_{KL}^{11}, \\ \mathfrak{g}_{(0,1)} &\ni Q_{\alpha K}^{01}, \bar{Q}_{\dot{\alpha} K}^{01}, & \mathfrak{g}_{(1,0)} &\ni Q_{\alpha K}^{10}, \bar{Q}_{\dot{\alpha} K}^{10}.\end{aligned}\quad (3.13)$$

Therefore the following theorem holds.

**Theorem 1.** *There exists the unique  $\mathbb{Z}_2^2$ -graded extension of supersymmetric algebra which is given by the  $\mathbb{Z}_2^2$ -graded LSA  $\langle P_\mu, Z_{KL}^{(1)}, \bar{Z}_{KL}^{(1)}, Z_{KL}^{(2)}, \bar{Z}_{KL}^{(2)}, Z_{KL}^{11}, \bar{Z}_{KL}^{11}, Q_{\alpha K}^{01}, \bar{Q}_{\dot{\alpha} K}^{01}, Q_{\alpha K}^{10}, \bar{Q}_{\dot{\alpha} K}^{10} \rangle$ . We call this  $\mathcal{N} = (K|L)$   $\mathbb{Z}_2^2$ -supersymmetric algebra, where  $K$  and  $L$  are the number of the  $(0, 1)$  and  $(1, 0)$ -graded supercharges.*

Specifically, if we set the number of each supercharges  $Q_{\alpha K}^{01}, Q_{\alpha L}^{10}$  to  $K = L = 1$ , it corresponds to the  $\mathcal{N} = 2$   $\mathbb{Z}_2^2$ -supersymmetric algebra, which was introduced by Bruce [22]. In the present case of  $\mathbb{Z}_2^2$ -grading, the internal symmetries are generated by ordinary Lie algebra  $B_l^{00}$  and  $B_l^{11}$ . This does not seem to be a big difference from the ordinary supersymmetry. However we will see that internal symmetries are generated by a graded Lie algebra when  $\mathbb{Z}_2^n$ -graded ( $n \geq 3$ ) extensions of supersymmetry are considered.

### 3.2 Possible $\mathbb{Z}_2^n$ -graded generators of symmetries

The discussion for the  $\mathbb{Z}_2^2$ -graded LSA in the previous section can also be applied to the  $\mathbb{Z}_2^n$ -graded LSA. We define two subsets of  $\mathbb{Z}_2^n$ :

$$\begin{aligned} (\mathbb{Z}_2^n)_{\text{ev.}} &:= \{\vec{b} \in \mathbb{Z}_2^n \mid \vec{b} \cdot \vec{b} = 0 \pmod{2}\} \\ (\mathbb{Z}_2^n)_{\text{od.}} &:= \{\vec{f} \in \mathbb{Z}_2^n \mid \vec{f} \cdot \vec{f} = 1 \pmod{2}\}, \end{aligned} \quad (3.14)$$

and the subalgebras of (2.1):

$$\mathfrak{h}_{\vec{b}} = \mathfrak{g}_{\vec{0}} \oplus \mathfrak{g}_{\vec{b}} \quad (\vec{b} \neq \vec{0}), \quad \mathfrak{h}_{\vec{f}} = \mathfrak{g}_{\vec{0}} \oplus \mathfrak{g}_{\vec{f}}, \quad (3.15)$$

where  $\vec{0}$  is the identity element of  $\mathbb{Z}_2^n$ . Then  $\mathfrak{h}_{\vec{b}}$  is a Lie algebra, while  $\mathfrak{h}_{\vec{f}}$  is a LSA. This shows that we can obtain generators by applying the Coleman-Mandula theorem to  $\mathfrak{h}_{\vec{b}}$  and the Haag-Lopuszański-Sohnius theorem to  $\mathfrak{h}_{\vec{f}}$ . It can be seen that the following generators of the  $S$ -matrix are allowed:

$$\mathfrak{g}_{\vec{0}} \ni P_\mu, M_{\mu\nu}, B_l^{\vec{0}}, \quad \mathfrak{g}_{\vec{b}} \ni B_l^{\vec{b}}, \quad \mathfrak{g}_{\vec{f}} \ni Q_{\alpha K}^{\vec{f}}, \bar{Q}_{\dot{\alpha} K}^{\vec{f}}, \quad (3.16)$$

and the algebraic relations become

$$\begin{aligned} \llbracket Q_{\alpha K}^{\vec{f}}, \bar{Q}_{\dot{\alpha} L}^{\vec{f}'} \rrbracket &= \sigma_{\alpha\dot{\alpha}}^\mu P_\mu \delta_{KL} \delta^{\vec{f}\vec{f}'}, & \llbracket Q_{\alpha K}^{\vec{f}}, Q_{\beta L}^{\vec{f}'} \rrbracket &= \varepsilon_{\alpha\beta} \sum_l a_{l, KL}^{(\vec{f}, \vec{f}')} B_l^{\vec{f}+\vec{f}'}, \\ \llbracket Q_{\alpha L}^{\vec{f}}, B_l^{\vec{b}} \rrbracket &= s_{l, LM}^{(\vec{f}, \vec{b})} Q_{\alpha M}^{\vec{b}+\vec{f}}, & \llbracket B_l^{\vec{b}}, B_m^{\vec{b}'} \rrbracket &= i \sum_k h_{lmk}^{(\vec{b}, \vec{b}')} B_k^{\vec{b}+\vec{b}'}, \\ \llbracket \bar{Q}_{\dot{\alpha} K}^{\vec{f}}, \bar{Q}_{\dot{\beta} L}^{\vec{f}'} \rrbracket &= -(-1)^{(\vec{f}, \vec{f}')} \varepsilon_{\dot{\alpha}\dot{\beta}} \sum_l \bar{a}_{l, KL}^{(\vec{f}, \vec{f}')} B_l^{\vec{f}+\vec{f}'}, \end{aligned} \quad (3.17)$$

where  $a$ ,  $s$ , and  $h$  are structure constants with the  $\vec{0}$ -grading, and  $a_{l, KL}^{(\vec{f}, \vec{f}')} = (-1)^{\vec{f} \cdot \vec{f}'} a_{l, LK}^{(\vec{f}', \vec{f})}$  and  $h_{lmk}^{(\vec{b}, \vec{b}')} = -(-1)^{\vec{b} \cdot \vec{b}'} h_{mlk}^{(\vec{b}', \vec{b})}$ .

In the same way as  $\mathbb{Z}_2^2$ -extension, we obtain the conditions for structure constants from the  $\mathbb{Z}_2^n$ -Jacobi identities (2.3). They are obtained from  $(A, B, C) = (Q, \bar{Q}, B)$ :

$$s_{l, LM}^{(\vec{f}, \vec{f}+\vec{f}')} = \bar{s}_{l, ML}^{(\vec{f}', \vec{f}+\vec{f}')}, \quad (3.18)$$

from  $(A, B, C) = (Q, Q, \bar{Q})$ :

$$\sum_l a_{l,LM}^{(\vec{f},\vec{f}')} \bar{s}_{l,NK}^{(\vec{f}'',\vec{f}+\vec{f}')} = 0, \quad (3.19)$$

from  $(A, B, C) = (Q, Q, Q)$ :

$$\sum_l a_{l,LM}^{(\vec{f},\vec{f}'')} s_{l,KN}^{(\vec{f},\vec{f}'+\vec{f}'')} - (-1)^{\vec{f},\vec{f}'} \sum_l a_{l,KM}^{(\vec{f},\vec{f}'')} s_{l,LN}^{(\vec{f}',\vec{f}+\vec{f}'')} = 0, \quad (3.20)$$

and from  $(A, B, C) = (Q, Q, B)$ :

$$s_{l,LM}^{(\vec{f}',\vec{b})} a_{m,KM}^{(\vec{f},\vec{f}'+\vec{b})} + (-1)^{\vec{f},\vec{f}'} s_{l,KM}^{(\vec{f},\vec{b})} a_{m,LM}^{(\vec{f}',\vec{f}+\vec{b})} = \sum_k i h_{klm}^{(\vec{f}+\vec{f}',\vec{b})} a_{k,KL}^{(\vec{f},\vec{f}')}. \quad (3.21)$$

Using these conditions and setting  $Z_{KL}^{\vec{f}+\vec{f}'} := \sum_l a_{l,KL}^{(\vec{f},\vec{f}')} B_l^{\vec{f}+\vec{f}'}$ , then it satisfies

$$\llbracket Z_{KL}^{\vec{f}+\vec{f}'}, T \rrbracket = \llbracket \bar{Z}_{KL}^{\vec{f}+\vec{f}'}, T \rrbracket = 0, \quad T \in \{P_\mu, Z_{KL}^{\vec{f}+\vec{f}'}, \bar{Z}_{KL}^{\vec{f}+\vec{f}'}, Q_{\alpha K}^{\vec{f}}, \bar{Q}_{\alpha K}^{\vec{f}}\}, \quad (3.22)$$

where  $Z_{KL}^{\vec{f}+\vec{f}'} = (-1)^{\vec{f},\vec{f}'} Z_{LK}^{\vec{f}+\vec{f}'}$ .

Therefore the following theorem holds.

**Theorem 2.** *There exists the unique  $\mathbb{Z}_2^n$ -graded extension of supersymmetric algebra which is given by the  $\mathbb{Z}_2^n$ -graded LSA  $\langle P_\mu, Z_{KL}^{\vec{f}+\vec{f}'}, \bar{Z}_{KL}^{\vec{f}+\vec{f}'}, Q_{\alpha K}^{\vec{f}}, \bar{Q}_{\alpha K}^{\vec{f}} \rangle$ .*

This algebra contains  $K_i$  supercharges  $Q_{\alpha K_i}^{\vec{f}}$  in each subspace  $\mathfrak{g}_{\vec{f}}$  and nontrivial centers in each subspace  $\mathfrak{g}_{\vec{b}}$ . We call this  $\mathcal{N} = (K_1|K_2|\dots|K_{2^{n-1}})$   $\mathbb{Z}_2^n$ -supersymmetric algebra. Similarly to the  $\mathbb{Z}_2^2$ -case, if we set the number of each  $Q_{\alpha,K}^{\vec{f}}$  to  $K_i = 1$ , it corresponds to the  $\mathcal{N} = 2^{n-1}$   $\mathbb{Z}_2^n$ -supersymmetric algebra, which was introduced by Bruce [22].

### 3.3 $\mathbb{Z}_2^n$ -Graded internal symmetries

In this section, we focus on the generators of internal symmetry with  $\mathbb{Z}_2^n$ -grading  $B_l^{\vec{b}}$ . The discussion in the previous section says there can exist  $B_l^{\vec{b}}$  in each subspace  $\mathfrak{g}_{\vec{b}}$  and they satisfy

$$\llbracket B_l^{\vec{b}}, B_m^{\vec{b}'} \rrbracket = i \sum_k h_{lmk}^{(\vec{b},\vec{b}')} B_k^{\vec{b}+\vec{b}'}. \quad (3.17)$$

This means the set of  $B_l^{\vec{b}}$  is a  $(\mathbb{Z}_2^n)_{\text{ev}}$ -graded LSA. Especially, if we set  $n = 2k + 1$ , we can prove that this algebra is isomorphic to the  $\mathbb{Z}_2^{2k}$ -graded Lie algebra, which is another type of  $\mathbb{Z}_2^n$ -graded extension of Lie algebra introduced by Rittenberg and Wyler [8, 9].

**Definition 3.** Let  $\mathfrak{g}_{LA}$  be a  $\mathbb{Z}_2^{2n}$ -graded vector space (2.1) and  $\vec{\alpha} = (\vec{\alpha}_1, \vec{\alpha}_2, \dots, \vec{\alpha}_n) \in \mathbb{Z}_2^{2n}$  where  $\vec{\alpha}_i = (\alpha_{2i-1}, \alpha_{2i}) \in \mathbb{Z}_2^2$ . Then  $\mathfrak{g}_{LA}$  is called  $\mathbb{Z}_2^n$ -graded Lie algebra if  $\mathfrak{g}_{LA}$  has the  $\mathbb{Z}_2^{2n}$ -Lie bracket defined by

$$[A, B] := AB - (-1)^{\sum_{i=1}^n |\vec{\alpha}_i \times \vec{\beta}_i|} BA, \quad A \in \mathfrak{g}_{\vec{\alpha}}, \quad B \in \mathfrak{g}_{\vec{\beta}}, \quad (3.23)$$

and  $[A, B]$  satisfies

$$\begin{aligned} [A, B] &\in \mathfrak{g}_{\vec{\alpha}+\vec{\beta}}, \\ [A, [B, C]] &= [[A, B], C] + (-1)^{\sum_{i=1}^n |\vec{\alpha}_i \times \vec{\beta}_i|} [B[A, C]] \end{aligned} \quad (3.24)$$

**Theorem 3.** *The  $(\mathbb{Z}_2^{2n+1})_{ev.}$ -graded LSA*

$$\mathfrak{g}_{ev.} := \bigoplus_{\vec{b} \in (\mathbb{Z}_2^{2n+1})_{ev.}} \mathfrak{g}_{\vec{b}} \quad (3.25)$$

is isomorphic to the  $\mathbb{Z}_2^{2n}$ -graded Lie algebra  $\mathfrak{g}_{LA}$ .

*Proof.* Let  $f$  be an injective map  $f : \vec{\alpha} \in \mathbb{Z}_2^{2n} \rightarrow \vec{a} \in (\mathbb{Z}_2^{2n+1})_{ev.}$ , defined by

$$\vec{a} = \left( \vec{\alpha}_1, \vec{\alpha}_2 + |\vec{\alpha}_1|(1, 1), \dots, \vec{\alpha}_i + \sum_{k=1}^{i-1} |\vec{\alpha}_k|(1, 1), \dots, \vec{\alpha}_n + \sum_{k=1}^{n-1} |\vec{\alpha}_k|(1, 1), \sum_{k=1}^n |\vec{\alpha}_k| \right)$$

where  $|\vec{a}_k|^2 := a_{2k-1}^2 + a_{2k}^2$ , and it is equal to  $|\vec{a}_k| = a_{2k-1} + a_{2k} \pmod{2}$ .

Since  $\mathbb{Z}_2^{2n}$ -graded Lie algebra  $\mathfrak{g}_{LA}$  and  $(\mathbb{Z}_2^{2n+1})_{ev.}$ -graded LSA  $\mathfrak{g}_{ev.}$  have the same number of the graded subspaces, the only difference of them is the Lie bracket, which comes from  $\sum_{i=1}^n |\vec{\alpha}_i \times \vec{\beta}_i|$  in (3.23) and  $\vec{a} \cdot \vec{b}$  in (2.2). However, taking into account that computations are performed in modulo 2, we have straightforwardly

$$\sum_{i=1}^n |\vec{\alpha}_i \times \vec{\beta}_i| = \vec{a} \cdot \vec{b}. \quad (3.26)$$

Therefore the map  $f$  is an isomorphism between  $\mathfrak{g}_{LA}$  and  $\mathfrak{g}_{ev.}$ .  $\square$

From this theorem, we can redefine  $\mathbb{Z}_2^{2n}$ -graded Lie algebra (Definition 3).

**Definition 4.** Let  $\mathfrak{g}$  be a  $\mathbb{Z}_2^n$ -graded LSA in Definition 2. We define  $\mathbb{Z}_2^{n-1}$ -graded Lie algebra as the subalgebra

$$\mathfrak{g}_{ev.} := \bigoplus_{\vec{b} \in (\mathbb{Z}_2^n)_{ev.}} \mathfrak{g}_{\vec{b}} \subset \mathfrak{g}. \quad (3.27)$$

This is a more general definition, which means we can define not only  $\mathbb{Z}_2^{2n}$ -graded Lie algebras, but also  $\mathbb{Z}_2^{2n+1}$ -graded Lie algebras. Finally, for the internal symmetry, the following theorem holds.

**Theorem 4.** *The  $\mathbb{Z}_2^n$ -graded Lie algebra is allowed as the generators of the internal symmetry of S-matrix.*

This is very interesting result because we can consider new types of gauge symmetries in relativistic QFT. Specifically,  $\mathbb{Z}_2^2$ -extension of  $\mathfrak{sl}_n$  has already been introduced in [9] and some representation of  $\mathbb{Z}_2^2$ - $\mathfrak{sl}_2$  was investigated in [60]. These will be helpful to construct  $\mathbb{Z}_2^2$ - $\mathfrak{sl}_n$  gauge field theory.

## 4 Concluding remarks

In this paper, we explored novel possible generators of symmetries of  $S$ -matrix in order to apply  $\mathbb{Z}_2^n$ -graded Lie superalgebra to relativistic quantum field theory. As a result, we found  $\mathcal{N} = (K_1|K_2|\cdots|K_{2^{n-1}})$   $\mathbb{Z}_2^n$ -graded supersymmetry is the only allowed extension of supersymmetric algebra as the  $\mathbb{Z}_2^n$ -graded extension. Specifically, if we set the number of each  $Q_{\alpha K}^{\bar{f}}$ 's to  $K_i = 1$ , they correspond to the  $\mathcal{N} = 2^{n-1}$   $\mathbb{Z}_2^n$ -supersymmetric algebra, which was introduced by Bruce [22]. However we cannot determine the number of supercharges purely from algebraic restrictions. This should be determined by the relationship between the degrees of freedom of particles within a model under consideration.

Nontrivial results of the present paper are summarized as follows.

1. It is possible to construct  $\mathbb{Z}_2^n$ -graded supersymmetric relativistic QFT.
2. Gauge symmetry is not necessary generated by a Lie algebra, but by a  $\mathbb{Z}_2^n$ -graded Lie algebra.

Thus an important future work is to construct a  $\mathbb{Z}_2^2$ - $\mathfrak{sl}_2$  gauge field theory.

Finally, in this paper, we have only considered the case of massive particles. According to [5, 7], conformal generators can exist in the case of massless particles. Therefore if we explore novel generators in the massless case, we could obtain more nontrivial  $\mathbb{Z}_2^n$ -graded conformal generators.

## Acknowledgment

This work was supported by JST SPRING, Grant Number JPMJSP2139 (RI and AN). RI and AN would like to express our sincere gratitude to Professor Naruhiko Aizawa and Professor Nobuhito Maru, at Osaka Metropolitan University, for their invaluable guidance and support throughout our research.

## References

- [1] F. J. Dyson, *Symmetry Groups in Nuclear and particle Physics*, W. A. Benjamin, New York (1966),
- [2] S. Weinberg, *The Quantum Theory of Fields: Volume III: Supersymmetry*, Cambridge University Press (2000),
- [3] M. Gell-Mann, *A schematic model of baryons and mesons*, Phys. Lett. **8**, 3 (1964),
- [4] S. Coleman, *Trouble with Relativistic  $SU(6)$* , Phys. Rev. **138**, B1262 (1965),
- [5] S. Coleman and J. Mandula, *All Possible Symmetry of  $S$ -matrix*, Phys. Rev. **159**, 1251 (1967),
- [6] J. Mandula, *Coleman-Mandula theorem*, Scholapedia, 10(2):7476 (2015),

- [7] R. Haag, J. T. Łopuszański, M. Sohnius, *All possible generators of supersymmetry of the S-matrix*, Nucl. Phys. B **88**, 257 (1975),
- [8] V. Rittenberg and D. Wyler, *Generalized superalgebras*, Nucl. Phys. B **139**, 189 (1978).
- [9] V. Rittenberg and D. Wyler, *Sequences of  $\mathbb{Z}_2 \otimes \mathbb{Z}_2$  graded Lie algebras and superalgebras*, J. Math. Phys. **19**, 2193 (1978).
- [10] R. Ree, *Generalized Lie elements*, Canad. J. Math. **12**, 493 (1960).
- [11] M. Scheunert, *Generalized Lie algebras*, J. Math. Phys. **20**, 712 (1979).
- [12] M. A. Vasiliev, *de Sitter supergravity with positive cosmological constant and generalised Lie superalgebras*, Class. Quantum Gravity **2**, 645 (1985).
- [13] P. D. Jarvis, M. Yang and B. G. Wybourne, *Generalized quasispin for supergroups*, J. Math. Phys. **28**, 1192 (1987).
- [14] V. N. Tolstoy, *Once more on parastatistics*, Phys. Part. Nucl. Lett. **11**, 933 (2014).
- [15] N. Aizawa, Z. Kuznetsova, H. Tanaka, F. Toppan,  *$\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded Lie Symmetries of the Lévy-Leblond Equations*, Prog. Theor. Exp. Phys. **2016**, 123A01 (2016).
- [16] N. Aizawa, Z. Kuznetsova, H. Tanaka, F. Toppan, *Generalized supersymmetry and Lévy-Leblond equation*, in “Physical and Mathematical Aspects of Symmetries,” J.-P. Gazeau, S. Faci, T. Micklitz, R. Scherer, F. Toppan (editors), Springer (2017) p.79.
- [17] A. A. Zheltukhin, *Para-Grassmann extension of the Neveu-Schwartz-Ramond algebra*, Theor. Math. Phys. **71**, 491 (1987) (Teor. Mat. Fiz. **71** 218 (1987)).
- [18] B. Le Roy,  *$\mathbb{Z}_n^3$ -Graded colored supersymmetry*, Czech. J. Phys. **47**, 47 (1997).
- [19] L. A. Wills-Toro,  *$(I, q)$ -graded Lie algebraic extensions of the Poincaré algebra, constraints on  $I$  and  $q$* , J. Math. Phys. **36**, 2085 (1995).
- [20] L. A. Wills-Toro, *Trefoil symmetries I. Clover extensions beyond Coleman-Mandula theorem*, J. Math. Phys. **42**, 3915 (2001).
- [21] V. N. Tolstoy, *Super-de Sitter and Alternative Super-Poincaré Symmetries*, In: Dobrev V. (eds) Lie Theory and Its Applications in Physics. Springer Proceedings in Mathematics & Statistics, vol. 111, Springer, Tokyo, 2014.
- [22] A. J. Bruce, *On a  $\mathbb{Z}_2^n$ -graded version of supersymmetry*, Symmetry **11**, 116 (2019).
- [23] A. J. Bruce and S. Duplij, *Double-graded supersymmetric quantum mechanics*, J. Math. Phys. **61**, 063503 (2020).
- [24] N. Aizawa, K. Amakawa, S. Doi,  *$\mathcal{N}$ -Extension of double-graded supersymmetric and superconformal quantum mechanics*, J. Phys. A: Math. Theor. **53**, 065205 (2020).

- [25] N. Aizawa, K. Amakawa, S. Doi,  $\mathbb{Z}_2^3$ -Graded extensions of supersymmetric quantum mechanics via Clifford algebras, *J. Math. Phys.* **61**, 052105 (2020).
- [26] N. Aizawa, Z. Kuznetsova and F. Toppan,  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded mechanics: the classical theory, *Eur. Phys. J. C* **80**, 668 (2020).
- [27] N. Aizawa, Z. Kuznetsova and F. Toppan,  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded mechanics: the quantization, *Nucl. Phys. B* **967**, 115426 (2021).
- [28] S. Doi and N. Aizawa,  $\mathbb{Z}_2^3$ -Graded extensions of Lie superalgebras and superconformal quantum mechanics, *SIGMA* **17** 071, (2021).
- [29] S. Doi and N. Aizawa, Comments of  $\mathbb{Z}_2^2$ -supersymmetry in superfield formalism, *Nucl. Phys.* **B974**, 115641 (2022).
- [30] N. Aizawa and S. Doi, Irreducible representations of  $\mathbb{Z}_2^2$ -graded  $\mathcal{N} = 2$  supersymmetry algebra and  $\mathbb{Z}_2^2$ -graded supermechanics, *J. Math. Phys.* **63**, 091704 (2022).
- [31] A. J. Bruce,  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded supersymmetry: 2-d sigma models, *J. Phys. A: Math. Theor.* **53**, 455201 (2020).
- [32] A. J. Bruce, Is the  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded sine-Gordon equation integrable ?, *Nucl. Phys. B* **971**, 115514 (2021).
- [33] N. Aizawa, R. Ito, Z. Kuznetsova, F. Toppan, New aspects of the  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded 1D superspace: induced strings and 2D relativistic models, *Nucl. Phys. B* **991**, 116202 (2023).
- [34] F. Toppan,  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded parastatics in multiparticle quantum Hamiltonians, *J. Phys. A: Math. Theor.* **54**, 115203 (2021).
- [35] F. Toppan, Inequivalent quantizations from gradings and  $\mathbb{Z}_2 \times \mathbb{Z}_2$  parabosons, *J. Phys. A, Math. Theor.* **54**, 355202 (2021).
- [36] N. Aizawa, R. Ito and T. Tanaka,  $\mathbb{Z}_2^2$ -graded supersymmetry via superfield on minimal  $\mathbb{Z}_2^2$ -superspace, *J. Phys. A: Math. Theor.* **57** 435201 (2024).
- [37] N. Aizawa, R. Ito and T. Tanaka,  $\mathcal{N} = 2$  Double-graded supersymmetric quantum mechanics via dimensional reduction, *AIMS Math.* **9** 10494 (2024).
- [38] M. Scheunert, Casimir elements of  $\epsilon$ -Lie algebras, *J. Math. Phys.* **24**, 2671 (1983).
- [39] M. Scheunert and R. B. Zhang, Cohomology of Lie superalgebras and their generalizations, *J. Math. Phys.* **39**, 5024 (1998).
- [40] S. D. Silvestrov, On the classification of 3-dimensional coloured Lie algebras, Banach Center Publications, **40**, 159 (1997).
- [41] X.-W. Chen, S. D. Silvestrov and F. Van Oystaeyen, Representations and cocycle twists of color Lie algebras, *Algebr. Represent. Theor.* **9**, 633 (2006).

- [42] G. Sigurdsson and S. D. Silvestrov, *Bosonic realizations of the colour Heisenberg Lie algebras*, Nonlinear Math. Phys. **13**, supplement 110 (2006).
- [43] D. Piontkovski and S. D. Silvestrov, *Cohomology of 3-dimensional color Lie algebras*, J. Algebra **316**, 499 (2007).
- [44] R. Campoamor-Stursberg and M. Rausch de Traubenberg, *Color Lie algebras and Lie algebras of order  $F$* , J. Gen. Lie Theory Appl. **3**, 113 (2009).
- [45] M. Mohammadi and H. Salmasian, *The Gelfand-Naimark-Segal construction for unitary representations of  $\mathbb{Z}_2^n$ -graded Lie supergroups*, Banach Cent. Publ. **113**, 263 (2017).
- [46] N. Aizawa and J. Segar,  *$\mathbb{Z}_2 \times \mathbb{Z}_2$  generalizations of  $\mathcal{N} = 2$  super Schrödinger algebras and their representations*, J. Math. Phys. **58**, 113501 (2017).
- [47] N. Aizawa, P. S. Isaac and J. Segar,  *$\mathbb{Z}_2 \times \mathbb{Z}_2$  generalizations of  $\mathcal{N} = 1$  superconformal Galilei algebras and their representations*, J. Math. Phys. **60**, 023507 (2019).
- [48] N. I. Stoilova, J. Van der Jeugt, *The  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded Lie superalgebra  $\mathfrak{psl}(2m+1|2n)$  and new parastatistics representations*, J. Phys. A:Math. Theor. **51**, 135201 (2018).
- [49] P. Meyer, *The Kostant invariant and special  $\epsilon$ -orthogonal representations for  $\epsilon$ -quadratic colour Lie algebras*, J. Algebra **572**, 337 (2021).
- [50] P. S. Isaac, N. I. Stoilova, J. van der Jeugt, *The  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded general linear Lie superalgebra*, J. Math. Phys. **61**, 011702 (2020).
- [51] N. Aizawa, P. S. Isaac and J. Segar,  *$\mathbb{Z}_2 \times \mathbb{Z}_2$  generalizations of infinite dimensional Lie superalgebra of conformal type with complete classification of central extensions*, Rep. Math. Phys. **85**, 351 (2020).
- [52] K. Amakawa and N. Aizawa, *A classification of lowest weight irreducible modules over  $\mathbb{Z}_2^2$ -graded extension of  $osp(1|2)$* , J. Math. Phys. **62**, 043502 (2021).
- [53] Z. Kuznetsova, F. Toppan, *Classification of minimal  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded Lie (super)algebras and some applications*, J. Math. Phys. **62**, 063512 (2021).
- [54] P. Meyer, *Cubic Dirac operators and the strange Freudenthal-de Vries formula for colour Lie algebras*, Transform. Groups **27**, 1307 (2022).
- [55] N. I. Stoilova, J. van der Jeugt, *The  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded Lie superalgebras  $\mathfrak{psl}(2n+1|2n)$  and  $\mathfrak{psl}(\infty|\infty)$ , and parastatistics Fock spaces*, J. Phys. A:Math. Theor. **55**, 045201 (2022).
- [56] N. I. Stoilova, J. van der Jeugt, *On classical  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded Lie algebras*, J. Math. Phys. **64**, 061702 (2023).
- [57] R. Lu and Y. Tan, *Construction of color Lie algebras from homomorphisms of modules of Lie algebras*, J. Algebra **620**, 1 (2023).



- [58] F. S. Alshammari, Md F. Hoque, J. Segar, *The  $osp(1|2)$   $\mathbb{Z}_2 \times \mathbb{Z}_2$  graded algebra and its irreducible representations*, arXiv:2306.12381 [math-ph].
- [59] N. Aizawa, J. Segar, *Affine extensions of  $\mathbb{Z}_2^2$ -graded  $osp(1|2)$  and Virasoro algebra*, arXiv:2409.07938 [math-ph]
- [60] N. Aizawa, R.Ito, Z. Kuznetsova, T. Tanaka, F, Toppan, *Integrable  $\mathbb{Z}_2^2$ -graded Extensions of the Liouville and Sinh-Gordon Theories*, arXiv:2406.13503 [math-ph].