SECOND QUANTIZATION FOR CLASSICAL NONLINEAR DYNAMICS

DIMITRIOS GIANNAKIS, MOHAMMAD JAVAD LATIFI JEBELLI, MICHAEL MONTGOMERY, PHILIPP PFEFFER, JÖRG SCHUMACHER, AND JOANNA SLAWINSKA

ABSTRACT. Using techniques from many-body quantum theory, we propose a framework for representing the evolution of observables of measure-preserving ergodic flows through infinite-dimensional rotation systems on tori. This approach is based on a class of weighted Fock spaces $F_w(\mathcal{H}_\tau)$ generated by a 1-parameter family of reproducing kernel Hilbert spaces \mathcal{H}_{τ} , and endowed with commutative Banach algebra structure under the symmetric tensor product using a subconvolutive weight w. We describe the construction of the spaces $F_w(\mathcal{H}_\tau)$ and show that their Banach algebra spectra, $\sigma(F_w(\mathcal{H}_\tau))$, decompose into a family of tori of potentially infinite dimension. Spectrally consistent unitary approximations U_{τ}^{t} of the Koopman operator acting on \mathcal{H}_{τ} are then lifted to rotation systems on these tori akin to the topological models of ergodic systems with pure point spectra in the Halmos-von Neumann theorem. Our scheme also employs a procedure for representing observables of the original system by polynomial functions on finite-dimensional tori in $\sigma(F_w(\mathcal{H}_\tau))$ of arbitrarily large degree, with coefficients determined from pointwise products of eigenfunctions of U_{τ}^{t} . This leads to models for the Koopman evolution of observables on L^2 built from tensor products of finite collections of approximate Koopman eigenfunctions. Numerically, the scheme is amenable to consistent data-driven implementation using kernel methods. We illustrate it with applications to Stepanoff flows on the 2-torus and the Lorenz 63 system. Connections with quantum computing are also discussed.

1. Introduction

The pursuit of connections between classical and quantum dynamics dates back to the origins of quantum mechanics and modern ergodic theory in the first few decades of the 20th century. Seminal work of Koopman and von Neumann [50,51] simultaneously established the foundations of operator-theoretic representations of classical dynamics by means of composition operators (now widely known as Koopman operators) and put forward a description of the Liouville evolution of classical probability densities through quantum mechanical wavefunctions (so-called Koopman–von Neumann waves [59]). The Koopman–von Neumann approach has found applications in areas including Hamiltonian mechanics [74,75], stochastic dynamics [24], and hybrid classical–quantum dynamics [12].

A related problem that has received significant attention in recent years is simulation of classical dynamical systems by quantum systems [5,6,27,37,44,46,47,54,56]. Example applications include viscous fluid flow [9,33], thermal convection [64], transport [60], wave propagation [19], climate dynamics [70], plasma dynamics [25,28], and energy science [45]. A primary motivation underpinning these efforts is the advent of quantum computing with its premise to deliver transformational advances in computing capabilities.

Yet another line of research has been on quantum-inspired techniques, i.e., methods based on the mathematical framework of quantum theory that are otherwise implemented classically. These methods leverage properties of spaces of operators associated with quantum systems that aid the design of structure-preserving approximation schemes (e.g., positivity-preserving projections) in applications such as data assimilation [31,34] and dynamical closure [32].

In broad terms, techniques for quantum simulation of classical dynamics are based on mappings of classical states and observables into states and observables of a quantum system, together with a corresponding mapping of the classical dynamical evolution maps into an evolution of quantum

states. Here, a major challenge stems from the fact that classical systems of interest are typically nonlinear, whereas quantum dynamics proceeds by unitary (linear) transformations of the quantum state. A popular strategy to overcome this challenge is to represent polynomial nonlinearities in the equations of motion as multilinear maps that can be treated linearly in a tensor product Hilbert space using appropriate time-stepping schemes. Variants of this strategy have been employed in schemes for solving nonlinear ordinary differential equations (ODEs) [54,56] and certain classes of partial differential equations (PDEs) after space discretization [33,55].

Rather than attempting to directly simulate the (nonlinear) evolution the classical state, an alternative approach to state-space-centric schemes employs operator techniques from ergodic theory [3,26] and the Koopman-von Neumann approach to simulate the evolution of observables and measures under the dynamics [37,44]. Indeed, for a measure-preserving, invertible transformation $\Phi \colon X \to X$ of a probability space (X, Σ, μ) , the Koopman and transfer (Perron-Frobenius) operators on $L^2(\mu)$, $Uf = f \circ \Phi$ and $Pf = f \circ \Phi^{-1}$, respectively, are unitary (and form a dual pair, $U^* = P$), and thus are natural candidates for approximation by quantum algorithms. By the Stone theorem on 1-parameter, strongly continuous unitary groups [68], the Koopman group $\{U^t \colon L^2(\mu) \to L^2(\mu)\}_{t \in \mathbb{R}}$ induced by a measure-preserving flow $\Phi^t \colon X \to X$, $t \in \mathbb{R}$, is completely characterized by its generator—a skew-adjoint operator $V \colon D(V) \to L^2(\mu)$ defined on a dense domain $D(V) \subseteq L^2(\mu)$ as the $L^2(\mu)$ -norm limit

$$Vf = \lim_{t \to 0} (U^t f - f)/t, \tag{1}$$

and giving the time-t Koopman operator by exponentiation, $U^t = e^{tV}$. In this continuous-time setting, the self-adjoint operator V/i plays an analogous role to a quantum mechanical Hamiltonian, and is therefore a natural object to study when making connections between classical and quantum systems.

1.1. Our contributions. Focusing on measure-preserving, ergodic flows in continuous time, the primary challenge we seek to address is to build a quantum mechanical representation of classical dynamics that consistently approximates the Koopman evolution generated by V from (1) for systems with non-trivial continuous spectra. Indeed, a hallmark result in ergodic theory is that a measure-preserving system is weak-mixing if and only if the unitary Koopman group on $L^2(\mu)$ has a single, simple eigenvalue at 1 (with a constant corresponding eigenfunction), and no other elements in the point spectrum [42]. In continuous time, this means that, aside from a simple eigenvalue at 0, the spectral measure of the generator is continuous. Thinking of weak-mixing as a signature of high dynamical complexity ("chaos") in a measure-theoretic sense, consistent approximation of the continuous spectrum is thus relevant to quantum simulation of a broad range of nonlinear systems encountered in applications.

Our approach to address this problem consists of the following principal elements:

- Spectral regularization techniques for Koopman operators [22, 39, 40] that approximate the generator V by a 1-parameter family of skew-adjoint, diagonalizable operators W_{τ} : $D(W_{\tau}) \to \mathcal{H}_{\tau}$, $\tau > 0$, each acting on a reproducing kernel Hilbert space (RKHS) \mathcal{H}_{τ} of continuous functions on state space X. Here, τ is a regularization parameter and W_{τ} converges to V as $\tau \to 0^+$ in an appropriate spectral sense.
- Constructions of so-called reproducing kernel Hilbert algebras (RKHAs) [20,21,36], which are RKHSs with coalgebra structure with respect to the Hilbert space tensor product and Banach algebra structure with respect to the pointwise product of functions.
- A class of weighted symmetric Fock spaces (described, to our knowledge, for the first time in this paper) with Banach algebra structure with respect to the tensor product. For our purposes, a key property of these spaces is that they are isomorphic to RKHAs of continuous functions on infinite-dimensional tori embedded in their spectra.

• Previously developed quantum algorithms for approximating the Koopman evolution of observables of rotation systems on tori [37].

Building on these tools, we propose a provably consistent framework for quantum simulation of observables of continuous-time measure-preserving ergodic flows with arbitrary (i.e., pure-point, continuous, and mixed) spectral characteristics. The mathematical framework of many-body quantum theory [53], also known as second quantization, plays a central role in our constructions, particularly through the use of Fock space theory.

In essence, our scheme approximates the unitary Koopman evolution on $L^2(\mu)$ by a family of rotation systems on tori, $\mathbb{T}_{\tau,x}$ of potentially infinite dimension, parameterized by the regularization parameter τ and the initial condition $x \in X$. The tori $\mathbb{T}_{\tau,x}$ are realized as weak-* compact subsets of the spectrum $\sigma(F_w(\mathcal{H}_\tau))$ of an abelian Banach algebra $F_w(\mathcal{H}_\tau)$ built as a weighted Fock space generated by the RKHS \mathcal{H}_τ that we employ for spectral approximation of the Koopman operator. Furthermore, the state space dynamics $\Phi^t \colon X \to X$ embeds naturally in $\sigma(F_w(\mathcal{H}_\tau))$ by means of a feature map. As a result, we can interpret the torus rotation system obtained by regularization at the Koopman/transfer operator level as an approximate topological model of the unperturbed state space dynamics. This construction should be of independent interest in broader contexts than quantum mechanical representation of classical dynamics.

Each torus $\mathbb{T}_{\tau,x}$ can be approximated by a sequence of d-dimensional tori $\mathbb{T}_{\tau,d,x} \subset \sigma(F_w(\mathcal{H}_\tau))$, with corresponding rotation systems generated by d basic frequencies. We put forward a procedure for representing continuous classical observables as polynomials of arbitrarily large degree m in the Fourier basis on $\mathbb{T}_{\tau,d,x}$. Increasing m allows to capture spectral information from large-dimensional spaces generated by tensor products of eigenfunctions of W_τ . This enhances prediction skill and should also aid the theoretical efficiency of implementations of our scheme on quantum computing platforms.

In this paper, we describe the mathematical formulation of our second-quantization approach. Moreover, we demonstrate this framework by means of numerical experiments involving measure-preserving ergodic dynamical systems with aperiodic behavior: a Stepanoff flow on the 2-torus [62] and the Lorenz 63 (L63) system [57] on \mathbb{R}^3 . These experiments are performed on classical hardware using classical numerical methods and demonstrate improved prediction skill over conventional Koopman operator approximation methods utilizing the same number of eigenfunctions. A gate-based implementation of our algorithms on simulated on actual quantum hardware is beyond the scope of this paper, and will be addressed elsewhere.

1.2. Plan of the paper. In section 2, we introduce the class of dynamical systems under study and establish basic notational conventions. This is followed by a survey of relevant previous work in section 3. In section 4, we describe our second quantization approach, relegating certain constructions and proofs to sections 5 and 6. Specifically, section 5 describes the construction and properties of the symmetric weighted Fock space $F_w(\mathcal{H}_\tau)$, and section 6 our schemes for representing observables of the original dynamical system as vectors in $F_w(\mathcal{H}_\tau)$, including the polynomial functions on the tori $\mathbb{T}_{\tau,d,x}$. Section 7 contains an overview of the data-driven formulation of our scheme. In section 8, we present the results from our numerical experiments testing these methods. A discussion and our primary conclusions are included in section 9. Appendices A and B collect auxiliary material on Markov semigroups employed in the construction of \mathcal{H}_τ and spectral approximation of Koopman operators, respectively.

2. Preliminaries and notation

2.1. **Dynamical system.** We consider a continuous flow $\Phi^t : X \to X$, $t \in \mathbb{R}$, on a metrizable space X with an ergodic invariant Borel probability measure μ of compact support. For $p \in [1, \infty]$, we let $U^t : L^p(\mu) \to L^p(\mu)$ denote the associated time-t Koopman operator that acts isometrically by composition with the flow, $U^t f = f \circ \Phi^t$. As noted in section 1, the Koopman group $\{U^t\}_{t \in \mathbb{R}}$ on

 $H := L^2(\mu)$ is strongly continuous and unitary, and thus has a skew-adjoint generator $V : D(V) \to H$ defined by (1). In addition, U^t acts as a *-isomorphism of the abelian von Neumann algebra $L^{\infty}(\mu)$ with respect to pointwise multiplication and complex conjugation,

$$U^t(fg) = (U^tf)(U^tg), \quad U^t(f^*) = (U^tf)^*, \quad \forall f,g \in L^\infty(\mu).$$

For every $p \in (1, \infty]$, U^t has a predual, $P^t : L^q(\mu) \to L^q(\mu)$ for $\frac{1}{p} + \frac{1}{q} = 1$, defined by $P^t g = g \circ \Phi^{-t}$ and satisfying

$$\langle U^t f, g \rangle = \langle f, P^t g \rangle, \quad \forall f \in L^p(\mu), \quad \forall g \in L^q(\mu),$$

where $\langle \cdot, \cdot \rangle$ denotes the natural pairing between $L^p(\mu)$ and $L^q(\mu)$. The operators P^t are known as transfer, or Perron–Frobenius, operators, and govern the evolution of densities with respect to the invariant measure μ under the dynamics. Specifically, given a finite Borel measure ν with density $g = \frac{d\nu}{d\mu} \in L^q(\mu)$, then P^tg is the density $\frac{d\nu_t}{d\mu}$ of the pushforward measure $\nu_t = \Phi^t_*\nu$ under the dynamics. The transfer operator $P^t \colon L^1(\mu) \to L^1(\mu)$ is a Markov operator with invariant measure μ ; i.e., (i) $P^t 1_X = 1_X$ for the constant function 1_X equal everywhere to 1; (ii) $P^t g \ge 0$ for $g \ge 0$; and (iii) $\int_X P^t g \, d\mu = \int_X g \, d\mu$ for all $g \in L^1(\mu)$. For further details on Koopman and transfer operators we refer the reader to [3, 26, 42].

Throughout the paper, we will assume that the support of μ is contained in a C^1 compact manifold $M \subseteq X$ that is forward-invariant under the flow, $\Phi^t(M) \subseteq M$ for all $t \ge 0$. We also assume that the restriction of Φ^t on M is generated by a continuous vector field $\vec{V}: M \to TM$. This means that

$$\iota \circ \vec{V} \cdot \nabla f = V \circ \iota f, \quad \forall f \in C^1(M),$$
 (2)

where $\iota \colon C(M) \to H$ is the map from continuous functions on M to their corresponding equivalence classes in H.

2.2. **Notation.** All vector spaces in this work will be over the complex numbers and all Hilbert spaces will be separable. For a normed space \mathbb{V} , $(\mathbb{V})_R \subset \mathbb{V}$ will denote the closed ball of radius R centered at the origin. Moreover, we will denote the space of bounded linear maps between \mathbb{V} and a Banach space \mathbb{E} as $B(\mathbb{V}, \mathbb{E})$, and $\|A\|$ will be the operator norm of $A \in B(\mathbb{V}, \mathbb{E})$. We will use the abbreviation $B(\mathbb{E}) \equiv B(\mathbb{E}, \mathbb{E})$. Given an operator $A \colon D(A) \to \mathbb{E}$ defined on a subspace $D(A) \subseteq \mathbb{E}$ of a Banach space \mathbb{E} , $\rho(A)$, $\sigma(A)$, and $\sigma_p(A)$ will denote the resolvent set, spectrum, and point spectrum of A, respectively. Moreover, for $z \in \rho(A)$, $R(z,A) = (z-A)^{-1} \in B(\mathbb{E})$ will denote the corresponding resolvent operator. An inner product on a vector space \mathbb{V} will be denoted as $\langle \cdot, \cdot \rangle_{\mathbb{V}}$ and will be taken to be conjugate-linear in the first argument. If \mathbb{V} is a closed subspace of a Hilbert space \mathbb{H} , $\operatorname{proj}_{\mathbb{V}} \in B(\mathbb{H})$ will be the orthogonal projection with range \mathbb{V} .

If \mathcal{H} is an RKHS of complex-valued functions on a set \mathbb{X} with reproducing kernel $k \colon \mathbb{X} \times \mathbb{X} \to \mathbb{C}$, we will let $k_x = k(x, \cdot) \in \mathcal{H}$ be the kernel section at $x \in \mathbb{X}$ and $\delta_x \in \mathcal{H}^*$ the corresponding pointwise evaluation functional, $\delta_x = \langle k_x, \cdot \rangle_{\mathcal{H}}$. Moreover, given a subset $S \subseteq \mathbb{X}$, $\mathcal{H}(S)$ will be the closed subspace of \mathcal{H} defined as $\mathcal{H}(S) = \overline{\operatorname{span}\{k_x : x \in S\}}^{\|\cdot\|_{\mathcal{H}}}$. Note that $\mathcal{H}(S)$ is an RKHS of functions on \mathbb{X} , whose kernel section at $x \in \mathbb{X}$ is equal to the orthogonal projection of k_x onto $\mathcal{H}(S)$. Moreover, $\mathcal{H}(S)$ is isomorphic as an RKHS to the restriction $\mathcal{H}|_S$ of \mathcal{H} onto S, equipped with $k|_{S\times S}$ as the reproducing kernel.

Given a set \mathbb{X} , $1_S \colon X \to \mathbb{R}$ will denote the characteristic function of a subset $S \subseteq X$. If \mathbb{X} is a topological space, $\mathcal{B}(\mathbb{X})$ will denote the Borel σ -algebra of subsets of \mathbb{X} . Given a Borel measure ν and RKHS \mathcal{H} on \mathbb{X} , we will use the abbreviated notation $\mathcal{H}(\nu) \equiv \mathcal{H}(\text{supp}(\nu))$, where $\text{supp}(\nu) \subseteq \mathbb{X}$ is the support of ν .

3. Background and previous work

3.1. Quantum simulation of systems with pure point spectra. In [37] a technique was developed for quantum simulation of a class of measure-preserving, ergodic flows with pure point

spectra; i.e., measure-preserving flows for which the union of eigenspaces of the Koopman operator is dense in $L^p(\mu)$, $p \in [1, \infty)$. These systems are examples of highly structured dynamics that in many ways is antithetical to weak-mixing.

By the Halmos-von Neumann theorem [43], every pure-point-spectrum system is isomorphic in a measure-theoretic sense to a rotation system on a compact abelian group. In discrete time, this implies that the point spectrum $\sigma_p(U)$ of the Koopman operator is a subgroup of \mathbb{T}^1 . In continuous time, the point spectrum $\sigma_p(V)$ of the generator is an additive subgroup of the imaginary line, giving, by the spectral mapping theorem, the point spectra of the Koopman operators as a group homomorphism, $\sigma_p(V) \ni i\omega \mapsto e^{i\omega t} \in \sigma_p(U^t)$.

The paper [37] studied the case of an ergodic torus rotation $R^t \colon \mathbb{T}^d \to \mathbb{T}^d$

$$R^t(\theta) = \theta + \alpha \cdot \theta \mod 2\pi, \quad t \in \mathbb{R},$$

where $\alpha=(\alpha_1,\ldots,\alpha_d)\in\mathbb{R}^d$ are rationally-independent frequency parameters. This system is a canonical representative in the measure-theoretic isomorphism class of pure-point-spectrum, continuous-time ergodic systems with spectra generated by d basic frequencies. Specifically, we have $\sigma_p(V)=\{i(j_1\alpha_1+\cdots+j_d\alpha_d):j_1,\ldots,j_d\in\mathbb{Z}\}$, so the point spectrum of the generator is isomorphic to the Pontryagin dual $\widehat{\mathbb{T}^d}\simeq\mathbb{Z}^d$ of the state space of the dynamics by rational independence of α_1,\ldots,α_d .

Here, the group structure of $\sigma_p(V)$ is a manifestation of the fact that V obeys the Leibniz rule,

$$V(fg) = (Vf)g + f(Vg), \tag{3}$$

for any $f,g \in D(V)$ for which the left-hand-side and right-hand side of the above equation are well-defined. In fact, [71] showed that satisfying (3) on the algebra $L^{\infty}(\mu) \cap D(V)$ is a necessary and sufficient condition for a skew-adjoint operator $V \colon D(V) \to L^2(\mu)$ to be the generator of a unitary Koopman group (i.e., a one-parameter unitary group of composition operators), so one can consider the Leibniz rule as a fundamental structural property of the generators of continuous-time classical dynamical systems.

3.1.1. Choice of Hilbert space. Using results from harmonic analysis [20, 21, 30], the approach of [37] was to build a quantum system on a function space $\mathfrak{A} \subset C(\mathbb{T}^d)$ that is simultaneously a reproducing kernel Hilbert space (RKHS) and a Banach *-algebra with respect to pointwise function multiplication and complex conjugation. On this space \mathfrak{A} , a rotation system $R^t \colon \mathbb{T}^d \to \mathbb{T}^d$ induces unitary Koopman operators $\hat{U}^t \colon \mathfrak{A} \to \mathfrak{A}$ analogously to U^t on $L^2(\mu)$, and the spectrum of the skewadjoint generator $\hat{W} \colon D(\hat{W}) \to \mathfrak{A}$ with $D(\hat{W}) \subseteq \mathfrak{A}$ is again $\sigma_p(\hat{W}) \simeq \mathbb{Z}^d$. Moreover, every classical observable $f \in \mathfrak{A}$ has an associated quantum observable $M_f \in B(\mathfrak{A})$ that acts as a multiplication operator, $M_f g = f g$.

The RKHS \mathfrak{A} has a translation-invariant reproducing kernel $k \colon \mathbb{T}^d \times \mathbb{T}^d \to \mathbb{R}_{>0}$ obtained from the inverse Fourier image of an integrable, strictly positive, symmetric, subconvolutive function on the dual group $\widehat{\mathbb{T}^d} \simeq \mathbb{Z}^d$; i.e., a function $\lambda \colon \mathbb{Z}^d \to \mathbb{R}_{>0}$ satisfying

$$\lambda \in L^1(\mathbb{Z}^d), \quad \lambda(j) > 0, \quad \lambda(-j) = \lambda(j), \quad \lambda * \lambda(j) \le C\lambda(j),$$

where $L^1(\mathbb{Z}^d)$ is defined with respect to the counting measure (equivalently, a Haar measure on $\widehat{\mathbb{T}^d}$). The generator \hat{W} admits the diagonalization

$$\hat{W}\psi_i = i\omega_i\psi_i, \quad \psi_i = \sqrt{\lambda(j)}\phi_i, \quad \omega_i = j_1\alpha_1 + \dots + j_d\alpha_d,$$

where $\phi_j : \mathbb{T}^d \to \mathbb{C}$ are characters (Fourier functions) in the dual group $\widehat{\mathbb{T}^d}$, $\phi_j(\theta) = e^{ij\dot{\theta}}$, indexed by $j = (j_1, \dots, j_d) \in \mathbb{Z}^d$. Moreover, the eigenfunctions ψ_j form an orthonormal basis of \mathfrak{A} , and $\omega_j \in \mathbb{R}$ are corresponding eigenfrequencies.

The simultaneous RKHA and Banach algebra structure of $\mathfrak A$ provides several useful properties for embedding classical dynamics into quantum dynamics and for building associated quantum

algorithms, which we outline below. For detailed treatments of RKHSs we refer the reader to [63,67]. Further details on weight functions in harmonic analysis and associated multiplication/convolution algebras can be found in [29,41,48].

3.1.2. Quantum embedding of classical dynamics. Recall that for the space $B(\mathbb{H})$ of bounded operators on a Hilbert space \mathbb{H} , viewed as a von Neumann algebra with respect to operator composition and adjoint, the space of normal states can be identified with the set of density operators on \mathbb{H} , i.e., the set of positive, trace-class operators $\rho \colon \mathbb{H} \to \mathbb{H}$ of unit trace [69]. We denote this set as $Q(\mathbb{H})$ and interpret it as the set of quantum states on \mathbb{H} . Every $\rho \in Q(\mathbb{H})$ induces a state $\mathbb{E}_{\rho} \colon B(\mathbb{H}) \to \mathbb{C}$ such that $\mathbb{E}_{\rho} a = \operatorname{tr}(\rho a)$. The number $\mathbb{E}_{\rho} a$ corresponds to the expectation of quantum observable a with respect to the quantum state ρ .

When $\mathbb{H} = \mathcal{H}$ is an RKHS of functions on a set X, we have a map $\varphi \colon X \to Q(\mathcal{H})$ of classical states into quantum states induced by the feature map,

$$\varphi \colon X \to \mathcal{H}, \quad \varphi(x) = k_x, \quad k_x = k(x, \cdot),$$
 (4)

where $k: X \times X \to \mathbb{C}$ is the reproducing kernel of \mathcal{H} . Specifically, defining $\varphi(x) \equiv \rho_x = \langle \xi_x, \cdot \rangle_{\mathcal{H}} \xi_x$ with $\xi_x = \varphi(x)/\sqrt{k(x,x)}$, one readily verifies that ρ_x is a rank-1 quantum state that projects along the unit vector $\xi_x \in \mathcal{H}$. If the feature map φ is injective (which is the case for many RKHS examples; e.g., [66]), then so is φ , so we have an embedding on classical states into quantum states. If, in addition, $\mathcal{H} = \mathfrak{A}$ is Banach algebra with respect to pointwise function multiplication, there is a faithful representation $\pi: \mathfrak{A} \to B(\mathfrak{A})$ of classical observables in \mathfrak{A} as multiplication operators in $B(\mathfrak{A})$,

$$(\pi f)g = fg, \quad \forall g \in \mathfrak{A}. \tag{5}$$

Let $\delta_x : \mathfrak{A} \to \mathbb{C}$ be the pointwise evaluation functional at $x \in X$, where $\delta_x = \langle k_x, \cdot \rangle_{\mathfrak{A}}$ since \mathfrak{A} is an RKHS [63]. One readily verifies that the quantum embeddings of states and observables through φ and π , respectively, are consistent with pointwise evaluation,

$$\mathbb{E}_{\varphi(x)}(\pi f) = \delta_x f = f(x), \quad \forall f \in \mathfrak{A}, \quad \forall x \in \mathbb{X}.$$

In the case of the rotation system on $X=\mathbb{T}^d$, these embeddings are also compatible with dynamical evolution. Defining the adjoint actions $\mathbf{R}^t\colon Q(\mathfrak{A})\to Q(\mathfrak{A})$ and $\mathbf{U}^t\colon B(\mathfrak{A})\to B(\mathfrak{A})$ of the Koopman group on quantum states and observables as $\mathbf{R}^t(\rho)=\hat{U}^{t*}\rho\hat{U}^t$ and $\mathbf{U}^tA=\hat{U}^tA\hat{U}^{t*}$, respectively, we have $\boldsymbol{\varphi}\circ R^t=\mathbf{R}^t\circ \boldsymbol{\varphi}$ and $\pi\circ \hat{U}^t=\mathbf{U}^t\circ \pi$; that is, the quantum feature map $\boldsymbol{\varphi}$ and multiplier representation π intertwine the classical and quantum evolutions of states and observables, respectively. Equivalently, the following two diagrams commute:

3.2. Spectral approximation for systems with continuous spectra. Let us recall the following fundamental results on unitary Koopman groups associated with measure-preserving ergodic flows (e.g., [42]):

Theorem 1. With the notation and assumptions of section 2.1, the Hilbert space H admits a U^t -invariant orthogonal decomposition $H = H_p \oplus H_c$, where the subspaces H_p and H_c have the following properties.

(i) H_p admits an orthonormal basis $\{\xi_i\}$ consisting of eigenvectors of the generator,

$$V\xi_j = i\omega_j\xi_j, \quad \omega_j \in \mathbb{R},$$

where all eigenvalues $i\omega_j$ are simple and can be indexed by integers j such that $\omega_{-j} = -\omega_j$ (in particular, $\omega_0 = 0$). Moreover, the set $\{i\omega_j\}$ constitutes the point spectrum $\sigma_p(V)$, and the corresponding eigenvectors can be chosen such that $\xi_{-j} = \xi_j^*$ with $\xi_0 = 1_X$.

(ii) Elements of H_c exhibit the following form of decay of correlations (also known as weak-mixing behavior),

$$\lim_{T \to \infty} \frac{1}{T} \int_0^T |C_{fg}(t)| dt = 0, \quad \forall f \in H, \quad \forall g \in H_c,$$

where $C_{fg}(t) = \langle f, U^t g \rangle_H$.

By Theorem 1(i), every element $f \in H_p$ evolves as an observable of a pure point spectrum system,

$$U^{t}f = \sum_{j} e^{i\omega_{j}t} \langle \xi_{j}, f \rangle_{H} \xi_{j}, \tag{6}$$

and is thus amenable to quantum simulation via the techniques described in section 3.1. On the other hand, it is not obvious how to apply these methods to observables in H_c , for this space does not admit an orthonormal basis consisting of generator eigenfunctions. To overcome this obstacle, we will approximate the generator V by a family of skew-adjoint operators that are diagonalizable on the entire Hilbert space H, and whose eigenfunctions behave as approximate Koopman eigenfunctions that we will use for building rotation systems approximating the dynamical flow Φ^t .

3.2.1. Spectrally accurate approximations. The development of analytical and computational techniques for spectral approximation of Koopman and transfer operators of measure-preserving systems has been a highly active research area in recent years; e.g., [17,61] and references therein. In this work, we employ a variant of the techniques developed in [22,39,40] that yield approximations of the generator that are spectrally accurate in the sense of strong convergence of resolvents.

Definition 2. A family of skew-adjoint operators $A_{\tau} \colon D(A_{\tau}) \to \mathbb{H}$, $\tau > 0$, on a Hilbert space \mathbb{H} is said to converge to a skew-adjoint operator $A \colon D(A) \to \mathbb{H}$ in strong resolvent sense if for some (and thus, every) $z \in \mathbb{C} \setminus i\mathbb{R}$ the resolvents $R(z, A_{\tau})$ converge strongly to R(z, A); that is, $\lim_{\tau \to 0^+} \|(R(z, A_{\tau}) - R(z, A))f\|_{\mathbb{H}} = 0$ for every $f \in \mathbb{H}$.

It can be shown that strong resolvent convergence $A_{\tau} \to A$ is equivalent to strong convergence of the corresponding unitaries, $e^{tA_{\tau}} \to e^{tA}$ for every $t \in \mathbb{R}$ (also known as strong dynamical convergence); e.g., [23, Proposition 10.1.8]. For our purposes, this implies that if a family of skew-adjoint operators converges to the Koopman generator V in strong resolvent sense, the unitary evolution groups generated by these operators consistently approximate the Koopman operators $U^t = e^{tV}$ generated by V.

Strong resolvent convergence and strong dynamical convergence imply the following form of spectral convergence; [22, Proposition 13].

Theorem 3. With the notation of Definition 2, let $\tilde{E} : \mathcal{B}(i\mathbb{R}) \to B(\mathbb{H})$ and $\tilde{E}_{\tau} : \mathcal{B}(i\mathbb{R}) \to B(\mathbb{H})$ be the spectral measures of A and A_{τ} , respectively, i.e., $A = \int_{i\mathbb{R}} \lambda \, d\tilde{E}(\lambda)$ and $A_{\tau} = \int_{i\mathbb{R}} \lambda \, d\tilde{E}_{\tau}(\lambda)$. Then, the following hold under strong resolvent convergence of A_{τ} to A.

- (i) For every element $\lambda \in \sigma(A)$ of the spectrum of A, there exists a sequence $\tau_1, \tau_2, \ldots \searrow 0$ and elements $\lambda_n \in \sigma(A_{\tau_n})$ of the spectra of A_{τ_n} such that $\lim_{n\to\infty} \lambda_n = \lambda$.
- (ii) For every bounded continuous function $h: i\mathbb{R} \to \mathbb{C}$, as $\tau \to 0^+$ the operators $h(A_\tau) = \int_{i\mathbb{R}} h(\lambda) d\tilde{E}_{\tau}(\lambda)$ converge strongly to $h(A) = \int_{i\mathbb{R}} h(\lambda) d\tilde{E}(\lambda)$.
- (iii) For every bounded Borel-measurable set $\Theta \in \mathcal{B}(i\mathbb{R})$ such that $\tilde{E}(\partial\Theta) = 0$ (i.e., the boundary of Θ does not contain eigenvalues of A_{τ}), as $\tau \to 0^+$ the projections $\tilde{E}_{\tau}(\Theta)$ converge strongly to $\tilde{E}(\Theta)$.

3.2.2. Markov smoothing operators. In broad terms, the schemes of [22, 39, 40] regularize the generator or its resolvent by composing it with Markov smoothing operators with associated RKHSs of continuous, complex-valued functions on the state space X.

These RKHSs, \mathcal{H}_{τ} , are defined for each $\tau > 0$ to have a bounded, continuous, strictly positivedefinite kernel $k_{\tau} \colon X \times X \to \mathbb{R}_{>0}$, built such that $k_{\tau}|_{M \times M}$ is C^1 . By standard results on RKHS theory (e.g., [63,67]), $K_{\tau} \colon H \to \mathcal{H}_{\tau}$ with

$$K_{\tau}f = \int_{X} k_{\tau}(\cdot, x) f(x) d\mu(x)$$

is a well-defined compact integral operator. Moreover, we have that (i) the range of K_{τ} is a dense subspace of $\mathcal{H}_{\tau}(\mu)$; (ii) $K_{\tau}^*: \mathcal{H}_{\tau} \to H$ implements the inclusion map (that is, $K_{\tau}^*|_{\mathcal{H}_{\tau}(M)} = \iota|_{\mathcal{H}_{\tau}(M)}$); and (iii) $\mathcal{H}_{\tau}(M)$ is a subspace of $C^1(M)$. In particular, the action of the generator on elements of \mathcal{H}_{τ} can be evaluated via (2).

Defining $G_{\tau} = K_{\tau}^* K_{\tau}$ for $\tau > 0$ and G_0 as the identity operator on H, we require that $\{G_{\tau}\}_{\tau \geq 0}$ is a strongly-continuous semigroup of strictly positive, Markov operators. This means:

- (K1) $\langle f, G_{\tau} f \rangle_H > 0$ whenever $f \neq 0$ (i.e., G_{τ} is strictly positive as an operator in B(H)).
- (K2) $G_{\tau}1_X = 1_X$ and $G_{\tau}f \geq 0$ whenever $g \geq 0$ (i.e., G_{τ} is a Markov operator).
- (K3) $G_{\tau} \circ G_{\tau'} = G_{\tau+\tau'}$ for every $\tau, \tau' \geq 0$ (i.e., $\{G_{\tau}\}_{\tau \geq 0}$ is a semigroup).
- (K4) $\lim_{\tau\to 0^+} G_{\tau}f = f$ for every $f\in H$ (i.e., $\{G_{\tau}\}_{\tau>0}$ is strongly continuous).

It can also be shown that K_{τ} admits the polar decomposition

$$K_{\tau} = T_{\tau} G_{\tau/2},\tag{7}$$

where $T_{\tau}: H \to \mathcal{H}_{\tau}$ is an isometry with range $\mathcal{H}_{\tau}(\mu)$.

Possible ways of constructing kernels k_{τ} satisfying properties (K1)–(K4) include normalization of Gaussian kernels on $X = \mathbb{R}^n$ and Fourier transform of positive functions on the dual group \widehat{X} when X is a compact abelian group and the invariant measure μ is a Haar measure (cf. λ from section 3.1). In appendix A we give an outline of the Markov normalization approach used in the numerical experiments of section 8. Examples of Markov kernel constructions on compact abelian groups via eight functions can be found, e.g., in [20, 35, 36].

- 3.2.3. Diagonalizable approximations of the generator. Using the smoothing operators G_{τ} , the papers [22,39] builds 1-parameter families of densely-defined operators $V_{\tau} : D(V_{\tau}) \to H$ with $z, \tau > 0$ and $D(V_{\tau}) \subseteq H$ with the following properties.
 - (V1) V_{τ} is skew-adjoint.
 - (V2) V_{τ} is compact [22] or has compact resolvent [39].
 - (V3) V_{τ} is real, $(V_{\tau}f)^* = V_{\tau}(f^*)$ for all $f \in D(V)$.
 - (V4) V_{τ} annihilates constant functions, $V_{\tau} \mathbf{1} = 0$.
 - (V5) V_{τ} converges to V in strong resolvent sense as $\tau \to 0^+$.

In [40] a related approach is developed whereby V is approximated by a 2-parameter family of operators $V_{z,\tau}$ satisfying properties (V1)–(V4) and converging to V in strong resolvent sense in the iterated limits $z \to 0^+$ after $\tau \to 0^+$. A key feature of this approach is that it is "physics-informed" in the sense of using known equations of motion to evaluate the action of the generator by means (2). We review this method in appendix B as it will be used in our numerical examples.

Here, as a concrete example we mention the (formulaically simpler) approach of [22], who define the compact approximations

$$V_{\tau} = G_{\tau/2} V G_{\tau/2}.$$

These operators are manifestly skew-adjoint. Their compactness follows by compactness of $G_{\tau/2}$, boundedness of $\vec{V} \cdot \nabla$ as an operator from $C^1(M)$ to C(M), and the relation

$$VG_{\tau/2} = \iota \circ (\vec{V} \cdot \nabla) \circ K_{\tau/2}.$$

Unless stated otherwise, in what follows we will use the abbreviated notation V_{τ} to denote any of the operators V_{τ} or $V_{z,\tau}$.

3.2.4. Approximate Koopman eigenfunctions. For every $\tau > 0$, V_{τ} is unitarily equivalent to a skew-adjoint operator $W_{\tau} : D(W_{\tau}) \to \mathcal{H}_{\tau}\mu$ with domain $D(W_{\tau}) = T_{\tau}(D(V_{\tau})) \cup \ker T_{\tau}^*$, defined as

$$W_{\tau} = T_{\tau} V_{\tau} T_{\tau}^*$$
.

Both V_{τ} and W_{τ} are unitarily diagonalizable, and there exist orthonormal bases $\{\xi_{j,\tau}\}$ and $\{\zeta_{j,\tau}\}$ of H and $\mathcal{H}_{\tau}(\mu)$, respectively, consisting of their eigenvectors,

$$V_{\tau}\xi_{j,\tau} = i\omega_{j,\tau}\xi_{j,\tau}, \quad W_{\tau}\zeta_{j,\tau} = i\omega_{j,\tau}\zeta_{j,\tau}, \quad \zeta_{j,\tau} = T_{\tau}\xi_{j,\tau}$$
(8)

where $\omega_{j,\tau} \in \mathbb{R}$ are corresponding eigenfrequency. By analogy of eigenvectors/eigenfrequencies of pure-point-spectrum systems in Theorem 1(i), we use indices $j \in \mathbb{Z}$ such that $\omega_{-j,\tau} = -\omega_{j,\tau}$ and choose the corresponding eigenvectors such that $\xi_{-j,\tau} = \xi_{j,\tau}^*$ and $\zeta_{j,\tau} = \zeta_{j,\tau}^*$ with $\xi_{0,\tau} = \zeta_{j,\tau} = 1_X$. Following [40], we order the eigenfunctions $\zeta_{0,\tau}, \zeta_{\pm 1,\tau}, \zeta_{\pm 2,\tau}$ in increasing order of a Dirichlet energy functional; see appendix B for more details.

On the basis of property (V5) and Theorem 3, we interpret $\xi_{j,\tau}/\zeta_{j,\tau}$ and $\omega_{j,\tau}$ as approximate Koopman eigenfunctions and eigenfrequencies, respectively. The corresponding unitaries satisfy (cf. (6))

$$e^{tV_{\tau}} = \sum_{j} e^{i\omega_{j,\tau}t} \langle \xi_{j,\tau}, \cdot \rangle_{H} \xi_{j,\tau}, \quad e^{tW_{\tau}} = \sum_{j} e^{i\omega_{j,\tau}t} \langle \zeta_{j,\tau}, \cdot \rangle_{\mathcal{H}_{\tau}} \zeta_{j,\tau} + \operatorname{proj}_{\mathcal{H}_{\tau}(\mu)^{\perp}},$$

and $e^{tV_{\tau}}f$ converges as $\tau \to 0^+$ to U^tf for every $f \in H$ by property (V5) and equivalence of strong resolvent convergence and strong dynamical convergence. It can also be shown [35, Lemma 6] that for every $f \in \mathcal{H}_{\tau'}$, the evolution under $e^{tW_{\tau}}$, $\tau \leq \tau'$, converges to the true Koopman evolution in L^2 sense,

$$\lim_{\tau \to 0^+} \left\| (K_{\tau}^* e^{tW_{\tau}} - U^t \iota) f \right\|_H = 0. \tag{9}$$

For the remainder of the paper we will use the notation $U_{\tau}^t := e^{tW_{\tau}}$.

3.3. Reproducing kernel Hilbert algebras. RKHAs are RKHSs equipped with coalgebra structure that will be central to the Fock space scheme studied in this paper. In this subsection, we give basic definitions and outline some of properties of RKHAs that are most relevant to our work, referring the reader to [36] for further details.

Definition 4. An RKHS \mathcal{H} on a set X with reproducing kernel $k: X \times X \to \mathbb{C}$ is a reproducing kernel Hilbert algebra (RKHA) if $k_x \mapsto k_x \otimes k_x$, $x \in X$, extends to a bounded linear map (comultiplication) $\Delta \colon \mathcal{H} \to \mathcal{H} \otimes \mathcal{H}$.

Since

$$\langle k_x, \Delta^*(f \otimes g) \rangle_{\mathcal{H}} = \langle \Delta k_x, f \otimes g \rangle_{\mathcal{H}} = \langle k_x \otimes k_x, f \otimes g \rangle_{\mathcal{H} \otimes \mathcal{H}} = \langle k_x, f \rangle_{\mathcal{H}} \langle k_x, g \rangle_{\mathcal{H}} = f(x)g(x),$$

it follows that Δ^* is a bounded linear map that implements pointwise multiplication. As a result, \mathcal{H} is simultaneously a Hilbert function space and commutative algebra with respect to pointwise function multiplication. It can further be shown that \mathcal{H} is a Banach algebra,

$$|||fg||_{\mathcal{H}} \le |||f||_{\mathcal{H}} |||g||_{\mathcal{H}}, \quad \forall f, g \in \mathcal{H},$$

for an operator norm $\|\cdot\|_{\mathcal{H}}$ induced by the multiplier representation $\pi\colon\mathcal{H}\to B(\mathcal{H})$ (defined analogously to (5)) that generates a coarser topology than the Hilbert space norm. By commutativity and associativity of pointwise multiplication, it follows that the comultiplication operator $\Delta:\mathcal{H}\to\mathcal{H}\otimes\mathcal{H}$ is cocommutative and coassociative, which implies in turn that $\Delta_n\colon\mathcal{H}\to\mathcal{H}^{\otimes n}$ with

$$\Delta_1 = \Delta, \quad \Delta_n = (\Delta \otimes \operatorname{Id}^{\otimes (n-1)}) \Delta_{n-1} \quad \text{for } n > 1,$$
(10)

is a well-defined amplification of Δ to the tensor product spaces $\mathcal{H}^{\otimes (n+1)}$ for any $n \in \mathbb{N}$.

If the reproducing kernel k is real-valued, \mathcal{H} becomes a *-algebra with isometric involution *: $\mathcal{H} \to \mathcal{H}$ given by the pointwise conjugation of functions. If \mathcal{H} contains the constant function $1_X : X \to \mathbb{R}$ that equals 1 everywhere on X, then it is a unital algebra with unit 1_X .

Given an RKHA \mathcal{H} , we will let $\sigma(\mathcal{H}) \subset \mathcal{H}^*$ denote its spectrum as a Banach algebra, i.e., the set of nonzero multiplicative linear functionals $\chi : \mathcal{H} \to \mathbb{C}$,

$$\chi(fg) = (\chi f)(\chi g), \quad \forall f, g \in \mathcal{H},$$

equipped with the weak-* topology of \mathcal{H}^* . The dual object to $\sigma(\mathcal{H})$ is the cospectrum $\sigma_{co}(\mathcal{H})$, which is defined as

$$\sigma_{\rm co}(\mathcal{H}) = \{ \xi \in \mathcal{H} : \langle \xi, \cdot \rangle_{\mathcal{H}} \in \sigma(\mathcal{H}) \}$$

and is equipped with the weak topology of \mathcal{H} . Equivalently, we have that $\sigma_{co}(\mathcal{H})$ is the subset of \mathcal{H} consisting of elements ξ such that $\Delta \xi = \xi \otimes \xi$.

It is clear that the kernel sections k_x and evaluation functionals $\delta_x = \langle k_x, \cdot \rangle_{\mathcal{H}}$ are elements of $\sigma_{\text{co}}(\mathcal{H})$ and $\sigma(\mathcal{H})$, respectively, for every $x \in X$. In particular, the feature map φ from (4) can be viewed as a map from X into the cospectrum $\sigma_{\text{co}}(\mathcal{H})$. This map is injective whenever \mathcal{H} has linearly independent kernel sections, and if X is a topological space it is continuous iff $\mathcal{H} \subseteq C(X)$. If \mathcal{H} is unital, $\sigma(\mathcal{H})$ and $\sigma_{\text{co}}(\mathcal{H})$ are compact Hausdorff spaces.

The paper [36] developed constructions that can be used to build many examples of RKHAs. These examples include the spaces \mathfrak{A} on $X=\mathbb{T}^d$ introduced in section 3.1 which are unital RKHAs of continuous functions with homeomorphic spectra and cospectra to X. More generally, for every compact subset $X \subset \mathbb{R}^n$ there exists a unital RKHA $\mathcal{H} \subset C(\mathbb{R}^n)$ with $X \cong \sigma(\mathcal{H}) \cong \sigma_{co}(\mathcal{H})$.

3.4. **Tensor network approximation.** As mentioned in section 3.1, a key structural property of skew-adjoint generators of unitary Koopman groups is that they act as derivations on algebras of observables via the Leibniz rule (3). While this property can considerably aid the efficiency of quantum simulation algorithms, aside from special cases (e.g., a priori known rotation systems as in [37]), one is compelled to work with regularizations of the generator that fail to satisfy the Leibniz rule. Examples include the methods of [22, 39, 40] outlined in section 3.2, as well several other methods for discrete approximation of the typically continuous spectral measures exhibited by unitary Koopman groups under complex measure-preserving dynamics; e.g., [18, 22].

To our knowledge, there is currently no operator approximation methodology for unitary Koopman groups associated with measure-preserving flows that simultaneously preserves skew-adjointness and the Leibniz rule for the generator (but note the recent MultDMD technique [13] that yields non-unitary approximations of the Koopman operator that preserve multiplicativity). In the transfer operator literature, a powerful approach for spectral analysis of hyperbolic dynamics is to work in anisotropic Banach spaces adapted to the stable/unstable directions of the dynamics, where the transfer operator is quasicompact (and thus exhibits isolated eigenvalues in the unit disc that are separated from an essential spectrum) [4,11,14]. In general, however, these spaces lack the Hilbert space structure and unitary dynamics required of quantum systems.

3.4.1. Dilation to Fock space. As an effort to overcome these challenges, [35] developed a dilation scheme that lifts a given regularized generator to a Fock space generated by an RKHA, in which the Leibniz rule is recovered. In more detail, they consider a family of approximations $V_{\tau} \colon D(V_{\tau}) \to H$ of the generator satisfying properties (V1)–(V5) and the convergence property (9) for the associated operators $W_{\tau} \colon D(W_{\tau}) \to \mathcal{H}_{\tau}$ acting on the RKHSs $\mathcal{H}_{\tau} \subseteq C(X)$.

Assuming that X is a compact Hausdorff space, the Hilbert spaces \mathcal{H}_{τ} are built as unital RKHAs \mathcal{H}_{τ} with isomorphic (co)spectrum to X. Each RKHA \mathcal{H}_{τ} in this family generates an associated Fock space, $F(\mathcal{H}_{\tau})$, defined using standard constructions from many-body quantum theory as the

Hilbert space closure of the tensor algebra $T(\mathcal{H}_{\tau}) := \mathbb{C} \oplus \mathcal{H}_{\tau} \oplus \mathcal{H}_{\tau}^{\otimes 2} \oplus \ldots$ with respect to the inner product $\langle \cdot, \cdot \rangle_{F(\mathcal{H}_{\tau})}$ satisfying

$$\langle a, b \rangle_{F(\mathcal{H}_{\tau})} = \bar{a}b, \quad a, b \in \mathbb{C},$$

$$\langle a, f \rangle_{F(\mathcal{H}_{\tau})} = 0, \quad a \in \mathbb{C}, \quad f \in \mathcal{H}_{\tau},$$

$$\langle f_1 \otimes \cdots \otimes f_n, g_1 \otimes \cdots \otimes g_n \rangle_{F(\mathcal{H}_{\tau})} = \prod_{i=1}^n \langle f_i, g_i \rangle_{\mathcal{H}_{\tau}}, \quad f_i, g_i \in \mathcal{H}_{\tau}.$$

$$(11)$$

The scheme of [35] lifts the regularized generator W_{τ} to a skew-adjoint operator $\tilde{W}_{\tau} \colon D(W_{\tau}) \to F(\mathcal{H}_{\tau})$, defined by linear extension of

$$\tilde{W}_{\tau}(f_1 \otimes f_2 \otimes \cdots \otimes f_n) = (W_{\tau}f_1) \otimes f_2 \otimes \cdots \otimes f_n + f_1 \otimes (W_{\tau}f_2) \otimes \cdots \otimes f_n + \cdots + f_1 \otimes f_2 \otimes \cdots \otimes (W_{\tau}f_n)$$

for $f_1, \ldots, f_n \in D(W_\tau)$. By construction, \tilde{W}_τ satisfies the Leibniz rule

$$\tilde{W}_{\tau}(f \otimes g) = (\tilde{W}_{\tau}f) \otimes g + f \otimes (\tilde{W}_{\tau}g)$$

with respect to the tensor product for all $f, g \in D(\tilde{W}_{\tau})$ such that the left- and right-hand sides of the above equation are well-defined. As a result, we have:

- (1) The point spectrum $\sigma_p(\tilde{W}_{\tau})$ is an abelian group generated by $\sigma_p(W_{\tau})$.
- (2) \tilde{W}_{τ} generates a 1-parameter group of unitary operators $\tilde{U}_{\tau}^{t} = e^{tW_{\tau}}$, $t \in \mathbb{R}$, that act multiplicatively with respect to the tensor product,

$$\tilde{U}_{\tau}^{t}(f \otimes g) = (\tilde{U}_{\tau}^{t}f) \otimes g + f \otimes (\tilde{U}_{\tau}^{t}g), \quad \forall, f, g \in F(\mathcal{H}_{\tau}).$$

Putting together the above, it follows that for every vector $q \in L^2(\mu)$ with a representative $\xi \in \mathcal{H}_{\tau_0}$ for some $\tau_0 > 0$ and a multiplicative decomposition of the form $\xi = \xi_1 \dots \xi_n$ for some $\xi_1, \dots, \xi_n \in \mathcal{H}_{\tau_0}$, we have

$$\Delta_n^* \tilde{U}_\tau^t \xi = \Delta_n^* \left(\bigotimes_{i=1}^n U_\tau^t \xi_i \right) = \prod_{i=1}^n U_\tau^t \xi_i \xrightarrow{\tau \to 0^+} U^t q \tag{12}$$

in $L^2(\mu)$ norm, where Δ_n is the amplified comultiplication operator defined in (10). Intuitively, the Fock space $F(\mathcal{H}_{\tau})$ generated by the RKHA \mathcal{H}_{τ} allows one to "distribute" the Koopman evolution of observables over tensor products in the Fock space for potentially arbitrarily high grading $n \in \mathbb{N}$.

3.4.2. Quantum representation of statistical evolution. The approach of [35] leverages (12) to build an approximation of the classical expectation $\mathbb{E}_p(U^t f) \equiv \int_X (U^t f) p \, d\mu$ for an observable $f \in L^{\infty}(\mu)$ and a probability density $p \in L^1(\mu)$. To any desired error tolerance, the square root \sqrt{p} may be approximated by an element $q \in L^2(\mu)$ with a strictly positive representative $\xi \in \mathcal{H}_{\tau_0}$. The latter may be written in turn as $\xi = (\xi^{1/n})^n$, $n \in \mathbb{N}$ (using the holomorphic functional calculus on \mathcal{H}_{τ_0} to compute the n-th root $\xi^{1/n} \in \mathcal{H}_{\tau}$), and dynamically evolved using the unitaries \tilde{U}_{τ}^t on the Fock space as in (12).

To cast this in the form of a quantum state evolution, we map ξ to a pure quantum state $\rho_{\tau} = \langle \eta_{\tau}, \cdot \rangle_{F(\mathcal{H}_{\tau})} \eta_{\tau} \in Q(F(\mathcal{H}_{\tau})), \ \tau \leq \tau_0$, with state vector

$$\eta_{\tau} = s_1 \frac{\xi}{\|\xi\|_{\mathcal{H}_{\tau}}} + s_2 \frac{\xi^{1/2} \otimes \xi^{1/2}}{\|\xi^{1/2}\|_{\mathcal{U}}^2} + \ldots \in F(\mathcal{H}_{\tau}).$$

Here, s_1, s_2, \ldots is an arbitrary sequence of strictly positive numbers such that $\sum_{i=1}^{\infty} s_i^2 = 1$.

On the side of the observable, we first smooth f to an element $f_{\tau} = K_{\tau}f \in \mathcal{H}_{\tau}$ where K_{τ} is the kernel integral operator induced by the reproducing kernel of \mathcal{H}_{τ} ,

$$K_{\tau}f = \int_X k_{\tau}(\cdot, x) \, d\mu(x).$$

Then, for $n \in \mathbb{N}$ we map the multiplication operator $M_{f_{\tau}} \in B(\mathcal{H}_{\tau})$ (which is a bounded quantum observable by virtue of the RKHA structure of \mathcal{H}_{τ} , as in section 3.1.2) to a quantum observable $A_{f,\tau,n} \in B(F(\mathcal{H}_{\tau}))$ on the Fock space given by the following amplification of $M_{f_{\tau}}$:

$$A_{f,\tau,n} = \Delta_n M_{f_\tau} \Delta_n^*$$
.

Proposition 9 in [35] then shows that $\mathbb{E}_p(U^t f)$ is approximated to arbitrarily high accuracy as $\tau \to 0^+$ by the normalized quantum expectation

$$f_{\tau,n}^{(t)} = \frac{\mathbb{E}_{\rho_{\tau}}(\tilde{\mathbf{U}}_{\tau}^t A_{f,\tau,n})}{\mathbb{E}_{\rho_{\tau}}(\tilde{\mathbf{U}}_{\tau}^t A_{1_X,\tau,n})},\tag{13}$$

where $\tilde{\boldsymbol{U}}_{\tau}^{t} \colon B(F(\mathcal{H}_{\tau})) \to B(F(\mathcal{H}_{\tau}))$ is the adjoint operator acting on quantum observables on the Fock space via $\tilde{\boldsymbol{U}}_{\tau}^{t} A = \tilde{U}_{\tau}^{t} A \tilde{U}_{\tau}^{t*}$.

3.4.3. Finite-rank approximation. An interesting aspect of the approximation (13) is that it converges as $\tau \to 0^+$ for any $n \in \mathbb{N}$. We can take advantage of this fact by projecting the state vector η_{τ} to a Fock subspace $F(\mathcal{Z}_{\tau,d}) \subset F(\mathcal{H}_{\tau})$ generated by a finite-dimensional subspace $\mathcal{Z}_{\tau,d} \subset \mathcal{H}_{\tau}$ of dimension 2d+1, spanned by d linearly independent eigenfunctions of W_{τ} with nonzero corresponding eigenvalue, their complex conjugates, and the constant eigenfunction 1_X such that $W_{\tau}1_X = 0$. Even though $F(\mathcal{Z}_{\tau,d})$ is infinite-dimensional, the fact that the range of $A_{f,\tau,n}$ lies in the subspace $\mathcal{H}_{\tau}^{\otimes n} \subset F(\mathcal{H}_{\tau})$ of fixed grading n means that the quantum expectation of $\tilde{U}_{\tau}^t A_{f,\tau,n}$ appearing in (13) with respect to the projected state onto $F(\mathcal{Z}_{\tau,d})$ is equal to the expectation of a finite-rank operator $\tilde{U}_{\tau}^t A_{f,\tau,n,d}$ with respect to ρ_{τ} , where ran $A_{f,\tau,n,d} \subseteq \mathcal{Z}_{\tau,d}^{\otimes n} \subset F(\mathcal{Z}_{\tau,d})$.

Letting $f_{\tau,n,d}^{(t)} \in \mathbb{C}$ denote the analog of $f_{\tau,n}^{(t)}$ resulting from this approximation,

$$f_{\tau,n,d}^{(t)} = \frac{\mathbb{E}_{\rho_{\tau}}(\tilde{\boldsymbol{U}}_{\tau}^{t} A_{f,\tau,n,d})}{\mathbb{E}_{\rho_{\tau}}(\tilde{\boldsymbol{U}}_{\tau}^{t} A_{1_{X},\tau,n,d})},\tag{14}$$

it was shown [35, Theorem 12] that $f_{\tau,n,d}^{(t)}$ converges to $f_{\tau,n}^{(t)}$ as $n \to \infty$ for fixed sufficiently large $d \in \mathbb{N}$.

Intuitively, this result signifies that due to the multiplicative nature of the underlying Koopman operator U^t , the correspondence between tensor products and pointwise products induced by the RKHA structure of \mathcal{H}_{τ} , and the strong convergence $U^t_{\tau} \to U^t$ as $\tau \to 0^+$, the quantum evolution $t \mapsto \tilde{U}^t_{\tau} A_{f,\tau,n,d}$ taking place in subspaces of $F(\mathcal{Z}_{\tau,d})$ of increasingly large dimension $(2d+1)^n$ captures sufficient information about the true Koopman evolution $t \mapsto U^t f$ so as to recover it fully as $\tau \to 0^+$ and $n \to \infty$ (at appropriate rates), even if d is held fixed to a finite value.

Practically, this allows one to devote resources to compute a collection of approximate Koopman eigenfunctions of modest size 2d+1, and algebraically amplify them to access tensor product approximation spaces $\mathcal{Z}_{\tau,d}^{\otimes n}$ of dimension $(2d+1)^n$.

In [35], it was shown that (14) can be efficiently evaluated by a tensor network with tree structure, alleviating the cost of brute-force computations in $\mathcal{Z}_{\tau,d}^{\otimes n}$ that increases exponentially with n. Numerical experiments with low-dimensional dynamical systems (including the Stepanoff flow example studied in this paper) found significant improvement in prediction skill for data-driven implementations of the tensor network approach over conventional models based on linear combinations of Koopman eigenfunctions.

4. SECOND QUANTIZATION FRAMEWORK

Our second-quantization approach framework the measure-preserving dynamical system introduced in section 2.1 by a rotation system on a family of tori embedded in the spectrum of a commutative Banach algebra, built as a weighted symmetric Fock space. In this section, we describe the main steps of our approach, using as a starting point an approximation of the skew-adjoint generator $V: D(V) \to H$ by a family of skew-adjoint, diagonalizable operators $V_{\tau}: D(V_{\tau}) \to H$ and their RKHS counterparts $W_{\tau}: D(W_{\tau}) \to \mathcal{H}_{\tau}$ satisfying properties (V1)–(V5) and (9). As noted in section 3.2, the numerical results in this paper are based in the method of [40] (summarized in appendices A and B), but any other technique satisfying these properties may also be employed.

Similarly to the tensor network approximation scheme [35] outlined in section 3.4, to recover the Leibniz rule lost through regularization, we construct an amplification $\tilde{W}_{\tau} : D(\tilde{W}_{\tau}) \to F_w(\mathcal{H}_{\tau})$ of W_{τ} to a Fock space, $F_w(\mathcal{H}_{\tau})$, generated by \mathcal{H}_{τ} on which it acts as an algebra derivation. Two major differences between our approach and the spaces $F(\mathcal{H}_{\tau})$ from [35] are that (i) $F_w(\mathcal{H}_{\tau})$ is based on the symmetric tensor algebra of \mathcal{H}_{τ} ; and (ii) Hilbert space closure is taken with respect to a weighted inner product. As a result, $F_w(\mathcal{H}_{\tau})$ becomes a Banach algebra with respect to the symmetric tensor product that is isomorphic to an RKHA of continuous functions on its spectrum. In addition, unlike [35], we do not require that the \mathcal{H}_{τ} are RKHAs. This affords the second-quantization approach presented in this paper greater flexibility. The Fock space $F_w(\mathcal{H}_{\tau})$, generator \tilde{W}_{τ} , and related RKHAs to $F_w(\mathcal{H}_{\tau})$ will be the foci of sections 4.1–4.4.

Next, in section 4.5, we employ the unitary evolution group generated by W_{τ} to build a rotation system on the Banach algebra spectrum $\sigma(F_w(\mathcal{H}_{\tau}))$ that will serve as an approximate topological model of the unitary Koopman evolution of observables generated by V. In sections 4.6 and 4.7, we couple this model with a procedure for embedding observables of the original system into $F_w(\mathcal{H}_{\tau})$ and its associated RKHAs. The outcome of this construction is an asymptotically consistent approximation of the unitary Koopman evolution of observables under potentially mixing dynamics by trigonometric (Fourier) polynomials of arbitrarily large degree on suitably chosen tori. A key aspect of this approximation is that it captures information from products of approximate Koopman eigenfunctions (the eigenfunctions of W_{τ}). In addition, the approximation is positivity preserving.

4.1. Weighted Fock space. The space $F_w(\mathcal{H}_{\tau})$ is constructed as the closure of the symmetric tensor algebra $T^{\vee}(\mathcal{H}_{\tau}) := \mathbb{C} \oplus \mathcal{H}_{\tau} \oplus \mathcal{H}_{\tau}^{\vee 2} \oplus \dots$ with respect to the inner product satisfying (cf. (11))

$$\langle a, b \rangle_{F_{w}(\mathcal{H}_{\tau})} = \bar{a}b, \quad a, b \in \mathbb{C},$$

$$\langle a, f \rangle_{F_{w}(\mathcal{H}_{\tau})} = 0, \quad a \in \mathbb{C}, \quad f \in \mathcal{H}_{\tau},$$

$$\langle f_{1} \vee \dots \vee f_{n}, g_{1} \vee \dots \vee g_{n} \rangle_{F_{w}(\mathcal{H}_{\tau})} = \frac{w^{2}(n)}{n!^{2}} \sum_{\sigma, \sigma' \in S} \prod_{i=1}^{n} \langle f_{\sigma(i)}, g_{\sigma'(i)} \rangle_{\mathcal{H}_{\tau}}, \quad f_{i}, g_{i} \in \mathcal{H}_{\tau},$$

$$(15)$$

for a strictly positive weight function $w \colon \mathbb{N}_0 \to \mathbb{R}_{>0}$. Here, \vee denotes the symmetric tensor product, defined as the average

$$f_1 \vee \cdots \vee f_n = \frac{1}{n!} \sum_{\sigma \in S_n} f_{\sigma(1)} \otimes \cdots \otimes f_{\sigma(n)}, \quad f_i \in \mathcal{H}_{\tau},$$

over the *n*-element permutation group S_n , and $\mathcal{H}_{\tau}^{\vee n}$ is the closed subspace of $\mathcal{H}_{\tau}^{\otimes n}$ consisting of symmetric tensors. The map $f_1 \otimes \cdots \otimes f_n \mapsto f_1 \vee \cdots \vee f_n$ defines, by linear extension, the orthogonal projection from $\mathcal{H}_{\tau}^{\otimes n}$ to $\mathcal{H}_{\tau}^{\vee n}$. We will use $\Omega \equiv 1 \in \mathbb{C} \subset F_w(\mathcal{H}_{\tau})$ to denote the "vacuum" vector of the Fock space. By convention, we will always choose w such that w(0) = 1.

In Theorem 8, we show that if w^{-2} is summable and subconvolutive,

$$w^{-2} \in \ell^1(\mathbb{N}_0), \quad w^{-2} * w^{-2}(n) \le Cw^{-2}(n),$$
 (16)

 $F_w(\mathcal{H}_{\tau})$ becomes a unital Banach algebra with respect to the symmetric tensor product for a norm $\|\cdot\|_{F_w(\mathcal{H}_{\tau})}$ equivalent to the Hilbert space norm,

$$|||f \vee g||_{F_w(\mathcal{H}_\tau)} \le |||f||_{F_w(\mathcal{H}_\tau)} |||g||_{F_w(\mathcal{H}_\tau)}, \quad \forall f, g \in F_w(\mathcal{H}_\tau),$$

and with Ω as the unit. Moreover, associated with $F_w(\mathcal{H}_\tau)$ is a coproduct, i.e., a bounded operator $\Delta \colon F_w(\mathcal{H}_\tau) \to F_w(\mathcal{H}_\tau) \otimes F_w(\mathcal{H}_\tau)$ such that

$$\Delta^*(f \otimes g) = f \vee g.$$

Boundedness of Δ allows us to characterize the spectrum of $F_w(\mathcal{H}_\tau)$ (i.e., the set of nonzero multiplicative functionals $\chi \colon F_w(\mathcal{H}_\tau) \to \mathbb{C}$) as the set

$$\sigma(F_w(\mathcal{H}_\tau)) = \left\{ \chi = \langle \xi, \cdot \rangle_{F_w(\mathcal{H}_\tau)} : \xi = \sum_{n=1}^\infty w^{-2}(n) \eta^{\vee n} : \eta \in (\mathcal{H}_\tau)_{R_w} \right\} \subset F_w(\mathcal{H}_\tau)^*,$$

where R_w is the radius of convergence of the series $\sum_{n=1}^{\infty} w^{-2}(n)z^n$, $z \in \mathbb{C}$; see Proposition 7. Since $w \in \ell^2(\mathbb{N})$, we have $R_w \geq 1$ and the set of admissible vectors η in the definition above includes the unit ball of \mathcal{H}_{τ} .

The dual object to $\sigma(F_w(\mathcal{H}_\tau))$ is the cospectrum of $F_w(\mathcal{H}_\tau)$, which is defined as

$$\sigma_{\rm co}(F_w(\mathcal{H}_\tau)) = \left\{ \xi \in F_w(\mathcal{H}_\tau) : \langle \xi, \cdot \rangle_{F_w(\mathcal{H}_\tau)} \in \sigma(F_w(\mathcal{H}_\tau)) \right\} \subset F_w(\mathcal{H}_\tau).$$

Equivalently, we have that $\sigma_{co}(F_w(\mathcal{H}_{\tau}))$ is the subset of $F_w(\mathcal{H}_{\tau})$ consisting of elements ξ such that $\Delta \xi = \xi \otimes \xi$. We equip $\sigma(F_w(\mathcal{H}_{\tau}))$ and $\sigma_{co}(F_w(\mathcal{H}_{\tau}))$ with the weak-* topology on $F_w(\mathcal{H}_{\tau})$ * and the weak topology on $F_w(\mathcal{H}_{\tau})$, respectively. With these topologies, they become compact Hausdorff spaces.

The weighted Fock space $F_w(\mathcal{H}_\tau)$ has an associated feature map $\varphi_\tau \colon X \to F_w(\mathcal{H}_\tau)$, where

$$\varphi_{\tau}(x) = \sum_{n=1}^{\infty} \frac{w^{-2}(n)}{\varpi_{\tau}^{2n}} k_{\tau,x}^{\vee n}, \quad \varpi_{\tau} = \sup_{x \in X} \sqrt{k_{\tau}(x,x)}.$$

$$\tag{17}$$

The range of φ_{τ} lies in the cospectrum $\sigma_{co}(F_w(\mathcal{H}_{\tau}))$, which implies that $\hat{\varphi}_{\tau} \colon X \to \sigma(F_w(\mathcal{H}_{\tau}))$ with

$$\hat{\varphi}_{\tau}(x) = \langle \varphi_{\tau}(x), \cdot \rangle_{F_{w}(\mathcal{H}_{\tau})}$$
(18)

is a well-defined map of state space into the spectrum of the weighted Fock space. The choice of the RKHS family \mathcal{H}_{τ} in our examples will ensure that this map is injective.

Among many possible constructions, in this paper we use weights from the subexponential family

$$w(n) = e^{\sigma n^p}, \quad \sigma > 0, \quad p \in (0, 1),$$

which is a prototypical class of subconvolutive weights used in harmonic analysis on locally compact abelian groups (e.g., [29,41]).

Next, using the convention $f^{\vee 0} = \Omega$, the $\{\zeta_{j,\tau}\}_{j\in\mathbb{Z}}$ eigenbasis of \mathcal{H}_{τ} from (8) induces an orthonormal basis of $F_w(\mathcal{H}_{\tau})$ consisting of elements

$$\zeta_{\tau}^{\vee J} = \frac{\zeta_{0,\tau}^{\vee j_0} \vee \zeta_{-1,\tau}^{\vee j_{-1}} \vee \zeta_{1,\tau}^{\vee j_1} \vee \zeta_{-2,\tau}^{\vee j_{-2}} \vee \zeta_{2,\tau}^{\vee j_2} \vee \cdots}{w(\|J\|)\sqrt{j_0! \, j_{-1}! \, j_1! \, j_{-2}! \, j_{2!} \cdots}}.$$
(19)

Here, $J = (j_0, j_{-1}, j_1, j_{-2}, j_2, ...)$ is a multi-index of non-negative integers j_i , finitely many of which are nonzero, and $||J|| = \sum_{j_i \in j} j_i$. Henceforth, we will use the symbol \mathbb{J} to denote the set of such multi-indices.

In second-quantization terminology, the basis $\{\zeta_{\tau}^{\vee J}\}_{J\in\mathbb{J}}$ is referred to as an occupation number basis. Intuitively, we think of each basis vector $\zeta_{i,\tau}$ as being associated with a "particle" of a certain type. A density operator $\rho = \langle \zeta_{\tau}^{\vee J}, \cdot \rangle_{F_w(\mathcal{H}_{\tau})} \zeta_{\tau}^J \in Q(F_w(\mathcal{H}_{\tau}))$ then induces a pure quantum state (see section 3.1) comprising of j_0 particles of type $\zeta_{\tau,0}$, j_{-1} particles of type $\zeta_{\tau,-1}$, and so on, for a total number of $\|J\|$ particles.

4.2. Reproducing kernel Hilbert algebras. The weighted Fock space $F_w(\mathcal{H}_{\tau})$ has two associated unital RKHAs: One, denoted as $\hat{F}_w(\mathcal{H}_{\tau})$, is a space of continuous functions on the spectrum $\sigma(F_w(\mathcal{H}_{\tau}))$, and another one, denoted as $\tilde{\mathcal{H}}_{\tau}$, is a space of continuous functions on the state space X

The space $\hat{F}_w(\mathcal{H}_\tau)$ is built using the Gelfand transform $\Gamma \colon F_w(\mathcal{H}_\tau) \to C(\sigma(F_w(\mathcal{H}_\tau)))$, defined as $(\Gamma f)\chi = \chi f$. We show, by Corollary 9 below, that the image $\hat{F}_w(\mathcal{H}_\tau) \subseteq C(\sigma(F_w(\mathcal{H}_\tau)))$ of $F_w(\mathcal{H}_\tau)$ under Γ has RKHA structure for the reproducing kernel $\hat{k}_\tau \colon \sigma(F_w(\mathcal{H}_\tau)) \times \sigma(F_w(\mathcal{H}_\tau)) \to \mathbb{C}$,

$$\hat{k}_{\tau}(\chi_1, \chi_2) = \langle \xi_1, \xi_2 \rangle_{F_w(\mathcal{H}_{\tau})}, \quad \chi_i = \langle \xi_i, \cdot \rangle_{F_w(\mathcal{H}_{\tau})},$$

and the Gelfand map Γ acts as a Banach algebra isomorphism. Using the Gelfand map, we obtain an orthonormal basis $\{\Gamma\zeta_{\tau}^{\vee J}\}_{J\in\mathbb{J}}$ of $\hat{F}_w(\mathcal{H}_{\tau})$ from the basis elements $\zeta_{\tau}^{\vee J}$ in (19).

Next, \mathcal{H}_{τ} is given by the pullback of $\hat{F}_w(\mathcal{H}_{\tau})$ onto X under the feature map, $\mathcal{H}_{\tau} = \{f \circ \hat{\varphi}_{\tau}\} \subset C(X)$. One then readily verifies that \mathcal{H}_{τ} has the reproducing kernel $k_{\tau} \colon X \times X \to \mathbb{R}_{>0}$, where

$$\tilde{k}_{\tau}(x,y) = \sum_{n=0}^{\infty} \frac{w^{-2}(n)}{\varpi_{\tau}^{2n}} k_{\tau}(x,y)^{n}.$$

Moreover, the maps $\hat{\pi}: \hat{F}_w(\mathcal{H}_{\tau}) \to \tilde{\mathcal{H}}_{\tau}$ and $\tilde{\pi}: F_w(\mathcal{H}_{\tau}) \to \tilde{\mathcal{H}}_{\tau}$ defined by $\hat{\pi}f = f \circ \hat{\varphi}_{\tau}$ and $\tilde{\pi} = \hat{\pi} \circ \Gamma$ are algebra homomorphisms,

$$\hat{\pi}(fg) = (\hat{\pi}f)(\hat{\pi}g), \quad \tilde{\pi}(f \vee g) = (\tilde{\pi}f)(\tilde{\pi}g).$$

Similarly to \mathcal{H}_{τ} , the RKHAs $\tilde{\mathcal{H}}_{\tau}$ have associated integral operators $\tilde{K}_{\tau} \colon H \to \tilde{\mathcal{H}}_{\tau}$, where

$$\tilde{K}_{\tau}f = \int_{X} \tilde{k}_{\tau}(\cdot, x) f(x) d\mu(x),$$

and \tilde{K}_{τ}^* implements the inclusion map from $\tilde{\mathcal{H}}_{\tau}$ into H.

4.3. **Spectral tori.** Let \mathbb{A}_w be the subset of $\ell^2(\mathbb{N}_0)$ consisting of vectors $a = (a_j)_j$ with norm $\|a\|_{\ell^2(\mathbb{N})} \leq R_w$ and non-negative elements a_j . The structure of the weighted Fock spaces and RKHAs introduced above can be further characterized by defining, for each $a \in \mathbb{A}_w$ and each sequence $z = (z_j)_j \in \ell^{\infty}(\mathbb{N})$ with unimodular elements z_j , the vectors

$$\xi_{\tau,a,z} = \sum_{n=0}^{\infty} w^{-2}(n) \left(a_0 + \sum_{j=1}^{\infty} a_j \frac{z_j \zeta_{j,\tau} + \overline{z_j \zeta_{j,\tau}}}{\sqrt{2}} \right)^{\vee n} \in \sigma_{\text{co}}(F_w(\mathcal{H}_\tau)), \tag{20}$$

and the subsets $\mathbb{T}_{\tau,a}$ of the spectrum $\sigma(F_w(\mathcal{H}_{\tau}))$ as

$$\mathbb{T}_{\tau,a} = \left\{ \chi_{\tau,a,z} \equiv \langle \xi_{\tau,a,z}, \cdot \rangle_{F_w(\mathcal{H}_\tau)} : z = (z_j)_j \in \ell^\infty(\mathbb{N}), \ z_j \in \mathbb{T}^1 \subset \mathbb{C} \right\}.$$

Each set $\mathbb{T}_{\tau,a}$ has the topology of a torus of dimension equal to the number of nonzero elements of (a_1, a_2, \ldots) . In what follows, $\mathbb{S}_{\tau} = \bigcup_{a \in \mathbb{A}_w} \mathbb{T}_{\tau,a} \subset \sigma(F_w(\mathcal{H}_{\tau}))$ will be the (disjoint) union of these tori.

For $x \in X$, let $a_{\tau}(x) = (a_{0,\tau}(x), a_{1,\tau}(x), \ldots) \in \mathbb{A}_w$ and $z_{\tau}(x) = (z_{\tau,1}(x), z_{\tau,2}(x), \ldots) \in \ell^{\infty}(\mathbb{N})$, where $a_{0,\tau}(x) = 1/\varpi_{\tau}$, and

$$a_{j,\tau}(x) = \frac{\sqrt{2}}{\varpi_{\tau}} |\zeta_{j,\tau}(x)|, \quad z_{j,\tau}(x) = \frac{\overline{\zeta_{j,\tau}(x)}}{|\zeta_{j,\tau}(x)|}, \quad j \in \mathbb{N}.$$

Observe that

$$k_{\tau,x} = a_{0,\tau}(x) + \sum_{j=1}^{\infty} a_{j,\tau}(x) \frac{z_{j,\tau}(x)\zeta_{j,\tau} + \overline{z_{j,\tau}(x)\zeta_{j,\tau}}}{\sqrt{2}}$$
(21)

and $||a_{\tau}(x)||_{\ell^{2}(\mathbb{N}_{0})} = ||k_{\tau,x}||_{\mathcal{H}_{\tau}}/\varpi_{\tau} \leq 1$. Comparing the above expression with (20), it follows that the feature vector $\varphi_{\tau}(x)$ is equal to $\xi_{\tau,a(x),z_{\tau}(x)}$, and thus that $\hat{\varphi}_{\tau}(x)$ is a point in the torus $\mathbb{T}_{\tau,a_{\tau}(x)}$

with coordinates $z_{\tau}(x)$. Henceforth, we will use the abbreviated notation $\mathbb{T}_{\tau,x} \equiv \mathbb{T}_{\tau,a_{\tau}(x)}$ for $x \in X$. Letting $\hat{X}_{\tau} = \hat{\varphi}_{\tau}(X) \subseteq \sigma(F_w(\mathcal{H}_{\tau}))$ be the image of the state space in the Fock space spectrum under the feature map $\hat{\varphi}_{\tau}$, we have that \hat{X}_{τ} is a subset of \mathbb{S}_{τ} .

By injectivity of the feature map, $\hat{\varphi}_{\tau}(x) \mapsto \hat{\varphi}_{\tau}(\Phi^{t}(x))$ defines a flow $\hat{\Phi}_{\tau}^{t} : \hat{X}_{\tau} \to \hat{X}_{\tau}$. This flow is topologically conjugate to Φ^{t} in the sense of the following commuting diagram involving continuous maps:

$$\begin{array}{ccc}
X & \xrightarrow{\Phi^t} X \\
\hat{\varphi}_{\tau} \downarrow & & \downarrow \hat{\varphi}_{\tau} \\
\hat{X}_{\tau} & \xrightarrow{\hat{\Phi}_{\tau}^t} \hat{X}_{\tau}
\end{array}$$

4.4. Fock space amplification. For $\tau > 0$ we define the strongly continuous, one-parameter evolution group $\{\tilde{U}_{\tau}^t \colon F_w(\mathcal{H}_{\tau}) \to F_w(\mathcal{H}_{\tau})\}_{t \in \mathbb{R}}$ by linear extension of

$$\tilde{U}_{\tau}^{t}(f_1 \vee \cdots \vee f_n) = (U_{\tau}^{t}f_1) \vee \cdots \vee (U_{\tau}^{t}f_n), \quad f_i \in \mathcal{H}_{\tau}.$$

The generator of this group is a skew-adjoint operator $\tilde{W}_{\tau} \colon D(\tilde{W}_{\tau}) \to F_w(\mathcal{H}_{\tau})$ satisfying

$$\tilde{W}_{\tau}(f_1 \vee \cdots \vee f_n) = (W_{\tau}f_1) \vee f_2 \cdots \vee f_n + \ldots + f_1 \vee \cdots \vee f_{n-1} \vee (W_{\tau}f_n)$$

for $f_1, \ldots, f_n \in D(W_\tau)$. Thus, \tilde{U}_τ^t acts multiplicatively on $F_w(\mathcal{H}_\tau)$, and \tilde{W}_τ satisfies the Leibniz rule on a suitable subspace. By virtue of the Leibniz rule, \tilde{W}_τ admits the eigendecomposition

$$\tilde{W}_{\tau}\zeta_{\tau}^{\vee J} = i\tilde{\omega}_{J,\tau}\zeta_{\tau}^{\vee J}, \quad \tilde{\omega}_{J,\tau} = \sum_{i=-\infty}^{\infty} j_i\omega_{i,\tau}, \quad J = (j_0, j_{-1}, j_1, j_{-2}, j_2, \ldots) \in \mathbb{J},$$

and the point spectrum $\sigma_p(\tilde{W}_{\tau})$ has the structure of a subgroup of $i\mathbb{R}$. Thus, lifting U_{τ}^t into the Fock space recovers key structural properties of one-parameter unitary Koopman groups that were lost through regularization.

4.5. Topological models of regularized Koopman evolution. The unitary operators U_{τ}^t induce a flow R_{τ}^t : $\sigma(F_w(\mathcal{H}_{\tau})) \to \sigma(F_w(\mathcal{H}_{\tau}))$, $t \in \mathbb{R}$, where $R_{\tau}^t(\chi) = \chi \circ \tilde{U}_{\tau}^t$. Each torus $\mathbb{T}_{\tau,a}$ is an invariant set under R_{τ}^t . On these sets, R_{τ}^t takes the form of a rotation system generated by the eigenfrequencies $\omega_{j,\tau}$:

$$R_{\tau}^t(\chi_{\tau,a,z}) = \chi_{\tau,a,z^{(t)}},$$

where $z = (z_j)_{j \in \mathbb{N}}$ and $z^{(t)} = (e^{-i\omega_{j,t}}z_j)_{j \in \mathbb{N}}$. Equivalently, we have

$$R_{\tau}^{t}(\langle \xi_{\tau,a,z}, \cdot \rangle_{F_{w}(\mathcal{H}_{\tau})}) = \langle \tilde{U}_{\tau}^{-t} \xi_{\tau,a,z}, \cdot \rangle_{F_{w}(\mathcal{H}_{\tau})},$$

so the vector $\xi_{\tau,a,z}$ evolves under the adjoint ("Perron–Frobenius") operators $\tilde{U}_{\tau}^{t*} = \tilde{U}_{\tau}^{-t}$.

The rotation system R_{τ}^t constitutes a topological model of the regularized Koopman dynamics U_{τ}^t as a rotation system on the spectrum of the weighted Fock space $F_w(\mathcal{H}_{\tau})$. This is non-trivial since U_{τ}^t is not a composition operator induced by a flow on the original state space X. Since \hat{X}_{τ} is a subset of $\mathbb{S}_{\tau} \subset \sigma(F_w(\mathcal{H}_{\tau}))$ (and \mathbb{S}_{τ} is invariant under R_{τ}^t), the union of tori \mathbb{S}_{τ} provides a common topological setting for studying the dynamical system associated with the regularized Koopman operators U_{τ}^t (represented by R_{τ}^t) in relation to the original dynamical system Φ^t (represented by $\hat{\Phi}_{\tau}^t$).

Letting \hat{U}_{τ}^t : $C(\sigma(F_w(\mathcal{H}_{\tau}))) \to C(\sigma(F_w(\mathcal{H}_{\tau})))$, $\hat{U}_{\tau}^t f = f \circ R_{\tau}^t$, be the Koopman operator on continuous functions induced by R_{τ}^t , we have

$$\Gamma \circ \tilde{U}_{\tau}^t = \hat{U}_{\tau}^t \circ \Gamma.$$

As a result, $\hat{F}_w(\mathcal{H}_\tau)$ is a \hat{U}_τ^t -invariant subspace of $C(\sigma(F_w(\mathcal{H}_\tau)))$.

Our approach is to employ the rotation system R_{τ}^t in conjunction with an embedding of observables in H into the weighted Fock space $F_w(\mathcal{H}_{\tau})$ and associated RKHA $\hat{F}_w(\mathcal{H}_{\tau})$ to build approximations of the unitary Koopman evolution of observables under U^t . Since the RKHS $\tilde{\mathcal{H}}_{\tau}$ spans a dense subspace of H and for a general observable $f \in \mathcal{H}_{\tau} \subset \tilde{\mathcal{H}}_{\tau}$ there exist multiple elements $\tilde{f} \in F_w(\mathcal{H}_{\tau})$ of the Fock space lying above it (i.e., $\tilde{\pi}\tilde{f} = f$) there are different possible strategies for representing observables in H by elements of $F_w(\mathcal{H}_{\tau})$ or, equivalently, $\hat{F}_w(\mathcal{H}_{\tau})$.

One such strategy is to employ a continuous lifting map $\mathcal{L} \colon \mathcal{H}_{\tau} \to F_w(\mathcal{H}_{\tau})$ satisfying

$$\tilde{\pi} \circ \mathcal{L}f = f. \tag{22}$$

Defining $\hat{\mathcal{L}} = \Gamma \circ \mathcal{L}$ then yields

$$f_{\tau}^{(t)} = \hat{\pi} \circ \hat{U}_{\tau}^{t} \circ \hat{\mathcal{L}}$$

as an approximation of the Koopman evolution $U^t f$ based on the rotation system on $\sigma(F_w(\mathcal{H}_\tau))$. A basic choice in that direction is to choose $\mathcal{L} \colon \mathcal{H}_\tau \hookrightarrow F_w(\mathcal{H}_\tau)$ as the inclusion map. In section 6.4 we consider aspects of "nonlinear" approximations wherein \mathcal{L} yields a representation of f as a product–sum of functions, thus utilizing higher gradings of the Fock space.

Yet another approach is to relax the requirement that \mathcal{L} yields an exact representation, and consider embeddings that satisfy (22) approximately while still utilizing higher gradings of the Fock space and converging to the true Koopman evolution in an appropriate asymptotic limit. The construction and analysis of such an approximation will be the focus of sections 4.6 and 4.7 below.

4.6. Fock space embedding of observables. We use integral operators to represent $f \in H$ by elements of the RKHA $\hat{F}_w(\mathcal{H}_\tau)$, whose restrictions on finite-dimensional tori $\mathbb{T}_{\tau,a} \subset \mathbb{S}_\tau$ are polynomials of arbitrarily large degree $m \in \mathbb{N}$ of the coordinates z_j .

Let $\kappa: X \times X \to \mathbb{R}_{>0}$ be a strictly positive kernel such that for every $\tau > 0$ and $x \in X$, $\kappa(x, \cdot)$ lies in \mathcal{H}_{τ} . For $m \in \mathbb{N}$ define $\kappa_m: X \times X \to \mathbb{R}_{>0}$, $d_m: X \to \mathbb{R}_{>0}$, and $p_m: X \times X \to \mathbb{R}_{>0}$ such that

$$\kappa_m(x,y) = \kappa(x,y)^m, \quad d_m = \int_X \kappa_m(\cdot,x) \, d\mu(x), \quad p_m(x,y) = \frac{\kappa_m(x,y)}{d_m(x)}.$$

Note that $\kappa_m(x,\cdot)$ lies in the RKHA $\tilde{\mathcal{H}}_{\tau}$ for every $x \in X$ and $\kappa_{\tau}^{\vee m}(x,\cdot) \in F_w(\mathcal{H}_{\tau})$ lies above $\kappa_m(x,\cdot)$,

$$\tilde{\pi}(\kappa^{\vee m}(x,\cdot)) = \kappa_m(x,\cdot).$$

Similarly, d_m lies in $\tilde{\mathcal{H}}_{\tau}$, and because it is bounded away from zero on the compact set M, the inverse $1/(d_m|_M)$ lies in $\tilde{\mathcal{H}}_{\tau}(M)$ by the holomorphic functional calculus on $\tilde{\mathcal{H}}_{\tau}(M)$. We can then conclude that $p_m(x,\cdot)|_M$ lies in $\tilde{\mathcal{H}}_{\tau}(M)$ for every $x \in M$.

On the basis of the above, $\mathcal{K}_m \colon H \to F_w(\mathcal{H}_\tau)$, $\tilde{\mathcal{K}}_m \colon H \to C(M)$, and $\tilde{P}_m \colon H \to C(M)$, where

$$\mathcal{K}_m f = \int_X \kappa^{\vee m}(\cdot, x) f(x) \, d\mu(x), \quad \tilde{\mathcal{K}}_m f = \int_X \kappa_m(\cdot, x) f(x) \, d\mu(x),$$
$$\tilde{P}_m f = \int_X p_m(\cdot, x) f(x) \, d\mu(x)$$

are well-defined integral operators. We also define $\hat{\mathcal{K}}_m \colon H \to \hat{F}_w(\mathcal{H}_\tau)$ and $P_m \colon H \to H$ as

$$\hat{\mathcal{K}}_m = \Gamma \tilde{\mathcal{K}}_m, \quad P_m = \iota \tilde{P}_m.$$

Note that P_m is a Markov operator on H; that is, (i) it preserves constant functions, $P_m 1_X = 1_X$; and (ii) it preserves positive functions, $P_m f \ge 0$ whenever $f \ge 0$. We will assume that the kernel κ is chosen such that the Markov operators P_m have the following approximation property for continuous functions,

$$\lim_{m \to \infty} ||P_m \iota f - \iota f||_H, \quad \forall f \in C(M).$$
 (23)

Given a metric $d: X \times X \to \mathbb{R}$ that metrizes the topology of X, examples of kernels satisfying (23) are radially decaying kernels such as

$$\kappa(x,y) = \exp\left(-\frac{d^2(x,y)}{\varepsilon^2}\right), \quad \varepsilon > 0.$$
(24)

With these constructions, we embed observables $f \in C(M)$ to elements of the RKHA $\hat{F}_w(\mathcal{H}_{\tau})$ by means of the maps $\hat{\mathcal{K}}_m$; specifically, $\hat{g}_m := \hat{\mathcal{K}}_m \iota f$. Defining, in addition,

$$\hat{h}_m = \hat{\mathcal{K}}_m 1_X, \quad \hat{f}_m = \frac{\hat{g}_m}{\hat{h}_m},$$

and

$$\hat{f}_{m,\tau}^{(t)} = \hat{U}_{\tau}^{t} \hat{f}_{m} \equiv \frac{\hat{g}_{m,\tau}^{(t)}}{\hat{h}_{m,\tau}^{(t)}}, \quad \hat{g}_{m,\tau}^{(t)} = \hat{U}_{\tau}^{t} \hat{g}_{m}, \quad \hat{h}_{m,\tau}^{(t)} = \hat{U}_{\tau}^{t} \hat{h}_{m}, \tag{25}$$

our approximation of the Koopman evolution $U^t f$ is given by the RKHA element

$$f_{m,\tau}^{(t)} = \hat{\pi}_{\tau} \hat{f}_{m,\tau}^{(t)} \equiv \hat{f}_{m,\tau}^{(t)} \circ \hat{\varphi}_{\tau} \in \tilde{\mathcal{H}}_{\tau}. \tag{26}$$

In section 6.1, we will show:

Proposition 5. With notation as above and under assumption (23), $f_{m,\tau}^{(t)}$ converges to $f^{(t)}$ in the norm of H in the iterated limit of $m \to \infty$ after $\tau \to 0^+$,

$$\lim_{m \to \infty} \lim_{\tau \to 0^+} \left\| \iota f_{m,\tau}^{(t)} - U^t f \right\|_H = 0.$$

4.7. **Finite-dimensional approximation.** To arrive at a practical approximation algorithm, we approximate $f_{m,\tau}^{(t)}(x) = \hat{f}_{m,\tau}^{(t)}(\hat{\phi}_{\tau}(x))$ by evaluating $\hat{f}_{m,\tau}^{(t)}$ on points in finite-dimensional tori in \mathbb{S}_{τ} lying near $\hat{\varphi}_{\tau}(x)$.

To that end, for a parameter $d \in \mathbb{N}$, and point $x \in X$, define $a_{\tau}^{(d)}(x) = (a_{0,\tau}(x), \dots a_{d,\tau}(x), 0, \dots) \in \mathbb{A}_w$,

$$\eta_{\tau,d,x} = a_{\tau,0}(x) + \sum_{j=1}^{d} a_{\tau,j}(x) \frac{z_{j,\tau}(x)\zeta_{j,\tau} + \overline{z_{j,\tau}(x)\zeta_{j,\tau}}}{\sqrt{2}},$$
(27)

and the truncated feature maps $\varphi_{\tau,d} \colon X \to F_w(\mathcal{H}_\tau)$ and $\hat{\varphi}_{\tau,d} \colon X \to \sigma(F_w(\mathcal{H}_\tau))$ as

$$\varphi_{\tau,d}(x) = \sum_{n=1}^{\infty} w^{-2}(n) \eta_{\tau,d,x}^{\vee n}, \quad \hat{\varphi}_{\tau,d}(x) = \langle \varphi_{\tau,d}(x), \cdot \rangle_{F_w(\mathcal{H}_{\tau})}.$$

Note that the image of X under $\hat{\varphi}_{\tau,d}$ lies in the finite-dimensional torus $\mathbb{T}_{\tau,d,x} \equiv T_{\tau,a^{(d)}(x)}$. We approximate the prediction function $f_{m,\tau}^{(t)}$ from (26) by pulling back $\hat{f}_{m,\tau}^{(t)}$ from this torus to the state space X, viz.

$$f_{m,\tau,d}^{(t)} = \hat{f}_{m,\tau}^{(t)} \circ \hat{\varphi}_{\tau,d} \in \tilde{\mathcal{H}}_{\tau}.$$

As $d \to \infty$ at fixed τ and m, $f_{m,\tau,d}^{(t)}$ converges to $f_{m,\tau}^{(t)}$ pointwise on X and in the norm of H,

$$\lim_{d \to \infty} \|\iota(f_{m,\tau,d}^{(t)} - f_{m,\tau}^{(t)})\|_{H} = 0; \tag{28}$$

see section 6.3 for further details. Combining (28) with Proposition 5, we deduce that the finite-dimensional Fock space approximation $f_{m,\tau,d}^{(t)}$ converges to the true Koopman evolution $U^t f$ in L^2 sense in the iterated limit of $m \to \infty$ after $\tau \to 0^+$ after $d \to \infty$.

For simplicity of exposition, we will henceforth assume that the coefficients $a_{1,\tau}(x), \ldots, a_{d,\tau}(x)$ are all nonzero at the chosen $x \in X$, so that $\mathbb{T}_{\tau,d,x}$ is d-dimensional. We then have that the

restrictions of the functions $\hat{g}_{m,\tau}^{(t)}$ and $\hat{h}_{m,\tau}^{(t)}$ in (25) on $\mathbb{T}_{\tau,d,x}$ are polynomials of degree m in the variables $z_1,\ldots,z_d,\overline{z_1},\ldots,\overline{z_d}$.

To build these polynomials, define for $i \in \{-d, \dots, d\}$ the functions

$$\varrho_{j,\tau} = \int_X \kappa(\cdot, x) \zeta_{j,\tau}(x) \, d\mu(x) \in \mathcal{H}_\tau. \tag{29}$$

Define also the index set $\mathbb{J}_{d,m} = \{j = (j_{-d}, \dots, j_d) \in \mathbb{N}_0^{2d+1} : |j| = m\}$, and for each $j \in \mathbb{J}_{d,m}$, the moment coefficients

$$C_{j,\tau,d}^{(g)} = \int_{X} f(x) \prod_{i=-d}^{d} \varrho_{i,\tau}^{j_{i}}(x) \, d\mu(x), \quad C_{j,\tau,d}^{(h)} = \int_{X} \prod_{i=-d}^{d} \varrho_{i,\tau}^{j_{i}}(x) \, d\mu(x), \tag{30}$$

and the amplitude functions

$$\alpha_{j,\tau,d}(x) = \prod_{i=-d}^{d} a_{i,\tau}(x).$$

We then have that for every $\chi_{\tau,a_{\tau}^{(d)}(x),z} \in \mathbb{T}_{\tau,d,x}$,

$$\hat{g}_{m,\tau}^{(t)}(\chi_{\tau,a_{\tau}^{(d)}(x),z}) = w^{-2}(m) \sum_{j \in \mathbb{J}_{d,m}} \binom{m}{j_{-d}\cdots j_{d}} \alpha_{j,\tau,d}(x) C_{j,\tau,d}^{(g)} \prod_{r=1}^{d} e^{-i(j_{r}-j_{-r})\omega_{r}t} z_{r}^{j_{r}-j_{-r}},$$

$$\hat{h}_{m,\tau}^{(t)}(\chi_{\tau,a_{\tau}^{(d)}(x),z}) = w^{-2}(m) \sum_{j \in \mathbb{J}_{d,m}} \binom{m}{j_{-d}\cdots j_{d}} \alpha_{j,\tau,d}(x) C_{j,\tau,d}^{(h)} \prod_{r=1}^{d} e^{-i(j_{r}-j_{-r})\omega_{r}t} z_{r}^{j_{r}-j_{-r}}.$$
(31)

Thus, on $\mathbb{T}_{\tau,d,x}$, $\hat{g}_{m,\tau}^{(t)}$ and $\hat{h}_{m,\tau}^{(t)}$ reduce to degree-m polynomials in the coordinates z_r and their conjugates. Correspondingly, the prediction function $\hat{f}_{m,\tau}^{(t)}$ is a rational function of the z_r .

Pulling back to X via the feature map $\hat{\varphi}_{\tau,d}$, we arrive at the approximation

$$f_{m,\tau,d}^{(t)} = \frac{\hat{g}_{m,\tau}^{(t)} \circ \hat{\varphi}_{\tau,d}}{\hat{h}_{m,\tau}^{(t)} \circ \hat{\varphi}_{\tau,d}} = \frac{\sum_{j \in \mathbb{J}_{d,m}} \binom{m}{j-d\cdots j_d} C_{j,\tau,d}^{(g)} \prod_{r=1}^d e^{i(j_{-r}-j_r)\omega_{r,\tau}t} \zeta_{r,\tau}^{j-r-j_r}}{\sum_{j \in \mathbb{J}_{d,m}} \binom{m}{j-d\cdots j_d} C_{j,\tau,d}^{(h)} \prod_{r=1}^d e^{i(j_{-r}-j_r)\omega_{r,\tau}t} \zeta_{r,\tau}^{j-r-j_r}}.$$
(32)

A key aspect of this approximation, which is evident from both (31) and (32), is that it captures spectral information from the entire set of eigenfrequencies $\{\tilde{\omega}_{j,\tau}\}_{j\in\mathbb{J}_{d,m}}$, containing $\binom{m+2d}{2d}$ elements. Thus, similarly to the tensor network scheme outlined in section 3.4, the second-quantization approach presented in this paper provides a route for algebraic amplification of (2d+1)-dimensional approximation spaces $\mathcal{Z}_{\tau,d} = \{\zeta_{-d,\tau},\ldots,\zeta_{d,\tau}\} \in \mathcal{H}_{\tau}$. As noted in the beginning of this section, an advantageous aspect of the second-quantization approach is that it does not require \mathcal{H}_{τ} to have RKHA structure. In addition, it provides a framework for building topological models of regularized Koopman evolution as rotation systems on tori.

5. Weighted Fock space

In this section, we discuss in more detail the structure of the weighted Fock spaces $F_w(\mathcal{H}_{\tau})$ as commutative Banach algebras (section 5.1), and establish the decomposition of their spectra in terms of the tori $\mathbb{T}_{\tau,a}$ (section 5.2).

5.1. Banach algebra structure. We will take an indirect approach to proving that $F_w(\mathcal{H}_{\tau})$ is a commutative Banach algebra that is isomorphic to an RKHA of continuous functions on its spectrum. We will pass through the full weighted Fock space, $F_w^{\otimes}(\mathcal{H}_{\tau})$, as a noncommutative Banach algebra with a comultiplication map and pass to the symmetric Fock space by taking a quotient.

In what follows, given a Hilbert space \mathcal{H} (not necessarily an RKHS), and a weight $w \colon \mathbb{N}_0 \to \mathbb{R}_{>0}$ with w(0) = 1, $F_w^{\otimes}(\mathcal{H})$ will be the weighted Fock space defined as the closure of the tensor algebra $T(\mathcal{H}) = \mathbb{C} \oplus \mathcal{H} \oplus \mathcal{H}^{\otimes 2} \oplus \ldots$ with respect to the inner product (cf. (15))

$$\langle a, b \rangle_{F_w^{\otimes}(\mathcal{H})} = \bar{a}b, \quad a, b \in \mathbb{C},$$

$$\langle a, f \rangle_{F_w^{\otimes}(\mathcal{H})} = 0, \quad a \in \mathbb{C}, \quad f \in \mathcal{H},$$

$$\langle f_1 \otimes \cdots \otimes f_n, g_1 \otimes \cdots \otimes g_n \rangle_{F_w^{\otimes}(\mathcal{H})} = w^2(n) \prod_{i=1}^n \langle f_i, g_i \rangle_{\mathcal{H}_{\tau}}, \quad f_i, g_i \in \mathcal{H}.$$

Moreover, we will use \boxtimes to denote the usual tensor product between two Hilbert spaces in order to distinguish it from the tensor product \otimes within $F_w^{\otimes}(\mathcal{H})$. We define the symmetric Fock space $F_w(\mathcal{H})$ similarly to $F_w(\mathcal{H}_{\tau})$ from section 4.1 as the closed subspace of $F_w^{\otimes}(\mathcal{H})$ spanned by symmetric tensors. We let Sym: $F_w^{\otimes}(\mathcal{H}) \to F_w(\mathcal{H})$ be the orthogonal projection onto that subspace.

Proposition 6. With notation as above, let $w: \mathbb{N}_0 \to \mathbb{R}_{>0}$ be inverse square-summable, $w^{-1} \in \ell^2(\mathbb{N})$. Then, there exists a unique bounded operator $\Delta: F_w^{\otimes}(\mathcal{H}) \to F_w^{\otimes}(\mathcal{H}) \boxtimes F_w^{\otimes}(\mathcal{H})$ such that $\Delta^*(f \boxtimes g) = f \otimes g$ iff w^{-2} is subconvolutive, $w^{-2} * w^{-2} \leq Cw^{-2}$.

Proof. Fix an orthonormal basis $\{\phi_i\}_{i=0}^{\infty} \subset \mathcal{H}$, and write $\phi^{\otimes J} = \phi_{j_1} \otimes \cdots \otimes \phi_{j_m}$ for $j \in \mathbb{J}^{\otimes} := \bigcup_{m=0}^{\infty} \mathbb{N}_0^m$ with the convention that $\mathbb{N}_0^0 = \{0\}$ and $f^{\otimes 0} = \Omega$ for every $f \in F_w^{\otimes}(\mathcal{H})$ (cf. the basis vectors $\zeta_{\tau}^{\vee J}$ for the symmetric Fock space from section 4.1). Note that the multi-indices $J \in \mathbb{J}^{\otimes}$ form a graded semigroup with grading |J| = m if $J \in \mathbb{N}_0^m$ and $J \cdot J'$ given by concatenation. Then $\{\psi_J := w^{-1}(|J|)\phi^{\otimes J}\}_{J \in \mathbb{J}^{\otimes}}$ is an orthonormal basis of $F_w^{\otimes}(\mathcal{H})$.

Define

$$\Delta\psi_{J} = \sum_{J',J'' \in \mathbb{J}^{\otimes}} \langle \psi_{J'} \otimes \psi_{J''}, \psi_{J} \rangle_{F_{w}^{\otimes}(\mathcal{H})} \psi_{J'} \boxtimes \psi_{J''} = \sum_{J',J'' = J} \frac{w^{-1}(|J'|)w^{-1}(|J''|)}{w^{-1}(|J|)} \psi_{J'} \boxtimes \psi_{J''},$$

and observe that

$$\langle \Delta \psi_J, \Delta \psi_L \rangle_{F_w^{\otimes}(\mathcal{H})} = \delta_{JL} \sum_{a+b=|J|} \frac{w^{-2}(a)w^{-2}(b)}{w^{-2}(a+b)} = \delta_{JL} \frac{w^{-2} * w^{-2}(|J|)}{w^{-2}(|J|)}.$$

Clearly, Δ extends to a bounded linear operator on $F_w^{\otimes}(\mathcal{H})$ iff w^{-2} is subconvolutive. Moreover, we have

$$\begin{split} &\langle \psi_{J}, \Delta^{*}(\phi^{\otimes J'}\boxtimes\phi^{\otimes J''})\rangle_{F_{w}^{\otimes}(\mathcal{H})} \\ &= \sum_{K'\cdot K''=J} \frac{\langle w^{-1}(|K'|)w^{-1}(|K''|)\phi^{\otimes K'}\boxtimes\phi^{\otimes K''}, w^{-1}(|K'|)w^{-1}(|K''|)\phi^{\otimes J'}\boxtimes\phi^{\otimes J''}\rangle_{F_{w}^{\otimes}(\mathcal{H})\boxtimes F_{w}^{\otimes}(\mathcal{H})}}{w^{-1}(|J|)} \\ &= \delta_{J,J'\cdot J''}w(|J|) \\ &= \langle \psi_{J}, \phi^{\otimes J'}\otimes\phi^{\otimes J''}\rangle_{F_{w}^{\otimes}(\mathcal{H})}, \end{split}$$

which implies that $\Delta^*(f \boxtimes g) = f \otimes g$. This also implies that Δ is independent of the choice of basis $\{\phi_i\}_{i=0}^{\infty} \subset \mathcal{H}$.

Boundedness of Δ implies that $(F_w^{\otimes}(\mathcal{H}), \otimes, 1_{\mathbb{C}})$ is a unital Banach algebra with respect to the operator norm (cf. section 4.2),

$$|||f \otimes g||_{F_w^{\otimes}(\mathcal{H})} \leq |||f||_{F_w^{\otimes}(\mathcal{H})} |||g||_{F_w^{\otimes}(\mathcal{H})},$$

where

$$|||f|||_{F_w^{\otimes}(\mathcal{H})} = \sup_{h \in F_w^{\otimes}(\mathcal{H}) \setminus \{0\}} \frac{||f \otimes h||_{F_w^{\otimes}(\mathcal{H})}}{||h||_{F_w^{\otimes}(\mathcal{H})}}.$$

Because $F_w^{\otimes}(\mathcal{H})$ is unital, this norm is equivalent to the Hilbert space norm [36],

$$||f||_{F_w^{\otimes}(\mathcal{H})} \le |||f|||_{F_w^{\otimes}(\mathcal{H})} \le ||\Delta|| ||f||_{F_w^{\otimes}(\mathcal{H})}.$$

Similarly to section 3.3, we will refer to $\Delta \colon F_w^{\otimes}(\mathcal{H}) \to F_w^{\otimes}(\mathcal{H}) \boxtimes F_w^{\otimes}(\mathcal{H})$ as a coproduct.

Proposition 7. The spectrum of $F_w^{\otimes}(\mathcal{H})$ is given by

$$\sigma(F_w^{\otimes}(\mathcal{H})) = \left\{ \chi = \langle \xi, \cdot \rangle_{F_w^{\otimes}(\mathcal{H})} : \xi = \sum_{n=0}^{\infty} w^{-2}(n) \eta^{\otimes n} : \eta \in (\mathcal{H})_{R_w} \right\},\,$$

where R_w is the radius of convergence of $\sum_{n=0}^{\infty} w^{-2}(n)z^{2n}$, $z \in \mathbb{C}$, and the spectrum is endowed with the topology of weak-* convergence. Moreover, the weak-* topology on $\sigma(F_w^{\otimes}(\mathcal{H}))$ is equivalent to the weak topology on $(\mathcal{H})_{R_w}$ with the identification of $\eta \mapsto \chi$ as above.

Proof. Let $\chi \colon F_w^{\otimes}(\mathcal{H}) \to \mathbb{C}$ be a nonzero multiplicative linear functional. Note that χ is necessarily unital since $F_w^{\otimes}(\mathcal{H})$ is unital, $\chi\Omega = 1$. Since $\chi(\phi^{\otimes J}) = \prod_{i=1}^{|j|} \chi(\phi_{j_i})$, χ is determined by its restriction to $\mathcal{H} \subset F_w^{\otimes}(\mathcal{H})$. Since characters are automatically continuous and the Hilbert space and Banach algebra norms, respectively $\|\cdot\|_{F_w^{\otimes}(\mathcal{H})}$ and $\|\cdot\|_{F_w^{\otimes}(\mathcal{H})}$, are equivalent, we have $\chi|_{\mathcal{H}} = \langle \eta, \cdot \rangle_{\mathcal{H}}$ for some $\eta \in \mathcal{H}$. As a consequence, $\chi = \langle \sum_{n=0}^{\infty} w^{-2}(n)\eta^{\otimes n}, \cdot \rangle_{F_w^{\otimes}(\mathcal{H})}$. The restriction of $\|\eta\|_{\mathcal{H}} \in [0, R_w]$ follows from finiteness of $\|\sum_{n=0}^{\infty} w^{-2}(n)\eta^{\otimes n}\|_{F_w^{\otimes}(\mathcal{H})}$. The converse is easily verified by checking multiplicativity of $\langle \sum_{n=0}^{\infty} w^{-2}(n)\eta^{\otimes n}, \cdot \rangle_{F_w^{\otimes}(\mathcal{H})}$.

Let $\tilde{R}_w = \sum_{n=0}^{\infty} w^{-2}(n) R_w^{2n}$. Since w^{-2} is strictly positive and summable, weak convergence of $\eta_i \to \eta$ in $(\mathcal{H})_{R_w}$ is equivalent to weak-* convergence of $\chi_i = \langle \xi_i, \cdot \rangle_{F_w^{\otimes}(\mathcal{H})} \to \chi = \langle \xi, \cdot \rangle_{F_w^{\otimes}(\mathcal{H})}$ in $\sigma(F_w^{\otimes}(\mathcal{H}))$, where $\xi_i = \sum_{n=0}^{\infty} w^{-2}(n) \eta_i^{\otimes n}$ and $\xi = \sum_{n=0}^{\infty} w^{-2}(n) \eta^{\otimes n}$ lie in $(F_w^{\otimes}(\mathcal{H}))_{\tilde{R}_w}$. Indeed, if $\eta_i \to \eta$ weakly, then for every $n \in \mathbb{N}_0$ and all $f_j \in \mathcal{H}$, $\chi_i(\bigotimes_{j=1}^n f_j) = \langle \eta_i^{\otimes n}, \bigotimes_{j=0}^n f_j \rangle_{F_w^{\otimes}(\mathcal{H})} \to \langle \eta^{\otimes n}, \bigotimes_{j=1}^n f_j, \rangle_{F_w^{\otimes}(\mathcal{H})} = \chi(\bigotimes_{j=1}^n f_j)$. Uniform boundedness of η_i then implies weak-* convergence of χ_i to χ . For the opposite direction, $\langle \eta_i, f \rangle_{\mathcal{H}} = \langle \sum_{n=0}^{\infty} w^{-2}(n) \eta_i^{\otimes n}, f \rangle_{F_w^{\otimes}(\mathcal{H})} = \chi_i f \to \chi f = \langle \eta, f \rangle_{\mathcal{H}}$.

By Proposition 7, for every $f, g \in F_w^{\otimes}(\mathcal{H})$ and $\chi = \langle \xi, \cdot \rangle_{F_w^{\otimes}(\mathcal{H})} \in \sigma(F_w^{\otimes}(\mathcal{H}))$,

$$\langle \Delta \xi, f \boxtimes g \rangle_{F_w^{\otimes}(\mathcal{H}) \boxtimes F_w^{\otimes}(\mathcal{H})} = \chi(\Delta^*(f \boxtimes w)) = (\chi f)(\chi g) = \langle \xi \boxtimes \xi, f \boxtimes g \rangle_{F_w^{\otimes}(\mathcal{H}) \boxtimes F_w^{\otimes}(\mathcal{H})}.$$

Thus, we can equivalently characterize the spectrum of $F_w^{\otimes}(\mathcal{H})$ as

$$\sigma(F_w(\mathcal{H})) = \left\{ \chi = \langle \xi, \cdot \rangle_{F_w^{\otimes}(\mathcal{H})} : \Delta \xi = \xi \boxtimes \xi \right\}. \tag{33}$$

The set of vectors ξ in (33) is then the cospectrum $\sigma_{co}(F_w^{\otimes}(\mathcal{H})) \subset F_w^{\otimes}(\mathcal{H})$.

We may now pass to a quotient Banach algebra and Hilbert space to recover the symmetric Fock space $F_w(\mathcal{H})$.

Theorem 8. With notation as above and $w^{-1} \in l^2(\mathbb{N}_0)$ strictly positive and w^{-2} subconvolutive, we have the following isomorphisms as Hilbert spaces and Banach algebras:

$$F_w^{\otimes}(\mathcal{H}) \Big/ \bigcap_{\chi \in \sigma(F_w^{\otimes}(\mathcal{H}))} \ker(\chi) \cong \overline{\operatorname{span}\left\{ \sum_{n=0}^{\infty} w^{-2}(n) \eta^{\otimes n} : \eta \in (\mathcal{H})_{R_w} \right\}} \cong F_w(\mathcal{H}).$$

Moreover, $F_w^{\otimes}(\mathcal{H})$ and $F_w(\mathcal{H})$ have the same spectra as Banach algebras.

Proof. Let $\mathcal{X} = \bigcap_{\chi \in \sigma(F_w^{\otimes}(\mathcal{H}))} \ker(\chi) \subset F_w^{\otimes}(\mathcal{H})$. The first isomorphism as Hilbert spaces follows from the orthogonal decomposition

$$F_w^{\otimes}(\mathcal{H}) = \mathcal{X} \oplus \operatorname{span} \left\{ \sum_{n=0}^{\infty} w^{-2}(n) \eta^{\otimes n} : \eta \in (\mathcal{H})_{R_w} \right\}.$$

The quotient is also equipped with a Banach algebra structure with the norm

$$|||f + \mathcal{X}||| = \inf_{g \in \mathcal{X}} |||f + g|||_{F_w^{\otimes}(\mathcal{H})},$$

which is equivalent to the Hilbert space norm

$$||f + \mathcal{X}|| = \inf_{g \in \mathcal{X}} ||f + g||_{F_w^{\otimes}(\mathcal{H})}.$$

The second isomorphism as Hilbert spaces follows from two observations. For $f \in (\mathcal{H})_{R_w}$ and $z \in \mathbb{C}$ with $|z| \leq 1$ we can isolate individual tensor powers using

$$f^{\otimes n} = \frac{1}{w^{-2}(n)n!} \frac{d^n}{dz^n} \Big|_{z=0} \left(\sum_{n=0}^{\infty} w^{-2}(n)(zf)^{\otimes n} \right).$$

Second, the symmetric part of $\mathcal{H}^{\otimes n}$ is densely spanned by the vectors $f^{\otimes n}$ for $f \in \mathcal{H}$. Let $n_i \in \mathbb{N}$ for $i = 1, \ldots, m$ and $\{\phi_j\}_{j=1}^{\infty} \subset \mathcal{H}$ an orthonormal basis. Then by a combinatorial argument,

$$\frac{1}{n_1!\cdots n_m!}\sum_{\sigma\in S_n}\phi_{j_{\sigma(1)}}\otimes\cdots\otimes\phi_{j_{\sigma(n)}}=\frac{1}{n_1!\cdots n_m!}\left.\frac{\partial^{n_1}\cdots\partial^{n_m}}{\partial z_1^{n_1}\cdots\partial z_m^{n_m}}\right|_{z_i=0}\left(\sum_{i=1}^m z_i\phi_i\right)^{\otimes n},$$

where (j_1, \ldots, j_n) is a vector with n_l instances of l and $\sum_{i=1}^m n_i = n$. The left hand side is a sum of every unique reordering of $\phi_{j_1} \otimes \cdots \otimes \phi_{j_n}$ while the right hand side is a sum of all sequences of n numbers drawn from $1, \ldots, m$ with n_l instances of l. Since the left hand side forms a basis for $\mathcal{H}^{\vee n} \subset \mathcal{H}^{\otimes n}$, we conclude that

$$\overline{\operatorname{span}\left\{\sum_{n=0}^{\infty}w^{-2}(n)\eta^{\otimes n}:\eta\in(\mathcal{H})_{R_w}\right\}}\cong F_w(\mathcal{H}).$$

Meanwhile, the algebra structure on $F_w(\mathcal{H})$ from the quotient and the symmetric tensor product are both given by $\operatorname{Sym} \circ \Delta^*$. Combining the above facts, we conclude that the spaces $F_w^{\otimes}(\mathcal{H})/\mathcal{X}$, $\operatorname{span} \{\sum_{n=0}^{\infty} w^{-2}(n)\eta^{\otimes n}: \eta \in (\mathcal{H})_{R_w}\}$, and $F_w(\mathcal{H})$ are isomorphic as Hilbert spaces and Banach algebras where all of the Banach algebra norms and Hilbert space norms are equivalent.

Finally, we identify the spectrum of $F_w(\mathcal{H})$. By density of the linear span of $\sigma_{co}(F_w^{\otimes}(\mathcal{H}))$ in $F_w^{\otimes}(\mathcal{H})$, the restriction of the coproduct to $F_w(\mathcal{H})$ is well-defined as a linear map

$$\Delta|_{F_w(\mathcal{H})} \colon F_w(\mathcal{H}) \to F_w(\mathcal{H}) \boxtimes F_w(\mathcal{H}) \subset F_w^{\otimes}(\mathcal{H}) \boxtimes F_w^{\otimes}(\mathcal{H}).$$

Thus, we have $(\Delta|_{F_w(\mathcal{H})})^* = \operatorname{Sym} \circ \Delta^*$, where the adjoint in the left-hand side is taken on $B(F_w(\mathcal{H}), F_w(\mathcal{H}) \boxtimes F_w(\mathcal{H}))$ and in the right-hand side it is taken on $B(F_w^{\otimes}(\mathcal{H}), F_w^{\otimes}(\mathcal{H}) \boxtimes F_w^{\otimes}(\mathcal{H}))$. Now, for every $\chi \in \sigma(F_w(\mathcal{H}))$, there exists $\xi \in F_w(\mathcal{H})$ such that $\chi = \langle \xi, \cdot \rangle_{F_w(\mathcal{H})}$, giving

$$\begin{split} \langle \xi \boxtimes \xi, f \boxtimes g \rangle_{F_w^{\otimes}(\mathcal{H}) \boxtimes F_w^{\otimes}(\mathcal{H})} &= \langle \xi, f \rangle_{F_w(\mathcal{H})} \langle \xi, g \rangle_{F_w(\mathcal{H})} = (\chi f)(\chi g) \\ &= \langle \xi, f \vee g \rangle_{F_w(\mathcal{H})} &= \langle \Delta \xi, f \boxtimes g \rangle_{F_w^{\otimes}(\mathcal{H}) \boxtimes F_w^{\otimes}(\mathcal{H})} \,. \end{split}$$

Thus, we have $\Delta \xi = \xi \boxtimes \xi$ and it follows from (33) that $\sigma(F_w(\mathcal{H})) \subseteq \sigma(F_w^{\otimes}(\mathcal{H}))$. Since $\sigma(F_w^{\otimes}(\mathcal{H})) \subseteq \sigma(F_w(\mathcal{H}))$ by the subspace relation $F_w(\mathcal{H}) \subset F_w^{\otimes}(\mathcal{H})$, we deduce that $\sigma(F_w(\mathcal{H})) = \sigma(F_w^{\otimes}(\mathcal{H}))$. \square

Letting $\Gamma \colon F_w(\mathcal{H}) \to C(\sigma(F_w(\mathcal{H})))$ denote the Gelfand map on the commutative Banach algebra $F_w(\mathcal{H})$, the following is a corollary of Theorem 8 and equality of the spectra of $F_w^{\otimes}(\mathcal{H})$ and $F_w(\mathcal{H})$.

Corollary 9. The Gelfand map $\Gamma \colon F_w(\mathcal{H}) \to C(\sigma(F_w(\mathcal{H})))$ is injective. As a result, the space $\hat{F}_w(\mathcal{H}) := \Gamma(F_w(\mathcal{H}))$ equipped with the inner product $\langle \hat{f}, \hat{g} \rangle_{\hat{F}_w(\mathcal{H})} = \langle \Gamma^{-1} \hat{f}, \Gamma^{-1} \hat{g} \rangle_{F_w(\mathcal{H})}$ is a unital RKHA with reproducing kernel $\hat{k} \colon \sigma(F_w(\mathcal{H})) \times \sigma(F_w(\mathcal{H})) \to \mathbb{C}$ defined as

$$\hat{k}(\chi_1, \chi_2) = \langle \xi_1, \xi_2 \rangle_{F_w(\mathcal{H})}, \quad \chi_i = \langle \xi_i, \cdot \rangle_{F_w(\mathcal{H})}.$$

Moreover, $F_w(\mathcal{H})$ and $\hat{F}_w(\mathcal{H})$ are isomorphic Banach algebras under the Gelfand map.

Proof. Suppose that $\Gamma f = \Gamma g$ for some $f, g \in F_w(\mathcal{H})$. Then $f - g \in \ker \chi$ for every $\chi \in \sigma(F_w(\mathcal{H}))$. Since $\sigma(F_w(\mathcal{H})) = \sigma(F_w^{\otimes}(\mathcal{H}))$, it follows that f - g lies in $\mathcal{X} \cap F_w(\mathcal{H})$ and thus f - g = 0 since $F_w(\mathcal{H}) \cong F_w^{\otimes}(\mathcal{H})/\mathcal{X}$ by Theorem 8. This proves injectivity of Γ and thus well-definition of $(\hat{F}_w(\mathcal{H}), \langle \cdot, \cdot \rangle_{\hat{F}_w(\mathcal{H})})$ as a Hilbert space.

Next, for every $\hat{f} = \Gamma f \in \hat{F}_w(\mathcal{H})$ and $\chi = \langle \xi, \cdot \rangle_{F_w(\mathcal{H})} \in \sigma(F_w(\mathcal{H}))$ we have

$$\hat{f}(\chi) = \chi f = \langle \xi, f \rangle_{F_w(\mathcal{H})} = \langle \Gamma \xi, \hat{f} \rangle_{\hat{F}_w(\mathcal{H})} = \langle \hat{k}_{\chi}, \hat{f} \rangle_{\hat{F}_w(\mathcal{H})}.$$

Thus, $\hat{F}_w(\mathcal{H})$ is an RKHS with reproducing kernel \hat{k} , as claimed.

To verify that $\hat{F}_w(\mathcal{H})$ is an RKHA, observe that

$$\|\hat{k}_{\chi} \otimes \hat{k}_{\chi}\|_{\hat{F}_{w}(\mathcal{H})}^{2} = \langle \xi, \xi \rangle_{F_{w}(\mathcal{H})}^{2} = \langle \Delta \xi, \Delta \xi \rangle_{F_{w}(\mathcal{H}) \boxtimes F_{w}(\mathcal{H})} \leq \|\Delta^{2}\| \|\xi\|_{F_{w}(\mathcal{H})}^{2} = \|\Delta\|^{2} \|\hat{k}_{\chi}\|_{\hat{F}_{w}(\mathcal{H})}^{2}.$$

This implies that $\hat{\Delta} : \hat{k}_{\chi} \mapsto \hat{k}_{\chi} \boxtimes k_{\chi}$ extends to a bounded linear operator $\Delta : \hat{F}_w(\mathcal{H}) \to \hat{F}_w(\mathcal{H}) \boxtimes \hat{F}_w(\mathcal{H})$, so $\hat{F}_w(\mathcal{H})$ is an RKHA according to Definition 4. This RKHA is unital since $\Gamma\Omega = 1_{\sigma(F_w(\mathcal{H}))}$. Finally, for every $f, g \in F_w(\mathcal{H})$ we have

$$\Gamma(f \otimes g)(\chi) = \chi(f \otimes g) = (\chi f)(\chi g) = (\Gamma f)(\Gamma g)$$

and we conclude that Γ is a Banach algebra isomorphism.

Suppose now that \mathcal{H} is an RKHS with reproducing kernel $k: X \times X \to \mathbb{C}$ with feature map $\varphi: X \times X \to \mathbb{C}$. Then we may define feature maps $\tilde{\varphi}: X \to F_w(\mathcal{H})$ and $\hat{\varphi}: X \to \hat{F}_w(\mathcal{H})$ analogously to (17) and (18), respectively. Pulling back $\hat{F}_w(\mathcal{H})$ to X then defines an RKHA $\tilde{\mathcal{H}} = \{\hat{f} \circ \hat{\varphi}: \hat{f} \in \hat{F}_w(\mathcal{H})\}$ with kernel

$$\tilde{k}(x,y) = \hat{k}(\hat{\varphi}(x), \hat{\varphi}(y)) = \sum_{n=0}^{\infty} \frac{w^{-2}(n)}{\varpi^{2n}} k(x,y)^n.$$

Since $\tilde{k}(x,y) - (\varpi w(1))^{-2}k(x,y)$ is a positive-definite kernel, we have the inclusion $\mathcal{H} \subset \tilde{\mathcal{H}}$ as shown in [1]. Note that $\tilde{\mathcal{H}}_w$ is unital even if 1_X is not an element of \mathcal{H} .

5.2. **Decomposition of the spectrum and** *-structure. Let \mathcal{H} be a Hilbert space and $J: \mathcal{H} \to \mathcal{H}$ be a conjugate-linear unitary such that $J^2 = \mathrm{Id}_{\mathcal{H}}$. Then we may also define a conjugate-linear unitary and Banach algebra involution * for $F_w^{\otimes}(\mathcal{H})$ and $F_w(\mathcal{H})$ by extending the following definition by conjugate linearity:

$$\Omega^* = \Omega, \quad (f_1 \otimes \cdots \otimes f_n)^* = J(f_n) \otimes \cdots \otimes J(f_1), \quad (f_1 \vee \cdots \vee f_n)^* = J(f_n) \vee \cdots \vee J(f_1).$$

Since $\langle J(v), J(u) \rangle_{\mathcal{H}} = \overline{\langle v, u \rangle_{\mathcal{H}}}$, these involutions satisfy $\langle \xi^*, \eta^* \rangle_{F_w^{\otimes}(\mathcal{H})} = \overline{\langle \xi, \eta \rangle_{F_w^{\otimes}(\mathcal{H})}}$ and $\langle \xi^*, \eta^* \rangle_{F_w(\mathcal{H})} = \overline{\langle \xi, \eta \rangle_{F_w(\mathcal{H})}}$.

In what follows, $\mathbb{J} \subset \mathbb{N}_0^{\mathbb{N}}$ will be the index set for basis vectors of $F_w(\mathcal{H})$ introduced in section 4.1, and $\mathbb{A}_w \subset \ell^2(\mathbb{N}_0)$ the set of square-integrable sequences with positive elements defined in section 4.3.

Proposition 10. Fix an orthonormal basis $\{\phi_j\}_{j=0}^{\infty} \subset \mathcal{H}$, a sequence $a \in \mathbb{A}_w$, and consider the subset

$$\mathbb{T}_a^{complex} = \left\{ \langle \xi, \cdot \rangle_{F_w(\mathcal{H})} : \xi = \sum_{n=0}^{\infty} w^{-2}(n) \left(\sum_{j=0}^{\infty} a_j z_j \phi_j \right)^{\vee n} : (z_j)_j \in l^{\infty}(\mathbb{N}_0), \ |z_j| = 1 \right\} \subset \sigma(F_w(\mathcal{H})).$$

Then, $\hat{F}_w(\mathcal{H})|_{\mathbb{T}_a^{complex}}$ is an RKHA of continuous functions on $\mathbb{T}_a^{complex}$, and if $a_j > 0$, then $f|_{\mathbb{T}_a^{complex}}$ uniquely determines $f \in \hat{F}_w(\mathcal{H})$.

Proof. Recall from section 2.2 that $\hat{F}_w(\mathcal{H})|_{\mathbb{T}_a^{\text{complex}}}$ is isomorphic as a Hilbert space to the norm closure of $\text{span}\{\hat{k}_\chi:\chi\in\mathbb{T}_a^{\text{complex}}\}$ in $\hat{F}_w(\mathcal{H})$, which is in turn isomorphic to the norm closure of $\Xi_a:=\text{span}\{\xi\in F_w(\mathcal{H}):\langle\xi,\cdot\rangle_{F_w(\mathcal{H})}\in\mathbb{T}_a^{\text{complex}}\}$ in $F_w(\mathcal{H})$ by Corollary 9. Thus, to prove the proposition it suffices to show that Ξ_a is a dense subspace of $F_w(\mathcal{H})$. To that end, we will show that for every sequence $(j_0,j_1,\ldots)\in\mathbb{J}$ we can recover the symmetric tensor $\phi_0^{\vee j_0}\cdots\phi_{i_0}^{\vee j_{i_0}}$ in the closure $\Xi_a^{-\|\cdot\|_{F_w(\mathcal{H})}}$, where j_i is constantly zero after i_0 . Since the collection of such tensors forms an orthogonal basis of $F_w(\mathcal{H})$ (see section 4.1), we can then deduce that $\Xi_a^{-\|\cdot\|_{F_w(\mathcal{H})}}=F_w(\mathcal{H})$.

Let $M = \sum_i j_i$ and ν_N denote the Haar measure on \mathbb{T}^N . Pick $N > i_0$ large enough such that $\left\|\sum_{i=N+1}^{\infty} a_i z_i \phi_i\right\|_{F_w(\mathcal{H})} = \delta_N < 1/2$. Then,

$$\begin{split} & \int_{\mathbb{T}^N} z_0^{-j_0} \cdots z_N^{-j_N} \sum_{n=0}^{\infty} w^{-2}(n) \left(\sum_{i=0}^{\infty} a_i z_i \phi_i \right)^{\vee n} d\nu_N(z_0, \dots, z_N) \\ & = \int_{\mathbb{T}^N} z_0^{-j_0} \cdots z_N^{-j_N} \sum_{n=0}^{\infty} w^{-2}(n) \sum_{k=0}^{n} \binom{n}{k} \left(\sum_{i=0}^{N} a_i z_i \phi_i \right)^{\vee k} \vee \left(\sum_{i=N+1}^{\infty} a_i z_i \phi_i \right)^{\vee n-k} d\nu_N(z_0, \dots, z_N) \\ & = \sum_{n=M}^{\infty} w^{-2}(n) a_0^{j_0} \cdots a_{i_0}^{j_{i_0}} \binom{n}{M} \left(\phi_0^{\vee j_0} \cdots \phi_N^{\vee j_N} \right) \vee \left(\sum_{i=N+1}^{\infty} a_i z_i \phi_i \right)^{\vee n-M} \\ & = w^{-2}(M) a_0^{j_0} \cdots a_{i_0}^{j_{i_0}} \left(\phi_0^{\vee j_0} \cdots \phi_{i_0}^{\vee j_{i_0}} \right) + r_N, \end{split}$$

where

If we can show that the remainder r_N above converges to zero as N increases, then we will be able to isolate $\phi_0^{\vee j_0} \cdots \phi_N^{\vee j_{i_0}}$. Since $\binom{n}{M} \leq n^M/M!$,

$$||r_N||_{F_w(\mathcal{H})}^2 = \left\| \sum_{n=M+1}^{\infty} w^{-2}(n) \binom{n}{M} \left(\sum_{i=N+1}^{\infty} a_i z_i \phi_i \right)^{\vee n-M-1} \right\|_{F_w(\mathcal{H})}^2$$

$$\leq \sum_{n=M+1}^{\infty} w^{-2}(n) \frac{n^{2M}}{(M!)^2} \delta_N^{2(n-M-1)}$$

$$\leq \frac{||w^{-2}||_{\ell^{\infty}(\mathbb{N}_0)}}{(M!)^2} \sum_{n=M+1}^{\infty} \frac{n^{2M}}{4^{n-M-1}} = \operatorname{Const}(w, M) < \infty.$$

By boundedness of the symmetric product in $F_w(\mathcal{H})$ and summability of w^{-2} , the remainder r_N converges to zero as $N \to \infty$.

Therefore, by strict positivity of $w^{-2}(M)$ and all of the a_i 's, $\phi_0^{\vee j_0} \cdots \phi_{i_0}^{\vee j_{i_0}} \in \overline{\Xi_a}^{\|\cdot\|_{F_w(\mathcal{H})}}$ for every sequence of nonnegative integers $(j_i)_i \in \mathbb{J}$, making Ξ_a a dense subspace of $F_w(\mathcal{H})$.

In fact, $\sigma(F_w(\mathcal{H})) = \bigcup_{a \in \mathbb{A}_w} \mathbb{T}_a^{\text{complex}}$, and we can identify $F_w(\mathcal{H}) \cong \hat{F}_{w,a}(\mathcal{H})$ for any $a \in \mathbb{A}_w$ with strictly positive elements. Here, $\hat{F}_{w,a}(\mathcal{H}) \subset C(\mathbb{T}_a)$ is the RKHA with reproducing kernel

 $\hat{k}^{(a)} \colon \mathbb{T}_a^{\text{complex}} \times \mathbb{T}_a^{\text{complex}} \to \mathbb{C},$

$$\hat{k}^{(a)}(\vec{z}, \vec{y}) = \sum_{n=0}^{\infty} w^{-2}(n) \left(\sum_{j=0}^{\infty} a_j^2 \overline{z_j} y_j \right)^n,$$

where $\vec{z} \equiv \chi_{a,z} = \langle \xi_{a,z}, \cdot \rangle_{F_w(\mathcal{H})}$ and $\vec{y} \equiv \chi_{a,z} = \langle \xi_{a,y}, \cdot \rangle_{F_w(\mathcal{H})}$ for $\xi_{a,z}, \xi_{a,y} \in \sigma_{co}(F_w(\mathcal{H}))$. The isomorphism $v_a : \hat{F}_{w,a}(\mathcal{H}) \to F_w(\mathcal{H})$ realizing this identification is defined by linear extension of

$$v_a\left(\hat{k}^{(a)}(\vec{z},\cdot)\right) = \xi_{a,z}.$$

In section 4, we specialize to tori in $\sigma(F_w(\mathcal{H}_\tau))$ with real kernel. If $\mathcal{H} \subset C(X)$ is an RKHS with a real kernel $k \colon X \times X \to \mathbb{R}$ and a self-conjugate orthonormal basis $\{\phi_j\}_{j \in \mathbb{Z}}$ containing the unit function $1_X = \phi_0$, then

$$k_x = a_0(x)1_X + \sum_{j=1}^{\infty} a_j(x) \frac{z_j(x)\phi_j + \overline{z_j(x)\phi_j}}{\sqrt{2}}$$

analogously to (21). The associated tori

$$\mathbb{T}_{a(x)} = \left\{ \langle \xi, \cdot \rangle_{F_w(\mathcal{H})} : \xi = \sum_{n=0}^{\infty} w^{-2}(n) \left(a_0(x) 1_X + \sum_{j=1}^{\infty} a_j(x) \frac{z_j \phi_j + \overline{z_j \phi_j}}{\sqrt{2}} \right)^{\vee n} : |z_j| = 1 \right\},$$

have real-valued kernels,

$$k^{(a)}(\vec{z}, \vec{y}) = \sum_{n=0}^{\infty} w^{-2}(n) \left(a_0^2(x) + \sum_{j=1}^{\infty} a_j^2(x) \operatorname{Re}(\overline{z_j} y_j) \right)^n.$$

Even between points in two different tori, $\chi = \langle \xi, \cdot \rangle_{F_w(\mathcal{H})} \in \mathbb{T}_a$ and $\psi = \langle \eta, \cdot \rangle_{F_w(\mathcal{H})} \in \mathbb{T}_b$, the kernel $\hat{k}(\chi, \psi) = \langle \xi, \eta \rangle_{F_w(\mathcal{H})}$ is real-valued.

If we use $J(\phi_i) = \overline{\phi_i}$ to define a *-structure, then for every $\langle \xi, \cdot \rangle_{F_w(\mathcal{H})} \in \mathbb{T}_a$, $\xi^* = \xi$. Since the *-structure is a conjugate-linear unitary involution, for $\hat{f} = \Gamma f$ and $\chi = \langle \xi, \cdot \rangle_{F_w(\mathcal{H})}$ we have

$$\widehat{f^*}(\chi) = \langle \xi, f^* \rangle_{F_w(\mathcal{H})} = \langle \xi^*, f^* \rangle_{F_w(\mathcal{H})} = \overline{\langle \xi, f \rangle_{F_w(\mathcal{H})}} = \overline{\widehat{f}(\chi)}.$$

6. Fock space embedding of observables and dynamics

This section addresses the remaining material on Fock space representation of observables and dynamics from sections 4.5–4.7. In section 6.1, we prove the L^2 convergence of the Fock space approximation $f_{m,\tau}^{(t)}$ to the true Koopman evolution U^tf claimed in Proposition 5, assuming that the results (9) (dynamical convergence) and (23) (approximation of the identity). In section 6.2, we show that (23) holds for a class of positive, continuous, decaying kernels that includes the Gaussian kernels from (24). In section 6.3 we verify the pointwise and L^2 convergence of the approximations $f_{m,\tau,d}^{(t)}$ on finite-dimensional tori from section 4.7. In section 6.4 we discuss embedding approaches utilizing the complex tori $\mathbb{T}_a^{\text{complex}}$ from Proposition 10.

6.1. **Proof of Proposition 5.** Observe that the prediction function $f_{m,\tau}^{(t)}$ from (26) can be expressed as

$$f_{m,\tau}^{(t)} = \frac{g_{m,\tau}^{(t)}}{h_{m,\tau}^{(t)}},$$

where $g_{m,\tau}^{(t)} = \tilde{\pi} \tilde{g}_{m,\tau}^{(t)}$, $h_{m,\tau}^{(t)} = \tilde{\pi} \tilde{h}_{m,\tau}^{(t)}$, and $\tilde{g}_{m,\tau}^{(t)}$, $\tilde{h}_{m,\tau}^{(t)} \in F_w(\mathcal{H}_\tau)$ are elements of the weighted Fock space given by

$$\tilde{g}_{m,\tau}^{(t)} = \tilde{U}_{\tau}^{-t} \tilde{\mathcal{K}}_m f, \quad \tilde{h}_{m,\tau}^{(t)} = \tilde{U}_{\tau}^{-t} \tilde{\mathcal{K}}_m 1_X.$$

Focusing first on $\tilde{g}_{m,\tau}^{(t)}$, it follows from multiplicativity of \tilde{U}_{τ}^{t} (see section 4.4) that

$$\tilde{g}_{m,\tau}^{(t)} = \tilde{U}_{\tau}^{-t} \int_{X} \kappa^{\vee m}(\cdot, x) f(x) \, d\mu(x) = \int_{X} (U_{\tau}^{-t} \kappa(\cdot, x))^{\vee m} f(x) \, d\mu(x).$$

As a result, since $\tilde{\pi}$ and the inclusion map $\tilde{K}_{\tau}^* \colon \tilde{\mathcal{H}}_{\tau} \to H$ are also multiplicative, we have

$$\tilde{K}_{\tau}^* \tilde{\pi} \tilde{g}_{m,\tau}^{(t)} = \tilde{K}_{\tau}^* \int_{X} (U_{\tau}^{-t} \kappa(\cdot, x))^m f(x) \, d\mu(x) = \int_{Y} (K_{\tau}^* U_{\tau}^{-t} \kappa(\cdot, x))^m f(x) \, d\mu(x).$$

Since $\kappa(\cdot, x)$ lies in \mathcal{H}_{τ} for all $\tau > 0$ by assumption, it follows from (9) in conjunction with unitarity and multiplicativity of U^t that

$$\begin{split} \lim_{\tau \to 0^+} \tilde{K}_{\tau}^* \tilde{\pi} \tilde{g}_{m,\tau}^{(t)} &= \int_X \left(\lim_{\tau \to 0^+} K_{\tau}^* U_{\tau}^{-t} \kappa(\cdot, x) \right)^m f(x) \, d\mu(x) = \int_X (U^{-t} \iota \kappa(\cdot, x))^m f \, d\mu \\ &= \int_X U^{-t} \iota \kappa_m(\cdot, x) f \, d\mu = \int_X \iota \kappa_m(\cdot, x) U^t f \, d\mu = \iota \tilde{\mathcal{K}}_m U^t f. \end{split}$$

Proceeding as above, we get

$$\lim_{\tau \to 0^+} \tilde{K}_{\tau}^* \tilde{\pi} \tilde{h}_{m,\tau}^{(t)} = \iota \tilde{\mathcal{K}}_m U^t 1_X = \iota d_m,$$

and thus

$$\lim_{\tau \to 0^+} \iota f_{m,\tau}^{(t)} = \lim_{\tau \to 0^+} \frac{\tilde{K}_{\tau}^* \tilde{\pi} \tilde{g}_{m,\tau}^{(t)}}{\tilde{K}_{\tau}^* \tilde{\pi} \tilde{h}_{m,\tau}^{(t)}} = \iota \frac{\tilde{\mathcal{K}}_m U^t f}{d_m} = P_m U^t f.$$

Therefore, under assumption (23),

$$\lim_{m \to \infty} \lim_{\tau \to 0^+} \iota f_{m,\tau}^{(t)} = U^t f$$

in the norm of H, proving the proposition.

6.2. Approximation of the identity on continuous functions. We will use the following lemma to deduce that that the radial Gaussian kernel κ from (24) provides an approximation of the identity in the sense of (23).

Lemma 11. Let X be a compact Hausdorff space and μ a Borel probability measure with full support on X. Let $\kappa \colon X \times X \to \mathbb{R}_{>0}$ be a continuous kernel such that $\kappa(x,x) > \kappa(x,y)$ for all $x,y \in X$ and $x \neq y$. For $m \in \mathbb{N}$, define $d_m(x) = \int_X \kappa(x,\cdot)^m d\mu$ and $p_m(x,y) = \frac{\kappa(x,y)^m}{d_m(x)}$. Then for $f \in C(X)$,

$$\lim_{m \to \infty} \left\| f - \int_X p_m(\cdot, y) f(y) \, d\mu(y) \right\|_{L^2(\mu)} = 0.$$

Proof. Set $\kappa_x = \kappa(x, \cdot)$ and define the sets $E_{x,\epsilon} = \kappa_x^{-1}((\kappa(x, x) - \epsilon, \kappa(x, x)])$. First, $E_{x,\epsilon}$ provide a basis for the topology of x. That is, for every open neighborhood $U \ni x$, $\kappa(x, X \setminus U) \subset [0, \kappa(x, x))$ is compact hence there is an $\epsilon > 0$ such that $x \in E_{x,\epsilon} \subset U$.

We also have the estimates

$$\int_{E_{x,\epsilon}} \kappa_x^m d\mu \ge \mu(E_{x,\epsilon} \setminus E_{x,\epsilon/2}) (\kappa(x,x) - \epsilon)^m + \mu(E_{x,\epsilon/2}) (\kappa(x,x) - \epsilon/2)^m,$$

$$\int_{X \setminus E_{x,\epsilon}} \kappa_x^m d\mu \le \mu(X \setminus E_{x,\epsilon}) (\kappa(x,x) - \epsilon)^m.$$

Therefore, there is a constant C_x depending on x and an $m_{x,0}$ such that $d_m(x) \ge C_x(\kappa(x,x) - \epsilon/2)^m$ for $m \ge m_{x,0}$. As a result, for every $m > m_{x,0}$, we have

$$p_m(x,\cdot)|_{X\setminus E_{x,\epsilon}} \le \mu(X\setminus E_{x,\epsilon}) \frac{(\kappa(x,x)-\epsilon)^m}{C_x(\kappa(x,x)-\epsilon/2)^m},$$

so $p_m(x,\cdot)|_{X\setminus E_{x,\epsilon}}$ converges uniformly to zero as $m\to\infty$.

We will first show pointwise convergence. Fix $x \in X$, $\delta > 0$, and pick $\epsilon > 0$ such that $\sup_{y \in E_{x,\epsilon}} |f(y) - f(x)| < \delta$. Choose also $n \in \mathbb{N}$ such that $p_m(x,\cdot)|_{X \setminus E_{x,\epsilon}} < \delta$ for all $m \geq n$. Define $f_m = \int_X p_m(\cdot,y) f(y) \, d\mu(y)$ and observe that

$$|f_m(x) - f(x)| \le \delta ||f||_{C(X)} \mu(X \setminus E_{x,\epsilon}) + \left| f(x) - \int_{E_{x,\epsilon}} p_m(x,y) f(y) \, d\mu(y) \right|, \quad \forall m \ge n.$$

Set $p_{x,\epsilon,m} = \int_{E_{x,\epsilon}} p_m(x,\cdot) d\mu$. Since $\int_X p_m(x,\cdot) d\mu = 1$ and $\int_{X \setminus E_{x,\epsilon}} p_m(x,\cdot) d\mu \leq \delta$, $1 - p_{x,\epsilon,m} \leq \delta$. Therefore,

$$\left| f(x) - \int_{E_{x,\epsilon}} p_m(x,y) f(y) \, d\mu(y) \right|$$

$$= \left| f(x) - \int_{E_{x,\epsilon}} p_m(x,y) (f(y) - f(x)) \, d\mu(y) - f(x) \int_{E_{x,\epsilon}} p_m(x,y) \, d\mu(y) \right|$$

$$\leq \delta + |f(x)| (1 - p_{x,\epsilon,m}) \leq \delta + ||f||_{C(X)} \delta,$$

giving

$$|f_m(x) - f(x)| \le (2||f||_{C(X)} + 1)\delta.$$

Pointwise convergence as $m \to \infty$ follows since δ was arbitrary,

$$\lim_{m \to \infty} |f_m(x) - f(x)| = 0, \quad \forall x \in \text{supp}(\mu).$$

Finally, we conclude $L^2(\mu)$ convergence by the dominated convergence theorem as

$$|f_m(x)| \le \int_X |f(y)| p_m(x,y) \, d\mu(y) \le ||f||_{C(X)}.$$

6.3. Approximation on finite-dimensional tori. Observe that for every $x \in X$ the feature vectors $\eta_{\tau,d,x}$ from (27) lie in $(\mathcal{H}_{\tau})_1 \subseteq (\mathcal{H}_{\tau})_{R_w}$, and norm-converge to $\eta_{\tau,x}$ as $d \to \infty$. By Proposition 7 and Theorem 8, the corresponding spectral elements $\hat{\varphi}_{\tau,d}(x)$ converge to $\hat{\varphi}_{\tau}(x)$ in the weak-* topology of $\sigma(F_w(\mathcal{H}))$. This implies pointwise convergence of the prediction function,

$$\lim_{d \to \infty} f_{m,\tau,d}^{(t)}(x) = f_{m,\tau}^{(t)}(x), \quad \forall x \in X, \tag{34}$$

as claimed in section 4.7.

Next, the square residual $|f_{m,\tau,d}^{(t)}(x) - f_{m,\tau}^{(t)}(x)|^2 = |\hat{f}_{m,\tau}^{(t)}(\hat{\varphi}_{\tau,d}(x)) - \hat{f}_{m,\tau}^{(t)}(\hat{\varphi}_{\tau}(x))|^2$ is dominated by the constant function on X equal to $||\hat{f}_{m,\tau}^{(t)}||^2_{C(\sigma(F_w(\mathcal{H}_{\tau})))}$. This function is μ -integrable since μ is a probability measure. Hence, by the dominated convergence theorem, the pointwise convergence (34) implies norm convergence in $H = L^2(\mu)$, verifying (28).

6.4. Topological models on complex tori. The topological models of regularized Koopman dynamics from section 4.5 were defined on the union of tori $\mathbb{S}_{\tau} \subset \sigma(\hat{F}_w(\mathcal{H}_{\tau}))$ that contain the image $\hat{X}_{\tau} \subset \mathbb{S}_{\tau}$ of state space under the feature map $\hat{\varphi}_{\tau}$. In this subsection, we consider an alternative formulation that uses Proposition 10 to build a topological model that is defined on a single complex torus $\mathbb{T}_{\tau,a}^{\text{complex}} \subset \sigma(\hat{F}_w(\mathcal{H}_{\tau}))$.

First, we choose the approximate Koopman eigenfunctions $\zeta_{j,\tau}$ as the orthonormal basis functions of \mathcal{H}_{τ} , indexed by $j \in \mathbb{Z}$. We also choose the complex torus $\mathbb{T}_a^{\text{complex}}$ associated with this basis and an arbitrary weight vector $a = (a_j)_{j \in \mathbb{Z}} \in \mathbb{A}_w$ with strictly positive elements.

Next, consider a lifting map $\mathcal{L} \colon \tilde{\mathcal{H}}_{\tau} \to F_w(\mathcal{H}_{\tau})$; i.e., a norm-continuous map satisfying $\tilde{\pi} \circ \mathcal{L} = \operatorname{Id}_{\tilde{\mathcal{H}}_{\tau}}$. Since \mathcal{H}_{τ} is a subspace of $\tilde{\mathcal{H}}_{\tau}$, a possible choice for \mathcal{L} is the inclusion map $\mathcal{H}_{\tau} \hookrightarrow \tilde{\mathcal{H}}_{\tau}$. More generally, $\tilde{f} = \mathcal{L}f$ could yield a representation of $f \in \tilde{\mathcal{H}}_{\tau}$ as a product–sum of functions $g_{i,j} \in \mathcal{H}_{\tau}$, i.e., $\tilde{f} = \sum_{i=1}^{N} c_i \bigvee_{j=0}^{M_i} g_{i,j} \in F_w(\mathcal{H}_{\tau})$ where $f = \sum_{i=1}^{N} c_i \prod_{j=0}^{M_i} g_{i,j}$. The embedding via \mathcal{K}_m from section 4.6 can be viewed as an approximate version of such an \mathcal{L} up to normalization.

With any such lifting map, define $\hat{\mathcal{L}}_a : \tilde{\mathcal{H}}_{\tau} \to \hat{F}_{w,a}(\mathcal{H}_{\tau})$ as $\hat{\mathcal{L}}_a = v_a^{-1} \circ \Gamma \circ \mathcal{L}$ and let $\hat{f}_a = \hat{\mathcal{L}}_a f$. Due to Proposition 10, we can evolve \hat{f}_a by the torus rotation $R_{\tau}^t|_{\mathbb{T}_a^{\text{complex}}}$ to get $\hat{f}_{\tau,a}^{(t)} = \hat{f}_a \circ R_{\tau}^t|_{\mathbb{T}_a^{\text{complex}}}$, and $\hat{f}_{\tau}^{(t)} = v_a \hat{f}_a^{(t)}$ will be the unique function in $\hat{F}_w(\mathcal{H}_{\tau})$ such that $\hat{f}_{\tau}^{(t)}|_{\mathbb{T}_a^{\text{complex}}} = \hat{f}_{\tau,a}^{(t)}$. Hence, for any lifting of f, there is a unique approximate Koopman evolution $f_{\tau}^{(t)} = \hat{\pi} \hat{f}_{\tau}^{(t)}$ induced from the rotation system on \mathbb{T}_a such that $f_{\tau}^{(0)} = f$.

7. Data-driven formulation

The finite-dimensional approximation scheme described in section 4.7 is amenable to data-driven approximation using kernel methods [7, 22, 34, 38–40]. In this subsection, we give a high-level overview of the data-driven scheme, relegating a detailed convergence analysis to these references.

We consider training data $y_0, y_1, \ldots, y_{N-1}$ in a data space Y. The data is sampled along a dynamical trajectory $x_0, x_1, \ldots, x_{N-1} \in X$ with $x_i = \Phi^{i \Delta t}(x_0)$, for an initial condition x_0 in the forward-invariant set M, sampling interval $\Delta t > 0$, and observation map $F: X \to Y$ such that $y_i = F(x_i)$. Letting $\mu_N = \sum_{i=0}^{N-1} \delta_{x_i}/N$ be the sampling measure associated with the training dataset, we have

$$\lim_{N \to \infty} \int_{M} f \, d\mu_{N} = \int_{M} f \, d\mu, \quad \forall f \in C(M), \tag{35}$$

for μ -a.e. $x \in M$ by ergodicity. For systems with so-called physical invariant measures [10], (35) holds for initial conditions x_0 drawn from a set of positive ambient measure (e.g., Lebesgue measure) on M.

Based on the above, our data-driven methods replace the Hilbert space $H = L^2(\mu)$ associated with the invariant measure with finite-dimensional Hilbert spaces $H_N := L^2(\mu_N)$ associated with the sampling measures. The various kernel integral operators employed in previous sections, including K_{τ} and G_{τ} from section 3.2, \tilde{K}_{τ} from section 4.2, and K_m , \tilde{K}_m , and \tilde{P}_m from section 4.6, are then approximated by analogous (finite-rank) operators defined on H_N . We will denote these operators using N subscripts; e.g., $\tilde{K}_{m,N} : H_N \to C(M)$ is the data-driven analog of \tilde{K}_m and is defined as

$$\tilde{\mathcal{K}}_{m,N}f = \int_X \kappa_m(\cdot, x) f(x) \, d\mu_N(x) \equiv \frac{1}{N} \sum_{i=0}^{N-1} \kappa_m(\cdot, x_i) f(x_i).$$

In our constructions, the reproducing kernel k_{τ} of \mathcal{H}_{τ} is built by Markov normalization of an un-normalized kernel function (see appendix A). In the data-driven setting, normalization leads to a data-dependent kernel $k_{\tau,N} \colon X \times X \to \mathbb{R}_+$ that is continuous, strictly positive-definite, and Markov-normalized with respect to μ_N . The kernels $k_{\tau,N}$ thus have associated RKHSs $\mathcal{H}_{\tau,N} \subset C(X)$, and the data-driven analog of $K_{\tau} \colon H \to \mathcal{H}_{\tau}$ is an integral operator $K_{\tau,N} \colon H_N \to \mathcal{H}_{\tau,N}$ associated with

the data-dependent kernel $k_{\tau,N}$. In the large data limit, $N \to \infty$, the restrictions of $k_{\tau,N}$ to $M \times M$ converge uniformly to k_{τ} for every initial condition $x_0 \in M$ satisfying (35).

Using an eigendecomposition of $G_{\tau,N} = K_{\tau,N}^* K_{\tau,N}$,

$$G_{\tau,N}\phi_{j,N} = \lambda_{j,\tau,N}\phi_{j,N}, \quad j \in \mathbb{N}_0,$$

we have, by analogous properties to (K1)–(K4), that the eigenvalues $\lambda_{j,\tau,N}$ lie in the interval (0,1] and can be ordered as $1=\lambda_{0,\tau,N}>\lambda_{1,\tau,N}\geq\lambda_{2,\tau,N}\geq\cdots\lambda_{N-1,\tau,N}>0$. Moreover, the corresponding eigenvectors $\phi_{j,N}$ form an orthonormal basis of H_N that does not depend on τ by a semigroup property analogous to (K3). These eigenvectors induce orthonormal vectors $\psi_{j,\tau,N}\in\mathcal{H}_{\tau}$, given by

$$\psi_{j,\tau,N} = \frac{1}{\sqrt{\lambda_{j,\tau,N}}} K_{\tau,N} \phi_{j,N}.$$

For $l \in \mathbb{N}$ such that $2l+1 \leq N$, define the subspace $\Psi_{\tau,l,N} = \operatorname{span}\{\psi_{0,\tau,N},\ldots,\psi_{2l,\tau,N}\} \subseteq \mathcal{H}_{\tau,N}$. We approximate the regularized Koopman generator $W_{\tau} \colon D(W_{\tau}) \to \mathcal{H}_{\tau,N}$ by skew-adjoint, finite-rank operators $W_{\tau,l,N} \colon \mathcal{H}_{\tau,N} \to \mathcal{H}_{\tau,N}$ with ran $W_{\tau,l,N} \subseteq \Psi_{\tau,L,N}$. These operators have associated eigendecomposition

$$W_{\tau,l,N}\zeta_{j,\tau,l,N} = i\omega_{j,\tau,l,N}\zeta_{j,\tau,l,N}, \quad j \in \{-l,\ldots,l\},$$

where $\omega_{j,\tau,l,N}$ are real eigenfrequencies satisfying $\omega_{-j,\tau,l,N} = -\omega_{j,\tau,l,N}$, and $\zeta_{j,\tau,l,N} \in \mathcal{H}_{\tau,l,N}$ are normalized eigenvectors that form complex-conjugate pairs, $\zeta_{-j,\tau,l,N} = \zeta_{j,\tau,l,N}^*$, with $\zeta_{0,\tau,l,N} = 1_X$. Computationally, we obtain the eigenpairs $(\omega_{j,\tau,l,N},\zeta_{j,\tau,l,N})$ by solving an $l \times l$ matrix eigenvalue problem giving eigenpairs $(\omega_{j,\tau,l,N},\xi_{j,\tau,l,N})$ for an approximate generator $V_{\tau,l,N} \colon H_N \to H_N$, and then mapping the eigenvectors $\xi_{j,\tau,l,N} \in H_N$ to $\zeta_{j,\tau,l,N} = T_{\tau,N}\xi_{j,\tau,l,N}$ using the isometry $T_{\tau,N} \colon H_N \to \mathcal{H}_{\tau,N}$ defined via the polar decomposition of $K_{\tau,N}$ as in (7). We order the computed eigenpairs in order of increasing Dirichlet energy as in section 3.2. Further details can be found in appendix B. The results converge to the eigendecomposition of W_{τ} in an iterated limit of $l \to \infty$ after $N \to \infty$.

Having obtained $W_{\tau,l,N}$ and its eigendecomposition, the rest of the data-driven scheme proceeds in a very similar manner to what was described in sections 4.2–4.7:

- We build the weighted symmetric Fock space $F_w(\mathcal{H}_{\tau,N})$ analogously to $F_w(\mathcal{H}_{\tau})$, and lift the unitaries $U^t_{\tau,l,N} \colon \mathcal{H}_{\tau,N} \to \mathcal{H}_{\tau,N}$ to unitaries $\tilde{U}^t_{\tau,l,N} \colon F_w(\mathcal{H}_{\tau,N}) \to F_w(\mathcal{H}_{\tau,N})$ that act multiplicatively with respect to the symmetric tensor product. Embedded in the spectrum $\sigma(F_w(\mathcal{H}_{\tau,N}))$ is a union of tori $\mathbb{S}_{\tau,N}$ analogous to \mathbb{S}_{τ} , in which the state space X is mapped by means of a feature map $\hat{\varphi}_{\tau,N} \colon X \to \sigma(F_w(\mathcal{H}_{\tau,N}))$.
- We define a rotation system $R_{\tau,l,N}^t$: $\sigma(F_w(\mathcal{H}_{\tau,N})) \to \sigma(F_w(\mathcal{H}_{\tau,N}))$ on the spectrum of $F_w(\mathcal{H}_{\tau,N})$ that is induced by the unitaries $\tilde{U}_{\tau,l,N}^t$, i.e., $R_{\tau,l,N}^t(\chi) = \chi \circ \tilde{U}_{\tau,l,N}^t$. This will serve as an approximation of the rotation system R_{τ}^t on $\sigma(F_w(\mathcal{H}_{\tau}))$.
- Letting $\hat{F}_w(\mathcal{H}_{\tau,N}) \subset C(\sigma(F_w(\mathcal{H}_{\tau,N})))$ be the Gelfand image of $F_w(\mathcal{H}_{\tau,N})$ as an RKHA on the spectrum of $F_w(\mathcal{H}_{\tau,N})$, we embed observables $f \in C(M)$ into elements $\hat{g}_{m,\tau,N} \in \hat{F}_w(\mathcal{H}_{\tau,N})$, $m \in \mathbb{N}$, by means of data-driven integral operators $\hat{\mathcal{K}}_{m,N} \colon H_N \to \hat{F}_w(\mathcal{H}_{\tau,N})$. The restrictions of these functions to d-dimensional tori in $\mathbb{S}_{\tau,N}$, $d \leq \lfloor (N-1)/2 \rfloor$, are degree-m polynomials in the coordinates z_1, \ldots, z_d and their complex conjugates. These polynomials evolve under the rotation system $R_{\tau,l,N}^t$ analogously to $\hat{g}_{m,\tau}^{(t)}$ in (31).
- Proceeding similarly to the construction of $f_{m,\tau,d}^{(t)}$ in (32), we approximate the Koopman evolution $U^t f$ by means of the function

$$f_{m,\tau,d,l,N}^{(t)} = \frac{\sum_{j \in \mathbb{J}_{d,m}} {m \choose j_{-d} \cdots j_d} C_{j,\tau,d,l,N}^{(g)} \prod_{r=1}^d e^{i(j_{-r} - j_r)\omega_{r,\tau,l,N} t} \zeta_{r,\tau,l,N}^{j_{-r} - j_r}}{\sum_{j \in \mathbb{J}_{d,m}} {m \choose j_{-d} \cdots j_d} C_{j,\tau,d,l,N}^{(h)} \prod_{r=1}^d e^{i(j_{-r} - j_r)\omega_{r,\tau,l,N} t} \zeta_{r,\tau,l,N}^{j_{-r} - j_r}} \in \tilde{\mathcal{H}}_{\tau,N}.$$
 (36)

Here, $\tilde{\mathcal{H}}_{\tau,N} \subset C(X)$ is the RKHA on X defined analogously to $\tilde{\mathcal{H}}_{\tau}$ using the kernel $k_{\tau,N}$. Moreover, $C_{j,\tau,d,l,N}^{(g)}$ and $C_{j,\tau,d,l,N}^{(h)}$ are moment coefficients computed using similar formulas to (30), replacing the invariant measure μ by the sampling measure μ_N and the regularized generator eigenfunctions $\zeta_{j,\tau}$ by their data-driven counterparts $\zeta_{j,\tau,l,N}$.

The final approximation $f_{m,\tau,d,l,N}^{(t)}$ converges to $U^t f$ in the norm of H in the iterated limits of $\lim_{t\to 0^+} \lim_{t\to \infty} \lim_{t\to \infty} \lim_{t\to \infty} \lim_{t\to \infty} t$, taken in that order.

8. Numerical experiments

We apply the second quantization framework presented in sections 2–7 to two measure-preserving dynamical systems: a Stepanoff flow on \mathbb{T}^2 [62] and the L63 system on \mathbb{R}^3 . Both systems exhibit dynamical complexity from an operator-theoretic standpoint: The Stepanoff flow is a topologically weak-mixing system that is characterized by absence of continuous nonconstant Koopman functions, and the L63 system is measure-theoretically mixing [58] with an associated continuous spectrum of the Koopman operator on H. A primary difference between the two examples is that the ergodic invariant measure μ is the Haar probability measure on \mathbb{T}^2 (and therefore has a smooth density in local coordinates), whereas in the L63 case μ is an SRB measure supported on a fractal set of zero Lebesgue measure (the Lorenz attractor) [72].

Both systems satisfy the assumptions laid out in section 2.1: For the Stepanoff flow, the state space X, forward-invariant compact manifold M, and support of the invariant measure μ all coincide, supp $\mu = M = X = \mathbb{T}^2$. For the L63 system, we have supp $\mu \subset M \subset X = \mathbb{R}^3$ where supp μ is the Lorenz attractor and M an absorbing ball containing the attractor [52].

The Stepanoff flow was also employed in numerical experiments in [35] using the tensor network approximation scheme summarized in section 3.4. Koopman spectral computations for the two systems using the methods outlined in section 3.2 and appendices A and B were performed in [40].

8.1. **Experimental procedure.** We generate (unobserved) states $x_0, x_1, \ldots, x_{N-1} \in X$ and associated training data $y_n = F(x_n)$ in a Euclidean data space Y using a smooth observation map $F: X \to Y$. In the case of the Stepanoff flow, the states x_n are sampled on a uniform grid on \mathbb{T}^2 , and F is the flat embedding of the 2-torus into $Y = \mathbb{R}^4$,

$$F(x) = (\cos x^1, \sin x^1, \cos x^2, \sin x^2), \quad x = (x^1, x^2), \quad x^1, x^2 \in [0, 2\pi).$$

The observation map for the L63 system is the identity map on $X = \mathbb{R}^3$.

We choose a prediction observable $f \in C(X)$ for each system and consider that the values $f(x_n) \in \mathbb{R}$ on the training states are known. The prediction observable for the Stepanoff experiments is a von Mises probability density function,

$$f(x) = \frac{e^{-\gamma(\cos x^1 + \cos x^2)}}{I_0^2(\gamma)}, \quad x = (x^1, x^2) \in [0, 2\pi)^2, \tag{37}$$

where $\gamma > 0$ is a concentration parameter and $I_0 \in C^{\infty}(\mathbb{R})$ the modified Bessel function of the first kind of order 0. For the L63 experiments, f is the component function for the first component component of the state vector,

$$f(x) = x^1, \quad x = (x^1, x^2, x^3) \in \mathbb{R}^3.$$
 (38)

We compute approximate Koopman eigenfunctions $\zeta_{j,\tau,l,N} \in \mathcal{H}_{\tau,N}$ and corresponding eigenfrequencies $\omega_{j,\tau,l,N} \in \mathbb{R}$ by processing the training data y_n with the Koopman spectral approximation technique from [40], summarized in sections 3.2 and 7 and appendix B. Note that this scheme makes use of equations of motion through the dynamical vector field \vec{V} (given in (39) and (40) below for the two systems). In both of the Stepanoff and L63 experiments, we build the RKHS $\mathcal{H}_{\tau,N}$ using a bistochastic kernel normalization, described in appendix A. The associated unitary evolution $U_{\tau,l,N}^t$ defines a rotation system on the tori $\mathbb{T}_{\tau,d,x} \subset \sigma(F_w(\mathcal{H}_{\tau,N}))$ that we will use for

4

Stepanoff flow Lorenz 63 system $(\beta, \rho, \sigma) = (8/3, 28, 10)$ $\alpha = \sqrt{20}$ System parameters Dimension of state space XDimension of data space Y4 $256 \times 256 = 65.536$ Number of training samples N80,000 Training sampling timestep Δt N/A5.0Prediction observable ffirst state vector component von Mises, $\gamma = 1$ 10^{-4} 5×10^{-7} Regularization parameter τ Koopman approximation space dimension l4096 2048 Torus dimension d50 50 Smoothing kernel bandwidth ε 0.05 0.1

4

Table 1. Dataset attributes and numerical parameters used in the Stepanoff flow and L63 experiments.

our dynamical prediction experiments. In all cases, we use the torus dimension d=50. We select the tori $\mathbb{T}_{\tau,d,x}$ using the complex-conjugate pairs of eigenfunctions $\zeta_{j,\tau,l,N}$ with the d largest L^2 projection coefficient amplitudes $c_j := |\langle \xi_{j,z,\tau,l,N}, f \rangle_{H_N}|$ with respect to the eigenbasis $\xi_{j,z,\tau,l,N}$ (that is, the sequence $a^{(d)}(x)$ parameterizing $\mathbb{T}_{\tau,d,x} \equiv T_{\tau,a^{(d)}(x)}$ has nonzero elements only at the positions corresponding to the constant eigenfunction $\zeta_{0,\tau,l,N} = 1_X$ and eigenfunctions $\zeta_{j,\tau,k,N}$ with the 50 largest corresponding values of c_j , for a total of 101 nonzero elements). Note that the weight function w cancels from our final formula (36) for the time-evolved prediction observable, so we do not need to specify it explicitly for numerical computations.

Fock space grading m

Next, to embed the prediction observable f into the Fock space $F_w(\mathcal{H}_{\tau,N})$, we use the grading parameter m=4 and a Gaussian smoothing kernel κ from (24) induced by the Euclidean metric $d(x,x') = ||F(x) - F(x')||_2$ on X. We compute smoothed eigenfunctions $\varrho_{j,\tau,l,N}$ and the associated moments $C_{j,\tau,d,l,N}^{(g)}$ and $C_{j,\tau,d,l,N}^{(h)}$ using data-driven analogs of (29) and (30). Note that the moment computations utilize the values $\zeta_{j,\tau,l,N}(x_n)$ and $f(x_n)$ of the eigenfunctions and prediction observable, respectively, on the states x_n . For our chosen parameter values of d=50 and m=4 the number of moments is $\binom{m+2d}{2d} = 4,598,126$. We then build the prediction function $f_{m,\tau,d,l,N}^{(t)}$ using (36). A listing of the dataset attributes and numerical parameters used in the Stepanoff and L63 experiments is included in table 1.

We test the accuracy of the scheme by comparing $f_{m,\tau,d,l,N}^{(t)}(x_n)$ with a high-fidelity numerical approximation of the true Koopman evolution $U^t f(x_n) = f(\Phi^t(x_n))$ based on a ordinary differential equation solver to approximate the flow map $\Phi^t(x_n)$. In addition, we compare the Fock space approach with a conventional RKHS approximation [22], utilizing the same eigenfunctions and eigenfrequencies as those generating the d-dimensional rotation system on $\mathbb{T}_{\tau,d,x}$. Hereafter, we refer to this method as classical approximation. The associated prediction function $f_{\mathrm{cl},\tau,d,l,N}^{(t)} \in \mathcal{H}_{\tau}$ is defined as

$$f_{\text{cl},\tau,d,l,N}^{(t)} = \sum_{j=0}^{2d} C_{j,\tau,l,N}^{(f)} e^{i\omega_{j,\tau,l,N}} \zeta_{j,\tau,l,N}, \quad C_{j,\tau,l,N}^{(f)} = \int_X \overline{\zeta_{j,\tau,l,N}} f \, d\mu_N.$$

The convergence properties of $f_{\mathrm{cl},\tau,d,l,N}^{(t)}$ are analogous to those of $f_{m,\tau,d,l,N}^{(t)}$; i.e., we have convergence in the norm of H in the iterated limit $\lim_{\tau\to 0^+} \lim_{d\to\infty} \lim_{l\to\infty} \lim_{N\to\infty}$; see [22] for further details. Let $f^{(t)}$ stand for either $f_{\mathrm{cl},\tau,d,l,N}^{(t)}$ or $f_{m,\tau,d,l,N}^{(t)}$. To quantitatively assess prediction skill, we compute normalized root mean square error (RMSE) and anomaly correlation (AC) scores, defined

as

$$RMSE(t) = \frac{\|f^{(t)} - U^t f\|_{H_N}}{\|U^t f\|_{H_N}}, \quad AC(t) = \frac{\langle g^{(t)}, h^{(t)} \rangle_{H_N}}{\|g^{(t)}\|_{H_N} \|h^{(t)}\|_{H_N}}$$

where $g^{(t)}, h^{(t)} \in H_N$ are the predicted and true anomalies relative to the empirical mean,

$$g^{(t)} = f^{(t)} - \int_X f^{(t)} d\mu_N, \quad h^{(t)} = U^t f - \int_X U^t f d\mu_N,$$

respectively. By construction, AC(t) lies in the interval [0,1]. Values of RMSE(t) close to 0 and AC(t) close to 1 indicate high prediction skill.

For the remainder of section 8, we will use the shorthand notations $f_{\rm cl}^{(t)} \equiv f_{{\rm cl},\tau,d,l,N}^{(t)}$ and $f_{\rm Fock}^{(t)} \equiv f_{m,\tau,d,l,N}^{(t)}$ for the classical and Fock space approximations, respectively.

8.2. **Stepanoff flow.** The dynamical vector field $\vec{V}: \mathbb{T}^2 \to T\mathbb{T}^2$ generating the Stepanoff flow has the coordinate representation $\vec{V}(x) = (V^1(x), V^2(x))$, where

$$V^{1}(x) = V^{2}(x) + (1 - \alpha)(1 - \cos x^{2}), \quad V^{2}(x) = \alpha(1 - \cos(x^{1} - x^{2})), \tag{39}$$

 $x=(x^1,x^2)\in[0,2\pi)^2$, and α is a real parameter. This vector field has zero divergence with respect to the Haar measure μ ,

$$\operatorname{div}_{\mu} \vec{V} = \frac{\partial V^{1}}{\partial x^{1}} + \frac{\partial V^{2}}{\partial x^{2}} = 0,$$

which implies that μ is an invariant measure under the associated flow Φ^t . In addition, the system has a fixed point at x = 0, $\vec{V}(0) = 0$. In [62] it is shown that the normalized Haar measure is the unique invariant Borel probability measure of this flow that assigns measure 0 to the singleton set $\{0\} \subset \mathbb{T}^2$ containing the fixed point.

Since any continuous, nonconstant Koopman eigenfunction of an ergodic flow induces a semiconjugacy with a circle rotation of nonzero frequency, the existence of the fixed point at x = 0implies that the system has no continuous Koopman eigenfunctions; i.e., it is topologically weakmixing. In fact, [62] shows that when α is irrational the Stepanoff flow is topologically conjugate to a time-reparameterized irrational rotation on \mathbb{T}^2 with frequency parameters $(1, \alpha)$, for a timereparameterization function that vanishes at x = 0. While, to our knowledge, there are no results in the literature on the measure-theoretic mixing properties of Stepanoff flows, the existence of fixed points is a necessary condition for a smooth ergodic flow on \mathbb{T}^2 to be (strong) mixing [49] and the vector field (39) meets at least that requirement.

Here, we work with the parameter value $\alpha = \sqrt{20}$. The resulting dynamical vector field is depicted as a quiver plot in (39). As prediction observable, we consider the von Mises density function f from (37) with concentration parameter $\gamma = 1$. This is shown as a heat map on \mathbb{T}^2 in the top-left panel of fig. 2. Note that f has a peak at (π, π) which is well-separated from the fixed point at (0,0). The remaining panels in the left-hand column of fig. 2 show the Koopman evolution $U^t f$ at times $t \in \{0.5, 1, 2, 4\}$. As t increases, the initially radially symmetric structure of f becomes stretched and folded, developing sharp gradients that are suggestive of sensitive dependence on initial conditions. The fact that f is strictly positive makes this observable suitable for testing the ability of approximation schemes to preserve positivity.

Our training data is sampled on a uniform 256×256 grid $\{x_n\} \subset \mathbb{T}^2$, amounting to a total of $N=65,\!536$ points. We build the RKHS $\mathcal{H}_{\tau,N}$ and the l-dimensional approximation space $\Psi_{\tau,l,N} \subset \mathcal{H}_{\tau,N}$ using the regularization parameter $\tau=10^{-4}$ and dimension parameter l=4096 (see section 7). Representative eigenfrequencies $\omega_{j,\tau,l,N}$ and the corresponding eigenfunctions are shown in figs. 3 and 4, respectively. Notice the characteristic "S-shaped" pattern of the eigenfunction level sets that is broadly aligned with the dynamical vector field (see again fig. 1). Also noteworthy in the eigenfunction plots is the oscillatory nature of the eigenfunctions near the fixed point. This




FIGURE 1. Generating vector field (39) of the Stepanoff flow for $\alpha = \sqrt{20}$.

behavior is qualitatively consistent with the slowing down of dynamical trajectories near the fixed point, which needs to be compensated by small-scale spatial oscillations in order to maintain a consistent evolution with the periodic phase evolution implied by the corresponding eigenfrequency, $\zeta_{j,\tau,l,N}(\Phi^t(x)) \approx e^{i\omega_{j,\tau,l,N}}\zeta_{j,\tau,l,N}(x)$.

Next, in the center column of fig. 2 we show the evolution $f_{\rm cl}^{(t)}$ under the classical Koopman approximation obtained for d=50. The evolution times are the same as those shown for the true evolution in the left-hand column (i.e., $t \in \{0, 0.5, 1, 2, 4\}$) and $f_{\rm cl}^{(t)}$ is plotted at the gridpoints x_n . Figure 5(center column) shows the approximation error of $f_{\rm cl}^{(t)}$ relative to the true evolution $U^t f$ at the gridpoints.

Starting from time t=0, it is clear that $f_{\rm cl}^{(0)}$ carries imprints of the characteristic S-shaped spatial structure of the eigenfunctions $\zeta_{j,\tau,d,l,N}$ employed in the approximation (see fig. 4), and as a result provides a poor reconstruction of the radial symmetry of the von Mises density function. Moreover, $f_{\rm cl}^{(0)}$ exhibits strongly oscillatory behavior near the fixed point with significant departures to negative values. This is a manifestation of the fact that, being an orthogonal subspace projection method, the classical approximation is not positivity-preserving. At later times, the classical approximation appears to track some of the large-scale qualitative features of the true evolution such as the large-amplitude bands seen at t=0.5,1 due to stretching of the originally radial von Mises density function by the Stepanoff flow. However, $f_{\rm cl}^{(t)}$ continues to exhibit significant oscillatory behavior and departures to negative values, particularly in regions near the fixed point.

We now consider the Fock space approximation $f_{\rm Fock}^{(t)}$ for torus dimension d=50, grading m=4,

We now consider the Fock space approximation $f_{\text{Fock}}^{(t)}$ for torus dimension d=50, grading m=4, and smoothing kernel bandwidth parameter $\varepsilon=0.05$. As noted in section 8.1, with this choice of parameters the approximation space dimension is amplified to 4,598,126 so we expect higher reconstruction accuracy than the 101-dimensional classical approximation. Moreover, with m being even, the Fock space approximation is positivity preserving. Indeed, as is evident from the top-right panels in figs. 2 and 5, the Fock space scheme leads to significant increase of reconstruction accuracy at t=0 compared to the classical approximation. While imprints from the eigenfunctions are still visible, the radial symmetry of the von Mises density is better represented, and the small-scale oscillations and negative values near the fixed point are eliminated.

Figure 6(left) shows RMSE and AC scores for the Stepanoff flow experiments, computed for the same prediction times t as in figs. 2 and 5. As the evolution time t increases, the error in $f_{\rm Fock}^{(t)}$ remains consistently lower than $f_{\rm cl}^{(t)}$ at least through $t \simeq 2.2$ and $t \simeq 3$ for the RMSE and




FIGURE 2. Time evolution of the von Mises density function from (37) with $\gamma=1$ under the Stepanoff flow $(U^tf;$ left column), the classical approximation based on 2d+1 eigenfunctions $(f_{\rm cl}^{(t)};$ center column), and the Fock space approximation for torus dimension d=50 and degree m=4 $(f_{\rm Fock}^{(t)};$ right column). Rows from top to bottom show snapshots at the evolution times t=0,0.5,1,2,4, respectively.




FIGURE 3. Spectrum of the regularized generator $W_{\tau,l,N}$ for the Stepanoff flow (left) and L63 system (right). Eigenfrequencies $\omega_{j,\tau,l,N}$ are plotted versus their corresponding Dirichlet energies from the RKHS \mathcal{H}_{τ} . The plotted points are colored by the logarithms of the expansion coefficient amplitudes $|\langle f, \xi_{j,\tau,l,N} \rangle_{H_N}|$ of the prediction observable f in the eigenbasis $\xi_{j,\tau,l,N}$.

AC metrics, respectively. At later times, the error in $f_{\rm Fock}^{(t)}$ is somewhat higher than the classical approximation and appears to be predominantly concentrated in narrow bands where the true evolution U^tf exhibits sharp gradients. As illustrated by the t=4 results in the bottom row of fig. 5, the maximal pointwise error in $f_{\rm Fock}^{(t)}$ eventually outgrows that in $f_{\rm cl}^{(t)}$.

8.3. Lorenz 63 system. The L63 system on \mathbb{R}^3 is generated by the vector field $\vec{V}: \mathbb{R}^3 \to T\mathbb{R}^3 \cong \mathbb{R}^3$, where

$$\vec{V}(x) = (V^1(x), V^2(x), V^3(x)) = (-\sigma(x^2 - x^1), x^1(\rho - x^3) - x^2, x^1x^2 - \beta x^3)$$
 (40)

and $x=(x^1,x^2,x^3)$. We use the standard parameter values $\beta=8/3,\ \rho=28,$ and $\sigma=10,$ leading to the chaotic Lorenz "butterfly" attractor.

The left-hand column in fig. 7 shows the evolution $U^t f$ of first state vector component (38) for evolution times $t \in \{0, 0.5, 1, 2, 4\}$. Due to chaotic mixing between the two lobes of the attractor, as t increases $U^t f$ exhibits increasingly finer-scale oscillations. We recall the approximate numerical value $\Lambda \simeq 0.91$ for the positive exponent of the standard L63 system [65], which implies that the pointwise predictability horizon of f is of order $1/\Lambda \simeq 1.10$ model time units.

The training dataset used in our L63 experiments consists of N=80,000 samples x_n , taken at a sampling interval of $\Delta t=5.0$ model time units along a numerical L63 trajectory starting from an arbitrary initial condition x_0 near the Lorenz attractor. The sampling interval Δt was chosen to be larger than the Lyapunov timescale $1/\Lambda$ to reduce correlations between the samples. This should aid in the convergence of the sampling measure μ_N to the invariant measure μ . As noted in appendix B, we are able to work with large sampling intervals by making use of known equations of motion to evaluate the action of the Koopman generator on functions. This is in contrast to, e.g., finite-difference methods that require small Δt to yield accurate approximations of the generator [22,34].

We compute approximate Koopman eigenfrequencies $\omega_{j,\tau,l,N}$ and eigenfunctions $\zeta_{j,\tau,l,N}$ similarly to the Stepanoff flow experiments, setting the regularization parameter and approximation space dimension to $\tau = 5 \times 10^{-7}$ and l = 2048, respectively. Eigenfrequency spectra and representative corresponding eigenfunctions are displayed in figs. 3 and 8, respectively. Using the eigenpairs $(\omega_{j,\tau,l,N}, \zeta_{j,\tau,l,N})$ and the samples $f(x_n)$ of the prediction observable, we build classical and Fock




FIGURE 4. Real and imaginary parts of representative eigenfunctions $\zeta_{j,\tau,l,N}$ for the Stepanoff flow. The eigenfunctions shown have index j=1,8,97 with respect to the Dirichlet energy ordering, and are members of the complex-conjugate (non-constant) eigenfunction pairs with the 7th, 1st, 5th largest projection amplitudes $|\langle \xi_{j,\tau,l,N}, f \rangle_{H_N}|$, respectively.

space approximations $f_{\rm cl}^{(t)}$ and $f_{\rm Fock}^{(t)}$, respectively, again via a similar procedure to that used for the Stepanoff flow. The torus dimension, Fock space grading, and kernel bandwidth parameters are d=50, m=4, and $\varepsilon=0.1$, respectively. Scatterplots of $f_{\rm cl}^{(t)}(x_n)$ and $f_{\rm Fock}^{(t)}(x_n)$ on the training data are displayed in the center and right columns of fig. 7, respectively, and the corresponding errors relative to the true evolution $U^t f$ are shown in fig. 9.

The general conclusions stemming from these results are broadly similar to what was previously observed for the Stepanoff flow. That is, amplification to the Fock space markedly improves prediction skill over a classical approximation utilizing the same number of eigenfunctions. At initialization time, t=0 (top rows in figs. 7 and 9), the Fock space reconstruction $f_{\rm Fock}^{(0)}$ captures more accurately the linear dependence of f on the x^1 component of the state vector than the classical reconstruction $f_{\rm cl}^{(0)}$. At later times, $f_{\rm Fock}^{(t)}$ provides a more accurate representation of the oscillations seen in the true evolution $U^t f$ along the radial direction in each lobe of the attractor due to chaotic mixing. Of course, since both methods utilize approximation spaces of fixed, finite dimension, they eventually fail to capture the small-scale spatial oscillatory behavior of $U^t f$ under the true dynamics. Still, even at t=4 (bottom rows in figs. 7 and 9), $f_{\rm Fock}^{(t)}$ is arguably closer to




FIGURE 5. Error in the classical and Fock space approximations from fig. 2 (center and right columns, respectively) relative to the true Stepanoff evolution. The true evolution is plotted in the left column for reference.




FIGURE 6. Normalized RMSE (top row) and anomaly correlation scores (bottom row) as a function of prediction time t for the Stepanoff (left column) and L63 experiments (right column).

the spatial structure of U^tf than $f_{\rm cl}^{(t)}$. Figure 6(right) shows the RMSE and AC scores for the same prediction times as in figs. 7 and 9. Aside from the approximate interval $t \in [0.5, 1.5]$, where the computed RMSE score is slightly larger for $f_{\rm Fock}^{(t)}$ than $f_{\rm cl}^{(t)}$, the Fock space approximation outperforms the classical approximation with respect to both the RMSE and AC metrics, particularly at early times, $t \lesssim 0.5$.

9. Discussion

Combining techniques from many-body quantum theory, RKHS theory, and Koopman operator theory, we have developed a framework for consistently approximating measure-preserving ergodic flows as infinite-dimensional topological rotation systems. A principal element of our approach is a family of weighted symmetric Fock spaces, $F_w(\mathcal{H}_\tau)$, generated from a 1-parameter family of RKHSs \mathcal{H}_τ , $\tau > 0$, and endowed with coalgebra structure and commutative Banach algebra with respect to the symmetric tensor product. On the RKHSs \mathcal{H}_τ , we build skew-adjoint, diagonalizable approximations W_τ of the Koopman generator on L^2 (which is generally a non-diagonalizable operator with non-trivial continuous spectrum). We dilate the unitary evolution groups generated by W_τ into infinite-dimensional rotation systems on the Banach algebra spectra of $F_w(\mathcal{H}_\tau)$, equipped with the weak-* topology making them compact Hausdorff spaces.

On the one hand, this construction realizes regularized approximations $e^{tW_{\tau}}$ of Koopman operators (which are not composition operators by flows on the original state space) as topological rotation systems on the spectrum of an algebra, with associated Koopman composition operators acting on continuous functions on the spectrum. This construction is therefore similar in spirit to the Halmos-von Neumann theorem that realizes a measure-preserving ergodic flow with pure point




FIGURE 7. Time evolution of the x component of the L63 state vector under the true dynamics (left column), a classical Koopman eigenfunction approximation using 2d+1 eigenfunctions (center column), and a Fock space approximation for torus dimension d=50 and degree m=4. Rows from top to bottom show snapshots at the evolution times t=0,0.5,1,2,4, respectively.




FIGURE 8. Real and imaginary parts of representative eigenfunctions $\zeta_{j,\tau,l,N}$ for the L63 system. The eigenfunctions shown have index j=1,5,22 with respect to the Dirichlet energy ordering, and are members of the complex-conjugate (nonconstant) eigenfunction pairs with the 1st, 2nd, and 6th largest projection amplitudes $|\langle \xi_{j,\tau,l,N}, f \rangle_{H_N}|$, respectively.

spectrum (i.e., diagonalizable generator) as a topological rotation system on the spectrum of an abelian C^* -algebra generated by the eigenspaces of the Koopman operator.

Simultaneously, the second quantization approach provides a flexible framework for approximating the Koopman evolution of observables of measure-preserving ergodic flows. The resulting approximations are positivity-preserving, amenable to data-driven numerical implementation, and consistent with the true Koopman evolution in appropriate asymptotic limits.

In more detail, we have shown that the spectrum $\sigma(F_w(\mathcal{H}_\tau))$ can be partitioned into a collection of tori (of potentially infinite dimension) that are invariant under the rotational dynamics induced by W_τ . Moreover, $F_w(\mathcal{H}_\tau)$ is isomorphic to a reproducing kernel Hilbert algebra (RKHA) $\hat{F}_w(\mathcal{H}_\tau)$ of continuous functions (i.e., an RKHS that is simultaneously a coalgebra and a Banach algebra with respect to pointwise function multiplication) on appropriately chosen such tori. We developed a scheme for lifting continuous observables of the original dynamical system to polynomials of arbitrarily large degree m in the Fourier bases of the spectral tori. A key aspect of this scheme is that it allows to systematically generate high-dimensional approximation spaces from tensor products of a given collection of approximate Koopman eigenfunctions (eigenfunctions of W_τ),




FIGURE 9. Errors in the classical and Fock space approximations from fig. 2 (center and right columns, respectively) relative to the true L63 evolution. The true evolution is plotted in the left column for reference.

capturing information from the product structure of these eigenfunctions. Conventional subspace projection methods for Koopman operator can be viewed as m = 1 versions of this scheme.

We numerically applied our approach to two examples of non-integrable measure-preserving ergodic systems: a Stepanoff flow on \mathbb{T}^2 as an example with smooth invariant measure and topological weak mixing (absence of continuous nonconstant Koopman eigenfunctions) and the L63 system on \mathbb{R}^3 as an example with invariant measure supported on a fractal attractor and measure-theoretic mixing. Both examples demonstrated improved prediction skill resulting from the Fock space construction over conventional subspace projection methods for Koopman operator approximation.

Previous work [35], developed tensor network approximation techniques for Koopman and transfer operators that similarly yield algebraic amplification of regularized Koopman operators to Fock spaces. From a high-level standpoint the tensor network approach can be viewed as being "dual" to the second quantization scheme presented in this paper: the former is based on a lift of states to the Fock space whereas the latter focuses on a lift of observables. An advantageous aspect of the tensor network scheme is high computational efficiency when implemented classically. On the other hand, the second quantization scheme in this paper offers greater flexibility as it can be implemented for larger families of RKHSs \mathcal{H}_{τ} , and it also yields a topological model that provides an arena for different types of approximation schemes (the polynomial lifts employed in this paper being just an example). Another noteworthy difference between the two methods is that the tensor network approach employs the full Fock space without algebra structure, whereas a centerpiece of the approach presented in this paper is a symmetric weighted Fock space with coalgebra and abelian Banach algebra structure.

In terms of future work, a natural next step would be to develop gate-based implementations of our scheme running on quantum computing platforms. One possibility in that direction would be to employ the quantum algorithm for pure point spectrum systems developed in [37] as a black box algorithm to simulate the evolution of the polynomial observables under the rotational dynamics on the tori within $\sigma(F_w(\mathcal{H}_\tau))$. Essentially, this would involve defining an RKHA $\mathfrak A$ on each torus of interest, and using $\mathfrak A$ to build a quantum mechanical representation of the rotation system as described in section 3.1. A more direct (but likely more challenging) approach would be to build quantum computational algorithms on the RKHAs $\hat{F}_w(\mathcal{H}_\tau)$ without invoking auxiliary algebras such as $\mathfrak A$. Such an approach would hinge upon availability of efficient algorithms for preparing quantum states associated with the RKHA feature vectors $\varphi_{\tau,d}(x)$ from section 4.7. Efforts in these directions are currently underway. In a broader context, we hope that this work will stimulate further research at the interface of ergodic theory and quantum theory.

Acknowledgments

Dimitrios Giannakis acknowledges support from the U.S. Department of Defense, Basic Research Office under Vannevar Bush Faculty Fellowship grant N00014-21-1-2946 and the U.S. Office of Naval Research under MURI grant N00014-19-1-242. Mohammad Javad Latifi Jebelli and Michael Montgomery were supported as postdoctoral fellows from these grants. Philipp Pfeffer is supported by the project no. P2018-02-001 "DeepTurb – Deep Learning in and of Turbulence" of the Carl Zeiss Foundation, Germany. Jörg Schumacher is supported by the European Union (ERC, MesoComp, 101052786). Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

APPENDIX A. MARKOV SEMIGROUP CONSTRUCTION

This appendix summarizes the construction of the 1-parameter family of Markov kernels $k_{\tau} \colon X \times X \to \mathbb{R}$ associated with the Markov semigroup $\{G_{\tau}\}_{{\tau} \geq 0}$ introduced in section 3.2. Our exposition follows closely [40], where we refer the reader for further details.

The starting point of the construction is a kernel function $k: X \times X \to \mathbb{R}_{>0}$ that is continuous, symmetric, strictly positive-valued, and integrally strictly positive-definite. The latter condition means [66] that

$$\int_{X \times X} k(x, y) d(\nu \times \nu)(x, y) > 0 \tag{41}$$

for every finite Borel measure ν . These conditions imply that k is strictly positive definite. In addition, $K_{\nu} : L^{2}(\nu) \to L^{2}(\nu)$ defined as

$$K_{\nu}f = \int_{X} k(\cdot, x) f(x) \, d\nu(x)$$

is a self-adjoint, strictly positive integral operator of trace class that preserves positive functions. In this work, we are interested in integral operators K_{ν} associated with the invariant measure of the dynamical system, $\nu = \mu$, and the corresponding sampling measures on dynamical trajectories, $\nu = \mu_N$.

When X can be continuously injected into \mathbb{R}^d under a map $F: X \to \mathbb{R}^d$, it inherits a metric $d: X \times X \to \mathbb{R}$ from the Euclidean metric, $d(x,y) = ||F(x) - F(y)||_2$. Then, a prototypical example satisfying (41) is the Gaussian radial basis function (RBF) kernel,

$$k_{\varepsilon}^{\text{RBF}}(x,y) = \exp\left(-\frac{d^2(x,y)}{\varepsilon^2}\right),$$
 (42)

where $\varepsilon > 0$ is a bandwidth parameter. Given a continuous, strictly positive function $\rho: X \to \mathbb{R}_{>0}$, a generalization of (42) is the so-called variable-bandwidth or self-tuning Gaussian kernel [8,76],

$$k_{\varepsilon}^{\text{vb}}(x,y) = \exp\left(-\frac{d^2(x,y)}{\varepsilon^2 \rho(x)\rho(y)}\right).$$
 (43)

In the numerical experiments of section 8, we use (43) with the bandwidth function

$$\rho(x) = \left(\int_X k_{\tilde{\varepsilon}}(x, y) \, d\nu(y)\right)^{-1/m_{\nu}}.$$

In the above, the bandwidth parameter $\tilde{\varepsilon}$ is independent from ε , and m_{ν} is a dimension parameter for the support of ν . In the data-driven computations of section 8, we determine ε and $\tilde{\varepsilon}$ automatically using a variant of a method proposed in [16]; see [7, Appendix A] and [34, Algorithm 1] for further details and pseudocode. The same method produces a numerical estimate of m_{ν} that we use in our computations.

Intuitively, the role of ρ in (43) is to decrease (increase) the effective bandwidth of the kernel in regions of data space with high (low) concentration of the data distribution, $F_*\nu$. This is particularly important in the case of the L63 experiments where $F_*\mu$ associated with the invariant measure μ is singular relative to the ambient Lebesgue measure on \mathbb{R}^3 .

Remark 12. Bandwidth functions ρ computed for the sampling measures μ_N are data-dependent, $\rho = \rho_N$, making the variable-bandwidth kernel $k_\varepsilon^{\rm vb}$ also data-dependent. In such cases, weak-* convergence (35) of μ_N in the large-data limit $(N \to \infty)$ implies uniform convergence of ρ_N on the compact set M to the bandwidth function ρ computed for the invariant measure μ (e.g., [73]). This implies in turn uniform convergence of $k_\varepsilon^{\rm vb}$ on $M \times M$, and the latter is sufficient to ensure convergence of our data-driven scheme in the large-data limit. For simplicity of exposition, in what follows we will not explicitly distinguish between data-dependent and data-independent kernels in our notation.

Assume that ν is a probability measure with compact support. Using a variant of the bistochastic kernel formulation proposed in [15], we normalize the kernel k to build a symmetric Markov kernel

 $p_{\nu} \colon X \times X \to \mathbb{R}_{>0}$, defined as

$$p_{\nu}(x,y) = \int_{X} \frac{k(x,z)k(z,y)}{d_{\nu}(x)q_{\nu}(z)d_{\nu}(y)} d\nu(z),$$

for $d_{\nu} = K_{\nu} 1_X$ and $q_{\nu} = K_{\nu} (1/d_{\nu})$. Clearly, $p_{\nu}(x,y) > 0$ and $p_{\nu}(x,y) = p_{\nu}(y,x)$, and the normalization $\int_X p_{\nu}(x,y) d\nu(y) = 1$ holds by construction. It can also be shown that p_{ν} is strictly positive definite, and thus has an associated RKHS $\mathcal{H} \subset C(X)$. It then follows that $G_{\nu} : L^2(\nu) \to L^2(\nu)$,

$$G_{\nu}f = \int_{X} p_{\nu}(\cdot, x) f(x) \, d\nu(x),$$

is a self-adjoint, strictly positive-definite, Markov operator of trace class. It can also be shown by strict positivity of p_{ν} that G_{ν} is ergodic; i.e., G_{ν} has a simple eigenvalue at 1 with corresponding eigenvector that is ν -a.e. equal to 1.

Let us consider an eigendecomposition of G_{ν} ,

$$G_{\nu}\phi_{j,\nu} = \lambda_{j,\nu}\phi_{j,\nu},$$

where the eigenvalues $1 = \lambda_{0,\nu} > \lambda_{1,\nu} \ge \lambda_{2,\nu}$ are strictly positive and have finite multiplicities, and the corresponding eigenvectors form an orthonormal basis $\{\phi_{j,\nu}\}_j$ of $L^2(\nu)$ with $\phi_{0,\nu} = 1_X$. Each eigenvector $\phi_{j,\nu}$ has a continuous representative $\varphi_{j,\nu} \in C(X)$, where

$$\varphi_{j,\nu} = \frac{1}{\lambda_{j,\nu}} \int_X p_{\nu}(\cdot, x) f(x) \, d\nu(x). \tag{44}$$

Defining $\psi_{j,\nu} = \sqrt{\lambda_{j,\nu}}\phi_{j,\nu}$, we have that $\{\psi_{j,\nu}\}$ is an orthonormal basis of \mathcal{H} . The RKHS \mathcal{H} also induces a Dirichlet energy functional, $\mathcal{E}_{\nu} \colon D(\mathcal{E}_{\nu}) \to \mathbb{R}_{\geq 0}$, providing a notion of regularity of $L^2(\nu)$ elements in a dense domain $D(\mathcal{E}_{\nu}) \subseteq L^2(\nu)$ with representatives in \mathcal{H} . For every $f \in D(\mathcal{E}_{\nu})$, we define

$$\mathcal{E}_{\nu}(f) = \sum_{i} \frac{|\langle \phi_{j,\nu}, f \rangle_{L^{2}(\nu)}|^{2}}{\lambda_{j}} - 1, \tag{45}$$

and we have $\mathcal{E}_{\nu}(f) \geq 0$ with equality iff f is constant ν -a.e.

Next, define the self-adjoint operator $\Delta_{\nu} \colon D(\Delta_{\nu}) \to L^2(\nu)$, $D(\Delta_{\nu}) \subseteq L^2(\nu)$, via the eigendecomposition

$$\Delta_{\nu}\phi_{j,\nu} = \eta_{j,\nu}\phi_{j,\nu}, \quad \eta_{j,\nu} = \frac{\lambda_{j,\nu}^{-1} - 1}{\lambda_{1,\nu}^{-1} - 1}.$$

This operator is positive, it annihilates constant functions, and it is unbounded whenever $L^2(\nu)$ is infinite-dimensional. Moreover, Δ_{ν} is normalized by convention so that its smallest nonzero eigenvalue $\eta_{1,\nu}$ is equal to 1. We interpret Δ_{ν} as a Laplace-type operator that generates a Markov semigroup $\{G_{\tau,\nu}\}_{\tau\geq 0}$ on $L^2(\nu)$ with $G_{\tau,\nu}=e^{-\tau\Delta_{\nu}}$. The following result is adapted from [22, Theorem 1].

Proposition 13. With notation and assumptions as above, $\{G_{\tau,\nu}\}_{\tau\geq 0}$ is a strongly continuous semigroup of self-adjoint, strictly positive, Markov operators on $L^2(\nu)$. Moreover, the following hold for $\tau>0$ and any compact set $M\subseteq X$ containing the support of ν :

(i) $G_{\tau,\nu}$ is a trace class integral operator induced by a continuous transition kernel $k_{\tau,\nu} \colon M \times M \to \mathbb{R}_{>0}$,

$$G_{\tau,\nu}f = \int_X k_{\tau,\nu}(\cdot,x)f(x)\,d\nu(x), \quad k_{\tau,\nu}(x,y) = \sum_j \lambda_{j,\tau,\nu}\varphi_{j,\nu}(x)\varphi_{j,\nu}(y), \quad \lambda_{j,\tau,\nu} = e^{-\tau\eta_{j,\nu}},$$

where the sum over j in the expression for $k_{\tau,\nu}(x,y)$ converges uniformly for $(x,y) \in M \times M$.

(ii) The kernel $k_{\tau,\nu}$ is strictly positive-definite, and has an associated RKHS $\mathcal{H}_{\tau,\nu} \subseteq C(M)$ whose restriction to $\operatorname{supp}(\nu)$ forms a dense subspace of $L^2(\nu)$.

- (iii) $\mathcal{H}_{\tau,\nu}$ is a subspace of \mathcal{H} and a subspace of $\mathcal{H}_{\tau',\nu}$ for every $\tau' < \tau$.
- (iv) If the kernel $k|_{M\times M}$ is C^r , then $\mathcal{H}_{\tau,\nu}$ is a subspace of $C^r(M)$.

Note that under Proposition 13 $G_{\tau,\nu}$ admits the factorization $G_{\tau,\nu} = K_{\tau,\nu}^* K_{\tau,\nu}$, where $K_{\tau,\nu} : L^2(\nu) \to \mathcal{H}_{\tau,\nu}$ is the integral operator with (integral) kernel $k_{\tau,\nu}$ and dense range in $\mathcal{H}_{\tau,\nu}(\nu)$, and the adjoint $K_{\tau,\nu}^*$ implements the inclusion map (see section 3.2). Moreover, $\{\psi_{i,\tau,\nu}\}_i$ with

$$\psi_{j,\tau,\nu} = \lambda_{j,\tau,\nu}^{-1/2} K_{\tau,\nu} \phi_{j,\nu} = \lambda_{j,\tau,\nu}^{1/2} \varphi_{j,\nu}$$

forms an orthonormal basis of $\mathcal{H}_{\tau,\nu}(\nu)$.

Setting $\nu = \mu$ leads to the kernel $k_{\tau} \equiv k_{\tau,\mu}$, RKHS $\mathcal{H}_{\tau} \equiv \mathcal{H}_{\tau,\mu}$, and integral operators $K_{\tau} \equiv K_{\tau,\mu}$ and $G_{\tau} \equiv G_{\tau,\mu}$ used in the main text. Properties (K1)–(K4) are then satisfied as a consequence of Proposition 13. In data-driven contexts we set $\nu = \mu_N$, leading to $k_{\tau,N} \equiv k_{\tau,\mu_N}$, $\mathcal{H}_{\tau,N} = \mathcal{H}_{\tau,\mu_N}$, $K_{\tau,N} \equiv K_{\tau,\mu_N}$, and $G_{\tau,N} \equiv G_{\tau,\mu_N}$. Spectral convergence of $G_{\tau,N}$ to G_{τ} as $N \to \infty$ occurs in the form of convergence of eigenvalues (including multiplicities), $\lim_{\tau \to 0^+} \lambda_{j,\tau,N} = \lambda_{j,\tau}$ where $\lambda_{j,\tau,N} \equiv \lambda_{j,\tau,\mu_N}$ and $\lambda_{j,\tau} \equiv \lambda_{j,\tau}$, and uniform convergence of the continuous representatives $\varphi_{j,\tau,N} \equiv \varphi_{j,\tau,\mu_N}$ to $\varphi_{j,\tau} \equiv \varphi_{j,\tau,\mu}$ on M for appropriately chosen eigenvectors $\phi_{j,\tau} \equiv \phi_{j,\tau,\mu}$ and $\phi_{j,\tau,N} = \varphi_{j,\tau,\mu_N}$ of G_{τ} and $G_{\tau,N}$, respectively.

APPENDIX B. SPECTRAL APPROXIMATION OF THE GENERATOR

In this appendix, we summarize the spectral approximation scheme for the Koopman generator from the paper [40], used here in the numerical experiments in section 8.

Let $\tilde{H} = \{f \in H : \int_X f \, d\mu = 0\}$ be the subspace of H consisting of zero-mean functions with respect to the invariant measure. By ergodicity, $\tilde{H} = \operatorname{span}\{1_X\}^{\perp}$ has codimension 1. Define also $H_V = D(V) \cap \tilde{H}$, and equip this space with the "graph" inner product

$$\langle f, g \rangle_{H_V} = \langle f, g \rangle_H + \langle V f, V g \rangle_H.$$

Since V is a closed operator mapping span $\{1_X\}$ to itself, $(H_V, \langle \cdot, \cdot \rangle_{H_V})$ is a Hilbert space. Next, for z > 0, define the bounded, continuous, antisymmetric function $q_z : i\mathbb{R} \to i\mathbb{R}$, where

$$q_z(i\omega) = \frac{i\omega}{z^2 + \omega^2}, \quad \operatorname{ran} q_z = i\left[-\frac{1}{2z}, \frac{1}{2z}\right].$$

Define also the bounded operator $Q_z = q_z(V|_{\tilde{H}}) \in B(\tilde{H})$ via the Borel functional calculus. Observe that $Q_z = R_z^* V R_z$, where $R_z = (z - V|_{\tilde{H}})^{-1}$ is the resolvent of $V|_{\tilde{H}} : \tilde{H} \to \tilde{H}$ at z.

For $\tau > 0$ and $G_{\tau} \colon H \to H$ constructed as in section 3.2 and appendix A, define the compact, skew-adjoint operators

$$Q_{z,\tau} = R_z^* V_\tau R_z, \quad V_\tau = G_{\tau/2} V G_{\tau/2}.$$

Note that well-definition and compactness of $Q_{z,\tau}$ follows from the fact that $G_{\tau/2}$ is a Markov operator induced by a C^1 kernel k_{τ} (in particular, ran $G_{\tau}|_{\tilde{H}} \subset H_V$), making $VG_{\tau/2}|_{\tilde{H}}$ a bounded operator. Since $G_{\tau} \to \operatorname{Id}$ as $\tau \to 0^+$, we view $Q_{z,\tau}$ as a compact approximation of $Q_z = q_z(V)$.

Define now the domain $\Omega_z = (-\infty, -z] \cup [z, \infty)$, and observe that $\tilde{q}_z := q_z|_{\Omega_z}$ is invertible with (unbounded) inverse $\tilde{q}_z^{-1} : i[-(2z)^{-1}, (2z)^{-1}] \setminus \{0\} \to i\mathbb{R}$,

$$\tilde{q}_z^{-1}(i\omega) = i \frac{1 + \sqrt{1 - 4z^2\omega^2}}{2\omega}.$$

As $z \to 0^+$, the domain Ω_z increases to $i\mathbb{R}$, so we view $\tilde{q}_z^{-1} \circ q_z$ as an approximation of the identity. Correspondingly, we view $\tilde{V}_z := \tilde{q}_z^{-1}(q_z(V|_{\tilde{H}})) \equiv \tilde{q}_z^{-1}(Q_z)$ as an unbounded, skew-adjoint approximation of $V|_{\tilde{H}}$.

Let $\tilde{V}_{z,\tau} = b_z(Q_{z,\tau})$ where $b_z : i\mathbb{R} \to i\mathbb{R}$ is any continuous extension of \tilde{q}_z^{-1} . Using the orthogonal projection $\operatorname{proj}_{\tilde{H}} : H \to \tilde{H}$, extend $\tilde{V}_{z,\tau}$ to the skew-adjoint operator $V_{z,\tau} = \tilde{V}_{z,\tau} \operatorname{proj}_{\tilde{H}}$ with dense domain $D(V_{z,\tau}) \subset H$. In [39, Theorem 6] it is shown:

Proposition 14. $V_{z,\tau}$ is a skew-adjoint operator with compact resolvent that converges to V in strong resolvent sense in the iterated limit of $z \to 0^+$ after $\tau \to 0^+$. In particular, properties (V1)–(V5) hold for $V_{z,\tau}$ in that limit.

Let us now consider an eigendecomposition of the compact, skew-adjoint operator $Q_{z,\tau}$:

$$Q_{z,\tau}\xi_{j,z,\tau} = \beta_{j,z,\tau}\xi_{j,z,\tau},\tag{46}$$

where $\beta_{j,z,\tau} \in i\mathbb{R}$ and the eigenvectors $\xi_{j,z,\tau}$ form an orthonormal basis of \tilde{H} . We index the eigenvalues by $j \in \mathbb{Z} \setminus \{0\}$ so that $\beta_{-j,z,\tau} = \overline{\beta_{j,z,\tau}}$ and $\xi_{-j,z,\tau} = \overline{\xi_{j,z,\tau}}$. We can then build an eigendecomposition

$$V_{z,\tau}\xi_{j,z,\tau} = i\omega_{j,z,\tau}\xi_{j,z,\tau},\tag{47}$$

with $i\omega_{j,z,\tau} = b_z(\beta_{j,z,\tau})$, and $\xi_{j,z,\tau}$ as above for $j \in \mathbb{Z} \setminus \{0\}$, and $\omega_{0,z,\tau} = 0$, $\xi_{0,z,\tau} = 1_X$. The eigendecomposition (47) completely characterizes $V_{z,\tau}$.

Remark 15. The extension $b_z \supset \tilde{q}_z^{-1}$ is introduced since the spectrum of $Q_{z,\tau}$ is not guaranteed to lie within the domain of definition $i[-(2z)^{-1},(2z)^{-1}]$ of \tilde{q}_z^{-1} for any $\tau > 0$. In practical applications, we have not observed any instances of (numerical approximations of) eigenvalues of $Q_{z,\tau}$ that do not lie in $i[-(2z)^{-1},(2z)^{-1}]$, so we work with \tilde{q}_z^{-1} without making an explicit choice of extension b_z .

To compute the eigenpairs $(\beta_{j,z,\tau}, \xi_{j,z,\tau})$, [40] formulates an associated variational eigenvalue problem to (46). Define the sesquilinear forms $A_{\tau} \colon H_{V} \times H_{V} \to \mathbb{C}$ and $B_{z} \colon H_{V} \times H_{V} \to \mathbb{C}$ such that

$$A_{\tau}(u,v) = \langle G_{\tau/2}u, VG_{\tau/2}v \rangle_H, \quad B_z(u,v) = \langle (z-V)u, (z-V)v \rangle_H.$$

It is shown that $(\beta_{j,z,\tau}, \xi_{j,z,\tau})$ solves (46) for $\beta_{j,z,\tau} \neq 0$ iff $\xi_{j,z,\tau} = (z - V)v_{j,z,\tau}$ and $(\beta_{j,z,\tau}, v_{j,z,\tau})$ solves the following variational problem.

Definition 16 (variational eigenvalue problem for $Q_{z,\tau}$). Find $\beta_{j,z,\tau} \in i\mathbb{R}$ and $v_{j,z,\tau} \in \tilde{H} \setminus \{0\}$ such that

$$A_{\tau}(u, v_{j,z,\tau}) = \beta_{j,z,\tau} B_z(u, v_{j,z,\tau}), \quad \forall u \in H_V.$$

The variational problem in Definition 16 is "physics-informed", in the sense that the sesquilinear forms A_{τ} and B_z can both be evaluated on elements $u = \iota f \in H_V$ and $v = \iota g \in H_V$ with continuous representatives $f, g \in C^1(M)$ using the generating vector field $\vec{V}: M \to TM$ of the dynamics. In particular, using (2), we get

$$A_{\tau}(u,v) = \int_{X \times X \times X} k'_{\tau}(w,x)k'_{\tau}(w,y)u(x)v(y) d(\mu \times \mu \times \mu)(w,x,y),$$

$$B_{z}(u,v) = \int_{X} \left((z - \vec{V} \cdot \nabla)f \right) \left((z - \vec{V} \cdot \nabla)g \right) d\mu,$$
(48)

where $k'_{\tau}(\cdot, y) = \vec{V} \cdot \nabla k(\cdot, x)$. In addition, if $f, g \in \text{ran } K_{\tau}$, i.e., $f = K_{\tau}r$ and $g = K_{\tau}s$ for $r, s \in H$, we have

$$B_z(u,v) = \int_{X \times X \times X} (zk_\tau(w,x) - k'_\tau(w,x))(zk_\tau(w,y) - k'_\tau(w,y))r(x)s(y) d(\mu \times \mu \times \mu)(w,x,y).$$
(49)

Thus, in such cases, $A_{\tau}(u, v)$ and $B_z(u, v)$ can be computed by "pushing" the action of the generator V to the directional derivatives $k'_{\tau}(\cdot, x)$ of the kernel sections $k_{\tau}(\cdot, x)$ with respect to the dynamical vector field \vec{V} .

The approach of [40] utilizes automatic differentiation to compute k'_{τ} without discretization errors that affect finite-difference approximations of the generator (e.g., [22, 34]). Using $E_l = \text{span}\{\phi_{1,\tau},\ldots,\phi_{l,\tau}\} \subset H_V$ as Galerkin approximation spaces, approximate solutions $(\beta_{j,\tau,lj,v_{j,z,\tau,l}})$

are computed by restricting the trial and test functions in Definition 16 to lie in E_l . Numerically, this is equivalent to solving a matrix generalized eigenvalue problem

$$A_{\tau}c_{j,z,\tau,l} = \beta_{j,z,\tau,l}B_{z}c_{j,z,\tau,l}, \quad \beta_{j,z,\tau,l} \in i\mathbb{R}, \quad c_{j,z,\tau,l} \in \mathbb{C}^{l} \setminus \{0\}.$$

Here $\mathbf{A}_{\tau} = [A_{\tau}(\phi_i, \phi_j)]_{i,j=1}^l$ and $\mathbf{B}_z = [B_z(\phi_i, \phi_j)]_{i,j=1}^l$ are $l \times l$ matrices whose elements are computed using (48) and (49), respectively, in conjunction with the kernel integral representation (44) of the $C^1(M)$ representatives $\varphi_{j,\tau}$ of the basis functions. Moreover, the generalized eigenvector $\mathbf{c}_{j,z,\tau,l} = (c_{1j,z,\tau,l}, \ldots, c_{lj,z,\tau,l})$ contains the expansion coefficients of $v_{j,z,\tau,l}$ in the kernel eigenbasis,

$$v_{j,z,\tau,l} = \sum_{i=1}^{l} c_{ij,z,\tau,l} \phi_i.$$

Specific formulas for the matrix elements of A_{τ} and B_z when using the Markov kernel construction from appendix A can be found in [40, section 4.3]. By convention, we order solutions in increasing order of Dirichlet energy (45), viz.

$$\mathcal{E}(v_{j,z,\tau,l}) = \sum_{i=1}^{l} \frac{|c_{ij,z,\tau,l}|^2}{\lambda_j}.$$

By results on Galerkin approximation of variational eigenvalue problems [2], as $l \to \infty$ $\beta_{j,z,\tau,l}$ converges to $\beta_{j,z,\tau}$ and, for an appropriate choice of eigenvectors, $v_{j,z,\tau,l}$ converges to $v_{j,z,\tau}$ in the norm of H_V for an appropriate choice of eigenvectors. The latter, implies convergence of $\xi_{j,z,\tau,l} := (z-V)v_{j,z,\tau,l}$ in the norm of H. Note that if $\beta_{j,z,\tau}$ is nonzero, $v_{j,z,\tau}$ lies in ran $G_{\tau/2}$ and thus has a representative in $\mathcal{H}_{\tau} \subseteq \mathcal{H}$. Thus, the Dirichlet energy $\mathcal{E}(v_{j,z,\tau})$ is finite so long as $\beta_{j,z,\tau} \neq 0$.

The scheme also has a data-driven formulation wherein all kernel integral operators are replaced by their counterparts induced from the sampling measures μ_N , as described in section 7 and appendix A. These methods converge in the limit of large data, $N \to \infty$; see [40, section 4.4] for further details.

References

- [1] N. Aronszajn, Theory of reproducing kernels, Trans. Amer. Math. Soc. 68 (1950), no. 3, 337–404.
- [2] I. Babuška and J. Osborn, Eigenvalue problems, Finite element methods (part 1), 1991, pp. 641–787.
- [3] V. Baladi, Positive transfer operators and decay of correlations, Advanced Series in Nonlinear Dynamics, vol. 16, World Scientific, Singapore, 2000.
- [4] V. Baladi and M. Tsujii, Anisotropic Hölder and Sobolev spaces for hyperbolic diffeomorphisms, Ann. Inst. Fourier 57 (2008), no. 1.
- [5] G. Benenti, G. Casati, S. Montangero, and D. L. Shepelyansky, Efficient quantum computing of complex dynamics, Phys. Rev. Lett. 87 (2001), 227901.
- [6] D. W. Berry, A. M. Childs, A. Ostrander, and G. Wang, Quantum algorithm for linear differential equations with exponentially improved dependence on precision, Commun. Math. Phys. **356** (2017), 1057–1081.
- [7] T. Berry, D. Giannakis, and J. Harlim, Nonparametric forecasting of low-dimensional dynamical systems, Phys. Rev. E. 91 (2015).
- [8] T. Berry and J. Harlim, Semiparametric modeling: Correcting low-dimensional model error in parametric models,
 J. Comput. Phys. 308 (2016), 305–321.
- [9] S. Bharadwaj and K. R. Sreenivasan, Quantum computation of fluid dynamics, Indian Acad. Sci. Conf. Ser. 3 (2020), no. 1, 77–96.
- [10] M. Blank, Egodic averaging with and without invariant measures, Nonlinearity 30 (2017), 4649–4664.
- [11] M. Blank, G. Keller, and C. Liverani, Ruelle-Perron-Frobenius spectrum for Anosov maps, Nonlinearity 15 (2002), no. 6, 1905–1973.
- [12] D. I. Bondar, F. Gay-Balmaz, and C. Tronci, Koopman wavefunctions and classical-quantum correlation dynamics, Proc. Roy. Soc. A 475 (2019), 20180879.
- [13] N. Boullé and M. Colbrook, Multiplicative Dynamic Mode Decomposition, 2024.
- [14] O. Butterley and C. Liverani, Smooth Anosov flows: Correlation spectra and stability, J. Mod. Dyn. 1 (2007), no. 2, 301–322.

- [15] R. Coifman and M. Hirn, Bi-stochastic kernels via asymmetric affinity functions, Appl. Comput. Harmon. Anal. **35** (2013), no. 1, 177–180.
- [16] R. R. Coifman, Y. Shkolnisky, F. J. Sigworth, and A. Singer, Graph Laplacian tomography from unknown random projections, IEEE Trans. Image Process. 17 (2008), no. 10, 1891–1899.
- [17] M. Colbrook, The multiverse of dynamic mode decomposition algorithms, Handbook of numerical analysis, 2024, pp. 88.
- [18] M. J. Colbrook and A. Townsend, Rigorous data-driven computation of spectral properties of Koopman operators for dynamical systems, Commun. Pure Appl. Math. 77 (2024), 221–283.
- [19] P. C. S. Costa, S. Jornan, and A. Ostrander, Quantum algorithm for simulating the wave equation, Phys. Rev. A 99 (2019).
- [20] S. Das and D. Giannakis, On harmonic Hilbert spaces on compact abelian groups, J. Fourier Anal. Appl. 29 (2023), no. 1, 12.
- [21] S. Das, D. Giannakis, and M. Montgomery, Correction to: On harmonic Hilbert spaces on compact abelian groups, J. Fourier Anal. Appl. 29 (2023), no. 6, 67.
- [22] S. Das, D. Giannakis, and J. Slawinska, Reproducing kernel Hilbert space compactification of unitary evolution groups, Appl. Comput. Harmon. Anal. 54 (2021), 75–136.
- [23] C. R. de Oliveira, Intermediate spectral theory and quantum dynamics, Progress in Mathematical Physics, vol. 54, Birkhäuser, Basel, 2009.
- [24] G. Della Ricia and N. Wiener, Wave mechanics in classical phase space, Brownian motion, and quantum theory, J. Math. Phys. 7 (1966), no. 8, 1732–1383.
- [25] I. Y. Dodi and E. A. Startsev, On applications of quantum computing to plasma simulations, Phys. Plasmas 28 (2021).
- [26] T. Eisner, B. Farkas, M. Haase, and R. Nagel, Operator theoretic aspects of ergodic theory, Graduate Texts in Mathematics, vol. 272, Springer, Cham, 2015.
- [27] T. J. Elliott and M. Gu, Superior memory efficiency of quantum devices for the simulation of continuous-time stochastic processes, npj Quantum Inf. 4 (2018), 18.
- [28] A. Engel, G. Smith, and S. E. Parker, Quantum algorithm for the Vlasov equation, Phys. Rev. A 100 (2019).
- [29] H. G. Feichtinger, Gewichtsfunktionen auf lokalkompakten Gruppen, Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II 188 (1979), no. 8–10, 451–471.
- [30] H. G. Feichtinger, S. S. Pandey, and T. Werther, Minimal norm interpolation in harmonic Hilbert spaces and Wiener amalgam spaces on locally compact abelian groups, J. Math. Kyoto Univ. 47 (2007), no. 1, 65–78.
- [31] D. C. Freeman, D. Giannakis, B. Mintz, A. Ourmazd, and J. Slawinska, Data assimilation in operator algebras, Proc. Natl. Acad. Sci. 120 (2023), no. 8.
- [32] D. C. Freeman, D. Giannakis, and J. Slawinska, Quantum mechanics for closure of dynamical systems, Multiscale Model. Simul. 22 (2024), no. 1, 283–333.
- [33] F. Gaitan, Finding flows of a Navier-Stokes fluid through quantum computing, npj Quantum Inf. 6 (2020), 61.
- [34] D. Giannakis, Data-driven spectral decomposition and forecasting of ergodic dynamical systems, Appl. Comput. Harmon. Anal. 47 (2019), no. 2, 338–396.
- [35] D. Giannakis, M. J. Latifi Jebelli, M. Montgomery, P. Pfeffer, J. Schumacher, and J. Slawinska, Tensor network approximation of Koopman operators, 2024.
- [36] D. Giannakis and M. Montgomery, An algebra structure for reproducing kernel Hilbert spaces, 2024.
- [37] D. Giannakis, A. Ourmazd, P. Pfeffer, J. Schumacher, and J. Slawinska, *Embedding classical dynamics in a quantum computer*, Phys. Rev. A **105** (2022).
- [38] D. Giannakis, J. Slawinska, and Z. Zhao, Spatiotemporal feature extraction with data-driven Koopman operators, Proceedings of the 1st international workshop on feature extraction: Modern questions and challenges at nips 2015, 2015, pp. 103–115.
- [39] D. Giannakis and C. Valva, Consistent spectral approximation of Koopman operators using resolvent compactification, Nonlinearity 37 (2024), no. 7.
- [40] _____, Physics-informed spectral approximation of Koopman operators, 2024.
- [41] K. Gröchenig, Weight functions in time-frequency analysis, Pseudodifferential operators: Partial differential equations and time-frequency analysis, 2007, pp. 343–366.
- [42] P. R. Halmos, Lectures on ergodic theory, American Mathematical Society, Providence, 1956.
- [43] P. R. Halmos and J. von Neumann, Operator methods in classical mechanics, II, Ann. Math. 43 (1942), no. 2, 332–350.
- [44] I. Joseph, Koopman-von Neumann approach to quantum simulation of nonlinear classical dynamics, Phys. Rev. Research 2 (2020), 043102.
- [45] I. Joseph, Y. Shi, M. D. Porter, A. R. Castelli, V. I. Geyko, F. R. Graziani, S. B. Libby, and J. L. DuBois, Quantum computing for fusion energy science applications, Phys. Plasmas 30 (2023), no. 1.

- [46] B. Kacewicz, Almost optimal solution of initial-value problems by randomized and quantum algorithms, J. Complex. 22 (2006), 676–690.
- [47] A. Kalev and I. Hen, Quantum algorithm for simulating Hamiltonian dynamics with an off-diagonal series expansion, Quantum 5 (2021), 426–449.
- [48] E. Kaniuth, A course in commutative Banach algebras, Graduate Texts in Mathematics, vol. 246, Springer Science+Media, 2009.
- [49] A. V. Kočergin, On mixing in special flows over a shifting of segments and in smooth flows on surfaces, Math. USSR Sbornik 25 (1975), no. 3, 441–469.
- [50] B. O. Koopman, Hamiltonian systems and transformation in Hilbert space, Proc. Natl. Acad. Sci. 17 (1931), no. 5, 315–318.
- [51] B. O. Koopman and J. von Neumann, Dynamical systems of continuous spectra, Proc. Natl. Acad. Sci. 18 (1932), no. 3, 255–263.
- [52] K. Law, A. Shukla, and A. M. Stuart, Analysis of the 3DVAR filter for the partially observed Lorenz'63 model, Discrete Contin. Dyn. Syst. 34 (2013), no. 3, 1061–10178.
- [53] D. Lehmann, Mathematical methods of many-body quantum field theory, Research Notes in Mathematics, vol. 436, Chapman & Hall/CRC, Boca Raton, 2004.
- [54] S. K. Leyton and T. J. Osborne, A quantum algorithm to solve nonlinear differential equations, 2008.
- [55] J.-P. Liu, H. Ø. Kolden, H. K. Krovi, and A. M. Childs, Efficient quantum algorithm for dissipative nonlinear differential equations, Proc. Natl. Acad. Sci. 118 (2021), no. 35, e2026805118.
- [56] S. Lloyd, G. DePalma, C. Gokler, B. Kiani, Z.-W. Liu, M. Marvian, F. Tennie, and T. Palmer, Quantum algorithm for nonlinear differential equations, 2020.
- [57] E. N. Lorenz, Deterministic nonperiodic flow, J. Atmos. Sci. 20 (1963), 130–141.
- [58] S. Luzzatto, I. Melbourne, and F. Paccaut, The Lorenz attractor is mixing, Comm. Math. Phys. 260 (2005), no. 2, 393–401.
- [59] D. Mauro, On Koopman-von Neumann waves, Int. J. Mod. Phys. A 17 (2002), 1301–1325.
- [60] A. Mezzacapo, M. Sanz, L. Lamata, I.L. Egusquiza, S. Succi, and E. Solano, Quantum simulator for transport phenomena in fluid flows, Sci. Rep. 5 (2015), 13153.
- [61] S. E. Otto and C. W. Rowley, Koopman operators for estimation and control of dynamical systems, Annu. Rev. Control Robot. Auton. Syst. 4 (2021), 59–87.
- [62] J. C. Oxtoby, Stepanoff flows on the torus, Proc. Amer. Math. Soc. 4 (1953), 982–987.
- [63] V. I. Paulsen and M. Raghupathi, An introduction to the theory of reproducing kernel Hilbert spaces, Cambridge Studies in Advanced Mathematics, vol. 152, Cambridge University Press, Cambridge, 2016.
- [64] P. Pfeffer, F. Heyder, and J. Schumacher, Hybrid quantum-classical reservoir computing of thermal convection flow, Phys. Rev. Research 4 (2022).
- [65] J. C. Sprott, Chaos and time-series analysis, Oxford University Press, Oxford, 2003.
- [66] B. K. Sriperumbudur, K. Fukumizu, and G. R. Lanckriet, Universality, characteristic kernels and RKHS embedding of measures, J. Mach. Learn. Res. 12 (2011), 2389–2410.
- [67] I. Steinwart and A. Christmann, Support vector machines, Information Science and Statistics, Springer, New York, 2008.
- [68] M. H. Stone, On one-parameter unitary groups in Hilbert space, Ann. Math 33 (1932), no. 3, 643-648.
- [69] M. Takesaki, Theory of operator algebras I, Encyclopaedia of Mathematical Sciences, vol. 124, Springer, Berlin, 2001.
- [70] F. Tennie and T. N. Palmer, Quantum computers for weather and climate prediction, Bull. Amer. Math. Soc. 104 (2023), no. 2, E488–E500.
- [71] A. F. M. ter Elst and M. Lemańczyk, On one-parameter Koopman groups, Ergodic Theory Dyn. Syst. 37 (2017), 1635–1656.
- [72] W. Tucker, The Lorenz attractor exists, C. R. Acad. Sci. Paris, Ser. I 328 (1999), 1197–1202.
- [73] U. von Luxburg, M. Belkin, and O. Bousquet, Consitency of spectral clustering, Ann. Stat. 26 (2008), no. 2, 555–586.
- [74] J. Wilkie and P. Brumer, Quantum-classical correspondence via Liouville dynamics. II. Correspondence for chaotic Hamiltonian systems, Phys. Rev. A 55 (1997), no. 1, 43–61.
- [75] _____, Quanum-classical correspondence via Liouville dynamics. I. Integrable systems and chaotic spectral decomposition, Phys. Rev. A 55 (1997), no. 1, 27–42.
- [76] L. Zelnik-Manor and P. Perona, Self-tuning spectral clustering, Advances in neural information processing systems, 2004, pp. 1601–1608.

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NH 03755, USA.

 $Email\ address: \verb|dimitrios.giannakis@dartmouth.edu|\\$

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NH 03755, USA.

 $Email\ address: {\tt mohammad.javad.latifi.jebelli@dartmouth.edu}$

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NH 03755, USA.

Email address: michael.r.montgomery@dartmouth.edu

Institut für Thermo- und Fluiddynamik, Technische Universität Ilmenau, D-98684 Ilmenau, Germany.

 $Email\ address \hbox{: philipp.pfeffer@tu-ilmenau.de}$

Institut für Thermo- und Fluiddynamik, Technische Universität Ilmenau, D-98684 Ilmenau, Germany.

 $Email\ address: \verb"joerg.schumacher@tu-ilmenau.de"$

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NH 03755, USA.

 $Email\ address: {\tt joanna.m.slawinska@dartmouth.edu}$