A New Cloud of Strings

G. Alencar,^{*} R. R. Landim,[†] and R. N. Costa Filho[‡]

Departamento de Física, Universidade Federal do Ceará,

Caixa Postal 6030, Campus do Pici, 60455-760 Fortaleza, Ceará, Brazil

(Dated: January 15, 2025)

In this manuscript, we generalize the cloud of strings model originally introduced by Letelier [1] by incorporating a magnetic-like component, Σ_{23} , alongside the electric-like component, Σ_{01} , considered in the original model. This extension yields a uniquely determined stress-energy tensor of the form diag $(\rho, -\rho, p, p)$, governed by a specific equation of state and parameterized by two constants. The resulting black hole solution exhibits significant modifications, including the presence of two independent hairs, distinguishing it from classical no-hair solutions. We analyze the thermodynamic properties of this black hole and demonstrate the existence of a remnant mass and finite entropy at the end of the Hawking evaporation process, suggesting a stable end-state. Furthermore, the presence of two hairs introduces new features that could influence black hole shadows and gravitational wave signals, offering valuable observational signatures for testing these exotic configurations.

INTRODUCTION

Recent years have witnessed significant advances in General Relativity, driven by groundbreaking observations such as gravitational waves from binary mergers involving black holes [2] and black hole-neutron star systems [3], as well as direct imaging of supermassive black hole environments in M87 [4] and the Milky Way [5]. These achievements mark the dawn of the multimessenger era, offering unprecedented opportunities to probe compact objects and test the strong-field regime of gravity [6].

In another direction, black holes interacting with surrounding matter provide critical insights into the interplay between geometry and matter. Among these, the cloud of strings model, introduced by Letelier [1], offers an elegant framework for describing spacetimes influenced by a continuous distribution of strings. This model has been extensively applied to explore the impact of extended matter sources on black hole spacetimes [7–13], highlighting its importance as a tool for understanding such interactions.

In this work, we generalize the cloud of strings model by introducing an additional magnetic-like component, Σ_{23} . This modification results in a complete stressenergy tensor of the form diag $(\rho, -\rho, p, p)$, parameterized by two constants and governed by a unique equation of state. We investigate the resulting black hole solutions, examining how these changes influence the spacetime geometry and thermodynamics. This provides a new perspective on the interaction between extended matter distributions and gravitational fields.

The starting point is the Nambu–Goto action, expressed as:

$$S_{NG} = \int \sqrt{-\gamma} \mathcal{M} d\lambda^0 d\lambda^1, \ \gamma_{AB} = g_{\mu\nu} \frac{\partial x^\mu}{\partial \lambda^A} \frac{\partial x^\nu}{\partial \lambda^B} \ . \tag{1}$$

Here, λ^0 and λ^1 are timelike and spacelike parameters, respectively, \mathcal{M} is a dimensionless constant characterizing the string, and γ is the determinant of the induced metric γ_{AB} on the submanifold. Using the following definitions:

$$\Sigma^{\mu\nu} = \epsilon^{AB} \frac{\partial x^{\mu}}{\partial \lambda^{A}} \frac{\partial x^{\nu}}{\partial \lambda^{B}}, \ \gamma = \frac{1}{2} \Sigma^{\mu\nu} \Sigma_{\mu\nu} \ , \qquad (2)$$

the stress-energy tensor is written as:

$$T^{\mu}_{\nu} = \rho \frac{\sum \mu \alpha \sum_{\alpha \nu}}{8\pi \sqrt{-\gamma}} , \qquad (3)$$

where $\Sigma^{\mu\nu}$ satisfies the following equations:

$$\nabla_{\mu} \left(\rho \Sigma^{\mu\nu} \right) = 0, \ \Sigma^{\mu\beta} \nabla_{\mu} \left[\frac{\Sigma_{\beta}{}^{\nu}}{(-\gamma)^{1/2}} \right] = 0 \ . \tag{4}$$

For spherically symmetric solutions, Letelier showed that the only component satisfying $\gamma < 0$ is the "electric-like" component, Σ_{01} . The corresponding black hole solution is:

$$f(r) = 1 - |a| - \frac{2M}{r}$$

Depending on the value of a, this solution may feature one or no horizons. Moreover, it lacks a remnant mass or entropy. However, by introducing the "magnetic-like" component Σ_{23} , the condition $\gamma < 0$ is preserved, and a more general stress-energy tensor is obtained while maintaining the desired physical properties.

To find the complete solution, we consider a general spherically symmetric metric of the form:

$$ds^{2} = -A(r)dt^{2} + B(r)dr^{2} + C^{2}(r)\left(d\theta^{2} + \sin^{2}\theta d\phi^{2}\right).$$
(5)

In this scenario, Eq. (4) leads to:

$$\partial_{\mu} \left(C^2 \sqrt{AB} \sin \theta \rho \Sigma^{\mu \nu} \right) = 0 . \qquad (6)$$

The general solutions of Eq. (6), consistent with spherical symmetry, are:

$$\rho \Sigma_{01} = \frac{a\sqrt{AB}}{C^2}, \ \rho \Sigma_{23} = b(r)\sin\theta \ , \tag{7}$$

where a is a constant and b(r) is an arbitrary function. The non-zero components of T^{μ}_{ν} are:

$$T^{0}_{0} = T^{1}_{1} = -\frac{a^{2}}{C^{2}\sqrt{a^{2} - b^{2}}} ,$$

$$T^{2}_{2} = T^{3}_{3} = \frac{b^{2}}{C^{2}\sqrt{a^{2} - b^{2}}} .$$
 (8)

From the conservation law, b(r) is determined as:

$$2C'\left(T^{0}_{\ 0} - T^{2}_{\ 2}\right) + \frac{dT^{0}_{\ 0}}{dr} = 0 \to b(r) = \frac{ac_{0}^{2}}{\sqrt{c_{0}^{4} + C^{4}}} , \quad (9)$$

where c_0 is a constant. The unique stress-energy tensor is then:

$$T^{0}_{\ 0} = -\frac{|a|}{C^{4}} \sqrt{c^{4}_{0} + C^{4}} = T^{1}_{\ 1} ,$$

$$T^{2}_{\ 2} = \frac{|a|c^{4}_{0}}{C^{4}\sqrt{c^{4}_{0} + C^{4}}} = T^{3}_{\ 3} .$$
(10)

In the limit $c_0 = 0$, this stress tensor reduces to Letelier's original model, with $a \rightarrow |a|$ [1]. Negative energy densities are excluded in both cases, and all energy conditions are satisfied. Finally, Eq. (9) ensures:

$$\gamma = \frac{-a^2}{\rho^2 (c_0^4 + C^4)} < 0 , \qquad (11)$$

as required, providing a fully consistent model.

The black hole solution sourced by the above stressenergy tensor is given by:

$$f(r) = 1 - \frac{2M}{r} + \frac{|a|c_0^2}{r^2} \,_2F_1\left(-\frac{1}{2}, -\frac{1}{4}, \frac{3}{4}, -\frac{r^4}{c_0^4}\right), \quad (12)$$

where the horizon is located at the roots of:

$$2M - r_h = \frac{|a|c_0^2}{r_h} \, _2F_1\left(-\frac{1}{2}, -\frac{1}{4}, \frac{3}{4}, -\frac{r_h^4}{c_0^4}\right). \tag{13}$$

While it is not possible to solve for r_h explicitly, the analysis shows that the solution features one horizon for $|a| \ge 1$ and two horizons for |a| < 1.

The asymptotic behavior of the solution is:

$$f(r) = 1 - \frac{2M}{r} + \frac{|a|c_0^2}{r^2}, \quad r \ll c_0, \tag{14}$$

$$f(r) = 1 - \frac{2M}{r} - |a| + \frac{|a|c_0^4}{6r^4}, \quad r \gg c_0.$$
(15)

This demonstrates that the solution is asymptotically flat, possesses a singularity at r = 0, and reduces to the Letelier solution in the limit $c_0 \rightarrow 0$.

The Hawking temperature is given by:

$$T_H = \frac{1}{4\pi r_h} \left(1 - \frac{|a|}{r_h^2} \sqrt{c_0^4 + r_h^4} \right).$$
(16)

For $|a| \ge 1$, $T_H \le 0$, which is non-physical, whereas for |a| < 1, $T_H > 0$, making this the relevant case. From Eq. (16), the remnant radius is:

$$r_h = r_0 = \frac{|c_0||a|^{1/2}}{(1-a^2)^{1/4}}.$$
(17)

The entropy of the remnant is:

$$S = \pi r_h^2 \to S_{rem} = \pi \frac{|c_0|^2 |a|}{(1-a^2)^{1/2}}.$$
 (18)

The derivative of the mass function satisfies:

$$M' = \frac{1}{2} \left(1 - \frac{|a|}{r_h^2} \sqrt{c_0^4 + r_h^4} \right), \tag{19}$$

which reaches a minimum at r_0 . The stability of the remnant is analyzed via the specific heat, $C = M'/T'_H$. The derivative of T_H is:

$$T'_{H} = \frac{3ac_{0}^{4} + ar_{h}^{4} - r_{h}^{2}\sqrt{c_{0}^{4} + r_{h}^{4}}}{4\pi r_{h}^{4}\sqrt{c_{0}^{4} + r_{h}^{4}}}.$$
 (20)

From Eqs. (19) and (20), it is observed that C = 0 and changes sign at $r = r_0$, confirming that the remnant is stable.

For cylindrical symmetry, the metric is:

$$ds^{2} = -f(r)dt^{2} + \frac{dr^{2}}{f(r)} + \alpha^{2}r^{2}dz^{2} + r^{2}d\theta^{2}.$$
 (21)

The stress-energy tensor components are:

$$T^{0}_{\ 0} = -\frac{|a|}{r^{4}}\sqrt{c_{0}^{4} + r^{4}} = T^{1}_{\ 1},$$

$$T^{2}_{\ 2} = \frac{|a|c_{0}^{4}}{r^{4}\sqrt{c_{0}^{4} + r^{4}}} = T^{3}_{\ 3}.$$
 (22)

The solution for f(r) is:

$$f(r) = \frac{2M}{r} + \frac{|a|c_0^2}{r^2} {}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}, \frac{3}{4}, -\frac{r^4}{c_0^4}\right), \quad (23)$$

and, as expected, black holes are not allowed in cylindrical symmetry. As a concluding remark, our results reveal that the extended model preserves key physical properties, such as $\gamma < 0$, while exhibiting novel features not present in the original Letelier solution. The stress tensor is uniquely determined and differs from the fluid of strings of Letelier in the sense that the pressures are uniquely fixed. For instance, the perfect fluid case, as considered by Soleng in Ref. [14], is not allowed.

The black hole solution sourced by this generalized stress-energy tensor is asymptotically flat and includes two horizons for |a| < 1. Importantly, we identified the existence of a remnant mass and entropy at the end of the Hawking evaporation process, providing a stable end-state. Furthermore, we explored the thermodynamic properties of the solution and demonstrated the stability of the remnant through an analysis of the specific heat. The results suggest that remnants arising in this context could serve as candidates for addressing longstanding issues in black hole physics, such as the information loss paradox [15]. Finally, we considered the cylindrical symmetry case, showing that black holes are not supported in this configuration, consistent with physical expectations.

An interesting extension of this work involves analyzing the rotating case. A rotating counterpart of this black hole solution would allow the investigation of shadows. The impact of the additional hair parameters on the shape and size of the shadow could provide valuable insights into the observational signatures of such exotic solutions.

Future work could also explore potential observational imprints of these remnants, including gravitational lensing, accretion disk dynamics, gravitational wave signals, and connections to Dark Matter. Another avenue is studying these solutions in the context of modified gravity theories and cosmology.

ACKNOWLEDGMENTS

The authors thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico.

- * geova@fisica.ufc.br
- [†] renan@fisica.ufc.br
- ‡ rai@fisica.ufc.br
- P. S. Letelier, CLOUDS OF STRINGS IN GENERAL RELATIVITY, Phys. Rev. D 20, 1294 (1979).
- [2] B. P. Abbott *et al.* (LIGO Scientific, Virgo), Observation of Gravitational Waves from a Binary Black Hole Merger, Phys. Rev. Lett. **116**, 061102 (2016), arXiv:1602.03837 [gr-qc].
- [3] B. P. Abbott et al. (LIGO Scientific, Virgo), GW170817: Observation of Gravitational Neutron Waves from а Binary Star In-Phys. Rev. Lett. 119, 161101 (2017), spiral, arXiv:1710.05832 [gr-qc].

- [4] K. Akivama et al.(Event Horizon Telescope). First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Astrophys. J. Lett. 875, L1 (2019), Black Hole, arXiv:1906.11238 [astro-ph.GA].
- [5] K. Akiyama *et al.* (Event Horizon Telescope), First Sagittarius A* Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole in the Center of the Milky Way, Astrophys. J. Lett. **930**, L12 (2022), arXiv:2311.08680 [astro-ph.HE].
- [6] A. Addazi *et al.*, Quantum gravity phenomenology at the dawn of the multi-messenger era—A review, Prog. Part. Nucl. Phys. **125**, 103948 (2022), arXiv:2111.05659 [hep-ph].
- [7] A. Ganguly, S. G. Ghosh, and S. D. Maharaj, Accretion onto a black hole in a string cloud background, Phys. Rev. D 90, 064037 (2014), arXiv:1409.7872 [gr-qc].
- [8] G. Panotopoulos and Α. Rincón, Minimal Geometric Deformation in cloud а Eur. Phys. J. C 78, 851 (2018), of strings, arXiv:1810.08830 [gr-qc].
- [9] D. V. Singh, S. G. Ghosh, and S. D. Maharaj, Clouds of strings in 4D Einstein–Gauss–Bonnet black holes, Phys. Dark Univ. **30**, 100730 (2020), arXiv:2003.14136 [gr-qc].
- [10] G. Mustafa, F. Atamurotov, I. Hussain, S. Shaymatov, and A. Övgün, Shadows and gravitational weak lensing by the Schwarzschild black hole in the string cloud background with quintessential field*, Chin. Phys. C 46, 125107 (2022), arXiv:2207.07608 [gr-qc].
- Ghosh and S. D. Maharaj, [11] S. G. Cloud of strings for radiating black holes in Lovelock gravity, Phys. Rev. D 89, 084027 (2014), arXiv:1409.7874 [gr-qc].
- [12] J. de M. Toledo and V. B. Bezerra, Black holes with cloud of strings and quintessence in Lovelock gravity, Eur. Phys. J. C 78, 534 (2018).
- [13] J. M. Toledo and V. B. Bezerra, The Reissner-Nordström black hole surrounded by quintessence and a cloud of strings: Thermodynamics and quasinormal modes, Int. J. Mod. Phys. D 28, 1950023 (2018).
- [14] H. H. Soleng, Dark matter and nonNewtonian gravity from general relativity coupled to a fluid of strings, Gen. Rel. Grav. 27, 367 (1995), arXiv:gr-qc/9412053.
- [15] Y. C. Ong, The Case For Black Hole Remnants: A Review (2024) arXiv:2412.00322 [gr-qc].