Reality conditions for sine-Gordon equation and quasi-periodic solutions in finite phase spaces

Julia Bernatska

Abstract. Quasi-periodic solutions of the sine-Gordon equation are expressed in terms of $\wp_{1,2g-1}$ -function, and reality conditions are completely determined. This new result leads to numerical computation and graphical representation of non-linear wave solutions. A way of obtaining finite-gap solutions by means of algebro-geometric integration is also explained, and a construction of the sine-Gordon hierarchy on coadjoint orbits of SU(2) loop group is addressed.

1. Introduction

The present paper is a continuation of the research started in [[7](#page-26-0)], and focuses on the sine-Gordon equation

$$
\phi_{\text{tx}} = -4\sin\phi.
$$

The sine-Gordon equation is known to be completely integrable, and possesses soliton solutions. The equation has various applications in physics, see [[19](#page-27-0)].

In [[13](#page-27-1)], Hirota applied his method to the sine-Gordon equation, and obtained N-soliton solutions; also interaction of two solitons was analyzed in detail. In [[1](#page-26-1)], a similar solution was obtained by the inverse scattering method.

The sine-Gordon equation is associated with a hierarchy of integrable hamiltonian systems. Each system is finite, and produces a finite-gap solution of the equation. Periodic solutions of [\(1\)](#page-0-0), expressed in terms of theta functions, are suggested in [[12](#page-27-2)], where also the reality conditions are addressed. These solutions have the form, see $[6, Eq. (4.2.25)],$ $[6, Eq. (4.2.25)],$ $[6, Eq. (4.2.25)],$

(2)
$$
\phi(x,t) = 2i \log \frac{\theta(i(\mathbf{U}x + \mathbf{W}t) + \mathbf{D} + i\pi\Delta)}{\theta(i(\mathbf{U}x + \mathbf{W}t) + \mathbf{D})},
$$

where U, W, D, Δ are vectors defined through periods of the corresponding spectral curve. How such a solution arises from an integrable system of the sine-Gordon hierarchy is explained in [[6](#page-26-2), §4.2].

Recently, some improvement on computing multi-soliton solutions by the inverse scattering method is suggested in [[4](#page-26-3)]; a review of solutions of the sine-Gordon equation in elementary and elliptic functions can be found therein. In addition to what mentioned in [[4](#page-26-3)], two- and three-gap elliptic solutions are constructed on spectral curves which are two- and three-sheeted coverings of the elliptic curve, see $[16, 17]$ $[16, 17]$ $[16, 17]$ $[16, 17]$ $[16, 17]$.

In the present paper we consider the sine-Gordon hierarchy which arises on coadjoint orbits of a loop group. This method differs from the one used in $[6]$ $[6]$ $[6]$, and so the reader will find a detailed explanation. The proposed method leads to algebraic integration in terms of \wp -functions, and the $\wp_{1,2g-1}$ -solution mentioned in [[10](#page-27-5), Theorem 4.13] is accurately derived. Though similarity in the structure with [[7](#page-26-0)] can be observed, the sine-Gordon hierarchy arise on different coadjoint orbits, and so the seeming repetition is an essentially different exposition.

The new results consist of reality conditions for the sine-Gordon equation, and a real-valued solution in terms of a particular \wp -function. The obtained reality conditions are more narrow than the conditions presented in [[12](#page-27-2)]. Actually, solutions of the sine-Gordon equation arise on hyperelliptic curves with one zero branch point and all other pairwise conjugate branch points. On hyperelliptic curves with all real branch points solutions of the sinh-Gordon equation arise.

The obtained real-valued quasi-periodic solutions are easily computed and illustrated. Analysis of the phase spaces of hamiltonian systems in the hierarchy allows to find proper values of integrals of motion, and examine different regimes of wave propagation.

The paper is organized as follows. In Preliminaries we recall basic knowledge on homologies and cohomologies of hyperelliptic curves, definitions of theta, sigma, and \wp -functions, and also representation of characteristics in terms of partitions on the set of indices of branch points. In Section 3 integrable systems of the sine(sinh)-Gordon hierarchy are constructed on coadjoint orbits of an affine Lie algebra. In Section 4 variables of separation are derived and proved to be quasicanonical. Section 5 is devoted to algebraic integration, here expressions in terms of ℘-functions are obtained for all dynamical variables of a hamiltonian system in the hierarchy. Also relations between x, t and coordinates of the Jacobian variety of the spectral curve, which serve as arguments of \wp -functions, are discovered. In Section 6 behaviour of \wp -functions on the Jacobian variety is analyzed, and particular subspaces which serve as the domain of solutions of the sine-Gordon equation are singled out. Finally, in Section 7 examples of systems with spectral curves of genera 1 and 2 are presented, with analysis of the phase space, and numerical computation of non-linear waves for several cases with different mutual positions of branch points.

2. Preliminaries

2.1. Hyperelliptic curves. We work with the canonical form of a hyperelliptic curve V of genus g, namely

(3)
$$
f(x,y) \equiv -y^2 + x^{2g+1} + \sum_{i=0}^{2g} \lambda_{2i+2} x^{2g-i} = 0.
$$

In terms of branch points $(e_i, 0)$, or simply e_i , $i = 1, \ldots, 2g + 1$, f has the form

(4)
$$
f(x,y) = -y^2 + \prod_{i=1}^{2g+1} (x - e_i).
$$

One more branch point e_0 is located at infinity, which serves as the base-point. We suppose that all branch points are distinct, and so the curve is not degenerate, that is the genus equals g. Finite branch points are enumerated in the ascending order of real and imaginary parts. The Riemann surface of $\mathcal V$ is constructed by means of the monodromy path through all branch points, according to the order, which starts at infinity and ends at infinity, see the orange line on fig. [1.](#page-2-0)

A homology basis is defined after H. Baker [[5](#page-26-4), p. 297], which is a kind of standard on a hyperelliptic curve. Cuts are made between points e_{2k-1} and e_{2k} with k from 1 to g. One more cut starts at e_{2g+1} and ends at infinity. Canonical homology cycles are defined as follows. Each \mathfrak{a}_k -cycle encircles the cut $(e_{2k-1}, e_{2k}), k = 1, \ldots, g$, and each \mathfrak{b}_k -cycle emerges from the cut (e_{2k-1}, e_{2k}) and enters the cut (e_{2g+1}, ∞) , see fig. [1.](#page-2-0)

Figure 1. Cuts and cycles on a hyperelliptic curve.

Fig. [1](#page-2-0) represents a spectral curve of a g-gap system in the sine-Gordon hierarchy, see section [6](#page-15-0) for more details. Parameters of the curve λ_k are real, and all finite branch points but one are complex conjugate.

Let first kind differentials $du = (du_1, du_3, \ldots, du_{2g-1})^t$ and second kind differentials $dr = (dr_1, dr_3, \ldots, dr_{2g-1})^t$ form a system of associated differentials, see $[5, \S 138]$ $[5, \S 138]$ $[5, \S 138]$, and be defined as in $[5, Ex. i, p. 195]$. Actually,

(5a)
$$
\mathrm{d}u_{2n-1} = \frac{x^{g-n} \mathrm{d}x}{\partial_y f(x, y)}, \quad n = 1, \dots, g,
$$

(5b)
$$
\mathrm{d}r_{2n-1} = \frac{\mathrm{d}x}{\partial_y f(x,y)} \sum_{j=1}^{2n-1} (2n-j) \lambda_{2j-2} x^{g+n-j}, \ \lambda_0 = 1, \quad n = 1, \dots, g.
$$

Indices of du_{2n-1} display the orders of zeros at infinity, and indices of dr_{2n-1} display the orders of poles.

The first kind differentials are not normalized. The corresponding periods along the canonical cycles a_k , b_k , $k = 1, \ldots, g$, are defined as follows

(6)
$$
\omega_k = \oint_{\mathfrak{a}_k} \mathrm{d}u, \qquad \omega'_k = \oint_{\mathfrak{b}_k} \mathrm{d}u.
$$

The vectors ω_k , ω'_k form first kind period matrices ω , ω' , respectively. Second kind period matrices η , η are composed of columns

(7)
$$
\eta_k = \oint_{\mathfrak{a}_k} dr, \qquad \eta'_k = \oint_{\mathfrak{b}_k} dr.
$$

Normalized first kind period matrices are 1_q , τ , where 1_q denotes the identity matrix of size g, and $\tau = \omega^{-1} \omega'$. Matrix τ is symmetric with a positive imaginary part: $\tau^t = \tau$, Im $\tau > 0$, that is τ belongs to the Siegel upper half-space. The normalized holomorphic differentials are denoted by

$$
dv = \omega^{-1} du.
$$

2.2. Abel's map. The vectors ω_k , ω'_k form a period lattice $\{\omega, \omega'\}$, and Then $Jac(\mathcal{V}) = \mathbb{C}^g / {\{\omega, \omega'\}}$ is the Jacobian variety of the curve [\(3\)](#page-1-0). Let $u = (u_1, u_3, \dots, u_n)$ $(u_{2g-1})^t$ be not normalized coordinates of Jac.

The Abel map is defined by

$$
\mathcal{A}(P) = \int_{\infty}^{P} du, \qquad P = (x, y) \in \mathcal{V},
$$

and on a positive divisor $D = \sum_{i=1}^{n} (x_i, y_i)$ by $\mathcal{A}(D) = \sum_{i=1}^{n} \mathcal{A}(P_i)$. The Abel map is one-to-one on the g-th symmetric power of the curve.

2.3. Theta function. The Riemann theta function is defined by

$$
\theta(v;\tau) = \sum_{n \in \mathbb{Z}^g} \exp\left(i\pi n^t \tau n + 2i\pi n^t v\right).
$$

where $v = \omega^{-1}u$ are normalized coordinates, and $\tau = \omega^{-1}\omega'$. Theta function with characteristic $[\varepsilon]$ is defined by

$$
\theta[\varepsilon](v;\tau) = \exp\left(i\pi(\varepsilon'^{t}/2)\tau(\varepsilon'/2) + 2i\pi(v+\varepsilon/2)^{t}\varepsilon'/2\right)\theta(v+\varepsilon/2 + \tau\varepsilon'/2;\tau),
$$

where $[\varepsilon] = (\varepsilon', \varepsilon)^t$ is a $2 \times g$ matrix, all components of ε , and ε' are real values within the interval $[0, 2)$. Modulo (mod) 2 addition is defined on characteristics.

Every point u within a fundamental domain of $Jac(\mathcal{V})$ can be represented by its characteristic $[\varepsilon]$ as follows

$$
u = \frac{1}{2}\omega\varepsilon + \frac{1}{2}\omega'\varepsilon'.
$$

Characteristics with values 0 and 1 correspond to half-periods, which are Abel images of divisors composed of branch points. Such a characteristic $[\varepsilon]$ is odd whenever $\varepsilon^t \varepsilon' = 0 \pmod{2}$, and even whenever $\varepsilon^t \varepsilon' = 1 \pmod{2}$. Theta function with characteristic has the same parity as its characteristic.

2.4. Sigma function and \wp **-functions.** The modular invariant entire function on $Jac(\mathcal{V})$ is called the sigma function, which we define by, see [[10](#page-27-5), Eq.(2.3)]:

(8)
$$
\sigma(u) = C \exp\left(-\frac{1}{2}u^t \times u\right) \theta[K](\omega^{-1}u; \omega^{-1}\omega'),
$$

where [K] is the characteristic of the vector K of Riemann constants, and $\varkappa = \eta \omega^{-1}$ is a symmetric matrix.

In what follows we use multiply periodic \wp -functions

$$
\wp_{i,j}(u) = -\frac{\partial^2 \log \sigma(u)}{\partial u_i \partial u_j}, \qquad \wp_{i,j,k}(u) = -\frac{\partial^3 \log \sigma(u)}{\partial u_i \partial u_j \partial u_k},
$$

which are defined on $Jac(V)\Sigma$, where $\Sigma = \{u \mid \sigma(u) = 0\}$ denotes the theta divisor in not normalized coordinates.

REALITY CONDITIONS FOR THE SINE-GORDON EQUATION AND QUASI-PERIODIC SOLUTIONS5

2.5. Jacobi inversion problem. A solution of the Jacobi inversion problem on a hyperelliptic curve is proposed in [[5](#page-26-4), Art. 216], see also [[10](#page-27-5), Theorem 2.2]. Let $u = A(D)$ be the Abel image of a non-special positive divisor $D \in \mathcal{V}^g$. Then D is uniquely defined by the system of equations

(9a)
$$
\mathcal{R}_{2g}(x, y; u) \equiv x^g - \sum_{i=1}^g x^{g-i} \wp_{1,2i-1}(u) = 0,
$$

(9b)
$$
\mathcal{R}_{2g+1}(x,y;u) \equiv 2y + \sum_{i=1}^{g} x^{g-i} \wp_{1,1,2i-1}(u) = 0.
$$

2.6. Characteristics and partitions. Let $S = \{0, 1, 2, \ldots, 2g+1\}$ be the set of indices of all branch points of a hyperellipitic curve of genus g , and 0 stands for the branch point at infinity. According to [[5](#page-26-4), § 202], all characteristics of half-periods are represented by partitions of S of the form $\mathcal{I}_{m} \cup \mathcal{J}_{m}$ with $\mathcal{I}_{m} = \{i_1, \ldots, i_{q+1-2m}\}\$ and $\mathcal{J}_{\mathfrak{m}} = \{j_1, \ldots, j_{g+1+2\mathfrak{m}}\}$, where \mathfrak{m} runs from 0 to $[(g+1)/2]$, and $[\cdot]$ denotes the integer part. Index 0, corresponding to infinity, is usually omitted in sets, it is also omitted in computation of cardinality of a set.

Denote by $[\varepsilon(\mathcal{I})] = \sum_{i \in \mathcal{I}} [\varepsilon_i] \pmod{2}$ the characteristic of

$$
\mathcal{A}(\mathcal{I}) = \sum_{i \in \mathcal{I}} \mathcal{A}(e_i) = \omega \left(\frac{1}{2} \varepsilon(\mathcal{I}) + \frac{1}{2} \tau \varepsilon'(\mathcal{I}) \right).
$$

Characteristics corresponding to $2q+1$ branch points serve as a basis for constructing all 2^{2g} half-period characteristics. Below a partition is referred to by the part of less cardinality, denoted by \mathcal{I} . Let

$$
[\mathcal{I}] = [\varepsilon(\mathcal{I})] + [K] \; (\text{mod } 2),
$$

and $|K|$ equals the sum of g odd characteristics of branch points. In the basis of canonical cycles introduced by fig. [1](#page-2-0) we have

$$
[K] = \sum_{k=1}^{g} [\varepsilon_{2k}].
$$

Let $\mathcal{I}_{m} \cup \mathcal{J}_{m}$ be a partition introduced above, then $[\mathcal{I}_{m}] = [\mathcal{J}_{m}]$. Characteristics $[\mathcal{I}_{m}]$ of even m are even, and of odd m are odd. According to the Riemann vanishing theorem, $\theta(v + \mathcal{A}(\mathcal{I}_m) + K; \tau)$ vanishes to order m at $v = 0$. Number m is called multiplicity. Characteristics of multiplicity 0 are called non-singular even characteristics. Characteristics of multiplicity 1 are called non-singular odd. All other characteristics are called singular.

3. Integrable systems on coadjoint orbits of a loop group

The sine-Gordon equation arises within the hierarchy of integrable Hamiltonian systems on coadjoint orbits of loop SU(2)-group. We briefly recall this construction, as presented in [[14](#page-27-6)] and recalled in [[9](#page-27-7)]. Such a construction is based on the results of [[2](#page-26-5),[3](#page-26-6)].

3.1. Affine Lie algebra. Let $\widetilde{\mathfrak{g}} = \mathfrak{g} \otimes \mathfrak{L}(z, z^{-1})$, where $\mathfrak{L}(z, z^{-1})$ denotes the algebra of Laurent series in z, and $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{C})$ has the standard basis

$$
\mathsf{H} = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathsf{X} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \mathsf{Y} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.
$$

The algebra $\tilde{\mathfrak{g}}$ is *principally graded*, with the grading operator

(10)
$$
\mathfrak{d} = 2z \frac{\mathrm{d}}{\mathrm{d}z} + \mathrm{ad}_{\mathsf{H}},
$$

where ad denotes the adjoint operator in g, that is $\forall Z \in \mathfrak{g}$ ad_H Z = [H, Z].

Let $\{X_{2m-1} = z^{m-1}X = z^mY, Y_{2m-1}, H_{2m} = z^mH, | m \in \mathbb{Z}\}\)$ form a basis of $\tilde{\mathfrak{g}}$. Let \mathfrak{g}_{ℓ} be the eigenspace of \mathfrak{d} of degree ℓ . Actually, $\mathfrak{g}_{2m-1} = \text{span}\{X_{2m-1}, Y_{2m-1}\},$ and $\mathfrak{g}_{2m} = \text{span}\{\mathsf{H}_{2m}\}.$

The bilinear form, with a fixed positive integer N ,

(11)
$$
\forall A(z), B(z) \in \widetilde{\mathfrak{g}} \qquad \langle A(z), B(z) \rangle = \mathop{\mathrm{res}}_{z=0} z^{-N} \operatorname{tr} A(z) B(z)
$$

introduces the dual algebra $\tilde{\mathfrak{g}}^*$, with the basis $\{X_{2m-1}^* = z^m X^*, Y_{2m-1}^* = z^{m-1} Y^*,\}$ $H_{2m}^* = z^m H^*$, $m \in \mathbb{Z}$, where

$$
\mathsf{H}^* = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathsf{X}^* = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad \mathsf{Y}^* = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}
$$

.

By means of [\(11\)](#page-5-0), we find

$$
X_{2m-1}^* = X_{2N-2m-1}, \quad Y_{2m-1}^* = Y_{2N-2m-1}, \quad H_{2m}^* = H_{2N-2m-2}.
$$

According to the Adler—Kostant—Symes scheme, see [[3](#page-26-6)], let

$$
\widetilde{\mathfrak{g}}_{+}=\oplus \sum_{\ell\geqslant -1}\mathfrak{g}_{\ell},\qquad \widetilde{\mathfrak{g}}_{-}=\oplus \sum_{\ell<-1}\mathfrak{g}_{\ell}.
$$

Then the dual subalgebras $\tilde{\mathfrak{g}}^*_+$ and $\tilde{\mathfrak{g}}^*_-$ with respect to the bilinear form [\(11\)](#page-5-0) are

$$
\widetilde{\mathfrak{g}}^*_+ = \oplus \sum_{\ell \leqslant 2N} \mathfrak{g}_\ell, \qquad \widetilde{\mathfrak{g}}^*_- = \oplus \sum_{\ell > 2N} \mathfrak{g}_\ell.
$$

Note, that $\mathfrak{g}_{N-1}^* = \mathfrak{g}_{N-1}$, and \mathfrak{g}_{ℓ}^* is dual to $\mathfrak{g}_{2N-\ell-2}$.

In what follows, we denote elements of the basis of $\tilde{\mathfrak{g}}$ by $\mathsf{Z}_{a,\ell}$, $a = 1, 2, 3$, where $\mathfrak{d}Z_{a,\ell} = \ell Z_{a,\ell}$, and $Z_{1,\ell} = H_{\ell}$, $Z_{2,\ell} = Y_{\ell}$, $Z_{3,\ell} = X_{\ell}$.

3.2. The phase space of sine-Gordon hierarchy. The sine-Gordon hierarchy of hamiltonian systems arise on coadjoint orbits of the group $G_{+} = \exp(\widetilde{\mathfrak{g}}_{+}).$ Indeed, the subspace $\mathcal{M}_N = \tilde{\mathfrak{g}}^*_+ / (\sum_{\ell < -1} \mathfrak{g}_\ell)$ is ad^{*}-invariant under the coadjoint action of $\widetilde{\mathfrak{g}}_+$. Actually,

$$
\mathcal{M}_N = \left\{ \Psi = \sum_{\ell=-1}^{2N} \sum_{a=1,2,3} \psi_{a,\ell} \mathsf{Z}_{a,\ell}^* \right\},\,
$$

where $\psi_{a,\ell}$ are dynamic variables and coordinates on \mathcal{M}_N ,

$$
\psi_{a,\ell} = \langle \Psi, \mathsf{Z}_{a,2N-\ell-2} \rangle.
$$

Let $\psi_{1,\ell} = \alpha_{\ell}, \psi_{2,\ell} = \beta_{\ell}, \psi_{3,\ell} = \gamma_{\ell}$, then every element $\Psi \in \mathcal{M}_N$ has the form

(12a)
$$
\Psi(z) = \begin{pmatrix} \alpha(z) & \beta(z) \\ \gamma(z) & -\alpha(z) \end{pmatrix},
$$

REALITY CONDITIONS FOR THE SINE-GORDON EQUATION AND QUASI-PERIODIC SOLUTIONS

(12b)
$$
\alpha(z) = \sum_{m=0}^{N} \alpha_{2m} z^{m}, \quad \beta(z) = \sum_{m=0}^{N} \beta_{2m-1} z^{m-1}, \quad \gamma(z) = \sum_{m=0}^{N} \gamma_{2m-1} z^{m},
$$

$$
\alpha_{2N} = 0, \qquad \beta_{2m-1} = \gamma_{2m-1} = b = \text{const.}
$$

The coordinates of \mathcal{M}_N , of number 3N, are ordered as follows:

(13)
$$
\{\beta_{2m-1}, \gamma_{2m-1}, \alpha_{2m}\}_{m=0}^{N-1}.
$$

The symplectic manifold \mathcal{M}_N is equipped with the Lie-Poisson bracket

(14a)
$$
\forall \mathcal{F}, \mathcal{H} \in \mathcal{C}(\mathcal{M}_N) \qquad \{\mathcal{F}, \mathcal{H}\} = \sum_{i,j=-1}^{2N-2} \sum_{a,b=1,2,3} W_{i,j}^{a,b} \frac{\partial \mathcal{F}}{\partial \psi_{a,i}} \frac{\partial \mathcal{H}}{\partial \psi_{b,j}},
$$
(4.1)

(14b)
$$
W_{i,j}^{a,b} = \langle \Psi, [Z_{a,2N-i-2}, Z_{a,2N-j-2}] \rangle,
$$

or in terms of the dynamic variables:

(15)
$$
\{\beta_{2m-1}, \alpha_{2n}\} = \beta_{2(n+m-N)+1}, \{\gamma_{2m-1}, \alpha_{2n}\} = -\gamma_{2(n+m-N)+1}, \quad N-1 \leq m+n, \{\beta_{2m-1}, \gamma_{2n-1}\} = -2\alpha_{2(m+n-N)}, \quad N \leq m+n.
$$

The Lie-Poisson structure $\mathsf{W} = (W_{i,j}^{a,b})$ has the form

$$
W = \begin{pmatrix} 0 & 0 & \dots & 0 & w_0 \\ \vdots & \vdots & \dots & w_0 & w_1 \\ 0 & 0 & \vdots & w_1 & w_2 \\ 0 & w_0 & \vdots & \vdots & \vdots \\ w_0 & w_1 & \dots & w_{N-2} & w_{N-1} \end{pmatrix},
$$

\n
$$
w_0 = \begin{pmatrix} 0 & 0 & \beta_{-1} \\ 0 & 0 & -\gamma_{-1} \\ -\beta_{-1} & \gamma_{-1} & 0 \end{pmatrix},
$$

\n
$$
w_n = \begin{pmatrix} 0 & -2\alpha_{2n-2} & \beta_{2n-1} \\ 2\alpha_{2n-2} & 0 & -\gamma_{2n-1} \\ -\beta_{2n-1} & \gamma_{2n-1} & 0 \end{pmatrix}, \quad n = 1, \dots, N-1.
$$

The Poisson bracket is degenerate, and not canonical.

The action of \tilde{G}_+ splits \mathcal{M}_N into orbits

$$
\mathcal{O} = \{ \Psi = \mathrm{Ad}_g^* \, \Phi \mid g \in \widetilde{G}_+ \}, \qquad \Phi \in \mathcal{M}_N.
$$

Orbits are of dimension 2N, and serve as phase spaces of hamiltonian systems.

Physically meaningful hamiltonian systems arise when g is one of the real forms of $\mathfrak{sl}(2,\mathbb{C})$, namely $\mathfrak{sl}(2,\mathbb{R})$ or $\mathfrak{su}(2)$. In the case of $\mathfrak{su}(2)$, the sine-Gordon equation is obtained, and in the case of $\mathfrak{sl}(2,\mathbb{R})$, the sinh-Gordon equation.

REMARK 1. In \mathcal{M}_N with Ψ of the form [\(12\)](#page-5-1), coadjoint orbits of \tilde{G}_- = exp($\tilde{\mathfrak{g}}_-$) serve as phase spaces for the hierarchy of +mKdV in the case of $\mathfrak{su}(2)$, and $-\mathrm{mKdV}$ in the case of $\mathfrak{sl}(2,\mathbb{R})$.

3.3. Integrals of motion. Let

(16)
$$
H(z) = \frac{1}{2} \operatorname{tr} \Psi^2(z) = \alpha(z)^2 + \beta(z)\gamma(z)
$$

$$
= h_{2N-1} z^{2N-1} + \dots + h_1 z + h_0 + h_{-1} z^{-1},
$$

where

(17)
\n
$$
h_{2N-1} = b^2,
$$
\n
$$
h_{2N-2} = \alpha_{2N-2}^2 + b(\beta_{2N-3} + \gamma_{2N-3}),
$$
\n
$$
\dots
$$
\n
$$
h_0 = \alpha_0^2 + \beta_1 \gamma_{-1} + \beta_{-1} \gamma_1,
$$
\n
$$
h_{-1} = \beta_{-1} \gamma_{-1}.
$$

Evidently, $H(z)$ is invariant under the action of \widetilde{G}_+ . Therefore, every h_n is an integral of motion, and $h_{2N-1} = b^2$ is an absolute constant. With respect to the symplectic structure [\(15\)](#page-6-0), $h_{N-1}, \ldots h_{2N-2}$ give rise to non-trivial hamiltonian flows, we call them hamiltonians.

On the other hand, $h_{-1}, h_0, \ldots, h_{N-2}$ annihilate the Poisson bracket [\(15\)](#page-6-0), since

$$
\sum_{i=-1}^{2N-2} \sum_{a=1,2,3} W_{i,j}^{a,b} \frac{\partial h_n}{\partial \psi_{a,i}} = 0, \quad n = -1, 0, \dots, N-2,
$$

and so $\{h_n, \mathcal{F}\}=0$ for any $\mathcal{F} \in \mathcal{C}(\mathcal{M}_N)$. Thus,

(18)
$$
h_{-1} = c_{-1}, \quad h_0 = c_0, \quad \dots, \quad h_{N-2} = c_{N-2}
$$

serve as *constraints* on the symplectic manifold M_N . They fix an orbit $\mathcal O$ of dimension 2N, which serves as the finite phase space of a hamiltonian system in the sine(sinh)-Gordon hierarchy.

3.4. Real forms of $\mathfrak{sl}(2,\mathbb{C})$ **.** The algebra $\mathfrak{sl}(2,\mathbb{C})$ has two real forms: $\mathfrak{su}(2)$, and $\mathfrak{sl}(2,\mathbb{R})$. In the both cases, $h_{-1}, h_0, \ldots, h_{2N-2}$, and $h_{2N-1} = b^2$ are real.

If $\mathfrak{g} = \mathfrak{sl}(2,\mathbb{R})$, then all coordinates $\beta_{2m-1}, \gamma_{2m-1}, \alpha_{2m}, m = 0, \ldots N-1$, and also b are real. Thus, $h_{2N-1} = b^2 > 0$.

If $\mathfrak{g} = \mathfrak{su}(2)$, then $\alpha_{2m} = ia_{2m}$, $a_{2m} \in \mathbb{R}$, and $\beta_{2m-1} = -\overline{\gamma}_{2m-1}$, $b = ib$, where $b \in \mathbb{R}$. Thus, $h_{2N-1} = b^2 = -b^2 < 0$. We also have

(19)
$$
h_{-1} = \beta_{-1} \gamma_{-1} = -|\gamma_{-1}|^2 < 0.
$$

3.5. Sine(sinh)-Gordon equation. Let h_{2N-2} give rise to a stationary flow with parameter x, and h_{N-1} give rise to an evolutionary flow with parameter t:

(20)
$$
\frac{d\psi_{a,\ell}}{dx} = \{h_{2N-2}, \psi_{a,\ell}\}, \qquad \frac{d\psi_{a,\ell}}{dt} = \{h_{N-1}, \psi_{a,\ell}\}.
$$

In more detail, the stationary flow is

(21a) $\frac{d\beta_{-1}}{dx} = 2\alpha_{2N-2}\beta_{-1}, \qquad \frac{d\gamma_{-1}}{dx} = -2\alpha_{2N-2}\gamma_{-1},$

(21b)
$$
\frac{d\alpha_{2m}}{dx} = b(\gamma_{2m-1} - \beta_{2m-1}), \quad m = 0, ..., N-1,
$$

(21c)
$$
\frac{d\beta_{2m-1}}{dx} = 2\alpha_{2N-2}\beta_{2m-1} - 2b\alpha_{2m-2},
$$

(21d)
$$
\frac{d\gamma_{2m-1}}{dx} = -2\alpha_{2N-2}\gamma_{2m-1} + 2b\alpha_{2m-2}, \quad m = 1, ..., N.
$$

The evolutionary flow is

(22a)
$$
\frac{d\alpha_{2m-2}}{dt} = \gamma_{-1}\beta_{2m-1} - \beta_{-1}\gamma_{2m-1}, \quad m = 1, ..., N-1,
$$

(22b)
$$
\frac{d\beta_{2m-1}}{dt} = 2\alpha_{2m}\beta_{-1}, \qquad \frac{d\gamma_{2m-1}}{dt} = -2\alpha_{2m}\gamma_{-1}, \quad m = 0, \dots, N-1,
$$

(22c)
$$
\frac{d\alpha_{2N-2}}{dt} = b(\gamma_{-1} - \beta_{-1}).
$$

The sine(sinh)-Gordon equation is obtained from the equality

(23)
$$
\frac{d\alpha_{2N-2}}{dt} = \frac{d\alpha_0}{dx}.
$$

Indeed, from [\(21a\)](#page-7-0) one can see that $\alpha_{2N-2} = \frac{1}{2} d \ln \beta_{-1}/dx = -\frac{1}{2} d \ln \gamma_{-1}/dx$. On the other hand, [\(22b\)](#page-8-0) implies $\alpha_0 = \frac{1}{2} d \ln \beta_{-1}/dt = -\frac{1}{2} d \ln \gamma_{-1}/dt$.

In the case of $\mathfrak{g} = \mathfrak{su}(2)$, we have $\beta_{-1} = -\overline{\gamma}_{-1}$, and so $-|\gamma_{-1}|^2 = c_{-1} = \text{const}$, as follows from [\(18\)](#page-7-1). Let $\gamma_{-1} = \imath r \exp(i\phi)$, and $\beta_{-1} = \imath r \exp(-i\phi)$, where ϕ is a function of x and t, and $r = \sqrt{-c_{-1}} = |\gamma_{-1}|$ is a constant. Then [\(21a\)](#page-7-0) and [\(22a\)](#page-8-1) implies, respectively,

$$
\alpha_{2N-2} = \frac{i}{2} \frac{d\phi}{dx}, \qquad \alpha_0 = \frac{i}{2} \frac{d\phi}{dt}.
$$

Finally, [\(22c\)](#page-8-2) gives the sine-Gordon equation (b = $ib, b \in \mathbb{R}$)

(24)
$$
\frac{d^2 \phi}{dt dx} = 4br \sin \phi.
$$

In the case of $\mathfrak{g} = \mathfrak{sl}(2,\mathbb{R})$, let $\gamma_{-1} = r \exp(\phi)$, $\beta_{-1} = r \exp(-\phi)$, where ϕ is a function of x and t, and $r = \sqrt{c-1} = \text{const}$, where c_{-1} is supposed to be positive. Then [\(22c\)](#page-8-2) gives the sinh-Gordon equation ($b \in \mathbb{R}$):

(25)
$$
\frac{\mathrm{d}^2 \phi}{\mathrm{d} t \mathrm{d} x} = -4 \mathrm{b} r \sinh \phi.
$$

REMARK 2. Note, that the sine(sinh)-Gordon equation arises when $N \geq 2$. If $N = 1$, there exists the only hamiltonian $h_0 = \alpha_0^2 + b(\beta_{-1} + \gamma_{-1})$, which produces the stationary flow

$$
\frac{d\beta_{-1}}{dx} = 2\alpha_0 \beta_{-1}, \qquad \frac{d\gamma_{-1}}{dx} = -2\alpha_0 \gamma_{-1}, \qquad \frac{d\alpha_0}{dx} = b(\gamma_{-1} - \beta_{-1}).
$$

From the flow of h_0 we find: $\alpha_0 = \frac{1}{2} d \ln \beta_{-1}/dx = -\frac{1}{2} d \ln \gamma_{-1}/dx$. With the same substitution for γ_{-1} and β_{-1} , where ϕ is a function of x we come to the stationary equation, which is the *equation of motion of a simple pendulum* if $\mathfrak{g} = \mathfrak{su}(2)$:

$$
\frac{\mathrm{d}^2 \phi}{\mathrm{d} x^2} = 4br \sin \phi.
$$

3.6. Zero curvature representation. The system of dynamic equations [\(20\)](#page-7-2) admits the matrix form

$$
\frac{\mathrm{d}\Psi}{\partial x} = [\Psi, \nabla h_{2N-2}], \qquad \frac{\mathrm{d}\Psi}{\partial t} = [\Psi, \nabla h_{N-1}],
$$

where ∇h_n denotes the matrix gradient of h_n , namely,

$$
\nabla h_n = \sum_{i=-1}^{2N-1} \sum_{a=1,2,3} \frac{\partial h_n}{\partial \psi_{a,i}} \mathbf{Z}_{a,2N-i-2}.
$$

The matrix gradient of each flow has a complementary matrix A, such that

$$
[\Psi,\nabla h_n]=[\Psi,\mathbf{A}].
$$

Unlike ∇h_n , the complementary matrix A is defined uniquely in all $\mathcal{M}_{\tilde{N}}$, $\tilde{N} \geqslant N$. Actually,

(26)
$$
\frac{d\Psi}{\partial x} = [\Psi, A_{st}], \qquad \frac{d\Psi}{\partial t} = [\Psi, A_{ev}],
$$

$$
A_{st} = -\begin{pmatrix} \alpha_{2N-2} & b \\ b_{2} & -\alpha_{2N-2} \end{pmatrix}, \qquad A_{ev} = \begin{pmatrix} 0 & \frac{1}{z}\beta_{-1} \\ \gamma_{-1} & 0 \end{pmatrix}.
$$

The zero curvature representation for the KdV hierarchy has the form

$$
\frac{\mathrm{dA}_{\mathrm{st}}}{\partial t} - \frac{\mathrm{dA}_{\mathrm{ev}}}{\partial x} = [A_{\mathrm{st}}, A_{\mathrm{ev}}].
$$

3.7. Summary. The affine algebra $\tilde{\mathfrak{g}} = \mathfrak{su}(2) \otimes \mathcal{L}(z, z^{-1})$ with the principal grading [\(10\)](#page-5-2) and the bilinear form [\(11\)](#page-5-0) is associated with the sine-Gordon hierarchy. Let \mathcal{M}_N be the manifold $\widetilde{\mathfrak{g}}^*/(\sum_{\ell<-1}^{\infty}\mathfrak{g}_{\ell})$, $\dim \mathcal{M}_N = 3N, N \in \mathbb{N}$. Evidently, $M_1 \subset M_2 \subset \cdots \subset M_N \subset M_{N+1} \subset \cdots$. Each manifold M_N is equipped with the symplectic structure [\(15\)](#page-6-0). Under the action of the loop group $\tilde{G}_{+} = \exp(\tilde{\mathfrak{g}}_{+})$ a manifold \mathcal{M}_N splits into orbits \mathcal{O} , each defined by the system of N constraints [\(18\)](#page-7-1). Each orbit serves as a phase space of dimension 2N for a hamiltonian system integrable in the Liouville sense. On orbits within \mathcal{M}_N , $N \geq 2$, there exist two hamiltonians whose flows give rise to the sine-Gordon equation [\(25\)](#page-8-3). We call these flows stationary and evolutionary with parameters x and t, correspondingly.

4. Separation of variables

4.1. Spectral curve. The sine-Gordon hierarchy is associated with the family of hyperelliptic curves

(27)
$$
-w^2 + z^2 H(z) = 0,
$$

where H is defined by [\(16\)](#page-7-3). This equation of a genus N curve is obtained from the characteristic polynomial of Ψ , namely det $(\Psi(z) - (w/z)) = 0$. All parameters of the curve are integrals of motion: hamiltonians $h_{N-1}, \ldots, h_{2N-2}$, constraints h_{-1} , ..., h_{N-2} , and $h_{2N-1} = b^2 = \text{const.}$

4.2. Canonical coordinates. As shown in [[15](#page-27-8)], variables of separation in the hierarchy of the sine-Gordon equation are given by a positive non-special divisor of degree equal to the genus of a spectral curve. In fact, pairs of coordinates of the points serve as quasi-canonical variables, and so lead to separation of variables. Below, we briefly explain how to find the required points in the sine-Gordon hierarchy, and prove that pairs of coordinates of N points serve as quasi-canonical variables.

Recall, that the symplectic manifold \mathcal{M}_N , with 3N coordinates [\(13\)](#page-6-1) on it, splits into orbits O fixed by N constraints, and so dim $O = 2N$. Thus, β_{2m-1} , $m = 0$, \ldots , $N-1$ are eliminated with the help of the constraints. Since the expressions [\(17\)](#page-7-4) are linear with respect to β_{2m-1} , it is convenient to present them in the matrix form. The constraints are written as

$$
\Gamma_{\rm c}\boldsymbol{\beta} + \mathsf{A}_{\rm c} = \boldsymbol{c},
$$

where

$$
\Gamma_{c} = \begin{pmatrix}\n b & 0 & 0 & \dots & 0 \\
 0 & \gamma_{-1} & \gamma_{1} & \dots & \gamma_{2N-3} \\
 \vdots & 0 & \gamma_{-1} & \ddots & \vdots \\
 0 & \vdots & \ddots & \ddots & \gamma_{1} \\
 0 & 0 & \dots & 0 & \gamma_{-1}\n\end{pmatrix}, \quad \beta = \begin{pmatrix}\n b \\
 \beta_{2N-3} \\
 \vdots \\
 \beta_{1} \\
 \beta_{-1}\n\end{pmatrix},
$$
\n
$$
\mathsf{A}_{c} = \begin{pmatrix}\n 0 \\
 \sum_{k=0}^{N-2} \alpha_{2(N-2-k)} \alpha_{2k} \\
 \vdots \\
 \sum_{k=0}^{n} \alpha_{2(n-k)} \alpha_{2k} \\
 \vdots \\
 \alpha_{0}^{2} \\
 0\n\end{pmatrix}, \quad c = \begin{pmatrix}\n b^{2} \\
 c_{N-2} \\
 \vdots \\
 c_{n} \\
 c_{0} \\
 c_{-1}\n\end{pmatrix}.
$$

The first equation is an identity. From [\(28\)](#page-9-0) we find

(29)
$$
\beta = \Gamma_c^{-1}(c - A_c).
$$

Expressions for the hamiltonians are

(30) $\Gamma_{h}\beta + A_{h} = h,$

where

$$
\Gamma_{\rm h} = \begin{pmatrix}\n\gamma_{2N-3} & b & 0 & \dots & 0 \\
\vdots & \gamma_{2N-3} & \ddots & \ddots & \vdots \\
\gamma_1 & \vdots & \ddots & b & 0 \\
\gamma_{-1} & \gamma_1 & \dots & \gamma_{2N-3} & b\n\end{pmatrix},
$$
\n
$$
A_{\rm h} = \begin{pmatrix}\n\alpha_{2N-2}^2 \\
\sum_{k=n}^{N-1} \alpha_{2(N-1-k+n)} \alpha_{2k} \\
\vdots \\
\sum_{k=0}^{N-1} \alpha_{2(N-1-k)} \alpha_{2k}\n\end{pmatrix}, \quad \mathbf{h} = \begin{pmatrix}\nh_{2N-2} \\
\vdots \\
h_{N-1+n} \\
\vdots \\
h_{N-1}\n\end{pmatrix}.
$$

Substituting [\(29\)](#page-10-0) into [\(30\)](#page-10-1), we obtain

(31)
$$
\boldsymbol{h} = \Gamma_{\mathrm{h}} \Gamma_{\mathrm{c}}^{-1} (\boldsymbol{c} - \mathsf{A}_{\mathrm{c}}) + \mathsf{A}_{\mathrm{h}}.
$$

On the other hand, hamiltonians h_{2N-2} , ..., h_{N-1} can be found from the equation of the spectral curve (27) , taken at N points which form a non-special^{[1](#page-10-2)} divisor. Namely, with $i = 1, \ldots, N$,

 $-w_i^2 + b^2 z_i^{2N+1} + h_{2N-2} z_i^{2N} + \cdots + h_{N-1} z_i^{N+1} + c_{N-2} z_i^{N} + \cdots + c_{-1} z_i = 0,$

or in the matrix form

$$
-\boldsymbol{w}+Z_c\boldsymbol{c}+Z_h\boldsymbol{h}=0,
$$

¹A non-special divisor on a hyperelliptic curve of genus g is a degree g positive divisor which contains no pair of points in involution.

where

$$
Z_{c} = \begin{pmatrix} z_1^{2N+1} & z_1^N & \dots & z_1 \\ z_2^{2N+1} & z_2^N & \dots & z_2 \\ \vdots & \vdots & \ddots & \vdots \\ z_N^{2N+1} & z_N^N & \dots & z_N \end{pmatrix}, \quad Z_{h} = \begin{pmatrix} z_1^{2N} & \dots & z_1^{N+2} & z_1^{N+1} \\ z_2^{2N} & \dots & z_2^{N+2} & z_2^{N+1} \\ \vdots & \ddots & \vdots & \vdots \\ z_N^{2N} & \dots & z_N^{N+2} & z_N^{N+1} \end{pmatrix}, \quad \mathbf{w} = \begin{pmatrix} w_1^2 \\ w_2^2 \\ \vdots \\ w_N^2 \end{pmatrix}.
$$

The matrix Z_h is square and invertible. Thus,

(32)
$$
h = Z_h^{-1}(w - Z_c c).
$$

Equations [\(31\)](#page-10-3) and [\(32\)](#page-11-0) give the same hamiltonians. Therefore,

$$
\Gamma_h\Gamma_c^{-1}(\boldsymbol{c}-\boldsymbol{A}_c)+\boldsymbol{A}_h=\boldsymbol{Z}_h^{-1}\big(\boldsymbol{w}-\boldsymbol{Z}_c\boldsymbol{c}\big).
$$

Moreover, constants c can be taken arbitrarily, and so the corresponding coefficients coincide, as well as the remaining terms:

(33a)
$$
\Gamma_h \Gamma_c^{-1} = -Z_h^{-1} Z_c,
$$

(33b)
$$
-\Gamma_h \Gamma_c^{-1} A_c + A_h = Z_h^{-1} \boldsymbol{w}.
$$

From [\(33a\)](#page-11-1) we find

$$
Z_h\Gamma_h+Z_c\Gamma_c=0,
$$

which is equivalent to $\gamma(z_i) = 0$. Then from [\(33b\)](#page-11-2) we obtain

$$
Z_c A_c + Z_h A_h = \boldsymbol{w},
$$

which is equivalent to $w_i^2 - z_i^2 \alpha(z_i)^2 = 0$. Thus, points (z_i, w_i) are defined by

$$
\gamma(z_i) = 0,
$$
 $w_i^2 - z_i^2 \alpha(z_i)^2 = 0,$ $i = 1, ..., N.$

This result was firstly discovered in [[15](#page-27-8)].

THEOREM 1. Suppose an orbit $\mathcal{O} \subset \mathcal{M}_N$ has the coordinates $(\gamma_{2m-1}, \alpha_{2m}),$ $m = 0, \ldots, N-1$, as above. Then the new coordinates $(z_i, w_i), i = 1, \ldots, N$, defined by the formulas

(34)
$$
\gamma(z_i) = 0, \qquad w_i = \epsilon z \alpha(z_i), \quad i = 1, \ldots, N,
$$

where $\epsilon^2 = 1$, have the following properties:

- 1) a pair (z_i, w_i) is a point of the spectral curve (27) .
- $2)$ a pair (z_i, w_i) is quasi-canonically conjugate with respect to the Lie-Poisson bracket [\(15\)](#page-6-0):

(35)
$$
\{z_i, z_j\} = 0, \qquad \{z_i, w_j\} = \epsilon z_i^N \delta_{i,j}, \qquad \{w_i, w_j\} = 0.
$$

3) the canonical 1-form is

(36)
$$
\epsilon \sum_{i=1}^{N} z_i^{-N} w_i \mathrm{d} z_i.
$$

PROOF. Since z_i depend only on γ_{2m-1} , $m = 0, \ldots, N-1$, and the latter commute, we have $\{z_i, z_j\} = 0$. Next,

$$
\{z_i, w_j\} = \sum_{n+m=N-1}^{2N-2} \left(\frac{\partial z_i}{\gamma_{2m-1}} \frac{\partial w_j}{\alpha_{2n}} - \frac{\partial z_i}{\alpha_{2n}} \frac{\partial w_j}{\gamma_{2m-1}} \right) \{ \gamma_{2m-1}, \alpha_{2n} \}
$$

REALITY CONDITIONS FOR THE SINE-GORDON EQUATION AND QUASI-PERIODIC SOLUTIONS

$$
= \frac{\epsilon}{\gamma'(z_i)} \sum_{m+n=N-1}^{2N-2} z_i^m z_j^{n+1} \gamma_{2(m+n-N)+1} = \frac{\epsilon z_j}{\gamma'(z_i)} \frac{z_i^{N-1} \gamma(z_i) - z_j^{N-1} \gamma(z_j)}{z_i - z_j},
$$

since from [\(34\)](#page-11-3) we have

$$
\frac{\partial z_i}{\partial z_{m-1}} = -\frac{z_i^m}{\gamma'(z_i)}, \qquad \frac{\partial z_i}{\partial z_n} = 0, \qquad \frac{\partial w_i}{\partial z_n} = \epsilon z_i^{n+1},
$$

$$
\frac{\partial w_i}{\partial z_{m-1}} = -\frac{\epsilon z_i^m}{\gamma'(z_i)} (\alpha(z_i) + z_i \alpha'(z_i)).
$$

As $i \neq j$, it is evident that $\{z_i, w_j\} = 0$, due to $\gamma(z_i) = \gamma(z_j) = 0$. As $i = j$, we get

$$
\{z_i, w_i\} = \lim_{z_j \to z_i} \frac{\epsilon z_j}{\gamma'(z_i)} \frac{z_i^{N-1} \gamma(z_i) - z_j^{N-1} \gamma(z_j)}{z_i - z_j} = \epsilon z_i^N.
$$

Finally, we find

$$
\{w_i, w_j\} = \sum_{n+m=N-1}^{2N-2} \left(\frac{\partial w_i}{\gamma_{2m-1}} \frac{\partial w_j}{\alpha_{2n}} - \frac{\partial w_i}{\alpha_{2n}} \frac{\partial w_j}{\gamma_{2m-1}} \right) \{ \gamma_{2m-1}, \alpha_{2n} \}
$$

$$
= \epsilon^2 \sum_{m+n=N-1}^{2N-2} \left(z_i^m z_j^{n+1} \frac{\alpha(z_i) + z_i \alpha'(z_i)}{\gamma'(z_i)} - z_i^{n+1} z_j^m \frac{\alpha(z_j) + z_j \alpha'(z_j)}{\gamma'(z_j)} \right) \gamma_{2(m+n-N)+1}
$$

$$
= \epsilon^2 \frac{z_i^{N-1} \gamma(z_i) - z_j^{N-1} \gamma(z_j)}{z_i - z_j} \left(z_j \frac{\alpha(z_i) + z_i \alpha'(z_i)}{\gamma'(z_i)} - z_i \frac{\alpha(z_j) + z_j \alpha'(z_j)}{\gamma'(z_j)} \right).
$$

Thus, $\{w_i, w_j\} = 0$, due to $\gamma(z_i) = \gamma(z_j) = 0$.

Then [\(36\)](#page-11-4) follows from the fact that pairs (z_i, w_i) , $i = 1, \ldots, N$, are quasicanonically conjugate and obey (35) .

In what follows we assign $\epsilon = 1$.

4.3. Summary. An orbit $\mathcal{O} \subset \mathcal{M}_N$, which serves as a phase space of dimension 2N, is completely parameterized by non-canonical variables $\gamma_{2n-1}, \alpha_{2n}, n = 0$, ..., $N-1$. Points $\{(z_i, w_i)\}_{i=1}^N$ of the spectral curve [\(27\)](#page-9-1) serve as variables of separation, which are quasi-canonical, see Theorem [1.](#page-11-6) The relation [\(34\)](#page-11-3) between non-canonical and quasi-canonical variables is closely connected to the solution [\(9\)](#page-4-0) of the Jacobi inversion problem. Indeed, the both systems of equations define the same point within the fundamental domain of the Jacobian variety of the spectral curve, as we see below.

5. Algebro-geometric integration

5.1. Uniformization of the spectral curve. Separation of variables leads to the Jacobi inversion problem for a non-special divisor of N points $\{(z_k, w_k)\}_{k=1}^N$ on the spectral curve (27) , which is a hyperelliptic curve of genus N

(37)
$$
0 = F(z, w) \equiv -w^2 + b^2 z^{2N+1} + h_{2N-2} z^{2N} + \dots + h_{N-1} z^{N+1} + c_{N-2} z^N + \dots + c_0 z^2 + c_{-1} z.
$$

Not normalized differentials of the first and second kinds acquire the form

(38a)
$$
\mathrm{d}u_{2n-1} = \frac{\mathrm{b}z^{N-n}\mathrm{d}z}{\partial_w F(z,w)}, \quad n = 1,\ldots,g,
$$

(38b)
$$
\mathrm{d}r_{2n-1} = \frac{\mathrm{d}z}{\mathrm{b}\partial_w F(z,w)} \sum_{j=1}^{2n-1} (2n-j) h_{2N-j} z^{N+n-j},
$$

which follow from $(5a)$, $(5b)$, after reducing (37) to the form (3) by applying the transformation: $z \mapsto x$, $w \mapsto by$, $F(z, w) \mapsto f(x, y) = F(x, by)/b^2$. Note, that one of the branch points of [\(37\)](#page-12-0) is fixed at 0, and so the corresponding canonical form [\(3\)](#page-1-0) has $\lambda_{4N+2} = 0$.

On the curve [\(37\)](#page-12-0), the Abel pre-image of $\mathcal{A}(D) = u \in \text{Jac}(\mathcal{V})$, which is a non-special positive divisor $D = \sum_{k=1}^{N} (z_k, w_k)$, is obtained from the system

(39a)
$$
z^{N} - \sum_{k=1}^{N} z^{N-k} \wp_{1,2k-1}(u) = 0,
$$

(39b)
$$
\frac{2w}{b} + \sum_{k=1}^{N} z^{N-k} \wp_{1,1,2k-1}(u) = 0.
$$

On the other hand, the N values z_i are zeros of the polynomial $\gamma(z)$, and the N values w_i are obtained from $w_i = z_i \alpha(z_i)$, cf. [\(34\)](#page-11-3). Thus,

(40a)
$$
\gamma_{2(N-k)-1} = -b\wp_{1,2k-1}(u), \quad k = 1,\ldots,N;
$$

(40b)
$$
\alpha_{2N-2} = -\frac{b\wp_{1,1,2N-1}(u)}{2\wp_{1,2N-1}(u)},
$$

(40c)
$$
\alpha_{2(N-k)} = -\frac{b}{2}\wp_{1,1,2k-3}(u) + \frac{b}{2}\frac{\wp_{1,2k-3}(u)\wp_{1,1,2N-1}(u)}{\wp_{1,2N-1}(u)}, \quad k = 2,\ldots,N.
$$

Taking into account the relations between \wp -functions, obtained from [\(22a\)](#page-8-1), we find

(40c')
$$
\alpha_{2(N-k)} = -\frac{b\wp_{1,2k-1,2N-1}(u)}{2\wp_{1,2N-1}(u)}, \quad k = 1,\ldots,N,
$$

and also

(40d)
$$
\beta_{-1} = \frac{-bc_{-1}}{\wp_{1,2N-1}(u)},
$$

(40e)
$$
\beta_{2(N-k)-1} = b \frac{\wp_{2k-1,2N-1}(u)}{\wp_{1,2N-1}(u)}, \quad k = 1, ..., N-1.
$$

Finally, we obtain the relation

(41)
$$
\gamma_{-1} = \imath r \exp(-\imath \phi) = -\mathbf{b}\wp_{1,2N-1}(u),
$$

which produces a finite-gap solution ϕ of the sine-Gordon equation [\(25\)](#page-8-3) in the 2Ndimensional phase space. This solution coincides with the one proposed in [[10](#page-27-5), Theorem 4.13].

Remark 3. The sine-Gordon equation comes from [\(22c\)](#page-8-2), which in terms of ℘-functions acquires the form

(42)
$$
\frac{d}{du_{2N-1}} \frac{\wp_{1,1,2N-1}(u)}{\wp_{1,2N-1}(u)} = 2\left(\wp_{1,2N-1}(u) - \frac{\lambda_{4N}}{\wp_{1,2N-1}(u)}\right).
$$

This relation corresponds to a curve of the form [\(3\)](#page-1-0) with $\lambda_{4N+2} = 0$.

REMARK 4. Parameters $\lambda_2, \ldots, \lambda_{2N}$ of [\(3\)](#page-1-0) serve as hamiltonians $h_{2N-2}, \ldots,$ h_{N-1} , since $h_n = b^2 \lambda_{4N-2-2n}$. Let $\mathcal{J}_{2N+2+2n}$ be functions of $\wp_{1,2i-1}, \wp_{1,1,2i-1}$, $i = 1, \ldots, N$, and $\lambda_{2k}, k = 1, \ldots, 2N + 1$, such that the equations

(43)
$$
\mathcal{J}_{2N+2+2n} = 0, \qquad n = 1, ..., N,
$$

give an algebraic model of $Jac(\mathcal{V})$ of the curve [\(3\)](#page-1-0), see [[8](#page-26-7), Sect. 4]. Then the hamiltonians, given by [\(17\)](#page-7-4), expressed in terms of \wp -functions by means of [\(40\)](#page-13-1), coincide with [\(43\)](#page-14-0) solved for $\lambda_2, \ldots, \lambda_{2N}$, provided $\lambda_{4N+2} = 0$.

5.2. Equation of motion in variables of separation. From [\(26\)](#page-9-2) we find

$$
\frac{d}{dx}\gamma(z) = -2\alpha_{2N-2}\gamma(z) + 2bz\alpha(z),
$$

$$
\frac{d}{dt}\gamma(z) = -2\gamma_{-1}\alpha(z),
$$

where all dynamic variables are functions of x and t. Therefore, zeros of $\gamma(z)$ are functions of x and t as well, namely $\gamma(z) = b \prod_{k=1}^{N} (z - z_k(x, t))$. Then

$$
\frac{\mathrm{d}}{\mathrm{d}x}\log\gamma(z) = -\frac{1}{z-z_k}\frac{\mathrm{d}z_k}{\mathrm{d}x} = -2\alpha_{2N-2} + 2bz \frac{\alpha(z)}{\gamma(z)},
$$

$$
\frac{\mathrm{d}}{\mathrm{d}t}\log\gamma(z) = -\frac{1}{z-z_k}\frac{\mathrm{d}z_k}{\mathrm{d}t} = -2\gamma_{-1}\frac{\alpha(z)}{\gamma(z)}, \quad k = 1, \dots, N.
$$

Taking into account [\(34\)](#page-11-3), we find as $z \to z_k$, $k = 1, \ldots, N$,

(46)
$$
\frac{dz_k}{dx} = -\frac{2w_k}{\prod_{j \neq k}^{N} (z_k - z_j)}, \qquad \frac{dz_k}{dt} = -\frac{2w_k \prod_{j \neq k} (-z_j)}{\prod_{j \neq k} (z_k - z_j)}.
$$

Let $D = \sum_{k=1}^{N} (z_k, w_k)$, and the N points satisfy [\(34\)](#page-11-3). The Abel image of D

$$
u = \mathcal{A}(D) = \sum_{k=1}^{N} \int_{\infty}^{(z_k, w_k)} \mathrm{d}u = \sum_{k=1}^{N} \int_{\infty}^{(z_k, w_k)} \begin{pmatrix} 1 \\ z \\ \vdots \\ z^{N-1} \end{pmatrix} \frac{\mathrm{bd}z}{-2w}
$$

depends on x and t. Applying [\(46\)](#page-14-1), we find

$$
\frac{du_{2n-1}}{dx} = \sum_{k=1}^{N} \frac{bz_k^{N-n}}{-2w_k} \frac{dz_k}{dx} = \sum_{k=1}^{N} \frac{bz_k^{N-n}}{\prod_{j \neq k}^{N} (z_k - z_j)} = b\delta_{n,1},
$$

$$
\frac{du_{2n-1}}{dt} = \sum_{k=1}^{N} \frac{bz_k^{N-n}}{-2w_k} \frac{dz_k}{dt} = \sum_{k=1}^{N} \frac{bz_k^{N-n} \prod_{j \neq k} (-z_j)}{\prod_{j \neq k} (z_k - z_j)} = b\delta_{n,N}.
$$

And so, we obtain

(48)
$$
u_1 = bx + C_1,
$$

$$
u_{2n-1} = bc_{2n-1} + C_{2n-1} = \text{const}, \quad n = 2, ..., N - 1,
$$

$$
u_{2N-1} = bt + C_{2N-1}.
$$

Here $c_{2n-1}, n = 2, \ldots, N-1$, are arbitrary real constants, and $\mathbf{C} = (C_1, C_3, \ldots, C_n)$ C_{2N-1}) is a constant vector chosen in a particular way, in order to guarantee desired properties of $\wp_{1,2N-1}$ -function. Thus, [\(41\)](#page-13-2) acquires the form

(49)
$$
\gamma_{-1}(x,t) = -b\wp_{1,2N-1}(b(x,c_3,\ldots,c_{2N-3},t)^t + \mathbf{C}),
$$

which produces a finite-gap solution of the sine(sinh)-Gordon equations.

In the sine-Gordon hierarchy, we have b = ib , and $r = \sqrt{-c-1}$. Note that $r/b = \sqrt{\lambda_{4N}}$, and according to [\(19\)](#page-7-5), and so values of γ_{-1} draw a circle of radius $\sqrt{\lambda_{4N}}$. Recalling the substitution for γ_{-1} , we find

(50a)
$$
\phi(x,t) = i \log \left(-\frac{b}{r} \wp_{1,2N-1} \left(i b(x,c_3,\ldots,c_{2N-3},t)^t + C \right) \right)
$$

In the sinh-Gordon hierarchy, b is real, and $r = \sqrt{c-1}$. Recalling the substitution for γ_{-1} , we find

(50b)
$$
\phi(x,t) = \log\left(-\frac{b}{r}\wp_{1,2N-1}(b(x,c_3,\ldots,c_{2N-3},t)^t + C)\right).
$$

In what follows we focus on the sine-Gordon hierarchy. Reality conditions for the sinh-Gordon hierarchy, associated with the affine algebra $\mathfrak{sl}(2,\mathbb{R})\otimes \mathfrak{L}(z,z^{-1}),$ are similar to reality conditions for the KdV equation, see [[7](#page-26-0)].

Below, we find the constant vector C such that $\wp_{1,2N-1}$ is bounded, and prove that $\wp_{1,2N-1}$ with the arguments as in [\(50a\)](#page-15-1) attains values on a circle of radius $\sqrt{\lambda_{4N}}$.

5.3. Summary. Uniformization of the spectral curve [\(37\)](#page-12-0) is given by [\(39\)](#page-13-3) in an implicit form. On the other hand, it produces explicit expressions [\(40\)](#page-13-1) for dynamic variables $\gamma_{2n-1}, \alpha_{2n}$. Coordinates u_1 and u_{2N-1} of the Jacobian variety of the spectral curve, up to a constant multiple b, serve as parameters x and t of the stationary and evolutionary flows, correspondingly.

6. Reality conditions

In this section we work with the spectral curve [\(37\)](#page-12-0) reduced to the canonical form [\(3\)](#page-1-0) with purely imaginary y, due to $w \mapsto \iota by$. Assuming that y is real, we consider the canonical form

(51)
$$
y^{2} + x^{2g+1} + \sum_{i=0}^{2g-1} \lambda_{2i+2} x^{2g-i} = 0,
$$

of the spectral curve, or shortly $y^2 + \Lambda(x) = 0$. Note that $\lambda_{4g+2} = 0$, since one branch point is fixed at 0.

All integrals of motion h_n in [\(37\)](#page-12-0) are required to be real. Therefore, parameters λ_n of the curve [\(51\)](#page-15-2) are real, which implies that branch points are complex conjugate or real. Let all but one finite branch points be complex conjugate, and this one is $(0, 0)$. In what follows, we prove that this requirement to spectral curves in the sine-Gordon hierarchy is a part of the reality conditions.

We enumerate branch points in the ascending order of real and imaginary parts, as shown on fig. [1.](#page-2-0)

PROPOSITION 1. Let the curve [\(51\)](#page-15-2) possesses the branch points $\{(e_k, 0)\}_{k=1}^{2g+1}$ such that $e_{2n+1} = 0$, and g pairs $\{e_{2i-1}, e_{2i} \mid i = 1, ..., n\}$, $\{e_{2i}, e_{2i+1} \mid i =$ $n+1,\ldots,g$ are complex conjugate. Then

(52)
$$
\lambda_{4g} = \prod_{i=1}^{n} e_{2i-1} e_{2i} \prod_{i=n+1}^{g} e_{2i} e_{2i+1} = \prod_{i=1}^{g} |e_{2i}|^2.
$$

PROOF. The equality is obtained by direct computations. Let the curve (51) has the form $y^2 + \Lambda(x) = 0$, and $\{e_k\}_{k=1}^{2g+1}$ be roots of Λ . Note, that each pair contains a root with an even index, say e_{2i} . The other root in each pair is the complex conjugate of e_{2i} , that is \bar{e}_{2i} . Namely, $e_{2i-1} = \bar{e}_{2i}$ as $i = 1, \ldots, n$, and $e_{2i+1} = \overline{e}_{2i}$ as $i = n+1, \ldots, g$. Thus, the product of roots in the *i*-th pair equals $|e_{2i}|^2$.

In the sine-Gordon hierarchy, the parameter $\lambda_{4g} = c_{-1}/b^2$ is positive, since $\mathfrak{su}(2)$ -algebra implies negative $c_{-1} = -r^2$ and $b^2 = -b^2$. Then

(53)
$$
\sqrt{\lambda_{4g}} = \prod_{i=1}^{g} |e_{2i}|.
$$

PROPOSITION 2. There are 2^g ways to compose the product [\(53\)](#page-16-0) from g branch points: by choosing one point from each pair.

6.1. Singularities of \wp **-functions.** Each collection of g branch points form a divisor whose Abel image is an even non-singular half-period. Odd non-singular and singular half-periods are Abel images of divisors of degrees less than q . As explained in subsection [2.6,](#page-4-1) each half-period is associated with a partition $\mathcal{I} \cup \mathcal{J}$ of indices of all branch points, where cardinality of $\mathcal I$ is equal to or less than q, that is $|\mathcal{I}| \leq g$. \wp -Functions have singularities at all half-periods with $|\mathcal{I}| < g$. Therefore, at half-periods corresponding to the collections of g branch points singled out by Proposition [2](#page-16-1) φ -functions have finite values.

The branch point divisors singled out by Proposition [2,](#page-16-1) by necessity, belong to the domain of $\wp_{1,2N-1}$ in [\(50a\)](#page-15-1). Indeed, an N-gap hamiltonian system in terms of variables of separation (z_i, w_i) splits into N independent systems, each with the coordinate z_i , the momenta w_i , and the hamiltonian given by the spectral curve equation [\(37\)](#page-12-0). Finite trajectories contain branch points as turning points.

THEOREM 2. Let a hyperelliptic curve (51) of genus g have branch points as defined in Proposition [1.](#page-15-3) Then with a choice of cycles as on fig. [1](#page-2-0) and the standard not normalized holomorphic differentials [\(5a\)](#page-2-1), the period lattice formed by each pair of periods ω_k , ω'_k is rhombic, with generators:

1) ω'_k and $\omega_k - \omega'_k$ as $k = 1, \ldots, n$, where

(54a)
$$
\omega_k \in i \mathbb{R}, \qquad \text{Im}\,\omega_k = 2\,\text{Im}\,\omega'_k;
$$

2) ω_k and $\omega_k + \omega'_k$ as $k = n + 1, \ldots, g$, where

(54b)
$$
\omega'_{k} \in \mathbb{R}, \qquad 2 \operatorname{Re} \omega_{g} = -2\omega'_{g} + \omega'_{g-1},
$$

$$
2 \operatorname{Re} \omega_{k} = -2\omega'_{k} + \omega'_{k-1} + \omega'_{k+1}, \quad k < g.
$$

PROOF. Let a hyperelliptic curve [\(51\)](#page-15-2) have the form $y^2 + \Lambda(x) = 0$. If branch points are as described in Proposition [1,](#page-15-3) then $\Lambda(x) < 0$ at $x < e_{2n-1}$ and $\Lambda(x) > 0$ at $x > e_{2n-1}$. As $k = 1, \ldots, n$, we have $\omega_k = \mathcal{A}(e_{2k}) - \mathcal{A}(\bar{e}_{2k})$. Expanding the expression for ω_k in the Taylor series about the point $P_0 = (\text{Re } e_{2k}, 2\sqrt{-\Lambda(\text{Re } e_{2k})})$, where $du_{2n-1} = x^{g-n} dx/(2\sqrt{-\Lambda(x)})$ is real-valued, and the integral $\mathcal{A}(P_0)$ is taken along the real axis, we immediately obtain that $\omega_k \in \mathbb{R}$.

Next, consider $\omega'_k = \sum_{i=k}^g \left(\mathcal{A}(\bar{e}_{2i}) - \mathcal{A}(e_{2i}) \right)$ as $k = n+1, \ldots, g$. Expanding each term of the sum on the right-hand side in the Taylor series about $P_0 = (\text{Re }e_{2i}, 2\sqrt{\Lambda(\text{Re }e_{2i})})$, where $du_{2n-1} = x^{g-n}dx/(2i\sqrt{\Lambda(x)})$ is purely imaginary, we see that $\omega'_k \in \mathbb{R}$.

As $k = 1, \ldots, n$, we have

$$
\omega'_{k} = -\bigg(\sum_{j=k}^{n} \Big(\mathcal{A}(e_{2j+1}) - \mathcal{A}(e_{2j})\Big) - \sum_{j=n+1}^{g} \Big(\mathcal{A}(\bar{e}_{2j}) - \mathcal{A}(e_{2j})\Big)\bigg).
$$

The second sum is real, as shown above. Each term of the first sum is computed as follows

(55)
$$
\mathcal{A}(e_{2j+1}) - \mathcal{A}(e_{2j}) = (\mathcal{A}(e_{2j+1}) - \mathcal{A}(Re\,e_{2j+1})) + (\mathcal{A}(Re\,e_{2j+1}) - \mathcal{A}(Re\,e_{2j})) + (\mathcal{A}(Re\,e_{2j}) - \mathcal{A}(e_{2j})).
$$

Note that $e_{2n+1} = 0$, and so the last term vanish as $j = n$. As we see from fig. [1,](#page-2-0)

Im
$$
(\mathcal{A}(Re e_{2j}) - \mathcal{A}(e_{2j})) = -\frac{1}{2} Im \omega_j
$$
, $j = k, ..., n$
\nIm $(\mathcal{A}(e_{2j+1}) - \mathcal{A}(Re e_{2j+1})) = \frac{1}{2} Im \omega_{j+1}$, $j = k, ..., n - 1$.

and the middle term on the right hand side of [\(55\)](#page-17-0) is real-valued. Thus, we obtain $\text{Im}\,\omega'_k = \frac{1}{2}\,\text{Im}\,\omega_k$, cf. [\(54a\)](#page-16-2).

Finally, as $k = n+1, \ldots, g$, we have $\omega_k = \mathcal{A}(e_{2k}) - \mathcal{A}(e_{2k-1})$, which is computed similar to [\(55\)](#page-17-0). Namely,

(56)
$$
\mathcal{A}(e_{2k}) - \mathcal{A}(e_{2k-1}) = (\mathcal{A}(e_{2k}) - \mathcal{A}(Re\,e_{2k})) + (\mathcal{A}(Re\,e_{2k}) - \mathcal{A}(Re\,e_{2k-1})) + (\mathcal{A}(Re\,e_{2k-1}) - \mathcal{A}(e_{2k-1})).
$$

Now we are interested in the real part of [\(56\)](#page-17-1), and we find that the middle term on the right hand side is purely imaginary, since du_{2n-1} is purely imaginary in this case. Then

$$
\operatorname{Re}\left(\mathcal{A}(e_{2k}) - \mathcal{A}(\operatorname{Re} e_{2k})\right) = -\left(\frac{1}{2}\omega'_k - \sum_{j=k+1}^g \frac{1}{2}\omega'_j\right), \qquad k = n+1,\ldots,g
$$

$$
\operatorname{Re}\left(\mathcal{A}(\operatorname{Re} e_{2k-1}) - \mathcal{A}(e_{2k-1})\right) = \frac{1}{2}\omega'_{k-1} - \sum_{j=k}^g \frac{1}{2}\omega'_j, \qquad k = n+2,\ldots,g.
$$

This implies $\text{Re}\,\omega_k = -\omega'_k + \frac{1}{2}\omega'_{k-1} + \frac{1}{2}\omega'_{k+1}$ as $k < g$, and $\text{Re}\,\omega_g = -\omega'_g + \frac{1}{2}\omega'_{g-1}$, cf. [\(54b\)](#page-16-3).

The relations [\(54\)](#page-16-4) guarantee that the period lattice formed by each pair of periods ω_k , ω'_k is rhombic, with the indicated generators.

Let $\bar{\mathfrak{J}} = \mathbb{C}^g$ be the vector space where the Jacobian variety of the curve [\(51\)](#page-15-2) is embedded. Let $\text{Re}\,\bar{\mathfrak{J}}\sim\mathbb{R}^g$ be the span of real axes of $\bar{\mathfrak{J}}$ over \mathbb{R} , and $\text{Im}\,\bar{\mathfrak{J}}\sim\mathbb{R}^g$ be the span of imaginary axes of $\bar{3}$ over R. Then $\bar{3} = \text{Re } \bar{3} \oplus \text{Im } \bar{3}$. As seen from [\(50a\)](#page-15-1), the domain of $\wp_{1,2N-1}$ is a subspace of $\bar{3}$ parallel to Im $\tilde{3}$ where \wp -functions are bounded.

Theorem 3. All half-periods obtained from the branch point divisors singled out by Proposition [2](#page-16-1) are located on the same subspace $C + \text{Im} \, \mathfrak{J}$, where

(57)
$$
C = \sum_{i=0}^{[(g-1)/2]} \frac{1}{2} \omega_{g-2i} + \sum_{k=1}^{g} \frac{1}{2} \omega'_k = \mathcal{A} \left(\sum_{i=1}^{g} e_{2i} \right).
$$

PROOF. With the choice of cycles as on fig. [1,](#page-2-0) we have the following correspondence between sets $\mathcal I$ of cardinality 1 and half-periods:

(58)
$$
\{2k-1\} \sim \frac{1}{2}\omega'_k + \sum_{i=1}^{k-1} \frac{1}{2}\omega_i, \quad \{2k\} \sim \frac{1}{2}\omega'_k + \sum_{i=1}^k \frac{1}{2}\omega_i, \quad k = 1, ..., g,
$$

$$
\{2g+1\} \sim \sum_{i=1}^g \frac{1}{2}\omega_i.
$$

Here and below \sim denotes the congruence relation. Eq. [\(58\)](#page-18-0) implies

$$
\{2, 4, \ldots, 2g\} \sim \sum_{i=0}^{[(g-1)/2]} \frac{1}{2}\omega_{g-2i} + \sum_{k=1}^{g} \frac{1}{2}\omega'_k.
$$

All points of the subspace $C + \text{Im } \overline{\mathfrak{J}}$ have the same ReC. Indeed, using [\(58\)](#page-18-0) we find, that in the first *n* complex conjugate pairs $e_{2i-1} = \overline{e}_{2i}, e_{2i}$

$$
\operatorname{Re}\left(\frac{1}{2}\omega'_i+\sum_{j=1}^{i-1}\frac{1}{2}\omega_j\right)=\operatorname{Re}\left(\frac{1}{2}\omega'_i+\sum_{j=1}^{i}\frac{1}{2}\omega_j\right)=\operatorname{Re}\left(\frac{1}{2}\omega'_i\right), \quad i=1,\ldots,n,
$$

due to ω_i are purely imaginary, according to Proposition [2.](#page-16-5) In the next $g - n$ complex conjugate pairs $e_{2i}, e_{2i+1} = \overline{e}_{2i}$ we have

$$
\operatorname{Re}\left(\frac{1}{2}\omega'_i+\sum_{j=1}^i\frac{1}{2}\omega_j\right)\sim\operatorname{Re}\left(\frac{1}{2}\omega'_{i+1}+\sum_{j=1}^i\frac{1}{2}\omega_j\right)\sim\operatorname{Re}\left(\sum_{j=1}^i\frac{1}{2}\omega_j\right),\quad i=n+1,\ldots,g,
$$

due to each Re $\frac{1}{2}\omega'_i$ equals a linear combination of Re ω_i as follows from [\(54b\)](#page-16-3), and so congruent to 0. Therefore, the branch point divisors introduced by Proposition [2](#page-16-1) have the same real part, and so belong to the subspace $C + \text{Im } \tilde{\mathfrak{J}}$.

COROLLARY [1](#page-15-3). On a hyperelliptic curve defined in Proposition 1 all 2^{2g} halfperiods are located on 2^g subspaces parallel to $\text{Im} \bar{\mathfrak{J}}$, each subspace contains 2^g halfperiods. The half-periods on $C+\operatorname{Im} \overline{\mathfrak{J}}$ are defined by Proposition [2.](#page-16-1) Other subspaces are obtained by shifts generated from $\{\frac{1}{2}\omega_k \mid k = 1,\ldots,n\}$, and $\{\frac{1}{2}\omega'_k \mid k = n +$ $1, \ldots, g$.

At the same time, 2^{2g} half-periods are located on 2^g subspaces parallel to Re $\bar{3}$, each contains 2^g half-periods. The half-periods on $C + \text{Re } \overline{\mathfrak{J}}$ are defined by Proposition [2.](#page-16-1) Other subspaces are obtained by the same shifts.

COROLLARY 2. \wp -Functions are bounded on the subspaces $C+\text{Im}\,\mathfrak{J}$ and $C+\text{Re}\,\mathfrak{J}$ with C defined by (57) .

REMARK 5. Note, that $C = \omega K$, where K denotes the vector of Riemann constants, and corresponds to the divisor of branch points $\{e_{2i}\}_{i=1}^g$.

6.2. Hyperelliptic addition law. Below, we briefly recall the addition laws on hyperelliptic curves, formulated in [[11](#page-27-9)].

The addition operation on the Jacobian variety of a plane algebraic curve is defined by the entire rational function \mathcal{R}_{3g} of weight 3g. On a hyperelliptic curve V of genus q it has the form

$$
\mathcal{R}_{3g}(x,y) \equiv y\nu_y(x) + \nu_x(x),
$$

where with $\mathfrak{k} \in \mathbb{N}$ and $\nu_0 = 1$

$$
\nu_y(x) \equiv \sum_{i=1}^{\mathfrak{k}} \nu_{2i-2} x^{\mathfrak{k}-i}, \quad \nu_x(x) \equiv \sum_{i=1}^{3\mathfrak{k}-1} \nu_{2i-1} x^{3\mathfrak{k}-1-i}, \quad \text{if} \quad g = 2\mathfrak{k} - 1, \text{ or}
$$

$$
\nu_y(x) \equiv \sum_{i=1}^{\mathfrak{k}} \nu_{2i-1} x^{\mathfrak{k}-i}, \quad \nu_x(x) \equiv \sum_{i=1}^{3\mathfrak{k}+1} \nu_{2i-2} x^{3\mathfrak{k}+1-i}, \quad \text{if} \quad g = 2\mathfrak{k}.
$$

The polynomial function \mathcal{R}_{3g} contains $2g$ arbitrary coefficients, and so is uniquely defined by fixing 2g points in its divisor of zeros. Let 2g points of V with no pairs of points in involution form two non-special divisors D_I and D_{II} , and $\mathcal{A}(D_I) = u_I$, $\mathcal{A}(D_{\text{II}}) = u_{\text{II}}$. Let D_{III} be such a divisor that $(\mathcal{R}_{3g})_0 = D_{\text{I}} + D_{\text{II}} + D_{\text{III}}$, and $A(D_{\text{III}}) = u_{\text{III}}$. This implies $u_{\text{I}} + u_{\text{II}} + u_{\text{III}} = 0$.

Equations [\(9\)](#page-4-0) allow to reduce \mathcal{R}_{3g} to a degree $g-1$ polynomial in x with coefficients expressed in terms of ν_k and basis \wp -functions, namely: $\wp_{1,2i-1}, \wp_{1,1,2i-1}$, $i = 1, \ldots, g$. Pre-images of u_{I} , u_{II} , u_{III} are supposed to form the divisor of zeros of \mathcal{R}_{3g} , which leads to the system of equations

(59)
$$
\begin{pmatrix} 1_g & \mathbf{Q}(u_{\mathrm{I}}) \\ 1_g & \mathbf{Q}(u_{\mathrm{II}}) \\ 1_g & \mathbf{Q}(u_{\mathrm{III}}) \end{pmatrix} \begin{pmatrix} \bar{\nu} \\ \nu \end{pmatrix} = - \begin{pmatrix} \mathbf{q}(u_{\mathrm{I}}) \\ \mathbf{q}(u_{\mathrm{II}}) \\ \mathbf{q}(u_{\mathrm{III}}) \end{pmatrix},
$$

where $\bar{\nu} = (\nu_{g+2}, \nu_{g+4}, \dots, \nu_{3g})^t$, and $\nu = (\nu_g, \dots, \nu_2, \nu_1)^t$.

Let u_{I} and u_{II} be given, then $u_{\text{III}} = -u_{\text{I}} - u_{\text{II}}$, as follows from the fact that $(\mathcal{R}_{3g})_0 \sim 3g\infty$. Thus, equations for u_I and u_{II} in [\(59\)](#page-19-0) are used to find ν and $\bar{\nu}$ by means of Cramer's rule. Values $\wp_{1,2i-1}(u_{\text{III}})$ are obtained from the equality

$$
\nu_y^2 f(x, -\nu_x/\nu_y) = \mathcal{R}_{2g}(x, y; u_1) \mathcal{R}_{2g}(x, y; u_{II}) \mathcal{R}_{2g}(x, y; u_{III}).
$$

Next, values $\wp_{1,1,2i-1}(u_{\text{III}})$ are computed from the equations for u_{III} in [\(59\)](#page-19-0). In more detail, hyperelliptic laws are presented in [[7](#page-26-0)], as well as explicit expression for $\wp_{1,2i-1}(u_{\text{III}})$ and $\wp_{1,1,2i-1}(u_{\text{III}})$.

For the purpose of the present paper, we need an expression for $\wp_{1,2g-1}$ -function of the form

(60)
$$
\varphi_{1,2g-1}(u_{\text{III}}) = (-1)^{g-1} \frac{\nu_{3g}^2 - \lambda_{4g+2} \nu_{g-1}^2}{\varphi_{1,2g-1}(u_1)\varphi_{1,2g-1}(u_{\text{II}})}.
$$

Note, that $\lambda_{4g+2} = 0$ in the case of sine(sinh)-Gordon hierarchy.

6.3. Real-valued \wp -functions. Recall, that the function [\(49\)](#page-14-2) in the sine-Gordon hierarchy satisfies the constraint [\(19\)](#page-7-5), which is equivalent to

$$
|\wp_{1,2g-1}(ib(x,c_3,\ldots,c_{2g-3},t)^t + \mathbf{C})|^2 = \lambda_{4g},
$$

where $N = g$. Below we prove, that this equality holds if C is defined by [\(57\)](#page-17-2). Evidently, values of $\wp_{1,2i-1}(C)$ are symmetric functions on $\{e_{2i}\}_{i=1}^g$ computed from [\(9a\)](#page-4-2), and $\wp_{1,1,2i-1}(C) = 0$ as follows from [\(9b\)](#page-4-3). Let $u_I \equiv C$, and $u_{II} =$ $\iota b(\mathbf{x}, \mathbf{c}_3, \dots, \mathbf{c}_{2N-3}, \mathbf{t})^t \equiv \iota s$, where $s \in \mathbb{R}^g$.

LEMMA 1. Let C be defined by [\(57\)](#page-17-2), and $s \in \mathbb{R}^g$. Then

(61)
$$
\nu_{3g}\bar{\nu}_{3g} = (-1)^{g-1}\lambda_{4g}\wp_{1,2g-1}(s).
$$

PROOF. By Cramer's rule, from (59) we find

(62)
$$
\nu_{3g} = \frac{\begin{vmatrix} C_g(C) & Q(C) \\ C_g(s) & Q(s) \end{vmatrix}}{\begin{vmatrix} 1_g & Q(C) \end{vmatrix}} = \frac{\begin{vmatrix} C_g(C) & P(C) & 0 \\ C_g(s) & P(s) & R(s) \end{vmatrix}}{\begin{vmatrix} 1_g & P(C) & 0 \end{vmatrix}} = \frac{|N|}{|D|}.
$$

where $C_g(u)$ is the identity matrix with the last column replaced with $-q(u)$, N and D denote the matrices in the numerator and denominator, respectively. The matrix Q(u) contains $\left[\frac{1}{2}g\right]$ columns linear in functions $\wp_{1,1,2i-1}(u)$ — we denote this block by R(u), and $g - \left[\frac{1}{2}g\right]$ columns which contain only functions $\wp_{1,2i-1}(u)$ this block is denoted by $\overline{P}(u)$. Entries of $P(C)$, $C_g(C)$ are expressed in terms of elementary symmetric functions in $\{e_{2i}\}_{i=1}^g$, and $\mathsf{R}(C) = 0$, since $\wp_{1,1,2i-1}$ vanish at half-periods.

Note, that the numerator and the denominator in [\(62\)](#page-20-0) contain the same block $R(s)$ of size $g \times [\frac{1}{2}g]$. Let $[R]_I$ denote a principal minor of $R(s)$ of order $[\frac{1}{2}g]$, where I shows selected rows and runs all combinations of $\left[\frac{1}{2}g\right]$ from g. Each block $\left[\mathsf{R}\right]_I$ contains full columns of R. Let $(\text{adj N})_I$ be the cofactor of N complementary to $\lbrack \text{R} \rbrack_I$, and $(\text{adj } D)_I$ be the cofactor of D complementary to $[R]_I$. The both numerator and denominator are computed by means of the Cauchy—Binet formula:

(63)
$$
|\mathsf{N}| = \sum_{I} \left(\mathrm{adj} \, \mathsf{N} \right)_{I} [\mathsf{R}]_{I}, \qquad |\mathsf{D}| = \sum_{I} \left(\mathrm{adj} \, \mathsf{D} \right)_{I} [\mathsf{R}]_{I}.
$$

Next, we compute

(64)
$$
\nu_{3g}\bar{\nu}_{3g} = \frac{|\mathcal{N}||\bar{\mathcal{N}}|}{|\mathcal{D}||\bar{\mathcal{D}}|},
$$

where the bar denotes the complex conjugate, which effects only entries of $P(C)$, $\mathsf{C}_g(\mathbf{C})$, where values e_i are replaced with the complex conjugate \bar{e}_i . Note, that $P(s)$, $C_g(s)$, $R(s)$ are real-valued.

Note, that $|\mathsf{N}||\bar{\mathsf{N}}|$ and $|\mathsf{D}||\bar{\mathsf{D}}|$ are homogeneous in $\wp_{1,1,2i-1}$ of degrees $2\left[\frac{1}{2}(g+1)\right]$, and $2\left[\frac{1}{2}g\right]$, respectively. Then, we apply the fundamental cubic relations, see [[10](#page-27-5), Theorem 3.2, (the argument s of φ -functions is omitted for simplicity):

(65)
$$
\wp_{1,1,2i-1}\wp_{1,1,2j-1} = 4(\wp_{1,1} + \lambda_2)\wp_{1,2i-1}\wp_{1,2j-1} + 4(\wp_{1,2i-1}\wp_{1,2j+1} + \wp_{1,2i+1}\wp_{1,2j-1}) + 4\wp_{2i+1,2j+1} - 2(\wp_{3,2i-1}\wp_{1,2j-1} + \wp_{1,2i-1}\wp_{3,2j-1}) - 2(\wp_{2i-1,2j+3} + \wp_{2i+3,2j-1}) + \lambda_4(\wp_{1,2j-1}\delta_{1,i} + \wp_{1,2i-1}\delta_{1,j}) + 4\lambda_{2+4i}\delta_{i,j} + 2(\lambda_{4i}\delta_{i-1,j} + \lambda_{4j}\delta_{i,j-1}).
$$

This completely eliminates $\wp_{1,1,2i-1}$ from the expression for $\nu_{3q}\bar{\nu}_{3q}$, and leads to the identity

(66)
$$
|\mathsf{N}||\bar{\mathsf{N}}| + (-1)^{g} \lambda_{4g} \wp_{1,2g-1}(s)|\mathsf{D}||\bar{\mathsf{D}}| = 0,
$$

which proves [\(62\)](#page-20-0). Direct computations of [\(66\)](#page-20-1) uses identities which define the Kummer variety of \mathcal{V} , see [[10](#page-27-5), § 4.1], [[8](#page-26-7), Sect. 5]. Such computations are done in genera 1, 2, 3. The identity [\(66\)](#page-20-1) holds in higher genera, which follows from [\(19\)](#page-7-5). \Box

LEMMA 2. Let C be defined by [\(57\)](#page-17-2), and $s \in \mathbb{R}^g$. Then

(67)
$$
\nu_{3g}\bar{\nu}_{3g} = (-1)^{g-1}\lambda_{4g}\wp_{1,2g-1}(is).
$$

PROOF. The statement of the Lemma follows immediately from the identity

(68)
$$
|\mathsf{N}||\bar{\mathsf{N}}| + (-1)^{g} \lambda_{4g} \wp_{1,2g-1}(i s)|\mathsf{D}||\bar{\mathsf{D}}| = 0,
$$

Note that $\wp_{i,j}(s)$ are even functions, and so real-valued. All terms in the funda-mental cubic relations [\(65\)](#page-20-2) remain real-valued after the substitution $s \mapsto \iota s$, as well as the identities which define the Kummer variety of $\mathcal V$. The proof of Lemma [1](#page-19-1) works Lemma 2 works Lemma [2.](#page-20-3)

THEOREM 4. Let C be defined by [\(57\)](#page-17-2), and $s \in \mathbb{R}^g$. Then

(69a)
$$
|\wp_{1,2g-1}(s+C)|^2 = \lambda_{4g},
$$

(69b)
$$
|\wp_{1,2g-1}(is+C)|^2 = \lambda_{4g}.
$$

PROOF. Using [\(60\)](#page-19-2) with $-u_{\text{III}} = s + C$, $u_{\text{I}} = C$, $u_{\text{II}} = s$, and $\lambda_{4g+2} = 0$, taking into account that $\wp_{1,2g-1}(C) = \prod_{i=1}^g e_{2i}$, and so $\wp_{1,2g-1}(C)\wp_{1,2g-1}(\bar{C}) = \lambda_{4g}$, cf. (52) , we find

$$
|\wp_{1,2g-1}(s+C)|^2 = \frac{(\nu_{3g}\bar{\nu}_{3g})^2}{\lambda_{4g}\wp_{1,2g-1}^2(s)},
$$

$$
|\wp_{1,2g-1}(s+C)|^2 = \frac{(\nu_{3g}\bar{\nu}_{3g})^2}{\lambda_{4g}\wp_{1,2g-1}^2(s)}
$$

.

Then, applying [\(61\)](#page-19-3) and [\(67\)](#page-20-4), we immediately obtain [\(69\)](#page-21-0). \Box

6.4. Summary. Quasi-periodic solutions of the sine-Gordon equation arise on the spectral curve [\(51\)](#page-15-2) with $\lambda_{4q+2} = 0$, positive λ_{4q} , and all finite non-zero branch points pairwise conjugate. In $\bar{\mathfrak{J}} = \mathbb{C}^g$, where the Jacobian variety of the spectral curve is embedded, all half-periods are located on 2^g affine subspaces parallel to the real axes subspace Re \mathfrak{J} , within the fundamental domain. There exists one subspace $C + \text{Re}\bar{3}$, with C defined by [\(57\)](#page-17-2), where $\wp_{1,2g-1}$ -function is bounded, and runs along the circle of radius $\sqrt{\lambda_{4g}}$. At the same time, all half-periods are located on 2^g affine subspaces parallel to the imaginary axes subspace Im $\bar{3}$, within the fundamental domain, and there exists one subspace $C + \text{Im } \mathfrak{J}$ where $\wp_{1,2g-1}$ function is bounded, and runs along the circle of radius $\sqrt{\lambda_{4g}}$. The subspace $C+\text{Im }\mathfrak{J}$ serves as the domain of the finite-gap solution of the sine-Gordon equation in a 2g-dimensional phase space, namely

(70)
$$
\phi = i \log \left(i \lambda_{4g}^{-1/2} \wp_{1,2g-1}(\boldsymbol{C} + i s) \right).
$$

Quasi-periodic solutions of the sinh-Gordon equation arise on the spectral curve [\(3\)](#page-1-0) with all real branch points, $\lambda_{4q+2} = 0$ and positive λ_{4q} . Reality conditions in this case are the same as presented in [[7](#page-26-0)].

7. Non-linear waves

In this section we present effective numerical computation of quasi-periodic finite-gap solutions of the sine-Gordon equation.

7.1. Numerical computation. In the case of sine-Gordon hierarchy, the curve [\(37\)](#page-12-0) is transformed to the canonical form [\(51\)](#page-15-2). The second kind differentials [\(5b\)](#page-2-2) change the sign:

(5b')
$$
\mathrm{d}r_{2n-1} = \frac{-\mathrm{d}x}{\partial_y f(x,y)} \sum_{j=1}^{2n-1} (2n-j) \lambda_{2j-2} x^{g+n-j}.
$$

Note, that $\lambda_{4g+2-2n} = h_{n-2}/b^2$, and $\lambda_{4N+2} = 0$. As a result, the function \mathcal{R}_{2g} , which defines the solution of the Jacobi inversion problem, transforms into

(9a')
$$
\mathcal{R}_{2g}(x,y;u) \equiv x^g + \sum_{i=1}^g x^{g-i} \wp_{1,2i-1}(u) = 0.
$$

℘-Functions are computed by the formulas

(72)

$$
\wp_{i,j}(u) = \varkappa_{i,j} - \frac{\partial^2}{\partial u_i \partial u_j} \theta[K](\omega^{-1} u; \omega^{-1} \omega'),
$$

$$
\wp_{i,j,k}(u) = -\frac{\partial^3}{\partial u_i \partial u_j \partial u_k} \theta[K](\omega^{-1} u; \omega^{-1} \omega'),
$$

where $x_{i,j}$ are entries of the symmetric matrix $x = \eta \omega^{-1}$, and the period matrices ω, ω', η are obtained from the differentials [\(5a\)](#page-2-1), [\(5b](#page-2-2)['\)](#page-22-0) along the canonical cycles, see fig. [1.](#page-2-0) Actually, columns of the matrices ω, ω', η are computed as follows

$$
\omega_k = 2 \int_{e_{2k-1}}^{e_{2k}} du, \qquad \eta_k = 2 \int_{e_{2k-1}}^{e_{2k}} dr,
$$

$$
\omega'_k = -2 \sum_{i=1}^k \int_{e_{2i-2}}^{e_{2i-1}} du = 2 \sum_{i=k}^g \int_{e_{2i}}^{e_{2i+1}} du.
$$

7.2. Genus 1. The hamiltonian system of the sine-Gordon equation in \mathcal{M}_1 with variables $\gamma_{-1}, \beta_{-1}, \alpha_0, (\beta_{-1} = -\bar{\gamma}_{-1}, \alpha_0 = ia_0)$ lives on the submanifold defined by the constraint

$$
|\gamma_{-1}|^2 = c^2 = -c_{-1}.
$$

The system is governed by the hamiltonian

$$
h_0 = \alpha_0^2 + b(\gamma_{-1} - \bar{\gamma}_{-1}) = -a_0^2 - 2b \operatorname{Im} \gamma_{-1},
$$

and possesses the spectral curve

(73)
$$
-w^2 - b^2 z^3 + h_0 z^2 + c_{-1} z = 0,
$$

which has complex conjugate branch points if $|h_0| < 2bc$.

On fig. [2a](#page-23-0) several trajectories of the sine-Gordon equation in \mathcal{M}_1 are displayed, $b = 1/2$, $c = 1/2$. In the 3-dimensional space with Re γ_{-1} , Im γ_{-1} , a_0 -axes, one can see a vertical circular cylinder $|\gamma_{-1}| = c$, and several parabolic cylinders which represent different level surfaces of the hamiltonian h_0 . Dark lines show trajectories, which arise on the intersection. When $|h_0| < 2bc$, each trajectory is one-loop (marked by green on fig. [2b](#page-23-0)). At $h_0 = 2bc$ the intersection collapses to the point $\gamma_{-1} = -\iota c, a_0 = 0$. When $h_0 > 2bc$, there is no intersection, that is no solution exists. When h_0 approaches $-2bc$, the trajectory twists so that two points with Im $\gamma_{-1} = -h_0/(2b)$, $a_0 = 0$ coincide at $\gamma_{-1} = ic$, $a_0 = 0$ (marked by blue on fig. [2b](#page-23-0)). At $h_0 < -2bc$ one loop splits into two (marked by orange on fig. [2b](#page-23-0)).

Recall that $\gamma_{-1} = i c \exp(i \phi)$, where ϕ obeys the sine-Gordon equation, which coincides with the equation of motion of a simple pendulum in the case of $N = 1$. On fig. [2b](#page-23-0) the reader can see a phase portrait of the 1-gap system from the sine-Gordon hierarchy, with the hamiltonian

$$
h_0 = -a_0^2 - 2bc \cos \phi.
$$

According to [\(40\)](#page-13-1), we have

(74)
$$
\gamma_{-1} = -ib\wp_{1,1}(u), \qquad a_0 = -\frac{b}{2} \frac{\wp_{1,1,1}(u)}{\wp_{1,1}(u)},
$$

where $u = ibx + \frac{1}{2}\omega + \frac{1}{2}\omega'$ as we combine [\(49\)](#page-14-2) and [\(57\)](#page-17-2). Therefore, the phase ϕ is defined by

.

(75)
$$
\phi(x) = i \log \left(-(b/c)\wp_{1,1}(bx + \frac{1}{2}\omega + \frac{1}{2}\omega') \right)
$$

As an example, we choose five values of h_0 which correspond to different mutual positions of branch points:

$$
h_0 = -0.49 \qquad e_1 = -0.98 - 0.199i,
$$

\n
$$
e_2 = -0.98 + 0.199i,
$$

\n
$$
e_3 = 0
$$

\n
$$
h_0 = -0.17 \qquad e_1 = -0.33 - 0.943i,
$$

\n
$$
e_2 = -0.33 + 0.943i,
$$

\n
$$
e_3 = 0
$$

\n
$$
h_0 = 0.17 \qquad e_1 = -0.33 - 0.943i,
$$

\n
$$
e_2 = -0.33 + 0.943i,
$$

\n
$$
e_3 = 0
$$

The corresponding fundamental domains and shapes of ϕ with $c = 1/2$, $b = 1/2$ are shown on fig. [3.](#page-24-0)

Figure 3.

7.3. Genus 2. The 2-gap hamiltonian system of the sine-Gordon hierarchy arises in \mathcal{M}_2 with variables γ_{-1} , β_{-1} , α_0 , γ_1 , β_1 , α_2 . In fact, the system lives in the submanifold defined by two constraints

$$
\begin{aligned} \gamma_{-1}\beta_{-1} &= c_{-1}, \\ \alpha_0^2 + \gamma_{-1}\beta_1 + \beta_{-1}\gamma_1 &= c_0, \end{aligned}
$$

which allow to eliminate β_{-1} and β_1 . The system is governed by two hamiltonians

$$
h_1 = 2\alpha_0\alpha_2 + b\gamma_{-1} + (c_0 - \alpha_0^2)\frac{\gamma_1}{\gamma_{-1}} + c_{-1}\left(\frac{b}{\gamma_{-1}} - \frac{\gamma_1^2}{\gamma_{-1}^2}\right),
$$

$$
h_2 = \alpha_2^2 + b\gamma_1 + (c_0 - \alpha_0^2) \frac{b}{\gamma_{-1}} - c_{-1} \frac{b\gamma_1}{\gamma_{-1}^2}.
$$

After substitution

$$
\gamma_{-1} = b z_1 z_2, \qquad \gamma_1 = -b(z_1 + z_2),
$$

\n
$$
\alpha_0 = -\frac{w_1 z_2^2 - w_2 z_1^2}{z_1 z_2 (z_1 z_2)}, \qquad \alpha_2 = \frac{w_1 z_2 - w_2 z_1}{z_1 z_2 (z_1 z_2)},
$$

induced by [\(34\)](#page-11-3), the hamiltonians acquire the form

$$
h_1 = -\frac{z_2}{z_1^3(z_1 - z_2)} (w_1^2 - b^2 z_1^5 - c_0 z_1^2 - c_{-1} z_1)
$$

+
$$
\frac{z_1}{z_2^3(z_1 - z_2)} (w_2^2 - b^2 z_2^5 - c_0 z_2^2 - c_{-1} z_2),
$$

$$
h_2 = \frac{1}{z_1^3(z_1 - z_2)} (w_1^2 - b^2 z_1^5 - c_0 z_1^2 - c_{-1} z_1)
$$

-
$$
\frac{1}{z_2^3(z_1 - z_2)} (w_2^2 - b^2 z_2^5 - c_0 z_2^2 - c_{-1} z_2).
$$

In the new variables z_1, z_2, w_1, w_2 separation into two samples of the spectral curve is achieved:

$$
-w_i^2 + b^2 z_i^5 + h_2 z_i^4 + h_1 z_i^3 + c_0 z_i^2 + c_{-1} z_i = 0, \qquad i = 1, 2.
$$

Taking into account, that $\beta_{2i-1} = -\overline{\gamma}_{2i-1}$, $\alpha_{2i} = ia_{2i}$, $i = 0, 1$, and $b = ib$, we obtain the constraints in the form

(76a)
$$
|\gamma_{-1}|^2 = c^2 = -c_{-1},
$$

(76b)
$$
a_0^2 + 2 \operatorname{Im} \gamma_1 \operatorname{Im} \gamma_{-1} + 2 \operatorname{Re} \gamma_1 \operatorname{Re} \gamma_{-1} = -c_0.
$$

In the projection to the plane $(Re \gamma_{-1})O(Im \gamma_{-1})$ the first equation is a circle of radius c = $\sqrt{c_{-1}}$, and the second one is a line with the normal 2(Re γ_1 , Im γ_1), and the distance from the origin $d = -\frac{1}{2}(c_0 + a_0^2)/|\gamma_1|$. An intersection exists if

$$
\frac{1}{2}|c_0 + a_0^2| \leqslant c|\gamma_1|.
$$

This means, $c_{-1} < 0$, and c_0 changes from $-2c|\gamma_1| - a_0^2$ to $2c|\gamma_1| - a_0^2$.

The hamiltonians acquire the form

(77a)
$$
h_1 = -2a_0a_2 - 2b \operatorname{Im} \gamma_{-1} - |\gamma_1|^2,
$$

(77b)
$$
h_2 = -a_2^2 - 2b \operatorname{Im} \gamma_1.
$$

Equations [\(77\)](#page-25-0) together with [\(76\)](#page-25-1) define a 2-dimensional surface parametrized by real variables $(x, t)^t = s$. Let $u = ibs + C$ with $C = \frac{1}{2}\omega_2 + \frac{1}{2}\omega'_1 + \frac{1}{2}\omega'_2$ as follows from [\(49\)](#page-14-2) and [\(57\)](#page-17-2). Then [\(40\)](#page-13-1) gives a parametrization of the surface:

(78)
$$
\gamma_{-1} = -\iota b \wp_{1,3}(u), \qquad \gamma_1 = -\iota b \wp_{1,1}(u),
$$

(79)
$$
a_0 = -\frac{b}{2} \frac{\wp_{1,3,3}(u)}{\wp_{1,3}(u)}, \qquad a_2 = -\frac{b}{2} \frac{\wp_{1,1,3}(u)}{\wp_{1,3}(u)}.
$$

The system possesses a spectral curve of the form

(80)
$$
-w^2 - b^2 z^5 + h_2 z^4 + h_1 z^3 c_0 z^2 + c_{-1} z = 0,
$$

By fixing c_{-1} , c_0 and choosing h_1 , h_2 such that all non-zero branch points of [\(80\)](#page-25-2) are complex conjugate, all possible 2-gap solutions of the sine-Gordon equation can be obtained, that is the matter of a separate investigations.

Therefore, the phase ϕ is defined by

(81)
$$
\phi(x) = i \log (- (b/c) \wp_{1,3}(ibs + C)).
$$

Let $b = 1$. We consider several cases with different mutual positions of branch points.

References

- [1] Ablowitz M. J., Kaup D. J., Newell A. C., Segur H., On symplectic structures and integrable systems on symmetric spaces, Phys, Rev. Lett. 30:25 (1973) 1262–1264.
- [2] Adler, M. On a trace functional for formal pseudo-differential operators and the symplectic structure of the Korteweg-devries type equations. Invent. Math. 50, 219–248 (1978).
- [3] Adler, M., Moerbeke P. Completely integrable systems, Euclidean Lie algebras, and curves. Advances in Math., 38:3, 267–317 (1980)
- [4] Aktosun T., Demontis F., van der Mee C., Exact solutions to the sine-Gordon equation, J. Math. Phys. 51 (2010) 123521.
- [5] Baker H.F., Abelian functions: Abel's theorem and the allied theory of theta functions, Cambridge, University press, Cambridge, 1897.
- [6] Belokolos E. D., Bobenko A. I., Enolski V. Z., Its A. R., Matveev. V.B., Algebro-geometric approach to nonlinear integrable equations., Springer-Verhag, 1994
- [7] Bernatska J., Reality conditions for the KdV equation and exact quasi-periodic solutions in finite phase spaces, J. Geom. Phys. 206 (2024) 105322.
- [8] Bernatska J., Abelian function fields on Jacobian varieties, [arXiv:2412.05455.](http://arxiv.org/abs/2412.05455)

- [9] Bernatska J, Holod P. On Separation of variables for Integrable equations of soliton type, J. Nonlin. Math. Phys. 14:3 (2007) 353–374
- [10] Buchstaber V. M., Enolskii V. Z., and Leykin D. V., Hyperelliptic Kleinian functions and applications, preprint ESI 380 (1996), Vienna
- [11] V. M. Buchstaber, D. V. Leikin, Addition laws on Jacobian varieties of plane algebraic curves, Nonlinear dynamics, Collected papers, Tr. Mat. Inst. Steklova, 251 (2005), pp. 54–126.
- [12] Dubrovin B. A., Natanzon S. M., Real two-zone solutions of the sine-Gordon equation, Functional Analysis and Its Applications, 16 (1982) 21-33.
- [13] Hirota R., Exact solution of the sine-Gordon equation for multiple collisions of solitons, J. Phys. Soc. Japan, 33:5 (1972) 1459–1463.
- [14] Holod P., Hamiltonian systems on the orbits of affine Lie groups and finite-band integration of nonlinear equation, pp.1361 E^{4} 1367; In Nonlinear and Turbulent Processes in Physics: Nonlinear effects in plasma physics, astrophysics, and elementary particle theory. Ed. Sagdeev R. Z., Harwood Academic Publishers, 1984.
- [15] Kozel V. A. and Kotlyarov V. P. Explicit almost periodic solutions of the sine-Gordon equation Dokl. Akad. Nauk Ukr. SSR, Ser. A, 10 (1976) 878–881 (in Russian); translation in English in [arXiv:1401.4410](http://arxiv.org/abs/1401.4410)
- [16] Smirnov, A. O., Real elliptic solutions of the sine-Gordon equation, Mathematics of the USSR-Sbornik, 70:1 (1991) 231–240
- [17] Smirnov, A. O., 3-Elliptic solutions of the sine-Gordon equation MathematicalNotes, 6:3 (1997) 368–376.
- [18] Steuerwald R., Über Enneper $\mathbb{B}^{\mathbb{N}}$ sche Flächen und Bäcklund $\mathbb{B}^{\mathbb{N}}$ sche Transformation. Abh. Bayer. Akad. Wiss. (München) 40:1 (1936).
- [19] The sine-Gordon model and its applications. From pendula and Josephson junctions to gravity and high-energy physics. Eds.: Cuevas-Maraver, J., Kevrekidis, P. G. Williams, F., Springer Cham, New York, 2014

University of Connecticut, Department of Mathematics Email address: jbernatska@gmail.com

