

Exotic massive fermionic systems with huge vacuum degeneracy at boundaries

Hiroki Kawakami and Satoshi Yamaguchi

*Department of Physics, Graduate School of Science,
Osaka University, Toyonaka, Osaka 560-0043, Japan*

Abstract

We investigate a massive non-relativistic fermionic system exhibiting exotic features. When the mass parameter is set to zero, the system acquires the fermionic subsystem symmetry. Introducing the mass term explicitly breaks this symmetry, resulting in a trivially gapped system in the absence of boundaries. We demonstrate that with the introduction of boundaries, the system remains gapped, but it has a huge vacuum degeneracy. The residual entropy is proportional to the area of the boundary.

1 Introduction

Recently, there has been increasing interest in the study of non-relativistic systems with exotic features, including fractons [1, 2]. For nice reviews on the topic, see [3, 4, 5]. Not only bosonic systems, fermionic systems are also studied. Notably, a fermionic model with fermionic subsystem symmetry was introduced in [6] by considering the supersymmetric version of the bosonic models studied in [7, 8, 9, 10]. Additional studies of similar fermionic models can be found in [11, 12]. Furthermore, lattice fermionic fracton systems have been explored in [13, 14, 15, 16].

Introducing boundaries to such exotic systems presents an intriguing avenue of study. For example, the anomaly inflow mechanism [17] associated with subsystem symmetry gives rise to robust, gapless boundary degrees of freedom in the systems that are otherwise trivially gapped in the absence of boundaries. These systems, commonly referred to as subsystem symmetry protected topological (SSPT) phases, have been extensively studied in works such as [18, 19, 20, 21, 22, 23, 24, 25].

In this paper, we investigate a different type of boundary degrees of freedom in a system that is trivially gapped in the bulk. Specifically, we study the ψ theory, an exotic fermionic system

introduced in [6]. By adding a mass term to the ψ theory, the fermionic subsystem symmetry is explicitly broken, resulting in a system that is trivially gapped in the absence of boundaries. This behavior is analogous to the $(1 + 1)$ -dimensional Majorana fermion [26], which is also trivially gapped in a space without boundaries when a mass term is present. For the Majorana fermion, the presence of boundaries may lead to robust ground state degeneracy depending on the mass term and the boundary condition. We examine a similar phenomenon in the massive ψ theory.

We observe that when boundaries are introduced to the massive ψ theory, the system remains gapped; however, it exhibits a huge number of ground states. The residual entropy, defined as $\log(\text{the number of ground states})$, is proportional to the area of the boundary. Our findings differ from those in previous works [21, 22] in the following ways:

- The system remains gapped even with the introduction of boundaries.
- The (fermionic) subsystem symmetry is explicitly broken by the mass term. Consequently, the boundary ground state degeneracy is not protected by this symmetry. If the degeneracy is robust, it must be protected by another, as-yet unidentified, symmetry.

Investigating these observations within the framework of SSPT phases is an interesting future work.

The rest of this paper is organized as follows. In Sec. 2, we introduce the massive ψ theory and demonstrate that it is trivially gapped in the absence of boundaries. In Sec. 3, we examine the effects of introducing boundaries, identify possible boundary conditions, and the spectrum of the system. Sec. 4 is devoted to conclusions and discussions.

2 The bulk system

In this section, we set up the model of the bulk system.

The system is defined on a $(3 + 1)$ -dimensional spacetime. The three-dimensional space is a torus T^3 , whose coordinates x, y, z satisfy the following identifications:

$$(x, y, z) \sim (x + l_x, y, z) \sim (x, y + l_y, z) \sim (x, y, z + l_z), \quad (2.1)$$

where $l_x, l_y, l_z > 0$ are the periods in the x, y, z directions, respectively. The temporal coordinate is denoted by t .

The theory considered in this paper is a massive fermionic theory, formulated as follows. The fundamental fields of the theory are two real fermionic fields $\psi_{\pm}(t, x, y, z)$. The action is given by

$$S = \int d^3x dt (i\psi_+ \partial_- \psi_+ + i\psi_- \partial_+ \psi_- - im\psi_- \psi_+), \quad (2.2)$$

where $m \in \mathbb{R}$ is the mass parameter. ∂_{\pm} are defined as:

$$\partial_{\pm} := \frac{1}{2}(\partial_t \pm \partial_x \partial_y \partial_z). \quad (2.3)$$

The theory described by (2.2) with $m = 0$ has been first introduced in [6] and referred as the ψ theory. In this case, the theory has so called the fermionic subsystem symmetry, which is a shift of ψ_{\pm} by a function of any two spatial coordinates among x , y , and z . This ψ theory is gapless or equivalently “massless.” If periodic boundary conditions are imposed on ψ_{\pm} , the system has a huge vacuum degeneracy as discussed in [6]. In this paper, for simplicity, we impose anti-periodic boundary conditions on ψ_{\pm} . While the system remains gapless under the anti-periodic boundary conditions, the vacuum is unique.

On the other hand, if $m \neq 0$, the mass term explicitly breaks the fermionic subsystem symmetry. The system becomes trivially gapped in the absence of boundaries, as we will demonstrate below.

Since the massive ψ theory (2.2) is a free field theory, we can easily solve it by the mode-expansion. The expressions of ψ_{\pm} in terms of eigen-modes are given by

$$\begin{aligned}\psi_+ &= \sum_{\vec{k}} \sqrt{\frac{1}{2V} \frac{\omega_{\vec{k}} + k_x k_y k_z}{\omega_{\vec{k}}}} \left(a_{\vec{k}} \exp(i\vec{k} \cdot \vec{x} - i\omega_{\vec{k}} t) + a_{\vec{k}}^{\dagger} \exp(-i\vec{k} \cdot \vec{x} + i\omega_{\vec{k}} t) \right), \\ \psi_- &= i \text{sign } m \sum_{\vec{k}} \sqrt{\frac{1}{2V} \frac{\omega_{\vec{k}} - k_x k_y k_z}{\omega_{\vec{k}}}} \left(a_{\vec{k}} \exp(i\vec{k} \cdot \vec{x} - i\omega_{\vec{k}} t) - a_{\vec{k}}^{\dagger} \exp(-i\vec{k} \cdot \vec{x} + i\omega_{\vec{k}} t) \right), \\ V &:= l_x l_y l_z,\end{aligned}\tag{2.4}$$

where the wave numbers $\vec{k} = (k_x, k_y, k_z)$ take the following values due to the anti-periodic boundary condition:

$$k_i = \frac{2\pi}{l_i} \left(n_i + \frac{1}{2} \right), \quad n_i \in \mathbb{Z}, \quad i = x, y, z.\tag{2.5}$$

The dispersion relation is given by

$$\omega_{\vec{k}} = \sqrt{k_x^2 k_y^2 k_z^2 + m^2}.\tag{2.6}$$

The anti-commutation relations of the mode operators $a_{\vec{k}}, a_{\vec{k}}^{\dagger}$ are given by

$$\{a_{\vec{k}}, a_{\vec{k}'}^{\dagger}\} = \delta_{\vec{k}, \vec{k}'}.\tag{2.7}$$

The Hamiltonian of this system is given by

$$H = \int d^3x \left(\frac{1}{2} i\psi_+ \partial_x \partial_y \partial_z \psi_+ - \frac{1}{2} i\psi_- \partial_x \partial_y \partial_z \psi_- + im\psi_- \psi_+ \right) = \sum_{\vec{k}} \omega_{\vec{k}} a_{\vec{k}}^{\dagger} a_{\vec{k}}.\tag{2.8}$$

From dispersion relation (2.6), we find

$$\omega_{\vec{k}} \geq |m|,\tag{2.9}$$

indicating that the system is gapped. The ground state is annihilated by $a_{\vec{k}}$ for all \vec{k} and it is unique. Thus, the system is trivially gapped in the absence of boundaries.

3 The system with boundary

In this section, we examine the effects of introducing boundaries to the system. First, we explore the possible boundary conditions. Subsequently, we analyze the spectrum of the system in the presence of boundaries.

3.1 Boundary conditions

The space we consider here is $I \times T^2$. Here, I is an interval defined by $0 \leq x \leq l_x$ with boundaries at $x = 0$ and $x = l_x$. The coordinates of the torus T^2 , denoted by y, z , are identified as $y \sim y + l_y$ and $z \sim z + l_z$.

The action of the massive ψ theory is given by

$$S = \int d^3x dt (i\psi_+ \partial_- \psi_+ + i\psi_- \partial_+ \psi_- - im\psi_- \psi_+). \quad (3.1)$$

For simplicity, we impose anti-periodic boundary conditions on ψ_{\pm} along the y and z directions:

$$\begin{aligned} \psi_{\pm}(x, y, z) &= -\psi_{\pm}(x, y + l_y, z), \\ \psi_{\pm}(x, y, z) &= -\psi_{\pm}(x, y, z + l_z). \end{aligned} \quad (3.2)$$

To determine the possible boundary conditions at $x = 0$ and $x = l_x$, we apply the variational principle. The variation of the action (3.1) is given by

$$\begin{aligned} \delta S &= \int d^3x dt [2i\delta\psi_+ \partial_- \psi_+ + 2i\delta\psi_- \partial_+ \psi_- - im\delta\psi_- \psi_+ - im\psi_- \delta\psi_+] \\ &\quad + \int dy dz dt [\delta\psi_+ \partial_y \partial_z \psi_+ - \delta\psi_- \partial_y \partial_z \psi_-]_{x=0}^{x=l_x}. \end{aligned} \quad (3.3)$$

The boundary conditions are chosen to ensure that the boundary term in (3.3) vanishes:

$$\int dy dz dt [\delta\psi_+ \partial_y \partial_z \psi_+ - \delta\psi_- \partial_y \partial_z \psi_-]_{x=0}^{x=l_x}. \quad (3.4)$$

From this, we identify the possible boundary conditions as:

$$\partial_y \partial_z \psi_{\pm} = \pm \partial_y \partial_z \psi_{\mp} |_{x=0, l_x}. \quad (3.5)$$

Actually, the boundary conditions (3.5) imply

$$\begin{aligned} \int dy dz [\delta\psi_+ \partial_y \partial_z \psi_+ - \delta\psi_- \partial_y \partial_z \psi_-]_{x=0} &= \int dy dz [(\delta\psi_+ \mp \delta\psi_-) \partial_y \partial_z \psi_+]_{x=0} \\ &= \int dy dz [\partial_y \partial_z (\delta\psi_+ \mp \delta\psi_-) \psi_+]_{x=0} = 0. \end{aligned} \quad (3.6)$$

The same holds at $x = l_x$. Therefore, the boundary conditions (3.5) satisfy (3.4) and the variational principle is satisfied.

There are four possible boundary conditions based on the signs in (3.5):

$$(i). \quad \partial_y \partial_z \psi_+ = \partial_y \partial_z \psi_-|_{x=0, l_x}.$$

$$(ii). \quad \partial_y \partial_z \psi_+ = \partial_y \partial_z \psi_-|_{x=0}, \quad \partial_y \partial_z \psi_+ = -\partial_y \partial_z \psi_-|_{x=l_x}.$$

$$(iii). \quad \partial_y \partial_z \psi_+ = -\partial_y \partial_z \psi_-|_{x=0, l_x}.$$

$$(iv). \quad \partial_y \partial_z \psi_+ = -\partial_y \partial_z \psi_-|_{x=0}, \quad \partial_y \partial_z \psi_+ = +\partial_y \partial_z \psi_-|_{x=l_x}.$$

However, the boundary conditions (iii) and (iv) are derived by substituting $m \rightarrow -m$, $\psi_{\pm} \rightarrow \pm\psi_{\pm}$ to the boundary conditions (i) and (ii), respectively. Thus, we focus on the boundary conditions (i) and (ii) in the following.

3.2 Spectrum of the system with boundary

Let us find the spectrum of the system with boundary. The bulk theory is a free theory and the boundary conditions (i) and (ii) are linear. Therefore, we can solve the system by the mode expansion. In order to find the spectrum, it is enough to find the eigen-modes and the dispersion relation.

We now examine the eigen-modes of the system in detail with the boundary conditions (i) and (ii).

3.2.1 Case (i)

Here, we analyze the system with the boundary condition (i). There are two classes of mode functions to consider: the bulk modes and the boundary-localized modes. Both of them satisfy the equations of motion derived from (3.3):

$$\begin{aligned} 2\partial_- \psi_+ + m\psi_- &= 0, \\ 2\partial_+ \psi_- - m\psi_+ &= 0. \end{aligned} \tag{3.7}$$

Let us first consider the bulk modes ϕ_{\pm} . A plane wave solution of the equations of motion (3.7) is given by

$$\begin{aligned} \phi_+ &= A \exp(i\vec{k} \cdot \vec{x} - i\omega t), \\ \phi_- &= \frac{1}{m}(i\omega - ik_x k_y k_z) A \exp(i\vec{k} \cdot \vec{x} - i\omega t), \\ \omega &= \pm\omega_{\vec{k}}, \quad \omega_{\vec{k}} := \sqrt{k_x^2 k_y^2 k_z^2 + m^2}, \end{aligned} \tag{3.8}$$

where A is a constant and $\vec{k} = (k_x, k_y, k_z)$ represents the wave vector.

To satisfy the boundary condition, we make an ansatz that the bulk modes are given by a superposition of incident and reflected waves:

$$\begin{aligned}\phi_+ &= A \exp(i\vec{k} \cdot \vec{x} - i\omega t) + B \exp(-ik_x x + ik_y y + ik_z z - i\omega t), \\ \phi_- &= \frac{1}{m}(i\omega - ik_x k_y k_z)A \exp(i\vec{k} \cdot \vec{x} - i\omega t) + \frac{1}{m}(i\omega + ik_x k_y k_z)B \exp(-ik_x x + ik_y y + ik_z z - i\omega t),\end{aligned}\quad (3.9)$$

where A and B are constants.

We substitute this ansatz for the boundary condition (i) and obtain

$$\begin{aligned}(m - i\omega + ik_x k_y k_z)A + (m - i\omega - ik_x k_y k_z)B &= 0, \\ (m - i\omega + ik_x k_y k_z) \exp(ik_x l_x)A + (m - i\omega - ik_x k_y k_z) \exp(-ik_x l_x)B &= 0.\end{aligned}\quad (3.10)$$

In order for the above equations to have non-trivial solutions, the determinant of the coefficient matrix must vanish.

$$\det \begin{pmatrix} m - i\omega + ik_x k_y k_z & m - i\omega - ik_x k_y k_z \\ (m - i\omega + ik_x k_y k_z) \exp(ik_x l_x) & (m - i\omega - ik_x k_y k_z) \exp(-ik_x l_x) \end{pmatrix} = 0. \quad (3.11)$$

This equation is equivalent to

$$(m - i\omega + ik_x k_y k_z)(m - i\omega - ik_x k_y k_z) (\exp(ik_x l_x) - \exp(-ik_x l_x)) = 0. \quad (3.12)$$

Since $m \neq 0$ and ω, k_x, k_y, k_z are real, the above equation is equivalent to

$$\exp(ik_x l_x) - \exp(-ik_x l_x) = 0. \quad (3.13)$$

From this condition, the possible values of k_x are given by

$$k_x = \frac{n_x \pi}{l_x}, \quad n_x \in \mathbb{Z}. \quad (3.14)$$

Actually, $n_x = n$ and $n_x = -n$ give the same mode function as seen in Eq. (3.9). Thus, we choose $n_x > 0$ as the representative. Moreover, $n_x = 0$ is not valid since the mode function (3.9) vanishes in this case. Therefore, we obtain

$$\begin{aligned}\omega &= \pm \sqrt{k_x^2 k_y^2 k_z^2 + m^2}, \\ k_x &= \frac{n_x \pi}{l_x}, \quad k_y = \frac{(2n_y + 1)\pi}{l_y}, \quad k_z = \frac{(2n_z + 1)\pi}{l_z}, \quad n_x, n_y, n_z \in \mathbb{Z}, \quad n_x > 0,\end{aligned}\quad (3.15)$$

for the spectrum corresponding to the bulk modes.

Next, let us consider the boundary-localized modes, denoted by χ_{\pm} . We propose the following ansatz for these modes:

$$\begin{aligned}\chi_+ &= A \exp(-\Lambda x + ik_y + ik_z z - i\omega t) + B \exp(\Lambda x + ik_y + ik_z z - i\omega t) \\ \chi_- &= \frac{1}{m}(i\omega + \Lambda k_y k_z)A \exp(-\Lambda x + ik_y + ik_z z - i\omega t) \\ &\quad + \frac{1}{m}(i\omega - \Lambda k_y k_z)B \exp(\Lambda x + ik_y + ik_z z - i\omega t), \\ \omega &= \pm \omega_{\text{local}}(\Lambda, k_y, k_z), \quad \omega_{\text{local}}(\Lambda, k_y, k_z) := \sqrt{-\Lambda^2 k_y^2 k_z^2 + m^2},\end{aligned}\quad (3.16)$$

where A, B are constants and Λ is a positive real number.

The boundary condition (i) leads to the following relations:

$$\begin{aligned}(m - i\omega - \Lambda k_y k_z)A + (m - i\omega + \Lambda k_y k_z)B &= 0, \\ (m - i\omega - \Lambda k_y k_z) \exp(-\Lambda l_x)A + (m - i\omega + \Lambda k_y k_z) \exp(\Lambda l_x)B &= 0.\end{aligned}\quad (3.17)$$

For non-trivial solutions to exist, the determinant of the coefficient matrix in (3.17) must vanish. This condition is equivalent to:

$$(m - i\omega + \Lambda k_y k_z)(m - i\omega - \Lambda k_y k_z)(\exp(\Lambda l_x) - \exp(-\Lambda l_x)) = 0. \quad (3.18)$$

The solutions of this equation are given by $\Lambda = 0, \pm \frac{m}{k_y k_z}$.

- First, the solution $\Lambda = 0$ is invalid, as the mode function vanishes in this case.
- Second, For $\Lambda = \frac{m}{k_y k_z}$, we find $B = 0$ and $\omega = 0$ from (3.17) and (3.16), respectively. Since $\Lambda > 0$, this solution is valid only when $\frac{m}{k_y k_z} > 0$. The corresponding mode function is:

$$\begin{aligned}\chi_+ &= A \exp\left(-\frac{m}{k_y k_z}x + ik_y + ik_z z\right), \\ \chi_- &= A \exp\left(-\frac{m}{k_y k_z}x + ik_y y + ik_z z\right).\end{aligned}\quad (3.19)$$

This solution is localized at $x = 0$.

- Third, For $\Lambda = -\frac{m}{k_y k_z}$, we find $A = 0$ and $\omega = 0$ from (3.17) and (3.16), respectively. Since $\Lambda > 0$, this solution is valid only when $\frac{m}{k_y k_z} < 0$. The corresponding mode function is:

$$\begin{aligned}\chi_+ &= B \exp\left(-\frac{m}{k_y k_z}x + ik_y + ik_z z\right), \\ \chi_- &= B \exp\left(-\frac{m}{k_y k_z}x + ik_y y + ik_z z\right).\end{aligned}\quad (3.20)$$

This solution is localized at $x = l_x$.

The frequencies of these modes are exactly zero.

It follows that the system with this boundary exhibits exponentially large vacuum degeneracy. As demonstrated above, there is one Majorana zero mode for each pair (k_y, k_z) . Let N be the number of such pairs, i.e., the number of Majorana zero modes. This number scales as $N \sim \frac{l_y l_z}{a^2}$ with the lattice spacing a . The vacuum degeneracy is given by $2^{\frac{1}{2}N}$. Consequently, the ground states of the system with this boundary are exponentially degenerate. Furthermore, the residual entropy is proportional to the area of the boundary.

Recall that the analogous boundary condition of the $(1 + 1)$ -dimensional Majorana fermion is forbidden due to the existence of an odd number of Majorana zero modes. However, the boundary condition in our theory is not forbidden since the number of the Majorana zero modes is even, as shown below. Actually, the complex conjugates of the mode functions (3.19) and (3.20) for (k_y, k_z) are identical to the mode functions for $(-k_y, -k_z)$. Therefore, precisely speaking, there is a Dirac zero mode or two Majorana zero modes for a pair (k_y, k_z) and $(-k_y, -k_z)$ if $(k_y, k_z) \neq (0, 0)$. Additionally, note that $(k_y, k_z) \neq (0, 0)$ for our anti-periodic boundary condition (3.2). Consequently, the number of Majorana zero modes are even.

3.2.2 Case (ii)

Here, we analyze the system with boundary condition (ii).

The ansatz for the bulk modes remains the same as in the previous case (3.9). The boundary condition (ii) leads to the following relations:

$$\begin{aligned} (m - i\omega + ik_x k_y k_z)A + (m - i\omega - ik_x k_y k_z)B &= 0, \\ (m + i\omega - ik_x k_y k_z) \exp(ik_x l_x)A + (m + i\omega + ik_x k_y k_z) \exp(-ik_x l_x)B &= 0. \end{aligned} \quad (3.21)$$

In order for the above equations to have non-trivial solutions, the determinant of the coefficient matrix must vanish:

$$\det \begin{pmatrix} m - i\omega + ik_x k_y k_z & m - i\omega - ik_x k_y k_z \\ (m + i\omega - ik_x k_y k_z) \exp(ik_x l_x) & (m + i\omega + ik_x k_y k_z) \exp(-ik_x l_x) \end{pmatrix} = 0. \quad (3.22)$$

This condition simplifies to:

$$\tan k_x l_x = \frac{k_x k_y k_z}{m}. \quad (3.23)$$

The general solution of this equation cannot be expressed in terms of elementary functions. Approximated solutions are available in the limiting cases:

$$\begin{aligned} k_x &\approx \frac{n\pi}{l_x}, & \text{if } |k_x| &\ll \left| \frac{m}{k_y k_z} \right|, \\ k_x &\approx \frac{(2n+1)\pi}{2l_x}, & \text{if } |k_x| &\gg \left| \frac{m}{k_y k_z} \right|, \end{aligned} \quad (3.24)$$

$n \in \mathbb{Z}$.

Although $k_x = 0$ is an exact solution of (3.23), the corresponding mode function is $\chi_{\pm} = 0$, and therefore not suitable as a valid mode function.

The ansatz for the boundary-localized modes is the same as the previous case (3.16). The boundary condition (ii) leads to the following relations:

$$\begin{aligned} (m - i\omega - \Lambda k_y k_z)A + (m - i\omega + \Lambda k_y k_z)B &= 0, \\ (m + i\omega + \Lambda k_y k_z) \exp(-\Lambda l_x)A + (m + i\omega - \Lambda k_y k_z) \exp(\Lambda l_x)B &= 0. \end{aligned} \quad (3.25)$$

For non-trivial solutions to exist, the determinant of the coefficient matrix must vanish:

$$\det \begin{pmatrix} m - i\omega - \Lambda k_y k_z & m - i\omega + \Lambda k_y k_z \\ (m + i\omega + \Lambda k_y k_z) \exp(-\Lambda l_x) & (m + i\omega - \Lambda k_y k_z) \exp(\Lambda l_x) \end{pmatrix} = 0. \quad (3.26)$$

This condition simplifies to:

$$\tanh \Lambda l_x = \frac{\Lambda k_y k_z}{m}. \quad (3.27)$$

$\Lambda = 0$ is an exact solution of (3.27), but the corresponding mode function for this solution is $\chi_{\pm} = 0$, and not suitable as a mode function. A non-zero solution exists for Eq. (3.27) if and only if $1 < \frac{m l_x}{k_y k_z}$. Assuming that the system size is large enough to satisfy $1 \ll \frac{m l_x}{k_y k_z}$,¹ we can evaluate the frequency of the boundary-localized mode. The solution of Eq. (3.27) is expressed as

$$\Lambda = \Lambda_0 + O(e^{-2\Lambda_0 l_x}), \quad \Lambda_0 := \frac{m}{k_y k_z}. \quad (3.28)$$

Note that the reminder term $O(e^{-2\Lambda_0 l_x})$ in the expression of Λ is negative, leading to $\Lambda < \Lambda_0$. The frequency of the boundary-localized mode is given by

$$\omega = \pm \omega_{\text{local}}(\Lambda, k_y, k_z), \quad \omega_{\text{local}}(\Lambda, k_y, k_z) := \sqrt{-\Lambda^2 k_y^2 k_z^2 + m^2} = O(e^{-\Lambda_0 l_x}). \quad (3.29)$$

Since $\omega_{\text{local}}(\Lambda, k_y, k_z) > 0$, the boundary-localized modes have exponentially small positive and negative frequencies.

The system with this boundary also exhibits exponentially large vacuum degeneracy. As shown above, there are two approximate zero modes for a pair (k_y, k_z) that satisfy $\frac{k_y k_z}{m} > 0$ and no approximate zero modes for (k_y, k_z) that satisfy $\frac{k_y k_z}{m} < 0$. The total number of approximate zero modes are the same as in Case (i). Consequently, the ground state degeneracy is also the same as that in Case (i).

This system is similar to the $(1 + 1)$ -dimensional Majorana fermion with the Neveu-Schwarz (NS) boundary condition. In the $(1 + 1)$ -dimensional case, the number of Majorana zero modes depends on the sign of the mass term and can be either zero or two. In contrast, for the present system, a large number of Majorana zero modes exist regardless of the sign of m .

¹We implicitly assume the existence of a cutoff, ensuring an upper bound for $|k_y k_z|$. Thus, this condition is satisfied if l_x is chosen sufficiently large.

4 Conclusion and discussion

In this paper, we investigated the massive ψ theory (2.2) with boundaries. While the system is trivially gapped in the absence of boundaries, we explored the effects of introducing boundary conditions (i) and (ii). For Case (i), there are a large number of exact zero modes, resulting a huge vacuum degeneracy. The residual entropy is proportional to the area of the boundary. Non-zero modes exhibit a gap of $|m|$, ensuring that the system remains gapped. For Case (ii), although the zero modes are not exact but approximate, the vacuum degeneracy and the gap are the same as in Case (i).

This behavior is reminiscent of symmetry protected topological (SPT) phases, where the system is trivially gapped in the absence of boundaries but becomes gapless or exhibits vacuum degeneracy in the presence of boundaries. One may suspect that the massive ψ theory belongs to an SPT phase or an SSPT phase. However, there are notable differences:

- Our system on the space with boundary has a vacuum degeneracy, regardless of the sign of m . In contrast, typical SPT phases become trivial phase if the sign of the mass is reversed.
- The symmetry that protects the vacuum degeneracy of the massive ψ theory is not yet identified. It is not the fermionic subsystem symmetry, as it is explicitly broken by the mass term.
- No inconsistencies were found in the boundary condition (i). This contrasts with the analogous boundary condition for the $(1+1)$ -dimensional Majorana fermion, which is inconsistent due to the existence of odd number of Majorana zero modes.

Identifying the symmetry that protects the vacuum degeneracy of the massive ψ theory is an intriguing direction for future research. Possible candidates include discrete symmetries such as discrete translations or 90-degree rotations.

In this paper, we consider only the boundary orthogonal to the x -direction. It would be interesting to introduce the boundaries orthogonal to the y or z direction and investigate the behavior of the system at corners. Furthermore, studying oblique boundaries is also promising direction, as our system does not have the continuous rotational symmetry. These questions are left for future work.

Acknowledgement

We would like to thank Hiromi Ebisu, Ken Shiozaki for useful discussions. The work of S.Y. was supported in part by JSPS KAKENHI Grant Number 21K03574.

References

- [1] C. Chamon, “Quantum Glassiness,” *Phys. Rev. Lett.* **94** no. 4, (2005) 040402, [arXiv:cond-mat/0404182](#).
- [2] J. Haah, “Local stabilizer codes in three dimensions without string logical operators,” *Phys. Rev. A* **83** no. 4, (2011) 042330, [arXiv:1101.1962 \[quant-ph\]](#).
- [3] R. M. Nandkishore and M. Hermele, “Fractons,” *Ann. Rev. Condensed Matter Phys.* **10** (2019) 295–313, [arXiv:1803.11196 \[cond-mat.str-el\]](#).
- [4] M. Pretko, X. Chen, and Y. You, “Fracton Phases of Matter,” *Int. J. Mod. Phys. A* **35** no. 06, (2020) 2030003, [arXiv:2001.01722 \[cond-mat.str-el\]](#).
- [5] A. Gromov and L. Radzihovsky, “Colloquium: Fracton matter,” *Rev. Mod. Phys.* **96** no. 1, (2024) 011001, [arXiv:2211.05130 \[cond-mat.str-el\]](#).
- [6] S. Yamaguchi, “Supersymmetric quantum field theory with exotic symmetry in 3+1 dimensions and fermionic fracton phases,” *PTEP* **2021** no. 6, (2021) 063B04, [arXiv:2102.04768 \[hep-th\]](#).
- [7] N. Seiberg, “Field Theories With a Vector Global Symmetry,” *SciPost Phys.* **8** no. 4, (2020) 050, [arXiv:1909.10544 \[cond-mat.str-el\]](#).
- [8] N. Seiberg and S.-H. Shao, “Exotic Symmetries, Duality, and Fractons in 2+1-Dimensional Quantum Field Theory,” [arXiv:2003.10466 \[cond-mat.str-el\]](#).
- [9] N. Seiberg and S.-H. Shao, “Exotic $U(1)$ Symmetries, Duality, and Fractons in 3+1-Dimensional Quantum Field Theory,” *SciPost Phys.* **9** no. 4, (2020) 046, [arXiv:2004.00015 \[cond-mat.str-el\]](#).
- [10] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, “More Exotic Field Theories in 3+1 Dimensions,” *SciPost Phys.* **9** (2020) 073, [arXiv:2007.04904 \[cond-mat.str-el\]](#).
- [11] H. Katsura and Y. Nakayama, “Spontaneously broken supersymmetric fracton phases with fermionic subsystem symmetries,” *JHEP* **08** (2022) 072, [arXiv:2204.01924 \[hep-th\]](#).

- [12] M. Honda and T. Nakanishi, “Scalar, fermionic and supersymmetric field theories with subsystem symmetries in $d + 1$ dimensions,” *JHEP* **03** (2023) 188, [arXiv:2212.13006 \[hep-th\]](#).
- [13] Y. You and F. von Oppen, “Building fracton phases by majorana manipulation,” *Physical Review Research* **1** no. 1, (2019) 013011.
- [14] N. Tantivasadakarn, “Jordan-Wigner Dualities for Translation-Invariant Hamiltonians in Any Dimension: Emergent Fermions in Fracton Topological Order,” *Phys. Rev. Res.* **2** no. 2, (2020) 023353, [arXiv:2002.11345 \[cond-mat.str-el\]](#).
- [15] W. Shirley, “Fractonic order and emergent fermionic gauge theory,” [arXiv:2002.12026 \[cond-mat.str-el\]](#).
- [16] W. Cao, M. Yamazaki, and Y. Zheng, “Boson-fermion duality with subsystem symmetry,” *Phys. Rev. B* **106** no. 7, (2022) 075150, [arXiv:2206.02727 \[cond-mat.str-el\]](#).
- [17] C. G. Callan, Jr. and J. A. Harvey, “Anomalies and Fermion Zero Modes on Strings and Domain Walls,” *Nucl. Phys. B* **250** (1985) 427–436.
- [18] Y. You, T. Devakul, F. J. Burnell, and S. L. Sondhi, “Subsystem symmetry protected topological order,” *Phys. Rev. B* **98** no. 3, (2018) 035112, [arXiv:1803.02369 \[cond-mat.str-el\]](#).
- [19] T. Devakul, D. J. Williamson, and Y. You, “Classification of subsystem symmetry-protected topological phases,” *Phys. Rev. B* **98** no. 23, (2018) 235121, [arXiv:1808.05300 \[cond-mat.str-el\]](#).
- [20] T. Devakul, W. Shirley, and J. Wang, “Strong planar subsystem symmetry-protected topological phases and their dual fracton orders,” *Phys. Rev. Res.* **2** no. 1, (2020) 012059, [arXiv:1910.01630 \[cond-mat.str-el\]](#).
- [21] F. J. Burnell, T. Devakul, P. Gorantla, H. T. Lam, and S.-H. Shao, “Anomaly inflow for subsystem symmetries,” *Phys. Rev. B* **106** no. 8, (2022) 085113, [arXiv:2110.09529 \[cond-mat.str-el\]](#).
- [22] S. Yamaguchi, “Gapless edge modes in (4+1)-dimensional topologically massive tensor gauge theory and anomaly inflow for subsystem symmetry,” *PTEP* **2022** no. 3, (2022) 033B08, [arXiv:2110.12861 \[hep-th\]](#).
- [23] S. Shimamura, “Anomaly of subsystem symmetries in exotic and foliated BF theories,” *JHEP* **06** (2024) 002, [arXiv:2404.10601 \[cond-mat.str-el\]](#).

- [24] T. Okuda, A. Parayil Mana, and H. Sukeno, “Anomaly inflow for CSS and fractonic lattice models and dualities via cluster state measurement,” *SciPost Phys.* **17** no. 4, (2024) 113, [arXiv:2405.15853 \[quant-ph\]](#).
- [25] H. Ebisu, M. Honda, and T. Nakanishi, “Anomaly inflow for dipole symmetry and higher form foliated field theories,” *JHEP* **09** (2024) 061, [arXiv:2406.04919 \[cond-mat.str-el\]](#).
- [26] A. Kitaev, “Unpaired Majorana fermions in quantum wires,” *Phys. Usp.* **44** no. 10S, (2001) 131–136, [arXiv:cond-mat/0010440](#).