# The Clifford hierarchy for one qubit or qudit

Nadish de Silva and Oscar Lautsch

Department of Mathematics, Simon Fraser University, Burnaby, B.C., Canada

### Abstract

The Clifford hierarchy is a nested sequence of sets of quantum gates that can be fault-tolerantly performed using gate teleportation within standard quantum error correction schemes. The groups of Pauli and Clifford gates constitute the first and second 'levels', respectively. Non-Clifford gates from the third level or higher, such as the T gate, are necessary for achieving fault-tolerant universal quantum computation.

Since it was defined twenty-five years ago by Gottesman-Chuang, two questions have been studied by numerous researchers. First, precisely which gates constitute the Clifford hierarchy? Second, which subset of the hierarchy gates admit efficient gate teleportation protocols?

We completely solve both questions in the practically-relevant case of the Clifford hierarchy for gates of one qubit or one qudit of prime dimension. We express every such hierarchy gate uniquely as a product of three simple gates, yielding also a formula for the size of every level. These results are a consequence of our finding that all such hierarchy gates can be expressed in a certain form that guarantees efficient gate teleportation. Our decomposition of Clifford gates as a unique product of three elementary Clifford gates is of broad applicability.

## Contents

1	Introduction	<b>2</b>
	1.1 Prior work	2
	1.2 Summary of main results	2
<b>2</b>	Background	3
	2.1 Notational conventions	3
	2.2 Pauli gates	
	2.3 The Pauli basis	
	2.4 Conjugate pairs	
	2.5 Clifford gates	
	2.6 The Clifford hierarchy	
	2.7 Diagonal hierarchy gates	6
	2.8 Semi-Clifford gates	
3	Technical preliminaries	7
	3.1 A normal form for Clifford gates	7
	3.2 Almost diagonal, simplified, and simplifiable gates	8
	3.3 Checking Clifford-conjugate equivalence	
4	Main Results	9
	4.1 Overview	
	4.2 Semi-Clifford gates and simplifiability	
	4.3 Characterising semi-Cliffordness via Pauli supports	11
	4.4 Every hierarchy gate is semi-Clifford	12
	4.5 A normal form for Clifford hierarchy gates	15
<b>5</b>	Conclusions	16
6	Acknowledgments	16

## 1 Introduction

The challenges of manipulating quantum data in the presence of environmental noise necessitated the development of quantum error-correcting codes [30]. Quantum gates can be performed directly on encoded data; doing so in a way that suppresses errors to a manageable level is the subject of quantum fault tolerance.

The most commonly studied and utilised family of quantum error-correcting codes are stabiliser codes [16]. Members of the group of *Clifford gates* are special in that they can typically be easily applied fault-tolerantly to data protected by stabiliser codes. Quantum universality, however, further requires the ability to perform non-Clifford gates. Gottesman-Chuang introduced the now-standard technique of *gate teleportation* to fault-tolerantly implement certain non-Clifford gates using only Clifford gates supplemented with ancillary *magic state* resources [17]. In the same work, they introduced the *Clifford hierarchy* (Definition 10), a nested sequence of sets of gates that can be implemented in this way.

A significant practical barrier to achieving quantum universality using this method is the need to prepare magic states for every desired application of a non-Clifford gate. The need to reduce the burden of this substantial resource overhead cost led to the study of more efficient protocols. Hierarchy gates that are diagonal in the computational basis can be implemented using magic states that are half the size of the magic states required in the standard gate teleportation protocol [36]. For example, the T gate, which is a diagonal third-level gate, can be implemented using the magic state  $(\mathbb{I} \otimes T) |\Psi_{\text{Bell}}\rangle$  using the standard protocol or using  $T |+\rangle$  with the efficient one. The efficient gate teleportation protocol for diagonal gates was generalised to also implement 'nearly diagonal' *semi-Clifford gates*, i.e. those gates G such that  $G = C_1 DC_2$  for  $C_1, C_2$  being Clifford gates and D a diagonal hierarchy gate [35]. This decomposition of semi-Clifford gates is highly nonunique.

The important role that the Clifford hierarchy and semi-Clifford gates play in fault-tolerant quantum computation motivates two key questions.

**Question 1.** Precisely which gates belong to the  $k^{th}$  level of the Clifford hierarchy?

**Question 2.** For which (d, n, k) are all  $k^{th}$ -level gates of n qubits or qudits (of dimension d) semi-Clifford?

### 1.1 Prior work

Research into the structure of the Clifford hierarchy and semi-Clifford gates has remained active from their discovery twenty-five years ago to the present [1, 7, 10, 12, 17, 19, 25, 27, 35, 36].

The most progress towards answering the first question was made by Cui-Gottesman-Krishna [10], who gave an explicit description of the diagonal hierarchy gates in terms of certain polynomials over finite fields. Their results apply to both the case of qubit gates and, more generally, to qudit gates of prime dimension.

Zeng-Chen-Chuang [35] gave a proof, aided by exhaustive computations, that all hierarchy gates of one or two qubits are semi-Clifford. They also showed that there exist three-qubit gates in the  $k^{\text{th}}$  level that are *not* semi-Clifford whenever  $k \geq 4$ . Beigi-Shor [4] reported a construction of Gottesman-Mochon showing that there exist *n*-qubit gates in the third level that are not semi-Clifford whenever  $n \geq 7$ .

More recent work of the first author [12] generalised the efficient gate teleportation protocol to higher prime dimensions and initiated the study of qudit semi-Clifford gates. In this work, it was proved that all third-level gates of one qudit are semi-Clifford. Further work of the first author and Chen utilised tools of algebraic geometry to establish that all third-level gates of two qudits are semi-Clifford [7].

The result of the present work is the first to allow both d and k to vary.

### **1.2** Summary of main results

In this work, we give a complete answer to Questions 1 and 2 in the case of one qubit or qudit of prime dimension.

- We show that all one-qudit gates from any level of the Clifford hierarchy are semi-Clifford (Theorem 6).
- To establish the above result, we show how to uniquely decompose a Clifford gate as a product of three elementary Clifford gates: *MDP* (Lemma 9). Here *M* is chosen from a finite set (see Definition 17), *D* is diagonal, and *P* is a permutation gate. This normal form for Clifford gates is of broader applicability.
- Combining the above two results yields a unique decomposition of any non-Clifford one-qubit or one-qudit gate from any level of the Clifford hierarchy as a product of three simple gates: *MDC* (Theorem 7). Here, *M* is as above, *D* is a diagonal hierarchy gate, and *C* is any Clifford gate; every *MDC* is a hierarchy gate.
- This yields a formula that counts the size of every level of the Clifford hierarchy (Corollary 2).

## 2 Background

Below, we review the mathematical definitions that we will require to state and prove our results.

This work builds upon an algebraic framework for studying the Clifford hierarchy based on the discrete Stonevon Neumann theorem that was articulated in [12]. It contains a more detailed introduction to the basic concepts studied below; it explicitly treats the general multiqudit case. Since, in this work, we are concerned with the onequbit or one-qudit case, our presentation below is restricted accordingly for simplicity. In this section, we assume that d is an odd prime but all of the background material extends to qubits with minor modifications; we explicitly mention the qubit case modifications only in the instances where we will require them.

### 2.1 Notational conventions

Let  $d \in \mathbb{N}$  be an odd prime. Denote by  $\omega = e^{i2\pi/d}$  the *d*-th primitive root of unity. The *computational basis* of  $\mathbb{C}^d$  is the standard basis and is denoted by the kets  $|z\rangle$  for  $z \in \mathbb{Z}_d$  where  $\mathbb{Z}_d$  is the field of integers modulo *d*.

The unitary group of  $\mathbb{C}^d$  is denoted  $\mathcal{U}$ . When we refer to a unitary U up to phase we mean its equivalence class [U] under the equivalence relation  $U \sim V \iff U = e^{i\theta}V$  for some  $\theta \in \mathbb{R}$ .

We denote by  $\mathbb{T}$  the complex unit circle group. For a function  $f : \mathbb{Z}_d \to \mathbb{C}$  we denote by D[f] the diagonal matrix with f(z) as its  $z^{\text{th}}$  diagonal entry:  $D[f] |z\rangle = f(z) |z\rangle$ .

### 2.2 Pauli gates

The Pauli gates form the basis of the stabiliser codes used for quantum error correction. Quantum data are encoded as simultaneous eigenvectors of commuting sets of Pauli gates; projections onto eigenspaces of Pauli gates represent the elementary measurements of stabiliser theory.

The basic Pauli gates for a single qudit are  $Z, X \in \mathcal{U}$ :  $Z |z\rangle = \omega^z |z\rangle$  and  $X |z\rangle = |z+1\rangle$  where the addition is taken modulo d. These unitaries have order d. They also satisfy the Weyl commutation relations:

$$ZX = \omega XZ.$$
 (1)

**Definition 1.** The group of Pauli gates on one qudit is the subgroup of  $\mathcal{U}$  generated by the basic Pauli gates:

$$\mathcal{C}_1 = \{ \omega^c Z^p X^q \mid c \in \mathbb{Z}_d, (p,q) \in \mathbb{Z}_d^2 \}.$$

$$\tag{2}$$

### 2.3 The Pauli basis

Using the above commutation relations, it is straightforward to check that Pauli gates from different phase classes are orthogonal with respect to the Hilbert-Schmidt inner product. For each choice of  $p, q \in \mathbb{Z}_d$ , one representative from each phase class is carefully chosen.

**Definition 2.** The Pauli basis for  $M_d(\mathbb{C})$  is a set of Pauli gates defined for each  $p, q \in \mathbb{Z}_d$  as

$$W(p,q) = \omega^{-2^{-1}pq} Z^p X^q.$$
(3)

The pairs  $(p,q) \in \mathbb{Z}_d^2$  admit a symplectic product calculated within  $\mathbb{Z}_d$ :

$$[(p,q),(p',q')] = p \cdot q' - p' \cdot q.$$
(4)

The phases for the Pauli basis were chosen so that  $W(p,q)^* = W(-p,-q)$  and to satisfy the commutation relation:

$$W(p,q)W(p',q') = \omega^{[(p,q),(p',q')]}W(p',q')W(p,q).$$
(5)

We will frequently decompose gates using the Pauli basis and collate the resulting coefficients.

**Definition 3.** Let  $M \in M_d(\mathbb{C})$  be a matrix. We define the function  $f_M : \mathbb{Z}_d^2 \to \mathbb{C}$  by

$$f_M(p,q) = d^{-1} \operatorname{Tr}(W(-p,-q)M).$$
 (6)

Cyclicity and linearity of trace straightforwardly yield a convolution formula for products of matrices.

**Lemma 1.** For any  $U, V \in M_d(\mathbb{C})$  and  $p, q \in \mathbb{Z}_d$ , we have

$$f_{UV}(p,q) = \sum_{i,j \in \mathbb{Z}_d} \omega^{2^{-1}(qi-pj)} f_U(i,j) f_V(p-i,q-j)$$
(7)

$$= \sum_{i,j\in\mathbb{Z}_d} \omega^{-2^{-1}(qi-pj)} f_U(p-i,q-j) f_V(i,j).$$
(8)

We will be particularly concerned with the set of Pauli basis elements for which a given matrix has nonzero coefficients. Pllaha et al. previously studied Clifford and third-level gates via their Pauli support in the qubit case [25].

**Definition 4.** The Pauli support of a matrix  $M \in M_d(\mathbb{C})$ , denoted  $\operatorname{supp}(f_M)$ , is the set of points  $(p,q) \in \mathbb{Z}_d^2$  for which  $f_M(p,q) \neq 0$ .

We can represent the Pauli support of  $M \in M_d(\mathbb{C})$  by creating a diagram for the *phase plane*  $\mathbb{Z}_d^2$  and highlighting the points where  $f_M(p,q)$  is nonzero. An example with d = 5 is shown below.



Figure 1: The Pauli support of the linear combination  $X^2 + 3Z^2X - Z^3X^3$ . The powers of Z range from 0 to d-1 along the horizontal axis, while the powers of X range along the vertical axis.

Lemma 2. A gate of order d is a Pauli gate if and only if its Pauli support is of size 1.

### 2.4 Conjugate pairs

If we conjugate  $Z, X \in \mathcal{U}$  by any unitary gate G, the gates  $G_Z = GZG^*$  and  $G_X = GXG^*$  also satisfy the Weyl commutation relations:  $G_Z{}^d = G_X{}^d = \mathbb{I}$  and  $G_ZG_X = \omega G_XG_Z$ . Remarkably, if (U, V) is any pair of unitaries satisfying these relations, there exists a gate G, unique up to phase, such that  $U = G_Z$  and  $V = G_X$ .

**Definition 5.** An ordered pair of unitaries  $(U, V) \in \mathcal{U}^2$  is a conjugate pair if

- 1.  $U^d = \mathbb{I}$  and  $V^d = \mathbb{I}$ ,
- 2.  $UV = \omega VU$ .

As mentioned, conjugate pairs are in bijective correspondence with one-qudit gates up to phase [12, Lemma 3.4].

**Lemma 3** (Discrete Stone-von Neumann theorem, single-qudit version). Suppose  $(U, V) \in \mathcal{U}^2$  is a conjugate pair. There is a unitary G, unique up to phase, such that  $U = GZG^*$  and  $V = GXG^*$ .

We call  $(GZG^*, GXG^*)$  the pair corresponding to the gate G. Below, we study gates via their conjugate pairs.

### 2.5 Clifford gates

Clifford gates typically have a simple fault-tolerant implementation on stabiliser codes.

**Definition 6.** The group of Clifford gates is the normaliser of the group of Pauli gates as a subgroup of  $\mathcal{U}$ :

$$\mathcal{C}_2 = \{ C \in \mathcal{U} \mid C\mathcal{C}_1 C^* \subseteq \mathcal{C}_1 \}.$$
(9)

**Definition 7.** A gate  $G_1 \in \mathcal{U}$  is Clifford-conjugate to  $G_2 \in \mathcal{U}$  if there exists a Clifford gate  $C \in \mathcal{C}_2$  such that  $G_1 = CG_2C^*$ .

**Definition 8.** A conjugate pair  $(U, V) \in \mathcal{U}^2$  is jointly Clifford-conjugate to a conjugate pair  $(U', V') \in \mathcal{U}^2$  if there exists a Clifford gate  $C \in \mathcal{C}_2$  such that  $U' = CUC^*$  and  $V' = CVC^*$ .

We define the group of Clifford gates up to phase:  $[C_2] = C_2/\mathbb{T}$ . Clifford gates of one qubit or qudit, up to phase, are in correspondence with affine symplectic transformations of  $\mathbb{Z}^2_d$  [18, 26].

The group of symplectic linear transformations  $\text{Sp}(1, \mathbb{Z}_d)$  coincides with  $\text{SL}(2, \mathbb{Z}_d)$ . The group  $\text{Sp}(1, \mathbb{Z}_d) \ltimes \mathbb{Z}_d^2$  of affine symplectic transformations of  $\mathbb{Z}_d^2$  are pairings of  $2 \times 2$  symplectic matrices over  $\mathbb{Z}_d$  and translations in  $\mathbb{Z}_d^2$  with the composition law:

$$(A, v) \circ (B, w) = (AB, Aw + v).$$
 (10)

There is a projective representation  $\rho : \operatorname{Sp}(1, \mathbb{Z}_d) \ltimes \mathbb{Z}_d^2 \to [\mathcal{C}_2]$  that is an isomorphism between the groups of affine symplectic transformations and Clifford gates up to phase.

We will require an explicit representation of the symplectic group given by Neuhauser [24]; we have slightly changed convention by associating a pair  $(p,q) \in \mathbb{Z}_d$  with  $Z^p X^q$  rather than  $X^{-p} Z^q$ . First, note that  $\operatorname{Sp}(1,\mathbb{Z}_d)$  is generated by the set of matrices of the form

$$\begin{pmatrix} a^{-1} & 0\\ 0 & a \end{pmatrix}, \quad \text{and} \quad \begin{pmatrix} 1 & b\\ 0 & 1 \end{pmatrix}, \quad \text{and} \quad \begin{pmatrix} 0 & 1\\ -1 & 0 \end{pmatrix}, \quad (11)$$

where  $a \in \mathbb{Z}_d^*$  and  $b \in \mathbb{Z}_d$ .

**Theorem 1** (Neuhauser's representation of  $\text{Sp}(1, \mathbb{Z}_d)$ ). There is a projective representation  $\mu : \text{Sp}(1, \mathbb{Z}_d) \to [\mathcal{C}_2]$ , defined by sending generators of  $\text{Sp}(1, \mathbb{Z}_d)$  to the following gates, up to phase:

- For all  $a \in \mathbb{Z}_d^*$ ,  $\mu\left(\begin{pmatrix} a^{-1} & 0 \\ 0 & a \end{pmatrix}\right)$  is the Clifford permutation gate that sends  $|z\rangle$  to  $|az\rangle$ .
- For all  $b \in \mathbb{Z}_d$ ,  $\mu(\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix})$  is the diagonal gate D[f], where in the qudit case  $f : \mathbb{Z}_d \to \mathbb{C}$  is the map  $f(z) = \omega^{2^{-1}bz^2}$ . In the qubit case, we instead have  $f(z) = i^{bz}$ , as noted in [26].
- $\mu\left(\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}\right)$  is the Hadamard gate H with  $H_{ij} = d^{-2^{-1}}\omega^{ij}$ .

We can use  $\mu$  to define the isomorphism  $\rho : \operatorname{Sp}(1, \mathbb{Z}_d) \ltimes \mathbb{Z}_d^2 \to [\mathcal{C}_2]$  by:

$$\rho((S, (p, q))) = [W(p, q)\mu(S)].$$
(12)

**Definition 9.** A symplectic Clifford gate  $C \in C_2$  is one such that there exists  $S \in \text{Sp}(1, Z_d)$  such that  $\mu(S) = [C]$ .

If C is a symplectic Clifford gate and  $S \in \text{Sp}(1, \mathbb{Z}_d)$  is the symplectic transformation corresponding to [C] under Neuhauser's representation, then for all  $p, q \in \mathbb{Z}_d$ ,

$$CW(p,q)C^* = W(S(p,q)).$$
 (13)

The group of Clifford gates is a maximal nondense subgroup of the unitary group, and thus, approximately performing arbitrary computations requires the ability to fault-tolerantly perform a non-Clifford gate.

### 2.6 The Clifford hierarchy

The standard choices for non-Clifford gates to supplement the group of Clifford gates in order to achieve universal quantum computation come from the *Clifford hierarchy* [17]. This is a recursively-defined and nested sequence of subsets of  $\mathcal{U}$ . The groups of Pauli and Clifford gates form the first and second *levels* respectively:  $\mathcal{C}_1 \subset \mathcal{C}_2 \subset \mathcal{C}_3 \subset \dots$ 

Non-Clifford gates G that are in the third level or higher can be implemented fault-tolerantly on encoded data indirectly via a gate teleportation protocol. Such a protocol takes as input an arbitrary data state  $|\psi\rangle$  and a resource magic state  $(\mathbb{I} \otimes G) |\Psi_{\text{Bell}}\rangle$  and as output produces, using only gates from lower levels of the hierarchy and standard measurements,  $G |\psi\rangle$ : the data state with the desired gate applied to it.

**Definition 10** (Gottesman-Chuang, 1999). The Clifford hierarchy is an inductively defined sequence of sets of gates. For  $k \ge 2$ , the k<sup>th</sup> level of the Clifford hierarchy is the set:

$$\mathcal{C}_k = \{ G \in \mathcal{U} \mid G\mathcal{C}_1 G^* \subseteq \mathcal{C}_{k-1} \}.$$
(14)

While the first two levels form groups, higher levels do not. The sets  $C_k$  are closed under left or right multiplication by Clifford gates: for  $k \ge 2$ ,  $C_2 C_k C_2 = C_k$  [35].

The following definition is used to characterise those conjugate pairs that correspond to gates of the Clifford hierarchy [12, Theorem 3.11].

**Definition 11.** A conjugate pair  $(U, V) \in \mathcal{U}^2$  is k-closed if it generates a group of  $k^{th}$ -level gates. Equivalently:

$$\{U^p V^q \mid (p,q) \in \mathbb{Z}_d\} \subseteq \mathcal{C}_k.$$
(15)

**Theorem 2.** Gates of the k + 1<sup>th</sup> level of the Clifford hierarchy, up to phase, are in bijective correspondence (via Lemma 3) with k-closed conjugate pairs.

### 2.7 Diagonal hierarchy gates

Of particular interest are the *diagonal* Clifford hierarchy gates. Every level restricted in this way forms a group.

**Definition 12.** The group of phaseless diagonal  $k^{\text{th}}$ -level gates is:

$$\mathcal{D}_k = \{ D \in \mathcal{C}_k \mid D \text{ is a diagonal gate and } D \mid 0 \rangle = \mid 0 \rangle \}.$$
(16)

We will denote the group of diagonal  $k^{\text{th}}$ -level gates by  $\overline{\mathcal{D}}_k = \mathbb{T} \mathcal{D}_k$ .

The condition that D fixes  $|0\rangle$  ensures that  $\mathcal{D}_k$  contains precisely one representative from every phase class. Cui-Gottesman-Krishna [10] characterised the diagonal hierarchy gates exactly for any number of qubits or qudits. We give a simplified restatement of their result restricted to the one-qudit case. This appeared, along with a concise proof of its correctness, in earlier work of the first author [12].

First, note that any integer  $k \ge 1$  can be uniquely expressed as k = (m-1)(d-1) + a with  $a \in \{1, ..., d-1\}$ . Given  $m \ge 1$ , denote the  $d^m$ -th primitive root of unity:  $\omega_m = e^{i 2\pi/d^m}$ . Denote by  $\mathcal{R}_k$  the set of rank-k polynomials:

$$\mathcal{R}_{k} = \left\{ \begin{array}{c} \phi : \mathbb{Z}_{d^{m}} \to \mathbb{Z}_{d^{m}} \\ \phi(z) = \sum_{j=1}^{d-1} \phi_{j} z^{j} \end{array} \middle| \begin{array}{c} \phi \text{ has degree at most } d-1, \ \phi(0) = 0 \\ \phi_{a+1}, \dots, \phi_{d-1} \equiv 0 \pmod{d} \end{array} \right\}.$$
(17)

**Theorem 3.**  $\mathcal{D}_k = \{D[\omega_m^{\phi}] \mid \phi \in \mathcal{R}_k\} \text{ for all } k \geq 1.$ 

One of our main results below (Theorem 7) extends this classification from  $\mathcal{D}_k$  to all of  $\mathcal{C}_k$ .

### 2.8 Semi-Clifford gates

Zhou-Leung-Chuang [36] introduced a simplified gate teleportation protocol, based on Bennett-Gottesman's onequbit teleportation, capable of fault-tolerantly implementing certain qubit Clifford hierarchy gates using half the ancillary resources required in the original Gottesman-Chuang protocol. This class of gates includes the diagonal Clifford hierarchy gates. Zeng-Chen-Chuang [35] introduced the notion of semi-Clifford gates which are 'nearly diagonal' in the sense of being within Clifford corrections of diagonal gates:

**Definition 13.** A gate  $G \in C_k$  is semi-Clifford if  $G = C_1DC_2$  where  $C_1, C_2 \in C_2$  and D is diagonal.

**Definition 14.** For  $k \ge 1$  the  $k^{th}$ -level semi-Clifford gates are:

$$\mathcal{SC}_k = \{ G \in \mathcal{C}_k \mid G = C_1 D C_2 \text{ where } C_1, C_2 \in \mathcal{C}_2, D \in \mathcal{D}_k \}.$$
(18)

The following gate teleportation protocol for implementing the qudit semi-Clifford gate  $G = C_1 D C_2$  using the magic state  $|M\rangle = D |+\rangle$  was introduced in [12, §5(a)]. It ensures that the notion of semi-Clifford gate is still relevant in the qudit setting.



Figure 2: The compact gate teleportation protocol for semi-Clifford gates.

The following two lemmas are a direct consequence of Definition 13.

**Lemma 4.** If  $G \in SC_k$  is semi-Clifford then, for any Clifford  $C \in C_2$ , GC and CG are also semi-Clifford. **Lemma 5.** If  $G_1 \in \mathcal{U}$  is Clifford-conjugate to  $G_2 \in \mathcal{U}$ , then  $G_1$  is semi-Clifford if and only if  $G_2$  is.

#### 3 Technical preliminaries

In this section, we develop some tools and techniques necessary for our proof that are either novel or build substantially on prior work. These have wider applicability beyond their use below.

#### A normal form for Clifford gates 3.1

In this section, we develop a normal form for  $C_2$ , which is a way to write each gate in  $C_2$  uniquely as the product of three elementary Clifford gates. For simplicity, we focus on the case of one qubit or qudit. Our normal form extends to the multi-qubit or -qudit case; this is the subject of forthcoming work.

**Definition 15.** A permutation gate  $P \in \mathcal{U}$  is one defined by  $P|z\rangle = |\sigma(z)\rangle$  for some permutation  $\sigma : \mathbb{Z}_d \to \mathbb{Z}_d$ .

**Definition 16.** The group of Clifford permutation gates, denoted  $\mathcal{N} \subset \mathcal{U}$ , is the intersection of the group of Clifford gates with the group of permutation gates.

Two examples include those arising from  $\sigma(z) = az$  for any  $a \in \mathbb{Z}_d^*$  (see Theorem 1) and  $\sigma(z) = z + b$  for any  $b \in \mathbb{Z}_d$  (which correspond to elements of  $\langle X \rangle$ ). These, in fact, generate all examples.

**Lemma 6.** Every Clifford permutation gate arises from  $\sigma(z) = az + b$  for some  $a \in \mathbb{Z}_d^*, b \in \mathbb{Z}_d$ .

*Proof.* Given  $P \in \mathcal{U}$  arising from  $\sigma : \mathbb{Z}_d \to \mathbb{Z}_d$ ,  $PZP^* |z\rangle = \omega^{\sigma^{-1}(z)} |z\rangle$ . Thus  $PZP^*$  is diagonal; it is a Pauli gate only if it is a power of Z multiplied by a power of  $\omega$ . Thus  $\sigma$  is an affine linear map.  $\square$ 

**Lemma 7.**  $\langle X \rangle$  is a normal subgroup of  $\mathcal{N}: \langle X \rangle \trianglelefteq \mathcal{N}$ .

We construct from  $\overline{\mathcal{D}}_k \leq \mathcal{U}$  two other groups at each level of the hierarchy.

Lemma 8. Let  $k \in \mathbb{N}$ ,

$$\overline{\mathcal{D}}_k \langle X \rangle = \{ DX^c \mid D \in \overline{\mathcal{D}}_k, c \in \mathbb{Z}_d \}$$
(19)

and, when  $k \geq 2$ ,

$$\overline{\mathcal{D}}_k \mathcal{N} = \{ DP \mid D \in \overline{\mathcal{D}}_k, P \in \mathcal{N} \}.$$
(20)

Both of these subsets of  $C_k$  are groups. Furthermore, for all applicable k, we have:

$$\overline{\mathcal{D}}_k\langle X\rangle \trianglelefteq \overline{\mathcal{D}}_{k+1}\langle X\rangle, \qquad \overline{\mathcal{D}}_k\langle X\rangle \trianglelefteq \overline{\mathcal{D}}_k\mathcal{N}, \qquad \overline{\mathcal{D}}_k\langle X\rangle \trianglelefteq \overline{\mathcal{D}}_{k+1}\mathcal{N}.$$
(21)

*Proof.* That  $\overline{\mathcal{D}}_k\langle X\rangle$ ,  $\overline{\mathcal{D}}_k\mathcal{N}$  are subsets of  $\mathcal{C}_k$  follows from the closure of  $\mathcal{C}_k$  under multiplication by Clifford gates [35]. That they are groups, i.e. closed under inverses and products, follows straightforwardly by the fact that, for any  $P \in \mathcal{N}$ ,  $PDP^* \in \overline{\mathcal{D}}_k$  if and only if  $D \in \overline{\mathcal{D}}_k$ .

Since the Clifford hierarchy is nested,  $\overline{\mathcal{D}}_k \leq \overline{\mathcal{D}}_{k+1}$  for all k. Then as  $\langle X \rangle \leq \mathcal{N}$ , both  $\overline{\mathcal{D}}_{k+1} \langle X \rangle$  and  $\overline{\mathcal{D}}_k \mathcal{N}$  are subgroups of  $\overline{\mathcal{D}}_{k+1}\mathcal{N}$ . Thus to show that  $\overline{\mathcal{D}}_k\langle X\rangle$  is a normal subgroup of each of  $\overline{\mathcal{D}}_{k+1}\langle X\rangle$ ,  $\overline{\mathcal{D}}_k\mathcal{N}$ , and  $\overline{\mathcal{D}}_{k+1}\mathcal{N}$ , it suffices to only show that  $\overline{\mathcal{D}}_k\langle X\rangle$  is a normal subgroup of  $\overline{\mathcal{D}}_{k+1}\mathcal{N}$ .

Let  $D_1 X^b \in \overline{\mathcal{D}}_k \langle X \rangle$  and  $D_2 P \in \overline{\mathcal{D}}_{k+1} \mathcal{N}$ . Then

$$(D_2 P)(D_1 X^b)(D_2 P)^* = D_2 (P D_1 P^*)(P X^b P^*) D_2^*$$
(22)

$$= D_2(PD_1P^*)X^cD_2^*$$
(23)

$$= D_2(PD_1P^*)X^cD_2^*$$
(23)  
=  $(PD_1P^*)(D_2X^cD_2^*)$ (24)

$$= (PD_1P^*)D'X^c \in \overline{\mathcal{D}}_k\langle X\rangle.$$
(25)

The second equality follows from  $\langle X \rangle \leq \mathcal{N}$ . The third equality follows from the fact that the diagonal gates  $D_2$ and  $PD_1P^*$  commute. The final equality follows by noting that  $D_2X^cD_2^* = D'X^c$  for some diagonal  $D' \in \overline{\mathcal{D}}_k$  and that  $P \in \mathcal{C}_2$  implies that  $PD_1P^* \in \overline{\mathcal{D}}_k$ . 

By choosing convenient coset representatives for the subgroup  $\overline{\mathcal{D}}_2 \mathcal{N}$  of  $\mathcal{C}_2$ , we construct a normal form for Clifford gates.

**Definition 17.** In the qubit case, let  $E \in \mathcal{U}$  be the phase gate S; in the qudit case, let E = D[h], where  $h : \mathbb{Z}_d \to \mathbb{C}$  is the map  $h(z) = \omega^{(z^2)}$ .

We define  $\mathcal{M} \subset \mathcal{U}$  to be the set of d+1 gates  $\mathcal{M} = \{\mathbb{I}\} \cup \{E^c H \mid c \in \mathbb{Z}_d\}.$ 

Note that by Theorem 3, E is a generator for the group of diagonal symplectic Clifford gates.

Lemma 9 (Normal form for Clifford gates). The following is a normal form for Clifford gates:

$$\mathcal{C}_2 = \mathcal{M}\overline{\mathcal{D}}_2 \mathcal{N}.$$
 (26)

That is, every Clifford gate may be written uniquely, as MDP, where  $M \in \mathcal{M}$ ,  $D \in \overline{\mathcal{D}}_2$ , and  $P \in \mathcal{N}$ .

*Proof.* We begin by giving a normal form for  $\operatorname{Sp}(1, \mathbb{Z}_d)$ . By Theorem 3.1 of [24],  $\binom{s_{11} \ s_{12}}{s_{21} \ s_{22}} \in \operatorname{Sp}(1, \mathbb{Z}_d)$  if and only if  $s_{11}s_{22} - s_{12}s_{21} = 1$ . Let  $\binom{s_{11} \ s_{12}}{s_{21} \ s_{22}} \in \operatorname{Sp}(1, \mathbb{Z}_d)$ . This matrix may be written as

$$\begin{pmatrix} s_{11} & s_{12} \\ s_{21} & s_{22} \end{pmatrix} = \begin{pmatrix} 1 & e \\ 0 & 1 \end{pmatrix} \begin{pmatrix} f & 0 \\ 0 & f^{-1} \end{pmatrix}$$
(27)

if and only if  $s_{21} = 0$ ,  $e = s_{11}s_{12}$ , and  $f = s_{11}$ . On the other hand, this matrix may be written as

$$\begin{pmatrix} s_{11} & s_{12} \\ s_{21} & s_{22} \end{pmatrix} = \begin{pmatrix} 1 & g \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & e \\ 0 & 1 \end{pmatrix} \begin{pmatrix} f & 0 \\ 0 & f^{-1} \end{pmatrix}$$
(28)

if and only if  $s_{21} \neq 0$ ,  $e = s_{21}s_{22}$ ,  $f = -s_{21}$ , and  $g = s_{11}s_{21}^{-1}$ . Letting  $A \subseteq \text{Sp}(1, \mathbb{Z}_d)$  denote the set of matrices of the form  $\begin{pmatrix} f & 0 \\ 0 & f^{-1} \end{pmatrix}$  and  $B \subseteq \text{Sp}(1, \mathbb{Z}_d)$  denote the set of matrices of the form  $\begin{pmatrix} 1 & e \\ 0 & 1 \end{pmatrix}$ , it follows that

$$\left(\{\mathbb{I}\} \cup B\begin{pmatrix} 0 & 1\\ -1 & 0 \end{pmatrix}\right) BA \tag{29}$$

is a normal form for  $\text{Sp}(1, \mathbb{Z}_d)$ . By Theorem 1, under the Neuhauser representation, elements of A correspond to symplectic Clifford permutation gates, elements of B correspond to diagonal symplectic Clifford gates, and  $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$  corresponds to H.

Thus, any symplectic Clifford gate may be written uniquely in the form  $MD_1P$ , where  $D_1$  is a diagonal symplectic Clifford gate, P is a symplectic Clifford permutation gate, and  $M = \mathbb{I}$  or  $M = D_2H$  for a diagonal symplectic Clifford gate  $D_2$ . Since any Clifford gate may be written uniquely as a Pauli gate multiplied by a symplectic Clifford gate, it follows that any Clifford gate may be written uniquely up to phase as

$$MD_1 P Z^p X^q \tag{30}$$

for some  $p, q \in \mathbb{Z}_d$ . Note that  $D_1 P Z^p X^q \in \overline{\mathcal{D}}_2 \mathcal{N}$ , so it may be written uniquely as D'P' for some  $D \in \overline{\mathcal{D}}_2$  and  $P \in \mathcal{N}$ . Hence, any Clifford gate may be written uniquely up to phase as MD'P', where  $D' \in \mathcal{D}_2$ ,  $P' \in \mathcal{N}$ , and  $M = \mathbb{I}$  or  $M = D_2 H$  for a symplectic diagonal Clifford gate  $D_2$ . As E is a generator for the group of symplectic diagonal Clifford gates, the claimed normal form follows.

An identical argument shows that  $C_2 = \overline{D}_2 \mathcal{N} \mathcal{M}^{-1}$  is also a normal form for Clifford gates, and we will use both of these normal forms in the remainder of this work.

### 3.2 Almost diagonal, simplified, and simplifiable gates

Here, we generalise two definitions made in the two-qudit third-level case in [7] to higher levels; while these definitions are easily extended to more qudits, we restrict them to one qudit for simplicity.

**Definition 18.** A hierarchy gate  $G \in C_k$  is almost diagonal if  $G \in \overline{D}_k \langle X \rangle$ .

**Definition 19.** A hierarchy gate  $G \in C_k$  is simplified if its conjugate pair contains only almost diagonal gates.

We will show below in Lemma 17 that a gate is simplified if and only if it is of the form DC with D a diagonal hierarchy gate and C a Clifford gate.

**Definition 20.** A hierarchy gate  $G \in C_k$  is simplifiable if it is Clifford-conjugate to a simplified gate.

It will immediately follow from Lemma 17 that a gate is semi-Clifford if and only if it is simplifiable. This reduces our primary goal to showing that every hierarchy gate is simplifiable.

That every third-level *n*-qudit gate is simplifiable [12, Lemma 5.7] was previously shown using a classification of the maximal abelian subgroups of the symplectic group over finite fields due to Barry [3]. In Section 4.2, we will show this to be true of a one-qudit gate from any level.

### 3.3 Checking Clifford-conjugate equivalence

In the qudit case, the Pauli basis decomposition and projective representation of the Clifford group give a convenient characterisation of when two matrices are Clifford-conjugate. This characterisation does not seem to appear explicitly in the literature, so we state it here. It extends naturally to higher numbers of qudits. We omit the proofs as they are relatively straightforward.

**Lemma 10.** Let  $M, M' \in M_d(\mathbb{C})$ . Then M is Clifford-conjugate to M' if and only if there exists  $(S, (p', q')) \in$ Sp $(1, \mathbb{Z}_d) \ltimes \mathbb{Z}_d^2$  such that, for all  $p, q \in \mathbb{Z}_d$ ,

$$f_M(p,q) = w^{[(p,q),(p',q')]} f_{M'}(S(p,q)).$$
(31)

**Lemma 11.** If  $M, M' \in M_d(\mathbb{C})$  are Clifford-conjugate, then there exists  $S \in \text{Sp}(1, \mathbb{Z}_d)$  that restricts to a bijection between the Pauli support of M and that of M'.

## 4 Main Results

Our main technical goal is to prove, in the one-qudit case, that all hierarchy gates are semi-Clifford. This was already shown in the one-qubit case by Zeng-Chen-Chuang [35]. We combine these results with the normal form for Clifford gates established above in both qubit and qudit cases, to find normal forms for the Clifford hierarchy.

### 4.1 Overview

- 1. In Section 4.2, we establish an equivalence between a gate being semi-Clifford and simplifiable.
- 2. In Section 4.3, we find that a one-qudit gate is semi-Clifford if and only if the Pauli supports of its conjugate pair lie within parallel lines in the phase plane.
- 3. In Section 4.4, we inductively prove that every one-qudit hierarchy gate is semi-Clifford via Pauli supports.
- 4. In Section 4.5 we use that all hierarchy gates are semi-Clifford, and our normal form for Clifford gates, to explicitly describe the Clifford hierarchy of one qubit or qudit and find a formula for the size of every level.

### 4.2 Semi-Clifford gates and simplifiability

Lemma 12. Every semi-Clifford gate is simplifiable.

*Proof.* Every Pauli gate is simplified. Thus, let  $k \ge 1$  and  $G = C_1 D C_2 \in \mathcal{C}_{k+1}$  be semi-Clifford, where  $C_1, C_2 \in \mathcal{C}_2$  and  $D \in \mathcal{D}_{k+1}$ . Then  $C_1^* G C_1 = D C$  with  $C = C_2 C_1$ , and Lemma 8 ensures that both elements of the pair corresponding to D C are in  $\overline{\mathcal{D}}_{k-1}\langle X \rangle$ , so D C is simplified.

In this section, we show that the converse of this also holds: that every simplifiable gate is semi-Clifford. The following sequence of lemmas allows us to precisely characterise simplified gates as those of the form DC; a gate is semi-Clifford if and only if is Clifford-conjugate to a gate of the form DC.

**Lemma 13.** Suppose  $(U, V_1)$  and  $(U, V_2)$  are both conjugate pairs. Then U commutes with  $V_1V_2^*$ .

This observation allows us to show that in a conjugate pair of the form  $(D_1 X^{q_1}, D_2 X^{q_2})$  with  $D_1$  and  $D_2$  diagonal,  $D_2$  is almost completely determined by  $D_1$ .

**Lemma 14.** Suppose that  $(D_1X^{q_1}, D_2X^{q_2})$  and  $(D_1X^{q_1}, D_3X^{q_2})$  are conjugate pairs, where  $D_1, D_2$ , and  $D_3$  are diagonal gates and  $q_1 \neq 0$ . Then  $D_3 = \omega^c D_2$  for some  $c \in \mathbb{Z}_d$ .

Proof. By the previous lemma,  $D_1 X^{q_1}$  must commute with  $(D_2 X^{q_2})(D_3 X^{q_2})^* = D_2 D_3^*$ . Then  $X^{q_1}$  must commute with  $D_2 D_3^*$ . As  $q_1 \neq 0$  and d is prime, this implies that  $D_2 D_3^*$  is a constant multiple of  $\mathbb{I}$ . Since both  $D_2 X^{q_2}$  and  $D_3 X^{q_2}$  have order d, this constant multiple must be a power of  $\omega$ .

The next lemma is used to construct the semi-Clifford gate corresponding to a conjugate pair of gates in  $\overline{\mathcal{D}}_k\langle X \rangle$ .

**Lemma 15.** Let  $D' \in \overline{\mathcal{D}}_k$  be a diagonal gate with  $\det(D') = 1$ . Then for any fixed  $c \in \mathbb{Z}_d^*$ , there exists a diagonal gate  $D \in \mathcal{D}_{k+1}$  such that  $D' = DX^c D^* X^{-c}$ .

Proof. Fix  $c \neq 0$  in  $\mathbb{Z}_d$ . As D' is a unitary diagonal, we have D' = D[f] for some  $f : \mathbb{Z}_d \to \mathbb{T}$ . For any function  $g(z) : \mathbb{Z}_d \to \mathbb{C}$ , if D = D[g(z)], then  $X^c D^* X^{-c} = D[g(z-c)^*]$ , and so  $DX^c D^* X^{-c} = D[g(z)g(z-c)^*]$ . Hence,  $D' = DX^c D^* X^{-c}$  if and only if

$$f(z) = g(z)g(z-c)^*$$
 (32)

for all  $z \in \mathbb{Z}_d$ . We will show that a function g satisfying (32) exists. Note that as c is nonzero, we may replace z with cz in (32) to find the equivalent equation

$$f(cz) = g(cz)g(c(z-1))^*.$$
(33)

We construct such a g by defining

$$g(cz) = \prod_{j=0}^{z} f(cj) \tag{34}$$

for all  $z \in \mathbb{Z}_d$ . This ensures that (33) is satisfied for all  $1 \leq z \leq d-1$ . Furthermore, since  $\det(D') = 1$  the product of all the diagonal entries of D' is 1. We thus have  $\prod_{j=0}^{d-1} f(cz) = 1$ , and so g(c(d-1)) = 1. Then  $g(0)g(c(d-1))^* = f(0)$ , so (33) also holds for z = 0. Since  $f(z) \in \mathbb{T}$  for all  $z \in \mathbb{Z}_d$ , we have  $g(z) \in \mathbb{T}$  for all z as well, so D = D[g] is unitary. Thus D is a diagonal gate such that  $D' = DX^c D^* X^{-c}$ .

To see that this  $D \in \overline{\mathcal{D}}_{k+1}$ , note that  $DX^cD^* = D'X^c \in \overline{\mathcal{D}}_k\langle X \rangle$ . As D and Z commute, D conjugates both Z and  $X^c$  into  $\overline{\mathcal{D}}_k\langle X \rangle$ . Since Z and  $X^c$  generate the Pauli group and  $\overline{\mathcal{D}}_k\langle X \rangle$  forms a group, it follows that D conjugates every Pauli into  $\overline{\mathcal{D}}_k\langle X \rangle$ , and so that  $D \in \overline{\mathcal{D}}_{k+1}$ . We choose the phase of D so that  $D \in \mathcal{D}_{k+1}$ .  $\Box$ 

The requirement that D' have determinant 1 in this lemma may appear to restrict its utility. However, we will only need to apply this lemma to gates D' such that  $D'Z^pX^q$  appears in a conjugate pair. As we observe below, in this scenario, requiring that D' have determinant 1 poses no issues.

**Lemma 16.** Suppose that  $D'Z^pX^q$  is conjugate to Z, where D' is a diagonal gate. Then det(D') = 1.

This follows from the fact that det(Z) = det(X) = 1. We can now precisely characterise simplified gates.

**Lemma 17.** A gate  $G \in C_k$  is simplified if and only if it is of the form DC, with  $D \in D_k$  and  $C \in C_2$ .

*Proof.* The case of Pauli gates is immediate; thus let  $k \ge 2$ .

The proof of Lemma 12 establishes that G = DC is simplified. Conversely, suppose that G is simplified. Then

$$(U,V) = (GZG^*, GXG^*) = (D_1X^{q_1}, D_2X^{q_2})$$
(35)

is a conjugate pair with  $U, V \in \overline{\mathcal{D}}_{k-1}\langle X \rangle$ . Since U, V do not commute,  $q_1$  and  $q_2$  cannot both be 0. Assume without loss of generality that  $q_1 \neq 0$ ; the proof of the case with  $q_2 \neq 0$  is completely analogous. There exist  $p_1, p_2 \in \mathbb{Z}_d$ such that  $p_1q_2 - p_2q_1 = 1$ , and there exist  $D'_1, D'_2 \in \overline{\mathcal{D}}_{k-1}$  such that  $D_1 = D'_1Z^{p_1}$  and  $D_2 = D'_2Z^{p_2}$ . Hence,

$$(U,V) = (D_1'Z^{p_1}X^{q_1}, D_2'Z^{p_2}X^{q_2}).$$
(36)

Combining the preceding two lemmas, we have  $D'_1 = DX^{q_1}D^*X^{-q_1}$  for some diagonal gate  $D \in \mathcal{D}_k$ . Thus

$$U = D_1' Z^{p_1} X^{q_1} = D X^{q_1} D^* X^{-q_1} Z^{p_1} X^{q_1} = D Z^{p_1} X^{q_1} D^*.$$
(37)

As  $p_1q_2 - p_2q_1 = 1$ ,  $(Z^{p_1}X^{q_1}, Z^{p_2}X^{q_2})$  is a conjugate pair, so

$$(DZ^{p_1}X^{q_1}D^*, DZ^{p_2}X^{q_2}D^*) = (U, DX^{q_2}D^*X^{-q_2}Z^{p_2}X^{q_2})$$
(38)

is as well. By Lemma 14, it follows that  $D'_2 = \omega^c D X^{q_2} D^* X^{-q_2}$  for some  $c \in \mathbb{Z}_d$ . Hence

$$(U,V) = (DZ^{p_1}X^{q_1}D^*, D\omega^c Z^{p_2}X^{q_2}D^*).$$
(39)

As  $(Z^{p_1}X^{q_1}, \omega^c Z^{p_2}X^{q_2})$  is a conjugate pair of Paulis, there exists a Clifford C such that

$$(CZC^*, CXC^*) = (Z^{p_1}X^{q_1}, \omega^c Z^{p_2}X^{q_2}).$$
(40)

Then  $(U, V) = (DCZC^*D^*, DCXC^*D^*)$ , so DC is the gate corresponding to the pair (U, V).

Our desired equivalence follows immediately.

**Theorem 4** (Equivalence of semi-Cliffordness and simplifiability). A gate  $G \in \mathcal{U}$  is semi-Clifford if and only if it is simplifiable.

*Proof.* We established one direction of this equivalence in Lemma 12. For the converse, suppose that  $G = C_1 G' C_1^*$  where  $G' \in \mathcal{C}_k$  is simplified. Then  $G = C_1 DCC_1^*$  with  $D \in \mathcal{D}_k$ ,  $C \in \mathcal{C}_2$  and is thus semi-Clifford with  $C_2 = CC_1^*$ .  $\Box$ 

### 4.3 Characterising semi-Cliffordness via Pauli supports

Here, we study the Pauli supports of the gates in conjugate pairs (U, V) arising from semi-Clifford gates. Recall, for any  $M \in M_d(\mathbb{C})$ , the definition of  $f_M$  and the Pauli support of M from Definitions 3 and 4.

We begin by understanding the Pauli supports of gates in  $\overline{\mathcal{D}}_k\langle X\rangle$ . By a *line* in  $\mathbb{Z}_d^2$ , we refer to a 1-dimensional affine subspace of the phase plane  $\mathbb{Z}_d^2$ . Two lines are *parallel* if they are affine translates of the same 1-dimensional linear subspace of  $\mathbb{Z}_d^2$ . A line is *horizontal* if it is of the form  $\{(p, q_0) \mid p \in \mathbb{Z}_d\} \subseteq \mathbb{Z}_d^2$  for some  $q_0 \in \mathbb{Z}_d$ .

**Lemma 18.** Let  $G \in \mathcal{U}$ . Then  $G = DX^{q_0}$  for some diagonal gate  $D \in \mathcal{U}$  and  $q_0 \in \mathbb{Z}_d$  if and only if the Pauli support of G is contained in a horizontal line  $\{(p, q_0) \mid p \in \mathbb{Z}_d\} \subseteq \mathbb{Z}_d^2$  for some  $q_0 \in \mathbb{Z}_d$ .

*Proof.* If  $G = DX^{q_0}$  for some diagonal gate  $D \in \overline{\mathcal{D}}_k$  and  $q_0 \in \mathbb{Z}_d$ , then for all  $p, q \in \mathbb{Z}_d$ ,

$$GW(-p,-q) = DX^{q_0}\omega^{-2^{-1}pq}Z^{-p}X^{-q} = D'X^{q_0-q}$$
(41)

for some diagonal D'. If  $q \neq q_0$ , then the matrix  $D'X^{q_0-q}$  has all zeros along its diagonal, and so has trace 0. Hence,  $f_G(p,q) = d^{-1}\text{Tr}(GW(-p,-q))$  is 0 as long as  $q \neq q_0$ . The Pauli support of G is then contained in  $\{(p,q_0) \mid p \in \mathbb{Z}_d\}$ .

The converse follows by noting that if  $\operatorname{supp}(f_G) \subset \{(p, q_0) \mid p \in \mathbb{Z}_d\}$ , then for some constants  $c_p \in \mathbb{C}$ ,

$$G = \sum_{p \in \mathbb{Z}_d} c_p Z^p X^{q_0} = \left(\sum_{p \in \mathbb{Z}_d} c_p Z^p\right) X^{q_0}.$$
(42)

The Pauli supports of gates in  $\overline{\mathcal{D}}_k\langle X\rangle$  can be visualised in Figure 3.



Figure 3: The Pauli support of a diagonal gate is contained in the linear subspace  $\{(p,0) \mid p \in \mathbb{Z}_d\}$ , and in general, the Pauli support of  $DX^{q_0}$  is contained in  $\{(p,q_0) \mid p \in \mathbb{Z}_d\}$  if D is diagonal.

There is an action of  $\text{Sp}(1, \mathbb{Z}_d)$  on the set of lines in  $\mathbb{Z}_d^2$ , with  $S \in \text{Sp}(1, \mathbb{Z}_d)$  sending L to S(L). This induces an action of  $\mathcal{C}_2$  on these lines: for any  $C \in \mathcal{C}_2$ , we may write [C] uniquely as [W(p,q)C'] for some symplectic Clifford C'. Letting the symplectic transformation corresponding to C' be S, C acts on a line  $L \subseteq \mathbb{Z}_d^2$  by sending L to S(L).

Lemma 18 means that horizontal lines will be especially important for us. We thus wish to understand which Clifford gates C send horizontal lines to horizontal lines under the above action. This is equivalent to C mapping the linear subspace  $L_Z = \{(p,0) \mid p \in \mathbb{Z}_d\}$  to itself under this action, since all horizontal lines are affine translates of  $L_Z$ . Finally, as  $L_Z$  is spanned by (1,0), this is equivalent to C sending (1,0) to (p,0) for some  $p \in \mathbb{Z}_d^*$ . The Clifford gate C does this if and only if the corresponding symplectic transformation S is of the form

$$S = \begin{pmatrix} s_{11} & s_{12} \\ 0 & s_{22} \end{pmatrix}. \tag{43}$$

As shown in the proof of our Clifford normal form (Lemma 9), S has the above form if and only if  $C \in \overline{\mathcal{D}}_2 \mathcal{N}$ . We have thus proven the following lemma.

**Lemma 19.** Let  $C \in C_2$ . Then C maps horizontal lines to horizontal lines under the above action if and only if  $C \in \overline{\mathcal{D}}_2 \mathcal{N}$ .

We also wish to have a standard set of Clifford gates to take  $L_Z$  to each 1-dimensional linear subspace of  $\mathbb{Z}_d$ . This is done in the following lemma, in which we use the set  $\mathcal{M}$  of d+1 gates from Definition 17.

**Lemma 20.** Under the above action, each  $M \in \mathcal{M}$  maps  $L_Z = \{(p,0) \mid p \in \mathbb{Z}_d\}$  to a 1-dimensional linear subspace of  $\mathbb{Z}_d^2$ . Furthermore, for any 1-dimensional subspace L, there is a unique  $M \in \mathcal{M}$  that carries  $L_Z$  to L.

*Proof.* It suffices to show that each such M maps (1,0) into a distinct 1-dimensional subspace. The identity I maps (1,0) into  $L_Z$ , and by Theorem 1, the Clifford gate  $E^c H$  corresponds to the symplectic matrix

$$S = \begin{pmatrix} 1 & 2c \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} -2c & 1 \\ -1 & 0 \end{pmatrix}.$$
(44)

S maps (1,0) to (-2c, -1), which never lies in  $L_Z$  and lies in a distinct 1-dimensional subspace for distinct values of c. Furthermore, any 1-dimensional subspace of  $\mathbb{Z}_d^2$  is spanned by either (1,0) or (-2c, -1) for some  $c \in \mathbb{Z}_d$ .  $\Box$ 

This also shows that  $\text{Sp}(1, \mathbb{Z}_d)$  acts transitively on the 1-dimensional subspaces of  $\mathbb{Z}_d^2$ . We can now understand exactly when a conjugate pair (U, V) arises from a semi-Clifford gate in terms of the Pauli supports of U and V.

**Theorem 5.** Let  $G \in C_k$  and (U, V) be the pair corresponding to G. Then G is semi-Clifford if and only if the Pauli supports of U and V are contained in two parallel lines  $L_U$  and  $L_V$  in  $\mathbb{Z}^2_d$ .

*Proof.* By Theorem 4, G is semi-Clifford if and only if

$$(U,V) = (CU'C^*, CV'C^*), (45)$$

for some  $C \in \mathcal{C}_2$  and  $U', V' \in \overline{\mathcal{D}}_k \langle X \rangle$ . By the Lemmas 18 and 10, this is equivalent to the existence of two horizontal lines  $L_{U'}$  and  $L_{V'}$  (containing the Pauli supports of U' and V'), and the existence of a symplectic transformation  $S \in \operatorname{Sp}(1, \mathbb{Z}_d)$  (corresponding to the symplectic part of C), such that the Pauli supports of U and V are contained in  $S(L_{U'})$  and  $S(L_{V'})$ . If such  $L_{U'}, L_{V'}$ , and S exist, then  $L_U = S(L_{U'})$  and  $L_V = S(L_{V'})$  are parallel since S is linear and  $L_{U'}$  and  $L_{V'}$  are parallel. Conversely, if the Pauli supports of U and V are contained in parallel lines  $L_U$  and  $L_V$ , then there exists  $S \in \operatorname{Sp}(1, \mathbb{Z}_d)$  such that  $L_{U'} = S^{-1}(L_U)$  and  $L_{V'} = S^{-1}(L_V)$  are horizontal, since  $\operatorname{Sp}(1, \mathbb{Z}_d)$  acts transitively on the 1-dimensional linear subspaces of  $\mathbb{Z}_d^2$  by the above lemma.

The proof of this corollary can be visualised with Figure 4.



Figure 4: Proof of Theorem 5. The horizontal lines in the left diagram contain the Pauli supports of U' and V', which are mapped to the parallel lines on the right side of the diagram by C.

### 4.4 Every hierarchy gate is semi-Clifford

Recall from Definition 11 that a pair (U, V) is k-closed if the group  $\langle U, V \rangle$  generated by U and V is contained in  $C_k$ . In this section, we will prove our main lemmas (23 and 24), which will show that every k-closed conjugate pair is jointly Clifford-conjugate (recall Definition 8) to a pair of gates in  $\overline{D}_k \langle X \rangle$ . Since gates at level k + 1 correspond to k-closed pairs by Theorem 2, the bijective correspondence established in Theorem 4 will then ensure that every gate at level k + 1 is semi-Clifford. Using induction, we will then be able to show that for every odd prime d, every gate at every level of the one-qudit Clifford hierarchy is semi-Clifford.

We begin by proving that in two special cases, certain order-d gates are Clifford-conjugate to gates in  $\overline{\mathcal{D}}_k\langle X \rangle$ .

**Lemma 21.** Any Clifford gate  $C \in \mathcal{C}_2$  of order d is Clifford-conjugate to a gate in  $\overline{\mathcal{D}}_2\langle X \rangle$ .

*Proof.* If C appears in a conjugate pair, it has order d, so [C] has order d as well. Thus [C] is in a Sylow d-subgroup of  $[\mathcal{C}_2]$ . The order of  $[\mathcal{C}_2]$  is that of  $\operatorname{Sp}(1,\mathbb{Z}_d) \ltimes \mathbb{Z}_d^2$ , which as shown in [24] is  $d^3(d^2-1)$ . Furthermore, the order of  $[\overline{\mathcal{D}}_2\langle X \rangle]$  is  $d^3$ , since  $[D_2]$  has order  $d^2$  and  $\langle X \rangle$  has order d. This implies that  $[\overline{\mathcal{D}}_2\langle X \rangle]$  is a Sylow d-subgroup of  $[\mathcal{C}_2]$ . All Sylow d-subgroups are conjugate, so this means C is Clifford-conjugate to an element of  $\overline{\mathcal{D}}_2\langle X \rangle$ .

### **Lemma 22.** If $DP \in \overline{\mathcal{D}}_k \mathcal{N}$ is of order d, then $DP \in \overline{\mathcal{D}}_k \langle X \rangle$ .

*Proof.* We first show that if  $P \in \mathcal{N}$  has order d, then  $P \in \langle X \rangle$ . Our discussion of the structure of  $\mathcal{N}$  at the beginning of Section 3.1 shows that  $\mathcal{N}$  has order d(d-1), while  $\langle X \rangle$  has order d. Thus  $\langle X \rangle$  is a Sylow d-subgroup of  $\mathcal{N}$ . As  $\langle X \rangle$  is also a normal subgroup of  $\mathcal{N}$ , it follows that  $\langle X \rangle$  is the only Sylow-d subgroup of  $\mathcal{N}$ . Hence, any order-d element of  $\mathcal{N}$  is in  $\langle X \rangle$ .

If  $DP \in \overline{\mathcal{D}}_k \mathcal{N}$  has order d, note that we may write  $(DP)^d = D'P^d$  for some diagonal D'. Thus in order for DP to have order d, the permutation matrix P must itself have order d. But then  $P \in \langle X \rangle$  and  $DP \in \overline{\mathcal{D}}_k \langle X \rangle$ .  $\Box$ 

**Lemma 23** (Main Lemma 1). Let  $k \ge 1$  and assume that  $C_k = SC_k$ : every  $k^{th}$ -level gate is semi-Clifford. Suppose  $G \in C_k$  is of order d and  $G^2 \in C_k$ . Then G is Clifford-conjugate to a gate in  $\overline{\mathcal{D}}_k(X)$ .

*Proof.* As every Pauli gate is in  $\overline{\mathcal{D}}_1$ , we can assume that  $k \geq 2$ .

Since G is semi-Clifford, it is Clifford-conjugate to DC for some  $D \in \mathcal{D}_k$  and  $C \in \mathcal{C}_2$ .

Without loss of generality, we can assume that  $C \notin \overline{\mathcal{D}}_2 \mathcal{N}$ . Otherwise, if  $C \in \overline{\mathcal{D}}_2 \mathcal{N}$ , by Lemma 22,  $C \in \overline{\mathcal{D}}_2 \langle X \rangle$ and  $DC \in \overline{\mathcal{D}}_k \langle X \rangle$ .

To finish the proof, it suffices to show that  $W = DXD^*$  is a Pauli gate. If this holds, since  $DZD^* = Z$  is a Pauli gate, D is a Clifford gate. Thus, DC is a Clifford gate of order d and Lemma 21 establishes our claim.

Since  $G^2 \in \mathcal{C}_k$ , we have that  $DCDC \in \mathcal{C}_k = \mathcal{SC}_k$ . As C is a Clifford gate, it follows that  $DCD \in \mathcal{SC}_k$  is semi-Clifford. From DCD, we obtain its (k-1)-closed conjugate pair

$$(U,V) = (DCDZD^*C^*D^*, DCDXD^*C^*D^*) = (DCZC^*D^*, DCDXD^*C^*D^*).$$
(46)

Since  $C \notin \overline{\mathcal{D}}_2 \mathcal{N}$ , by Lemma 19, we have that  $CZC^* = \omega^c Z^p X^q$ , for some  $p, q, c \in \mathbb{Z}_d$ , where  $q \neq 0$ . Hence,

$$U = \omega^c D Z^p X^q D^* = \omega^c Z^p (D X^q D^*), \tag{47}$$

which is a Pauli gate only if  $W = (DX^qD^*)^{q^{-1}}$  is a Pauli gate.

Assume for contradiction that U is not a Pauli gate. The computation above shows that  $U \in \overline{\mathcal{D}}_{k-1}\langle X \rangle$ , so the support of  $f_U$  is contained in a horizontal line  $L_U$  in the phase plane  $\mathbb{Z}_d^2$ . Since U is not a Pauli gate and is in a conjugate pair, Lemma 2 ensures that the Pauli support of  $f_U$  has size at least 2. As two points in the phase plane define a unique line, it follows that  $L_U$  is the unique line in the phase plane that contains the support of  $f_U$ .

Since (U, V) is a conjugate pair corresponding to the semi-Clifford gate DCD, Theorem 5 ensures that the Pauli supports of U and V are contained in two parallel lines. Hence, as  $L_U$  is horizontal and is the *unique* line containing the Pauli support of U, the Pauli support of V must be contained in a horizontal line. By Lemma 18, V must then itself be in  $\overline{\mathcal{D}}_{k-1}\langle X \rangle$ . That is,

$$DCDXD^*C^*D^* \in \overline{\mathcal{D}}_{k-1}\langle X \rangle.$$
(48)

As conjugating an element of  $\overline{\mathcal{D}}_{k-1}\langle X \rangle$  by  $D^*$  yields another element of  $\overline{\mathcal{D}}_{k-1}\langle X \rangle$ , it follows that  $CDXD^*C^* \in \overline{\mathcal{D}}_{k-1}\langle X \rangle$ . Since  $W = DXD^*$  is in  $\overline{\mathcal{D}}_k\langle X \rangle$ , the support of  $f_W$  is contained in a horizontal line  $L_W$  in the phase plane. The support of  $f_{CWC^*}$  is then contained in another line  $L'_W$  in the phase plane. Since  $C \notin \overline{\mathcal{D}}_2 \mathcal{N}$ , Lemma 19 implies that the symplectic transformation corresponding to C does not carry horizontal lines to horizontal lines, so  $L'_W$  is not horizontal. However,  $CWC^*$  is in  $\overline{\mathcal{D}}_{k-1}\langle X \rangle$ , so its support must also be contained in a horizontal line. Hence, the support of  $CWC^*$  is contained in the intersection of  $L'_W$  and a horizontal line, which ensures that the support of  $f_{CWC^*}$  consists of only one point. As  $CWC^*$  is conjugate to X, it has order d, so it then follows from Lemma 2 that  $CWC^*$  is Pauli. As C is a Clifford gate, both  $W = DXD^*$  and U are then also Pauli gates, which completes the proof.

This lemma makes an assumption that is equivalent to asserting that (k-1)-closed conjugate pairs are *jointly* Clifford-conjugate to a pair of almost diagonal gates. However, the conclusion proves something weaker than this about a k-closed conjugate pair (U, V): U and V are merely both *individually* Clifford-conjugate to almost diagonal gates. The following lemma resolves this issue and will allow us to inductively establish that  $C_k = SC_k$  for all k. Interestingly, it does not actually require the inductive hypothesis that the preceding lemma did. **Lemma 24** (Main Lemma 2). Let (U, V) be the conjugate pair of a gate  $G \in \mathcal{U}$  and suppose that both U and V are individually Clifford-conjugate to elements of  $\overline{\mathcal{D}}_k(X)$ . Then G is simplifiable.

*Proof.* Note that if U is a Pauli gate, the claim holds, as we may then conjugate V by a Clifford gate to map it into  $\overline{\mathcal{D}}_k\langle X \rangle$ , and this Clifford conjugation sends U to another Pauli gate, which is in  $\overline{\mathcal{D}}_k\langle X \rangle$ . By the same reasoning, the claim holds if V is a Pauli gate. We now assume that neither U nor V is a Pauli gate.

By conjugating by an appropriate Clifford gate, we may assume without loss of generality that  $U \in \overline{\mathcal{D}}_k \langle X \rangle$ , while  $V \in C\overline{\mathcal{D}}_k \langle X \rangle C^*$  for some Clifford gate C. Using our normal form for Clifford gates, either  $C \in \overline{\mathcal{D}}_2 \mathcal{N}$  or  $C = DPH^*E^c$  for some  $DP \in \overline{\mathcal{D}}_2 \mathcal{N}$ ,  $c \in \mathbb{Z}_d$ , and E as in Definition 17. Since  $\mathcal{D}_2 \mathcal{N}$  normalises  $\overline{\mathcal{D}}_k \langle X \rangle$ , if  $C \in \overline{\mathcal{D}}_2 \mathcal{N}$ then  $V \in \overline{\mathcal{D}}_k \langle X \rangle$  and we are done. We will show that the other case leads to a contradiction. If  $C = DPH^*E^c$ , then as conjugating by  $E^c$  sends  $\overline{\mathcal{D}}_k \langle X \rangle$  to itself, we have

$$V \in C\overline{\mathcal{D}}_k \langle X \rangle C^* = DPH^*\overline{\mathcal{D}}_k \langle X \rangle (DPH^*)^*.$$
<sup>(49)</sup>

Furthermore, as  $(DP)^*$  normalises  $\overline{\mathcal{D}}_k\langle X \rangle$ , we may conjugate both U and V by  $(DP)^*$  to create a conjugate pair with the first gate in  $\overline{\mathcal{D}}_k\langle X \rangle$  and the other in  $H^*\overline{\mathcal{D}}_k\langle X \rangle H$ . We now redefine (U, V) to be this conjugate pair.

Since  $U \in \overline{\mathcal{D}}_k \langle X \rangle$ , the support of  $f_U$  is contained in the horizontal line  $\{(p, q_0) \mid p \in \mathbb{Z}_d\}$  for some  $q_0 \in \mathbb{Z}_d$ . Similarly, as  $V \in H^*\overline{\mathcal{D}}_k \langle X \rangle H$  and conjugation by  $H^*$  maps horizontal lines in the phase plane to vertical lines, the support of  $f_V$  is contained in the vertical line  $\{(p_0, q) \mid q \in \mathbb{Z}_d\}$  for some  $p_0 \in \mathbb{Z}_d$ . Using Lemma 1, for any  $p, q \in \mathbb{Z}_d$  we have

$$f_{UV}(p,q) = \sum_{i,j \in \mathbb{Z}_d} \omega^{2^{-1}(qi-pj)} f_U(i,j) f_V(p-i,q-j).$$
(50)

Since  $f_U(i,j)$  vanishes except when  $j = q_0$  and  $f_V(p-i, q-j)$  vanishes except when  $p-i = p_0$ , this simplifies to

$$f_{UV}(p,q) = \omega^{2^{-1}(q(p-p_0)-pq_0)} f_U(p-p_0,q_0) f_V(p_0,q-q_0).$$
(51)

By the same reasoning, we compute that

$$f_{VU}(p,q) = \omega^{-2^{-1}(q(p-p_0)-pq_0)} f_U(p-p_0,q_0) f_V(p_0,q-q_0).$$
(52)

Then as  $UV = \omega VU$ , we have  $f_{UV} = \omega f_{VU}$ . The above equations thus ensure that whenever  $f_U(p - p_0, q_0)$  and  $f_V(p_0, q - q_0)$  are both nonzero,

$$\omega^{2^{-1}(q(p-p_0)-pq_0)} = \omega^{-2^{-1}(q(p-p_0)-pq_0)+1},\tag{53}$$

or equivalently,

$$pq - pq_0 - qp_0 - 1 = 0. (54)$$

As U is in a conjugate pair and is not Pauli, Lemma 2 ensures that the support of  $f_U$  has size at least 2. We may then choose  $p_1 \neq p_0$  such that  $f_U(p_1 - p_0, q_0) \neq 0$ . By the same reasoning, since V is not a Pauli gate, we may choose distinct  $q_1, q_2$  such that  $f_V(p_0, q_1 - q_0)$  and  $f_V(p_0, q_2 - q_0)$  are both nonzero. Then both  $(p, q) = (p_1, q_1)$ and  $(p, q) = (p_1, q_2)$  must be solutions to (54). However, for any fixed value of p other than  $p = p_0$ , (54) is a linear polynomial in q, which has exactly one solution. This gives the desired contradiction.

We may now prove the main theorem.

**Theorem 6** (Main Theorem). For all odd primes d, every gate at every level of the single-qudit Clifford hierarchy is semi-Clifford.

Proof. We claim that for all  $k \geq 1$ , every gate at level k is semi-Clifford. We proceed by induction on k. The base case of k = 1 holds, as all Pauli gates are semi-Clifford. Assume now that  $k \geq 2$  and that the claim holds at level k - 1. Let  $G \in \mathcal{C}_k$  and (U, V) be the (k - 1)-closed conjugate pair corresponding to G. As (U, V) is (k - 1)-closed,  $U^2$  and  $V^2$  are also in  $\mathcal{C}_{k-1}$ , so the inductive hypothesis combines with Lemma 23 to ensure that U and V are both individually Clifford-conjugate to elements of  $\overline{\mathcal{D}}_{k-1}\langle X \rangle$ . Lemma 24 then ensures that (U, V) is jointly Clifford-conjugate to a pair of gates in  $\overline{\mathcal{D}}_{k-1}\langle X \rangle$ . Thus G is simplifiable, and so semi-Clifford by Theorem 4. By induction, every gate in  $\mathcal{C}_k$  is semi-Clifford for  $k \geq 1$ .

### 4.5 A normal form for Clifford hierarchy gates

Our proof of the conjecture also has several nice consequences for the structure of the hierarchy. First, as the inverse of a semi-Clifford gate in  $C_k$  is also in  $C_k$ , we have the following corollary.

### **Corollary 1.** If $U \in C_k$ , then $U^* \in C_k$ .

Thus, while  $C_k$  is not a group for  $k \ge 3$ , it is at least closed under inversion for all k.

The fact that every hierarchy gate is semi-Clifford tells us a great deal about the structure of the hierarchy, as we now know that each hierarchy gate has the form  $C_1DC_2$ . However, the expression  $C_1DC_2$  is highly nonunique. It would be useful to have a unique way to write each gate in the hierarchy. In the theorem below, we extend our Clifford normal form from Lemma 9 to a normal form for all semi-Clifford gates, and hence, for all gates in the hierarchy by Theorem 6. For  $k \ge 2$ , we let  $\mathcal{D}_k/\mathcal{D}_2$  denote a fixed set of representatives for the quotient group  $\mathcal{D}_k/\mathcal{D}_2$ , and we may assume that I is one of these representatives. Explicit representatives are easily specified using the classification of diagonal hierarchy gates of Cui-Gottesman-Krishna [10].

**Theorem 7** (Normal Form for Semi-Clifford Gates). Let  $k \ge 2$ . Then any gate G in  $C_k$  may be written uniquely in one of the following forms:

$$G = \begin{cases} C & G \in \mathcal{C}_2 \\ MDC & G \notin \mathcal{C}_2 \end{cases}$$
(55)

Here,  $\mathcal{M}$  is the finite set of gates of Definition 17,  $D \in (\mathcal{D}_k/\mathcal{D}_2) \setminus \{\mathbb{I}\}$ , and  $C \in \mathcal{C}_2$ .

*Proof.* Let  $k \ge 2$  and  $G \in \mathcal{C}_k$ . By Theorem 6, G is semi-Clifford. Let (U, V) be the conjugate pair corresponding to G. If U and V both have Pauli supports of size 1, Lemma 2 implies that U and V are both Pauli, and so that G is a Clifford gate. In this case, we are done. Assume now that G is not a Clifford gate. Then at least one of U and V has a Pauli support of size greater than 1, and we assume without loss of generality that U does.

By Corollary 5, as G is semi-Clifford, the Pauli supports of U and V are contained in two parallel lines in  $\mathbb{Z}_d^2$ . Since U has a Pauli support of size greater than 1, there is then a unique line  $L_U$  containing  $\operatorname{supp}(f_U)$ . By Lemma 20, there is a unique  $M \in \mathcal{M}$  such that the symplectic transformation S corresponding to M carries horizontal lines to lines parallel to  $L_U$ . Hence, this is the unique M such that the Pauli supports of  $M^*UM$  and  $M^*VM$  are contained in two horizontal lines. By Lemma 19, this is the unique M such that  $M^*G = DC$  for some  $D \in \mathcal{D}_k$  and  $C \in \mathcal{C}_2$ .

Since  $G \notin C_2$ , we have  $D \notin D_2$ . We may then write D uniquely as  $D_1D_2$ , where  $D_1 \in (\mathcal{D}_k/\mathcal{D}_2) \setminus \{\mathbb{I}\}$  and  $D_2 \in \mathcal{D}_2$ . Then  $D_2C = C_2 \in C_2$  and  $DC = D_1C_2$ . Note that if we also have  $DC = D'_1C'_2$  for some  $D'_1 \in (\mathcal{D}_k/\mathcal{D}_2) \setminus \{\mathbb{I}\}$  and  $C'_2 \in C_2$ , then

$$(D_1')^* D_1 = C_2' C_2^* \in \mathcal{C}_2, \tag{56}$$

so  $(D'_1)^*D_1 \in \mathcal{D}_2$ . As  $\mathcal{D}_k/\mathcal{D}_2$  contains a unique representative from each coset of  $\mathcal{D}_2$ , it follows that  $D_1 = D'_1$ , and so that  $C_2 = C'_2$ . Thus  $G = MD_1C_2$  is in the claimed normal form, and each of M,  $D_1$ , and  $C_2$  are unique.  $\Box$ 

If we wished, we could use our Clifford normal form from Lemma 9 to express the Clifford gate C in our semi-Clifford normal form more explicitly. The normal form thus allows us to understand exactly which gates are at each level of the hierarchy. It also enables us to precisely count the number of gates in  $C_k$ , up to phase.

**Corollary 2.** The number of gates in  $C_k$ , up to phase, is exactly

$$|[\mathcal{C}_k]| = \begin{cases} d^2 & k = 1, \\ d^3(d^2 - 1)(d^{k-1} + d^{k-2} - d) & k \ge 2. \end{cases}$$
(57)

Proof. Up to phase, each Pauli gate is of the form W(p,q) for a unique  $(p,q) \in \mathbb{Z}_d^2$ , so  $[\mathcal{C}_1]$  has size  $d^2$ . For  $k \geq 2$ , Theorem 7 reduces the problem of determining the size of  $[\mathcal{C}_k]$  to counting the number of options for the gates M, D, and C used in the normal form. By definition, there are d + 1 possible choices for M. The characterisation of  $\mathcal{D}_k$ [10] shows that  $[\mathcal{D}_k]$  contains  $d^k$  elements. It follows that  $(\mathcal{D}_k/\mathcal{D}_2) \setminus \{\mathbb{I}\}$  contains  $d^{k-2} - 1$  gates, up to phase, and so there are  $d^{k-2} - 1$  options for D. The order of  $[\mathcal{C}_2]$  equals that of  $\operatorname{Sp}(1, \mathbb{Z}_d) \ltimes \mathbb{Z}_d^2$ , which from [24] is  $d^3(d^2 - 1)$ , giving that many choices for C. Since every hierarchy gate may be written uniquely up to phase as either C or MDC by Theorem 7, it follows that

$$|[\mathcal{C}_k]| = d^3(d^2 - 1) + (d+1)(d^{k-2} - 1)d^3(d^2 - 1)$$
(58)

$$= d^{3}(d^{2} - 1)(d^{k-1} + d^{k-2} - d).$$

## 5 Conclusions

This work significantly advances the program of classifying gates of the Clifford hierarchy and semi-Clifford gates. We have solved, in the practically-relevant case of one qubit/qudit, the problem of specifying precisely the gates of the Clifford hierarchy. Contributions to the classification problem have been made by many researchers, including founding figures of the field of quantum computing, over the past twenty-five years.

In future work, we will extend our classification of two-qubit/qudit hierarchy gates.

**Conjecture 1** (Classification of two-qubit/qudit hierarchy gates). Every hierarchy gate of two qudits is either a Clifford gate or can be uniquely expressed as  $M_1M_2DC$ . Here,  $M_1$  is  $H_1$  or  $H_1D_1H_1^*$  and  $M_2$  is  $H_2$  or  $H_1H_2D_2H_2^*H_1^*$  with  $H_i$  being the Hadamard gate on the i<sup>th</sup> qudit,  $D_1 = (E_1)^a$ ,  $D_2 = (CZ)^b(E_2)^c$  for  $a, b, c \in \mathbb{Z}_d$ . The gates D, C are as in Theorem 7. In the qubit case,  $D_1 = (S_1)^a$  and  $D_2 = (CZ)^b(S_2)^c$  for  $a, b, c \in \mathbb{Z}_2$ .

This conjectured classification can be extended to semi-Clifford gates of more than two qubits/qudits. We intend also to give a complete classification for hierarchy gates of any number of qubits/qudits.

Deeper understanding of the Clifford hierarchy and semi-Clifford gates will lead to progress in error correction and fault-tolerance and more efficient circuit and gate synthesis (e.g. [15, 20, 22, 33, 34]). It is also relevant to the theoretical study of quantum advantage. Hierarchy gates and their magic states are a key resource for powering quantum computation and hierarchy level provides a natural ranking or measure of their power/complexity which can be in turn correlated with advantage (e.g. [11, 13, 14]). For this reason, the Clifford hierarchy is relevant to many seemingly unconnected topics such as classical simulation [5] and learning circuits [2]. By clarifying the algebraic structure of magic states, we contribute to the understanding of the resources that power quantum computation.

We further bolster the viability of qudit-based fault-tolerant universal quantum computers by providing many more pathways towards efficient fault-tolerant qudit quantum computing. This is practically important as qudit magic state distillation has been proposed as a significantly more efficient alternative to the qubit case [6]. The advantages of qudit-based computation [31] has led to rapidly accelerating development by experimentalists [8, 9, 21, 23, 28, 29, 32].

## 6 Acknowledgments

ND acknowledges support from the Canada Research Chair program, NSERC Discovery Grant RGPIN-2022-03103, the NSERC-European Commission project FoQaCiA, and the Faculty of Science of Simon Fraser University. OL was supported by NSERC Undergraduate Student Research Awards (USRA).

## References

- [1] J. T. Anderson. On groups in the qubit Clifford hierarchy. Quantum, 8:1370, 2024. 1.1
- [2] A. Anshu and S. Arunachalam. A survey on the complexity of learning quantum states. Nature Reviews Physics, 6(1):59–69, 2024.
   5
- M. J. J. Barry. Large abelian subgroups of Chevalley groups. Journal of the Australian Mathematical Society, 27(1):59–87, 1979.
   3.2
- [4] S. Beigi and P. W. Shor. C<sub>3</sub>, semi-Clifford and generalized semi-Clifford operations. Quantum Information and Computation, 10(1):41-59, 2010. 1.1
- [5] K. Bu and D. E. Koh. Efficient classical simulation of Clifford circuits with nonstabilizer input states. *Physical Review Letters*, 123(17):170502, 2019. 5
- [6] E. T. Campbell, H. Anwar, and D. E. Browne. Magic-state distillation in all prime dimensions using quantum Reed-Muller codes. *Physical Review X*, 2:041021, 2012. 5
- [7] I. Chen and N. de Silva. Characterising semi-Clifford gates using algebraic sets. Communications in Mathematical Physics, 405(9):201, 2024. 1.1, 3.2
- [8] Y. Chi, J. Huang, Z. Zhang, J. Mao, Z. Zhou, X. Chen, C. Zhai, J. Bao, T. Dai, and H. Yuan. A programmable qudit-based quantum processor. *Nature Communications*, 13(1):1166, 2022. 5
- [9] M. Chizzini, L. Crippa, A. Chiesa, F. Tacchino, F. Petiziol, I. Tavernelli, P. Santini, and S. Carretta. Molecular nanomagnets with competing interactions as optimal units for qudit-based quantum computation. *Physical Review Research*, 4(4):043135, 2022. 5
- [10] S. X. Cui, D. Gottesman, and A. Krishna. Diagonal gates in the Clifford hierarchy. Physical Review A, 95(1):012329, 2017. 1.1, 2.7, 4.5, 4.5
- [11] M. de Oliveira, L. S. Barbosa, and E. F. Galvão. Quantum advantage in temporally flat measurement-based quantum computation. *Quantum*, 8:1312, 2024. 5

- [12] N. de Silva. Efficient quantum gate teleportation in higher dimensions. Proceedings of the Royal Society A, 2021. 1.1, 2, 2.4, 2.6, 2.7, 2.8, 3.2
- [13] M. Frembs, C. Okay, and H. Y. Chung. No quantum solutions to linear constraint systems in odd dimension from Pauli group and diagonal Cliffords. Quantum, 9:1583, 2025. 5
- [14] M. Frembs, S. Roberts, E. T. Campbell, and S. D. Bartlett. Hierarchies of resources for measurement-based quantum computation. New Journal of Physics, 25(1):013002, 2023. 5
- [15] A. Glaudell, N. J. Ross, J. van de Wetering, and L. Yeh. Qutrit metaplectic gates are a subset of Clifford+T. Proceedings TQC 2022., 2022. 5
- [16] D. Gottesman. Stabilizer Codes and Quantum Error Correction. PhD thesis, California Institute of Technology, Pasadena, CA, USA, 1997. 1
- [17] D. Gottesman and I. L. Chuang. Quantum teleportation is a universal computational primitive. Nature, 402(6760):390–393, 1999.
   1, 1.1, 2.6
- [18] D. Gross. Hudson's theorem for finite-dimensional quantum systems. Journal of Mathematical Physics, 47(12):122107, 2006. 2.5
- [19] Z. He, L. Robitaille, and X. Tan. Permutation gates in the third level of the Clifford hierarchy. arXiv preprint arXiv:2410.11818, 2024. 1.1
- [20] T. Jochym-O'Connor and T. J. Yoder. Four-dimensional toric code with non-Clifford transversal gates. Physical Review Research, 3(1):013118, 2021. 5
- [21] M. Karácsony, L. Oroszlány, and Z. Zimborás. Efficient qudit based scheme for photonic quantum computing. arXiv:2302.07357, 2023. 5
- [22] A. Krishna and J.-P. Tillich. Towards low overhead magic state distillation. Physical Review Letters, 123(7):070507, 2019. 5
- [23] P. J. Low, B. M. White, A. A. Cox, M. L. Day, and C. Senko. Practical trapped-ion protocols for universal qudit-based quantum computing. *Physical Review Research*, 2(3):033128, 2020. 5
- [24] M. Neuhauser. An explicit construction of the metaplectic representation over a finite field. Journal of Lie Theory, 12(1):15–30, 2002. 2.5, 3.1, 4.4, 4.5
- [25] T. Pllaha, N. Rengaswamy, O. Tirkkonen, and R. Calderbank. Un-Weyl-ing the Clifford hierarchy. Quantum, 4:370, 2020. 1.1, 2.3
- [26] T. Pllaha, K. Volanto, and O. Tirkkonen. Decomposition of Clifford gates. In 2021 IEEE Global Communications Conference, pages 01–06, 2021. 2.5, 1
- [27] N. Rengaswamy, R. Calderbank, and H. D. Pfister. Unifying the Clifford hierarchy via symmetric matrices over rings. *Physical Review A*, 100(2):022304, 2019. 1.1
- [28] M. Ringbauer, M. Meth, L. Postler, R. Stricker, R. Blatt, P. Schindler, and T. Monz. A universal qudit quantum processor with trapped ions. *Nature Physics*, 18(9):1053–1057, 2022. 5
- [29] L. M. Seifert, J. Chadwick, A. Litteken, F. T. Chong, and J. M. Baker. Time-efficient qudit gates through incremental pulse re-seeding. In 2022 IEEE International Conference on Quantum Computing and Engineering (QCE), pages 304–313. IEEE, 2022. 5
- [30] P. W. Shor. Fault-tolerant quantum computation. In Proceedings of the 37th Conference on Foundations of Computer Science, pages 56–65. IEEE, 1996. 1
- [31] H. S. Tonchev and N. V. Vitanov. Quantum phase estimation and quantum counting with qudits. *Physical Review A*, 94(4):042307, 2016.
- [32] S. Wang, Z.-Q. Yin, H. Chau, W. Chen, C. Wang, G.-C. Guo, and Z.-F. Han. Proof-of-principle experimental realization of a qubit-like qudit-based quantum key distribution scheme. *Quantum Science and Technology*, 3(2):025006, 2018. 5
- [33] M. A. Webster, A. O. Quintavalle, and S. D. Bartlett. Transversal diagonal logical operators for stabiliser codes. New Journal of Physics, 25(10):103018, 2023. 5
- [34] L. Yeh. Scaling w state circuits in the qudit Clifford hierarchy. In Companion Proceedings of the 7th International Conference on the Art, Science, and Engineering of Programming, pages 90–100, 2023. 5
- [35] B. Zeng, X. Chen, and I. L. Chuang. Semi-Clifford operations, structure of  $C_k$  hierarchy, and gate complexity for fault-tolerant quantum computation. *Physical Review A*, 77(4):042313, 2008. 1, 1.1, 2.6, 2.8, 3.1, 4
- [36] X. Zhou, D. W. Leung, and I. L. Chuang. Methodology for quantum logic gate construction. Physical Review A, 62(5):052316, 2000. 1, 1.1, 2.8