ISOMETRIES BETWEEN SPACES OF METRICS

KATSUHISA KOSHINO

ABSTRACT. Given a metrizable space Z, denote by PM(Z) the space of continuous bounded pseudometrics on Z, and denote by AM(Z) the one of continuous bounded admissible metrics on Z, the both of which are equipped with the sup-norm $\|\cdot\|$. Let Pc(Z) be the subspace of AM(Z) satisfying the following:

• for every $d \in Pc(Z)$, there exists a compact subset $K \subset Z$ such that if d(x, y) = ||d||, then $x, y \in K$.

Moreover, set

 $\operatorname{Pp}(Z) = \{ d \in \operatorname{AM}(Z) \mid \text{ there only exists } \{z, w\} \subset Z \text{ such that } d(z, w) = \|d\| \},\$

and let M(Z) be Pc(Z) or Pp(Z). In this paper, we shall prove the Banach-Stone type theorem on spaces of metrics, that is, for metrizable spaces X and Y, the following are equivalent:

(1) X and Y are homeomorphic;

- (2) there exists a surjective isometry $T : PM(X) \to PM(Y)$ with T(M(X)) = M(Y);
- (3) there exists a surjective isometry $T : AM(X) \to AM(Y)$ with T(M(X)) = M(Y);
- (4) there exists a surjective isometry $T : M(X) \to M(Y)$.

Then for each surjective isometry $T : PM(X) \to PM(Y)$ with T(M(X)) = M(Y), there is a homeomorphism $\phi : Y \to X$ such that for any $d \in PM(X)$ and for any $x, y \in Y$, $T(d)(x, y) = d(\phi(x), \phi(y))$. Except for the case where the cardinality of X or Y is equal to 2, the homeomorphism ϕ can be chosen uniquely.

1. INTRODUCTION

Isometries between function spaces have been studied in functional analysis. The Banach-Stone theorem [1, 14] is one of the most important results among those research, and its developments have been obtained until now, refer to [3] as a historical note. Throughout the paper, an isometry means a surjective isometry. For a metrizable space Z, let C(Z) be the space of continuous bounded real-valued functions on Z with the sup-norm $\|\cdot\|$: for any $f \in C(Z)$, $\|f\| = \sup\{|f(z)| \mid z \in Z\}$. Denote the positive cone by $C_+(Z) \subset C(Z)$. Recently, L. Sun, Y. Sun and D. Dai [15] showed the Banach-Stone type theorem on positive cones of continuous function spaces as follows:

Theorem 1.1. Suppose that X and Y are compact metrizable spaces. Then X and Y are homeomorphic if and only if $C_+(X)$ and $C_+(Y)$ are isometric.

D. Hirota, I. Matsuzaki and T. Miura [5] generalized the above theorem in the non-compact case. In this paper, we shall establish the Banach-Stone type theorem on spaces of metrics. Let $PM(Z) \subset C_+(Z^2)$ be the subspace consisting of continuous bounded pseudometrics on Z, and let $AM(Z) \subset PM(Z)$ be the subspace consisting of continuous bounded admissible metrics. As is easily observed, PM(X) and PM(Y) (respectively, AM(X) and AM(Y)) are isometric if metrizable spaces X and Y are homeomorphic. When X and Y are compact, M.E. Shanks [13] showed the converse of it and established the following:

Theorem 1.2. Let X and Y be compact metrizable spaces. The following are equivalent:

²⁰²⁰ Mathematics Subject Classification. Primary 46B04; Secondary 46E15, 54C35, 54E35.

Key words and phrases. isometric, pseudometric, admissible metric, sup-metric, the Banach-Stone theorem.

(1) X and Y are homeomorphic;

(2) PM(X) and PM(Y) are isometric;

(3) AM(X) and AM(Y) are isometric.

Shanks focused on certain lattice structure on equivalent classes of PM(X), which determined the topology of X, and the method was different from those of S. Banach and M.H. Stone. However, Shanks did not give descriptions of isometries by using homeomorphisms like the canonical formula (*) as in Main Theorem, which appeared in the Banach-Stone theorem. Set Pc(Z) be the subspace of AM(Z) that satisfies the following condition:

• for every $d \in Pc(Z)$, there is a compact set K in Z such that if d(x, y) = ||d||, then $x, y \in K$. Notice that AM(Z) = Pc(Z) when Z is compact. Moreover, put

 $\operatorname{Pp}(Z) = \{ d \in \operatorname{AM}(Z) \mid \text{ there only exists } \{z, w\} \subset Z \text{ such that } d(z, w) = ||d|| \},\$

and let M(Z) be Pc(Z) or Pp(Z). We shall generalize this result and determine isometries by homeomorphisms as follows:

Main Theorem. Let X and Y be metrizable spaces. The following are equivalent:

- (1) X and Y are homeomorphic;
- (2) there exists an isometry $T : PM(X) \to PM(Y)$ with T(M(X)) = M(Y);
- (3) there exists an isometry $T : AM(X) \to AM(Y)$ with T(M(X)) = M(Y);
- (4) there exists an isometry $T : M(X) \to M(Y)$.

In this case, for each isometry $T : PM(X) \to PM(Y)$ with T(M(X)) = M(Y), there exists a homeomorphism $\phi : Y \to X$ such that for any $d \in PM(X)$ and for any $x, y \in Y$,

$$T(d)(x,y) = d(\phi(x),\phi(y)).$$
 (*)

Except for the case where the cardinality of X or Y is equal to 2, the homeomorphism ϕ can be chosen uniquely.

2. Spaces of metrics

In this section, we shall review the study on spaces of metrics. Recently, Y. Ishiki [6, 7, 8] have researched topologies of spaces of metrics. The author [11, 12] investigated their Borel hierarchy, complete metrizability and topological types, and proved the following:

Theorem 2.1. Let κ be a cardinal and $\ell_2(\kappa)$ be the Hilbert space of density κ . When a metrizable space Z is of density κ , the space PM(Z) is homeomorphic to

- (i) $[0,1)^{\kappa(\kappa-1)/2}$ if Z is finite;
- (ii) $\ell_2(2^{<\kappa})$ if Z is infinite and generalized compact;

(iii) $\ell_2(2^{\kappa})$ if Z is not generalized compact.

Additionally, when Z is infinite and σ -compact, the subspace AM(Z) is homeomorphic to

- (i) $\ell_2(\aleph_0)$ if Z is compact;
- (ii) $\ell_2(2^{\aleph_0})$ if Z is not compact.

This means that for metrizable spaces X and Y, even if PM(X) and PM(Y) (respectively, AM(X) and AM(Y)) are homeomorphic, X and Y are not necessarily homeomorphic. On metric structures of spaces of metrics, Y. Ishiki and the author [10] studied their isometric universality.

As a basic property on metrics, we have the following, see [10, Lemma 2.1].

Lemma 2.2. For a metrizable space Z, for every $d \in PM(Z)$ and every $\rho \in AM(Z)$, their sum $d + \rho \in AM(Z)$.

F. Hausdorff [4] showed the metric extension theorem, which states that for every metrizable space Z and its closed subset $A \subset Z$, any $d \in AM(X)$ can be extended over X. We may obtain the pseudometric version of it, that is preserving their norms, as follows, refer to [9] for example.

Theorem 2.3. Suppose that Z is a metrizable space and $A \subset Z$ is a closed subset. For each $d \in PM(A)$, there is $d \in PM(Z)$ such that $d|_{A^2} = d$ and ||d|| = ||d||.

This result will play a key role in the present paper instead of Urysohn's lemma, which have been frequently used in proving the various Banach-Stone type theorems.

For a metrizable space Z with a pseudometric $d \in PM(Z)$, denote the open ball centered at $z \in Z$ of radius r > 0 by

$$B_d(z, r) = \{ w \in Z \mid d(z, w) < r \}$$

and the closed ball by

$$\overline{B}_d(z,r) = \{ w \in Z \mid d(z,w) \le r \}.$$

Note that since d is continuous, $B_d(z,r)$ is open and $\overline{B}_d(z,r)$ is closed in Z. Recall that the supmetric induced by $\|\cdot\|$ is complete on $C(Z^2)$ and PM(Z), which is closed in $C(Z^2)$. Since AM(Z)is dense in PM(Z), see [11, Proposition 5], an isometry $T: AM(X) \to AM(Y)$ can be extended to an isometry $T: \mathrm{PM}(X) \to \mathrm{PM}(Y)$. Furthermore, we will prove that $\mathrm{Pc}(Z)$ is also a dense subset of PM(Z).

Proposition 2.4. For every metrizable space Z, the subset Pp(Z) is dense in PM(Z), and hence so is Pc(Z).

Proof. The cases where $Z = \emptyset$ and where Z is a singleton are trivial, so suppose that Z is nondegenerate. Fix any $d \in PM(Z)$ and any $\epsilon > 0$. Since AM(Z) is dense in PM(X), we may assume that $d \in AM(Z)$. Note that d is bounded, so choose distinct points $x, y \in Z$ so that $d(x,y) \geq ||d|| - \epsilon$. Let a = d(x,y), where we may also assume that $a \geq \epsilon$ by replacing ϵ with a sufficient small positive number. Applying Theorem 2.3, we can find a pseudometric $\rho_n \in PM(Z)$ for every natural number $n \ge 1$ such that

- (i) $\rho(z, w) = 4\epsilon$ if $z \in \overline{B}_d(x, \epsilon/2^{n+1})$ and $w \in \overline{B}_d(y, \epsilon/2^{n+1})$;
- (ii) $\rho(z,w) = 2\epsilon$ if $z \in Z \setminus (B_d(x,\epsilon/2^n) \cup B_d(y,\epsilon/2^n))$ and $w \in \overline{B}_d(x,\epsilon/2^{n+1}) \cup \overline{B}_d(y,\epsilon/2^{n+1})$; (iii) $\rho(z,w) = 0$ if $z,w \in Z \setminus (B_d(x,\epsilon/2^n) \cup B_d(y,\epsilon/2^n))$, if $z,w \in \overline{B}_d(x,\epsilon/2^{n+1})$, or if $z,w \in C$ $\overline{B}_d(y,\epsilon/2^{n+1});$
- (iv) $\rho(z, w) \leq 4\epsilon$ if otherwise.

Define $\rho = \sum_{n=1}^{\infty} \rho_n/2^n$, so $\rho \in PM(Z)$ because PM(Z) is complete. Observe that

$$||(d+\rho) - d|| = ||\rho|| \le \sum_{n=1}^{\infty} ||\rho_n|| \le \sum_{n=1}^{\infty} 4\epsilon/2^n = 4\epsilon.$$

It is only needed to verify that $d + \rho \in Pp(Z)$. Remark that $d + \rho \in AM(Z)$ by Lemma 2.2. Define the closed subsets

$$Z_1 = Z \setminus (B_d(x, \epsilon/2) \cup B_d(y, \epsilon/2)),$$

and for every $n \geq 2$,

$$Z_n = (\overline{B}_d(x, \epsilon/2^{n-1}) \cup \overline{B}_d(y, \epsilon/2^{n-1})) \setminus (B_d(x, \epsilon/2^n) \cup B_d(y, \epsilon/2^n)).$$

(1) When z = x or z = y, and $w \in Z_n$,

$$d(z,w) + \rho(z,w) = d(z,w) + \sum_{n=1}^{\infty} \rho_n(z,w)/2^n \le a + \epsilon/2^{n-1} + \sum_{i=1}^{n-1} 4\epsilon/2^i + \sum_{i=n}^{\infty} 2\epsilon/2^i \le a + 4\epsilon/2^{n+1} + \sum_{i=1}^{n-1} 4\epsilon/2^i + \sum_{i=n+1}^{\infty} 4\epsilon/2^i < a + \sum_{i=1}^{\infty} 4\epsilon/2^i = a + 4\epsilon.$$

(2) When $z, w \in Z_n$,

$$d(z,w) + \rho(z,w) = d(z,w) + \sum_{n=1}^{\infty} \rho_n(z,w)/2^n \le a + \epsilon/2^{n-2} + \sum_{i=1}^{n-1} 4\epsilon/2^i$$
$$\le a + 4\epsilon/2^n + \sum_{i=1}^{n-1} 4\epsilon/2^i < a + \sum_{i=1}^{\infty} 4\epsilon/2^i = a + 4\epsilon.$$

(3) When $z \in Z_n$ and $w \in Z_{n+1}$,

$$\begin{aligned} d(z,w) + \rho(z,w) &= d(z,w) + \sum_{n=1}^{\infty} \rho_n(z,w)/2^n \le a + \epsilon/2^{n-1} + \epsilon/2^n + \sum_{i=1}^n 4\epsilon/2^i \\ &\le a + 4\epsilon/2^{n+1} + 4\epsilon/2^{n+2} + \sum_{i=1}^n 4\epsilon/2^i < a + \sum_{i=1}^{\infty} 4\epsilon/2^i = a + 4\epsilon. \end{aligned}$$

(4) When $z \in Z_n$ and $w \in Z_m$, $m \ge n+2$,

$$\begin{split} d(z,w) + \rho(z,w) &= d(z,w) + \sum_{n=1}^{\infty} \rho_n(z,w)/2^n \\ &\leq a + \epsilon/2^{n-1} + \epsilon/2^{m-1} + \sum_{i=1}^{n-1} 4\epsilon/2^i + \sum_{i=n}^{m-2} 2\epsilon/2^i + 4\epsilon/2^{m-1} \\ &\leq a + 4\epsilon/2^{n+1} + 4\epsilon/2^{m+1} + \sum_{i=1}^{n-1} 4\epsilon/2^i + \sum_{i=n+1}^{m-1} 4\epsilon/2^i + 4\epsilon/2^{m-1} \\ &= a + 4\epsilon/2^n - 4\epsilon/2^{n+1} + 4\epsilon/2^{m+1} + \sum_{i=1}^{n-1} 4\epsilon/2^i + \sum_{i=n+1}^{m-1} 4\epsilon/2^i + 4\epsilon/2^{m-1} \\ &< a + \sum_{i=1}^{\infty} 4\epsilon/2^i = a + 4\epsilon. \end{split}$$

To sum up, we have that $d(z, w) + \rho(z, w) < a + 4\epsilon$ for all pairs $(z, w) \in \mathbb{Z}^2 \setminus \{(x, y), (y, x)\}$. By the definition of ρ ,

$$d(x,y) + \rho(x,y) = d(x,y) + \sum_{n=1}^{\infty} \rho_n(x,y)/2^n = a + \sum_{n=1}^{\infty} 4\epsilon/2^n = a + 4\epsilon.$$

The proof is completed. \Box

3. The peaking function argument

We shall prove Main Theorem by using the peaking function argument, which is based on [5], and traces its history back to Stone's method, that is, our strategy is different from Shanks' one. From now on, let X and Y be non-degenerate metrizable spaces and $T : PM(X) \to PM(Y)$ be an isometry. Assume that **0** is the zero function, that is a pseudometric. By the same argument as Lemmas 2.3 and 2.4 of [5], we have the following:

Lemma 3.1. For every $d \in PM(X)$, if

$$\max\{\|T(\mathbf{0})\|, \|T^{-1}(\mathbf{0})\|\} < \|d\|,\$$

then ||T(d)|| = ||d||. Similarly, for every $\rho \in PM(Y)$, if $\max\{||T(\mathbf{0})||, ||T^{-1}(\mathbf{0})||\} < ||\rho||,$

then $||T^{-1}(\rho)|| = ||\rho||.$

It follows from the above lemma that the isometry T is norm-preserving.

Proposition 3.2. The equalities $T(\mathbf{0}) = \mathbf{0}$ and $T^{-1}(\mathbf{0}) = \mathbf{0}$ hold. Hence ||T(d)|| = ||d|| for any $d \in PM(X)$ and $||T^{-1}(\rho)|| = ||\rho||$ for any $\rho \in PM(Y)$.

Proof. Assume that $T(\mathbf{0}) \neq \mathbf{0}$, so there exist points $x, y \in Y$ such that $T(\mathbf{0})(x, y) > 0$. Then

$$\max\{\|T(\mathbf{0})\|, \|T^{-1}(\mathbf{0})\|\} < 2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\| = \frac{2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\|}{\|T(\mathbf{0})\|} \|T(\mathbf{0})\|$$
$$= \left\|\frac{2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\|}{\|T(\mathbf{0})\|} T(\mathbf{0})\right\|.$$

Due to Lemma 3.1,

$$\left\|T^{-1}\left(\frac{2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\|}{\|T(\mathbf{0})\|}T(\mathbf{0})\right)\right\| = \left\|\frac{2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\|}{\|T(\mathbf{0})\|}T(\mathbf{0})\right\|.$$

Since T is an isometry,

$$2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\| > 2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\| - \|T(\mathbf{0})\| = \left(\frac{2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\|}{\|T(\mathbf{0})\|} - 1\right)\|T(\mathbf{0})\|$$

$$= \left\| \left(\frac{2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\|}{\|T(\mathbf{0})\|} T(\mathbf{0}) - T(\mathbf{0})\right)\right\|$$

$$= \left\| T^{-1} \left(\frac{2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\|}{\|T(\mathbf{0})\|} T(\mathbf{0})\right) - T^{-1}(T(\mathbf{0}))\right\|$$

$$= \left\| T^{-1} \left(\frac{2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\|}{\|T(\mathbf{0})\|} T(\mathbf{0})\right) - \mathbf{0} \right\|$$

$$= \left\| T^{-1} \left(\frac{2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\|}{\|T(\mathbf{0})\|} T(\mathbf{0})\right) - \mathbf{0} \right\|$$

$$= \left\| T^{-1} \left(\frac{2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\|}{\|T(\mathbf{0})\|} T(\mathbf{0})\right) \right\| = \left\| \frac{2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\|}{\|T(\mathbf{0})\|} T(\mathbf{0}) \right\|$$

$$= 2\|T(\mathbf{0})\| + \|T^{-1}(\mathbf{0})\|.$$

This is a contradiction. We conclude that $T(\mathbf{0}) = \mathbf{0}$. Moreover, for each $d \in PM(X)$,

$$\|T(d)\| = \|T(d) - \mathbf{0}\| = \|T(d) - T(\mathbf{0})\| = \|d - \mathbf{0}\| = \|d\|.$$

Similarly, $T^{-1}(\mathbf{0}) = \mathbf{0}$ and $\|T^{-1}(\rho)\| = \|\rho\|$ for any $\rho \in \text{PM}(Y)$. \Box

For a metrizable space Z, let $\operatorname{Fin}_2(Z)$ be the hyperspace consisting of singletons and doubletons of Z endowed with the Vietoris topology, and let

$$\mathcal{D}(Z) = \{\{x, y\} \in \operatorname{Fin}_2(Z) \mid x \neq y\}$$

For each $\{x, y\} \in \operatorname{Fin}_2(Z)$, put

$$\mathcal{P}(Z, \{x, y\}) = \{ d \in \text{PM}(Z) \mid d(x, y) = \|d\| \}$$

Given a pseudometric $d \in PM(Z)$, we define

$$\mathcal{F}(Z,d) = \{\{x,y\} \in \operatorname{Fin}_2(Z) \mid d(x,y) = \|d\|\}.$$

Lemma 3.3. Fix any $\{x, y\} \in \operatorname{Fin}_2(X)$ and any $d_i \in \mathcal{P}(X, \{x, y\}), 1 \leq i \leq n$. Then the sum $d = \sum_{i=1}^n d_i \in \mathcal{P}(X, \{x, y\})$. Furthermore, if $T(\operatorname{Pc}(X)) \subset \operatorname{Pc}(Y)$ and each $d_i \in \operatorname{Pc}(X)$, there exists $\{z, w\} \in \operatorname{Fin}_2(Y)$ such that T(d)(z, w) = d(x, y).

Proof. Observe that

$$d(x,y) \le ||d|| = \left\|\sum_{i=1}^{n} d_i\right\| \le \sum_{i=1}^{n} ||d_i|| = \sum_{i=1}^{n} d_i(x,y) = d(x,y).$$

so d(x, y) = ||d||. Therefore $d \in \mathcal{P}(X, \{x, y\})$. To show that the latter part, take a compact subset $K_i \subset X, 1 \leq i \leq n$, such that for any $u, v \in X$ with $d_i(u, v) = ||d_i||, u, v \in K_i$. If d(u, v) = ||d||, then

$$\sum_{i=1}^{n} d_i(u, v) = d(u, v) = ||d|| = d(x, y) = \sum_{i=1}^{n} d_i(x, y)$$

Since for each $i \in \{1, \ldots, n\}$, $d_i(u, v) \leq ||d_i|| = d_i(x, y)$, we can get that $d_i(u, v) = d_i(x, y) = ||d_i||$. Hence the points u and v are contained in any K_i , which implies that $d \in Pc(X)$. By the assumption of T, $T(d) \in Pc(Y)$, that is, there is a compact set $L \subset Y$ such that for any $z, w \in Y$, if T(d)(z, w) = ||T(d)||, then $z, w \in L$. Since L is compact and T(d) is continuous, there is $\{z, w\} \in \mathcal{F}(Y, T(d))$. Then according to Proposition 3.2,

$$T(d)(z,w) = ||T(d)|| = ||d|| = d(x,y).$$

The proof is finished. \Box

Moreover, we have the following:

Lemma 3.4. Suppose that $T(\operatorname{Pc}(X)) \subset \operatorname{Pc}(Y)$. For any $\{x, y\} \in \operatorname{Fin}_2(X)$ and any $d_i \in \mathcal{P}(X, \{x, y\}) \cap \operatorname{Pc}(X)$, $1 \leq i \leq n$, the intersection $\bigcap_{i=1}^n \mathcal{F}(Y, T(d_i)) \neq \emptyset$.

Proof. Let $d = \sum_{i=1}^{n} d_i$ and take a doubleton $\{z, w\} \in \operatorname{Fin}_2(Y)$ such that T(d)(z, w) = d(x, y) as in Lemma 3.3. For each $i \in \{1, \ldots, n\}$, set $\rho_i = d - d_i$, so $\rho_i \in \mathcal{P}(X, \{x, y\})$ and $\rho_i(x, y) = \|\rho_i\|$ by Lemma 3.3. Since T is isometric, we have that

$$T(d)(z,w) - T(d_i)(z,w) \le ||T(d) - T(d_i)|| = ||d - d_i|| = ||\rho_i|| = \rho_i(x,y),$$

and that according to Proposition 3.2,

 $\|T(d_i)\| = \|d_i\| = d_i(x, y) = d(x, y) - \rho_i(x, y) = T(d)(z, w) - \rho_i(x, y) \leq T(d_i)(z, w) \leq \|T(d_i)\|.$ Thus $T(d_i)(z, w) = \|T(d_i)\|$, which implies that $\{z, w\} \in \mathcal{F}(Y, T(d_i))$. Consequently, the intersection $\bigcap_{i=1}^n \mathcal{F}(Y, T(d_i))$ is not empty. \Box

Due to the similar method to Proposition 2.4, we can prove the following:

Lemma 3.5. Let Z be a non-degenerate metrizable space. For each $d \in PM(Z)$ and each $\{x, y\} \in D(Z)$, there is $\rho \in \mathcal{P}(Z, \{x, y\})$ such that $d + \rho \in \mathcal{P}(Z, \{x, y\}) \cap Pp(Z)$.

Proof. Adding an admissible metric in $\mathcal{P}(X, \{x, y\})$, we may assume that $d \in AM(Z)$ according to Lemma 2.2. Let a = d(x, y) > 0 and

$$b = \min \Big\{ \max_{z \in \mathbb{Z}} d(x, z), \max_{z \in \mathbb{Z}} d(y, z) \Big\}.$$

Remark that $a \leq b$. Using Theorem 2.3, we can obtain $\rho_n \in PM(Z)$ for each natural number $n \geq 1$ such that

- (i) $\rho(z, w) = 4b$ if $z \in \overline{B}_d(x, a/2^{n+1})$ and $w \in \overline{B}_d(y, a/2^{n+1})$;
- (ii) $\rho(z,w) = 2b$ if $z \in Z \setminus (B_d(x, a/2^n) \cup B_d(y, a/2^n))$ and $w \in \overline{B}_d(x, a/2^{n+1}) \cup \overline{B}_d(y, a/2^{n+1});$ (iii) $\rho(z,w) = 0$ if $z, w \in Z \setminus (B_d(x, a/2^n) \cup B_d(y, a/2^n)),$ if $z, w \in \overline{B}_d(x, a/2^{n+1}),$ or if $z, w \in \overline{B}_d(y, a/2^{n+1});$
- (iv) $\rho(z, w) \leq 4b$ if otherwise.

By the same argument as Proposition 2.4, $\rho = \sum_{n=1}^{\infty} \rho_n / 2^n$ is the desired pseudometric. Indeed, $\rho(x,y) = 4b = \|\rho\|$, and hence $\rho \in \mathcal{P}(Z, \{x,y\})$. Moreover, for every pair $(z,w) \in Z^2 \setminus \{(x,y), (y,x)\},$

$$d(x,y) + \rho(x,y) = a + 4b = ||d + \rho|| > d(z,w) + \rho(z,w),$$

so $d + \rho \in \mathcal{P}(Z, \{x, y\}) \cap \operatorname{Pp}(Z)$. We complete the proof. \Box

Using the finite intersection property in compact spaces, see [2, Theorem 3.1.1], we can obtain the following lemma.

Lemma 3.6. Assume that $T(Pc(X)) \subset Pc(Y)$. For every doubleton $\{x, y\} \in D(X)$,

$$\bigcap_{d \in \mathcal{P}(X, \{x, y\}) \cap \operatorname{Pc}(X)} \mathcal{F}(Y, T(d)) \neq \emptyset$$

Proof. Remark that $\mathcal{P}(X, \{x, y\}) \cap \mathrm{Pc}(X) \neq \emptyset$ by virtue of Lemma 3.5. Fix any $d_0 \in \mathcal{P}(X, \{x, y\}) \cap \mathrm{Pc}(X)$, so $T(d_0) \in \mathrm{Pc}(Y)$ and there exists a compact subset L of Y such that for any $z, w \in Y$ with $T(d_0)(z, w) = ||T(d_0)||, z, w \in L$. Since L is compact and

$$i: L^2 \ni (z, w) \mapsto \{z, w\} \in \operatorname{Fin}_2(L)$$

is surjective and continuous due to [16, Lemma 5.3.4], $\operatorname{Fin}_2(L)$ is also compact. Observe that $\mathcal{F}(Y, T(d))$ is closed in $\operatorname{Fin}_2(Y)$ for every $d \in \mathcal{P}(X, \{x, y\})$. Indeed, fix any $\{z, w\} \in \operatorname{Fin}_2(Y) \setminus \mathcal{F}(Y, T(d))$. Then T(d)(z, w) < ||T(d)||. Since T(d) is continuous, we can find open neighborhoods U of z and V of w such that if $u \in U$ and $v \in V$, then T(d)(u, v) < ||T(d)||. Remark that the subset

$$\mathcal{U} = \{\{u, v\} \in \operatorname{Fin}_2(Y) \mid \{u, v\} \cap U \neq \emptyset, \{u, v\} \cap V \neq \emptyset, \text{ and } \{u, v\} \subset U \cup V\}$$

is an open neighborhood of $\{z, w\}$ in Fin₂(Y). If $\{u, v\} \in \mathcal{U}$, then we may assume that $u \in U$ and $v \in V$, and hence T(d)(u, v) < ||T(d)||. Therefore $\{u, v\} \in \text{Fin}_2(Y) \setminus \mathcal{F}(Y, T(d))$, which means that $\mathcal{F}(Y, T(d))$ is closed in Fin₂(Y). Thus the set $\mathcal{F}(Y, T(d_0)) \subset \text{Fin}_2(L)$ is compact. According to Lemma 3.4, the family

$$\{\mathcal{F}(Y,T(d)) \cap \mathcal{F}(Y,T(d_0)) \mid d \in \mathcal{P}(X,\{x,y\})\}$$

has the finite intersection property, and hence

$$\bigcap_{d \in \mathcal{P}(X, \{x,y\}) \cap \operatorname{Pc}(X)} \mathcal{F}(Y, T(d)) = \bigcap_{d \in \mathcal{P}(X, \{x,y\}) \cap \operatorname{Pc}(X)} \left(\mathcal{F}(Y, T(d)) \cap \mathcal{F}(Y, T(d_0)) \right) \neq \emptyset.$$

We finish the proof. \Box

We have certain uniqueness of peaks of metrics as follows:

Lemma 3.7. Suppose that Z is a non-degenerate metrizable space. For all doubletons $\{x, y\}, \{x', y'\} \in D(Z)$, if

$$\mathcal{P}(Z, \{x, y\}) \cap \operatorname{Pp}(Z) \subset \mathcal{P}(Z, \{x', y'\}),$$

then $\{x, y\} = \{x', y'\}.$

Proof. Suppose not, so we may assume that $x \notin \{x', y'\}$ or $x' \notin \{x, y\}$. By Theorem 2.3, we can obtain $d \in PM(X)$ such that d(x, y) = 1, d(x', y') = 0, and ||d|| = 1. Moreover, there exists an admissible metric $\rho \in PM(X)$ such that $\rho(z, w) < \rho(x, y) = ||\rho||$ for any $(z, w) \in Z^2 \setminus \{(x, y), (y, x)\}$ by the same argument as Lemma 3.5. Then due to Lemma 2.2,

$$d + \rho \in (\mathcal{P}(Z, \{x, y\}) \cap \operatorname{Pp}(Z)) \setminus \mathcal{P}(Z, \{x', y'\}),$$

which is a contradiction. Hence $\{x, y\} = \{x', y'\}$. \Box

Now the following corresponding between D(X) and D(Y) will be given.

Proposition 3.8. Suppose that T(Pc(X)) = Pc(Y). There exists a bijection $\Phi : D(Y) \to D(X)$ such that

$$T(\mathcal{P}(X, \{x, y\}) \cap \operatorname{Pc}(X)) = \mathcal{P}(Y, \Phi^{-1}(\{x, y\})) \cap \operatorname{Pc}(Y)$$

for every $\{x, y\} \in D(X)$ and

$$T^{-1}(\mathcal{P}(Y, \{z, w\}) \cap \operatorname{Pc}(Y)) = \mathcal{P}(X, \Phi(\{z, w\})) \cap \operatorname{Pc}(X)$$

for every $\{z, w\} \in D(Y)$.

Proof. For each $\{z, w\} \in D(Y)$, by virtue of Lemma 3.6, we can choose $\{x, y\} \in Fin_2(X)$ so that

 $\{x,y\} \in \bigcap_{\rho \in \mathcal{P}(Y,\{z,w\}) \cap \mathrm{Pc}(Y)} \mathcal{F}(X,T^{-1}(\rho)).$

Note that $T^{-1}(\rho)(x,y) = ||T^{-1}(\rho)||$ for all $\rho \in \mathcal{P}(Y, \{z,w\}) \cap \operatorname{Pc}(Y)$, so $T^{-1}(\mathcal{P}(Y, \{z,w\}) \cap \operatorname{Pc}(Y)) \subset \mathcal{P}(X, \{x,y\}) \cap \operatorname{Pc}(X).$

Furthermore, due to Proposition 3.2,

$$T^{-1}(\rho)(x,y) = ||T^{-1}(\rho)|| = ||\rho|| > 0,$$

and hence $x \neq y$. Similarly, there exists $\{u, v\} \in D(Y)$ such that

$$T(\mathcal{P}(X, \{x, y\}) \cap \operatorname{Pc}(X)) \subset \mathcal{P}(Y, \{u, v\}) \cap \operatorname{Pc}(Y).$$

Then we get that

$$\mathcal{P}(Y, \{z, w\}) \cap \operatorname{Pc}(Y) = T(T^{-1}(\mathcal{P}(Y, \{z, w\}) \cap \operatorname{Pc}(Y)))$$

$$\subset T(\mathcal{P}(X, \{x, y\}) \cap \operatorname{Pc}(X)) \subset \mathcal{P}(Y, \{u, v\}) \cap \operatorname{Pc}(Y),$$

and hence $\{z, w\} = \{u, v\}$ due to Lemma 3.7. Therefore

$$T^{-1}(\mathcal{P}(Y, \{z, w\}) \cap \Pr(Y)) = T^{-1}(T(\mathcal{P}(X, \{x, y\}) \cap \Pr(X))) = \mathcal{P}(X, \{x, y\}) \cap \Pr(X).$$

Assume that

$$\{x', y'\} = \bigcap_{\rho \in \mathcal{P}(Y, \{z, w\}) \cap \operatorname{Pc}(Y)} \mathcal{F}(X, T^{-1}(\rho)),$$

so we get that

$$\mathcal{P}(X, \{x, y\}) \cap \Pr(X) = T^{-1}(\mathcal{P}(Y, \{z, w\}) \cap \Pr(Y)) = \mathcal{P}(X, \{x', y'\}) \cap \Pr(X).$$

Using Lemma 3.7 again, we have that $\{x, y\} = \{x', y'\}$. Hence we can define a map $\Phi : D(Y) \to D(X)$ by $\Phi(\{z, w\}) = \{x, y\}$.

On the other hand, fix any $\{x, y\} \in D(X)$. By the same argument as the above, we can uniquely choose $\{z, w\} \in D(Y)$ so that

$$T(\mathcal{P}(X, \{x, y\}) \cap \operatorname{Pc}(X)) = \mathcal{P}(Y, \{z, w\}) \cap \operatorname{Pc}(Y)$$

Then observe that

$$\mathcal{P}(X, \{x, y\}) \cap \Pr(X) = T^{-1}(T(\mathcal{P}(X, \{x, y\}) \cap \Pr(X)))$$

= $T^{-1}(\mathcal{P}(Y, \{z, w\}) \cap \Pr(Y)) = \mathcal{P}(X, \Phi(\{z, w\})) \cap \Pr(X),$

which implies that $\{x, y\} = \Phi(\{z, w\})$ by Lemma 3.7. Consequently, Φ is a bijection. The proof is completed. \Box

Remark 3.9. Due to the same argument, we can also obtain a bijection $\Phi : D(Y) \to D(X)$ as in the above proposition under the assumption T(Pp(X)) = Pp(Y). If T(M(X)) = M(Y), then the cardinality of X is coincident with the one of Y.

4. Constructing a bijection between X and Y

From now on, assume that T(M(X)) = M(Y), and let $\Phi : D(Y) \to D(X)$ be a bijection as in Proposition 3.8. Define $d(\Phi\{z, w\}) = d(x, y)$ for any $d \in PM(X)$ and any $\{z, w\} \in D(Y)$ with $\Phi(\{z, w\}) = \{x, y\} \in D(X)$. In this section, we will construct a bijection $\phi : Y \to X$ such that $\Phi(\{x, y\}) = \{\phi(x), \phi(y)\}$ for any $x, y \in Y$ and $\Phi^{-1}(\{z, w\}) = \{\phi^{-1}(z), \phi^{-1}(w)\}$ for any $z, w \in X$, and more, such that

$$T(d)(x,y) = d(\phi(x),\phi(y)) \tag{(*)}$$

holds for every $d \in PM(X)$. The map Φ induces the following equality between $d \in PM(X)$ and $T(d) \in PM(Y)$.

Proposition 4.1. The equality $T(d)(x, y) = d(\Phi(\{x, y\}))$ holds for every $d \in PM(X)$ and every $\{x, y\} \in D(Y)$.

Proof. First, we prove that $T(d)(x,y) \leq d(\Phi(\{x,y\}))$. Using Lemma 3.5, we can take $\rho \in \mathcal{P}(Y, \{x,y\})$ so that $T(d) + \rho \in \mathcal{P}(Y, \{x,y\}) \cap M(Y)$. By Proposition 3.8,

$$T^{-1}(\mathcal{P}(Y, \{x, y\}) \cap \mathcal{M}(Y)) = \mathcal{P}(X, \Phi(\{x, y\})) \cap \mathcal{M}(X),$$

and hence $T^{-1}(T(d) + \rho) \in \mathcal{P}(X, \Phi(\{x, y\}))$. Then due to Proposition 3.2,

$$T(d)(x,y) + \rho(x,y) = (T(d) + \rho)(x,y) = ||T(d) + \rho|| = ||T^{-1}(T(d) + \rho)||$$

= $T^{-1}(T(d) + \rho)(\Phi(\{x,y\})).$

Since T is isometry,

$$T(d)(x,y) + \rho(x,y) - d(\Phi(\{x,y\})) = T^{-1}(T(d) + \rho)(\Phi(\{x,y\})) - d(\Phi(\{x,y\}))$$

$$\leq \|T^{-1}(T(d) + \rho) - d\| = \|T^{-1}(T(d) + \rho) - T^{-1}(T(d))\|$$

$$= \|T(d) + \rho - T(d)\| = \|\rho\| = \rho(x,y),$$

which implies that $T(d)(x,y) \leq d(\Phi(\{x,y\}))$. Similarly, we get that $T(d)(x,y) \geq d(\Phi(\{x,y\}))$. The proof is completed. \Box

Now we shall construct a bijection from Y to X that is compatible with the isometry T.

Lemma 4.2. Assume that the cardinality of Y is greater than 2. For each point $y \in Y$, there uniquely exists a point $x \in X$ such that $\{x\} = \bigcap_{z \in Y \setminus \{y\}} \Phi(\{y, z\})$.

KATSUHISA KOSHINO

Proof. First, fix distinct points z_1 and z_2 in $Y \setminus \{y\}$, so we can uniquely find a point $x \in X$ such that $\{x\} = \Phi(\{y, z_1\}) \cap \Phi(\{y, z_2\})$. In fact, since Φ is an injection, the cardinality of $\Phi(\{y, z_1\}) \cap \Phi(\{y, z_2\})$ is less than 2. Suppose that $\Phi(\{y, z_1\})$ does not meet $\Phi(\{y, z_2\})$. It follows from Theorem 2.3 that there is $d \in PM(X)$ such that $d(\Phi(\{y, z_1\})) = d(\Phi(\{y, z_2\})) = 1$ and $d(\Phi(\{z_1, z_2\})) = 3$. Then by Proposition 4.1,

$$3 = d(\Phi(\{z_1, z_2\})) = T(d)(z_1, z_2) \le T(d)(y, z_1) + T(d)(y, z_2)$$

= $d(\Phi(\{y, z_1\})) + d(\Phi(\{y, z_2\})) = 2,$

which is a contradiction. Thus the doubletons $\Phi(\{y, z_1\})$ and $\Phi(\{y, z_2\})$ intersect at the only point $x \in X$. Then we can choose distinct points $w_1, w_2 \in X \setminus \{x\}$ so that $\{x, w_1\} = \Phi(\{y, z_1\})$ and $\{x, w_2\} = \Phi(\{y, z_2\})$. Suppose that there exists $z \in Y \setminus \{y\}$ such that $\Phi(\{y, z\})$ does not contain the point x, so by the above argument, $\{w_1, w_2\} = \Phi(\{y, z_2\})$. According to Theorem 2.3, taking a pseudometric $d \in PM(Y)$ such that $d(y, z_1) = d(y, z_2) = 1$ and d(y, z) = 3, we have that by Proposition 4.1,

$$3 = d(y, z) = T^{-1}(d)(\Phi(\{y, z\})) = T^{-1}(d)(w_1, w_2) \le T^{-1}(d)(x, w_1) + T^{-1}(d)(x, w_2)$$

= $T^{-1}(d)(\Phi(\{y, z_1\})) + T^{-1}(d)(\Phi(\{y, z_2\})) = d(y, z_1) + d(y, z_2) = 2.$

This is a contradiction. Hence $\{x\} = \bigcap_{z \in Y \setminus \{y\}} \Phi(\{y, z\})$. We complete the proof. \Box

By virtue of the above lemma, we can define a map $\phi: Y \to X$ by $\{\phi(y)\} = \bigcap_{z \in Y \setminus \{y\}} \Phi(\{y, z\})$ for every $y \in Y$.

Proposition 4.3. Suppose that the cardinality of X or Y is greater than 2. The map ϕ is a bijection, and $\Phi(\{x, y\}) = \{\phi(x), \phi(y)\}$ for any $x, y \in Y$ and $\Phi^{-1}(\{z, w\}) = \{\phi^{-1}(z), \phi^{-1}(w)\}$ for any $z, w \in X$.

Proof. Recall that the cardinality of X is coincident with the one of Y according to Remark 3.9. By the same argument as Lemma 4.2, we can define a map $\psi : X \to Y$ by $\{\psi(x)\} = \bigcap_{z \in X \setminus \{x\}} \Phi^{-1}(\{x, z\})$ for every point $x \in X$. Then $y = \psi(\phi(y))$ for any $y \in Y$. Indeed, letting $z_1, z_2 \in Y \setminus \{y\}$ be distinct points, we have that $\{\phi(y)\} = \Phi(\{y, z_1\}) \cap \Phi(\{y, z_2\})$ and

$$\{y\} = \{y, z_1\} \cap \{y, z_2\} = \Phi^{-1}(\Phi(\{y, z_1\})) \cap \Phi^{-1}(\Phi(\{y, z_2\})) = \{\psi(\phi(y))\}$$

Similarly, $\phi(\psi(x)) = x$ for each $x \in X$. Therefore ϕ is a bijection and $\phi^{-1} = \psi$. The latter part follows from the definition of ϕ and ψ . \Box

5. Proof of Main Theorem

Now we shall show Main Theorem.

Proof of Main Theorem. First, the implications $(1) \Rightarrow (2)$, $(1) \Rightarrow (3)$ and $(1) \Rightarrow (4)$ follow from [10, Lemma 2.4]. Indeed, taking any homeomorphism $\psi : Y \to X$, we can define an isometry $S : PM(X) \to PM(Y)$ by $S(d)(x, y) = d(\psi(x), \psi(y))$ for each $d \in PM(X)$ and for any $x, y \in Y$. Then S(AM(X)) = AM(Y) and S(M(X)) = M(Y). Second, since AM(X) and M(X) are dense in PM(X) by Proposition 2.4, each isometry in (3) and (4) can be extended to the one in (2), which means that $(3) \Rightarrow (2)$ and $(4) \Rightarrow (2)$ hold. Third, we shall show the implication $(2) \Rightarrow (1)$. In the case where the cardinality of X or Y is less than or equal to 2, obviously this implication holds, see Remark 3.9. In the other case, we need only to prove that the map $\phi : Y \to X$ as in Proposition 4.3 is a homeomorphism. To investigate that ϕ is continuous, fix any point $y \in Y$ and

any open neighborhood U of $\phi(y)$ in X. According to Theorem 2.3, we can find $d \in PM(X)$ such that $d(\phi(y), x) = 2$ if $x \in X \setminus U$, and d(x, x') = 0 if $x, x' \in X \setminus U$. Let

$$V = \{ z \in Y \mid T(d)(y, z) < 1 \},\$$

so it is an open neighborhood of y in Y. Then for every $z \in V$, $d(\phi(y), \phi(z)) = T(d)(y, z) < 1$, and hence $\phi(z) \in U$. It follows that ϕ is continuous. Similarly, ϕ^{-1} is also continuous. We conclude that $\phi: Y \to X$ is a homeomorphism. Finally, we shall investigate that the canonical formula (*) holds. When the cardinality of X or Y is less than or equal to 2, it is clear. When the cardinality of X or Yis greater than 2, we will show the uniqueness of the above homeomorphism ϕ . Let $\psi: Y \to X$ be a homeomorphism such that for every $d \in PM(X)$ and for any $x, y \in Y$, $T(d)(x, y) = d(\psi(x), \psi(y))$. Suppose that $\phi(x) \neq \psi(x)$ for some point $x \in Y$. Fix any $y \in Y \setminus \{x, \phi^{-1}(\psi(x))\}$ and take a pseudometric $\rho \in PM(X)$ such that $\rho(\phi(x), \phi(y)) = 0$ and $\rho(\psi(x), \psi(y)) = 1$, using Theorem 2.3. Then

$$0 = \rho(\phi(x), \phi(y)) = d(x, y) = \rho(\psi(x), \psi(y)) = 1.$$

which is a contradiction. We complete the proof. \Box

References

- [1] S. Banach, Théorie des opérations linéaires, Chelsea, Warsaw, (1932).
- [2] R. Engelking, General Topology, Revised and Complete Edition, Sigma Ser. in Pure Math., 6, Heldermann Verlag, Berlin, 1989.
- [3] R.J. Fleming and J.E. Jamison, Isometries on Banach spaces: function spaces, Chapman & Hall/CRC Monogr. Surv. Pure Appl. Math., 129, Boca Raton, FL, (2003).
- [4] F. Hausdorff, Erweiterung einer Homöomorphie, Fund. Math. 16 (1930), 353–360.
- [5] D. Hirota, I. Matsuzaki and T. Miura, Phase-isometries between the positive cones of the Banach spaces of continuous real-valued functions, Ann. Funct. Anal. 15 (2024), no. 4, Paper No. 77, 11pp.
- [6] Y. Ishiki, An interpolation of metrics and spaces of metrics, arXiv: 2003.13227 [math.MG].
- [7] Y. Ishiki, On dense subsets in spaces of metrics, Colloq. Math. 170 (2022), no. 1, 27–39.
- [8] Y. Ishiki, Spaces of metrics are Baire, arXiv: 2402.04565 [math.MG].
- [9] Y. Ishiki, An isometric extensor of metrics, arXiv: 2407.03030 [math.MG].
- [10] Y. Ishiki and K. Koshino, On isometric universality of spaces of metrics, arXiv: 2409.17701 [math.MG].
- [11] K. Koshino, Recognizing the topologies of spaces of metrics with the topology of uniform convergence, Bull. Pol. Acad. Sci. Math. 70 (2022), 165–171.
- [12] K. Koshino, On the Borel complexity and the complete metrizability of spaces of metrics, Anal. Geom. Metr. Spaces (2024), Article number: 20240014.
- [13] M.E. Shanks, The space of metrics on a compact metrizable space, Amer. J. Math. 66 (1944), 461–469.
- [14] M.H. Stone, Applications of the theory of Boolean rings to general topology, Trans. Amer. Math. Soc. 41 (1937), no. 3, 375–481.
- [15] L. Sun, Y. Sun and D. Dai, On phase-isometries between the positive cones of continuous function spaces, Ann. Funct. Anal. 14, (2023), No. 1, Paper No. 17, 12 pp.
- [16] J. van Mill, Infinite-Dimensional Topology, Prerequisites and Introduction, North-Holland Math. Library 43, Elsevier Sci. Publ., Amsterdam, 1989.

(Katsuhisa Koshino) Faculty of Engineering, Kanagawa University, 3-27-1 Rokkakubashi, Kanagawaku, Yokohama-shi, 221-8686, Japan

Email address: ft160229no@kanagawa-u.ac.jp