

TIME-FRACTIONAL GRADIENT FLOWS FOR NONCONVEX ENERGIES IN HILBERT SPACES

GORO AKAGI AND YOSHIHITO NAKAJIMA

ABSTRACT. This article is devoted to presenting an abstract theory on time-fractional gradient flows for nonconvex energy functionals in Hilbert spaces. Main results consist of local and global in time existence of (continuous) strong solutions to time-fractional evolution equations governed by the difference of two subdifferential operators in Hilbert spaces. To prove these results, fractional chain-rule formulae, a Lipschitz perturbation theory for convex gradient flows and Gronwall-type lemmas for nonlinear Volterra integral inequalities are developed. They also play a crucial role to cope with the lack of continuity (in time) of energies due to the subdiffusive nature of the issue. Moreover, the abstract theory is applied to the Cauchy–Dirichlet problem for some p -Laplace subdiffusion equations with blow-up terms complying with the so-called Sobolev (sub)critical growth condition.

1. INTRODUCTION

Let H be a real Hilbert space equipped with an inner product $(\cdot, \cdot)_H$ and the norm $\|\cdot\|_H = \sqrt{(\cdot, \cdot)_H}$. For $i = 1, 2$, let $\varphi^i : H \rightarrow (-\infty, \infty]$ be a proper (i.e., $\varphi^i \not\equiv \infty$) lower-semicontinuous convex functional with the *effective domain*,

$$D(\varphi^i) := \{w \in H : \varphi^i(w) < \infty\} \neq \emptyset,$$

and define the *subdifferential operator* $\partial\varphi^i : H \rightarrow 2^H$ of φ^i by

$$\partial\varphi^i(z) := \{\xi \in H : \varphi^i(v) - \varphi^i(z) \geq (\xi, v - z)_H \text{ for } v \in D(\varphi^i)\}$$

Date: January 15, 2025.

2020 Mathematics Subject Classification. *Primary:* 47J35; *Secondary:* 35K61.

Key words and phrases. Time-fractional gradient flows ; subdifferential operator ; fractional chain-rule formula ; p -Laplace subdiffusion equation ; local and global existence of strong solutions.

G.A. is supported by JSPS KAKENHI Grant Numbers JP24H00184, JP21KK0044, JP21K18581, JP20H01812 and JP20H00117. Y.N. is supported by JST SPRING, Grant Number JPMJSP2114. This work was also supported by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University.

for $z \in D(\varphi^i)$ with the domain $D(\partial\varphi^i) = \{w \in D(\varphi^i) : \partial\varphi^i(w) \neq \emptyset\}$. In this article, we are concerned with the abstract Cauchy problem for time-fractional evolution equations of the form,

$$\frac{d}{dt} [k * (u - u_0)](t) + \partial\varphi^1(u(t)) - \partial\varphi^2(u(t)) \ni f(t) \text{ in } H, \quad 0 < t < T, \quad (1.1)$$

where $T \in (0, \infty]$, $u_0 \in D(\varphi^1) \cap D(\varphi^2)$ and $f : (0, T) \rightarrow H$ are prescribed, and the convolution $k * (u - u_0)$ with a kernel $k \in L^1_{\text{loc}}([0, \infty))$ is defined by

$$(k * w)(t) := \int_0^t k(t-s)w(s) ds \quad \text{for } w \in L^1_{\text{loc}}([0, \infty); H), \quad t \geq 0.$$

Here and henceforth, we assume:

- (K) The kernel $k \in L^1_{\text{loc}}([0, \infty))$ is nonnegative and nonincreasing. There exists a nonnegative and nonincreasing kernel $\ell \in L^1_{\text{loc}}([0, \infty))$ such that

$$(k * \ell)(t) = \int_0^t k(t-s)\ell(s) ds = 1 \quad \text{for } t \geq 0.$$

Hence k is a *completely positive kernel* (see [12, Theorem 2.2]). A typical example of k satisfying (K) is the so-called *Riemann–Liouville kernel*,

$$k_\alpha(t) = \frac{t^{-\alpha}}{\Gamma(1-\alpha)} \quad \text{for } t > 0 \text{ and } 0 < \alpha < 1, \quad (1.2)$$

and then, the nonlocal time-derivative $(d/dt)[k_\alpha * (u - u_0)]$ corresponds to the α -th order Riemann–Liouville derivative $\partial_t^\alpha(u - u_0)$ of $u - u_0$, which also coincides with the α -th order Caputo derivative of u if it is smooth.

The study of classical gradient flows dates back to the early work of Haïm Brézis (see, e.g., [9]) concerning the evolution equation

$$\frac{du}{dt}(t) + \partial\varphi(u(t)) \ni f(t) \text{ in } H, \quad 0 < t < T, \quad u(0) = u_0$$

for a proper lower-semicontinuous *convex* functional φ on a Hilbert space H . The work of Brézis is nowadays known as *Brézis–Kōmura theory* (see also [20]) and has been extended in various directions. Among those, Ôtani [25] significantly relaxed the convexity assumption of the functional φ by studying the Cauchy problem for evolution equations governed by the difference of two subdifferential operators such as

$$\frac{du}{dt}(t) + \partial\varphi^1(u(t)) - \partial\varphi^2(u(t)) \ni f(t) \text{ in } H, \quad 0 < t < T, \quad u(0) = u_0, \quad (1.3)$$

which can be regarded as gradient flows for (essentially) *nonconvex* energies (see also [24] for long-time dynamics and [6] for a variational analysis). Actually, in contrast with gradient flows for *semiconvex* energies, the abstract Cauchy problem (1.3) can cover (even degenerate or singular) parabolic equations whose solutions blow up in finite time, and hence, one cannot generally expect global (in time) existence of solutions to (1.3). We further

refer the reader to [26, 27] for a nonmonotone perturbation theory for (convex) gradient flows. Extensions of these results on classical gradient flows to time-fractional variants have been pursued only for gradient flows of *convex* energies. In [3], the well-posedness of the abstract Cauchy problem

$$\frac{d}{dt} [k * (u - u_0)](t) + \partial\varphi(u(t)) \ni f(t) \text{ in } H, \quad 0 < t < T$$

is proved for proper lower-semicontinuous (semi)convex functionals $\varphi : H \rightarrow (-\infty, \infty]$ and $f \in L^2(0, T; H)$, and moreover, a Lipschitz perturbation theory is developed. Furthermore, the abstract theory is also applied to time-fractional variants of *nonlinear diffusion equations* as well as *Allen-Cahn equations*. We also refer the reader to [1] and references therein for related works (see also [2, 7, 16, 21, 29, 33, 34, 35, 36, 38, 39, 40, 42, 43, 44]). On the other hand, time-fractional gradient flows for (essentially) *nonconvex* energies such as (1.1) have never been studied so far.

The main purpose of the present paper is to establish an abstract theory on existence of strong solutions (continuous on $[0, T)$) to the Cauchy problem (1.1). To this end, there arise several significant difficulties from the presence of time-fractional derivatives. We immediately face a difficulty from the defect of chain-rule formula for time-fractional derivatives. Indeed, the chain-rule formula is a crucial device to analyze gradient flow equations. There is an alternative formula established in [3], from which a fractional chain-rule *inequality* for subdifferentials is derived in a practical form instead of usual identities; however, it is still insufficient to analyze the present issue (1.1). Moreover, the semigroup property does not hold true for the time-fractional evolution equations, that is, the concatenation of two solutions may not be a solution any more. As a result, one cannot construct a global solution to (1.1) by concatenating some local solutions (see Remark 2.2 below). We further realize another difficulty due to the lack of continuity (in time) of the energy $t \mapsto \varphi^1(u(t))$, which is actually from the subdiffusive nature of the issue. In the previous studies on non-fractional equations, various arguments based on the continuity are potentially used here and there, e.g., to construct local (in time) solutions to evolution equations (see, e.g., [18]). Hence we may need to develop an alternative scheme to establish a local existence result without exploiting the continuity of the energy.

Another aim of this paper is to apply the preceding abstract theory to the following Cauchy–Dirichlet problem for *p-Laplace subdiffusion equations with blow-up terms*:

$$\partial_t^\alpha(u - u_0) - \Delta_p u - |u|^{q-2}u = f \text{ in } \Omega \times (0, \infty), \tag{1.4}$$

$$u = 0 \text{ on } \partial\Omega \times (0, \infty), \tag{1.5}$$

where $1 < p, q < \infty$, Ω is a smooth bounded domain of \mathbb{R}^d with boundary $\partial\Omega$, $\partial_t^\alpha(u - u_0) = \partial_t[k_\alpha * (u - u_0)]$ denotes the Riemann–Liouville derivative of $u - u_0$ (i.e., Caputo derivative of u) of order $0 < \alpha < 1$ (see (1.2)), Δ_p is

the so-called *p-Laplacian* given as

$$\Delta_p w = \nabla \cdot (|\nabla w|^{p-2} \nabla w)$$

and $u_0 = u_0(x)$ and $f = f(x, t)$ are prescribed. Equation (1.4) is a time-fractional variant of *p*-Laplace diffusion equations with blow-up terms of the form

$$\partial_t u - \Delta_p u - |u|^{q-2} u = f \quad \text{in } \Omega \times (0, \infty), \quad (1.6)$$

along with the initial condition $u|_{t=0} = u_0$ in Ω (cf. see [15]). Equation (1.6) was proposed in [22] and then studied in [32], where local and global (in time) existence of weak solutions to the Cauchy–Dirichlet problem for (1.6) is proved for $u_0 \in W_0^{1,p}(\Omega)$ based on Galerkin’s method. Although the global existence is proved under the so-called Sobolev subcritical condition $q < p^* := dp/(d-p)_+$, the local existence is verified under a somewhat restrictive assumption on q . The abstract results established in [25, 18, 24, 26] are also applied to (1.6) for studying existence and asymptotic behavior of its L^2 -solutions under the assumption that $q \leq p^*/2 + 1$, which is strictly less than p^* (on the other hand, it can be relaxed to $q < 2^*$ for the semilinear case $p = 2$ in [26, 27]). Moreover, in [5], an abstract theory for (1.3) is developed in a reflexive Banach space setting and it is also used to prove local (in time) existence of weak solutions to the Cauchy–Dirichlet problem for (1.6) under the subcritical condition $q < p^*$ for $u_0 \in W_0^{1,p}(\Omega)$. Furthermore, in [4], local existence of weak solutions to the Cauchy–Dirichlet problem for (1.6) is eventually proved for any $u_0 \in L^r(\Omega)$ and $q > p \geq 2$, provided that $r > d(q-p)/p$; this result can cover all the local existence results mentioned above and even be regarded as an extension of Weissler’s results [37] on the (semilinear) Fujita equation. We also refer the reader to [23, 14]. On the other hand, to the best of the authors’ knowledge, there have been no contribution to the study of the Cauchy–Dirichlet problem (1.4), (1.5) for the *p*-Laplace subdiffusion equation so far. In this paper, thanks to the abstract theory on (1.1) along with a nonlinear Calderón–Zygmund theory, we prove local (in time) existence of L^2 -solutions to (1.4), (1.5) under the Sobolev subcritical condition $q < p^*$ for any $2d/(d+2) < p < \infty$. Moreover, global (in time) existence of L^2 -solutions for small data is also verified for $q \leq p^*$ even including the Sobolev critical exponent p^* .

Plan of the paper. This paper is composed of eight sections. In Section 2, we shall exhibit main results of the present paper, which consist of local and global (in time) existence of strong solutions to the abstract Cauchy problem (1.1) under certain assumptions. We also stress that the strong solutions are continuous on $[0, T)$. Section 3 is concerned with preliminary material on subdifferential operators (see §3.1) as well as nonlocal time-differential operators (see §3.2). In Section 4, we shall develop time-fractional chain-rule formulae for subdifferentials (see §4.1), continuous representatives of some convolutions (see §4.2) and a Lipschitz perturbation theory for convex

gradient flows in Hilbert spaces (see §4.3) in order to prove the main results. Sections 5–7 are devoted to giving proofs of the main results: Section 5 gives a proof of Theorem 2.3 concerning the local (in time) existence result. In Section 6, we shall prove Theorem 2.6 and Corollary 2.8, which are concerned with the global (in time) existence of strong solutions for small data. Moreover, another global existence result (see Theorem 2.9) will be proved in Section 7. Section 8 is dedicated to discussing applications of the abstract theory to the Cauchy–Dirichlet problem for p -Laplace subdiffusion equations with blow-up terms. In Appendix, we shall give an outline of a proof for the solvability of some approximate problems which will be used only in §7.

Notation. For each $1 < r < \infty$, we denote by r' the Hölder conjugate of r , that is, $1/r + 1/r' = 1$. Moreover, we use the same letter I for identity mappings defined on any spaces when no confusion can arise. We denote by C a generic nonnegative constant which may vary from line to line.

2. MAIN RESULTS

In this section, we present main results concerning existence of strong solutions to the abstract Cauchy problem (1.1). Throughout this paper, let H be a real Hilbert space equipped with an inner product $(\cdot, \cdot)_H$ and the norm $\|\cdot\|_H = \sqrt{(\cdot, \cdot)_H}$, and moreover, we are concerned with strong solutions to the Cauchy problem (1.1) in the following sense:

DEFINITION 2.1 (Strong solutions to (1.1)). Let $T \in (0, \infty]$. For any $S \in (0, T)$ (and $S \in (0, T]$ if $T < \infty$), a function $u \in L^2(0, S; H)$ is called a *strong solution on $[0, S]$* to the Cauchy problem (1.1), if the following conditions are all satisfied:

- (i) It holds that $k * (u - u_0) \in W^{1,2}(0, S; H)$ and $[k * (u - u_0)](0) = 0$. Moreover, $u(t) \in D(\partial\varphi^1) \cap D(\partial\varphi^2)$ for a.e. $t \in (0, S)$.
- (ii) There exist $\xi, \eta \in L^2(0, S; H)$ such that

$$\begin{cases} \xi(t) \in \partial\varphi^1(u(t)), & \eta(t) \in \partial\varphi^2(u(t)), \\ \frac{d}{dt} [k * (u - u_0)](t) + \xi(t) - \eta(t) = f(t) \end{cases} \quad (2.1)$$

for a.e. $t \in (0, S)$.

Moreover, a function $u \in L^2_{\text{loc}}([0, \infty); H)$ is called a *strong solution on $[0, \infty)$* to the Cauchy problem (1.1) with $T = \infty$, if the following conditions are all satisfied:

- (i)' It holds that $k * (u - u_0) \in W^{1,2}_{\text{loc}}([0, \infty); H)$ and $[k * (u - u_0)](0) = 0$. Moreover, $u(t) \in D(\partial\varphi^1) \cap D(\partial\varphi^2)$ for a.e. $t \in (0, \infty)$.
- (ii)' There exist $\xi, \eta \in L^2_{\text{loc}}([0, \infty); H)$ such that (2.1) holds for a.e. $t \in (0, \infty)$.

REMARK 2.2 (Lack of semigroup property). We emphasize that the semigroup property fails for strong solutions to the Cauchy problem (1.1) due to the nonlocal nature of the equation, that is, the concatenation

$$u(t) := \begin{cases} u_1(t) & \text{if } t \in [0, T], \\ u_2(t - T) & \text{if } t \in [T, 2T] \end{cases}$$

of two strong solutions u_1, u_2 on $[0, T]$ to (1.1) does not always solve (1.1) on $[0, 2T]$ even if $u_1(T) = u_2(0)$. Hence, the lack of semigroup property prevents us to construct (global) strong solutions on $[0, \infty)$ by concatenation. Moreover, it is noteworthy that, in contrast with classical gradient flows, the existence of strong solutions to (1.1) on $[0, \infty)$ does not follow immediately from the existence of those on $[0, T]$ for an arbitrary $T > 0$, unless solutions are uniquely determined by initial data.

In order to state main results of the present paper, we set up the following assumptions:

- (A1) For each $r \in \mathbb{R}$, the sublevel set $\{w \in H : \varphi^1(w) + \|w\|_H \leq r\}$ is (pre)compact in H .
- (A2) It holds that $D(\partial\varphi^1) \subset D(\partial\varphi^2)$, and moreover, there exist a constant $\nu_1 \in [0, 1)$ and a nondecreasing function $M_1 : \mathbb{R} \rightarrow [0, \infty)$ such that

$$\|\partial\overset{\circ}{\varphi}^2(w)\|_H \leq \nu_1 \|\partial\overset{\circ}{\varphi}^1(w)\|_H + M_1(\varphi^1(w) + \|w\|_H)$$

for $w \in D(\partial\varphi^1)$. Here $\partial\overset{\circ}{\varphi}^i$ denotes the minimal section of $\partial\varphi^i$ for $i = 1, 2$, that is, $\partial\overset{\circ}{\varphi}^i(w)$ is the unique element of the set $\partial\varphi^i(w)$ such that $\|\partial\overset{\circ}{\varphi}^i(w)\|_H = \min\{\|\xi\|_H : \xi \in \partial\varphi^i(w)\}$ for $w \in D(\partial\varphi^i)$.

We are now ready to state main results. The following theorem is concerned with local existence of strong solutions to (1.1).

THEOREM 2.3 (Local existence of strong solutions to (1.1)). *In addition to (K), assume that (A1) and (A2) hold and let $T \in (0, \infty)$. Then for every $u_0 \in D(\varphi^1)$ and $f \in L^2(0, T; H)$ satisfying $\ell * \|f(\cdot)\|_H^2 \in L^\infty(0, T)$, there exists $T_0 \in (0, T]$ such that the Cauchy problem (1.1) admits a strong solution $u \in L^2(0, T_0; H)$ on $[0, T_0]$ such that*

$$u \in C([0, T_0]; H), \quad u(0) = u_0 \quad \text{and} \quad \varphi^1(u(\cdot)) \in L^\infty(0, T_0).$$

Here T_0 depends only on $\ell, \nu_1, M_1(\cdot), \|u_0\|_H, \varphi^1(u_0)$ and $\|\ell * \|f(\cdot)\|_H^2\|_{L^\infty(0, T)}$.

REMARK 2.4. Let $T \in (0, \infty)$. Since $\ell \in L^1(0, T)$, every $f \in L^\infty(0, T; H)$ fulfills $\ell * \|f(\cdot)\|_H^2 \in C([0, T])$. In particular, if $k = k_\alpha$ with $0 < \alpha < 1$, since $\ell = k_{1-\alpha}$ belongs to $L^{1/(1-\alpha), \infty}(0, T)$, one can check $\ell * \|f(\cdot)\|_H^2 \in C([0, T])$ for any $f \in L^{2/\beta}(0, T; H)$ with some $\beta \in (0, \alpha)$ due to Young's convolution inequality.

We next present a theorem on global existence of strong solutions to (1.1) for ‘‘small’’ initial data by assuming the following conditions instead of (A1) and (A2):

(A1)' For each $r \in \mathbb{R}$, the sublevel set $\{w \in H : \varphi^1(w) \leq r\}$ is (pre)compact in H .

(A2)' It holds that $D(\partial\varphi^1) \subset D(\partial\varphi^2)$ and $\varphi^1 \geq 0$, and moreover, there exist a constant $\nu_2 \in [0, 1)$, a nonnegative Borel measurable locally bounded function $m_2 : [0, \infty) \rightarrow [0, \infty)$ and a nondecreasing function $M_2 : [0, \infty) \rightarrow [0, \infty)$ satisfying

$$m_2(r) > 0 \text{ for } 0 < r < r_0 \quad \text{and} \quad \lim_{r \rightarrow 0_+} \frac{M_2(r)}{m_2(r)} = 0 \quad (2.2)$$

for some $r_0 \in (0, \infty]$ such that

$$m_2(\varphi^1(w)) \leq \|\partial\dot{\varphi}^1(w)\|_H, \quad (2.3)$$

$$\|\partial\dot{\varphi}^2(w)\|_H \leq \nu_2 \|\partial\dot{\varphi}^1(w)\|_H + M_2(\varphi^1(w)) \quad (2.4)$$

for all $w \in D(\partial\varphi^1)$.

REMARK 2.5 (Coercivity of $\varphi^1 \Rightarrow$ condition (2.3)). If φ^1 fulfills the following coercivity condition:

$$\varphi^1(0) = 0 \quad \text{and} \quad \alpha \|w\|_H^p \leq \varphi^1(w) \quad \text{for } w \in D(\varphi^1) \quad (2.5)$$

for some constants $\alpha, p > 0$ (hence $\varphi^1(w) > 0$ if and only if $w \neq 0$), then one has

$$\begin{aligned} \varphi^1(w) &\leq (\partial\dot{\varphi}^1(w), w)_H \\ &\leq \|\partial\dot{\varphi}^1(w)\|_H \|w\|_H \stackrel{(2.5)}{\leq} \|\partial\dot{\varphi}^1(w)\|_H \alpha^{-1/p} \{\varphi^1(w)\}^{1/p}, \end{aligned}$$

which yields (2.3) with $m_2(s) := \alpha^{1/p} s^{1-1/p} > 0$ for $s > 0$ and $m_2(0) := 0$. Moreover, (A1)' also follows from (2.5) and (A1).

THEOREM 2.6 (Global existence of strong solutions to (1.1) for small data). *In addition to (K), assume that (A1)' and (A2)' hold. Then there exists a constant $\delta_0 > 0$ depending only on $M_2(\cdot)$, $m_2(\cdot)$ and ν_2 such that the following (i) and (ii) hold true:*

(i) *In case $T \in (0, \infty)$, for every $u_0 \in D(\varphi^1)$ and $f \in L^2(0, T; H)$ satisfying*

$$E_T(u_0, f) := \varphi^1(u_0) + \|\ell * \|f(\cdot)\|_H^2\|_{L^\infty(0, T)} < \delta_0,$$

the Cauchy problem (1.1) admits a strong solution $u \in L^2(0, T; H)$ on $[0, T]$ satisfying

$$u \in C([0, T]; H), \quad u(0) = u_0 \quad \text{and} \quad \varphi^1(u(\cdot)) \in L^\infty(0, T) \quad (2.6)$$

as well as

$$\sup_{t \in [0, T]} \varphi^1(u(t)) \leq C E_T(u_0, f)$$

for some constant $C > 0$ depending only on ν_2 .

(ii) In case $T = \infty$, for every $u_0 \in D(\varphi^1)$ and $f \in L^2_{\text{loc}}([0, \infty); H)$ satisfying

$$E_\infty(u_0, f) := \varphi^1(u_0) + \|\ell * \|f(\cdot)\|_H^2\|_{L^\infty(0, \infty)} < \delta_0, \quad (2.7)$$

the Cauchy problem (1.1) with $T = \infty$ admits a strong solution $u \in L^2_{\text{loc}}([0, \infty); H)$ on $[0, \infty)$ such that

$$u \in C([0, \infty); H), \quad u(0) = u_0 \quad \text{and} \quad \varphi^1(u(\cdot)) \in L^\infty(0, \infty)$$

and

$$\sup_{t \in [0, \infty)} \varphi^1(u(t)) \leq CE_\infty(u_0, f)$$

for some constant $C > 0$ depending only on ν_2 .

REMARK 2.7. When $\varphi^1(u_0) \ll 1$, a sufficient condition for $f \in L^2_{\text{loc}}([0, \infty); H)$ satisfying (2.7) reads,

$$\|f(t)\|_H^2 \leq \varepsilon k(t) \quad \text{for a.e. } t \in (0, \infty)$$

for some constant $\varepsilon > 0$ small enough.

We also give a corollary concerning global existence of strong solutions to (1.1) for small data under the following critical growth condition $(A2)'_c$ instead of $(A2)'$:

$(A2)'_c$ It holds that $D(\partial\varphi^1) \subset D(\partial\varphi^2)$ and $\varphi^1 \geq 0$, and moreover, there exists a nondecreasing function $M_3 : [0, \infty) \rightarrow [0, \infty)$ satisfying

$$M_3(r) > 0 \quad \text{for } r > 0 \quad \text{and} \quad \lim_{r \rightarrow 0^+} M_3(r) = 0 \quad (2.8)$$

such that

$$\|\partial\dot{\varphi}^2(w)\|_H \leq \|\partial\dot{\varphi}^1(w)\|_H M_3(\varphi^1(w)) \quad (2.9)$$

for all $w \in D(\partial\varphi^1)$.

COROLLARY 2.8 (Critical growth case). *In addition to (K), assume $(A1)'$ and $(A2)'_c$ above. Then there exists a constant $\delta_0 > 0$ depending only on $M_3(\cdot)$ such that (i) and (ii) of Theorem 2.6 hold true with $C = 1$.*

Finally, the following theorem asserts global existence of strong solutions to (1.1) even for “large” initial data under an additional assumption,

(A3) There exist constants $\nu_3 \in [0, 1)$, $r \in [0, 2)$ and $c_1 \geq 0$ such that

$$\varphi^2(w) \leq \nu_3 \varphi^1(w) + c_1 (\|w\|_H^r + 1) \quad \text{for } w \in D(\varphi^1),$$

which in particular yields $D(\varphi^1) \subset D(\varphi^2)$.

THEOREM 2.9 (Global existence of strong solutions to (1.1)). *In addition to (K), assume that (A1), (A2) and (A3) hold. In case $T \in (0, \infty)$, for every $u_0 \in D(\varphi^1)$ and $f \in W^{1,2}(0, T; H)$, the Cauchy problem (1.1) admits a strong solution $u \in L^2(0, T; H)$ on $[0, T]$ satisfying (2.6).*

In case $T = \infty$, for every $u_0 \in D(\varphi^1)$ and $f \in W^{1,2}_{\text{loc}}([0, \infty); H)$, then the Cauchy problem (1.1) with $T = \infty$ admits a strong solution $u \in L^2_{\text{loc}}([0, \infty); H)$ on $[0, \infty)$ satisfying (2.6) for any $T > 0$.

3. PRELIMINARIES

3.1. Subdifferential operators. In this subsection, we give a brief summary on *subdifferential operators* and *maximal monotone operators*. We refer the reader to, e.g., [9, 8] for the details of the following material.

Let $\varphi : H \rightarrow (-\infty, \infty]$ be a proper (i.e., $\varphi \not\equiv \infty$) lower-semicontinuous convex functional (see, e.g., [10]). Set the *effective domain* of φ as

$$D(\varphi) := \{w \in H : \varphi(w) < \infty\} \neq \emptyset.$$

Then the *subdifferential operator* $\partial\varphi : H \rightarrow 2^H$ of φ is defined by

$$\partial\varphi(w) := \{\xi \in H : \varphi(v) - \varphi(w) \geq (\xi, v - w)_H \text{ for all } v \in D(\varphi)\}$$

for $w \in D(\varphi)$ with the domain

$$D(\partial\varphi) := \{w \in D(\varphi) : \partial\varphi(w) \neq \emptyset\}.$$

It is known that every subdifferential operator is maximal monotone in H .

An operator $A : H \rightarrow 2^H$ is said to be *maximal monotone* (or *m-accretive*) in the Hilbert space H , if the following two properties hold:

- $(\xi_1 - \xi_2, u_1 - u_2)_H \geq 0$ for any $(u_1, \xi_1), (u_2, \xi_2) \in G(A)$,
- $R(I + \lambda A) = H$ for any (or some) $\lambda > 0$,

where $G(A)$ stands for the graph of the operator A and $R(I + \lambda A)$ denotes the range of the operator $I + \lambda A$. We also denote by $\mathring{A}(w)$ the *minimal section* of $A(w)$, that is, $\mathring{A}(w) = \operatorname{argmin}\{\|\xi\|_H : \xi \in A(w)\}$, for $w \in D(A) := \{w \in H : A(w) \neq \emptyset\}$. Such a minimal section is uniquely determined for each $w \in D(A)$. Moreover, as every maximal monotone operator is *demiclosedness*, we have the following useful property: Let $(u_n, \xi_n) \in G(A)$ be such that $u_n \rightarrow u$ strongly in H and $\xi_n \rightarrow \xi$ weakly in H . Then $(u, \xi) \in G(A)$.

For each maximal monotone operator $A : H \rightarrow 2^H$, the *resolvent* $J_\lambda : H \rightarrow H$ and *Yosida approximation* $A_\lambda : H \rightarrow H$ are defined by

$$J_\lambda = (I + \lambda A)^{-1}, \quad A_\lambda = \frac{I - J_\lambda}{\lambda}$$

for $\lambda > 0$. The following properties are well known:

PROPOSITION 3.1 (Resolvents and Yosida approximations). *Let $A : D(A) \subset H \rightarrow H$ be a maximal monotone operator in a Hilbert space H and let $J_\lambda : H \rightarrow H$ and $A_\lambda : H \rightarrow H$ be the resolvent and Yosida approximation of A , respectively, for $\lambda > 0$. Then the following (i)–(iv) are all satisfied:*

- (i) $J_\lambda : H \rightarrow H$ is non-expansive (i.e., 1-Lipschitz continuous) in H and $A_\lambda : H \rightarrow H$ is Lipschitz continuous in H with the Lipschitz constant $1/\lambda$. Moreover, A_λ is maximal monotone in H .
- (ii) $J_\lambda w \rightarrow w$ strongly in H as $\lambda \rightarrow 0_+$ for every $w \in \overline{D(A)}^H$.
- (iii) $A_\lambda(w) \in A(J_\lambda w)$ for $w \in H$ and $\lambda > 0$.
- (iv) $\|A_\lambda(w)\|_H \leq \|\mathring{A}(w)\|_H$ for $w \in D(A)$ and $\lambda > 0$, and moreover, $A_\lambda(w) \rightarrow \mathring{A}(w)$ strongly in H as $\lambda \rightarrow 0_+$ for $w \in D(A)$.

We next define the *Moreau–Yosida regularization* $\varphi_\lambda : H \rightarrow \mathbb{R}$ of φ for $\lambda > 0$ by

$$\begin{aligned}\varphi_\lambda(w) &:= \inf_{z \in H} \left(\frac{1}{2\lambda} \|w - z\|_H^2 + \varphi(z) \right) \\ &= \frac{1}{2\lambda} \|w - J_\lambda w\|_H^2 + \varphi(J_\lambda w) = \frac{\lambda}{2} \|A_\lambda(w)\|_H^2 + \varphi(J_\lambda w) \quad \text{for } w \in H,\end{aligned}$$

where J_λ and A_λ stand for the resolvent and the Yosida approximation of $A := \partial\varphi$, respectively. Then we recall that

PROPOSITION 3.2 (Moreau–Yosida regularizations). *Let $\varphi : H \rightarrow (-\infty, \infty]$ be a proper lower-semicontinuous convex functional defined in a Hilbert space H and let $\varphi_\lambda : H \rightarrow \mathbb{R}$ be the Moreau–Yosida regularization of φ for $\lambda > 0$. Then the following (i)–(iii) are all satisfied:*

- (i) *The Moreau–Yosida regularization $\varphi_\lambda : H \rightarrow \mathbb{R}$ is convex and Fréchet differentiable in H , and moreover, its derivative $\partial(\varphi_\lambda)$ coincides with the Yosida approximation $(\partial\varphi)_\lambda$ of $\partial\varphi$. Hence we shall simply write $\partial\varphi_\lambda$ instead of both $\partial(\varphi_\lambda)$ and $(\partial\varphi)_\lambda$ in what follows.*
- (ii) *$\varphi(J_\lambda w) \leq \varphi_\lambda(w) \leq \varphi(w)$ for $w \in H$ and $\lambda > 0$. Here $J_\lambda : H \rightarrow H$ stands for the resolvent of $\partial\varphi$.*
- (iii) *$\varphi_\lambda(w) \rightarrow \varphi(w)$ as $\lambda \rightarrow 0_+$ for $w \in H$.*

Define a functional $\Phi : L^2(0, T; H) \rightarrow (-\infty, \infty]$ by

$$\Phi(w) := \begin{cases} \int_0^T \varphi(w(t)) \, dt & \text{if } \varphi(w(\cdot)) \in L^1(0, T), \\ \infty & \text{otherwise} \end{cases}$$

for $w \in L^2(0, T; H)$. Then Φ is proper, lower-semicontinuous and convex in $L^2(0, T; H)$, and moreover, it is known that

$$\xi \in \partial\Phi(w) \quad \text{if and only if} \quad \xi(t) \in \partial\varphi(w(t)) \quad \text{for a.e. } t \in (0, T)$$

for $w \in D(\Phi)$ and $\xi \in L^2(0, T; H)$. Furthermore, it also holds that

$$\Phi_\lambda(w) = \int_0^T \varphi_\lambda(w(t)) \, dt, \quad \partial\Phi_\lambda(w) = \partial\varphi_\lambda(w(\cdot))$$

for $w \in L^2(0, T; H)$ and $\lambda > 0$ (see also [30, IV.Example 2.C]).

Finally, we recall the following chain-rule formula for subdifferentials:

PROPOSITION 3.3 (See [9, p.73, Lemma 3.3]). *Let $T \in (0, \infty)$ and let $u \in W^{1,2}(0, T; H)$ be such that $u(t) \in D(\partial\varphi)$ for a.e. $t \in (0, T)$. Assume that there exists $g \in L^2(0, T; H)$ such that $g(t) \in \partial\varphi(u(t))$ for a.e. $t \in (0, T)$. Then the function $t \mapsto \varphi(u(t))$ is absolutely continuous on $[0, T]$, and moreover, it holds that*

$$\left(\frac{du}{dt}(t), h \right)_H = \frac{d}{dt} \varphi(u(t)) \quad \text{for any } h \in \partial\varphi(u(t))$$

for a.e. $t \in (0, T)$.

3.2. Nonlocal time-differential operators. Let $T \in (0, \infty)$ and $p \in [1, \infty]$ be fixed and let X be a Banach space. We denote by

$$\mathcal{A} : D(\mathcal{A}) \subset L^p(0, T; X) \rightarrow L^p(0, T; X)$$

the (standard) *time-differential operator* defined by

$$\begin{aligned} D(\mathcal{A}) &:= \{w \in W^{1,p}(0, T; X) : w(0) = 0\}, \\ \mathcal{A}(w) &:= \frac{dw}{dt} \quad \text{for } w \in D(\mathcal{A}). \end{aligned}$$

Then \mathcal{A} is a linear m -accretive operator in $L^p(0, T; X)$.

Let $k \in L^1_{\text{loc}}([0, \infty))$ satisfy (K) (hence, k is a completely positive kernel). Then the *nonlocal time-differential operator*

$$\mathcal{B} : D(\mathcal{B}) \subset L^p(0, T; X) \rightarrow L^p(0, T; X)$$

is defined by

$$\begin{aligned} D(\mathcal{B}) &:= \{w \in L^p(0, T; X) : k * w \in D(\mathcal{A})\} \\ &= \{w \in L^p(0, T; X) : k * w \in W^{1,p}(0, T; X), (k * w)(0) = 0\}, \\ \mathcal{B}(w) &:= \mathcal{A}(k * w) = \frac{d}{dt}(k * w) \quad \text{for } w \in D(\mathcal{B}). \end{aligned}$$

Then we observe that

$$D(\mathcal{A}) \subset D(\mathcal{B}).$$

It is also known that \mathcal{B} is a linear m -accretive operator in $L^p(0, T; X)$ under the assumption (K) (see, e.g., [11], [17], [13], [33], [41]). Hence one can define the resolvent $\mathcal{J}_{1/n} : L^p(0, T; X) \rightarrow L^p(0, T; X)$ and the Yosida approximation $\mathcal{B}_{1/n} : L^p(0, T; X) \rightarrow L^p(0, T; X)$ of \mathcal{B} for $n \in \mathbb{N}$ by

$$\begin{aligned} \mathcal{J}_{1/n}(w) &:= (I + n^{-1}\mathcal{B})^{-1}(w), \\ \mathcal{B}_{1/n}(w) &:= n(w - \mathcal{J}_{1/n}(w)) = \mathcal{B}(\mathcal{J}_{1/n}w) = \frac{d}{dt}(k_n * w) \end{aligned}$$

for $w \in L^p(0, T; X)$. Here $k_n \in W^{1,1}(0, T)$ is a nonincreasing nonnegative kernel given by $k_n = ns_n$, where s_n is a unique solution to the Volterra integral equation,

$$s_n + n(\ell * s_n) = 1 \quad \text{in } (0, \infty).$$

Then k_n depends only on ℓ and n , and it is in particular independent of the choices of X and p . Due to the general theory on linear m -accretive operators (see, e.g., [28], [8]), we assure that

$$\mathcal{B}_{1/n}(w) \rightarrow \mathcal{B}(w) \quad \text{strongly in } L^p(0, T; X) \quad \text{as } n \rightarrow \infty, \quad (3.1)$$

provided that $w \in D(\mathcal{B})$. Indeed, \mathcal{B} is densely defined in $L^p(0, T; X)$, and hence, for $w \in D(\mathcal{B})$, we deduce that $\mathcal{B}_{1/n}(w) = \mathcal{B}(\mathcal{J}_{1/n}(w)) = \mathcal{J}_{1/n}(\mathcal{B}(w)) \rightarrow \mathcal{B}(w)$ strongly in $L^p(0, T; X)$ as $n \rightarrow \infty$. In particular, setting $X = \mathbb{R}$ and $p = 1$, we find that $w \equiv 1 \in D(\mathcal{B})$, and hence, it follows from (3.1) that

$$k_n \rightarrow k \quad \text{strongly in } L^1(0, T) \quad \text{as } n \rightarrow \infty. \quad (3.2)$$

In what follows, we always set

$$X = H \quad \text{and} \quad p = 2,$$

unless noted otherwise, to consider the operators \mathcal{A} and \mathcal{B} for $k \in L^1_{\text{loc}}([0, \infty))$ satisfying (K). Then \mathcal{A} and \mathcal{B} are maximal monotone in $L^2(0, T; H)$, and hence, thanks to Mazur's lemma along with the (demi)closedness of \mathcal{A} and \mathcal{B} , the graphs $G(\mathcal{A})$ and $G(\mathcal{B})$ of \mathcal{A} and \mathcal{B} , respectively, are weakly closed in $L^2(0, T; H) \times L^2(0, T; H)$.

Finally, we recall the following

PROPOSITION 3.4 (See [3, Proposition 3.5 and Corollary 3.6]). *Let $T \in (0, \infty)$. Under the assumption (K), it holds that*

$$\begin{aligned} & \int_0^t (\mathcal{A}(u)(\tau), \mathcal{B}(u)(\tau))_H \, d\tau \\ & \geq \frac{1}{2} (\ell * \|\mathcal{B}(u)(\cdot)\|_H^2)(t) + \frac{1}{2} \int_0^t \ell(\tau) \|\mathcal{B}(u)(\tau)\|_H^2 \, d\tau \end{aligned}$$

for all $u \in D(\mathcal{A})$ and $t \in [0, T]$. It further implies the maximal monotonicity of $\mathcal{A} + \mathcal{B}$ in $L^2(0, T; H)$.

4. SOME DEVICES

This section is mainly devoted to developing *integral forms of fractional chain-rule formulae* and a *Lipschitz perturbation theory* for proving the main results stated in §2. Moreover, we also give a sufficient condition for the existence of continuous representatives of convolutions.

4.1. Fractional chain-rule formulae. In this subsection, we develop some integral forms of (time-)fractional chain-rule formulae. To this end, we first recall the following nonlocal chain-rule formula for regular kernels, which is used in [3] to derive differential forms of fractional chain-rule formulae:

PROPOSITION 4.1 (Nonlocal chain-rule formula [3, Proposition 3.4]). *Let $T \in (0, \infty)$. Let $h \in W^{1,1}(0, T)$ be a nonnegative nonincreasing function and let $\varphi : H \rightarrow (-\infty, \infty]$ be a proper (i.e., $\varphi \not\equiv \infty$) lower-semicontinuous convex functional. Let $u \in L^1(0, T; H)$ be such that $\varphi(u(\cdot)) \in L^1(0, T)$. Then for each $t \in (0, T)$ satisfying $u(t) \in D(\partial\varphi)$, it holds that*

$$\left(\frac{d}{dt} [h * (u - u_0)](t), g \right)_H \geq \frac{d}{dt} [h * (\varphi(u(\cdot)) - \varphi(u_0))](t)$$

for any $u_0 \in D(\varphi)$ and $g \in \partial\varphi(u(t))$.

In particular, if we choose $\varphi(\cdot) = (1/2)\|\cdot\|_H^2$, then we also have

$$\left(\frac{d}{dt} [h * w](t), w(t) \right)_H \geq \frac{1}{2} \frac{d}{dt} [h * \|w(\cdot)\|_H^2](t)$$

for a.e. $t \in (0, T)$ and $w \in L^2(0, T; H)$ (see [41]).

We next present some integral forms of fractional chain-rule formulae, which will play a crucial role to derive a priori estimates for approximate solutions to (1.1).

PROPOSITION 4.2 (Integral forms of fractional chain-rule formulae). *Let $k \in L^1_{\text{loc}}([0, \infty))$ be a function satisfying (K). Let $\varphi : H \rightarrow (-\infty, \infty]$ be a proper (i.e., $\varphi \not\equiv \infty$) lower-semicontinuous convex functional. Let $T \in (0, \infty)$, $u_0 \in D(\varphi)$ and let $u \in L^2(0, T; H)$ satisfy*

- $k * (u - u_0) \in W^{1,2}(0, T; H)$ and $[k * (u - u_0)](0) = 0$,
- $u(t) \in D(\partial\varphi)$ for a.e. $t \in (0, T)$.

Assume that there exists $g \in L^2(0, T; H)$ such that $g(t) \in \partial\varphi(u(t))$ for a.e. $t \in (0, T)$. Then the following holds true:

(i) *It holds that*

$$\int_0^t \left(\frac{d}{dt} [k * (u - u_0)](\tau), g(\tau) \right)_H d\tau \geq [k * (\varphi(u(\cdot)) - \varphi(u_0))] (t)$$

for a.e. $t \in (0, T)$.

(ii) *Let ℓ be the conjugate kernel of k (see (K)). It holds that*

$$\left[\ell * \left(\frac{d}{dt} [k * (u - u_0)](\cdot), g(\cdot) \right)_H \right] (t) \geq \varphi(u(t)) - \varphi(u_0)$$

for a.e. $t \in (0, T)$.

Proof. First of all, it follows from the assumption that $\varphi(u(\cdot)) \in L^1(0, T)$. Indeed, we see that

$$\varphi(u(t)) \leq \varphi(u_0) + (g(t), u(t) - u_0)_H$$

for a.e. $t \in (0, T)$, and moreover, u and g belong to $L^2(0, T; H)$. Furthermore, φ is affinely bounded from below (see, e.g., [9]), we conclude that $\varphi(u(\cdot)) \in L^1(0, T)$.

Assertion (i) can be proved as in [3]; however, we give a proof for the convenience of the reader. For $n \in \mathbb{N}$, let $k_n \in W^{1,1}(0, T)$ be a regularized kernel (see §3.2). Thanks to Proposition 4.1, we have

$$\left(\frac{d}{dt} [k_n * (u - u_0)](t), g(t) \right)_H \geq \frac{d}{dt} [k_n * (\varphi(u(\cdot)) - \varphi(u_0))] (t) \quad (4.1)$$

for a.e. $t \in (0, T)$. Integrate both sides over $(0, t)$, we infer that

$$\int_0^t \left(\frac{d}{dt} [k_n * (u - u_0)](\tau), g(\tau) \right)_H d\tau \geq [k_n * (\varphi(u(\cdot)) - \varphi(u_0))] (t)$$

for $t \in [0, T]$. Since $\mathcal{B}_{1/n}(u - u_0) = (d/dt)[k_n * (u - u_0)] \rightarrow \mathcal{B}(u - u_0) = (d/dt)[k * (u - u_0)]$ strongly in $L^2(0, T; H)$ (by $u - u_0 \in D(\mathcal{B})$ and (3.1)) and $k_n \rightarrow k$ strongly in $L^1(0, T)$ (see (3.2)), passing to the limit as $n \rightarrow \infty$, we deduce that

$$\int_0^t \left(\frac{d}{dt} [k * (u - u_0)](\tau), g(\tau) \right)_H d\tau \geq [k * (\varphi(u(\cdot)) - \varphi(u_0))] (t)$$

for a.e. $t \in (0, T)$.

As for (ii), we convolve both sides of (4.1) with ℓ to see that

$$\begin{aligned} & \left[\ell * \left(\frac{d}{dt} [k_n * (u - u_0)](\cdot), g(\cdot) \right)_H \right] (t) \\ & \geq \ell * \frac{d}{dt} [k_n * (\varphi(u(\cdot)) - \varphi(u_0))] (t) \\ & = \frac{d}{dt} [\ell * k_n * (\varphi(u(\cdot)) - \varphi(u_0))] (t) \end{aligned} \quad (4.2)$$

for a.e. $t \in (0, T)$. Here we used the fact that $[k_n * (\varphi(u(\cdot)) - \varphi(u_0))](0) = 0$. Moreover, we claim that

$$\ell * \left(\frac{d}{dt} [k_n * (u - u_0)](\cdot), g(\cdot) \right)_H \rightarrow \ell * \left(\frac{d}{dt} [k * (u - u_0)](\cdot), g(\cdot) \right)_H \quad (4.3)$$

strongly in $L^1(0, T)$ as $n \rightarrow \infty$. Indeed, we see that

$$\begin{aligned} & \left\| \ell * \left(\frac{d}{dt} [k_n * (u - u_0)](\cdot), g(\cdot) \right)_H - \ell * \left(\frac{d}{dt} [k * (u - u_0)](\cdot), g(\cdot) \right)_H \right\|_{L^1(0, T)} \\ & \leq \|\ell\|_{L^1(0, T)} \|\mathcal{B}_{1/n}(u - u_0) - \mathcal{B}(u - u_0)\|_{L^2(0, T; H)} \|g\|_{L^2(0, T; H)} \rightarrow 0. \end{aligned}$$

We next test both sides of (4.2) by a nonnegative test function $\phi \in C_c^\infty(0, T)$. It then follows that

$$\begin{aligned} & \int_0^T \phi(t) \left[\ell * \left(\frac{d}{dt} [k_n * (u - u_0)](\cdot), g(\cdot) \right)_H \right] (t) dt \\ & \geq \int_0^T \phi(t) \frac{d}{dt} [\ell * k_n * (\varphi(u(\cdot)) - \varphi(u_0))] (t) dt \\ & = - \int_0^T \phi'(t) [\ell * k_n * (\varphi(u(\cdot)) - \varphi(u_0))] (t) dt \\ & \rightarrow - \int_0^T \phi'(t) [\ell * k * (\varphi(u(\cdot)) - \varphi(u_0))] (t) dt \\ & \stackrel{(K)}{=} - \int_0^T \phi'(t) \left(\int_0^t [\varphi(u(\tau)) - \varphi(u_0)] d\tau \right) dt \\ & = \int_0^T \phi(t) [\varphi(u(t)) - \varphi(u_0)] dt. \end{aligned}$$

Hence combining the above with (4.3), we conclude that

$$\begin{aligned} & \int_0^T \phi(t) \left[\ell * \left(\frac{d}{dt} [k * (u - u_0)](\cdot), g(\cdot) \right)_H \right] (t) dt \\ & \geq \int_0^T \phi(t) [\varphi(u(t)) - \varphi(u_0)] dt, \end{aligned}$$

whence it follows from the arbitrariness of $\phi \geq 0$ that

$$\left[\ell * \left(\frac{d}{dt} [k * (u - u_0)](\cdot), g(\cdot) \right) \right]_H(t) \geq \varphi(u(t)) - \varphi(u_0)$$

for a.e. $t \in (0, T)$. This completes the proof. \square

4.2. Continuous representatives of convolutions. In this subsection, we give some criterion on existence of continuous representatives of convolutions, which will be used to prove continuity of strong solutions to the Cauchy problem (1.1) and to check initial conditions in a classical sense. Moreover, it will also be used in the next subsection concerned with the Lipschitz perturbation theory.

LEMMA 4.3 (Continuous representatives of convolutions). *Let $T \in (0, \infty)$, $a \in H$ and let $g \in L^1(0, T)$ be nonnegative and nonincreasing. Let $G \in L^1(0, T; H)$ be such that $g * \|G(\cdot)\|_H^2 \in L^\infty(0, T)$ and define $u \in L^1(0, T; H)$ by*

$$u(t) = a + (g * G)(t) \quad \text{for a.e. } t \in (0, T). \quad (4.4)$$

Then u has a continuous representative $\tilde{u} \in C([0, T]; H)$, that is, $u(t) = \tilde{u}(t)$ for a.e. $t \in (0, T)$, such that

$$\begin{aligned} \|\tilde{u}(t) - \tilde{u}(s)\|_H &\leq \|g\|_{L^1(0, |t-s|)}^{1/2} \|g * \|G(\cdot)\|_H^2\|_{L^\infty(0, T)}^{1/2} \\ &\quad + \|g(|t-s| + \cdot) - g(\cdot)\|_{L^1(0, T-|t-s|)}^{1/2} \|g * \|G(\cdot)\|_H^2\|_{L^\infty(0, T)}^{1/2} \end{aligned}$$

for all $t, s \in [0, T]$. Moreover, it holds that $\tilde{u}(t) \rightarrow a$ strongly in H as $t \rightarrow 0_+$.

Proof. By assumption, there exists $I \subset (0, T)$ such that $|I| = 0$ and (4.4) holds for all $t \in (0, T) \setminus I$. Then letting $s, t \in (0, T) \setminus I$ satisfy $s \leq t$ and using Hölder's inequality, one observes that

$$\begin{aligned} \|u(t) - u(s)\|_H &= \|(g * G)(t) - (g * G)(s)\|_H \\ &= \left\| \int_0^t g(t-\tau)G(\tau) \, d\tau - \int_0^s g(s-\tau)G(\tau) \, d\tau \right\|_H \\ &\leq \int_s^t g(t-\tau)\|G(\tau)\|_H \, d\tau + \int_0^s |g(t-\tau) - g(s-\tau)|\|G(\tau)\|_H \, d\tau \\ &= \int_s^t g(t-\tau)^{1/2}g(t-\tau)^{1/2}\|G(\tau)\|_H \, d\tau \\ &\quad + \int_0^s |g(t-\tau) - g(s-\tau)|^{1/2}|g(t-\tau) - g(s-\tau)|^{1/2}\|G(\tau)\|_H \, d\tau \\ &\leq \|g\|_{L^1(0, t-s)}^{1/2} \|g * \|G(\cdot)\|_H^2\|_{L^\infty(0, T)}^{1/2} \\ &\quad + \|g(t-s + \cdot) - g(\cdot)\|_{L^1(0, T-(t-s))}^{1/2} \|g * \|G(\cdot)\|_H^2\|_{L^\infty(0, T)}^{1/2}. \end{aligned}$$

Here we also used $|g(t-\tau) - g(s-\tau)| = g(s-\tau) - g(t-\tau) \leq g(s-\tau)$ from the nonincrease and nonnegativity of g to derive the last inequality.

For each $t \in [0, T]$ one can take a sequence (t_n) in $(0, T) \setminus I$ converging to t , and then, $(u(t_n))$ forms a Cauchy sequence in H . Hence $\tilde{u}(t)$ can be defined as its limit, since the limit is also uniquely determined. Therefore there exists a continuous representative $\tilde{u} \in C([0, T]; H)$ of u such that \tilde{u} fulfills the desired inequality for any $s, t \in [0, T]$. Moreover, repeating the same calculation above, we also find that

$$\|u(t) - a\|_H \leq \|g\|_{L^1(0,t)}^{1/2} \|g * \|G(\cdot)\|_H^2\|_{L^\infty(0,T)}^{1/2}$$

for all $t \in (0, T) \setminus I$. Hence $\tilde{u}(t) \rightarrow a$ strongly in H as $t \rightarrow 0_+$. This completes the proof. \square

4.3. Lipschitz perturbation theory. A Lipschitz perturbation theory to time-fractional gradient flows for convex energies is established in [3, §5]. In this subsection, it is further developed in order to derive better regularity (which can enable us to guarantee the initial condition in a classical sense) of strong solutions to the Cauchy problem,

$$\frac{d}{dt} [k * (u - u_0)](t) + \partial\varphi(u(t)) + B(u(t)) \ni f(t) \text{ in } H, \quad 0 < t < T, \quad (4.5)$$

where $T \in (0, \infty]$, $\varphi : H \rightarrow (-\infty, \infty]$ is a proper lower-semicontinuous convex functional, $B : H \rightarrow H$ is a Lipschitz continuous operator, i.e., there exists a constant $L_B \geq 0$ such that

$$\|B(w_1) - B(w_2)\|_H \leq L_B \|w_1 - w_2\|_H \quad \text{for } w_1, w_2 \in H, \quad (4.6)$$

for every $u_0 \in D(\varphi)$ and $f \in L^2(0, T; H)$ satisfying $\ell * \|f(\cdot)\|_H^2 \in L^\infty(0, T)$. Strong solutions to (4.5) are defined analogously to Definition 2.1. The following theorem will be used to construct approximate solutions to (1.1).

THEOREM 4.4 (Lipschitz perturbation theory (cf. see [3])). *Assume that (K) and (4.6) are satisfied and let $T \in (0, \infty]$. In case $T \in (0, \infty)$, for every $u_0 \in D(\varphi^1)$ and $f \in L^2(0, T; H)$ satisfying $\ell * \|f(\cdot)\|_H^2 \in L^\infty(0, T)$, the Cauchy problem (4.5) admits a unique strong solution u on $[0, T]$ such that*

$$u \in C([0, T]; H), \quad u(0) = u_0 \quad \text{and} \quad \varphi(u(\cdot)) \in L^\infty(0, T). \quad (4.7)$$

*In case $T = \infty$, for every $u_0 \in D(\varphi^1)$ and $f \in L_{\text{loc}}^2([0, \infty); H)$ satisfying $\ell * \|f(\cdot)\|_H^2 \in L_{\text{loc}}^\infty([0, \infty))$, the Cauchy problem (4.5) with $T = \infty$ admits a unique strong solution u on $[0, \infty)$ satisfying (4.7) for every $T > 0$.*

Proof. Let $T \in (0, \infty)$ be fixed. With the aid of Theorem 5.1 of [3], for every $u_0 \in D(\varphi)$ and $f \in L^2(0, T; H)$, one can assure that there exists a unique strong solution $u \in L^2(0, T; H)$ on $[0, T]$ to (4.5), that is, there is $\xi \in L^2(0, T; H)$ such that

$$\frac{d}{dt} [k * (u - u_0)](t) + \xi(t) + B(u(t)) = f(t), \quad \xi(t) \in \partial\varphi(u(t)) \quad (4.8)$$

for a.e. $t \in (0, T)$. Hence it suffices to show (4.7) under the (additional) assumption $\ell * \|f(\cdot)\|_H^2 \in L^\infty(0, T)$. We assume $\varphi \geq 0$ without loss of

generality. Multiplying both sides of (4.8) by $(d/dt)[k * (u - u_0)](t)$, we have

$$\begin{aligned}
 & \left\| \frac{d}{dt} [k * (u - u_0)](t) \right\|_H^2 + \left(\xi(t), \frac{d}{dt} [k * (u - u_0)](t) \right)_H \\
 &= - \left(B(u(t)), \frac{d}{dt} [k * (u - u_0)](t) \right)_H + \left(f(t), \frac{d}{dt} [k * (u - u_0)](t) \right)_H \\
 &\leq \frac{1}{2} \left\| \frac{d}{dt} [k * (u - u_0)](t) \right\|_H^2 + \|B(u(t))\|_H^2 + \|f(t)\|_H^2 \\
 &\leq \frac{1}{2} \left\| \frac{d}{dt} [k * (u - u_0)](t) \right\|_H^2 + (\|B(u_0)\|_H + L_B \|u(t) - u_0\|_H)^2 \\
 &\quad + \|f(t)\|_H^2
 \end{aligned}$$

for a.e. $t \in (0, T)$. Convolving both sides with ℓ and using (ii) of Proposition 4.2, we infer that

$$\begin{aligned}
 & \frac{1}{2} \left(\ell * \left\| \frac{d}{dt} [k * (u - u_0)](\cdot) \right\|_H^2 \right) (t) + \varphi(u(t)) - \varphi(u_0) \\
 &\leq \|\ell * \|f(\cdot)\|_H^2\|_{L^\infty(0,T)} + 2\|B(u_0)\|_H^2 \|\ell\|_{L^1(0,T)} \\
 &\quad + 2L_B^2 (\ell * \|u(\cdot) - u_0\|_H^2) (t)
 \end{aligned} \tag{4.9}$$

for a.e. $t \in (0, T)$. Since φ is nonnegative, one has

$$\begin{aligned}
 & \left(\ell * \left\| \frac{d}{dt} [k * (u - u_0)](\cdot) \right\|_H^2 \right) (t) + \varphi(u(t)) \\
 &\leq C \left[\varphi(u_0) + \|\ell * \|f(\cdot)\|_H^2\|_{L^\infty(0,T)} + \|B(u_0)\|_H^2 \|\ell\|_{L^1(0,T)} \right] \\
 &\quad + C (\ell * \|u(\cdot) - u_0\|_H^2) (t)
 \end{aligned}$$

for a.e. $t \in (0, T)$. Here we note the following fact:

$$\begin{aligned}
 \|u(t) - u_0\|_H^2 &= \left\| \frac{d}{dt} [k * \ell * (u - u_0)](t) \right\|_H^2 \\
 &= \left\| \left(\ell * \frac{d}{dt} [k * (u - u_0)] \right) (t) \right\|_H^2 \\
 &= \left\| \int_0^t \ell^{1/2}(t-s) \ell^{1/2}(t-s) \left(\frac{d}{dt} [k * (u - u_0)] \right) (s) ds \right\|_H^2 \\
 &\leq \left(\int_0^t \ell(t-s) ds \right) \left(\int_0^t \ell(t-s) \left\| \frac{d}{dt} [k * (u - u_0)](s) \right\|_H^2 ds \right) \\
 &\leq \|\ell\|_{L^1(0,T)} \left(\ell * \left\| \frac{d}{dt} [k * (u - u_0)](\cdot) \right\|_H^2 \right) (t)
 \end{aligned}$$

for a.e. $t \in (0, T)$. It follows that

$$\begin{aligned} & \|u(t) - u_0\|_H^2 \\ & \leq C \|\ell\|_{L^1(0,T)} \left[\varphi(u_0) + \|\ell * \|f(\cdot)\|_H^2\|_{L^\infty(0,T)} + \|B(u_0)\|_H^2 \|\ell\|_{L^1(0,T)} \right] \\ & \quad + C \|\ell\|_{L^1(0,T)} (\ell * \|u(\cdot) - u_0\|_H^2)(t) \end{aligned}$$

for a.e. $t \in (0, T)$. Thus applying a Gronwall-type lemma for a Volterra integral inequality (see Lemma 4.5 below), we obtain

$$\operatorname{ess\,sup}_{t \in (0, T)} \|u(t)\|_H \leq C,$$

which along with (4.9) yields

$$\operatorname{ess\,sup}_{t \in (0, T)} \left[\left(\ell * \left\| \frac{d}{dt} [k * (u - u_0)](\cdot) \right\|_H^2 \right)(t) + \varphi(u(t)) \right] \leq C.$$

Moreover, we recall that

$$u(t) = u_0 + \left(\ell * \frac{d}{dt} [k * (u - u_0)] \right)(t) \quad \text{for a.e. } t \in (0, T).$$

Therefore Lemma 4.3 asserts that there exists a continuous representative of u (still denoted by the same letter) and $u(0) = u_0$.

Finally, we prove existence of strong solutions on $[0, \infty)$ to the Cauchy problem (4.5) with $T = \infty$ for $u_0 \in D(\varphi^1)$ and $f \in L_{\text{loc}}^2([0, \infty); H)$ satisfying $\ell * \|f(\cdot)\|_H^2 \in L_{\text{loc}}^\infty([0, \infty))$. For each finite $T > 0$, from the fact which we have proved so far, the Cauchy problem (4.5) admits a unique strong solution $u^{(T)} \in C([0, T]; H)$ on $[0, T]$ satisfying (4.7). Hence due to the uniqueness of strong solutions to (4.5) (i.e., $u^{(T)}|_{[0, S]} = u^{(S)}$ if $S \leq T$), one can define $u \in C([0, \infty); H)$ such that $u|_{[0, T]} = u^{(T)}$ for $T > 0$. We also set $\xi := f - B(u(\cdot)) - (d/dt)[k * (u - u_0)] \in L_{\text{loc}}^2([0, \infty); H)$ and $\eta := B(u(\cdot)) \in C([0, \infty); H)$. Therefore it follows that $u \in C([0, \infty); H)$ is a strong solution on $[0, \infty)$ to (4.5) with $T = \infty$ and $u(0) = u_0$. This completes the proof. \square

We close this subsection with the following Gronwall-type lemma for a Volterra integral inequality:

LEMMA 4.5. *Let $S \in (0, \infty)$ and $1 \leq r \leq \infty$. Let $\phi \in L^1(0, S)$, $h \in L^r(0, S)$ and $g \in L^1(0, S)$ be nonnegative functions such that*

$$\phi(t) \leq h(t) + (g * \phi)(t) \quad \text{for a.e. } t \in (0, S). \quad (4.10)$$

Then it holds that $\phi \in L^r(0, S)$, and moreover, there exists a constant $C_0 \geq 0$ depending only on S and g such that

$$\|\phi\|_{L^r(0, S)} \leq C_0 \|h\|_{L^r(0, S)}.$$

Proof. Let (ρ_n) be a mollifier¹ and convolve both sides of (4.10) with $\rho_n \geq 0$. It then follows that

$$\phi_n(t) \leq h_n(t) + (g * \phi_n)(t) \quad \text{for a.e. } t \in (0, S), \quad (4.11)$$

where $\phi_n := \rho_n * \phi \in L^\infty(0, S)$ and $h_n := \rho_n * h \in L^\infty(0, S)$. We observe that

$$\begin{aligned} (g * \phi_n)(t) &= \int_0^t g(t-s)\phi_n(s) \, ds \\ &= e^{Mt} \int_0^t g(t-s)e^{-M(t-s)}\phi_n(s)e^{-Ms} \, ds \\ &= e^{Mt} [(g e^{-M\cdot}) * (\phi_n e^{-M\cdot})](t) \quad \text{for } t \in [0, S], \end{aligned}$$

where M is a constant which will be determined later and $e^{-M\cdot}$ stands for the function $t \mapsto e^{-Mt}$ from $[0, \infty) \rightarrow \mathbb{R}$, and which along with the Young's convolution inequality yields

$$\|(g * \phi_n) e^{-M\cdot}\|_{L^r(0,S)} \leq \|g e^{-M\cdot}\|_{L^1(0,S)} \|\phi_n e^{-M\cdot}\|_{L^r(0,S)}.$$

Hence it follows from (4.11) that

$$\begin{aligned} \|\phi_n e^{-M\cdot}\|_{L^r(0,S)} &\leq \|h_n e^{-M\cdot}\|_{L^r(0,S)} + \|(g * \phi_n) e^{-M\cdot}\|_{L^r(0,S)} \\ &\leq \|h_n e^{-M\cdot}\|_{L^r(0,S)} + \|g e^{-M\cdot}\|_{L^1(0,S)} \|\phi_n e^{-M\cdot}\|_{L^r(0,S)}. \end{aligned}$$

We take a positive constant M large enough so that

$$\|g e^{-M\cdot}\|_{L^1(0,S)} \leq \frac{1}{2}.$$

Here we note that M depends only on S and g . Therefore we deduce that

$$\|\phi_n e^{-M\cdot}\|_{L^r(0,S)} \leq 2\|h_n e^{-M\cdot}\|_{L^r(0,S)}$$

for $n \in \mathbb{N}$. We pass to the limit as $n \rightarrow \infty$. Note that $\phi_n \rightarrow \phi$ strongly in $L^1(0, S)$ and $h_n \rightarrow h$ strongly in $L^r(0, S)$. Hence as for $r > 1$, one finds that $\phi_n \rightarrow \phi$ weakly in $L^r(0, S)$ up to a subsequence. Thus for $r \in [1, \infty)$ we can reach

$$\|\phi e^{-M\cdot}\|_{L^r(0,S)} \leq 2\|h e^{-M\cdot}\|_{L^r(0,S)},$$

which yields the desired conclusion with the choice $C_0 = 2e^{MS}$. \square

¹The mollifier (ρ_n) may be supposed to enjoy the following properties: $\rho_n \in C_c^\infty([0, \infty))$, $\rho_n \geq 0$, $\int_0^\infty \rho_n(s) \, ds = 1$ and $\text{supp } \rho_n = [0, 1/n]$ for $n \in \mathbb{N}$. Then for $1 \leq p < \infty$ and $T \in (0, \infty)$, one can check in a standard way that $\rho_n * w \in L^\infty(0, T)$ and $\rho_n * w \rightarrow w$ strongly in $L^p(0, T)$ for any $w \in L^p(0, T)$.

5. PROOF OF THEOREM 2.3

In this section, we give a proof for Theorem 2.3. To this end, we first set up a Gronwall-type lemma for nonlinear Volterra integral inequalities.

LEMMA 5.1. *Let $S \in (0, \infty)$ and let $\phi \in L^\infty(0, S)$ be a nonnegative function. Suppose that there exist a constant $a \geq 0$, a nonnegative function $g \in L^1_{\text{loc}}([0, \infty))$ and a nondecreasing function $M : [0, \infty) \rightarrow [0, \infty)$ such that*

$$\phi(t) \leq a + [g * M(\phi(\cdot))](t) \quad \text{for a.e. } t \in (0, S).$$

Then there exists a constant $R > 0$ which depends only on $M(a + 1)$ and g such that

$$\|\phi\|_{L^\infty(0, \min\{R, S\})} \leq a + 1.$$

Here it is noteworthy that ϕ is not supposed to be continuous in Lemma 5.1. Actually, in order to prove Theorem 2.3, we shall apply Lemma 5.1 for some possibly *discontinuous* functions $\phi \in L^\infty(0, T)$ due to the subdiffusive nature of the problem.

Proof. We can assume $M > 0$ in $(0, \infty)$ without loss of generality, and we denote by the same letter the zero extension of ϕ onto $[0, \infty)$. Since $g \in L^1_{\text{loc}}([0, \infty))$, there exists a constant $R > 0$ such that

$$\int_0^R g(t) dt < \frac{1}{4M(a + 1)}.$$

Here R depends only on $M(a + 1)$ and g . Moreover, we note that

$$\begin{aligned} \phi(t) &\leq a + [g * M(\phi(\cdot))](t) \\ &\leq a + [g * M(\|\phi\|_{L^\infty(0, S)})](t) = a + \|g\|_{L^1(0, t)} M(\|\phi\|_{L^\infty(0, S)}) \end{aligned}$$

for a.e. $t \in (0, \infty)$. Hence we similarly observe that $\|\phi\|_{L^\infty(0, t)} \leq a + 1$ for $t > 0$ small enough. Set

$$T_\phi := \sup\{t \in (0, \infty) : \|\phi\|_{L^\infty(0, t)} \leq a + 1\} > 0.$$

We claim that $R \leq T_\phi$. Suppose to the contrary that $T_\phi < R$. Then since $g \in L^1_{\text{loc}}([0, \infty))$, there exists $t_\phi \in (T_\phi, R)$ such that

$$\begin{cases} a + 1 < \phi(t_\phi) \leq a + [g * M(\phi(\cdot))](t_\phi), \\ M(\|\phi\|_{L^\infty(0, S)}) \int_0^{t_\phi - T_\phi} g(t) dt < \frac{1}{4}. \end{cases} \quad (5.1)$$

Therefore we derive from the monotonicity of M and the choices of t_ϕ and T_ϕ that

$$\begin{aligned} a + 1 &< \phi(t_\phi) \leq a + [g * M(\phi(\cdot))](t_\phi) \\ &= a + \int_0^{T_\phi} g(t_\phi - t) M(\phi(t)) dt + \int_{T_\phi}^{t_\phi} g(t_\phi - t) M(\phi(t)) dt \end{aligned}$$

$$\begin{aligned}
 &\leq a + M(a + 1) \int_0^{T_\phi} g(t_\phi - t) dt + M (\|\phi\|_{L^\infty(0,S)}) \int_{T_\phi}^{t_\phi} g(t_\phi - t) dt \\
 &= a + M(a + 1) \int_{t_\phi - T_\phi}^{t_\phi} g(\tau) d\tau + M (\|\phi\|_{L^\infty(0,S)}) \int_0^{t_\phi - T_\phi} g(\tau) d\tau \\
 &\leq a + M(a + 1) \int_0^R g(\tau) d\tau + M (\|\phi\|_{L^\infty(0,S)}) \int_0^{t_\phi - T_\phi} g(\tau) d\tau \\
 &\leq a + M(a + 1) \frac{1}{4M(a + 1)} + \frac{1}{4} = a + \frac{1}{2},
 \end{aligned}$$

which yields a contradiction. Thus we conclude that $R \leq T_\phi$, which implies the desired conclusion. \square

We are now ready to prove Theorem 2.3.

Proof of Theorem 2.3. Thanks to the affine boundedness of proper lower-semicontinuous convex functionals (see, e.g., [9]), one can assume $\varphi^i \geq 0$ for $i = 1, 2$ without loss of generality.² Let $T \in (0, \infty)$ be fixed and consider the following approximate problems:

$$\frac{d}{dt} [k * (u_\lambda - u_0)](t) + \partial\varphi^1(u_\lambda(t)) - \partial\varphi_\lambda^2(u_\lambda(t)) \ni f(t) \quad \text{in } H, \quad 0 < t < T, \tag{5.2}$$

where φ_λ^2 stands for the Moreau–Yosida regularization of φ^2 , for $\lambda \in (0, 1)$ (see §3.1). Since $\partial\varphi_\lambda^2 : H \rightarrow H$ is Lipschitz continuous, due to Theorem 4.4, for $\lambda \in (0, 1)$, $u_0 \in D(\varphi^1)$ and $f \in L^2(0, T; H)$ satisfying $\ell * \|f(\cdot)\|_H^2 \in L^\infty(0, T)$, we can prove that the Cauchy problem (5.2) admits a strong solution $u_\lambda \in L^2(0, T; H)$ on $[0, T]$ such that

$$u_\lambda \in C([0, T]; H), \quad u_\lambda(0) = u_0, \quad \varphi^1(u_\lambda(\cdot)) \in L^\infty(0, T).$$

Let $g_\lambda \in L^2(0, T; H)$ be such that $g_\lambda(t) \in \partial\varphi^1(u_\lambda(t))$ and

$$\frac{d}{dt} [k * (u_\lambda - u_0)](t) + g_\lambda(t) - \partial\varphi_\lambda^2(u_\lambda(t)) = f(t) \tag{5.3}$$

for a.e. $t \in (0, T)$.

We next establish a priori estimates uniformly for $\lambda \in (0, 1)$. Test (5.3) by $g_\lambda(t)$ to see that

$$\begin{aligned}
 &\left(\frac{d}{dt} [k * (u_\lambda - u_0)](t), g_\lambda(t) \right)_H + \|g_\lambda(t)\|_H^2 \\
 &= (\partial\varphi_\lambda^2(u_\lambda(t)), g_\lambda(t))_H + (f(t), g_\lambda(t))_H \\
 &\leq \|\partial\varphi_\lambda^2(u_\lambda(t))\|_H \|g_\lambda(t)\|_H + \|f(t)\|_H \|g_\lambda(t)\|_H
 \end{aligned}$$

²Indeed, for each $i = 1, 2$, there exist $a_i \in \mathbb{R}$ and $z_i \in H$ such that $\varphi^i(w) \geq (z_i, w)_H + a_i$ for $w \in H$. Set $\hat{\varphi}^i(w) := \varphi^i(w) - (z_i, w)_H - a_i \geq 0$. We note that $\partial\hat{\varphi}^i(w) = \partial\varphi^i(w) - z_i$. Hence (1.1) is rewritten as $(d/dt)[k * (u - u_0)](t) + \partial\hat{\varphi}^1(u(t)) - \partial\hat{\varphi}^2(u(t)) \ni f(t) - z_1 + z_2 =: \hat{f}(t)$. Moreover, $\hat{\varphi}^1$ and $\hat{\varphi}^2$ fulfill (A1) and (A2) as well.

for a.e. $t \in (0, T)$. From (iv) of Proposition 3.1 and (A2), using Young's inequality, we find that

$$\begin{aligned} \|\partial\varphi_\lambda^2(u_\lambda(t))\|_H &\leq \|\partial\dot{\varphi}^2(u_\lambda(t))\|_H \\ &\leq \nu_1\|g_\lambda(t)\|_H + M_1(\varphi^1(u_\lambda(t)) + \|u_\lambda(t)\|_H) \\ &\leq \nu_1\|g_\lambda(t)\|_H \\ &\quad + M_1\left((1/2)\|u_\lambda(t) - u_0\|_H^2 + \varphi^1(u_\lambda(t)) + \|u_0\|_H + 1\right). \end{aligned} \quad (5.4)$$

Hence we have

$$\begin{aligned} &\left(\frac{d}{dt}[k * (u_\lambda - u_0)](t), g_\lambda(t)\right)_H + \|g_\lambda(t)\|_H^2 \\ &\leq \nu_1\|g_\lambda(t)\|_H^2 \\ &\quad + M_1\left((1/2)\|u_\lambda(t) - u_0\|_H^2 + \varphi^1(u_\lambda(t)) + \|u_0\|_H + 1\right)\|g_\lambda(t)\|_H \\ &\quad + \|f(t)\|_H\|g_\lambda(t)\|_H \end{aligned}$$

for a.e. $t \in (0, T)$. Therefore employing Young's inequality, one can take a constant $C_1 = C_1(\nu_1) > 0$ such that

$$\begin{aligned} &\left(\frac{d}{dt}[k * (u_\lambda - u_0)](t), g_\lambda(t)\right)_H + \frac{1 - \nu_1}{2}\|g_\lambda(t)\|_H^2 \\ &\leq C_1 M_1\left((1/2)\|u_\lambda(t) - u_0\|_H^2 + \varphi^1(u_\lambda(t)) + \|u_0\|_H + 1\right)^2 + C_1\|f(t)\|_H^2 \end{aligned}$$

for a.e. $t \in (0, T)$. Convolving both sides with ℓ and employing (ii) of Proposition 4.2, we have

$$\begin{aligned} &\varphi^1(u_\lambda(t)) - \varphi^1(u_0) + \frac{1 - \nu_1}{2}(\ell * \|g_\lambda(\cdot)\|_H^2)(t) \\ &\leq C_1\left[\ell * M_1\left((1/2)\|u_\lambda(\cdot) - u_0\|_H^2 + \varphi^1(u_\lambda(\cdot)) + \|u_0\|_H + 1\right)^2\right](t) \\ &\quad + C_1(\ell * \|f(\cdot)\|_H^2)(t) \end{aligned} \quad (5.5)$$

for a.e. $t \in (0, T)$.

Moreover, testing (5.3) by $u_\lambda(t) - u_0$, we derive from (5.4) that

$$\begin{aligned} &\left(\frac{d}{dt}[k * (u_\lambda - u_0)](t), u_\lambda(t) - u_0\right)_H + (g_\lambda(t), u_\lambda(t) - u_0)_H \\ &\leq \|\partial\varphi_\lambda^2(u_\lambda(t))\|_H\|u_\lambda(t) - u_0\|_H + \|f(t)\|_H\|u_\lambda(t) - u_0\|_H \end{aligned}$$

for a.e. $t \in (0, T)$. Recalling the definition of subdifferential and employing Young's inequality again, one can take a constant $C_2 = C_2(\nu_1) > 0$ such that

$$\begin{aligned} &\left(\frac{d}{dt}[k * (u_\lambda - u_0)](t), u_\lambda(t) - u_0\right)_H + \varphi^1(u_\lambda(t)) - \varphi^1(u_0) \\ &\leq \frac{1 - \nu_1}{4}\|g_\lambda(t)\|_H^2 \\ &\quad + C_2\left\{M_1\left((1/2)\|u_\lambda(t) - u_0\|_H^2 + \varphi^1(u_\lambda(t)) + \|u_0\|_H + 1\right)^2\right. \end{aligned}$$

$$+ \|f(t)\|_H^2 + (1/2)\|u_\lambda(t) - u_0\|_H^2 \Big\}$$

for a.e. $t \in (0, T)$. Convolving both sides with ℓ and applying (ii) of Proposition 4.2 to $\varphi(\cdot) := (1/2)\|\cdot - u_0\|_H^2$, we deduce from the nonnegativity of φ^1 that

$$\begin{aligned} & \frac{1}{2}\|u_\lambda(t) - u_0\|_H^2 \\ & \leq \|\ell\|_{L^1(0,T)}\varphi^1(u_0) + \frac{1-\nu_1}{4} (\ell * \|g_\lambda(\cdot)\|_H^2)(t) \\ & \quad + C_2 \left[\ell * M_1 \left((1/2)\|u_\lambda(\cdot) - u_0\|_H^2 + \varphi^1(u_\lambda(\cdot)) + \|u_0\|_H + 1 \right)^2 \right](t) \\ & \quad + C_2 (\ell * \|f(\cdot)\|_H^2)(t) + C_2 [\ell * (1/2)\|u_\lambda(\cdot) - u_0\|_H^2](t) \end{aligned} \quad (5.6)$$

for a.e. $t \in (0, T)$. Combining (5.5) and (5.6), we obtain

$$\begin{aligned} & \underbrace{\frac{1}{2}\|u_\lambda(t) - u_0\|_H^2 + \varphi^1(u_\lambda(t))}_{=: \phi(t)} + \frac{1-\nu_1}{4} (\ell * \|g_\lambda(\cdot)\|_H^2)(t) \\ & \leq a + \left[\ell * M \left(\underbrace{(1/2)\|u_\lambda(t) - u_0\|_H^2 + \varphi^1(u_\lambda(\cdot))}_{=\phi(\cdot)} \right) \right](t) \end{aligned} \quad (5.7)$$

for a.e. $t \in (0, T)$. Here a and M are given as

$$\begin{aligned} a & := (\|\ell\|_{L^1(0,T)} + 1) \varphi^1(u_0) + (C_1 + C_2) \|\ell * \|f(\cdot)\|_H^2\|_{L^\infty(0,T)}, \\ M(s) & := C_2 s + (C_1 + C_2) M_1 (s + \|u_0\|_H + 1)^2 \quad \text{for } s \in \mathbb{R}. \end{aligned}$$

Hence applying Lemma 5.1, we can take a constant $T_0 \in (0, T]$ such that

$$\operatorname{ess\,sup}_{t \in (0, T_0)} [(1/2)\|u_\lambda(t) - u_0\|_H^2 + \varphi^1(u_\lambda(t))] \leq C.$$

Here and henceforth, C denotes a generic constant being independent of t and λ . We also note that T_0 depends only on ℓ and $M(a + 1)$. Moreover, since u_λ belongs to $C([0, T]; H)$ and φ^1 is lower-semicontinuous in H , one can further verify that

$$\sup_{t \in [0, T_0]} [(1/2)\|u_\lambda(t) - u_0\|_H^2 + \varphi^1(u_\lambda(t))] \leq C. \quad (5.8)$$

Recalling (5.7), we also get

$$\operatorname{ess\,sup}_{t \in (0, T_0)} (\ell * \|g_\lambda(\cdot)\|_H^2)(t) \leq C,$$

which along with (K) yields

$$\int_0^{T_0} \|g_\lambda(t)\|_H^2 dt \leq C.$$

Moreover, we deduce from (A2) and (5.8) that

$$\operatorname{ess\,sup}_{t \in (0, T_0)} \left(\ell * \|\partial\varphi_\lambda^2(u_\lambda(\cdot))\|_H^2 \right)(t) \leq C,$$

which leads us to obtain

$$\int_0^{T_0} \|\partial\varphi_\lambda^2(u_\lambda(t))\|_H^2 dt \leq C. \quad (5.9)$$

It further follows from (5.2) that

$$\operatorname{ess\,sup}_{t \in (0, T_0)} \left(\ell * \left\| \frac{d}{dt} [k * (u_\lambda - u_0)](\cdot) \right\|_H^2 \right) (t) \leq C, \quad (5.10)$$

which also gives

$$\int_0^{T_0} \left\| \frac{d}{dt} [k * (u_\lambda - u_0)](t) \right\|_H^2 dt \leq C.$$

Furthermore, noting that

$$\left(\ell * \frac{d}{dt} [k * (u_\lambda - u_0)] \right) (t) = u_\lambda(t) - u_0 \quad \text{for } t \in (0, T), \quad (5.11)$$

we deduce from Lemma 4.3 along with (5.10) that

$$\sup_{\lambda \in (0, 1)} \sup_{t \in [0, T_0 - h]} \|u_\lambda(t + h) - u_\lambda(t)\|_H \rightarrow 0$$

as $h \rightarrow 0_+$. On the other hand, it follows from (5.8) and (A1) that $(u_\lambda(t))_{\lambda \in (0, 1)}$ is precompact in H for each $t \in [0, T_0]$.

Combining all these facts and using Ascoli's compactness lemma (see, e.g., [31, Lemma 1]), one can take a sequence (λ_n) in $(0, 1)$ converging to 0 such that

$$u_{\lambda_n} \rightarrow u \quad \text{strongly in } C([0, T_0]; H), \quad (5.12)$$

$$g_{\lambda_n} \rightarrow \xi \quad \text{weakly in } L^2(0, T_0; H), \quad (5.13)$$

$$\partial\varphi_{\lambda_n}^2(u_{\lambda_n}(\cdot)) \rightarrow \eta \quad \text{weakly in } L^2(0, T_0; H),$$

$$(d/dt)[k * (u_{\lambda_n} - u_0)] \rightarrow \zeta \quad \text{weakly in } L^2(0, T_0; H)$$

for some $u \in C([0, T_0]; H)$ and $\xi, \eta, \zeta \in L^2(0, T_0; H)$. Moreover, (5.12) along with $u_\lambda(0) = u_0$ also yields $u(0) = u_0$. Recall that the map $\partial\Phi^1 : u \mapsto \partial\varphi^1(u(\cdot))$ is maximal monotone in $L^2(0, T_0; H)$ (see [9, p. 25]) and every maximal monotone operator is demiclosed (see §3.1). Since (u_λ, g_λ) lies on the graph $G(\partial\Phi_1)$ of $\partial\Phi_1$ (equivalently, $(u_\lambda(t), g_\lambda(t)) \in G(\partial\varphi^1)$ for a.e. $t \in (0, T_0)$), we find from (5.12) and (5.13) that $(u, \xi) \in D(\partial\Phi_1)$, that is,

$$u(t) \in D(\partial\varphi^1) \quad \text{and} \quad \xi(t) \in \partial\varphi^1(u(t))$$

for a.e. $t \in (0, T_0)$. Furthermore, let $J_\lambda : H \rightarrow H$ be the resolvent of $\partial\varphi^2$, that is, $J_\lambda = (1 + \lambda\partial\varphi^2)^{-1}$, for $\lambda \in (0, 1)$. Note that

$$\|u_\lambda(t) - J_\lambda u_\lambda(t)\|_H = \lambda \|\partial\varphi_\lambda^2(u_\lambda(t))\|_H$$

for all $t \in [0, T]$. It follows from (5.9) that

$$u_\lambda - J_\lambda u_\lambda \rightarrow 0 \quad \text{strongly in } L^2(0, T_0; H) \quad \text{as } \lambda \rightarrow 0_+,$$

which along with (5.12) implies

$$J_{\lambda_n} u_{\lambda_n} \rightarrow u \quad \text{strongly in } L^2(0, T_0; H).$$

Recalling that $(J_\lambda u_\lambda(t), \partial\varphi_\lambda^2(u_\lambda(t)))$ lies on the graph of $\partial\varphi^2$ for a.e. $t \in (0, T_0)$, we can similarly conclude that

$$u(t) \in D(\partial\varphi^2) \quad \text{and} \quad \eta(t) \in \partial\varphi^2(u(t))$$

for a.e. $t \in (0, T_0)$. Moreover, due to the linear maximal monotonicity (in particular, weak closedness) of \mathcal{B} in $L^2(0, T_0; H)$, we further obtain

$$u - u_0 \in D(\mathcal{B}) \quad \text{and} \quad \zeta = \mathcal{B}(u - u_0) = \frac{d}{dt}[k * (u - u_0)],$$

the former of which yields the initial condition, that is, $k * (u - u_0) \in W^{1,2}(0, T_0; H)$ and $[k * (u - u_0)](0) = 0$. Finally, from (5.8) along with the lower-semicontinuity of φ^1 on H , one can verify that

$$\varphi^1(u(\cdot)) \in L^\infty(0, T_0).$$

This completes the proof. \square

6. PROOFS OF THEOREM 2.6 AND COROLLARY 2.8

This section is devoted to proving Theorem 2.6 and Corollary 2.8. To this end, we first set up the following lemma, which is a variant of Lemma 5.1.

LEMMA 6.1. *Let $S \in (0, \infty)$ and let $\phi \in L^\infty(0, S)$ be a nonnegative function. Suppose that there exist constants $\delta > b \geq 0$, a nonnegative function $g \in L^1(0, S)$ and a locally bounded Borel measurable function $N : [0, \infty) \rightarrow \mathbb{R}$ satisfying*

$$N(r) \leq 0 \quad \text{for } r \in [0, \delta] \tag{6.1}$$

such that

$$\phi(t) \leq b + [g * N(\phi(\cdot))](t) \quad \text{for a.e. } t \in (0, S).$$

Then $\|\phi\|_{L^\infty(0, S)} \leq b$.

Proof. Fix $\epsilon > 0$ satisfying $0 < \epsilon < (\delta - b)/2$ (hence $b + 2\epsilon < \delta$). As in the proof of Lemma 5.1, we set

$$T_{\phi, \epsilon} := \sup\{s \in (0, S) : \|\phi\|_{L^\infty(0, s)} \leq b + 2\epsilon\} > 0.$$

From the arbitrariness of $\epsilon > 0$, it suffices to show that $T_{\phi, \epsilon} = S$. Suppose to the contrary that $T_{\phi, \epsilon} < S$. Then there exists $t_{\phi, \epsilon} \in (T_{\phi, \epsilon}, S)$ such that

$$b + 2\epsilon < \phi(t_{\phi, \epsilon}) \leq b + [g * N(\phi(\cdot))](t_{\phi, \epsilon}),$$

$$\|N\|_{L^\infty(0, R_\phi)} \int_0^{t_{\phi, \epsilon} - T_{\phi, \epsilon}} g(t) dt < \epsilon,$$

where $R_\phi := \|\phi\|_{L^\infty(0, S)}$. As $\phi(t) \leq \delta$ for a.e. $t \in (0, T_{\phi, \epsilon})$, we find from (6.1) that

$$b + 2\epsilon < \phi(t_{\phi, \epsilon}) \leq b + [g * N(\phi(\cdot))](t_{\phi, \epsilon})$$

$$\begin{aligned}
&= b + \int_0^{t_{\phi,\epsilon}} g(t_{\phi,\epsilon} - t)N(\phi(t)) dt \\
&\leq b + \int_{T_{\phi,\epsilon}}^{t_{\phi,\epsilon}} g(t_{\phi,\epsilon} - t)N(\phi(t)) dt \\
&\leq b + \|N\|_{L^\infty(0,R_\phi)} \int_0^{t_{\phi,\epsilon}-T_{\phi,\epsilon}} g(t) dt < b + \epsilon,
\end{aligned}$$

which yields a contradiction. Thus we obtain $S = T_{\phi,\epsilon}$. The desired conclusion follows from the arbitrariness of $\epsilon > 0$. \square

We are in a position to prove Theorem 2.6.

Proof of Theorem 2.6. Let $T \in (0, \infty)$ be fixed. Set

$$E_T(u_0, f) := \varphi^1(u_0) + \|\ell * \|f(\cdot)\|_H^2\|_{L^\infty(0,T)}$$

and recall the approximate problems (5.2), which admit strong solutions u_λ on $[0, T]$ such that $u_\lambda \in C([0, T]; H)$, $u_\lambda(0) = u_0$ and $\varphi^1(u_\lambda(\cdot)) \in L^\infty(0, T)$. Moreover, let $g_\lambda \in L^2(0, T; H)$ be such that $g_\lambda(t) \in \partial\varphi^1(u_\lambda(t))$ and (5.3) holds for a.e. $t \in (0, T)$.

From (iv) of Proposition 3.1 and (A2)', we have

$$\|\partial\varphi_\lambda^2(u_\lambda(t))\|_H \leq \|\partial\varphi^2(u_\lambda(t))\|_H \leq \nu_2 \|g_\lambda(t)\|_H + M_2(\varphi^1(u_\lambda(t))) \quad (6.2)$$

for a.e. $t \in (0, T)$. By the use of (6.2) instead of (5.4), testing (5.3) by $g_\lambda(t)$ and convolving it with ℓ , as in the proof of Theorem 2.3, we can derive

$$\begin{aligned}
&\varphi^1(u_\lambda(t)) - \varphi^1(u_0) + \frac{1 - \nu_2}{2} (\ell * \|g_\lambda(\cdot)\|_H^2)(t) \\
&\leq C_1 \left[\ell * M_2(\varphi^1(u_\lambda(\cdot)))^2 \right](t) + C_1 \|\ell * \|f(\cdot)\|_H^2\|_{L^\infty(0,T)}
\end{aligned}$$

for a.e. $t \in (0, T)$ and a constant $C_1 = C_1(\nu_2) \geq 0$ (see (5.5)). Thus one can take a constant $C_3 = C_3(\nu_2) \geq 1$ such that

$$\begin{aligned}
&\varphi^1(u_\lambda(t)) + \frac{1 - \nu_2}{2} (\ell * \|g_\lambda(\cdot)\|_H^2)(t) \\
&\leq C_3 E_T(u_0, f) + C_3 \left[\ell * M_2(\varphi^1(u_\lambda(\cdot)))^2 \right](t)
\end{aligned}$$

for a.e. $t \in (0, T)$. Here we note from (2.3) that

$$\|g_\lambda(t)\|_H \geq m_2(\varphi^1(u_\lambda(t)))$$

for a.e. $t \in (0, T)$. Hence it follows that

$$\begin{aligned}
&\varphi^1(u_\lambda(t)) + \frac{1 - \nu_2}{4} (\ell * \|g_\lambda(\cdot)\|_H^2)(t) \\
&\leq C_3 E_T(u_0, f) + [\ell * N(\varphi^1(u_\lambda(\cdot)))](t) \quad \text{for a.e. } t \in (0, T), \quad (6.3)
\end{aligned}$$

where N is a function given by

$$N(r) := C_3 M_2(r)^2 - \frac{1 - \nu_2}{4} m_2(r)^2 \quad \text{for } r \geq 0.$$

Here due to (2.2) one can take a constant $\delta_0 > 0$ small enough depending only on ν_2 , $m_2(\cdot)$ and $M_2(\cdot)$ that $N(r) \leq 0$ for any $r \in [0, \delta_0]$. Thanks to Lemma 6.1, we obtain

$$\sup_{t \in [0, T]} \varphi^1(u_\lambda(t)) \leq C_3 E_T(u_0, f), \quad (6.4)$$

provided that $C_3 E_T(u_0, f) < \delta_0$ (cf. see (5.8)). Furthermore, it follows from (6.3) that

$$\operatorname{ess\,sup}_{t \in (0, T)} (\ell * \|g_\lambda(\cdot)\|_H^2)(t) \leq C, \quad (6.5)$$

which in particular implies

$$\int_0^T \|g_\lambda(t)\|_H^2 dt \leq C.$$

Here and henceforth, C denotes a generic constant being independent of t and λ . Hence combining (2.4) with (6.4) and (6.5), we deduce that

$$\operatorname{ess\,sup}_{t \in (0, T)} (\ell * \|\partial\varphi_\lambda^2(u_\lambda(\cdot))\|_H^2)(t) \leq C,$$

which also yields

$$\int_0^T \|\partial\varphi_\lambda^2(u_\lambda(t))\|_H^2 dt \leq C,$$

and hence, (5.3) gives

$$\operatorname{ess\,sup}_{t \in (0, T)} \left(\ell * \left\| \frac{d}{dt} [k * (u_\lambda - u_0)](\cdot) \right\|_H^2 \right)(t) \leq C.$$

Due to (6.4) and (A1)', one can also deduce that $(u_\lambda(t))_{\lambda \in (0, 1)}$ is precompact in H for each $t \in [0, T]$. Hence repeating the same argument as in the proof of Theorem 2.3, we can conclude that

$$u_{\lambda_n} \rightarrow u \quad \text{strongly in } C([0, T]; H)$$

for some sequence $\lambda_n \rightarrow 0_+$ and $u \in C([0, T]; H)$ complying with $u(0) = u_0$, and moreover, the limit u turns out to be a strong solution on $[0, T]$ to (1.1). Furthermore, (6.4) and the lower-semicontinuity of φ^1 yield

$$\varphi^1(u(t)) \leq C_3 E_T(u_0, f)$$

for all $t \in [0, T]$.

Finally, we show existence of strong solutions on $[0, \infty)$ to the Cauchy problem (1.1) with $T = \infty$ for $u_0 \in D(\varphi^1)$ and $f \in L_{\text{loc}}^2([0, \infty); H)$ satisfying $C_3 E_\infty(u_0, f) < \delta_0$. Thanks to Theorem 4.4, for each $\lambda > 0$ the approximate problem (5.2) with $T = \infty$ admits a unique strong solution $u_\lambda \in L_{\text{loc}}^2([0, \infty); H)$ on $[0, \infty)$ satisfying (4.7) for any $T > 0$. From the arguments so far, for each $N \in \mathbb{N}$, one can inductively take a subsequence $(\lambda_n^{(N)})_{n \in \mathbb{N}}$ of $(\lambda_n^{(N-1)})_{n \in \mathbb{N}}$ such that

$$u_{\lambda_n^{(N)}} \rightarrow u^{(N)} \quad \text{strongly in } C([0, N]; H)$$

for some strong solution $u^{(N)} \in C([0, N]; H)$ on $[0, N]$ to (1.1) with $T = N$ and $u^{(N)}(0) = u_0$. Then one observes by definition that the restriction $u^{(N)}|_{[0, N-1]}$ coincides with $u^{(N-1)}$. Due to a diagonal argument, one can take a (not relabeled) subsequence of $(\lambda_n^{(n)})_{n \in \mathbb{N}}$ such that

$$u_{\lambda_n^{(n)}} \rightarrow u \quad \text{strongly in } C([0, T]; H) \quad \text{for any } T > 0$$

for some $u \in C([0, \infty); H)$ satisfying $u(0) = u_0$. Moreover, recalling the relation $g_\lambda := f - (\text{d}/\text{d}t)[k * (u_\lambda - u_0)] + \partial\varphi_\lambda^2(u_\lambda(\cdot))$ (see (5.3) with $T = \infty$) along with the uniqueness of u_λ , one can construct $\xi, \eta \in L_{\text{loc}}^2([0, \infty); H)$ satisfying (2.1) for a.e. $t \in (0, \infty)$. Indeed, for each $N \in \mathbb{N}$, one can similarly construct $\xi^{(N)}, \eta^{(N)} \in L^2(0, N; H)$ such that (2.1) holds for a.e. $t \in (0, N)$, $\xi^{(N)}|_{[0, N-1]} = \xi^{(N-1)}$ and $\eta^{(N)}|_{[0, N-1]} = \eta^{(N-1)}$. Define a strongly measurable function $\xi : (0, \infty) \rightarrow H$ as a limit

$$\xi(t) := \lim_{N \rightarrow \infty} \bar{\xi}^{(N)}(t) \quad \text{for a.e. } t \in (0, \infty),$$

where $\bar{\xi}^{(N)} \in L^2(0, \infty; H)$ is a function given as

$$\bar{\xi}^{(N)}(t) = \begin{cases} \xi^{(N)}(t) & \text{if } t \in [0, N], \\ 0 & \text{if } t \in (N, \infty) \end{cases} \quad \text{for } t \in [0, \infty).$$

Then we observe that $\xi|_{[0, N]} = \xi^{(N)}$ for $N \in \mathbb{N}$. Hence $\xi \in L_{\text{loc}}^2([0, \infty); H)$. By a similar argument, we can also construct $\eta \in L_{\text{loc}}^2([0, \infty); H)$ such that $\eta|_{[0, N]} = \eta^{(N)}$ for $N \in \mathbb{N}$. Therefore u is a strong solution to (1.1) on $[0, \infty)$ in the sense of Definition 2.1, since $u|_{[0, N]} = u^{(N)}$ for each N . This completes the proof. \square

We now move on to a proof of Corollary 2.8.

Proof of Corollary 2.8. Let $T \in (0, \infty)$ be fixed. Due to (2.8), one can take constants $\delta_0, \epsilon > 0$ small enough depending only on $M_3(\cdot)$ that

$$0 < M_3(r + \epsilon) \leq \frac{1}{2} \quad \text{for } r \in [0, \delta_0].$$

Testing (5.3) by $g_\lambda(t)$, using Young's inequality, and convolving it with ℓ , one can deduce that

$$\begin{aligned} & \varphi^1(u_\lambda(t)) - \varphi^1(u_0) + \frac{1}{2} (\ell * \|g_\lambda(\cdot)\|_H^2)(t) \\ & \leq (\ell * \|\partial\varphi_\lambda^2(u_\lambda(\cdot))\|_H^2)(t) + \|\ell * \|f(\cdot)\|_H^2\|_{L^\infty(0, T)} \end{aligned}$$

for a.e. $t \in (0, T)$. From (iv) of Proposition 3.1 and (2.9) of (A2)'_c, we have

$$\|\partial\varphi_\lambda^2(u_\lambda(t))\|_H \leq \|\partial\varphi^2(u_\lambda(t))\|_H \leq \|g_\lambda(t)\|_H M_3(\varphi^1(u_\lambda(t))),$$

which yields

$$\|\partial\varphi_\lambda^2(u_\lambda(t))\|_H M_3(\varphi^1(u_\lambda(t)) + \epsilon)^{-1} \leq \|g_\lambda(t)\|_H \quad (6.6)$$

for a.e. $t \in (0, T)$. Using (6.6), we obtain

$$\begin{aligned} & \varphi^1(u_\lambda(t)) + \frac{1}{4} (\ell * \|g_\lambda(\cdot)\|_H^2)(t) \\ & \leq E_T(u_0, f) + [\ell * N(\varphi^1(u_\lambda(\cdot)))](t) \end{aligned}$$

for a.e. $t \in (0, T)$. Here N is a function given by

$$N(r) := \begin{cases} 0 & \text{if } 1 - (1/4)M_3(r + \epsilon)^{-2} \leq 0, \\ \|\partial\varphi_\lambda^2(u_\lambda(\cdot))\|_{L^\infty(0, T; H)}^2 & \text{if } 1 - (1/4)M_3(r + \epsilon)^{-2} > 0 \end{cases}$$

for $r \geq 0$. Then we note that $N(r) \leq 0$ for any $r \in [0, \delta_0]$. Thanks to Lemma 6.1, we obtain

$$\operatorname{ess\,sup}_{t \in (0, T)} \varphi^1(u_\lambda(t)) \leq E_T(u_0, f),$$

provided that $E_T(u_0, f) < \delta_0$. The rest of the proof runs as before. \square

7. PROOF OF THEOREM 2.9

In this section, we prove Theorem 2.9.

Proof of Theorem 2.9. Let $T \in (0, \infty)$ be fixed. In this proof, we consider the following (slightly different) approximate problems:

$$\left. \begin{aligned} & \lambda \frac{du_\lambda}{dt}(t) + \frac{d}{dt} [k * (u_\lambda - u_0)](t) + \partial\varphi^1(u_\lambda(t)) - \partial\varphi_\lambda^2(u_\lambda(t)) \ni f(t), \\ & 0 < t < T, \\ & u_\lambda(0) = u_0, \end{aligned} \right\} \quad (7.1)$$

where φ_λ^2 stands for the Moreau–Yosida regularization of φ^2 , for $\lambda \in (0, 1)$. For $u_0 \in D(\varphi^1)$, $f \in W^{1,2}(0, T; H)$ and $\lambda \in (0, 1)$, one can prove existence of a unique strong solution $u_\lambda \in W^{1,2}(0, T; H)$ on $[0, T]$ to the Cauchy problem (5.2) as in [3] (see §A in Appendix). Testing (7.1) by $(du_\lambda/dt)(t)$ and employing Proposition 3.3, we have

$$\begin{aligned} & \lambda \left\| \frac{du_\lambda}{dt}(t) \right\|_H^2 + \left(\frac{d}{dt} [k * (u_\lambda - u_0)](t), \frac{du_\lambda}{dt}(t) \right)_H \\ & \quad + \frac{d}{dt} [\varphi^1(u_\lambda(t)) - \varphi_\lambda^2(u_\lambda(t))] \\ & = \left(f(t), \frac{du_\lambda}{dt}(t) \right)_H \\ & = \frac{d}{dt} (f(t), u_\lambda(t))_H - \left(\frac{df}{dt}(t), u_\lambda(t) \right)_H \end{aligned}$$

for a.e. $t \in (0, T)$. Note that

$$\int_0^t \left(\frac{d}{dt} [k * (u_\lambda - u_0)](\tau), \frac{du_\lambda}{dt}(\tau) \right)_H d\tau$$

$$\geq \frac{1}{2} \left(\ell * \left\| \frac{d}{dt} [k * (u_\lambda - u_0)] (\cdot) \right\|_H^2 \right) (t)$$

for all $t \in [0, T]$ (see Proposition 3.4). Integrating both sides over $(0, t)$, we find that

$$\begin{aligned} & \lambda \int_0^t \left\| \frac{du_\lambda}{dt}(\tau) \right\|_H^2 d\tau + \frac{1}{2} \left(\ell * \left\| \frac{d}{dt} [k * (u_\lambda - u_0)] (\cdot) \right\|_H^2 \right) (t) \\ & \quad + \varphi^1(u_\lambda(t)) - \varphi_\lambda^2(u_\lambda(t)) \\ & \leq \varphi^1(u_0) - \varphi_\lambda^2(u_0) + (f(t), u_\lambda(t))_H - (f(0), u_0)_H \\ & \quad - \int_0^t \left(\frac{df}{dt}(\tau), u_\lambda(\tau) \right)_H d\tau \end{aligned}$$

for $t \in [0, T]$. Using (A3), we observe that

$$\begin{aligned} \varphi^1(u_\lambda(t)) - \varphi_\lambda^2(u_\lambda(t)) & \geq \varphi^1(u_\lambda(t)) - \varphi^2(u_\lambda(t)) \\ & \geq (1 - \nu_3) \varphi^1(u_\lambda(t)) - c_1 (\|u_\lambda(t)\|_H^r + 1) \end{aligned}$$

for $t \in [0, T]$. Moreover, exploiting Jensen's inequality, we note that

$$\begin{aligned} & \left(\ell * \left\| \frac{d}{dt} [k * (u_\lambda - u_0)] (\cdot) \right\|_H^2 \right) (t) \\ & = \int_0^t \ell(t-s) \left\| \frac{d}{dt} [k * (u_\lambda - u_0)] (s) \right\|_H^2 ds \\ & \geq \|\ell\|_{L^1(0,T)}^{-1} \left\| \left(\ell * \frac{d}{dt} [k * (u_\lambda - u_0)] \right) (t) \right\|_H^2 \\ & \stackrel{(5.11)}{=} \|\ell\|_{L^1(0,T)}^{-1} \|u_\lambda(t) - u_0\|_H^2 \end{aligned}$$

for $t \in [0, T]$. Thus we obtain

$$\begin{aligned} & \lambda \int_0^t \left\| \frac{du_\lambda}{dt}(\tau) \right\|_H^2 d\tau + \frac{1}{4} \left(\ell * \left\| \frac{d}{dt} [k * (u_\lambda - u_0)] (\cdot) \right\|_H^2 \right) (t) \\ & \quad + \frac{1}{4} \|\ell\|_{L^1(0,T)}^{-1} \|u_\lambda(t) - u_0\|_H^2 + (1 - \nu_3) \varphi^1(u_\lambda(t)) \\ & \leq \varphi^1(u_0) - \varphi_\lambda^2(u_0) + c_1 (\|u_\lambda(t)\|_H^r + 1) + (f(t), u_\lambda(t))_H - (f(0), u_0)_H \\ & \quad - \int_0^t \left(\frac{df}{dt}(\tau), u_\lambda(\tau) \right)_H d\tau \end{aligned}$$

for $t \in [0, T]$. Therefore exploiting Young's inequality, one finds that

$$\begin{aligned} & \lambda \int_0^t \left\| \frac{du_\lambda}{dt}(\tau) \right\|_H^2 d\tau + \frac{1}{4} \left(\ell * \left\| \frac{d}{dt} [k * (u_\lambda - u_0)] (\cdot) \right\|_H^2 \right) (t) \\ & \quad + \frac{1}{8} \|\ell\|_{L^1(0,T)}^{-1} \|u_\lambda(t) - u_0\|_H^2 + (1 - \nu_3) \varphi^1(u_\lambda(t)) \end{aligned}$$

$$\begin{aligned} &\leq \varphi^1(u_0) - \varphi_\lambda^2(u_0) + C (\|f(t)\|_H^2 + \|f(0)\|_H^2 + \|u_0\|_H^2 + 1) \\ &\quad + \int_0^t \left\| \frac{df}{dt}(\tau) \right\|_H \|u_\lambda(\tau)\|_H d\tau \end{aligned}$$

for $t \in [0, T]$. Here and henceforth, C denotes a generic constant being independent of t and λ . Since $\varphi_\lambda^2(u_0)$ is bounded as $\lambda \rightarrow 0_+$, using Gronwall's inequality, we conclude that

$$\sup_{t \in [0, T]} (\varphi^1(u_\lambda(t)) + \|u_\lambda(t)\|_H^2) \leq C, \quad (7.2)$$

$$\begin{aligned} &\lambda \int_0^T \left\| \frac{du_\lambda}{dt}(\tau) \right\|_H^2 d\tau \leq C, \\ &\sup_{t \in [0, T]} \left(\ell * \left\| \frac{d}{dt} [k * (u_\lambda - u_0)](\cdot) \right\|_H^2 \right) (t) \leq C \end{aligned} \quad (7.3)$$

for $\lambda \in (0, 1)$. Furthermore, since φ^1 is affinely bounded from below, we infer that

$$\sup_{t \in [0, T]} |\varphi^1(u_\lambda(t))| \leq C.$$

Hence as in the proof of Theorem 2.3, we can deduce from (7.3) along with Lemma 4.3, (A1) and Ascoli's theorem that

$$u_{\lambda_n} \rightarrow u \quad \text{strongly in } C([0, T]; H)$$

for some sequence $\lambda_n \rightarrow 0_+$, and therefore, we have $u(0) = u_0$. Moreover, the convolution of (7.3) with k also yields

$$\int_0^T \left\| \frac{d}{dt} [k * (u_\lambda - u_0)](t) \right\|_H^2 dt \leq C.$$

Furthermore, it follows from (A2) and (7.2) that

$$\|\partial \varphi_\lambda^2(u_\lambda(t))\|_H \leq \nu_1 \|g_\lambda(t)\|_H + C \quad (7.4)$$

for a.e. $t \in (0, T)$. Testing (7.1) by $g_\lambda(t)$, one finds that

$$\begin{aligned} &\lambda \frac{d}{dt} \varphi^1(u_\lambda(t)) + \left(\frac{d}{dt} [k * (u_\lambda - u_0)](t), g_\lambda(t) \right)_H + \|g_\lambda(t)\|_H^2 \\ &\quad \leq \|\partial \varphi_\lambda^2(u_\lambda(t))\|_H \|g_\lambda(t)\|_H + \|f(t)\|_H \|g_\lambda(t)\|_H \\ &\quad \leq \nu_1 \|g_\lambda(t)\|_H^2 + C \|g_\lambda(t)\|_H + \|f(t)\|_H \|g_\lambda(t)\|_H \end{aligned}$$

for a.e. $t \in (0, T)$. With the aid of Young's inequality, we have

$$\begin{aligned} &\lambda \frac{d}{dt} \varphi^1(u_\lambda(t)) + \left(\frac{d}{dt} [k * (u_\lambda - u_0)](t), g_\lambda(t) \right)_H + \frac{1 - \nu_1}{2} \|g_\lambda(t)\|_H^2 \\ &\quad \leq C (\|f(t)\|_H^2 + 1) \end{aligned}$$

for a.e. $t \in (0, T)$. Integrating both sides over $(0, T)$ and using (i) of Proposition 4.2, we conclude that

$$\begin{aligned} & \lambda \varphi^1(u_\lambda(t)) + [k * (\varphi^1(u_\lambda(\cdot)) - \varphi^1(u_0))](t) + \frac{1 - \nu_1}{2} \int_0^t \|g_\lambda(t)\|_H^2 dt \\ & \leq C \left(\int_0^T \|f(t)\|_H^2 dt + 1 \right) \quad \text{for } t \in [0, T], \end{aligned}$$

which along with (7.4) implies

$$\int_0^T \|g_\lambda(t)\|_H^2 dt + \int_0^T \|\partial \varphi_\lambda^2(u_\lambda(t))\|_H^2 dt \leq C.$$

The rest of the proof runs as in §6. \square

8. APPLICATIONS

In this section, we apply the abstract theory developed so far in the present paper to the Cauchy–Dirichlet problem for p -Laplace subdiffusion equations with blow-up terms of the form,

$$\partial_t^\alpha(u - u_0) - \Delta_p u - |u|^{q-2}u = f \quad \text{in } \Omega \times (0, \infty), \quad (8.1)$$

$$u = 0 \quad \text{on } \partial\Omega \times (0, \infty), \quad (8.2)$$

where $0 < \alpha < 1$, $1 < p, q < \infty$, Ω is a bounded domain of \mathbb{R}^d with smooth boundary $\partial\Omega$ and $u_0 = u_0(x)$ and $f = f(x, t)$ are prescribed. Moreover, $\partial_t^\alpha(u - u_0) := \partial_t[k_\alpha * (u - u_0)]$ denotes the Riemann–Liouville derivative of $u - u_0$ (i.e., Caputo derivative of u) of order $0 < \alpha < 1$ and Δ_p is the so-called p -Laplacian given as

$$\Delta_p u = \nabla \cdot (|\nabla u|^{p-2} \nabla u).$$

Here and henceforth, we are concerned with L^2 -solutions to (8.1), (8.2) defined as follows:

DEFINITION 8.1 (L^2 -solutions to (8.1), (8.2)). For each $S \in (0, \infty)$, a function $u \in L^2(0, S; L^2(\Omega))$ is called an L^2 -solution on $[0, S]$ to the Cauchy–Dirichlet problem (8.1), (8.2), if the following conditions are all satisfied:

- (i) It holds that $k_\alpha * (u - u_0) \in W^{1,2}(0, S; L^2(\Omega))$, $[k_\alpha * (u - u_0)](0) = 0$ and $u(t) \in W_0^{1,p}(\Omega) \cap L^q(\Omega)$ for a.e. $t \in (0, S)$.
- (ii) It holds that $\Delta_p u, |u|^{q-2}u \in L^2(0, S; L^2(\Omega))$ and

$$\frac{d}{dt} [k_\alpha * (u - u_0)](\cdot, t) - \Delta_p u(\cdot, t) - |u|^{q-2}u(\cdot, t) = f(\cdot, t) \quad \text{in } L^2(\Omega)$$

for a.e. $t \in (0, S)$.

Moreover, a function $u \in L_{\text{loc}}^2([0, \infty); L^2(\Omega))$ is called an L^2 -solution on $[0, \infty)$ to the Cauchy–Dirichlet problem (8.1), (8.2), if the restriction $u|_{[0, S]}$ of u is an L^2 -solution on $[0, S]$ to (8.1), (8.2) for every $S > 0$.

We set $H = L^2(\Omega)$ and define $\varphi^1, \varphi^2 : H \rightarrow [0, \infty]$ by

$$\varphi^1(w) = \begin{cases} \frac{1}{p} \int_{\Omega} |\nabla w|^p dx & \text{if } w \in W_0^{1,p}(\Omega), \\ \infty & \text{otherwise,} \end{cases}$$

$$\varphi^2(w) = \begin{cases} \frac{1}{q} \int_{\Omega} |w|^q dx & \text{if } w \in L^q(\Omega), \\ \infty & \text{otherwise} \end{cases}$$

for $w \in H = L^2(\Omega)$. Then φ^1 and φ^2 turn out to be proper, lower-semicontinuous and convex in H . Moreover, $\partial\varphi^1(w)$ and $\partial\varphi^2(w)$ coincide with $-\Delta_p w$ equipped with the homogeneous Dirichlet boundary condition in the distributional sense for $w \in D(\partial\varphi^1) = \{w \in W_0^{1,p}(\Omega) \cap L^2(\Omega) : \Delta_p w \in L^2(\Omega)\}$ and $|w|^{q-2}w$ for $w \in D(\partial\varphi^2) = \{w \in L^2(\Omega) \cap L^q(\Omega) : |w|^{q-2}w \in L^2(\Omega)\}$, respectively. Hence the Cauchy–Dirichlet problem (8.1), (8.2) is reduced into the abstract Cauchy problem,

$$\frac{d}{dt} [k_{\alpha} * (u - u_0)](t) + \partial\varphi^1(u(t)) - \partial\varphi^2(u(t)) = f(t) \quad \text{in } H \quad (8.3)$$

for $t \in (0, \infty)$.

We now set up the following:

LEMMA 8.2 (Calderón–Zygmund estimates for p -Laplacian). *Let Ω be a bounded $C^{1,1}$ domain of \mathbb{R}^d with boundary $\partial\Omega$. Let $p \in (1, \infty)$ be such that $p \geq 2d/(d+2)$. Let $r = 2^*(p-1)$ for $d \geq 3$ and $r \in (p, \infty)$ arbitrarily for $d \leq 2$. Then for each $f \in L^2(\Omega)$, the Dirichlet problem*

$$-\Delta_p u = f \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega \quad (8.4)$$

admits a unique weak solution $u \in W_0^{1,p}(\Omega)$ such that $\nabla u \in L^r(\Omega; \mathbb{R}^d)$. Moreover, there exists a constant $C \geq 0$ depending only on d, p and $\partial\Omega$ such that

$$\|\nabla u\|_{L^r(\Omega; \mathbb{R}^d)}^{p-1} \leq C \|f\|_{L^2(\Omega)}$$

for any $f \in L^2(\Omega)$. Here $u \in W_0^{1,p}(\Omega)$ denotes the unique weak solution to (8.4).

Proof. We only treat the case where $d \geq 3$, but the case where $d \leq 2$ can also be treated analogously with the continuous embedding $H^1(\Omega) \hookrightarrow L^\rho(\Omega)$ for any $1 \leq \rho < \infty$. Let $\phi \in H^2(\Omega) \cap H_0^1(\Omega)$ be the unique strong solution to the Dirichlet problem,

$$\Delta\phi = f \quad \text{in } \Omega, \quad \phi = 0 \quad \text{on } \partial\Omega.$$

Then it follows from (8.4) that

$$-\Delta_p u = f = \operatorname{div}(\nabla\phi) = \operatorname{div}(|F|^{p-2}F), \quad u = 0 \quad \text{on } \partial\Omega,$$

where $F \in L^{2^*(p-1)}(\Omega; \mathbb{R}^d)$ is given by

$$F := |\nabla\phi|^{p'-2} \nabla\phi.$$

Here we note that $p \leq 2^*(p-1)$ is equivalent to $p \geq 2d/(d+2)$. Therefore by virtue of nonlinear Calderón–Zygmund estimates established in [19, Theorem 1.6], we obtain $\nabla u \in L^{2^*(p-1)}(\Omega; \mathbb{R}^d)$ and

$$\|\nabla u\|_{L^{2^*(p-1)}(\Omega; \mathbb{R}^d)} \leq C\|F\|_{L^{2^*(p-1)}(\Omega; \mathbb{R}^d)} = C\|\nabla \phi\|_{L^{2^*(p-1)}(\Omega; \mathbb{R}^d)}^{p'-1}$$

for some constant $C \geq 0$ independent of f . Furthermore, we observe that

$$\|\nabla \phi\|_{L^{2^*(p-1)}(\Omega; \mathbb{R}^d)} \lesssim \|\nabla \phi\|_{H^1(\Omega; \mathbb{R}^d)} \lesssim \|\phi\|_{H^2(\Omega)} \lesssim \|\Delta \phi\|_{L^2(\Omega)} = \|f\|_{L^2(\Omega)}$$

due to (linear) Calderón–Zygmund estimates. Here $L \lesssim R$ means $L \leq CR$ for some constant $C \geq 0$ independent of f . Thus the desired conclusion follows. \square

We first apply Theorem 2.3 to obtain local (in time) existence of L^2 -solutions to the Cauchy–Dirichlet problem (8.1), (8.2).

THEOREM 8.3 (Local existence). *Suppose that*

$$p > \frac{2d}{d+2} \quad \text{and} \quad q < p^* := \frac{dp}{(d-p)_+} \quad (8.5)$$

and let $T \in (0, \infty)$. For every $u_0 \in W_0^{1,p}(\Omega)$ and $f \in L^2(0, T; L^2(\Omega))$ satisfying

$$\operatorname{ess\,sup}_{t \in (0, T)} \left(\int_0^t k_{1-\alpha}(t-s) \|f(\cdot, s)\|_{L^2(\Omega)}^2 ds \right) < \infty,$$

there exists $T_0 \in (0, \infty)$ such that the Cauchy–Dirichlet problem (8.1), (8.2) admits an L^2 -solution $u \in C([0, T_0]; L^2(\Omega)) \cap L^\infty(0, T_0; W_0^{1,p}(\Omega))$ on $[0, T_0]$ such that $u(\cdot, 0) = u_0$.

Proof. We note that $k = k_\alpha$ satisfies (K) with $\ell = k_{1-\alpha}$ (see §1). We next check (A1) and (A2) for φ^1 and φ^2 defined above. First of all, (A1) follows immediately from Rellich–Kondrachov’s theorem (i.e., $W_0^{1,p}(\Omega)$ is compactly embedded in $L^2(\Omega)$, provided that $p > 2d/(d+2)$). We next check (A2). In case $2(q-1) \leq p^*$, one finds that

$$\|\partial \varphi^2(w)\|_H = \| |w|^{q-2} w \|_{L^2(\Omega)} \leq C \|\nabla w\|_{L^p(\Omega; \mathbb{R}^d)}^{q-1} = C \{\varphi^1(w)\}^{(q-1)/p}$$

for $w \in W_0^{1,p}(\Omega)$. In case $2(q-1) > p^*$, let us consider $d \geq 3$. As $q \leq p^*$ and $2d/(d+2) < p$, we see that $2(q-1) \leq 2(p^*-1) < [2^*(p-1)]^* = 2d(p-1)/(d-2p)_+$. Hence by virtue of Gagliardo–Nirenberg’s inequalities along with Lemma 8.2, one can take a constant $\theta \in (0, 1)$ such that

$$\begin{aligned} \|\partial \varphi^2(w)\|_H &= \|w\|_{L^{2(q-1)}(\Omega)}^{q-1} \\ &\leq C \|\nabla w\|_{L^{2^*(p-1)}(\Omega; \mathbb{R}^d)}^{\theta(q-1)} \|w\|_{L^{p^*}(\Omega)}^{(1-\theta)(q-1)} \\ &\leq C \|\Delta_p w\|_{L^2(\Omega)}^{\theta(q-1)/(p-1)} \{\varphi^1(w)\}^{(1-\theta)(q-1)/p} \end{aligned} \quad (8.6)$$

for $w \in D(\partial \varphi^1)$. Here one can observe that

$$\theta(q-1)/(p-1) < 1 \quad \text{if and only if} \quad q < p^*.$$

Hence (A2) follows from (8.5), and analogously for $d \leq 2$. Thus applying Theorem 2.3 to (8.3), we obtain the desired conclusion. \square

In case $p < q$, the following theorem assures global (in time) existence of L^2 -solutions to (8.1), (8.2) for small data for $q \in (p, p^*]$ even including the Sobolev critical exponent p^* .

THEOREM 8.4 (Global existence for small data and $p < q$). *In addition to $p < q$, suppose that*

$$p > \frac{2d}{d+2} \quad \text{and} \quad q \leq p^* = \frac{dp}{(d-p)_+}.$$

Then there exists $\delta_0 > 0$ such that, for every $u_0 \in W_0^{1,p}(\Omega)$ and $f \in L_{\text{loc}}^2([0, \infty); L^2(\Omega))$, the Cauchy–Dirichlet problem (8.1), (8.2) possesses an L^2 -solution u on $[0, \infty)$ satisfying

$$u \in C([0, \infty); L^2(\Omega)) \cap L^\infty(0, \infty; W_0^{1,p}(\Omega)), \quad u(\cdot, 0) = u_0,$$

provided that

$$\|u_0\|_{W_0^{1,p}(\Omega)}^p + \text{ess sup}_{t>0} \left(\int_0^t k_{1-\alpha}(t-s) \|f(\cdot, s)\|_{L^2(\Omega)}^2 ds \right) < \delta_0.$$

Proof. We start with the case where $q < p^*$. We observe that

$$\begin{aligned} \|\partial\varphi^1(w)\|_H &= \|\Delta_p w\|_H \\ &\geq C \|\Delta_p w\|_{W^{-1,p'}(\Omega)} \\ &= C \|\nabla w\|_{L^p(\Omega; \mathbb{R}^d)}^{p-1} = C \{\varphi^1(w)\}^{(p-1)/p} \end{aligned}$$

for $w \in D(\partial\varphi^1)$. Hence (2.3) of (A2)' holds with $m_2(r) := Cr^{(p-1)/p} > 0$ for $r > 0$. In case $2(q-1) \leq p^*$, (A2)' holds with $M_2(r) := Cr^{(q-1)/p}$ for $r \geq 0$. Hence $M_2(r)/m_2(r) = Cr^{(q-p)/p} \rightarrow 0$ as $r \rightarrow 0_+$ because of $q > p$. In case $2(q-1) > p^*$, let us consider $d \geq 3$. Then one can check (2.4) of (A2)' with

$$M_2(r) = Cr^{\frac{1}{p} \frac{(1-\theta)(q-1)}{1-\theta(q-1)/(p-1)}} \quad \text{for } r \geq 0,$$

where $\theta \in (0, 1)$ is same as in the proof of Theorem 8.3. Then we find from $p < q < p^*$ that

$$\lim_{r \rightarrow 0_+} \frac{M_2(r)}{m_2(r)} = C \lim_{r \rightarrow 0_+} r^{\frac{1}{p} \frac{q-p}{1-\theta(q-1)/(p-1)}} = 0,$$

which yields (A2)'. Thus the desired conclusion follows from Theorem 2.6. As for the Sobolev critical case where $q = p^*$, we can still use (8.6), and moreover, we observe that

$$\theta(q-1)/(p-1) = 1.$$

Hence (A2)'_c follows from (8.6). Furthermore, the case $d \leq 2$ can also be treated analogously. Therefore the same conclusion follows from Corollary 2.8. \square

In case $p > q$, one can apply Theorem 2.9 to prove existence of L^2 -solutions on $[0, \infty)$ to (8.1), (8.2).

THEOREM 8.5 (Global existence for $p > q$). *In addition to $p > q$, suppose that (8.5) holds true. For every $u_0 \in W_0^{1,p}(\Omega)$ and $f \in W_{\text{loc}}^{1,2}([0, \infty); L^2(\Omega))$, the Cauchy–Dirichlet problem (8.1), (8.2) admits an L^2 -solution u on $[0, \infty)$ satisfying*

$$u \in C([0, \infty); L^2(\Omega)) \cap L_{\text{loc}}^\infty([0, \infty); W_0^{1,p}(\Omega)), \quad u(\cdot, 0) = u_0.$$

Proof. We have already checked (A1) and (A2) in the proof of Theorem 8.3. Moreover, we find that

$$\varphi^2(w) = \frac{1}{q} \|w\|_{L^q(\Omega)}^q \leq C \|\nabla w\|_{L^p(\Omega; \mathbb{R}^d)}^q = C \{\varphi^1(w)\}^{q/p}$$

for $w \in W_0^{1,p}(\Omega)$, which along with the assumption $p > q$ yields (A3). Thus applying Theorem 2.9 to (8.3), we obtain the desired conclusion. \square

We close this section with the following

REMARK 8.6 (On classical p -Laplace Fujita equation). The nonlinear Calderón–Zygmund estimates (see Lemma 8.2) also enable us to upgrade all the existence results obtained in [18, 24, 26, 27] concerning (1.6) to the Sobolev subcritical range $1 < q < p^*$ even for $p \neq 2$. Furthermore, following the same argument as in the proof of Corollary 2.8, one may also prove global (in time) existence of L^2 -solutions for small data even for the Sobolev critical exponent $q = p^*$.

APPENDIX A. EXISTENCE OF STRONG SOLUTIONS TO (7.1)

This appendix is devoted to exhibiting an outline of a proof for the existence of strong solutions to the Cauchy problem (7.1). We also refer the reader to [3], where the case $\lambda = 0$ is treated and which may also be useful to handle the present case $\lambda > 0$.³ Let $T \in (0, \infty)$ be fixed. Then the Cauchy problem (7.1) is rewritten as

$$\lambda \mathcal{A}(u - u_0) + \mathcal{B}(u - u_0) + \partial \Phi(u) + B(u(\cdot)) \ni f \quad (\text{A.1})$$

for positive $\lambda \in (0, 1)$, $u_0 \in D(\varphi)$, $f \in L^2(0, T; H)$ and $B : H \rightarrow H$ satisfying (4.5) (see Subsections 3.1 and 3.2 for the definition of \mathcal{A} , \mathcal{B} and Φ). To prove the existence of solutions to (A.1), we set $\mathcal{X} = C([0, T]; H)$ equipped with the norm $\|u\|_{\mathcal{X}} := \sup_{t \in [0, T]} e^{-\omega t} \|u(t)\|_H$ for some $\omega > 0$ which will be determined later and define a mapping $S : \mathcal{X} \rightarrow \mathcal{X}$ given by

$$S(v) = u \quad \text{for } v \in \mathcal{X},$$

where $u \in D(\mathcal{A}) + u_0$ is the unique solution of

$$\lambda \mathcal{A}(u - u_0) + \mathcal{B}(u - u_0) + \partial \Phi(u) \ni f - B(v(\cdot)). \quad (\text{A.2})$$

³The present case $\lambda > 0$ is much simpler to the case $\lambda = 0$. Actually, due to the presence of the first-order derivative, one can get better a priori estimates for (approximate) solutions in a standard manner.

Hence it suffices to find a fixed point of S . We first show that, for each $v \in L^2(0, T; H)$, (A.2) admits a unique solution $u \in D(\mathcal{A}) + u_0$. To this end, let us introduce

$$\lambda \mathcal{A}(u_\mu - u_0) + \mathcal{B}(u_\mu - u_0) + \partial \Phi_\mu(u) \ni f - B(v(\cdot)). \quad (\text{A.3})$$

Then for each $\mu \in (0, 1)$ and $v \in L^2(0, T; H)$, the Cauchy problem (A.3) always possesses a unique solution $u_\mu \in D(\mathcal{A}) + u_0$. Indeed, since $\lambda \mathcal{A} + \mathcal{B}$ is maximal monotone in $L^2(0, T; H)$ (see Proposition 3.4) and $D(\partial \Phi_\mu) = L^2(0, T; H)$, one finds that $\lambda \mathcal{A} + \mathcal{B} + \partial \Phi_\mu(\cdot + u_0)$ is surjective on $L^2(0, T; H)$ (see, e.g., [9], [8]). Hence as $f - B(v(\cdot)) \in L^2(0, T; H)$, there exists $u_\mu \in D(\mathcal{A}) + u_0$ such that (A.3) holds. As in the proof of Theorem 2.3 of [3], testing (A.3) by du/dt and noting that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|w(t)\|_H^2 &= \left(\frac{dw}{dt}(t), w(t) \right)_H \\ &\leq \left\| \frac{dw}{dt}(t) \right\|_H \|w(t)\|_H \quad \text{for a.e. } t \in (0, T) \end{aligned} \quad (\text{A.4})$$

for $w \in W^{1,2}(0, T; H)$, we can derive

$$\sup_{\mu \in (0, 1)} \left(\lambda \int_0^t \left\| \frac{du_\mu}{dt}(\tau) \right\|_H^2 d\tau + \varphi(u_\mu(t)) \right) \leq C$$

for some $C \geq 0$, which may depend on $\lambda > 0$ but is independent of $\mu \in (0, 1)$. Furthermore, as λ is positive, one can prove that (u_μ) forms a Cauchy sequence in $C([0, T]; H)$ by using (A.4) again. Thus we can verify that

$$\begin{aligned} u_{\mu_n} &\rightarrow u \quad \text{strongly in } C([0, T]; H), \\ u_{\mu_n} &\rightarrow u \quad \text{weakly in } W^{1,2}(0, T; H) \end{aligned}$$

for some $\mu_n \rightarrow 0_+$ and $u \in W^{1,2}(0, T; H)$ satisfying $u(0) = u_0$, and then, as in the proof of Theorem 2.3 of [3], the limit u turns out to be a solution to (A.2).

We next claim that choosing $\omega > 0$ large enough, one can take a constant $\kappa \in (0, 1)$ such that

$$\|S(v_1) - S(v_2)\|_{\mathcal{X}} \leq \kappa \|v_1 - v_2\|_{\mathcal{X}} \quad \text{for } v_1, v_2 \in \mathcal{X}. \quad (\text{A.5})$$

Then by virtue of Banach's contraction mapping principle, we can deduce that S admits a unique fixed point $u \in \mathcal{X}$, i.e., $S(u) = u$, which implies that u is a solution to (A.1). Moreover, the uniqueness of solutions to (A.1) follows as well. Indeed, set $u_1 = S(v_1)$ and $u_2 = S(v_2)$. Subtracting equations and multiplying it by $u_1 - u_2$, we have

$$\begin{aligned} &\frac{\lambda}{2} \frac{d}{dt} \|u_1(t) - u_2(t)\|_H^2 + (\mathcal{B}(u_1 - u_2)(t), u_1(t) - u_2(t))_H \\ &\quad + (\partial \varphi(u_1(t)) - \partial \varphi(u_2(t)), u_1(t) - u_2(t))_H \\ &= - (B(v_1(t)) - B(v_2(t)), u_1(t) - u_2(t))_H \end{aligned}$$

for a.e. $t \in (0, T)$. Integrating both sides over $(0, t)$ and using the monotonicity of \mathcal{B} and $\partial\varphi$, we find that

$$\begin{aligned} \frac{\lambda}{2} \|u_1(t) - u_2(t)\|_H^2 &\leq L_B \int_0^t \|v_1(s) - v_2(s)\|_H \|u_1(s) - u_2(s)\|_H \, ds \\ &\leq L_B \|v_1 - v_2\|_{\mathcal{X}} \|u_1 - u_2\|_{\mathcal{X}} \int_0^t e^{2\omega s} \, ds \\ &\leq \frac{L_B}{2\omega} e^{2\omega t} \|v_1 - v_2\|_{\mathcal{X}} \|u_1 - u_2\|_{\mathcal{X}} \quad \text{for } t \in [0, T], \end{aligned}$$

which yields

$$\|u_1 - u_2\|_{\mathcal{X}} \leq \frac{L_B}{\omega\lambda} \|v_1 - v_2\|_{\mathcal{X}}.$$

Therefore choosing $\omega > L_B/\lambda$ and setting $\kappa := L_B/(\omega\lambda) \in (0, 1)$, we obtain (A.5). Thus we conclude that

PROPOSITION A.1. *Let $T \in (0, \infty)$ and $\lambda \in (0, 1)$ be fixed. For every $f \in L^2(0, T; H)$ and $u_0 \in D(\varphi)$, the Cauchy problem (A.1) admits a unique strong solution $u \in W^{1,2}(0, T; H) \cap D(\partial\Phi)$ satisfying $u(0) = u_0$.*

Therefore for each $T \in (0, \infty)$, $\lambda \in (0, 1)$, $u_0 \in D(\varphi)$ and $f \in L^2(0, T; H)$, the Cauchy problem (7.1) admits a unique strong solution $u_\lambda \in W^{1,2}(0, T; H)$ on $[0, T]$. Concerning the case where $T = \infty$, as in the proof of Theorem 4.4, one can verify that for each $\lambda \in (0, 1)$, $u_0 \in D(\varphi)$ and $f \in L^2_{\text{loc}}([0, \infty); H)$, the Cauchy problem (7.1) also admits a unique strong solution $u_\lambda \in W^{1,2}_{\text{loc}}([0, \infty); H)$ on $[0, \infty)$.

REFERENCES

- [1] Franz Achleitner, Goro Akagi, Christian Kuehn, Jens Markus Melenk, Jens D. M. Rademacher, Cinzia Soresina, and Jichen Yang. Fractional dissipative PDEs. In *Fractional dispersive models and applications—recent developments and future perspectives*, volume 37 of *Nonlinear Syst. Complex.*, pages 53–122. Springer, Cham, [2024] ©2024.
- [2] E. Affili and E. Valdinoci. Decay estimates for evolution equations with classical and fractional time-derivatives. *J. Differential Equations*, 266:4027–4060, 2019.
- [3] G. Akagi. Fractional flows driven by subdifferentials in Hilbert spaces. *Israel J. Math.*, 234:809–862, 2019.
- [4] Goro Akagi. Local existence of solutions to some degenerate parabolic equation associated with the p -Laplacian. *J. Differential Equations*, 241(2):359–385, 2007.
- [5] Goro Akagi and Mitsuharu Ôtani. Evolution inclusions governed by the difference of two subdifferentials in reflexive Banach spaces. *J. Differential Equations*, 209(2):392–415, 2005.
- [6] Goro Akagi and Ulisse Stefanelli. A variational principle for gradient flows of non-convex energies. *J. Convex Anal.*, 23(1):53–75, 2016.
- [7] C.O. Alves and T. Boudjeriou. Existence and uniqueness of solution for some time fractional parabolic equations involving the 1-Laplace operator. *Partial Differential Equations and Applications*, 4, 2023.
- [8] V. Barbu. *Nonlinear Semigroups and Differential Equations in Banach spaces*. Noordhoff, 1976.

- [9] H. Brézis. *Opérateurs Maximaux Monotones et Semi-Groupes de Contractions dans les Espaces de Hilbert*, volume 5 of *Math Studies*. North-Holland, Amsterdam/New York, 1973.
- [10] Haim Brezis. *Functional analysis, Sobolev spaces and partial differential equations*. Universitext. Springer, New York, 2011.
- [11] Ph. Clément. *On abstract Volterra equations in Banach spaces with completely positive kernels*, volume 1076 of *Lecture Notes in Math.*, pages 32–40. Springer, Berlin, 1984.
- [12] Ph. Clément and J.A. Nohel. Asymptotic behavior of solutions of nonlinear volterra equations with completely positive kernels. *SIAM J. Math. Anal.*, 12:514–535, 1981.
- [13] Ph. Clément and J. Prüss. Completely positive measures and Feller semigroups. *Math. Ann.*, 287:73–105, 1990.
- [14] Ataru Fujii and Masahito Ohta. Asymptotic behavior of blowup solutions of a parabolic equation with the p -Laplacian. *Publ. Res. Inst. Math. Sci.*, 32(3):503–515, 1996.
- [15] Hiroshi Fujita. On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$. *J. Fac. Sci. Univ. Tokyo Sect. I*, 13:109–124, 1966.
- [16] C.G. Gal and M. Warma. *Fractional-in-time semilinear parabolic equations and applications*, volume 84 of *Mathematics & Applications*. Springer, Cham, 2020.
- [17] G. Gripenberg. Volterra integro-differential equations with accretive nonlinearity. *J. Differential Equations*, 60:57–79, 1985.
- [18] Hitoshi Ishii. Asymptotic stability and blowing up of solutions of some nonlinear equations. *J. Differential Equations*, 26(2):291–319, 1977.
- [19] Juha Kinnunen and Shulin Zhou. A boundary estimate for nonlinear equations with discontinuous coefficients. *Differential Integral Equations*, 14(4):475–492, 2001.
- [20] Y. Kōmura. Nonlinear semi-groups in Hilbert space. *J. Math. Soc. Japan*, 19:493–507, 1967.
- [21] A. Kubica, K. Ryszevska, and M. Yamamoto. *Time-fractional differential equations—a theoretical introduction*. Springer Briefs Math. Springer, Singapore, 2020.
- [22] J.-L. Lions. *Quelques méthodes de résolution des problèmes aux limites non linéaires*. Dunod, Paris; Gauthier-Villars, Paris, 1969.
- [23] Mitsuhiro Nakao. Global solutions for some nonlinear parabolic equations with non-monotonic perturbations. *Nonlinear Anal.*, 10(3):299–314, 1986.
- [24] M. Ôtani. Existence and asymptotic stability of strong solutions of nonlinear evolution equations with a difference term of subdifferentials. In *Qualitative theory of differential equations, Vol. I, II (Szeged, 1979)*, volume 30 of *Colloq. Math. Soc. János Bolyai*, pages 795–809. North-Holland, Amsterdam-New York, 1981.
- [25] Mitsuharu Ôtani. On the existence of strong solutions for $du/dt(t) + \partial\psi^1(u(t)) - \partial\psi^2(u(t)) \ni f(t)$. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 24(3):575–605, 1977.
- [26] Mitsuharu Ôtani. Nonmonotone perturbations for nonlinear parabolic equations associated with subdifferential operators, Cauchy problems. *J. Differential Equations*, 46(2):268–299, 1982.
- [27] Mitsuharu Ôtani. Nonmonotone perturbations for nonlinear parabolic equations associated with subdifferential operators, periodic problems. *J. Differential Equations*, 54(2):248–273, 1984.
- [28] A. Pazy. *Semigroups of linear operators and applications to partial differential equations*, volume 44 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1983.
- [29] Kerstin Schmitz and Petra Wittbold. Entropy solutions for time-fractional porous medium type equations. *Differential Integral Equations*, 37(5-6):309–322, 2024.
- [30] R. E. Showalter. *Monotone operators in Banach space and nonlinear partial differential equations*, volume 49 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997.
- [31] Jacques Simon. Compact sets in the space $L^p(0, T; B)$. *Ann. Mat. Pura Appl. (4)*, 146:65–96, 1987.

- [32] Masayoshi Tsutsumi. Existence and nonexistence of global solutions for nonlinear parabolic equations. *Publ. Res. Inst. Math. Sci.*, 8:211–229, 1972.
- [33] V. Vergara and R. Zacher. Lyapunov functions and convergence to steady state for differential equations of fractional order. *Math. Z.*, 259:287–309, 2008.
- [34] V. Vergara and R. Zacher. A priori bounds for degenerate and singular evolutionary partial integro-differential equations. *Nonlinear Anal.*, 73:3572–3585, 2010.
- [35] V. Vergara and R. Zacher. Optimal decay estimates for time-fractional and other nonlocal subdiffusion equations via energy methods. *SIAM J. Math. Anal.*, 47:210–239, 2015.
- [36] V. Vergara and R. Zacher. Stability, instability, and blowup for time fractional and other nonlocal in time semilinear subdiffusion equations. *J. Evol. Equ.*, 17:599–626, 2017.
- [37] Fred B. Weissler. Local existence and nonexistence for semilinear parabolic equations in L^p . *Indiana Univ. Math. J.*, 29(1):79–102, 1980.
- [38] P. Wittbold, P. Wolejko, and R. Zacher. Bounded weak solutions of time-fractional porous medium type and more general nonlinear and degenerate evolutionary integro-differential equations. *J. Math. Anal. Appl.*, 499:20 pp, 2021.
- [39] R. Zacher. Maximal regularity of type l_p for abstract parabolic Volterra equations. *J. Evol. Equ.*, 5:79–103, 2005.
- [40] R. Zacher. Boundedness of weak solutions to evolutionary partial integro-differential equations with discontinuous coefficients. *J. Math. Anal. Appl.*, 348:137–149, 2008.
- [41] R. Zacher. Weak solutions of abstract evolutionary integro-differential equations in Hilbert spaces. *Funkcial. Ekvac.*, 52:1–18, 2009.
- [42] R. Zacher. A De Giorgi–Nash type theorem for time fractional diffusion equations. *Math. Ann.*, 356:99–146, 2013.
- [43] R. Zacher. A weak harnack inequality for fractional evolution equations with discontinuous coefficients. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 12:903–940, 2013.
- [44] R. Zacher. *Time fractional diffusion equations: solution concepts, regularity, and long-time behavior*, volume 2, pages 159–179. De Gruyter, Berlin, 2019.

(Goro Akagi) MATHEMATICAL INSTITUTE AND GRADUATE SCHOOL OF SCIENCE, TOHOKU UNIVERSITY, AOBA, SENDAI 980-8578, JAPAN

Email address: goro.akagi@tohoku.ac.jp

(Yoshihito Nakajima) GRADUATE SCHOOL OF SCIENCE, TOHOKU UNIVERSITY, AOBA, SENDAI 980-8578, JAPAN

Email address: yoshihito.nakajima.p8@dc.tohoku.ac.jp