Mixed State Entanglement Entropy in CFT

Xin Jiang,^{*} Peng Wang,[†] Houwen Wu,[‡] and Haitang Yang[§] College of Physics, Sichuan University, Chengdu, 610065, China

How to calculate the entanglement entropy between two subsystems for a mixed state has remained an important problem. In this paper, we provide a straightforward method, namely the subtraction approach, to solve this problem for generic covariant bipartite mixed states, where time dependence is explicitly included. We further demonstrate that the mixed state entanglement entropy $S_{\rm vN}$ can be calculated in a more abstract yet powerful way. Within the context of the AdS₃/CFT₂ and AdS₅/CFT₄ correspondences, we show that $S_{\rm vN}$ exactly matches the corresponding entanglement wedge cross section in the AdS bulk, respectively.

^{*} domoki@stu.scu.edu.cn

[†] pengw@scu.edu.cn

[‡] iverwu@scu.edu.cn

[§] hyanga@scu.edu.cn

CONTENTS

I.	Introduction	2
II.	Covariant entanglement entropy for mixed states	5
	A. Symmetric configuration and Replica trick	6
	B. Generic covariant configurations	8
	C. Two gauge parameters	10
III.	Holographic dual of the mixed state entanglement entropy	11
IV.	Renormalization group flow induced entanglement entropy	11
	A. Two-dimensional CFT	13
	B. Four-dimensional CFT	13
V.	Conclusion	15
	Acknowledgments	15
	References	15

I. INTRODUCTION

Quantum entanglement has unveiled profound connections between quantum gravity and quantum information theory. Within the framework of the AdS/CFT correspondence [1–3], the entanglement entropy [4–6] provides significant insights into the relationship between entanglement and the emergence of spacetime [7–9]. The entanglement entropy, also known as the von Neumann entropy, is defined as

$$S_{\rm vN}\left(A\right) = -\mathrm{Tr}\rho_A \log \rho_A,\tag{1}$$

where ρ_A is the reduced density matrix of a subsystem A. The entanglement entropy satisfactorily characterizes the quantum entanglement between two subsystems in a *pure state*. However, for the bipartite systems in a *mixed state*, such as two disjoint intervals in CFT₂, the traditional entanglement entropy is no longer a suitable measure of quantum entanglement. Several approaches have been proposed to quantify the degree of entanglement in a mixed state, such as the entanglement negativity [10], the entanglement of purification [11], the odd entanglement entropy [12], the



Figure 1. The bulk dual of the EoP (E_W in left panel) and the bulk dual of the reflected entropy (S_R in right panel).

reflected entropy [13], and the balanced partial entanglement [14].

The key idea of the entanglement of purification (EoP) is to purify a mixed state ρ_{AB} by introducing an auxiliary system \overline{AB} . The entanglement entropy between the purified state $A\overline{A}$ and $B\overline{B}$ is then defined as the minimum von Neumann entropy over all possible purifications: $E_p = \min_{\overline{AB}} S_{\rm vN} (A\overline{A} : B\overline{B})$. The holographic dual of EoP is supposed to be a geodesic in the bulk of AdS₃, known as the entanglement wedge cross-section (EWCS or E_W), as shown by the left panel of Figure 1. However, optimization over purifications is, in practice, not workable in CFTs. As a compromise, canonical purification is often employed, which involves directly duplicating the mixed state AB into a state denoted AA^*BB^* . The resulting entanglement entropy, called the reflected entropy, is given by $S_R(A : B) = -\text{Tr}\rho_{AA^*} \log \rho_{AA^*}$. In holographic interpretation, it has been shown that $S_R(A : B) = 2E_W(A : B)$. The bulk dual of the reflected entropy is a closed curve comprising two identical geodesics, as illustrated by the right panel of Figure 1.

In [15], we proposed an alternative approach, namely SUBTRACTION, to purification in CFT₂. Instead of purifying the mixed state by adding auxiliary systems, our method subtracts the undetectable regions of the system, as illustrated in Figure 2. The remaining regions, which are disjoint but mutually complementary, form a pure entangled state ψ_{AB} , satisfying $A^C = B$. This process is equivalent to enlarging the cutoff regulators, effectively situating the boundary CFT (BCFT) on the edge of the excluded region (dashed circle). Using the replica trick [16, 17] and the annulus CFT techniques [18–20], we computed the von Neumann entropy $S_{\rm vN}$ (A : B) for two disjoint intervals A and B living at the time t = 0, as illustrated in Figure 3. Notably, it was found that



Figure 2. The subtraction method could be understood as introducing a finite regulator. Left panel: Two subsystems with an infinitesimal regulator. Right panel: Two subsystems with a finite regulator.

for a *static* time slice of bulk AdS₃. This proposal is independent of purifications and four-point conformal block data, making it very easy to compute. In this paper, our aim is to extend the subtraction method to the covariant case and compute the corresponding entropy $S_{\rm vN}(A:B)$ for two generic intervals A and B, as illustrated in Figure 3.

In the subtraction method, enlarging the regulator is reminiscent of the renormalization group (RG) flow. Using the Callan-Symanzik RG equation, we can derive an RG flow-induced entangle-



Figure 3. Two different configurations of mixed states in CFT living on the complex plane, where the blue shaded regions represent the Euclidean path integral. Left panel: At $\tau = 0$, two subsystems A and B on the x-axis in a mixed state ρ_{AB} . Right panel: Two generic subsystems that are determined by four ending points z_i at different time.

ment entropy, which turns out to be exactly $S_{vN}(A : B)$. Furthermore, since the RG equation is universal, we can apply it to higher dimensional cases. As an example, we calculate the RG flow induced entanglement entropy in holographic CFT₄ and find that it agrees with the entanglement wedge cross-section in the AdS₅ bulk.

The remainder of this paper is structured as follows: In Section II, we provide a rigorous calculation of covariant mixed state $S_{\rm vN}(A:B)$ using the replica trick in the subtraction method. We clarify that there are two gauge parameters in the entanglement entropy $S_{\rm vN}(A:B)$. In Section III, we discuss the holographic dual of $S_{\rm vN}(A:B)$. In Section IV, we define the RG flow induced entanglement entropy through the Callan-Symanzik RG equation and apply it to the CFT₂ and CFT₄, respectively. Section V contains the conclusion.

II. COVARIANT ENTANGLEMENT ENTROPY FOR MIXED STATES

In previous work [15], we introduced the SUBTRACTION approach, as an alternative to purification in CFT. The computed quantity, $S_{\rm vN}(A:B)$, is the entropy of entanglement between two disjoint intervals A and B in a mixed state ρ_{AB} on a time slice. To complete the story, it is of importance to include the time dependence in the expression of $S_{\rm vN}(A:B)$.



Figure 4. Static symmetric configuration. Left panel: At $\tau = 0$, two subsystems A and B on the x-axis are in a mixed state ρ_{AB} , separated by segments C and D. Right panel: After introducing two finite regulators (subtracting C and D with two discs), we obtain an annulus, in which two subsystems A and B are now in a pure entangled state ψ_{AB} .

A. Symmetric configuration and Replica trick

We begin by considering the symmetric case and will extend to generic configurations using conformal transformations. As explained in the introduction, in ref. [15], we considered two disjoint segments A and B on the x-axis at $\tau = 0$, which are separated by undetectable segments C and D, as illustrated in Figure 4. Segments C and D are then removed by two discs with conformal invariant boundary conditions, which encode the irrelevant information coming from C and D.



Figure 5. Covariant symmetric configuration. Left panel: Density matrix of the mixed state ρ_{AB} , segments A and B are symmetric with respect to the origin. Right panel: The complimentary parts of $A \cup B$ in the system are removed by two discs with conformal invariant boundary conditions. The annulus is a pure state ψ_{AB} .

To include the time dependence, referring to the Euclidean path integral region in Figure 5, we now consider $z_i = x_i + \tau_i$ for i = 1, 2, 3, 4 and $\tau_i \neq 0$. If we define $Z_{ij} = z_j - z_i = R_{ij}e^{i\theta_{ij}}$ and use the analytic continuation $z_i = x_i - \tau_i$ and $\bar{z}_i = x_i + \tau_i$ to a Lorentzian signature, the angle θ_{ij} maps to a boost parameter $\theta_{ij} = i\kappa_{ij}$. For simplicity, we denote $R_1 \equiv R_{23}$ and $R_2 \equiv R_{14}$. We first address the symmetric configuration by setting

$$z_1 = -z_4, \quad z_2 = -z_3. \tag{3}$$

Following the similar fashion as the static case, after removing two discs representing the regions separating A and B, we obtain an annulus indicating a pure state ψ_{AB} , as illustrated in Figure

5. The boundaries of the annulus are imposed with conformal invariant boundary conditions, corresponding to boundary states caused by the removed regions.

It is helpful to define the width of the annulus

$$W = \log \frac{R_2}{R_1},\tag{4}$$

where R_1 and R_2 denote the inner and outer radii, respectively. The entanglement entropy $S_{\rm vN}(A)$ in the annulus CFT could be defined through the Rényi entropy:

$$S^{(n)}(A) = \frac{1}{1-n} \log \operatorname{Tr}_A \rho_A^n,$$
 (5)

$$S_{\rm vN}(A) = \lim_{n \to 1} S^{(n)}(A).$$
 (6)

 $\operatorname{Tr}_A \rho_A^n$ is given by

$$\operatorname{Tr}_A \rho_A^n = \frac{Z_n}{Z_1^n}.$$
(7)

Here, Z_1 is the partition function on the original annulus. Z_n is the partition function on the *n*-sheeted cover \mathcal{M}_n obtained by cyclically gluing *n* copies of \mathcal{M} along *A*, see Figure 6. Note that $\operatorname{Tr}_A \rho_A^n$ is independent of the shape of the subsystem *A* but depends only on the endpoints of *A*.



Figure 6. The cut-and-glue procedure in the replica trick. Each annulus is cut along the subsystem A and is glued with others cyclically. Red lines represent gluing operations. Note that the resulted manifold is also an annulus.

By the generator of scale transformations, the annulus partition function Z_1 can be written as [18–20]:

$$Z_1 = e^{cW/12} \sum_k \langle a|k \rangle \langle k|b \rangle e^{-2\delta_k W}, \tag{8}$$

where c is the central charge, k is all allowed scalar operators with dimensions δ_k , and $|a,b\rangle$ are boundary states on the annulus two boundaries. The replicated manifold \mathcal{M}_n is conformally equivalent to an annulus with the width $W_n = W/n$, implying that

$$Z_n = e^{cW/12n} \sum_k \langle a|k \rangle \langle k|b \rangle e^{-2\delta_k W/n}.$$
(9)

Since we are calculating the vacuum entanglement (k = 0), we have

$$Z_1 = e^{cW/12} \langle a|0\rangle \langle 0|b\rangle, \tag{10}$$

$$Z_n = e^{cW/12n} \langle a|0\rangle \langle 0|b\rangle, \tag{11}$$

and

$$\operatorname{Tr}_{A}\rho_{A}^{n} = e^{\frac{c}{12}(\frac{1}{n}-n)W} \left(\langle a|0\rangle\langle 0|b\rangle\right)^{1-n}.$$
(12)

Substituting $\text{Tr}_A \rho_A^n$ into Equation (5), we obtain the Rényi entropy

$$S^{(n)}(A) = \frac{c}{12} \left(1 + \frac{1}{n} \right) W + g_a + g_b,$$
(13)

where $g_{a,b} = \log \langle a, b | 0 \rangle$ are the Affleck-Ludwig boundary entropies [21]. The boundary entropies encode the information from the removed subsystems. They are model dependent and irrelevant to the entanglement between A and B. So, one can simply choose $g_a = g_b = 0$ to get

$$S_{\rm vN}(A) = \lim_{n \to 1} S^{(n)}(A) = \frac{c}{6}W.$$
 (14)

From the derivation, it is straightforward to see $S_{vN}(A) = S_{vN}(B)$ since the roles of A and B are symmetric. Therefore, the entropy of entanglement between A and B is naturally defined by the universal term,

$$S_{\rm vN}(A:B) := \frac{c}{6}W = \frac{c}{6}\log\frac{R_2}{R_1}.$$
(15)

B. Generic covariant configurations

Next, we consider a generic covariant configuration where two subsystems are selected as $A \in [z_1, z_2]$ and $B \in [z_3, z_4]$ without specific constraints. Through the following conformal transformations:

$$v(z) = 2\frac{z - z_1}{z_4 - z_1} - 1, \quad w(v) = \frac{\gamma - v}{\gamma v - 1},$$
(16)



Figure 7. Two generic intervals determined by z_1, z_2, z_3 and z_4 on the z-plane could be conformally mapped to two symmetric intervals on the w-plane.

with

$$\gamma = \frac{1 + \alpha\beta + \sqrt{(1 - \alpha^2)(1 - \beta^2)}}{\alpha + \beta}, \quad \alpha = v(z_2), \quad \beta = v(z_3), \tag{17}$$

we can map a generic covariant configuration on the z-plane to a symmetric one on the w-plane:

$$(z_1, z_2, z_3, z_4) \to (-1, -\delta, \delta, 1),$$
 (18)

with

$$\delta = \frac{1 - \alpha\beta + \sqrt{(1 - \alpha^2)(1 - \beta^2)}}{\alpha - \beta},\tag{19}$$

as illustrated in Figure 7. On the *w*-plane, we could apply the subtraction method, where the inner radius is $R_1 = 1$ and the outer radius is $R_2 = |\delta| > 1$. The corresponding entanglement entropy in this annulus is thus given by:

$$S_{\rm vN}(A:B) = \frac{c}{6} \log |\delta|, \qquad (20)$$

$$= \frac{c}{6} \log \left| 1 + \frac{2(z_1 - z_2)(z_3 - z_4)}{(z_1 - z_4)(z_2 - z_3)} + 2\sqrt{\frac{(z_1 - z_2)(z_1 - z_3)(z_2 - z_4)(z_3 - z_4)}{(z_1 - z_4)^2(z_2 - z_3)^2}} \right|.$$
(21)

In terms of the cross ratio

$$\eta = \frac{z_{12}z_{34}}{z_{13}z_{24}},\tag{22}$$

the covariant entanglement entropy of mixed state ρ_{AB} can be written as

$$S_{\rm vN}(A:B) = \frac{c}{6} \log \left| \frac{1+\sqrt{\eta}}{1-\sqrt{\eta}} \right|,\tag{23}$$

$$= \frac{c}{12} \log \left(\frac{1 + \sqrt{\eta}}{1 - \sqrt{\eta}} \right) + \frac{c}{12} \log \left(\frac{1 + \sqrt{\eta}}{1 - \sqrt{\eta}} \right).$$
(24)

Using Equation (24), it is straightforward to obtain the result for a finite temperature system or a finite size system. To this end, one only needs to apply the following conformal transformations to z while leaving the form of the equation (24) invariant:

Finite temperature :
$$z = \exp \frac{2\pi w}{\beta}$$
,
Finite size : $z = \tan \frac{\pi u}{L}$. (25)

Therefore, the covariant mixed state entanglement entropy for a finite temperature system or a finite size system is given by:

$$S_{\rm vN}\left(A:B\right) = \frac{c}{12}\log\left(\frac{1+\sqrt{\eta}}{1-\sqrt{\eta}}\right) + \frac{c}{12}\log\left(\frac{1+\sqrt{\eta}}{1-\sqrt{\eta}}\right),$$

with

Finite temperature :
$$\eta = \frac{\sinh(\pi w_{12}/\beta)\sinh(\pi w_{34}/\beta)}{\sinh(\pi w_{13}/\beta)\sinh(\pi w_{24}/\beta)},$$

Finite size :
$$\eta = \frac{\sin(\pi u_{12}/L)\sin(\pi u_{34}/L)}{\sin(\pi u_{13}/L)\sin(\pi u_{24}/L)},$$
(26)

respectively.

C. Two gauge parameters

Notably, two generic intervals are uniquely fixed by eight real numbers. However, when considering the entropy of entanglement between two generic intervals, there are two gauge parameters. To see this, it suffices to consider the symmetric case,

$$z_1 = -z_4, \quad z_2 = -z_3,$$
 (27)

where the corresponding entanglement entropy is given by (15):

$$S_{\rm vN}(A:B) = \frac{c}{6} \log \frac{|z_1 - z_4|}{|z_2 - z_3|}.$$
(28)

Note that $S_{vN}(A:B)$ is invariant under the following *local* transformations:

$$(z_1, z_2, z_3, z_4) \to (z_1 e^{i\theta}, z_2, z_3, z_4 e^{i\theta}), \quad (z_1, z_2, z_3, z_4) \to (z_1, z_2 e^{i\theta'}, z_3 e^{i\theta'}, z_4),$$
 (29)

with θ, θ' two real numbers. Up to these local transformations, two intervals in different configurations that cannot be related by global conformal transformations, may still share the same entanglement entropy. In other words, even though two generic intervals are determined by eight real numbers, two of them are gauge redundancy. So, there are really only six independent parameters for generic covariant $S_{\rm vN}(A:B)$.

III. HOLOGRAPHIC DUAL OF THE MIXED STATE ENTANGLEMENT ENTROPY

For completeness, we briefly discuss the bulk dual of the entanglement entropy $S_{\rm vN}(A:B)$, i.e. the covariant EWCS, which is also addressed in [22, 23]. The two gauge parameters discussed above play a crucial role when considering the bulk dual of the covariant entanglement entropy. In the AdS₃ bulk, the covariant EWCS is determined by two endpoints, which are fixed by six real numbers. This precisely matches the six independent real numbers in $S_{\rm vN}(A:B)$ after fixing the two gauge parameters. Thus, a one-to-one correspondence between bulk points and boundary points can be established.

For example, we could choose $(z_1, z_2, z_3, z_4) = (0, 1, 3, z_4)$. Following [23], the covariant EWCS is determined by two endpoints (t_a, x_a, y_a) and (t_b, x_b, y_b) through the following transformations:

$$t_{a} = 0,$$

$$x_{a} = 1 + \frac{2}{1 + 9\sqrt{\eta\bar{\eta}}},$$

$$y_{a} = \left(\frac{36\sqrt{\eta\bar{\eta}}}{(9\sqrt{\eta\bar{\eta}} + 1)^{2}}\right)^{1/2},$$
(30)

and

$$t_{b} = -\frac{3(\eta - \bar{\eta})}{9\eta\bar{\eta} - 3\bar{\eta} + 4\sqrt{\eta\bar{\eta}} - 3\eta + 1},$$

$$x_{b} = \frac{3(3\eta\bar{\eta} - 2\bar{\eta} - 2\eta + 1)}{9\eta\bar{\eta} - 3\bar{\eta} + 4\sqrt{\eta\bar{\eta}} - 3\eta + 1},$$

$$y_{b} = \left(\frac{36(\eta - 1)\sqrt{\eta\bar{\eta}}(\bar{\eta} - 1)}{(-3\bar{\eta} + \eta(9\bar{\eta} - 3) + 4\sqrt{\eta\bar{\eta}} + 1)^{2}}\right)^{1/2}.$$
(31)

The relations between the four boundary points of the cross ratio η and the two bulk points (t_a, x_a, y_a) and (t_b, x_b, y_b) are illustrated in Figure 8. By applying these coordinate transformations, the entanglement entropy (24) becomes:

$$S_{\rm vN}(A:B) = \frac{c}{6} \operatorname{arccosh}\left(\frac{-(t_a - t_b)^2 + (x_a - x_b)^2 + (y_a - y_b)^2}{2y_a y_b} + 1\right),\tag{32}$$

which is precisely the geodesic length between (t_a, x_a, y_a) and (t_b, x_b, y_b) in the AdS₃ geometry.

IV. RENORMALIZATION GROUP FLOW INDUCED ENTANGLEMENT ENTROPY

In this section, we develop a more abstract but powerful approach to obtain the mixed state entanglement entropy $S_{\rm vN}$. By recognizing that the essence of the subtraction method lies in



Figure 8. Two generic intervals on the boundary and their corresponding EWCS in the AdS_3 bulk.

introducing finite regulators, it is natural to define $S_{\rm vN}$ by directly using the RG flow equation in a finite region.

In local quantum field theories (QFTs), the entanglement entropy for an entangling surface Σ is typically invariant under a global rescaling of all dimensional parameters. Especially for a CFT the universal part ¹ of entanglement entropy is invariant under conformal transformations, allowing us to calculate entanglement entropy through the trace anomaly [5]. Consider a classically scale-invariant theory living on a *d*-dimensional manifold \mathcal{M} with the metric $ds^2 = \gamma_{ab} dx^a dx^b$, where the trace anomaly is given by

$$\langle T_a^a \rangle = 2 \frac{\gamma^{ab}}{\sqrt{\gamma}} \frac{\delta}{\delta \gamma^{ab}} \log Z_{\rm CFT}.$$
 (33)

In this set-up, the Callan-Symanzik RG equation is known as [24]:

$$\left[\ell \frac{\partial}{\partial \ell} - 2 \int_{\mathcal{M}} \gamma^{ab} \frac{\delta}{\delta \gamma^{ab}}\right] \log Z_{\rm CFT} = 0, \tag{34}$$

with the renormalization scale ℓ that has the Length dimension. Incorporating with the replica trick [16], and the Rényi entropy

$$S_{\rm EE}^{(n)} = \frac{1}{1-n} \log \left[\frac{Z_{\mathcal{M}_n}}{(Z_{\mathcal{M}_1})^n} \right],\tag{35}$$

where \mathcal{M}_n is the replicated manifold and \mathcal{M}_1 is the original manifold with entangling surface Σ , we can define a RG equation for the Rényi entropy:

$$\ell \frac{\partial}{\partial \ell} S_{\rm EE}^{(n)} = \frac{\int_{\mathcal{M}_n} \langle T_a^a \rangle_{\mathcal{M}_n} - n \int_{\mathcal{M}_1} \langle T_a^a \rangle_{\mathcal{M}_1}}{1 - n}.$$
(36)

¹ Thereafter, we disregard the non-universal terms and refer to $S_{\rm EE}$ as its universal part.

A. Two-dimensional CFT

When d = 2, the trace anomaly simplifies to

$$\langle T_a^a \rangle = \frac{c}{24\pi} \mathcal{R},\tag{37}$$

with \mathcal{R} as the Ricci scalar and c as the central charge. For a replicated manifold with conical singularities, it was shown [25] that

$$\int_{\mathcal{M}_n} \mathcal{R}^{(n)} = n \int_{\mathcal{M}_1} \mathcal{R} + 4\pi \left(1 - n\right) \int_{\Sigma} 1.$$
(38)

In the two-dimensional case, the entangling surface Σ is point-like, and $\int_{\Sigma} 1 = \mathcal{A}$ represents the number of points. In what follows we assume $\mathcal{A} = 1$. Plugging equations (37) and (38) into equation (36) and taking the limit $n \to 1$, we obtain the Callan-Symanzik equation for the entanglement entropy

$$\ell \frac{\partial}{\partial \ell} S_{\rm EE} = \frac{c}{6}.\tag{39}$$

Therefore, the RG flow induced entanglement entropy, denoted as $S_{\rm RG}$, is

$$S_{\rm RG} = \frac{c}{6} \log \frac{R_2}{R_1},$$
 (40)

where R_1 and R_2 are two renormalization scales. In the complex coordinates $(z, \bar{z}) = (x + it, x - it)$, the renormalization scale (Weyl scaling) ℓ is identified with the radius $r = |z| = \sqrt{x^2 + t^2}$ on the complex plane. The renormalization scale region $r \in [R_1, R_2]$ actually defines an annular region on the complex plane, which precisely is the right panel of Figure 5, representing the pure state obtained from a mixed state by subtraction. Therefore, we conclude $S_{\rm RG} = S_{\rm vN}$.

B. Four-dimensional CFT

It is challenging to calculate the entanglement measure for mixed state in higher-dimensional CFTs. However, this can be easily done in our proposal for some configurations. Let us consider the holographic CFT living on a four-dimensional spacetime $ds^2 = -dt^2 + dx^2 + d\vec{x}_2^2$, where $d\vec{x}_2^2$ represents the transverse directions. The trace anomaly for such a CFT₄ is given by

$$\langle T_a^a \rangle = \frac{c}{16\pi^2} W^2 - \frac{a}{16\pi^2} E^2,$$
(41)

where the central charge c and a are equal in our case, and the curvature square terms are expressed as

$$W^{2} = R_{abcd}R^{abcd} - 2R_{ab}R^{ab} + \frac{1}{3}\mathcal{R}^{2},$$

$$E^{2} = R_{abcd}R^{abcd} - 4R_{ab}R^{ab} + \mathcal{R}^{2}.$$
(42)

If we utilize the RG equation for the *n*-th Rényi entropy (36) and take the $n \to 1$ limit, we find

$$\ell \frac{\partial}{\partial \ell} S_{\rm RG} = a\lambda_1 \cdot \frac{\operatorname{Area}\left(\Sigma\right)}{\ell^2} + a\lambda_2,\tag{43}$$

as previously found in [5]. Here, both λ_1 and λ_2 are numerical constants. For simplicity, we choose the entangling surface Σ to be a two-dimensional plane of transverse directions $d\vec{x}_2^2$, with an area denoted as V. In this case, the constant λ_2 vanishes and Eq. (43) is simplified

$$\ell \frac{\partial}{\partial \ell} S_{\rm RG} = a \lambda_1 \cdot \frac{V}{\ell^2}.$$
(44)

Integrating this equation gives us:

$$S_{\rm RG} = \frac{a\lambda_1 V}{2} \left(\frac{1}{s^2} - \frac{1}{(2l+s)^2} \right),\tag{45}$$

where we choose the UV cutoff to be s and the IR cutoff to be 2l + s.

Within the context of AdS_{d+1}/CFT_d correspondence, the higher-dimensional EWCS has been investigated in AdS_{d+1} bulk [26], where the metric is given by:

$$ds^{2} = \frac{R_{\text{AdS}}^{2}}{\rho^{2}} \left(-dt^{2} + d\rho^{2} + dx^{2} + d\vec{x}_{d-2}^{2} \right), \tag{46}$$

Here, $d\vec{x}_{d-2}^2$ represents the transverse directions, and the EWCS for two parallel strips with equal widths l, separated by a distance s, is:

$$E_W = \frac{VR_{\text{AdS}}^{d-1}}{4G_N^{(d+1)}(d-2)} \frac{2^{d-2}\pi^{\frac{d-2}{2}}\Gamma\left(\frac{d}{2(d-1)}\right)^{d-2}}{\Gamma\left(\frac{1}{2(d-1)}\right)^{d-2}} \left(\frac{1}{s^{d-2}} - \frac{1}{(2l+s)^{d-2}}\right),\tag{47}$$

with V the area of transverse directions. For d = 4, the EWCS is simply

$$E_W = aV \frac{\Gamma\left(\frac{2}{3}\right)^2}{\Gamma\left(\frac{1}{6}\right)^2} \left(\frac{1}{s^2} - \frac{1}{(2l+s)^2}\right),$$
(48)

where the central charge *a* is determined to be $a = \frac{R_{AdS}^3 \pi}{2G_N^{(5)}}$ [27, 28]. Remarkably, the S_{RG} (45) perfectly matches the EWCS (48), provided that $\lambda_1 = 2\Gamma\left(\frac{2}{3}\right)^2/\Gamma\left(\frac{1}{6}\right)^2$, in four-dimensional case.

V. CONCLUSION

In conclusion, we have explored the subtraction approach and the RG flow method for calculating the entanglement entropy $S_{\rm vN}$ of mixed states in CFT. By considering different configurations and applying the subtraction method, we derived expressions for the $S_{\rm vN}$ of two disjoint generic intervals in CFT₂. We also established connections between the mixed state covariant entanglement entropy and the bulk duals, and demonstrated the significance of the gauge parameters in determining the correspondence. Finally, we developed the RG flow approach to calculate $S_{\rm vN}$ and validated it within the AdS₅/CFT₄ correspondence.

Our proposal demonstrates that in CFT_2 , the covariant mixed state entanglement entropy can be calculated exactly. Therefore, with the same fashion presented in [8], the dual full spacetime geometry and dynamics, that is, the Einstein equation, can be derived from CFT_2 .

ACKNOWLEDGMENTS

Acknowledgments. This work is supported in part by NSFC (Grant No. 12105191, 12275183 and 12275184).

- Juan Martin Maldacena. The Large N limit of superconformal field theories and supergravity. Adv. Theor. Math. Phys., 2:231-252, 1998. arXiv:hep-th/9711200, doi:10.1023/A:1026654312961.
- [2] Edward Witten. Anti-de Sitter space and holography. Adv. Theor. Math. Phys., 2:253-291, 1998. arXiv:hep-th/9802150, doi:10.4310/ATMP.1998.v2.n2.a2.
- S. S. Gubser, Igor R. Klebanov, and Alexander M. Polyakov. Gauge theory correlators from noncritical string theory. *Phys. Lett. B*, 428:105-114, 1998. arXiv:hep-th/9802109, doi:10.1016/ S0370-2693(98)00377-3.
- Shinsei Ryu and Tadashi Takayanagi. Holographic derivation of entanglement entropy from AdS/CFT. *Phys. Rev. Lett.*, 96:181602, 2006. arXiv:hep-th/0603001, doi:10.1103/PhysRevLett.96.181602.
- [5] Shinsei Ryu and Tadashi Takayanagi. Aspects of holographic entanglement entropy. Journal of High Energy Physics, 2006(08):045–045, aug 2006. doi:10.1088/1126-6708/2006/08/045.
- Veronika E. Hubeny, Mukund Rangamani, and Tadashi Takayanagi. A Covariant holographic entanglement entropy proposal. JHEP, 07:062, 2007. arXiv:0705.0016, doi:10.1088/1126-6708/2007/ 07/062.

- [7] Mark Van Raamsdonk. Building up spacetime with quantum entanglement. Gen. Rel. Grav., 42:2323–2329, 2010. arXiv:1005.3035, doi:10.1142/S0218271810018529.
- [8] Xin Jiang, Peng Wang, Houwen Wu, and Haitang Yang. Einstein Equation Governs the Dynamics of Entanglement Entropy in CFT. 10 2024. arXiv:2410.19711.
- [9] Xin Jiang, Peng Wang, Houwen Wu, and Haitang Yang. Realization of "ER=EPR". 11 2024. arXiv: 2411.18485.
- [10] Pasquale Calabrese, John Cardy, and Erik Tonni. Entanglement negativity in extended systems: A field theoretical approach. J. Stat. Mech., 1302:P02008, 2013. arXiv:1210.5359, doi:10.1088/1742-5468/2013/02/P02008.
- [11] Tadashi Takayanagi and Koji Umemoto. Entanglement of purification through holographic duality. Nature Phys., 14(6):573-577, 2018. arXiv:1708.09393, doi:10.1038/s41567-018-0075-2.
- Kotaro Tamaoka. Entanglement Wedge Cross Section from the Dual Density Matrix. Phys. Rev. Lett., 122(14):141601, 2019. arXiv:1809.09109, doi:10.1103/PhysRevLett.122.141601.
- [13] Souvik Dutta and Thomas Faulkner. A canonical purification for the entanglement wedge cross-section. JHEP, 03:178, 2021. arXiv:1905.00577, doi:10.1007/JHEP03(2021)178.
- [14] Qiang Wen. Balanced Partial Entanglement and the Entanglement Wedge Cross Section. JHEP, 04:301, 2021. arXiv:2103.00415, doi:10.1007/JHEP04(2021)301.
- [15] Xin Jiang, Peng Wang, Houwen Wu, and Haitang Yang. An alternative to purification in CFT. 6 2024. arXiv:2406.09033.
- [16] Pasquale Calabrese and John L. Cardy. Entanglement entropy and quantum field theory. J. Stat. Mech., 0406:P06002, 2004. arXiv:hep-th/0405152, doi:10.1088/1742-5468/2004/06/P06002.
- [17] Pasquale Calabrese and John L. Cardy. Entanglement entropy and conformal field theory. J. Phys. A, 42:504005, 2009. arXiv:0905.4013, doi:10.1088/1751-8113/42/50/504005.
- [18] John L. Cardy. Boundary Conditions, Fusion Rules and the Verlinde Formula. Nucl. Phys. B, 324:581–596, 1989. doi:10.1016/0550-3213(89)90521-X.
- [19] John L. Cardy. Boundary conformal field theory. 11 2004. arXiv:hep-th/0411189.
- [20] John Cardy and Erik Tonni. Entanglement hamiltonians in two-dimensional conformal field theory. J. Stat. Mech., 1612(12):123103, 2016. arXiv:1608.01283, doi:10.1088/1742-5468/2016/12/123103.
- [21] Ian Affleck and Andreas W. W. Ludwig. Universal noninteger 'ground state degeneracy' in critical quantum systems. *Phys. Rev. Lett.*, 67:161–164, 1991. doi:10.1103/PhysRevLett.67.161.
- [22] Debarshi Basu, Himanshu Parihar, Vinayak Raj, and Gautam Sengupta. Entanglement negativity, reflected entropy, and anomalous gravitation. *Phys. Rev. D*, 105(8):086013, 2022. [Erratum: Phys.Rev.D 105, 129902 (2022)]. arXiv:2202.00683, doi:10.1103/PhysRevD.105.086013.
- [23] Qiang Wen and Haocheng Zhong. Covariant entanglement wedge cross-section, balanced partial entanglement and gravitational anomalies. *SciPost Phys.*, 13(3):056, 2022. arXiv:2205.10858, doi:10.21468/SciPostPhys.13.3.056.
- [24] H. Osborn. Weyl consistency conditions and a local renormalization group equation for general renor-

malizable field theories. Nucl. Phys. B, 363:486-526, 1991. doi:10.1016/0550-3213(91)80030-P.

- [25] Dmitri V. Fursaev and Sergey N. Solodukhin. On the description of the Riemannian geometry in the presence of conical defects. *Phys. Rev. D*, 52:2133-2143, 1995. arXiv:hep-th/9501127, doi: 10.1103/PhysRevD.52.2133.
- [26] Niko Jokela and Arttu Pönni. Notes on entanglement wedge cross sections. JHEP, 07:087, 2019. arXiv:1904.09582, doi:10.1007/JHEP07(2019)087.
- [27] Robert C. Myers and Aninda Sinha. Seeing a c-theorem with holography. Phys. Rev. D, 82:046006, 2010. arXiv:1006.1263, doi:10.1103/PhysRevD.82.046006.
- [28] Robert C. Myers and Aninda Sinha. Holographic c-theorems in arbitrary dimensions. JHEP, 01:125, 2011. arXiv:1011.5819, doi:10.1007/JHEP01(2011)125.