ON THE LYAPUNOV EXPONENT FOR THE RANDOM FIELD ISING TRANSFER MATRIX, IN THE CRITICAL CASE

ORPHEE COLLIN, GIAMBATTISTA GIACOMIN, RAFAEL L. GREENBLATT AND YUEYUN HU ´

ABSTRACT. We study the top Lyapunov exponent of a product of random 2×2 matrices appearing in the analysis of several statistical mechanical models with disorder, extending a previous treatment of the critical case (Giacomin and Greenblatt, ALEA 19 (2022), 701-728) by significantly weakening the assumptions on the disorder distribution. The argument we give completely revisits and improves the previous proof. As a key novelty we build a probability that is close to the Furstenberg probability, i.e. the invariant probability of the Markov chain corresponding to the evolution of the direction of a vector in \mathbb{R}^2 under the action of the random matrices, in terms of the ladder times of a centered random walk which is directly related to the random matrix sequence. We then show that sharp estimates on the ladder times (renewal) process lead to a sharp control on the probability measure we build and, in turn, to the control of its distance from the Furstenberg probability.

AMS subject classification (2010 MSC): 60K37, 82B44, 60K35, 37H15

Keywords: product of random matrices, singular behavior of Lyapunov exponents, disordered systems, renewal theory

1. INTRODUCTION

1.1. The model we consider. Let

$$
M = M(\varepsilon, Z) := \begin{pmatrix} 1 & \varepsilon \\ \varepsilon Z & Z \end{pmatrix}, \tag{1.1}
$$

with $|\varepsilon| < 1$ and $Z \in (0, \infty)$. Consider now an IID sequence of positive random variables $(Z_j)_{j=1,2,...}$ with $\mathbb{E}[|\log Z_1|] < \infty$ and let us assume that the distribution of Z is not trivial. The top Lyapunov exponent of the product of the random matrices $(M(\varepsilon, Z_j))_{j=1,2,...}$ is

$$
\mathcal{L}_Z(\varepsilon) := \lim_{n \to \infty} \frac{1}{n} \mathbb{E} \log \left\| M(\varepsilon, Z_1) M(\varepsilon, Z_2) \cdots M(\varepsilon, Z_n) \right\|, \tag{1.2}
$$

where $\|\cdot\|$ is an arbitrary matrix norm. $\mathcal{L}_Z(\varepsilon)$ is well defined and independent of the matrix norm. Existence of the limit is a straightforward consequence of a well-known property of subadditive sequences (Fekete's Lemma) that can be applied if $\|\cdot\|$ is a submultiplicative norm. The fact that all matrix norms are equivalent then yields that $\mathcal{L}_Z(\varepsilon)$ does not depend on the choice of the norm. This result holds for every $\varepsilon \in \mathbb{R}$, however it will be soon important to focus on invertible matrices, i.e. $|\varepsilon| \neq 1$ and we are interested in the behavior of $\mathcal{L}_Z(\varepsilon)$ when ε approaches 0, that is why we restrict to $|\varepsilon| < 1$.

Let us remark that if D is the diagonal matrix with $(1, -1)$ on the diagonal, we have $D = D^{-1}$ and $M(-\varepsilon, Z) = DM(\varepsilon, Z)D$, which implies $\mathcal{L}_Z(\varepsilon) = \mathcal{L}_Z(-\varepsilon)$ and so without loss of generality we may consider only $\varepsilon \in [0,1)$. Moreover

Date: January 16, 2025.

- (1) it is straightforward to verify that the conditions of Furstenberg Theorem (e.g., [\[3,](#page-32-0) Th. 4.1, p. 30) are verified for $\varepsilon \in (0,1)$ for the sequence of unit determinant matrices $(M(\varepsilon, Z_j)/\sqrt{(1-\varepsilon^2)Z_j})_{j=1,2,...}$ and this implies a number of consequences, in particular $\varepsilon \mapsto \mathcal{L}_Z(\varepsilon)$ is C^0 and that there is a formula for $\mathcal{L}_Z(\varepsilon)$ that we give and (crucially) exploit below.
- (2) For $\varepsilon \in (0,1)$ the problem we consider falls into the cone contraction theory of [\[30\]](#page-33-0) and $\varepsilon \mapsto \mathcal{L}_Z(\varepsilon)$ is real analytic.

The case $\varepsilon = 0$ is trivial in the sense that we explicitly compute $\mathcal{L}_Z(0) = \max(0, \mathbb{E}[\log Z])$. But the case $\varepsilon = 0$ fails to satisfy the hypotheses of Furstenberg Theorem: in fact the irreducibility property [\[3,](#page-32-0) Th. 4.1] fails because $M(0, Z)v \propto v$ both for $v = (1, 0)^t$ and for $v = (0, 1)^t$. So $\varepsilon = 0$ is a singular point in the sense that Furstenberg Theorem cannot be applied when $\varepsilon = 0$ and this motivates, at least from the mathematical viewpoint, the study of $\mathcal{L}_Z(\varepsilon)$ as $\varepsilon \to 0$: we will soon see that the motivation goes well beyond this.

We notice that if furthermore $\mathbb{E}[\log Z] = 0$, then not only the matrix is diagonal and fails the irreducibility property, but the exponential growth rate of the terms on the diagonal is the same (zero in both cases), that suggests that this case is particularly singular (lack of hyperbolicity). The $\mathbb{E}[\log Z] = 0$ case is naturally referred as critical: we will come back to discuss and confirm this aspect in Sec. [1.3.](#page-2-0)

A change of perspective on the problem is notationally useful, particularly in the critical case: we set

$$
\Gamma := -\log \varepsilon \quad \text{and} \quad z := \log Z, \tag{1.3}
$$

so the original matrix M of (1.1) becomes

$$
\begin{pmatrix} 1 & \exp(-\Gamma) \\ \exp(-\Gamma + z) & \exp(z) \end{pmatrix}, \tag{1.4}
$$

and we set

$$
\mathcal{L}(\Gamma) := \mathcal{L}_Z(\varepsilon). \tag{1.5}
$$

In analogy with before, $z_j = \log Z_j$, so $(z_j)_{j=1,2,...}$ is an IID sequence. With this notation the critical case means

$$
\mathbf{z} \in \mathbb{L}^{1} \quad \text{and} \quad \mathbb{E}[\mathbf{z}] = \int_{\mathbb{R}} x \zeta(\mathrm{d}x) = 0, \tag{1.6}
$$

where ζ is the law of z.

1.2. Our main result. In [\[19\]](#page-33-1) two of the authors proved the following result:

Theorem 1.1. Choose ζ that satisfies [\(1.6\)](#page-1-0) and such that

(h-1): ζ has a density and its density is Hölder continuous uniformly over \mathbb{R} ; (h-2): there exists $c > 0$ such that $\int_{\mathbb{R}} \exp(c|x|) \zeta(\mathrm{d}x) < \infty$. Then there exists $\kappa_1 > 0$, $\kappa_2 \in \mathbb{R}$ and $\delta > 0$ such that

$$
\mathcal{L}(\Gamma) \stackrel{\Gamma \to \infty}{=} \frac{\kappa_1}{\Gamma + \kappa_2} + O\left(\exp\left(-\delta\Gamma\right)\right). \tag{1.7}
$$

Here we prove the following result: for $n = 1, 2, \ldots$ denote by ζ^{*n} the law of $z_1 + \ldots + z_n$.

Theorem 1.2. Choose ζ that satisfies [\(1.6\)](#page-1-0) and such that **(H-1):** there exists $n_0 \geq 1$ such that ζ^{*n_0} has a density;

and either

(H-2): there exists $\xi > 5$ such that $\int_{\mathbb{R}} |x|^{\xi} \zeta(\mathrm{d}x) < \infty$; or

(H-2'): there exists $c > 0$ such that $\int_{\mathbb{R}} \exp(c|x|) \zeta(\mathrm{d}x) < \infty$.

Then there exists $\kappa_1 > 0$ and $\kappa_2 \in \mathbb{R}$ such that

$$
\mathcal{L}(\Gamma) \stackrel{\Gamma \to \infty}{=} \frac{\kappa_1}{\Gamma + \kappa_2} + R(\Gamma). \tag{1.8}
$$

where for $\Gamma \to \infty$

$$
R(\Gamma) = \begin{cases} O(\Gamma^{-(\xi-4)}) & under (\mathbf{H-2}), \\ O(\exp(-\delta \Gamma)) & under (\mathbf{H-2}'), \end{cases}
$$
(1.9)

for a suitable choice of $\delta > 0$.

Let us stress the elementary fact that

$$
\frac{\kappa_1}{\Gamma + \kappa_2} = \kappa_1 \sum_{j=1}^{\infty} (-1)^{j-1} \kappa_2^{j-1} \Gamma^{-j}, \qquad (1.10)
$$

so that [\(1.7\)](#page-1-1) provides a development of $\mathcal{L}(\Gamma)$ to all orders. This is also the case for [\(1.8\)](#page-2-1) if (H-2) holds for ξ arbitrarily large, but of course this only gives $R(\Gamma) = O(\Gamma^{-C})$ for every $C > 0$. An exponential control on the rest is recovered under hypothesis $(H-2') = (h-2)$.

When $\xi < \infty$, i.e. $\xi \in (5, \infty)$ then (1.8) is equivalent to

$$
\mathcal{L}(\Gamma) \stackrel{\Gamma \to \infty}{=} \kappa_1 \sum_{j=1}^{\lfloor \xi \rfloor - 4} (-1)^{j-1} \kappa_2^{j-1} \Gamma^{-j} + O\left(\Gamma^{-(\xi - 4)}\right). \tag{1.11}
$$

with [\cdot] the lower integer part of \cdot . Therefore for $\xi \in (5,6]$ we only have $\mathcal{L}(\Gamma) = \kappa_1 \Gamma^{-1} +$ $O(\Gamma^{-1-\epsilon})$, with $\epsilon = \xi - 5$, and κ_2 enters the game only for $\xi > 6$.

An important point of these results is that we do have expressions for κ_1 and κ_2 – they are in (6.4) and (3.13) – and they are not simple expressions because they depend on the invariant measure of a suitable null recurrent Markov Chain (MC). The dependence on invariant measures of MC's is not surprising because this is in the nature of the Furstenberg expression for $\mathcal{L}(\Gamma)$ (see [\(2.4\)](#page-6-0) below). However one of us proved [\[6\]](#page-32-1), by using the statistical mechanics origin of the question and avoiding the Furstenberg expression for $\mathcal{L}(\Gamma)$, that κ_1 is equal to the variance of the external magnetic field (see also Sec. [1.3\)](#page-2-0), which in the variables we use here means that $4\kappa_1$ is the variance of z, i.e. $\int_{\mathbb{R}} x^2 \zeta(dx)$. This very intriguing point, proven in [\[6\]](#page-32-1) under weaker assumptions than Theorem [1.2,](#page-1-2) is picked up again in Sec. [1.3.](#page-2-0) Moreover in App. [A](#page-29-0) we show that the precise estimates we obtain can be used to provide an alternative proof (under stronger assumptions) of the result in [\[6\]](#page-32-1).

1.3. Origin of the question and discussion of the literature. The key matrix M in (1.1) is proportional to the transfer matrix of the one dimensional Ising model (see e.g. [\[23,](#page-33-2) Ch. 3] or [\[12\]](#page-32-2)):

$$
\begin{pmatrix} e^{J+h} & e^{-J+h} \ e^{-J-h} & e^{J-h} \end{pmatrix} = e^{J+h} \begin{pmatrix} 1 & e^{-2J} \ e^{-2J-2h} & e^{-2h} \end{pmatrix} = e^{J+h} M \left(e^{-2J}, e^{-2h} \right), \quad (1.12)
$$

where h is the external field and J the interaction. Explicitly, the partition function of the Ising model of length n with periodic boundary conditions and with inhomogeneous external field h_1, h_2, \ldots, h_n is equal to $\exp(JN + \sum_{j=1}^n h_j)$ times the trace of

 $M\left(e^{-2J},e^{-2h_1}\right)\cdots M\left(e^{-2J},e^{-2h_n}\right)$, where we consider $(h_n)_{n\geq 1}$ as an IID sequence of random variables, distributed as h . Hence the Lyapunov exponent we consider is the free energy density of the random external field one dimensional Ising model minus J. The same transfer matrix product and Lyapunov exponent enter the exact expression for the free energy density of the two dimensional Ising model with zero external field and *colum*nar disorder [\[24,](#page-33-3) [25,](#page-33-4) [9\]](#page-32-3). The seminal works in the analysis of this matrix in statistical mechanics of disordered systems are [\[24\]](#page-33-3) and [\[12\]](#page-32-2);

- (1) In the first work, McCoy and Wu provide a solution to a continuum model arising in the weak disorder limit of the matrix product model we consider (a mathematical approach to this limit can be found in [\[9\]](#page-32-3) and solvable continuum limits for a much greater class of models may be found in [\[10\]](#page-32-4)). The Lyapunov exponent of the continuum model is explicitly expressed in terms of Bessel functions, and one easily obtains an arbitrarily precise analysis, notably in the $J \to \infty$ limit: the e^{-2J} appearing in the matrix in the 2d Ising model considered in [\[24\]](#page-33-3) is not an interaction term, but rather a *phase* parameter and key to understanding the phase transition is the singularity arising near phase zero, which explains the link with the $J \to \infty$ limit of the 1d case.
- (2) In the second work, Derrida and Hilhorst provide an analysis of the $J \to \infty$ limit of the Lyapunov exponent in the case where the expectation of h is nonzero, targeting directly the one dimensional case, but mentioning several other cases in which this limit is relevant, see [\[12\]](#page-32-2), but also the monograph [\[11\]](#page-32-5). The case of centered h has been first taken up in [\[28\]](#page-33-5).

For the physical relevance of the analysis of the $J \to \infty$ and of why the $\mathbb{E}[h] = 0$ case is critical we refer in particular to $[12, 11, 25, 9]$ $[12, 11, 25, 9]$ $[12, 11, 25, 9]$ $[12, 11, 25, 9]$. Here we insist on the fact that our main result [\(1.8\)](#page-2-1) arises in the exactly solvable McCoy-Wu continuum model, with exponentially small rest term $R(\Gamma)$ (in fact, the sharp behavior of this term is known in this case): in the continuum model the noise, or disorder, is Brownian, hence with Gaussian tails. A priori it is not obvious to which extent the result obtained in the continuum limit corresponds to the behavior of the original matrix product: in the off-critical case, i.e. $\mathbb{E}[h] \neq 0$, the leading behavior of $\mathcal{L}(\Gamma)$ [\[12\]](#page-32-2) corresponds to the leading order of the continuum case [\[9\]](#page-32-3), and this has been mathematically established in [\[18\]](#page-33-6) when the disorder is compactly supported and has a C^1 density. For the $\mathbb{E}[h] = 0$ the sharp result [\(1.8\)](#page-2-1), i.e. a development to all orders in powers of $1/\Gamma$, is claimed in [\[28\]](#page-33-5) for a special compactly supported disorder law for which some exact computations are possible, but the analysis is not rigorous and no control on the error term $R(\Gamma)$ is given. This naturally sets forth the issue of understanding, at a mathematical level and for matrix products, the validity of the result emerging in the continuum model.

A step in this direction has been performed in [\[19\]](#page-33-1) where Theorem [1.1](#page-1-3) is established: [\(1.8\)](#page-2-1) holds with an exponential control on $R(\Gamma)$ under the assumption that disorder has exponential tails and uniformly Hölder- C^0 density of the disorder.

Here we greatly generalize the result in [\[19\]](#page-33-1): in particular, we establish the result without assuming regularity of the density and, above all, we work with disorder power law tails. In this case the error term $R(\Gamma)$ has a power law behavior and therefore the free energy development holds only up to a finite order, see (1.11) .

Like in [\[18,](#page-33-6) [19\]](#page-33-1), our method is based on constructing an explicit measure, the Derrida-Hilhorst probability proposed in [\[12\]](#page-32-2) – we introduce it in [\(3.11\)](#page-10-1) and we call it γ_{Γ} – that is expected to be close to the Furstenberg measure (that we call ν_{Γ}), which is the unique invariant probability associated to the action of the matrix product in the projective space [\[3\]](#page-32-0), from which the Lyapunov exponent is computed in a straightforward way, see [\(2.4\)](#page-6-0). The construction of the Derrida-Hilhorst probability and showing that it is indeed close to the Furstenberg measure is the mathematical challenge. The Derrida-Hilhorst probability is built by using the invariant measures of two associated Markov chains: when $\mathbb{E}[h] \neq 0$ one chain is positive recurrent and the other is transient, while in the $\mathbb{E}[h] = 0$ both chains are null recurrent, so the invariant measures are not finite measures. These Markov chains constitute a classical and much studied problem in probability and statistics, the random difference equation " $X = AX + B$ ", see for example [\[4,](#page-32-6) [22\]](#page-33-7). In this context, the case $\mathbb{E}[h] = 0$ is naturally considered critical in the sense of random dynamical systems [\[2,](#page-32-7) [4\]](#page-32-6). A contribution of our work is also providing a very precise solution to this problem in a special case.

The main novelty of the current work, presented in Section [4,](#page-11-0) is in the analysis of the invariant measure of the null recurrent Markov chains $Q_{n+1} = Z_{n+1}(Q_n + 1)$, with $Q_0 \ge 0$ and $Z_{n+1} = \exp(-2h_{n+1}),$ with $\mathbb{E}[h] = 0$: the process Y introduced in Sec. [2.3](#page-7-0) is log Q. In the critical case $Y = \log Q$ becomes a centered random walk – the increments are $(-2h_i)$ – with a force term that biases the walk against entering the negative half-line. The key to this treatment is exploiting renewal estimates to obtain a precise control on the ladder process of the centered random walk and use it to control the invariant measure of the Markov chain Y (or, equivalently, Q).

Once the invariant measure of the Y process is under control the Derrida-Hilhorst probability γ_{Γ} can be built and we are left with showing that γ_{Γ} is close to the Furstenberg probability ν_{Γ} . This part follows the two step procedure used in [\[19\]](#page-33-1), and before that in [\[18\]](#page-33-6) in the noncritical case. Both steps are improved with respect to [\[19\]](#page-33-1):

- (1) One step consists in establishing a residual contraction property of the action of the Furstenberg Markov chain in the critical case: at criticality the spectral gap vanishes with the strength of the interaction $\Gamma = 2J$ and in [\[19\]](#page-33-1) an estimate $O(\log \Gamma/\Gamma^2)$ is established via diffusion approximation. Here we exploit a very different, more microscopic argument that yields $O(1/\Gamma^2)$ and it is a probabilistic version of an argument due to Q. Moulard [\[27\]](#page-33-8).
- (2) The other step consists in showing, in a quantitative way and in a precise sense, that the one step action of the Furstenberg Markov chain does not modify much the Derrida-Hilhorst probability γ_{Γ} : the point here is generalizing the one-step bound developed in [\[19\]](#page-33-1) from exponential to power law disorder tail.

The result in [\[6\]](#page-32-1), i.e. that $\mathcal{L}(\Gamma) \sim \text{var}(\mathbf{z})/(4\Gamma)$ for $\xi > (3+\sqrt{5})/2 = 2.618...$, shows that our moment condition $\xi > 5$ is not optimal. An improvement in this direction appears to be rather challenging, since it seems to require new ideas about how to construct the invariant law of the Y process, or a new approach, i.e. not involving the Derrida-Hilhorst construction, in proving properties of the invariant probability ν_{Γ} . We signal however that our approach does provide additional information, in particular a by-product is that if [\(1.6\)](#page-1-0) and (**H-1**) hold along with $z \in \mathbb{L}^{\xi}$ for $\xi > 4$ we have

$$
d_{W,1}(\nu_{\Gamma}, \gamma_{\Gamma}) = O\left(\Gamma^{-(\xi - 4)}\right), \qquad (1.13)
$$

where $d_{W,1}$ is the Wasserstein–1 distance: [\(1.13\)](#page-4-0) is a direct consequence of Corollary [3.3](#page-9-0) and [\(3.16\)](#page-11-1) of Proposition [3.5.](#page-10-2)

We remark also that the fact that $\kappa_1 = \text{var}(\mathbf{z})/4$ is strongly suggested by Fisher Renormalization Group (RG) approach [\[16\]](#page-32-8). In [\[7,](#page-32-9) [8\]](#page-32-10) we have proven results that confirm the application of Fisher RG to the critical random field $1d$ Ising model in [\[17\]](#page-32-11). The main results in $[7, 8]$ $[7, 8]$ aim at understanding how the critical random field 1d Ising configurations look like when $\Gamma = 2J \to \infty$. In particular, in [\[8\]](#page-32-10) the results are shown assuming only that $z \in \mathbb{L}^2$ (i.e. $\xi = 2$) centered and nontrivial. Moreover, as already discussed at length in [\[7,](#page-32-9) [8\]](#page-32-10), the application of Fisher RG in [\[17\]](#page-32-11) does not yield the correct behavior of $\mathcal{L}(\Gamma)$ beyond the leading order. This is probably not surprising because the subleading correction is not expected to be universal. And even restricting to the 1d Ising model, we see that the subleading correction involves κ_2 , which appears to depend on details of the law of z (even if this is only based on numerical evidence at this stage). Nonetheless we stress that our result sets forth another, somewhat surprising, form of universality: given the variance of z, the subleading corrections to all order in powers of $1/\Gamma$ depend only on the constant κ_2 .

1.4. Organisation of the rest of the paper. In section [2](#page-5-0) we recall the necessary tools from Furstenberg theory and we introduce the MC whose invariant measure is used to build the probability γ_{Γ} that will be shown to be a good approximation of the Furstenberg probability ν_{Γ} . In Sec. [3](#page-8-0) we state three results – Theorem [3.1,](#page-8-1) Theorem [3.2](#page-9-1) and Proposition [3.5](#page-10-2) – that, combined, yield a proof of Theorem [1.2.](#page-1-2) The next three sections are devoted to the proof of these three results.

2. Tools from Furstenberg theory

2.1. The Furstenberg expression for $\mathcal{L}(\Gamma)$. Key to the Furstenberg theory is understanding the action of our random matrices [\(1.4\)](#page-1-4) on the projective space which in our case is simply one dimensional because $P(\mathbb{R}^2) \cong \mathbb{R}$. If we parametrize $P(\mathbb{R}^2) \cong \mathbb{R}$ with the coordinates $(1, \exp(x))^T$ we readily see that the action of the matrix (1.4) is

$$
x \mapsto z + \log\left(\frac{e^{-\Gamma} + e^x}{1 + e^{x - \Gamma}}\right) = z + h_{\Gamma}(x). \tag{2.1}
$$

We observe that $h_{\Gamma}(\cdot)$ is odd and $x \mapsto x - h_{\Gamma}(x)$, which is also odd, is small if $x \in (-\Gamma, \Gamma)$ and far from the boundary points $\pm \Gamma$:

$$
x - h_{\Gamma}(x) = \log\left(\frac{1 + e^{x - \Gamma}}{1 + e^{-x - \Gamma}}\right) = O\left(e^{-(\Gamma - x)}\right) + O\left(e^{-(\Gamma + x)}\right),\tag{2.2}
$$

so $h_{\Gamma}(x)$ is very close to x on an interval that approaches R in the limit $\Gamma \to \infty$ (Fig. [1\)](#page-5-1).

FIGURE 1. For $\Gamma = 10$: on the left the plot of $h_{\Gamma}(\cdot)$ (solid line) and of $x \mapsto h_{\Gamma}(x) =$ $h(x + \Gamma) - \Gamma$ (dotted line); on the right the derivative of the same functions.

Denote by $X = (X_n)_{n=0,1,\dots}$ the MC generated by the map [\(2.1\)](#page-5-2), that is

$$
X_{n+1} = \mathbf{z}_{n+1} + h_{\Gamma}(X_n) \tag{2.3}
$$

Since the image of h_{Γ} is $(-\Gamma,\Gamma)$ the process (X_n) hardly leaves $(-\Gamma,\Gamma)$: in fact we may remark that, no matter what is the value taken by X_n , $|X_{n+1}| \leq \Gamma + |z_{n+1}|$. Moreover when the process is in $(-\Gamma, \Gamma)$ and far from the boundaries it is *close to being* a random walk with centered increments.

The MC X has a unique invariant probability ν_{Γ} : this can be established just under the assumption that z is not trivial either by checking the assumptions of the Furstenberg Theorem [\[3\]](#page-32-0) or by remarking the contractive property of this MC (this is detailed in [\[8,](#page-32-10) Lemma 2.2]) and we have that

$$
\mathcal{L}(\Gamma) = \int_{\mathbb{R}} \log(1 + \exp(-\Gamma + x)) \nu_{\Gamma}(\mathrm{d}x). \tag{2.4}
$$

Remark 2.1. Formula [\(2.4\)](#page-6-0) is what follows by specializing the Furstenberg formula for the top Lyapunov exponent [\[3,](#page-32-0) Th. 3.6] to our context. More precisely, this is one of the possible equivalent formulas that one finds applying Furstenberg formula with different matrix norms, or even semi-norms, because it is a byproduct of the theory that for positive entry matrices, the exponential rate of growth of every entry of the matrix product is given by the top Lyapunov exponent. In particular we have also

$$
\mathcal{L}(\Gamma) = \int_{\mathbb{R}} \log(1 + \exp(-\Gamma - x)) \nu_{\Gamma}(\mathrm{d}x), \qquad (2.5)
$$

that a priori may look surprising, but from [\(2.3\)](#page-6-1) one obtains that $\int_{\mathbb{R}} |x| \nu_{\Gamma}(\mathrm{d}x) \leq \Gamma +$ $\mathbb{E}[\vert \mathbf{z} \vert] < \infty$ and $\int_{\mathbb{R}} (x - h_{\Gamma}(x)) \nu_{\Gamma}(\mathrm{d}x) = 0$, and this identity offers a direct way to see that the two expressions for the Lyapunov exponent in [\(2.4\)](#page-6-0) and [\(2.5\)](#page-6-2) coincide.

2.2. Building the framework to estimate $\mathcal{L}(\Gamma)$. In general, if A is a real-valued random variable, we use the notations $G_A(x) = \mathbb{P}(A > x)$ and $F_A(x) = 1 - G_A(x)$: G_A is the risk (distribution) function of A and F_A is the cumulative (distribution) function of A. Likewise we speak of risk and cumulative functions of a probability on $\mathbb R$ and we use this notation and terminology also for finite measures μ on \mathbb{R} : $G_{\mu}(x) = \mu((x,\infty))$ and $F_{\mu}(x) = \mu((-\infty, x]).$

Remark 2.2. Later on, we will work also with nonnormalizable (σ -finite) measures, notably with measures μ such that $\mu((-\infty, x]) < \infty$ for every x, and the notation $F_{\mu}(x)$ will be used in this case as well. However, unless specified otherwise, μ is a finite measure.

For $G = G_\mu$, we define $T_\Gamma G$ by

$$
T_{\Gamma}G(x) := G(-\infty)G_{\mathbf{z}}((\Gamma + x)_{-}) + \int_{(-\Gamma + x,\Gamma + x)} G\left(h_{\Gamma}^{-1}(x - z)\right)\zeta(\mathrm{d}z), \quad \forall x \in \mathbb{R}, \quad (2.6)
$$

where $G_{\mathbf{z}}(y_-) := \lim_{x \uparrow y} G_{\mathbf{z}}(x)$ for $y \in \mathbb{R}$ and as before, ζ denotes the law of \mathbf{z} . Similarly for $F = F_{\mu}$, we define

$$
T_{\Gamma}F(x) := F(\infty)F_{\mathbf{z}}(-\Gamma + x) + \int_{(-\Gamma + x,\Gamma + x)} F\left(h_{\Gamma}^{-1}(x - z)\right)\zeta(\mathrm{d}z), \quad \forall x \in \mathbb{R}.
$$
 (2.7)

Lemma 2.3. We have for $n = 0, 1, ...,$

$$
G_{X_{n+1}} = T_{\Gamma} G_{X_n},\tag{2.8}
$$

and

$$
F_{X_{n+1}} = T_{\Gamma} F_{X_n} \,. \tag{2.9}
$$

Proof. By using that $h_{\Gamma}(\cdot)$ is an increasing bijection from $\mathbb R$ to $(-\Gamma,\Gamma)$ we see that

$$
G_{X_{n+1}}(x) = \mathbb{P}(\mathbf{z}_{n+1} + h_{\Gamma}(X_n) > x)
$$

\n
$$
= \mathbb{P}(x - \mathbf{z}_{n+1} \leq -\Gamma) + \mathbb{P}(h_{\Gamma}(X_n) > x - \mathbf{z}_{n+1}; |x - \mathbf{z}_{n+1}| < \Gamma)
$$

\n
$$
= \mathbb{P}(\mathbf{z} \geq x + \Gamma) + \mathbb{P}(X_n > h_{\Gamma}^{-1}(x - \mathbf{z}_{n+1}); |x - \mathbf{z}_{n+1}| < \Gamma)
$$

\n
$$
= G_{\mathbf{z}}((\Gamma + x)_{-}) + \int_{(-\Gamma + x, \Gamma + x)} G_{X_n} (h_{\Gamma}^{-1}(x - z)) \zeta(\mathrm{d}z).
$$
\n(2.10)

The proof for (2.9) is analogous. \Box

The map T_{Γ} can then be extended by linearity to act on G defined by $G(x) = G_{\nu+}(x) G_{\nu_{-}}(x), \nu_{\pm}$ finite measures (so $|G(-\infty)| < \infty$):

$$
T_{\Gamma}G(x) = G(-\infty)G_{\mathbf{z}}((\Gamma + x)_{-}) + \int_{(-\Gamma + x, \Gamma + x)} G\left(h_{\Gamma}^{-1}(x - z)\right)\zeta(\mathrm{d}z). \tag{2.11}
$$

In the applications we consider ν is the difference of two probability measures, so $G(-\infty)$ = 0 and T_{Γ} reduces to $T_{\Gamma,0}$, with

$$
T_{\Gamma,0}G(x) := \int_{(-\Gamma+x,\Gamma+x)} G\left(h_{\Gamma}^{-1}(x-z)\right)\zeta(\mathrm{d}z), \quad \forall x \in \mathbb{R},\tag{2.12}
$$

which is well defined also in the slightly different set-up of $G \in \mathbb{L}^1$. In fact, by the Fubini-Tonelli Theorem and since h'_{Γ} takes values in $(0, 1)$ we have

$$
\begin{split} \left\|T_{\Gamma,0}G\right\|_{1} &\leq \int_{\mathbb{R}} \left(\int_{\left(-\Gamma+x,\Gamma+x\right)} \left|G\left(h_{\Gamma}^{-1}(x-z)\right)\right|\zeta(\mathrm{d}z) \right) \mathrm{d}x \\ &= \int_{\mathbb{R}} \left(\int_{\mathbb{R}} \left|G\left(y\right)\right|h_{\Gamma}'(y)\mathrm{d}y \right) \zeta(\mathrm{d}z) \leq \|G\|_{1} \,. \end{split} \tag{2.13}
$$

2.3. Looking from the edge: the reduced chain Y. If we sit on $-\Gamma$, that is if we make it our new origin, in the limit as $\Gamma \to \infty$ the MC becomes

$$
Y_{n+1} = \mathbf{z}_{n+1} + h(Y_n) \quad \text{with } h(y) = y + \log(1 + \exp(-y)). \tag{2.14}
$$

Note in particular that, since $h(\cdot)$ is positive and increasing, we have

$$
F_{Y_{n+1}}(x) = \mathbb{P}(\mathbf{z}_{n+1} + h(Y_n) \le x, x - \mathbf{z}_{n+1} > 0)
$$

= $\mathbb{P}(Y_n \le h^{-1} (x - \mathbf{z}_{n+1}), \mathbf{z}_{n+1} < x)$
= $\int \mathbf{1}_{(-\infty,x)}(z) F_{Y_n} (h^{-1} (x - z)) \zeta(\mathrm{d}z).$ (2.15)

Proposition 2.4. Assume [\(1.6\)](#page-1-0) and that there exists $p > 2$ such that $z \in \mathbb{L}^p$. Then the MC Y has a unique nonzero invariant Radon measure ν . It satisfies that $\nu((-\infty, x]) < \infty$ for every $x \in \mathbb{R}$, $\nu((-\infty, x]) \leq \nu((-\infty, 0]) \mathbb{P}(\mathbf{z} < x)$ for $x \leq 0$ and $\nu(\mathbb{R}) = \infty$. Moreover, Y is recurrent in the sense that if $O \subset \mathbb{R}$ is an open set with $\nu(O) > 0$, then, given any $Y_0, \, \mathbb{P}(Y_n \in O \, i.o.) = 1.$

For nonnormalizable measures uniqueness is of course meant up to a multiplicative *constant* and we note that ν is characterized by

$$
\int g(z)\nu(\mathrm{d}z) = \iint g(z+h(y))\zeta(\mathrm{d}z)\,\nu(\mathrm{d}y) \quad \text{for every measurable } g \ge 0. \tag{2.16}
$$

We recall that a measure ν on R, equipped with the Borel σ -algebra $\mathcal{B}(\mathbb{R})$, is Radon if the measure of compact sets is finite and if it is regular, i.e. if for every $B \in \mathcal{B}(\mathbb{R})$ we have $\nu(B)$ is equal both to the infimum of $\nu(O)$ taken over O open that contains B, and to the supremum of $\nu(K)$ taken over K compact contained in B.

Proof of Prop. [2.4.](#page-7-2) If we set $R_n := \exp(Y_n)$ then, with $Z_{n+1} = \exp(z_{n+1}), R_{n+1} =$ $Z_{n+1}(R_n + 1)$ and R is a MC with state space $(0, \infty)$. This is a special case of the process considered in [\[2\]](#page-32-7). So by [\[2,](#page-32-7) Proposition 1.1 and Theorem 3.3] we know that R has a unique nonzero invariant Radon measure m and that this measure satisfies $m((0,\infty)) = \infty$. Of course the MC R with state space $(0, \infty)$ is equivalent to the MC Y with state space R, in particular if we introduce the measure ν by setting $\nu((s,t)) = m((\exp(s), \exp(t)))$ for every $s < t$, it is straightforward to see that ν is invariant for Y and that ν inherits the properties of m. By the bound [\[19,](#page-33-1) (A.2)] one infers that $\nu((-\infty, t]) < \infty$ for every $t \in \mathbb{R}$. Recurrence of the process is proven in [\[2,](#page-32-7) Cor. 4.2(*i*)]. Finally, the explicit bound on $F_{\nu}(x) := \nu((-\infty, x])$ holds because F_{ν} satisfies the analogue of [\(2.15\)](#page-7-3), that is

$$
F_{\nu}(x) = \int_{(-\infty, x)} F_{\nu} \left(h^{-1} \left(x - z \right) \right) \zeta(\mathrm{d} z) \,. \tag{2.17}
$$

Using that $F_{\nu}(\cdot)$ and $h^{-1}(\cdot)$ are nondecreasing and that $h^{-1}(y) < y$, for every y, we see that for $z < x \leq 0$ we have $F_{\nu} (h^{-1} (x - z)) \leq F_{\nu} (h^{-1} (-z)) \leq F_{\nu} (-z) \leq F_{\nu} (0)$, hence $F_{\nu}(x) \leq F_{\nu}(0)\mathbb{P}(\mathbf{z}$

3. The proof of the main result

We give here a series of three statements that we prove in the next sections. We then prove Theorem [1.2](#page-1-2) assuming the validity of these statements.

3.1. On the invariant measure of the Y process. Recall that $(z_n)_{n=1,2,...}$ is an IID sequence with $z_1 \stackrel{\text{(law)}}{=} z$. We introduce (S_n) the corresponding random walk starting at 0: $S_n := \sum_{j=1}^n z_j$ for $n = 1, 2, \ldots$ and $S_0 = 0$. Of course this random walk is null recurrent.

Recall [\(2.14\)](#page-7-4) for the definition of the Markov chain (Y_n) .

Theorem 3.1. Assume [\(1.6\)](#page-1-0) and $(H-1)$ of Theorem [1.2](#page-1-2) plus the following two conditions: (T-1): there exists $\xi > 3$ such that $\mathbb{E}[(z^+)^{\xi}] < \infty$; $(T-2): \mathbb{E}[z^2] \in (0,\infty).$

Then any nonzero invariant measure ν of Y satisfies that

$$
\nu((-\infty, x]) \stackrel{x \to \infty}{=} c_{\nu} x + d_{\nu} + O(x^{-(\xi - 3)}).
$$
\n(3.1)

for appropriately chosen constants $c_{\nu} > 0$ and $d_{\nu} \in \mathbb{R}$. Moreover, if $(T-1)$ is replaced by

(T-1'): there exists $c > 0$ such that $\mathbb{E}[\exp(c z^+)] < \infty$;

then [\(3.1\)](#page-8-2) holds with error term $O(\exp(-\delta x))$, for some positive constant δ .

The proof of Theorem [3.1](#page-8-1) is in Section [4.](#page-11-0) When applying Theorem [3.1](#page-8-1) we will set $c_{\nu} = 1$, see [\(3.10\)](#page-10-3).

3.2. Contraction estimates. Next is the following improvement of [\[19,](#page-33-1) Prop. 2.2], based on a result and approach developed by Q. Moulard [\[27\]](#page-33-8).

Theorem 3.2. Choose ζ satisfying [\(1.6\)](#page-1-0) and such that $\sigma^2 := \mathbb{E}[z^2] < \infty$. Then there exists Γ_0 such that for every $\Gamma \geq \Gamma_0$ and every $G = G_1 - G_2$, with G_1 and G_2 risk functions of \mathbb{L}^1 random variables, we have

$$
\sum_{n=0}^{\infty} \left\| T_{\Gamma,0}^n G \right\|_1 \le \frac{75}{\sigma^2} \Gamma^2 \left\| G \right\|_1 ,\tag{3.2}
$$

where $T_{\Gamma,0}$ is defined in [\(2.12\)](#page-7-5).

If G is the risk function of a random variable we introduce

$$
L_{\Gamma}[G] := \int_{\mathbb{R}} \frac{1}{1 + e^{\Gamma - x}} G(x) dx, \qquad (3.3)
$$

and one readily sees that, from [\(2.4\)](#page-6-0), the Lyapunov exponent $\mathcal{L}(\Gamma)$ is equal to $L_{\Gamma}[G_{\nu_{\Gamma}}],$ and that

$$
|L_{\Gamma}[G_1] - L_{\Gamma}[G_2]| \leq ||G_1 - G_2||_1, \qquad (3.4)
$$

whenever G_1 and G_2 are the risk functions of two random variables. We thus have the following important corollary to Theorem [3.2:](#page-9-1)

Corollary 3.3. With Γ_0 as in Theorem [3.2,](#page-9-1) for $\Gamma \geq \Gamma_0$ and for any probability γ such that $\int_{\mathbb{R}} |x| \gamma(\mathrm{d}x) < \infty$ (γ may depend on Γ), we have

$$
|\mathcal{L}(\Gamma) - L_{\Gamma}[G_{\gamma}]| \leq ||G_{\nu_{\Gamma}} - G_{\gamma}||_1 \leq \frac{75}{\sigma^2} \Gamma^2 ||T_{\Gamma} G_{\gamma} - G_{\gamma}||_1.
$$
 (3.5)

Remark 3.4. The alternative expression for $\mathcal{L}(\Gamma)$ in [\(2.5\)](#page-6-2) suggests introducing

$$
L_{\Gamma}^{\star}[G] := \int_{\mathbb{R}} \frac{1}{1 + e^{\Gamma + x}} (1 - G(x)) dx
$$
 (3.6)

instead of L_{Γ} . In general $L_{\Gamma}[G] \neq L_{\Gamma}^{\star}[G]$, but $L_{\Gamma}[G_{\nu_{\Gamma}}] = L_{\Gamma}^{\star}[G_{\nu_{\Gamma}}]$. Moreover Corollary [3.3](#page-9-0) applies verbatim to $L_{\Gamma}^{\star}[\,\cdot\,].$

Proof. Choose X_0 of law γ and X_1 as in [\(2.3\)](#page-6-1). By construction $G_{X_1} = T_{\Gamma} G_{\gamma}$ and, by the observation made right after [\(2.3\)](#page-6-1), we have that $|X_1| \leq \Gamma + |\mathbf{z}_1| \in \mathbb{L}^2$. Since the Wasserstein-1 distance between the two \mathbb{L}^1 random variables X_0 and X_1 is equal to $||T_{\Gamma}G_{\gamma}-G_{\gamma}||_1$, we have that $||T_{\Gamma}G_{\gamma}-G_{\gamma}||_1 < \infty$.

Since the MC (X_n) is positive recurrent, $\lim_{N\to\infty} T_{\Gamma}^N G_{\gamma}(x) = G_{\nu_{\Gamma}}(x)$ for every $x \in \mathbb{R}$ which is a continuity point of $G_{\nu_{\Gamma}}(\cdot)$, hence this convergence holds out of a set that is at most countable. Therefore, by Fatou's Lemma and Theorem [3.2,](#page-9-1) we have that

$$
\|G_{\nu_{\Gamma}} - G_{\gamma}\|_{1} \leq \liminf_{N \to \infty} \|T_{\Gamma}^{N} G_{\gamma} - G_{\gamma}\|_{1}
$$
\n(3.7)

$$
\leq \sum_{n=0}^{\infty} \left\| T_{\Gamma,0}^{n} \left(T_{\Gamma} G_{\gamma} - G_{\gamma} \right) \right\|_{1} \tag{3.8}
$$

$$
\leq \frac{75}{\sigma^2} \Gamma^2 \|T_{\Gamma} G_{\gamma} - G_{\gamma}\|_1 , \qquad (3.9)
$$

and the proof is complete by Equation (3.4) .

$$
\sqcup
$$

3.3. Approximating the Furstenberg probability and the proof of Theorem [1.2.](#page-1-2) We are now going to patch together the (infinite!) invariant measure of the Y process, translated to the left of Γ, and the (infinite too, of course) invariant measure of the Y process driven by −z instead of z, reflected with respect to the origin and translated to the right of Γ, see Fig. [2,](#page-10-4) to form a probability measure that we call γ_{Γ} . We expect (and will show) that γ_{Γ} is close to the invariant probability ν_{Γ} of the X MC.

FIGURE 2. For $\Gamma = 10$ we have drawn in black (respectively, in grey) the density of the invariant measure of the Y process translated of $-\Gamma$ (respectively, of the Y process driven by $-z$, reflected with respect to the origin and then translated of $+\Gamma$): one can readily see that the densities of these two measure exist if z has a density. Note also that, in particular,the cumulative function associated to the black density function is $F_{\leq}(\cdot + \Gamma)$, Therefore, up to a multiplicative factor, the density of the probability γ_{Γ} is given by the black line on the left of the origin and by the grey one on the right.

Let $F_{\lhd}(x) := c_{\nu}^{-1} \int \mathbf{1}_{(-\infty,x]}(y) \nu(\mathrm{d}y)$ for $x \in \mathbb{R}$, where ν and c_{ν} are as in Theorem [3.1.](#page-8-1) So $F_{\leq}(\cdot)$ is the cumulative function of the (suitably normalized) stationary measure of Y and Theorem [3.1](#page-8-1) yields

$$
F_{\lhd}(x) \stackrel{x \to \infty}{=} x + c_{\lhd} + \begin{cases} O\left(x^{-(\xi - 3)}\right) & \text{under (T-1)},\\ O\left(\exp(-\delta x)\right) & \text{under (T-1')}.\end{cases} (3.10)
$$

Let F_{\triangleright} be the same with z replaced by $-z$, which describes the related object for the right edge instead of the left edge, which then has comparable asymptotic behavior. Using this, we define an approximate stationary distribution by

$$
G_{\gamma_{\Gamma}}(x) := \begin{cases} F_{\triangleright}(\Gamma - x)/C_{\Gamma} & \text{if } x \ge 0, \\ 1 - (F_{\lhd}(x + \Gamma)/C_{\Gamma}) & \text{if } x \le 0, \end{cases} \quad \text{with } C_{\Gamma} := F_{\lhd}(\Gamma) + F_{\rhd}(\Gamma). \tag{3.11}
$$

Note that C_{Γ} has been selected so that $G_{\gamma_{\Gamma}}(0)$ is coherently chosen. We register also that for the cumulative probability $F_{\gamma_{\Gamma}}(\cdot) = 1 - G_{\gamma_{\Gamma}}(\cdot)$ we have

$$
F_{\gamma_{\Gamma}}(x) := \begin{cases} 1 - (F_{\triangleright}(\Gamma - x)/C_{\Gamma}) & \text{if } x \ge 0, \\ F_{\triangleleft}(x + \Gamma)/C_{\Gamma} & \text{if } x \le 0. \end{cases}
$$
(3.12)

Note also that, by the explicit bound on $\nu((-\infty, x])$ for $x \leq 0$ in Prop. [2.4,](#page-7-2) the tail behavior(s) of γ_{Γ} is controlled by the tail behavior(s) of z, in particular we have $\int_{\mathbb{R}} |x| \gamma_{\Gamma}(\mathrm{d}x)$ ∞ , which is required to apply Corollary [3.3.](#page-9-0)

Set

$$
\kappa_2 := \frac{c_{\lhd} + c_{\rhd}}{2} \,. \tag{3.13}
$$

Proposition 3.5. Assume that both z and $-z$ satisfy the assumptions of Theorem [3.1](#page-8-1) (in particular, there exists $\xi > 3$ such that $z \in \mathbb{L}^{\xi}$, so (T-2) becomes superfluous). Then for a suitably chosen $\delta > 0$

$$
C_{\Gamma} \stackrel{\Gamma \to \infty}{=} 2\Gamma + 2\kappa_2 + \begin{cases} O\left(\Gamma^{-(\xi - 3)}\right) & under \textbf{(T-1) for both } \pm z, \\ O\left(\exp(-\delta\Gamma)\right) & under \textbf{(T-1') for both } \pm z. \end{cases} (3.14)
$$

Moreover

$$
L_{\Gamma}[G_{\gamma_{\Gamma}}] = \frac{1}{C_{\Gamma}} \int_{\mathbb{R}} \frac{F_{\triangleright}(y)}{1 + e^{y}} dy + O(\exp(-\Gamma)), \qquad (3.15)
$$

and

$$
\|T_{\Gamma}G_{\gamma_{\Gamma}} - G_{\gamma_{\Gamma}}\|_{1} = \begin{cases} O\left(\Gamma^{-\xi+2}\right) & under \textbf{(T-1) for both } \pm z, \\ O\left(\exp(-\delta\Gamma)\right) & under \textbf{(T-1') for both } \pm z. \end{cases}
$$
(3.16)

The proof of Proposition [3.5](#page-10-2) is in Section [6.](#page-25-1) We have now all the ingredients for the proof of Theorem [1.2.](#page-1-2)

Proof. Theorem [1.2](#page-1-2) follows immediately by combining Corollary [3.3,](#page-9-0) with the choice of $\gamma = \gamma_{\Gamma}$, and Proposition [3.5:](#page-10-2) note that the value of δ in Theorem [1.2](#page-1-2) may be chosen equal to $\delta/2$, with δ the constant appearing in Proposition [3.5.](#page-10-2) In particular by [\(3.15\)](#page-11-2) we obtain

$$
\kappa_1 = \frac{1}{2} \int_{\mathbb{R}} \frac{F_{\rhd}(y)}{1 + e^y} \mathrm{d}y. \tag{3.17}
$$

We note that κ_1 can also be evaluated using F_{\lhd} instead of F_{\rhd} , as shown in the forthcoming Remark [6.1.](#page-25-2) \square

4. THE STATIONARY MEASURE OF THE Y PROCESS

This section is devoted to the proof of Theorem [3.1.](#page-8-1) Let ν be a nonzero invariant measure of Y. By proposition [2.4,](#page-7-2) we have $\nu((-\infty, x]) < \infty$ for every $x \in \mathbb{R}$.

We state a first lemma which rephrases the result of Theorem [3.1.](#page-8-1)

Lemma 4.1. Let $\xi > 3$ and $c_{\nu} > 0$. Then, the existence of $d_{\nu} \in \mathbb{R}$ such that equation [\(3.1\)](#page-8-2) holds is equivalent to the fact that, as $x \to \infty$, uniformly in $x \le y \le 2x$ we have

$$
\nu((x, y]) = c_{\nu}(y - x) + O(x^{-(\xi - 3)}).
$$
\n(4.1)

Similarly, let $c_{\nu} > 0$, then the existence of $d_{\nu} \in \mathbb{R}$ and $\delta > 0$ such that equation [\(3.1\)](#page-8-2) with error term replaced by $O(e^{-\delta x})$ holds, is equivalent to the existence of $\delta > 0$, such that, as $x \to \infty$, uniformly in $x \leq y \leq 2x$:

$$
\nu((x, y]) = c_{\nu}(y - x) + O\left(\exp\left(-\delta x\right)\right). \tag{4.2}
$$

Proof. We prove the first statement. Writing $\nu((x, y)) = \nu((-\infty, y]) - \nu((-\infty, x])$, it is clear that [\(3.1\)](#page-8-2) implies [\(4.1\)](#page-11-3). Conversely, assume that (4.1) holds. For any $x \ge 1$, consider the (unique) integer *n* such that $2^n < x \leq 2^{n+1}$ and write

$$
\nu((-\infty, x]) - c_{\nu}x = \nu((-\infty, 1]) - c_{\nu}
$$

+
$$
\sum_{i=0}^{n-1} \nu((2^i, 2^{i+1}]) - c_{\nu}(2^{i+1} - 2^i)
$$

+
$$
\nu((2^n, x]) - c_{\nu}(x - 2^n),
$$
 (4.3)

and note that, due to [\(4.1\)](#page-11-3), the series $\sum_{i\geq 0} (\nu([2^i, 2^{i+1}]) - c_{\nu}(2^{i+1} - 2^i))$ converges and its partial remainder $\sum_{i\geq n} (\nu([2^i, 2^{i+1}]) - c_{\nu}(2^{i+1} - 2^i)) = O(2^{-n(\xi-3)})$. By [\(4.1\)](#page-11-3), the term $\nu((2^n,x]) - c_\nu(x-2^n)$ is also a $O(2^{-n(\xi-3)})$. Using finally that $O(2^{-n(\xi-3)}) = O(x^{-(\xi-3)})$, we get that [\(3.1\)](#page-8-2) holds with $d_{\nu} := \nu((-\infty, 1]) - c_{\nu} + \sum_{i=0}^{\infty} \nu((2^i, 2^{i+1}]) - c_{\nu}(2^{i+1} - 2^i).$ The proof is similar in the case of error term $O(e^{-\delta x})$. □

Our focus will now be proving (4.1) . There will be four preliminary steps:

- (1) By elaborating on the well-known representation of the invariant measure of a re-current MC (see for example [\[14,](#page-32-12) Sec. 6 of Ch. 3]), we show that $\nu((x, y))$ can be represented by using the excursion occupation times $V_a(b,(x,y))$ of Y, see the equa-tion [\(4.5\)](#page-12-0) below. In Lemma [4.2,](#page-13-0) we give an expression of $V_a(b,(x,y))$ in terms of the strictly descending epochs of the random walk S and a crucial function F , that links $V_a(b,(x,y))$ and the Y process observed at the *ladder epochs* of S (see [\(4.7\)](#page-13-1)).
- (2) By considering the weak ascending ladder epochs of S, we rewrite F in terms of a new process $(J_k(\theta))_{k>0}$, defined in [\(4.18\)](#page-14-0), which is now an increasing Markov chain.
- (3) We determine the asymptotic behavior of F , see Proposition [4.6:](#page-16-0) this is the main technical estimate in the proof.
- (4) We perform some technical estimates that are central in controlling the error arising when replacing F by its asymptotic behavior into the representation formula for the cumulative function of ν .

We recall that we work under the assumptions of Theorem [3.1,](#page-8-1) that is the criticality condition (1.6) , $(H-1)$ of Theorem [1.2,](#page-1-2) $(T-1)$ or $(T-1')$, and $(T-2)$ explicitly stated in Theorem [3.1.](#page-8-1)

4.1. Step 1: representation of the invariant measure. First, we observe that Y can be explicitly solved. By induction we see that, when $Y_0 = \theta$, for $n = 0, 1, 2, \ldots$

$$
Y_n = \log \left(e^{\theta + S_n} + \sum_{i=0}^{n-1} e^{S_n - S_i} \right) = \theta + S_n + \log \left(1 + e^{-\theta} \sum_{i=0}^{n-1} e^{-S_i} \right), \quad (4.4)
$$

where as before $S_n = \sum_{i=1}^n z_i$ and we adopt the convention $\sum_{\emptyset} := 0$.

Since ν is nonzero, let us fix an arbitrary $a > 0$ such that $\nu((-\infty, a]) > 0$ and denote by $\tau_Y(a) := \inf\{n \geq 1 : Y_n \in (-\infty, a]\}\$ the first return time to $(-\infty, a]$ of Y. By applying [\[14,](#page-32-12) Theorem 3.6.5 to the recurrent Markov chain Y , and by applying the Markov property at time 1, we have that for any Borel set $I \subset (a,\infty)$,

$$
\nu(I) = \int_{(-\infty, a]} \mathbb{E}_v \left[\sum_{n=0}^{\tau_Y(a)-1} \mathbf{1}_{\{Y_n \in I\}} \right] \nu(dv)
$$

$$
= \int_{(-\infty, a]} \mathbb{E}_v \left[V_a(Y_1, I) \mathbf{1}_{\{Y_1 > a\}} \right] \nu(dv), \tag{4.5}
$$

where, for $b > a$ and any Borel set $I \subset (a, \infty]$, we set

$$
V_a(b, I) := \mathbb{E}_b \left[\sum_{n=0}^{\tau_Y(a)-1} \mathbf{1}_{\{Y_n \in I\}} \right] = \sum_{n=0}^{\infty} \mathbb{P}_b \left(\tau_Y(a) > n, Y_n \in I \right).
$$
 (4.6)

Later on, we will take I of the form $(x, y]$, with $a < x \le y \le 2x$, but let us remain general for now.

We are now going to give a representation of $V_a(b, I)$ using the fluctuation theory for the one-dimensional random walk S. Consider the strictly descending ladder times $\varrho_0 := 0$ and, for $j \geq 1$

$$
\varrho_j := \min \left\{ n \ge \varrho_{j-1} : S_n < S_{\varrho_{j-1}} \right\} \tag{4.7}
$$

Furthermore, consider the process $\Theta = (\Theta_j)_{j \geq 0}$ defined by $\Theta_j := Y_{\varrho_j}$. Let

$$
\eta_j := \sum_{n=\varrho_{j-1}}^{\varrho_j-1} e^{S_{\varrho_j}-S_n} \quad \text{and} \quad \Delta_j = S_{\varrho_{j-1}}-S_{\varrho_j} > 0. \quad (4.8)
$$

By the strong Markov property of S, $((\eta_k, \Delta_k))_{k\geq 1}$ is an IID sequence adapted to the filtration $(\mathcal{F}_{\varrho_j})_{j\geq 0}$, where $\mathcal{F}_n := \sigma(\{S_i, i \leq n\})$ for $n \geq 0$. Using [\(4.4\)](#page-12-1), we have that

$$
\Theta_{j+1} = \log \left(e^{-\Delta_{j+1}} e^{\Theta_j} + \eta_{j+1} \right),\tag{4.9}
$$

which, in view of the fact that $(\eta_i, \Delta_j)_{i\geq 1}$ is an IID sequence, implies that $(\Theta_j)_{j\geq 0}$ is a Markov chain. For completeness, let us mention that, if $\Theta_0 = Y_0 = b$, then Θ_j is given by:

$$
\Theta_j = Y_{\varrho_j} = \log \left(e^{b + S_{\varrho_j}} + \sum_{k=1}^j e^{S_{\varrho_j} - S_{\varrho_k}} \eta_k \right).
$$
 (4.10)

Lemma 4.2. For $b > a$, for any Borel set $I \subset (a, \infty)$,

$$
V_a(b,I) = \mathbb{E}_b \left[\sum_{j=0}^{\tau_{\Theta}(a)-1} F(\Theta_j, I) \right] = \sum_{j=0}^{\infty} \mathbb{E}_b \left[\mathbf{1}_{\{\tau_{\Theta}(a) > j\}} F(\Theta_j, I) \right],
$$
(4.11)

where $\tau_{\Theta}(a)$ is the first return time to $(-\infty, a]$ of Θ :

$$
\tau_{\Theta}(a) := \inf\{j \ge 1 : \Theta_j \le a\},\tag{4.12}
$$

and $F(\theta, I)$ is defined, for any $\theta \in \mathbb{R}$ and any Borel set $I \subset \mathbb{R}$, by:

$$
F(\theta, I) := \mathbb{E}_{\theta} \left[\sum_{n=0}^{\rho_1 - 1} \mathbf{1}_{\{Y_n \in I\}} \right] = \sum_{n=0}^{\infty} \mathbb{P}_{\theta}(\rho_1 > n, Y_n \in I).
$$
 (4.13)

Proof. We use the definition of Y to see that $Y_n \geq Y_{\varrho_i} + S_n - S_{\rho_i} \geq Y_{\varrho_i} = \Theta_j$ for $\varrho_j \leq n < \varrho_{j+1}$. This means that in fact $\tau_Y(a)$ must be equal to $\varrho_{\tau_{\Theta}(a)}$, with $\tau_{\Theta}(a)$ defined in [\(4.12\)](#page-13-2). Hence

$$
V_{a,b}(I) = \mathbb{E}_b \left[\sum_{j=0}^{\tau_{\Theta}(a)-1} \sum_{n=\varrho_j}^{\varrho_{j+1}-1} \mathbf{1}_{\{Y_n \in I\}} \right]
$$

=
$$
\sum_{j=0}^{\infty} \mathbb{E}_b \left[\mathbf{1}_{\{\tau_{\Theta}(a) > j\}} \sum_{n=\varrho_j}^{\varrho_{j+1}-1} \mathbf{1}_{\{Y_n \in I\}} \right].
$$
 (4.14)

Observe that the event $\{\tau_{\Theta}(a) > j\}$ is measurable with respect to \mathcal{F}_{ϱ_j} , and that, due to the Markov property of Y, the conditional expectation of $\sum_{n=\varrho_j}^{\varrho_{j+1}-1} \mathbf{1}_{\{Y_n \in I\}}$ with respect to \mathcal{F}_{ϱ_j} , is equal to $F(Y_{\varrho_j}, I)$. We obtain [\(4.11\)](#page-13-3) and the proof of Lemma [4.2](#page-13-0) is complete. \Box 4.2. Step 2: representation of the function F in terms of an auxiliary process. Recall [\(4.13\)](#page-13-4) for the definition of $F(\theta, I)$. We are going to use again the fluctuation theory for the random walk S to give a representation of F in terms of an *increasing* process. Let us introduce some notations.

Let $H_k := S_{\alpha_k}$ and $\alpha_k, k \geq 0$, be the weak ascending ladder heights and epochs of S: $\alpha_0 := 0$ and for any $k \geq 1$

$$
\alpha_k := \inf \left\{ n > \alpha_{k-1} : S_n \ge S_{\alpha_{k-1}} \right\} . \tag{4.15}
$$

Let us also introduce for $k \geq 1$:

$$
\widetilde{\Delta}_k := H_k - H_{k-1} \ge 0, \qquad \widetilde{\eta}_k := \sum_{n = \alpha_{k-1} + 1}^{\alpha_k} e^{S_n - H_k}.
$$
\n(4.16)

By the Markov property of S, the sequence $((\widetilde{\Delta}_k, \widetilde{\eta}_k))_{k>1}$ is IID. Let us define a new Markov chain $(J_k(\theta))_k$, adapted to the filtration $(\mathcal{F}_{\alpha_k})_{k\geq 0}$, by $J_0(\theta) := \theta$, and for all $k = 0, 1, 2, \ldots$

$$
J_{k+1}(\theta) := J_k(\theta) + \widetilde{\Delta}_{k+1} + \log(1 + \widetilde{\eta}_{k+1}e^{-J_k(\theta)}).
$$
\n(4.17)

Note that $J_k(\theta)$ is increasing in k. Explicitly, $J_k(\theta)$ is given by:

$$
J_k(\theta) = \log \left(e^{\theta + H_k} + \sum_{j=1}^k \widetilde{\eta}_j \, e^{H_k - H_{j-1}} \right) = \theta + H_k + \log \left(1 + e^{-\theta} \sum_{j=1}^k \widetilde{\eta}_j \, e^{-H_{j-1}} \right). \tag{4.18}
$$

Lemma 4.3. For any $\theta \in \mathbb{R}$, any Borel set $I \subset \mathbb{R}$, we have

$$
F(\theta, I) = \sum_{k=0}^{\infty} \mathbb{P}(J_k(\theta) \in I).
$$
 (4.19)

Proof. The proof is based on the duality lemma for the random walk S . By the definitions of $F(\theta, I)$ and ρ_1 , we have

$$
F(\theta, I) = \sum_{n=0}^{\infty} \mathbb{P}\left(S_1 \ge 0, ..., S_n \ge 0, \log\left(e^{\theta + S_n} + \sum_{j=0}^{n-1} e^{S_n - S_j}\right) \in I\right)
$$

=
$$
\sum_{n=0}^{\infty} \mathbb{P}\left(S_1 \ge 0, ..., S_n \ge 0, \log\left(e^{\theta + S_n} + \sum_{j=1}^{n} e^{S_n - S_{n-j}}\right) \in I\right).
$$

Since $(S_n - S_{n-j})_{0 \leq j \leq n}$ has the same distribution as $(S_j)_{0 \leq j \leq n}$, we obtain that

$$
F(\theta, I) = \sum_{n=0}^{\infty} \mathbb{P}\left(S_n \ge S_{n-1}, ..., S_n \ge S_1, S_n \ge 0, \log\left(e^{\theta + S_n} + \sum_{j=1}^n e^{S_j}\right) \in I\right)
$$

=
$$
\sum_{k=0}^{\infty} \mathbb{P}\left(\log\left(e^{\theta + H_k} + \sum_{j=1}^{\alpha_k} e^{S_j}\right) \in I\right)
$$

=
$$
\sum_{k=0}^{\infty} \mathbb{P}\left(\theta + H_k + \log\left(1 + e^{-\theta} \sum_{j=1}^{\alpha_k} e^{S_j - H_k}\right) \in I\right).
$$

Note that

$$
\sum_{j=1}^{\alpha_k} e^{S_j - H_k} = \sum_{\ell=1}^k \sum_{j=\alpha_{\ell-1}+1}^{\alpha_{\ell}} e^{S_j - H_{\ell}} e^{H_{\ell} - H_k} = \sum_{\ell=1}^k \widetilde{\eta}_{\ell} e^{H_{\ell} - H_k}, \tag{4.20}
$$

by definition of $\widetilde{\eta}$ in [\(4.16\)](#page-14-1). The strong Markov property of S yields that $(\widetilde{\eta}_{\ell}, H_{\ell} - H_{\ell-1})_{\ell \geq 1}$
see ID. This vial a that $(H_{\ell}, \sum_{\ell}^{k} \widetilde{\chi}_{\ell} - H_{\ell-1})_{\ell}$ is distributed as $(H_{\ell}, \sum_{\ell}^{k} \widetilde{\chi}_{$ are IID. This yields that $(H_k, \sum_{\ell=1}^k \widetilde{\eta}_{\ell} e^{H_{\ell} - H_k})$ is distributed as $(H_k, \sum_{j=1}^k \widetilde{\eta}_{j} e^{-H_{j-1}})$ and therefore that $\theta + H_k + \log(1 + e^{-\theta} \sum_{j=1}^{\alpha_k} e^{S_j - H_k})$ is distributed as $J_k(\theta)$. This concludes the proof. \Box

4.3. Step 3: the asymptotic behavior of the function F. We now take $I = (x, y)$, with $0 \le x \le y \le 2x$. We are going to determine the asymptotic behavior of $F(\theta, (x, y))$ as $x \to \infty$ by using the fact that $J_k(\theta)$ behaves asymptotically like a random walk with increments having the law of H_1 , which is a nonnegative random variable. The renewal function of H is defined as

$$
R(x) := \sum_{k=0}^{\infty} \mathbb{P}\left(H_k \le x\right), \qquad x \ge 0,
$$
\n(4.21)

and $R(x) := 0$ for $x < 0$.

The following Lemma, due to Rogozin [\[29\]](#page-33-9) and Stone [\[31\]](#page-33-10), plays a crucial role in the study of F.

Lemma 4.4 ([\[29\]](#page-33-9), formula (28), and [\[31\]](#page-33-10)). Assume that

(R-1): there exists $n_0 \geq 1$ such that the law of H_{n_0} has an absolutely continuous part: this property is normally referred to as "the law of H_1 is spread out"; and either

(R-2): there exists $\kappa \geq 2$ such that $\mathbb{E}[H_1^{\kappa}] < \infty$; or

(**R-2'**): there exists $c > 0$ such that $\mathbb{E}[e^{cH_1}] < \infty$.

Then for any $x > 0$,

$$
\left| R(x) - (c_R x + c'_R) \right| \le \varphi(x) := \begin{cases} C_{\varphi} (1+x)^{-(\kappa-2)}, & \text{under } (\mathbf{R-2}), \\ C_{\varphi} e^{-C'_{\varphi} x}, & \text{under } (\mathbf{R-2'}) \end{cases}
$$
(4.22)

where $C_{\varphi}, C'_{\varphi}$ are two (unimportant) positive constants and

$$
c_R := \frac{1}{\mathbb{E}(H_1)}, \qquad c'_R := \frac{\mathbb{E}\left[H_1^2\right]}{2\mathbb{E}\left[H_1\right]^2}.
$$
\n(4.23)

Remark 4.5. (i) By the main theorem in Doney [\[13\]](#page-32-13),

• (R-2) is satisfied if $\mathbb{E}[(z^+)^{\kappa+1}] < \infty$, hence it is satisfied under hypothesis (T-1) of Theorem [3.1](#page-8-1) with $\kappa = \xi - 1 \geq 2$;

 \bullet (R-2') is satisfied under (T-1').

(ii) The case $(R-2')$ of Lemma [4.4](#page-15-0) is stated in Stone [\[31\]](#page-33-10) with an implicit constant that must be equal to c'_R . See also [\[1,](#page-32-14) Sec. VII.2 and Ex. VII.2.2].

The case (R-2) is also obtained in Carlsson [\[5,](#page-32-15) Corollary 2 and Remark 2] under the slightly weaker assumption that H_1 is strongly nonlattice, instead of $(R-1)$. Here, we follow the formalism of Rogozin [\[29\]](#page-33-9), which states that under $(R-2)$,

$$
R(x) \stackrel{x \to \infty}{=} c_R x + c'_R - \frac{1}{2(\mathbb{E}(H_1))^2} \mathbb{E}\left[(H_1 - x)^2 \mathbf{1}_{\{H_1 \ge x\}} \right] + o(x^{-\kappa+1}). \tag{4.24}
$$

Notice that the condition $(R-1)$ is equivalent to [\[29,](#page-33-9) Condition (6)], by Remark 4 there. Using (**R-2**) we see that $\mathbb{E}[H_1^2 \mathbf{1}_{\{H_1 \ge x\}}] \le \mathbb{E}(H_1^{\kappa})x^{2-\kappa}$, hence there exists $C_R > 0$ such that

$$
|R(x) - (c_R x + c'_R)| \le C_R x^{2-\kappa}
$$
 and $R(x) \le C_R x$ for $x \ge 1$. (4.25)

Of course [\(4.25\)](#page-15-1) is also valid in the case $(\mathbb{R}-2')$, and C_R is just a constant involved in upper bounds on error terms, while c_R and c'_R are the constants in [\(4.23\)](#page-15-2).

The following Proposition gives the key estimate in the proof of Theorem [3.1.](#page-8-1) Recall (4.13) for the definition of F.

Proposition 4.6. Assume (R-1) and (R-2) with $\kappa \geq 2$. Then, for $x \to \infty$ and uniformly in $0 \le \theta \le x/2$ and $x \le y \le 2x$, we have

$$
F(\theta, (x, y]) = c_R (y - x) + O(x^{-(\kappa - 2)}).
$$
\n(4.26)

Assuming $(R-2')$ instead of $(R-2)$, there exists a positive constant δ such that [\(4.26\)](#page-16-1) holds with error term replaced by $O(e^{-\delta x})$.

Before presenting the proof of Proposition [4.6,](#page-16-0) we state two lemmas. The first lemma will allow us to jump from initial point θ to some larger (random) initial point. From this larger initial point, it will become possible to compare F with R : upper and lower bounds are given in the second lemma, which is useful really for large θ .

Lemma 4.7. For every $\theta \in \mathbb{R}$, any Borel set $I \subset \mathbb{R}$ and any $m \geq 1$, we have

$$
F(\theta, I) = F_m(\theta, I) + \mathbb{E}\left[F(J_m(\theta), I)\right]
$$
\n(4.27)

where

$$
F_m(\theta, I) := \sum_{k=0}^{m-1} \mathbb{P}(J_k(\theta) \in I).
$$
 (4.28)

Proof. This is just an application of the Markov property at time m for the Markov chain $(J_k(\theta))_{k\geq 0}$.

Lemma 4.8. For every x and y such that $0 \le x \le y \le 2x$, for every $\theta \ge 0$ and $t > 0$, we have

$$
F(\theta, (x, y]) \le R(y - \theta) - R(x - \theta - t) + \Lambda(2x, e^{\theta}t)
$$

\n
$$
F(\theta, (x, y]) \ge R(y - \theta - t) - R(x - \theta) - \Lambda(2x, e^{\theta}t),
$$
\n(4.29)

where

$$
\Lambda(x,t) := \sum_{k=1}^{\infty} \mathbb{P}\left(H_k \le x, \sum_{j=1}^{\infty} \widetilde{\eta}_j e^{-H_{j-1}} > t\right).
$$
\n(4.30)

Proof. We use [\(4.19\)](#page-14-2) for an expression of $F(\theta, (x, y])$. By [\(4.18\)](#page-14-0), we have

$$
H_k + \theta \le J_k(\theta) \le H_k + \theta + e^{-\theta} \sum_{j=1}^{\infty} \widetilde{\eta}_j e^{-H_{j-1}}.
$$
\n(4.31)

Hence for $z \in \{x, y\},\$

$$
\mathbb{P}(J_k(\theta) \le z) \le \mathbb{P}(H_k \le z - \theta)
$$
\n(4.32)

and

$$
\mathbb{P}\left(J_k(\theta) \leq z\right) \geq \mathbb{P}\left(H_k + \theta + e^{-\theta} \sum_{j=1}^{\infty} \widetilde{\eta}_j e^{-H_{j-1}} \leq z\right)
$$
\n
$$
\geq \mathbb{P}\left(H_k + \theta + t \leq z, \sum_{j=1}^{\infty} \widetilde{\eta}_j e^{-H_{j-1}} \leq e^{\theta}t\right) \qquad (4.33)
$$
\n
$$
\geq \mathbb{P}\left(H_k + \theta + t \leq z\right) - \mathbb{P}\left(H_k \leq 2x, \sum_{j=1}^{\infty} \widetilde{\eta}_j e^{-H_{j-1}} > e^{\theta}t\right),
$$

using $z - \theta - t \leq 2x$ in the last line. Applying these two inequalities with $z = y$ then $z = x$, and summing over k, we get the lower and the upper bounds in the lemma. \Box

Proof of Proposition [4.6.](#page-16-0) The asymptotic behavior of F will follow from Lemmas [4.7](#page-16-2) and [4.8](#page-16-3) once we have obtained the estimates on $F_m(\theta,(x,y])$, $\mathbb{E}[R(y-J_m(\theta))-R(x-J_m(\theta))$ t)], $\mathbb{E}[R(y - J_m(\theta) - t) - R(x - J_m(\theta))]$ and $\mathbb{E}[\Lambda(2x, e^{J_m(\theta)}t)]$. This will be done separately in what follows.

For ease of exposition we introduce the symbol $O_u(f(s))$ to denote a quantity whose absolute value is bounded by $C|f(s)|$ with $C > 0$ independent of the parameter s. In general s may be of dimension larger than one, and belong to a domain that will be specified.

Estimate of $F_m(\theta,(x,y])$. Since $F_m(\theta,(x,y]) \leq \sum_{k=0}^{m-1} \mathbb{P}(J_k(\theta) > x) \leq m \mathbb{P}(J_m(\theta) > x)$, our goal is to estimate $\mathbb{P}(J_m(\theta) > x)$. We observe that for all $x \geq 4$, $\theta \in [0, x/2]$ and $m \geq 1$, if $H_m \leq \frac{x}{4}$ $\frac{x}{4}$ and $\sum_{j=1}^{\infty} \tilde{\eta}_j e^{-H_{j-1}} \leq e^{x/2}$, then

$$
J_m(\theta) \le H_m + \log \left(e^{\theta} + \sum_{j=1}^{\infty} \widetilde{\eta}_j e^{-H_{j-1}} \right) \le \frac{x}{4} + \log \left(2e^{\frac{x}{2}} \right) \le x.
$$

Hence, for all for all $x \ge 4$, $\theta \in [0, x/2]$ and $m \ge 1$,

$$
\mathbb{P}(J_m(\theta) > x) \le \mathbb{P}\left(H_m > \frac{x}{4}\right) + \mathbb{P}\left(\sum_{j=1}^{\infty} \widetilde{\eta}_j \, e^{-H_{j-1}} > e^{x/2}\right) \tag{4.34}
$$

We first estimate the second term in [\(4.34\)](#page-17-0), as the estimation of first term will depend on whether the case $(R-2)$ or $(R-2')$ applies.

Remark that $\widetilde{\eta}_1 = 1 + \sum_{i=1}^{\alpha_1-1} e^{S_i - H_1} \leq 1 + \sum_{i=1}^{\infty} e^{S_i} \mathbf{1}_{\{\max_{0 \leq j \leq i} S_j < 0\}}$ by using the definition of α_1 and the fact that $H_1 \geq 0$. By [\[8,](#page-32-10) Lemma A.2], $\mathbb{E}[\mathbf{z}^2] < \infty$ (i.e., hypothesis (T-2)) implies that for some $\delta_1 > 0$,

$$
\mathbb{E}[\exp(\delta_1 \widetilde{\eta}_1)] < \infty. \tag{4.35}
$$

Recall that $(\tilde{\eta}_{\ell}, H_{\ell} - H_{\ell-1})_{\ell \geq 1}$ is an IID sequence. By [\(4.35\)](#page-17-1), we can apply [\[20,](#page-33-11) Theorem 2.1] to obtain some $\delta_0 > 0$ such that 2.1 to obtain some $\delta_2 > 0$ such that

$$
C_{\delta_2} := \mathbb{E} \, e^{\delta_2 \sum_{j=1}^{\infty} \widetilde{\eta}_j \, e^{-H_{j-1}}} < \infty. \tag{4.36}
$$

We deduce from [\(4.36\)](#page-17-2) and the Markov inequality that for every $\lambda > 0$

$$
\mathbb{P}\left(\sum_{j=1}^{\infty} \widetilde{\eta}_j e^{-H_{j-1}} > \lambda\right) \le C_{\delta_2} e^{-\delta_2 \lambda}.
$$
\n(4.37)

In particular,

$$
\mathbb{P}\left(\sum_{j=1}^{\infty} \tilde{\eta}_j \, e^{-H_{j-1}} > e^{x/2}\right) \le C_{\delta_2} e^{-\delta_2 \, e^{x/2}}.\tag{4.38}
$$

It remains to estimate $\mathbb{P}\left(H_m > \frac{x}{4}\right)$ $\frac{x}{4}$) according to whether the hypothesis (**R-2**) or (**R-2**[']) applies. We start with the case $(R-2)$. As discussed right after Lemma [4.4,](#page-15-0) hypothesis (T-1) of Theorem [3.1](#page-8-1) yields that the $\kappa = \xi - 1$ moment of H_1 is finite, hence, using Rosenthal inequality, there exists a constant $C_{\kappa} > 0$ such that

$$
\mathbb{E}\left[H_n^{\kappa}\right] \le C_{\kappa} n^{\kappa}, \qquad \text{for } n = 1, 2, \dots. \tag{4.39}
$$

Consequently, using the Markov inequality, for any $x > 0$ and $m \ge 1$

$$
\mathbb{P}\left(H_m > \frac{x}{4}\right) \le 4^{\kappa} C_{\kappa} m^{\kappa} x^{-\kappa}, \qquad \text{in the case (\mathbf{R-2}).} \tag{4.40}
$$

In the case $(\mathbf{R-2}')$, $\mathbb{E}[e^{cH_m}] = e^{c_H m}$ with $c_H := \log \mathbb{E}[e^{cH_1}] \in (0,\infty)$. The Markov inequality yields that for any $x \geq 0$ and $m \geq 1$,

$$
\mathbb{P}\left(H_m > \frac{x}{4}\right) \le e^{c_H m - cx/4}, \qquad \text{in the case } (\mathbf{R-2}'). \tag{4.41}
$$

Applying [\(4.38\)](#page-18-0), [\(4.40\)](#page-18-1) and [\(4.41\)](#page-18-2) to [\(4.34\)](#page-17-0), we get that for all $x \ge 4$, $\theta \in [0, x/2]$ and $m \geq 1$,

$$
\mathbb{P}(J_m(\theta) > x) = \begin{cases} O_u(m^{\kappa} x^{-\kappa}), & \text{in the case (\mathbf{R-2}),} \\ O_u(e^{c_H m - cx/4}), & \text{in the case (\mathbf{R-2'})}, \end{cases}
$$
(4.42)

so

$$
F_m(\theta, x) = \begin{cases} O_u(m^{\kappa+1}x^{-\kappa}), & \text{in the case (\mathbf{R-2}),} \\ O_u(m e^{c_H m - cx/4}), & \text{in the case (\mathbf{R-2').} \end{cases}
$$
(4.43)

Estimate of $\mathbb{E}[R(y - J_m(\theta)) - R(x - J_m(\theta) - t)].$ We note that, by [\(4.25\)](#page-15-1), $R(x) = O_u(x)$ for $x \geq 1$, and we observe that R is nondecreasing.

Let $x \geq 2, y \in [x, 2x], m \geq 1$ and $t \in (0, 1)$. Whenever $J_m(\theta) \leq x/2$, we have, as a consequence of Lemma [4.4:](#page-15-0)

$$
R(y - J_m(\theta)) - R(x - J_m(\theta) - t) = c_R(y - x + t) + O_u\left(\varphi\left(\frac{x}{2} - 1\right)\right),\tag{4.44}
$$

with function φ defined in [\(4.22\)](#page-15-3). Hence, for $x \geq 2, y \in [x, 2x]$, $m \geq 1$ and $t \in (0, 1)$

$$
\mathbb{E}\left[R(y-J_m(\theta)) - R(x-J_m(\theta) - t)\right]
$$
\n
$$
= \mathbb{E}\left[\left(c_R(y-x+t) + O_u\left(\varphi\left(\frac{x}{2}-1\right)\right)\right) \mathbf{1}_{\{J_m(\theta) \le \frac{x}{2}\}}\right] + \mathbb{E}\left[O_u(x) \mathbf{1}_{\{J_m(\theta) > \frac{x}{2}\}}\right]
$$
\n
$$
= c_R(y-x+t) + O_u\left(\varphi\left(\frac{x}{2}-1\right)\right) + O_u\left(x \mathbb{P}\left(J_m(\theta) > \frac{x}{2}\right)\right).
$$
\n(4.45)

Using [\(4.42\)](#page-18-3), we conclude that for all for $x \ge 8, y \in [x, 2x], \theta \in [0, x/2], m \ge 1$ and $t \in (0, 1)$

$$
\mathbb{E}[R(y - J_m(\theta)) - R(x - J_m(\theta) - t)] =
$$

\n
$$
c_R(y - x + t) + \begin{cases} O_u(x^{-(\kappa - 2)} + m^{\kappa} x^{-(\kappa - 1)}) , & \text{in the case (\mathbf{R-2}),} \\ O_u(e^{c_H m - \delta_3 x}), & \text{in the case (\mathbf{R-2'})}, \end{cases}
$$
(4.46)

where δ_3 is an arbitrarily fixed constant, satisfying $0 < \delta_3 < \min(c/8, C'_{\varphi})$.

Estimate of $\mathbb{E}[R(y - J_m(\theta) - t) - R(x - J_m(\theta))]$. Exactly in the same manner, we obtain that for all $x \ge 8, y \in [x, 2x], \theta \in [0, x/2], m \ge 1$ and $t \in (0, 1)$

$$
\mathbb{E}[R(y - J_m(\theta) - t) - R(x - J_m(\theta))] =
$$

$$
c_R(y - x - t) + \begin{cases} O_u(x^{-(\kappa - 2)} + m^{\kappa} x^{-(\kappa - 1)}), & \text{in the case (\mathbf{R-2}),} \\ O_u(e^{c_H m - \delta_3 x}), & \text{in the case (\mathbf{R-2'})}. \end{cases}
$$
(4.47)

Estimate of $\mathbb{E}[\Lambda(2x, e^{J_m(\theta)}t)]$. Recall [\(4.30\)](#page-16-4) for the definition of $\Lambda(\cdot, \cdot)$. For any $n \ge 1$ and $s > 0$, we have:

$$
\Lambda(x,s) \le \sum_{k=n}^{\infty} \mathbb{P}[H_k \le x] + n \mathbb{P}\left(\sum_{j=1}^{\infty} \widetilde{\eta}_j e^{-H_{j-1}} > s\right).
$$
 (4.48)

The second term in [\(4.48\)](#page-19-0) is controlled using [\(4.37\)](#page-17-3): it is bounded by $C_{\delta_2} n e^{-\delta_2 s}$. For the first term in [\(4.48\)](#page-19-0) observe that it is equal to

$$
\mathbb{E}[R(x - H_n)] \le R(x)\,\mathbb{P}(H_n \le x) = O_u(x\,\mathbb{P}(H_n \le x)),\tag{4.49}
$$

assuming that $x \geq 1$. We complete this estimate by observing that, since $\mathbb{E}[H_1] > 0$, by the classical Cramér large deviation theorem there exist $C' > 0$ and some small $\delta_4 > 0$ such that

$$
\mathbb{P}\left(H_k \le \delta_4 k\right) \le C' e^{-\delta_4 k} \qquad \text{for } k = 1, 2, \dots \tag{4.50}
$$

Consequently, choosing $n := [x/\delta_4]$ in [\(4.48\)](#page-19-0), we obtain, for $x \ge 1$ and $s > 0$,

$$
\Lambda(x,s) = O_{u}(x(e^{-x} + e^{-\delta_{2}s})).
$$
\n(4.51)

Furthermore, for $\theta \ge 0$, by using the fact that $J_m(\theta) \ge H_m$ and by using [\(4.50\)](#page-19-1) again, we have

$$
\mathbb{E}\left[e^{-\delta_2 t e^{J_m(\theta)}}\right] \le \mathbb{E}\left[e^{-\delta_2 t e^{H_m}}\right] \le e^{-\delta_2 t e^{\delta_4 m}} + C' e^{-\delta_4 m},\tag{4.52}
$$

which, combined with [\(4.51\)](#page-19-2), shows that for all $x \geq 1$, $y \in [x, 2x]$, $\theta \geq 0$, $m \geq 1$ and $t \in (0, 1),$

$$
\mathbb{E}\left[\Lambda\big(2x, e^{J_m(\theta)}t\big)\right] = O_{\mathbf{u}}\left(x\left(e^{-2x} + e^{-\delta_2 t e^{\delta_4 m}} + e^{-\delta_4 m}\right)\right). \tag{4.53}
$$

Completion of the proof of Proposition [4.6.](#page-16-0) Now we are ready to give the proof of (4.26) . We make our choice for m and t. In the case (R-2), we take $m := |x^{\delta}|$ with some $\delta \in (0,1/\kappa)$. In the case $(\mathbf{R-2'})$, we take $m := \lfloor \delta_3 x/(2c_H) \rfloor$, where we recall the constants c_H , δ_3 appeared in [\(4.46\)](#page-18-4). In both cases, we set $t := \exp(-\delta_4 m/2)$.

By Lemmas [4.7](#page-16-2) and [4.8,](#page-16-3) together with estimates (4.43) , (4.46) , (4.47) and (4.53) , for all $x \ge 8, y \in [x, 2x], \theta \in [0, x/2],$

$$
F(\theta, (x, y]) = c_R(y - x) + \begin{cases} O_u\left(x^{-(\kappa - 2)}\right), & \text{in the case (\mathbf{R-2}),} \\ O_u\left(e^{-\delta_5 x}\right), & \text{in the case (\mathbf{R-2'})}, \end{cases}
$$
(4.54)

with $\delta_5 := \min(\frac{\delta_3}{2}, \frac{\delta_3 \delta_4}{4c_H})$ $\frac{\delta_3 \delta_4}{4c_H}$, 1), and the proof of Proposition [4.6](#page-16-0) is complete.

□

4.4. Step 4: technical estimates for the control of the cumulative function of $ν$. Recall that we have chosen $a > 0$ in order to ensure that $ν((−∞, a]) > 0$. Recall [\(4.5\)](#page-12-0), i.e., that for any $a < x \le y \le 2x$,

$$
\nu((x,y]) = \int_{(-\infty,a]} \mathbb{E}_v[V_a(Y_1,(x,y])\mathbf{1}_{\{Y_1>a\}}]\nu(\mathrm{d}v). \tag{4.55}
$$

In the representation of $V_a(b,(x, y))$ given in Lemma [4.2,](#page-13-0) we are going to replace F by its asymptotic expression in the right-hand side of [\(4.26\)](#page-16-1). To this end, we need to control the error terms, this is the goal of the next lemma. Recall the definition of process Θ just after [\(4.7\)](#page-13-1) and the notation τ_{Θ} introduced in Lemma [4.2.](#page-13-0)

Lemma 4.9. If a is large enough, there exists $C > 0$ such that for every $b > a$:

$$
\mathbb{E}_b\left[\tau_{\Theta}(a)\right] \le C b,\tag{4.56}
$$

and, for every $x \geq 0$,

$$
\mathbb{E}_b \left[\sum_{j=0}^{\tau_{\Theta}(a)-1} \mathbf{1}_{\{\Theta_j \ge x\}} \right] \le C e^{b-x}.
$$
 (4.57)

Proof. Recall [\(4.9\)](#page-13-5). If $\Theta_j > a$, then

$$
\Theta_{j+1} - \Theta_j = \log(e^{-\Delta_{j+1}} + \eta_{j+1}e^{-\Theta_j})
$$

$$
\leq \log(e^{-\Delta_{j+1}} + \eta_{j+1}e^{-a}) =: W_{j+1} - W_j.
$$
 (4.58)

In other words, if we consider a real-valued random walk (W_j) with step distribution given as above and starting from $W_0 := b > a$, then for any $j \leq \tau_{\Theta}(a)$, $\Theta_j \leq W_j$, hence

$$
\mathbb{E}_b [\tau_{\Theta}(a)] \leq \mathbb{E}_b [\tau_W(a)] ,
$$

where $\tau_W(a) := \inf\{n \geq 1 : W_n \leq a\}$. Similar to [\(4.35\)](#page-17-1), η_1 also has some exponential moments; in particular its expectation is finite. Note that for a large enough

$$
\mathbb{E}_0[e^{W_1}] = \mathbb{E}[e^{-\Delta_1} + \eta_1 e^{-a}] = \mathbb{E}[e^{-\Delta_1}] + e^{-a} \mathbb{E}[\eta_1] < 1,
$$

which yields in particular that $\mathbb{E}_0[W_1] < 0$. Then it is well known, see [\[21,](#page-33-12) Theorem 3.6.1], that

$$
\mathbb{E}_b\left[\tau_W(a)\right] = O_u(1+b-a),\tag{4.59}
$$

which gives [\(4.56\)](#page-20-0). To show [\(4.57\)](#page-20-1), we use again the comparison $\Theta_j \leq W_j$ for all $j \leq \tau_{\Theta}(a)$:

$$
\mathbb{E}_b\left[\sum_{j=0}^{\tau_{\Theta}(a)-1} \mathbf{1}_{\{\Theta_j \geq x\}}\right] = \sum_{j=0}^{\infty} \mathbb{P}_b(j < \tau_{\Theta}(a), \Theta_j \geq x) \leq \sum_{j=0}^{\infty} \mathbb{P}_b(W_j \geq x).
$$

By the Markov bound, $\mathbb{P}_b(W_j \geq x) = \mathbb{P}_0(W_j \geq x - b) \leq e^{b-x} (\mathbb{E}_0[e^{W_1}])^j$. Taking the sum over j yields [\(4.57\)](#page-20-1) and the proof of Lemma [4.9](#page-20-2) is complete. \Box

We are now ready for:

Proof of Theorem [3.1.](#page-8-1) We have already remarked that $(T-1)$ of Theorem [3.1](#page-8-1) yields $(R-2)$ of Lemma [4.4](#page-15-0) by [\[13\]](#page-32-13), with $\kappa = \xi - 1$. Moreover the condition (H-1) of Theorem [1.2](#page-1-2) – one of the hypotheses in Theorem [3.1](#page-8-1) – implies $(R-1)$ by the following argument. By (H-1), S_{n_0} has a density with respect to the Lebesgue measure, hence S_k , $k \geq n_0$ has a density too. Recall the definitions [\(4.15\)](#page-14-3) and [\(4.21\)](#page-15-4): if H_{n_0} is singular with respect to the Lebesgue measure there exists a Borel set $B \subset [0,\infty)$ of zero Lebesgue measure

and full measure under the law of H_{n_0} , that is $1 = \mathbb{P}(H_{n_0} \in [0, \infty)) = \mathbb{P}(H_{n_0} \in B)$. But $\mathbb{P}(H_{n_0}\in B)=\sum_{k\geq n_0}\mathbb{P}(S_k\in B, \alpha_{n_0}=k)\leq \sum_{k\geq n_0}\mathbb{P}(S_k\in B)=0.$ We have reached a contradiction, hence H_{n_0} is not singular. Hence $(R-1)$ holds and we can use the conclusions of Lemma [4.4.](#page-15-0)

Assume that a is large enough so that Lemma [4.9](#page-20-2) holds. By [\(4.5\)](#page-12-0), for every $a < x \leq$ $y \leq 2x$

$$
\nu((x,y]) = \int_{(-\infty,a]} \mathbb{E}_v \left[V_a(Y_1,(x,y]) \mathbf{1}_{\{Y_1 > a\}} \right] \nu(\mathrm{d}v) \tag{4.60}
$$

We begin with some estimates on $V_a(b,(x,y))$. Recall [\(4.11\)](#page-13-3) for its definition. The first estimate will be useful for large b. Using the fact that $J_k(\theta) \ge H_k + \theta$, we see that for any $x \geq 1, y \in [x, 2x], \theta \geq 0,$

$$
0 \le F(\theta, (x, y]) \le R(y - \theta) \le R(2x) = O_{\mathbf{u}}(x),\tag{4.61}
$$

and thus, using [\(4.56\)](#page-20-0),

$$
V_a(b,(x,y]) = O_u(x \mathbb{E}_b[\tau_\Theta(a)]) = O_u(xb). \qquad (4.62)
$$

Now let us focus on b small, using Proposition [4.6.](#page-16-0) In fact we claim that uniformly in $a < b \leq x/4, y \in [x, 2x]$

$$
V_a(b,(x,y]) = c_R(y-x) \mathbb{E}_b [\tau_{\Theta}(a)] + O_{\mathbf{u}} \left(b \, x^{-(\kappa - 2)} \right). \tag{4.63}
$$

To show [\(4.63\)](#page-21-0) we use, whenever $\Theta_i \leq x/2$, the asymptotic development of $F(\Theta_i, (x, y))$ given by (4.26) . We have:

$$
V_a(b,(x,y)) =
$$

$$
\mathbb{E}_b \left[\sum_{j=0}^{\tau_{\Theta}(a)-1} \left(c_R(y-x) + O_u\left(x^{-(\kappa-2)}\right) \right) \mathbf{1}_{\{\Theta_j \le \frac{x}{2}\}} \right] + \mathbb{E}_b \left[\sum_{j=0}^{\tau_{\Theta}(a)-1} F(\Theta_j,(x,y]) \mathbf{1}_{\{\Theta_j > \frac{x}{2}\}} \right]
$$

$$
= \mathbb{E}_b \left[\sum_{j=0}^{\tau_{\Theta}(a)-1} \left(c_R(y-x) + O_{\mathbf{u}} \left(x^{-(\kappa-2)} \right) \right) \right] + \mathbb{E}_b \left[\sum_{j=0}^{\tau_{\Theta}(a)-1} O_{\mathbf{u}}(x) \mathbf{1}_{\{\Theta_j > \frac{x}{2}\}} \right], \quad (4.64)
$$

where we used (4.61) for the second equality. Using (4.56) , the first term is equal to $c_R(y-x) \mathbb{E}_b \left[\tau_{\Theta}(a) \right] + O_u \left(b x^{-(\kappa-2)} \right)$, while [\(4.57\)](#page-20-1) yields that the second term is $O_u(xe^{b-x/2})$, hence negligible (since $b \leq x/4$). Thus, [\(4.63\)](#page-21-0) is established.

Going back to the estimation of [\(4.60\)](#page-21-2), we use [\(4.63\)](#page-21-0) whenever $a < Y_1 \leq x/4$ and then [\(4.62\)](#page-21-3) to obtain that for $x \ge 1$, uniformly in $y \in [x, 2x]$ and in $v \in (-\infty, a]$

$$
\mathbb{E}_{v}[V_{a}(Y_{1},(x,y])\mathbf{1}_{\{Y_{1}>a\}}] = \n\mathbb{E}_{v}\left[\left(c_{R}(y-x)\mathbb{E}_{Y_{1}}[\tau_{\Theta}(a)]+O_{u}\left(Y_{1}x^{-(\kappa-2)}\right)\right)\mathbf{1}_{\{ax/4\}}] \n= \mathbb{E}_{v}\left[\left(c_{R}(y-x)\mathbb{E}_{Y_{1}}[\tau_{\Theta}(a)]+O_{u}\left(Y_{1}x^{-(\kappa-2)}\right)\right)\mathbf{1}_{\{Y_{1}>a\}}\right] + \mathbb{E}_{v}[O_{u}(Y_{1}x)\mathbf{1}_{\{Y_{1}>x/4\}}] \n= c_{R}(y-x)\mathbb{E}_{v}\left[\mathbb{E}_{Y_{1}}[\tau_{\Theta}(a)]\mathbf{1}_{\{Y_{1}>a\}}\right] + O_{u}\left(\mathbb{E}_{v}[Y_{1}\mathbf{1}_{\{Y_{1}>a\}}]x^{-(\kappa-2)}\right) + O_{u}\left(x\mathbb{E}_{v}[Y_{1}\mathbf{1}_{\{Y_{1}>x/4\}}]\right).
$$
\n(4.65)

Finally let us argue that, uniformly over $v \in (-\infty, a]$, $\mathbb{E}_v[Y_1 \mathbf{1}_{\{Y_1 > a\}}]$ is bounded and $\mathbb{E}_v[Y_1\mathbf{1}_{\{Y_1\geq x/4\}}] = O(x^{-\kappa}).$ Using the assumption $(T-1)$, $\mathbb{E}[(z^+)^{\kappa+1}] < \infty$ (recall that $\kappa = \xi - 1$) and using the fact that under \mathbb{P}_v , $Y_1 = z_1 + h(v) \leq z_1 + h(a)$ if $v \leq a$, we see that the $\kappa + 1$ moment of Y_1 is bounded (by a constant that depends on a). Hence the claim.

Integrating now over $(-\infty, a]$ against the measure ν yields that, as $x \to \infty$, uniformly in $y \in [x, 2x]$

$$
\nu((x, y]) = c_{\nu}(y - x) + O_{\mathbf{u}}\left(x^{-(\kappa - 2)}\right), \tag{4.66}
$$

with

$$
c_{\nu} = c_R \int_{(-\infty,a]} \mathbb{E}_v \left[\mathbb{E}_{Y_1}[\tau_{\Theta}(a)] \mathbf{1}_{\{Y_1 > a\}} \right] \nu(\mathrm{d}v).
$$
 (4.67)

If we know that z_+ is exponentially integrable, thus superseding $(T-1)$, we get the claimed improved control on the error by following the same argument we have just developed. The crucial changes are:

- (1) The main theorem in [\[13\]](#page-32-13) yields that $\mathbb{E}[\exp(c\mathbf{z}^+)] < \infty$ implies $\mathbb{E}[\exp(cH_1)] < \infty$.
- (2) In [\(4.63\)](#page-21-0), the term $O_u(bx^{-(\kappa-2)})$ becomes $O_u(e^{-\delta_6x})$, with some positive constant δ_6 . This can be done exactly in the same way, by applying the case (**R-2'**) of Proposition [4.6.](#page-16-0)
- (3) The expectation $\mathbb{E}_{v}[Y_1 \mathbf{1}_{\{Y_1 \ge x/4\}}]$ in [\(4.65\)](#page-21-4) is now dominated uniformly over $v \in$ $(-\infty, a]$ by $e^{-\delta_6 x}$, by eventually choosing a smaller constant δ_6 , instead of $x^{-\kappa}$.

The proof of Theorem [3.1](#page-8-1) is therefore complete. \Box

Remark 4.10. The formula [\(4.67\)](#page-22-0) provides an expression for c_{ν} , the constant of the leading term in the right-hand-side of (3.1) . We can express d_{ν} , the other constant in (3.1) , as follows:

$$
d_{\nu} = \nu((-\infty, a]) + \int_{(-\infty, a]} \mathbb{E}_{v} \left[\mathbf{1}_{\{Y_1 > a\}} \mathbb{E}_{Y_1} \left(\sum_{j=0}^{\tau_{\Theta}(a) - 1} \psi(\Theta_j) \right) \right] \nu(\mathrm{d}v), \tag{4.68}
$$

where $\psi(\theta) := c_R' - c_R \mathbb{E}[\log(e^{\theta} + \sum_{k=1}^{\infty} \tilde{\eta}_k e^{-H_{k-1}})], c_R$ and c_R' are given in [\(4.23\)](#page-15-2) and $\tilde{\eta}_j$
in (4.16). Note that defended to generally (3.13) in [\(4.16\)](#page-14-1). Note that d_{ν}/c_{ν} is related to κ_2 through [\(3.13\)](#page-10-0).

5. Contraction properties: proofs

In this section, we prove Theorem [3.2,](#page-9-1) that is, we show that the dynamical system defined by T_{Γ} is contractive with a suitably explicit rate. The techniques we use are lightly adapted from a recent and so far unpublished work by Quentin Moulard [\[27\]](#page-33-8).

Let \mathcal{M}_+ be the set of two-by-two matrices with positive elements and positive determinant, i.e. of

$$
A = \begin{pmatrix} a & b \\ c & d \end{pmatrix},\tag{5.1}
$$

with a, b, c, d, ad – bc > 0. This includes matrices of the form in Equation [\(1.1\)](#page-0-0), so long as $\varepsilon \in (0,1)$, and importantly also arbitrary products of such matrices. With the components labelled as in Equation [\(5.1\)](#page-22-1),

$$
A\begin{pmatrix} 1\\ e^x \end{pmatrix} = \begin{pmatrix} a+be^x\\ c+de^x \end{pmatrix} = (a+be^x) \begin{pmatrix} 1\\ e^{\mathcal{T}_A(x)} \end{pmatrix} \text{ with } \mathcal{T}_A(x) := \log \left(\frac{c+de^x}{a+be^x} \right),\tag{5.2}
$$

then

$$
\mathcal{T}'_A(x) = \frac{ad - bc}{(ae^{-x} + b)(c + de^x)} \quad \implies \quad 0 < \mathcal{T}'_A(x) < \frac{\det(A)}{A_{11}A_{22}} =: \tau(A) < 1,\tag{5.3}
$$

for all $A \in \mathcal{M}_+$, $x \in \mathbb{R}$; in particular \mathcal{T}_A is increasing, and so invertible on $\mathcal{T}_A(\mathbb{R}) =$ $(\log \frac{c}{a}, \log \frac{d}{b})$. We define for G of bounded total variation

$$
\mathcal{T}_A G(x) := \begin{cases} \lim_{y \to -\infty} G(y), & x \le \log(c/a) \\ G \circ \mathcal{T}_A^{-1}(x), & \log(c/a) < x < \log(d/b) \\ \lim_{y \to \infty} G(y), & x \ge \log(d/b) \end{cases} \tag{5.4}
$$

 $(\lim_{y\to\pm\infty} G(y)$ exist because G is the difference of two nonincreasing functions). Observe that if G is the risk function of some variable X, then $\mathcal{T}_A G = G_{\mathcal{T}_A(X)}$. Furthermore, note that when $M = M(e^{-\Gamma}, e^{-z})$, we have $\mathcal{T}_M(x) = z + h_\Gamma(x)$, so T_Γ introduced in [\(2.6\)](#page-6-3) can be expressed as follows: when G is the risk function of some variable X , we have

$$
T_{\Gamma}G(x) = \mathbb{E}[\mathcal{T}_M G(x)]. \qquad (5.5)
$$

and this extends to the case when G is the difference of the risk functions of two random variables. It is obvious from Equation [\(5.2\)](#page-22-2) that $\mathcal{T}_A \circ \mathcal{T}_B = \mathcal{T}_{AB}$, as a result of which

$$
(T_{\Gamma})^n G(x) = \mathbb{E}\left[\mathcal{T}_{M_1\ldots M_n} G(x)\right].
$$
\n(5.6)

Lemma 5.1. Let G_j be the risk function of $X_j \in \mathbb{L}^1$, $j = 1, 2$. Then for $A \in \mathcal{M}_+$ we have that $T_A G_j$, $j = 1, 2$, are the risk functions of two \mathbb{L}^1 random variables, in fact \mathbb{L}^{∞} , and

$$
\|\mathcal{T}_A G_1 - \mathcal{T}_A G_2\|_1 \le \tau(A) \|G_1 - G_2\|_1. \tag{5.7}
$$

Proof. We use a well-known identity: the \mathbb{L}^1 distance between the risk functions (or cumulative functions) of two real random variables is equal to the Wasserstein distance between the laws of these variables, in our case: $||G_1 - G_2||_1 = d_{W,1}(X_1, X_2)$, where with a slight abuse of notation, we write $d_{W,1}(X_1, X_2)$ for the Wasserstein distance between the laws of X_1 and of X_2 . Similarly, $||\mathcal{T}_A G_1 - \mathcal{T}_A G_2||_1 = d_{W,1}(\mathcal{T}_A(X_1), \mathcal{T}_A(X_2))$. Hence, our aim is to prove that

$$
d_{W,1}(\mathcal{T}_A(X_1), \mathcal{T}_A(X_2)) \le \tau(A) d_{W,1}(X_1, X_2). \tag{5.8}
$$

This is immediate using the Taylor inequality

$$
\forall x, y \in \mathbb{R}, \quad |\mathcal{T}_A(x) - \mathcal{T}_A(y)| \le \tau(A)|x - y|,\tag{5.9}
$$

which follows from (5.3) . \Box

Recall [\(2.12\)](#page-7-5) for $T_{\Gamma,0}G$. Combining the above lemma with Equation [\(5.6\)](#page-23-1), we have

Corollary 5.2. For any $n \in \mathbb{N}$ and any G the difference of the risk functions of two \mathbb{L}^1 random variables,

$$
||T_{\Gamma,0}^n G||_1 \le \mathbb{E}\left[\tau(M_1 \dots M_n)\right]||G||_1. \tag{5.10}
$$

Consequently, we can relate Theorem [3.2](#page-9-1) to a bound on $\sum_{n=0}^{\infty} \mathbb{E}[\tau(M_1 \dots M_n)]$. We obtain such a bound as follows: set

$$
n_{\Gamma,\sigma} := \left[37 \left(\Gamma/\sigma\right)^2\right].\tag{5.11}
$$

Lemma 5.3. There exists $\Gamma_0 < \infty$ such that

$$
\mathbb{E}\left[\tau\left(M_1 \dots M_{n_{\Gamma,\sigma}}\right)\right] \leq \frac{1}{2},\tag{5.12}
$$

for all $\Gamma \geq \Gamma_0$.

Proof. Note that $\tau(A) = \tau(\lambda A)$ for any $A \in \mathcal{M}_+$, $\lambda > 0$, so instead of M in [\(1.1\)](#page-0-0) we can equivalently consider

$$
Q_j := \begin{pmatrix} e^{-z_j/2} & e^{-\Gamma - z_j/2} \\ e^{-\Gamma + z_j/2} & e^{z_j/2} \end{pmatrix}.
$$
 (5.13)

Abbreviating $\widetilde{Q}^{(n)} = Q_1 \dots Q_n$, we have

$$
\tau(M_1 \dots M_n) = \tau\left(\widetilde{Q}^{(n)}\right) \le 1 / \left[\widetilde{Q}_{11}^{(n)} \widetilde{Q}_{22}^{(n)}\right]. \tag{5.14}
$$

Since all of the matrices involved are positive, we can bound the matrix products involved from below by single products of elements; most obviously

$$
\widetilde{Q}_{11}^{(n)}\widetilde{Q}_{22}^{(n)} \ge (Q_1)_{11} \dots (Q_n)_{11} \times (Q_1)_{22} \dots (Q_n)_{22} = 1,\tag{5.15}
$$

but also

$$
\widetilde{Q}_{11}^{(n)}\widetilde{Q}_{22}^{(n)} \ge (Q_1)_{11} \dots (Q_n)_{11} \times (Q_1)_{21} (Q_2)_{11} \dots (Q_{n-1})_{11} (Q_n)_{12} = e^{-2\Gamma} e^{-\sum_{j=2}^n z_j} \tag{5.16}
$$

and

$$
\widetilde{Q}_{11}^{(n)}\widetilde{Q}_{22}^{(n)} \ge (Q_1)_{12}(Q_2)_{22}\dots (Q_{n-1})_{22}(Q_n)_{21} \times (Q_1)_{22}\dots (Q_n)_{22} = e^{-2\Gamma}e^{\sum_{j=2}^n \mathbf{z}_j}.
$$
 (5.17)

Assembling these three estimates and combining them with Equation [\(5.14\)](#page-24-0),

$$
\tau(M_1 \dots M_n) \le \min\left(1, e^{2\Gamma} e^{\sum_{j=2}^n \mathbf{z}_j}, e^{2\Gamma} e^{-\sum_{j=2}^n \mathbf{z}_j}\right) = \min\left(1, e^{2\Gamma} e^{-|\sum_{j=2}^n \mathbf{z}_j|}\right). \tag{5.18}
$$

Since the $z_j \in \mathbb{L}^2$ are centered and IID, we can use the Central Limit Theorem together with $3\Gamma/(\sigma\sqrt{n_{\Gamma,\sigma}}) \leq 1/2$ for Γ sufficiently large to see that with N denoting a standard Gaussian variable

$$
\mathbb{P}\left(\left|\sum_{j=2}^{n_{\Gamma,\sigma}} \mathbf{z}_{j}\right| \geq 3\,\Gamma\right) \geq \mathbb{P}\left(|\mathcal{N}| \geq 1/2\right) =: \delta > \frac{3}{5}.
$$
\n
$$
(5.19)
$$

Combining this with Equation [\(5.14\)](#page-24-0) we see that by suitably choosing Γ_0 we have that for every $\Gamma \geq \Gamma_0$

$$
\mathbb{E}\left[\tau(M_1 \dots M_{n_{\Gamma,\sigma}}\right] \le (1-\delta) + \delta e^{-\Gamma} \le 1 - \delta(1 - e^{-\Gamma_0}) \le \frac{1}{2}.
$$
 (5.20)

Proof of Theorem [3.2.](#page-9-1) Combining this with Corollary [5.2](#page-23-2) and Lemma [5.3](#page-24-1) we obtain

$$
\left\|T_{\Gamma,0}^{n_{\Gamma,\sigma}}G\right\|_{1} \leq \frac{1}{2} \|G\|_{1}.
$$
\n(5.21)

Writing $n = n_{\Gamma,\sigma} m + r$ with $r = n - n_{\Gamma,\sigma} m \in [0, n_{\Gamma,\sigma})$ we have

$$
T_{\Gamma_0}^n G = T_{\Gamma,0}^{n_{\Gamma,\sigma}} \left[T_{\Gamma}^{n_{\Gamma,\sigma}} \dots [T_{\Gamma}^r G] \right];\tag{5.22}
$$

noting that $||T_{\Gamma,0}^rG||_1 \leq ||G||_1$ (as can be seen from either Equation [\(2.13\)](#page-7-6) or Equation [\(5.18\)](#page-24-2)), applying Corollary [5.2](#page-23-2) and Lemma [5.3](#page-24-3) iteratively leads to

$$
\left\|T_{\Gamma,0}^n G\right\|_1 \le \left(\frac{1}{2}\right)^m \|G\|_1,
$$
\n(5.23)

so that

$$
\sum_{n=0}^{\infty} \|T_{\Gamma,0}^n G\|_1 \le n_{\Gamma,\sigma} \sum_{m=0}^{\infty} \left(\frac{1}{2}\right)^m = 2 n_{\Gamma,\sigma} \le \frac{75}{\sigma^2} \Gamma^2,\tag{5.24}
$$

which completes the proof of Theorem [3.2.](#page-9-1) \Box

6. The one-step bound

This section is devoted to the proof of Proposition [3.5.](#page-10-2)

Proof. We start by observing that (3.14) is a direct consequence of (3.10) . Moreover (3.15) follows by the following simple calculation: by (3.3) and (3.11) we have

$$
L_{\Gamma}[G_{\Gamma}] = \frac{1}{C_{\Gamma}} \int_0^{\infty} \frac{F_{\triangleright}(\Gamma - x)}{1 + e^{\Gamma - x}} dx + \int_{-\infty}^0 \frac{G_{\Gamma}(x)}{1 + e^{\Gamma - x}} dx, \qquad (6.1)
$$

where for notational brevity, we will write in this proof G_{Γ} in lieu of $G_{\gamma_{\Gamma}}$, and analogous notation for the cumulative function. Using $G_{\Gamma}(x) \in [0,1]$ we readily see that the second addendum is smaller than $\exp(-\Gamma)$. Moreover

$$
\int_0^\infty \frac{F_{\rhd}(\Gamma - x)}{1 + e^{\Gamma - x}} dx = \int_{\mathbb{R}} \frac{F_{\rhd}(y)}{1 + e^y} dy - \int_{\Gamma}^\infty \frac{F_{\rhd}(y)}{1 + e^y} dy = \int_{\mathbb{R}} \frac{F_{\rhd}(y)}{1 + e^y} dy + O\left(\Gamma \exp(-\Gamma)\right),\tag{6.2}
$$

where in the last step we used [\(3.10\)](#page-10-3) for F_{\triangleright} instead of F_{\triangleleft} .

Remark 6.1. Repeating the same argument using instead the functional $L_{\Gamma}^{\star}[\cdot]$ defined in [\(3.6\)](#page-9-4) of Remark [3.4](#page-9-5) one obtains

$$
L_{\Gamma}^{\star}[G_{\gamma_{\Gamma}}] = \frac{1}{C_{\Gamma}} \int_{\mathbb{R}} \frac{F_{\lhd}(y)}{1 + e^{y}} dy + O(\exp(-\Gamma)). \tag{6.3}
$$

Then, similarly to (3.17) , we arrive at another formula for κ_1 :

$$
\kappa_1 = \frac{1}{2} \int_{\mathbb{R}} \frac{F_{\triangleleft}(y)}{1 + e^y} dy = \frac{1}{2} \int \log(1 + e^{-y}) \nu(dy), \qquad (6.4)
$$

where the second equality follows from an integration by parts, and we recall that F_{\lhd} is the cumulative function of the (normalized) stationary measure ν of Y. This highlights a symmetry with respect to exchanging F_{\lhd} and F_{\rhd} .

Now for the proof of [\(3.16\)](#page-11-1). We refer to assumption "(T-1) for $\pm z$ " as " $z \in \mathbb{L}^{\xi}$ " and to "(T-1') for $\pm z$ " as "exponential tails assumption" and, when the assumption is omitted, $z \in \mathbb{L}^{\xi}$ is assumed. We have

$$
||T_{\Gamma}G_{\Gamma} - G_{\Gamma}||_1 = ||(T_{\Gamma}G_{\Gamma} - G_{\Gamma})\mathbf{1}_{(-\infty,0)}||_1 + ||(T_{\Gamma}G_{\Gamma} - G_{\Gamma})\mathbf{1}_{(0,\infty)}||_1
$$

=
$$
||(T_{\Gamma}F_{\Gamma} - F_{\Gamma})\mathbf{1}_{(-\infty,0)}||_1 + ||(T_{\Gamma}G_{\Gamma} - G_{\Gamma})\mathbf{1}_{(0,\infty)}||_1.
$$
 (6.5)

If ζ is symmetric the two terms coincide; in general, they do not coincide but they can be treated in the same way because one can be obtained from the other by replacing z with −z. So we focus on the first one and use [\(2.7\)](#page-6-4) to obtain

$$
\left\| (T_{\Gamma} F_{\Gamma} - F_{\Gamma}) \mathbf{1}_{(-\infty,0)} \right\|_{1} = \left\| (F_{\mathbf{z}}(-\Gamma + \cdot) + T_{\Gamma,0} F_{\Gamma} - F_{\Gamma}) \mathbf{1}_{(-\infty,0)} \right\|_{1}
$$

\$\leq \int_{-\infty}^{-\Gamma} F_{\mathbf{z}}(x) dx + \left\| (T_{\Gamma,0} F_{\Gamma} - F_{\Gamma}) \mathbf{1}_{(-\infty,0)} \right\|_{1} .\$ (6.6)

Note that the first addendum in the right-most term is $O(\Gamma^{-\xi+1})$ when $z \in \mathbb{L}^{\xi}$ (and $O(\exp(-c\Gamma))$ assuming exponential tails). So we focus on the second one, which is equal to

$$
A_{\Gamma,1} := \int_{-\infty}^{0} \left| \int \mathbf{1}_{(x-\Gamma,x+\Gamma)}(z) F_{\Gamma}(h_{\Gamma}^{-1}(x-z)) \zeta(\mathrm{d}z) - F_{\Gamma}(x) \right| \mathrm{d}x, \tag{6.7}
$$

with $F_{\Gamma}(\cdot)$ given in [\(3.12\)](#page-10-5).

The first step in controlling $A_{\Gamma,1}$ is to remark that we can avoid the nuisance of the fact that $F_{\Gamma}(y)$ has two different expressions according to the sign of y. Namely, we want to relate it to

$$
A_{\Gamma,2} := \frac{1}{C_{\Gamma}} \int_{-\infty}^{0} \left| \int \mathbf{1}_{(x-\Gamma,x+\Gamma)}(z) F_{\lhd}(h_{\Gamma}^{-1}(x-z)+\Gamma) \zeta(\mathrm{d}z) - F_{\lhd}(x+\Gamma) \right| \mathrm{d}x. \tag{6.8}
$$

We can do this because

$$
|A_{\Gamma,2} - A_{\Gamma,1}| \leq \int_{-\infty}^{0} \int \mathbf{1}_{(x-\Gamma,x)}(z) \left| \frac{F_{\lhd}(\Gamma+y) + F_{\rhd}(\Gamma-y)}{C_{\Gamma}} - 1 \right| \Big|_{y=h_{\Gamma}^{-1}(x-z)} \zeta(\mathrm{d}z) \mathrm{d}x \,, \tag{6.9}
$$

(note h_{Γ}^{-1} $_{\Gamma}^{-1}(0) = 0, h_{\Gamma}^{-1}$ $\Gamma^{-1}(\Gamma^-) = +\infty$, so $\mathbf{1}_{(x-\Gamma,x+\Gamma)}(z)\mathbf{1}_{(0,\infty)}(h_{\Gamma}^{-1})$ $\Gamma^{-1}(x-z)$ = $\mathbf{1}_{(x-\Gamma,x)}(z)$ and, by (3.10) and (3.14) , we have

$$
\left|\frac{F_{\lhd}(\Gamma+y)+F_{\rhd}(\Gamma-y)}{C_{\Gamma}}-1\right| \leq C \times \begin{cases} \Gamma^{-\xi+2} & \text{if } y \in [0,\Gamma/2],\\ y/\Gamma & \text{if } y > \Gamma/2, \end{cases} \tag{6.10}
$$

for suitably chosen $C > 0$ and for Γ sufficiently large: in the exponential case the first line of the estimate may be replaced by $\exp(-\delta \Gamma/2)/\Gamma$. So $|A_{\Gamma,2} - A_{\Gamma,1}|$ is bounded above by a constant times

$$
\Gamma^{-\xi+2} \int_{-\infty}^{0} \int \mathbf{1}_{(x-\Gamma,x)}(z) \zeta(\mathrm{d}z) \mathrm{d}x + \Gamma^{-1} \int_{-\infty}^{0} \int \mathbf{1}_{(x-\Gamma,x-h_{\Gamma}(\Gamma/2))}(z) h_{\Gamma}^{-1}(x-z) \zeta(\mathrm{d}z) \mathrm{d}x \leq \Gamma^{-\xi+2} \int_{-\infty}^{0} F_{\zeta}(x) \mathrm{d}x + \Gamma^{-1} F_{\zeta}(-h_{\Gamma}(\Gamma/2)) \int_{\mathbb{R}} |y| h_{\Gamma}'(y) \mathrm{d}y,
$$
\n(6.11)

where the second addendum is treated by a change of variable of integration from x to $y=h_{\Gamma}^{-1}$ $\Gamma^{-1}(x-z)$, taking the absolute value of the integrand and by releasing the integration constraints on y (in the exponential case the prefactors $\Gamma^{-\xi+2}$ in [\(6.11\)](#page-26-0) may be replaced by $\exp(-\delta \Gamma/2)/\Gamma$). The assumption on $z \in \mathbb{L}^{\xi}$ together with the Markov inequality imply

 $F_{\zeta}(x) = O(|x|^{-\xi})$ for $x \to -\infty$ $(F_{\zeta}(x)) = O(\exp(-c|x|))$ in the exponential case). Since $h_{\Gamma}(\Gamma/2) \sim \Gamma/2$, $h'_{\Gamma}(y) < 1$ and the fact that $h'_{\Gamma}(y)$ decays exponentially for $y > \Gamma$ (so $\int |y|h'_{\Gamma}(y)dy = O(\Gamma^2)$, we have $|A_{\Gamma,2} - A_{\Gamma,1}| = O(\Gamma^{-\xi+2}) + O(\Gamma^{-\xi+1})$ (in the exponential case: $O(\exp(-\delta \Gamma/2)/\Gamma) + O(\Gamma \exp(-c\Gamma/2))),$ so

$$
|A_{\Gamma,2} - A_{\Gamma,1}| = \begin{cases} O(\Gamma^{-\xi+2}) & \text{if } z \in \mathbb{L}^{\xi}, \\ O(\Gamma \exp(-(\delta \wedge c)\Gamma/2)) & \text{under exponential tails assumption.} \end{cases}
$$
(6.12)

Having shown this, we now need to control $A_{\Gamma,2}$. To do this we introduce

$$
\mathbf{h}_{\Gamma}(x) := h(x + \Gamma) - \Gamma \tag{6.13}
$$

which approximates h_{Γ} for a certain range of arguments in the sense that

$$
0 \leq \mathbf{h}_{\Gamma}(x) - h_{\Gamma}(x) = \log(1 + \exp(x - \Gamma)) = \begin{cases} O(\exp(x - \Gamma)) & \text{for } x - \Gamma \to -\infty, \\ O(x - \Gamma) & \text{for } x - \Gamma \to +\infty, \end{cases}
$$
(6.14)

$$
0 \leq \mathbf{h}'_{\Gamma}(x) - h'_{\Gamma}(x) = 1/(1 + \exp(\Gamma - x)) = \begin{cases} O(\exp(x - \Gamma)) & \text{for } x - \Gamma \to -\infty, \\ \leq 1 & \text{for every } x \text{ and } \Gamma, \end{cases}
$$
(6.15)

and

$$
0 \le h_{\Gamma}^{-1}(y) - \mathbf{h}_{\Gamma}^{-1}(y) = -\log(1 - e^{y - \Gamma}) \qquad \forall y \in (-\Gamma, \Gamma). \tag{6.16}
$$

Moreover we point out that $h'_{\Gamma}(x)$ and $h'_{\Gamma}(x)$ are in $(0,1)$ for every x and, as shown in Figure [1,](#page-5-1) the two functions almost coincide for $x \ll \Gamma$. They start to differ when x approaches Γ from the left, where $\mathbf{h}'_{\Gamma}(x)$ keeps being very close to one while $\mathbf{h}_{\Gamma}(x) \sim x$.

Using the characterization of $F_{\lhd}(\cdot)$ as the cumulative function of a stationary measure,

$$
\int \mathbf{1}_{(-\infty,x+\Gamma)}(z) F_{\lhd}(\mathbf{h}_{\Gamma}^{-1}(x-z)+\Gamma) \zeta(\mathrm{d}z) - F_{\lhd}(x+\Gamma) =
$$
\n
$$
\int \mathbf{1}_{(-\infty,x)}(z) F_{\lhd}(h^{-1}(x-z)) \zeta(\mathrm{d}z) - F_{\lhd}(x) = 0, \quad (6.17)
$$

and so

$$
A_{\Gamma,2} = \frac{1}{C_{\Gamma}} \int_{-\infty}^{0} \left| \int \left[\mathbf{1}_{(x-\Gamma,x+\Gamma)}(z) F_{\lhd}(h_{\Gamma}^{-1}(x-z)+\Gamma) - \mathbf{1}_{(-\infty,x+\Gamma)}(z) F_{\lhd}(\mathbf{h}_{\Gamma}^{-1}(x-z)+\Gamma) \right] \zeta(\mathrm{d}z) \right| \mathrm{d}x
$$

\n
$$
\leq \frac{1}{C_{\Gamma}} \int_{-\infty}^{0} \int \left\{ \mathbf{1}_{(-\infty,x-\Gamma/2]}(z) F_{\lhd}(\mathbf{h}_{\Gamma}^{-1}(x-z)+\Gamma) + \mathbf{1}_{(x-\Gamma,x-\Gamma/2]}(z) F_{\lhd}(h_{\Gamma}^{-1}(x-z)+\Gamma) \right. \\ \left. + \mathbf{1}_{(x-\Gamma/2,x+\Gamma)}(z) \left[F_{\lhd}(h_{\Gamma}^{-1}(x-z)+\Gamma) - F_{\lhd}(\mathbf{h}_{\Gamma}^{-1}(x-z)+\Gamma) \right] \right\} \zeta(\mathrm{d}z) \mathrm{d}x
$$

\n=: $T_{1} + T_{2} + T_{3}$. (6.18)

For T_1 we first note that, since \mathbf{h}_{Γ}^{-1} $\Gamma^{-1}(x') \sim x'$ and $F_{\lhd}(x') \sim x'$ for $x' \to +\infty$, we have for Γ large

$$
\begin{split} \mathbf{1}_{(-\infty,x-\Gamma/2]}(z)F_{\lhd}(\mathbf{h}_{\Gamma}^{-1}(x-z)+\Gamma) &= \mathbf{1}_{[\Gamma/2,\infty)}(x-z)F_{\lhd}(\mathbf{h}_{\Gamma}^{-1}(x-z)+\Gamma) \\ &\leq 2(x-z+\Gamma)\mathbf{1}_{[\Gamma/2,\infty)}(x-z), \end{split} \tag{6.19}
$$

and integrating by parts (keep in mind that $x < 0$)

$$
\int (x-z)\mathbf{1}_{[\Gamma/2,\infty)}(x-z)\zeta(\mathrm{d}z) = \frac{\Gamma}{2}F_{\zeta}(x-\Gamma) + \int_{-\infty}^{x-\Gamma/2} F_{\zeta}(z)\mathrm{d}z = O\left(|x-\Gamma|^{-\xi+1}\right),\tag{6.20}
$$

where we have used that $z \in \mathbb{L}^{\xi}$: in the exponential case the estimate is $O(\exp(-c|x-\Gamma|/2))$. Regarding the remaining term in Equation [\(6.19\)](#page-27-0), note that

$$
\Gamma \int \mathbf{1}_{\left[\Gamma/2,\infty\right)} (x-z)\zeta(\mathrm{d}z) = \Gamma F_{\zeta} (x-\Gamma/2) \,, \tag{6.21}
$$

and we can bound this by the Markov inequality, combining with Equation [\(6.20\)](#page-27-1) to give

$$
\int (x - z + \Gamma) \mathbf{1}_{[\Gamma/2, \infty)} (x - z) \zeta(\mathrm{d}z) = O\left(|x - \Gamma|^{-\xi + 1}\right),\tag{6.22}
$$

for $z \in \mathbb{L}^{\xi}$ and $O(|x-\Gamma|\exp(-c|x-\Gamma|/2))$ in the exponential case. Therefore integrating in x and dividing by C_{Γ} we obtain $T_1 = O(\Gamma^{-\xi+1})$ and $O(\exp(-c\Gamma/3))$ in the exponential case.

For T_2 we use $F_{\lhd}(x') \sim x'$ for $x' \to \infty$ (note that $x - z \in [\Gamma/2, \Gamma)$) so

$$
F_{\lhd}(h_{\Gamma}^{-1}(x-z)+\Gamma) \le 2(h_{\Gamma}^{-1}(x-z)+\Gamma). \tag{6.23}
$$

Moreover

$$
\mathbf{1}_{(-\infty,0)}(x)\mathbf{1}_{(x-\Gamma,x-\Gamma/2]}(z) = \mathbf{1}_{(-\infty,-\Gamma]}(z)\mathbf{1}_{(z+\Gamma/2,z+\Gamma]}(x) + \mathbf{1}_{(-\Gamma,-\Gamma/2)}(z)\mathbf{1}_{(z+\Gamma/2,0]}(x),\tag{6.24}
$$

and we remark that

$$
\int_{z+\frac{\Gamma}{2}}^{z+\Gamma} (h_{\Gamma}^{-1}(x-z)+\Gamma) dx = \int_{\frac{\Gamma}{2}}^{\Gamma} (h_{\Gamma}^{-1}(x)+\Gamma) dx = O(\Gamma^2), \qquad (6.25)
$$

because when $x \nearrow \Gamma$ we have h_{Γ}^{-1} $\Gamma^{-1}_\Gamma(x) - \Gamma \sim \log(\Gamma - x)$ and the integral converges. For the other term we have by using that $-z \in (-\Gamma/2, \Gamma)$ and the behavior of h_{Γ}^{-1} $^{-1}$ approaching Γ

$$
\int_{z+\frac{\Gamma}{2}}^{0} (h_{\Gamma}^{-1}(x-z)+\Gamma) dx = \int_{\frac{\Gamma}{2}}^{-z} (h_{\Gamma}^{-1}(x)+\Gamma) dx \le \int_{\frac{\Gamma}{2}}^{\Gamma} (h_{\Gamma}^{-1}(x)+\Gamma) dx = O(\Gamma^{2}), \tag{6.26}
$$

By integrating over z, which is required to be smaller that $-\Gamma/2$ in both terms, and dividing by C_{Γ} we see that $T_2 = O(\Gamma^{-\xi+1})$ and $O(\Gamma \exp(-c\Gamma/2))$ in the exponential case.

For T_3 we note that by (6.16)

$$
0 \le h_{\Gamma}^{-1}(y) - \mathbf{h}_{\Gamma}^{-1}(y) = O\left(e^{y-\Gamma}\right),\tag{6.27}
$$

when $y \in (-\Gamma, \Gamma/2)$. We then note that $C_{\Gamma}T_3$ is equal to

$$
\int_{-\infty}^{0} dx \int_{\mathbb{R}} \zeta(\mathrm{d}z) \mathbf{1}_{(x-\Gamma/2, x+\Gamma)}(z) \int_{\mathbf{1}_{(h_{\Gamma}^{-1}(x-z), h_{\Gamma}^{-1}(x-z))}(y-\Gamma) \nu(\mathrm{d}y) =
$$
\n
$$
\int_{-\infty}^{\Gamma} \zeta(\mathrm{d}z) \int_{z-\Gamma}^{(z+\Gamma/2)\wedge 0} dx \int_{\mathbf{1}_{(h_{\Gamma}^{-1}(x-z), h_{\Gamma}^{-1}(x-z))}(y-\Gamma) \nu(\mathrm{d}y). \quad (6.28)
$$

We then change variable from x to $x - z$ and ignore the constraint $x \leq 0$ to obtain

$$
C_{\Gamma}T_3 \leq \int_{-\infty}^{\Gamma} \zeta(\mathrm{d}z) \int_{-\Gamma}^{\Gamma/2} \mathrm{d}x \int \mathbf{1}_{(\mathbf{h}_{\Gamma}^{-1}(x-z), h_{\Gamma}^{-1}(x-z))} (y-\Gamma) \nu(\mathrm{d}y)
$$

\n
$$
\leq \int_{-\Gamma}^{\Gamma/2} \mathrm{d}x \int \mathbf{1}_{(\mathbf{h}_{\Gamma}^{-1}(x-z), h_{\Gamma}^{-1}(x-z))} (y-\Gamma) \nu(\mathrm{d}y)
$$

\n
$$
= \int_{-\infty}^{h_{\Gamma}^{-1}(\Gamma/2)+\Gamma} \nu(\mathrm{d}y) \int_{h_{\Gamma}(y-\Gamma)\wedge(-\Gamma)}^{h_{\Gamma}(y-\Gamma)\vee(\Gamma/2)} \mathrm{d}x
$$

\n
$$
\leq \int_{-\infty}^{h_{\Gamma}^{-1}(\Gamma/2)+\Gamma} \nu(\mathrm{d}y) \left(\mathbf{h}_{\Gamma}(y-\Gamma) - h_{\Gamma}(y-\Gamma)\right) \mathrm{d}x,
$$
\n(6.29)

so, by [\(6.27\)](#page-28-0), we see that $C_{\Gamma}T_3$ is bounded above by a constant times

$$
\int_{-\infty}^{h_{\Gamma}^{-1}(\Gamma/2)+\Gamma} e^{y-2\Gamma} \nu(\mathrm{d}y) \le e^{h_{\Gamma}^{-1}(\Gamma/2)-\Gamma} \nu\left((-\infty, h_{\Gamma}^{-1}(\Gamma/2)+\Gamma]\right) = O\left(\Gamma e^{-\Gamma/2}\right). \tag{6.30}
$$

Therefore T_3 is negligible.

By collecting all the estimates, and properly redefining the value of $\delta > 0$ in the exponential case, we obtain (3.16) and this completes the proof of Proposition [3.5.](#page-10-2) \Box

APPENDIX A. $\kappa_1 = \text{var}(\mathbf{z})/4$: A PROOF Recall that $\kappa_1 = \frac{1}{2}$ $\frac{1}{2} \int \log(1 + e^{-x}) \nu(\mathrm{d}x)$ (Equation [\(6.4\)](#page-25-0)).

Proposition A.1. Under the assumptions of Theorem [3.1,](#page-8-1) but requiring that $\xi > 4$, we have that $\int_{\mathbb{R}} \log(1 + e^{-x}) \nu(\mathrm{d}x) = \mathbb{E}[z^2]/2$.

Proof. For conciseness we treat only the case of assumption $(T-1)$ of Theorem [3.1](#page-8-1) and we remind the reader that the assumptions include $z_-\in\mathbb{L}^2$, so $z\in\mathbb{L}^2$. Note that these assumptions are weaker than those of our main result, Theorem [1.2.](#page-1-2)

We recall that Theorem [3.1](#page-8-1) allows us to choose ν such that

$$
\nu((-\infty, x]) = x + d_{\nu} + r_{\nu}(x) \quad \text{with } r_{\nu}(x) = \begin{cases} O\left(x^{-(\xi - 3)}\right) & \text{for } x \to \infty \\ O(|x|) & \text{for } x \to -\infty, \end{cases} \tag{A.1}
$$

with a proper choice of $d_{\nu} \in \mathbb{R}$. The (very weak) bound for $x \to -\infty$ comes from the boundedness of $F_{\nu}(x) = \nu((-\infty, x])$ in this limit. In fact, we have $F_{\nu}(x) = O(|x|^{-2})$ for $x \to -\infty$, as a consequence of $z_-\in \mathbb{L}^2$ and Proposition [2.4.](#page-7-2) We remark that $F_\nu(x) =$ $O(|x|^{-2})$ for $x \to -\infty$ and $\nu((-\infty, x]) \sim x$ for $x \to \infty$ yield that $\int_{\mathbb{R}} \log(1 + e^{-x}) \nu(\mathrm{d}x) =$ $\int_{\mathbb{R}} \frac{1}{1+h}$ $\frac{1}{1+e^x}F_{\nu}(\mathrm{d}x)$ is finite.

By [\(2.16\)](#page-8-3) applied to $g(x) = x\mathbf{1}_{(-\infty,b]}(x)$ we have that for every $b \in \mathbb{R}$

$$
\int x\mathbf{1}_{(-\infty,b]}(x)\,\nu(\mathrm{d}x) = \iint \left(z+h(x)\right)\mathbf{1}_{(-\infty,b]}(z+h(x))\nu(\mathrm{d}x)\zeta(\mathrm{d}z). \tag{A.2}
$$

Both sides diverge when $b \to \infty$, so we rearrange to obtain the form $I_1 = I_2 - I_3$, with:

$$
I_1 := \iint (h(x) - x) \mathbf{1}_{(-\infty, b]}(z + h(x)) \nu(\mathrm{d}x) \zeta(\mathrm{d}z),
$$

\n
$$
I_2 := -\iint z \mathbf{1}_{(-\infty, b]}(z + h(x)) \nu(\mathrm{d}x) \zeta(\mathrm{d}z),
$$

\n
$$
I_3 := \iint (\mathbf{1}_{(-\infty, b]}(z + h(x)) - \mathbf{1}_{(-\infty, b]}(x)) x \nu(\mathrm{d}x) \zeta(\mathrm{d}z).
$$
\n(A.3)

We pass to the $b \to \infty$ limit in the three terms I_1 , I_2 and I_3 separately. In all three cases we exploit the properties of $h(\cdot)$, in particular that $h(x) = x + O(\exp(-x))$ for $x \to \infty$, to replace $h(x)$ with x in the argument of the indicator function.

For I_1 we have

$$
I_1 = \iint \log (1 + e^{-x}) \mathbf{1}_{(-\infty, b]}(z + h(x)) \nu(\mathrm{d}x) \zeta(\mathrm{d}z)
$$

=
$$
\int_{(-\infty, b)} \int_{(-\infty, b^{-1}(b-z)]} \log (1 + e^{-x}) \nu(\mathrm{d}x) \zeta(\mathrm{d}z)
$$

=
$$
\mathbb{P}(\mathbf{z} < b) \int_{\mathbb{R}} \log (1 + e^{-x}) \nu(\mathrm{d}x) - \int_{(-\infty, b)} \int_{(h^{-1}(b-z), \infty)} \log (1 + e^{-x}) \nu(\mathrm{d}x) \zeta(\mathrm{d}z),
$$
(A.4)

so, recalling that $\int_{\mathbb{R}} \log(1 + e^{-x}) \nu(\mathrm{d}x) < \infty$,

$$
\left| I_1 - \int_{\mathbb{R}} \log(1 + e^{-x}) \nu(\mathrm{d}x) \right| = O\left(b^{-\xi} \right) + \int_{(-\infty, b)} \int_{(h^{-1}(b-z), \infty)} \log\left(1 + e^{-x}\right) \nu(\mathrm{d}x) \zeta(\mathrm{d}z), \tag{A.5}
$$

and the last integral is bounded from above, by separating the case $z \leq b/2$ and the case $z \in (b/2, b)$, by

$$
\int_{(h^{-1}(b/2),\infty)} e^{-x} \nu(\mathrm{d}x) + \mathbb{P}\left(z \in (b/2,b)\right) \int_{\mathbb{R}} \log\left(1 + e^{-x}\right) \nu(\mathrm{d}x),\tag{A.6}
$$

and we see that the first addendum is $O(\exp(-b/3))$ and the second is $O(b^{-\xi})$. Therefore

$$
I_1 = \int_{\mathbb{R}} \log(1 + e^{-x}) \nu(\mathrm{d}x) + O(b^{-\xi}). \tag{A.7}
$$

For the term I_2 , we start by observing that, since $h(x) > x$ and $h(\mathbb{R}) = (0, \infty)$,

$$
\int \left(\mathbf{1}_{(-\infty,b]}(z+x)-\mathbf{1}_{(-\infty,b]}(z+h(x))\right)\nu(\mathrm{d}x) = \begin{cases} \nu\left(\left(h^{-1}(b-z),b-z\right]\right) & \text{if } z < b, \\ \nu\left((-\infty,b-z]\right) & \text{if } z \ge b, \\ (A.8) & \end{cases}
$$

which is $O(b^{-(\xi-3)})$ if $z < b/2$ and $O(b)$ otherwise. Therefore we see that the expression in [\(A.8\)](#page-30-0) multiplied by z and integrated with respect to $\zeta(\mathrm{d}z)$ is $O(b^{-(\xi-3)})$ and so

$$
I_2 = -\int z \nu ((-\infty, b-z]) \zeta(\mathrm{d}z) + O(b^{-(\xi-3)}) = \mathbb{E}[z^2] - \int z r_{\nu}(b-z) \zeta(\mathrm{d}z) + O(b^{-(\xi-3)}),
$$
\n(A.9)

where in the second step we used [\(A.1\)](#page-29-1). By separating once again the cases $z < b/2$ and $z \ge b/2$ we see that $\int zr_{\nu}(b-z)\zeta(\mathrm{d}z) = O(b^{-(\xi-3)})$. Therefore

$$
I_2 = \mathbb{E}[\mathbf{z}^2] + O(b^{-(\xi - 3)}).
$$
 (A.10)

Before handling the estimation of I_3 , we establish the following estimate:

$$
\int_{(-\infty,y]} x \nu(\mathrm{d}x) = \frac{1}{2}y^2 + C_\nu + R_\nu(y), \quad \text{with } R_\nu(y) = \begin{cases} O\left(y^{-(\xi-4)}\right) & \text{for } y \to \infty, \\ O(y^2) & \text{for } y \to -\infty, \end{cases} \tag{A.11}
$$

with C_{ν} a real constant that is implicitly defined by $(A.11)$ and will be given explicitly below. In fact the integration by parts formula for measures yields

$$
\int_{(-\infty,y]} x \nu(\mathrm{d}x) = y F_{\nu}(y) - \int_{-\infty}^{y} F_{\nu}(x) \mathrm{d}x, \tag{A.12}
$$

because the identity function $x \mapsto x$ is differentiable, so by [\(A.1\)](#page-29-1) we obtain for $y > 0$

$$
\int_{(-\infty,y]} x \nu(\mathrm{d}x) = y F_{\nu}(y) - \int_0^y F_{\nu}(x) \mathrm{d}x - \int_{-\infty}^0 F_{\nu}(x) \mathrm{d}x \n= \frac{y^2}{2} + y r_{\nu}(y) + \int_y^{\infty} r_{\nu}(x) (x) \mathrm{d}x + \left[- \int_{-\infty}^0 F_{\nu}(x) \mathrm{d}x - \int_0^{\infty} r_{\nu}(x) \mathrm{d}x \right],
$$
\n(A.13)

Considering the asymptotics for $y \to \infty$ we directly obtain the desired estimate, with C_{ν} equal to the expression in square brackets in the last line; note that the terms with d_{ν} cancel. We then complete the proof of [\(A.11\)](#page-31-0) by observing that $F_{\nu}(x) = O(1/x^2)$ for $x \to -\infty$ yields $R_{\nu}(y) \sim -y^2/2$ for $y \to -\infty$.

We now estimate I_3 . We start by establishing that for a suitable choice of $c > 0$, for all $b \geq 1$

$$
\left| \int \left[\mathbf{1}_{(-\infty, b]}(z + x) - \mathbf{1}_{(-\infty, b]}(z + h(x)) \right] x \, \nu(\mathrm{d}x) \right| \le c \begin{cases} b^{-(\xi - 4)} & \text{if } z < b/2 \,, \\ b^2 & \text{if } b/2 \le z < b \,, \\ 1 & \text{if } z \ge b \,. \end{cases} \tag{A.14}
$$

If $z \geq b$ we have that $\mathbf{1}_{(-\infty,b]}(z+h(x))=0$ for every x because $h(x) > 0$ so the term in [\(A.14\)](#page-31-1) is bounded by $\int_{(-\infty,b-z]} |x| \nu(\mathrm{d}x) \leq \int_{(-\infty,0]} |x| \nu(\mathrm{d}x) < \infty$. If instead $z < b$ we have

$$
0 \leq \int \left[\mathbf{1}_{(-\infty, b]}(z + x) - \mathbf{1}_{(-\infty, b]}(z + h(x)) \right] x \nu(\mathrm{d}x)
$$

=
$$
\int_{(h^{-1}(b-z), b-z]} x \nu(\mathrm{d}x)
$$

\$\leqslant (b-z)\nu ((h^{-1}(b-z), b-z])\$ (A.15)

We estimate this last term by considering separately $z < b/2$ and $z \in [b/2, b)$. In the first case $h^{-1}(b-z) = b - z + O(\exp(-b/2))$ so, by [\(A.1\)](#page-29-1), we see that $\nu((h^{-1}(b-z), b-z)) =$ $O(b^{-(\xi-3)})$. In the other case, $z \in [b/2, b)$, we use instead $\nu((h^{-1}(b-z), b-z]) \le$ $\nu ((-\infty, b/2]) = O(b)$. This yields [\(A.14\)](#page-31-1).

From [\(A.14\)](#page-31-1) we easily infer that

$$
\iint \left[\mathbf{1}_{(-\infty,b]}(z+x) - \mathbf{1}_{(-\infty,b]}(z+h(x)) \right] x \, \nu(\mathrm{d}x) \zeta(\mathrm{d}z) \, = \, O\left(b^{-(\xi-4)}\right) \,, \tag{A.16}
$$

so to estimate I_3 it suffices to consider

$$
\iint \left(\mathbf{1}_{(-\infty,b]}(z+x) - \mathbf{1}_{(-\infty,b]}(x) \right) x \nu(\mathrm{d}x) \zeta(\mathrm{d}z) \stackrel{(A.11)}{=} \\
\int \left(\frac{1}{2} \left((b-z)^2 - b^2 \right) + R_\nu(b-z) - R_\nu(b) \right) \zeta(\mathrm{d}z) = \frac{1}{2} \mathbb{E}[z^2] + O\left(b^{-(\xi-4)} \right), \quad (A.17)
$$

where the estimate of the term containing $R_{\nu}(b-z)$ is performed as before, by separately considering two cases: for $z > b/2$, we use the fact that $|R_{\nu}(b-z)| = O(z^2)$, obtaining a bound $O(b^{-(\xi-2)})$; whereas for $z \leq b/2$, we obtain $O(b^{-(\xi-4)})$. Therefore

$$
I_3 = \frac{1}{2} \mathbb{E}[\mathbf{z}^2] + O\left(b^{-(\xi - 4)}\right). \tag{A.18}
$$

By putting together the estimates on I_1 , I_2 and I_3 we complete the proof of Proposi-tion [A.1.](#page-29-2) \Box

Acknowledgements

The work of R.L.G. was supported in part by the MIUR Excellence Department Project MatMod@TOV awarded to the Department of Mathematics, University of Rome Tor Vergata, CUP E83C23000330006. G.G. acknowledges the support of the Cariparo Foundation. Y.H. acknowledges the support of ANR LOCAL.

REFERENCES

- [1] S. Asmussen, Applied Probability and Queues, Second edition, Springer-Verlag, New York, 2003.
- [2] M. Babillot, P. Bougerol and L. Elie, The random difference equation $X_n = A_n X_{n-1} + B_n$ in the critical case, Ann. Probab. 25(1997), 478-493.
- [3] P. Bougerol and J. Lacroix, *Products of random matrices with applications to Schrödinger operators*, Progress in Probability and Statistics, 8. Birkhäuser, 1985.
- [4] D. Buraczewski, E. Damek and T. Mikosch, Stochastic models with power-law tails. The equation $X=AX+B$, Springer Series in Operations Research and Financial Engineering. Springer, 2016.
- [5] H. Carlsson, Remainder term estimates of the renewal function, Ann. Probab. 11 (1983), 143-157.
- [6] O. Collin, On the large interaction asymptotics of the free energy density of the Ising chain with disordered centered external field, [arXiv:2403.19626](http://arxiv.org/abs/2403.19626)
- [7] O. Collin, G. Giacomin and Y. Hu, Infinite disorder renormalization fixed point for the continuum random field Ising chain, Probab. Theory Related Fields 190 (2024), 881-939.
- [8] O. Collin, G. Giacomin and Y. Hu, The random field Ising chain domain-wall structure in the large interaction limit, [arXiv:2401.03927](http://arxiv.org/abs/2401.03927)
- [9] F. Comets, G. Giacomin and R. L. Greenblatt, Continuum limit of random matrix products in statistical mechanics of disordered systems, Commun. Math. Phys. 369 (2019), 171-219.
- [10] A. Comtet, J.-M. Luck, C. Texier and Y. Tourigny, The Lyapunov exponent of products of random 2×2 matrices close to the identity, J. Statist. Phys. 150 (2013), 13-65.
- [11] A. Crisanti, G. Paladin and A. Vulpiani, Products of random matrices in statistical physics, Springer Series in Solid-State Sciences 104, 1993.
- [12] Derrida and H.J. Hilhorst, Singular behaviour of certain infinite products of random 2×2 matrices, J. Phys. A 16 (1983), 2641-2654.
- [13] R. A. Doney, Moments of ladder heights in random walks, J. Appl. Probab. 17 (1980), 248-252.
- [14] R. Douc, E. Moulines, P. Priouret and P. Soulier, Markov Chains, Springer Series in Operation Research and Financial Engineering, Springer, 2018.
- [15] W. Feller, An Introduction to Probability Theory and Its Applications, Volume II, Second Edition, John Wiley & Sons, 1971.
- [16] D. S. Fisher, Critical behavior of random transverse-field Ising spin chains, Phys. Rev. B 51 (1995), 6411-6461.
- [17] D. S. Fisher, P. Le Doussal and C. Monthus, Nonequilibrium dynamics of random field Ising spin chains: exact results via real space renormalization group, Phys. Rev. E 64 (2001), 066107, 41 pp..
- [18] G. Genovese, G.Giacomin and R. L. Greenblatt, Singular behavior of the leading Lyapunov exponent of a product of random 2×2 matrices, Commun. Math. Phys. **351** (2017), 923-958.
- [19] G. Giacomin and R. L. Greenblatt, Lyapunov exponent for products of random Ising transfer matrices: the balanced disorder case, ALEA 19 (2022), 701-728.
- [20] C.M. Goldie and R. Grübel, *Perpetuities with thin tails*, Adv. Appl. Prob. 28 (1996), 463-480.
- [21] A. Gut, Stopped random walks. Limit theorems and applications. 2nd ed. Springer Series in Operations Research and Financial Engineering. (2009) New York, NY, Springer. F. Iglói and C. Monthus, *Strong disorder RG approach – a short review of recent developments*, Eur. Phys. J. B 91, (2018) article number 290.
- [22] H. Kesten, Random difference equations and renewal theory for products of random matrices, Acta Math. 131 (1973), 207-248.
- [23] J. M. Luck, Sysèmes désordonnés unidimensionnels, Publications CEA Saclay, 1992.
- [24] B. M. McCoy and T. T. Wu, Theory of a two-dimensional Ising model with random impurities. I. Thermodynamics. Phys. Rev. 176 (1968), 631-643.
- [25] B. M. McCoy and T. T. Wu, *The Two-Dimensional Ising Model*. Harvard University Press, 1973.
- [26] S.P. Meyn and R.L. Tweedie, MCs and stochastic stability, Second edition, Cambridge University Press, 2009.
- [27] Q. Moulard, Absence of Griffiths singularities for smoothly disordered statistical mechanics systems, Master thesis, Univeité de Saclay, 2023.
- [28] T. M. Nieuwenhuizen and J.-M. Luck, Exactly soluble random field Ising models in one dimension, J. Phys. A, 19 (1986), 1207-1227.
- [29] B. A. Rogozin, Asymptotic analysis of the renewal function, Theory Probab. Appl. 21 (1976), 669-686.
- [30] D. Ruelle, Analycity properties of the characteristic exponents of random matrix products. Adv. Math 32 (1979), 68-80.
- [31] C. Stone, On Absolutely Continuous Components and Renewal Theory. Annals Math. Stat. 37 (1966), 271-275.

(OC) UNIVERSITÉ PARIS CITÉ, LABORATOIRE DE PROBABILITÉS, STATISTIQUES ET MODÉLISATION, UMR 8001, F-75205 Paris, France

(GG) Universita di Padova, Dipartimento di Matematica "Tullio Levi-Civita", Via Trieste ` 63, I-35121 Padova, Italy

(RG) Universita di Roma "Tor Vergata", Dipartimento di Matematica, Rome, Italy `

(YH) UNIVERSITÉ PARIS XIII, LAGA, INSTITUT GALILÉE, F-93430 VILLETANEUSE, FRANCE