Physical characteristics and causality of cosmological models in generalized matter-geometry coupling gravity theory with observational constraints

Dinesh Chandra Maurya^{1,a}, Javlon Rayimbaev^{2,b,c,d,e}, Inomjon Ibragimov^{3,f}, Sokhibjan Muminov^{4,g}

^aCentre for Cosmology, Astrophysics and Space Science, GLA University, Mathura-281 406, Uttar Pradesh, India.

^bInstitute of Fundamental and Applied Research, National Research University TIIAME, Kori Niyoziy 39, Tashkent 100000, Uzbekistan.

^cUniversity of Tashkent for Applied Sciences, Str. Gavhar 1, Tashkent 100149, Uzbekistan.

^dUrgench State University, Kh. Alimjan Str. 14, Urgench 221100, Uzbekistan.

^eShahrisabz State Pedagogical Institute, Shahrisabz Str. 10, Shahrisabz 181301, Uzbekistan.

^{*t*}Kimyo International University in Tashkent, Shota Rustaveli street 156, Tashkent 100121, Uzbekistan.

^gMamun university, Bolkhovuz Street 2, Khiva 220900, Uzbekistan.

¹E-mail:dcmaurya563@gmail.com

²E-mail:javlon@astrin.uz

³E-mail:i.ibragimov@kiut.uz

⁴E-mail:sokhibjan.muminov@mamunedu.uz

Abstract

In the generalized matter-geometry coupling theory, we investigate the physical characteristics and causality of some new cosmological models for a flat, homogeneous, and isotropic spacetime filled with stiff, radiation, dust, and curvature fluid sources. We obtain a particular cosmological model corresponding to each source fluid, called Models I, II, III, and IV, respectively. We make observational constraints on each model using the joint analysis of 31 Cosmic Chronometer (CC) Hubble dataset and 1048 Pantheon datasets to estimate the current values of model parameters. Using these statistical results, we have analyzed the information criteria, effective EoS parameter, causality of the models, and viability of this generalized gravity theory. Subsequently, we investigate the effective equation of state and deceleration parameter for each model. We found that all models in the late-time universe exhibit transit-phase acceleration, and Models I and II show both the early as well as late-time accelerating phase of the expanding universe. We found the current values of the deceleration parameter in the range $-0.886 \leq q_0 \leq -0.54$ with transition redshift $0.5137 \leq z_t \leq 0.6466$ and the effective EoS parameter in the range $-0.924 \leq \omega_{eff} \leq -0.6933$. We analyzed the square sound speed condition $c_s^2 \leq c^2$ for each model.

Keywords: generalized matter-geometry coupling gravity; causality; accelerating universe; equation of state; observational constraints.

PACS number: 98.80-k, 98.80.Jk, 04.50.Kd

1 Introduction

The discovery of the accelerating phase of the expanding universe [1–5], the cosmological constant problem [6–11], and the fundamental laws of physics force to modify Einstein's general theory of gravity (GR). Although, several modifications to Einstein's general relativity theory of gravity (GR) have been introduced in the literature, but we will only discuss some of these modifications within the Riemannian framework here, since, we are interested in investigating the generalized matter-geometry coupling gravity theory. In 1970, Buchdahl [12] suggested a wider version of GR theory by changing the Lagrangian L = R to L = f(R) in the Einstein-Hilbert action. He focused on an open universe model that changed between non-singular states, and this idea was later studied in [13–16]. Recently, cosmologists found that the universe is expanding faster than they thought. They use this theory to explain this and to solve problems with the cosmological constant [17–31]. The modified f(R)-gravity theory presents initial cosmological models that include both cosmic phenomena, inflation, and cosmic speed-up in the expansion of the universe. Recently, we have investigated several cosmological models in the context of a Riemannian framework to explain the accelerating scenarios and evolution of expanding universe properties in [32–40].

The extended f(R)-gravity theories are introduced in [41-43] which create a relationship between matter and curvature by including them externally using the expression $S = \int [f_1(R) + (1 + \lambda f_2(R))L_m]\sqrt{-g}d^4x$. The extra relationship between matter and curvature results in additional forces. This theory is later improved by Harko [44], and this time, he proposed the matter-geometry relationship with action $S = \int [\frac{1}{2}f_1(R) + G(L_m)f_2(R)]\sqrt{-g}d^4x$. After that, this idea was developed into the $f(R, L_m)$ gravity theory by Harko and Lobo [45]. In this improved form of matter-geometry coupling theory, the Lagrangian function f is an arbitrary function of the Ricci curvature scalar R and the matter Lagrangian L_m . This matter-geometry coupling theory is used by several cosmologists [46–57] to explore the late-time cosmic accelerating scenarios and observable universe physical phenomena in recent literature. We have, also recently investigated transit cosmological models in the $f(R, L_m)$ gravity theory, using observational constraints in [58–62].

Subsequently, a further study [63] approached matter-geometry coupling in a different form called f(R,T) gravity theory. At this time, the f is an arbitrary function of the Ricci scalar R and trace T of the energy-momentum tensor T_{ij} . Furthermore, this geometry creates a non-minimal correlation between the trace of the matter-energy-momentum tensor. In [64], a thermodynamic interpretation of matter-geometry coupling models is given, and [65] investigated cosmic evolution of f(R,T) gravity models with collisional matter. [66] has presented a scalar tensor approach of f(R,T) gravity models. Stability of f(R,T) gravity models is investigated in [67], and [68] has studied the in-viability of f(R,T) gravity theory. In [69–71], some more cosmological implications and late-time cosmic acceleration in f(R,T) gravity models are investigated. Recently, we have investigated the late-time scenarios of the cosmological models in f(R,T) gravity [72–76]. Particle creation [77] and variable Chaplygin gas [78] models are investigated in f(R,T) gravity theory.

Further unification of f(R), $f(R, L_m)$, and f(R, T) gravity theories is proposed by Haghani and Harko [79] in the generalized form of matter-geometry coupling, which is referred to as $f(R, L_m, T)$ gravity. The main objective of developing this theory is to integrate modified gravity theories in Riemannian geometry, including the f(R), $f(R, L_m)$, and f(R, T) gravity theories. Although theories such as f(R)-gravity, f(T)-gravity, and f(Q)-gravity have their own advantages in describing the universe's history, our specific focus is on the more comprehensive and less investigated theory of $f(R, L_m, T)$ gravity. This theory guarantees that the modification is substantial for all types of matter fields. Physical factors are the reason for including this background in the examination of cosmological theories. Recently, [80] has discussed the energy conditions in $f(R, L_m, T)$ gravity theory, and [81] investigated dusty universe in $f(R, L_m, T)$ gravity to achieve accelerating cosmological scenarios using observational constraints. The causality and its violation in $f(R, L_m, T)$ gravity theory is discussed in [82]. Recently, we have developed some constrained accelerating cosmological models and its stability analysis in [83,84]. Here, we investigate physical characteristics and causality of the expanding universe in this unified $f(R, L_m, T)$ gravity in the presence of perfect fluid with constant equation of state (EoS) parameter. We use the flat, homogeneous, and isotropic Friedmann-Lemaître-Robertson-Walker (FLRW) spacetime universe because, in this case, we are not to fix the properties openness or closedness of the universe [85].

We organize the present research paper into six sections. A brief introduction of the development of generalized matter-geometry coupling theory is given in Sect. 1. The field equations in this generalized gravity theory are derived in Sect. 2. We obtain the cosmological solutions in Sect. 3, while the observational constraints made on the models are in Sect. 4. The result discussions with causality analysis are in Sect. 5, and in the last Sect. 6, concluding remarks are given.

2 Field equations in $f(R, L_m, T)$ gravity theory

We consider the action for generalized matter-geometry coupling $f(R, L_m, T)$ gravity theory in the following form [79]

$$I = \frac{1}{16\pi} \int f(R, L_m, T) \sqrt{-g} d^4 x + \int L_m \sqrt{-g} d^4 x,$$
 (1)

where $f(R, L_m, T)$ is an arbitrary function of the Ricci scalar, R, the trace T of the stress-energy momentum tensor of the matter, T_{ij} , and of the matter Lagrangian density L_m . We define the stress-energy momentum tensor of the matter as [86]

$$T_{ij} = -\frac{2}{\sqrt{-g}} \frac{\delta(\sqrt{-g}L_m)}{\delta g^{ij}},\tag{2}$$

and its trace by $T = g^{ij}T_{ij}$, respectively. By assuming that the Lagrangian density L_m of matter depends only on the metric tensor components g_{ij} , and not on its derivatives, we obtain

$$T_{ij} = g_{ij}L_m - 2\frac{\partial L_m}{\partial g^{ij}}.$$
(3)

By varying the action I of the gravitational field with respect to the metric tensor components g^{ij} provides the following relationship

$$\delta I = \int \left[f_R \delta R + f_T \frac{\delta T}{\delta g^{ij}} \delta g^{ij} + f_{L_m} \frac{\delta L_m}{\delta g^{ij}} \delta g^{ij} - \frac{1}{2} g_{ij} f \delta g^{ij} - 8\pi T_{ij} \delta g^{ij} \right] \sqrt{-g} d^4 x, \tag{4}$$

where we have denoted $f_R = \partial f / \partial R$, $f_T = \partial f / \partial T$ and $f_{L_m} = \partial f / \partial L_m$, respectively. For the variation of the Ricci scalar, we obtain

$$\delta R = \delta(g^{ij}R_{ij}) = R_{ij}\delta g^{ij} + g^{ij}(\nabla_\lambda\delta\Gamma^\lambda_{ij} - \nabla_j\delta\Gamma^\lambda_{i\lambda}),\tag{5}$$

where ∇_{λ} is the covariant derivative with respect to the symmetric connection Γ associated to the metric g. The variation of the Christoffel symbols yields

$$\delta\Gamma_{ij}^{\lambda} = \frac{1}{2}g^{\lambda\alpha}(\nabla_i\delta g_{j\alpha} + \nabla_j\delta g_{\alpha i} - \nabla_\alpha\delta g_{ij}),\tag{6}$$

and the variation of the Ricci scalar provides the expression

$$\delta R = R_{ij} \delta g^{ij} + g_{ij} \Box \delta g^{ij} - \nabla_i \nabla_j \delta g^{ij}.$$
(7)

Therefore, for the variation of the action of the gravitational field, we obtain

$$\delta R = \int \left[f_R R_{ij} + (g_{ij} \Box - \nabla_i \nabla_j) f_R + f_T \frac{\delta(g^{\alpha\beta} T_{\alpha\beta})}{\delta g^{ij}} + f_{L_m} \frac{\delta L_m}{\delta g^{ij}} - \frac{1}{2} g_{ij} f - 8\pi T_{ij} \right] \delta g^{ij} \sqrt{-g} d^4 x.$$
(8)

We define the variation of T with respect to the metric tensor as

$$\frac{\delta(g^{\alpha\beta}T_{\alpha\beta})}{\delta g^{ij}} = T_{ij} + \Theta_{ij},\tag{9}$$

where

$$\Theta_{ij} = g^{\alpha\beta} \frac{\delta T_{\alpha\beta}}{\delta g^{ij}} = L_m g_{ij} - 2T_{ij}, \tag{10}$$

for the perfect-fluid matter source.

Taking $\delta I = 0$, we obtain the field equations of $f(R, L_m, T)$ gravity model as

$$f_R R_{ij} - \frac{1}{2} [f - (f_{L_m} + 2f_T) L_m] g_{ij} + (g_{ij} \Box - \nabla_i \nabla_j) f_R = \left[8\pi + \frac{1}{2} (f_{L_m} + 2f_T) \right] T_{ij}.$$
 (11)

By contracting Eq. (11) gives the following relation between the Ricci scalar R and the trace T of the stress-energy tensor,

$$f_R R - 2[f - (f_{L_m} + 2f_T)L_m] + 3\Box f_R = \left\lfloor 8\pi + \frac{1}{2}(f_{L_m} + 2f_T) \right\rfloor T,$$
(12)

where we have denoted $\Theta = \Theta_i^i$.

The problem of the perfect fluids, described by an energy density ρ , pressure p and four-velocity u^i is more complicated, since there is no unique definition of the matter Lagrangian. However, in the present study we assume that the stress-energy tensor of the matter is given by

$$T_{ij} = (\rho + p)u_i u_j + pg_{ij},\tag{13}$$

for the flat FLRW homogeneous and isotropic spacetime metric

$$ds^{2} = -dt^{2} + a(t)^{2}(dx^{2} + dy^{2} + dz^{2}),$$
(14)

and the matter Lagrangian can be taken as $L_m = -\rho$. The four-velocity u_i satisfies the conditions $u_i u^i = -1$ and $u^i \nabla_j u_i = 0$, respectively.

The energy-conservation equation is obtained as

$$\dot{\rho} + 3H(\rho + p) = \frac{(\rho + L_m)(\dot{f}_T + \frac{1}{2}\dot{f}_{L_m}) - \frac{1}{2}f_T(\dot{T} - 2\dot{L}_m)}{8\pi + f_T + \frac{1}{2}f_{L_m}}.$$
(15)

or

$$\dot{\rho} + 3H(\rho + p) = D_{\mu}u^{\mu} \tag{16}$$

The source term $D_{\mu}u^{\mu}$ represents the energy production or destruction within the system. The conservation of total energy in a gravitational system is attained alone when the condition $D_{\mu}u^{\mu} = 0$ is fulfilled at every point in spacetime. If $D_{\mu}u^{\mu} \neq 0$, an energy transfer process or particle formation occurs within the system. Comparing Eqs. (15) and (16), we obtain

$$D_{\mu}u^{\mu} = \frac{(\rho + L_m)(\dot{f}_T + \frac{1}{2}\dot{f}_{L_m}) - \frac{1}{2}f_T(\dot{T} - 2\dot{L}_m)}{8\pi + f_T + \frac{1}{2}f_{L_m}}.$$
(17)

Equations (16) and (17) indicate that the source term $D_{\mu}u^{\mu} \neq 0$, signifying the formation or destruction of matter within the system. In the present study, we assume $D_{\mu}u^{\mu} = 0$ and solve the field equations for different fluid sources which satisfies this condition.

3 Cosmological solutions

To investigate the cosmological properties of the above-proposed modified gravity, we consider the following form of the matter-geometry coupling function $f(R, L_m, T)$ with matter Lagrangian $L_m = -\rho$, (as suggested in [79]),

$$f(R, L_m, T) = R + \mu T L_m - \nu,$$
 (18)

where μ and ν are the arbitrary constants.

Then

$$f_R = 1, \quad f_T = \mu L_m, \quad f_{L_m} = \mu T.$$
 (19)

By applying (18) in (11), we get the following field equations:

$$R_{ij} - \frac{1}{2}Rg_{ij} = (8\pi + \frac{\mu}{2}T + \mu L_m)T_{ij} - (\mu L_m^2 + \frac{\nu}{2})g_{ij}.$$
(20)

Now, using (13) and (14) in (20), we obtain the following equations:

$$3H^2 = 8\pi\rho - \frac{\mu}{2}\rho^2 + \frac{3\mu}{2}\rho p + \frac{\nu}{2},\tag{21}$$

$$2\dot{H} + 3H^2 = -8\pi p + \frac{3\mu}{2}(\rho - p)p + \mu\rho^2 + \frac{\nu}{2}.$$
(22)

From Eqs. (15) and (18), we obtain the equation of continuity as follows

$$\dot{\rho} + 3H(\rho + p) = \frac{\mu\rho(\dot{\rho} + 3\dot{p})}{16\pi - 3\mu(\rho - p)}.$$
(23)

Since the $f(R, L_m, T)$ theory is a non-conservative theory and hence, generally, it does not satisfy the energy conservation equation i.e., $D_{\mu}u^{\mu} \neq 0$. Therefore, we consider the conserved matter fluid sources that satisfies the continuity equations $\dot{\rho} + 3H(1 + \omega_i)\rho = 0$ for i = s, r, m, c, which stand for stiffness, radiation, dust, and curvature fluids, respectively. Now, we solve the highly nonlinear Eqs. (21) and (22) for different phase of perfect fluid $(p = \omega_i \rho)$ as stiff, radiation, dust and curvature in four subsections as follows:

3.1 Model-I

To investigate the physical characteristics and accelerating scenarios in the context of barotropic fluid $\omega \ge 0$, we consider the perfect fluid source as stiff fluid ($\omega = 1$) for which the above Eqs. (21) and (22) reduced to

$$3H^2 = 8\pi\rho + \mu\rho^2 + \frac{\nu}{2},$$
(24)

$$2\dot{H} + 3H^2 = -8\pi\rho + \mu\rho^2 + \frac{\nu}{2}.$$
(25)

The energy density for stiff fluid is obtained after solving the continuity equation $\dot{\rho} + 3H(1 + \omega_s)\rho = 0$, as

$$\rho = \rho_{s0} (1+z)^6, \tag{26}$$

where ρ_{s0} is the current value of energy density and $1 + z = \frac{a_0}{a}$ as given in [87]. Now, we derive the Hubble function from Eqs. (24) and (26) as

$$H(z) = H_0 \sqrt{\Omega_{s0}(1+z)^6 + \Omega_{\mu 0}(1+z)^{12} + \Omega_{\nu 0}},$$
(27)

where $\Omega_{s0} = \frac{8\pi\rho_{s0}}{3H_0^2}$, $\Omega_{\mu0} = \frac{\mu\rho_{s0}^2}{3H_0^2}$ and $\Omega_{\nu0} = \frac{\nu}{6H_0^2}$. At z = 0, $\Omega_{s0} + \Omega_{\mu0} + \Omega_{\nu0} = 1$. The physical behaviour of model is characterized by the equation of state parameter ω , and hence, we derive the effective equation of state (EoS) parameter ω_{eff} is from Eqs. (24) and (25) as

$$\omega_{eff} = -1 + \frac{2\Omega_{s0}(1+z)^6}{\Omega_{s0}(1+z)^6 + \Omega_{\mu0}(1+z)^{12} + \Omega_{\nu0}}.$$
(28)

The phase of the expanding universe is characterized by the deceleration parameter q(z), and the deceleration parameter q(z) is obtained as

$$q(z) = \frac{1}{2} + \frac{3}{2} \frac{\Omega_{s0}(1+z)^6 - \Omega_{\mu0}(1+z)^{12} - \Omega_{\nu0}}{\Omega_{s0}(1+z)^6 + \Omega_{\mu0}(1+z)^{12} + \Omega_{\nu0}}.$$
(29)

3.2 Model-II

This model is derived for the perfect fluid as radiation fluid ($\omega = \frac{1}{3}$) and $p = \frac{1}{3}\rho$, to get the characteristics of the model under radiation fluid sources, and hence, the above Eqs. (21) and (22) reduced to

$$3H^2 = 8\pi\rho + \frac{\nu}{2},\tag{30}$$

$$2\dot{H} + 3H^2 = -\frac{8\pi}{3}\rho + \frac{4\mu}{3}\rho^2 + \frac{\nu}{2}.$$
(31)

The energy density for radiation fluid source is obtained after solving the continuity equation $\dot{\rho} + 3H(1+\omega_r)\rho = 0$ as given by

$$\rho = \rho_{r0} (1+z)^4, \tag{32}$$

where ρ_{r0} is the current value of the energy density.

We derive the Hubble function from the Eqs. (30) and (32) as

$$H(z) = H_0 \sqrt{\Omega_{r0}(1+z)^4 + \Omega_{\nu 0}},$$
(33)

where $\Omega_{r0} = \frac{8\pi\rho_{r0}}{3H_0^2}$ and $\Omega_{\nu 0} = \frac{\nu}{6H_0^2}$. At z = 0, $\Omega_{r0} + \Omega_{\nu 0} = 1$. The effective EoS parameter ω_{eff} for this model, is derived as

$$\omega_{eff} = -1 + \frac{4}{3} \frac{\Omega_{r0} (1+z)^4 - \Omega_{\mu 0} (1+z)^8}{\Omega_{r0} (1+z)^4 + \Omega_{\nu 0}},\tag{34}$$

where $\Omega_{\mu 0} = \frac{\mu \rho_{r0}^2}{3H_0^2}$. The deceleration parameter is obtained as

$$q(z) = \frac{1}{2} + \frac{1}{2} \frac{\Omega_{r0}(1+z)^4 - 4\Omega_{\mu0}(1+z)^8 - \Omega_{\nu0}}{\Omega_{r0}(1+z)^4 + \Omega_{\nu0}}.$$
(35)

3.3 Model-III

In this model, we consider the perfect fluid source as dust ($\omega = 0$) and p = 0 to get the characteristics of the cosmological models in this modified generalized theory of gravity, and hence, the above Eqs. (21) and (22) reduced to

$$3H^2 = 8\pi\rho - \frac{\mu}{2}\rho^2 + \frac{\nu}{2},\tag{36}$$

$$2\dot{H} + 3H^2 = \mu\rho^2 + \frac{\nu}{2}.$$
(37)

The energy density for dusty fluid source is obtained from the equation of continuity $\dot{\rho} + 3H(1 + \omega_m)\rho = 0$ as given by

$$\rho = \rho_{m0} (1+z)^3, \tag{38}$$

where ρ_{m0} is the current value of energy density. The Hubble function is derived from Eqs. (36) and (38) as

$$H(z) = H_0 \sqrt{\Omega_{m0}(1+z)^3 + \Omega_{\mu 0}(1+z)^6 + \Omega_{\nu 0}},$$
(39)

where $\Omega_{m0} = \frac{8\pi\rho_{m0}}{3H_0^2}$, $\Omega_{\mu0} = -\frac{\mu\rho_{m0}^2}{6H_0^2}$ and $\Omega_{\nu0} = \frac{\nu}{6H_0^2}$. At z = 0, $\Omega_{m0} + \Omega_{\mu0} + \Omega_{\nu0} = 1$. The effective EoS parameter ω_{eff} for this model is derived as

$$\omega_{eff} = -1 + \frac{\Omega_{m0}(1+z)^3 + 3\Omega_{\mu0}(1+z)^6}{\Omega_{m0}(1+z)^3 + \Omega_{\mu0}(1+z)^6 + \Omega_{\nu0}}.$$
(40)

The deceleration parameter is obtained as

$$q(z) = \frac{1}{2} + \frac{3}{2} \frac{2\Omega_{\mu 0} (1+z)^6 - \Omega_{\nu 0}}{\Omega_{m 0} (1+z)^3 + \Omega_{\mu 0} (1+z)^6 + \Omega_{\nu 0}}.$$
(41)

3.4 Model-IV

The present model is derived for perfect fluid source as curvature ($\omega = -\frac{1}{3}$) and $p = -\frac{1}{3}\rho$, to investigate the cosmological properties of the universe, and hence, the Eqs. (21) and (22) reduced to

$$3H^2 = 8\pi\rho - \mu\rho^2 + \frac{\nu}{2},\tag{42}$$

$$2\dot{H} + 3H^2 = \frac{8\pi}{3}\rho + \frac{\mu}{3}\rho^2 + \frac{\nu}{2}.$$
(43)

The equation of continuity is obtained as

$$\dot{\rho} + 2H\rho = 0. \tag{44}$$

Hence, the energy density for curvature is obtained as

$$\rho = \rho_{c0}(1+z)^2, \tag{45}$$

where ρ_{c0} is the current value of energy density

We derive the Hubble function from Eqs. (42) and (45) as

$$H(z) = H_0 \sqrt{\Omega_{c0}(1+z)^2 - \Omega_{\mu 0}(1+z)^4 + \Omega_{\nu 0}},$$
(46)

where $\Omega_{c0} = \frac{8\pi\rho_{c0}}{3H_0^2}$, $\Omega_{\mu0} = \frac{\mu\rho_{c0}^2}{3H_0^2}$ and $\Omega_{\nu0} = \frac{\nu}{6H_0^2}$. At z = 0, $\Omega_{c0} - \Omega_{\mu0} + \Omega_{\nu0} = 1$. The effective EoS parameter ω_{eff} for this model is derived as

$$\omega_{eff} = -1 + \frac{2}{3} \frac{\Omega_{c0}(1+z)^2 - 4\Omega_{\mu0}(1+z)^4}{\Omega_{c0}(1+z)^2 - \Omega_{\mu0}(1+z)^4 + \Omega_{\nu0}}.$$
(47)

The deceleration parameter is obtained as

$$q(z) = \frac{1}{2} - \frac{1}{2} \frac{\Omega_{c0}(1+z)^2 + \Omega_{\mu 0}(1+z)^4 + 3\Omega_{\nu 0}}{\Omega_{c0}(1+z)^2 - \Omega_{\mu 0}(1+z)^4 + \Omega_{\nu 0}}.$$
(48)

4 Observational Constraints

This section aims to constrain the free model parameters of our derived models (27), (33), (39), and (46) utilizing accessible datasets concerning the evolution of the expanding universe. We utilize 31 non-correlated cosmic chronometer (CC) Hubble data points [88,89], obtained by the differential age technique, along with 1048 apparent magnitude m(z) data points from the Pantheon sample of SNe Ia [90]. We examine the observational datasets employing the conventional Bayesian methodology and utilize the Monte Carlo Markov Chain (MCMC) technique to ascertain the posterior variations of the parameters. We also employ the emcee software for MCMC analysis, which is available at no cost [91]. Moreover, the probability function $\mathcal{L} \propto e^{-\chi^2/2}$ yields optimal values for the free parameters by maximizing \mathcal{L} or reducing the χ^2 -value.

4.1 Hubble datasets

We commence with the 31 Hubble data points of the Hubble parameter H(z) associated with different redshifts z from the cosmic chronometer (CC) sample. The CC dataset of H(z) is derived using the differential age method. The Hubble parameter may directly signify the rate of cosmic expansion in the universe. The decision to employ CC data is mostly contingent upon age difference measurements between two passively evolving galaxies that arose simultaneously, characterized by a small redshift interval. This interval facilitates the calculation of $\delta z/\delta t$. CC data has demonstrated greater reliability than previous methods reliant on absolute age determinations for galaxies [92]. To assess the model's goodness of fit with the data, we utilize the following χ^2 formula defined as:

$$\chi^2_{CC} = \sum_{i=1}^{31} \frac{[H_{ob}(z_i) - H_{th}(\phi, z_i)]^2}{\sigma^2_{H(z_i)}}.$$
(49)

where H_{ob} refers to the observed values, while H_{th} represents the theoretical values of the Hubble function. $\sigma_{H(z_i)}$ denotes the standard deviations associated with the Hubble data points and ϕ denote the parameter space.

4.2 Pantheon datasets

We utilize the 1048 apparent magnitude $m(z_i)$ data points at different redshifts z_i from the Pantheon sample of SNe Ia. The importance of supernovae is in their capacity to elucidate the history of the expanding cosmos and reveal the characteristics of dark energy. The SuperNova Legacy Survey (SNLS), Sloan Digital Sky Survey (SDSS), Hubble Space Telescope (HST) survey, and Panoramic Survey Telescope and Rapid Response System (Pan-STARRS1) all acquire spectroscopic data on supernovae that supports this conclusion. The Pantheon dataset, the latest sample of supernova data, comprises 1048 apparent magnitudes m_b recorded across the interval $0.01 \leq z \leq 2.26$ for the redshift z [90]. We take the theoretical definition of apparent magnitude, as delineated in [90–93]

$$m(z) = M + 5 \log_{10}\left(\frac{D_L}{Mpc}\right) + 25.$$
 (50)

where M represents the absolute magnitude, and the luminosity distance D_L is quantified in terms of length and is defined as

$$D_L = c(1+z) \int_0^z \frac{dz'}{H(z')}.$$
(51)

We define the Hubble-free luminosity distance d_L as $d_L \equiv \frac{H_0}{c}D_L$, a dimensionless quantity. Consequently, the apparent magnitude m(z) expressed as

$$m(z) = M + 5\log_{10} d_L + 5\log_{10} \left(\frac{c/H_0}{Mpc}\right) + 25.$$
(52)

The degeneracy between H_0 and M in the aforementioned equation, remains constant within the Λ CDM framework [90,93]. We shall consolidate these degenerate parameters by redefining them as follows:

$$\mathcal{M} \equiv M + 5\log_{10}\left(\frac{c/H_0}{Mpc}\right) + 25.$$
(53)

Here, \mathcal{M} denotes a dimensionless parameter, may also be written as $\mathcal{M} = M - 5 \log_{10}(h) + 42.39$, where $H_0 = h \times 100 \text{ km/s/Mpc}$. It is worthwhile to note that in most instances [94,95], this degenerate combination is marginalized; however, recent studies [96–100] suggests that this process may result in the loss of physical information, as a physical model exhibiting an abrupt transition in the absolute magnitude M at a low-redshift z_t has the capacity to simultaneously address the H_0 and growth tensions [101–106]. Consequently, we opt to employ this degenerate parameter in the estimation procedure. In the context of Λ CDM model the value of this parameter is $\mathcal{M} = 23.809 \pm 0.011$ [90]. Note that \mathcal{M} changes in the different cosmological models (see [90,96–110]). We employ the subsequent χ^2 formula for the analysis of the Pantheon data, as referenced in [90]:

$$\chi_P^2 = V_P^i C_{ij}^{-1} V_P^j \tag{54}$$

The formula V_P^i represents the discrepancy between the observed $m_{ob}(z_i)$ and the theoretical $m(\phi, \mathcal{M}, z_i)$ as outlined in equation (52), while C_{ij}^{-1} signifies the inverse of the covariance matrix of the Pantheon sample. In order to get the joint estimates on model parameters, we use the 31 CC data points Hubble parameter with 1048 Pantheon datasets. We use the following χ^2_{CC+P} formula to perform joint MCMC analysis of Pantheon and CC data points to get combined constraints on the model parameters of all derived models.

$$\chi^2_{CC+P} = \chi^2_{CC} + \chi^2_P \tag{55}$$

Model	Parameter	Prior	Value
	H_0	(50, 100)	66.2 ± 2.2
Ι	Ω_{s0}	(0, 0.05)	$0.0380^{+0.0033}_{-0.0038}$
	$\Omega_{\mu 0}$	Fixed	0.0000001
	\mathcal{M}	—	23.7302 ± 0.0077
	χ^2	—	1140.1806
	H_0	(50, 100)	67.8 ± 1.9
II	Ω_{r0}	(0, 0.3)	0.160 ± 0.012
	\mathcal{M}	—	23.7875 ± 0.0094
	χ^2	—	1049.2001
	H_0	(50, 100)	$68.4^{+1.4}_{-1.6}$
III	Ω_{m0}	(0, 0.6)	0.29 ± 0.02
	$\Omega_{\mu 0}$	(0, 0.005)	0.0024 ± 0.0013
	\mathcal{M}	—	23.8087 ± 0.0092
	χ^2	—	1041.1368
	H_0	(50, 100)	68.6 ± 1.9
IV	Ω_{c0}	(0, 1)	0.29 ± 0.10
	$\Omega_{\mu 0}$	(-0.2, 0.01)	-0.085 ± 0.027
	\mathcal{M}	—	23.810 ± 0.012
	χ^2	—	1041.2018
	H_0	(50, 100)	68.9 ± 1.8
ΛCDM	Ω_{m0}	(0, 1)	0.303 ± 0.020
	\mathcal{M}	—	23.811 ± 0.010
	χ^2	—	1041.2708

Table 1: The MCMC estimates with errors in different models.




Figure 1: The contour plots of H_0 , Ω_{s0} and \mathcal{M} and the contour plots of H_0 , Ω_{r0} and \mathcal{M} , respectively, for joint analysis of CC+Pantheon.




Figure 2: The contour plots of H_0 , Ω_{m0} , $\Omega_{\mu0}$ and \mathcal{M} and the contour plots of H_0 , Ω_{c0} , $\Omega_{\mu0}$ and \mathcal{M} , respectively, for joint analysis of CC+Pantheon.




Figure 3: The contour plots of H_0 , Ω_{m0} and \mathcal{M} for Λ CDM model for joint analysis of CC+Pantheon.

Model	$\rho_{i0} \ ({\rm in} \ gm/cm^3)$	μ (in s^2)	$\nu (\text{in } s^{-2})$
Ι	2.0867×10^{-38}	3.1694×10^{33}	2.6554×10^{-35}
II	9.2161×10^{-38}	1.7043×10^{32}	2.4320×10^{-35}
III	1.7001×10^{-37}	-6.1170×10^{35}	2.0851×10^{-35}
IV	1.7100×10^{-37}	-4.3076×10^{37}	1.8525×10^{-35}

Table 2: The values of other model parameters for CC+Pantheon datasets.

Utilizing the minimum χ^2 value (maximizing the likelihood function), we conduct MCMC analysis on the CC + Pantheon datasets for Models I, II, III, and IV to estimate the model parameters associated with their respective Hubble functions. The contour plots of the estimated parameters at $1 - \sigma$ and $2 - \sigma$ confidence levels are illustrated in Figures 1a, 1b, 2a, and 2b, while the MCMC analysis for the Λ CDM is depicted in Figure 3. The estimated values of the model parameters are presented in Table 1. Utilizing the estimated values of Table 1, we assessed the approximated values of other model parameters, which are shown in Table 2.

4.3 Information Criteria

In this part, we employ the Akaike Information Criterion (AIC) and the Bayesian Information Criterion (BIC) as model selection criteria to statistically compare our derived models fitting with the standard cosmological Λ CDM model. The AIC criteria can be computed using the following formula, as given in references [111,112], under the assumption of Gaussian errors.

$$AIC = \chi^2_{\text{tot,min}} + \frac{2kN_{\text{tot}}}{N_{\text{tot}} - k - 1},\tag{56}$$

The symbol $\chi^2_{\text{tot,min}}$ reflects the minimum value of the χ^2_{CC+P} for joint analysis of CC+Pantheon datasets being studied. The variable N_{tot} represents the total number of data points utilized in the analysis, whereas k indicates the number of parameters that have been fitted in a particular model. Minimizing the χ^2 -value is comparable to maximizing the likelihood function \mathcal{L} . The BIC criteria can be computed using the formula specified in references [111–113].

$$BIC = \chi^2_{\text{tot,min}} + k \ln N_{\text{tot}}$$
(57)

Our objective is to categorize the models based on their capacity to effectively match the provided data, while considering a range of scenarios that depict the same type of event. When evaluating a group of models, we detect bias in the information criteria (IC) value. To do this, we utilize two specific ICs: AIC and BIC. The expression $\Delta IC_{model} = IC_{model} - IC_{min}$ represents the difference between the IC value of the derived model (IC_{model}) and the minimal IC value corresponding to the standard model (IC_{min}). To evaluate the suitability of each model, we employ Jeffrey's scale [114]. More precisely, if the value of ΔIC is less than or equal to 2, it indicates that the data strongly supports the conventional model. When the disparity value, ΔIC , falls within the range of 2 to 6, it signifies a substantial level of disagreement between the two models. Ultimately, when the disparities reach or exceed 10, it signifies a moderate level of tension between the two models [115].

Model	$\chi^2_{ m tot,min}$	$N_{\rm tot}$	k	AIC	ΔAIC	BIC	ΔBIC
ΛCDM	1041.2708	1079	3	1047.2931	0	1062.2221	0
Model I	1140.1806	1079	3	1146.2029	98.9098	1161.1319	98.9098
Model II	1049.2001	1079	3	1055.2224	7.9293	1070.1514	7.9293
Model III	1041.1368	1079	4	1049.1740	1.8809	1069.0719	6.8498
Model IV	1041.2018	1079	4	1049.2390	1.9459	1069.1369	6.9148

Table 3: The information criteria

We found the model parameters by fitting them to the CC+Pantheon datasets to get the lowest χ^2 value. The minimum χ^2 value for each model is shown in Table 1. Here, the total number of datasets is denoted by $N_{\text{tot}} = 1079$ for the CC+Pantheon datasets, and the number of fitted parameters with CC+Pantheon datasets are k = 3, 3, 4, 4 for Models I, II, III, and IV, respectively. In the case of the standard Λ CDM model, k = 3 for CC+Pantheon datasets. Using these values in the formulas (56) and (57), we have determined the values of AIC, BIC, Δ AIC, and Δ BIC, which are presented in Table 3.

5 Result discussions

In this part, we present a comparative study of the analysis of all four derived universe models, I, II, III, and IV. The Hubble function has been determined for each model in terms of z and free parameters. Subsequently, the best-fit values of these free parameters have been estimated using MCMC analysis with joint analysis of 31 cosmic chronometer (CC) Hubble datasets and 1048 Pantheon datasets. The minimum χ^2 values and values of the fitted

parameters along with their corresponding confidence levels of $1 - \sigma$ and $2 - \sigma$, are presented in Table 1. As a result, we have obtained the AIC, Δ AIC, and BIC, Δ BIC values for each model fitting (see Table 3). We have calculated the effective equation of the state parameter, denoted as ω_{eff} , and the deceleration parameter for each model to explore the physical characteristics of the models.

Based on the information provided in Table 1, it is evident that the minimum χ^2 value for Model I is very distant, whereas for Model III, it is equivalent to the least χ^2 value for the Λ CDM. We found that the Hubble constants for Models I, II, III, and IV are $H_0 = 66.2 \pm 2.2$, 67.8 ± 1.9 , $68.4^{+1.4}_{-1.6}$, 68.6 ± 1.9 Km/s/Mpc along the joint datasets CC+Pantheon. The estimated values of some other model parameters are shown in Table 4. Furthermore, from Table 3, it becomes evident that the differences in AIC and BIC values for Model I exceed 10, indicating a mild tension between Model I and Λ CDM. The Δ AIC and Δ BIC values for Model II fall in the range of 1–2, which means that it fits the Λ CDM estimates better. For the model II, the Δ IC values for the fall in the range of 6–10, which means that there is some tension between Model II and Λ CDM. However, in Models III and IV, the values of Δ AIC and Δ BIC fall within the range (1, 2) that reveals the closeness of the model fittings with standard model, and the values of Δ BIC fall within the range (6, 10) that reveals a mild tension among the models from standard Λ CDM model. Thus, the Models II, III, and IV are substantially closed to Λ CDM standard model statistically while there is a mild tension between Model I and Λ CDM.




Figure 4: The variations of effective EoS parameter ω_{eff} versus z.

Figure 4 displays the variations of the effective equation of state parameter ω_{eff} for Models I, II, III, and IV across different redshift values over $-1 \leq z \leq 30$. The mathematical formulas for these parameters are provided in equations (28), (34), (40), and (47), respectively. Figure 4 shows that the effective equation of state (EoS) parameter for Models I and II starts out negative in the early stages of the universe. It then goes up to its highest positive value, then goes down and slowly approaches -1 as the universe gets older. On the other hand, the equation of state parameter ω_{eff} for Models III and IV initially has a positive value and approaches -1 in the later stages of the universe. The calculated present value of the effective equation of state (EoS) parameter is determined for models I, II, III, and IV, and these are shown in Table 4. One can observe that the present values of effective EoS parameter falls in the range $-0.9240 \leq \omega_{eff0} \leq -0.6933$ which are compatible with recent estimations in different investigations. From Table 4, we can see that Model I, II have higher dark energy as $0.8399 \leq \Omega_{\nu 0} \leq 0.9619$ while in Model III and IV $0.7076 \leq \Omega_{\nu 0} \leq 0.7950$ which may in good agreement with standard Λ CDM value $\Omega_{\Lambda 0} = 0.70$. Figures 4 and 5 demonstrate that models I, II, III, and IV exhibit accelerated

expansion within the specified range of the effective equation of state parameter, $-1 \leq \omega_{eff} < -\frac{1}{3}$, where model I and II show two phases of accelerating scenarios early and late-time universe. This demonstrates the viability of our generated models. The models I, II, and III, behaves just like dark energy late-time accelerating cosmological models for the barotropic fluid source $0 \leq \omega_i \leq 1$.

Model	ω_{eff0}	q_0	z_t	$\Omega_{\nu 0}$
Ι	-0.9240	-0.8860	8.5530	0.9619
			0.5266	
II	-0.7866	-0.6800	31.1370	0.8399
			0.5137	
III	-0.7028	-0.5542	0.5832	0.7076
IV	-0.6933	-0.5400	0.6466	0.7950

Table 4: The value of parameters.




Figure 5: The variations of deceleration parameter q(z) versus z.

The estimated present values of the energy density parameters Ω_{s0} , Ω_{r0} , Ω_{m0} , Ω_{c0} , and $\Omega_{\mu0}$ are shown in Table 1, while the present values of the dark energy density parameter $\Omega_{\nu0}$ are shown in Tables 4. Based on these measurements, we can see that universe models I and II, which have stiff and radiation fluid sources, have a lot more dark energy from the $f(R, L_m, T)$ function in the form of $\omega_{\nu0}$ than universe models III and IV. The second part of the Lagrangian function $f(R, L_m, T)$ makes a lot more of a difference in the last two models, III and IV, compared to the first two models, I and II. The non-zero values of $\Omega_{\mu0}$ and $\Omega_{\nu0}$ represent the significant role of modified terms in Einstein's Lagrangian. You can see the estimated values of the model parameters $\rho_{i0}(i = s, r, m, c)$, μ , and ν in Table 2.

The mathematical formulas for the deceleration parameter are given in Eqs. (29), (35), (41), and (48) for Models I, II, III, and IV, respectively. Figure 5 show how the deceleration parameter q(z) changes shape for Models I, II, III, and IV, using the estimated values of the model parameters along observational datasets. It is evident that the plot of q(z) for Models I and II exhibits a negative value at the beginning and at the end within the redshift range of $-1 \le z \le 32$ and within this range once q(z) reaches its maximum positive value. This observation indicates that Models I and II show two phases: transitioning from accelerating to decelerating (early stage) and then decelerating to accelerating (late-time scenarios) in each model. On the other hand, q(z) for Models III and IV shows transition from decelerating to accelerating phase. The estimated present values of q(z) for each model are shown in Tables 4, which are as follows: $q_0 = -0.8860, -0.68, -0.5542, -0.54$. The transition redshift z_t is measured to be $z_t = 0.5266, 0.5137, 0.5832, 0.6466$ for Models I, II, III, and IV, respectively, for late-time transition, while Models I and II show a transition in its early stages at $z_t = 8.553, 31.137$, respectively. Therefore, it can be concluded that all of the generated models are transit phase universes that experience late-time acceleration during the expansion of the universe, while Models I and II show two phases of acceleration and deceleration. The value of the transition redshift was recently estimated to be $z_t = 0.8596^{+0.2886}_{-0.2722}$ for SNIa datasets, and the value of $z_t = 0.6320^{+0.1605}_{-0.1403}$ is obtained for Hubble datasets in f(R) gravity models [116]. The research [117] presents a transition redshift of $z_t = 0.643^{+0.034}_{-0.030}$ in the framework of f(T) gravity. The transition value z_t was measured by [118] using a model-independent technique as $z_t = 0.646^{+0.020}_{-0.158}$. The transition value reported in another study by [119] is $z_t = 0.702^{+0.094}_{-0.044}$. In addition, the transition redshift value was measured by [120, 121] to be $z_t = 0.684^{+0.136}_{-0.092}$. Therefore, our estimated values of z_t for Models I, II, III, and IV align with the aforementioned values.

5.1 Causality

To analyze the stability of derived cosmological models, we investigate the square sound speed c_s^2 defined as [122]

$$c_s^2 = \frac{dp_e}{d\rho_e} \tag{58}$$

where $p_e = p_{eff}$ and $\rho_e = \rho_{eff}$. For a perfect fluid source the square speed of sound can not exceed the square velocity of light i.e., $c_s^2 \leq c^2$ [122]. Hence, for a stable cosmological model, the square speed of sound should be less than unity (for c = 1), i.e., $0 \leq c_s^2 \leq 1$. Thus, we can derive the speed of sound for effective pressure and energy density of each model using equation (58), respectively, as follows:

$$c_s^2 = \frac{\Omega_{s0} - 2\Omega_{\mu0}(1+z)^6}{\Omega_{s0} + 2\Omega_{\mu0}(1+z)^6}$$
(59)

$$c_s^2 = \frac{1}{3} - \frac{8}{3} \frac{\Omega_{\mu 0}}{\Omega_{r0}} (1+z)^4 \tag{60}$$

$$c_s^2 = \frac{4\Omega_{\mu 0}(1+z)^3}{\Omega_{m0} + 2\Omega_{\mu 0}(1+z)^3}$$
(61)

$$c_s^2 = \frac{-\Omega_{c0} - 2\Omega_{\mu0}(1+z)^2}{3\Omega_{c0} - 6\Omega_{\mu0}(1+z)^2}$$
(62)

The equations (59), (60), (61), and (62) represent the expressions of square sound speed for the models I, II, III, and IV, respectively, in terms of energy density parameters and redshift z. Figure 6 depicts the evolution of square sound speed c_s^2 versus redshift z. One can see that the models for barotropic fluid have square sound speed values within the given range $0 \le c_s^2 \le 1$, whereas model IV violates this condition in late-time stages. Therefore, the models I, II, and III are stable, but model IV is unstable at late-time.




Figure 6: The plot of square sound speed c_s^2 versus z.

5.2 $f(R, L_m, T)$ function

Within this part, we analyze the Lagrangian function $f(R, L_m, T)$. In order to accomplish this, we can derive the expression for the function $f(R, L_m, T)$ for each model separately, as follows:

$$f_I(R, L_m, T) = -16\pi \rho_{s0} (1+z)^6 - 10\mu \rho_{s0}^2 (1+z)^{12} + \nu,$$
(63)

$$f_{II}(R, L_m, T) = \nu, \tag{64}$$

$$f_{III}(R, L_m, T) = 8\pi\rho_{m0}(1+z)^3 + 2\mu\rho_{m0}^2(1+z)^6 + \nu,$$
(65)

$$f_{IV}(R, L_m, T) = 16\pi\rho_{c0}(1+z)^2 + 2\mu\rho_{c0}^2(1+z)^4 + \nu.$$
(66)

The partial derivatives of the function $f(R, L_m, T)$ with respect to L_m and T are calculated and presented in Table 5 for each model.

Models	$f_{L_m} = \mu T$	$f_T = \mu L_m$	Value of μ
Ι	$2\mu\rho$	$-\mu\rho$	$\mu > 0$
II	0	$-\mu\rho$	$\mu > 0$ or $\mu < 0$
III	$-\mu\rho$	$-\mu\rho$	$\mu < 0$
IV	$-2\mu\rho$	$-\mu\rho$	$\mu < 0$

Table 5: The derivatives of function $f(R, L_m, T)$.

By examining equations (63)-(66), it becomes evident that the function $f(R, L_m, T)$ approaches ν as z approaches -1, with ν being greater than zero. Conversely, $f(R, L_m, T)$ approaches negative infinity as z approaches infinity, regardless of the model. Furthermore, according to the information provided in Table 5, it can be observed that $f_{L_m} \geq 0$ for all models while $f_T > 0$ for models II, III, and IV but $f_T < 0$ for model I. This demonstrates the feasibility of our derived universe models.

6 Conclusions

In the generalized matter-geometry coupling theory, we investigated the physical characteristics and causality of some new cosmological models for a flat, homogeneous, and isotropic spacetime filled with stiff, radiation, dust, and curvature fluid sources. We have obtained a particular cosmological model corresponding to each source fluid, called Models I, II, III, and IV, respectively. We have made observational constraints on each model using joint analysis of the 31 CC Hubble dataset and 1048 Pantheon datasets to estimate the current values of model parameters. Using these statistical results, we have analyzed the information criteria, effective EoS parameter, causality of the models, and viability of this generalized gravity theory. We have measured the values of the Hubble constant in the range $66.2 \leq H_0 \leq 68.9$, which are compatible with recent observed values. We have found that all the derived models are transit phase-accelerating in late-time universes, while models I and II also depict the accelerating phase in their early stages. We have obtained the current values of the deceleration parameter q(z) in the range $-0.8860 \leq q_0 \leq -0.54$ with the transition redshift $0.5137 \leq z_t \leq 0.6466$ for all Models I, II, III, and IV, in their late-time stages, while Models I and II have a transition point from accelerating to decelerating in the transition redshift range $8.553 \leq z_t \leq 31.137$ (for detail see Table 4).

We observed that the effective equation of state (EoS) parameter for Models I and II initially had a negative value during the early stages of the universe, then increased to a maximum positive value, and then decreased to -1 in the later stages of the universe. On the other hand, the equation of state parameter ω_{eff} for Models III and IV initially has a positive value and approaches -1 in the later stages of the universe. We found the present value of the effective EoS parameter to be $-0.924 \leq \omega_{eff} \leq -0.6933$. We have concluded that all models exhibit an accelerated expansion phase within the specified range of the effective equation of state parameter, $-1 \leq \omega_{eff} < -\frac{1}{3}$, whereas models I and II show two phases of accelerating scenarios: early and late-time universes. The models I, II, and III behave just like dark energy late-time accelerating cosmological models for the barotropic fluid source $0 \leq \omega_i \leq 1$. We have measured the present values of $\Omega_{\mu 0}$, $\Omega_{\nu 0}$, μ , ν , and $\rho_{i0}(i = s, r, m, c)$, which are mentioned in Tables 1, 2, 4. The non-zero values of $\Omega_{\mu 0}$ and $\Omega_{\nu 0}$ represent the significant role of modified terms in Einstein's Lagrangian.

We have found that statistical analysis of joint datasets represents a mild tension between Model I and Λ CDM in their analysis of AIC and BIC (see Table 3). Furthermore, in Models II, III, and IV, the values of Δ AIC and Δ BIC fall below 6, suggesting a minor deviation from the Λ CDM model. We found the square sound speed values for Models I, II, and III within the range $0 \le c_s^2 \le 1$, which illustrates that they are in stable stages, while Model IV violates this condition of stability. For each model, the Lagrangian function $f(R, L_m, T)$ and its derivatives f_{L_m} and f_T have been computed. The positive derivatives $f_{L_m} \ge 0$ and $f_T > 0$ indicate that our derived models are feasible. In this way, the cosmological models in the generalized matter-geometry coupling gravity theory can explain the physical features of a universe that can be seen. In general, this theory doesn't meet the energy conditions, and more research is needed to see how the formation or destruction of matter takes place in this $f(R, L_m, T)$ gravity theory.

Acknowledgments

The authors are thankful to renowned and honorable reviewers and editors for their valuable suggestions to improve the quality of this paper. The author Dinesh Chandra Maurya is thankful to IUCAA - Center for Astronomy Research and Development (ICARD), CCASS, GLA University, Mathura, India for providing facilities and support where part of this work is carried out.

7 Data Availability Statement

No data associated in the manuscript.

8 Declarations

Funding and/or Conflicts of interests/Competing interests

The authors of this article have no conflict of interests. The authors have no competing interests to declare that are relevant to the content of this article. The authors did not receive support from any organization for the submitted work.

References

- A. G. Riess, A. V. Filippenko, P. Challis, et al., Observational evidence from supernovae for an accelerating universe and a cosmological constant, Astron. J. 116 (1998) 1009-1038, [arXiv:astro-ph/9805201]
- S. Perlmutter, G. Aldering, G. Goldhaber, et al., Measurements of Omega and Lambda from 42 High-Redshift Supernovae, Astrophys. J. 517 (1999) 565-586, [arXiv:astro-ph/9812133]
- [3] R. A. Knop, G. Aldering, R. Amanullah, *et al.*, New Constraints on Ω_M , Ω_Λ , and ω from an Independent Set of 11 High-Redshift Supernovae Observed with the Hubble Space Telescope, *Astrophys. J.* **598** (2003) 102, [arXiv:astro-ph/0309368]
- [4] R. Amanullah, C. Lidman, D. Rubin, et al., Spectra and Hubble Space Telescope light curves of six type Ia supernovae at 0.511 < z < 1.12 and the Union2 compilation Astrophys. J. 716 (2010) 712-738, [arXiv:1004.1711 [astro-ph.CO]]
- [5] D. H. Weinberg, M. J. Mortonson, D. J. Eisenstein, et al., Observational probes of cosmic acceleration, Phys. Rep. 530 (2013) 87-255, [arXiv:1201.2434 [astro-ph.CO]]
- [6] A. Einstein, The Principle of General Relativity, Annalen Physik Leipzig 55 (1918) 241-245.
- [7] P. Salucci, N. Turini, and C. Di Paolo, Paradigms and scenarios for the dark matter phenomenon, Universe 6 (2020) 118, [arXiv:2008.04052 [astro-ph.CO]]
- [8] S. Alam, M. Ata, S. Bailey, et al. (BOSS Collaboration), The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: cosmological analysis of the DR12 galaxy sample, Mon. Not. R. Astron. Soc. 470 (2017) 2617-2652, [arXiv:1607.03155 [astro-ph.CO]]
- [9] T. M. C. Abbott, F. B. Abdalla, A. Alarcon, et al. (DES Collaboration), Dark Energy Survey year 1 results: Cosmological constraints from galaxy clustering and weak lensing, Phys. Rev. D 98 043526 (2018), [arXiv:1708.01530 [astro-ph.CO]]
- [10] M. Tanabashi, K. Hagiwara, K. Hikasa, et al. (Particle Data Group), Review of Particle Physics: particle data groups, Phys. Rev. D 98, 030001 (2018),
- [11] N. Aghanim, Y. Akrami, M. Ashdown, et al. (Planck Collaboration), Planck 2018 results. VI. Cosmological parameters, Astron. Astrophys. 641, A6 (2020), [arXiv:1807.06209 [astro-ph.CO]]
- [12] H. A. Buchdahl, Non-linear Lagrangians and cosmological theory, Mon. Not. R. Astron. Soc. 150, 1-8 (1970),
- [13] R. Kerner, Cosmology without singularity and nonlinear gravitational Lagrangians, Gen. Relativ. Gravit. 14 453-469 (1982),
- [14] J. P. Duruisseau, R. Kerner, P. Eysseric, Non-Einsteinian gravitational Lagrangians assuring cosmological solutions without collapse, *Gen. Relativ. Gravit.* 15 797-807 (1983),
- [15] J. D. Barrow, A. C. Ottewill, The stability of general relativistic cosmological theory, J. Phys. A Math. Gen. 16, 2757 (1983),

- [16] H. Kleinert, H.-J. Schmidt, Cosmology with Curvature-Saturated Gravitational Lagrangian R, Gen. Relativ. Gravit. 34, 1295-1318 (2002), [arXiv:gr-qc/0006074]
- [17] S.M. Carroll, V. Duvvuri, M. Trodden, M.S. Turner, Is cosmic speed-up due to new gravitational physics?, *Phys. Rev. D* 70, 043528 (2004). [arXiv:astro-ph/0306438]
- [18] W. Hu, I. Sawicki, Models of f(R) cosmic acceleration that evade solar system tests, *Phys. Rev. D* **76**, 064004 (2007), [arXiv:0705.1158 [astro-ph]]
- [19] S. A. Appleby, R. A. Battye, Do consistent F(R) models mimic general relativity plus Λ ?, *Phys. Lett. B* **654**, 7-12 (2007), [arXiv:0705.3199 [astro-ph]]
- [20] A. A. Starobinsky, Disappearing cosmological constant in f(R) gravity, *JETP Lett.* 86, 157-163 (2007), [arXiv:0706.2041 [astro-ph]]
- [21] V. Faraoni, de Sitter space and the equivalence between f(R) and scalar-tensor gravity, *Phys. Rev. D* **75**, 067302 (2007), [arXiv:gr-qc/0703044]
- [22] C. G. Böhmer, L. Hollenstein, F. S. N. Lobo, Stability of the Einstein static universe in f(R) gravity, *Phys. Rev. D* **76**, 084005 (2007), [arXiv:0706.1663 [gr-qc]]
- [23] C. G. Böhmer, T. Harko, F. S. N. Lobo, Dark matter as a geometric effect in f(R) gravity, Astropart. Phys. 29, 386-392 (2008), [arXiv:0709.0046 [gr-qc]]
- [24] S. A. Appleby, R. A. Battye, A. A. Starobinsky, Curing singularities in cosmological evolution of F(R) gravity, JCAP 1006, 005 (2010), [arXiv:0909.1737 [astro-ph.CO]]
- [25] S. Chakraborty, K. MacDevette, P. Dunsby, A Model independent approach to the study of f(R) cosmologies with expansion histories close to Λ CDM, *Phys. Rev. D* **103**, 124040 (2021), [arXiv:2103.02274 [gr-qc]]
- [26] V. K. Oikonomou, Unifying inflation with early and late dark energy epochs in axion F(R) gravity, *Phys. Rev. D* 103, 044036 (2021), [arXiv:2012.00586 [astro-ph.CO]]
- [27] S. Nojiri and S. D. Odintsov, Modified gravity with negative and positive powers of the curvature: Unification of the inflation and of the cosmic acceleration, *Phys. Rev. D* 68 (2003), 123512 [arXiv:hep-th/0307288 [hep-th]].
- [28] S. Nojiri and S. D. Odintsov, Modified f(R) gravity consistent with realistic cosmology: From matter dominated epoch to dark energy universe, *Phys. Rev. D* 74 (2006), 086005 [arXiv:hep-th/0608008 [hep-th]].
- [29] G. Cognola, E. Elizalde, S. Nojiri, S. D. Odintsov, L. Sebastiani and S. Zerbini, A Class of viable modified f(R) gravities describing inflation and the onset of accelerated expansion, *Phys. Rev. D* 77 (2008), 046009 [arXiv:0712.4017 [hep-th]].
- [30] S. Nojiri and S. D. Odintsov, Unified cosmic history in modified gravity: from F(R) theory to Lorentz non-invariant models, *Phys. Rept.* **505** (2011), 59-144 [arXiv:1011.0544 [gr-qc]].
- [31] S. Nojiri and S. D. Odintsov, Unified cosmic history in modified gravity: from F(R) theory to Lorentz non-invariant models, *Phys. Rept.* **505** (2011), 59-144 [arXiv:1011.0544 [gr-qc]].
- [32] A. Dixit, D. C. Maurya, A. Pradhan, Transit cosmological models coupled with zero-mass scalar field with high redshift in higher derivative theory, New Astronomy 87, 101587 (2021), [arXiv:2005.11139 [physics.gen-ph]]
- [33] A. Pradhan, De Avik, Tee How Loo, D. C. Maurya, A flat FLRW model with dynamical Λ as function of matter and geometry, New Astronomy 89, 101637 (2021), [arXiv:2010.09444 [physics.gen-ph]]
- [34] D. C. Maurya, R. Zia, A. Pradhan, Anisotropic string cosmological model in Brans–Dicke theory of gravitation with time-dependent deceleration parameter, J. Exp. Theor. Phys. 123, 617-622 (2016),

- [35] D. C. Maurya, R. Zia, A. Pradhan, Dark energy models in LRS Bianchi type-II space-time in the new perspective of time-dependent deceleration parameter, *Int. J. Geom. Meth. Mod. Phys.* **14**(05) 1750077 (2017),
- [36] D. C. Maurya, Anisotropic dark energy transit cosmological models with time-dependent $\omega(t)$ and redshiftdependent $\omega(z)$ EoS parameter, Int. J. Geom. Meth. Mod. Phys. 15(02) 1850019 (2018),
- [37] R. Zia, D. C. Maurya, Brans-Dicke scalar field cosmological model in Lyra's geometry with time-dependent deceleration parameter, Int. J. Geom. Meth. Mod. Phys. 15(11) 1850186 (2018),
- [38] D. C. Maurya, R. Zia, Brans-Dicke scalar field cosmological model in Lyra's geometry, *Physical Review D* 100(2) 023503 (2019), [arXiv:1907.07135 [gr-qc]]
- [39] R. Zia, U. K. Sharma, D. C. Maurya, Transit two-fluid models in anisotropic Bianchi type-III space-time, New Astronomy 72, 83-91 (2019),
- [40] D. C. Maurya, R. Zia, Reply to "Comment on 'Brans-Dicke scalar field cosmological model in Lyra's geometry", Phys. Rev. D 102(10) 108302 (2020),
- [41] O. Bertolami, C. G. Boehmer, T. Harko, F. S. N. Lobo, Extra force in f(R) modified theories of gravity, *Phys. Rev. D* **75**, 104016 (2007), [arXiv:0704.1733 [gr-qc]]
- [42] T. P. Sotiriou, V. Faraoni, f(R) theories of gravity, Rev. Mod. Phys. 82, 451 (2010), [arXiv:0805.1726 [gr-qc]]
- [43] A. De Felice, S. Tsujikawa, f(R) theories, Living Rev. Relativ. 13, 3 (2010), [arXiv:1002.4928 [gr-qc]]
- [44] T. Harko, Modified gravity with arbitrary coupling between matter and geometry, Phys. Lett. B 669, 376 (2008), [arXiv:0810.0742 [gr-qc]]
- [45] T. Harko, F. S. N. Lobo, $f(R, L_m)$ gravity, Eur. Phys. J. C 70, 373 (2010), [arXiv:1008.4193 [gr-qc]]
- [46] T. Harko, The matter Lagrangian and the energy-momentum tensor in modified gravity with nonminimal coupling between matter and geometry, *Phys. Rev. D* 81, 044021 (2010), [arXiv:1001.5349 [gr-qc]]
- [47] T. Harko, F. S. N. Lobo, Geodesic deviation, Raychaudhuri equation, and tidal forces in modified gravity with an arbitrary curvature-matter coupling, *Phys. Rev. D* 86, 124034 (2012), [arXiv:1210.8044 [gr-qc]]
- [48] J. Wang, K. Liao, Energy conditions in $f(R, L_m)$ gravity, Class. Quantum Gravity 29, 215016 (2012), [arXiv:1212.4656 [physics.gen-ph]]
- [49] O. Minazzoli, T. Harko, New derivation of the Lagrangian of a perfect fluid with a barotropic equation of state, *Phys. Rev. D* 86, 087502 (2012), [arXiv:1209.2754 [gr-qc]]
- [50] T. Harko, F. S. N. Lobo, O. Minazzoli, Extended $f(R, L_m)$ gravity with generalized scalar field and kinetic term dependences, *Phys. Rev. D* 87, 047501 (2013), [arXiv:1210.4218 [gr-qc]]
- [51] D. W. Tian, I. Booth, Lessons from (R, R_c^2, R_m^2, L_m) gravity: Smooth Gauss-Bonnet limit, energy-momentum conservation, and nonminimal coupling, *Phys. Rev. D* **90**, 024059 (2014), [arXiv:1404.7823 [gr-qc]]
- [52] T. Harko, F. S. N. Lobo, J. P. Mimoso, D. Pavón, Gravitational induced particle production through a nonminimal curvature-matter coupling, *Eur. Phys. J. C* 75, 386 (2015), [arXiv:1508.02511 [gr-qc]]
- [53] R. P. L. Azevedo, P. P. Avelino, Big-bang nucleosynthesis and cosmic microwave background constraints on nonminimally coupled theories of gravity, *Phys. Rev. D* 98, 064045 (2018), [arXiv:1807.00798 [gr-qc]]
- [54] S. Bahamonde, Generalised nonminimally gravity-matter coupled theory, Eur. Phys. J. C 78, 326 (2018), [arXiv:1709.05319 [gr-qc]]

- [55] R. March, O. Bertolami, M. Muccino, et al., Constraining a nonminimally coupled curvature-matter gravity model with ocean experiments, *Phys. Rev. D* 100, 042002 (2019), [arXiv:1904.12789 [gr-qc]]
- [56] O. Bertolami, C. Gomes, Nonminimally coupled Boltzmann equation: foundations, Phys. Rev. D 102, 084051 (2020), [arXiv:2002.08184 [gr-qc]]
- [57] G. Allemandi, A. Borowiec, M. Francaviglia and S. D. Odintsov, Dark energy dominance and cosmic acceleration in first order formalism, *Phys. Rev. D* 72 (2005), 063505 [arXiv:gr-qc/0504057 [gr-qc]].
- [58] A. Pradhan, D. C. Maurya, G. K. Goswami, A. Beesham, Modeling Transit Dark Energy in $F(R, L_m)$ -gravity, Int. J. Geom. Meth. Mod. Phys., **20**(06) 2350105 (2023), [arXiv:2209.14269 [gr-qc]]
- [59] D. C. Maurya, Accelerating scenarios of massive universe in $f(R, L_m)$ -gravity, New Astronomy 100, 101974 (2023),
- [60] D. C. Maurya, Exact Cosmological Models in Modified $F(R, L_m)$ -Gravity with Observational Constraints, Grav. & Cosmo. 29(3) 315-325 (2023),
- [61] D. C. Maurya, Bianchi-I dark energy cosmological model in $f(R, L_m)$ -Gravity, Int. J. Geom. Meth. Mod. Phys. **21**(04) 2450072-194 (2024),
- [62] D. C. Maurya, Constrained ACDM Dark Energy Models in Higher Derivative $F(R, L_m)$ -Gravity Theory, Phys. Dark Universe 42 (2023) 101373,
- [63] T. Harko, F. S. N. Lobo, S. Nojiri, S. D. Odintsov, f(R,T) gravity, *Phys. Rev. D* 84, 024020 (2011), [arXiv:1104.2669 [gr-qc]]
- [64] T. Harko, Thermodynamic interpretation of the generalized gravity models with geometry-matter coupling, *Phys. Rev. D* **90**, 044067 (2014), [arXiv:1408.3465 [gr-qc]]
- [65] E. H. Baffou, M. J. S. Houndjo, M. E. Rodrigues, et al., Cosmological evolution in f(R,T) theory with collisional matter, *Phys. Rev. D* **92**, 084043 (2015), [arXiv:1504.05496 [gr-qc]]
- [66] T. B. Gonçalves, J. L. Rosa, F. S. N. Lobo, Cosmology in scalar-tensor f(R,T) gravity, *Phys. Rev. D* 105, 064019 (2022), [arXiv:2112.02541 [gr-qc]]
- [67] H. Shabani, A.H. Ziaie, Stability of the Einstein static universe in f(R,T) gravity, Eur. Phys. J. C 77, 31 (2017), [arXiv:1606.07959 [gr-qc]]
- [68] H. Velten, T. R. P. Caramês, Cosmological inviability of f(R, T) gravity, *Phys. Rev. D* **95**, 123536 (2017), [arXiv:1702.07710 [gr-qc]]
- [69] P. H. R. S. Moraes, R. A. C. Correa, G. Ribeiro, Evading the non-continuity equation in the f(R, T) cosmology, Eur. Phys. J. C 78, 192 (2018).
- [70] S. K. Maurya, A. Errehymy, Ksh. Newton Singh, et al., Gravitational decoupling minimal geometric deformation model in modified f(R,T) gravity theory, Phys. Dark Universe **30**, 100640 (2020).
- [71] T. Harko, P. H. R. S. Moraes, Comment on "Reexamining f(R, T) gravity", Phys. Rev. D 101, 108501 (2020).
- [72] R. Zia, D. C. Maurya, A. Pradhan, Transit dark energy string cosmological models with perfect fluid in F(R,T)-gravity, Int. J. Geom. Meth. Mod. Phys. 15(10) 1850168 (2018).
- [73] D. C. Maurya, Modified F(R,T) cosmology with observational constraints in Lyra's geometry, Int. J. Geom. Meth. Mod. Phys. 17(01) 2050001 (2020).

- [74] D. C. Maurya, A. Pradhan, A. Dixit, Domain walls and quark matter in Bianchi type-V universe with observational constraints in F(R,T) gravity, Int. J. Geom. Meth. Mod. Phys. 17(01) 2050014 (2020).
- [75] D. C. Maurya, Transit cosmological model with specific Hubble parameter in F(R, T) gravity, New Astronomy 77, 101355 (2020).
- [76] D. C. Maurya, J. singh, L. K. Gaur, Dark Energy Nature in Logarithmic f(R,T) Cosmology, Int. J. Geom. Meth. Mod. Phys. **20**(11) 2350192 (2023).
- [77] G. P. Singh, N. Hulke, A. Singh, Cosmological study of particle creation in higher derivative theory, *Indian J. Phys.* 94 127-141 (2020).
- [78] N. Hulke, G. P. Singh, B. K. Bishi, A. Singh, Variable Chaplygin gas cosmologies in f(R,T) gravity with particle creation, New Astronomy 77 101357 (2020).
- [79] Z Haghani and T. Harko, Generalizing the coupling between geometry and matter: $f(R, L_m, T)$ gravity, Eur. Phys. J. C (2021) 81:615.
- [80] S. Arora, P.H.R.S. Moraes, and P.K. Sahoo, Energy conditions in the f(R, L, T) theory of gravity, *Eur. Phys. J. Plus* **139** (2024) 542.
- [81] A. Pradhan, M. Zeyauddin, A. Dixit, and S. Krishnannair, Dust-fluid accelerating flat cosmological models in $f(R, T, L_m)$ -gravity with observational constraints, Int. J. Geom. Meth. Mod. Phys., (2024), https://doi.org/10.1142/S0219887825500604.
- [82] J.S. Gonçalves, A.F. Santos, On causality and its violation in $f(R, L_m, T)$ gravity, Nuclear Physics B 1010 (2025) 116751.
- [83] D. C. Maurya, Late-time accelerating cosmological models in $f(R, L_m, T)$ -gravity with observational constraints, *Phys. Dark Universe* **46** (2024) 101722.
- [84] D. C. Maurya, Constrained transit cosmological models in $f(R, L_m, T)$ -gravity, Int. J. Geom. Meth. Mod. Phys., (2024), https://doi.org/10.1142/S0219887825500288.
- [85] S. Capozziello, O. Luongo, R. Pincak, et al., Cosmic acceleration in non-flat f(T) cosmology, Gen. Relativ. Gravit. 50, 53 (2018), [arXiv:1804.03649v1 [gr-qc]].
- [86] L. D. Landau, E. M. Lifshitz, The Classical Theory of Fields (Butterworth-Heinemann, Oxford, 1998).
- [87] E. J. Copeland, M. Sami and S. Tsujikawa, Dynamics of dark energy, Int. J. Mod. Phys. D 15, 1753 (2006), [arXiv:hep-th/0603057].
- [88] J. Simon, L. Verde, R. Jimenez, Phys. Rev. D 71, 123001 (2005)
- [89] G. S. Sharov, V. O. Vasiliev, Math. Model. Geom. 6, 1-20 (2018)
- [90] K. Asvesta, L. Kazantzidis, L. Perivolaropoulos, C.G. Tsagas, Observational constraints on the deceleration parameter in a tilted universe, Mon. Not. R. Astron. Soc. 513, 2394-2406 (2022).
- [91] D.W. Hogg and D.F. Mackey, Data analysis recipes: Using Markov Chain Monte Carlo, ApJS 236 (2018)
 18. [arXiv:1710.06068 [astro-ph.IM]].
- [92] R. Jimenez and A. Loeb, Constraining Cosmological Parameters Based on Relative Galaxy Ages, ApJ 573 (2002) 37.
- [93] G. Ellis, R. Maartens, M. MacCallum, Relativistic Cosmology (Cambridge University Press, Cambridge, 2012). https://doi.org/10.1017/CBO9781139014403.

- [94] A. Conley et al., Supernova constrants and systemetic uncertainities from the first three years of the supernova legacy survey^{*}, ApJ **192** (2011) 1.
- [95] D. M. Scolnic et al., The Complete Light-curve Sample of Spectroscopically Confirmed SNe Ia from Pan-STARRS1 and Cosmological Constraints from the Combined Pantheon Sample, ApJ 859 (2018) 101.
- [96] Dong Zhao, Yong Zhou, Zhe Chang, Anisotropy of the Universe via the Pantheon supernovae sample revisited, MNRAS 486 (2019) 5679-5689, https://doi.org/10.1093/mnras/stz1259.
- [97] L. Kazantzidis, L. Perivolaropoulos, Hints of a local matter underdensity or modified gravity in the low z Pantheon data, Phys. Rev. D 102 (2020) 023520, https://link.aps.org/doi/10.1103/PhysRevD.102.023520.
- [98] D. Sapone, et al., Is there any measurable redshift dependence on the SN Ia absolute magnitude?, *Phys. Dark Univ.* 32 (2021) 100814.
- [99] L. Kazantzidis, H. Koo, S. Nesseris, L. Perivolaropoulos, A. Shafieloo, Hints for possible low redshift oscillation around the best-fitting ACDM model in the expansion history of the Universe, MNRAS 501 (2021) 3421-3426, https://doi.org/10.1093/mnras/staa3866.
- [100] M. G. Dainotti et al., On the Hubble Constant Tension in the SNe Ia Pantheon Sample, ApJ **912** (2021) 150.
- [101] M. G. Dainotti et al., On the Evolution of the Hubble Constant with the SNe Ia Pantheon Sample and Baryon Acoustic Oscillations: A Feasibility Study for GRB-Cosmology in 2030, *Galaxies* 10 (2022) 24; https://doi.org/10.3390/galaxies10010024.
- [102] G. Alestas, L. Kazantzidis, L. Perivolaropoulos, w M phantom transition at $z_t \simeq 0.1$ as a resolution of the Hubble tension, *Phys. Rev. D* 103 (2021) 083517.
- [103] D. Camarena, V. Marra, On the use of the local prior on the absolute magnitude of Type Ia supernovae in cosmological inference, MNRAS 504 (2021) 5164-5171, https://doi.org/10.1093/mnras/stab1200.
- [104] V. Marra, L. Perivolaropoulos, Rapid transition of G_{eff} at $z_t \simeq 0.01$ as a possible solution of the Hubble and growth tensions, *Phys. Rev. D* **104** (2021) L021303.
- [105] G. Alestas, I. Antoniou L. Perivolaropoulos, Hints for a Gravitational Transition in Tully-Fisher Data, Universe 7 (2021) 366; https://doi.org/10.3390/universe7100366.
- [106] L. Perivolaropoulos, Is the Hubble crisis connected with the extinction of dinosaurs?, arXiv:2201.08997 [astro-ph.EP].
- [107] D. C. Maurya, Accelerating cosmological models in Hoyle-Narlikar gravity with observational constraints, Int. J. Geom. Meth. Mod. Phys., (2025). https://doi.org/10.1142/S0219887825500896.
- [108] A.R. Lalke, G.P. Singh, A. Singh, Cosmic dynamics with late-time constraints on the parametric deceleration parameter model. Eur. Phys. J. Plus 139 288 (2024). https://doi.org/10.1140/epjp/s13360-024-05091-5
- [109] S. Mandal, A. Singh, R. Chaubey, Late-time constraints on barotropic fluid cosmology, Phys. Lett. A 519 (2024) 129714, https://doi.org/10.1016/j.physleta.2024.129714.
- [110] A. Singh, S. Krishnannair, Affine EoS cosmologies: Observational and dynamical system constraints, Astronomy and Computing 47 (2024) 100827, https://doi.org/10.1016/j.ascom.2024.100827.
- [111] K. Anderson, Model selection and multimodel inference: a practical information-theoretic approach, 2nd edn. Springer, New York (2002).

- [112] K. P. Burnham, D. R. Anderson, Multimodel Inference: Understanding AIC and BIC in Model Selection, Sociological Methods and Research 33, 261 (2004).
- [113] A. R. Liddle, Information criteria for astrophysical model selection, Mon. Not. Roy. Astron. Soc. 377, (2007) L74, [arXiv:astro-ph/0701113].
- [114] R. E. Kass and A. E. Raftery, Bayes Factors, J. Am. Statist. Assoc. 90 773 (1995).
- [115] F. K. Anagnostopoulos, S. Basilakos, E.N. Saridakis, Observational constraints on Myrzakulov gravity, *Phys. Rev. D* 103, 104013 (2021), [arXiv:2012.06524 [gr-qc]].
- [116] S. Capozziello, O. Farooq, O. Luongo, *et al.*, Cosmographic bounds on the cosmological decelerationacceleration transition redshift in f(R) gravity, *Phys. Rev. D* **90**, 044016 (2014), [arXiv:1403.1421v1 [gr-qc]].
- [117] S. Capozziello, O. Luongo, E. N. Saridakis, Transition redshift in f(T) cosmology and observational constraints, *Phys. Rev. D* **91** 124037 (2015), [arXiv:1503.02832v2 [gr-qc]].
- [118] S. Capozziello, P. K. S. Dunsby, O. Luongo, Model independent reconstruction of cosmological accelerateddecelerated phase, MNRAS 509 (2022) 5399-5415, [arXiv:2106.15579v2 [astro-ph.CO]].
- [119] M. Muccino, O. Luongo, D. Jain, Constraints on the transition redshift from the calibrated Gamma-ray Burst Ep-Eiso correlation, MNRAS 523 (2023) 4938-4948, [arXiv:2208.13700v3 [astro-ph.CO]].
- [120] A. C. Alfano, S. Capozziello, O. Luongo, et al., Cosmological transition epoch from gamma-ray burst correlations, (2024) [arXiv:2402.18967v1 [astro-ph.CO]].
- [121] A. C. Alfano, C. Cafaro, S. Capozziello, et al., Dark energy-matter equivalence by the evolution of cosmic equation of state, Phys. Dark Univ. 42 101298 (2023), [arXiv:2306.08396v2 [astro-ph.CO]].
- [122] G. Ellis, R. Maartens, M. MacCallum, Causality and the speed of sound, Gen. Rel. Grav. 39 1651-1660 (2007), [arXiv:gr-qc/0703121]