Edge Exchangeable Graphs: Connectedness, Gaussianity and Completeness

Edward Eriksson

17th January 2025

1 Introduction

1.1 Model (Informal)

Following [1][2][3][4][5][6] among others, we study models of the following type. Start with an empty (no vertices, no edges) graph and a probability measure μ on \mathbb{N}_2 , the set of unordered pairs of distinct natural numbers. We think of the natural numbers as the set of possible vertices and \mathbb{N}_2 as the set of possible edges. Define a (multi-)graph-valued process by repeatedly sampling an edge according to μ and adding it, along with any needed vertices to the graph. We will be interested in properties (e.g. connectedness, number of vertices) which are invariant under identifying parallel edges, which we may therefore do. There are closely related (poissonized) versions both before and after identifying parallel edges. For the former define a poisson process with intensity $\mu_e := \mu(\{e\})$ for each edge $e \in \mathbb{N}_2$ which counts the number of copies of that edge and for the latter let the arrival time of e, denoted τ_e , be exponentially distributed with parameter μ_e . In both cases take the vertex set to be the minimal set of vertices necessary. The unfamiliar reader may benefit from reviewing the more careful introduction of these models in Section 2 before proceeding.

1.2 Contributions

Our main contributions are giving a complete characterization of eventual forever connectedness in these models (see Theorem 14) and resolving a conjecture of Janson [1] on the asymptotic normality of the number of vertices,

in this connected regime (see Theorem 24). We also resolve an open problem of Janson [1] regarding when the produced graphs are almost complete (see Theorem 39). We also obtain an apparently new result characterizing when the urns in a fairly classical, albeit much less so than its finite variants, urn scheme [7][8][9][10] are eventually forever filled in order (see Proposition 37).

1.3 Structure

The remainder of Section 1 consists of two parts. The former, Subsection 1.4, will motivate the model of interest and our contribution to the theory thereof. The second, Subsection 1.5, will discuss previous work on the model we are studying, as well as on other related models. In Section 2 we will introduce the models of interest formally. In Section 3 we introduce some preliminaries and notation. Each of the following three sections, namely Sections 4, 5 and 6 are dedicated to one of the three main theorems. The first the characterization of eventual forever connectedness, the second the asymptotic Gaussianity of the vertex count in the connected regime and the third the characterization of eventual forever connectedness. We finish by discussing some possible directions for future work in Section 7.

1.4 Motivation

1.4.1 Motivating Edge Exchangeable Models

A sequence of random variables is said to be exchangeable if its law is invariant to finite permutations. This is weaker than being iid from some distribution, instead capturing the intuition that the random variables are observed in no particular order. The classical graphon models are vertex-exchangeable [3] meaning that their law is invariant under permuting the order of arrival of the vertices. As an example, if our graphs represent the friend network on a social media site then assuming vertex exchangability would correspond to assuming that people join the network in no particular order [2][3]. When using a vertex exchangeable model we end up tacitly assuming that we have seen the all interactions between the observed vertices and are only able to produce dense (or empty graphs) which is often not realistic [2][11]. In some applications it seems quite plausible then to instead believe that we are observing *interactions* rather than *agents* in no particular order, which would call for edge exchangeable models.

1.4.2 Motivating our Contribution

The models we study are motivated by the study of edge exchangeable graphs, from which they arise once we fix μ instead of having μ be random. One would like to ensure that edge exchangeable models can produce graphs which resemble those encountered in practice. This calls for understanding how the qualities of the produced graphs depend on μ . Moreover, in some applications one may have a choice of (prior over) μ . In that case results of the type we provide may help inform the choice of this infinite-dimensional parameter. In addition to being motivated by statistical applications, the model is quite natural and displays some interesting phenomena which are not typical of random graph models, making them probabilistically interesting. For example, the vertex set is not deterministic nor generated randomly and then the graph generated conditionally on it, we prove results which are really about the entire history of the graph-valued process, not just its marginals and we find that a topological assumption has distributional consequences.

1.5 Related Work

1.5.1 Previous Work on Edge Exchangeable Models

Edge exchangeable models go back to at least [2][4]. The former places significant emphasis on hyper-graphs, but we restrict ourselves to ordinary graphs. The same paper introduces a particular model, called the Hollywood model as a particular edge exchangeable model aimed at modeling associations between actors and studies inference in this context. It moreover gives a De Finetti-type characterization theorem which morally states that all edge exchangeable graph-valued processes are of the type we described, with μ random. See Section 2 for more details. Both [2] and [4] consider sparsity both before and after identifying parallel edges. They both show that sparsity is possible for edge exchangeable graphs by showing that a particular model is sparse. A more general study of sparsity as well as other graph properties after identifying parallel edges was undertaken by Janson in [1]. The present paper can be seen as a continuation of the research program started in [1]. Janson gives asymptotic formulas for the number of edges and vertices, albeit the asymptotics of those formulas themselves are in general not trivial to understand. Moreover, asymptotic normality for the number of edges is established under the assumption that the variance of the number of edges goes to infinity. Example μ are given which produce everything

from extremely sparse to extremely dense graphs. A sufficient condition on μ is given such that the graphs are essentially complete (see Definition 32). Statistical applications of edge exchangeability are also still being developed, for instance, anomaly detection [5]. The theory of simulation (sampling) and inference for first rank μ (see Definition 21) was developed in [6]. The textbook by Crane [12] contains an introduction to edge-exchangeable models and motivates them better than we can do here.

1.5.2 Comparisons to other Models

Another type of exchangability for graph-valued stochastic processes is that of Caron and Fox's exchangeable measures [13]. These models can also produce sparse graphs, but it is much less clear what their exchangability corresponds to in terms of modeling the real-world and hence when it can be expected (see Crane's comment in [13]).

Models which resemble the ones we are interested in arise in various contexts. For instance, we may imagine an inference problem in which there is some true unknown finite graph and for each edge there is an exponential time (independent but not necessarily identically distributed) until we observe it. This was considered, and suitable statistical estimators developed, in [14], based on the second author's PhD thesis [15]. The same model, although therein viewed from the perspective of percolation theory more so than statistics, is considered in [16],[17] where some weak concentration results for the emergence of a giant component are established. This has some rough parallels to our characterization of connectedness.

One general type of graph model is the inhomogeneous random graph. There one works with a finite deterministic vertex set and assign a probability p_e (not necessarily identical) to each possible edge and decide whether to include each edge independently of all other edges. At all times our exponential arrival time graph can be viewed as an instantiation of this model with an infinite vertex set, $p_e = 1 - e^{-\mu_e t}$ and the additional rule that all isolated vertices are discarded.

In this work we are agnostic as to how μ and consequently the p_e :s arise. However, every way to obtain p_e :s for an inhomogeneous random graph can also be combined with the edge exchangeable framework and produce μ_e :s. For instance, one may imagine that each vertex is associated to a vector in some latent space and that the μ_e :s arise as dot products of these vectors like the classical dot product random graphs, a survey of which can be found here [18]. Note that in this case some technical care needs to be taken to ensure that μ can be normalized.

The Norros-Reittu model [19] is another model which has an independent poisson number of edges between each pair of vertices. However, in that model the intensity is (up to a a common scaling) the product of intensities associated to the vertices. Thus it roughly corresponds to our first rank (see Definition 21) case. Moreover, the vertex set is deterministic unlike ours.

2 Model (Formal)

Throughout we assume that we work on a probability space rich enough to support all processes and random variables we describe but make no further mention of it.

2.1 Discrete Time

Let $G_0 = (\{\}, \{\})$ be the empty un-directed multigraph and recursively define G_n by

$$G_{n+1} := G_n \cup e_{n+1}, e_{n+1} \sim \mu,$$

where the edge $e_{n+1} = \{i, j\}$ sampled according to μ is identified with the undirected (multi)-graph ($\{i,j\},\{\{i,j\}\}$) and multi-graph unions are defined by

$$G_1 \cup G_2 := (V_1 \cup V_2, E_1 + E_2)_2$$

where the sum is understood to be a multi-set sum. We abuse notation and use G_n also for the graph-valued stochastic process obtained by identifying parallel edges.

2.2 Continuous Time

For each $e \in \mathbb{N}_2$ let $N_e(t)$ be a Poisson process, all independent, with intensity μ_e .

For a set A and a non-negative integer k define kA to be the multiset which consists of k copies of each element of A. Let $E_t := \sum_{e \in \mathbb{N}_2} N_e(t) \{e\}$, where the sum is a multiset sum. Since the sets are disjoint one could equivalently take a multiset union. Let $V_t = \{v \in \mathbb{N} : \exists e \in E_t \text{ s.t. } v \in e\}$ and finally define

$$G_t := (V_t, E_t).$$

We abuse notation and use G_t also for the graph-valued stochastic process obtained by identifying parallel edges.

Remark 1. These are not the most general edge exchangeable random graphs possible, indeed one can allow a random μ , loops, hyperedges and directed edges. There is also a slight bit of additional generality by allowing a different latent vertex set than N. See [2][3][12][1] for more general definitions and characterization theorems.

Remark 2. Note that we no longer need μ to be normalized for the construction in Subsection 2.2 to make sense. It suffices that $\mu(\mathbb{N}_2) < \infty$. By a standard property of poisson processes rescaling μ corresponds to rescaling time. Thus when studying properties which are invariant to rescaling time one need not normalize μ . This can save us from computing normalizing constants.

3 Preliminaries and Notation

For a set \mathcal{X} we use $\mathcal{P}(\mathcal{X})$ to denote the set of all probability measures over \mathcal{X} . When clear from context we drop limits of summation, preferring for example to write $\sum_i a_i$ rather than $\sum_{i=1}^{\infty} a_i$. For $\mu \in \mathcal{P}(\mathbb{N}_2)$ define $\mu_{ij} := \mu(\{i, j\})$ and define $M_i = \sum_j \mu_{ij}$ for $i \in \mathbb{N}$. For $e = \{i, j\}$ define $M_e := M_{ij} :=$ $M_i + M_j - \mu_{ij}$.

Definition 3. For a stochastic process X indexed by I taking values in some set \mathcal{X} and a property $A : \mathcal{X} \to \{\text{True}, \text{False}\}$ we say that X has or satisfies A **eventually forever** iff there exists an $i(\omega)$ such that A(X(j)) = True for all $j \in I$ such that j > i.

Remark 4. In general whether X(i) has A eventually forever is random. However, it will transpire that for the processes and properties we are interested in whether the process satisfies the property eventually forever has probability either zero or one. We may therefore omit the almost surely and simply write X has A eventually forever to mean that X almost surely has A eventually forever.

Lemma 5. If A is a property that depends only on the order of appearance of edges then G_t has A eventually forever iff G_n has A eventually forever.

Proof. This is immediate from the fact that we can sample G_n by waiting until n events have occurred in G_t .

For a finite collection of random variables $C = {\tau_i}_{i \in I}$ we use $\check{C} := \max_{i \in I} \tau_i$ and $\hat{C} := \min_{i \in I} \tau_i$ to denote the (random) maximum and minimum, respectively.

Definition 6. Let X_t be a stochatic process in either discrete or continuous time which takes values in (a subset of) \mathbb{R} . We call such a process **asymptotically Gaussian** if there exist deterministic functions $f, g : \mathbb{R} \to \mathbb{R}$ such that

$$\frac{X_t - f(t)}{g(t)} \to \phi$$

where ϕ is the standard normal distribution and the convergence is in distribution as $t \to \infty$.

Remark 7. Of course, the most natural candidates for f and g in Definition 6 are $\mathbb{E}[X_t]$ and $\sqrt{\mathbb{V}[X_t]}$, respectively.

4 Connectedness

We will now build up towards characterizing eventual forever connectedness.

Definition 8. For a $\mu \in \mathcal{P}(\mathbb{N}_2)$ consider the (typically infinite) graph ($\{v \in \mathbb{N} : \exists e \in \mathbb{N}_2 \text{ s.t. } v \in e, \mu_e > 0\}, \{e \in \mathbb{N}_2 : \mu_e > 0\}$). We call this graph the **support** of μ and when it is connected we say that μ has **connected support**.

Remark 9. Many natural μ assign positive probability to each edge, in which case the support is the complete graph on \mathbb{N} .

By Lemma 5 it will suffice to consider G_t , so we state the following only for G_t , even when the same holds for G_n .

Remark 10. If μ does not have connected support, then G_t is not eventually forever connected since edges that cannot be connected will eventually appear.

Let I_e be the event that the the edge e brings two new vertices to G_t when it arrives. This is well-defined since almost surely edges are not added at the same time.

Remark 11. When an edge arrives it adds zero, one or two new vertices. A new component is created if and only if the edge brings two new vertices.

Lemma 12. If only finitely many I_e occur and μ has connected support, then G_t is eventually forever connected. One the other hand, if infinitely many I_e occur then G_t is not eventually forever connected.

Proof. If only finitely many I_e occur there exists a last time that a new component is added by Remark 11. Since μ has connected support edges which will cause all the existing components to join up will subsequently appear, after which point G_t will be connected. On the other hand, immediately following an edge bringing two new vertices the graph cannot be connected. If this happens infinitely many times the graph cannot be eventually forever connected as by standard properties of poisson processes it cannot be that there exists a finite time interval in which all of the edges for which I_e occurred appear.

Lemma 12 reduces the problem of determining eventual forever connectedness to deciding whether infinitely many I_e occur. We do this by computing $P(I_e)$ and $P(I_e \wedge I_f)$ and applying the first Borel-Cantelli Lemma and a suitable pseudo-inverse. We now define the notation that will be required for the computation of $P(I_e \wedge I_f)$. For $e, f \in \mathbb{N}_2$ with $e \cap f = \emptyset$ let $C_{ef} = \{g \in \mathbb{N}_2 : g \cap e \neq \emptyset, g \cap f \neq \emptyset\}$, $C_e := \{g : |e \cap g| = 1\} \setminus C_{ef}$ and define C_f analogously. Note that, $\{e\}, \{f\}, C_{ef}, C_e, C_f$ are disjoint and that $\tau_e, \tau_f, \{\hat{\tau}_g\}_{g \in C_e f}, \{\hat{\tau}_g\}_{g \in C_e}, \{\hat{\tau}_g\}_{g \in C_f}$ are therefore independent and exponentially distributed with parameters $\mu_e, \mu_f, b_{ef} := \sum_{g \in C_e f} \mu_g, r_e := \sum_{g \in C_f} \mu_g$ respectively. For compactness of notation we define $a_e := r_e + \mu_e, c_e := 1 - \frac{r_e}{r_e + \mu_e} = \frac{\mu_e}{a_e}$. We include Figure 1 to help with parsing this notation.

Lemma 13. For $e, f \in \mathbb{N}_2$ we have $|e \cap f| \in \{0, 1, 2\}$ and

•
$$|e \cap f| = 0 \implies P(I_e \wedge I_f) = c_e c_f \left(1 - \frac{b_{ef}}{a_e + b_{ef}} - \frac{b_{ef}}{a_f + b_{ef}} + \frac{b_{ef}}{a_e + a_f + b_{ef}}\right).$$

• $|e \cap f| = 1 \implies P(I_e \wedge I_f) = 0.$



Figure 1: Edge Notation Some selected edges labeled by the sets they belong to.

• $|e \cap f| = 2 \implies e = f$ and $P(I_e \wedge I_f) = P(I_e) = \frac{\mu_e}{M_e}$.

Proof. $|e \cap f| \in \{0, 1, 2\}$ is immediate from the fact that |e| = |f| = 2. If either μ_e or μ_f is zero, it is clear that $P(I_e \wedge I_f) = 0$ in which case the Lemma holds so we may assume both to be non-zero. First we deal with the case $|e \cap f| = 0$. We will need the following sub-lemma.

$$\begin{split} P(I_e|\{\hat{\tau}_g\}_{g\in C_{ef}} = t) &= \int_0^\infty P(I_e|\{\hat{\tau}_g\}_{g\in C_{ef}} = t, \{\hat{\tau}_g\}_{g\in C_e} = \tau)r_e e^{-r_e\tau}d\tau, \\ &= \int_0^\infty (1 - e^{-\mu_e \min(t,\tau)})r_e e^{-r_e\tau}d\tau, \\ &= 1 - \int_0^\infty e^{-\mu_e \min(t,\tau)}r_e e^{-r_e\tau}d\tau, \\ &= 1 - \int_0^t e^{-\mu_e\tau}r_e e^{-r_e\tau} - e^{-\mu_e t}\int_t^\infty r_e e^{-r_e\tau}d\tau, \\ &= 1 - \frac{r_e}{r_e + \mu_e} \left(1 - e^{-(r_e + \mu_e)t}\right) - e^{-(\mu_e + r_e)t}, \\ &= \left(1 - \frac{r_e}{r_e + \mu_e}\right) \left(1 - e^{-(r_e + \mu_e)t}\right), \\ &= c_e (1 - e^{-a_e t}). \end{split}$$

Now we compute

$$\begin{split} P(I_e \wedge I_f) &= \int_0^\infty P(I_e \wedge I_f | \{ \hat{\tau}_g \}_{g \in C_{ef}} = t) b_{ef} e^{-b_{ef} t} dt, \\ &= \int_0^\infty P(I_e | \{ \hat{\tau}_g \}_{g \in C_{ef}} = t) P(I_f | \{ \hat{\tau}_g \}_{g \in C_{ef}} = t) b_{ef} e^{-b_{ef} t} dt, \\ &= \int_0^\infty c_e (1 - e^{-a_e t}) c_f (1 - e^{-a_f t}) b_{ef} e^{-b_{ef} t} dt, \\ &= c_e c_f \left(1 - \frac{b_{ef}}{a_e + b_{ef}} - \frac{b_{ef}}{a_f + b_{ef}} + \frac{b_{ef}}{a_e + a_f + b_{ef}} \right). \end{split}$$

where the third equality is obtained by applying the sub-lemma twice.

If $|e \cap f| = 1$ let $v \in e \cap f$ be the shared vertex. For both I_e and I_f to occur both edges have to bring v. Since edges almost surely do not arrive at the same time and the vertex set is increasing this is not possible so $P(I_e \wedge I_f) = 0$.

If $|e \cap f| = 2$, then, since |e| = |f| = 2 we have e = f and $I_e \wedge I_f = I_e$. The event I_e is precisely the event that $\{\hat{\tau}_g\}_{g:e \cap g \neq \emptyset} = \tau_e$ and so, by a standard property of exponential random variables,

$$P(I_e) = \frac{\mu_e}{M_e}.$$

Theorem 14. G_t and G_n are eventually forever connected with probability one if $\sum_{e \in \mathbb{N}_2} \frac{\mu_e}{M_e} < \infty$ and μ has connected support. Else they are almost surely not eventually forever connected.

Proof. By Lemma 5 it suffices to consider G_t . Recall that the support of μ being connected is necessary by Remark 10. We may therefore henceforth assume that μ has connected support. By Lemma 12 it remains to resolve whether infinitely many I_e occur. If

$$\sum_{e \in \mathbb{N}_2} \frac{\mu_e}{M_e} < \infty,$$

then almost surely only finitely many I_e occur by the first Borel-Cantelli lemma and the third case of Lemma 13. Now assume instead that $\sum_{e \in \mathbb{N}_2} \frac{\mu_e}{M_e} =$

 ∞ .

$$\begin{aligned} \frac{P(I_e \wedge I_f)}{P(I_e)P(I_f)} &= \frac{a_e + b_{ef}}{\mu_e} \frac{a_f + b_{ef}}{\mu_f} c_e c_f \left(1 - \frac{b_{ef}}{a_e + b_{ef}} - \frac{b_{ef}}{a_f + b_{ef}} + \frac{b_{ef}}{a_e + a_f + b_{ef}} \right) \\ &= \frac{a_e + a_f + 2b_{ef}}{a_e + a_f + b_{ef}}, \\ &< 2. \end{aligned}$$

where we used the first and third cases in Lemma 13 in the first equality, along with the observation that $M_e = a_e + b_{ef}$ and lengthy but elementary algebra in the second and the last inequality is obtained from an elementary calculus exercise. Namely, one notices that for a constant k, $h(x) := \frac{k+2x}{k+x}$ satisfies $\lim_{x\to\infty} h(x) = 2$ and $h'(x) = \frac{k}{(k+x)^2} > 0$. Taking reciprocals gives

$$\frac{P(I_e)P(I_f)}{P(I_e \wedge I_f)} > \frac{1}{2}.$$
(1)

Since \mathbb{N}_2 is countable we may order it in some arbitrary but fixed way. In the following we take e < n to mean that $e \in \mathbb{N}_2$ is one of the first n - 1 edges under the ordering.

$$P(I \text{ i.o}) \ge \limsup_{n \to \infty} \frac{\sum_{e, f < n} P(I_e) P(I_f)}{\sum_{e, f < n} P(I_e \land I_f)} \ge \limsup_{n \to \infty} \min_{e, f < n} \frac{1}{2} = \frac{1}{2}$$

where we applied the Kochen-Stone Lemma [20] in the first inequality and Lemma 41 as well as (1) in the second. Next notice that I i.o is measurable with respect to $\sigma(\bigcup_{e \in \mathbb{N}_2} \sigma(\tau_e))$, that the τ_e :s are independent and that whether I i.o occurs cannot be changed by changing the values of the τ_e :s in a fixed finite collection. That is to say, I i.o is a tail event in the sense of Kolmogorov's 0-1 law, the invoking of which implies the theorem.

Corollary 15. Let $\mu_{ij} \propto (ij)^{-\gamma}$ for $i \neq j$. Then G_n and G_t are eventually forever connected iff $\gamma > 2$.

Proof. The support of μ is the complete graph which is connected. Note that $\frac{\mu_e}{M_e}$ is invariant under rescaling of μ so we may assume that the proportionality

is in fact an equality. We compute

$$M_{ij} = \sum_{k} \mu_{kj} + \sum_{k} \mu_{ik} - \mu_{ij}$$

= $\sum_{k \neq j} (kj)^{-\gamma} + \sum_{k \neq i} (ik)^{-\gamma} - (ij)^{-\gamma}$
= $j^{-\gamma} \zeta(\gamma) - (jj)^{-\gamma} + i^{-\gamma} \zeta(\gamma) - (ii)^{-\gamma} - (ij)^{-\gamma},$

where ζ denotes the Riemann zeta function. This implies that

$$\sum_{ij} \frac{\mu_{ij}}{M_{ij}} = \sum_{i \neq j} \frac{(ij)^{-\gamma}}{j^{-\gamma} \zeta(\gamma) - (jj)^{-\gamma} + i^{-\gamma} \zeta(\gamma) - (ii)^{-\gamma} - (ij)^{-\gamma}}, \qquad (2)$$

$$=\sum_{i\neq j}\frac{1}{i^{\gamma}(\zeta(\gamma)-j^{-\gamma})+j^{\gamma}(\zeta(\gamma))-i^{-\gamma})-1}.$$
(3)

Since one can easily show that the diagonal terms are negligible and that the row and column sums converge and since

$$\lim_{i,j\to\infty}\frac{i^{\gamma}(\zeta(\gamma)-j^{-\gamma})+j^{\gamma}(\zeta(\gamma))-i^{-\gamma})-1}{i^{\gamma}+j^{\gamma}}=\zeta(\gamma)>0,$$

by the limit comparison test for double sums [21], $\sum_{e} \frac{\mu_e}{M_e}$ converges if and only if

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \frac{1}{i^{\gamma} + j^{\gamma}},$$

converges. Finally, this second series converges iff $\gamma > 2$ [22].

Remark 16. From [1] it is known that μ of this type produce sparse graphs regardless of γ . We thus see that even natural models ("natural" being used colloquially) can simultaneously have good connectedness and good sparsity properties.

Eventual forever connectedness is quite a strong notion of connectedness so one can of course straightforwardly pass to the usual marginal notion. However, one does need to de-Poissonize slightly differently than we have seen before as Lemma 5 does not apply. **Corollary 17.** Let C_n be the event that G_n is connected and let C_t be the event that G_t is connected. Suppose that μ satisfies the assumptions of Theorem 14. Then

$$\lim_{n \to \infty} P(C_n) = 1,$$
$$\lim_{t \to \infty} P(C_t) = 1.$$

Proof. Eventual forever connectedness is equivalent to $\{1_{C_n}\}_{n=1}^{\infty}$ converging almost surely to 1. The first part now follows from almost sure convergence being stronger than convergence in probability. To avoid any subtleties in convergence of a sequence of uncountably many random variables, we obtain the continuous time result by first generating the number of edges that have been drawn and then conditioning on it. Fix $\epsilon > 0$. Choose an N such that $P(C_n) > 1 - \epsilon$ for all n > N. For t sufficiently large,

$$\sum_{k=0}^{N} \frac{t^k e^{-t}}{k!} < \epsilon.$$

For such t we have

$$P(C_t) = \sum_{k=0}^{\infty} P(C_k) \frac{t^k e^{-t}}{k!} > \sum_{k=N}^{\infty} P(C_k) \frac{t^k e^{-t}}{k!},$$
$$> \sum_{k=N}^{\infty} (1-\epsilon) \frac{t^k e^{-t}}{k!},$$
$$= (1-\epsilon)^2.$$

Since epsilon was arbitrary, this completes the proof.

5 Gaussinaity

Janson [1] showed asymptotic Gaussianity for the number of edges by coupling to certain urn schemes and we take a similar approach for the vertices. The key difference is that the arrival of edges are, after poissonizing, independent while the arrival of vertices are not. We will assume that the graph is eventually forever connected. Our key insight is that this assumption will allow us to control the dependence.

We will now introduce two urn schemes. For the first scheme, fix a sequence of probabilities $\{p_i\}_{i=1}^{\infty}$ such that $\sum_i p_i = 1$. For each $i \in \mathbb{N}$ define an urn. At times t = 1, 2, 3, ... throw a ball with probability p_i of landing in urn *i*. Let U_n be the set of occupied urns at time *n* so that $|U_n|$ is the number of occupied urns at time n. For the second scheme, fix a sequence of non-negative reals $\{\lambda_i\}_{i=1}^{\infty}$ such that $\sum_i \lambda_i < \infty$. For each urn define a poisson process with intensity λ_i and add a ball to the urn whenever there is an event in the associated poisson process. Let U_t be the set of occupied urns at time t so that $|U_t|$ is the number of occupied urns at time t. When $\sum_i \lambda_i = 1$ the latter scheme is the poisonized version of the former with $p_i = \lambda_i$. We will abuse notation and write U_t even when we are evaluating the continuous time scheme at the integer time n, in this sense t and n should be thought of as interchangeable and which one we use merely distinguishes continuous and discrete time. Let ϕ be the law of a standard normal random variable. Convergence to ϕ will always be in distribution as t (or equivalently, n) goes to infinity.

Theorem 18. Let U_n, U_t be as above. If $\mathbb{V}[|U_t|] \to \infty$ then,

$$\frac{|U_t| - \mathbb{E}[|U_t|]}{\sqrt{\mathbb{V}[|U_t|]}} \to \phi,$$
$$\frac{|U_n| - \mathbb{E}[|U_n|]}{\sqrt{\mathbb{V}[|U_t|]}} \to \phi.$$

Proof. If $\sum_j \lambda_j = 1$ this is precisely Theorems 1 and 2 of [9]. The case $\sum_j \lambda_j \neq 1$ is handled by a simple scaling argument.

Asymptotic Gaussianity of the number of edges is a corollary of Theorem 18 by defining an urn for each edge. In fact, one can and Janson [1] indeed does even obtain a local limit theorem from [8]. However, we will settle for asymptotic Gaussianity as, for one, we do not know how to obtain rate estimates from the coupling.

Conjecture 19. ([1], Problem 6.9). Let $\mu \in \mathcal{P}(\mathbb{N}_2)$ be such that $\mathbb{V}[|V_n|] \to \infty$, then $|V_n|$ is asymptotically Gaussian.

Janson only conjectures the result in discrete time we will also consider continuous time. As Janson pointed out "The number of vertices corresponds to an occupancy problem where balls are thrown in pairs, with a dependency inside each pair" [1]. Thus the indicators of the vertices are dependent. However, we will show that actually one can couple to the standard urn scheme, if one assumes eventual forever connectedness. We begin with some special cases where the asymptotic Gaussianity of the number of vertices is trivial.

Proposition 20. Let the support of μ consist only of isolated edges and assume that $\mathbb{V}[|V_t|] \to \infty$ or equivalently $\mathbb{V}[|E_t|] \to \infty$. Then

$$\frac{|V_t| - \mathbb{E}[|V_t|]}{\sqrt{\mathbb{V}[|V_t|]}} \to \phi.$$

Proof. Simply noting that the number of vertices is just twice the number of edges gives

$$\frac{|V_t| - \mathbb{E}[|V_t|]}{\sqrt{\mathbb{V}[|V_t|]}} = \frac{\frac{1}{2}|E_t| - \frac{1}{2}\mathbb{E}[|E_t|]}{\sqrt{\frac{1}{4}\mathbb{V}[|E_t|]}} \to \phi,$$

by Theorem 18.

In defining our models we insisted on edges being between distinct vertices, however, one may allow for self-loops. In this extended context the following definition makes sense.

Definition 21. We call $\mu \in \mathcal{P}(\{\{i, j\} | i, j \in \mathbb{N}\})$ first rank if there exists a $\sigma \in \mathcal{P}(\mathbb{N})$ such that $\mu(\{i, j\}) = \sigma(\{i\})\sigma(\{j\})$ for all $i, j \in \mathbb{N}$.

Proposition 22. Let μ be first rank and be such that $\mathbb{V}[|V_t|] \to \infty$ then

$$\frac{|V_n| - \mathbb{E}[|V_n|]}{\sqrt{\mathbb{V}[V_t]}} \to \phi,$$
$$\frac{|V_t| - \mathbb{E}[|V_t|]}{\sqrt{\mathbb{V}[|V_t|]}} \to \phi.$$

Proof. Define an urn for each vertex, setting $\lambda_i = M_i$. These urns are independent as a consequence of the first rank assumption. Hence the result follows from Theorem 18.

Remark 23. In some sense Propositions 20 and 22 are opposite extremes. In the former there is perfect dependence between the vertices in each edge, in the latter there is perfect independence but asymptotic Gaussianity holds all the same. This seems to us like good evidence for Conjecture 19.

Theorem 24. Let G_t be eventually forever connected and suppose that $\mathbb{V}[|U_t|] \to \infty$. Then the following hold.

$$\begin{split} \frac{|V_n| - \mathbb{E}[|V_n|]}{\sqrt{\mathbb{V}[|U_t|]}} &\to \phi, \\ \frac{|V_t| - \mathbb{E}[|V_t|]}{\sqrt{\mathbb{V}[|U_t|]}} \to \phi. \end{split}$$

We begin by sketching the proof. We will couple to the urn scheme with intensities $\lambda_j = M_j$. By choosing an appropriate coupling, vertices will be added to V_t but not U_t only when a pair of new vertices are simultaneously added to V_t . Since we know this happens only finitely many times in the connected regime, this will not cause us any trouble. We will need to fill some extra urns in U_t but no more often than the doubles in V_t . Thus there will only be finitely many vertices that are ever in one but not the other. Eventually these vertices will be present in both and from that point onward U_t and V_t will be equal. The coupling inequality will then complete the proof.

Proof. We define a coupling $(\tilde{V}_t, \tilde{U}_t)$ of U_t and V_t . Repeatedly wait an exp(3) amount of time then do the following

For every $i \in \mathbb{N} \setminus \tilde{U}_t$ compute

$$\lambda^{(i)}(\tilde{V}_t, \tilde{U}_t) := \begin{cases} \frac{1}{2} \sum_{j \in \overline{\tilde{V}_t} \land \overline{\tilde{U}_t}} \mu_{ij} & \text{if } i \notin \tilde{V}_t \\ \frac{1}{2} \sum_{j \in \overline{\tilde{U}}_t \land \tilde{V}_t} \mu_{ij} + \sum_{j \in \overline{\tilde{U}}_t \land \overline{\tilde{V}}_t} \mu_{ij} & \text{if } i \in \tilde{V}_t \end{cases}$$

Then generate a $\mathbb{N} \cup \mathbb{N}_2 \cup \{\emptyset\}$ -valued random variable χ by generating $\eta \sim$ Uniform[0, 1] independent of everything else and thresholding appropriately to obtain:

- $P(\chi = i) = \frac{\lambda^{(i)}}{3}$ for all $i \in \mathbb{N}$.
- $P(\chi = e) = \frac{\mu_e}{3}$ for all $e \in \mathbb{N}_2$.
- $P(\chi = \emptyset) = \frac{2}{3} \frac{1}{3} \sum_i \lambda^{(i)}$.

Then increment the processes according to

- If $\chi = \emptyset$: pass
- If $\chi = i$: Add *i* to \tilde{U}_t .

- If $\chi = \{i, j\}$ for some $\{i, j\} \in \mathbb{N}_2$,
 - If $|\{i, j\} \setminus \tilde{V}_t| = 0$:
 - * If $|\{i, j\} \setminus \tilde{U}_t| = 0$: pass.
 - * If $|\{i, j\} \setminus \tilde{U}_t| = 1$: add $\{i, j\} \setminus \tilde{U}_t$ to \tilde{U}_t .
 - * If $|\{i, j\} \setminus \tilde{U}_t| = 2$: Choose *i* or *j* according to a coin toss, independent of everything else, and add that one to \tilde{U}_t .
 - If $|\{i, j\} \setminus \tilde{V}_t| = 1$: Let $x \in \{i, j\} \setminus \tilde{V}_t$ be that one element and let $y \in \{i, j\} \cap \tilde{V}_t$ be the other. Then:
 - * If $x \notin \tilde{U}_t$: Add x to \tilde{U}_t .
 - * If $x \in \tilde{U}_t$ and $y \notin \tilde{U}_t$: add y to \tilde{U}_t .
 - * If $x \in \tilde{U}_t$ and $y \in \tilde{U}_t$: pass.

$$- If |\{i, j\} \setminus V_t| = 2$$

- * If $|\{i, j\} \setminus \tilde{U}_t| = 0$: pass.
- * If $|\{i, j\} \setminus \tilde{U}_t| = 1$: add $\{i, j\} \setminus \tilde{U}_t$ to \tilde{U}_t .
- * If $|\{i, j\} \setminus \tilde{U}_t| = 2$: Choose *i* or *j* according to a coin toss, independent of everything else, and add that one to \tilde{U}_t .
- Add $\{i, j\} \setminus \tilde{V}_t$ to \tilde{V}_t

To ensure that the above construction is well-defined we need to ensure that χ is well-defined. To see this note that $\lambda^{(i)} \leq \sum_{j} \mu_{ij} = M_i$ and that $\sum_{i} M_i = 2$, so that therefore

$$P(\chi = \emptyset) = \frac{2}{3} - \frac{1}{3} \sum_{i} \lambda^{(i)} \ge 0.$$

Note also that

$$\sum_{i} P(\chi = i) + \sum_{e} P(\chi = e) + P(\chi = \emptyset) = \sum_{i} \frac{\lambda^{(i)}}{3} + \frac{1}{3} + \frac{2}{3} - \sum_{i} \frac{\lambda^{(i)}}{3} = 1.$$

From the thinning property of Poisson processes and the third bullet point in the increment rule we see that \tilde{V}_t has the same distribution as V_t .

Let p be the probability that $i \notin \tilde{U}_t$ is added to \tilde{U}_t after a particular exp(3) wait. By summing over the probabilities of all the different ways i can be added to \tilde{U}_t this can be seen to be given by

$$\begin{split} 3p =& 1_{i\notin \tilde{V}_{t}} \frac{1}{2} \sum_{j\in \overline{\tilde{V}_{t}} \wedge \overline{\tilde{U}_{t}}} \mu_{ij} + 1_{i\in \tilde{V}_{t}} \frac{1}{2} \sum_{j\in \tilde{V}_{t} \wedge \overline{\tilde{U}_{t}}} \mu_{ij} + 1_{i\in \tilde{V}_{t}} \sum_{j\in \overline{\tilde{V}_{t}} \wedge \overline{\tilde{U}_{t}}} \mu_{ij} \\ &+ 1_{i\in \tilde{V}_{t}} \frac{1}{2} \sum_{j\in \overline{\tilde{V}_{t}} \wedge \overline{\tilde{U}_{t}}} \mu_{ij} + 1_{i\in \tilde{V}_{t}} \sum_{j\in \overline{\tilde{V}_{t}} \wedge \tilde{U}_{t}} \mu_{ij} + 1_{i\notin \tilde{V}_{t}} \sum_{j\in \overline{\tilde{V}_{t}} \wedge \overline{\tilde{U}_{t}}} \mu_{ij} \\ &+ 1_{i\in \tilde{V}_{t}} \sum_{j\in \overline{\tilde{V}_{t}} \wedge \tilde{U}_{t}} + 1_{i\notin \tilde{V}_{t}} \sum_{j\in \overline{\tilde{V}_{t}} \wedge \tilde{U}_{t}} + 1_{i\notin \tilde{V}_{t}} \frac{1}{2} \sum_{j\in \overline{\tilde{V}_{t}} \wedge \overline{\tilde{U}_{t}}} \mu_{ij} \\ &= 1_{i\in \tilde{V}_{t}} \sum_{j} \mu_{ij} + 1_{i\notin \tilde{V}_{t}} \sum_{j} \mu_{ij}, \\ &= M_{i}. \end{split}$$

Note that this does not depend on t, \tilde{V}_t and \tilde{U}_t . Moreover, the η are independent, as are the exp(3) waiting times, so looking only at \tilde{U}_t we observe an urn filling process, that is to say the coupling has the correct second marginal and so constitutes a valid coupling. Therefore $(|\tilde{U}_t|, |\tilde{V}_t|)$ is a valid coupling of $|V_t|, |U_t|$.

Next we show that $\tilde{V}_t = \tilde{U}_t$ eventually forever by showing eventual forever containment both ways. We begin by showing that $\tilde{V}_t \subset \tilde{U}_t$ eventually forever. First, notice that due to how we have set up the coupling the only way for a vertex v to be added to $\tilde{V}_t \setminus \tilde{U}_t$ is by an edge χ adding both vand another vertex to \tilde{V}_t . Since this happens only finitely many times, by the connectedness assumption, eventually, it will cease to happen and, since those vertices will eventually appear in \tilde{U}_t , they will no longer be in $\tilde{V}_t \setminus \tilde{U}_t$. Thus, eventually forever $\tilde{V}_t \subset \tilde{U}_t$. Next we show $\tilde{U}_t \subset \tilde{V}_t$ eventually forever. Immediately following an exp(3) wait, the probability that an urn is filled in \tilde{U}_t without the corresponding vertex being added to or already being in \tilde{V}_t is $\frac{1}{6} \sum_{u \notin \tilde{V}_t} \sum_{v \in \tilde{U}_t \wedge \tilde{V}_t} \mu_{uv}$. This is because we need a $\chi = i$ event where $i \notin \tilde{V}_t$. Now observe that after one of the exp(3) waits the probability of adding two new vertices to \tilde{V}_t at the same time is $\frac{1}{3} \sum_{i,j \in \tilde{V}_t} \mu_{ij}$ and that

$$\frac{1}{6} \sum_{i \notin \tilde{V}_t} \sum_{j \in \overline{\tilde{U}}_t \land \overline{\tilde{V}_t}} \mu_{ij} \le \frac{1}{3} \sum_{i,j \in \overline{\tilde{V}}_t} \mu_{ij}.$$
(4)

Since the η are independent, (4) is sufficient to see that the distribution of the number of positive integers are ever in $\tilde{U}_t \setminus \tilde{V}_t$ is dominated by the number of vertices that appear at the same time as another vertex in \tilde{V}_t . Since the latter is a.s. finite, so is the former. This is perhaps easier to see in terms of processes, in which case it corresponds to the obvious fact that if a Cox-process only ever has finitely many events and another process has a lower intensity, the so does the second. Since any vertices added to $\tilde{U}_t \setminus \tilde{V}_t$ eventually leave, one has $\tilde{U}_t \setminus \tilde{V}_t = \emptyset$ eventually forever. Thus we have two way containment and $\tilde{V}_t = \tilde{U}_t$ eventually forever. In particular $|\tilde{V}_t| = |\tilde{U}_t|$ eventually forever almost surely.

By the coupling inequality and since almost sure convergence is stronger than convergence in probability it follows that

$$d_{TV}(\mathcal{L}[V_t], \mathcal{L}[U_t]) \le P(|\tilde{V}_t| \ne |\tilde{U}_t|) \to 0,$$
(5)

where \mathcal{L} denotes the law of a random variable. Now combining the fact that the Levy-Prokhov distance, which we denote by π , metrizes convergence in distribution [23], that the total variation distance upper bounds the Levy-Prokhov distance [23], the fact that the total variation distance between laws of random variables is invariant to shifting and rescaling the random variables, equation (5) and Theorem 18 will complete the proof in the continuous time case.

$$\begin{aligned} \pi \left(\mathcal{L} \left[\frac{|V_t| - \mathbb{E}[|V_t|]}{\sqrt{\mathbb{V}[|U_t|]}} \right], \phi \right) \\ &\leq \pi \left(\mathcal{L} \left[\frac{|V_t| - \mathbb{E}[|V_t|]}{\sqrt{\mathbb{V}[|U_t|]}} \right], \mathcal{L} \left[\frac{|U_t| - \mathbb{E}[|U_t|]}{\sqrt{\mathbb{V}[|U_t|]}} \right] \right) + \pi \left(\mathcal{L} \left[\frac{|U_t| - \mathbb{E}[|U_t|]}{\sqrt{\mathbb{V}[|U_t|]}} \right], \phi \right) \\ &\leq d_{TV} \left(\mathcal{L} \left[\frac{|V_t| - \mathbb{E}[|V_t|]}{\sqrt{\mathbb{V}[|U_t|]}} \right], \mathcal{L} \left[\frac{|U_t| - \mathbb{E}[|U_t|]}{\sqrt{\mathbb{V}[|U_t|]}} \right] \right) + \pi \left(\mathcal{L} \left[\frac{|U_t| - \mathbb{E}[|U_t|]}{\sqrt{\mathbb{V}[|U_t|]}} \right], \phi \right) \\ &= d_{TV} \left(\mathcal{L}[|V_t|], \mathcal{L}[|U_t|] \right) + \pi \left(\mathcal{L} \left[\frac{|U_t| - \mathbb{E}[|U_t|]}{\sqrt{\mathbb{V}[|U_t|]}} \right], \phi \right) \to 0. \end{aligned}$$

To de-Poissonize we use the exact same argument as in Theorem 2 of [9], which we avoid repeating here. One needs to verify that the proofs of Lemmas 1,3,4,5,6 and Theorem 2 in [9] all work in our setting, in particular that the sum of the intensities being two and the vertex presence indicators not being independent does not cause any issues. We spare the reader the

details, but morally this works because linearity of expectation does not need independence and that therefore $\mathbb{E}[|V_t|] = \sum_i 1 - e^{-M_i t} = \mathbb{E}[|U_t|]$. The fact that we are in any case normalizing by $\mathbb{V}[|U_t|]$ and not $\mathbb{V}[|V_t|]$ means that the fact that $\mathbb{V}[|V_t|] \neq \mathbb{V}[|U_t|] = \sum_i e^{-M_i t} (1 - e^{-M_i t})$ does not pose any issues. \Box

Remark 25. We interpret Corollary 15 as suggesting that the assumption of eventual forever connectedness is not very restrictive as in this case it corresponds to σ in Definition 21 having sub-quadratic tails, something which in general is comparable to having a finite mean.

Remark 26. Note that both 20 and 22 allow μ which produce graphs that are not eventually forever connected, so eventual forever connectedness is certainly not necessary for asymptotic Gaussianity of the vertex count.

Remark 27. In the proof of Theorem 24 and elsewhere we only make use of there being finitely many I_e which we have seen is equivalent to $\sum_e \frac{\mu_e}{M_e} < \infty$ so one could give a theorem with this assumption instead, but we prefer our version as it is more immediately interpretable.

Remark 28. Without the assumption of eventual forever connectedness we do not know how to obtain something like (5). As a next step towards proving Conjecture 19 one could imagine wanting to generalize our result to cover μ where vertices added by events of the type I_e are asymptotically a vanishingly small fraction of all vertices. Indeed, if we think of vertices added in such a way as contributing an error term then in this case this error term should not be enough to destroy the Gaussianity. However, we do not know how to push our techniques to cover this setting as the coupling inequality is rather demanding, requiring exact equality. Proving Conjecture 19 in full generality will likely require a rather different approach.

In the proof of the Theorem 24 the scaling that naturally emerges is that involving the urn scheme. This variance is convenient because it is straightforward to compute. However, one may desire a theorem statement which is in terms of only the vertex process. We will allow this by showing that in the connected regime the two variances are within a constant and hence asymptotically equivalent.

Proposition 29.

$$\mathbb{V}[|U_t|] \le \mathbb{V}[|V_t|] \le \mathbb{V}[|U_t|] + \sum_e \frac{\mu_e}{M_e}.$$

Proof. Let X_i be the event that $i \in V_t$ so that $|V_t| = \sum_i 1_{X_i}$.

$$\mathbb{V}[|V_t|] = \mathbb{V}\left[\sum_i 1_{X_i>0}\right],$$

= $\sum_i \mathbb{V}[1_{X_i}] + \sum_{i\neq j} P(X_i \wedge X_j) - P(X_i)P(X_j),$
= $\mathbb{V}[|U_t|] + \sum_{i\neq j} P(X_i \wedge X_j) - P(X_i)P(X_j),$

where in the last equality we recalled that the urns are independent and that $P(i \in V_t) = 1 - e^{-M_i t} = P(i \in U_t)$. For compactness of notation, let $M_{i\setminus j} := M_i - \mu_{ij}$.

$$P(X_i \wedge X_j) = P(X_i \wedge X_j | \tau_{ij} \le t) P(\tau_{ij} \le t) + P(X_i \wedge X_j | \tau_{ij} > t) P(\tau_{ij} > t),$$

= 1 - e^{-\mu_{ij}t} + (1 - e^{-M_{i\backslash j}t})(1 - e^{-M_{j\backslash i}t})e^{-\mu_{ij}t},
= 1 - e^{-M_it} - e^{-M_jt} + e^{-M_{ij}t}.

Since

$$P(X_i)P(X_j) = (1 - e^{-M_i t})(1 - e^{M_j t}),$$

straightforward algebra gives

$$P(X_i \wedge X_j) - P(X_i)P(X_j) = e^{-M_{ij}t}(1 - e^{-\mu_{ij}t}).$$

We note that this is non-negative and apply Lemma 40 to obtain

$$0 \le P(X_i \land X_j) - P(X_i)P(X_j) < \frac{\mu_{ij}}{M_{ij}},$$

which completes the proof.

Corollary 30. If μ is such that G_t is eventually forever connected and at least one of $\mathbb{V}[|U_t|]$ and $\mathbb{V}[|V_t|]$ goes to infinity, then so does the other and

$$\frac{\mathbb{V}[|V_t|]}{\mathbb{V}[|U_t|]} \to 1.$$

Proof. By Theorem 4 eventual forever connectedness implies that $\sum_{e} \frac{\mu_e}{M_e} < \infty$. The result then follows from Proposition 29 by dividing by $\mathbb{V}[|U_t|]$ and taking limits.

Remark 31. The proof of Lemma 29 gives some more intuition for why one might expect the connectedness assumption to help with showing Gaussianity. Namely, the calculations in the proof show that $\frac{\mu_{ij}}{M_{ij}}$, the quantity we have already seen is $P(I_{\{i,j\}})$ also serves as an upper bound on the amount of dependence, as measured by the covariance, between the indicators of the vertices i and j.

6 Completeness

It is not possible for the graphs to be eventually forever complete. This is because when a new vertex is added the graph is necessarily not complete and new vertices are added infinitely often, assuming μ has unbounded support. The closest one could hope to get to eventual forever completeness is captured in Definition 32.

Definition 32. We say that a graph G = (V, E) is **essentially complete** iff it is connected and there is an $n \in \mathbb{N}$ such that $V = \{1, \ldots, n, n+1\}$ and the induced subgraph obtained by taking the vertex subset $\{1, \ldots, n\}$ is the complete graph on $\{1, \ldots, n\}$.

We illustrate this definition by giving some examples and non-examples in Figure 2.

Remark 33. Clearly eventual forever essential completeness is a much stronger notion than eventual forever connectedness.

In our terminology, Janson [1] showed the following.

Proposition 34. ([1], Example 8.1) If for all $k \ge 2$,

$$0 < \sup_{l} \mu_{k+1,l} \le k^{-4} \min_{i < k} \mu_{k,i},$$

then G_t is eventually forever essentially complete. Moreover, $\mu_{ij} \propto (\max(i, j)!)^{-4}$ and $\mu_{ij} \propto e^{-3^i} e^{-3^j}$ satisfy this assumption.

Janson [1] left it as an open problem to find a necessary and sufficient condition for eventual forever essential completeness. We solve this problem.



Figure 2: Examples and non-examples of essentially complete graphs. The first two are not essentially complete as they fail the induced subgraph condition. The former is missing edges, the latter a vertex (and its associated edges). The third is an essentially complete graph because the subgraph induced by vertices 1, 2, 3, 4 is the complete graph on four vertices, the fifth vertex is the only extra vertex, and that vertex is not isolated.

Definition 35. Let $\mathcal{F} = {\tau_i}_{i \in I}$ be a countable family of random variables, interpreted as arrival times. Let $C = {C_n}_{n=1}^{\infty}$ be a partition of I into finite sets. Let $S_n = {\tau_i}_{i \in C_n}$. We define the event that \mathcal{F} respects C to be

$$\bigcup_{N=1}^{\infty} \cap_{n=N}^{\infty} \{ \check{S}_n < \hat{S}_{n+1} \}.$$

Note that even if the τ_i are independent exponential random variables the events $\{\check{S}_n < \hat{S}_{n+1}\}, \{\check{S}_m < \hat{S}_{m+1}\}$ are not independent. However, by using the chain rule of probability and some conditioning we can compute the probability that a family of exponential random variables respects a partitioning. Let $F_n \subset I$ be the (random) set of indices of the first *n* arrivals, let $K_n = \bigcup_{m=1}^{n-1} C_m, L_n := I \setminus K_n$. Note that K_n and are L_n deterministic.

Lemma 36. Let $\mathcal{F} = {\tau_i}_{i \in I}$ be a countable family of independent exponential random variables indexed by I with intensities ${\lambda_i}_{i \in I}$. Let C be a partition of I into finite sets. Then \mathcal{F} respects C with probability one if

$$\lim_{N \to \infty} \prod_{n=1}^{N} \int_{0}^{\infty} \prod_{i \in C_n} (1 - e^{-\lambda_i t}) \left(\sum_{i \in L_n} \lambda_i \right) e^{-\sum_{i \in L_n} \lambda_i t} dt > 0,$$

else \mathcal{F} respects C with probability zero.

Proof. Let R be the event that \mathcal{F} respects C. Notice that the exponential random variables are independent and that changing finitely many of them cannot change whether R occurs. Therefore Kolmogorov's 0-1 law applies. We calculate,

$$P(R) = \lim_{N \to \infty} P(\bigcap_{n=N}^{\infty} \{\check{S}_n < \hat{S}_{n+1}\}),$$

= $\lim_{N \to \infty} P(\bigcap_{n=N}^{\infty} \{\check{S}_n < \hat{S}_{n+1}\} \cap \{F_{|K_N|} = K_N\}),$
= $\lim_{N \to \infty} (P(\bigcap_{n=N}^{\infty} \{\check{S}_n < \hat{S}_{n+1}\} | \{F_{|K_N|} = K_N\}))$
 $\times P(\{F_{|K_N|} = K_N\})).$

Since probabilities are at most one, if the limit of either factor is not one, then the limit of the product is not one. By Kolmogorov's 0-1 law P(R) = 0 in that case. We turn our attention to the the first factor and exploit memorylessness of exponential random variables.

$$\lim_{N \to \infty} P(\bigcap_{n=N}^{\infty} \{\check{S}_n < \hat{S}_{n+1}\} | \{F_{|K_N|} = K_N\}),$$

$$= \lim_{N \to \infty} P(\bigcap_{n=N}^{\infty} \{\check{S}_n < \hat{L}_n\} | \{F_{|K_N|} = K_N\}),$$

$$= \lim_{N \to \infty} \prod_{n=N}^{\infty} P(\{\check{S}_n < \hat{L}_n\} | \{F_{|K_N|} = K_N\}, \bigcap_{m=N}^{n-1} \{\check{S}_m < \hat{L}_m\}),$$

$$= \lim_{N \to \infty} \prod_{n=N}^{\infty} P(\{\check{S}_n < \hat{L}_n\} | \{F_{|K_n|} = K_n\}),$$

$$= \lim_{N \to \infty} \prod_{n=N}^{\infty} \int_0^{\infty} \prod_{i \in C_n} (1 - e^{-\lambda_i t}) \left(\sum_{i \in L_n} \lambda_i\right) e^{-\sum_{i \in L_n} \lambda_i t} dt.$$

It is a standard property of infinite products of strictly positive terms that the tail converging to one is equivalent to the full product converging to a non-zero value. Thus we have as a necessary condition,

$$\lim_{N \to \infty} \prod_{n=1}^{N} \int_{0}^{\infty} \prod_{i \in C_n} (1 - e^{-\lambda_i t}) \left(\sum_{i \in L_n} \lambda_i \right) e^{-\sum_{i \in L_n} \lambda_i t} dt > 0.$$

It remains to show that this condition is sufficient. Returning to the

second factor,

$$P(\{F_{|K_N|} = K_N\}) = P(\{\check{\tau_i}\}_{i \in K_N} < \{\check{\tau_i}\}_{i \in L_N}),$$

$$= \int_0^\infty \prod_{i \in K_N} (1 - e^{-\lambda_i t}) \left(\sum_{i \in L_N} \lambda_i\right) e^{-\sum_{i \in L_N} \lambda_i t} dt,$$

$$= \int_0^\infty \prod_{n=1}^N \prod_{i \in C_n} (1 - e^{-\lambda_i t}) \left(\sum_{i \in L_N} \lambda_i\right) e^{-\sum_{i \in L_N} \lambda_i t} dt$$

Notice that $d\nu_N := \left(\sum_{i \in L_N} \lambda_i\right) e^{-\sum_{i \in L_N} \lambda_i t} dt$ is a probability measure. Note also that $\prod_{i \in C_n} 1 - e^{-\lambda_i t}$ is monotonically increasing for every n as it is a product of monotonically increasing functions. By Chebyshev's other inequality [24] we therefore have

$$\int_0^\infty \prod_{n=1}^N \prod_{i \in C_n} (1 - e^{-\lambda_i t}) d\nu_N \ge \prod_{n=1}^N \int_0^\infty \prod_{i \in C_n} (1 - e^{-\lambda_i t}) d\nu_N.$$

Recalling that we had already assumed the limit of the right hand side as $N \to \infty$ to be strictly positive we see that the same must be true of the left side. This completes the proof.

The following result regarding the urn schemes is a simple corollary of Lemma 36.

Proposition 37. Consider a discrete or continuous time urn scheme with probabilities or intensities (respectively) $\{\lambda_i\}_{i \in \mathbb{N}}$. Then the urns are eventually forever filled in order if and only if

$$\prod_{i=1}^{\infty} \frac{\lambda_i}{\sum_{j \ge i} \lambda_j} > 0.$$
(6)

Proof. By a de-Poissonization argument similar to the proof of Lemma 5 it suffices to consider the continuous time case. The stated event is equivalent to the family \mathcal{F} with intensities $\{\lambda_i\}_{i\in\mathbb{N}}$, (indexed by $I = \mathbb{N}$) respects $C = \{C_n = \{n\}\}_{n\in\mathbb{N}}$. We check the condition of Lemma 36, which essentially amounts to showing the standard fact that the probability that a particular exponential random variable is smallest in its family is the ratio of its intensity to the

sum of the intensities.

$$\lim_{N \to \infty} \prod_{n=1}^{N} \int_{0}^{\infty} \prod_{\lambda \in C_{n}} (1 - e^{-\lambda t}) \left(\sum_{\lambda \in L_{n}} \lambda\right) e^{-\sum_{\lambda \in L_{n}} \lambda t} dt,$$
$$= \lim_{N \to \infty} \prod_{n=1}^{N} \int_{0}^{\infty} (1 - e^{-\lambda_{n} t}) \left(\sum_{m=n+1}^{\infty} \lambda_{m}\right) e^{-\sum_{m=n+1}^{\infty} \lambda_{m} t} dt,$$
$$= \lim_{N \to \infty} \prod_{n=1}^{N} \frac{\lambda_{n}}{\sum_{m=n}^{\infty} \lambda_{m}}.$$

Remark 38. Note that if the λ :s decay geometrically then the factors we take the product over in (6) are all equal and the limit is zero, so intuitively, the condition may be interpreted as measuring how much faster than geometrically the decay is occurring.

We turn to the main motivation behind Lemma 36. Namely, to answer Janson's question about completeness.

Proposition 39. If μ is such that

$$\lim_{N \to \infty} \prod_{n=1}^{N} \int_{0}^{\infty} \prod_{\{i,j\} \in \mathbb{N}_{2}: \max(i,j) = n} \left(1 - e^{-\mu_{ij}t} \right) d\nu_{N} > 0,$$

where

$$d\nu_N := \left(\sum_{\{i,j\}\in\mathbb{N}_2:\max(i,j)>n}\mu_{ij}\right)e^{-\sum_{\{i,j\}\in\mathbb{N}_2:\max(i,j)>n}\mu_{ij}t}dt,$$

then G_t and G_n are almost surely eventually forever essentially complete. Else G_t and G_n are almost surely not eventually forever essentially complete.

Proof. By Lemma 5 it suffices to consider G_t . With a bit of thought one sees that G_t is eventually forever essentially complete if and only if \mathcal{F} , indexed by $I = \mathbb{N}_2$ respects $C = \{C_n = \{\{i, j\} \in \mathbb{N}_2 : \max(i, j) = n\}\}_{n \in \mathbb{N}}$. Applying Lemma 36 completes the proof.

7 Conclusions and Future Directions

If Janson's paper is a "second step" [1] in understanding edge exchangeable graphs, then the present paper may be considered a third. However, much work still remains. We begin with a discussion of the state of Conjecture 19. As we note in Remark 25, Theorem 24 resolves Conjecture 19 under a fairly mild extra assumption. However, as we observe in Remark 23 it really seems that it should be possible to do away with this assumption. However, as discussed in Remark 28 getting rid of this assumption is likely to require new ideas. It would also be desirable to give a convergence rate in the connected regime.

As we discussed, it would be interesting to characterize not only eventual forever connectedness but also the marginal notion, sharpening Corollary 17 as well as more geometric notions like the size of the largest component. This seems hard for a general μ , but may be tractable in the first rank case. Possibly the connections to the Norros-Reittu model discussed in Section 1.5.2 could be exploited since the geometry of the Norros-Reittu model has been studied with much success [25].

The connection to inhomogeneous random graphs discussed in Subsection 1.5.2, along with the fact that concentration results for the graph Laplacian exist for such graphs [26][27][28] suggest looking for a more quantitative connectedness-type result than Theorem 14 in terms of the spectral gap. A closely related open problem is "Spectral gap of Bayesian graph Laplacian" in Aldous' collection of open problems [29]. That problem arises in [14] and is closely related to the graph inference problem discussed in Subsection 1.5.2. Other related issue are understanding clustering, dependence [30] and the range of deg distributions that are possible [1]. In short, there are many interesting probabilistic questions left to answer regarding edge exchangeable random graphs. We are hopeful that the study of such questions and other questions regarding edge exchangeable random graphs will continue to foster fruitful interactions between probabilists, statisticians and practitioners.

References

 Svante Janson. 'On Edge Exchangeable Random Graphs'. In: Journal of Statistical Physics 173 (June 2017), pp. 448–484. DOI: 10.1007/s10955-017-1832-9.

- Harry Crane and Walter Dempsey. Edge exchangeable models for network data. arXiv.org, Oct. 2016. DOI: 10.48550/arXiv.1603.04571.
 URL: https://arxiv.org/abs/1603.04571.
- [3] Harry Crane and Walter Dempsey. 'Edge Exchangeable Models for Interaction Networks'. In: Journal of the American Statistical Association 113 (June 2018), pp. 1311–1326. DOI: 10.1080/01621459.2017.1341413. (Visited on 16/01/2025).
- [4] Diana Cai, Trevor Campbell and Tamara Broderick. Edge-exchangeable graphs and sparsity. Neural Information Processing Systems, 2016. URL: https://papers.nips.cc/paper_files/paper/2016/hash/1a0a283bfe7c549dee6c638a05 (visited on 05/08/2024).
- [5] Rui Luo, Buddhika Nettasinghe and Vikram Krishnamurthy. 'Anomalous Edge Detection in Edge Exchangeable Social Network Models'. In: *PMLR* 204 (Aug. 2023), pp. 287–310. URL: https://proceedings.mlr.press/v204/luo23a. (visited on 04/12/2024).
- [6] Xinglong Li and Trevor Campbell. 'Truncated simulation and inference in edge-exchangeable networks'. In: *Electronic Journal of Statistics* 15 (Jan. 2021). DOI: 10.1214/21-ejs1916. (Visited on 04/12/2024).
- [7] Anna Ben-Hamou, Stéphane Boucheron and Mesrob I. Ohannessian.
 'Concentration inequalities in the infinite urn scheme for occupancy counts and the missing mass, with applications'. In: *Bernoulli* 23 (Feb. 2017), pp. 249–287. DOI: 10.3150/15-bej743. URL: https://www.jstor.org/stable/44075 (visited on 08/10/2024).
- [8] Hsien-Kuei Hwang and Svante Janson. 'Local limit theorems for finite and infinite urn models'. In: *The Annals of Probability* 36 (May 2008), pp. 992–1022. DOI: 10.1214/07-aop350. (Visited on 19/01/2020).
- [9] Michael Dutko. 'Central Limit Theorems for Infinite Urn Models'. In: The Annals of Probability 17 (July 1989), pp. 1255–1263. DOI: 10.1214/aop/1176991268. (Visited on 15/06/2020).
- [10] SAMUEL KARLIN. 'Central Limit Theorems for Certain Infinite Urn Schemes'. In: Journal of Mathematics and Mechanics 17 (1967), pp. 373– 401. DOI: 10.2307/24902077. URL: https://www.jstor.org/stable/24902077.
- [11] Daniel M. Busiello et al. 'Explorability and the origin of network sparsity in living systems'. In: Scientific Reports 7 (Sept. 2017). DOI: 10.1038/s41598-017-12521-1. (Visited on 16/01/2025).

- [12] Harry Crane. Probabilistic Foundations of Statistical Network Analysis. CRC Press, Apr. 2018.
- [13] François Caron and Emily B. Fox. 'Sparse Graphs Using Exchangeable Random Measures'. In: Journal of the Royal Statistical Society Series B: Statistical Methodology 79 (Nov. 2017), pp. 1295–1366. DOI: https://doi.org/10.1111/rssb.12233. URL: https://academic.oup.com/jrsssb/articl (visited on 30/04/2024).
- [14] David Aldous and Xiang Li. 'A Framework for Imperfectly Observed Networks'. In: *Journal of Statistical Physics* 173 (Oct. 2021), pp. 1303– 1320. DOI: 10.1007/s10955-017-1838-3. (Visited on 05/08/2024).
- [15] Xiang Li. 'Inference on Graphs: From Probability Methods to Deep Neural Networks'. PhD thesis. 2017. URL: https://escholarship.org/content/qt29k2m4p2 (visited on 04/12/2024).
- [16] David J. Aldous. 'Weak Concentration for First Passage Percolation Times on Graphs and General Increasing Set-valued Processes'. In: *Latin American Journal of Probability and Mathematical Statistics* 13 (2016), p. 925. DOI: 10.30757/alea.v13-35. URL: https://arxiv.org/pdf/1604.06418 (visited on 01/08/2024).
- [17] David Aldous. 'The incipient giant component in bond percolation on general finite weighted graphs'. In: *Electronic Communications in Probability* 21 (Feb. 2019), pp. 1–9. DOI: 10.1214/16-ecp21. (Visited on 05/08/2024).
- [18] Avanti Athreya et al. 'Statistical inference on random dot product graphs: a survey'. In: Journal of Machine Learning Research 18 (Sept. 2017), pp. 1–92. URL: http://jmlr.org/papers/v18/17-448.html (visited on 07/10/2024).
- [19] Ilkka Norros and Hannu Reittu. 'On a conditionally Poissonian graph process'. In: Advances in Applied Probability 38 (Mar. 2006), pp. 59-75.
 DOI: 10.1239/aap/1143936140. URL: https://www.cambridge.org/core/services/aop-ca (visited on 27/08/2024).
- [20] Jia-An Yan. 'A Simple Proof of Two Generalized Borel-Cantelli Lemmas'. In: Springer eBooks (Oct. 2006), pp. 77–79. DOI: 10.1007/978-3-540-35513-7_7. (Visited on 07/10/2024).

- [21] M. Mursaleen and S A Mohiuddine. 'Double Series and Convergence Tests'. In: Springer eBooks (Oct. 2013), pp. 149–166. DOI: 10.1007/978-81-322-1611-7_9. (Visited on 08/10/2024).
- [22] double. Convergence of double sum $\sum_{m,n} \frac{1}{m^p + n^k}$. Mathematics Stack Exchange, Oct. 2013. URL: https://math.stackexchange.com/questions/520854/converge (visited on 08/10/2024).
- [23] Alison L. Gibbs and Francis Edward Su. 'On Choosing and Bounding Probability Metrics'. In: International Statistical Review / Revue Internationale de Statistique 70 (Dec. 2002), pp. 419–435. DOI: 10.2307/1403865. URL: https://www.jstor.org/stable/1403865 (visited on 05/12/2024).
- [24] A.M. Fink and Max Jodeit. 'On Chebyshev's Other Inequality'. In: Inequalities in Statistics and Probability 5 (1984). URL: https://www.jstor.org/stable/43554 (visited on 08/10/2024).
- [25] Remco van der Hofstad. Random Graphs and Complex Networks: Volume
 2. Cambridge University Press, Feb. 2024. URL: https://rhofstad.win.tue.nl/NotesRGCNI (visited on 07/10/2024).
- [26] Roberto Imbuzeiro Oliveira. Concentration of the adjacency matrix and of the Laplacian in random graphs with independent edges. arXiv.org. URL: https://arxiv.org/abs/0911.0600 (visited on 07/10/2024).
- [27] Can M Le, Elizaveta Levina and Roman Vershynin. 'Concentration and regularization of random graphs'. In: *Random Structures and Algorithms* 51 (Mar. 2017), pp. 538-561. DOI: https://doi.org/10.1002/rsa.20713. URL: https://onlinelibrary.wiley.com/doi/10.1002/rsa.20713 (visited on 07/10/2024).
- [28] Linyuan Lu and Xing Peng. 'Spectra of edge-independent random graphs'. In: arXiv.org 20 (Nov. 2013). URL: https://arxiv.org/abs/1204.6207 (visited on 04/08/2024).
- [29] David Aldous. Open problems. Berkeley.edu, 2018. URL: https://www.stat.berkeley.edu/~ (visited on 04/08/2024).
- [30] Michael Schweinberger et al. 'Exponential-Family Models of Random Graphs: Inference in Finite, Super and Infinite Population Scenarios'. In: *Statistical Science* 35 (Nov. 2020). DOI: 10.1214/19-sts743. (Visited on 14/01/2025).

8 Appendix

Lemma 40. Let $x, a, b \in \mathbb{R}$ with a, b > 0. Then

$$e^{-ax}(1-e^{-bx}) \le \frac{b}{a}.$$

Proof. Define $f(x) := e^{-ax}(1 - e^{-bx})$. Differentiating and studying the derivative one finds that the unique global maximum is achieved at $\hat{x} = \frac{1}{b} \ln \frac{a+b}{a}$. Evaluating f at this point gives

$$f(\hat{x}) = \left(\frac{a}{a+b}\right)^{\frac{a}{b}} \frac{b}{a+b} < \frac{b}{a}.$$

The following is a well-known result for positive sequences and essentially the same proof can no doubt be found elsewhere. However, we give a detailed proof to justify that it extends to non-negative sequences in the obvious way.

Lemma 41. For any collections of non-negative reals $\{a_i\}_{i \in I}, \{b_i\}_{i \in I}$ not both identically zero such that the sums exist one has

$$\inf_{i \in I} \frac{a_i}{b_i} \le \frac{\sum_{i \in I} a_i}{\sum_{i \in I} b_i} \le \sup_{i \in I} \frac{a_i}{b_i},$$

where we define $\frac{r}{0} := \infty$ for any r > 0.

Proof. If b is identically zero then the stated inequality holds trivially so we may assume that b is not identically zero and thus divide by $\sum_{i \in I} b_i$ without trouble. Letting \tilde{I} be the set of indices such that a and b are not both simultaneously zero we notice that

$$\frac{\sum_{i\in I} a_i}{\sum_{i\in I} b_i} = \frac{\sum_{i\in \tilde{I}} a_i}{\sum_{i\in \tilde{I}} b_i}.$$

Thus we may assume that a and b are never simultaneously zero and hence that all ratios are well-defined in our extended sense.

Let $J := \{i \in I : b_i > 0\}.$

$$\sum_{i \in I} a_i \ge \sum_{i \in J} \frac{b_i}{b_i} a_i \ge \left(\inf_{i \in J} \frac{a_i}{b_i} \right) \sum_{i \in J} b_i = \left(\inf_{i \in I} \frac{a_i}{b_i} \right) \sum_{i \in I} b_i,$$

where the last equality comes from noticing that the ratios in the extra indices over which we take the infimum are infinite and that the extra terms in the sum are zero by definition of the index set. Now dividing by $\sum_{i \in I} b_i$ gives the first inequality.

For the second inequality, if there is an index i such that $b_i = 0$ then since we have ensured that a and b are never simultaneously zero, the inequality holds trivially. Thus we may assume b is never zero. We may then write,

$$\sum_{i \in I} a_i = \sum_{i \in I} \frac{b_i}{b_i} a_i \le \left(\sup_{i \in I} \frac{a_i}{b_i} \right) \sum_{i \in I} b_i.$$