H^{∞} and Complex Interpolation.

Jaak Peetre

Per G. Nilsson (Typist)

LUND INSTITUTE OF TECHNOLOGY, LUND, SWEDEN

Nilsson: Stockholm, Sweden Email address: pgn@plntx.com

ABSTRACT. This note is an (exact) copy of the report of Jaak Peetre, " H^{∞} and Complex Interpolation". Published as Technical Report, Lund (1981). Some more recent general references have been added, some references updated though (in italics) and some misprints corrected.

H^{∞} and complex interpolation.

Jaak Peetre

0. Introduction

In the theory of complex interpolation one usually defines the interpolation spaces in question using holomorphic functions taking continuous boundary values in the classical sense. (For the complex interpolation the main source is still Calderon's classic, albeit (helas !) tough reading paper [[5](#page-15-0)]. For an excellent introduction we recommend chap. 4 in the book $[3]$ $[3]$ $[3]$.)^{[1](#page-1-0)} That is, we have to deal with the space A.

The main thesis advanced in this paper is that in many questions is much more advantageous to consider the space H^{∞} (of bounded holomorphic functions) and further that it is in this connection often easier to work with distributional boundary values rather than with classical pointwise (a.e.) ones.

Our main result in this direction (sec. 4, theorem) says that one gets the same interpolation space \overline{X}_{θ} no matter whether one uses A or H^{∞} . (Here and in the sequel $\overline{X} = (X_0, X_1)$ denotes any fixed (compatible) pair of Banach spaces; see [[3](#page-15-1)].) We likewise obtain (sec. 5, theorem) a similar characterization of the second Calderon space \overline{X}^{θ} .

Our new description of \overline{X}^{θ} is formally obtained from the one for \overline{X}_{θ} by substituting the functor hom $(L^1,-)$ for the functor $L^{\infty}(-)$. As a consequence we obtain the result (sec. 6, proposition) that if either X_0 or X_1 has the RN (Radon-Nikodym) property (see e.g. [[11](#page-15-2)]) then $\overline{X}_{\theta} = \overline{X}^{\theta}$. Calderon [[5](#page-15-0)] proved this with "reflexive" instead of "RN".

In fact our new proof of his result is almost trivial, modulo that fact that "reflexive" entails "RN". We likewise give a new proof of the Calderon's duality theorem [[5](#page-15-0)] and of Janson's recent characterization [[23](#page-15-3)] of the complex interpolation spaces in terms of orbit spaces.

Apart from these applications to essentially known results of the Calderon theory we can use our new insight to treat vector valued versions of various problems traditionally considered for scalar H^{∞} . In particular we consider a natural analogue of the (free) interpolation problem: Given a sequence of points $\{z_n\}_{n=1}^{\infty}$ in the strip $\{0 < Re z < 1\}$ of the complex plane, which is uniformly separating in the sense of Carleson [[5](#page-15-0)]. To find for a given sequence ${w_n}_{n=1}^{\infty}$ of elements of $X_0 + X_1$, satisfying the appropriate boundedness condition, a function $f \in H^{\infty}(\overline{X})$ (definition in sec. 3) such that $f_n(z_n) = w_n$ $(n = 1, 2, ...)$. Again this problem is but a special case of a corresponding problem for the $\overline{\delta}$ operator (cf. [[21](#page-15-4)]). In this connection a recent paper by Jones [[24](#page-15-5)] was decisive.

Finally, to conclude this Introduction, we would like to point out what we are really interested in is interpolation of infinitely many Banach spaces, not just two. (Concerning interpolation of infinitely many spaces see the works of Coifman - Cwikel - Rochberg - Sagher - Weiss (the "western" branch") and Krein - Nicolova (the "eastern" branch) respectively, for instance

¹For references, post 1981, see notably Brudnyi-Krugljak [[27](#page-15-6), Chap. 2, Chap. 4], Brudnyi-Krein-Semenov [[28](#page-15-7)], Ovchinnikov [[29](#page-15-8)], Lindemulder-Lorist [[30](#page-15-9)] and the references listed.

the references [[8](#page-15-10)], [[9](#page-15-11)], [[25](#page-15-12)] listed here). It is mainly in order to keep the difficulties apart that we presently have restricted ourselves to just \underline{two} spaces. We hope however to be able to return to the general case in the near future. Let us only mention here that the leading underlying idea, which has motivated us in the present work too but which will find its true significance only in these later developments, is that the true setting for the theory of interpolation of infinitely many spaces is provided by Banach bundles in the sense of Fell [[15](#page-15-13)], [[16](#page-15-14)]. (Thus what we are really concerned with here is the Banach bundle valued $\overline{\delta}$ problem, rather than the vector valued one.) As a further indication of that we are now firmly on the right track let us point out that whereas Calderon's definition of \overline{X}^{θ} depends on special features of the underlying domain (viz. the strip ${0 < Re z < 1}$), the invariance with respect to (vertical) translations, and does not generalize, our's works a priori for any plane domain with a sufficiently smooth boundary; indeed we can formally even consider generalizations with complex manifolds (with a distinguished boundary) in any number of dimensions (cf. Favini [[15](#page-15-15)], Fernandez [[17](#page-15-16)]).

Having assimilated the particulars of this Introduction the reader can now get a rough idea of how this paper is organized by glancing at the titles of the various sections.

Acknowledgement. My thanks are due to the following persons for helpful suggestions, in particular what concerns bibliographic references, in connection with this research: Lars Gårding, Svante Janson, Mario Milman, Per Nilsson, Jan-Erik Roos, the last mentioned also for his hospitably during the recent meeting of the Swedish Mathematical Society in Stockholm (May 25-26, 1981).

1. The space $H^{\infty}(\Sigma)$.

Let us denote by S the (closed) strip $\{0 \leq Re z \leq 1\}$ in the complex plane. Put further $S^i = \{0 < Rez < 1\}, \ \delta S = \delta_0 S \cup \delta_1 S = \{Rez = 0\} \cup \{Rez = 1\}.$

We usually set $z = x + iy$.

Let $\overline{X} = (X_0, X_1)$ be any (compatible) pair of Banach spaces and write $\Sigma = \Sigma \overline{X}$ $X_0 + X_1$; Σ is a Banach space (cf. [[3](#page-15-1)], chap. 2). We say $f \in H^{\infty}(\Sigma)$ if f is bounded holomorphic function defined in S^i with values in Σ . Clearly $H^{\infty}(\Sigma)$ is a Banach space in the norm $||f|| = \sup_{z \in S^i} ||f(z)||^2$ $||f|| = \sup_{z \in S^i} ||f(z)||^2$ We denote by H^{∞} the corresponding space of scalar (= complex valued) functions.

Claim. Boundary values of functions in $H^{\infty}(\Sigma)$ exists in the distribution sense.

To substantiate this claim we first make a long detour and consider in all generality the problem of existence of distributional boundary values. Sec. 2 will be devoted to it.

2. Digression on the existence of distributional boundary values.

First we perform a rotation by an angle of 90° .

We begin with the scalar case. Here we follow the most elegant treatment of Hörmander [[22](#page-15-17)]. (The result in itself goes back to antiquity (Archimedes?).

Let thus f be any (scalar) function holomorphic in an open $"\frac{1}{2}$ neighborhood" of 0, that is, a set of the form $\{|z| < r, y > 0\}$ $(r > 0)$. Assume that $f(z) = O(y^{-A})$ uniformly in x on compact sets, for some $A \geq 0.3$ $A \geq 0.3$. Then f can be continued to a distribution in the full neighborhood $\{|z| < r\}$, still denoted by $f = f(z)$, with the support contained in the "closed" $\frac{1}{2}$ -neighborhood $\{|z| < r, y \ge 0\}$, such that for a suitable distribution $g = g(x)$ (in one variable) holds

(1)
$$
\int g(x) \psi(x,0) dx = -2 \int_{y>0} f(x+iy) \frac{\delta \psi(x,y)}{\delta \overline{z}} dx dy
$$

²See note $\langle 1 \rangle$.

 3 See note $\langle 2 \rangle$.

for any (compactly supported) testfunction $\psi(x, y)$. (The integrals in [\(1\)](#page-2-2) are of course interpreted in distribution sense; $\frac{\delta}{\delta \overline{z}} = \frac{1}{2} \left(\frac{\delta}{\delta x} + i \frac{\delta}{\delta y} \right)$ is the Cauchy-Riemann operator.)

To prove this we first assume that f takes continuous boundary values in the classical sense (pointwise limits). Then [\(1\)](#page-2-2) holds trivially with bona fide integrals (just integrate by parts). Given any test function $\phi = \phi(x)$ (in one variable we can determine ψ with $\psi(x, 0) = \phi(x)$ such that $\frac{\delta \psi(x,y)}{\delta \overline{z}} = O(y^A)$. E.g. we take $\psi(x,y) = \phi(x) + \frac{iy}{1!} \phi'(x) + ... + \frac{(iy)^k}{k!}$ $\frac{y)^k}{k!} \phi^{(k)}(x)$ for y close to 0, with $k \geq A$. Then we obtain from [\(1\)](#page-2-2) the estimate

$$
(2) \qquad \qquad \left| \int g(x) \, \phi(x) \, dx \right| \leq C \, \|\phi\|_{k+1}
$$

where $\|\phi\|_{k+1}$ stands for a suitable norm involving the maxima of the partial derivatives of ϕ up to order $k + 1$ and C a constant depending on the bound in $f(z) = O(y^{-A})$ only. The general case (distributional boundary values) is now easily handled using [\(2\)](#page-3-0) by replacing $f(z)$ by the functions $f(z + i\epsilon)$ ($\epsilon > 0$) and then letting ϵ tend to 0.

In particular we see thus that $g(x)$ arises as the distributional limit ($\epsilon \to 0$) of the functions $f(x+i\epsilon)$. (We therefore allow ourselves in the sequel to write f (x) instead of $g(x)$, whenever this is convenient.) Notice also that it follows that any relation between holomorphic functions can by continuity be translated into a similar relation between the corresponding boundary values. (E.g. if $f_1 = f_2$ and g_1 and g_2 are the boundary values of f_1 and f_2 respectively then $g'_1 = g_2$, in the distribution sense of course.) We will somewhat pretentiously refer to this property as the principle of permanence.

We give now some additional comments on the above construction.

Remark 1. It is clear that the "uniform" condition $f(z) = O(y^{-A})$ can be replaced by an integral condition of the type continuous $\int \int |f(z)| y^A dx dy < \infty$, with integration over smaller $\frac{1}{2}$ -neighborhoods.

Remark 2 . For later use (sec. 8) we record that the argument given also applies to functions f satisfying the inhomogeneous Cauchy-Riemann equation $\frac{\delta f}{\delta \overline{z}} = \mu$, μ a measure with finite mass on smaller $\frac{1}{2}$ -neighborhoods.

Remark 3. It is further easy to see that the condition $f(z) = O(y^{-A})$ (or its integral analogue; see remark 1) values of a holomorphic function f exists in the sense of (1) then we must have $f(z) = O(y^{-A})$ for some $A \ge 0$ uniformly in x on compact sets.

Remark 4. The proceeding discussion can formally be extended to any partial differential operator with C^{∞} coefficients, preferably elliptic though, even in n variables. Notice that to the special test function $\psi(x, y)$ constructed then corresponds the formal solution of the Cauchy problem truncated after sufficiently many steps.

We are now ready for the <u>vector valued</u> case.

First of all we nail down that by a distribution with values in Σ (or any other Banach space for that matter) we mean a continuous linear map f from $\mathcal D$ (the space of all testfunctions) into Σ. For the value $f(\phi)$ of f at the element $\phi \in \mathcal{D}$ we again use the notation $\int f(x) \phi(x) dx$, in the case of one dimension, and similarly in two dimensions.

Let thus f be a holomorphic function in the same $\frac{1}{2}$ - neighborhood, however with values in Σ. Assume that $|| f(z) ||_{Σ} = O(y^{-A})$ uniformly in x on compacts sets, for some $A ≥ 0$. Then the proceeding considerations carry over $\frac{mutatis \ mutandi}{m}$. We conclude that f admits vector values distributional boundary values g in the sense that the inequality [\(1\)](#page-2-2) holds true for any test function ψ .

Finally we state a simple lemma which will be needed in the sequel.

Lemma, Assume that g in [\(1\)](#page-2-2) happens to be a continuous function. Then f has a continuous extension "up to the boundary" (i.e. to the set $\{z : |z| < r, y \ge 0\}$ and consequently the boundary values exists in the classical sense.

Proof: The proof is equally simple in the scalar as in the vector valued case. It suffices to establish the continuity at the origin 0. Introduce to this end a suitable cut-off function κ equal to 1 near 0. Then κf clearly can be represented as the sum of a function which is C^{∞} near 0 and the Poisson integral of κq . Since κq too is continuous the latter term has the desired continuity properties.

Remark. From this proof it is clear that this is a result for harmonic functions, rather that for holomorphic ones.

3. Return to $H^{\infty}(\Sigma)$. The space $H(\overline{X})$.

After this long digression we return back to the space $H^{\infty}(\Sigma)$.

Let thus again $f \in H^{\infty}(\Sigma)$. From the discussion in sec. 2 (after first rotating an angle 90° in the opposite sense) we conclude that the boundary values $f(i.)$ and $f(1+i.)$ are well-defined distributions in the variable y with values in Σ .

Remark. For future use (see sec. 4) we remark that the functions in $H^{\infty}(\Sigma)$ also take the same boundary values in the sense of the weak topology of Σ . Indeed let $f \in H^{\infty}(\Sigma)$. For any continuous linear functional l on Σ $(l \in \Sigma')$ the scalar function $h(z) = l(f(z))$ is in (scalar) H^{∞} . Thus h takes (scalar) distributional boundary values (and the latter agree of course with the classical pointwise a.e. boundary values, known to exist by Fatou's theorem). It is clear that the distributions $l(f (k + i.))$ and $h (k + i.)$ coincide.

Now we define $H^{\infty}(\overline{X})$ to be the subspace of $H^{\infty}(\Sigma)$ of those functions $f \in H^{\infty}(\Sigma)$ such that $f(i.) \in L^{\infty}(X_0)$ and $f(1+i.) \in L^{\infty}(X_1)$. $(L^{\infty}(X), X$ any Banach space, is realized as a space of vector valued distributions in the obvious way.^{[4](#page-4-0)}) $H^{\infty}(\overline{X})$ is a Banach space in the norm $||f|| = \max\left(||f(i)||_{L^{\infty}(X_0)}, ||f(1+i)||_{L^{\infty}(X_1)} \right)$, indeed a Banach subspace of $H^{\infty}(\Sigma)$. Both of these statements are embodied in the inequality

$$
||f||_{H^{\infty}(\Sigma)} \leq ||f||_{H^{\infty}(\overline{X})}.
$$

Let us prove [\(1\)](#page-4-1). Since $L^{\infty}(X_k) \subseteq L^{\infty}(\Sigma)$ $(k = 0, 1)$ it is obvious that

$$
\|f\|_{H^{\infty}\left(\overline{X}\right)} \geq \max\left(\|f(i.)\|_{L^{\infty}(\Sigma)}, \|f(1+i.)\|_{L^{\infty}(\Sigma)}\right).
$$

It thus suffices to prove that

(1')
$$
||f||_{H^{\infty}(\Sigma)} \leq \max \{ ||f(i)||_{L^{\infty}(\Sigma)}, ||f(1+i)||_{L^{\infty}(\Sigma)} \}
$$

that is, the maximum principle. The simplest way of establishing $(1')$ $(1')$ $(1')$ is perhaps via the scalar case, in which case we take $(1')$ $(1')$ $(1')$ for granted. Let l be any (continuous) linear functional on Σ of norm ≤ 1 . Then the scalar function $h(z) = l(f(z))$, as we have already noticed ultra (see the remark), certainly belongs to H^{∞} and in addition holds $||h (k + i.)||_{L^{\infty}} \leq ||f (k + i.)||_{L^{\infty}(\Sigma)}$ $(k = 0, 1)$ $(k = 0, 1)$ $(k = 0, 1)$, since $||l|| \leq 1$. It follows that $||h||_{H^{\infty}} \leq$ the right hand side of $\binom{1'}{l}$. Since for any $z \in S$ holds $|| f (z) ||_{\Sigma} = \sup |l (f (z))| \left(1 \right)$ $|| f (z) ||_{\Sigma} = \sup |l (f (z))| \left(1 \right)$ $|| f (z) ||_{\Sigma} = \sup |l (f (z))| \left(1 \right)$ now follows.

Let further $A(\Sigma)$ be the space of those functions in $H^{\infty}(\Sigma)$ which take continuous boundary values (belonging to Σ). By the lemma in Sec. 2 this is to say that f has a continuous extension to S. We can then also define $A(\overline{X})$ to be the subspace of $A(\Sigma)$ such that $f(i)$ is a continuous

 4 See note $\langle 3 \rangle$.

function with values in X_0 and $f(1+i)$ a continuous function with values in X_1 , in symbols: $f(i.) \in C(X_0), f(1+i.) \in C(X_1)$. Obviously $A(\overline{X})$ is a closed subspace of $H^{\infty}(\overline{X})$.

4. A new characterization of the Calderon(-Lions) space \overline{X}_{θ} .

By definition the image in Σ of $A(\overline{X})$ under the evaluation map $f \mapsto f(\theta)$ is the (first) Calderon space \overline{X}_{θ} . (Here $\theta \in (0,1)$; we identify of course $(0,1)$ with a closed subset of S^i .) As is well-known \overline{X}_{θ} is a Banach space in the natural quotient norm.

Now we verify that we get the same space \overline{X}_{θ} if we in this definition substitute H^{∞} for A. Theorem. The image of the evaluation map $H(\overline{X}) \to \Sigma : f \mapsto f(\theta)$ $(\theta \in (0,1))$ is \overline{X}_{θ} .

Proof: For convince, let us denote provisionally the image of $H(\overline{X})$ by $\overline{X}_{\theta}^{\infty}$ $\widetilde{\theta}$. Thus our concern is to show that $\overline{X}_{\theta} = \overline{X}_{\theta}^{\infty}$ $\widetilde{\theta}$, indeed with equality of norms, if we again agree to equip $\overline{X}_{\theta}^{\infty}$ with the quotient norm.

One direction is easy. Indeed if $a \in \overline{X}_{\theta}$ then by definition $a = f(\theta)$ for some function $f \in A(\overline{X})$. Since $A(\overline{X}) \subseteq H^{\infty}(\overline{X})$ this shows that $\overline{X}_{\theta} \subseteq \overline{X}_{\theta}^{\infty}$ $\overset{\infty}{\theta}$.

For the opposite direction we first recall Calderon's famous inequality: If $f \in A(\overline{X})$ then

(1)
$$
\|f(\theta)\|_{\overline{X}_{\theta}} \leq \exp\left(\int \log\|f(i.)\|_{X_0} d\pi_0^{\theta} + \int \log\|f(1+i.)\|_{X_1} d\pi_1^{\theta}\right)
$$

where π_k^{θ} stands for the <u>harmonic measure</u> of $\delta_K S$ at the point θ ; $d\pi_k^{\theta}/dy$ is thus the corresponding part of the <u>Poisson kernel</u> (evaluated at θ) : $d\pi_k^{\theta}/dy = P(\theta, k + iy)$; explicit expressions for the latter can be found in [[5](#page-15-0)] or in [[3](#page-15-1)], chap. 4.

Let now $a \in \overline{X}_{\theta}^{\infty}$ ϕ^{∞} so that $a = f(\theta)$ for some $f \in H^{\infty}(\overline{X})$.

Set $f_n(z) = n \int_0^{1/n} f(z+it) dt$, $n = 1, 2, ...$ We claim $f_n \in A(\overline{X})$. Indeed it is at once clear that f_n is at least a function in $H^{\infty}(\Sigma)$, whose boundary values by our principle of permanence (sec. 2) are given by an analogous formula: $f_n (k + iy) = n \int_0^{1/n} f (k + iy + it) dt$, $(k = 0, 1)$. But the latter fact shows that $f_n (k + iy)$ is a continuous function with values in X_k : in symbols $f_n (k + i.) \in C(X_k)$ $(k = 0, 1)$. (In fact $f_n (k + i.)$ is even in $Lip(X_k)$ with the Lipschitz constant bounded by $O(n)$. Because we can write

$$
f_n (k + i (y + h)) - f_n (k + iy) = n \left(\int_{1/n}^{1/n + h} - \int_0^h f (k + i (y + t)) dt \right)
$$

for h small). Thus by the lemma in sec. 2 f must be in $A(\Sigma)$ and so in $A(\overline{X})$ too. Our claim is substantiated. We infer now that $a_n = f_n(\theta) \in \overline{X}_{\theta}$. Next we apply [\(1\)](#page-5-0) to the difference $f_n - f_m$. Then by Fatou's lemma we see that $\{a_n\}$ is a Cauchy sequence in \overline{X}_{θ} . But \overline{X}_{θ} is complete and continuously embedded in Σ . Since the limit of $\{a_n\}$ in Σ clearly is $a = f(\theta)$ we therefore finally get $a \in \overline{X}_{\theta}$. \Box

As a first application of this result (cf. infra sec. 6) we get that we still have the same space \overline{X}_{θ} if we assume that the boundary values are taken in the sense of the weak topology.

Corollary (generalizing partial results by Janson [[23](#page-15-3)], see e.g. th. 27). Let $a \in \Sigma$ and assume $a = f(\theta)$ for some $f \in H^{\infty}(\Sigma)$ such that for every $l \in \Sigma'$ holds $\lim_{z \to k + iy} l(f(z)) =$ $l(g_k(y))$ a.e. where g_k is some function in $L^{\infty}(X_k)$ $(k = 0, 1)$. Then $a \in \overline{X}_{\theta}$.

Proof: It suffices to apply the remark in sec. 3. \Box

Remark. In this connection recall that a function with values in a Banach space is holomorphic if and only if it is "weakly holomorphic" (Dunford's theorem; see e.g. [[20](#page-15-18)], p.93).

5. The second Calderon space $\overline{X}^{\theta}.$

We now turn to the analogous question for the second Calderon space \overline{X}^{θ} .

First recall the definition: $a \in \overline{X}^{\theta}$ if and only if $a \in \Sigma$ and there exists a function g in $B(\overline{X})$ such that $g'(\theta) = a$. Again $g \in B(\overline{X})$ means that g is holomorphic in S^i and continuous in S taking values in Σ , with $||g(z)||_{\Sigma} = O(y)$, the boundary values $g(k+i.)$ being continuous functions with values in X_k (at least mod Σ) subject to the condition

$$
||g (k + iy + it) - g (k + iy)||_{X_k} \le c |t|,
$$

t real, $k = 0, 1$.

We also denote by $H_{\infty}\left(\overline{X}\right)$ the space of functions f which we obtain if we in the definition of $H^{\infty}(\overline{X})$ (sec. 3) replace the functor $L^{1}(-)$ by the functor hom $(L^{1}, -)$; if X is any Banach space hom (L^1, X) consist of all continuous linear maps from L^1 into X. (Clearly $L^{\infty}(X) \subseteq$ hom (L^1, X) for any X but the converse holds true if and only if X has the RN (Radon-Nikodym) property; this is practically a definition; see $[15]^5$ $[15]^5$ $[15]^5$ $[15]^5$.) Thus $f \in H_{\infty}(\overline{X})$ means that f is in $H^{\infty}(\Sigma)$ with $f(k+i.) \in \text{hom}(L^1, X_k)$ $(k = 0, 1)$.

Theorem. The image of the evaluation map $H_{\infty}\left(\overline{X}\right) \to \Sigma : f \mapsto f(\theta)$ is the Calderon space \overline{X}^θ .

<u>Proof:</u> In analogy with the corresponding proof in sec. 4 we denote the image in question by $\overline{X}^{\theta}_{\infty}$. We want to show that $\overline{X}^{\theta} = \overline{X}^{\theta}_{\infty}$.

Let $a \in \overline{X}^{\theta}_{\infty}$ so that $a = f(\theta)$ for some $f \in H_{\infty}(\overline{X})$. Put $g(z) \stackrel{def}{=} \int_{z_0}^{z} f(\zeta) d\zeta$, z_0 some fixed point in S. Because of the analyticity the integral is independent of contour of integration, to be taken inside S of course.

Claim. $g \in B(\overline{X})$. (Then will follows that $a \in \overline{X}^{\theta}$ since it is clear that $g'(\theta) = a$.)

<u>Proof</u> (of the claim): It is clear that g is holomorphic in S^i with values in Σ and that $||g(z)||_{\Sigma} = O(y)$. There remains to investigate the boundary values of g. To fix the ideas consider those on $\delta_0 S$ (x = 0) taking for convenience $z_0 = 0$. Since $g' = f$ the permanence principle (see sec. 2) shows that the distributional derivative of $g (iy)$ is $if (iy)$. That is, for every test function ϕ holds

(1)
$$
\int g(iy) \phi'(y) dy = - \int f(iy) \phi(y) dy.
$$

Now put

$$
F(y) \stackrel{def}{=} \int_0^y f(i\eta) d\eta = \int_{-\infty}^\infty \chi_{(0,y)}(\eta) f(i\eta) d\eta.
$$

(If (a, b) , $a < b$ is an interval on the real line we let $\chi_{(a,b)}$ be its characteristic function; if $a > b$ we set $\chi_{(a,b)} = -\chi_{(b,a)}$. Then $F(0) = 0$ and

(2)
$$
F(y+t) - F(y) = \int_{-\infty}^{\infty} \chi_{(y,y+t)}(\eta) f(i\eta) d\eta.
$$

Since by assumption $f(i) \in \text{hom}(L^1, X_0)$ we see that $F \in C(X_0)$. Moreover by our choice of z_0 it is easy to see that $g(iy) = -F(t)$. (Take $\phi(y) = \int_{-\infty}^{y} \psi(\eta) d\eta$ in [\(1\)](#page-6-1), where $\int_{-\infty}^{\infty} \psi(\eta) d\eta =$ 0.) So $g(i)$ is a continuous function with values in X_0 . Using [\(2\)](#page-6-2) and once more $f(i) \in$ hom (L^1, X_0) we get the crucial estimate

$$
||g(iy + it) - g(iy)||_{X_0} \leq C ||\chi_{(y,y+t)}||_{L^1} \leq C |t|.
$$

In the same way we treat the boundary values on $\delta_1 S$ ($x = 1$). Therefore we can safely contend that $g \in B(\overline{X})$. \Box

 5 See note $\langle 4 \rangle$.

Proof (of the theorem/continued/): We have thus shown that $\overline{X}^{\theta}_{\infty} \subset \overline{X}^{\theta}$. To prove the opposite inclusion consider any $a \in \overline{X}^{\theta}$ and let g be in $B(\overline{X})$ such that $g'(\theta) = a$. Set $f = g'$.

Claim. $f \in H_{\infty}(\overline{X})$. (From which at once follows that $a \in \overline{X}_{\infty}^{\theta}$ since $f(\theta) = g'(\theta) = a$.)

<u>Proof</u> (of the claim): It is clear that f is a holomorphic function with values in Σ . Since $g \in B(\overline{X})$ there holds in particular $||g(k+iy+it) - g(k+iy)||_{\Sigma} \leq C |t|$ $(k = 0, 1)$. Therefore by the maximum principle $||g(z + it) - g(z)||_{\Sigma} \leq C |t| (z \in S)$. $f(z) = g'(z) =$ $\lim_{t\to 0} (g(z+it)-g(z))/it$ in the norm topology of Σ . Therefore $||f(z)||_{\Sigma} \leq C$ and $f \in$ $H^{\infty}(\Sigma).$

Let us investigate the boundary values on $\delta_0 S$, say. Again by the permanence principle the distributional derivative of $g(iy)$ is if (iy) . That is, [\(1\)](#page-6-1) holds as before.

But the left hand side is the limit in X_0 of

=

$$
\int_{-\infty}^{\infty} g(iy) \frac{\phi(y+t) - \phi(y)}{t} dy
$$

$$
= \int_{-\infty}^{\infty} \frac{g(i(y-t)) - g(iy)}{t} \phi(y) dy.
$$

 $g \in B\left(\overline{X}\right)$ again shows that the norm (in X_0) of the latter expression for any t is majorized by $C \|\phi\|_{L^1}$. Therefore $\left\| \int f(iy) \phi(y) dy \right\|_{X_0} \leq C \|\phi\|_{L^1}$ and $f(i.) \in \text{hom} (L^1, X_0)$.

In exactly the same way we handle the boundary values on δ_1S so we get $f(1+i.) \in$ hom (L^1, X_1) too. This proves $f \in H_{\infty}(\overline{X})$. \Box

The proof is now complete. \square

6. Applications to the Calderon theory.

N.B. - The reading of this sec. is not needed for sec. 7-9.

6.1. Calderon [[5](#page-15-0)] proved that if one of the spaces, say X_0 , is reflexive then with equality of norms $\overline{X}_{\theta} = \overline{X}^{\theta}$ for any $\theta \in (0, 1)$. We now generalize this result.

Proposition. Assume that X_0 has the RN property, that is hom $(L^1, X_0) = L^{\infty}(X_0)$ if the natural identifications are made. Then $\overline{X}_{\theta} = \overline{X}^{\theta}$ with equality of norms.

Remark. This is generalization because by a theorem of Phillips's (see [[11](#page-15-2)], p. 82) reflexive spaces have the RN property. Notice also by a theorem of Bergh's [[2](#page-15-19)] we know that \overline{X}_{θ} always is isometrically a subspace of \overline{X}^{θ} .

If X_1 to has the RN property then $H^{\infty}(\overline{X}) = H_{\infty}(\overline{X})$ and there is not much to prove. For the general case we need again Calderon's inequality (see formula [1](#page-5-0) in sec. 4).

Proof (of the proposition): Let $a \in \overline{X}^{\theta}$ and pick up $f \in H_{\infty}(\overline{X})$ such that $f(\theta) = a$. (Here we use the theorem in sec. 5.) As in a previous proof (sec. 4, theorem) we set $f_n(z)$ = $n \int_0^{1/n} f(z+it) dt$ $(n = 1, 2, ...)$ or more properly $f_n(z) = n \int \chi_{(y,y+1/n)}(\eta) f(x+i\eta) d\eta$ where we as usual write $z = x + iy$. It is easily seen that $f \in H^{\infty}(\overline{X})$ and even $\in A(\overline{X})$. Therefore $a_n \stackrel{def}{=} f_n(\theta) \in \overline{X}_{\theta}.$

Because of the RN property of the space X_0 we have $f(i) \in L^{\infty}(X_0)$ and $f_n(iy) \to f(iy)$ a.e. as $n \to \infty$. Also $||f_n(k+i)||_{L^{\infty}(X_k)} \leq C$ $(k = 0, 1)$ for some C. Therefore using now Calderon's inequality (formula [1](#page-5-0) of sec. $4)$ and Fatou's lemma, in a manner already familiar to us, we conclude that $\{a_n\}$ is a Cauchy sequence in \overline{X}_{θ} and from this again $a \in \overline{X}_{\theta}$. This proves $\overline{X}^{\theta} \subseteq \overline{X}_{\theta}$. The reverse inclusion is trivial. \Box

Remark. On the functor hom $(L^1, -)$. Let $f \in \text{hom}(L^1, X)$ where X is any given Banach space; that is, f is a continuous linear mapping from L^1 into X ; L^1 (and L^{∞}) is now taken with respect to some fixed measure space Ω with measure μ. The adjoint operator f^t maps X'into L^{∞} . For every element $l \in X'$ with $||l|| \leq 1$ we consider the element

 $f^t(l) = l \circ f$ of L^{∞} . Let now $f_X^{\#}$ denote the supremum of all the $f^t(l)$ considered as element of the Riesz space of (Radon) measure on Ω (cf. e.g. [[4](#page-15-20)]). This supremum clearly exists and it is easily seen that it is a measure absolutely continuous with respect to the given measure μ . So $f_X^{\#}$ can be identified with an element of L^{∞} and one readily verifies that $||f|| = ||f_X^{\#}||_{L^{\infty}}$. In order to demonstrate the usefulness of this notion let us indicate a quick proof of one of the simplest positive results of the RN property, namely Dunford's theorem to the effect that every Banach space with a conditionally bounded basis $(u_n)_1^{\infty}$ $\frac{\infty}{1}$ has the RN property (see [[11](#page-15-2)], p. 64). With no loss of generality we can assume that $(u_n)_1^{\infty}$ in addition is <u>monotone</u>. Let $(v_n)_1^{\infty}$ \sum_{1}^{∞} be the dual "basis" in X'. For $\phi \in L^{1}$ we can write $f(\phi) = \sum_{n=1}^{\infty} v_n(f(\phi)) u_n = \sum_{n=1}^{\infty} f^t(v_n(\phi)) u_n$. Each $f^n v_n$ can be identified with a function in L[∞]. Consider $\sum_{1}^{N} v_n(f(\phi)) u_n = \int_{\Omega} \sum_{1}^{N} f^t v_n(\omega) u_n \phi(\omega) d\mu(\omega) = \int_{\Omega} f_N(\omega) \phi(\omega) d\mu(\omega)$. By the monotonicity we see that $||f_N(\omega)||_X \leq f_X^{\#}(\omega)$ a.e. Therefore by the conditional boundedness $f_N(\omega) \to f(\omega)$ a.e. for some function $f \in L^{\infty}(X)$. Clearly f "represents" f and the proof is complete. Perhaps there are other applications too (to the RN theory). Here we content to point out returning to the arena of complex interpolation that with the aid of this notion we can formally generalize Calderon's inequality to the space \overline{X}^{θ} too. If $f \in H_{\infty}(\overline{X})$ then holds

$$
\|f(\theta)\|_{\overline{X}^{\theta}} \leq \exp\left(\int \log f_{X_0}^{\#}(i)\right) d\pi_0^{\theta} + \int \log f_{X_1}^{\#}(1+i)\,d\pi_1^{\theta}
$$

with the same meaning of π_K^{θ} as in sec. 4 $(k = 0, 1)$; $f_{X_k}^{\#}(k + i)$ is of course formed with respect to Lebesque measure considering $f(k+i)$ as an X_k valued function.

The proof is about the same as for the classical Calderon's inequality in the case of the space \overline{X}_{θ} (see [[5](#page-15-0)], p. 134); instead of $||f(k+iy)||_{X_k}$, which does not "exists", use $f_{X_k}^{\#}(k+iy)$. Unfortunately we know of no application of this new inequality.

6.2. Having returned to the Calderon theory for good, we sketch a new and conceptually perhaps simpler proof of the duality theorem (see [[5](#page-15-0)] or [[3](#page-15-1)]), to the effect that $(\overline{X}_{\theta})^{'} \cong (\overline{X}')^{\theta}$ if the dual pair $\overline{X'} = \left(X'_0\right)$ $\left(0, X_1'\right)$ "exists". It will be based on the following representation for the dual of $L^1(X)$, X any Banach space: $(L^1(X))^{T} \cong \text{hom}(L^1, X^{T})$, which is different from the one used by Calderon (and, what is important, not tied to the properties of the real line).

Let thus l be a continuous linear functional on \overline{X}_{θ} . We wish to identify l with an element of $(\overline{X}')^{\theta}$. (The other inclusion is trivial and will not be considered here.) By Calderon's inequality once more the relation $f \mapsto l(f(\theta))$ defines a continuous linear functional on $H^{\infty}(\overline{X})$, equipped with the norm $|||f||| = \int ||f(iy)||_{X_0} P(\theta, iy) dy + \int ||f(1+iy)||_{X_1} P(\theta, 1+iy) dy$. So by the Hahn-Banach theorem and by the above representation of the dual of $L^1(X)$ we see that there exists elements h_0 and h_1 of hom $(L^1, X_0^{'})$ and hom $(L^1, X_1^{'})$ respectively such that (formally)

$$
l(f(\theta)) = \int \langle h_0(y), f(iy) \rangle P(\theta, iy) dy + \int \langle h_1(y), f(1+iy) \rangle P(\theta, 1+iy) dy.
$$

The proof is completed by observing that h_0 and h_1 are the (distributional) boundary values on $\delta_0 S$ and $\delta_1 S$ respectively of a suitable holomorphic function g with values in $\Sigma\left(\overrightarrow{X}\right) = X_0 + X_1$ $\frac{1}{1}$. This is done more or less as in Calderon's case (see [[5](#page-15-0)] or [[3](#page-15-1)]) and one then also finds that $g \in H_{\infty}\left(\overline{X}'\right)$. So we get $l(x) = \langle y, x \rangle$ where $y = g(\theta)$ thus is an element of $\left(\overline{X}'\right)^{\theta}$.

 6.3 . We next turn our attention to Janson's characterization ([[23](#page-15-3)], th. 22) of the complex interpolation functors as orbit functors in the sense of the Aronszajn-Gagliardo theorem [[1](#page-15-21)] (cf. [[3](#page-15-1)], chap. 2). Here we offer an alternative proof which again from the conceptual point of view might have some advantages. (What we have in mind is of course possible extensions. E.g. the same procedure should be applicable in the case of infinitely many spaces; cf. Introduction.)

First we reformulate Janson's theorem in a way suitable for our purposes. (Janson uses the discrete version of the Calderon spaces (Cwikel's theorem [[10](#page-15-22)]) but this is not really the point.)

Consider the following (compatible) pair of Banach spaces $\overline{F} = (F_0, F_1)^6$ $\overline{F} = (F_0, F_1)^6$. The "containing" space is simply the dual of (scalar) H^{∞} and the space F_k consists of those elements μ of (H^{∞}) ′ which can be represented in the form $\mu(\phi) = \int_{-\infty}^{\infty} \psi(y) \phi(k+iy) dy$, $(\phi \in H^{\infty})$ with $\psi \in L^{1}$ $(k = 0, 1).$

It follows that an element $\mu \in (H^{\infty})^{'}$ is in the hull (sum) $F_0 + F_1$ if and only if it can be represented in the form $\mu(\phi) = \int_{-\infty}^{\infty} \psi_0(y) \phi(iy) dy + \int_{-\infty}^{\infty} \psi_1(y) \phi(1+iy) dy$ with $\psi_k \in L^1$ $(k = 0, 1)$. Let further $F = F_{\theta}$ $(\theta \in (0, 1))$ denote the subspace of $(H^{\infty})^{'}$ consisting of the linear functionals of the form $\mu(\phi) = \int_{-\infty}^{\infty} \psi(y) \phi(\theta + iy) dy$ with $\psi \in L^1$ and by δ_z ($z \in S$) the linear functional defined by $\delta_z(\phi) = \phi(z)$ (in other words, the evaluation map).

Then Janson's results says that

 1° \overline{X}_{θ} is the orbit of F in \overline{X} .

 $2^{\circ} \overline{X}^{\theta}$ is the orbit of δ_{θ} in \overline{X} .

 (\overline{X}) denotes as before an arbitrary Banach couple.).

Proof: . One way is easy. By general principles connected with the Aronszajn-Gagliardo theorem [[1](#page-15-21)] it suffices to show $F \subseteq \overline{F}_{\theta}$ and $\delta_{\theta} \in \overline{F}^{\theta}$ respectively.

In the first case if $\mu \in F$, $\mu(\phi) = \int \psi(y) \phi(\theta + iy) dy$ $(\phi \in H^{\infty})$ for some $\psi \in L^{1}$, we get a vector valued function f by defining $f(z)$ for $z \in S^i$ as the linear functional $\phi \mapsto$ $\int \psi(\eta) \phi(z + i\eta) d\eta$. It is readily seen $f \in H^{\infty}(\overline{F})$ and that $f(\theta) = \mu$. This proves $\mu \in \overline{F}_{\theta}$.

In the second case set $f(z) = \delta_z$. Then $f \in H_{\infty}(\overline{F})$ and trivially $f(\theta) = \delta_{\theta}$ which gives $\delta_{\theta} \in \overline{F}^{\theta}$. The fact that $f \in H_{\infty}(\overline{F})$ requires a proof. Let us just indicate how one sees that $f(i) \in \text{hom}(L^1, F_0)$. The point is that the pointwise boundary values δ_{iy} do not belong to the sum $F_0 + F_1$. But if we smear them out with a test function ψ in we get the linear functional $\phi \mapsto \int_{-\infty}^{\infty} \psi(y) \phi(iy) dy \, (\phi \in H^{\infty})$ which obviously is in F_0 . This fixes the matter.

Now we turn to the opposite inclusion. If a is 1° in \overline{X}_{θ} or 2° in \overline{X}^{θ} let f be in $H^{\infty}(\overline{X})$ or in $H_{\infty}\left(\overline{X}\right)$ respectively with $a = f(\theta)$. First define a linear mapping $U : \overline{F} \to \overline{X}$ as follows. If $\mu \in \Sigma(\overline{F}) = F_0 + F_1$, $\mu(\phi) = \int_{-\infty}^{\infty} \psi_0(y) \phi(iy) dy + \int_{-\infty}^{\infty} \psi_1(y) \phi(1+iy) dy \, (\phi \in H^{\infty})$ with $\psi_k \in L^1$ $(k = 0, 1)$ we set $U(\mu) = \int_{-\infty}^{\infty} \psi_0(y) f(iy) dy + \int_{-\infty}^{\infty} \psi_1(y) f(1 + iy) dy$. This definition makes sense, since we know at least that $f(k+iy) \in \text{hom}(L^1, X_k)$ $(k = 0, 1)$. It is also practically obvious that $U(\mu)$ does not depend on the particular representation of μ in terms of functions ψ_0 and ψ_1 . Moreover clearly $U: F_k \to X_k$ $(k = 0, 1)$, that is, in abbrevia $U: \overline{F} \to \overline{X}$.

In case 2[°] we simply put $\mu = \delta_{\theta}$. Then we can take $\psi_k(y) = P(\theta, k + iy)$ (Poisson kernel) so $U(\delta_{\theta}) = f(\theta) = a$ and a is in the orbit of δ_{θ} .

In the case 1[°] we must (as in [[23](#page-15-3)]) use an approximation device. Set $a_n = n \int_0^{1/n} f(\theta + it) dt$. Then each $a_n \in \overline{X}_{\theta}$ and a_n tends to a in \overline{X}_{θ} (Here we used $f (k + iy) \in L^{\infty}(X_k)$.). Also $a_n = U(\mu_n)$ where μ_n is the linear functional defined by $\mu_n(\phi) = n \int_0^{1/n} \phi(\theta + it) dt$. Since, as is readily seen, $\mu_n \in F$ each a_n lies in the orbit of F and its (orbit) norm can be uniformly bounded by the one of a (in \overline{X}_{θ}). This shows (apply the usual iterative procedure) that a too is in the orbit.

 6 See note $\langle 5 \rangle$.

In conclusion let us point out that as a consequence of his theorem Janson also obtains a simple proof of the reiteration theorem (see [[23](#page-15-3)], th. 25).

7. A vector valued (or better Banach bundle valued) interpolation problem.

Let $\{z_n\}_{n=1}^{\infty} \subseteq S^i$ be any (fixed) sequence <u>uniformly separating</u> in the sense of Carleson [[6](#page-15-23)]. Then by Carleson's theorem [[6](#page-15-23)] for any (scalar) sequence ${w_n}_{n=1}^{\infty} \subseteq l^{\infty}$ we can find a (scalar) function $f \in H^{\infty}$ such that $f(z_n) = w_n$ $(n = 1, 2...).$

Now we turn to the corresponding vector valued problem: Given a (vector valued) sequence ${w_n}_{n=1}^{\infty}$ to find a function $f \in H^{\infty}(\overline{X})$ such that $f(z_n) = w_n$ $(n = 1, 2, ...)$. Clearly a necessary condition for this to be possible is that $w_n \in X_{z_n}$ $(n = 1, 2, ...)$, $\sup_n \|w_n\|_{\overline{X}_{z_n}} < \infty$. $(\overline{X}_z \ (z \in S^i)$ denotes of course the space \overline{X}_{Rez} .

Claim. This necessary condition is sufficient too.

To substantiate this claim we will make use of the functions constructed by Per Beurling (see [[7](#page-15-24)]): There exists a sequence of (scalar) functions ${F_n}_{n=1}^{\infty} \subseteq H^{\infty}$ such that $F_n(z_k) = \delta_{nk}$ $(n, k = 1, 2, ...)$ and $\sum_{n=1}^{\infty} |F_n(z)| \leq C < \infty$ $(z \in S^i)$. Then in the scalar case a particular solution of our interpolation problem is provided by

(1)
$$
f(z) = \sum_{n=1}^{\infty} w_n F_n(z).
$$

In the vector valued case we must modify [\(1\)](#page-10-0) somewhat. Indeed by definition (and by the theorem in sec. 4) we can find functions $f_n \in H^{\infty}(\overline{X})$ such that $f_n(z_n) = w_n$ $(n = 1, 2, ...)$ and such that $C' \stackrel{def}{=} \sup_n ||f_n||_{H^\infty(\overline{X})} < \infty$. Then we put

(1')
$$
f(z) = \sum_{n=1}^{\infty} f_n(z) F_n(z).
$$

Sub-claim. The function f defined by $(1')$ $(1')$ $(1')$ is in $H^{\infty}(\overline{X})$ and solves our interpolation problem, i.e. $f(z_n) = w_n$ $(n = 1, 2, ...).$

Proof (of the sub-claim). It is clear that for any $z \in S^i$ holds $||f_n(z)||_{\Sigma} \le ||f_n||_{H^{\infty}(\Sigma)} \le$ $||f_n||_{H^{\infty}}(\overline{X}) \leq C'$ $(n = 1, 2, ...)$. Therefore the series $\sum_{n=1}^{\infty} f_n(z) F_n(z)$ is normally convergent in Σ for $z \in S^i$ and its sum $f(z)$ is clearly a holomorphic function with $||f(z)||_{\Sigma} \leq CC'$. In particular at least $f \in H^{\infty}(\Sigma)$ and $f(z_n) = w_n$ $(n = 1, 2, ...).$

It remains as usual to investigate the boundary values. By the principle of permanence (sec. 2) we have for the boundary functions on $\delta_0 S$, say,

$$
f(i.) = \sum f_n(i.) F_n(i.),
$$

$$
\sum |F_n(i.)| \leq C < \infty, ||f_n(i.)||_{X_0} \leq C.
$$

Therefore $f(i) \in L^{\infty}(X_0)$. Similarly we prove that $f(1+i) \in L^{\infty}(X_1)$. So that indeed $f\in H^{\infty}\left(\overline{X}\right).$

8. A Banach bundle valued $\overline{\delta}$ problem.

As is well-known (see $\vert 21 \vert$ $\vert 21 \vert$ $\vert 21 \vert$) the interpolation problem in the previous sec. is a special case of a problem for the $\bar{\delta}$ -operator which in the scalar case can be formulated as follows: Let μ be a Carleson measure (see [[6](#page-15-23)], [[21](#page-15-4)], [[24](#page-15-5)]), to fix the ideas positive, and consider any function $w \in L^{\infty}(\mu)$. To find a function u in S^i such that $\overline{\delta}u = w\mu \ (\overline{\delta} = \delta/\delta\overline{z})$ and such that its distributional boundary values (in the sense of sec. 2) lies in L^{∞} . (To see the connection with the interpolation problem of sec. 7 we remark that if μ is discrete then u is related to the

previous f by $u = f/B$ where B is the Blaschke product formed with the sequence $\{z_n\}_{n=1}^{\infty}$. We can assume that at least $u \in L^1$, in which case we know for sure (sec. 2) that the boundary distribution exists.

This problem always has a solution, in view of the Hahn-Banach theorem (see [[21](#page-15-4)]), but it is of course not unique. Once an appropriate substitute for the Per Beurling functions has been found we can however write down a particular solution. To this end let us consider a kernel $K(z,\zeta)$ assumed to be at least measurable in both variables (measurable in ζ with respect to μ) such that

- (i) : $K(z,\zeta)$ is analytic in z for ζ fixed μ a.e.,
- (ii) : $K(\zeta, \zeta) = 1$,

$$
(iii): \int K(z,\zeta) P(z,\zeta) d\mu(\zeta) \leq C < \infty \ (z \in \delta S).
$$

Here $P(z,\zeta)$ ($z \in \delta S, \zeta \in S$) stands for the Poisson kernel. We define the kernel $H(z,\zeta)$ $(z \in S, \zeta \in S)$ by stipulating that 1[°] it should for ζ fixed be analytic in z except at $z = \zeta$ where it should have a simple pole with the residue $1/2\pi i$ and that 2° $|H(z,\zeta)| = P(z,\zeta)$ $(z \in \delta S, \zeta \in S)$. (See [[24](#page-15-5)] where a fairly explicit construction of such a kernel $K(z, \zeta)$ is given.^{[7](#page-11-0)}) Now the desired solution is obtained as

(1)
$$
u(z) = \int K(z,\zeta) H(z,\zeta) w(\zeta) d\mu(\zeta).
$$

Next let us consider the generalization to the vector valued case. It seems natural to ask for a solution u of $\overline{\delta}u = w\mu$ such that $u(i) \in L^{\infty}(X_0), u(1+i) \in L^{\infty}(X_1)$. What do we then have to require from w? (A priori we assume only that w is Σ valued.) A minute's reflection gives that we should assume $w(\zeta) \in X_{\zeta}$ μ -a.e., sup $||w(\zeta)||_{\overline{X}_{\zeta}} < \infty$ - to be exact, sup here means μ −ess sup - along with some measurability condition too. We postpone for a moment the discussion of the exact form of the latter. Since $w(\zeta) \in \overline{X}_{\zeta}$ we can for each ζ ($\mu - a.e.$) find a function $W(\cdot,\zeta) \in H^{\infty}(\overline{X})$ such that $w(\zeta) = W(\zeta,\zeta)$. We can also assume that $\|W(\cdot,\zeta)\|_{\overline{X}_{\zeta}} \leq C' < \infty \ (\zeta \in S)$. Then we set (formally)

(2)
$$
u(z) = \int_{S} K(z,\zeta) H(z,\zeta) W(z,\zeta) d\mu(\zeta) (z \in S^{i}).
$$

Note that this formula [\(2\)](#page-11-1) generalizes both [\(1\)](#page-11-2) above and $\binom{1'}{1}$ $\binom{1'}{1}$ $\binom{1'}{1}$ of sec. 7, the discrete case. If everything works out smoothly we expect [\(2\)](#page-11-1) to provide us with a solution of $\overline{\delta}y = w\mu$ of the desired kind. Indeed by the properties $(i) - (ii)$ of $K(z, \zeta)$ and by property 1° of $H(z, \zeta)$ we have (formally) $\overline{\delta}K(\cdot,\zeta) H(\cdot,\zeta) W(\cdot,\zeta) = w(\eta) \delta(\cdot-\zeta)$, so $\overline{\delta}u(z) = \int \delta(z-\zeta) w(\zeta) d\mu(\zeta) =$ $w(z) \mu(z)$. Also if we can pass to the limit in [\(2\)](#page-11-1) we expect that the same identity holds for $z \in \delta S$. From this using property *(iii)* and 2° we infer that

$$
\|u(z)\|_{X_k} \leq \int |K(z,\zeta)| P(z,\zeta) \|W(z,\zeta)\|_{X_k} d\mu(\zeta)
$$

$$
\leq CC' \ (k=0,1,z\in\delta S)
$$

so that we get the correct boundary behavior.

To make all this rigorous we have to make precise the assumptions on w . Let us denote by \overline{X} the family of Banach spaces $(\overline{X}_{\zeta})_{\zeta \in S^i}$. Let us also introduce the norm

$$
||w|| = \mu - \text{ess sup}_{\zeta \in S^i} ||w(\zeta)||_{\overline{X}_{\zeta}}.
$$

 7 See note $\langle 6 \rangle$.

Definition. We say that w is in $L^{\infty}(\mu,\overline{\overline{X}})$ is w can be approximated in the preceding metric $||w||$ with a sequence of functions w_n of the form

$$
w_n(\zeta) = \sum_{v=1}^{N_n} \chi_{e_v^n}(\zeta) f_v^n(\zeta)
$$

where each e_v^n is a μ measurable subset of S^i , $\chi_{e_v^n}$ standing for its characteristic function, and the f_v^n belong to $H^{\infty}(\overline{X})$.

With this definition it is an easy matter to prove rigorously the following result.

Proposition. For any $w \in L^{\infty}\left(\mu, \overline{\overline{X}}\right)$ there exists a function $u \in L^{1}_{loc}(\Sigma)$ such that $\overline{\delta}u = w\mu$ and such that in the sense of distributions $u (k + i.) \in L^{\infty} (X_k)$ $(k = 0, 1)$.

<u>Proof:</u> We assume first that w itself is of the form $w(\zeta) = \sum_{v=1}^{N} \chi_{e_v}(\zeta) f_v(\zeta)$ where the e_v are μ measurable sets and $f_v \in H^{\infty}(\overline{X})$. Then we set

$$
u(z) = \sum_{v=1}^{N} f_v(z) \int_{e_v} K(z, \zeta) H(z, \zeta) d\mu(\zeta) =
$$

$$
\sum_{v=1}^{n} f_v(z) u_v(z).
$$

(This is formally [\(2\)](#page-11-1) with $W(z,\zeta) = \sum \chi_{e_v}(\zeta) f_v(z)!$) By Jones's results [[24](#page-15-5)] we certainly have for each $v \overline{\delta} u_v = \chi_{e_v} \mu$. Therefore we get $\overline{\delta} u = \sum f_v \overline{\delta} u_v = \sum f_v \chi_{e_v} \mu = w \mu$ (since $\overline{\delta}f_v = 0!$). It is clear that u is in $L^1_{loc}(\Sigma)$ and satisfies all the requirements of the proposition. In particular we have

$$
||u (k + i.)||_{L^{\infty}(X_k)} \leq C_k ||w|| \quad (k = 0, 1).
$$

Moreover it is easy to see that we have for each R an estimate of the type

$$
\int_{|y|\leq R} \|u\|_{\Sigma} dx dy \leq C_R \|w\|.
$$

It is now obvious how to treat the general case too. If ${w_n}_{n=1}^{\infty}$ is an approximating sequence for w in the sense of the above definition, $\|w - w_n\| \to 0$ as $n \to \infty$ and if $\{u_n\}$ is the corresponding sequence of solutions of $\overline{\delta}u_n = w_n$ obtained by the previous procedure then u_n tends to a solution u of $\overline{\delta}u = w\mu$ solving our problem.

Remark. A final comment is in order. The definition of $L^{\infty}\left(\mu,\overline{\overline{X}}\right)$ ultra (as well as the notation) might seem quite ad hoc, but it is not; in fact, we insist, it is the most natural thing to be thought of. To see this we first have to change slightly our point of view. Instead of considering the family of Banach spaces $(\overline{X}_{\zeta})_{\zeta \in S^i}$ (until now denoted by \overline{X}) we consider their set theoretic union, using for this object the previous symbol $\overline{\overline{X}}$. Then $\overline{\overline{X}}$ might be considered a vector bundle over S^i (with no topology yet, however):

We have a natural projection $\pi : \overline{X} \to S^i$ and each fiber $\pi^{-1}(\zeta) = \overline{X}_{\zeta}$ is a Banach space, thus <u>a fortiori</u> a vector space. But more, \overline{X} is in fact in a technical sense (see Fell [[15](#page-15-13)], [[16](#page-15-14)]) a Banach bundle. In particular \overline{X} thus indeed carriers a natural topology by its own right. By a classical procedure by Godement's [[18](#page-15-25)], [[19](#page-15-26)] the Banach bundle structure can quite generally (any vector bundle over a (usually) locally compact space Ω) be defined by first specifying a suitable family of "principal sections". In our case $(\Omega = S^i)$ there is a canonical choice of the principal sections. These are simply the sections of the type $S^i \to \overline{X}$: $\zeta \mapsto f(\zeta)$, f a function in $H^{\infty}(\overline{X})$. For any Banach bundle B and any positive measure μ over its base space Ω we can construct a theory of L^p spaces $L^p(\mu, B)$ (apart from the works already listed see the books by

NOTES. 13

Dixmier [[13](#page-15-27)], p. 186-194 and by Dinculeanu [[12](#page-15-28)], p. 413-414 where additional references can be found). If we specialize we see that our $L^{\infty}\left(\mu,\overline{\overline{X}}\right)$ (μ is now again a Carleson measure) is just a special case of these general spaces. We reader can also convince himself of the fact that in the scalar case our definitions reduces to (one of) the usual definitions of $L^{\infty}(\mu)$. Also it is now finally plain why the word "Banach bundle" appears in the title of this and the previous sec.

9. Other problems for H^{∞} .

After this initial success it is now natural to pause and to ask what other traditional problems for H^{∞} can be generalized to this new setting of Banach bundles. In the first place what comes to ones mind is the Corona Problem but we have not obtained any positive results in that direction so presently we are bound to think that this might be a quite hopeless thing. One can also ask if one could do something with H^p . If we understand correctly a remark in [[24](#page-15-5)] it should be quite easy to prove vector valued analogous of the interpolation theorem of Shapiro and Shields [[26](#page-15-29)] but we have not tried to carry out the details.

Notes.

 $\langle 1 \rangle$. Convention: If V is any normed space, fixed under the discussion, we denote its norm by $\|\cdot\|$. If there are several spaces involved, in order to avoid confusion, we use V as subscript, thus writing $\left\| \cdot \right\|_V$ for $\left\| \cdot \right\|$.

 $\langle 2 \rangle$. We are in particular interested in the (rather trivial) special case $A = 0$ (bounded functions) but also in derivatives of bounded functions. If f is bounded then $f'(z) = O(y^{-1})$ by an easy application of Cauchy's theorem.

 $\langle 3 \rangle$. Unless otherwise specified by L^{∞} we mean L^{∞} with respect to Lebesque measure on the real line. Similarly for L^1 .

 $\langle 4 \rangle$. This work is entitled "Vector measures" but is in large portions devoted to a study of Banach space having the RN property.

 $\langle 5 \rangle$. F stands for Fourier.

 $\langle 6 \rangle$. In [[24](#page-15-5)] there is considered of course not the strip S but the upper (Poincare) halfplane $\Pi = \{y \geq 0\}$, the whole set-up being essentially invariant for biholomorphic (conformal) transformations; in particular we have the canonical map $z \to \exp(i\pi z)$ of S^i onto Π^i . In the case of the half-plane Π one has

$$
H(z,\zeta) = \frac{1}{2\pi i} \frac{\zeta - \overline{\zeta}}{(z-\zeta)(z-\overline{\zeta})} (z \in \Pi^{i}),
$$

$$
P(x,\zeta) = \frac{1}{\eta i} \frac{\operatorname{Im}\zeta}{|x-\zeta|^{2}} (x = z \in \delta \Pi).
$$

Bibliography

- [1] N. Aronszajn - E. Gagliardo, Interpolation spaces and interpolation methods. Ann. Math. Pura Appl. 68 (1965), 51-118.
- [2] J. Bergh, On the relation between the two complex methods of interpolation. Indiana U. Math. J. 28 (1979), 775-778.
- [3] J. Bergh - J. Löfström, Interpolation spaces. An introduction. Grundlehren der mathematischen Wissenschaften 223, Springer-Verlag, Berlin-Heidelberg-New York, 1976.
- [4] N. Bourbaki, Integration, chap. II. Actualites scientifiques et industrielles 1175, Hermann, Paris, 1952.
- [5] A.P. Calderon, Intermediate spaces and interpolation, the complex method. Studia Math. 24 (1964), 113- 190.
- [6] L. Carleson, Interpolations of bounded analytic functions and the Corona problem. Ann. Math. 76 (1962), 547-559.
- [7] L. Carleson, Interpolations by bounded analytic functions. Proc. Internat. Congr. Math., 15-22 Aug. 1962, Inst. Mittag-Leffter, Stockholm, 1963, 314-316.
- [8] R. Coifman - M. Cwikel - R. Rochberg - Y. Sagher - G. Weiss. The complex method for interpolation of operators acting on families of Banach spaces. Lecture notes in mathematics 779, Springer-Verlag, Berlin-Heidelberg-New York, 1980, 123-153.
- [9] R. Coifman - M. Cwikel - R. Rochberg - Y. Sagher - G. Weiss, Complex interpolation for families of Banach spaces. Proc. Symp. Pure Math. 35, part 2, A.M.S., 1979, 269-282.
- [10] M. Cwikel, Complex interpolation spaces, a discrete definition and reiteration. Indiana J. Math. 27 (1978), 1005-1009.
- [11] J. Diestel - J.J. Uhl, Vector measures. Mathematical Surveys 15, A.M.S., Providence, 1977.
- [12] N. Dinculeanu, Vector measures. Pergamon Press, Oxford etc., 1967.
- [13] J. Dixmier, Les C∗−algebres et leurs representations. Deuxieme ed., Gauthier Villiars, Paris, 1969.
- [15] A. Favini, Su una extensione del metodo d'interpolazione complesso. Rend. Sem. Mat. Univ. Padova 47 (1972), 244-298.
- [15] J.M.G. Fell, Induced representations and Banach ∗-algebraic bundles. Lecture notes in mathematics 582, Springer-Verlag, Berlin-Heidelberg-New York, 1977.
- [16] J.M.G. Fell, An extension of Mackey's method to ∗-algebraic bundles. Mem. Amer. Math. Soc. 90 (1969).
- [17] D.L. Fernandez, An extension of the complex method of interpolation. Univ. Estudual de Campinas, Relatorio Interno No. 120, 1978. (To appear in Boll.U.M.I. in a revised and shortened version.)
- [18] R. Godement, Sur la theorie des representations unitaires. Ann. Math. 53 (1951), 68-124.
- [19] R. Godement, Theorie generale des sommes continues d'espaces de Banach. C.R. Acad. Sci. Paris 228 (1949), 1321-1323.
- [20] E. Hille - R. Phillips, Functional analysis and semi-groups. Revised ed., Amer. Math. Soc., Providence, 1957.
- [21] L. Hörmander, Generators for some rings of analytic functions. Bull. Amer. Math. Soc. 73 (1976), 943-949.
- [22] L. Hörmander, personal communication.
- [23] S. Janson, Minimal and maximal methods of interpolation. Institut Mittag-Leffler, Report No. 6, 1980, J. Functional Analysis 44, 50-73 (1981).
- [24] P. Jones, L^{∞} estimates for the δ problem in a half-plane. Pre-print. Acta. Math. 150, 137-152 (1983).
- [25] S.G. Krein - L.I. Nikolova, Holomorphic functions in a family of Banach spaces, interpolation. Dokl. Akad. Nauk. SSSR 250 (1980), 547-550.
- [26] H. Shapiro - A. Shields, On some interpolation problems for analytic functions. Amer. J. Math. 83 (1961), 513-532.
- [27] Y.A. Brudnyi - N.Y. Kruglyak, Interpolation Functors and Interpolation Spaces 1. Elsevier, North-Holland (1991).
- [28] Y.A. Brudnyi - S.G. Krein - E.M. Semenov, Interpolation of linear operators. J. Soviet Math 42, 2009– 2113 (1988).
- [29] V.I. Ovchinnikov, The Method of Orbits in Interpolation Theory, Math. Reports 1 (1984), 349–516.
- [30] N. Lindemulder - E. Lorist, A discrete framework for the interpolation of Banach spaces. Adv. Math. 440, 109506 (2024).