

STATISTICAL INFERENCE FOR INTERACTING INNOVATION PROCESSES AND RELATED GENERAL RESULTS

GIACOMO ALETTI, IRENE CRIMALDI, AND ANDREA GHIGLIETTI

ABSTRACT. Given the importance of understanding how different innovation processes affect each other, we have introduced a model for a finite system of interacting innovation processes. The present work focuses on the second-order asymptotic properties of the model and illustrates how to leverage the theoretical results in order to make statistical inference on the intensity of the interaction. We apply the proposed tools to two real data sets (from Reddit and Gutenberg).

Key-words: Innovation, Interaction, Reinforcement, Statistical inference for Poisson-Dirichlet processes.

1. INTRODUCTION

The study of the mechanisms through which novelties emerge, diffuse and trigger further novelties is very important in many research fields (biology, linguistics, social science and others [7, 8, 14, 29, 30, 31]). We can define a *novelty* (or *innovation*) as the first time occurrence of some event of interest. A widely used mathematical object that models an *innovation process* is an *urn model with infinitely many colors* [32], also known as *species sampling sequence* [13, 17, 22, 23, 26]. Let C_1 be the first observed color, then, given the colors C_1, \dots, C_t of the first t extractions, the color of the $(t+1)$ -th extracted ball is new (i.e. not already drawn in the previous extractions) with a probability Z_t^* which is a function of C_1, \dots, C_t (sometimes called “birth probability”) and it is equal to the already observed color c with probability $P_{c,t} = \sum_{n=1}^t Q_{n,t} I_{\{C_n=c\}}$, where $Q_{n,t}$ is a function of C_1, \dots, C_t . The quantities Z_t^* and $Q_{n,t}$ specify the model: precisely, Z_t^* describes the probability of having a new color (that is a novelty) at time-step $t+1$ and $Q_{n,t}$ is the weight at time-step t associated to extraction n , with $1 \leq n \leq t$, so that the probability of having at time-step $t+1$ the “old” color c is proportional to the total weight at time-step t associated to that color (according to a *reinforcement mechanism*). Note that the number of possible colors is not fixed a priori, but new colors continuously enter the system. We can see the urn with infinitely many colors as the space of possibilities, while the sequence of extracted balls with their colors represents the history which has been actually realized.

The *Poisson-Dirichlet model* [20, 27, 28] provides the most used example of innovation process. For this model we have

$$(1) \quad \begin{aligned} Z_t^* &= \frac{\theta + \gamma D_t}{\theta + t}, & Q_{n,t} &= \frac{1 - \gamma/K_{C_n,t}}{\theta + t}, \\ \text{and so } P_{c,t} &= \frac{K_{c,t} - \gamma}{\theta + t}, \end{aligned}$$

where $0 \leq \gamma < 1$, $\theta > -\gamma$, D_t denotes the number of distinct extracted colors until time-step t and $K_{c,t}$ indicates the number of times color c has been drawn until time-step t . In particular, the probability $P_{c,t}$ to observe color c is proportional to $K_{c,t}$, according to a “preferential attachment” or “popularity” principle. From an applicative point of view, as an innovation process, the Poisson-Dirichlet process has the merit to reproduce in many cases the correct basic statistics, namely the

Heaps' [18] and the (generalized) Zipf's laws [34], which quantify, respectively, the rate at which new elements appear and the frequency distribution of the elements. In particular, the *Heaps' law* states that the number D_t of distinct observed elements (i.e. colors, according to the metaphor of the urn) when the system consists of t elements (i.e. after t extractions from the urn) follows a power law with an exponent smaller than or equal to 1 and, for the Poisson-Dirichlet model, we have $D_t \propto t^\gamma$ for $0 < \gamma < 1$ (while $D_t \propto \ln(t)$ for $\gamma = 0$). A very clear representation of the evolution dynamics of the Poisson-Dirichlet process is given by the *urn with triggering* model [1, 33]. An urn initially contains $N_0 > 0$ distinct balls of different colors. Then, at each time-step $t + 1$, a ball is randomly drawn with replacement from the urn and

- if the color of the extracted ball is new, i.e. it was not been extracted in the previous extractions, then we add $(\hat{\rho} - 1)$ balls of the same color as the extracted ball plus $(\nu + 1)$ balls of distinct new colors, i.e. not already present in the urn;
- if the color of the extracted ball is old, i.e. it has been already extracted in the previous extractions, we add ρ balls of the same color as the extracted one.

It is easy to verify that, when the *balance condition* $\hat{\rho} + \nu = \rho$ is satisfied (this means that at each time-step the number of balls added to the urn is always ρ , regardless of the outcome of the extraction), the above updating rule gives rise to the above probabilities (1), taking $\rho > \nu \geq 0$, $\theta = N_0/\rho$ and $\gamma = \nu/\rho$. The above reformulation of the Poisson-Dirichlet model has the merit to point out another principle, besides the *reinforcement*, that drives the dynamics: the *Kauffman's principle of the adjacent possible* [21]. Indeed, the model starts with an urn with a finite number of balls with distinct colors and, whenever a color is extracted for the first time, a set of balls with new colors is added to the urn. This represents Kauffman's idea that, when a novelty occurs, it triggers further potential novelties.

Given the importance of understanding how different innovation processes affect each other, in [5] we have introduced and analyzed a model for a finite network of innovation processes. In the proposed model, for each node h , the probability of observing a new or an old item depends, not only on the path of observations recorded for h itself, but also on the outcomes registered for the other nodes $j \neq h$. More precisely, we introduce a system of N urns with triggering representing N different innovation processes that interact each other as follows:

- (i) the probability of *exploitation* of an old item c by node h , i.e. the probability of extracting from urn h a color c already drawn in the past from an urn of the system (not necessarily from h itself), has an increasing dependence not only on the number of times c has been observed in node h itself (that could be even zero), but also on the number of times c has been observed in each of the other nodes;
- (ii) the probability of *production* (or *exploration*) of a novelty for the entire system by node h , i.e. the probability of extracting from urn h a color never extracted before from any of the urns in the system, has an increasing dependence not only on the number of novelties produced by h itself in the past, but also on the number of novelties produced by each of the other nodes in the past.

In particular, (ii) means that Kauffman's principle of the adjacent possible is at the "system level": that is, when node h produces a novelty for the system, this fact triggers further potential novelties in all the nodes of the system, not only in node h itself. The two different dependencies described above ((i) and (ii)) are tuned by two different matrices (called Γ and W in the sequel). More specifically, suppose to have N urns, labeled from 1 to N . *At time-step 0, the colors inside each urn are different from those in the other urns.* Let $N_{0,h} > 0$ be the number of distinct balls with distinct colors inside the urn h . Then, at each time-step $t \geq 1$, one ball is randomly drawn with

replacement from each urn and, for any $h = 1, \dots, N$, urn h is so updated according to the colors extracted from urn h itself and from all the other urns $j \neq h$:

- if the color of the ball extracted from urn h is “new” (i.e., it appears for the first time in the system), then into urn h we add $(\widehat{\rho}_{h,h} - 1)$, with $\widehat{\rho}_{h,h} > 0$, balls of the same color as the extracted one plus $(\nu_{h,h} + 1)$, with $\nu_{h,h} \geq 0$, balls of distinct “new” colors (i.e. not already present in the system);
- if the color of the ball extracted from urn h is “old” (i.e., it has been already extracted in the system), we add $\rho_{h,h} > 0$ balls of the same color into urn h ;
- for each $j \neq h$, if the color of the ball extracted from urn j is “new” (i.e., it appears for the first time in the system), then into urn h we add $\widehat{\rho}_{j,h} \geq 0$ balls of the same color as the one extracted from urn j plus $\nu_{j,h} \geq 0$ balls of distinct “new” colors (i.e. not already present in the system);
- for each $j \neq h$, if the color of the ball extracted from urn j is “old” (i.e., it has been already extracted in the system), then into urn h we add $\rho_{j,h} \geq 0$ balls of the same color as the one extracted from urn j .

The terms “new” and “old” refer to the entire system, that is a “new” color is a color that has never been extracted from an urn of the system. On the contrary, an “old” color is a color that has already been extracted from at least one urn of the system. We assume that the “*new*” colors added to a certain urn are always different from those added to the other urns (at the same time-step or in the past). By means of this fact, together with the assumption that initially the colors in the urns are different from each other, we cannot have the same new color extracted simultaneously from different urns. In other words, *we cannot have the same novelty observed simultaneously in different processes of the system*. Therefore, for each observed new color (novelty) c , there exists a unique urn (process), say $j^*(c)$, in the system that produced it. However, in a time-step following its first extraction, color c could be also extracted from another urn $h \neq j^*(c)$, as a consequence of the interaction among the urns (processes). Indeed, the “contamination” of the color-set of the urn h with the colors present in the other urns is possible by means of the interaction terms $\widehat{\rho}_{j,h}$ and/or $\rho_{j,h}$ in the above model dynamics. Furthermore, as in the standard Poisson-Dirichlet model, we assume the *balance condition*

$$(2) \quad \widehat{\rho}_{j,h} + \nu_{j,h} = \rho_{j,h}, \quad \text{i.e. } \widehat{\rho}_{j,h} = \rho_{j,h} - \nu_{j,h},$$

so that, at each time-step, each urn j contributes to increase the number of balls inside urn h by $\rho_{j,h} \geq 0$, with $\rho_{h,h} > 0$. Moreover, letting $\rho_h = \sum_{j=1}^N \rho_{j,h}$ and setting (without loss of generality)

$$(3) \quad \theta_h = N_{0,h}/\rho_h, \quad \gamma_{j,h} = \nu_{j,h}/\rho_h, \quad \lambda_{j,h} = \widehat{\rho}_{j,h}/\rho_h \quad \text{and} \quad w_{j,h} = \rho_{j,h}/\rho_h,$$

we obtain

$$(4) \quad Z_{t,h}^* = P(C_{t+1,h} = \text{“new”} \mid \mathbf{C}_1, \dots, \mathbf{C}_t) = \frac{\theta_h + \sum_{j=1}^N \gamma_{j,h} D_{t,j}^*}{\theta_h + t},$$

where $\mathbf{C}_k = (C_{k,1}, \dots, C_{k,N})^\top$ with $k = 1, \dots, t$, $D_{t,j}^*$ denotes the number, until time-step t , of distinct observed colors extracted for their first time from urn j , that is the number of distinct novelties for the whole system “produced” by urn (process) j until time-step t , and, for each “old” color c ,

$$(5) \quad \begin{aligned} P_t(h, c) &= P(C_{t+1,h} = c \mid \mathbf{C}_1, \dots, \mathbf{C}_t) = \frac{\sum_{j \neq j^*(c)} w_{j,h} K_t(j, c) + w_{j^*(c),h} (K_t(j^*(c), c) - 1) + \lambda_{j^*(c),h}}{\theta_h + t} \\ &= \frac{\sum_{j=1}^N w_{j,h} K_t(j, c) - \gamma_{j^*(c),h}}{\theta_h + t}, \end{aligned}$$

where $K_t(j, c)$ denotes the number of times the color c has been extracted from urn j until time-step t and $j^*(c)$ denotes the urn from which the color c has been extracted for the first time. Note that the probability that urn (process) h will produce at time-step $t + 1$ a novelty for the entire system has an increasing dependence on the number $D_{t,j}^*$ of novelties produced by the urn (process) j until time-step t and the parameter $\gamma_{j,h}$ regulates this dependence. In other words, Kauffman's principle of the adjacent possible is at the "system level": that is, for each pair (j, h) of urns in the system, the parameter $\gamma_{j,h}$ quantifies how much the production of a novelty by urn j induces potential novelties in urn h . Moreover, on the other hand, the probability that from urn h we will extract at time-step $t + 1$ an old color c has an increasing dependence on the number $K_t(j, c)$ of times the color c has been drawn from urn j until time-step t and the parameter $w_{j,h}$ quantifies how much the number $K_t(j, c)$ leads toward a future extraction of a ball of color c from urn h . In [5] we proved the first-order asymptotic properties of the model:

Theorem 1.1. *Suppose that the matrix $\Gamma = (\gamma_{j,h})_{j,h=1,\dots,N}$ is irreducible and denote by $\gamma^* \in (0, 1)$ the Perron-Frobenius eigenvalue of Γ . Then, for each $h = 1, \dots, N$, we have*

$$(6) \quad t^{-\gamma^*} D_{t,h}^* \xrightarrow{a.s.} D_{\infty,h}^{**},$$

where $D_{\infty,h}^{**}$ is a finite strictly positive random variable. Moreover, we have

$$\frac{D_{t,h}^*}{D_{t,j}^*} \xrightarrow{a.s.} \frac{u_h}{u_j} \quad \text{and} \quad \frac{D_{t,h}^*}{\sum_{j=1}^N D_{t,j}^*} \xrightarrow{a.s.} \frac{u_h}{\sum_{j=1}^N u_j},$$

where $\mathbf{u} = (u_h)_h$ is the (up to a multiplicative constant) left eigenvalue of Γ associated to γ^* .

Theorem 1.2. *Suppose that the matrix $W = (w_{j,h})_{j,h=1,\dots,N}$ is irreducible. Then, for each $h = 1, \dots, N$ and each observed color c in the system, we have*

$$(7) \quad P_t(h, c) \xrightarrow{a.s.} \tilde{P}_\infty(c) \mathbf{1},$$

$$(8) \quad \frac{1}{t} K_t(h, c) \xrightarrow{a.s.} \tilde{P}_\infty(c) \quad \text{and} \quad \frac{K_t(h, c)}{\sum_{j=1}^N K_t(j, c)} \rightarrow \frac{1}{N},$$

where $\tilde{P}_\infty(c)$ is a random variable that takes values in $(0, 1)$ and does not depend on h ¹.

The first result states that, if Γ is irreducible, that is the graph with the innovation processes as nodes and with Γ as the adjacency matrix is strongly connected, then $D_{t,h}^* \propto t^{\gamma^*}$ a.s. for all $h = 1, \dots, N$, that is all the $D_{t,h}^*$ grow with the same Heaps' exponent $\gamma^* \in (0, 1)$. This means that, at the steady state, all the processes of the network produce innovations for the system at the same rate. In addition, the ratio $D_{t,h}^*/D_{t,j}^*$ provides a strongly consistent estimator of the ratio u_h/u_j of the relative eigenvector centrality scores (with respect to Γ^\top) of the two nodes h and j and the share of innovations produced by process h for all the system converges almost surely to the absolute eigenvector centrality score of h . The second result affirms that if W is irreducible, that is the graph with the innovation processes as nodes and W as the adjacency matrix is strongly connected, then, for each observed item c , the number of times item c has been observed in process h grows linearly. Moreover, the number of times item c has been observed is asymptotically uniformly distributed among the processes of the system and, at the steady state, the probability that an item c is observed in a process is the same for each process.

¹Note that in Th 3.2 of [5] we state that this random variable takes values in $(0, 1]$, but, applying Theorem S1.3 of [5] also replacing $P_t(h, c)$ by $1 - P_t(h, c)$, we immediately obtain $(1 - \tilde{P}_\infty(c)) > 0$ a.s., that is $\tilde{P}_\infty(c) < 1$ a.s..

The present work aims at completing the study of the model dynamics providing the second-order asymptotic behaviour of the quantities in the above Theorems 1.1 and 1.2. Furthermore, we will show how to leverage the obtained theoretical results in order to get statistical tools for making inference on the interaction matrices Γ and W and on the limit probabilities $\tilde{P}_\infty(c)$ of observing an old item c .

It is worthwhile to note that we deduce the convergence results for the considered system of interacting innovation processes from general theoretical results, that we here prove and that can be applied also in other frameworks. Indeed, they regard a pair of multidimensional stochastic processes following a “stochastic approximation” dynamics. A dynamics similar to the one studied in the present work has already been considered in [24], where the authors established a central limit theorem for a pair of recursive procedures having two different step-sizes. However, this result does not apply to our situation. Indeed, the covariance matrices in their main result (Theorem 1) are deterministic, while the asymptotic covariance matrices in our central limit theorems are random. This is why we do not use the simple convergence in distribution, but we employ the notion of stable convergence, which is, among other things, essential for the considered statistical applications.

1.1. Previous literature. Despite the amount of scientific works regarding interacting urns with a finite set of colors (see, for instance, [2, 6] and the references therein), in the existing literature we have found only a few papers about a collection of interacting (in the same sense of [5] and the present work) urns with infinitely many colors, that is [15, 19]. In the model provided in [15] (see Example 3.8 in that paper), there is a finite collection of Dirichlet processes with random reinforcement. More precisely, in that model we have a random weight $W_{t,h}$ associated to the extraction at time-step t from the urn h so that, the probability of extracting from urn h an old color c (here, the term “old” refers to urn h , that is a color never extracted before from urn h) is proportional to the weight associated to that color, specifically $\sum_{n=1}^t W_{n,h} I_{C_{n,h}=c} / (\theta + \sum_{n=1}^t W_{n,h})$. The interaction across the urns is introduced by means of the weights, which could be stochastically dependent: each $W_{n,h}$ may be the same for each urn h , or a function of the observed outcomes of the other urns, or a function of some common (observable or latent) variables. It is easy to understand that this model is different from the one introduced in [5] and considered in this work: we consider Poisson-Dirichlet processes, not only Dirichlet processes, and, differently from the model in [15], for us, the notion of “old” or “new” color refers to the entire system, not to each single urn, and Kauffman’s principle of the adjacent possible is at the system level as explained above. The present work and [19] share the fact that the considered models are both a collection of urns with triggering with an interacting dynamics that brings the Kauffman’s principle of the adjacent possible from the single agent to the network of agents. Adopting the terminology of [19], we can say that both interacting mechanisms are based on the construction and the updating of a “social” urn for each network node from which the extractions take place, but the construction and the updating rules of the social urns are deeply different in the two models. In particular, differently from [19], we introduce the notion of “new” and “old” at the system level. In [19] the authors focus on the novelties in each sequence (novelty in h = first appearance in h of a new item), that they call “discoveries”; while we also study the sequence of the novelties for whole the system produced by each agent. Furthermore, in [19] the extraction of an “old” item in a certain network node does not affect the other nodes, in our model we also have an interacting reinforcement mechanism for the “old” items: the probability of the extraction of an old item depends on the number of times it has been observed in all the nodes. This allows us to get a specific result on the distribution of the observations in the system among the different items observed.

1.2. Structure of the work. In Section 2 we present the main results of this paper, which consist in the Central Limit Theorems (CLTs) describing the second-order asymptotic behaviour of the considered system of interacting innovation processes, that is of the quantities in Theorems 1.1 and 1.2. Then, in Section 3 we illustrate some statistical tools derived from these CLTs in order to make inference on the interaction matrices Γ and W and the limit probability $\tilde{P}_\infty(c)$. Section 4 provides two applications to real data sets, where, performing some suitable hypothesis tests, we try to evaluate the interaction intensities related to the interaction matrices in those empirical cases. The appendix contains the general results, with their proofs, and the general statistical methodology.

2. CENTRAL LIMIT THEOREMS

Assume $N \geq 2$. Let Γ, W, Λ be equal to the *non-negative* $N \times N$ square matrices with elements equal to the model parameters $\gamma_{j,h}, w_{j,h}$ (with $w_{h,h} > 0$) and $\lambda_{j,h}$ (with $\lambda_{h,h} > 0$), respectively. We recall that, by the balance condition (2) and the reparametrization (3), we have

$$(9) \quad W = \Gamma + \Lambda, \quad \mathbf{1}^\top \Gamma < \mathbf{1}^\top \quad \text{and} \quad \mathbf{1}^\top W = \mathbf{1}^\top,$$

where $\mathbf{1}$ denotes the vector with all the components equal to 1. As observed above, the matrix Γ rules the production of potential novelties and, in particular, its elements out of the diagonal regulate the interaction among the innovation processes with respect to this issue; while, the matrix W rules the interaction among the innovation processes with respect to the choice of an old item.

Moreover, set $\mathbf{D}_t^* = (D_{t,1}^*, \dots, D_{t,N}^*)^\top$, $\mathbf{D}_\infty^{**} = (D_{\infty,1}^{**}, \dots, D_{\infty,N}^{**})^\top$, $\mathbf{P}_t(c) = (P_t(1,c), \dots, P_t(N,c))^\top$ and finally $\mathbf{K}_t(c) = (K_t(1,c), \dots, K_t(N,c))^\top$.

In order to simplifying the statements of the following theorems, we summarize the required notation and conditions in the following two sets of assumptions:

Assumption 1. *Assume that the non-negative matrix $\Gamma = (\gamma_{j,h})_{j,h=1,\dots,N}$ is irreducible. Denote by $\gamma^* \in (0,1)$ the Perron-Frobenius eigenvalue of Γ , by \mathbf{v} the corresponding right eigenvector such that $\mathbf{v}^\top \mathbf{1} = 1$ and denote by \mathbf{u} the corresponding left eigenvector such that $\mathbf{v}^\top \mathbf{u} = 1$. Finally, let γ_2^* be the second eigenvalue of Γ with highest real part and assume that $\mathcal{R}e(\gamma_2^*)/\gamma^* < 1/2$.*

Assumption 2. *Assume that the non-negative matrix $W = (w_{j,h})_{j,h=1,\dots,N}$ is irreducible. Denote by \mathbf{v}_W the right eigenvector associated to the leading eigenvalue $w^* = 1$ such that $\mathbf{v}_W^\top \mathbf{1} = 1$. Finally, let w_2^* be the second eigenvalue of W with highest real part and assume that $\mathcal{R}e(w_2^*) < \frac{1}{2}$.*

We are now ready to provide the central limit theorems for the considered model, that describe the second-order behaviour of the quantities in Theorems 1.1 and 1.2. The symbol $\mathbf{0}$ denotes the vector with all the components equal to 0.

Theorem 2.1. *Under Assumption 1, we have*

$$t^{\gamma^*/2} \left(\frac{\mathbf{D}_t^*}{t^{\gamma^*}} - \mathbf{D}_\infty^{**} \right) \xrightarrow{\text{stably}} \mathcal{N}(\mathbf{0}, \tilde{Z}_\infty^{**} C_{\det,\Gamma}),$$

where \tilde{Z}_∞^{**} is an integrable strictly positive random variable such that $\mathbf{D}_\infty^{**} = \frac{\tilde{Z}_\infty^{**}}{\gamma^*} \mathbf{u}$ and $C_{\det,\Gamma}$ is a deterministic matrix depending on the eigen-structure of Γ , which is possible to write explicitly.

Proof. Denote by $X_{t,h}^*$ the random variable that takes value 1 when the ball extracted from urn h at time-step t has a new (for all the system) color and is equal to 0 otherwise. Then $Z_{t,h}^*$ coincides

with $P(X_{t+1,h}^* = 1 \mid \mathbf{C}_1, \dots, \mathbf{C}_t) = E[X_{t+1,h}^* \mid \mathbf{C}_1, \dots, \mathbf{C}_t]$ and $D_{t,h}^*$ can be written as $\sum_{n=1}^t X_{n,h}^*$. Since we have

$$Z_{t,h}^* = \frac{\theta_h + \sum_{n=1}^t \sum_{j=1}^N \gamma_{j,h} X_{n,j}^*}{\theta_h + t},$$

we obtain the following dynamics for $Z_{t,h}^*$:

$$Z_{0,h}^* = 1, \quad Z_{t+1,h}^* = (1 - r_{t,h})Z_{t,h}^* + r_{t,h} \sum_{j=1}^N \gamma_{j,h} X_{t+1,j}^* \quad \text{for } t \geq 0,$$

where $r_{t,h} = 1/(\theta_h + t + 1) = 1/(t + 1) + O(1/t^2)$ ². The corresponding vectorial dynamics for $\mathbf{Z}_t^* = (Z_{t,1}^*, \dots, Z_{t,N}^*)^\top$ is

$$\begin{aligned} \mathbf{Z}_0^* &= \mathbf{1} \\ (10) \quad \mathbf{Z}_{t+1}^* &= \left(1 - \frac{1}{t+1}\right) \mathbf{Z}_t^* + \frac{1}{t+1} \Gamma^\top \mathbf{X}_{t+1}^* + O(1/t^2) \mathbf{1} \\ &= \left[1 - \frac{1}{t+1} (I - \Gamma^\top)\right] \mathbf{Z}_t^* + \frac{1}{t+1} \Gamma^\top \Delta \mathbf{M}_{t+1}^* + O(1/t^2) \mathbf{1} \quad \text{for } t \geq 0, \end{aligned}$$

where $\Delta \mathbf{M}_{t+1}^* = \mathbf{X}_{t+1}^* - \mathbf{Z}_t^*$. Now, we fix $x > 0$ and set

$$\zeta_0(x) = 1, \quad \zeta_t(x) = \frac{\Gamma(t+x)}{\Gamma(t)} \sim t^x \uparrow + \infty.$$

More precisely, from Lemma 4.1 in [16] we have

$$\zeta_t(x) = t^x + O(t^{x-1}) \quad \text{and} \quad \frac{1}{\zeta_t(x)} = \frac{1}{t^x} + O(1/t^{x+1}),$$

and so

$$\begin{aligned} (11) \quad \frac{1}{\zeta_{t+1}(x) \zeta_{t+1}(1-x)} &= \left(\frac{1}{(t+1)^x} + O(1/t^{x+1}) \right) \left(\frac{1}{(t+1)^{1-x}} + O(1/t^{2-x}) \right) \\ &= \frac{1}{t+1} + O(1/t^2). \end{aligned}$$

Hence, multiplying (10) by $\zeta_{t+1}(1 - \gamma^*)$ and using the relation

$$\frac{\zeta_{t+1}(x)}{\zeta_t(x)} = \frac{\Gamma(t+x+1)}{\Gamma(t+1)} \frac{\Gamma(t)}{\Gamma(t+x)} = 1 + \frac{x}{t} = 1 + \frac{x}{t+1} + O(1/t^2),$$

with $x = 1 - \gamma^*$, we get the following dynamics for $\mathbf{Z}_t^{**} = \zeta_t(1 - \gamma^*) \mathbf{Z}_t^*$, where we set $\Delta \mathbf{M}_{t+1}^{**} = \zeta_t(1 - \gamma^*) \Delta \mathbf{M}_{t+1}^*$:

$$\begin{aligned} \mathbf{Z}_0^{**} &= \mathbf{1} \\ \mathbf{Z}_{t+1}^{**} &= \left[1 - \frac{1}{t+1} (I - \Gamma^\top)\right] \frac{\zeta_{t+1}(1 - \gamma^*)}{\zeta_t(1 - \gamma^*)} \mathbf{Z}_t^{**} + \frac{1}{t+1} \Gamma^\top \Delta \mathbf{M}_{t+1}^{**} + O\left(\frac{\zeta_{t+1}(1 - \gamma^*)}{t^2}\right) \mathbf{1} \\ &= \left[1 - \frac{1}{t+1} (I - \Gamma^\top)\right] \left(1 + \frac{1 - \gamma^*}{t+1} + O(1/t^2)\right) \mathbf{Z}_t^{**} + \frac{1}{t+1} \Gamma^\top \Delta \mathbf{M}_{t+1}^{**} + O\left(\frac{\zeta_{t+1}(1 - \gamma^*)}{t^2}\right) \mathbf{1} \\ &= \mathbf{Z}_t^{**} - \frac{1}{t+1} (\gamma^* I - \Gamma^\top) \mathbf{Z}_t^{**} + \frac{1}{t+1} \Gamma^\top \Delta \mathbf{M}_{t+1}^{**} + O\left(\frac{\zeta_{t+1}(1 - \gamma^*)}{t^2}\right) \mathbf{1}. \end{aligned}$$

²In all the paper, the notation $O(s_t)$ denotes a generic (possibly random) remainder term R_t such that $|R_t| \leq s_t$ eventually.

Moreover, recalling that $D_{t,h}^* = \sum_{n=1}^t X_{n,h}^*$, we find

$$\begin{aligned} \frac{\mathbf{D}_0^*}{\zeta_0(\gamma^*)} &= \mathbf{0} \\ \frac{\mathbf{D}_{t+1}^*}{\zeta_{t+1}(\gamma^*)} &= \frac{\zeta_t(\gamma^*)}{\zeta_{t+1}(\gamma^*)} \frac{\mathbf{D}_t^*}{\zeta_t(\gamma^*)} + \frac{1}{\zeta_{t+1}(\gamma^*)} \mathbf{X}_{t+1}^* \\ &= \left(1 - \frac{\gamma^*}{t+1}\right) \frac{\mathbf{D}_t^*}{\zeta_t(\gamma^*)} + \frac{1}{\zeta_{t+1}(\gamma^*)} \Delta \mathbf{M}_{t+1}^* + \frac{1}{\zeta_{t+1}(\gamma^*)} \mathbf{Z}_t^* \\ &= \left(1 - \frac{\gamma^*}{t+1}\right) \frac{\mathbf{D}_t^*}{\zeta_t(\gamma^*)} + \frac{1}{\zeta_{t+1}(\gamma^*) \zeta_{t+1}(1-\gamma^*)} \Delta \mathbf{M}_{t+1}^{**} + \frac{1}{\zeta_{t+1}(\gamma^*) \zeta_{t+1}(1-\gamma^*)} \frac{\zeta_{t+1}(1-\gamma^*)}{\zeta_t(1-\gamma^*)} \mathbf{Z}_t^{**}. \end{aligned}$$

Using (11) and the relation $\zeta_{t+1}(x)/\zeta_t(x) = 1 + O(1/t)$, and combining together with the previous dynamics of $(\mathbf{Z}_t^{**})_t$ and $(\mathbf{D}_{t+1}^*/\zeta_{t+1}(\gamma^*))_t$, we obtain

$$(12) \quad \begin{aligned} \mathbf{Z}_{t+1}^{**} &= \mathbf{Z}_t^{**} - \frac{1}{t+1} (\gamma^* I - \Gamma^\top) \mathbf{Z}_t^{**} + \frac{1}{t+1} \Gamma^\top \Delta \mathbf{M}_{t+1}^{**} + O(\zeta_{t+1}(1-\gamma^*)/t^2) \mathbf{1} \\ \frac{\mathbf{D}_{t+1}^*}{\zeta_{t+1}(\gamma^*)} &= \frac{\mathbf{D}_t^*}{\zeta_t(\gamma^*)} - \frac{1}{t+1} \left(\gamma^* \frac{\mathbf{D}_t^*}{\zeta_t(\gamma^*)} - \mathbf{Z}_t^{**} \right) + \frac{1}{t+1} \Delta \mathbf{M}_{t+1}^{**} + O(\zeta_t(1-\gamma^*)/t^2) \mathbf{1}, \end{aligned}$$

where $O(\zeta_t(1-\gamma^*)/t^2) = O(1/t^{1+\gamma^*})$. Moreover, we recall that in [5] (Theorem S1.1 and Lemma S1.2), we proved

$$\tilde{\mathbf{Z}}_t^{**} = \zeta_t(1-\gamma^*) \tilde{\mathbf{Z}}_t^* = \zeta_t(1-\gamma^*) \mathbf{v}^\top \mathbf{Z}_t^* \sim t^{1-\gamma^*} \mathbf{v}^\top \mathbf{Z}_t^* = t^{1-\gamma^*} \tilde{\mathbf{Z}}_t^* \xrightarrow{a.s./mean} \tilde{\mathbf{Z}}_\infty^{**},$$

where $\tilde{\mathbf{Z}}_\infty^{**}$ is an integrable strictly positive random variable and $\mathbf{D}_\infty^{**} = \frac{\tilde{\mathbf{Z}}_\infty^{**}}{\gamma^*} \mathbf{u}$. Therefore, we can apply Theorem A.4 with $\mathbf{A}_t = \mathbf{Z}_t^{**} = \zeta_t(1-\gamma^*) \mathbf{Z}_t^*$, $\mathbf{B}_t = \frac{\mathbf{D}_t^*}{\zeta_t(\gamma^*)}$, $\Phi = \Gamma$, $\phi^* = \gamma^*$, $\Sigma_\infty = \tilde{\mathbf{Z}}_\infty^{**} \text{diag}(\mathbf{u})$, $\Delta \mathbf{M}_{t+1} = \Delta \mathbf{M}_{t+1}^{**} = \zeta_t(1-\gamma^*) \Delta \mathbf{M}_{t+1}^* = \zeta_t(1-\gamma^*) (\mathbf{X}_{t+1}^* - \mathbf{Z}_t^*)$. Finally, we can conclude by observing that, since $\gamma^* < 1$, we have

$$t^{\gamma^*/2} \left(\frac{\mathbf{D}_t^*}{t^{\gamma^*}} - \frac{\mathbf{D}_t^*}{\zeta_t(\gamma^*)} \right) = t^{\gamma^*/2} \mathbf{D}_t^* O(1/t^{\gamma^*+1}) = \frac{\mathbf{D}_t^*}{t^{\gamma^*}} O(1/t^{1-\gamma^*/2}) \xrightarrow{a.s.} \mathbf{0}.$$

Note that in Theorem A.4 we have the explicit formulas for the elements of the matrix $C_{det,\Gamma}$: that is, $C_{det,\Gamma} = 1/\gamma^*$ when $N = 1$ and

$$C_{det,\Gamma} = \frac{1}{\gamma^*} (\mathbf{v}^\top \text{diag}(\mathbf{u}) \mathbf{v}) \mathbf{u} \mathbf{u}^\top + \frac{1}{\tilde{\mathbf{Z}}_\infty^{**}} \mathcal{M}_\infty^{33},$$

where \mathcal{M}_∞^{33} is the matrix defined in Theorem A.3 when $N \geq 2$ with the matrix Φ equal to Γ . Note that, since the expression for \mathcal{M}_∞^{33} contains the matrix Σ_∞ , and so the term $\tilde{\mathbf{Z}}_\infty^{**}$, we have that the matrix $\frac{1}{\tilde{\mathbf{Z}}_\infty^{**}} \mathcal{M}_\infty^{33}$ is deterministic. \square

Theorem 2.2. *Under Assumption 1, we have*

$$t^{\gamma^*/2} \left(\frac{\mathbf{D}_t^*}{\sum_{h=1}^N D_{t,h}^*} - \frac{\mathbf{u}}{\sum_{h=1}^N u_h} \right) \xrightarrow{\text{stably}} \mathcal{N} \left(\mathbf{0}, \frac{1}{\tilde{\mathbf{Z}}_\infty^{**}} \frac{(\gamma^*)^2}{(\sum_{j=1}^N u_j)^2} Q_\Gamma C_{det,\Gamma} Q_\Gamma^\top \right),$$

and

$$\sqrt{\sum_{h=1}^N D_{t,h}^*} \left(\frac{\mathbf{D}_t^*}{\sum_{h=1}^N D_{t,h}^*} - \frac{\mathbf{u}}{\sum_{h=1}^N u_h} \right) \xrightarrow{\text{stably}} \mathcal{N} \left(\mathbf{0}, \frac{\gamma^*}{\sum_{j=1}^N u_j} Q_\Gamma C_{det,\Gamma} Q_\Gamma^\top \right),$$

where $Q_\Gamma = I - \mathbf{u}\mathbf{1}^\top/\mathbf{1}^\top\mathbf{u}$ and $C_{det,\Gamma}$ is defined in Theorem 2.1. Moreover, for each j we have

$$t^{\gamma^*/2} \left(\frac{\mathbf{D}_t^*}{D_{t,j}^*} - \frac{\mathbf{u}}{u_j} \right) \xrightarrow{\text{stably}} \mathcal{N} \left(\mathbf{0}, \frac{1}{\tilde{Z}_\infty^{**}} \frac{(\gamma^*)^2}{u_j^2} Q_j C_{det,\Gamma} Q_j^\top \right),$$

and

$$\sqrt{D_{t,j}^*} \left(\frac{\mathbf{D}_t^*}{D_{t,j}^*} - \frac{\mathbf{u}}{u_j} \right) \xrightarrow{\text{stably}} \mathcal{N} \left(\mathbf{0}, \frac{\gamma^*}{u_j} Q_j C_{det,\Gamma} Q_j^\top \right),$$

where $Q_j = I - \mathbf{u}\mathbf{e}_j^\top/u_j$ with \mathbf{e}_j the j th vector of the canonical base and $C_{det,\Gamma}$ is defined in Theorem 2.1.

Proof. We have to apply Corollary A.6 in appendix with $\mathbf{A}_t = \mathbf{Z}_t^{**} = \zeta_t(1 - \gamma^*)\mathbf{Z}_t^*$, $\mathbf{B}_t = \frac{\mathbf{D}_t^*}{\zeta_t(\gamma^*)}$, $\mathbf{x} = \mathbf{1}$, $\tilde{A}_\infty = \tilde{Z}_\infty^{**}$ and $\phi^* = \gamma^*$. Note that, as specified in the proof of this corollary, we have $\tilde{Z}_\infty^{**} Q_\Gamma C_{det,\Gamma} Q_\Gamma^\top = Q_\Gamma \mathcal{M}_\infty^{33} Q_\Gamma^\top$, where \mathcal{M}_∞^{33} is the matrix defined in Theorem A.3 when $N \geq 2$ with the matrix Φ equal to Γ . Similarly, we obtain the second part of the statement by means of Corollary A.6 with $\mathbf{x} = \mathbf{e}_j$. \square

Theorem 2.3. *Under Assumption 2, we have*

$$\sqrt{t} \left(\frac{\mathbf{K}_t(c)}{t} - \tilde{P}_\infty(c)\mathbf{1} \right) \xrightarrow{\text{stably}} \mathcal{N}(\mathbf{0}, \tilde{P}_\infty(c)(1 - \tilde{P}_\infty(c))C_{det,W}),$$

where $\tilde{P}_\infty(c) = a.s. - \lim_t P_t(h, c) = a.s. - \lim_t K_t(h, c)/t$ for each $h = 1, \dots, N$, $\tilde{P}_\infty(c) \in (0, 1)$ a.s. and $C_{det,W}$ is a deterministic matrix depending on the eigen-structure of W , which is possible to write explicitly.

Proof. For any color c already present in the network at time t , we can write

$$P_t(h, c) = \frac{\sum_{n=1}^t \sum_{j=1}^N w_{j,h} \Delta K_n(j, c)}{\theta_h + t} - \frac{\gamma_{j^*(c),h}}{\theta_h + t},$$

where $\Delta K_n(j, c) = K_n(j, c) - K_{n-1}(j, c)$ takes values in $\{0, 1\}$ and $E[\Delta K_{n+1}(j, c)|\text{past}] = P_n(j, c)$. Then, we obtain the following dynamics for $P_t(h, c)$:

$$P_{t+1}(h, c) = (1 - r_{t,h})P_t(h, c) + r_{t,h} \sum_{j=1}^N w_{j,h} \Delta K_{t+1}(j, c),$$

where $r_{t,h} = 1/(\theta_h + t + 1) = 1/(t + 1) + O(1/t^2)$. Thus the corresponding vectorial dynamics for $\mathbf{P}_t(c) = (P_t(1, c), \dots, P_t(N, c))^\top$ is

$$\begin{aligned} \mathbf{P}_{t^*(c)}(c) \neq \mathbf{0}, \quad \mathbf{P}_{t+1}(c) &= \left(1 - \frac{1}{t+1}\right) \mathbf{P}_t(c) + \frac{1}{t+1} W^\top \Delta \mathbf{K}_{t+1}(c) + O(1/t^2)\mathbf{1} \\ &= \mathbf{P}_t(c) - \frac{1}{t+1} (I - W^\top) \mathbf{P}_t(c) + \frac{1}{t+1} W^\top \Delta \mathbf{M}_{t+1}(c) + O(1/t^2)\mathbf{1} \quad \text{for } t \geq t^*(c), \end{aligned}$$

where $t^*(c)$ is the time-step of the first extraction of c in the system, $\Delta \mathbf{K}_t(c) = (\Delta K_t(1, c), \dots, \Delta K_t(N, c))^\top$ and $\Delta \mathbf{M}_{t+1}(c) = \Delta \mathbf{K}_{t+1}(c) - \mathbf{P}_t(c)$.

Considering also the dynamics of $\mathbf{K}_t(c)/t$, we obtain for $t \geq t^*(c)$

$$(13) \quad \begin{aligned} \mathbf{P}_{t+1}(c) &= \mathbf{P}_t(c) - \frac{1}{t+1} (I - W^\top) \mathbf{P}_t(c) + \frac{1}{t+1} W^\top \Delta \mathbf{M}_{t+1}(c) + O(1/t^2)\mathbf{1} \\ \frac{\mathbf{K}_{t+1}(c)}{t+1} &= \frac{\mathbf{K}_t(c)}{t} - \frac{1}{t+1} \left(\frac{\mathbf{K}_t(c)}{t} - \mathbf{P}_t(c) \right) + \frac{1}{t+1} \Delta \mathbf{M}_{t+1}(c). \end{aligned}$$

Moreover, from Theorem 1.2, we have $\tilde{P}_t(c) = \mathbf{v}_W^\top \mathbf{P}_t(c) \xrightarrow{a.s.} \tilde{P}_\infty(c)$, and this convergence is also in mean because $(\tilde{P}_t(c))_t$ is uniformly bounded. Therefore, we can apply Theorem A.4 with $\mathbf{A}_t = \mathbf{P}_t(c)$, $\mathbf{B}_t = \mathbf{K}_t(c)/t$, $\Phi = W$, $\phi^* = w^* = 1$, $\mathbf{u} = \mathbf{u}_W = \mathbf{1}$, $\mathbf{v} = \mathbf{v}_W$, $\Sigma_\infty = \tilde{P}_\infty(c)(1 - \tilde{P}_\infty(c))I$, $\Delta \mathbf{M}_{t+1} = \Delta \mathbf{M}_{t+1}(c) = \Delta \mathbf{K}_{t+1}(c) - \mathbf{P}_t(c)$. Note that in Theorem A.4 we have the explicit formulas for the elements of the matrix $C_{det,W}$: that is, $C_{det,W} = 1$ when $N = 1$ and

$$C_{det,W} = (\mathbf{v}_W^\top \mathbf{v}_W) \mathbf{1} \mathbf{1}^\top + \frac{1}{\tilde{P}_\infty(c)(1 - \tilde{P}_\infty(c))} \mathcal{M}_\infty^{33},$$

where \mathcal{M}_∞^{33} is the matrix defined in Theorem A.3 when $N \geq 2$ with the matrix Φ equal to W . Note that, since the expression for \mathcal{M}_∞^{33} contains the matrix Σ_∞ , and so the factor $\tilde{P}_\infty(c)(1 - \tilde{P}_\infty(c))$, we have that the matrix $\frac{1}{\tilde{P}_\infty(c)(1 - \tilde{P}_\infty(c))} \mathcal{M}_\infty^{33}$ is deterministic. \square

Theorem 2.4. *Under Assumption 2, we have*

$$\sqrt{t} \left(\frac{\mathbf{K}_t(c)}{\sum_{h=1}^N K_t(h, c)} - \frac{1}{N} \mathbf{1} \right) \xrightarrow{stably} \mathcal{N} \left(\mathbf{0}, \frac{(1 - \tilde{P}_\infty(c))}{N^2 \tilde{P}_\infty(c)} Q_W C_{det,W} Q_W^\top \right),$$

and

$$\sqrt{\frac{\sum_{h=1}^N K_t(h, c)}{1 - (Nt)^{-1} \sum_{h=1}^N K_t(h, c)}} \left(\frac{\mathbf{K}_t(c)}{\sum_{h=1}^N K_t(h, c)} - \frac{1}{N} \mathbf{1} \right) \xrightarrow{stably} \mathcal{N} \left(\mathbf{0}, \frac{1}{N} Q_W C_{det,W} Q_W^\top \right).$$

where $Q_W = (I - N^{-1} \mathbf{1} \mathbf{1}^\top)$ and $C_{det,W}$ is defined in Theorem 2.3.

Proof. We have to apply Corollary A.6 in appendix with $\mathbf{A}_t = \mathbf{P}_t(c)$, $\mathbf{B}_t = \mathbf{K}_t(c)/t$, $\mathbf{x} = \mathbf{1}$, $\tilde{A}_\infty = \tilde{P}_\infty(c)$, $\phi^* = w^* = 1$ and $\mathbf{u} = \mathbf{u}_W = \mathbf{1}$. Note that, as specified in the proof of this corollary, we have $\tilde{P}_\infty(c)(1 - \tilde{P}_\infty(c)) Q_W C_{det,W} Q_W^\top = Q_W \mathcal{M}_\infty^{33} Q_W^\top$, where \mathcal{M}_∞^{33} is the matrix defined in Theorem A.3 when $N \geq 2$ with the matrix Φ equal to W . \square

3. STATISTICAL INFERENCE

We here present some statistical tools based on the results of Section 2. More precisely, we provide a statistical test for each of the two interaction matrices, Γ and W , based on the observable processes \mathbf{D}_t^* and $\mathbf{K}_t(c)$. Furthermore, we use the latter process to provide also a confidence interval for the limit probability $\tilde{P}_\infty(c)$. The general procedure to be followed in order to obtain these statistical instruments is described in Sections B and C of the appendix. In particular, in our framework, we have to apply it with $\mathbf{B}_t = \frac{\mathbf{D}_t^*}{\zeta_t(\gamma^*)}$ (or, equivalently $\mathbf{B}_t = \frac{\mathbf{D}_t^*}{t\gamma^*}$) and with $\mathbf{B}_t = \mathbf{K}_t(c)/t$. To this end, we observe that Assumption 3 is true in both cases with $\Sigma_{det} = \Sigma_{det,\Gamma} = \text{diag}(\mathbf{u})$ in the first one and $\Sigma_{det} = \Sigma_{det,W} = I$ in the second one.

We firstly focus on the case $N = 2$ and, then, we consider a particular type of interaction (mean-field interaction) in the general case $N \geq 2$. It is worthwhile to underline that the statistical tools described in Section B are very general and so it is also possible to explicit the formulas for any family of interaction matrices.

3.1. The case $N = 2$.

3.1.1. *Hypothesis test on Γ .* As the parameters $\gamma^* \in (0, 1)$ and $r = u_1/u_2 > 0$ can be estimated with high accuracy by means of a suitable regression on \mathbf{D}_t^* (see [5] and references therein), in what follows we will assume them known. Therefore, we can use the reparametrization of the matrix Γ given by the constrains

$$r\gamma_{1,1} + \gamma_{2,1} = r\gamma^* \quad \text{and} \quad r\gamma_{1,2} + \gamma_{2,2} = \gamma^*.$$

and make inference on the other two parameters $\gamma_{1,2}$ and $\gamma_{2,1}$ (that have to be such that the assumptions on the matrix Γ are verified). If we use the parametrization $\iota = \gamma_{1,2} + \gamma_{2,1}$ and $\eta = \frac{\gamma_{1,2}}{\gamma_{1,2} + \gamma_{2,1}}$ so that ι rules the intensity of the whole interaction between the two processes due to matrix Γ , while η (resp. $(1 - \eta)$) denotes the percentage of this intensity due to the influence of process 1 on process 2 (resp., of process 2 on process 1), the expressions for the elements of the matrix Γ become

$$(14) \quad \gamma_{1,1} = \gamma^* - \frac{(1 - \eta)\iota}{r}, \quad \gamma_{1,2} = \eta\iota, \quad \gamma_{2,1} = (1 - \eta)\iota \quad \text{and} \quad \gamma_{2,2} = \gamma^* - r\eta\iota,$$

where (by the required conditions on Γ)

$$(15) \quad \eta \in (0, 1) \quad \text{and} \quad \iota \in J_\eta = \left(0, \min\left(\frac{\gamma^*r}{1-\eta}, \frac{\gamma^*}{r\eta}\right)\right] \cap \left(0, \frac{1-\gamma^*}{\eta(1-r)}I_{(r<1)} + \frac{1-\gamma^*}{(1-\eta)(1-1/r)}I_{(r>1)} + \infty I_{(r=1)}\right).$$

Therefore, for a two-sided test with $H_0 : \iota = \iota_0, \eta = \eta_0$, where ι_0 and η_0 satisfy (15), i.e. $\iota_0 \in J_{\eta_0}$, and the required condition in Assumption 1 for the second eigenvalue of Γ , that is $\iota_0 \frac{(1-\eta_0)+r^2\eta_0}{r} > \gamma^*/2$, we can use Theorem 2.1 and obtain the test statistic (see (35) from Remark 3 in the appendix, with $\phi = \gamma^*$, $\phi_2^* = \gamma_2^* = \gamma^* - [(1 - \eta_0)\iota_0 + r^2\eta_0\iota_0]/r$, $g(x) = x$, and $\mathbf{B}_t = \frac{\mathbf{D}_t^*}{\zeta_t(\gamma^*)}$ or, equivalently $\mathbf{B}_t = \frac{\mathbf{D}_t^*}{t\gamma^*}$)

$$(16) \quad q_{r,\eta_0} \Delta_0 \frac{(D_{1,t}^* - rD_{2,t}^*)^2}{\tilde{D}_t^*} \stackrel{d}{\sim} \chi^2(1) \quad \text{under } H_0,$$

where $\tilde{D}_t^* = \mathbf{v}^\top \mathbf{D}_t^* = \frac{\eta_0 r D_{1,t}^* + (1-\eta_0) D_{2,t}^*}{\eta_0 r + (1-\eta_0)}$, $q_{r,\eta_0} = \frac{2[(1-\eta_0)+\eta_0 r^2]}{r(1+r)[(1-\eta_0)+\eta_0 r]}$ and

$$\Delta_0 = \frac{1}{2} - \frac{\gamma_2^*}{\gamma^*} = \frac{\iota_0 (1 - \eta_0) + \eta_0 r^2}{r} - \frac{1}{2}.$$

It is immediate to see that the test statistic (16) has an increasing dependence on the parameter ι_0 , while its behaviour with respect the parameter η_0 is not clear. However, we can observe that it can be rewritten as

$$\frac{2}{r(1+r)} \frac{(1 - \eta_0) + \eta_0 r^2}{(1 - \eta_0) + \eta_0 r D_{1,t}^*/D_{2,t}^*} \Delta_0 D_{2,t}^* (D_{1,t}^*/D_{2,t}^* - r)^2,$$

where, for any value of η_0 , we have

$$\frac{(1 - \eta_0) + \eta_0 r^2}{(1 - \eta_0) + \eta_0 r D_{1,t}^*/D_{2,t}^*} \xrightarrow{a.s.} 1.$$

Hence, we can simplify the test statistic as

$$(17) \quad \frac{2}{r(1+r)} \Delta_0 D_{2,t}^* (D_{1,t}^*/D_{2,t}^* - r)^2 \stackrel{d}{\sim} \chi^2(1) \quad \text{under } H_0,$$

where the dependence on ι_0 and η_0 is only in the factor Δ_0 . Notice also that in the present framework the two eigenvalues γ^* and γ_2^* are both real and so the quantity Δ_0 represents the distance of the ratio between the second and the leading eigenvalue from the critical value $1/2$ (see condition $\mathcal{R}e(\gamma_2^*)/\gamma^* < 1/2$ in Assumption 1). Moreover, Δ_0 is the only quantity in the test

statistics (17) that contains the parameters ι_0 and η_0 . Moreover, it is neither random nor time-dependent. The random quantity $D_{2,t}^*(D_{1,t}^*/D_{2,t}^* - r)^2 = [\sqrt{D_{2,t}^*}(D_{1,t}^*/D_{2,t}^* - r)]^2$ describes the fluctuations of $D_{1,t}^*/D_{2,t}^*$ around r (note that (17) is coherent with the convergence result stated in Theorem 2.2). Since Δ_0 , and so the statistic (17), is increasing in ι_0 , we can replace the condition $\iota = \iota_0$ in H_0 by $\iota \geq \iota_0$. Further, the quantity Δ_0 is increasing in η_0 when $r > 1$ and decreasing in η_0 when $r < 1$. Hence, the statistic (17) works well also if we replace the condition $\eta = \eta_0$ in H_0 by $\eta \geq \eta_0$ when $r > 1$ or by $\eta \leq \eta_0$ when $r < 1$. Therefore, if we set $\eta_0 = 1/2$, we get

$$\Delta_0 = \frac{1}{2} - \frac{\gamma_2^*}{\gamma^*} = \frac{\iota_0}{\gamma^*} \frac{1+r^2}{2r} - \frac{1}{2}$$

and we can perform a one-sided test with one of the following two null hypotheses, according to the value of r :

- Case $r > 1$: $H_0 : \iota \geq \iota_0, \eta \geq 1/2$ (the interaction intensity is at least ι_0 and the processes equally influence each other or the interaction is mostly due to the influence of process 1 on process 2), where ι_0 satisfies (15) for some η in H_0 , i.e. $\iota_0 \in \cup_{\eta \geq 1/2} J_\eta$, and $\iota_0 > \gamma^* \frac{r}{1+r^2}$;
- Case $r < 1$: $H_0 : \iota \geq \iota_0, \eta \leq 1/2$ (the interaction intensity is at least ι_0 and the processes equally influence each other or the interaction is mostly due to the influence of process 2 on process 1), where ι_0 satisfies (15) for some η in H_0 , i.e. $\iota_0 \in \cup_{\eta \leq 1/2} J_\eta$, and $\iota_0 > \gamma^* \frac{r}{1+r^2}$.

Finally, in the very special case when $r = 1$, the test statistic (17) does not depend on the value of η_0 and it can be used for testing the intensity ι only. Regarding the power, following Remark 2 in the appendix, we find that, under the alternative $H_1 : \iota = \iota_1, \eta = \eta_1$, with $\iota_1 \in J_{\eta_1}$ and $\iota_1 \frac{(1-\eta_1)+r^2\eta_1}{r} > \gamma^*/2$, the statistic (17) is asymptotically distributed as

$$\frac{\Delta_0}{\Delta_1} \chi^2(1) = \frac{\frac{\iota_0}{\gamma^*} \frac{(1-\eta_0)+\eta_0 r^2}{r} - \frac{1}{2}}{\frac{\iota_1}{\gamma^*} \frac{(1-\eta_1)+\eta_1 r^2}{r} - \frac{1}{2}} \chi^2(1).$$

As a consequence, the power increases as the value of ι_1 decreases and it tends toward 1 as $\Delta_1 \rightarrow 0$, that is as ι_1 approaches its minimum admissible value $\frac{r}{(1-\eta_1)+\eta_1 r^2} \frac{\gamma^*}{2}$. Moreover, the power increases as η_1 decreases if $r > 1$ or as η_1 increases if $r < 1$. We must highlight that the term $\frac{\Delta_0}{\Delta_1}$ is time-invariant and this means that increasing the time t does not increase the power.

3.1.2. *Hypothesis test on W* . As $W^\top \mathbf{1} = \mathbf{1}$ (i.e. $w^* = 1$ and $r = 1$), we have

$$w_{1,1} = 1 - w_{2,1} \quad \text{and} \quad w_{2,2} = 1 - w_{1,2}$$

and so we can make inference on the two parameters $w_{2,1}$ and $w_{1,2}$. Using the parametrization $\iota = w_{1,2} + w_{2,1}$ for the intensity of the interaction due to the matrix W , and $\eta = \frac{w_{1,2}}{w_{1,2}+w_{2,1}}$ (resp. $(1-\eta)$) for the percentage of this intensity due to the influence of process 1 on process 2 (resp., of process 2 on process 1), we have

$$(18) \quad w_{1,1} = 1 - (1-\eta)\iota, \quad w_{1,2} = \eta\iota, \quad w_{2,1} = (1-\eta)\iota \quad \text{and} \quad w_{2,2} = 1 - \eta\iota,$$

where (by the required conditions on W)

$$(19) \quad \eta \in (0, 1) \quad \text{and} \quad \iota \in (0, \min(1/\eta, 1/(1-\eta))) \subseteq (0, 2).$$

Then, for a two-sided test with $H_0 : \iota = \iota_0, \eta = \eta_0$, where (19) and $\iota_0 > 1/2$ (the required condition in Assumption 2 for the second eigenvalue of W) are satisfied, we can use Theorem 2.3

and obtain the test statistic (see (35) from Remark 3 in the appendix, with $\phi = w^* = 1$, $r = 1$, $\phi_2^* = 1 - \iota_0$, $g(x) = x(1 - x)$ and $\mathbf{B}_t = \mathbf{K}_t(c)/t$ with c an arbitrary old color)

$$(20) \quad \Delta_0 \frac{(K_{1,t}(c) - K_{2,t}(c))^2}{\tilde{K}_t(c)(1 - \tilde{K}_t(c)/t)} \stackrel{d}{\sim} \chi^2(1) \quad \text{under } H_0, \quad \text{where} \quad \Delta_0 = \frac{1}{2} - (1 - \iota_0) = \iota_0 - \frac{1}{2},$$

and

$$(21) \quad \tilde{K}_t(c) = \mathbf{v}^\top \mathbf{K}_t(c) = \eta_0 K_{1,t}(c) + (1 - \eta_0) K_{2,t}(c).$$

Since, for any η_0 and η_1 , we have

$$\frac{\eta_1 K_{1,t}(c) + (1 - \eta_1) K_{2,t}(c)}{\tilde{K}_t(c)} \frac{(1 - \eta_1 K_{1,t}(c)/t + (1 - \eta_1) K_{2,t}(c)/t)}{(1 - \tilde{K}_t(c)/t)} \xrightarrow{a.s.} 1,$$

the choice of the value of η_0 in H_0 does not affect the asymptotic distribution of the test statistic. This means that we can use the test statistic (20) with an arbitrary value of η_0 in order to perform a test on the intensity ι . In particular, if we choose $\eta_0 = 1/2$, we simply have $\tilde{K}_t(c) = \frac{K_{1,t}(c) + K_{2,t}(c)}{2}$. In addition, note that the statistic (20) is increasing in ι_0 and so it works well also for a one-sided test with condition $\iota = \iota_0$ in H_0 replaced by $\iota \geq \iota_0$, where $\iota_0 \in (\frac{1}{2}, 2)$. Regarding the power, following Remark 2 in the appendix, we find that, under the alternative $H_1 : \iota = \iota_1$, where $\iota_1 \in (\frac{1}{2}, 2)$ with $\iota_1 \neq \iota_0$, the statistic (20) is asymptotically distributed as

$$\frac{\Delta_0}{\Delta_1} \chi^2(1) = \frac{\iota_0 - \frac{1}{2}}{\iota_1 - \frac{1}{2}} \chi^2(1).$$

Therefore, the power is monotone decreasing in ι_1 , and it tends to 1 as ι_1 approaches its lower bound $1/2$. However, the term $\frac{\Delta_0}{\Delta_1}$ is time-invariant and so, as before, increasing the time t does not increase the power.

3.1.3. Confidence interval for $\tilde{P}_\infty(c)$. Fix the confidence level $(1 - \alpha)$ and denote by z_α the quantile of order $(1 - \alpha/2)$ of the standard normal distribution, that is $\mathcal{N}(0, 1)(z_\alpha, +\infty) = \alpha/2$. Assuming $\iota > 1/2$, we can use Theorem 2.3 in order to construct a confidence interval for the random limit $\tilde{P}_\infty(c)$ associated to a given item c (see the interval (36) from Remark 3 in the appendix, with $\phi = w^* = 1$, $r = 1$, $g(x) = x(1 - x)$ and $\mathbf{B}_t = \mathbf{K}_t(c)/t$). Precisely, we obtain:

$$(22) \quad CI_{1-\alpha}(\tilde{P}_\infty(c)) = \frac{\tilde{K}_t(c)}{t} \pm \frac{z_\alpha}{t^{1/2}} \sqrt{\frac{\tilde{K}_t(c)}{t} \left(1 - \frac{\tilde{K}_t(c)}{t}\right)} c_\eta,$$

with $c_\eta = \eta^2 + (1 - \eta)^2$ and $\tilde{K}_t(c)$ defined as in (21). It is interesting to notice the absence of the dependence on the specific value of ι . On the other hand, while the dependence of $\tilde{K}_t(c)$ on η is not important, because $\frac{\eta_1 K_{1,t}(c)/t + (1 - \eta_1) K_{2,t}(c)/t}{\tilde{K}_t(c)/t} \xrightarrow{a.s.} 1$, the presence of the quantity c_η implies that we need to know the value of the parameter η . However, c_η is always smaller than 1 for $\eta \in (0, 1)$ and hence, if we do not know the value of η , we can use $\tilde{K}_t(c)$ with $\eta = 1/2$, so that we have $\tilde{K}_t(c) = \frac{K_{1,t}(c) + K_{2,t}(c)}{2}$, and take the largest interval obtained from (22) with c_η replaced by 1. As an example, we have simulated $S = 200$ independent pair of innovation processes with Γ and W defined with $r = 0.75$, $\gamma^* = 0.75$, $\eta = 1/2$, $\iota_\Gamma = 1$ and $\iota_W = 1.25$ and then, for any simulation, we have constructed the confidence intervals (22) of level $(1 - \alpha) = 95\%$ computed at time-step $t = 10^3$ associated to the item c with highest $(K_{t,1}(c) + K_{t,2}(c))$ (recall that the interval has the same expression for any given item c). In Figure 1 we can see these $S = 200$ confidence intervals (sorted according to the value of $K_{t,1}(c) + K_{t,2}(c)$) and the estimates of the corresponding $\tilde{P}_\infty(c)$ computed as $\frac{K_{t_\infty,1}(c) + K_{t_\infty,2}(c)}{2t_\infty}$ at time-step $t_\infty = 10^5$. The empirical coverage is represented by the




FIGURE 1. Confidence interval (see (22)) for the random variable $\tilde{P}_\infty(c)$ at time-step $t = 10^3$ and related to item c with highest $\mathbf{1}^\top \mathbf{K}_t(c)$ in the case $N = 2$, Γ defined as in (14) with $r = 0.75$, $\gamma^* = 0.75$, $\eta = 1/2$ and $\iota_\Gamma = 1$, and W defined as in (18) with $\eta = 1/2$ and $\iota_W = 1.25$. Number of simulations $S = 200$ (sorted according to the value of $\mathbf{1}^\top \mathbf{K}_t(c)$). The crosses represent the value of $\tilde{P}_\infty(c)$ estimated by $\mathbf{1}^\top \mathbf{K}_{t_\infty}(c)/(Nt_\infty)$ at time-step $t_\infty = 10^5$.

proportion of times the estimates of $\tilde{P}_\infty(c)$ belong to the corresponding confidence interval, and in this simulation this is equal to 0.955, which is very close to the nominal value of 0.95.

3.2. Mean-field interaction for $N \geq 2$.

3.2.1. *Hypothesis test on Γ .* Suppose that Γ is of the form

$$(23) \quad \gamma_{j,h} = \phi(\iota/N + \delta_{j,h}(1 - \iota)) \quad \phi, \iota \in (0, 1].$$

Note that the leading eigenvalue γ^* coincides with ϕ and so, since γ^* can be estimated with high accuracy by a suitable regression on \mathbf{D}_t^* (see [5] and references therein), we can assume that the parameter ϕ is known and we can make inference only on the parameter ι that rules the intensity of the interaction. For a two-sided test with $H_0 : \iota = \iota_0$, where $\iota_0 \leq 1$ and $\iota_0 > 1/2$ (the required condition in Assumption 1 for the second eigenvalue of Γ), we can use Theorem 2.1 and obtain the test statistic (see (38) from Remark 4 in the appendix, with $\phi = \gamma^*$, $\phi_2^* = \gamma^*(1 - \iota_0)$, $g(x) = x$, and $\mathbf{B}_t = \frac{\mathbf{D}_t^*}{\zeta_t(\gamma^*)}$ or, equivalently $\mathbf{B}_t = \frac{\mathbf{D}_t^*}{t\gamma^*}$)

$$(24) \quad 2\Delta_0 \frac{\|\mathbf{D}_t^* - \tilde{D}_t^* \mathbf{1}\|^2}{\tilde{D}_t^*} \stackrel{d}{\sim} \chi^2(N-1) \quad \text{under } H_0, \quad \text{where} \quad \Delta_0 = \frac{1}{2} - \frac{\gamma^*(1 - \iota_0)}{\gamma^*} = \iota_0 - \frac{1}{2}$$

and $\tilde{D}_t^* = \mathbf{v}^\top \mathbf{D}_t^* = \frac{\mathbf{1}^\top \mathbf{D}_t^*}{N}$. Note that this statistic is increasing in ι_0 and so it works well also for the one-sided test with $H_0 : \iota \geq \iota_0$, where $\iota_0 \in (1/2, 1]$. Regarding the power, following Remark 2 in the appendix, we observe that, under the alternative $H_1 : \iota = \iota_1$, with $\iota_1 \in (1/2, 1]$, the statistics (24)

is asymptotically distributed as

$$\frac{\Delta_0}{\Delta_1} \chi^2(N-1) = \frac{\iota_0 - \frac{1}{2}}{\iota_1 - \frac{1}{2}} \chi^2(N-1).$$

Therefore, the power is monotone decreasing in ι_1 , and it tends to 1 as ι_1 approaches its lower bound $1/2$. Moreover, as before, increasing the time t does not increase the power. We also highlight that the term $\frac{\Delta_0}{\Delta_1}$ does not depend on N and so increasing the number of innovations processes would not increase the power either.

3.2.2. Hypothesis test on W . Suppose that W is of the form

$$(25) \quad w_{j,h} = \iota/N + \delta_{j,h}(1-\iota) \quad \iota \in (0, 1].$$

For a two-sided test with $H_0 : \iota = \iota_0$, where $\iota_0 \leq 1$ and $\iota_0 > 1/2$ (the required condition in Assumption 2 for the second eigenvalue of W), we can use Theorem 2.3 and obtain the test statistic (see (38) from Remark 4 in the appendix, with $\phi = w^* = 1$, $\phi_2^* = w^*(1-\iota_0) = (1-\iota_0)$, $g(x) = x(1-x)$, and $\mathbf{B}_t = \mathbf{K}_t(c)/t$)

$$(26) \quad 2\Delta_0 \frac{\|\mathbf{K}_t(c) - \tilde{K}_t(c)\mathbf{1}\|^2}{\tilde{K}_t(c)(1 - \tilde{K}_t(c)/t)} \stackrel{d}{\sim} \chi^2(N-1) \quad \text{under } H_0, \quad \text{where} \quad \Delta_0 = \frac{1}{2} - (1-\iota_0) = \iota_0 - \frac{1}{2},$$

c is an arbitrary observed item and $\tilde{K}_t(c) = \mathbf{v}^\top \mathbf{K}_t(c) = \frac{\mathbf{1}^\top \mathbf{K}_t(c)}{N}$. This statistic is increasing in ι_0 and so it works well also for the one-sided test with $H_0 : \iota \geq \iota_0$, where $\iota_0 \in (1/2, 1]$. Regarding the power, following Remark 2 in the appendix, we observe that, under the alternative $H_1 : \iota = \iota_1$, with $\iota_1 \in (1/2, 1]$, the statistics (26) is asymptotically distributed exactly as the previous statistics (24) and so the same considerations on the power of the test hold true.

3.2.3. Confidence interval for $\tilde{P}_\infty(c)$. Fix the confidence level $(1-\alpha)$ and denote by z_α the quantile of order $(1-\alpha/2)$ of the standard normal distribution, that is $\mathcal{N}(0, 1)(z_\alpha, +\infty) = \alpha/2$. Assuming $\iota \in (1/2, 1]$, we can use Theorem 2.3 in order to construct a confidence interval for the random limit $\tilde{P}_\infty(c)$ associated to a given item c (see the interval (37) from Remark 4 in the appendix, with $\phi = w^* = 1$, $g(x) = x(1-x)$ and $\mathbf{B}_t = \mathbf{K}_t(c)/t$). Precisely, we obtain:

$$(27) \quad CI_{1-\alpha}(\tilde{P}_\infty(c)) = \frac{\tilde{K}_t(c)}{t} \pm \frac{z_\alpha}{t^{1/2}} \sqrt{\frac{\tilde{K}_t(c)}{t} \left(1 - \frac{\tilde{K}_t(c)}{t}\right) \frac{1}{N}},$$

where $\tilde{K}_t(c) = \mathbf{v}^\top \mathbf{K}_t(c) = \frac{\mathbf{1}^\top \mathbf{K}_t(c)}{N}$. Note that this interval does not depend on the value of the parameter ι .

As an example, we have simulated $S = 200$ independent triplets of innovation processes with Γ and W of the mean-field type defined with $N = 3$, $\gamma^* = 0.75$, $\iota_\Gamma = \iota_W = 0.8$, and we show the confidence intervals (27) of level $(1-\alpha) = 95\%$ at time-step $t = 10^3$, associated to the item c with highest $\mathbf{1}^\top \mathbf{K}_t(c)$ (see Figure 2). In this case, the empirical coverage is equal to 0.970, which is again very close to the nominal value 0.95.

4. APPLICATIONS

In this section we apply the proposed inferential methodology on the two real data sets, already considered in [5], to test the minimum level of interaction presented by the matrices Γ and W in these specific contexts. In particular, the first one is taken from the social content aggregation website Reddit, collected, elaborated and made freely available on the web by the authors of [25]




FIGURE 2. Confidence interval (see (27)) for the random variable $\tilde{P}_\infty(c)$ at time-step $t = 10^3$ and related to item c with highest $\mathbf{1}^\top \mathbf{K}_t(c)$ in the case $N = 3$, Γ defined as in (23) with $\phi = 0.75$ and $\iota_\Gamma = 0.8$, and W defined as in (25) with $\iota_W = 0.8$. Number of simulations $S = 200$ (sorted according to the value of $\mathbf{1}^\top \mathbf{K}_t(c)$). The crosses represent the value of $\tilde{P}_\infty(c)$ estimated by $\mathbf{1}^\top \mathbf{K}_{t_\infty}(c)/(Nt_\infty)$ at time-step $t_\infty = 10^5$.

(see <https://github.com/corradomonti/demographic-homophily>), while the second one is downloaded from the on-line library Project Gutenberg, which is a collection of public domain books (see <https://www.gutenberg.org>).

4.1. Reddit data set. This data set consists of a collection of news, and comments associated to each news, for the period 2016 – 2020, downloaded from the r/news community on the website Reddit [25]. Each news is associated with the author who posted it. Moreover, the data set contains the specific topic the news belongs to and, to each comment is also assigned a measurement of the sentiment, expressed as a real value in $(-1, 1)$. We consider only the comments to news belonging to the topic "Politics". Moreover, we categorize the sentiment variable: precisely, we define it as "positive" if the provided sentiment value was larger than $+0.35$ and "negative" if the provided sentiment value was lower than -0.35 . (We remove any comment with an original sentiment value that lies within $(-0.35, +0.35)$.) We finally take a data set of 2602173 comments for each of the two categories. Hence, we consider two innovation processes: one related to the authors who receive at least one comment with negative sentiment for the news they post and the similar one referred to the comments with positive sentiment. A first analysis of this data set can be found in [5], where we illustrate that these two sequences exhibit empirical behaviours that are in accordance with those predicted by Theorems 1.1 and 1.2, and we provide the estimates $\gamma^* = 0.781$ and $r = 10^{-0.727} = 0.187$. We here continue the study performing the two hypothesis tests on Γ and W described in Subsection 3.1. More precisely, we consider the two one-sided tests:

$$H_{\Gamma,0} : \iota \geq \iota_{\Gamma,0}, \eta \leq 1/2 \quad \text{and} \quad H_{W,0} : \iota \geq \iota_{W,0},$$

where $\iota_{\Gamma,0}$ must be selected in the interval $(0.141, 0.291]$ and $\iota_{W,0}$ must be selected in the interval $(\frac{1}{2}, 2)$. The p -values associated to the null hypothesis $H_{\Gamma,0}$ and $H_{W,0}$, for different values of $\iota_{\Gamma,0}$ and $\iota_{W,0}$, respectively, are collected in Table 1. In order to better describe the empirical distribution of the p -values for testing W , instead of taking a single item c , we take the 100 items c with highest $(K_{t,1}(c) + K_{t,2}(c))$, and report in Table 1 the summary statistics of this sample (i.e. the quantiles of order 50%, 75% and 95%). We can see that, considering a test on Γ performed at level $\alpha = 0.05$, we can reject the null hypothesis with $\iota_{\Gamma,0} = 0.21$, but we cannot reject it for $\iota_{\Gamma,0} = 0.18$. Regarding the test on W instead, we observe that the proportion of items c providing a p -value that let us reject the null hypothesis that the interaction intensity ι_W is higher than $\iota_{W,0}$ is from 75% to 95% for $\iota_{W,0} \leq 1.25$, while is even higher than 95% or $\iota_{W,0} = 1.50$.

$\iota_{\Gamma,0}$	p-value	$\iota_{W,0}$	p-value:	$q_{50\%}, q_{75\%}, q_{95\%}$
0.15	0.384	0.60		<0.001, 0.007, 0.486
0.18	0.064	0.80		<0.001, <0.001, 0.229
0.21	0.014	1.00		<0.001, <0.001, 0.121
0.24	0.003	1.25		<0.001, <0.001, 0.058
0.27	0.001	1.50		<0.001, <0.001, 0.029

TABLE 1. p -values associated to the hypothesis tests described in Subsection 3.1 with $H_{\Gamma,0} : \iota_{\Gamma} \geq \iota_{\Gamma,0}, \eta \leq 1/2$ and $H_{W,0} : \iota_W \geq \iota_{W,0}$, respectively. The table also contains the summary statistics related to the sample of the 100 items c with highest $(K_{t,1}(c) + K_{t,2}(c))$.

4.2. Gutenberg data set. This data set comes from the on-line library Project Gutenberg and it is related to seven books written in English and belonging to two particular literary genres: "Western" and "History". Precisely, it consists of 476 948 words for both genres. The two sequences of words are the innovation processes object of the study. Therefore, each word is new or old for a specific literary genre if, respectively, it has or has not already been used within that genre. Analogously, each word is new or old for the entire system if, respectively, it has or has not already been used within any considered genre. A first analysis of also this data set can be found in [5], where we show that the behaviors of these two sequences predicted by Theorems 1.1 and 1.2 match with the observed ones, and we provide the estimates $\gamma^* = 0.466$ and $r = 10^{-0.238} = 0.578$. We here continue the study performing the two hypothesis tests on Γ and W described in Subsection 3.1. More precisely, we consider the two one-sided tests:

$$H_{\Gamma,0} : \iota \geq \iota_0, \eta \leq 1/2 \quad \text{and} \quad H_{W,0} : \iota \geq \iota_0,$$

where $\iota_{\Gamma,0}$ must be selected in the interval $(0.202, 0.539]$ and $\iota_{W,0}$ must be selected in the interval $(\frac{1}{2}, 2)$. The p -values associated to the null hypothesis $H_{\Gamma,0}$ and $H_{W,0}$, for different values of $\iota_{\Gamma,0}$ and $\iota_{W,0}$, respectively, are collected in Table 2. Moreover, we take the 100 items c with highest $(K_{t,1}(c) + K_{t,2}(c))$ and report in the table the empirical quantiles of the corresponding sample of p -values. We can observe that, considering a test on Γ performed at level $\alpha = 0.05$, we can never reject the null hypothesis. Regarding the test on W instead, we observe that the proportion of items c providing a p -value that let us reject the null hypothesis that the interaction intensity ι_W is higher than $\iota_{W,0}$ is from 50% to 75% for $\iota_{W,0} = 0.60$, while is from 75% to 95% for $\iota_{W,0} \geq 0.80$.

$\iota_{\Gamma,0}$	p-value	$\iota_{W,0}$	p-value:	$q_{50\%}, q_{75\%}, q_{95\%}$
0.21	0.034	0.60		0.010, 0.209, 0.823
0.24	<0.001	0.80		<0.001, 0.030, 0.699
0.27	<0.001	1.00		<0.001, 0.005, 0.618
0.30	<0.001	1.25		<0.001, 0.001, 0.541
0.40	<0.001	1.50		<0.001, <0.001, 0.480

TABLE 2. p -values associated to the hypothesis tests described in Subsection 3.1 with $H_{\Gamma,0} : \iota_{\Gamma} \geq \iota_{\Gamma,0}$, $\eta \leq 1/2$ and $H_{W,0} : \iota_W \geq \iota_{W,0}$, respectively. The table also contains the summary statistics related to the sample of the 100 items c with highest $(K_{t,1}(c) + K_{t,2}(c))$.

APPENDIX A. GENERAL RESULTS

Consider two multi-dimensional real stochastic processes $(\mathbf{A}_t, \mathbf{B}_t)_t$, adapted to a filtration $(\mathcal{F}_t)_t$, with the following joint dynamics:

$$(28) \quad \begin{aligned} \mathbf{A}_{t+1} &= \mathbf{A}_t - \frac{1}{t+1}(\phi^* I - \Phi^\top) \mathbf{A}_t + \frac{1}{t+1} \Phi^\top \Delta \mathbf{M}_{t+1} + \mathbf{R}_{A,t+1} \\ \mathbf{B}_{t+1} &= \mathbf{B}_t - \frac{1}{t+1}(\phi^* \mathbf{B}_t - \mathbf{A}_t) + \frac{1}{t+1} \Delta \mathbf{M}_{t+1} + \mathbf{R}_{B,t+1}, \end{aligned}$$

where \mathbf{A}_0 and \mathbf{B}_0 are integrable and

- (i) Φ^\top is a non-negative irreducible matrix with leading eigenvalue $0 < \phi^* \leq 1$;
- (ii) $\mathbf{R}_{A,t+1} = O(t^{-(1+\beta)})\mathbf{1}$ and $\mathbf{R}_{B,t+1} = O(t^{-(1+\beta)})\mathbf{1}$ for some $\beta > \frac{\phi^*}{2}$;
- (iii) $\Delta \mathbf{M}_{t+1} = O(t^{1-\phi^*})\mathbf{1}$ and $t^{-(1-\phi^*)} E[\Delta \mathbf{M}_{t+1} \Delta \mathbf{M}_{t+1}^\top | \mathcal{F}_t] \xrightarrow{a.s.} \Sigma_\infty$ with Σ_∞ a positive semidefinite symmetric random matrix.

Let \mathbf{u} and \mathbf{v} be the right and the left eigenvectors of Φ^\top associated to ϕ^* and such that $\mathbf{v}^\top \mathbf{1} = 1$ and $\mathbf{v}^\top \mathbf{u} = 1$. (Recall that the components of these vectors are all strictly positive according to the Frobenius-Perron theory)

Let $\tilde{A}_t = \mathbf{v}^\top \mathbf{A}_t$ and get its dynamics by multiplying \mathbf{v}^\top to the dynamics of \mathbf{A}_t as follows:

$$\tilde{A}_{t+1} = \tilde{A}_t + \frac{1}{t+1} \phi^* \mathbf{v}^\top \Delta \mathbf{M}_{t+1} + \mathbf{v}^\top \mathbf{R}_{A,t+1}.$$

Define

$$(29) \quad \hat{\mathbf{A}}_t = \mathbf{A}_t - \tilde{A}_t \mathbf{u} = (I - \mathbf{u} \mathbf{v}^\top) \mathbf{A}_t \quad \text{and} \quad \hat{\mathbf{B}}_t = \mathbf{B}_t - \frac{1}{\phi^*} \tilde{A}_t \mathbf{u}.$$

Theorem A.1. (CLT for (\tilde{A}_t) when $N \geq 1$)

In addition to assumptions (i), (ii) and (iii), also assume:

- (iv) $\tilde{A}_t \xrightarrow{a.s./in\ mean} \tilde{A}_\infty \in (0, +\infty)$, where \tilde{A}_∞ is a strictly positive random variable.

Then, we have

$$t^{\phi^*/2} (\tilde{A}_t - \tilde{A}_\infty) \xrightarrow{stably} \mathcal{N}(0, \phi^* \mathbf{v}^\top \Sigma_\infty \mathbf{v}).$$

Moreover, the above convergence is in the sense of the almost sure conditional convergence with respect to the filtration (\mathcal{F}_t) .

Proof. Set $\check{M}_t = \sum_{n=1}^t \frac{1}{n} \mathbf{v}^\top \Delta \mathbf{M}_n$ and $\check{R}_{A,t+1} = \mathbf{v}^\top \mathbf{R}_{A,t+1}$. Therefore, we have

$$\begin{aligned} \check{A}_{t+1} - \check{A}_0 &= \sum_{n=0}^t \check{A}_{n+1} - \check{A}_n \\ &= \phi^* \sum_{n=0}^t \frac{1}{n+1} \mathbf{v}^\top \Delta \mathbf{M}_{n+1} + \sum_{n=0}^t \check{R}_{A,n+1} \\ &= \phi^* \check{M}_{t+1} + \sum_{n=0}^t \check{R}_{A,n+1}. \end{aligned}$$

Since $\check{A}_t \xrightarrow{\text{a.s./in mean}} \check{A}_\infty$ and $\sum_{n \geq 0} \check{R}_{A,n+1} = \sum_{n \geq 0} O(1/n^{1+\beta}) < +\infty$ then $\check{M}_t \xrightarrow{\text{a.s./in mean}} \check{M}_\infty$ for a certain integrable random variable \check{M}_∞ . Moreover, we have

$$\begin{aligned} t^{\phi^*/2} (\check{A}_t - \check{A}_\infty) &= t^{\phi^*/2} \sum_{n \geq t} (\check{A}_n - \check{A}_{n+1}) \\ &= t^{\phi^*/2} \phi^* (\check{M}_t - \check{M}_\infty) + t^{\phi^*/2} \sum_{n \geq t} \check{R}_{A,n+1} \\ &= t^{\phi^*/2} \phi^* (\check{M}_t - \check{M}_\infty) + t^{\phi^*/2} \sum_{n \geq t} O(1/n^{1+\beta}) \\ &= t^{\phi^*/2} \phi^* (\check{M}_t - \check{M}_\infty) + t^{\phi^*/2} O(1/t^\beta), \end{aligned}$$

where $\beta - \phi^*/2 > 0$. Summing up it is enough to prove the convergence of $t^{\phi^*/2}(\check{M}_t - \check{M}_\infty)$. For this purpose, we want to apply Theorem D.1. Let us consider, for each t the filtration $(\mathcal{F}_{t,k})_k$ and the process $(L_{t,k})_k$ defined by

$$\mathcal{F}_{t,0} = \mathcal{F}_{t,1} = \mathcal{F}_t, \quad L_{t,0} = L_{t,1} = 0$$

and, for $k \geq 2$,

$$\mathcal{F}_{t,k} = \mathcal{F}_{t+k-1}, \quad L_{t,k} = t^{\phi^*/2}(\check{M}_t - \check{M}_{t+k-1}).$$

The process $(L_{t,k})_k$ is a martingale w.r.t. $(\mathcal{F}_{t,k})_k$ which converges (for $k \rightarrow +\infty$) a.s. and in mean to the random variable $L_{t,\infty} = t^{\phi^*/2}(\check{M}_t - \check{M}_\infty)$. In addition, the increment $Y_{t,n} = L_{t,n} - L_{t,n-1}$ is equal to zero for $n = 1$ and, for $n \geq 2$, it coincides with a random variable of the form $t^{\phi^*/2}(\check{M}_k - \check{M}_{k+1})$ with $k \geq t$. Therefore, we have

$$\begin{aligned} \sum_{n \geq 1} Y_{t,n}^2 &= t^{\phi^*} \sum_{k \geq t} (\check{M}_k - \check{M}_{k+1})^2 \\ &= t^{\phi^*} \sum_{k \geq t} \frac{1}{(k+1)^2} (\mathbf{v}^\top \Delta \mathbf{M}_{k+1})^2 \\ &= t^{\phi^*} \sum_{k \geq t} \frac{k^{1-\phi^*}}{(k+1)^2} k^{-(1-\phi^*)} (\mathbf{v}^\top \Delta \mathbf{M}_{k+1})^2 \xrightarrow{\text{a.s.}} \mathbf{v}^\top \Sigma_\infty \mathbf{v} / \phi^*, \end{aligned}$$

where the almost sure convergence follows by applying [11, Lemma 4.1] and by the fact that

$$k^{-(1-\phi^*)} E[(\mathbf{v}^\top \Delta \mathbf{M}_{k+1})^2 | \mathcal{F}_t] = k^{-(1-\phi^*)} \mathbf{v}^\top E[\Delta \mathbf{M}_{k+1} \Delta \mathbf{M}_{k+1}^\top | \mathcal{F}_t] \mathbf{v} \xrightarrow{\text{a.s.}} \mathbf{v}^\top \Sigma_\infty \mathbf{v}.$$

Finally, we have

$$\sup_{n \geq 1} |Y_{t,n}| = t^{\phi^*/2} \sup_{k \geq t} |\check{M}_k - \check{M}_{k+1}| \leq \sup_{k \geq t} k^{\phi^*/2} \frac{|\mathbf{v}^\top \Delta \mathbf{M}_{k+1}|}{(k+1)} \rightarrow 0,$$

because $\Delta \mathbf{M}_{k+1} = O((k+1)^{1-\phi^*})$ by assumption (iii) and so $k^{\phi^*/2} \frac{|\mathbf{v}^\top \Delta \mathbf{M}_{k+1}|}{(k+1)} = O(k^{-\phi^*/2})$. Hence, the thesis follows by Theorem D.1. \square

Theorem A.2. (Convergence result for $(\widehat{A}_t, \widehat{B}_t)$ when $N = 1$)
Under the assumptions (i) and (ii), we have

$$\widehat{A}_t \equiv 0 \quad \text{and} \quad \widehat{B}_t = B_t - \frac{1}{\phi^*} A_t \stackrel{\text{a.s.}}{=} o(t^{-\frac{\phi^*}{2}}).$$

Proof. First, when $N = 1$ it is immediate to see $\widetilde{A}_t \equiv A_t$ and hence $\widehat{A}_t = A_t - \widetilde{A}_t \equiv 0$ for any $t \geq 1$. The dynamics of \widehat{B}_t can be obtained by subtracting the dynamics of $\frac{1}{\phi^*} \widetilde{A}_t = \frac{1}{\phi^*} A_t$ to the dynamics of B_t as follows:

$$\widehat{B}_{t+1} = \widehat{B}_t - \frac{1}{t+1} (\phi^* \widehat{B}_t + \phi^* \frac{1}{\phi^*} \widetilde{A}_t - A_t) + \frac{1}{t+1} \Delta M_{t+1} - \frac{1}{\phi^*} (\widetilde{A}_{t+1} - \widetilde{A}_t) + R_{B,t+1},$$

from which we can derive

$$\widehat{B}_{t+1} = \widehat{B}_t - \frac{1}{t+1} (\phi^* \widehat{B}_t) + \frac{1}{t+1} \Delta M_{t+1} - \frac{1}{\phi^*} \left(\frac{1}{t+1} \phi^* \Delta M_{t+1} \right) + R_{B,t+1} - \frac{1}{\phi^*} R_{A,t+1},$$

that is

$$\widehat{B}_{t+1} = \left(1 - \frac{\phi^*}{t+1} \right) \widehat{B}_t + O(t^{-(1+\beta)}).$$

Then, since $\sum_t \frac{\phi^*}{t+1} = +\infty$ and $\sum_t O(t^{-(1+\beta)}) < +\infty$, we can apply Lemma D.2 to obtain $\widehat{B}_t \stackrel{\text{a.s.}}{\rightarrow} 0$. Moreover, if we multiply the previous dynamics of \widehat{B}_t by $(t+1)^{\frac{\phi^*}{2}}$ and we use the relation $(\frac{t+1}{t})^{\frac{\phi^*}{2}} = (1 + \frac{1}{t})^{\frac{\phi^*}{2}} = 1 + \frac{\phi^*/2}{t+1} + O(1/t^2)$, we get the following dynamics for the process $\widehat{B}_t^* = t^{\frac{\phi^*}{2}} \widehat{B}_t$:

$$\widehat{B}_{t+1}^* = \left(1 - \frac{\phi^*}{t+1} \right) \left(1 + \frac{\phi^*/2}{t+1} \right) \widehat{B}_t^* + O(t^{-(1+\beta)}),$$

which is equivalent to

$$\widehat{B}_{t+1}^* = \left(1 - \frac{\phi^*/2}{t+1} \right) \widehat{B}_t^* + O(t^{-(1+\beta-\frac{\phi^*}{2})}).$$

Therefore, since $\sum_t O(t^{-(1+\beta-\frac{\phi^*}{2})}) < +\infty$ because $\beta > \frac{\phi^*}{2}$ by assumption, we can apply Lemma D.2 also to the sequence \widehat{B}_t^* to obtain $\widehat{B}_t^* \stackrel{\text{a.s.}}{\rightarrow} 0$, i.e. $\widehat{B}_t \stackrel{\text{a.s.}}{=} o(t^{-\frac{\phi^*}{2}})$. \square

Theorem A.3. (CLT for $(\widehat{\mathbf{A}}_t, \widehat{\mathbf{B}}_t)$ when $N \geq 2$)

In addition to assumptions (i), (ii) and (iii), let ϕ_2^* be the second eigenvalue of Φ with highest real part, i.e. $\phi_2^* \in Sp(\Phi) \setminus \{\phi^*\}$ such that $\text{Re}(\phi_2^*) = \max\{\text{Re}(\phi) : \phi \in Sp(\Phi) \setminus \{\phi^*\}\}$ and assume

$$(v) \quad \text{Re}(\phi_2^*)/\phi^* < 1/2.$$

Then, we have

$$t^{\phi^*/2} (\widehat{\mathbf{A}}_t, \widehat{\mathbf{B}}_t) \xrightarrow{\text{stably}} \mathcal{N}(\mathbf{0}, \mathcal{M}_\infty)$$

with

$$(30) \quad \mathcal{M}_\infty = \begin{pmatrix} \mathcal{M}_\infty^{11} & \mathcal{M}_\infty^{13} \\ \mathcal{M}_\infty^{13} & \mathcal{M}_\infty^{33} \end{pmatrix} = \begin{pmatrix} US_\infty^{11}U^\top & US_\infty^{13}U^\top \\ US_\infty^{13}U^\top & US_\infty^{33}U^\top \end{pmatrix},$$

where, denoting by ϕ_j the eigenvalues of Φ different from ϕ^* , we have for $j = 2, \dots, N$

$$\begin{aligned} [\mathcal{S}_\infty^{11}]_{h,j} &= \frac{\phi_j \phi_h}{\phi^* - \phi_j - \phi_h} (\mathbf{v}_j^\top \Sigma_\infty \mathbf{v}_h), \\ [\mathcal{S}_\infty^{33}]_{h,j} &= \frac{1}{\phi^* - \phi_j - \phi_h} (\mathbf{v}_j^\top \Sigma_\infty \mathbf{v}_h), \\ [\mathcal{S}_\infty^{13}]_{h,j} &= \frac{\phi_j}{\phi^* - \phi_j - \phi_h} (\mathbf{v}_j^\top \Sigma_\infty \mathbf{v}_h). \end{aligned}$$

Proof. We split the proofs in some steps.

Step 1 (Dynamics of $\theta_t = (\hat{A}_t, \hat{B}_t)$).

Firstly, we recall that the matrix Φ^\top can be decomposed as

$$\Phi^\top = \phi^* \mathbf{u} \mathbf{v}^\top + U D V^\top,$$

where D is the diagonal matrix whose elements are the eigenvalues of Φ (i.e. of Φ^\top) different from ϕ^* and U and V denote the matrices whose columns are the left (right) and the right (left) eigenvectors of Φ (of Φ^\top , respectively) associated to these eigenvalues, so that we have

$$(31) \quad V^\top \mathbf{u} = U^\top \mathbf{v} = 0, \quad V^\top U = U^\top V = I, \quad I = \mathbf{u} \mathbf{v}^\top + U V^\top.$$

Then, we have $\hat{\mathbf{A}}_t = U V^\top \mathbf{A}_t$ and the dynamics of $\hat{\mathbf{A}}_t$ can be obtained by multiplying by $U V^\top$ the dynamics of \mathbf{A}_t as follows:

$$\hat{\mathbf{A}}_{t+1} = \hat{\mathbf{A}}_t - \frac{1}{t+1} U (I \phi^* - D) V^\top \hat{\mathbf{A}}_t + \frac{1}{t+1} U D V^\top \Delta \mathbf{M}_{t+1} + U V^\top \mathbf{R}_{A,t+1}.$$

The dynamics of $\hat{\mathbf{B}}_t$ can be obtained by subtracting the dynamics of $\frac{1}{\phi^*} \tilde{A}_t \mathbf{u}$ to the dynamics of \mathbf{B}_t as follows:

$$\hat{\mathbf{B}}_{t+1} = \hat{\mathbf{B}}_t - \frac{1}{t+1} (\phi^* \hat{\mathbf{B}}_t + \phi^* \frac{1}{\phi^*} \tilde{A}_t \mathbf{u} - \hat{\mathbf{A}}_t - \tilde{A}_t \mathbf{u}) + \frac{1}{t+1} \Delta \mathbf{M}_{t+1} - \frac{1}{\phi^*} \mathbf{u} (\tilde{A}_{t+1} - \tilde{A}_t) + \mathbf{R}_{B,t+1},$$

from which we can derive

$$\hat{\mathbf{B}}_{t+1} = \hat{\mathbf{B}}_t - \frac{1}{t+1} (\phi^* \hat{\mathbf{B}}_t - \hat{\mathbf{A}}_t) + \frac{1}{t+1} (I - \mathbf{u} \mathbf{v}^\top) \Delta \mathbf{M}_{t+1} + \mathbf{R}_{B,t+1} - \frac{1}{\phi^*} \mathbf{u} \mathbf{v}^\top \mathbf{R}_{A,t+1}.$$

Then the dynamics of $\boldsymbol{\eta}_t = (\hat{\mathbf{A}}_t, \hat{\mathbf{B}}_t)$ can be expressed as

$$\boldsymbol{\eta}_{t+1} - \boldsymbol{\eta}_t = \frac{1}{t+1} Q \boldsymbol{\eta}_t + \frac{1}{t+1} R \Delta \mathbf{M}_{\boldsymbol{\eta},t+1} + \mathbf{R}_{\boldsymbol{\eta},t+1},$$

where³

$$Q = \begin{pmatrix} U(I\phi^* - D)V^\top & \mathcal{O} \\ -UV^\top & I\phi^* \end{pmatrix}, \quad \Delta \mathbf{M}_{\boldsymbol{\eta},t+1} = \begin{pmatrix} \Delta \mathbf{M}_{t+1} \\ \Delta \mathbf{M}_{t+1} \end{pmatrix}, \quad R = \begin{pmatrix} U D V^\top & \mathcal{O} \\ \mathcal{O} & U V^\top \end{pmatrix}.$$

Now, let us define the $(2N) \times (2N - 1)$ matrices

$$U_\boldsymbol{\eta} = \begin{pmatrix} U & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{u} & U \end{pmatrix}, \quad \text{and} \quad V_\boldsymbol{\eta}^\top = \begin{pmatrix} V^\top & \mathbf{0} \\ \mathbf{0}^\top & \mathbf{v}^\top \\ \mathbf{0} & V^\top \end{pmatrix},$$

so that $V_\boldsymbol{\eta}^\top U_\boldsymbol{\eta} = I$ and

$$U_\boldsymbol{\eta} V_\boldsymbol{\eta}^\top = \begin{pmatrix} U V^\top & \mathbf{0} \\ \mathbf{0}^\top & I \end{pmatrix}.$$

³The symbol \mathcal{O} denotes the null matrix.

Then, defining the $(2N - 1) \times (2N - 1)$ matrices

$$S_Q = \begin{pmatrix} I\phi^* - D & \mathbf{0} & 0 \\ \mathbf{0}^\top & \phi^* & \mathbf{0}^\top \\ -I & \mathbf{0} & I\phi^* \end{pmatrix}, \quad \text{and} \quad S_R = \begin{pmatrix} D & \mathbf{0} & 0 \\ \mathbf{0}^\top & 0 & \mathbf{0}^\top \\ 0 & \mathbf{0} & I \end{pmatrix},$$

we have that $Q = U_\eta S_Q V_\eta^\top$ and $R = U_\eta S_R V_\eta^\top$. Then, from the above relations on U_η and V_η^\top we get $U_\eta V_\eta^\top \boldsymbol{\eta}_t = \boldsymbol{\eta}_t$, and so multiplying the dynamics of $\boldsymbol{\eta}_t$ by $U_\eta V_\eta^\top$ we finally get

$$\boldsymbol{\eta}_{t+1} = \frac{1}{t+1} U_\eta \left(I - \frac{1}{t+1} S_Q \right) V_\eta^\top \boldsymbol{\eta}_t + \frac{1}{t+1} R \Delta \mathbf{M}_{\boldsymbol{\eta}, t+1} + \mathbf{R}_{\boldsymbol{\eta}, t+1}.$$

Step 2 (Explicit expression for $\boldsymbol{\eta}_t = (\widehat{\mathbf{A}}_t, \widehat{\mathbf{B}}_t)$).

Denoting the eigenvalues of Φ^\top different from ϕ^* by ϕ_j for $j = 2, \dots, N$, let us set $\alpha_j = \phi^* - \phi_j$. Recall that $\mathcal{R}e(\alpha_j) > 0$ since ϕ^* is the leading eigenvalue and so $\mathcal{R}e(\phi_j) < \phi^*$. Then, we can write

$$\begin{aligned} \boldsymbol{\eta}_{t+1} &= C_{m_0, t} \boldsymbol{\eta}_{m_0} + \sum_{k=m_0}^t \frac{1}{k+1} C_{k+1, t} R \Delta \mathbf{M}_{\boldsymbol{\eta}, k+1} + \sum_{k=m_0}^t C_{k+1, t} \mathbf{R}_{\boldsymbol{\eta}, k+1} \\ &= C_{m_0, t} \boldsymbol{\eta}_{m_0} + \sum_{k=m_0}^t \mathbf{T}_{t, k+1} + \boldsymbol{\rho}_{t+1}, \end{aligned}$$

where

$$\begin{aligned} C_{k+1, t} &= U_\eta A_{k+1, t} V_\eta^\top, \quad \text{and} \\ A_{k+1, t} &= \begin{cases} \prod_{m=k+1}^t \left(I - \frac{1}{m+1} S_Q \right) = \begin{pmatrix} A_{k+1, t}^{11} & \mathbf{0} & 0 \\ 0 & a_{k+1, t}^{22} & 0 \\ A_{k+1, t}^{31} & \mathbf{0} & A_{k+1, t}^{33} \end{pmatrix} & k = m_0, \dots, t-1 \\ I & k = t. \end{cases} \end{aligned}$$

Notice that the blocks $A_{k+1, t}^{11}$, $A_{k+1, t}^{31}$ and $A_{k+1, t}^{33}$ are all diagonal $(N-1) \times (N-1)$ matrices. Moreover, setting for any $x \in \mathbb{C}$, $p_{m_0-1}(x) = 1$ and $p_k(x) = \prod_{m=m_0}^k (1 - \frac{x}{m+1})$ for $k \geq m_0$ and $F_{k+1, t} = \frac{p_t(x)}{p_k(x)}$ for $m_0 - 1 \leq k \leq t-1$, from [3, Lemma A.5] we get

$$\begin{aligned} [A_{k+1, t}^{11}]_{jj} &= F_{k+1, t}(\alpha_j) \\ [A_{k+1, t}^{33}]_{jj} &= a_{k+1, t}^{22} = F_{k+1, t}(\phi^*) \\ [A_{k+1, t}^{31}]_{jj} &= \begin{cases} -\frac{F_{k+1, t}(\alpha_j) - F_{k+1, t}(\phi^*)}{\alpha_j - \phi^*} & \text{for } \alpha_j \neq \phi^*, \text{ i.e. } \phi_j \neq 0 \\ \frac{1 - \phi^*}{\phi^*} F_{k+1, t}(\phi^*) \ln\left(\frac{t}{k}\right) + O\left(\frac{1}{t}\right) & \text{for } \alpha_j = \phi^*, \text{ i.e. } \phi_j = 0. \end{cases} \end{aligned}$$

Step 3 (Study of the terms $C_{m_0, t} \boldsymbol{\eta}_{m_0}$ and $\boldsymbol{\rho}_{t+1}$).

We will prove that $t^{\frac{\phi^*}{2}} |C_{m_0, t} \boldsymbol{\eta}_{m_0}| \xrightarrow{a.s.} 0$ and $t^{\frac{\phi^*}{2}} |\boldsymbol{\rho}_{t+1}| \xrightarrow{a.s.} 0$. To this end, first notice that $O(|C_{k+1, t}|) =$

$O(|A_{k+1,t}|)$ and, setting $a_2^* = \phi^* - \mathcal{R}e(\phi_2^*)$, we have (see [2, Lemma A.4])

$$\begin{aligned} |A_{k+1,t}| &= O\left(\frac{|p_t(\alpha_2^*)|}{|p_k(\alpha_2^*)|}\right) + O\left(\frac{|p_t(\phi^*)|}{|p_k(\phi^*)|}\right) + O\left(\frac{|p_t(\phi^*)|}{|p_k(\phi^*)|} \ln\left(\frac{t}{k}\right)\right) + O\left(\frac{1}{t}\right) \\ &= O\left(\left(\frac{k}{t}\right)^{a_2^*}\right) + O\left(\left(\frac{k}{t}\right)^{\phi^*}\right) + O\left(\left(\frac{k}{t}\right)^{\phi^*} \ln\left(\frac{t}{k}\right)\right) + O\left(\frac{1}{t}\right) \\ &= O\left(\left(\frac{k}{t}\right)^{a_2^*}\right) + O\left(\left(\frac{k}{t}\right)^{\phi^*} \ln(t)\right) \\ &= O\left(\left(\frac{k}{t}\right)^{(\phi^* - \mathcal{R}e(\phi_2^*))}\right) + O\left(\left(\frac{k}{t}\right)^{\phi^*} \ln(t)\right) \quad \text{for } k = m_0, \dots, t-1 \end{aligned}$$

and simply $|A_{t+1,t}| = O(1)$ for $k = t + 1$. Therefore, since assumption (v), we have

$$t^{\frac{\phi^*}{2}} |C_{m_0,t} \boldsymbol{\eta}_{m_0}| = \left(\frac{1}{t}\right)^{\phi^*/2 - \mathcal{R}e(\phi_2^*)} + \left(\frac{1}{t}\right)^{\phi^*/2} \ln(t) \xrightarrow{a.s.} 0.$$

Moreover, recalling that $\mathbf{R}_{\boldsymbol{\eta},t+1} = O(t^{-(1+\beta)})\mathbf{1}$ for some $\beta > \frac{\phi^*}{2}$, we have

$$\begin{aligned} t^{\frac{\phi^*}{2}} |\boldsymbol{\rho}_{t+1}| &= t^{\frac{\phi^*}{2}} \left| \sum_{k=m_0}^t C_{k+1,t} \mathbf{R}_{\boldsymbol{\eta},k+1} \right| \\ &= t^{\frac{\phi^*}{2}} O\left(\sum_{k=m_0}^{t-1} \left(\frac{k}{t}\right)^{a_2^*} \frac{1}{k^{1+\beta}} + \sum_{k=m_0}^{t-1} \left(\frac{k}{t}\right)^{\phi^*} \ln(t) \frac{1}{k^{1+\beta}}\right) + t^{\phi^*/2} O(1/t^{1+\beta}) \\ &= t^{\frac{\phi^*}{2}} O\left(\frac{1}{t^{a_2^*}} \sum_{k=m_0}^{t-1} k^{a_2^* - 1 - \beta} + \frac{\ln(t)}{t^{\phi^*}} \sum_{k=m_0}^{t-1} k^{\phi^* - 1 - \beta}\right) + O(1/t^{1+\beta - \phi^*/2}) \\ &= t^{\frac{\phi^*}{2}} O\left(\frac{1}{t^\beta} + \frac{\ln(t)}{t^\beta}\right) + O(1/t^{1+\beta - \phi^*/2}) \\ &= O\left(\frac{\ln(t)}{t^{\beta - \frac{\phi^*}{2}}}\right), \end{aligned}$$

which tends to zero because $\beta > \frac{\phi^*}{2}$.

Step 4 (Study of the “delta-martingale” term).

We want to apply Theorem D.3. Therefore we have to check conditions (1) and (2) in Theorem D.3.

Since the relation $V_{\boldsymbol{\eta}}^\top U_{\boldsymbol{\eta}} = I$ implies $V_{\boldsymbol{\eta}}^\top R = S_R V_{\boldsymbol{\eta}}^\top$, we have that

$$\begin{aligned} \sum_{k=m_0}^t (t^{\frac{\phi^*}{2}} \mathbf{T}_{t,k+1}) (t^{\frac{\phi^*}{2}} \mathbf{T}_{t,k+1}^\top) &= t^{\phi^*} \sum_{k=m_0}^t \frac{1}{(k+1)^2} C_{k+1,t} R \Delta \mathbf{M}_{\boldsymbol{\eta},k+1} \Delta \mathbf{M}_{\boldsymbol{\eta},k+1}^\top R^\top C_{k+1,t}^\top \\ &= U_{\boldsymbol{\eta}} \left(t^{\phi^*} \sum_{k=m_0}^t \frac{1}{(k+1)^2} A_{k+1,t} V_{\boldsymbol{\eta}}^\top R \Delta \mathbf{M}_{\boldsymbol{\eta},k+1} \Delta \mathbf{M}_{\boldsymbol{\eta},k+1}^\top R^\top V_{\boldsymbol{\eta}} A_{k+1,t}^\top \right) U_{\boldsymbol{\eta}}^\top \\ &= U_{\boldsymbol{\eta}} \left(t^{\phi^*} \sum_{k=m_0}^t \frac{1}{(k+1)^2} A_{k+1,t} S_R V_{\boldsymbol{\eta}}^\top \Delta \mathbf{M}_{\boldsymbol{\eta},k+1} \Delta \mathbf{M}_{\boldsymbol{\eta},k+1}^\top V_{\boldsymbol{\eta}} S_R A_{k+1,t}^\top \right) U_{\boldsymbol{\eta}}^\top. \end{aligned}$$

Therefore, omitting the last term of the sum because $|A_{t+1,t}| = O(1)$ and, by assumption (iii), we have $(t^{\phi^*}/(t+1)^2)|\Delta \mathbf{M}_{t+1}| = O(t^{\phi^*-2+1-\phi^*}) = O(t^{-1}) \rightarrow 0$, it is enough to study the convergence of

$$(32) \quad t^{\phi^*} \sum_{k=m_0}^{t-1} \frac{1}{(k+1)^2} A_{k+1,t} S_R V_{\boldsymbol{\eta}}^{\top} \Delta \mathbf{M}_{\boldsymbol{\eta},k+1} \Delta \mathbf{M}_{\boldsymbol{\eta},k+1}^{\top} V_{\boldsymbol{\eta}} S_R A_{k+1,t}^{\top}.$$

To this purpose, setting $B_{\boldsymbol{\eta},k+1} = V_{\boldsymbol{\eta}}^{\top} \Delta \mathbf{M}_{\boldsymbol{\eta},k+1} \Delta \mathbf{M}_{\boldsymbol{\eta},k+1}^{\top} V_{\boldsymbol{\eta}}$, $B_{k+1} = V^{\top} \Delta \mathbf{M}_{k+1} \Delta \mathbf{M}_{k+1}^{\top} V$, $\mathbf{b}_{k+1} = V^{\top} \Delta \mathbf{M}_{k+1} \Delta \mathbf{M}_{k+1}^{\top} \mathbf{v}$ and $b_k = \mathbf{v}^{\top} \Delta \mathbf{M}_{k+1} \Delta \mathbf{M}_{k+1}^{\top} \mathbf{v}$, we observe that

$$B_{\boldsymbol{\eta},k+1} = \begin{pmatrix} B_{k+1} & \mathbf{b}_{k+1} & B_{k+1} \\ \mathbf{b}_{k+1}^{\top} & b_{k+1} & \mathbf{b}_{k+1}^{\top} \\ B_{k+1} & \mathbf{b}_{k+1} & B_{k+1} \end{pmatrix}.$$

Since in $B_{\boldsymbol{\eta},k+1}$ the first and the third row and column of blocks are the same, the matrix $(A_{k+1,t} S_R)$ can be replaced by a diagonal matrix, with the following diagonal blocks: $A_{k+1,t}^1 = A_{k+1,t}^{11} D$, $A_{k+1,t}^3 = A_{k+1,t}^{31} D + A_{k+1,t}^{33}$ and $a_{k+1,t}^2 = 0$. Hence, the above expression (32) can be rewritten as

$$t^{\phi^*} \sum_{k=m_0}^{t-1} \frac{1}{(k+1)^2} \begin{pmatrix} A_{k+1,t}^1 B_{k+1} A_{k+1,t}^1 & \mathbf{0} & A_{k+1,t}^1 B_{k+1} A_{k+1,t}^3 \\ & \mathbf{0}^{\top} & \mathbf{0} \\ A_{k+1,t}^3 B_{k+1} A_{k+1,t}^1 & \mathbf{0} & A_{k+1,t}^3 B_{k+1} A_{k+1,t}^3 \end{pmatrix}.$$

The elements of $A_{k+1,t}^1$ and $A_{k+1,t}^3$ in the above matrix can be rewritten in terms of $F_{k+1,t}(\cdot)$ as

$$\begin{aligned} [A_{k+1,t}^1]_{jj} &= (\phi^* - \alpha_j) F_{k+1,t}(\alpha_j) = \phi_j F_{k+1,t}(\alpha_j) \\ [A_{k+1,t}^3]_{jj} &= \begin{cases} F_{k+1,t}(\alpha_j) & \text{for } \alpha_j \neq \phi^*, \text{ i.e. } \phi_j \neq 0 \\ F_{k+1,t}(\phi^*) & \text{for } \alpha_j = \phi^*, \text{ i.e. } \phi_j = 0. \end{cases} \end{aligned}$$

Hence, the almost sure convergences of all the elements can be obtained by [4, Lemma A.3] together with [11, Lemma 4.1]). Indeed, we have

$$t^{-(1-\phi^*)} E[B_{t+1} | \mathcal{F}_t] = t^{-(1-\phi^*)} V^{\top} E[\Delta \mathbf{M}_{t+1} \Delta \mathbf{M}_{t+1}^{\top} | \mathcal{F}_t] V \rightarrow V^{\top} \Sigma_{\infty} V$$

and so, for complex numbers x, y such that $\Re(x+y) - \phi^* > 0$ (recall that, by assumption (v), we have $\Re(x+y) - \phi^* \geq \phi^* - 2\Re(\phi_2^*) > 0$), by the above quoted lemmas, we have

$$t^{\phi^*} \sum_{k=m_0}^{t-1} \frac{1}{(k+1)^{1+\phi^*}} \left((k+1)^{-(1-\phi^*)} B_{k+1} \right) F_{k+1,t}(x) F_{k+1,t}(y) \rightarrow \frac{1}{x+y-\phi^*} V^{\top} \Sigma_{\infty} V.$$

Hence, recalling that $\alpha_j = \phi^* - \phi_j$, we find

$$\begin{aligned} t^{\phi^*} \sum_{k=m_0}^{t-1} \frac{1}{(k+1)^2} [A_{k+1,t}^1 B_{k+1} A_{k+1,t}^1]_{h,j} &\rightarrow [\mathcal{S}_{\infty}^{11}]_{h,j} = \frac{\phi_j \phi_h}{\phi^* - \phi_j - \phi_h} (\mathbf{v}_j^{\top} \Sigma_{\infty} \mathbf{v}_h), \\ t^{\phi^*} \sum_{k=m_0}^{t-1} \frac{1}{(k+1)^2} [A_{k+1,t}^3 B_{k+1} A_{k+1,t}^3]_{h,j} &\rightarrow [\mathcal{S}_{\infty}^{33}]_{h,j} = \frac{1}{\phi^* - \phi_j - \phi_h} (\mathbf{v}_j^{\top} \Sigma_{\infty} \mathbf{v}_h), \\ t^{\phi^*} \sum_{k=m_0}^{t-1} \frac{1}{(k+1)^2} [A_{k+1,t}^1 B_{k+1} A_{k+1,t}^3]_{h,j} &\rightarrow [\mathcal{S}_{\infty}^{13}]_{h,j} = \frac{\phi_j}{\phi^* - \phi_j - \phi_h} (\mathbf{v}_j^{\top} \Sigma_{\infty} \mathbf{v}_h). \end{aligned}$$

Therefore condition (1) of Theorem D.3 is satisfied with the matrix \mathcal{M}_{∞} defined in (30).

Regarding condition (2) of Theorem D.3, we observe that, by assumption (iii), we have

$$\begin{aligned} |\mathbf{T}_{t,k+1}| &= O\left(\frac{1}{k+1}|A_{k+1,t}|\Delta\mathbf{M}_{\boldsymbol{\eta},k+1}\right) = O\left(|A_{k+1,t}|(k+1)^{-\phi^*}\right) \\ &= O\left(k^{-\phi^*}\left(\frac{k}{t}\right)^{(\phi^*-\mathcal{R}e(\phi_2^*))}\right) + O\left(k^{-\phi^*}\left(\frac{k}{t}\right)^{\phi^*}\ln(t)\right) \quad \text{for } k = m_0, \dots, t-1 \end{aligned}$$

and $|\mathbf{T}_{t,t+1}| = O\left(\frac{1}{t+1}|A_{t+1,t}|\Delta\mathbf{M}_{\boldsymbol{\eta},t+1}\right) = O(t^{-\phi^*})$ for $k = t$. Hence, for any u , we have

$$\begin{aligned} \left(\sup_{m_0 \leq k \leq n} |t^{\phi^*/2}\mathbf{T}_{t,k+1}|\right)^{2u} &\leq t^{\phi^*u} \sum_{k=m_0}^{t-1} |\mathbf{T}_{t,k+1}|^{2u} + t^{\phi^*u} |\mathbf{T}_{t,t+1}|^{2u} \\ &= O\left(t^{-(\phi^*-2\mathcal{R}e(\phi_2^*))u} \sum_{k=m_0}^{t-1} k^{-2\mathcal{R}e(\phi_2^*)u}\right) + O\left(\frac{\ln(t)^{2u}}{t^{\phi^*u-1}}\right) + O(1/t^{\phi^*u}) \\ &= O\left(t^{-(\phi^*-2\mathcal{R}e(\phi_2^*))u} t^{1-2\mathcal{R}e(\phi_2^*)u}\right) + O\left(\frac{\ln(t)^{2u}}{t^{\phi^*u-1}}\right) \\ &= O\left(1/t^{\phi^*u-1}\right) + O\left(\frac{\ln(t)^{2u}}{t^{\phi^*u-1}}\right) = O\left(\frac{\ln(t)^{2u}}{t^{\phi^*u-1}}\right) \rightarrow 0 \quad \text{for } u > 1/\phi^*, \end{aligned}$$

which, in particular, implies $(\sup_{m_0 \leq k \leq t} |t^{\phi^*/2}\mathbf{T}_{t,k+1}|)^{2u} \xrightarrow{L^1} 0$ for any $u > 1/\phi^*$. As a consequence, condition (2) of Theorem D.3 holds true.

By Theorem D.3, we can finally conclude that $t^{\phi^*/2} \sum_{k=m_0}^t \mathbf{T}_{t,k+1}$ converges stably to the Gaussian kernel $\mathcal{N}(\mathbf{0}, \mathcal{M}_\infty)$. \square

Theorem A.4. *When $N = 1$, under assumptions (i), (ii), (iii) and (iv), we have*

$$t^{\phi^*/2} \begin{pmatrix} A_t - \tilde{A}_\infty \\ B_t - \tilde{A}_\infty/\phi^* \end{pmatrix} \xrightarrow{\text{stably}} \mathcal{N}\left(0, \begin{pmatrix} \phi^*\Sigma_\infty & \Sigma_\infty \\ \Sigma_\infty & \Sigma_\infty/\phi^* \end{pmatrix}\right).$$

When $N \geq 2$, under assumptions (i), (ii), (iii), (iv) and (v), we have

$$t^{\phi^*/2} \begin{pmatrix} \mathbf{A}_t - \tilde{A}_\infty \mathbf{u} \\ \mathbf{B}_t - \tilde{A}_\infty/\phi^* \mathbf{u} \end{pmatrix} \xrightarrow{\text{stably}} \mathcal{N}\left(\mathbf{0}, \mathbf{v}^\top \Sigma_\infty \mathbf{v} \begin{pmatrix} \phi^* \mathbf{u} \mathbf{u}^\top & \mathbf{u} \mathbf{u}^\top \\ \mathbf{u} \mathbf{u}^\top & \mathbf{u} \mathbf{u}^\top/\phi^* \end{pmatrix} + \mathcal{M}_\infty\right),$$

where \mathcal{M}_∞ is given in (30).

Proof. When $N = 1$, we simply have $(A_t, B_t) = (\tilde{A}_t, \tilde{A}_t/\phi^* + o(t^{-\phi^*/2}))$ and so the statement follows from the clt for \tilde{A}_t . When $N \geq 2$, we have

$$\begin{aligned} &t^{\phi^*/2}(\mathbf{A}_t - \tilde{A}_\infty \mathbf{u}, \mathbf{B}_t - \tilde{A}_\infty/\phi^* \mathbf{u}) \\ &= t^{\phi^*/2}((\tilde{A}_t - \tilde{A}_\infty)\mathbf{u} + \hat{\mathbf{A}}_t, (\tilde{A}_t - \tilde{A}_\infty)/\phi^* \mathbf{u} + \hat{\mathbf{B}}_t) \\ &= t^{\phi^*/2}(\tilde{A}_t - \tilde{A}_\infty)(\mathbf{u}, \mathbf{u}/\phi^*) + t^{\phi^*/2}(\hat{\mathbf{A}}_t, \hat{\mathbf{B}}_t). \end{aligned}$$

The first term converges in the sense of the almost sure conditional convergence toward the Gaussian kernel with mean $\mathbf{0}$ and covariance matrix

$$\mathbf{v}^\top \Sigma_\infty \mathbf{v} \begin{pmatrix} \phi^* \mathbf{u} \mathbf{u}^\top & \mathbf{u} \mathbf{u}^\top \\ \mathbf{u} \mathbf{u}^\top & \mathbf{u} \mathbf{u}^\top/\phi^* \end{pmatrix}.$$

The second term converges stably toward the Gaussian kernel $\mathcal{N}(\mathbf{0}, \mathcal{M}_\infty)$. By Lemma D.4, we can combine the two convergences and obtain the desired convergence. \square

As an immediate corollary, we can get the central limit theorem for $\tilde{B}_t = \mathbf{v}^\top \mathbf{B}_t$ when $N \geq 2$.

Corollary A.5. (CLT for (\tilde{B}_t) when $N \geq 2$)

When $N \geq 2$, under assumptions (i), (ii), (iii), (iv) and (v), we have

$$t^{\phi^*/2}(\tilde{B}_t - \tilde{A}_\infty/\phi^*) \xrightarrow{\text{stably}} \mathcal{N}\left(0, \mathbf{v}^\top \Sigma_\infty \mathbf{v}/\phi^*\right).$$

Proof. Setting $\tilde{B}_t = \mathbf{v}^\top \mathbf{B}_t$ and using the relations $\mathbf{v}^\top \mathbf{u} = 1$ and $\mathbf{v}^\top U = \mathbf{0}$ in Theorem A.4, we find the desired convergence. \square

We can also derive the following result that consists in a CLT for a normalized version of $(\mathbf{A}_t, \mathbf{B}_t)$.

Corollary A.6. (CLT for $(\mathbf{A}_t/(\mathbf{x}^\top \mathbf{A}_t), \mathbf{B}_t/(\mathbf{x}^\top \mathbf{B}_t))$ when $N \geq 2$)

Under assumptions (i), (ii), (iii), (iv) and (v), for any given vector with $\mathbf{x}^\top \mathbf{u} \neq 0$, we have

$$t^{\phi^*/2} \begin{pmatrix} \mathbf{A}_t/(\mathbf{x}^\top \mathbf{A}_t) - \mathbf{u}/(\mathbf{x}^\top \mathbf{u}) \\ \mathbf{B}_t/(\mathbf{x}^\top \mathbf{B}_t) - \mathbf{u}/(\mathbf{x}^\top \mathbf{u}) \end{pmatrix} \xrightarrow{\text{stably}} \mathcal{N}\left(\mathbf{0}, \frac{1}{(\mathbf{x}^\top \mathbf{u})^2 \tilde{A}_\infty^2} \begin{pmatrix} Q_{\mathbf{x}} \mathcal{M}_\infty^{11} Q_{\mathbf{x}}^\top & \phi^* Q_{\mathbf{x}} \mathcal{M}_\infty^{13} Q_{\mathbf{x}}^\top \\ \phi^* Q_{\mathbf{x}} \mathcal{M}_\infty^{13} Q_{\mathbf{x}}^\top & (\phi^*)^2 Q_{\mathbf{x}} \mathcal{M}_\infty^{33} Q_{\mathbf{x}}^\top \end{pmatrix}\right),$$

where $Q_{\mathbf{x}} = I - \mathbf{u}\mathbf{x}^\top/(\mathbf{x}^\top \mathbf{u})$ and the matrices \mathcal{M}_∞^{11} , \mathcal{M}_∞^{13} , \mathcal{M}_∞^{33} are defined in (30). In particular, for $\mathbf{x} = \mathbf{v}$ we have

$$t^{\phi^*/2} \begin{pmatrix} \mathbf{A}_t/\tilde{A}_t - \mathbf{u} \\ \mathbf{B}_t/\tilde{B}_t - \mathbf{u} \end{pmatrix} \xrightarrow{\text{stably}} \mathcal{N}\left(\mathbf{0}, \frac{1}{\tilde{A}_\infty^2} \begin{pmatrix} \mathcal{M}_\infty^{11} & \phi^* \mathcal{M}_\infty^{13} \\ \phi^* \mathcal{M}_\infty^{13} & (\phi^*)^2 \mathcal{M}_\infty^{33} \end{pmatrix}\right).$$

Remark 1. Note that, by (29) and (31), we have $\hat{\mathbf{A}}_t = \mathbf{A}_t - \tilde{A}_t \mathbf{u} = UV^\top \mathbf{A}_t$ and, similarly, $\mathbf{B}_t - \tilde{B}_t \mathbf{u} = UV^\top \mathbf{B}_t$. Therefore, the processes in the above result in the case $\mathbf{x} = \mathbf{v}$ coincide, up to multiplicative (random) factors, with the processes $UV^\top \mathbf{A}_t$ and $UV^\top \mathbf{B}_t$ (note also that we have $Q_{\mathbf{v}} = I - \mathbf{u}\mathbf{v}^\top/(\mathbf{v}^\top \mathbf{u}) = UV^\top$ by (31)). These processes will play a fundamental role in the following section dedicated to the statistical tools.

Proof. The result follows from the CLT of Theorem A.4 once we note that

$$\begin{aligned} & t^{\phi^*/2} \begin{pmatrix} \mathbf{A}_t/(\mathbf{x}^\top \mathbf{A}_t) - \mathbf{u}/(\mathbf{x}^\top \mathbf{u}) \\ \mathbf{B}_t/(\mathbf{x}^\top \mathbf{B}_t) - \mathbf{u}/(\mathbf{x}^\top \mathbf{u}) \end{pmatrix} \\ &= t^{\phi^*/2} \begin{pmatrix} (\mathbf{x}^\top \mathbf{A}_t)^{-1} I & 0 \\ 0 & (\mathbf{x}^\top \mathbf{B}_t)^{-1} I \end{pmatrix} \begin{pmatrix} \mathbf{A}_t - \mathbf{u}\mathbf{x}^\top \mathbf{A}_t/(\mathbf{x}^\top \mathbf{u}) \\ \mathbf{B}_t - \mathbf{u}\mathbf{x}^\top \mathbf{B}_t/(\mathbf{x}^\top \mathbf{u}) \end{pmatrix} \\ &= t^{\phi^*/2} \begin{pmatrix} (\mathbf{x}^\top \mathbf{A}_t)^{-1} Q_{\mathbf{x}} & 0 \\ 0 & (\mathbf{x}^\top \mathbf{B}_t)^{-1} Q_{\mathbf{x}} \end{pmatrix} \begin{pmatrix} \mathbf{A}_t \\ \mathbf{B}_t \end{pmatrix} \\ &= t^{\phi^*/2} \begin{pmatrix} (\mathbf{x}^\top \mathbf{A}_t)^{-1} Q_{\mathbf{x}} & 0 \\ 0 & (\mathbf{x}^\top \mathbf{B}_t)^{-1} Q_{\mathbf{x}} \end{pmatrix} \begin{pmatrix} \mathbf{A}_t - \tilde{A}_\infty \mathbf{u} \\ \mathbf{B}_t - \tilde{A}_\infty/\phi^* \mathbf{u} \end{pmatrix}. \end{aligned}$$

Indeed, we have $Q_{\mathbf{x}} \mathbf{u}\mathbf{u}^\top = 0$ and so

$$\begin{aligned} & \begin{pmatrix} (\mathbf{x}^\top \mathbf{A}_t)^{-1} Q_{\mathbf{x}} & 0 \\ 0 & (\mathbf{x}^\top \mathbf{B}_t)^{-1} Q_{\mathbf{x}} \end{pmatrix} \left(\mathbf{v}^\top \Sigma_\infty \mathbf{v} \begin{pmatrix} \phi^* \mathbf{u}\mathbf{u}^\top & \mathbf{u}\mathbf{u}^\top \\ \mathbf{u}\mathbf{u}^\top & \mathbf{u}\mathbf{u}^\top/\phi^* \end{pmatrix} + \mathcal{M}_\infty \right) \begin{pmatrix} (\mathbf{x}^\top \mathbf{A}_t)^{-1} Q_{\mathbf{x}} & 0 \\ 0 & (\mathbf{x}^\top \mathbf{B}_t)^{-1} Q_{\mathbf{x}} \end{pmatrix}^\top = \\ & \begin{pmatrix} (\mathbf{x}^\top \mathbf{A}_t)^{-1} Q_{\mathbf{x}} & 0 \\ 0 & (\mathbf{x}^\top \mathbf{B}_t)^{-1} Q_{\mathbf{x}} \end{pmatrix} \mathcal{M}_\infty \begin{pmatrix} (\mathbf{x}^\top \mathbf{A}_t)^{-1} Q_{\mathbf{x}}^\top & 0 \\ 0 & (\mathbf{x}^\top \mathbf{B}_t)^{-1} Q_{\mathbf{x}}^\top \end{pmatrix} = \\ & \frac{1}{(\mathbf{x}^\top \mathbf{u})^2 \tilde{A}_\infty^2} \begin{pmatrix} Q_{\mathbf{x}} \mathcal{M}_\infty^{11} Q_{\mathbf{x}}^\top & \phi^* Q_{\mathbf{x}} \mathcal{M}_\infty^{13} Q_{\mathbf{x}}^\top \\ \phi^* Q_{\mathbf{x}} \mathcal{M}_\infty^{13} Q_{\mathbf{x}}^\top & (\phi^*)^2 Q_{\mathbf{x}} \mathcal{M}_\infty^{33} Q_{\mathbf{x}}^\top \end{pmatrix}. \end{aligned}$$

The result for $\mathbf{x} = \mathbf{v}$ simply follows by the fact that, since (31), we have the equalities $Q_{\mathbf{x}} = Q_{\mathbf{v}} = (I - \mathbf{u}\mathbf{v}^\top/(\mathbf{v}^\top \mathbf{u})) = UV^\top$ and so $UV^\top \mathcal{M}_\infty^{ij} VU^\top = \mathcal{M}_\infty^{ij}$. \square

APPENDIX B. STATISTICAL TOOLS

In this section we present some statistical tools, based on Theorem A.4, to provide an interval estimation of the random limit \tilde{A}_∞ and an hypothesis test on the structure of the matrix Φ . To this end, we need to assume that the asymptotic conditional covariance matrix Σ_∞ of the martingale vector $\Delta \mathbf{M}_t$ satisfies the following assumption:

Assumption 3. *Assume the matrix Σ_∞ can be factorized as*

$$\Sigma_\infty = g(\tilde{A}_\infty) \Sigma_{det},$$

with g being a continuous function and Σ_{det} a deterministic matrix determined by the eigen-structure of Φ .

We will provide statistical tools based on the process (\mathbf{A}_t) or on the process (\mathbf{B}_t) so that the suitable tool can be selected in the practical applications when only one of the two processes is observable.

B.1. Confidence intervals for the random limit \tilde{A}_∞ based on (\mathbf{A}_t) . First of all, we can observe that by means of the central limit theorem for the stochastic process $\tilde{A}_t = \mathbf{v}^\top \mathbf{A}_t$ described in Theorem A.1, it is possible to construct an asymptotic confidence interval for its limit \tilde{A}_∞ . Specifically, an asymptotic confidence interval for \tilde{A}_∞ with approximate level $(1-\alpha)$ is the following:

$$CI_{1-\alpha}(\tilde{A}_\infty) := \left(\tilde{A}_t - \frac{z_\alpha}{t^{\phi^*/2}} \sqrt{g(\tilde{A}_t) \phi^* \mathbf{v}^\top \Sigma_{det} \mathbf{v}}; \tilde{A}_t + \frac{z_\alpha}{t^{\phi^*/2}} \sqrt{g(\tilde{A}_t) \phi^* \mathbf{v}^\top \Sigma_{det} \mathbf{v}} \right),$$

where z_α is such that $\mathcal{N}(0, 1)(z_\alpha, +\infty) = \alpha/2$. Note that the term $g(\tilde{A}_\infty)$ in the confidence interval has been replaced by its strongly consistent estimator $g(\tilde{A}_t)$.

B.2. Confidence intervals for the random limit \tilde{A}_∞ based on (\mathbf{B}_t) . We here construct an asymptotic confidence interval for \tilde{A}_∞ based on the process $(\mathbf{B}_t)_t$. First notice that using Theorem A.4 we can obtain for $N = 1$

$$CI_{1-\alpha}(\tilde{A}_\infty) = \phi^* B_t \pm \frac{z_\alpha}{t^{\phi^*/2}} \sqrt{g(\phi^* B_t) \phi^* \Sigma_{det}},$$

where $g(\phi^* B_t)$ represents a consistent estimator of $g(\tilde{A}_\infty)$ and z_α is such that $\mathcal{N}(0, 1)(z_\alpha, +\infty) = \alpha/2$.

When $N \geq 2$, it is required the knowledge of $\mathcal{R}e(\phi_2^*)$. Indeed, when $\mathcal{R}e(\phi_2^*)/\phi^* < 1/2$, setting $\tilde{B}_t = \mathbf{v}^\top \mathbf{B}_t$ and using Corollary A.5, we obtain

$$CI_{1-\alpha}(\tilde{A}_\infty) = \phi^* \tilde{B}_t \pm \frac{z_\alpha}{t^{\phi^*/2}} \sqrt{g(\phi^* \tilde{B}_t) \phi^* \mathbf{v}^\top \Sigma_{det} \mathbf{v}},$$

where z_α is such that $\mathcal{N}(0, 1)(z_\alpha, +\infty) = \alpha/2$.

B.3. Hypothesis tests on the matrix Φ based on (\mathbf{A}_t) . We now focus on the case $N \geq 2$ and the inferential problem of testing the null hypothesis $H_0 : \Phi = \Phi_0$, using the multi-dimensional stochastic process $(\mathbf{A}_t)_t$. Since the distribution of \tilde{A}_∞ should be unknown in practice, we propose a test statistics whose limit does not involve \tilde{A}_∞ . First, we need to introduce some notation. Let us define the matrix \mathcal{M}_{det}^{11} as the matrix \mathcal{M}_∞^{11} defined in Theorem A.3 with Σ_∞ replaced by Σ_{det} , i.e. $\mathcal{M}_\infty^{11} = g(\tilde{A}_\infty) \mathcal{M}_{det}^{11}$. Then, notice that the matrix \mathcal{M}_{det}^{11} has rank $(N-1)$ and it admits a spectral decomposition as follows: $\mathcal{M}_{det}^{11} = O^{11} \mathcal{D}^{11} (O^{11})^\top$, where \mathcal{D}^{11} is the diagonal matrix of dimension $(N-1)$ containing the eigenvalues of \mathcal{M}_{det}^{11} different from 0 and O^{11} is the $N \times (N-1)$ matrix whose columns form an orthonormal basis of $Im(\mathcal{M}_{det}^{11})$ of corresponding eigenvectors. Then, we set $L^{11} = (\mathcal{D}^{11})^{-1/2}$ (diagonal matrix of dimension $(N-1)$) and take the product $L^{11} (O^{11})^\top$ (matrix

of dimension $(N - 1) \times N$). Fixed the matrix assumed under H_0 , i.e. Φ_0 , we can compute for it the eigenvectors \mathbf{u} and \mathbf{v} , the eigenvectors contained in the matrices U and V and the matrices L^{11} and O^{11} defined above. Hence, we can obtain under H_0 the real process $\tilde{A}_t = \mathbf{v}^\top \mathbf{A}_t$ and the multi-dimensional process $\hat{\mathbf{A}}_t = \mathbf{A}_t - \tilde{A}_t \mathbf{u} = UV^\top \mathbf{A}_t$. Then, from Theorem A.3 we have that, under H_0 and the assumption $\mathcal{R}e(\phi_2^*)/\phi^* < 1/2$,

$$(33) \quad \mathbf{T}_t = t^{\phi^*/2} g(\tilde{A}_t)^{-1/2} L^{11} (O^{11})^\top UV^\top \mathbf{A}_t = t^{\phi^*/2} g(\tilde{A}_t)^{-1/2} L^{11} (O^{11})^\top \hat{\mathbf{A}}_t \stackrel{d}{\sim} \mathcal{N}(\mathbf{0}, I_{N-1})$$

(in the above formulas the matrices L^{11} and O^{11} are related to \mathcal{M}_{det}^{11} computed under H_0 and so for Φ_0). Hence, under H_0 , the test statistics $\|\mathbf{T}_t\|^2$ is asymptotically chi-squared distributed with $(N - 1)$ degrees of freedom. This result lets us construct an asymptotic critical region for testing any Φ_0 .

Remark 2. (Power)

The performance in terms of power of this inferential procedure is strongly related to the matrix Φ_1 considered under the alternative hypothesis H_1 . For instance, the leading eigenvalue of Φ under H_0 may not be equal to the one under H_1 and so the test statistics may go to infinity with t at different rates under H_0 or H_1 . Moreover, the leading right eigenvector \mathbf{v}_0 computed under $H_0 : \Phi = \Phi_0$ may not be equal to the leading right eigenvector \mathbf{v}_1 of Φ_1 and so it is possible that $(\mathbf{v}_0^\top \mathbf{A}_t) \neq (\mathbf{v}_1^\top \mathbf{A}_t)$. In that case, we would have, under H_1 , that $\mathbf{A}_t \xrightarrow{a.s.} \tilde{A}_\infty \mathbf{u}_1$ but $(\mathbf{v}_0^\top \mathbf{A}_t) \xrightarrow{a.s.} (\mathbf{v}_0^\top \mathbf{u}_1) \tilde{A}_\infty$ and hence, since we use $\tilde{A}_t = \mathbf{v}_0^\top \mathbf{A}_t$ in the test statistics, we can be sure that $g(\tilde{A}_t) \xrightarrow{a.s.} g(\tilde{A}_\infty)$ remains valid under H_1 only if $(\mathbf{v}_0^\top \mathbf{u}_1) = 1$. This condition is guaranteed in a framework where the leading left eigenvector \mathbf{u} of Φ is known in advanced (e.g. under the popular balance condition $\Phi^\top \mathbf{1} = \mathbf{1}$) and so in that case we would always have $\mathbf{u}_0 = \mathbf{u}_1 = \mathbf{u}$. Analogously, the columns of U_1 and V_1 computed under $H_1 : \Phi = \Phi_1$ may not be equal to the corresponding U_0 and V_0 derived under $H_0 : \Phi = \Phi_0$. However, if we assume $V_0^\top \mathbf{u}_1 = \mathbf{0}$ (that is automatically true when $\mathbf{u}_0 = \mathbf{u}_1$) we can be sure that, under H_1 , we have $U_0 V_0^\top \mathbf{A}_t = U_0 V_0^\top \hat{\mathbf{A}}_t$, since

$$U_0 V_0^\top \mathbf{A}_t = U_0 V_0^\top (\mathbf{A}_t - \tilde{A}_t \mathbf{u}_1) = U_0 V_0^\top \hat{\mathbf{A}}_t.$$

In summary, whenever $\mathbf{u}_0 = \mathbf{u}_1$ we have under H_1 that \mathbf{T}_t is still asymptotically normally distributed with covariance matrix $L^{11} (O^{11})^\top UV^\top \mathcal{M}_{det}^{11} VU^\top O^{11} (L^{11})^\top$, where \mathcal{M}_{det}^{11} is referred to the eigen-structure of Φ_1 , while $L^{11} (O^{11})^\top UV^\top$ is referred to Φ_0 . Consequently, the distance between the identity matrix I and the matrix $L^{11} (O^{11})^\top UV^\top \mathcal{M}_{det}^{11} VU^\top O^{11} (L^{11})^\top$ describes the relation between the asymptotic distribution of $\|\mathbf{T}_t\|^2$ under H_0 and the one under H_1 , which determines the power of the test. For instance, note that $E[\|\mathbf{T}_t\|^2] = (N - 1)$ under H_0 , while $E[\|\mathbf{T}_t\|^2]$ is equal to the trace of $L^{11} (O^{11})^\top UV^\top \mathcal{M}_{det}^{11} VU^\top O^{11} (L^{11})^\top$ under H_1 .

B.4. Hypothesis tests on the matrix Φ based on (\mathbf{B}_t) . We here want to build for the case $N \geq 2$ a test statistic for $H_0 : \Phi = \Phi_0$, that is based only on the process (\mathbf{B}_t) . To this end, analogously as in Subsection B.3, let us define the matrix \mathcal{M}_{det}^{33} such that $\mathcal{M}_\infty^{33} = g(\tilde{A}_\infty) \mathcal{M}_{det}^{33}$ and the corresponding matrices L^{33} and O^{33} . Fixed the matrix Φ_0 assumed under H_0 , we can obtain the real process $\tilde{B}_t = \mathbf{v}^\top \mathbf{B}_t$ and the multi-dimensional process

$$\mathbf{B}'_t = \mathbf{B}_t - \tilde{B}_t \mathbf{u} = UV^\top \mathbf{B}_t.$$

By Corollary A.6 and Remark 1, we get that, under H_0 and the assumption $\mathcal{R}e(\phi_2^*)/\phi^* < 1/2$, we have

$$(34) \quad \mathbf{T}_t = t^{\phi^*/2} g(\phi^* \tilde{B}_t)^{-1/2} L^{33} (O^{33})^\top \mathbf{B}'_t = t^{\phi^*/2} g(\phi^* \tilde{B}_t)^{-1/2} L^{33} (O^{33})^\top UV^\top \mathbf{B}_t \stackrel{d}{\sim} \mathcal{N}(\mathbf{0}, I_{N-1})$$

(in the above formulas the matrices L^{33} and O^{33} are related to \mathcal{M}_{det}^{33} computed under H_0 and so for Φ_0). The considerations on the power made for the test based on \mathbf{A}_t in Remark 2 remain valid as well for the test based on \mathbf{B}_t .

APPENDIX C. STATISTICAL TOOLS IN SOME EXAMPLES

We here specify the previous general statistical tools in two particular cases: firstly, we consider the case $N = 2$ and, then, we take into account the case when Φ is of the ‘‘mean-field’’ type and $N \geq 2$.

C.1. Case $N = 2$. A matrix Φ with dimension 2×2 and satisfying condition (i) can be reparametrized imposing that the number $\phi \in (0, 1]$ is an eigenvalue with left eigenvector \mathbf{u} such that $r = u_1/u_2$, with $r > 0$, that is

$$r\phi_{11} + \phi_{21} = r\phi \quad \text{and} \quad r\phi_{12} + \phi_{22} = \phi.$$

This reparametrization has been inspired by the fact that we typically are able to estimate with high accuracy the leading eigenvalue $\phi^* \in (0, 1]$ and the ratio $r = u_1/u_2 > 0$ of the two componets of the leading left eigenvector \mathbf{u} . Hence, we may think that $\phi = \phi^*$ and r are known in practice and the only parameters that are object of the test are ϕ_{21} and ϕ_{12} . We can also calculate the second eigenvalue from the trace of Φ :

$$\phi_2^* = \phi_{1,1} + \phi_{2,2} - \phi = \phi - \frac{\phi_{2,1} + r^2\phi_{1,2}}{r},$$

so that the condition $\mathcal{R}e(\phi_2^*)/\phi^* < 1/2$ required in the previous results corresponds to the condition

$$2\frac{\phi_{2,1} + r^2\phi_{1,2}}{r} > \phi.$$

Moreover, since according to our notation we have $\mathbf{v}^\top \mathbf{1} = 1$ and $\mathbf{v}^\top \mathbf{u} = 1$, we obtain the following relations:

$$v_1 = \frac{r\phi_{1,2}}{r\phi_{1,2} + \phi_{2,1}}, \quad v_2 = 1 - v_1, \quad u_1 = \frac{r}{1 - (1 - r)v_1}, \quad u_2 = \frac{1}{1 - (1 - r)v_1},$$

which leads to

$$v_1 = \frac{r\phi_{1,2}}{r\phi_{1,2} + \phi_{2,1}}, \quad v_2 = \frac{\phi_{2,1}}{r\phi_{1,2} + \phi_{2,1}}, \quad u_1 = \frac{r\phi_{2,1} + r^2\phi_{1,2}}{\phi_{2,1} + r^2\phi_{1,2}}, \quad u_2 = \frac{\phi_{2,1} + r\phi_{1,2}}{\phi_{2,1} + r^2\phi_{1,2}}.$$

If we adopt the parametrization $\iota = \phi_{1,2} + \phi_{2,1}$ and $\eta = \frac{\phi_{1,2}}{\phi_{1,2} + \phi_{2,1}}$ so that ι represents the intensity of the whole ‘‘interaction’’ among the two components 1 and 2 due to matrix Φ and η (resp. $(1 - \eta)$) gives the percentage of this intensity due to the influence of the first component on the second one (resp., of the second component on the first one). Then, we have

$$\mathbf{u} = \frac{(1 - \eta) + r\eta}{(1 - \eta) + r^2\eta} \begin{pmatrix} r \\ 1 \end{pmatrix} \quad \text{and} \quad \mathbf{v} = \frac{1}{r\eta + (1 - \eta)} \begin{pmatrix} r\eta \\ 1 - \eta \end{pmatrix},$$

and so

$$\mathbf{u}\mathbf{v}^\top = \frac{1}{(1 - \eta) + r^2\eta} \begin{pmatrix} r \\ 1 \end{pmatrix} \begin{pmatrix} r\eta & 1 - \eta \end{pmatrix} \quad \text{and} \quad UV^\top = \frac{1}{(1 - \eta) + r^2\eta} \begin{pmatrix} 1 - \eta \\ -r\eta \end{pmatrix} \begin{pmatrix} 1 & -r \end{pmatrix}.$$

Finally, imposing $U^\top U = 1$, we get

$$U = \frac{1}{\sqrt{(1 - \eta)^2 + r^2\eta^2}} \begin{pmatrix} 1 - \eta \\ -r\eta \end{pmatrix} \quad \text{and} \quad V = \frac{\sqrt{(1 - \eta)^2 + r^2\eta^2}}{(1 - \eta) + r^2\eta} \begin{pmatrix} 1 \\ -r \end{pmatrix}.$$

We thus obtain

$$\phi^* \mathbf{v}^\top \Sigma_{det} \mathbf{v} = \frac{\phi}{(r\eta + (1-\eta))^2} (r\eta \quad 1-\eta) \Sigma_{det} \begin{pmatrix} r\eta \\ 1-\eta \end{pmatrix}$$

and, for $2\iota \frac{(1-\eta)+r^2\eta}{r} > \phi$,

$$\mathcal{M}_{det}^{11} = U \mathcal{S}_{det}^{11} U^\top = \frac{(\iota \frac{(1-\eta)+r^2\eta}{r} - \phi)^2}{2\iota \frac{(1-\eta)+r^2\eta}{r} - \phi} \frac{1}{((1-\eta) + r^2\eta)^2} \begin{pmatrix} 1-\eta \\ -r\eta \end{pmatrix} (1 \quad -r) \Sigma_{det} \begin{pmatrix} 1 \\ -r \end{pmatrix} (1-\eta \quad -r\eta),$$

$$\mathcal{M}_{det}^{33} = U \mathcal{S}_{det}^{33} U^\top = \frac{1}{2\iota \frac{(1-\eta)+r^2\eta}{r} - \phi} \frac{1}{((1-\eta) + r^2\eta)^2} \begin{pmatrix} 1-\eta \\ -r\eta \end{pmatrix} (1 \quad -r) \Sigma_{det} \begin{pmatrix} 1 \\ -r \end{pmatrix} (1-\eta \quad -r\eta).$$

(where \mathcal{S}_{det}^{ii} are the matrices defined in Theorem A.3 with Σ_∞ replaced by Σ_{det}). Therefore, recalling that

$$\tilde{A}_t = \mathbf{v}^\top \mathbf{A}_t = \frac{r\eta A_{1,t} + (1-\eta)A_{2,t}}{r\eta + (1-\eta)} \quad \text{and} \quad \tilde{B}_t = \mathbf{v}^\top \mathbf{B}_t = \frac{r\eta B_{1,t} + (1-\eta)B_{2,t}}{r\eta + (1-\eta)},$$

and defining the constant

$$c_{r,\eta} = \frac{(r\eta \quad 1-\eta) \Sigma_{det} \begin{pmatrix} r\eta \\ 1-\eta \end{pmatrix}}{(r\eta + (1-\eta))^2},$$

the confidence interval based on \mathbf{A}_t and \mathbf{B}_t are respectively

$$CI_{1-\alpha}(\tilde{A}_\infty) = \tilde{A}_t \pm \frac{z_\alpha}{t^{\phi/2}} \sqrt{\phi g(\tilde{A}_t) c_{r,\eta}}, \quad \text{and} \quad CI_{1-\alpha}(\tilde{B}_\infty) = \phi \tilde{B}_t \pm \frac{z_\alpha}{t^{\phi/2}} \sqrt{\phi g(\phi \tilde{B}_t) c_{r,\eta}}.$$

Analogously, the test statistics based on \mathbf{A}_t is

$$T_t = t^{\phi/2} \frac{1}{(1-\eta) + r^2\eta} L^{11} (O^{11})^\top \begin{pmatrix} 1-\eta \\ -r\eta \end{pmatrix} \frac{(A_{1,t} - rA_{2,t})}{g(\tilde{A}_t)^{1/2}},$$

and the one based on \mathbf{B}_t

$$T_t = t^{\phi/2} \frac{1}{(1-\eta) + r^2\eta} L^{33} (O^{33})^\top \begin{pmatrix} 1-\eta \\ -r\eta \end{pmatrix} \frac{(B_{1,t} - rB_{2,t})}{g(\phi \tilde{B}_t)^{1/2}}.$$

Remark 3. In the particular case $\Sigma_{det} = \text{diag}(\mathbf{u})$ we have

$$\mathcal{S}_{det}^{33} = \frac{1}{\left(2\iota \frac{(1-\eta)+r^2\eta}{r} - \phi\right)} \frac{(r+r^2)((1-\eta)+r\eta)((1-\eta)^2+r^2\eta^2)}{((1-\eta)+r^2\eta)^3},$$

which leads to $L^{33} (O^{33})^\top = (\mathcal{S}_{det}^{33})^{-1/2} U^\top$ and so the test statistics based on \mathbf{B}_t becomes

$$T_t = t^{\phi/2} \sqrt{\left(2\iota \frac{(1-\eta)+r^2\eta}{r} - \phi\right)} \frac{(1-\eta) + r^2\eta}{(r+r^2)((1-\eta)+r\eta)} \frac{(B_{1,t} - rB_{2,t})}{g(\phi \tilde{B}_t)^{1/2}}.$$

Then, denoting

$$\Delta = \frac{1}{2} - \frac{\phi_2^*}{\phi} = \frac{\iota}{\phi} \frac{(1-\eta) + r^2\eta}{r} - \frac{1}{2} \quad \text{and} \quad q_{r,\eta} = \frac{2((1-\eta) + r^2\eta)}{r(1+r)((1-\eta) + r\eta)},$$

we have

$$(35) \quad \|T_t\|^2 = t^\phi q_{r,\eta} \Delta \frac{(B_{1,t} - rB_{2,t})^2}{g(\phi \tilde{B}_t)/\phi}.$$

Moreover, the confidence interval based on \mathbf{B}_t becomes

$$(36) \quad CI_{1-\alpha}(\tilde{B}_\infty) = \phi \tilde{B}_t \pm \frac{z_\alpha}{t^{\phi/2}} \sqrt{\phi g(\phi \tilde{B}_t) c_{r,\eta}}, \quad \text{with} \quad c_{r,\eta} = \frac{(r^3\eta^2 + (1-\eta)^2)}{(r\eta + (1-\eta))(r^2\eta + (1-\eta))}.$$

C.2. Mean-field case for $N \geq 2$. This first case we consider is the ‘‘Mean-field’’ case, which refers to the family of matrices Φ that can be expressed as follows: for any $1 \leq h, j \leq N$

$$\phi_{j,h} = \phi \left(\frac{\iota}{N} + \delta_{j,h}(1 - \iota) \right) \quad \text{with } \phi, \iota \in (0, 1],$$

where $\delta_{j,h}$ is equal to 1 when $h = j$ and to 0 otherwise. Note that Φ is irreducible and doubly stochastic, we have $\mathbf{v} = N^{-1}\mathbf{1}$ and $\mathbf{u} = \mathbf{1}$ and so the random variable \tilde{A}_t coincides with the average of the processes $A_{t,h}$, i.e. $N^{-1}\mathbf{1}^\top \mathbf{A}_t$ and $\hat{\mathbf{A}}_t = (I - N^{-1}\mathbf{1}\mathbf{1}^\top) \mathbf{A}_t$. Furthermore, in this case, the leading eigenvalue ϕ^* coincides with ϕ , while all the eigenvalues of Φ different from ϕ^* are equal to $\phi(1 - \iota)$ and, consequently, the condition $\mathcal{R}e(\phi_2^*)/\phi^* < 1/2$ required in the previous results corresponds to the condition $2\iota > 1$. Finally, since Φ is also symmetric, we have $U = V$ and so $U^\top U = V^\top V = I$ and $UU^\top = VV^\top = I - N^{-1}\mathbf{1}\mathbf{1}^\top$. Notice that the mean-field matrix has two parameters, ϕ and ι , but, since the leading eigenvalue ϕ^* is typically easy to be estimated with high accuracy, we may think that the only parameter that is object of inference in this case is ι , which is the quantity that rules the intensity of the ‘‘interaction’’ among the different components due to matrix Φ . We thus obtain $\phi^* \mathbf{v}^\top \Sigma_{det} \mathbf{v} = \phi N^{-2} (\mathbf{1}^\top \Sigma_{det} \mathbf{1})$ and, for $2\iota > 1$,

$$\mathcal{M}_{det}^{11} = U \mathcal{S}_{det}^{11} U^\top = \frac{\phi(1 - \iota)^2}{2\iota - 1} UV^\top \Sigma_{det} VU^\top = \frac{\phi(1 - \iota)^2}{2\iota - 1} (I - N^{-1}\mathbf{1}\mathbf{1}^\top) \Sigma_{det} (I - N^{-1}\mathbf{1}\mathbf{1}^\top),$$

$$\mathcal{M}_{det}^{33} = U \mathcal{S}_{det}^{33} U^\top = \frac{1}{\phi(2\iota - 1)} UV^\top \Sigma_{det} VU^\top = \frac{1}{\phi(2\iota - 1)} (I - N^{-1}\mathbf{1}\mathbf{1}^\top) \Sigma_{det} (I - N^{-1}\mathbf{1}\mathbf{1}^\top),$$

Therefore, recalling $\tilde{A}_t = \frac{\mathbf{1}^\top \mathbf{A}_t}{N}$, $\tilde{B}_t = \frac{\mathbf{1}^\top \mathbf{B}_t}{N}$ and defining $c_N = \frac{(\mathbf{1}^\top \Sigma_{det} \mathbf{1})}{N^2}$, the confidence interval based on \mathbf{A}_t and \mathbf{B}_t are respectively

$$CI_{1-\alpha}(\tilde{A}_\infty) = \tilde{A}_t \pm \frac{z_\alpha}{t^{\phi/2}} \sqrt{\phi g(\tilde{A}_t) c_N} \quad \text{and} \quad CI_{1-\alpha}(\tilde{A}_\infty) = \phi \tilde{B}_t \pm \frac{z_\alpha}{t^{\phi/2}} \sqrt{\phi g(\phi \tilde{B}_t) c_N}.$$

Analogously, the test statistics based on \mathbf{A}_t is

$$\mathbf{T}_t = t^{\phi/2} L^{11} (O^{11})^\top UV^\top \frac{\mathbf{A}_t}{g(\tilde{A}_t)^{1/2}} = t^{\phi/2} L^{11} (O^{11})^\top (I - N^{-1}\mathbf{1}\mathbf{1}^\top) \frac{\mathbf{A}_t}{g(\tilde{A}_t)^{1/2}},$$

and the one based on \mathbf{B}_t is

$$\mathbf{T}_t = t^{\phi/2} L^{33} (O^{33})^\top UV^\top \frac{\mathbf{B}_t}{g(\phi \tilde{B}_t)^{1/2}} = t^{\phi/2} L^{33} (O^{33})^\top (I - N^{-1}\mathbf{1}\mathbf{1}^\top) \frac{\mathbf{B}_t}{g(\phi \tilde{B}_t)^{1/2}}.$$

Remark 4. In the particular case $\Sigma_{det} = I$ we have $\mathbf{1}^\top \Sigma_{det} \mathbf{1} = N$ and hence the following confidence interval based on \mathbf{B}_t :

$$(37) \quad CI_{1-\alpha}(\tilde{A}_\infty) = \phi \tilde{B}_t \pm \frac{z_\alpha}{t^{\phi/2}} \sqrt{\phi g(\phi \tilde{B}_t) c_N} \quad \text{with} \quad c_N = \frac{1}{N}.$$

Moreover, we have $\mathcal{M}_{det}^{33} = U \mathcal{S}_{det}^{33} U^\top$, with

$$UU^\top = (I - N^{-1}\mathbf{1}\mathbf{1}^\top) \quad \text{and} \quad \mathcal{S}_{det}^{33} = \frac{1}{\phi(2\iota - 1)} I,$$

which leads to $L^{33} (O^{33})^\top = \sqrt{\phi(2\iota - 1)} U^\top$ and so the test statistics based on \mathbf{B}_t becomes

$$\mathbf{T}_t = t^{\phi/2} \sqrt{\phi(2\iota - 1)} U^\top \frac{\mathbf{B}_t}{g(\phi \tilde{B}_t)^{1/2}},$$

from which we can get

$$\begin{aligned}\|\mathbf{T}_t\|^2 &= t^\phi(2\iota - 1) \frac{\mathbf{B}_t^\top (I - N^{-1}\mathbf{1}\mathbf{1}^\top) \mathbf{B}_t}{g(\phi\tilde{\mathbf{B}}_t)/\phi} = t^\phi(2\iota - 1) \frac{\|\mathbf{B}_t\|^2 - \frac{(\mathbf{1}^\top \mathbf{B}_t)^2}{N}}{g(\tilde{\mathbf{B}}_t)/\phi} \\ &= t^\phi(2\iota - 1) \frac{\|\mathbf{B}_t - \left(\frac{\mathbf{1}^\top \mathbf{B}_t}{N}\right) \mathbf{1}\|^2}{g(\phi\tilde{\mathbf{B}}_t)/\phi},\end{aligned}$$

that is

$$(38) \quad \|T_t\|^2 = t^\phi 2\Delta \frac{\|\mathbf{B}_t - \tilde{\mathbf{B}}_t \mathbf{1}\|^2}{g(\phi\tilde{\mathbf{B}}_t)/\phi} \quad \text{with} \quad \Delta = \frac{1}{2} - \frac{\phi_2^*}{\phi} = \iota - \frac{1}{2}.$$

APPENDIX D. TECHNICAL RESULTS

For the reader's convenience, we here recall some results used in the above proofs. (For a synthetic review about the notion of stable convergence and its variants, see for instance the appendix of [4] and the references recalled therein.)

Theorem D.1. (A consequence of [10, Theorem A.1])

On (Ω, \mathcal{A}, P) , let $(\mathcal{F}_t)_t$ be a filtration and, for each $t \geq 1$, let $(L_{t,k})_{k \in \mathbb{N}}$ be a real martingale with respect to $(\mathcal{F}_{t+k-1})_k$, with $L_{t,0} = 0$, which converges in mean to a random variable $L_{t,\infty}$. Set

$$Y_{t,n} = L_{t,n} - L_{t,n-1} \quad \text{for } n \geq 1, \quad U_t = \sum_{n \geq 1} Y_{t,n}^2, \quad Y_t^* = \sup_{n \geq 1} |Y_{t,n}|.$$

Further, suppose that $(U_t)_t$ converges almost surely to a positive real $(\mathcal{F}_\infty$ -measurable) random variable U and $Y_t^* \leq y(t)$ with $y(t)$ a deterministic function such that $y(t) \xrightarrow{a.s.} 0$.

Then, with respect to $(\mathcal{F}_t)_t$, the sequence $(L_{t,\infty})_t$ converges to the Gaussian kernel $\mathcal{N}(0, U)$ in the sense of the almost sure conditional convergence (and so also stably).

Lemma D.2. ([6, Supplementary material])

If $a_t \geq 0$, $a_t \leq 1$ for t large enough, $\sum_t a_t = +\infty$, $\delta_t \geq 0$, $\sum_t \delta_t < +\infty$, $b > 0$, $y_t \geq 0$ and $y_{t+1} \leq (1 - a_t)^b y_t + \delta_t$, then $\lim_{t \rightarrow +\infty} y_t = 0$.

Theorem D.3. (A consequence of [12, Proposition 3.1])

Let $(\mathbf{T}_{t,k+1})_{t \geq m_0, m_0 \leq k \leq t}$ be a triangular array of d -dimensional real random vectors, such that, for each fixed t , the finite sequence $(\mathbf{T}_{t,k+1})_{m_0 \leq k \leq t}$ is a martingale difference array with respect to a given filtration $(\mathcal{F}_{k+1})_{k \geq m_0}$. Moreover, let $(s_t)_t$ be a sequence of positive real numbers and assume that the following conditions hold:

- (1) $\sum_{k=m_0}^t (s_t \mathbf{T}_{t,k+1})(s_t \mathbf{T}_{t,k+1})^\top = s_t^2 \sum_{k=m_0}^t \mathbf{T}_{t,k+1} \mathbf{T}_{t,k+1}^\top \xrightarrow{P} \mathcal{M}_\infty$, where \mathcal{M}_∞ is a $(\mathcal{F}_\infty$ -measurable) random positive semidefinite matrix;
- (2) $\sup_{m_0 \leq k \leq t} |s_t \mathbf{T}_{t,k+1}| = s_t \sup_{m_0 \leq k \leq t} \sum_{i=1}^d |T_{t,k+1,i}| \xrightarrow{L^1} 0$.

Then $(s_t \sum_{k=m_0}^t \mathbf{T}_{t,k+1})_t$ converges stably to the Gaussian kernel $\mathcal{N}(\mathbf{0}, \mathcal{M}_\infty)$.

The following result combines together stable convergence and almost sure conditional convergence.

Lemma D.4. (A consequence of [9, Lemma 1])

Suppose that $(\mathbf{Y}_t^{(1)})$ and $(\mathbf{Y}_t^{(2)})$ are real random vectors such that the first is adapted to a filtration $(\mathcal{F}_t)_t$ and the second is measurable with respect to \mathcal{F}_∞ . If $(\mathbf{Y}_t^{(1)})$ converges stably to $\mathcal{N}(0, \mathcal{M}_\infty^{(1)})$ and $(\mathbf{Y}_t^{(2)})$ converges to $\mathcal{N}(0, \mathcal{M}_\infty^{(2)})$ in the sense of the almost sure conditional convergence with respect to $(\mathcal{F}_t)_t$, then

$$(\mathbf{Y}_t^{(1)} + \mathbf{Y}_t^{(2)}) \xrightarrow{\text{stably}} \mathcal{N}(\mathbf{0}, \mathcal{M}_\infty^{(1)} + \mathcal{M}_\infty^{(2)}).$$

ACKNOWLEDGMENTS

Giacomo Aletti is a member of the Italian Group “Gruppo Nazionale per il Calcolo Scientifico” of the Italian Institute “Istituto Nazionale di Alta Matematica”. Irene Crimaldi is a member of the Italian Group “Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni” of the Italian Institute “Istituto Nazionale di Alta Matematica”.

Funding Sources

Irene Crimaldi is partially supported by the European Union - NextGenerationEU through the Italian Ministry of University and Research under the National Recovery and Resilience Plan (PNRR) - M4C2 - Investment 1.3, title [Public Sector Indicators for Sustainability and Wellbeing (PUWELL)] - Program [Growing Resilient, INclusive and Sustainable (GRINS)] - PE18 - CUP J33C22002910001.

REFERENCES

- [1] G. Aletti and I. Crimaldi. Twitter as an innovation process with damping effect. *Sci. Rep.*, 11(1), 2021.
- [2] G. Aletti, I. Crimaldi, and A. Ghiglietti. Synchronization of reinforced stochastic processes with a network-based interaction. *Ann. Appl. Probab.*, 27(6):3787–3844, 2017.
- [3] G. Aletti, I. Crimaldi, and A. Ghiglietti. Networks of reinforced stochastic processes: asymptotics for the empirical means. *Bernoulli*, 25(4 B):3339–3378, 2019.
- [4] G. Aletti, I. Crimaldi, and A. Ghiglietti. Interacting reinforced stochastic processes: Statistical inference based on the weighted empirical means. *Bernoulli*, 26(2):1098–1138, 2020.
- [5] G. Aletti, I. Crimaldi, and A. Ghiglietti. Interacting innovation processes. *Sci. Rep.*, 13:17187, 2023.
- [6] G. Aletti, I. Crimaldi, and A. Ghiglietti. Networks of reinforced stochastic processes: A complete description of the first-order asymptotics. *Stochastic Process. Appl.*, 176:104427, 2024.
- [7] G. Armano and M. A. Javarone. The beneficial role of mobility for the emergence of innovation. *Sci. Rep.*, 7:1781, 2017.
- [8] W. Arthur. *The Nature of Technology: What It Is and How It Evolves*. Free Press, 2009.
- [9] P. Berti, I. Crimaldi, L. Pratelli, and P. Rigo. A central limit theorem and its applications to multicolor randomly reinforced urns. *J. Appl. Probab.*, 48(2):527–546, 2011.
- [10] I. Crimaldi. An almost sure conditional convergence result and an application to a generalized Pólya urn. *Int. Math. Forum*, 4(21-24):1139–1156, 2009.
- [11] I. Crimaldi, P. Dai Pra, and I. G. Minelli. Fluctuation theorems for synchronization of interacting Pólya’s urns. *Stoch. Process. Appl.*, 126(3):930–947, 2016.
- [12] I. Crimaldi and L. Pratelli. Convergence results for multivariate martingales. *Stoch. Process. Appl.*, 115(4):571–577, 2005.
- [13] S. Favaro, A. Lijoi, R. H. Mena, and I. Prünster. Bayesian non-parametric inference for species variety with a two-parameter poisson–dirichlet process prior. *J. R. Stat. Soc. Ser. B*, 71(5):993–1008, 2009.
- [14] T. M. A. Fink, M. Reeves, R. Palma, and R. S. Farr. Serendipity and strategy in rapid innovation. *Nat. Commun.*, 8(2002), 2017.
- [15] S. Fortini, S. Petrone, and P. Sporysheva. On a notion of partially conditionally identically distributed sequences. *Stoch. Process. Appl.*, 128(3):819–846, 2018.
- [16] R. Gouet. Martingale Functional Central Limit Theorems for a Generalized Polya Urn. *Ann. Probab.*, 21(3):1624 – 1639, 1993.
- [17] B. Hansen and J. Pitman. Prediction rules for exchangeable sequences related to species sampling. *Statist. Probab. Lett.*, 46:251–256, 2000.
- [18] H. S. Heaps. *Information Retrieval-Computational and Theoretical Aspects*. Academic Press, 1978.
- [19] I. Iacopini, G. Di Bona, E. Ubaldi, V. Loreto, and V. Latora. Interacting discovery processes on complex networks. *Phys. Rev. Lett.*, 125:248301, Dec 2020.
- [20] L. F. James, A. Lijoi, and I. Prünster. Distributions of linear functionals of two parameter Poisson–Dirichlet random measures. *Ann. Appl. Probab.*, 18(2):521 – 551, 2008.
- [21] S. A. Kauffman. *Investigations*. Oxford University Press, New York/Oxford, 2000.
- [22] A. Lijoi, R. Mena, and I. Prünster. Bayesian nonparametric estimation of the probability of discovering new species. *Biometrika*, 94:769–786, 08 2007.
- [23] A. Lijoi and I. Prünster. *Models beyond the Dirichlet process*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2010.

- [24] A. Mokkadem and M. Pelletier. Convergence rate and averaging of nonlinear two-time-scale stochastic approximation algorithms. *Ann. Appl. Probab.*, 16(3):1671–1702, 08 2006.
- [25] C. Monti, J. D’Ignazi, M. Starnini, and G. De Francisci Morales. Evidence of demographic rather than ideological segregation in news discussion on reddit. In *Proceedings of the ACM Web Conference 2023 (WWW ’23)*, 2023.
- [26] J. Pitman. Some developments of the blackwell-macqueen urn scheme. *Lecture Notes-Monograph Series*, 30:245–267, 1996.
- [27] J. Pitman. *Combinatorial Stochastic Processes*. Ecole d’Eté de Probabilités de Saint-Flour XXXII. Springer, 2006.
- [28] J. Pitman and M. Yor. The two-parameter poisson-dirichlet distribution derived from a stable subordinator. *Ann. Probab.*, 25(2):855–900, 1997.
- [29] A. Puglisi, A. Baronchelli, and V. Loreto. Cultural route to the emergence of linguistic categories. *Proceedings of the National Academy of Sciences*, 105(23):7936–7940, 2008.
- [30] E. Rogers, A. Singhal, and M. Quinlan. *Diffusion of innovations*. Taylor and Francis, 2019.
- [31] A. Rzhetsky, J. G. Foster, I. T. Foster, and J. A. Evans. Choosing experiments to accelerate collective discovery. *Proceedings of the National Academy of Sciences*, 112(47):14569–14574, 2015.
- [32] H. Sarioev, S. Fortini, and S. Petrone. Infinite-color randomly reinforced urns with dominant colors. *Bernoulli*, 29(1):132 – 152, 2023.
- [33] F. Tria, V. Loreto, V. D. P. Servedio, and S. H. Strogatz. The dynamics of correlated novelties. *Sci. Rep.*, 4, 2014.
- [34] G. K. Zipf. *Human Behavior and the Principle of Least Effort*. Addison-Wesley, Reading MA (USA), 1949.

ADAMSS CENTER, UNIVERSITÀ DEGLI STUDI DI MILANO, MILAN, ITALY
Email address: giacomo.aletti@unimi.it

IMT SCHOOL FOR ADVANCED STUDIES LUCCA, LUCCA, ITALY
Email address: irene.crimaldi@imtlucca.it

UNIVERSITÀ DEGLI STUDI DI MILANO-BICOCCA, MILAN, ITALY
Email address: andrea.ghiglietti@unimib.it (Corresponding author)