Comment on "Quantum coherence between mass eigenstates of a neutrino cannot be destroyed by its mass-momentum entanglement"

Shi-Biao Zheng*

College of Physics and Information Engineering, Fuzhou University,
Fuzhou 350108, China

In arXiv:2410.21850, I proved that the quantum coherence between the mass eigenstates of a neutrino will be destroyed if they are correlated with different momenta. In arXiv:2411.01190, James M. Cline claimed that I had made the unrealistic assumption that the neutrino is always in a nearly exact energy eigenstate, and ignored the spatial dependence of the wavefunction in my paper. However, I did not assume that the neutrino is in a nearly exact eigenstate of energy anywhere in my paper, and the wavefunction I wrote in the position representation has a spatial dependence. The argumentation of arXiv:2411.01190 is based on misinterpreting my claim, and on ignoring the critical fact that the neutrino's wavepacket has a finite size and the detector has a large volume.

In Ref. 1, I proved that the quantum coherence between the mass eigenstates of a neutrino will be destroyed if they are correlated with different momenta. This point can be well understood in the momentum representation, where the state of the neutrino can be written as

$$|\nu\rangle = \sum_{j} \int_{\sigma_{j}} d^{3}\mathbf{p}_{j} f(\mathbf{p}_{j}) |\mathbf{p}_{j}\rangle |\nu_{j}\rangle.$$
 (1)

Here $|\nu_j\rangle$ denotes the jth mass eigenstate, and σ_j represents the distribution region of the momentum associated with the mass eigenstate $|\nu_j\rangle$, with the probability amplitude distribution function $f(\mathbf{p}_j)$. As the flavor oscillations are assumed to arise from the quantum coherence among the mass eigenstates, the momentum degree of freedom needs to be traced out for the discussion of such oscillations. This leaves the mass degree of freedom in a classical mixture, given by the density operator

$$\rho = \sum_{j,k} D_{j,k} |\nu_j\rangle \langle \nu_k|, \qquad (2)$$

where

$$D_{j,k} = \int d^{3}\mathbf{p} \int_{\sigma_{j}} \int_{\sigma_{k}} f(\mathbf{p}_{j}) f^{*}(\mathbf{p}_{k}) \langle \mathbf{p} | \mathbf{p}_{j} \rangle \langle \mathbf{p}_{k} | \mathbf{p} \rangle$$

$$= \int_{\sigma_{k}} \int_{\sigma_{k}} f(\mathbf{p}_{j}) f^{*}(\mathbf{p}_{k}) \langle \mathbf{p}_{k} | \mathbf{p}_{j} \rangle.$$
(3)

When there is no overlapping between σ_j and σ_k , we have $\mathbf{p}_j \neq \mathbf{p}_k$, which leads to $D_{j,k} = 0$ for $j \neq k$. This implies that the coherence among the mass eigenstates is destroyed when they are entangled with different momenta, so that the flavor oscillations cannot occur.

The entanglement-induced decoherence can also be understood in terms of quantum-mechanical complementarity [3,4]. When the mass eigenstates are correlated with different momenta, one can know in which mass eigenstate the neutrino is by measuring its momentum in principle. This is sufficient to destroy the coherence among the mass eigenstates. It does not matter that the momentum is not measured actually [5]. Such a complementary principle, which has passed a number of experimental tests [5-11], cannot be violated.

In the above derivation, there is no restriction on the energy and momentum for each mass eigenstate. In Ref. 2, James M. Cline seriously misinterpreted my starting point by saying "The essence of the claim is that a flavor state emitted from a weak interaction will be in a nearly exact eigenstate of energy." As a matter of fact, I have assumed that the momentum associated with each mass eigenstate has a spread. When the spread is sufficiently large, the neutrino cannot be in a nearly exact eigenstate of energy. The statement that the spatial dependence of the neutrino wavefunction was ignored in my derivation is also incorrect.

^{*} E-mail: t96034@fzu.edu.cn

For simplicity, we here suppose that the neutrino travels along the z direction. Then, in the position representation, the evolution of the wave function is given by

$$|\varphi_{\nu}(z,t)\rangle = \sum_{j} g_{j}(z,t) |\nu_{j}\rangle,$$
 (4)

where

$$g_j(z,t) = (2\pi)^{-1/2} \int_{\sigma_j} dp_j f(p_j) e^{i(p_j z - E_j t)},$$
 (5)

and $E_j = \sqrt{p_j^2 + m_j^2}$ with m_j being the mass of the jth mass eigenstate. Contrary to what was said in Ref. 2, the function $g_j(z,t)$ has a spatial dependence. For a definite position z, the coherence between $|\nu_j\rangle$ and $|\nu_k\rangle$ is manifested by the relative phase factor $e^{i(p_j-p_k)z}$. However, the neutrino detector has a large volume, and it was not recorded at which point of the detector the detected neutrino reacts with the medium in neutrino experiments. The probability for detecting the electron flavor on the detector is given by

$$P_e = \int_D dz \left| \langle \nu_e | \varphi_\nu(z, t) \rangle \right|^2 \tag{6}$$

where

$$|\nu_e\rangle = \sum_j U_{ej} |\nu_j\rangle \tag{7}$$

is the electron flavor eigenstate, and D is the detection region. Substituting Eqs. (4) and (7) into (6), we obtain

$$P_e = (2\pi)^{-1} \sum_{i,k} U_{ej}^* U_{ek} \int_D dz \int_{\sigma_j} dp_j \int_{\sigma_k} dp_k f(p_j) f^*(p_k) e^{i(p_j - p_k)z} e^{i(E_k - E_j)t}.$$
 (8)

When the neutrino detector has a size much larger than the wavepacket size, it is reasonable to replace $\int_D dz e^{i(p_j - p_k)z}$ with $\int_{-\infty}^{\infty} dz e^{i(p_j - p_k)z}$. For $p_j \neq p_k$, this integral vanishes, and P_e is approximated by

$$P_e \simeq \sum_j |U_{ej}|^2 \,, \tag{9}$$

which does not show interference effects. This is due to the fact that the position-dependent phase factor $e^{i(p_j-p_k)z}$, which is responsible for the spatial interference, is averaged out when the size of the detection region is larger than that of the neutrino wavepacket. The argumentation of Ref. 2 is valid only when the wavepacket size of the neutrino is much larger than the detector size, which does not coincide with neutrino experiments, where the detector has a large volume.

In Ref. [12], I further pointed out that the 2/3 deficit of solar ⁸B electron neutrinos cannot be reasonably interpreted in terms of the inconsistency between the flavor and mass eigenstates even if the corresponding superposition of mass eigenstates can be produced. This is due to the fact that the states of a neutrino and an electron are entangled after their charged-current interaction, which has been ignored in previous investigations of the matter effect [13-15]. Due to this nonseparability, the effects of the electrons cannot be modeled as a static potential for solar ⁸B neutrinos. Consequently, during the propagation ⁸B neutrinos cannot adiabatically evolve to a pure mass eigenstate, which was supposed to have an 1/3 overlapping with the electron flavor. In Ref. 12, I proposed an alternative mechanism, where neutrino oscillations are caused by virtual excitation of the Z bosonic field, which can connect different neutrino flavors. Under the competition between the coherent coupling induced by the Z bosonic field and the decoherence effect caused by charged-current interactions, solar ⁸B neutrinos would finally evolve to a steady state, where the electron flavor has a 1/3 population.

Finally, it should be pointed out that it is unreasonable to use the Standard Model of particle physics to criticize the mechanism proposed in Ref. 12. It is a well known fact that neutrino oscillations themselves are a phenomenon that is beyond the Standard Model, and certainly cannot be interpreted in the framework of the Standard Model. In other words, if one insists to judge the correctness of an interpretation in terms of the Standard Model, then there does not exist any correct interpretation.

^[1] S.-B. Zheng, Quantum coherence between mass eigenstates of a neutrino can be destroyed by its mass-momentum entanglement, arXiv:2410.21850.

- [2] J. M. Cline, Quantum coherence between mass eigenstates of a neutrino cannot be destroyed by its mass-momentum entanglementar, arXiv:2411.01190.
- [3] M. O. Scully, B.-G. Englert, and H. Walther, Quantum optical tests of complementarity, Nature 351, 111 (1991).
- [4] B.-G. Englert, Fringe visibility and which-way information: an inequality, Phys. Rev. Lett. 77, 2154 (1996).
- [5] M. Brune et al., Observing the progressive decoherence of the "meter" in a quantum measurement, Phys. Rev. Lett. 77, 4887 (1996).
- [6] E. Buks, R. Schuster, M. Heiblum, D. Mahalu, and V. Umansky, Dephasing in electron interference by a 'which-path' detector, Nature 391, 871 (1998).
- [7] S. Dürr, T. Nonn, and G. Rempe, Origin of quantum-mechanical complementarity probed by a 'which-way' experiment in an atom interferometer, Nature 395, 33 (1998).
- [8] S. Dürr, T. Nonn, and G. Rempe, Fringe visibility and which-way information in an atom interferometer, Phys. Rev. Lett. 81, 5705 (1998).
- [9] P. Bertet et al., A complementarity experiment with an interferometer at the quantum-classical boundary, Nature 411,166 (2001).
- [10] K. Liu et al., A twofold quantum delayed-choice experiment in a superconducting circuit, Sci. Adv. 3, e1603159 (2017).
- [11] X.-J. Huang et al., Entanglement-interference complementarity and experimental demonstration in a superconducting circuit, npj Quantum Information 9, 43 (2023).
- [12] S.-B. Zheng, Revisiting the origin of neutrino flavor transformations, arXiv:2407.00954.
- [13] L. Wolfenstein, Neutrino oscillations in matter, Phys. Rev. D 17, 2369 (1978).
- [14] S.P. Mikheyev and A.Yu. Smirnov, Resonance amplification of oscillations in matter and spectroscopy of solar neutrinos, Sov. J. Nucl. Phys. 42, 913 (1985).
- [15] T. K. Kuo and James Pantaleone, Neutrino oscillations in matter, Rev. Mod. Phys. 61, 937 (1989).