Graphs with given automorphism group and large clique number

John Haslegrave

January 20, 2025

Abstract

Barbieri recently showed that the finite graphs realising any given finite automorphism group have unbounded genus, answering a question of Cornwell et al. In this note we give a short proof of a stronger result: they have unbounded clique number.

In a paper concerned primarily with the minimum genus of graphs realising a given automorphism group, Cornwell, Doring, Lauderdale, Morgan and Storr asked how to determine the maximum genus of such a graph, or more precisely the maximum genus of a surface on which such a graph has a cellular embedding [2, Open Question 3]. Barbieri recently showed that the genus of a connected graph with given automorphism group is unbounded, and so this maximum does not exist for any group [1]. Barbieri's construction has high genus by virtue of containing a high-dimensional hypercube. However, we note that this structure does not guarantee that other parameters of potential interest, such as the chromatic number, are also large. We give a simple proof that graphs realising any given automorphism group have unbounded clique number, and hence unbounded genus and chromatic number.

Proposition 1. Let G be a non-complete finite graph, and let G' be the graph obtained from G by adding a disjoint clique with the same number of vertices and a perfect matching between the two. Then $\operatorname{Aut}(G') \cong \operatorname{Aut}(G)$.

Proof. Let n = |V(G)|. The statement holds for the unique non-complete graph on at most 2 vertices, so we may assume $n \ge 3$. We partition V(G') as $V_1 \cup V_2$, where $|V_1| = |V_2| = n$ and $G'[V_1] \cong G$ and $G'[V_2] \cong K_n$. Furthermore, we may write $V_2 = \{v' \mid v \in V_1\}$ such that $vv' \in E(G')$ for each $v \in V_1$. Note that $G'[V_2]$ is the unique copy of K_n in G', and so any automorphism ϕ of G' must have $\phi(V_2) = V_2$ and therefore $\phi(V_1) = V_1$. Consequently, ϕ restricted to V_1 is an automorphism of G. Conversely, for any $\psi \in \operatorname{Aut}(G)$, there is a unique $\psi' \in \operatorname{Aut}(G')$ whose restriction to V_1 is ψ , obtained by setting $\psi'(v') = \psi(v)'$ for each $v \in V_1$. For any $\psi_1, \psi_2 \in \operatorname{Aut}(G)$ we have $(\psi_1\psi_2)' = \psi'_1\psi'_2$, since these agree on V_1 , and hence $\operatorname{Aut}(G') \cong \operatorname{Aut}(G')$.

Corollary 2. For any graph G there exist connected graphs with automorphism group Aut(G) and arbitrarily large clique number.

Proof. If G is not complete, this follows immediately by iterating the construction of Proposition 1. If G is complete then $\operatorname{Aut}(G) \cong S_n$ for some n, and we may apply the same argument to some non-complete graph with the same automorphism group: \overline{K}_n if $n \ge 2$ or the graph obtained from the 6-vertex path by adding an edge between the second and fourth vertices if n = 1.

Since the possible values of $\operatorname{Aut}(G)$ range over all finite groups [3], it follows that any finite group Γ is the automorphism group of graphs of unbounded clique number.

References

- [1] M. Barbieri, Every group is the automorphism group of a graph with arbitrarily large genus. Preprint, arXiv:2501.08747 (2025).
- [2] C. Cornwell, M. Doring, L.-K. Lauderdale, E. Morgan, and N. Storr, On a new invariant for finite groups. J. Comb. 16 (2025), no. 1, 113–139.
- [3] R. Frucht, Herstellung von Graphen mit vorgegebener abstrakter Gruppe. Compositio Math. 6 (1939), 239–250.