# Optical gravitational waves as signals of Gravitationally-Decaying Particles

Alessandro Strumia<sup>a</sup>, Giacomo Landini<sup>b</sup>

<sup>a</sup> Dipartimento di Fisica, Università di Pisa, Italia <sup>b</sup> Instituto de Física Corpuscular. Universitat de València-CSIC. Spain

### Abstract

Long-lived heavy particles present during the big bang could have a decay channel opened by gravitons. Such decays can produce gravitational waves with large enough abundance to be detectable, and a peculiar narrow spectrum peaked today around optical frequencies. We identify which particles can decay in one or two gravitons. The maximal gravitational wave abundance arises from theories with extra hidden strong gauge dynamics, such as a confining pure-glue group. An interesting abundance also arises in theories with perturbative couplings. Future observation might shed light on early cosmology and allow some spectroscopy of sub-Planckian gravitationally-decaying particles, plausibly present in a variety of theories such as gauge unification, supersymmetry, extra dimensions, strings.

## Contents

1	Introduction	<b>2</b>
<b>2</b>	A particle decaying into gravitons	3
	2.1 Production of $S$	 4
	2.2 Decay of $S$	 4
	2.3 Gravitational wave spectrum	 5
	2.4 Decay of $S$ as sub-dominant component	 7
	2.5 Decay of $S$ as dominant component $\ldots \ldots \ldots$	 8
3	Allowed decays into gravitons	9
	3.1 Decay of a massive particle into two gravitons	 10
	3.2 Decay of a massive particle into a graviton and a SM particle	 11
	3.3 Decay of a massive particle into a graviton and a massive particle	 11
4	Theories for decay into gravitons	12
	4.1 Theories with non-perturbative interactions	 12
	4.2 Theories with perturbative interactions	 14
<b>5</b>	Conclusions	19

### 1 Introduction

Gravitons interact much less than neutrinos or other SM particles, so relic cosmological gravitational waves would provide information on the very early universe. However, the tiny interaction renders challenging detecting gravitational waves. The average cosmological energy density  $d\rho_{\rm GW}/d\ln f$  of relic gravitons g with frequency f must satisfy CMB and BBN constraints that exclude an extra relativistic species with abundance comparable to neutrinos [1]:

$$\Delta N_{\nu}^{\text{eff}} \equiv 1.8 \ 10^5 \int h^2 \frac{d\Omega_{\text{GW}}}{d\ln f} d\ln f \lesssim 0.2 \quad \text{where} \quad \frac{d\Omega_{\text{GW}}}{d\ln f} \equiv \frac{1}{\rho_{\text{cr}}} \frac{d\rho_{\text{GW}}}{d\ln f} \tag{1}$$

and  $h \approx 0.67$ . Future CMB data are expected to improve the sensitivity by one order of magnitude thanks to higher precision. A cosmological detection of some extra radiation would not tell its identity nor its spectrum. More interestingly, future experiments based on graviton to photon conversion in magnetic fields aim at detecting gravitational waves with good spectral resolution around atomic frequencies  $f \sim 10^{15-19}$  Hz. For the moment, experiments planned to search for axions using magnets built for colliders are expected to be sensitive only to  $\Omega_{\rm GW} \sim 1$  [2–20], as illustrated in fig. 1. Opening a new observational window requires reaching sensitivities below the BBN/CMB bound of eq. (1).

Various processes could produce a cosmological background of relic gravitons with  $f \sim T_0 \sim 300 \,\mathrm{GHz}$ , the current CMB temperature. However, most processes produce a graviton abundance significantly lower than that of photons or neutrinos. This scarcity stems from the suppression of graviton couplings  $E/\bar{M}_{\rm Pl}$  at energies E below the reduced Planck mass  $\bar{M}_{\rm Pl} \approx 2.4 \, 10^{18} \,\mathrm{GeV}$ . The minimal contribution, from scatterings among SM particles, produces a gravitational cosmological background with abundance  $\Delta N_{\nu}^{\rm eff} \approx 0.01 T_{\rm RH}/\bar{M}_{\rm Pl} \lesssim 10^{-4.5}$  around microwave frequencies [21], as illustrated by the gray region in fig. 1. This is small because the reheating temperature must be below  $T_{\rm RH} \lesssim 0.003 \bar{M}_{\rm Pl}$  to avoid overproducing inflationary tensor modes [1].

This paper discusses an exception to the typical outcome of a small graviton relic background: some particle S with mass  $M \sim 10^{10-18}$  GeV might have been present during the big-bang and decayed slowly. Gravitons might have *opened* a channel for its decay (as opposed to having a fast decay channel already opened by other particles and adding a small branching ratio  $\sim (M/4\pi \bar{M}_{\rm Pl})^2$  into gravitons with smooth spectrum peaked around micro-waves [22–29]). In view of the assumed slow decay, small Planck-suppressed graviton couplings can lead to a big branching ratio into gravitons, as they only compete with other small couplings. Moreover, during the big bang, a weakly interacting long-lived massive particle acquires an enhanced abundance, because matter gets diluted more slowly than SM radiation. As a result, the resulting graviton abundance  $\Omega_{\rm GW}$  can easily approach the bound in eq. (1).

In section 2 we compute the assumed cosmology and the resulting peculiar spectrum of produced gravitons, peaked at  $f \gg T_0$  in view of the slow decay. In section 3 we apply helicity methods to classify which processes  $S \to gg$ ,  $S \to g SM$ ,  $S' \to Sg$  are allowed by Lorentz invariance and locality, where S, S' are massive particles, and SM denotes any SM particle.



Figure 1: Panoramic plot. The vertical axis is the gravitational wave characteristic strain  $h_c$ , connected to the energy density by  $d\Omega_{\rm GW}/d\ln f = 2\pi^2 f^2 h_c^2/3H_0^2$ . The red region is excluded by the CMB/BBN bound of eq. (1). The continuous curve is the LIGO/VIRGO GW observation. The dashed curves are sensitivities of possible future experiments with the indicated names. The gray region is the maximal cosmic microwave gravitational background. The hatching indicates expected astrophysical backgrounds, including the solar background at optical frequencies [30, 31]. The two green peaks at optical frequencies show an example of optical gravitational waves from decays of the spin 0 and spin 2 glue-balls S and S' in the theory of section 4.1.

In section 4 we discuss specific theories. An hidden sector with strong coupling provides the maximal graviton abundance. Theories with perturbative couplings (in particular, extensions of Einstein gravity) can plausibly contain one or more particles that slowly decay gravitationally producing detectable gravitational waves. Observing a multi-peaked spectrum of optical gravitational waves could allow a spectroscopy of such states, offering an experimental window into theories such as gauge unification, supersymmetry, extra dimensions, string theories, and more. Conclusions are given in section 5.

# 2 A particle decaying into gravitons

We here compute the cosmology during the big-bang at temperature T of a particle S with mass M, number abundance  $n_S$  and slow decay rate  $\Gamma_S = \Gamma_{\rm GW} + \Gamma_{\rm SM} \ll M$  with a significant  ${\rm BR}_{\rm GW} = \Gamma_{\rm GW}/\Gamma_S$  opened by one or more gravitons (as opposed to adding a graviton to an allowed decay into SM particles). We parameterise the S decay width into gravitons as

$$\Gamma_{\rm GW} = \frac{cM}{4\pi} \left(\frac{M}{\bar{M}_{\rm Pl}}\right)^p \tag{2}$$

where c and p are dimension-less constants and  $\bar{M}_{\rm Pl} = M_{\rm Pl}/\sqrt{8\pi} = 2.4 \ 10^{18} \,\text{GeV}$  is the reduced Planck mass. As discussed in section 4, the plausible value that maximise  $\Gamma_{\rm GW}$  seem  $c \sim 1$  and p = 2 (for one graviton production) or p = 4 (for two-graviton production).

The minimal gravitational decay rate into SM particles due to virtual one-graviton exchange or related gravitational operators is expect to be suppressed by 4 powers of the Planck mass,

$$\Gamma_{\rm SM} \gtrsim \frac{g_{\rm SM}M}{4\pi} \left(\frac{M}{\bar{M}_{\rm Pl}}\right)^4$$
(3)

where  $g_{\rm SM} \approx 106.75$  is the number of SM degrees of freedom. Extra particles could exist at energies around M. Of course, extra non-gravitational interactions can enhance  $\Gamma_{\rm SM}$ .

### 2.1 Production of S

We here consider plausible values of the cosmological number density  $n_S$  of S. It is convenient to consider the combination  $Y_S = n_S/s$  not affected by the expansion thanks to the entropy density  $s = 2\pi^2 g_{\rm SM} T^3/45$ . Thermal equilibrium corresponds to  $Y_S \sim 1/g_{\rm SM}$ . A minimal contribution to  $Y_S$  arises from the inverse process SM SM  $\rightarrow S$  with space-time rate density  $\gamma \sim s \Gamma_{\rm SM} e^{-M/T}$ . This inverse decay contributes to the S abundance as:

$$Y_S \equiv \frac{n_S}{s} \sim \max_T \frac{\gamma}{Hs} \sim \Gamma_{\rm SM} \frac{\bar{M}_{\rm Pl}}{M^2} e^{-M/T_{\rm RH}} \sim \left(\frac{M}{\bar{M}_{\rm Pl}}\right)^3 e^{-M/T_{\rm RH}} \tag{4}$$

having assumed eq. (3). The peculiar inverse decay kinematics does not allow to benefit from the possibility that the reheating temperature  $T_{\rm RH}$  after inflation is larger than M. So a larger contribution to the S number density can arise from gravitational SM SM  $\rightarrow SS$  pair production, with rate  $\gamma \sim T^8/\bar{M}_{\rm Pl}^4$  resulting in

$$Y_S \sim 10^{-2-3} \left(\frac{T_{\rm RH}}{\bar{M}_{\rm Pl}}\right)^3 e^{-M/T_{\rm RH}}.$$
 (5)

This contribution can be larger in specific theories with extra interactions. Furthermore, inflationary production (considered in [32]) can contribute to  $Y_S$ . We will keep  $Y_S$  as a free parameter.

### 2.2 Decay of S

We here compute the total gravitational wave abundance  $\rho_{\rm GW}$  produced from S decays. Since the decay is slow, S decays while non-relativistic and out of equilibrium. We assume that its interactions are small enough to be cosmologically negligible. The Hubble rate at temperature T and time t is

$$H^{2} = \frac{\rho_{\rm SM} + \rho_{S}}{3\bar{M}_{\rm Pl}^{2}}, \qquad \rho_{\rm SM} = \frac{\pi^{2}}{30}g_{\rm SM}(T)T^{4}.$$
 (6)

The cosmological evolution during S decays is described by

$$\frac{d\rho_S}{d\ln a} = -3\rho_S - \frac{\Gamma_S}{H}\rho_S, \qquad \frac{d\rho_{\rm SM}}{d\ln a} = -4\rho_{\rm SM} + \frac{\Gamma_{\rm SM}}{H}\rho_S, \qquad \frac{d\rho_{\rm GW}}{d\ln a} = -4\rho_{\rm GW} + \frac{\Gamma_{\rm GW}}{H}\rho_S. \tag{7}$$

The equations for  $\rho_S$  and  $\rho_{GW}$  are solved starting at an initial  $a = a_i$  by

$$\rho_S(a) = \rho_S(a_i) \left(\frac{a_i}{a}\right)^3 e^{-\Gamma_S t}, \qquad \rho_{\rm GW}(a) = \int_{a_i}^a d\ln a' \left(\frac{a'}{a}\right)^4 \frac{\Gamma_{\rm GW} \rho_S}{H}.$$
(8)

Eq.s (7) can be rewritten omitting the 2nd equation and switching as variable from scale factor a to temperature T (defined in terms of  $\rho_{\text{SM}}$  as in eq. (6)), by using (see e.g. [33]):

$$Z = -\frac{d\ln T}{d\ln a} = 1 - \frac{1}{4} \frac{\Gamma_{\rm SM} \rho_S}{H \rho_{\rm SM}}.$$
(9)

So  $T \propto 1/a^Z$ . Initially  $Z \simeq 1$  even when  $\rho_S > \rho_{\rm SM}$ . Next, after  $\rho_S/\rho_{\rm SM} > H/\Gamma_{\rm SM}$ , one has  $Z \simeq 3/8$  i.e.  $T \propto a^{-3/8}$  cools more slowly, due to reheating from S decays. Reheating ends when  $1/t \sim H \sim \Gamma_S$ , reverting to  $Z \simeq 1$ .

#### 2.3 Gravitational wave spectrum

While SM particles (including neutrinos at  $T \gtrsim \text{MeV}$ ) interacted and thermalised, gravitons would have travelled nearly free, such that the relic gravitational wave spectrum today still reflects early cosmology. We write as  $dN/d\ln E$  the differential spectrum of gravitons with energy E produced by each S decay. Assuming that it happens at scale factor a, gravitons have current energy  $E_0 = aE$ . So the current graviton energy density spectrum is

$$\frac{d\rho_{\rm GW}}{d\ln E_0} = E_0 \int \frac{da}{a} a^3 \frac{\rho_S}{M} \frac{\Gamma_{\rm GW}}{H} \frac{dN}{d\ln E}.$$
(10)

For example,  $S \to gg$  decays correspond to  $dN/dE = 2\delta(E - M/2)$ , such that the graviton spectrum is a line convoluted with the exponential distribution of decay times, that gives different redshift:

$$\frac{d\rho_{\rm GW}}{d\ln E_0} = a^4 \rho_S \frac{\Gamma_{\rm GW}}{H} \qquad \text{evaluated at} \qquad a = \frac{2E_0}{M}.$$
(11)

In view of the simplest particle physics (a line), the frequency  $f = E_0/2\pi$  spectrum of gravitational waves directly reflects different epochs of the cosmology during S decays. In our case  $d\Omega_{\rm GW}/d\ln f$  has a peak with order unity width at some frequency  $f_{\rm 0peak}$ .

The gravitational wave spectra can be computed in terms of three effective parameters:

1) the S abundance  $Y_S$ ;



Figure 2: Spectra of gravitational waves produced by a scalar S with mass M and initial abundance  $Y_S$  slowly decaying into two gravitons at the temperature  $T_{\text{peak}} = M/z_{\text{peak}}$  with branching ration into gravitons  $BR_{GW} = 0.01$ . The plot illustrates the full parameter space:  $z_{\text{peak}}$  accounts for both the scalar mass and decay width, and different values of the branching ratio in gravitons simply increase or decrease the overall gravitational wave abundance. The gravitational wave signals lie below the CMB/BBN bound of eq. (1) in all the plotted region. The numbers indicate  $\rho_S/\rho_{SM}$  at peak for each spectrum.

- 2) the S branching ratio into gravitons  $BR_{GW}$ , that acts as a proportionality constant;
- 3) the dimension-less ratio  $z_{\text{peak}} \equiv M/T_{\text{peak}} \gg 1$  between the S mass and the temperature  $T_{\text{peak}}$  that solves  $\Gamma_S \approx H(T_{\text{peak}})$ , at which the S decay dominantly happens.

At this temperature, S can either be a negligible or dominant component of the cosmological energy density, depending on whether the S density during the peak of S decays is dominant or sub-dominant compared to the energy density of the SM bath:

$$\frac{\rho_S}{\rho_{\rm SM}}\Big|_{\rm peak} \sim Y_S z_{\rm peak}.$$
(12)

Numerical spectra are shown in fig. 2 for  $BR_{GW} = 0.01$  and for different values of  $z_{peak}$  and of  $Y_S$ . The main features of the spectra can be understood as follows. Above the peak at  $f \gg f_{0peak}$  the spectrum is proportional to an exponential cut-off  $\exp[-f^{3(1+w)/2}]$ , with w = 0if S dominates and w = 1/3 if SM radiation dominates. Below the peak at  $f \ll f_{0peak}$  the GW spectrum gets suppressed as  $a^4 \rho_S / H \propto a / H$ . To evaluate this factor and compute  $f_{0\text{peak}}$  we need to consider the two limiting cases.

- If S decays while sub-dominant, the low-frequency tail of the gravitational wave spectrum is suppressed by  $a/H \propto a^3 \propto f^3$ . The total  $\rho_{\rm GW}$  will be computed in instantaneous decay approximation in section 2.4.
- If instead S decays as dominant component and reheats the SM plasma, the lower-frequency tail is initially suppressed as  $a/H \propto a^{5/2} \propto f^{5/2}$ , implying a mildly broader spectrum around the peak. The total  $\rho_{\rm GW}$  is approximated in section 2.5.

The half peak is at  $f = \{0.565, 1.51\} f_{0peak}$  if S decays while sub-dominant, and at  $f = \{0.519, 1.56\} f_{0peak}$  if S decays while dominant. If a positive signal will be seen, the peak frequency will tell  $z_{peak} = M/T_{peak}$ , but data alone will not allow to separately infer M and  $T_{peak}$ .

### 2.4 Decay of S as sub-dominant component

Assuming that S decays while  $\rho_{\rm SM} \gg \rho_S$  dominates the energy budget, the decay temperature  $T_{\rm peak}$  is determined by  $\Gamma_S \approx H \approx \sqrt{\rho_{\rm SM}/3}/\bar{M}_{\rm Pl}$ , giving

$$R \equiv \frac{\rho_S}{\rho_{\rm SM}} \approx 0.77 g_{\rm SM}^{1/4} \frac{Y_S M}{\sqrt{\bar{M}_{\rm Pl} \Gamma_S}} \ll 1 \qquad \text{at} \qquad T_{\rm peak} \approx M \frac{4Y_S}{3R}.$$
 (13)

The graviton abundance at peak is

$$\frac{\rho_{\rm GW}}{\rho_{\rm SM}}\Big|_{\rm peak} \approx \frac{{\rm BR}_{\rm GW}\rho_S}{\rho_{\rm SM}} \approx {\rm BR}_{\rm GW}R = {\rm BR}_{\rm GW}\frac{4Y_S}{3}z_{\rm peak}.$$
(14)

The exact solution of eq. (8) can be computed analytically finding that  $\rho_{\rm GW}/\rho_{\rm SM}$  is 1.25 higher than in the instantaneous decay approximation of eq. (14). Gravitons produced with energy E = M/2 and thereby frequency  $f = E/2\pi$  have today (at  $T = T_0 \approx 300 \,\text{GHz}$ ) a frequency redshifted down to

$$f_{0\text{peak}} \simeq \frac{M}{4\pi} \frac{T_0}{T_{\text{peak}}} = \frac{T_0}{4\pi} z_{\text{peak}}.$$
(15)

The gravitational decays of eq. (2) predict  $z_{\text{peak}} \approx 2g_{\text{SM}}^{1/4} (\bar{M}_{\text{Pl}}/M)^{(p-1)/2} / \sqrt{c}$ . This means that a slow decay,  $z_{\text{peak}} \gg 1$ , can enhance  $\rho_{\text{GW}}$  up to its maximal allowed value, and that the same  $z_{\text{peak}}$  increases the graviton frequency  $f_{0\text{peak}}$  above  $T_0$ .

To conclude we convert a generic  $\rho_{\rm GW}/\rho_{\rm SM}$  at a generic high temperature  $T_{\rm peak}$  into  $\Omega_{\rm GW}$  today. Gravitons redshift as radiation,  $\rho_{\rm GW} \propto 1/a^4$ , while the SM bath cools keeping its entropy  $sa^3$  constant with  $g_{s0} = 3.91$  today. So the current graviton energy density is

$$\frac{\rho_{\rm GW}(T_0)}{\rho_{\rm GW}(T)} = \left(\frac{g_{s0}}{g_{\rm SM}}\right)^{4/3} \frac{T_0^4}{T^4}, \qquad \text{i.e.} \qquad \Omega_{\rm GW} = \frac{\rho_{\rm GW}(T_0)}{\rho_{\rm cr}} = 3.6 \ 10^{-5} \left.\frac{\rho_{\rm GW}}{\rho_{\rm SM}}\right|_{\rm peak} \tag{16}$$

having used  $T_0 = 2.725 \text{ K}$ ,  $\rho_{\rm cr} = 3H_0^2 \overline{M}_{\rm Pl}^2$  with  $H_0 = h \, 100 \text{ km/sMpc}$  and  $h \approx 0.67$ .

### 2.5 Decay of S as dominant component

Assuming that S decays while dominating the energy budget, the decay temperature  $T_{\text{decay}}$  is determined by  $\Gamma_S \approx H \approx \sqrt{\rho_S/3}/\bar{M}_{\text{Pl}}$ . We obtain

$$\frac{\rho_S}{\rho_{\rm SM}} \approx R^{4/3} \gg 1$$
 at  $T_{\rm decay} \approx M \frac{4Y_S}{3R^{4/3}}$  (17)

with the same R as in eq. (13). The value of  $T_{\text{decay}}$  is however irrelevant, because S decays later reheat the SM plasma up to  $T_{\text{peak}}$  given by

$$\rho_{\rm SM}(T_{\rm peak}) \equiv \frac{\pi^2}{30} g_{\rm SM}(T_{\rm peak}) T_{\rm peak}^4 = \rho_{\rm SM}(T_{\rm decay}) + BR_{\rm SM} \rho_S(T_{\rm decay}).$$
(18)

So the reheating temperature after S decays is

$$\frac{T_{\text{peak}}}{M} \approx (\text{BR}_{\text{SM}} + R^{-4/3})^{1/4} \frac{4Y_S}{3R}.$$
(19)

It differs from what  $T_{\text{peak}}$  was in the previous case of section 2.4 only because the extra first factor. The single effective parameter  $z_{\text{peak}} = M/T_{\text{peak}} > 1$  controls the current frequency of gravitational waves produced with energy E = M/2 at  $T_{\text{peak}}$ :

$$f_{0\text{peak}} = \frac{M}{4\pi} \frac{T_0}{T_{\text{peak}}} \approx \frac{T_0 z_{\text{peak}}}{4\pi} \tag{20}$$

In the limit of strong S dominance the graviton density at  $T_{\text{peak}}$  is

$$\left. \frac{\rho_{\rm GW}}{\rho_{\rm SM}} \right|_{\rm peak} \simeq \frac{\rm BR_{\rm GW}}{\rm BR_{\rm SM}}.$$
(21)

and the current  $\Omega_{\rm GW}$  is obtained using eq. (16). BBN/CMB data demand  $\rho_{\rm GW}/\rho_{\rm SM} < 0.03$  at peak. This is close to the naive expectation  ${\rm BR}_{\rm GW}/{\rm BR}_{\rm SM} \sim 0.01$  based on assuming a common gravitational rate and counting the number of degrees of freedom is just below current BBN/CMB bounds.

We provided simple analytic expressions for the gravitational wave abundance and spectrum in terms of the effective parameters  $Y_S$ ,  $z_{\text{peak}}$ , BR<sub>GW</sub>. Finally, we assume the decay rates of eq. (2) and eq. (3) for different values of p = 2 (decay into a single-graviton), p = 4 (decay into 2 gravitons), p = 6 (decay into 3 gravitons, or into 2 if the effective operator of eq. (25) is pessimistically suppressed by  $\Lambda = \overline{M}_{\text{Pl}}$ ). Fig. 3 shows the resulting iso-contours of the overall abundance  $\Omega_{\text{GW}}$  of gravitational waves (computed numerically solving eq. (7)), and of the peak frequency  $f_{0\text{peak}}$ . The S-decay mechanism for GW production is so efficient that part of the parameter space (shaded in red) is already excluded. An  $\Omega_{\text{GW}}$  within two orders of magnitude below current bounds arises in the area shaded in green. Gravitationally decaying particles below the Planck scale down to about  $10^{10}$  GeV (depending on p) can lead to detectable



Figure 3: Parameter space for the decay of a particle S with mass M and cosmological abundance  $Y_S = n_S/s$  into into SM particles with rate  $\Gamma_{\rm SM} = g_{\rm SM}M^5/4\pi \bar{M}_{\rm Pl}^4$ , and into  $N_g$  gravitons with tree-level rate  $\Gamma_{\rm GW} = M(M/\bar{M}_{\rm Pl})^{2N_g}/4\pi$  (for  $N_g = 1$  half of the energy goes into SM particles). S decays while dominating the cosmological energy density below the grey dot-dashed line. The continuous curves are iso-contours of the graviton abundance  $\Omega_{\rm GW}$ , with the region excluded by the BBN/CMB bound of eq. (1) shaded in red. Signals are more prominent in the region shaded in green, just below the bound. The dashed horizontal lines are iso-contours of the current peak frequency of gravitons. The dashed green curve is the minimal S abundance, estimated as the sum of eq.s (4) and (5), dominated by the second. The dot-dashed green curve is the S abundance in the glue-ball model of section 4.1. Both are computed for the maximal  $T_{\rm RH} = 0.003\bar{M}_{\rm Pl}$ . The bottom corner of the middle plot is excluded by entropy injection after BBN.

gravitational waves around optical frequencies. For  $p = \{2, 4\}$  even the minimal S abundance, given by the sum of eq. (4) and eq. (5) (dashed green curve), allows for significant gravitational signals, provided that the reheating temperature is large enough. The dot-dashed green curve shows the larger  $Y_S$  arising in the theory of section 4.1. We demand  $T_{\text{peak}} \gtrsim \text{MeV}$  to avoid entropy injection after BBN. (In line of principle the opposite extreme where S is decaying today is allowed if BR<sub>SM</sub> is tiny enough).

# 3 Allowed decays into gravitons

We here consider which  $1 \rightarrow 2$  decays of a massive particle allow for graviton production compatibly with locality and conservation of quantum numbers related to Lorentz invariance. The Landau-Yang theorem forbids the decay of a massive vector into two identical massless vectors. The extension to gravitons of this kind of non-trivial restrictions can be computed with two different methods: effective field theory operators, or on-shell helicity amplitude techniques [34–37]. The second method appears simpler, as it avoids unnecessarily extending the little-group symmetries of the process into the full Lorentz group. The decay amplitude of a massive particle S with spin s and momentum P into two massless particles with helicities  $h_1$ and  $h_2$  and momenta  $p_1$  and  $p_2$  is found by writing the momenta  $P = p_1 + p_2$  as  $p_1^{\mu} = \lambda_1 \sigma^{\mu} \tilde{\lambda}_1$ ,  $p_2^{\mu} = \lambda_2 \sigma^{\mu} \tilde{\lambda}_2$  in terms of two-component spinors  $\lambda^a$ , so that  $\sigma^{\mu}$  are the usual relativistic Pauli matrices. The amplitude can be written as [35]

$$\mathscr{A} \propto (\lambda_1^{s+h_2-h_1} \lambda_2^{s+h_1-h_2}) [\tilde{\lambda}_1^a \epsilon_{ab} \tilde{\lambda}_2^b]^{s+h_1+h_2}, \qquad (22)$$

where the term in squared parentheses is the Lorentz scalar built using the anti-symmetric tensor  $\epsilon_{ab}$ , and the term in round parentheses indicates the completely symmetric spin s product of the number of  $\lambda_1$  and  $\lambda_2$  dictated by their exponents. Locality demands no negative powers of momenta  $\lambda_{1,2}$ , so helicities must satisfy

$$s \ge |h_1 - h_2|. \tag{23}$$

### 3.1 Decay of a massive particle into two gravitons

Inserting the graviton helicities  $h_1, h_2 = \pm 2$  in eq. (22) shows that:

0) A spin 0 particle can decay into two gravitons, that must have the same helicity  $h_1 = h_2$  because of eq. (23). The amplitude of eq. (22) arises from the effective operators<sup>1</sup>

$$\sqrt{|\det g|} S R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} / \Lambda, \qquad \sqrt{|\det g|} S R_{\mu\nu\rho\sigma} \tilde{R}^{\mu\nu\rho\sigma} / \tilde{\Lambda}$$
(24)

where  $\Lambda, \tilde{\Lambda}$  are mass scales and  $\tilde{R}$  is the dual Riemann tensor. The first operator contributes to S decay into two gravitons as  $[37]^2$ 

$$\mathscr{A}(S \to h^{++}h^{++}) = \frac{[\tilde{\lambda}_1 \tilde{\lambda}_2]^4}{2\Lambda \bar{M}_{\rm Pl}^2}, \qquad \Gamma_{\rm GW} = \frac{M^7}{64\pi \Lambda^2 \bar{M}_{\rm Pl}^4}.$$
 (25)

1) The decay amplitude for a spin s = 1 vector into two gravitons vanishes because of the anti-symmetry of the last factor in eq. (22). The only possible decay of a vector into two identical particles is into spin 1/2 fermions.

<sup>&</sup>lt;sup>1</sup>Simpler operators do not contribute for the following reasons. The tadpole operator  $\Lambda^4 \sqrt{|\det g|}S$  gets cancelled at the minimum of the *S* potential. The operator  $\Lambda \sqrt{|\det g|} SR$  does not contribute to *S* decays into gravitons, as it also induces a *S*/graviton kinetic mixing: one needs to diagonalise the action reaching mass eigenstates via a Weyl field redefinition to the Einstein frame, that removes such operator. Effective operators of the form  $\sqrt{|\det g|}SR^2/\Lambda$  or  $\sqrt{|\det g|}SR_{\mu\nu}R^{\mu\nu}/\Lambda$  do not contribute since on-shell gravitons in flat space satisfy  $R_{\mu\nu} = 0$  and thereby R = 0 [37,32]. The first operator of eq. (24) is mediated at one loop by matter fields with coefficient proportional to a  $\beta$  function of 4-derivative gravity. The second operator involving the dual Riemann tensor  $\tilde{R}$  can be mediated by one loop of chiral fermions.

<sup>&</sup>lt;sup>2</sup>Its gravitational wave effects have been studied in [32] in a parameter range with  $H_{\text{infl}} \sim T_{\text{RH}}$  that leads to a large  $f_{0\text{peak}} \sim 10^{23}$  Hz.

2) A spin 2 particle can decay into two same-helicity gravitons  $h_1 = h_2$ . The effective operator is a contraction of  $(\partial_{\mu}\partial_{\nu}S_{\alpha\beta})$  Riemann<sup>2</sup>/ $\Lambda^3$  and

$$\mathscr{A}(S \to h^{++}h^{++}) \sim \frac{[\tilde{\lambda}_1 \tilde{\lambda}_2]^6 \lambda_1^2 \lambda_2^2}{M^2 \Lambda^3 \bar{M}_{\rm Pl}^2}, \qquad \Gamma_{\rm GW} \sim \frac{M^{11}}{4\pi \Lambda^6 \bar{M}_{\rm Pl}^4}.$$
 (26)

- 3) A spin 3 particle cannot decay into two gravitons, similarly to a spin 1 particle.
- 4) A spin 4 particle can decay into two same-helicity gravitons and (via a lower dimensional operator) into opposite-helicity gravitons. We do not consider this possibility.

#### 3.2 Decay of a massive particle into a graviton and a SM particle

A process that produces one graviton g only has a rate suppressed by only two powers of the Planck scale. Given that we are interested in values of M much above the weak scale, we can approximate the SM particle as massless and apply eq. (22), inserting now  $h_1 = \pm 2$  and  $-1 \le h_2 \le 1$ , the possible helicities of a SM particle. The amplitude shows that:

- 0) A massive spin 0 or 1/2 particle cannot decay into a graviton and a SM particle.
- 1) A massive spin 1 vector can decay as  $S \to g\gamma$  where  $\gamma$  denotes an abelian SM vector such that  $h_1 = 2$  and  $h_2 = 1$ . Effective operators such as  $R^{\alpha\beta\mu\nu}F_{\alpha\beta}S_{\mu\nu}/\Lambda^d$  (where the anti-symmetric field  $S_{\mu\nu}$  could have mass dimension d = 2 or 1) lead to

$$\mathscr{A}(S \to h^{++}\gamma^{+}) \sim \frac{[\tilde{\lambda}_1 \tilde{\lambda}_2]^4 \lambda_2^2}{M^{3-d} \Lambda^d \bar{M}_{\rm Pl}}, \qquad \Gamma_{\rm GW} \sim \frac{M^{3+2d}}{4\pi \Lambda^{2d} \bar{M}_{\rm Pl}^2}.$$
 (27)

3/2) A massive spin 3/2 particle can decay into a graviton and a SM fermion  $\psi$  as

$$\mathscr{A}(S \to h\psi) \sim \frac{[\tilde{\lambda}_1 \tilde{\lambda}_2]^4 \lambda_2^3}{M^{5/2} \Lambda \bar{M}_{\rm Pl}}, \qquad \Gamma_{\rm GW} \sim \frac{M^5}{4\pi \Lambda^2 \bar{M}_{\rm Pl}^2}.$$
 (28)

2) A massive spin 2 vector can decay into  $g\gamma$  and into gh, where h is a massless neutral scalar (so the decay amplitude into the Higgs is suppressed by the weak scale v).

In both cases 1) and 2), S can also decay into two SM vectors. If the decay operator is mediated by a loop of particles with mass m and gauge charge g, one expects  $1/\Lambda^2 \sim g/(4\pi m)^2$  and  $\Gamma_{\rm GW}/\Gamma_{\rm SM} \sim (m/g\bar{M}_{\rm Pl})^2 \lesssim 1$  if gravity is the weakest interaction.

### 3.3 Decay of a massive particle into a graviton and a massive particle

Finally, we consider a decay  $S' \to Sg$  where S and S' are two massive particles with spins s and s' [35]. The graviton energy from the decay at rest is  $E = (M'^2 - M^2)/2M'$ . The decay is allowed if  $s + s' \ge 2$  as long as S and S' are either both bosonic or both fermionic. For low spins this reduces to the spin combinations listed in the previous section. At larger spins, decays with |s - s'| > 2 become allowed, altought suppressed in the non-relativistic limit.



Figure 4: Illustrative Feynman diagrams for gravitational glue-ball decays. Blue dots denotes gravitational couplings. SM denotes any Standard Model particle, including gravitons.

### 4 Theories for decay into gravitons

The next key issue is whether in motivated theories the suppression scales  $\Lambda$  of operators for decays into gravitons can have a sub-Planckian value, down to  $\Lambda \gtrsim M$  where the range of validity of the effective operator approximation breaks down. Section 4.1 presents theories based on strong gauge dynamics that achieve  $\Lambda \sim M$  and thereby the maximal gravitational wave abundance. Section 4.2 presents theories with weak couplings  $g \sim 1$ , resulting in  $\Lambda \sim (4\pi)^2 M/g$ a loop factor above M.

#### 4.1 Theories with non-perturbative interactions

Maximal gravitational decays arise in theories with gauge group structure  $G_{\rm SM} \otimes G$ , provided the extra factor G becomes strongly coupled and remains hidden. Hidden means that no sub-Planckian matter fields couple the two sectors at renormalizable level. Concretely, this means that no matter field can be charged under both factors  $G_{\rm SM}$  and G, and no scalar charged under G can have a quartic coupling to the SM Higgs. We will focus our discussion on the simplest possibility: a pure-gauge non-abelian gauge interaction G. This is automatically hidden, as gauge invariance forbids renormalizable interactions with SM particles. We thereby extend the SM adding an extra pure-glue gauge sector G. The action is

$$S = \int d^4x \sqrt{|\det g|} \left[ -\frac{\bar{M}_{\rm Pl}^2}{2} R - \frac{1}{4} G^a_{\mu\nu} G^{\mu\nu a} + \mathscr{L}_{\rm SM} + \mathscr{L}_{\rm NRO} \right]$$
(29)

where  $G^a_{\mu\nu} = \partial_{\mu}G^a_{\nu} - \partial_{\nu}G^a_{\mu} - gf^{abc}G^b_{\mu}G^c_{\nu}$ , g is the gauge coupling, and we left implicit the dark  $\theta$  term. The extra gauge vectors interact with SM particles only via gravitational interactions and via possible non-renormalizable operators  $\mathscr{L}_{\text{NRO}}$  that we assume to be Planck-suppressed, corresponding to the absence of light matter field charged both under G and under the SM. Planck-suppressed operators are unavoidably generated by RG running. Similarly to QCD, the gauge interaction G confines at some scale  $\Lambda$  naturally below the Planck scale. The confinement phase transition is expected to be of 1st order with slow bubbles [38], so it produces a negligible extra amount of gravitons. Being a pure glue theory, confinement forms a set of glue-ball

composite states with masses  $M \sim \Lambda$ . The abundance of glue-balls from pair production of dark massless vectors via gravitational interactions (gravitational freeze-in) followed by confinement is [39]:

- For low masses  $M \leq \bar{M}_{\rm Pl} (T_{\rm RH}/\bar{M}_{\rm Pl})^{15/4}$ , the dark vectors thermalize among themselves before confining into glue-balls with abundance  $Y_S \sim (T_{\rm RH}/\bar{M}_{\rm Pl})^{9/4}$ , neglecting logarithmic corrections from  $SSS \to SS$  scatterings. We omitted  $10^{-2-3}$  factors, that depend on the dimension of the gauge group.
- In the intermediate range  $\bar{M}_{\rm Pl}(T_{\rm RH}/\bar{M}_{\rm Pl})^{15/4} \lesssim M \lesssim T_{\rm RH}^2/\bar{M}_{\rm Pl}$ , the glue-balls self-thermalize. The abundance is similar to previous case.
- For larger masses, the dark vectors immediately hadronise and the glue-balls never reach an equilibrium distribution. The resulting abundance is  $Y_S \sim N(T_{\rm RH}/\bar{M}_{\rm Pl})^3 e^{-M/T_{\rm RH}}$ , where N is an enhancement due to the particle multiplicity in G-vector jets produced by collisions at energy  $T_{\rm RH} \gg \Lambda$  [39].

The lightest glue-ball corresponds to the operator  $\operatorname{Tr} G^2$ : it has spin s = 0 and we denote it as S. Its decay rates into gravitons and SM particles are estimated to be comparable and have the form of eq. (2) for p = 4:

$$\Gamma_{\rm SM} = \Gamma(S \to {\rm SM\,SM}) \sim g_{\rm SM} \frac{M^5}{4\pi \bar{M}_{\rm Pl}^4}, \qquad \Gamma_{\rm GW} = \Gamma(S \to gg) \sim \frac{M^5}{4\pi \bar{M}_{\rm Pl}^4} \tag{30}$$

where  $g_{\rm SM}$  is number of SM degrees of freedom. A precise prediction of the decay rates is prevented by uncertainties due to the non-perturbative nature of the glue-ball state, and by Planck-suppressed operators in  $\mathscr{L}_{\rm NRO}$ , that provide contributions comparable to Einstein gravity. The estimates in eq. (30) are justified as follows.

An Effective Field Theory approach restricts the coupling of the lightest glue-ball S to *soft* gravitons. The dominant operator would be the tadpole  $\sqrt{|\det g|}S$ , with coefficient computable from the covariantization of the matrix element  $\langle 0|\operatorname{Tr} G^2|S\rangle$ , known from lattice computations [40]. However, as discussed in section 3.1, the tadpole vanishes at the minimum and does not lead to a gravitational decay. Going to higher orders, the dominant effect is given by eq. (25) with expected suppression scale  $\Lambda \sim M$ , leading to the rates in eq. (30).

However, higher order operators are relevant and the EFT expansion breaks down, as S decays into hard gravitons with energy E = M/2. Their production probes the glue-ball inner structure and thereby depends on unknown non-perturbative form factors. Going beyond the EFT limit, the S decay rate is estimated as follows. If a non-relativistic description of S as a bound state with wave function  $\psi_S$  were applicable, the glue-ball gravitational decay would be computable as a gravitational collision of dark gluons, as illustrated by the Feynman diagrams in fig. 4. Given that non-perturbative couplings lead to relativistic constituents, we approximate the collision speed as the speed of light, obtaining

$$\Gamma(S \to gg, \text{SM SM}) \approx |\psi_S|^2 \sigma(GG \to gg, \text{SM SM}) \approx \frac{M^5}{4\pi \bar{M}_{\text{Pl}}^4}.$$
 (31)

Indeed the squared wave-function  $\psi_S$  of the bound state is roughly given by its inverse volume,  $|\psi_S|^2 \sim M^3$ , and the gravitational cross-section of two dark gluons G into two gravitons or two SM particles is  $\sigma(GG \leftrightarrow gg, \text{SM SM}) \sim s/4\pi \bar{M}_{\text{Pl}}^4$ . Graviton-mediated cross sections into SM particles are computed e.g. in [39]. No extra suppression is expected, as the decay of a massive spin 2 particle into two gravitons is allowed by symmetries as discussed in section 3.1. Non-renormalizable operators, such as  $|H|^2 G^a_{\mu\nu} G^{\mu\nu a} / \bar{M}_{\text{Pl}}^2$ , can contribute at comparable level. In conclusion, the expected amount of gravitational waves from glue-ball decays is shown by the middle panel of fig. 3. The dot-dashed green curve shows the expected glue-ball abundance.

Each group G predicts a specific glue-ball spectrum that contains extra heavier glue-balls. Those with mass  $\geq 2M$  can decay via gauge interactions into SS, negligibly contributing to gravitational waves. Some models and computations predict at least one exception: a resonance S' with spin s' = 2 and mass  $M' \approx 1.5M$  [43]. Its lightness would imply that  $S' \rightarrow SS$  is kinematically blocked, so that these two lighter glue-balls are co-stable. In such a case S' can decay emitting one graviton g as  $S' \rightarrow Sg$  (allowed according to section 3.3), with decay rate estimated as

$$\Gamma'_{\rm GW} = \Gamma(S' \to Sg) \sim \frac{M'^3}{4\pi \bar{M}_{\rm Pl}^2} \left(1 - \frac{M^2}{M'^2}\right).$$
 (32)

It is suppressed by only p = 2 powers of  $\overline{M}_{\text{Pl}}$  because one graviton only is emitted (a S' with spin 1 would instead decay as  $S' \to Sgg$ ). This kind of earlier S' decays produce extra gravitational waves at lower frequencies. The resulting spectrum gravitational waves can feature multiple peaks, observable in part of the parameter space. An example is shown in fig. 1: the faster  $S' \to Sg$  decay with p = 2 gives gravitational waves peaked around  $\sim 10^{13}$  Hz. These get mildly suppressed by the later  $S \to gg$ , SM SM decay with p = 4 that gives GW peaked around  $10^{15}$  Hz, about  $\overline{M}_{\text{Pl}}/M$  higher.

Some pure gauge theories predict extra special very long lived glue-balls S'', proposed in [39] as gravitational dark matter candidates. For example, the group  $G = \mathrm{SU}(N)$  predicts C-odd GGG glue-balls with slow gravitational decay rate  $\Gamma_{S''} \sim M^9/\bar{M}_{\mathrm{Pl}}^8$  arising from Planck-suppressed operators. The group  $G = \mathrm{SO}(N)$  with even N predicts glue-balls odd under an  $\mathrm{SO}(N)$  parity with decay rate  $\Gamma_{S''} \sim M(M/\bar{M}_{\mathrm{Pl}})^{2N-4}$ . Such longer-lived glue-balls match the dark matter density if their abundance is  $Y_{S''} \sim \mathrm{eV}/M''$ , much below the  $Y_{S,S'}$  that lead to gravitational waves at observable level. Naive hadronization models suggest  $Y_{S''}/Y_{S,S'} \sim 0.76^N$  [39]. Some other groups don't predict longer-lived glue-balls.

### 4.2 Theories with perturbative interactions

The previous discussion suggests that observing gravitational waves around optical frequencies might allow to discover 'gravitational particles', with mass a few orders of magnitude below the Planck scale, that slowly decayed with a non-negligible branching ratio into gravitons. We conclude discussing which theories contain plausible candidates for such states.



Figure 5: Feynman diagrams for gravitational decays of a Higgs-like particle S.

We start considering the pessimistic case where operators such as eq. (24) arise at loop level, from the first three Feynman diagrams in fig. 5. To compute the coefficient we recall that the coefficient  $\gamma$  of the Riemann squared operator

$$\sqrt{|\det g|}\gamma R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} \tag{33}$$

at one loop level runs as [44, 45, 32]

$$\frac{\partial \gamma}{\partial \ln \mu} = \beta_{\gamma}^{\text{gravity}} + \beta_{\gamma}^{\text{matter}} \quad \text{with} \quad \beta_{\gamma}^{\text{matter}} = \frac{52N_V - 7N_F - 4N_S}{720(4\pi)^2} \tag{34}$$

in the presence of  $N_S$  real scalars,  $N_F$  Weyl fermions,  $N_V$  vectors, generically denoted as 'matter'. Indeed, the operator of eq. (24) arises when S couples as a Higgs to these matter particles, giving them S-dependent masses that act as an IR cut-off in the RG running of  $\gamma$ . The maximal operator coefficient arises when the masses of these particle arise solely from the vacuum expectation value of S as  $m_F = y\langle S \rangle$ ,  $m_S = \sqrt{\lambda} \langle S \rangle$ ,  $m_V = g \langle S \rangle$ , where y generically denotes Yukawa couplings, g denotes gauge couplings and  $\lambda$  quartic couplings. In this case the operator arises with coefficient [32]

$$\frac{1}{\Lambda} = \frac{\beta_{\gamma}^{\text{mat}}}{\langle S \rangle} \sim \frac{g}{\left(4\pi\right)^2 m_V} + \cdots .$$
(35)

The  $S R_{\mu\nu\rho\sigma} \tilde{R}^{\mu\nu\rho\sigma}/\tilde{\Lambda}$  operator arises from one loop exchange of fermions with Yukawa couplings to a scalar S that affects the phase of the fermion mass. The operator coefficient can be extracted from the gravitational chiral anomaly, as a rephasing  $F_i \to e^{ic_i S/\langle S \rangle} F_i$  of Weyl spinors  $F_i$  can remove the coupling to S, while generating the operator with coefficient [46,32]

$$\frac{1}{\tilde{\Lambda}} = \frac{\sum_{i} c_{i}}{24(4\pi)^{2} \langle S \rangle}.$$
(36)

Eq.s (35) and (36) mean that the pure gauge theory of section 4.1 where  $g \sim (4\pi)^2$  and  $m_V \sim M$ provides the maximal S decay rate into gravitons. If instead  $1/\Lambda$  is loop suppressed, the decay width into gravitons  $\Gamma_{\rm GW}$  gets suppressed by  $(M/\Lambda)^2 \sim g^2/(4\pi)^4$ . The amount of gravitational waves can remain large, despite that  $\Gamma_{\rm GW}$  is now loop suppressed, provided that other decay channels such as  $\Gamma_{\rm SM}$  are suppressed down to a similar level. A first danger is that the Scouplings to matter can also allow for S decays into pairs of matter particles at tree level, that would suppress the S branching ratio into gravitons. Such decays are kinematically blocked if  $m_{S,F,V} \geq M/2$ . The condition is automatically satisfied in the minimal theory where S itself is the only matter particle in the loop. In such a case, the  $S \rightarrow gg$  decay arises provided that the S potential has a cubic  $S^3$  term when expanded around the minimum. The second danger is decays into SM particles, that are much lighter than M. Then, the possibility of a small  $\Gamma_{\rm SM}$ depends on whether the S, F, V matter in the loop directly interacts with SM particles.

Let us consider grand unified theories, where a unification gauge group breaks to the SM gauge group via the Higgs mechanism at mildly sub-Planckian energies  $\gtrsim 10^{15}$  GeV, as demanded by proton decay bounds. Candidates for gravitationally-decaying scalars S are the SM singlets in the 24 of SU(5), in the 16 of SO(10), in the 27 of  $E_6$ . Indeed unification theories predict, in addition to SM particles, extra heavy vectors and scalars (and possibly fermions) that receive masses from S and that thereby mediate at one loop level the operator in eq. (24). These heavy GUT particles have gauge-like couplings  $g_{\text{GUT}} \sim 1$  to S, but also to the SM particles. SM particles are not directly coupled to S, because they do not acquire GUT-scale masses. If  $m_{S,F,V} \geq M$  the decays of S into SM particles are partially suppressed, by a 4-body phase space or by a 1-loop factor. The resulting partially small  $\Gamma_{\text{SM}} \sim g^6 M/(4\pi)^5$  leads to an amount of gravitational waves that falls between the CMB/BBN bound  $\Omega_{\text{GW}} \sim 10^{-6}$  and the solar background  $\Omega_{\text{GW}} \sim 10^{-15}$ , depending on M. An Higgs-like scalar lighter than the particles to which it gives mass can arise e.g. if the unification symmetry is broken dynamically via the Coleman-Weinberg mechanism.

Better candidates for gravitationally-decaying particles can be found in extensions of Einstein gravity. Theories with (discrete or continuous) extra dimensions predict a (finite or infinite) number of massive Kaluza-Klein excitations of gravitons  $g_n$  as well as extra particles. These include extra scalars: the extra dimensional 'radion' components of the graviton  $g_{MN}$ , and extra 'moduli'.<sup>3</sup> The zero modes of such fields control the size and shape of the extra dimensions. Thereby they act as geometric Higgs bosons, affecting the masses of matter fields. Eq. (35) means that the  $S \to gg$  operator of eq. (24) arises at the compactification scale.

The lower Kaluza-Klein excitations  $g_n$  of the graviton  $g_0$  have spin 2, so they could decay into gravitons such as

$$g_1 \to g_0 g_0, \qquad g_2 \to g_1 g_0. \tag{37}$$

This kind of decays can happen under two conditions. First, the extra dimension must be not

 $<sup>^{3}</sup>$ A different phenomenon, production of gravitational waves from stochastic inhomogeneities in theories with extra dimensions, was considered in [47, 48].

translationally invariant, to avoid "KK number" conservation. Second, the higher dimensional action gravitational action cannot have the minimal Einstein form (curvature plus vacuum energy), and must instead contain 4-derivative Riemann<sup>2</sup> terms (again unavoidably generated by one loop corrections). These non-minimal gravitational terms are necessary because the effective operators identified in section 3 all involve 4 derivatives.<sup>4</sup>

Warping is one plausible source of breaking of extra-dimensional translational invariance. However, according to the AdS/CFT duality, Kaluza Klein graviton excitations in a warped geometry can be re-interpreted as spin 2 glue-balls of some conformal field theory. More general holographic techniques can similarly approximate the spectrum of scalar and vector glueballs [43] arising from 4-dimensional strong dynamics, similar to the theories of section 4.1.

A related possibility is that the SM fields are confined around some value of the extradimensional coordinates, thereby approximatively living along a 3+1 dimensional brane in extra dimensions. In such a case, the 4-dimensional effective field theory also contains a set of massive 'branon' scalars, describing the position of the brane in the extra dimensions. The branon potential energy can contain a self cubic term, leading to decays into gravitons at one loop level.

Given that decays into gravitons need Riemann squared operators, an interesting possibility is the most general action in 3+1 dimensions containing terms with positive mass dimension only, as it gives a renormalizable theory of quantum gravity [50]. This is known as 4-derivative gravity because  $R^2$  and  $R^2_{\mu\nu}$  terms in the action imply a graviton kinetic term with 4 derivatives. As a result the theory contains the usual graviton, a possibly problematic spin 2 ghost and the spin 0 gravi-scalar also present in Starobinsky  $R^2$  inflation. Such extra states could decay into gravitons. However these decay rates vanish at tree level [51], because the Gauss-Bonnet term is topological in 3+1 dimensions. A non-vanishing rate is expected at loop level. 4derivative theories predict a different big source of gravitational waves: UV divergences are made renormalizable at the price of enhancing IR divergences, such that energies above the ghost mass are released via graviton bremsstrahlung [52].

More in general, the Planck mass might be dynamically induced by the vacuum expectation value of a scalar S, dubbed *Planckion*. As it acts as the Higgs for gravity, one expects  $S \rightarrow gg$  similar to eq. (35), mediated at loop level by a purely gravitational loop (4th diagram in fig. 5). This loop is computable in renormalizable 4-derivative gravity that predicts  $\beta_{\gamma}^{\text{gravity}} = -413/180(4\pi)^2$  [44, 45].

<sup>&</sup>lt;sup>4</sup>Furthermore, if the gravitational action only contains minimal Einstein terms and vacuum energy terms (possibly including background contributions localised along sub-spaces such as branes and orbifold fixed points), graviton KK modes are described by eigenfunctions of the Laplacian, orthogonal when weighted by the background metric. The cubic couplings among different KK graviton modes  $n_{1,2,3}$  are proportional to wave-function overlaps  $I_{n_1n_2n_3}$ . We live in a special compactification, with energies tuned such that the 4-dimensional vacuum energy nearly vanishes (this might require a warping or negative-energy non-dynamical objects). Then the graviton 0 mode  $g_0$  is massless and its wave-function is constant, when weighted by the metric [49]. This implies the vanishing of some overlap integrals, such as  $I_{100} \propto I_{10} = 0$  and  $I_{210} \propto I_{21} = 0$ .

If supersymmetry exists and is broken around the Planck scale, the minimal N = 1 gravitational super-multiplet contains the massless graviton, a massive spin 3/2 gravitino and two scalars (remnants of the chiral super-multiplet 'eaten' by the massless gravitational supermultiplet), such that the number of bosonic and fermionic degrees of freedom match. The gravitino does not decay into a graviton and a SM fermion. Theories where supersymmetry is broken in an hidden sector are often considered to avoid problematic spectra of super-particles. Then, the two extra scalars can be gravitationally decaying particles.

Strings  $X_M(\tau, \sigma)$  in 10 dimensions have been proposed as theories of quantum gravity [53]. A huge number of possible compactifications can lead to the gravitationally-decaying particles typical of extra dimensions, but hugely reduce the string predictivity. For example one of the many possibilities is that the theories of section 4.1 with a maximal gravitational wave abundance could arise with hidden group  $G = E_8$  from the heterotic string [53]. To avoid getting lost in a plethora of possibilities, lets us focus one the most characteristic particles arising in the QFT limit of strings. These are fields with the quantum numbers of  $X_M X_N$ , that decompose into the graviton  $g_{MN}$  (symmetric two-index gauge field) accompanied by a scalar and by the Kalb-Ramond field  $B_{MN}$  (anti-symmetric two-index gauge field). It generalises the photon coupling to particle currents into coupling to string currents. A variety of sources can make it massive [53, 54]. Then, its 4-dimensional component is dual to a pseudo-vector  $S_{\mu\nu}$ , with suppressed couplings to matter that include axion-like couplings, magnetic-like couplings to fermions, and interactions with vectors and gravitons. The KR field could decay into a graviton and a SM vector as in eq. (27). The higher-dimensional components of  $B_{MN}$  are dual to pseudo-scalars, that can decay in two gravitons as in eq. (25). Other stringy candidates for gravitationally-decaying scalars are moduli. In particular, chiral fermions and CP violation in 3+1 dimension can arise in string theories from compactification on extra dimension with complex structure. Toroidal-like compactifications can lead to a QFT with a stringy  $SL(2,\mathbb{Z})$ symmetry [53]. Its modulus  $\tau$  is equivalent to a combination of scalars that spontaneously break CP. Thereby  $\tau$  affects the phase of fermion masses, and is expected to acquire a gravitational decay width as in eq. (36), plus a Planck-suppressed width into SM particles.

Trying to be more general than strings, graviton amplitudes with good properties seem to need heavy states along one or more *Regge trajectories*, see e.g. [55, 56]. This kind of states could decay gravitationally, be sub-Planckian and maybe generalise the string excitations.

Alternatively, these attempts could just be an approximation to quantum gravity analogous to how Regge trajectories approximate QCD. The analogy suggests that a deeper quantum gravity theory might feature gravitons and space-time emerging out of (unknown) fermionic dynamics. Such an hypothetical quantum gravity theory could feature, around the Planck scale, massive spin 3/2 'excitations' of SM fermions, that could decay into a graviton and a SM fermion as discussed in section 3.2.

Finally, a signal could arise from hypothetical primordial black holes with mass M mildly above the Planck scale. These would have evaporated with rate  $g_{\rm SM}\pi \bar{M}_{\rm Pl}^4/80M^3$  and branching ratio BR<sub>GW</sub>  $\approx 1/g_{\rm SM} \sim 0.01$  into gravitons, with a different spectrum [57]. In conclusion, while dedicated computations can be performed, the current discussion suffices to show that plausible candidates for mildly sub-Planckian states with substantial gravitational decays seem easily found in plausible theories.

### 5 Conclusions

We proposed theories where relic gravitational waves around optical frequencies can have densities mildly below the CMB/BBN bound on extra radiation,  $\Omega_{\rm GW} \lesssim 10^{-6}$ , despite that gravitons interact less than photons. This can happen if some long-lived particle S has a decay channel *opened* by gravitons, and thereby proceeding with a Planck-suppressed rate. The cosmology is computed in section 2, finding that a particle that decays at a temperature  $T_{\rm peak}$  much lower than its mass M gives a specific graviton spectrum, today peaked around optical frequencies  $\sim T_0 M/T_{\rm peak}$ . Simple analytic approximations are provided. The detailed shape of the gravitational wave peak depends on whether S decays as a sub-dominant component, or while dominating the cosmological abundance. In the latter case, a decay with branching ratio into gravitons given by the inverse of the number of SM degrees of freedom (as naively expected in various theories) implies a gravitational wave abundance just below current CMB/BBN bounds.

- Section 3.1 shows that a particle with spin s = 0 or 2 can decay into two gravitons, with rate suppressed by  $p \ge 4$  powers of the Planck mass. The middle panel of fig. 3 shows that gravitational wave signals can be observable even if the S abundance is low and/or M is mildly sub-Planckian.
- Section 3.2 shows that a particle with spin s = 1 or 2 can decay into a graviton and a SM particle with rate suppressed by  $p \ge 2$  powers of the Planck mass. The left panel of fig. 3 shows that gravitational wave signals can be observable provided that the S abundance is mildly small.

Section 4 presents plausible theories where such decays generate gravitational waves with observable abundance. The maximal abundance arises in theories with a hidden extra gauge interaction that becomes non-perturbative and confines at some scale  $\Lambda$ . A simple case is an extra pure gauge sector (thereby automatically hidden) that forms a spectrum of glue-ball states with mass  $M \sim \Lambda$ . Such states decay gravitationally into a characteristic gravitational wave spectrum with multiple peaks, as exemplified in fig. 1.

Moving from theories at strong coupling to perturbative SM-like couplings  $g \sim 1$ , the decay rate into gravitons gets loop suppressed. The amount of gravitational waves remains detectable provided that the decay rate into SM particles too is suppressed. Plausible candidates for gravitationally decaying particles are found, under favourable but plausible conditions, in a variety of plausible theories mildly below the Planck scale. As discussed in section 4.2, these include gauge unification, extra dimensions, supersymmetry, strings.

These theories were considered before the Large Hadron Collider, that however found no new physics around the weak scale. Thereby they remain in a limbo: these new physics plausibly lies around Planckian energies, many orders of magnitude above the energies where physics is an experimental discipline. Optical gravitational waves might offer a new window and maybe even some spectroscopical information on gravitationally decaying particles mildly below the Planck scale. This possibility appears as interesting as colliders. Detecting optical gravitational waves would need experiments able of reaching sensitivity to  $\Omega_{\rm GW} \sim 10^{-6}$ , and improving possibly down to the ultimate solar astrophysical background,  $\Omega_{\rm GW} \sim 10^{-15}$  [30,31]. Longo lo cammino ma grande la meta [58].

Acknowledgements We thank Gian Giudice, Andrew Long, Michele Redi, Alberto Salvio, Riccardo Torre, Michael Zantedeschi for discussions, and Anish Ghoshal who collaborated in an early phase. We do not thank Artificial Intelligences, because all most advanced systems gave the wrong answer to: "can a spin 0 particle decay into two gravitons?". A.S. thanks Duccio Pappadopulo for dedicated prompts (that allow to get also the correct answer, with roughy 50% probability) and Riccardo Rattazzi, who correctly answered to the prompt. G.L. is supported by the Generalitat Valenciana APOSTD/2023 grant CIAPOS/2022/193.

### References

- See e.g. PLANCK Collaboration, 'Planck 2018 results. VI. Cosmological parameters', Astron. Astrophys. 641 (2020) A6 [arXiv:1807.06209].
- [2] R. Ballantini et al., 'Microwave apparatus for gravitational waves observation' [arXiv:grqc/0502054].
- [3] N. Aggarwal et al., 'Challenges and opportunities of gravitational-wave searches at MHz to GHz frequencies', Living Rev.Rel. 24 (2021) 4 [arXiv:2011.12414].
- [4] A. Arvanitaki, A.A. Geraci, 'Detecting highfrequency gravitational waves with opticallylevitated sensors', Phys.Rev.Lett. 110 (2013) 071105 [arXiv:1207.5320].
- [5] A. Ejlli, D. Ejlli, A.M. Cruise, G. Pisano, H. Grote, 'Upper limits on the amplitude of ultra-high-frequency gravitational waves from graviton to photon conversion', Eur.Phys.J.C 79 (2019) 1032 [arXiv:1908.00232].
- [6] N. Aggarwal, G.P. Winstone, M. Teo, M. Baryakhtar, S.L. Larson, V. Kalogera, A.A. Geraci, 'Searching for New Physics with a Levitated-Sensor-Based Gravitational-Wave Detector', Phys.Rev.Lett. 128 (2022) 111101 [arXiv:2010.13157].

- [7] LSD Collaboration, 'Optical Trapping of High-Aspect-Ratio NaYF Hexagonal Prisms for kHz-MHz Gravitational Wave Detectors', Phys.Rev.Lett. 129 (2022) 053604 [arXiv:2204.10843].
- [8] A. Berlin et al., 'Detecting high-frequency gravitational waves with microwave cavities', Phys.Rev.D 105 (2022) 116011 [arXiv:2112.11465].
- [9] M. Goryachev, W.M. Campbell, I.S. Heng, S. Galliou, E.N. Ivanov, M.E. Tobar, 'Rare Events Detected with a Bulk Acoustic Wave High Frequency Gravitational Wave Antenna', Phys.Rev.Lett. 127 (2021) 071102 [arXiv:2102.05859].
- [10] W.M. Campbell, M. Goryachev, M.E. Tobar, 'The multi-mode acoustic gravitational wave experiment: MAGE', Sci.Rep. 14 (2024) 4762 [arXiv:2307.00715].
- [11] F. Sorge, 'High-Frequency Gravitational Waves in Electromagnetic Waveguides', Annalen Phys. 535 (2023) 2300228.
- [12] G. Tobar, S.K. Manikandan, T. Beitel, I. Pikovski, 'Detecting single gravitons with quantum sensing' [arXiv:2308.15440].
- [13] V. Domcke, C. Garcia-Cely, N.L. Rodd, 'Novel Search for High-Frequency Gravita-

tional Waves with Low-Mass Axion Haloscopes', Phys.Rev.Lett. 129 (2022) 041101 [arXiv:2202.00695].

- [14] T. Bringmann, V. Domcke, E. Fuchs, J. Kopp, 'High-frequency gravitational wave detection via optical frequency modulation', Phys.Rev.D 108 (2023) L061303 [arXiv:2304.10579].
- [15] G. Vacalis, G. Marocco, J. Bamber, R. Bingham, G. Gregori, 'Detection of high-frequency gravitational waves using high-energy pulsed lasers', Class.Quant.Grav. 40 (2023) 155006 [arXiv:2301.08163].
- [16] T. Liu, J. Ren, C. Zhang, 'Limits on High-Frequency Gravitational Waves in Planetary Magnetospheres', Phys.Rev.Lett. 132 (2024) 13 [arXiv:2305.01832].
- [17] A. Ito, T. Ikeda, K. Miuchi, J. Soda, 'Probing GHz gravitational waves with graviton-magnon resonance', Eur.Phys.J.C 80 (2020) 179 [arXiv:1903.04843].
- [18] A. Ito, J. Soda, 'Exploring high-frequency gravitational waves with magnons', Eur.Phys.J.C 83 (2023) 766 [arXiv:2212.04094].
- [19] A. Ito, R. Kitano, 'Macroscopic Quantum Response to Gravitational Waves' [arXiv:2309.02992].
- [20] N. Herman, L. Lehoucq, A. Fűzfa, 'Electromagnetic antennas for the resonant detection of the stochastic gravitational wave background', Phys.Rev.D 108 (2023) 124009 [arXiv:2203.15668].
- [21] J. Ghiglieri, G. Jackson, M. Laine, Y. Zhu, 'Gravitational wave background from Standard Model physics: Complete leading order', JHEP 07 (2020) 092 [arXiv:2004.11392].
- [22] K. Nakayama, Y. Tang, 'Stochastic Gravitational Waves from Particle Origin', Phys.Lett.B 788 (2019) 341 [arXiv:1810.04975].
- [23] A. Ghoshal, R. Samanta, G. White, 'Bremsstrahlung high-frequency gravitational wave signatures of high-scale nonthermal leptogenesis', Phys.Rev.D 108 (2023) 035019 [arXiv:2211.10433].

- [24] S. Kanemura, K. Kaneta, 'Gravitational waves from particle decays during reheating', Phys.Lett.B 855 (2024) 138807 [arXiv:2310.12023].
- [25] N. Bernal, S. Cléry, Y. Mambrini, Y. Xu, 'Probing reheating with graviton bremsstrahlung', JCAP 01 (2024) 065 [arXiv:2311.12694].
- [26] A. Tokareva, 'Gravitational waves from inflaton decay and bremsstrahlung', Phys.Lett.B 853 (2024) 138695 [arXiv:2312.16691].
- [27] W. Hu, K. Nakayama, V. Takhistov, Y. Tang, 'Gravitational wave probe of Planckscale physics after inflation', Phys.Lett.B 856 (2024) 138958 [arXiv:2403.13882].
- [28] Y. Xu, 'Ultra-high frequency gravitational waves from scattering, Bremsstrahlung and decay during reheating', JHEP 10 (2024) 174 [arXiv:2407.03256].
- [29] A. Datta, A. Sil, 'Probing Leptogenesis through Gravitational Waves' [arXiv:2410.01900].
- [30] G.S. Bisnovatyi-Kogan, V.N. Rudenko, 'Very high frequency gravitational wave background in the universe', Class.Quant.Grav. 21 (2004) 3347 [arXiv:gr-qc/0406089].
- [31] C. García-Cely, A. Ringwald, 'Complete Gravitational-Wave Spectrum of the Sun' [arXiv:2407.18297].
- [32] Y. Ema, K. Mukaida, K. Nakayama, 'Scalar field couplings to quadratic curvature and decay into gravitons', JHEP 05 (2022) 087 [arXiv:2112.12774].
- [33] G.F. Giudice, A. Notari, M. Raidal, A. Riotto, A. Strumia, 'Towards a complete theory of thermal leptogenesis in the SM and MSSM', Nucl.Phys.B 685 (2004) 89 [arXiv:hepph/0310123].
- [34] E. Conde, E. Joung, K. Mkrtchyan, 'Spinor-Helicity Three-Point Amplitudes from Local Cubic Interactions', JHEP 08 (2016) 040 [arXiv:1605.07402].
- [35] N. Arkani-Hamed, T.-C. Huang, Y.-. Huang, 'Scattering amplitudes for all masses and spins', JHEP 11 (2021) 070 [arXiv:1709.04891].

- [36] S.Y. Choi, J.H. Jeong, 'Selection rules for the decay of a particle into two identical massless particles of any spin', Phys.Rev.D 103 (2021) 096013 [arXiv:2102.11440].
- [37] A. Alonzo-Artiles, A. Avilez-López, J.L. Díaz-Cruz, B.O. Larios-López, 'The Higgs-Graviton Couplings: from Amplitudes to the Action' [arXiv:2105.11684].
- [38] For a summary and references see e.g. C. Gross et al., 'Dark Matter as dark dwarfs and other macroscopic objects: multiverse relics?', JHEP 09 (2021) 033 [arXiv:2105.02840].
- [39] C. Gross, S. Karamitsos, G. Landini, A. Strumia, 'Gravitational Vector Dark Matter', JHEP 03 (2021) 174 [arXiv:2012.12087].
- [40] Y. Chen et al., 'Glueball spectrum and matrix elements on anisotropic lattices', Phys.Rev.D 73 (2006) 014516 [arXiv:hep-lat/0510074].
- [41] H.B. Meyer, 'Glueball matrix elements: A Lattice calculation and applications', JHEP 01 (2009) 071 [arXiv:0808.3151].
- [42] A. Mitridate, M. Redi, J. Smirnov, A. Strumia, 'Dark Matter as a weakly coupled Dark Baryon', JHEP 10 (2017) 210 [arXiv:1707.05380].
- [43] A. Dymarsky, D. Melnikov, 'Spectrum of large N glueballs: holography vs lattice', JHEP 11 (2022) 164 [arXiv:2206.14826].
- [44] I.G. Avramidi, 'Covariant methods for the calculation of the effective action in quantum field theory and investigation of higher derivative quantum gravity' [arXiv:hepth/9510140].
- [45] N. Ohta, R. Percacci, 'Higher Derivative Gravity and Asymptotic Safety in Diverse Dimensions', Class.Quant.Grav. 31 (2014) 015024 [arXiv:1308.3398].
- [46] K. Fujikawa, 'Path Integral for Gauge Theories with Fermions', Phys.Rev.D 21 (1980) 2848.

- [47] D. Andriot, G. Lucena Gómez, 'Signatures of extra dimensions in gravitational waves', JCAP 06 (2017) 048 [arXiv:1704.07392].
- [48] O-K. Kwon, S. Lee, D.D. Tolla, 'Gravitational Waves as a Probe of the Extra Dimension', Phys.Rev.D 100 (2019) 084050 [arXiv:1906.11652].
- [49] G.B. De Luca, N. De Ponti, A. Mondino, A. Tomasiello, 'Harmonic functions and gravity localization', JHEP 09 (2023) 127 [arXiv:2306.05456].
- [50] K.S. Stelle, 'Renormalization of Higher Derivative Quantum Gravity', Phys.Rev.D 16 (1977) 953.
- [51] J.F. Donoghue, G. Menezes, 'Gauge Assisted Quadratic Gravity: A Framework for UV Complete Quantum Gravity', Phys.Rev.D 97 (2018) 126005 [arXiv:1804.04980].
- [52] A. Salvio, A. Strumia, H. Veermäe, 'New infrared enhancements in 4-derivative gravity', Eur.Phys.J.C 78 (2018) 842 [arXiv:1808.07883].
- [53] See e.g. the book L.E. Ibañez and A.M. Uranga, 'String theory and particle physics' (2012).
- [54] C. Capanelli, L. Jenks, E.W. Kolb, E. Mc-Donough, 'Cosmological implications of Kalb-Ramond-like particles', JHEP 06 (2024) 075 [arXiv:2309.02485].
- [55] N. Arkani-Hamed, C. Cheung, C. Figueiredo, G.N. Remmen, 'Multiparticle Factorization and the Rigidity of String Theory', Phys.Rev.Lett. 132 (2024) 091601 [arXiv:2312.07652].
- [56] J. Albert, W. Knop, L. Rastelli, 'Where is treelevel string theory?' [arXiv:2406.12959].
- [57] A. Ireland, S. Profumo, J. Scharnhorst, 'Primordial gravitational waves from black hole evaporation in standard and nonstandard cosmologies', Phys.Rev.D 107 (2023) 104021 [arXiv:2302.10188].
- [58] B. da Norcia and F. Zenone, 'Brancaleone alle crociate' (1970).