# More varieties of 4-d gauge theories: product representations

## Ben Gripaios<sup>a</sup> and Khoi Le Nguyen Nguyen<sup>b</sup>

<sup>a</sup>Cavendish Laboratory, University of Cambridge, J.J. Thomson Avenue, Cambridge, CB3 0HE, United Kingdom

<sup>b</sup>DAMTP, University of Cambridge, Wilberforce Road, Cambridge, CB3 0WA, United Kingdom

E-mail: gripaios@hep.phy.cam.ac.uk, k1518@cam.ac.uk

ABSTRACT: Recently, we used methods of arithmetic geometry to study the anomalyfree irreducible representations of an arbitrary gauge Lie algebra. Here we generalize to the case of products of irreducible representations, where it is again possible to give a complete description. A key result is that the projective variety corresponding to m-fold product representations of the Lie algebra  $\mathfrak{su}_n$  is a rational variety for every m and n. We study the simplest case of  $\mathfrak{su}_3$  (corresponding to the strong interaction) in detail. We also describe the implications of a number-theoretic conjecture of Manin (and related theorems) for the number of chiral representations of bounded size B (measured roughly by the Dynkin labels) compared to non-chiral ones, giving a precise meaning to the sense in which the former, which are those most relevant for phenomenology, are rare compared to the latter. As examples, we show that, for both irreducible representations of  $\mathfrak{su}_5$ and once-reducible product representations of  $\mathfrak{su}_3$  that are non-anomalous, the number of chiral representations is asymptotically between  $B(\log B)^5$  and  $B^{\frac{4}{3}}$ , while the number of non-chiral representations is asymptotically  $B^2$ .

## Contents

1	Introduction	1
<b>2</b>	Generalities of product representations	3
3	Binary products of $\mathfrak{su}(3)$	5
4	More products of $\mathfrak{su}(n)$	12
5	The distribution of bounded rational points	13

### 1 Introduction

For a gauge theory to be consistent up to high energies, anomalies must cancel, putting non-trivial constraints on the allowed fermionic matter content. The idea of applying ideas of arithmetic geometry to the problem of cancellation of anomalies in four-dimensional gauge theories that are local (*i.e.* those which depend only on the datum of the Lie algebra of the gauge theory rather than its Lie group) was introduced in [1], where it was used to study arbitrary representations of the simplest Lie algebra, namely  $\mathfrak{u}(1)$ . Recently, it was used [2] to study the simplest representations, namely the irreducible ones, of an arbitrary Lie algebra.<sup>1</sup>

These two cases are complementary, in that the only example which is common to both is the trivial representation of  $\mathfrak{u}(1)$  (since any other irreducible representation of an abelian Lie algebra is necessarily anomalous). Given that, we are emboldened to go further and ask whether it can be applied to other cases, with reducible representations of some Lie algebra other than  $\mathfrak{u}(1)$ .

Here we study a case that is a generalization of [2], namely where we have an arbitrary Lie algebra  $\mathfrak{g}$ , but we allow the fermions to transform in a representation that is an *m*fold (tensor) product of irreducible representations (henceforth: irreps), for some *m*. This case turns out to be a relatively straightforward generalization of the m = 1 case, namely irreps of  $\mathfrak{g}$ . We show that, just as for the m = 1 case, we obtain a projective variety over the field of rational numbers  $\mathbb{Q}$  that is, moreover, a rational variety (meaning that it is birationally equivalent to projective space). This makes it possible to not only merely 'solve' the equations (which means, since there are usually infinitely many solutions, to give a parameterization of them, which had already been achieved for the  $\mathfrak{u}(1)$  case in [5]) but to give a detailed geometric picture of them as well.

To get a more concrete idea of what we mean by this, the reader might glance ahead to Fig. 2, which shows the rational points corresponding to anomaly-free representations for

<sup>&</sup>lt;sup>1</sup>For an application to the Standard Model, see [3], and for an application to 2-d gauge theories, see [4].

an example we discuss in exhaustive detail, namely the case of binary products (so m = 2) of irreps of the Lie algebra  $\mathfrak{su}(3)$ .<sup>2</sup> This case has some relevance for phenomenology, since  $\mathfrak{su}(3)$  describes the strong interaction. In Nature, we currently observe only a non-chiral representation of  $\mathfrak{su}(3)$ , given by the six flavours of quarks and their antiquarks, and there are no chiral anomaly-free irreps. But as we will see, there are plenty of binary (and higher) products of irreps that are both chiral and anomaly-free, and which could therefore play a rôle in physics beyond the Standard Model.

The geometric approach also allows us to gain some further insights that are relevant for phenomenology. As we already observed for the case of irreps in [2], chiral anomaly-free representations (which are needed to build natural theories with light fermions) are infinite in number (in cases where they exist at all) and even dominate the non-chiral representations in a specific geometric sense (namely, the chiral ones are dense, in either the usual euclidean topology or the Zariski topology, in the underlying real variety, while the nonchiral ones are not). Nevertheless, if we organize the representations in size (as measured by, *e.g.*, their dimension or their largest Dynkin label), we find that chiral anomaly-free representations tend to be very large compared to non-chiral ones. To give one example, the smallest anomaly-free chiral irrep of  $\mathfrak{su}(5)$  has dimension 1357 824, but there are 35 non-chiral irreps of smaller dimension. To give another, the smallest anomaly-free chiral once-reducible rep of  $\mathfrak{su}(3)$  has dimension 4312, but there are 41 non-chiral once-reducible representations of smaller dimension.

This motivates studying the question of counting chiral vs. non-chiral solutions of bounded size. As it turns out, this too can be turned into a question that can be attacked (if not yet completely answered) geometrically. As we shall see, an easy argument shows that the number of non-chiral solutions grows asymptotically as  $B^2$  (where *B* is some measure of the size of the Dynkin labels to be defined later), while a conjecture of Manin [7] here boils down to the claim that the number of chiral solutions grows as  $B(\log B)^{\rho-1}$ , where  $\rho$  is the rank of the Picard group of the corresponding variety. Whilst Manin's conjecture has yet to be verified for any of the varieties we consider, there are a number of relevant bounds. For example, for both irreps of  $\mathfrak{su}(5)$  and binary products of irreps of  $\mathfrak{su}(3)$ , we can carry over existing results from number theory [8] to show that the number of chiral solutions is bounded above by  $B^{4/3}$ . (In both cases, Manin's conjectured number has also been shown [9] to be a lower bound.) So once we put a bound on the size of representations, as presumably we should when building physics models, we see that chiral anomaly-free representations, even when they are dense, are few-and-far-between.

The outline is as follows. In Section 2 we show that for products of irreps of an arbitrary  $\mathfrak{g}$ , the constraints imposed by anomaly cancellation define a projective variety (more specifically a cubic hypersurface) and that this variety is rational. In Section 3 we discuss the specific case of binary products of irreps of  $\mathfrak{su}(3)$  in detail. In Section 4 we give a selection of results for other cases. In the final Section we discuss the distribution of solutions via Manin's conjecture and related partial theorems.

 $<sup>^{2}</sup>$ This variety has been studied before in mathematics under the name of the *equianharmonic cubic* surface [6].

#### 2 Generalities of product representations

The case of anomaly-free irreps of an arbitrary gauge Lie algebra  $\mathfrak{g}$  was studied in [2]; we now wish to generalize to representations that are products of irreps. Given a compact Lie algebra  $\mathfrak{g}$  and a representation  $\rho$  thereof, the anomaly cancellation equations read

$$\mathrm{tr}\rho = \mathrm{tr}\rho^3 = 0. \tag{2.1}$$

As in [2], in solving these we are helped by a number of serendipitous facts.

The first of these is that every product of irreps of an abelian algebra is one-dimensional and that every such representation is anomalous unless it is the trivial representation. As a result, we need consider only the semisimple summand of  $\mathfrak{g}$ .

The second of these is that for such algebras  $tr\rho = 0$  holds automatically and that  $tr\rho^3$  decomposes into a sum of contributions from each of its simple summands.

The third is that  $\operatorname{tr} \rho^3 = 0$  holds automatically for all simple Lie algebras except  $\mathfrak{su}(n)$ , with  $n \geq 3$ .

The fourth is that in such cases, we can express  $tr\rho^3$  for an irrep in a particularly simple form. Namely, we can replace the usual Dynkin labels  $(m_1, \ldots, m_{n-1})$  (whose entries are nonnegative integers) with either the n-1 positive integers  $(q_1, \ldots, q_{n-1}) :=$  $(m_1 + 1, \ldots, m_{n-1} + 1)$  or the *n* integers

$$\sigma_i := -\sum_{k=1}^{i-1} kq_k + \sum_{k=i}^{n-1} (n-k)q_k, \qquad (2.2)$$

satisfying the constraint

$$\sum_{i=1}^{n} \sigma_i = 0 \tag{2.3}$$

along with

$$\sigma_i - \sigma_{i+1} = nq_i \in n\mathbb{Z}^+, i \in \{1, \dots, n-1\}.$$
(2.4)

Doing so, the anomaly takes the form [10, 11],

$$A(\rho) = \frac{2}{n^2(n^2 - 1)(n^2 - 4)} D(\rho) \sum_{i=1}^n (\sigma_i)^3,$$
(2.5)

where the dimension is given by

$$D(\rho) = \prod_{j=1}^{n-1} \left[ \frac{1}{j!} \prod_{k=j}^{n-1} \left( \sum_{i=k-j+1}^{k} q_i \right) \right] = \frac{\prod_{j< k}^{n} (\sigma_j - \sigma_k)}{n^{n(n-1)/2} \prod_{i=1}^{n-1} i!}.$$
 (2.6)

These are homogeneous polynomials (in either set of variables  $\{q_i\}$  or  $\{\sigma_j\}$ , but not in the Dynkin labels).

The fifth is that the usual formulæ for the anomaly and dimension of a product of representations, i.e.

$$D(\rho_1 \otimes \rho_2) = D(\rho_1) D(\rho_2), \qquad (2.7)$$

$$A(\rho_{1} \otimes \rho_{2}) = A(\rho_{1}) D(\rho_{2}) + D(\rho_{1}) A(\rho_{2}), \qquad (2.8)$$

which generalize to an m-fold product in an obvious way, show that each term in the formula for a product of m irreps contains a common factor of the product of the dimensions of each of the irreps. Since this factor cannot vanish (the dimension of an irrep is positive-definite), our problem reduces to solving a single equation of rather low degree, namely the cubic

$$\sum_{\alpha=1}^{m} \sum_{i=1}^{n} (\sigma_{i\alpha})^3 = 0, \qquad (2.9)$$

where the additional subscript  $\alpha$  indicates that the *n* values  $\sigma_{1\alpha}, \ldots, \sigma_{n\alpha}$  are those characterising the  $\alpha$ -th irrep factor ( $\alpha \in \{1, \ldots, m\}$ ) of the product, along with the *m* linear constraints

$$\sum_{i=1}^{n} \sigma_{i\alpha} = 0. \tag{2.10}$$

Here, the q-tuples of each irrep,  $(q_{1\alpha}, \ldots, q_{n-1,\alpha}) := (m_{1\alpha} + 1, \ldots, m_{n-1,\alpha} + 1)$  satisfy the following relations for  $i \in \{1, \ldots, n-1\}$  and  $\alpha \in \{1, \ldots, m\}$ :

$$\sigma_{i\alpha} = -\sum_{k=1}^{i-1} kq_{k\alpha} + \sum_{k=i}^{n-1} (n-k)q_{k\alpha}, \qquad (2.11)$$

$$\sigma_{i\alpha} - \sigma_{i+1,\alpha} = nq_{i\alpha} \in n\mathbb{Z}^+.$$
(2.12)

The sixth is that Eqs. (2.9) and (2.10) are homogeneous, so not only is solving them over the ring of integers equivalent (via clearing denominators) to solving them over the field of rational numbers, but also we can (try to) do so by studying the projective variety that they define.<sup>3</sup>

The seventh is that, because Eq. (2.10) are linear, this projective variety is a fairly simple one, namely a hypersurface (whose dimension is m(n-1) - 2). Because of our passage to projective geometry, a point on this variety corresponds to an equivalence class containing infinitely many representations, whose  $q_i$  or  $\sigma_j$  (but not the Dynkin labels  $m_i$ ) are all proportional to one another.

The eighth is that this cubic hypersurface features a number of projective linear subspaces (of dimension (m(n-1)-2)/2 for even m(n-1) and (m(n-1)-1)/2 for odd m(n-1)). The presence of these linear subspaces is hardly serendipitous in itself, since they correspond precisely to the non-chiral reps (which are automatically anomaly-free). But their presence makes it easy to prove not only that the variety is rational (which, in layman's terms, means that we can solve this cubic equation), but also to construct, using the method of secants, an explicit parameterization of at least a dense subset (in the Zariski topology) of the solutions.

The ninth is that our variety has a large degree of symmetry. Indeed, the equations Eqs. (2.9) and (2.10) that define it show immediately that its automorphisms contain

$$\sigma_{i\alpha} - \sigma_{i+1,\alpha} \in n\mathbb{Z}^+, i \in \{1, \dots, n-1\}, \alpha \in \{1, \dots, m\}$$

$$(2.13)$$

and will do so later on.

<sup>&</sup>lt;sup>3</sup>According to Eq. (2.12), it is not quite sufficient to solve Eqs. (2.9) and (2.10) over the integers; we must also take into account the restrictions

a subgroup isomorphic to the semi-direct product  $\underbrace{(S_n \times S_n \times \cdots \times S_n)}_{m \text{ factors}} \rtimes S_m$ , where the action of each  $S_n$  factor is to permute the coordinate  $\sigma_{i\alpha}$  for each  $\alpha \in \{1, \ldots, m\}$  and

action of each  $S_n$  factor is to permute the coordinate  $\sigma_{i\alpha}$  for each  $\alpha \in \{1, \ldots, m\}$  and the action of the  $S_m$  factor is to permute the tuples  $\sigma_{\alpha} := (\sigma_{1\alpha}, \ldots, \sigma_{n\alpha})$ . This large symmetry has at least two benefits. One is that we can exploit it to produce an explicit parameterization of every solution (not just a dense subset of solutions as above). The other is that we can use it to turn integer solutions of Eqs. (2.9) and (2.10) that do not satisfy Eq. (2.12) into ones that do, simply by permuting them, provided that for every  $\alpha$ , no two  $\sigma_{i\alpha}$  are equal, and scaling all the  $\sigma_{\alpha}$  by an appropriate positive integer if necessary. This results in a very efficient computational algorithm for finding anomaly-free products of irreps.

To begin describing all of this more explicitly, let us establish some notation. It will occasionally be convenient to work over the complex or real numbers as well as over the rationals, so we let  $\mathbb{k} \in \{\mathbb{C}, \mathbb{R}, \mathbb{Q}\}$  denote such a field. A point in the projective space  $\mathbb{k}P^{nm-1}$  has homogeneous co-ordinates  $[\sigma_{11}:\cdots:\sigma_{n1}:\sigma_{12}:\cdots:\sigma_{n2}:\cdots:\sigma_{1m}:\cdots:\sigma_{nm}]$  and is thus given by an equivalence class of  $(\sigma_{11},\ldots,\sigma_{nm}) \in \mathbb{k}^{nm}$ , where the equivalence relation is given by  $(\sigma_{11},\ldots,\sigma_{nm}) \sim (\lambda\sigma_{11},\ldots,\lambda\sigma_{nm})$  for any non-zero  $\lambda \in \mathbb{k}$ . The zero locus of Eqs. (2.9) and (2.10) defines a projective variety  $V_{n,\otimes m}$  in  $\mathbb{k}P^{nm-1}$ .

To deal with the restriction imposed by Eq. (2.12), we define the following four regions<sup>4</sup> on this variety:

- 1. The unorderable region, for which  $\exists i_1 \neq i_2 \in \{1, \ldots, n\}$  such that  $\sigma_{i_1\alpha} = \sigma_{i_2\alpha}$  for some  $\alpha \in \{1, \ldots, m\}$ ;
- 2. Its complement, the *orderable* region;
- 3. For  $\mathbb{k} \neq \mathbb{C}$ , the *weakly-ordered* region, for which either  $\sigma_{1\alpha} \geq \sigma_{2\alpha} \geq \cdots \geq \sigma_{n\alpha}$  or  $\sigma_{1\alpha} \leq \sigma_{2\alpha} \leq \cdots \leq \sigma_{n\alpha}$  for all  $\alpha \in \{1, \ldots, m\}$ ;
- 4. For  $\mathbb{k} \neq \mathbb{C}$ , the *ordered* region, in which we replace the weak inequalities in the previous definition by strict ones.

Before stating the results that are valid for general n and m, we first consider the simplest non-trivial case, which is that of products of two irreps of  $\mathfrak{su}(3)$ . Here our projective variety describes a surface in three dimensions, allowing us to draw pictures (at least over  $\mathbb{R}$ ).

#### **3** Binary products of $\mathfrak{su}(3)$

Since  $\mathfrak{su}(2)$  has no anomalous representations, the simplest non-trivial case is  $\mathfrak{su}(3)$ , which of course describes the strong interaction in Nature. There, we currently only observe a non-chiral representation of  $\mathfrak{su}(3)$ , made up of the six flavour of quarks and their antiquarks. But it is of interest in going beyond the Standard Model to ask whether we might in the

<sup>&</sup>lt;sup>4</sup>These regions correspond to (semi-)algebraic sets, a notion which (for varieties over  $\mathbb{C}$ ) has become important elsewhere in physics in the study of scattering amplitudes via so-called 'positive geometry'.

future be able to observe fermions carrying chiral representations. For m = 1, it is easy to see that there are no chiral irreps, because the equation describing the variety reduces to

$$\sigma_{11}\sigma_{21}(\sigma_{11} + \sigma_{21}) = 0, \tag{3.1}$$

so we have exactly 3 rational points. Of these, only the point  $\sigma_{21} = 0$  is in the ordered region and its equivalence class consists of irreps of the form  $(q_{11}, q_{11})$ , which are precisely the non-chiral irreps of  $\mathfrak{su}(3)$ . We will now see however that we can find chiral anomaly-free representations for m = 2 (and, *ergo*, for all  $m \ge 2$ ).

For m = 2, the variety  $V_{3,\otimes 2}$  can be described by the single equation

$$\sigma_{11}\sigma_{21}(\sigma_{11} + \sigma_{21}) + \sigma_{12}\sigma_{22}(\sigma_{12} + \sigma_{22}) = 0.$$
(3.2)

This describes a smooth cubic surface, so contains exactly 27 lines over  $\mathbb{k} = \mathbb{C}$ . Nine of these are given by

$$\sigma_{i1} + \sigma_{j1} = \sigma_{k1} = \sigma_{a2} + \sigma_{b2} = \sigma_{c2} = 0, \tag{3.3}$$

with  $\{i, j, k\} = \{a, b, c\} = \{1, 2, 3\}$ , and all of them are rational (that is, they also define lines in the variety over  $\mathbb{Q}$ ). The remaining eighteen are given by

$$\sigma_{i1} + \omega \sigma_{a2} = \sigma_{j1} + \omega \sigma_{b2} = \sigma_{k1} + \omega \sigma_{c2} = \sigma_{i1} + \sigma_{j1} + \sigma_{k1} = 0, \qquad (3.4)$$

again with  $\{i, j, k\} = \{a, b, c\} = \{1, 2, 3\}$ , where  $\omega$  is a cube root of 1. Only the six lines with  $\omega = 1$  are real and rational. Thus, there are 9 + 6 = 15 rational lines on  $V_{3,\otimes 2}$ , 12 of which can be seen on the affine patch  $\sigma_{31} + \sigma_{32} \neq 0$  that we use for our Figures.<sup>5</sup> On this affine patch, we define the coordinates

$$x_1 = -\frac{\sigma_{11}}{\sigma_{31} + \sigma_{32}}, \qquad x_2 = -\frac{\sigma_{21}}{\sigma_{31} + \sigma_{32}}, \qquad x_3 = -\frac{\sigma_{12}}{\sigma_{31} + \sigma_{32}}, \qquad (3.5)$$

such that  $\sigma_{22} = -(\sigma_{31} + \sigma_{32})(1 - x_1 - x_2 - x_3)$  and that  $V_{3,\otimes 2}$  is described by

$$x_1x_2(x_1+x_2) + x_3(1-x_1-x_2-x_3)(1-x_1-x_2) = 0.$$
(3.6)

Fig. 1 shows the variety, along with its rational lines. Two of these lines have special relevance for physics. Namely, the lines  $\sigma_{11} + \sigma_{31} = \sigma_{21} = \sigma_{12} + \sigma_{32} = \sigma_{22} = 0$  and  $\sigma_{11} + \sigma_{32} = \sigma_{21} + \sigma_{22} = \sigma_{31} + \sigma_{12} = \sigma_{11} + \sigma_{21} + \sigma_{31} = 0$ , which are coloured orange and green respectively in Fig. 1, intersect the ordered region, where they correspond to representations. The ordered rational points on the first line correspond to products of two non-chiral irreps, while those on the second line correspond to products of an irrep with its dual. Both classes of products are trivially anomaly-free and overall non-chiral; indeed, they comprise all non-chiral solutions to the anomaly cancellation for n = 3, m = 2. Evidently, these points are dense in the underlying real region, which is diffeomorphic to a union of two copies of  $\mathbb{R}^1$ . We shall soon see, however, that the chiral anomaly-free representations are dense in the two-dimensional region given by the complement of this

 $<sup>{}^{5}</sup>$ We chose this affine patch because it contains the ordered region.



**Figure 1.** The variety  $V_{3,\otimes 2}$  in the affine patch  $\sigma_{31} + \sigma_{32} \neq 0$ , along with its 12 rational lines. The ordered region is shaded in red. An ordered rational point on the orange line corresponds to a product of two non-chiral irreps, while one on the green line corresponds to the product of an irrep with its dual. The rational points contained in each of these real regions are dense in them, in the euclidean topology. The coordinates  $x_1$ ,  $x_2$  and  $x_3$  are defined in Eq. (3.5).

region in the ordered region, shown in red in the Figure. In this sense, the non-chiral binary products are overwhelmed by the chiral ones.

Just as in the case with the variety  $V_{5,\otimes 1}$  studied in [2], there are disjoint (*i.e.* skew) rational lines on  $V_{3,\otimes 2}$  (for example  $\sigma_{11} + \sigma_{21} = \sigma_{31} = \sigma_{12} + \sigma_{22} = \sigma_{32} = 0$  and  $\sigma_{11} = \sigma_{21} + \sigma_{31} = \sigma_{12} = \sigma_{22} + \sigma_{32} = 0$ ), and so the variety  $V_{3,\otimes 2}$  is rational, and we can use the method of secants described in [1, 2] to generate at least a dense subset of the rational points on it.

In brief, the method of secants can be described as follows. Given any two distinct points on a cubic, the line through those points must either intersect the cubic at a third point (which may coincide with one of the other two points), or must lie on the cubic. So by taking a pair of disjoint lines, we can construct a rational map, from  $\mathbb{k}P^1 \times \mathbb{k}P^1$  to the cubic, whose value is given by this third point. Since every point in the underlying projective space that is not on the disjoint lines (*ergo* every such point on the cubic) lies on exactly one line between the disjoint lines, it follows that one can construct an inverse to this map, exhibiting it as a birational map. (We refrain from giving an explicit description of the map here, since it is rather messy. The reader can easily construct it for themself, in analogy with the construction in [2].) In fact, it is easy to see that the only points we miss are those on the disjoint lines themselves, or those on any third line that intersects both lines. These additional points are easily found, since we have already found all of the rational lines on  $V_{3,\otimes 2}$ .

Just as for the case of  $V_{5,\otimes 1}$  studied in [2], it also follows from this construction that the ordered rational points on  $V_{3,\otimes 2}$  are dense (not just in the Zariski topology but also in the euclidean topology) in the ordered real points; the analogous statement also holds for the orderable points.

Fig. 2 shows the rational points produced in this way by a scan over the parameters, while Fig. 3 shows a close-up of the ordered region.

Of the rational points produced by the method of secants in this way, we must reject the unorderable ones, since we cannot use them to construct representations. We will now show that (just as for  $V_{5,\otimes 1}$  in [2]), there are many fewer such points than expected. In particular, unlike the ordered and the orderable points, the unorderable rational points are not dense (in either topology) in the unorderable real points, but rather are finite in number. To see this, we observe that such points have  $\sigma_{i\alpha} = \sigma_{j\alpha}$  for any  $\alpha \in \{1, 2\}$  and  $i \neq j \in \{1, 2, 3\}$ . Thus, they are given by the  $(S_3 \times S_3) \rtimes \mathbb{Z}/2$  orbit of the curve

$$2(\sigma_{11})^3 + \sigma_{12}\sigma_{22}(\sigma_{12} + \sigma_{22}) = 0, \qquad (3.7)$$

where  $\sigma_{21} = \sigma_{11}$  and  $\sigma_{31} = -2\sigma_{11}$ . The part of this curve with  $\sigma_{11} = \sigma_{21} \geq \sigma_{31}$  and  $\sigma_{12} \geq \sigma_{22} \geq \sigma_{32}$  belongs to the boundary of the weakly-ordered region on  $V_{3,\otimes 2}$ . It is not hard to see that  $[\sigma_{11} : \cdots : \sigma_{32}] = [0 : 0 : 0 : 1 : 0 : -1]$  is a rational point on the curve. Therefore, we have a smooth projective plane cubic equipped with a rational point, which defines an elliptic curve C. As with the case of  $V_{5,\otimes 1}$ , we can calculate the Mordell-Weil group of the k-rational points on C. On the one hand, over  $k = \mathbb{R}$ ,  $C(\mathbb{R}) \cong U(1)$  because the cubic form has negative determinant. In other words, the  $2 \times {}^{3}C_{2} = 6$  elliptic curves over  $\mathbb{R}$  in the  $S_{3} \times S_{3} \rtimes \mathbb{Z}/2$  orbit making up the real unorderable region each have one connected component containing infinitely many points. On the other hand, over  $k = \mathbb{Q}$  we can use the change of coordinates

$$X = 2\sigma_{11}, Y = \sigma_{12} - \sigma_{22}, Z = \sigma_{12} + \sigma_{22}, (3.8)$$

to show that C is projectively equivalent to a Mordell elliptic curve of the form

$$y^2 = x^3 + 1, (3.9)$$



**Figure 2**. Rational points on the variety  $V_{3,\otimes 2}$  in the affine patch  $\sigma_{31} + \sigma_{32} \neq 0$  produced by a parameter scan. The red points are in the ordered region and the blue points are in the orderable region, so yield anomaly-free representations after applying a suitable automorphism. The black points are the unorderable points which cannot be used to obtain such representations. The green and orange lines are described in the caption of Fig. 1, and the coordinates  $x_1, x_2$  and  $x_3$  are defined in Eq. (3.5).



Figure 3. A close-up of the ordered region in Fig. 2.

whose Mordell-Weil group is isomorphic to  $\mathbb{Z}/6$  [12, 13]; explicitly, the rational points on this curve are given by

$$C(\mathbb{Q}) = \{ [0:0:0:1:-1:0], [1:1:-2:2:-1:-1], [0:0:0:1:0:-1], [-1:-1:2:1:1:-2], [0:0:0:0:1:-1], [1:1:-2:-1:2:-1] \}, (3.10)$$

where the elements are ordered as the powers of a generator of  $\mathbb{Z}/6$ , beginning with the identity. Of each of these six points, three (the even powers of the generator) lie on three elliptic curves (for example, [0:0:0:1:-1:0] satisfies  $\sigma_{11} = \sigma_{21} = \sigma_{31}$ , while the other three (the odd powers) lie on two elliptic curves (for example, [1:1:-2:2:-1:-1] has  $\sigma_{11} = \sigma_{21}$  and  $\sigma_{22} = \sigma_{32}$ ). Thus, there are only  $6 \times 3/3 + 6 \times 3/2 = 15$  unorderable



Figure 4. The six elliptic curves making up the unorderable region on  $V_{3,\otimes 2}$ . Each curve has six rational points and there are 15 distinct rational points, of which eight are visible on this affine patch.

Q-rational points on  $V_{3,\otimes 2}$ , eight of which are visible on the affine patch  $\sigma_{31} + \sigma_{32} \neq 0$ . As promised, we see that the unorderable rational points are not dense in the unorderable real points, implying that our algorithm for finding representations is highly efficient. The curves and their rational points are shown in Fig. 4.

A plot of the ordered real points on this affine patch of  $V_{3,\otimes 2}$  seems to suggest that, just as with the case for  $V_5$ , they form a connected region that is diffeomorphic to  $\mathbb{R}^2$ , though we do not prove it here.

#### 4 More products of $\mathfrak{su}(n)$

By way of generalizing the results for n = 3, m = 2 to arbitrary n and m, we will prove the following:

- 1. The variety  $V_{n,\otimes m}$  is rational over  $\mathbb{Q}$ ,  $\mathbb{R}$  and  $\mathbb{C}$  (so we can parameterize the rational points);
- 2. The (ordered, respectively orderable) rational points in  $V_{n,\otimes m}$  are dense, in either the Zariski topology or the euclidean topology, in the (ordered, respectively orderable) real points;
- 3. The ordered rational points that correspond to overall non-chiral products are dense (in the euclidean topology) in a union of subspaces, each of dimension m(n-1)/2-1 for odd n or at most mn/2 1 for even n. Thus the chiral anomaly-free products overwhelm the non-chiral ones.

We show that the variety  $V_{n,\otimes m}$  (over  $\mathbb{k} \in \{\mathbb{Q}, \mathbb{R}, \mathbb{C}\}$ ) is rational by generalizing the method of secants described for n = 3, m = 2 in the previous Section. The *m* homogeneous linear constraints (Eq. (2.10)) and the cubic constraint (Eq. (2.9)) imply that the variety  $V_{n,\otimes m}$ can also be viewed as a cubic hypersurface in  $\mathbb{k}P^{m(n-1)-1}$  defined by

$$\sum_{j=1}^{m} \left[ \sum_{i=1}^{n-1} (\sigma_{ij})^3 - \left( \sum_{i=1}^{n-1} \sigma_{ij} \right)^3 \right] = 0.$$
(4.1)

The analogue of the proof in [2] now implies that the method of secants generalizes as follows: instead of picking a pair of disjoint rational lines as in the n = 3, m = 2 case, we now need to find a pair of disjoint linear subspaces of  $\Bbbk P^{m(n-1)-1}$  of dimensions  $d_1 = d_2 :=$ (m(n-1)-2)/2 if m(n-1) is even and  $d_1 := (m(n-1)-1)/2$ ,  $d_2 := d_1 - 1$  if m(n-1) is odd, such that the method of secants, including the step of permuting coordinates within each irrep factor to get ordered points from orderable ones, works as before. Note that m(n-1) is even if n is odd and vice versa, so these requirements do indeed reduce to those found in [2] for the case m = 1. To get a pair of disjoint linear subspaces for every n and m, we can assign to each of the m irrep factors a pair of disjoint linear subspaces of the appropriate dimensions for the case m = 1; examples of the latter can be found in [1, 2]. To make this concrete, consider the example of n = 5, m = 2, where the homogeneous coordinates  $[\sigma_{11}: \cdots : \sigma_{51}: \sigma_{12}: \cdots : \sigma_{52}]$  on two subspaces can be parameterized as

$$\Gamma^{1} = [k_{1}: k_{2}: 0: -k_{2}: -k_{1}: l_{1}: l_{2}: 0: -l_{2}: -l_{1}], \qquad (4.2)$$

$$\Gamma^2 = [0: m_1: -m_1: m_2: -m_2: 0: n_1: -n_1: n_2: -n_2].$$
(4.3)

Thus, the proof in [2] generalizes for arbitrary n and m: we always have a rational variety, where the birational map is given by the method of secants construction as described earlier.

Since the method of secants, which gives a manifestly continuous map with respect to either the Zariski topology or the usual euclidean topology on the real variety, works for our cubic hypersurfaces with arbitrary n and m, it follows that on all of these hypersurfaces, the (ordered, respectively orderable) rational points are dense in the (ordered, respectively orderable) real points in either topology. At this point, we recall that for the case n = 3, m = 2, we observed from a plot of an affine patch of  $V_{3,\otimes 2}$  that the ordered region on it appears to be diffeomorphic to  $\mathbb{R}^2$ . Based on this, and based on what we obtained for  $V_{n,\otimes 1}$ in [2], we conjecture that, in general, the ordered real region on  $V_{n,\otimes m}$  is contractible and is of dimension m(n-1) - 2.

Assuming this conjecture holds, we can compare the density of chiral vs. non-chiral points. To do so, suppose that we have a partition  $m = m_1 + 2m_2$ , where  $m_1$  and  $m_2$  are both non-negative integers, such that we form a non-chiral m-fold product representation of  $\mathfrak{su}_n$  from  $m_1$  non-chiral irreps and  $m_2$  pairs of dual chiral irreps. If n is odd, such a representation can be seen to lie on a rational linear subspace of dimension  $m_1(n-1)/2 + m_2(n-1) - 1 = m(n-1)/2 - 1$ , while if n is even, it lies on a rational linear subspace of dimension  $m_1n/2 + m_2(n-1) - 1 = mn/2 - m_2 - 1 \le mn/2 - 1$ . Thus, we see that indeed the chiral products always overwhelm the non-chiral ones for m = 1 and  $n \ge 5$ , or  $m \ge 2$  and  $n \ge 3$ .<sup>6</sup>

We conclude this Section by listing in Table 1 some low-dimensional anomaly-free products for various n and m that we obtained by the method of secants. The representations are labelled by variables  $q_{i\alpha}$  related to  $\sigma_{i\alpha}$  by Eq. (2.12). An implementation of the method of secants to obtain the results for n = 3, m = 2 as an example is included in an ancillary Mathematica notebook.

As one can see, even going from m = 1 to m = 2 allows us to decrease the dimension of chiral anomaly-free representations by a factor of roughly 1000, bringing us much closer to the realm that is palatable for the purposes of phenomenology.

## 5 The distribution of bounded rational points

In this paper and its prequel [2], we have seen that when anomaly-free chiral irreps (or products thereof) exist, they overwhelm the non-chiral ones, in the sense that they are dense (in the usual topology) in a real space of higher dimension. But we have also seen that such reps tend to have very large dimension. As such, they do not seem to be of much use for phenomenology.

In fact, a related observation has already been made in a much more general context in diophantine geometry, as we now describe. Given a rational point x on a variety V over  $\mathbb{Q}$ , we define its height H(x) by  $\max(|x_0|, \ldots, |x_N|)$ , where the integers  $x_0, \ldots, x_N$  define a representative of x and have no common factor (other than  $\pm 1$ ). Given a subset  $U \subset V$ , we may then consider the number N(U; B) of rational points of height at most  $B \geq 1$  in U.

<sup>&</sup>lt;sup>6</sup>In more detail, for odd n, we need m(n-1)-2 > m(n-1)/2-1, which gives m(n-1) > 2, so either m = 1 and  $n \ge 5$ , or  $m \ge 2$  and  $n \ge 3$ , (both with n odd); for even n, we need m(n-1)-2 > mn/2-1, which gives m(n-2) > 2, so either m = 1 and  $n \ge 6$ , or  $m \ge 2$  and  $n \ge 4$  (both with n even). Putting these together, we obtain the claimed result.

n	m	$\bigotimes_{\alpha=1}^{m}(q_{1,\alpha},\ldots,q_{n-1,\alpha})$			dim
3	2	$(1,7)\otimes(7,4)$	$28 \times 154$	=	4312
		$(2,7)\otimes(8,6)$	$63 \times 336$	=	21168
		$(1,8)\otimes(12,11)$	$36\times1518$	=	54468
3	3	$(2,1)\otimes(4,1)\otimes(4,5)$	$3\times10\times90$	=	2700
		$(1,4)\otimes(1,4)\otimes(5,2)$	$10\times10\times35$	=	3500
		$(1,2)\otimes(1,6)\otimes(6,3)$	$3\times21\times81$	=	5103
		$(1,2)\otimes(3,6)\otimes(6,2)$	$3\times81\times48$	=	11664
		$(1,3)\otimes(1,7)\otimes(8,7)$	$6\times28\times336$	=	56448
		$(1,3)\otimes(2,9)\otimes(9,3)$	$6\times99\times162$	=	92228
4	2	$(1,4,3)\otimes(4,1,2)$	$280 \times 70$	=	19600
		$(1,1,5)\otimes(5,2,3)$	$35 \times 875$	=	30625
		$(4,1,1)\otimes(3,4,4)$	$20 \times 2464$	=	49280
		$(1,1,7)\otimes(7,1,3)$	$85\times616$	=	51744
		$(1,1,5)\otimes(4,5,2)$	$35 \times 2310$	=	80850

**Table 1.** Some anomaly-free chiral products of m irreducible representations of  $\mathfrak{su}_n$  for some values of n and m, with dimension below  $10^5$ .

Suppose such a (projective) variety contains a (projective) line L. For large B, we have the asymptotic formula  $N(L;B) \sim B^2$ , a result known as Schanuel's theorem [14].<sup>7</sup> On the other hand, a conjecture of Manin [7] asserts that (when V is a smooth Fano variety whose rational points are Zariski dense) there exists a (non-empty) open set  $U \subset V$  such that

$$N(U;B) \sim B(\log B)^{\rho-1},\tag{5.1}$$

where  $\rho$  is the rank of the Picard group of V (namely the set of isomorphism classes of line bundles on V with group multiplication given by the tensor product of line bundles). Thus, there is a dense open set in V (which clearly should not contain any lines, or more generally hyperplanes) that contains far fewer points than a line (or hyperplane), once we bound their size.

Let us now consider how this can be applied to physics. The variety  $V_{n,\otimes m}$  is smooth if and only if n is odd. Being a cubic hypersurface, it is Fano if and only if

$$(n-1)m-1 \ge 3 \implies (n-1)m \ge 4. \tag{5.2}$$

We have seen that the rational points are always Zariski dense. Therefore, the conjecture applies for all odd n and all m except when (n, m) = (3, 1) (in which case we have seen that

<sup>&</sup>lt;sup>7</sup>A summary of the proof is as follows: It can be shown that calculating  $N(\mathbb{Q}P^n, B)$  is equivalent to counting the number of *integral* lattice points in an (n+1)-dimensional cube of side length proportional to B, and the number of such points scale as  $B^{n+1}$ . In particular, the increase in dimension from n to n+1 is because a point in n-dimensional projective space corresponds to a line in (n+1)-dimensional affine space. In the limit of large B, and after considering all such cubes, the claimed asymptotic behaviour follows as the special case n = 1.

all points are nonchiral).<sup>8</sup> In such cases, the height is given by removing any common factor from the shifted Dynkin labels  $q_i$  and then taking the maximum. The points in the set Ucorrespond to chiral representations and their orbits under the automorphism group of the variety. So indeed, there is a general expectation that there are few chiral anomaly-free representations of bounded size, corresponding to the non-chiral representations.

Unfortunately, Manin's conjecture has been proven in very few cases. Indeed, it has yet to be proven for any smooth cubic surface, including the varieties  $V_{5,\otimes 1}$  and  $V_{3,\otimes 2}$  considered here and in [2].

What we do have are corresponding upper and lower bounds. Specifically, we have an upper bound for any such variety V featuring a line over  $\mathbb{Q}$  given by [16]

$$N(V \setminus \{\text{lines on } V\}; B) \lesssim B^{12/7}$$
(5.3)

and for a variety V featuring three coplanar lines over  $\mathbb{Q}$  given by [8]

$$N(V \setminus \{\text{lines on } V\}; B) \lesssim B^{4/3}.$$
(5.4)

We observe that we can find three coplanar lines on both  $V_{5,\otimes 1}$  and  $V_{3,\otimes 2}$ , so that the stronger upper bound of Eq. (5.4) holds. The former, which is projectively defined by

$$V_{5,\otimes 1}: \sigma_1^3 + \sigma_2^3 + \sigma_3^3 + \sigma_4^3 - (\sigma_1 + \sigma_2 + \sigma_3 + \sigma_4)^3 = 0,$$
(5.5)

such that  $\sigma_5 := -(\sigma_1 + \sigma_2 + \sigma_3 + \sigma_4)$ , contains the lines

$$L_1: \sigma_1 + \sigma_2 = \sigma_3 + \sigma_4 = \sigma_5 = 0, \tag{5.6}$$

$$L_2: \sigma_1 + \sigma_2 = \sigma_3 + \sigma_5 = \sigma_4 = 0, \tag{5.7}$$

$$L_3: \sigma_1 + \sigma_2 = \sigma_3 = \sigma_4 + \sigma_5 = 0, \tag{5.8}$$

which all lie on the plane  $\sigma_1 + \sigma_2 = 0$ . The latter is the set of projective solutions to the equation

$$V_{3,\otimes 2}:\sigma_{11}\sigma_{21}(\sigma_{11}+\sigma_{21})+\sigma_{12}\sigma_{22}(\sigma_{12}+\sigma_{22})=0,$$
(5.9)

where we have made the substitutions  $\sigma_{31} = -(\sigma_{11} + \sigma_{21})$  and  $\sigma_{32} = -(\sigma_{12} + \sigma_{22})$ . In a similar way, it is evident that the plane  $\sigma_{11} + \sigma_{21} = \sigma_{31} = 0$  contains the three lines

$$L'_1: \sigma_{11} + \sigma_{21} = \sigma_{31} = \sigma_{12} + \sigma_{22} = \sigma_{32} = 0,$$
(5.10)

$$L'_{2}:\sigma_{11} + \sigma_{21} = \sigma_{31} = \sigma_{12} + \sigma_{32} = \sigma_{22} = 0,$$
(5.11)

$$L'_3: \sigma_{11} + \sigma_{21} = \sigma_{31} = \sigma_{12} = \sigma_{22} + \sigma_{32} = 0.$$
(5.12)

Thus, we conclude that chiral representations of bounded size are rarer than non-chiral representations, at least in these two examples.

Manin's relation has also been shown by Slater and Swinnerton-Dyer to be a lower bound if the smooth cubic surface contains two disjoint rational lines [9]; this is the case

<sup>&</sup>lt;sup>8</sup>There are also versions of Manin's conjecture that apply to singular varieties (see *e.g.* [15]); we therefore expect that a similar conjecture applies for all (n, m) except (3, 1) and (4, 1) (where there are also no nonchiral solutions).

for both  $V_{5,\otimes 1}$  and  $V_{3,\otimes 2}$ , so (at least asymptotically) there are nevertheless plenty of chiral anomaly-free representations available for us to do physics with. To see just how many, let us close with a computation of  $\rho$  in both cases. To do so, we follow [17]: given a smooth cubic surface with two disjoint lines in 3-dimensional projective space over any field of characteristic zero with co-ordinates  $[x_0: x_1: x_2: x_3]$ , we can do linear transformations to put the lines at  $x_0 = x_1 = 0$  and  $x_2 = x_3 = 0$ . It follows from the second of these that the cubic can be written in the form

$$a(x_0, x_1)x_2^2 + d(x_0, x_1)x_2x_3 + f(x_0, x_1)x_3^2 + b(x_0, x_1)x_2 + e(x_0, x_1)x_3,$$
(5.13)

where a, d, f are linear homogeneous and b, e are quadratic homogeneous functions. The rank  $\rho$  is then given by r + 2 where r is the number of irreducible factors over the ground field of the quintic discriminant

$$\Delta = ae^2 - bde + fb^2. \tag{5.14}$$

For  $V_{5,\otimes 1}$ , we do the change of variables

$$\sigma_1 = -x_1 - x_3, \quad \sigma_2 = x_0 + x_1 + x_3, \quad \sigma_3 = -x_2 - x_3, \quad \sigma_4 = x_1 + x_2 + x_3, \quad (5.15)$$

such that the lines  $L_1$  and  $L_3$  have  $x_0 = x_1 = 0$  and  $x_2 = x_3 = 0$  respectively in the new coordinates, to obtain

$$\Delta = 27x_0x_1(x_0 + x_1)(x_0^2 + 3x_0x_1 + x_1^2)$$
(5.16)

which has five reducible factors over  $\mathbb{C}$  and  $\mathbb{R}$ , but only 4 over  $\mathbb{Q}$ , to conclude that  $\rho = 6$ .

For  $V_{3,\otimes 2}$ , we do the change of variables

$$\sigma_{11} = x_2, \qquad \sigma_{21} = x_0 - x_2, \qquad \sigma_{12} = x_3, \qquad \sigma_{22} = x_1 - x_3, \qquad (5.17)$$

which maps  $L'_1$  to the line  $x_0 = x_1 = 0$  and the line

$$L'_4: \sigma_{11} = \sigma_{21} + \sigma_{31} = \sigma_{12} = \sigma_{22} + \sigma_{32} = 0 \tag{5.18}$$

to  $x_2 = x_3 = 0$ , to obtain

$$\Delta = -x_0 x_1 (x_0 + x_1) (x_0^2 - x_0 x_1 + x_1^2)$$
(5.19)

which has five reducible factors over  $\mathbb{C}$ , but only 4 over  $\mathbb{R}$  and  $\mathbb{Q}$ , to conclude again that  $\rho = 6$ .

## Acknowledgments

This work has been partially supported by STFC consolidated grants ST/T000694/1 and ST/X000664/1, a JSPS Invitational Fellowship, and a Trinity-Henry Barlow Scholarship.

#### References

- B.C. Allanach, B. Gripaios and J. Tooby-Smith, Geometric general solution to the U(1) anomaly equations, JHEP 05 (2020) 065. [arXiv:1912.04804].
- B. Gripaios and K. Le Nguyen Nguyen, Varieties of four-dimensional gauge theories, JHEP 12 (2024) 041. [arXiv:2409.15430].
- [3] B. Allanach, B. Gripaios and J. Tooby-Smith, Solving local anomaly equations in gauge-rank extensions of the Standard Model, Phys. Rev. D 101 (2020) 075015. [arXiv:1912.10022].
- [4] I. Camp, B. Gripaios and K. Le Nguyen Nguyen, Two-dimensional gauge anomalies and p-adic numbers, JHEP 06 (2024) 202. [arXiv:2403.10611].
- [5] D.B. Costa, B.A. Dobrescu and P.J. Fox, General solution to the U(1) anomaly equations, Phys. Rev. Lett **123** (2019) 151601. [arXiv:1905.13729].
- [6] B. Segre, The non-singular cubic surfaces, Oxford University Press (1942).
- [7] J. Franke, Y. Manin and Y. Tschinkel, Rational points of bounded height on Fano varieties, Invent. Math. 95 (1989) 421.
- [8] D.R. Heath-Brown, The density of rational points on cubic surfaces, Acta Arith. 79 (1997) 17.
- [9] J.B. Slater and P. Swinnerton-Dyer, Counting points on cubic surfaces I, Astérisque 251 (1998) 1.
- [10] J. Banks and H. Georgi, Comment on gauge theories without anomalies, Phys. Rev. D 14 (1976) 1159.
- [11] S. Okubo, Gauge groups without triangular anomaly, Phys. Rev. D 16 (1977) 3528.
- [12] L.J. Mordell, *Diophantine equations*, Academic Press (1969).
- [13] J.H. Silverman, *Rational points on elliptic curves*, Undergraduate texts in mathematics, Springer (1992).
- [14] S.H. Schanuel, Heights in number fields, Bull. Soc. Math. Fr. 107 (1979) 433.
- [15] V.V. Batyrev and Y. Tschinkel, Tamagawa numbers of polarized algebraic varieties, Astérisque (1998) 299. [arXiv:alg-geom/9712002].
- [16] P. Salberger, Uniform bounds for rational points on cubic hypersurfaces, in Arithmetic and Geometry, London Mathematical Society Lecture Note Series, p. 401–421, Cambridge University Press (2015).
- [17] C. Frei and E. Sofos, Counting rational points on smooth cubic surfaces, Math. Res. Lett. 23 (2016) 127. [arXiv:1409.6224].