A STRENGTHENING OF THE HARNACK INEQUALITY

MAREK SVETLIK

ABSTRACT. We prove the stronger version of Harnack's inequality for positive harmonic functions defined on the unit disc.

1. INTRODUCTION AND THE MAIN RESULT

Let $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$ be the unit disc. It is known (see e.g. [1, p. 47] or [4, p. 13]) that the harmonic function $u : \mathbb{U} \to (0, +\infty)$ satisfies Harnack's inequality

(1)
$$\frac{1-|z|}{1+|z|} \leqslant \frac{u(z)}{u(0)} \leqslant \frac{1+|z|}{1-|z|}$$

for all $z \in \mathbb{U}$.

In this paper we prove the following stronger version of this classical result.

Theorem 1. Let $u : \mathbb{U} \to (0, +\infty)$ be a harmonic function. Then

(2)
$$\left(\frac{1+|z|^2}{1-|z|^2}+\frac{|\nabla u(0)|}{u(0)}\frac{|z|}{1-|z|^2}\right)^{-1} \leq \frac{u(z)}{u(0)} \leq \left(\frac{1+|z|^2}{1-|z|^2}+\frac{|\nabla u(0)|}{u(0)}\frac{|z|}{1-|z|^2}\right),$$

for all $z \in \mathbb{U}$. Here $\nabla u = (u_x, u_y)$ is the gradient of the function u.

Since for harmonic function $u : \mathbb{U} \to (0, +\infty)$ we have $|\nabla u(0)| \leq 2u(0)$ (see inequality (12) below), it follows that

(3)
$$\frac{1+|z|^2}{1-|z|^2} + \frac{|\nabla u(0)|}{u(0)} \frac{|z|}{1-|z|^2} \leqslant \frac{1+|z|}{1-|z|}$$

and

(4)
$$\left(\frac{1+|z|^2}{1-|z|^2} + \frac{|\nabla u(0)|}{u(0)}\frac{|z|}{1-|z|^2}\right)^{-1} \ge \frac{1-|z|}{1+|z|}$$

for all $z \in \mathbb{U}$. Therefore, the inequality (2) is stronger than the inequality (1).

Fix $c \in \mathbb{U}$ and denote by φ_c the mapping given by

$$\varphi_c(z) = \frac{z+c}{1+\overline{c}z}.$$

It is known that φ_c is a conformal isomorphism of the unit disc \mathbb{U} onto itself. In particular, if c = 1, then we define $\varphi_c(z) = 1$ for all $z \in \mathbb{U}$.

The following example shows that there is function $u : \mathbb{U} \to (0, +\infty)$ and $z \in \mathbb{U}$, so that the equality in (1) holds.

Date: January 16, 2025.

²⁰²⁰ Mathematics Subject Classification. Primary 31A05; Secondary 30C80; 30F45.

Key words and phrases. The Harnack inequality; positive harmonic functions; the Schwarz-Pick lemma; hyperbolic density.

Example 1. Let $0 \leq c \leq 1$ and let $u_1, u_2 : \mathbb{U} \to (0, +\infty)$ be defined by

$$u_1(z) = \operatorname{Re} \frac{1 + z\varphi_c(z)}{1 - z\varphi_c(z)}$$

and

$$u_2(z) = \operatorname{Re} \frac{1 - z\varphi_c(z)}{1 + z\varphi_c(z)}.$$

Then $u_1(0) = u_2(0) = 1$ and $|\nabla u_1(0)| = |\nabla u_2(0)| = 2c$. Further,
 $\frac{u_1(x)}{u_1(0)} = \frac{1 + x^2}{1 - x^2} + \frac{|\nabla u_1(0)|}{u_1(0)} \frac{x}{1 - x^2}$

and

$$\frac{u_2(x)}{u_2(0)} = \left(\frac{1+x^2}{1-x^2} + \frac{|\nabla u_2(0)|}{u_2(0)}\frac{x}{1-x^2}\right)^{-1}$$

for all $x \in [0, 1)$.

To prove the Theorem 1, we use the hyperbolic densities on the unit disc \mathbb{U} and on the right half-plane $\mathbb{K} = \{z \in \mathbb{C} : \operatorname{Re} z > 0\}.$

The hyperbolic density on the unit disc \mathbb{U} is defined by

$$\rho_{\mathbb{U}}(z) = \frac{2}{1 - |z|^2}, \quad z \in \mathbb{U}.$$

Based on Riemann's Mapping Theorem we can define the hyperbolic density on an arbitrary simply connected domain $\Omega \subset \mathbb{C}$, Ω different from \mathbb{C} (we call these domains hyperbolic), in the following way

$$\rho_{\Omega}(z) = \rho_{\mathbb{U}}(\phi(z))|\phi'(z)|, \quad z \in \Omega,$$

where $\phi: \Omega \to \mathbb{U}$ is a conformal isomorphism from Ω onto \mathbb{U} . In particular, if ψ is defined by $\psi(z) = \frac{z-1}{z+1}$, then ψ is a conformal isomorphism from \mathbb{K} onto \mathbb{U} and by direct calculation we obtain

$$\rho_{\mathbb{K}}(z) = \frac{1}{\operatorname{Re} z}, \quad z \in \mathbb{K}$$

We define the hyperbolic distance on a hyperbolic domain Ω as follows

(5)
$$d_{\Omega}(z_1, z_2) = \inf \int_{\gamma} \rho_{\Omega}(z) |dz|$$

where the infimum is taken over all C^1 curves γ that connect z_1 with z_2 in Ω . Specifically, it can be shown that

$$d_{\mathbb{U}}(z,0) = \log \frac{1+|z|}{1-|z|} = 2 \operatorname{artanh} |z|.$$

Let $f: \Omega_1 \to \Omega_2$ (Ω_1 and Ω_2 are hyperbolic domains in \mathbb{C}) be a holomorphic function. The famous Schwarz-Pick lemma asserts that

(6)
$$\rho_{\Omega_2}(f(z))|f'(z)| \leq \rho_{\Omega_1}(z), \quad z \in \Omega_1$$

As a corollary of (6) we have

(7)
$$d_{\Omega_2}(f(z_1), f(z_2)) \leq d_{\Omega_1}(z_1, z_2), \text{ for all } z_1, z_2 \in \Omega_1$$

Moreover, if f is a conformal isomorphism from Ω_1 onto Ω_2 , then equalities hold in (6) and (7). Thus, if f is a conformal isomorphism from Ω_1 onto Ω_2 , then f is an isometry from the metric space (Ω_1, d_{Ω_1}) onto the metric space (Ω_2, d_{Ω_2}) . The hyperbolic derivative of f at $z \in \Omega_1$ (for motivation and details see section 5 in [3], cf. [2]) is defined as follows

$$f^{h}(z) = \frac{\rho_{\Omega_{2}}(f(z))}{\rho_{\Omega_{1}}(z)}f'(z).$$

From (6) it follows immediately that $|f^h(z)| \leq 1$ for all $z \in \Omega_1$.

Using this notion, A. F. Beardon and T. K. Carne proved the following theorem, which is a strengthening of the Schwarz-Pick inequality (7).

Theorem 2 ([2]). Let $f : \Omega_1 \to \Omega_2$ (Ω_1 and Ω_2 are hyperbolic domains in \mathbb{C}) be a holomorphic function. Then

(8)
$$d_{\Omega_2}(f(z_1), f(z_2)) \leq \log \Big(\cosh d_{\Omega_1}(z_1, z_2) + |f^h(z)| \sinh d_{\Omega_1}(z_1, z_2) \Big).$$

Note that in the paper [2] the previous theorem was formulated and proved for the case $\Omega_1 = \Omega_2 = \mathbb{U}$. The version of this theorem that we have stated follows directly from the theorem given in [2]. The inequality (8) plays a crucial role in the proof of the Theorem 1. In addition, the following elementary facts play an important role in this proof (for this approach see [6]).

- 1) If $u : \mathbb{U} \to (0, +\infty)$ is a harmonic function, then it is known that there exists a holomorphic function $f : \mathbb{U} \to \mathbb{K}$ such that $\operatorname{Re} f = u$ and f(0) = u(0).
- 2) Using the Schwarz-Pick inequality, we obtain

(9)
$$\rho_{\mathbb{K}}(f(z))|f'(z)| \leq \rho_{\mathbb{U}}(z), \quad z \in \mathbb{U}.$$

3) Due to the Cauchy-Riemann equations, we have $f' = u_x - iu_y$, i.e. $f' = (u_x, -u_y) = \overline{\nabla u}$. From this follows

(10)
$$|f'(z)| = |\overline{\nabla u}(z)| = |\nabla u(z)|, \quad z \in \mathbb{U}.$$

4) Since $\rho_{\mathbb{K}}(z) = \frac{1}{\operatorname{Re} z}, \ z \in \mathbb{K}$ we get

(11)
$$\rho_{\mathbb{K}}(f(z)) = \rho_{\mathbb{K}}(u(z)), \quad z \in \mathbb{U}$$

5) From (9), (10) and (11) we immediately obtain the following proposition.

Proposition 1. Let $u : \mathbb{U} \to (0, +\infty)$ be a harmonic function. Then

$$\rho_{\mathbb{K}}(u(z))|\nabla u(z)| \leqslant \rho_{\mathbb{U}}(z),$$

i.e.

(12)
$$|\nabla u(z)| \leqslant \frac{2u(z)}{1-|z|^2},$$

for all $z \in \mathbb{U}$. Moreover, if u is the real part of a conformal isomorphism from \mathbb{U} onto \mathbb{K} , then in (12) equality holds for all $z \in \mathbb{U}$. On the other hand, if in (12) equality holds for some $z \in \mathbb{U}$, then u is the real part of a conformal isomorphism \mathbb{U} onto \mathbb{K} .

We call the facts 1)-5) together with the Proposition 1 the *half-plane method*. This method can be seen as an analogous version of the strip method used in the paper [7] (see also [6]).

Note that the following theorem can be derived as a corollary of (12).

Theorem 3 ([5, Theorem 1.1]). Let $u : \mathbb{U} \to (0, +\infty)$ be a harmonic function. Then

(13)
$$d_{\mathbb{K}}(u(z_1), u(z_2)) \leq d_{\mathbb{U}}(z_1, z_2),$$

for all $z_1, z_2 \in \mathbb{U}$.

Remark 1. In [5], M. Marković proved the Theorem 3 for the case where the domain of the function u is $\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$, but this is not essential since \mathbb{H} and \mathbb{U} are conformally equivalent.

Remark 2. Considering the proof of the Theorem 3 from the paper [8], it is straightforward to prove that (13) is equivalent to Harnack's inequality.

For further results of this type of inequalities, we refer the reader to [9] and the literature cited there.

2. Proof of the main result

First we recall another notion and prove two lemmas.

For a hyperbolic domain Ω , $a \in \Omega$ and r > 0 we denote by $\overline{D}_{\Omega}(a, r)$ a hyperbolic closed disc with centre a and radius r, i.e. $\overline{D}_{\Omega}(a, r) = \{z \in \Omega : d_{\Omega}(z, a) \leq r\}$. It is simply checked that

(14)
$$\overline{D}_{\mathbb{U}}(0, 2\operatorname{artanh} r) = \{ z \in \mathbb{U} : |z| \leqslant r \},\$$

for all $r \in (0, 1)$.

Lemma 1. Let $b \in (0, +\infty)$ and $r \in (0, 1)$. If $z \in \overline{D}_{\mathbb{K}}(b, 2 \operatorname{artanh} r)$ then

$$\operatorname{Re} z \in \left[\frac{1-r}{1+r}b, \frac{1+r}{1-r}b\right]$$

Proof. Recall that the mapping ψ defined by $\psi(z) = \frac{z-1}{z+1}$ is a conformal isomorphism from \mathbb{K} onto \mathbb{U} . Set $\kappa = \psi^{-1}$, $a = \psi(b)$ and $\kappa_b = \kappa \circ \varphi_a$. It is clear that κ_b is a conformal isomorphism from \mathbb{U} onto \mathbb{K} and $\kappa_b(0) = b$. Therefore, κ_b is an isometry from the metric space $(\mathbb{U}, d_{\mathbb{U}})$ onto the metric space $(\mathbb{K}, d_{\mathbb{K}})$. Therefore

$$\kappa_b \Big(\overline{D}_{\mathbb{U}}(0, 2 \operatorname{artanh} r) \Big) = \overline{D}_{\mathbb{K}}(b, 2 \operatorname{artanh} r).$$

On the other hand, taking into account (14) we have

$$\kappa_b \Big(\overline{D}_{\mathbb{U}}(0, 2 \operatorname{artanh} r) \Big) = \kappa_b \Big(\{ z \in \mathbb{U} : |z| \leq r \} \Big).$$

Furthermore, we obtain by straightforward calculations that $\kappa_b \left(\{ z \in \mathbb{U} : |z| \leq r \} \right)$ is a closed Euclidean disc with centre $c = \frac{1+r^2}{1-r^2}b$ and radius $R = \frac{2r}{1-r^2}b$. So, if $z \in \kappa_b \left(\{ z \in \mathbb{U} : |z| \leq r \} \right)$ then $\begin{bmatrix} 1 - r & 1 + r \end{bmatrix}$

Re
$$z \in [c - R, c + R] = \left[\frac{1 - r}{1 + r}b, \frac{1 + r}{1 - r}b\right].$$

4

Lemma 2. Let $0 \leq c \leq 1$. Then, for all $z \in \mathbb{U}$,

(15)
$$\log(\cosh d_{\mathbb{U}}(z,0) + c \sinh d_{\mathbb{U}}(z,0)) = \log\left(\frac{1+|z|^2+2c|z|}{1-|z|^2}\right) = d_{\mathbb{U}}(|z|\varphi_c(|z|),0).$$

Proof. Let $z \in \mathbb{U}$ be arbitrary. The first equality in (15) holds due to the fact that

$$\cosh d_{\mathbb{U}}(z,0) = \frac{1+|z|^2}{1-|z|^2}$$
 i $\sinh d_{\mathbb{U}}(z,0) = \frac{2|z|}{1-|z|^2}.$

To prove the second equality in (15), we have to determine a point $R(z) \in [0, 1)$ such that

(16)
$$d_{\mathbb{U}}(R(z),0) = \log\left(\frac{1+|z|^2+2c|z|}{1-|z|^2}\right).$$

Note that the equality (16) is equivalent to the equality

$$\frac{1+R(z)}{1-R(z)} = \frac{1+|z|^2+2c|z|}{1-|z|^2},$$

we get $R(z) = |z|\frac{c+|z|}{1-|z|^2} = |z|\varphi_c(|z|).$

and from this we get $R(z) = |z| \frac{c + |z|}{1 + c|z|} = |z|\varphi_c(|z|).$

Proof of the Theorem 1. By using the half-plane method, we obtain that there is a unique holomorphic function $f : \mathbb{U} \to \mathbb{K}$ such that $\operatorname{Re} f = u$, f(0) = u(0), $|f'(0)| = |\nabla u(0)|$ and

$$|f^{h}(0)| = \frac{\rho_{\mathbb{K}}(f(0))}{\rho_{\mathbb{U}}(0)}|f'(0)| = \frac{|\nabla u(0)|}{2u(0)}.$$

Define $c = \frac{|\nabla u(0)|}{2u(0)}$. According to Theorem 2, if $\Omega_1 = \mathbb{U}$ and $\Omega_2 = \mathbb{K}$ we get

(17)
$$d_{\mathbb{K}}(f(z), f(0)) \leq \log\left(\cosh d_{\mathbb{U}}(z, 0) + c \sinh d_{\mathbb{U}}(z, 0)\right)$$

for all $z \in \mathbb{U}$. By Lemma 2 we thus have

$$d_{\mathbb{K}}(f(z), f(0)) \leq 2 \operatorname{artanh}\left(|z|\varphi_c(|z|)\right),$$

i.e

$$f(z) \in \overline{D}_{\mathbb{K}}\Big(f(0), 2 \operatorname{artanh}\Big(|z|\varphi_c(|z|)\Big)\Big).$$

Using Lemma 1 (note that f(0) = u(0) and $u = \operatorname{Re} f$) we now obtain

(18)
$$u(z) \in \left[\frac{1-|z|\varphi_c(|z|)}{1+|z|\varphi_c(|z|)}u(0), \frac{1+|z|\varphi_c(|z|)}{1-|z|\varphi_c(|z|)}u(0)\right]$$

But since $\frac{1+|z|\varphi_c(|z|)}{1-|z|\varphi_c(|z|)} = \frac{1+|z|^2+2c|z|}{1-|z|^2}$, it follows that (18) is equivalent to (1).

MAREK SVETLIK

References

- [1] S. Axler, P. Bourdon, W. Ramey, *Harmonic function theory*, Springer Verlag, New York, 2020.
- [2] A. F. Beardon, T. K. Carne, A strengthening of the Schwarz-Pick Inequality, Amer. Math. Monthly, 99 (1992), 216-217.
- [3] A. F. Beardon and D. Minda, *The Hyperbolic Metric and Geometric Function Theory*, Proceedings of the International Workshop on Quasiconformal Mappings and their Applications (New Delhi, India, 2007), Narosa Publishing House, pp. 10-56.
- [4] P. Duren, Harmonic mappings in the plane, Cambridge University Press, 2004.
- [5] M. Marković, On harmonic functions and the hyperbolic metric, Indag. Math. 26 (2015) 19-23.
- [6] M. Mateljević, Schwarz lemma and Kobayashi metrics for harmonic and holomorphic functions, J. Math. Anal. Appl. 464 (2018) 78-100.
- [7] M. Mateljević, M. Svetlik, Hyperbolic metric on the strip and the Schwarz lemma for HQR mappings, Appl. Anal. Discrete Math. 14 (2020), 150-168.
- [8] P. Melentijević, Invariant gradientin refinements of Schwarz and Harnack inequalities, Ann. Acad. Sci. Fenn. Math 43 (2018), pp. 391-399.
- M. Svetlik, The Schwarz-Pick type estimates for harmonic mappings and hyperbolic metric, Doctoral Dissertation, University of Belgrade, Faculty of Mathematics, 2020.

FACULTY OF MATHEMATICS, UNIVERSITY OF BELGRADE, STUDENTSKI TRG 16, BELGRADE, RE-PUBLIC OF SERBIA

Email address: svetlik@matf.bg.ac.rs