

A SIMPLE EXTRAPOLATION CRITERION WITH AN APPLICATION TO WAVELET CHARACTERIZATION OF VARIOUS FUNCTION SPACES

MITSUO IZUKI, TAKAHIRO NOI AND YOSHIHIRO SAWANO

ABSTRACT. The aim of this paper is to obtain an extrapolation result without using convexification. What is new about this criterion is that the convexification of Banach spaces does not come into play. As an application, a characterization of ball Banach function spaces in terms of wavelets can be obtained. The result can be formulated so that we can take into account the smoothness property of the function spaces under consideration. The same technique can be used for the proof of the vector-valued inequalities for example. Also, the result in the present paper refines a recent result on the extension operator.

2020 Classification 42B35, 41A17, 26B33

Keywords extrapolation, wavelet, Riesz transform, Hardy–Littlewood maximal operator, Muckenhoupt weight

1. INTRODUCTION

Wavelets were originally defined as functions that provide an orthonormal basis in the L^2 space, and their mathematical theory has developed over time. Currently, various wavelets with inherent properties such as sufficient smoothness, sufficient decay, compact support, and band-limitedness are known. By applying these properties according to the situation, it is possible to characterize the space by giving it a different norm, and to construct bases with excellent properties in various function spaces, not limited to the L^2 space. Wavelet characterization of function spaces is a fascinating topic in real analysis and applied mathematics. Among others, the theory related to weights and wavelets is strongly developed. Wavelet characterization of function spaces is useful when we analyze the boundedness property of operators. Here we connect wavelet and extrapolation by presenting a simple extrapolation result for ball Banach function spaces. Our assumptions are formulated exclusively in terms of the ball Banach function space and its Köthe dual. As an application, we obtain wavelet characterization of various function spaces.

As we mentioned, we can use wavelet characterization of function spaces to investigate the boundedness property of operators. Recently, it is extremely useful to use weights when we analyze operators appearing in harmonic analysis. By a “weight” we mean a non-negative measurable function that is positive almost everywhere. The characterization of weighted function spaces has a rich history, developed based on Muckenhoupt theory in real analysis. For more details, refer to [1, Introduction]. Notably, the method for characterizing weighted Lebesgue spaces has been established through the boundedness of integral operators [9, 25]. This approach is applicable to certain generalized function

spaces, such as weighted Lebesgue spaces with variable exponent [17] and Herz-type spaces [16, 21].

Furthermore, the wavelet characterization of Herz-type spaces is achieved through the φ -transform [12]. Another method for wavelet characterization of function spaces involves the application of the extrapolation theorem, which is employed in Lebesgue spaces with variable exponent [15, 23].

We recall the definition of ball Banach function spaces defined in [35]. The space $L^0(\mathbb{R}^n)$ denotes the space of all equivalence classes of measurable functions modulo null functions. Let ρ be a mapping which maps $L^0(\mathbb{R}^n)$ to $[0, \infty]$. Recall that it is called a *ball Banach function norm* (over \mathbb{R}^n) if, for all non-negative measurable functions f, g, f_j ($j = 1, 2, 3, \dots$), for all constants $a \geq 0$ and for all open balls B in \mathbb{R}^n , the following properties hold:

- (P1) $\rho(f) = 0 \Leftrightarrow f = 0$ a.e.; $\rho(af) = a\rho(f)$, $\rho(f + g) \leq \rho(f) + \rho(g)$,
- (P2) $\rho(g) \leq \rho(f)$ if $0 \leq g \leq f$ a.e.,
- (P3) the Fatou property; $\rho(f_j) \uparrow \rho(f)$ holds whenever $0 \leq f_j \uparrow f$ a.e.,
- (P4) $\rho(\chi_B) < \infty$,
- (P5) $\|\chi_B f\|_{L^1} \lesssim \rho(f)$ with the implicit constant depending on B and ρ but independent of f .

Accordingly, the space generated by such ρ , which is given by

$$X = X_\rho = \{f \in L^0(\mathbb{R}^n) : f \text{ is a measurable function satisfying } \|f\|_{X_\rho} := \rho(|f|) < \infty\},$$

is called a ball Banach function space. Furthermore its “associate norm” ρ' is defined for a non-negative measurable function g by $\rho'(g) := \sup \{\|f \cdot g\|_{L^1} : f \in X, \rho(f) \leq 1\}$. Likewise we can consider the Köthe dual $X' = (X_\rho)' = X_{\rho'}$ for Banach lattices.

We will use the Hardy–Littlewood maximal operator M to describe our assumption. For $f \in L^0(\mathbb{R}^n)$, one defines a function Mf by

$$Mf(x) := \sup_{r>0} \frac{1}{r^n} \int_{|y-x|<r} |f(y)| dy$$

for $x \in \mathbb{R}^n$. The mapping $f \mapsto Mf$ is called the Hardy–Littlewood maximal operator.

In [31] Rutsky characterized the condition for which the Riesz transforms, given by

$$\mathcal{R}_j f(x) := \lim_{\varepsilon \downarrow 0} \int_{|y-x|>\varepsilon} \frac{x_j - y_j}{|x - y|^{n+1}} f(y) dy \quad (x \in \mathbb{R}^n),$$

for $j = 1, 2, \dots, n$, are bounded on X . In fact, he showed that M is bounded on X and on X' if and only if \mathcal{R}_j is bounded on X for all $j = 1, 2, \dots, n$.

As we mentioned, weights are fundamental tools to investigate the boundedness of operators. Let w be a weight and $1 < p < \infty$. We also consider $L^p(w)$ which collects all $f \in L^0(\mathbb{R}^n)$ for which $\|f\|_{L^p(w)} := \|fw^{\frac{1}{p}}\|_{L^p}$ is finite. Motivated by this result, using the Muckenhoupt class A_p , which will be recalled in (1.2) below, we will provide an extrapolation theorem together with applications in this paper. This strongly shows that weights play the fundamental role when we analyze the boundedness properties of operators.

Theorem 1.1. *Let X be a ball Banach function space and $1 < p < \infty$. Assume that M is bounded on X and on X' . For a given an increasing function $N : [1, \infty) \rightarrow (0, \infty)$, write*

$$(1.1) \quad \mathcal{F} = \{(f, g) : \|f\|_{L^p(w)} \leq N([w]_{A_p}) \|g\|_{L^p(w)}\}.$$

Then there exists a constant $C > 0$ such that $\|f\|_X \leq C\|g\|_X$ for all $(f, g) \in \mathcal{F}$.

It is useful to state Theorem 1.1 in a different manner.

Corollary 1.2. *Let X be a Banach function space and $1 < p < \infty$. Then the following are equivalent:*

- (A) *Assume that M is bounded on X and on X' .*
- (B) *The Riesz transform \mathcal{R}_j is bounded on X for each $j = 1, 2, \dots, n$.*
- (C) *Define \mathcal{F} by (1.1). Then there exists a constant $C > 0$ such that $\|f\|_X \leq C\|g\|_X$ for all $(f, g) \in \mathcal{F}$.*

Corollary 1.2 follows directly from a known result and Theorem 1.1. As we mentioned, Rutsky established that (A) and (B) are equivalent. Theorem 1.1 states that (A) implies (C). Finally, since $(\mathcal{R}_j f, f) \in \mathcal{F}$ for $N(t) = \alpha t$ with $\alpha \gg 1$ [34, Theorem 305], it follows that (C) implies (B).

Here, we make a clarifying remark on the strength of the assumption in Theorem 1.1 regarding X .

Remark 1.3. If X satisfies the assumption in Theorem 1.1, meaning that M is bounded on both X and X' , then according to Lerner and Pérez [24], there exists a constant $\eta > 1$ such that the operator $f \mapsto (M[|f|^\eta])^{\frac{1}{\eta}}$ is bounded on both X and X' . In this scenario, we can establish $X \hookrightarrow L^\eta(w)$ for some $w \in A_1$. We refer to [29, Lemma 2.7]. See also [33, Lemma 3].

Curbera, García-Cuerva, Martell, and Pérez made a significant breakthrough by obtaining the extrapolation theorem for rearrangement invariant Banach function spaces [7]. The case where X is not rearrangement invariant has gained increasing attention. For instance, in [13, Theorem 3.3], Ho employed p -convexification to achieve a vector-valued extension of the maximal inequality. For a ball Banach function space X and $p > 0$, we consider its p -convexification given by $X^p := \{f \in L^0(\mathbb{R}^n) : |f|^{\frac{1}{p}} \in X\}$. The norm of X^p is given by $\|f\|_{X^p} := (\| |f|^{\frac{1}{p}} \|_X)^p$. Remarkably, Theorem 1.1 does not require p -convexification. Nor do we need the absolute continuity of the norm of X . Recall that X is said to have the absolutely continuous norm if $\|f\chi_{E_j}\|_X \rightarrow 0$ as $j \rightarrow \infty$ for any $f \in X$ and any sequence $\{E_j\}_{j=1}^\infty$ of sets decreasing to 0.

Here and below, in addition to the above notation, we use the following convention in this paper.

- By a “cube” we mean a compact cube whose edges are parallel to the *coordinate axes*. The metric closed ball defined by ℓ^∞ is called a *cube*. If a cube has center x and radius r , we denote it by $Q(x, r)$. Namely, we write

$$Q(x, r) \equiv \left\{ y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n : \max_{j=1,2,\dots,n} |x_j - y_j| \leq r \right\}$$

when $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ and $r > 0$. The symbol \mathcal{Q} stands for all cubes.

- Given $x \in \mathbb{R}^n$ and $r > 0$, we denote the open ball centered at x and having radius $r > 0$ by $B(x, r)$. We abbreviate $B(r) := B(0, r)$.
- Let $A, B \geq 0$. Then $A \lesssim B$ means that there exists a constant $C > 0$ such that $A \leq CB$, where C is usually independent of the functions we are considering. The symbol $A \sim B$ stands for the two-sided inequality $A \lesssim B \lesssim A$.

- Let $l \in \mathbb{N}$. Denote by M^l the l -fold composition of M . The operator M^0 stands for the operator $f \in L^0(\mathbb{R}^n) \mapsto |f| \in L^0(\mathbb{R}^n)$.
- Given a measurable set $E \subset \mathbb{R}^n$, we denote the characteristic function by χ_E . In addition, $|E|$ is the Lebesgue measure of E .
- For a measurable set E and a weight w , $w(E)$ stands for the integral of w over E :

$$w(E) := \int_E w(x) dx.$$

- A locally integrable weight w is an A_1 -weight or belongs to the A_1 -class, if $0 < w < \infty$ almost everywhere, and

$$[w]_{A_1} := \left\| \frac{Mw}{w} \right\|_{L^\infty} < \infty.$$

- Let $1 < p < \infty$. A locally integrable weight w is an A_p -weight or belongs to the A_p -class, if $0 < w < \infty$ almost everywhere, and

$$(1.2) \quad [w]_{A_p} := \sup_{Q \in \mathcal{Q}} \frac{w(Q)}{|Q|} \left(\frac{1}{|Q|} \int_Q w(x)^{-\frac{1}{p-1}} dx \right)^{p-1} < \infty.$$

- Denote by $\|M\|_{X \rightarrow X}$ the operator norm of M acting on X .
- Given two complex-valued functions f, g , we formally write the L^2 -inner product by

$$\langle f, g \rangle := \int_{\mathbb{R}^n} f(x) \overline{g(x)} dx.$$

Here is the organization of this paper. Section 2 proves Theorem 1.1. Section 3 is oriented to applications of Theorem 1.1. We obtain the boundedness properties of operators and the wavelet characterization in Section 3. We also take up the vector-valued inequality of M and the extension operator in Section 3. We will see that our method allows us to include more function spaces than are dealt with in [39]. We take up various function spaces in Section 4. We compare what we can say from Theorem 1.1 with known results in Sections 3 and 4.

2. PROOF OF THEOREM 1.1

Let $\varepsilon > 0$. It suffices to show that

$$(2.1) \quad \|f\|_X \leq C_1 \|g\|_X + C_2 \varepsilon \|f\|_X.$$

Here, $C_1 > 0$ depends on ε while $C_2 > 0$ is independent of ε or $(f, g) \in \mathcal{F}$. In fact, since, for any $r > 0$, $(\chi_{B(r)} \chi_{[0,r]}(|f|)f, g) \in \mathcal{F}$, we have

$$\|\chi_{B(r)} \chi_{[0,r]}(|f|)f\|_X \leq C_1 \|g\|_X + C_2 \varepsilon \|\chi_{B(r)} \chi_{[0,r]}(|f|)f\|_X$$

once we show (2.1). Observe that $\chi_{B(r)} \chi_{[0,r]}(|f|)f \in X$. By taking $\varepsilon = \frac{1}{2C_2}$, we obtain

$$\|\chi_{B(r)} \chi_{[0,r]}(|f|)f\|_X \leq 2C_1 \|g\|_X.$$

Letting $r \rightarrow \infty$, we obtain the desired result.

We will show (2.1). Since $(f, g) \in \mathcal{F}$ if and only if $(|f|, |g|) \in \mathcal{F}$ for any $f, g \in L^0(\mathbb{R}^n)$, we may assume that f, g are non-negative here and below. As before, we may assume that $f \in L_c^\infty(\mathbb{R}^n)$ by the truncation argument, where $L_c^\infty(\mathbb{R}^n)$ is the set of all bounded and compactly supported functions. Furthermore, by adding $\varepsilon M \chi_{B(1)}$ to g with $\varepsilon > 0$, we

may assume that g is positive everywhere. Finally, we may assume that $g \in X$; otherwise the conclusion is clear.

We write

$$R_k(x) := \sum_{l=0}^{\infty} \frac{M^l k(x)}{(\alpha \|M\|_{X \rightarrow X})^l}, \quad R'_k(x) := \sum_{l=0}^{\infty} \frac{M^l k(x)}{(\alpha \|M\|_{X' \rightarrow X'})^l} \quad (x \in \mathbb{R}^n).$$

for a function k . Here $\alpha \geq 2$ is a constant that is fixed shortly. See (2.13).

We use

$$(2.2) \quad ab^{p-1} \leq C_\varepsilon a^p + \varepsilon b^p$$

for $a, b, \varepsilon > 0$, where $C_\varepsilon > 0$ is a constant that depends only on ε and p . To estimate $\|f\|_X$, we fix a non-negative function $h \in X'$ with norm strictly less than 1 to dualize. It suffices to show that there exist a constant C_0 independent of ε, f, g and h , and a constant C_ε independent of f, g and h but dependent on ε such that

$$(2.3) \quad \int_{\mathbb{R}^n} f(x)h(x)dx \leq C_\varepsilon \|g\|_X + C_0 \varepsilon \|f\|_X.$$

Indeed, once (2.3) is proved, since $X = (X)'$ and $f \in L_c^\infty(\mathbb{R}^n) \subset X$, we have (2.1). Remark that h can be assumed positive by adding $\varepsilon M \chi_{B(1)}$ with $\varepsilon \ll 1$ if necessary. Let us concentrate on the proof of (2.1) assuming that $f \in L_c^\infty(\mathbb{R}^n)$ and that g, h is positive.

Assuming that $f \geq 0$ and $g, h > 0$, we see that $R_{g+f} > 0$ and $R'_h > 0$. Arithmetic shows

$$[R_{g+f}]_{A_1} \leq \alpha \|M\|_{X \rightarrow X}, \quad [R_h]_{A_1} \leq \alpha \|M\|_{X' \rightarrow X'}.$$

Thus,

$$(2.4) \quad [R_{g+f}^{1-p} R'_h]_{A_p} \leq ([R_{g+f}]_{A_1})^{p-1} [R_{g+f}^{1-p} R'_h]_{A_1} \lesssim 1.$$

As we mentioned $R_{g+f} > 0$. Thus the power function $(R_{g+f})^{p-1}$ makes sense. Since $h \leq R'_h$, we have

$$(2.5) \quad \begin{aligned} \int_{\mathbb{R}^n} f(x)h(x)dx &\leq \int_{\mathbb{R}^n} f(x)R'_h(x)dx \\ &= \int_{\mathbb{R}^n} f(x)R_{g+f}(x)^{p-1} R_{g+f}(x)^{1-p} R'_h(x)dx. \end{aligned}$$

Due to (2.2),

$$(2.6) \quad \begin{aligned} \int_{\mathbb{R}^n} f(x)R_{g+f}(x)^{p-1} R_{g+f}(x)^{1-p} R'_h(x)dx \\ \leq C_\varepsilon \int_{\mathbb{R}^n} f(x)^p R_{g+f}(x)^{1-p} R'_h(x)dx + \varepsilon \int_{\mathbb{R}^n} R_{g+f}(x)^p R_{g+f}(x)^{1-p} R'_h(x)dx. \end{aligned}$$

We estimate the first term on the right-hand side of (2.6). Since $(f, g) \in \mathcal{F}$, we deduce

$$(2.7) \quad \begin{aligned} \int_{\mathbb{R}^n} f(x)^p R_{g+f}(x)^{1-p} R'_h(x)dx &\leq N([R_{g+f}^{1-p} R'_h]_{A_p})^p \int_{\mathbb{R}^n} g(x)^p R_{g+f}(x)^{1-p} R'_h(x)dx \\ &\lesssim \int_{\mathbb{R}^n} g(x)^p R_{g+f}(x)^{1-p} R'_h(x)dx \end{aligned}$$

from (2.4). Since $R_{g+f} \geq g$, we obtain

$$(2.8) \quad \int_{\mathbb{R}^n} g(x)^p R_{g+f}(x)^{1-p} R'_h(x) dx \leq \int_{\mathbb{R}^n} g(x) R'_h(x) dx.$$

By the Hölder inequality for X and the inequality $\|R'_h\|_{X'} \lesssim \|h\|_{X'}$, we obtain

$$(2.9) \quad \int_{\mathbb{R}^n} g(x) R'_h(x) dx \lesssim \|g\|_X \|R'_h\|_{X'} \lesssim \|g\|_X \|h\|_{X'} \lesssim \|g\|_X.$$

Putting (2.7)–(2.9) together, we have

$$(2.10) \quad \int_{\mathbb{R}^n} f(x)^p R_{g+f}(x)^{1-p} R'_h(x) dx \lesssim \|g\|_X.$$

We embark on the estimate of the second term on the right-hand side of (2.6). Since

$$\begin{aligned} (a_1 + a_2 + \cdots + a_l + \cdots)^p &\leq 2^p a_1^p + 2^p (a_2 + a_3 + \cdots + a_l + \cdots)^p \\ &\leq 2^p a_1^p + 4^p a_2^p + 4^p (a_3 + a_4 + \cdots + a_l + \cdots)^p \\ &\leq \sum_{l=1}^{\infty} 2^{pl} a_l^p \end{aligned}$$

for any non-negative sequence $\{a_l\}_{l=1}^{\infty}$, we obtain

$$(2.11) \quad \begin{aligned} &\int_{\mathbb{R}^n} R_{g+f}(x)^p R_{g+f}(x)^{1-p} R'_h(x) dx \\ &\leq 2^p \sum_{l=0}^{\infty} \frac{2^{pl}}{(\alpha \|M\|_{X \rightarrow X})^{pl}} \int_{\mathbb{R}^n} M^l [f + g](x)^p R_{g+f}(x)^{1-p} R'_h(x) dx. \end{aligned}$$

Recall that $1 < p < \infty$. Since M is bounded on $L^p(\mathbb{R}^n)$, we can find a constant $\beta \in (1, \infty)$ such that

$$\int_{\mathbb{R}^n} M F(x)^p W(x) dx \leq \beta ([W]_{A_p})^{p'} \int_{\mathbb{R}^n} |F(x)|^p W(x) dx$$

for all $F \in L^0(\mathbb{R}^n)$ and $W \in A_p$. Thus, from (2.4) once again

$$(2.12) \quad \begin{aligned} &\int_{\mathbb{R}^n} M^l [f + g](x)^p R_{g+f}(x)^{1-p} R'_h(x) dx \\ &\leq \beta^l \int_{\mathbb{R}^n} (f(x) + g(x))^p R_{g+f}(x)^{1-p} R'_h(x) dx. \end{aligned}$$

Assuming that $\alpha \gg 2$, we may suppose

$$\alpha > \frac{2\beta^{\frac{1}{p}}}{\|M\|_{X \rightarrow X}}.$$

In this case,

$$(2.13) \quad \sum_{l=0}^{\infty} \frac{2^{pl} \beta^l}{(\alpha \|M\|_{X \rightarrow X})^{pl}} < \infty.$$

Since $R_{g+f} \geq f + g$, it follows from (2.11) and (2.12) that

$$(2.14) \quad \int_{\mathbb{R}^n} R_{g+f}(x)^p R_{g+f}(x)^{1-p} R'_h(x) dx$$

$$\begin{aligned}
 &\lesssim \sum_{l=0}^{\infty} \frac{2^{pl} \beta^l}{(\alpha \|M\|_{X \rightarrow X})^{pl}} \int_{\mathbb{R}^n} (f(x) + g(x)) R'_h(x) dx \\
 &\sim \int_{\mathbb{R}^n} (f(x) + g(x)) R'_h(x) dx \\
 &\lesssim \|f\|_X + \|g\|_X.
 \end{aligned}$$

Putting (2.6), (2.10) and (2.14) together, we have (2.1). This completes the proof of Theorem 1.1.

3. APPLICATIONS

We present applications. To this end, we consider X -based Sobolev spaces over domains. First we recall the one over \mathbb{R}^n . Let $s > 0$. Recall that the Bessel potential $(1 - \Delta)^{\frac{s}{2}}$ is an isomorphism in $\mathcal{S}'(\mathbb{R}^n)$; see [32, p. 251].

Definition 3.1. Let $s > 0$. Define the X -based Sobolev space $W_X^s(\mathbb{R}^n)$ of order s to be the set of all $f \in X(\mathbb{R}^n)$ for which $(1 - \Delta)^{\frac{s}{2}} f$ is represented by an element in X . Equip $W_X^s(\mathbb{R}^n)$ with the norm given by $\|f\|_{W_X^s} = \|(1 - \Delta)^{\frac{s}{2}} f\|_X$ for all $f \in W_X^s(\mathbb{R}^n)$.

Assume that M is bounded on X and on X' . Recall that the inverse operator $(1 - \Delta)^{-\frac{s}{2}}$ of $(1 - \Delta)^{\frac{s}{2}}$ has the convolution kernel K satisfying $K \in L^1(\mathbb{R}^n)$ (see [34, Corollary 197]). Hence $(1 - \Delta)^{-\frac{s}{2}}$ is bounded on X since we have a pointwise estimate $|(1 - \Delta)^{-\frac{s}{2}} f(x)| \lesssim Mf(x)$, $x \in \mathbb{R}^n$, for all $f \in X$; see [34, Proposition 159]. Thus, $W_X^s(\mathbb{R}^n)$ is a Banach space. Note that this is a natural extension of $L^{p,s}(w)$ considered in [14, 18] if $X = L^p(w)$ with $1 < p < \infty$ and $w \in A_p$. By definition $W_X^0(\mathbb{R}^n) = X$.

Our definitions can be extended to domains. Let D be a bounded Lipschitz domain. Let $k \in \mathbb{N}$, $1 < p < \infty$, and $w \in A_p$. Define $L^{p,k}(D, w)$ to be the set of all $f \in L^p(w)$ for which $\partial^\alpha f$, the weak partial derivative of order α , belongs to $L^p(w)$ for any multiindex α with $|\alpha| \leq k$.

For a ball Banach function space X , we write $X(D)$ to be the set of all $f \in L_{\text{loc}}^1(D)$ for which Zf belongs to X . Let Z be the zero extension operator. The norm of $f \in X(D)$ is given by $\|f\|_{X(D)} = \|Zf\|_X$.

We write $W_X^k(D)$ for the set of all $f \in L_{\text{loc}}^1(D)$ for which $\partial^\alpha f$ belongs to $X(D)$ for any multiindex α with $|\alpha| \leq k$.

Here is the organization of Section 3. Section 3.1 considers wavelet characterization. Section 3.2 deals with vector-valued inequalities of the Hardy–Littlewood maximal operator. We refine a recent result in [39] in Section 3.3.

3.1. Wavelet characterization. Based on the fundamental wavelet theory (see [11, 30] for example), we can choose compactly supported C^K -functions

$$(3.1) \quad \varphi \text{ and } \psi^l \quad (l = 1, 2, \dots, 2^n - 1)$$

so that the following conditions are satisfied:

- (1) For any $J \in \mathbb{Z}$, the system

$$\left\{ \varphi_{J,k}, \psi_{j,k}^l : k \in \mathbb{Z}^n, j \geq J, l = 1, 2, \dots, 2^n - 1 \right\}$$

is an orthonormal basis of $L^2(\mathbb{R}^n)$. Here, given a function F defined on \mathbb{R}^n , we write

$$F_{j,k} := 2^{\frac{jn}{2}} F(2^j \cdot -k)$$

for $j \in \mathbb{Z}$ and $k \in \mathbb{Z}^n$.

(2) Denote by \mathcal{P}_0^\perp the set of all integrable functions having 0 integral. We have

$$(3.2) \quad \psi^l \in \mathcal{P}_0^\perp \quad (l = 1, 2, \dots, 2^n - 1).$$

In addition, they are real-valued and compactly supported with

$$(3.3) \quad \text{supp}(\varphi) = \text{supp}(\psi^l) = [0, 2N - 1]^n$$

for some $N \in \mathbb{N}$.

We also consider cubes $Q_{j,k}$ and $Q_{j,k}^*$ defined by

$$Q_{j,k} := \prod_{m=1}^n [2^{-j}k_m, 2^{-j}(k_m + 1)], \quad Q_{j,k}^* := \prod_{m=1}^n [2^{-j}k_m, 2^{-j}(k_m + 2N - 1)],$$

where $N \in \mathbb{N}$ satisfies (3.3), and

$$(3.4) \quad \chi_{j,k} := 2^{\frac{jn}{2}} \chi_{Q_{j,k}}, \quad \chi_{j,k}^* := 2^{\frac{jn}{2}} \chi_{Q_{j,k}^*}.$$

We note that the L^2 -inner product $\langle f, \phi \rangle$ exists for the complex-valued functions $f \in L_{\text{loc}}^1(\mathbb{R}^n)$ and $\phi \in C_c^\infty$. Then, using the L^2 -inner product, we define two square functions Vf , $W_s f$ by

$$Vf := V^\varphi f := \left(\sum_{k \in \mathbb{Z}^n} |\langle f, \varphi_{J,k} \rangle \chi_{J,k}|^2 \right)^{\frac{1}{2}},$$

$$W_s f := W_s^{\psi^l} f := \left(\sum_{l=1}^{2^n-1} \sum_{j=J}^{\infty} \sum_{k \in \mathbb{Z}^n} \left| 2^{js} \langle f, \psi_{j,k}^l \rangle \chi_{j,k} \right|^2 \right)^{\frac{1}{2}}.$$

Here, $s \geq 0$ is a non-negative real number and J is a fixed integer. We abbreviate $Wf := W_0 f$.

The wavelet characterization of X can be obtained with ease.

Theorem 3.2. *Let $K > s > 0$. Suppose that we have $\varphi, \psi_j \in C^K(\mathbb{R}^n)$ as in (3.1). Assume that (3.2) holds, that M is bounded on X and on X' , or equivalently, each \mathcal{R}_j is bounded on X . Then the following are equivalent for any $f \in X$:*

- (1) $Vf + W_s f \in X$.
- (2) $f \in W_X^s(\mathbb{R}^n)$.

Furthermore, in this case $\|f\|_{W_X^s} \sim \|Vf + W_s f\|_X$.

Proof. We note that

$$(3.5) \quad ((1 - \Delta)^{\frac{s}{2}} f, Vf + W_s f) \in \mathcal{F}$$

and

$$(3.6) \quad (Vf + W_s f, (1 - \Delta)^{\frac{s}{2}} f) \in \mathcal{F}$$

follow [18, Theorem 4.6]. Although in [18] the operator $(1 - t_0^2 \Delta)^{-\frac{s}{2}}$ with $0 < t_0 \ll 1$ was considered, we can replace it with $(1 - \Delta)^{-\frac{s}{2}}$ since the Hardy–Littlewood maximal operator M is bounded on $L^p(w)$. \square

Despite the above observations, it is helpful to offer some words (3.5) and (3.6) when s is an integer.

Remark 3.3. Relations (3.5) and (3.6) are essentially proved by [11, 30] in the case that X is the non-weighted Lebesgue space. Lemarié-Rieusset [25] has proved it for the weighted case with $s = 0$. The first author [14] has generalized the result to the case that s is a non-negative integer. Thus, we are in the position of using Theorem 1.1 again.

Letting $s = 0$, we derive the following corollary. It suffices to assume *a priori* that f is locally integrable.

Corollary 3.4. *Assume that M is bounded on X and on X' , or equivalently, each \mathcal{R}_j is bounded on X . Then the following are equivalent for any $f \in L^1_{\text{loc}}(\mathbb{R}^n)$:*

- (1) $Vf + Wf \in X$.
- (2) $f \in X$.

Furthermore, in this case $\|f\|_X \sim \|Vf + Wf\|_X$.

The proof is the same as Theorem 3.2, although we assume that f is merely locally integrable.

It is interesting to compare Corollary 3.4 with the following result:

Theorem 3.5. [22, Theorem 8] *In addition to the assumption of Corollary 3.4, assume that X is separable. Let $f \in X$. Then*

$$(3.7) \quad f = \sum_{k \in \mathbb{Z}^n} \langle f, \varphi_{J,k} \rangle \varphi_{J,k} + \sum_{l=1}^{2^n-1} \sum_{j=J}^{\infty} \sum_{k \in \mathbb{Z}^n} \langle f, \psi_{J,k}^l \rangle \psi_{J,k}^l$$

holds unconditionally in the topology of X .

Once we use Corollary 3.4, Theorem 3.5 is easy to show. In fact, since X is separable, X has the absolutely continuous norm. Corollary 3.4 shows that (3.7) converges in X .

3.2. Vector-valued maximal inequality. A similar technique can be used to obtain the vector-valued inequality.

Example 3.6. Let $r \in (1, \infty]$. Assume that M is bounded on X and on X' , or equivalently, each \mathcal{R}_j is bounded on X . Then there exists a constant $C > 0$ such that

$$\left\| \left(\sum_{j=1}^{\infty} M f_j^r \right)^{\frac{1}{r}} \right\|_X \leq C \left\| \left(\sum_{j=1}^{\infty} |f_j|^r \right)^{\frac{1}{r}} \right\|_X$$

for all $\{f_j\}_{j=1}^{\infty} \subset X$ since

$$\left(\left(\sum_{j=1}^{\infty} M f_j^r \right)^{\frac{1}{r}}, \left(\sum_{j=1}^{\infty} |f_j|^r \right)^{\frac{1}{r}} \right) \in \mathcal{F}$$

for $N(t) = \alpha t$ with $t \gg 1$. See [34, Theorem 303]. This result extends [29, Theorem 2.1].

3.3. Extension operator. Let D be a bounded Lipschitz domain. Chua established that there exists a bounded linear operator $\Lambda : L^{p,k}(D, w) \rightarrow L^{p,k}(w)$ independent of w such that $\Lambda f|_D = f$ for all $f \in L^{p,k}(D, w)$. More precisely, letting Z the zero extension operator, Chua essentially proved that

$$\left(\Lambda f, \sum_{|\alpha| \leq k} |Z \partial^\alpha f| \right) \in \mathcal{F}$$

with some suitable choice of $N(\cdot)$. See [4, Theorem 1.1]. Based on this fact, we investigate the action of the extension operator Λ on X -based Sobolev spaces over D .

Example 3.7. Assume that M is bounded on X and on X' , or equivalently, each \mathcal{R}_j is bounded on X . Let $f \in W_X^k(D)$. Then $f \in L^{p,k}(w, D)$ for some $w \in A_1$. Thus, Λf makes sense and $\|\Lambda f\|_X \sim \sum_{|\alpha| \leq k} \|Z \partial^\alpha f\|_X$.

We compare Example 3.7 with an existing result [39, Theorem 5.4]

Remark 3.8.

- (1) A recent work [39, Theorem 5.4] requires that $X^{\frac{1}{p}}$ is a ball Banach space for some $p > 1$ and that M is bounded on $(X^{\frac{1}{p}})'$. This assumption implies that M is bounded on both X and X' , since it implies that \mathcal{R}_j is bounded on X .
- (2) A recent work [39, Theorem 5.4] requires that X has the absolutely continuous norm. According to Theorem 1.1, X need not have the absolutely continuous norm.

4. EXAMPLES OF BANACH LATTICES TO WHICH THEOREM 1.1 IS APPLICABLE

We will exhibit examples of Banach lattices to which Theorem 1.1 is applicable. We concentrate on wavelet characterization. However, it is worth mentioning that we have a counterpart to vector-valued inequalities.

4.1. Weighted Lebesgue spaces. Let $1 < q < \infty$ and $w \in A_q$. It is well known that M and \mathcal{R}_j , $j = 1, 2, \dots, n$, are bounded on $L^q(w)$ [34, Theorems 290 and 306]. Thus, Theorem 1.1 is applicable to $L^q(w)$.

Consider a more concrete case. Let $1 < q < \infty$ and $v(x) = |x|^\alpha$ be a power weight of order $\alpha \in \mathbb{R}$. According to [34, Theorem 283], $v \in A_q$ if and only if $-n < \alpha < n(q-1)$. This condition also applies to the weight W , defined by $W(x) = \max(1, |x|)^\alpha$ for $x \in \mathbb{R}^n$. Therefore, if $-n < \alpha < n(q-1)$, Theorem 1.1 is applicable to both $L^q(v)$ and $L^q(W)$.

4.2. Lorentz spaces. Let $1 \leq p < \infty$ and $1 \leq q \leq \infty$. Then the Lorentz space $L^{p,q}(\mathbb{R}^n)$ is the set of all $f \in L^0(\mathbb{R}^n)$ for which the quasi-norm

$$\|f\|_{L^{p,q}} := \left\{ \int_0^\infty (t^{\frac{1}{p}} f^*(t))^q \frac{dt}{t} \right\}^{\frac{1}{q}}$$

is finite. Here f^* stands for the decreasing rearrangement of f . Remark that if $1 < p < \infty$, then there exists a norm $\|\cdot\|$ on $L^{p,q}(\mathbb{R}^n)$ such that

$$\|f\|_{L^{p,q}} \sim \|f\|$$

for all $f \in L^{p,q}(\mathbb{R}^n)$ [34, Corollary 39].

Furthermore, since the space $L^{p,q}(\mathbb{R}^n)$ can be obtained through the real interpolation of the spaces $L^{p_0}(\mathbb{R}^n)$ and $L^{p_1}(\mathbb{R}^n)$ as shown in [2], the operators M and \mathcal{R}_j (for $j = 1, 2, \dots, n$) are bounded on $L^{p,q}(\mathbb{R}^n)$ for $1 < p < \infty$ and $1 \leq q \leq \infty$. Consequently, Theorem 1.1 can be applied to $L^{p,q}(\mathbb{R}^n)$.

To our knowledge, the wavelet characterization of the Lorentz spaces $L^{p,q}(\mathbb{R}^n)$ for $1 < p < \infty$ and $1 \leq q \leq \infty$ has been established by Soardi [37].

It is important to note that Lorentz spaces $L^{p,\infty}(\mathbb{R}^n)$, also known as the weak Lebesgue spaces $WL^p(\mathbb{R}^n)$, are not separable. Therefore, in this case, we cannot apply Theorem 3.5 to obtain norm equivalence. However, according to Remark 1.3, for any $\eta \in (1, p)$ and any $f \in WL^p(\mathbb{R}^n)$, there exists an A_1 -weight w (which is independent of f) such that the convergence of the expansion in Theorem 3.5 occurs in $L^\eta(w)$.

4.3. Herz spaces. For each $k \in \mathbb{Z}$, we set $C_k := B(2^k) \setminus B(2^{k-1})$. Let $p, q \in [1, \infty]$ and $\alpha \in \mathbb{R}$. We write $\chi_k = \chi_{C_k}$ for each $k \in \mathbb{Z}$. The non-homogeneous Herz space $K_p^{\alpha,q}(\mathbb{R}^n)$ consists of all $f \in L^0(\mathbb{R}^n)$ for which

$$\|f\|_{K_p^{\alpha,q}} := \|f\|_{L^p(B(1))} + \left(\sum_{k=1}^{\infty} (2^{k\alpha} \|f\chi_k\|_{L^p})^q \right)^{\frac{1}{q}} < \infty$$

and the homogeneous Herz space $\dot{K}_p^{\alpha,q}(\mathbb{R}^n)$ consists of all $f \in L^0(\mathbb{R}^n)$ for which

$$\|f\|_{\dot{K}_p^{\alpha,q}} := \left(\sum_{k=-\infty}^{\infty} (2^{k\alpha} \|f\chi_k\|_{L^p})^q \right)^{\frac{1}{q}} < \infty.$$

Let $\alpha \in \mathbb{R}$, $1 < p < \infty$ and $1 \leq q \leq \infty$ satisfy

$$(4.1) \quad -\frac{n}{p} < \alpha < n - \frac{n}{p}.$$

Then M and \mathcal{R}_j are bounded on $\dot{K}_p^{\alpha,q}(\mathbb{R}^n)$ and $K_p^{\alpha,q}(\mathbb{R}^n)$ for all $j = 1, 2, \dots, n$. See [26, 27, 28]. Thus, Theorem 1.1 is applicable to $K_p^{\alpha,q}(\mathbb{R}^n)$ and $\dot{K}_p^{\alpha,q}(\mathbb{R}^n)$. As far as we know, there are some results [12, 21] on wavelet characterization of Herz spaces. Since M is bounded on $\dot{K}_p^{\alpha,q}(\mathbb{R}^n)$, $K_p^{\alpha,q}(\mathbb{R}^n)$ and their preduals, it follows that many of these results fall within the scope of Corollary 3.4. Remark that $\dot{K}_p^{\alpha,\infty}(\mathbb{R}^n)$ and $K_p^{\alpha,\infty}(\mathbb{R}^n)$ are not separable. However, in view of Remark 1.3, we see that for any $\eta \in (1, p)$ and $f \in \dot{K}_p^{\alpha,\infty}(\mathbb{R}^n) \cup K_p^{\alpha,\infty}(\mathbb{R}^n)$ there exists an A_1 -weight w independent of f such that the convergence of the expansion in Theorem 3.5 takes place in $L^\eta(w)$.

4.4. Weighted Lebesgue spaces with variable exponents. Let w be a weight. The Lebesgue space $L^{p(\cdot)}(w)$ with variable exponent consists of all $f \in L^0(\mathbb{R}^n)$ for which

$$\|f\|_{L^{p(\cdot)}(w)} := \inf \left\{ \lambda > 0 : \int_{\mathbb{R}^n} \left(\frac{f(x)}{\lambda} \right)^{p(x)} w(x) dx \leq 1 \right\} < \infty.$$

We postulate some conditions on variable exponents. We use the following terminology: we use the following standard notation on exponents:

Definition 4.1.

(1) Let $r(\cdot)$ be a variable exponent. We write

$$r_- := \operatorname{ess\,inf}_{x \in \mathbb{R}^n} r(x), \quad r_+ := \operatorname{ess\,sup}_{x \in \mathbb{R}^n} r(x).$$

(2) The class $\mathcal{P}_0 = \mathcal{P}_0(\mathbb{R}^n)$ collects all the variable exponents $r(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ that satisfy $0 < r_- \leq r_+ < \infty$, while $\mathcal{P} = \mathcal{P}(\mathbb{R}^n)$ collects all the variable exponents $r(\cdot)$ in \mathcal{P}_0 that satisfy $r_- > 1$.

Definition 4.2. Let $r(\cdot) \in L^0(\mathbb{R}^n)$.

(1) The variable exponent $r(\cdot)$ satisfies the local log-Hölder continuity condition if

$$(4.2) \quad |r(x) - r(y)| \leq \frac{c_*}{\log(|x - y|^{-1})} \quad \text{for } x, y \in \mathbb{R}^n \quad \text{with } |x - y| \leq \frac{1}{2}.$$

(2) If (4.2) is satisfied only for $y = 0$, then $r(\cdot)$ is log-Hölder continuous at the origin. The class $\operatorname{LH}_0 = \operatorname{LH}_0(\mathbb{R}^n)$ stands for the class of exponents which are log-Hölder continuous at the origin.

(3) The exponent $r(\cdot)$ satisfies the log-Hölder-type decay condition at infinity if

$$(4.3) \quad |r(x) - r_\infty| \leq \frac{c^*}{\log(e + |x|)} \quad \text{for } x \in \mathbb{R}^n.$$

The class $\operatorname{LH}_\infty = \operatorname{LH}_\infty(\mathbb{R}^n)$ represents the class of exponents that satisfy (4.3).

(4) The class $\operatorname{LH} = \operatorname{LH}(\mathbb{R}^n)$ contains all exponents that satisfy (4.2) and (4.3).

Here c_* , c^* and r_∞ are constants independent of x and y .

For a variable exponent $r(\cdot)$ satisfying $1 < r_- \leq r_+ < \infty$, write $r'(\cdot) = \frac{r(\cdot)}{r(\cdot)-1}$. We employ the definition in [5].

Definition 4.3. Suppose $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$. A weight w is said to be an $A_{p(\cdot)}$ -weight if w satisfies $\sup_{B: \text{open ball}} \frac{1}{|B|} \left\| w^{\frac{1}{p(\cdot)}} \chi_B \right\|_{L^{p(\cdot)}} \left\| w^{-\frac{1}{p'(\cdot)}} \chi_B \right\|_{L^{p'(\cdot)}} < \infty$. The Muckenhoupt class $A_{p(\cdot)}$ adapted to $p(\cdot)$ consists of all $A_{p(\cdot)}$ weights.

Note that this definition extends the classical class A_p . In fact, when $p(\cdot)$ is a constant function p , the two definitions coincide. It is known that M is bounded on $L^{p(\cdot)}(w)$ if $p(\cdot) \in \operatorname{LH}$ satisfies $1 < p_- \leq p_+ < \infty$ and $w \in A_{p(\cdot)}$. Since this condition is symmetric; in this case $p'(\cdot) \in \operatorname{LH}$, $1 < p'_- \leq p'_+ < \infty$ and $w^{-\frac{1}{p'(\cdot)-1}} \in A_{p'(\cdot)}$, Theorem 1.1 is applicable to $L^{p(\cdot)}(w)$. This recaptures a result by [5, 6, 17].

4.5. Herz spaces with variable exponents. Modelled on the previous section, it seems useful to study Herz spaces with variable exponents. The first author [16] has established a method of wavelet characterization applying the boundedness of sublinear operators and norm estimates for characteristic functions with variable exponent. Recently more and more papers have appeared in this direction. In this subsection we establish a new method of wavelet characterization as an application of the extrapolation theorem.

We now set up some notation to recall the definition of Herz spaces with three variable exponents established by the authors' previous work in [20]. Let $p(\cdot)$, $q(\cdot)$ be variable exponents. The space $\ell^{q(\cdot)}(L^{p(\cdot)}(w))$ is the set of all sequences $\{f_j\}_{j \in \mathbb{Z}}$ in $L^0(\mathbb{R}^n)$ such that

$$\|\{f_j\}_{j \in \mathbb{Z}}\|_{\ell^{q(\cdot)}(L^{p(\cdot)}(w))} := \inf \left\{ \mu > 0 : \rho_{\ell^{q(\cdot)}(L^{p(\cdot)}(w))}(\{\mu^{-1} f_j\}_{j \in \mathbb{Z}}) \leq 1 \right\} < \infty,$$

where

$$\rho_{\ell^{q(\cdot)}(L^{p(\cdot)}(w))}(\{\mu^{-1}f_j\}_{j \in \mathbb{Z}}) := \sum_{j=-\infty}^{\infty} \|\mu^{-1}f_j\|_{L^{p(\cdot)/q(\cdot)}(w)}^{q(\cdot)}.$$

We sometimes consider the case where \mathbb{Z} is replaced by \mathbb{N} .

Now, we recall the definition of two-weighted Herz spaces with three variable exponents.

Definition 4.4. Let v, w be weights on \mathbb{R}^n , $p(\cdot), q(\cdot) \in \mathcal{P}_0(\mathbb{R}^n)$ and $\alpha(\cdot) \in L^\infty(\mathbb{R}^n)$.

- (1) The non-homogeneous two-weighted Herz space $K_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(v, w)$ consists of all $f \in L^0(\mathbb{R}^n)$ for which

$$\|f\|_{K_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(v, w)} := \|f\chi_{B_0}\|_{L^{p(\cdot)}(w)} + \left\| \left\{ [v(B_k)]^{\alpha(\cdot)/n} f\chi_{C_k} \right\}_{k=1}^{\infty} \right\|_{\ell^{q(\cdot)}(L^{p(\cdot)}(w))}$$

is finite.

- (2) The homogeneous two-weighted Herz space $\dot{K}_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(v, w)$ consists of all $f \in L^0(\mathbb{R}^n)$ for which

$$\|f\|_{\dot{K}_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(v, w)} := \left\| \left\{ [v(B_k)]^{\alpha(\cdot)/n} f\chi_k \right\}_{k=-\infty}^{\infty} \right\|_{\ell^{q(\cdot)}(L^{p(\cdot)}(w))} < \infty.$$

- (3) In the above, if $v = w = 1$, then we write $\dot{K}_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(\mathbb{R}^n)$ instead of $\dot{K}_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(v, w)$ and omit (v, w) in the norm $\|\cdot\|_{\dot{K}_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(v, w)}$. Analogously, we define $K_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(\mathbb{R}^n)$.

The space $\dot{K}_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(\mathbb{R}^n)$ is called homogeneous (non-weighted) Herz spaces with three variable exponents, while the space $K_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(\mathbb{R}^n)$ is called non-homogeneous (non-weighted) Herz spaces with three variable exponents.

Let $p(\cdot) \in \mathcal{P}(\mathbb{R}^n) \cap \text{LH}(\mathbb{R}^n)$, $q(\cdot) \in \mathcal{P}_0(\mathbb{R}^n) \cap \text{LH}_0(\mathbb{R}^n) \cap \text{LH}_\infty(\mathbb{R}^n)$, $\alpha(\cdot) \in L^\infty(\mathbb{R}^n) \cap \text{LH}_0(\mathbb{R}^n) \cap \text{LH}_\infty(\mathbb{R}^n)$, $v \in A_{p_v}$ for some $p_v \in [1, \infty)$, $w \in A_{p(\cdot)}$. We know that there exist constants $C > 0$ and $\delta \in (0, 1)$ such that

$$(4.4) \quad \frac{v(B)}{v(E)} \leq C \left(\frac{|B|}{|E|} \right)^{p_v},$$

$$(4.5) \quad \frac{v(E)}{v(B)} \leq C \left(\frac{|E|}{|B|} \right)^\delta$$

for all open balls B and all measurable sets $E \subset B$. See [8] for fundamental properties of the Muckenhoupt weights. Define

$$(4.6) \quad v^- := \begin{cases} \delta & \text{if } \alpha_- \geq 0, \\ p_v & \text{if } \alpha_- < 0, \end{cases} \quad v^+ := \begin{cases} p_v & \text{if } \alpha_+ \geq 0, \\ \delta & \text{if } \alpha_+ < 0. \end{cases}$$

In our earlier paper [19], the first and second authors established that there exist positive constants $\delta_1 = \delta_1(w, p(\cdot))$, $\delta_2 = \delta_2(w, p(\cdot))$, $\delta_3 = \delta_3(w, p(\cdot))$, $\delta_4 = \delta_4(w, p(\cdot)) \in (0, 1)$ and $C > 0$ such that

$$(4.7) \quad \frac{\|\chi_k\|_{L^{p(\cdot)}(w)}}{\|\chi_l\|_{L^{p(\cdot)}(w)}} \sim \frac{\|\chi_k\|_{(L^{p'(\cdot)}(w^{-p(\cdot)/p'(\cdot)})')'}}{\|\chi_l\|_{(L^{p'(\cdot)}(w^{-p(\cdot)/p'(\cdot)})')'}} \leq C \left(\frac{|C_k|}{|C_l|} \right)^{\delta_1},$$

that

$$(4.8) \quad \frac{\|\chi_k\|_{(L^{p(\cdot)}(w))'}}{\|\chi_l\|_{(L^{p(\cdot)}(w))'}} \leq C \left(\frac{|C_k|}{|C_l|} \right)^{\delta_2}$$

that

$$(4.9) \quad \chi_{\mathbb{N} \times \mathbb{N}}(k, l) \frac{\|\chi_k\|_{L^{p(\cdot)}(w)}}{\|\chi_l\|_{L^{p(\cdot)}(w)}} \leq C \left(\frac{|C_k|}{|C_l|} \right)^{\delta_3},$$

and that

$$(4.10) \quad \chi_{\mathbb{N} \times \mathbb{N}}(k, l) \frac{\|\chi_k\|_{(L^{p(\cdot)}(w))'}}{\|\chi_l\|_{(L^{p(\cdot)}(w))'}} \leq C \left(\frac{|C_k|}{|C_l|} \right)^{\delta_4}$$

for all $k, l \in \mathbb{Z}$ with $k \leq l$. Finally, assume that $\alpha(\cdot)$ satisfies

$$(4.11) \quad -n\delta_1 < v^- \alpha_-$$

and

$$(4.12) \quad v^+ \alpha_+ < n\delta_2.$$

Then in [20] the authors showed that there exists a constant $C > 0$ such that

$$\|\mathcal{R}_m f\|_{\dot{K}_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(v, w)} \leq C \|f\|_{\dot{K}_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(v, w)}$$

for all $f \in L_C^\infty(\mathbb{R}^n)$ and $m = 1, 2, \dots, n$. We have an analogy to the non-homogeneous space $K_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(v, w)$ if we replace (4.11) and (4.12) by

$$(4.13) \quad -n\delta_3 < v^- \alpha_-$$

and

$$(4.14) \quad v^+ \alpha_+ < n\delta_4.$$

Hence, in this case Theorem 1.1 is applicable to $\dot{K}_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(v, w)$ and $K_{p(\cdot)}^{\alpha(\cdot), q(\cdot)}(v, w)$.

4.6. Orlicz spaces. First, let us recall the notion of Young functions. A function $\Phi : [0, \infty) \rightarrow [0, \infty)$ is a *Young function*, if it satisfies the following conditions:

- (1) $\Phi(0) = 0$.
- (2) Φ is continuous.
- (3) Φ is convex. That is, $\Phi((1-\theta)t_1 + \theta t_2) \leq (1-\theta)\Phi(t_1) + \theta\Phi(t_2)$ for all $t_1, t_2 \in [0, \infty)$ and $0 < \theta < 1$.

Next, we recall the definition of Orlicz spaces which a Young function $\Phi : [0, \infty) \rightarrow [0, \infty)$ generates. We define the *Luxemburg–Nakano norm* $\|\cdot\|_{L^\Phi}$ by

$$\|f\|_{L^\Phi} := \inf \left(\left\{ \lambda \in (0, \infty) : \int_{\mathbb{R}^n} \Phi \left(\frac{f(x)}{\lambda} \right) dx \leq 1 \right\} \cup \{\infty\} \right)$$

for $f \in L^0(\mathbb{R}^n)$. The *Orlicz space* $L^\Phi(\mathbb{R}^n)$ over \mathbb{R}^n is the set of all $f \in L^0(\mathbb{R}^n)$ for which $\|f\|_{L^\Phi}$ is finite.

In order that M is bounded on both $L^\Phi(\mathbb{R}^n)$ and on its Köthe dual, we recall the following classes.

- (1) A function $\varphi : (0, \infty) \rightarrow [0, \infty)$ or $\varphi : [0, \infty) \rightarrow [0, \infty)$ is a doubling function, if there exists a constant $C > 0$, called a *doubling constant*, such that $C^{-1}\varphi(t) \leq \varphi(s) \leq C\varphi(t)$ for all t and s in the domain of φ satisfying $s \leq t \leq 2s$.
- (2) Denote by Δ_2 the set of all convex bijections $\Phi : [0, \infty) \rightarrow [0, \infty)$ satisfying the doubling condition; $\Phi(2r) \lesssim \Phi(r)$ for $r > 0$. The implicit constant is again called a *doubling constant*. In this case Φ also satisfies the Δ_2 -condition.
- (3) The set ∇_2 is the set of all convex bijections $\Phi : [0, \infty) \rightarrow [0, \infty)$ such that there exists $C > 1$ such that $\Phi(2t) \geq 2C\Phi(t)$ for all $t \geq 0$. In this case one says that Φ satisfies the ∇_2 -condition.

If $\Phi \in \Delta_2 \cap \nabla_2$, then M is bounded on both $L^\Phi(\mathbb{R}^n)$ and on its Köthe dual. See [34, Theorem 53] and [34, Example 71]. Thus, Theorem 1.1 is applicable to $L^\Phi(\mathbb{R}^n)$. See [37] for wavelet characterization of Orlicz spaces.

4.7. Morrey spaces. Let $1 \leq r \leq r_0 < \infty$. For an $L^r_{\text{loc}}(\mathbb{R}^n)$ -function f , its (classical) Morrey norm is defined by

$$(4.15) \quad \|f\|_{\mathcal{M}_r^{r_0}} := \sup_{x \in \mathbb{R}^n, R > 0} |B(x, R)|^{\frac{1}{r_0} - \frac{1}{r}} \left(\int_{B(x, R)} |f(y)|^r dy \right)^{\frac{1}{r}}.$$

The Morrey space $\mathcal{M}_r^{r_0}(\mathbb{R}^n)$ is the set of all $f \in L^r_{\text{loc}}(\mathbb{R}^n)$ for which the norm $\|f\|_{\mathcal{M}_r^{r_0}}$ is finite. This is a natural extension of $L^{r_0}(\mathbb{R}^n)$ since $\mathcal{M}_{r_0}^{r_0}(\mathbb{R}^n)$ and $L^{r_0}(\mathbb{R}^n)$ are the same with coincidence of norms.

Let $1 < r \leq r_0 < \infty$. Then M and \mathcal{R}_j , $j = 1, 2, \dots, n$, are bounded on $\mathcal{M}_r^{r_0}(\mathbb{R}^n)$. See [3]. Thus, Theorem 1.1 is applicable to $\mathcal{M}_r^{r_0}(\mathbb{R}^n)$.

Let $1 < r < r_0 < \infty$. Remark that the Morrey space $\mathcal{M}_r^{r_0}(\mathbb{R}^n)$ is not separable but that M is bounded on $\mathcal{M}_r^{r_0}(\mathbb{R}^n)$ [34, Theorem 382] and $\mathcal{M}_r^{r_0}(\mathbb{R}^n)'$ [36]. Thus, in general Morrey spaces do not fall within the scope of Theorem 3.5. Remark that $\mathcal{M}_r^{r_0}(\mathbb{R}^n)$ fails to be reflexive [34, Example 148]. Although $L^{r_0}(\mathbb{R}^n)$ is a subset of $\mathcal{M}_r^{r_0}(\mathbb{R}^n)$, it fails to be dense [34, Theorem 20], it is still possible to obtain the characterization of Morrey spaces in terms of wavelets.

4.8. Besov–Bourgain–Morrey spaces. Let $1 \leq q \leq p < \infty$ and $r, \tau \in [1, \infty]$. The Besov–Bourgain–Morrey space $B\mathcal{M}_{q,r}^{p,\tau}(\mathbb{R}^n)$ is defined to be the set of all the $f \in L^q_{\text{loc}}(\mathbb{R}^n)$ satisfying that

$$\|f\|_{B\mathcal{M}_{q,r}^{p,\tau}(\mathbb{R}^n)} := \left\| \left\{ \left\| \left\{ |Q_{\nu,m}|^{\frac{1}{p} - \frac{1}{q}} \left[\int_{Q_{\nu,m}} |f(y)|^q dy \right]^{\frac{1}{q}} \right\}_{m \in \mathbb{Z}^n} \right\|_{\ell^r(\mathbb{Z}_m^n)} \right\}_{\nu \in \mathbb{Z}} \right\|_{\ell^\tau(\mathbb{Z}_\nu)} < \infty.$$

Here the inner ℓ^r norm is taken with respect to m and the outer ℓ^τ norm is taken with respect to ν . See [38] for more. The Bourgain–Morrey space $\mathcal{M}_{q,r}^p(\mathbb{R}^n)$ is the Besov–Bourgain–Morrey space $B\mathcal{M}_{q,r}^{p,\tau}(\mathbb{R}^n)$ [10]. If $1 < q < p < r < \infty$ and $1 \leq \tau < \infty$ or if $1 < q < p \leq r < \tau = \infty$, then M and \mathcal{R}_j , $j = 1, 2, \dots, n$, are bounded on $B\mathcal{M}_{q,r}^{p,\tau}(\mathbb{R}^n)$. Thus, Theorem 1.1 is applicable to $B\mathcal{M}_{q,r}^{p,\tau}(\mathbb{R}^n)$.

Remark that Besov–Bourgain–Morrey spaces fall within the scope of Theorem 3.5 unlike Morrey spaces.

DECLARATIONS

- Funding: Yoshihiro Sawano was supported by Grant-in-Aid for Scientific Research (C) (19K03546), the Japan Society for the Promotion of Science.
- Conflict of interest/Competing interests (check journal-specific guidelines for which heading to use) Not applicable
- Ethics approval and consent to participate: Not applicable
- Consent for publication: Not applicable
- Data availability: Not applicable
- Materials availability: Not applicable
- Code availability: Not applicable
- Author contribution: The three authors contributed equally to the correctness of this article.

ACKNOWLEDGMENT

This work was partly supported by MEXT Promotion of Distinctive Joint Research Center Program JPMXP0723833165.

REFERENCES

- [1] H. A. Aimar, A. L. Bernardis, and F. J. Martín-Reyes, Multiresolution approximations and wavelet bases of weighted L^p spaces, *J. Fourier Anal. Appl.*, **9** (2003), 497–510. [1]
- [2] J. Bergh and J. Löfström, Interpolation spaces. An introduction. Grundlehren der Mathematischen Wissenschaften, No. 223. Springer-Verlag, Berlin-New York, 1976. [11]
- [3] F. Chiarenza and M. Frasca, Morrey spaces and Hardy–Littlewood maximal function, *Rend. Mat.*, **7** (1987), 273–279. [15]
- [4] S.K. Chua, Extension theorems on weighted Sobolev spaces, *Indiana Univ. Math. J.* **41** (1992), 1027–1076. [10]
- [5] D. Cruz-Uribe, A. Fiorenza, and C. J. Neugebauer, Weighted norm inequalities for the maximal operator on variable Lebesgue spaces, *J. Math. Anal. Appl.*, **394** (2012), 744–760. [12]
- [6] D. Cruz-Uribe, SFO and L.-W. Wang, Extrapolation and weighted norm inequalities in the variable Lebesgue spaces, *Trans. Amer. Math. Soc.*, **369** (2016), 1205–1235. [12]
- [7] G. P. Curbera, J. García-Cuerva, J. M. Martell, and C. Pérez, Extrapolation with weights, rearrangement-invariant function spaces, modular inequalities, and applications to singular integrals, *Adv. Math.*, **203** (2006), 256–318. [3]
- [8] J. Duoandikoetxea, Fourier Analysis, Graduate Studies in Mathematics, vol. 29, American Mathematical Society, Providence, RI, 2001. Translated and revised from the 1995 Spanish original by D. Cruz-Uribe. [13]
- [9] J. García-Cuerva and J. M. Martell, Wavelet characterization of weighted spaces, *J. Geom. Anal.*, **11** (2001), 241–264. [1]
- [10] N. Hatano, T. Nogayama, Y. Sawano, and D. I. Hakim, Bourgain–Morrey spaces and their applications to boundedness of operators, *J. Funct. Anal.*, **284** (2023), Paper No. 109720, 52 pp. [15]
- [11] E. Hernández and G. Weiss, A First Course on Wavelets, CRC Press, Boca Raton, FL., 1996. [7, 9]
- [12] E. Hernández, G. Weiss, and D. Yang, The φ -transform and wavelet characterizations of Herz-type spaces, *Collect. Math.*, **47** (1996), 285–320. [2, 11]
- [13] K.-P. Ho, Generalized Boyds indices and applications, *Analysis (Munich)*, **32** (2012), 97–106. [3]
- [14] M. Izuki, The characterization of weighted Sobolev spaces, *Taiwanese J. Math.* **13** (2009), 467–492. [7, 9]
- [15] M. Izuki, Wavelets and modular inequalities in variable L^p spaces, *Georgian Math. J.*, **15** (2008), 281–293. [2]
- [16] M. Izuki, Boundedness of sublinear operators on Herz spaces with variable exponent and application to wavelet characterization, *Analysis Mathematica*, **36** (2010), 33–50. [2, 12]

- [17] M. Izuki, E. Nakai, and Y. Sawano, Wavelet characterization and modular inequalities for weighted Lebesgue spaces with variable exponent, *Annales Academia Scientiarum Fennica Mathematica*, **40** (2015), 551–571. [2, 12]
- [18] M. Izuki, T. Nogayama, T. Noi and Y. Sawano, Wavelet characterization of local Muckenhoupt weighted Sobolev spaces with variable exponents, *Constructive Approx.* **57** (2022), 161–234. <https://doi.org/10.1007/s00365-022-09573-6> [7, 9]
- [19] M. Izuki and T. Noi, Two weighted Herz spaces with variable exponents, *Bull. Malays. Math. Sci. Soc.*, **43** (2020), No.1, 169–200. [13]
- [20] M. Izuki, T. Noi, and Y. Sawano, Extrapolation to two-weighted Herz spaces with three variable exponents, *Advances in Operator Theory*, **9(2)** (2024), 1–30. <https://doi.org/10.1007/s43036-024-00130-9> [12, 14]
- [21] M. Izuki and K. Tachizawa, Wavelet characterizations of weighted Herz spaces, *Sci. Math. Jpn.*, **67** (2008), 353–363. [2, 11]
- [22] A. Yu. Karlovich, Wavelet bases in Banach function spaces, *Bull. Malays. Math. Sci. Soc.*, **44** (2021), 3, 1669–1689. [9]
- [23] T. S. Kopaliani, Greediness of the wavelet system in $L^{p(t)}(\mathbb{R})$, *East J. Approx.*, **14** (2008), 59–67. [2]
- [24] A. K. Lerner and C. Pérez, A new characterization of the Muckenhoupt A_p weights through an extension of the Lorentz–Shimogaki theorem, *Indiana Univ. Math. J.*, **56** (2007), no. 6, 2697–2722. [3]
- [25] P. G. Lemarié-Rieusset, Ondelettes et poids de Muckenhoupt, *Studia Math.*, **108** (1994), 127–147. [1, 9]
- [26] S. Lu and D. Yang, Oscillatory singular integrals on Hardy spaces associated with Herz spaces, *Proc. Amer. Math. Soc.*, **123** (1995), 1695–1701. [11]
- [27] S. Lu and D. Yang, Some characterizations of weighted Herz-type Hardy spaces and their applications, *Acta Math. Sinica (N.S.)*, **13** (1997), 45–58. [11]
- [28] S. Lu and D. Yang, Boundedness of sublinear operators in Herz spaces on Vilenkin groups and its application, *Math. Nachr.*, **191** (1998), 229–246. [11]
- [29] M. Mastyo and Y. Sawano The Navier–Stokes equation in some general Banach spaces, in preparation. [3, 9]
- [30] Y. Meyer, *Wavelets and Operators*, Cambridge University Press, Cambridge, 1992. [7, 9]
- [31] D. V. Rutsky, A_1 -regularity and boundedness of Riesz transforms in Banach lattices of measurable functions, *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, **447** (2016), 113–122. [2]
- [32] Y. Sawano, *Theory of Besov Spaces*, Springer, 2018. [7]
- [33] Y. Sawano, Factorization of Hardy spaces using the ball Banach spaces, In: Conference Proceedings “The 50, 70, 80, . . . ∞ Conference in Mathematics” (Karlstad, Sweden, 2024), pp. 240–254, Element, Zagreb, 2024. [3]
- [34] Y. Sawano, G. Di Fazio, and D. I. Hakim, *Morrey Spaces. Vol. I. Introduction and applications to integral operators and PDE’s*, Monographs and Research Notes in Mathematics, CRC Press, Boca Raton, FL (2020), 479 pp. ISBN: 978-1-4987-6551-0; 978-0-429-08592-5. [3, 7, 9, 10, 11, 15]
- [35] Y. Sawano, K.-P. Ho, D. Yang and S. Yang, Hardy spaces for ball quasi-Banach function spaces. *Dissertationes Math. (Rozprawy Mat.)* **525**, 1–102 (2017) [2]
- [36] Y. Sawano and H. Tanaka, Predual spaces of Morrey spaces with non-doubling measures, *Tokyo J. Math.* **32** (2009), 471–486. [15]
- [37] P. Soardi, Wavelet bases in rearrangement invariant function spaces, *Proc. Am. Math. Soc.* **125** (1997), 3669–3673. [11, 15]
- [38] Y. Zhao, J. Tao, D. Yang, W. Yuan, and Y. Sawano, Bourgain–Morrey Spaces mixed with structure of Besov spaces, *Proc. Steklov Inst. Math.*, **323** (2024), 244–295. [15]
- [39] C.F. Zhu, D.C. Yang and W. Yuan, Bourgain–Brezis–Mironescu-type characterization of inhomogeneous ball Banach Sobolev spaces on extension domains, *J. Geom. Anal.* **34** (2024), no. 10, Paper No. 295, 70 pp. [4, 7, 10]

FACULTY OF LIBERAL ARTS AND SCIENCES, TOKYO CITY UNIVERSITY, 1-28-1, TAMADUTSUMI SETAGAYA-KU TOKYO 158-8557, JAPAN.

Email address: `izuki@tcu.ac.jp`

OTEMON GAKUIN UNIVERSITY
2-1-15 NISHIAI IBARAKI, OSAKA 567-8502

Email address: `taka.noi.hiro@gmail.com`

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE AND ENGINEERING, CHUO UNIVERSITY, 13-27 KASUGA 1-CHOME, BUNKYOKU, TOKYO, JAPAN

Email address: `yoshihiro-sawano@celery.ocn.ne.jp`