

MINIMUM NUMBERS OF DEHN COLORS OF KNOTS AND \mathcal{R} -PALETTE GRAPHS

ERI MATSUDO, KANAKO OSHIRO, AND GAISHI YAMAGISHI

ABSTRACT. In this paper, we consider minimum numbers of colors of knots for Dehn colorings. In particular, we will show that for any odd prime number p and any Dehn p -colorable knot K , the minimum number of colors for K is at least $\lfloor \log_2 p \rfloor + 2$. Moreover, we will define the \mathcal{R} -palette graph for a set of colors. The \mathcal{R} -palette graphs are quite useful to give candidates of sets of colors which might realize a nontrivially Dehn p -colored diagram. In Appendix, we also prove that for Dehn 5-colorable knot, the minimum number of colors is 4.

INTRODUCTION

In knot theory, minimum numbers of colors for arc colorings have been studied in many papers (see [1, 2, 4, 5, 9, 10, 11] for example). We denote by $\text{mincol}_p^{\text{Fox}}(K)$ the number for a Fox p -colorable knot K . As one of properties for $\text{mincol}_p^{\text{Fox}}(K)$, the following result is obtained in [8].

- For any odd prime number p and any Fox p -colorable knot K ,

$$\text{mincol}_p^{\text{Fox}}(K) \geq \lfloor \log_2 p \rfloor + 2.$$

Note that for the cases when p is a composite number or K is a link with multiple components, the similar results are known under some condition (see [6]).

On the other hand, a *Dehn p -coloring* is well-known as one of *region* colorings of knot diagrams which is corresponding to a Fox p -coloring. However, minimum numbers of colors for Dehn p -colorings almost have not been studied yet.

In this paper, we will consider minimum numbers of colors of Dehn p -colorable knots for an odd prime number p . The number for a Dehn p -colorable knot K is denoted by $\text{mincol}_p^{\text{Dehn}}(K)$. We show in Theorem 1.5 that

- for any odd prime number p and any Dehn p -colorable knot K ,

$$\text{mincol}_p^{\text{Dehn}}(K) \geq \lfloor \log_2 p \rfloor + 2.$$

Moreover, we will define the *\mathcal{R} -palette graph* for a set of colors. The \mathcal{R} -palette graphs are quite useful to give candidates of sets of colors which might realize a nontrivially Dehn p -colored diagram (see Theorem 5.1). In particular, in Theorem 5.3, we give the sets of $\lfloor \log_2 p \rfloor + 2$ colors which might realize a nontrivially Dehn p -colored diagram for an odd prime number p with $p < 2^5$. Furthermore, we show in Proposition 5.4 that there exists a knot K with $\text{mincol}_p^{\text{Dehn}}(K) = \lfloor \log_2 p \rfloor + 2$ for some p .

Date: January 20, 2025.

2020 Mathematics Subject Classification. 57K10, 57K12.

Key words and phrases. Knots, Dehn colorings, Minimum numbers of colors, \mathcal{R} -palette graphs.

This paper is organized as follows: In Section 1, we review the definitions and some basic properties of Dehn colorings and minimum numbers of colors for Dehn colorings. Besides, our main result (Theorem 1.5) is stated in this section. Section 2 is devoted to setting up some extended coloring matrices which are used in Section 3. Theorem 1.5 is proven in Section 3. In Section 4, the \mathcal{R} -palette graph of a set of colors and that of a Dehn p -colored diagram are defined. In Section 5, we give candidates of sets of colors each of which might realize a nontrivially Dehn p -colored diagram by using the \mathcal{R} -palette graphs. Moreover in Appendix A, we show that $\text{mincol}_5^{\text{Dehn}}(K) = 4$ for any Dehn 5-colorable knot K .

1. DEHN COLORINGS AND MINIMUM NUMBERS OF COLORS

In this paper, for a prime number p , we denote by \mathbb{Z}_p the cyclic group $\mathbb{Z}/p\mathbb{Z}$. When $p = 0$, read all the parts of this paper by replacing all \mathbb{Z}_p and p with \mathbb{Z} .

Let p be an odd prime number or $p = 0$. Let D be a diagram of a knot K and $\mathcal{R}(D)$ the set of regions of D . A Dehn p -coloring of D is a map $C : \mathcal{R}(D) \rightarrow \mathbb{Z}_p$ satisfying the following condition:

- for each crossing c with regions x_1, x_2, x_3 , and x_4 as depicted in Figure 1,

$$C(x_1) + C(x_3) = C(x_2) + C(x_4)$$

holds, where the region x_2 is adjacent to x_1 by an under-arc and x_3 is adjacent to x_1 by the over-arc.

We call $C(x)$ the *color* of a region x by C . In this paper, as shown in the right of Figure 1, we represent a Dehn p -coloring C of a knot diagram D by assigning the color $C(x)$ to each region x . We mean by (D, C) a diagram D given a Dehn p -coloring C , and call it a *Dehn p -colored diagram*. We denote by $\mathcal{C}(D, C)$ the set of colors assigned to a region of D by C , that is $\mathcal{C}(D, C) = \text{Im } C$. The set of Dehn p -colorings of D is denoted by $\text{Col}_p(D)$. We remark that the number $\#\text{Col}_p(D)$ is an invariant of the knot K .




FIGURE 1. A crossing on D and the one on (D, C) with $C(x_1) = a_1$, $C(x_2) = a_2$, $C(x_3) = a_3$, $C(x_4) = a_1 - a_2 + a_3$

Let c be a crossing of D with regions x_1, x_2, x_3 , and x_4 as depicted in Figure 1. We say that c of (D, C) is *trivially colored* if

$$C(x_1) = C(x_4), \text{ and } C(x_3) = C(x_2)$$

hold, and *nontrivially colored* otherwise. A Dehn p -coloring C of D is *trivial* if each crossing of (D, C) is trivially colored and *nontrivial* otherwise. We note that trivial p -colorings are separated into two types: monochromatic colorings and checkerboard colorings with two colors. We call the former *1-trivial* colorings and denote them by C^{1T} . We call the latter *2-trivial* colorings and denote them by C^{2T} .

A knot K is *Dehn p -colorable* if K has a Dehn p -colored diagram (D, C) such that C is nontrivial.

For a semiarc u of a Dehn p -colored diagram (D, C) , when the two regions x_1 and x_2 which are on the both side of u have $C(x_1) = a_1$ and $C(x_2) = a_2$ for some $a_1, a_2 \in \mathbb{Z}_p$, we call u an $\{a_1, a_2\}$ -*semiarc*, where $\{a_1, a_2\}$ is regarded as a multiset $\{a_1, a_1\}$ when $a_1 = a_2$. Moreover, we call an arc u including an $\{a_1, a_2\}$ -semiarc an $\overline{a_1 + a_2}$ -*arc*, where we note that the value $\overline{a_1 + a_2}$ does not depend on the choice of a semiarc included in u .

Remark 1.1. Fox colorings of knot diagrams have been well-studied for a long time. There is the p -to-1 correspondence, given in the upper picture of Figure 2, from Dehn colorings to Fox colorings, where $a_1, a_2 \in \mathbb{Z}_p$ represent the colors of regions of a Dehn p -colored diagram, $\overline{a_1 + a_2} \in \mathbb{Z}_p$ represents the colors of arcs of the corresponding Fox p -colored diagram, and we distinguish colors of regions for Dehn p -colorings and those of arcs for Fox p -colorings by putting a bar over a color of an arc as in Figure 2. Then a trivial (resp. nontrivial) Dehn p -coloring corresponds to a trivial (resp. nontrivial) Fox p -coloring as depicted in the lower picture of Figure 2. We also note that a knot K is Dehn p -colorable if and only if K is Fox p -colorable.




FIGURE 2. A correspondence between Dehn colorings and Fox colorings

The set $\text{Col}_p(D)$ can be regarded as a \mathbb{Z}_p -module with the scalar product sC and the addition $C + C'$ with

$$\begin{aligned} (sC) : \mathcal{R}(D) &\rightarrow \mathbb{Z}_p; (sC)(x) = sC(x), \\ (C + C') : \mathcal{R}(D) &\rightarrow \mathbb{Z}_p; (C + C')(x) = C(x) + C'(x). \end{aligned}$$

Moreover, since for any $t \in \mathbb{Z}_p$, the constant map $C_t^{1\text{T}} : \mathcal{R}(D) \rightarrow \mathbb{Z}_p; C_t^{1\text{T}}(x) = t$ is a kind of Dehn p -coloring, the map

$$(sC + t) : \mathcal{R}(D) \rightarrow \mathbb{Z}_p; (sC + t)(x) = (sC + C_t^{1\text{T}})(x) = sC(x) + t,$$

is also a Dehn p -coloring, and thus, the transformations $C \mapsto sC + t$ are closed in $\text{Col}_p(D)$. In particular, a transformation $C \mapsto sC + t$ with $s \in \mathbb{Z}_p^\times$ is called a *regular*

affine transformation on $\text{Col}_p(D)$. Two Dehn p -colorings $C, C' \in \text{Col}_p(D)$ are *affine equivalent* (written $C \sim C'$) if they are related by a regular affine transformation on $\text{Col}_p(D)$, that is, $C' = sC + t$ for some $s \in \mathbb{Z}_p^\times$ and $t \in \mathbb{Z}_p$.

A *regular affine transformation on the \mathbb{Z}_p -module \mathbb{Z}_p* is defined by $a \mapsto sa + t$ for some $s \in \mathbb{Z}_p^\times$ and $t \in \mathbb{Z}_p$. Two subsets $S, S' \subset \mathbb{Z}_p$ are *affine equivalent* (written $S \sim S'$) if they are related by a regular affine transformation on \mathbb{Z}_p , that is, $S' = sS + t$ for some $s \in \mathbb{Z}_p^\times$ and $t \in \mathbb{Z}_p$. We note that since any regular affine transformation is a bijection, $\#S = \#S'$ holds if $S \sim S'$.

Lemma 1.2. (1) *Let $C, C' \in \text{Col}_p(D)$. Then we have*

$$C \sim C' \implies \mathcal{C}(D, C) \sim \mathcal{C}(D, C').$$

Hence we have

$$C \sim C' \implies \#\mathcal{C}(D, C) = \#\mathcal{C}(D, C').$$

(2) *Let $S, S' \subset \mathbb{Z}_p$, and we assume that $S \sim S'$. Then there exists $C \in \text{Col}_p(D)$ such that $\mathcal{C}(D, C) = S$ if and only if there exists $C' \in \text{Col}_p(D)$ such that $\mathcal{C}(D, C') = S'$.*

Proof. (1) A regular affine transformation $C \mapsto C' = sC + t$ on $\text{Col}_p(D)$ induces the regular affine transformation $x \mapsto sx + t$ on \mathbb{Z}_p with $\mathcal{C}(D, C') = s\mathcal{C}(D, C) + t$.

(2) Assume that $S' = sS + t$ holds for some regular affine transformation $x \mapsto sx + t$ on \mathbb{Z}_p . When there exists $C \in \text{Col}_p(D)$ such that $\mathcal{C}(D, C) = S$, we have the Dehn p -coloring $C' = sC + t$. Then $\mathcal{C}(D, C') = s\mathcal{C}(D, C) + t = sS + t = S'$ holds. \square

Remark 1.3. (1) of Lemma 1.2 implies that when we focus on the number of colors used for each nontrivial Dehn p -coloring of a diagram D , we may consider only the number of colors used for a representative of each affine equivalence class for nontrivial Dehn p -colorings of D .

(2) of Lemma 1.2 implies that when we focus on the sets of colors each of which might be used for a nontrivial Dehn p -coloring of a diagram, we may consider only representatives of the affine equivalence classes on the subsets of \mathbb{Z}_p .

In this paper, we focus on the minimum number of colors.

Definition 1.4. The *minimum number of colors* of a knot K for Dehn p -colorings is the minimum number of distinct elements of \mathbb{Z}_p which produce a nontrivially Dehn p -colored diagram of K , that is,

$$\min \left\{ \#\mathcal{C}(D, C) \mid (D, C) \in \left\{ \begin{array}{l} \text{nontrivially Dehn } p\text{-colored} \\ \text{diagrams of } K \end{array} \right\} \right\}.$$

We denote it by $\text{mincol}_p^{\text{Dehn}}(K)$.

The next theorem is our main result, which will be proven in Section 3.

Theorem 1.5. *Let p be an odd prime number. For any Dehn p -colorable knot K , we have*

$$\text{mincol}_p^{\text{Dehn}}(K) \geq \log_2 p + 1,$$

that is,

$$\text{mincol}_p^{\text{Dehn}}(K) \geq \lfloor \log_2 p \rfloor + 2.$$

2. EXTENDED COLORING MATRICES

From now on, for an integer matrix M , $\text{rank}_p M$ means the rank of M regarded as a matrix on \mathbb{Z}_p . Note again that $\text{rank}_0 M$ is replaced with $\text{rank}_{\mathbb{Z}} M$.

Let D be a diagram of a knot K . Let x_1, \dots, x_{n+2} be the regions, and c_1, \dots, c_n the crossings of D . We set an arbitrary orientation for D .

For each crossing c_i , we set the equation

$$x_{i_1} - x_{i_2} + x_{i_3} - x_{i_4} = 0,$$

where $x_{i_1}, x_{i_2}, x_{i_3}, x_{i_4}$ are the regions around c_i such that x_{i_1} is in the right side of both of over- and under-arcs, x_{i_2} is adjacent to x_{i_1} by an under-arc and x_{i_3} is adjacent to x_{i_1} by the over-arc (see the left of Figure 3), and where we regard x_1, \dots, x_{n+2} as variables of the equation. We then have the simultaneous equations

$$\begin{cases} x_{1_1} - x_{1_2} + x_{1_3} - x_{1_4} = 0, \\ \vdots \\ x_{i_1} - x_{i_2} + x_{i_3} - x_{i_4} = 0, \\ \vdots \\ x_{n_1} - x_{n_2} + x_{n_3} - x_{n_4} = 0, \end{cases}$$

that is

$$M_D^{\text{col}} \begin{pmatrix} x_1 \\ \vdots \\ x_{n+2} \end{pmatrix} = \mathbf{0}. \quad (2.1)$$

The coefficient matrix M_D^{col} is called the *Dehn coloring matrix* of D as an unoriented diagram since the \mathbb{Z}_p -module of solutions of (2.1) for an odd prime p or $p = 0$ is isomorphic to the \mathbb{Z}_p -module $\text{Col}_p(D)$ by the isomorphism which sends a solution ${}^t(a_1 \cdots a_{n+2})$ to the Dehn coloring C such that $C(x_i) = a_i$ for each $i \in \{1, \dots, n+2\}$.




FIGURE 3.

It is known that the knot group G_K of K , which is the fundamental group of $\mathbb{R}^3 \setminus K$, has the presentation

$$G_K = \langle x_1, \dots, x_{n+2} \mid r_1, \dots, r_n, x_1 \rangle, \quad (2.2)$$

where for each crossing c_i ($1 \leq i \leq n$) as in the left of Figure 3, we set the relator r_i by

$$r_i = x_{i_3} x_{i_1}^{-1} x_{i_2} x_{i_4}^{-1}.$$

Let $A_D(t)$ denote the Alexander matrix of G_K obtained from the above presentation (2.2) by using the Fox calculus (see [3]), that is $A_D(t) = \left(\tilde{\alpha} \circ \tilde{\pi} \left(\frac{\partial r_i}{\partial x_j} \right) \right)$, where $\frac{\partial}{\partial x_j}$ is the Fox derivative $\mathbb{Z}F \rightarrow \mathbb{Z}F$ with $F = \langle x_1, \dots, x_{n+2} \rangle$, $\tilde{\pi}$ is the projection $\mathbb{Z}F \rightarrow \mathbb{Z}G_K$, and $\tilde{\alpha}$ is the abelianization $\mathbb{Z}G_K \rightarrow \mathbb{Z}\langle t \rangle = \mathbb{Z}[t, t^{-1}]$ such that a meridian element is sent to t . For the i th row vector of $A_D(-1)$ ($1 \leq i \leq n$), the j th entry is 1 if $j = i_1$ or i_3 , -1 if $j = i_2$ or i_4 , and 0 otherwise, which follows from

$$\begin{aligned} \frac{\partial r_i}{\partial x_{i_1}} &= -x_{i_3} x_{i_1}^{-1} \xrightarrow{\tilde{\alpha} \circ \tilde{\pi}} -t^{-(k+1)} t^{-k} = -t^{-2k-1} \xrightarrow{t \mapsto -1} 1, \\ \frac{\partial r_i}{\partial x_{i_2}} &= x_{i_3} x_{i_1}^{-1} \xrightarrow{\tilde{\alpha} \circ \tilde{\pi}} t^{-(k+1)} t^{-k} = t^{-2k-1} \xrightarrow{t \mapsto -1} -1, \\ \frac{\partial r_i}{\partial x_{i_3}} &= 1 \xrightarrow{\tilde{\alpha} \circ \tilde{\pi}} 1 \xrightarrow{t \mapsto -1} 1, \\ \frac{\partial r_i}{\partial x_{i_4}} &= -x_{i_3} x_{i_1}^{-1} x_{i_2} x_{i_4}^{-1} \xrightarrow{\tilde{\alpha} \circ \tilde{\pi}} -1 \xrightarrow{t \mapsto -1} -1 \end{aligned}$$

(refer to the right of Figure 3), where we here suppose that x_{i_1}, \dots, x_{i_4} differ from each other. Thus we have

$$A_D(-1) = \begin{pmatrix} M_D^{\text{col}} \\ \mathbf{1} \quad \mathbf{0} \end{pmatrix}.$$

The greatest common divisor of all the $(n+1)$ -minors of $A_D(-1)$ is called the *determinant* of K , which is denoted by $\det(K)$.

For an odd prime number p , we suppose that there exists a nontrivial Dehn p -coloring, say C_0 , satisfying that for some $i, j \in \{1, \dots, n+2\}$ such that $i < j$, $C_0(x_i) = C_0(x_j)$, $C^{2T}(x_i) \neq C^{2T}(x_j)$ holds. Then we set

$$B_{D, C_0; i, j} = \begin{pmatrix} M_D^{\text{col}} \\ \mathbf{0} \quad 1 \quad \mathbf{0} \quad -1 \quad \mathbf{0} \end{pmatrix},$$

where for the last row vector, the i th and j th entries are 1 and -1 , respectively, and the other entries are 0.

Let us consider the ranks of the matrices M_D^{col} , $A_D(-1)$ and $B_{D, C_0; i, j}$.

For an odd prime p or $p = 0$, we have at least two linearly independent solutions of (2.1) on \mathbb{Z}_p : one, denoted by \mathbf{a}^{1T} , is corresponding to the 1-trivial Dehn \mathbb{Z}_p -coloring C^{1T} with $C^{1T}(x_1) = 1$, and the other, denoted by \mathbf{a}^{2T} , is corresponding to a 2-trivial Dehn \mathbb{Z}_p -coloring C^{2T} with $C^{2T}(x_1) = 0$ and $\text{Im}C^{2T} = \{0, 1\}$. Thus we can see that $\text{rank}_p M_D^{\text{col}} \leq n$.

Besides, it is well-known that $\det(K) \neq 0$, which gives $\text{rank}_{\mathbb{Z}} A_D(-1) = n+1$, and hence, $\text{rank}_{\mathbb{Z}} M_D^{\text{col}} \geq n$. Considering this property together with the fact that $\text{rank}_{\mathbb{Z}} M_D^{\text{col}} \leq n$, we have $\text{rank}_{\mathbb{Z}} M_D^{\text{col}} = n$, and indeed, the \mathbb{Z} -module of solutions of (2.1) on \mathbb{Z} is spanned by two vectors \mathbf{a}^{1T} and \mathbf{a}^{2T} .

On the other hand, for an odd prime p , if D admits a nontrivial Dehn p -coloring C , then there exists a solution \mathbf{a} of (2.1) on \mathbb{Z}_p such that \mathbf{a}^{1T} , \mathbf{a}^{2T} and \mathbf{a} are linearly independent. Hence, for this case, $\text{rank}_p M_D^{\text{col}} \leq n-1$.

For the matrix $A_D(-1)$, $\text{rank}_{\mathbb{Z}} A_D(-1) = n+1$ as mentioned above. For an odd prime p or $p = 0$, note that the \mathbb{Z}_p -module of solutions of the simultaneous

equations

$$A_D(-1) \begin{pmatrix} x_1 \\ \vdots \\ x_{n+2} \end{pmatrix} = \mathbf{0} \quad (2.3)$$

on \mathbb{Z}_p is a submodule of that of (2.1) on \mathbb{Z}_p . Any vector $k\mathbf{a}^{1T} + \ell\mathbf{a}^{2T}$ ($k, \ell \in \mathbb{Z}$, $k \neq 0$) can not be a solution of (2.3) on \mathbb{Z}_p since any solution $\mathbf{a} = {}^t(a_1 \cdots a_{n+2})$ of (2.3) needs to satisfy $a_1 = 0$, while \mathbf{a}^{2T} is a linearly independent solution of (2.3) on \mathbb{Z}_p . This implies that the \mathbb{Z} -module of solutions of (2.3) on \mathbb{Z} is spanned by \mathbf{a}^{2T} .

On the other hand, for an odd prime p , if D admits a nontrivial Dehn p -coloring C such that $C(x_1) = 0$, then there exists a solution \mathbf{a} of (2.3) on \mathbb{Z}_p such that \mathbf{a}^{2T} and \mathbf{a} are linearly independent. Hence, for this case, $\text{rank}_p A_D(-1) \leq n$.

For the matrix $B_{D,C_0;i,j}$, we note that it is defined for a given odd prime p , a Dehn p -coloring C_0 , and $i, j \in \{1, \dots, n+2\}$. Note that the \mathbb{Z} - (resp. \mathbb{Z}_p -)module of solutions of the simultaneous equations

$$B_{D,C_0;i,j} \begin{pmatrix} x_1 \\ \vdots \\ x_{n+2} \end{pmatrix} = \mathbf{0} \quad (2.4)$$

on \mathbb{Z} (resp. \mathbb{Z}_p) is a submodule of that of (2.1) on \mathbb{Z} (resp. \mathbb{Z}_p).

Any vector $k\mathbf{a}^{1T} + \ell\mathbf{a}^{2T}$ ($k, \ell \in \mathbb{Z}$, $\ell \neq 0$) can not be a solution of (2.4) on \mathbb{Z} (resp. \mathbb{Z}_p) since any solution $\mathbf{a} = {}^t(a_1 \cdots a_{n+2})$ of (2.4) needs to satisfy $a_i = a_j$, while \mathbf{a}^{1T} is a linearly independent solution of (2.4) on \mathbb{Z} (resp. \mathbb{Z}_p). Hence the \mathbb{Z} -module of solutions of (2.4) on \mathbb{Z} is spanned by \mathbf{a}^{1T} , which implies that $\text{rank}_{\mathbb{Z}} B_{D,C_0;i,j} = n+1$.

On the other hand, since C_0 is a nontrivial p -coloring, there exists a solution $\mathbf{a}_0 = {}^t(C_0(x_1) \cdots C_0(x_{n+2}))$ of (2.4) on \mathbb{Z}_p , and \mathbf{a}^{1T} and \mathbf{a}_0 are linearly independent. Hence, $\text{rank}_p B_{D,C_0;i,j} \leq n$.

As a summary, we have the following properties:

Lemma 2.1. *Let p be an odd prime number.*

- (1) *Suppose that D admits a nontrivial Dehn p -coloring C . Then we have*

$$\text{rank}_{\mathbb{Z}} M_D^{\text{col}} = n, \text{ and } \text{rank}_p M_D^{\text{col}} \leq n - 1.$$

In particular, the module of solutions of (2.1) on \mathbb{Z} is spanned by \mathbf{a}^{1T} and \mathbf{a}^{2T} , and that on \mathbb{Z}_p is spanned by at least three linearly independent solutions: \mathbf{a}^{1T} , \mathbf{a}^{2T} and $\mathbf{a} = {}^t(C(x_1) \cdots C(x_{n+2}))$.

- (2) *Suppose that D admits a nontrivial Dehn p -coloring C with $C(x_1) = 0$. Then we have*

$$\text{rank}_{\mathbb{Z}} A_D(-1) = n + 1, \text{ and } \text{rank}_p A_D(-1) \leq n.$$

In particular, the module of solutions of (2.3) on \mathbb{Z} is spanned by \mathbf{a}^{2T} , and that on \mathbb{Z}_p is spanned by at least two linearly independent solutions: \mathbf{a}^{2T} and $\mathbf{a} = {}^t(C(x_1) \cdots C(x_{n+2}))$.

- (3) *Suppose that D admits a nontrivial Dehn p -coloring C satisfying that $C(x_i) = C(x_j)$, $C^{2T}(x_i) \neq C^{2T}(x_j)$ for some i, j with $1 \leq i < j \leq n+2$. Then we have*

$$\text{rank}_{\mathbb{Z}} B_{D,C;i,j} = n + 1, \text{ and } \text{rank}_p B_{D,C;i,j} \leq n.$$

In particular, the module of solutions of (2.4) on \mathbb{Z} is spanned by $\mathbf{a}^{1\text{T}}$, and that on \mathbb{Z}_p is spanned by at least two linearly independent solutions: $\mathbf{a}^{1\text{T}}$ and $\mathbf{a} = {}^t(C(x_1) \cdots C(x_{n+2}))$.

3. PROOF OF THEOREM 1.5

Lemma 3.1. *Let M be an integer square matrix of order k which satisfies the following condition:*

- (\star) for each $i \in \{1, \dots, k\}$, the multiset of nonzero entries of the i th row vector of M is $\{-2\}$, $\{-1\}$, $\{1\}$, $\{2\}$, $\{-2, 1\}$, $\{-2, 2\}$, $\{-1, -1\}$, $\{-1, 1\}$, $\{-1, 2\}$, $\{1, 1\}$, $\{-2, 1, 1\}$, $\{-1, -1, 1\}$, $\{-1, -1, 2\}$, $\{-1, 1, 1\}$ or $\{-1, -1, 1, 1\}$.

Then we have

$$|\det M| \leq 2^k.$$

Proof. We prove this lemma by the induction of k and the cofactor expansion of a matrix. In the case that $k = 1$, since $|\det M| = 0, 1$ or 2 , it holds that $|\det M| \leq 2$.

Let $k > 1$. We assume that any square matrix M' , of order $k - 1$, satisfying the condition (\star) has $|\det M'| \leq 2^{k-1}$.

We first note that the following transformations of matrices do not affect whether the condition (\star) is satisfied.

- Compose a matrix $(\mathbf{w}_1, \dots, \mathbf{w}_{j-1}, -\sum_{\ell=1}^k \mathbf{w}_\ell, \mathbf{w}_{j+1}, \dots, \mathbf{w}_k)$ from a matrix $(\mathbf{w}_1, \dots, \mathbf{w}_k)$ satisfying the condition (\star).
- Remove a row and a column from a matrix satisfying the condition (\star).

Put $M = (\mathbf{w}_1, \dots, \mathbf{w}_k)$. We will show by cases.

(i) In the case that M has a row whose multiset of nonzero entries is $\{-1\}$ or $\{1\}$, the cofactor expansion of the row induces $|\det M| \leq 1 \cdot 2^{k-1} < 2^k$.

(ii) In the case that M has a row whose multiset of nonzero entries is $\{-2\}$ or $\{2\}$, the cofactor expansion of the row induces $|\det M| \leq 2 \cdot 2^{k-1} = 2^k$.

(iii) In the case that M has a row whose multiset of nonzero entries is $\{-1, -1\}$, $\{-1, 1\}$ or $\{1, 1\}$, the cofactor expansion of the row induces $|\det M| \leq 2^{k-1} + 2^{k-1} = 2^k$.

(iv) In the case that M has a row \mathbf{v}_i whose multiset of nonzero entries is $\{-2, 1\}$, $\{-1, 2\}$, $\{-2, 1, 1\}$ or $\{-1, -1, 2\}$, assume that the (i, j) -entry is -2 or 2 for some j . Then we can see the determinant as follows.

$$\begin{aligned} |\det M| &= \left| \det (\mathbf{w}_1, \dots, \mathbf{w}_{j-1}, -\sum_{\ell=1}^k \mathbf{w}_\ell, \mathbf{w}_{j+1}, \dots, \mathbf{w}_k) \right| \\ &= \left| \det \begin{pmatrix} \pm 1 & \pm 1 & 0 & \cdots & 0 \\ \mathbf{w}'_1 & \mathbf{w}'_2 & \mathbf{w}'_3 & \cdots & \mathbf{w}'_k \end{pmatrix} \right|. \end{aligned}$$

Thus, this case comes down to the case (iii), and hence, we have $|\det M| \leq 2^k$.

(v) In the case that M has a row \mathbf{v}_i whose multiset of nonzero entries is $\{-2, 2\}$, assume that the (i, j) -entry is -2 or 2 for some j . Then we can see the determinant as follows.

$$\begin{aligned} |\det M| &= \left| \det (\mathbf{w}_1, \dots, \mathbf{w}_{j-1}, -\sum_{\ell=1}^k \mathbf{w}_\ell, \mathbf{w}_{j+1}, \dots, \mathbf{w}_k) \right| \\ &= \left| \det \begin{pmatrix} \pm 2 & 0 & \cdots & 0 \\ \mathbf{w}'_1 & \mathbf{w}'_2 & \cdots & \mathbf{w}'_k \end{pmatrix} \right|. \end{aligned}$$

Thus, this case comes down to the case (ii), and hence, we have $|\det M| \leq 2^k$.

(vi) In the case that M has a row \mathbf{v}_i whose multiset of nonzero entries is $\{-1, -1, 1\}$ or $\{-1, 1, 1\}$, we can see the determinant as follows.

$$\begin{aligned}
|\det M| &= \left| \det \begin{pmatrix} \pm 1 & 1 & -1 & 0 & \cdots & 0 \\ \mathbf{w}'_1 & \mathbf{w}'_2 & \mathbf{w}'_3 & \mathbf{w}'_4 & \cdots & \mathbf{w}'_k \end{pmatrix} \right| \\
&= \left| \det \begin{pmatrix} \pm 1 & 1 & 0 & 0 & \cdots & 0 \\ \mathbf{w}'_1 & \mathbf{w}'_2 & \mathbf{w}'_2 + \mathbf{w}'_3 & \mathbf{w}'_4 & \cdots & \mathbf{w}'_k \end{pmatrix} \right| \\
&\leq \left| \det \begin{pmatrix} \mathbf{w}'_2 & \mathbf{w}'_2 + \mathbf{w}'_3 & \mathbf{w}'_4 & \cdots & \mathbf{w}'_k \end{pmatrix} \right| \\
&\quad + \left| \det \begin{pmatrix} \mathbf{w}'_1 & \mathbf{w}'_2 + \mathbf{w}'_3 & \mathbf{w}'_4 & \cdots & \mathbf{w}'_k \end{pmatrix} \right| \\
&= \left| \det \begin{pmatrix} \mathbf{w}'_2 & \mathbf{w}'_3 & \mathbf{w}'_4 & \cdots & \mathbf{w}'_k \end{pmatrix} \right| \\
&\quad + \left| \det \begin{pmatrix} \mathbf{w}'_1 & -\sum_{\ell=1}^k \mathbf{w}'_\ell & \mathbf{w}'_4 & \cdots & \mathbf{w}'_k \end{pmatrix} \right|
\end{aligned}$$

Since the matrices $(\mathbf{w}'_2 \ \cdots \ \mathbf{w}'_k)$ and $(\mathbf{w}'_1 \ -\sum_{\ell=1}^k \mathbf{w}'_\ell \ \mathbf{w}'_4 \ \cdots \ \mathbf{w}'_k)$ satisfy the condition (\star) , we obtain $|\det M| \leq 2^{k-1} + 2^{k-1} = 2^k$.

(vii) Lastly we consider the case that M has a row \mathbf{v}_i whose multiset of nonzero entries is $\{-1, -1, 1, 1\}$. We then have

$$\begin{aligned}
|\det M| &= \left| \det \begin{pmatrix} 1 & 1 & -1 & -1 & 0 & \cdots & 0 \\ \mathbf{w}'_1 & \mathbf{w}'_2 & \mathbf{w}'_3 & \mathbf{w}'_4 & \mathbf{w}'_5 & \cdots & \mathbf{w}'_k \end{pmatrix} \right| \\
&= \left| \det \begin{pmatrix} 1 & 1 & -1 & 0 & 0 & \cdots & 0 \\ \mathbf{w}'_1 & \mathbf{w}'_2 & \mathbf{w}'_3 & -\sum_{\ell=1}^k \mathbf{w}'_\ell & \mathbf{w}'_5 & \cdots & \mathbf{w}'_k \end{pmatrix} \right|.
\end{aligned}$$

This case comes down to the case (vi), and hence, we have $|\det M| \leq 2^k$.

This completes the proof. \square

Let p be an odd prime integer. Let D be a diagram of a knot K . Let x_1, \dots, x_{n+2} be the regions, and c_1, \dots, c_n the crossings of D . Let C^{1T} and C^{2T} be the 1- and 2-trivial Dehn p -coloring, respectively, such that $C^{1T}(x_1) = 1$, $C^{2T}(x_1) = 0$ and $\text{Im}C^{2T} = \{0, 1\}$. Let C be a nontrivial Dehn p -coloring of D .

Lemma 3.2. *Suppose that for any $i, j \in \{1, \dots, n+2\}$ such that $C(x_i) = C(x_j)$, $C^{2T}(x_i) = C^{2T}(x_j)$ holds. Then we have*

$$\#\mathcal{C}(D, C) \geq \log_2 p + 1.$$

Proof. Put $\ell = \#\mathcal{C}(D, C)$ and $M_1 = A_D(-1)$, where $A_D(-1)$ is the matrix constructed in Section 2 by setting an arbitrary orientation for D . By Lemma 1.2 and Remark 1.3, we may assume that $C(x_1) = 0$ by replacing the Dehn p -coloring C by a regular affine transformation if necessary. Put $\mathbf{a}_1 = {}^t(C(x_1) \cdots C(x_{n+2}))$ and $\mathbf{a}_1^{2T} = {}^t(C^{2T}(x_1) \cdots C^{2T}(x_{n+2}))$. Then by (2) of Lemma 2.1,

$$\text{rank}_{\mathbb{Z}} M_1 = n + 1 \text{ and } \text{rank}_p M_1 \leq n,$$

and the simultaneous equations $M_1 \mathbf{x} = \mathbf{0}$ have the linearly independent solution \mathbf{a}_1^{2T} on \mathbb{Z} , and two linearly independent solutions \mathbf{a}_1^{2T} and \mathbf{a}_1 on \mathbb{Z}_p .

We put $M_1 = (\mathbf{m}_1 \cdots \mathbf{m}_{n+2})$ and $\mathbf{x} = {}^t(x_1 \cdots x_{n+2})$. We transform M_1 , \mathbf{x} , \mathbf{a}_1 and \mathbf{a}_1^{2T} as follows. Suppose that $i, j \in \{1, \dots, n+2\}$ satisfy $i < j$ and $C(x_i) = C(x_j)$. For M_1 , we add \mathbf{m}_j to \mathbf{m}_i and delete \mathbf{m}_j . For \mathbf{x} , \mathbf{a}_1 and \mathbf{a}_1^{2T} , we delete x_j , $C(x_j)$ and $C^{2T}(x_j)$, respectively. We then have the new equation

$$(\mathbf{m}_1 \cdots \mathbf{m}_i + \mathbf{m}_j \cdots \widehat{\mathbf{m}}_j \cdots \mathbf{m}_{n+2}) {}^t(x_1 \cdots x_i \cdots \widehat{x}_j \cdots x_{n+2}) = \mathbf{0},$$

and the solutions

$${}^t(C(x_1) \cdots C(x_i) \cdots \widehat{C(x_j)} \cdots C(x_{n+2}))$$

on \mathbb{Z}_p , and

$${}^t(C^{2\mathbb{T}}(x_1) \cdots C^{2\mathbb{T}}(x_i) \cdots \widehat{C^{2\mathbb{T}}(x_j)} \cdots C^{2\mathbb{T}}(x_{n+2}))$$

on \mathbb{Z} and \mathbb{Z}_p , where \widehat{x} means that the entry x is removed. For the resultant matrix obtained from M_1 and the resultant column vectors obtained from \mathbf{x} , \mathbf{a}_1 and $\mathbf{a}_1^{2\mathbb{T}}$, we repeat the same transformations unless the number of column vectors for the resultant matrix originally from M_1 is ℓ . Then, as a consequence, we have the $n \times \ell$ -matrix M_2 , the column vectors $\mathbf{y} = {}^t(y_1, \dots, y_\ell)$, $\mathbf{a}_2 = {}^t(C(y_1), \dots, C(y_\ell))$ and $\mathbf{a}_2^{2\mathbb{T}} = {}^t(C^{2\mathbb{T}}(y_1), \dots, C^{2\mathbb{T}}(y_\ell))$, where \mathbf{y} is the column vector originally from \mathbf{x} .

Remark that $\ell = \#\mathcal{C}(D, C)$. We can easily see that $\text{rank}_{\mathbb{Z}} M_2 = \ell - 1$ and $\text{rank}_p M_2 \leq \ell - 2$, and $\mathbf{a}_2^{2\mathbb{T}}$ is a linearly independent solution of $M_2 \mathbf{y} = \mathbf{0}$ on \mathbb{Z} , and \mathbf{a}_2 and $\mathbf{a}_2^{2\mathbb{T}}$ are linearly independent solutions on \mathbb{Z}_p . Hence, there exists an $(\ell - 1) \times (\ell - 1)$ -submatrix M_3 of M_2 such that $\det M_3 \neq 0 \in \mathbb{Z}$ and $\det M_3 = 0 \in \mathbb{Z}_p$, which implies that $p \leq |\det M_3|$.

Furthermore, since we have $|\det M_3| \leq 2^{\ell-1}$ by Lemma 3.1, it holds that $p \leq 2^{\ell-1}$. This gives the conclusion $\#\mathcal{C}(D, C) = \ell \geq \log_2 p + 1$. \square

Lemma 3.3. *Suppose that there exists $i, j \in \{1, \dots, n+2\}$ with $i < j$ such that $C(x_i) = C(x_j)$ and $C^{2\mathbb{T}}(x_i) \neq C^{2\mathbb{T}}(x_j)$. Then we have*

$$\#\mathcal{C}(D, C) \geq \log_2 p + 1.$$

Proof. We prove this lemma in much the same way as Lemma 3.2. Put $\ell = \#\mathcal{C}(D, C)$ and $M_1 = B_{D, C; i, j}(-1)$, where $B_{D, C; i, j}(-1)$ is the matrix constructed in Section 2 by setting an arbitrary orientation for D . Put $\mathbf{a}_1 = {}^t(C(x_1) \cdots C(x_{n+2}))$ and $\mathbf{a}_1^{1\mathbb{T}} = {}^t(C^{1\mathbb{T}}(x_1) \cdots C^{1\mathbb{T}}(x_{n+2})) = {}^t(1, \dots, 1)$. Then by (3) of Lemma 2.1,

$$\text{rank}_{\mathbb{Z}} M_1 = n + 1 \text{ and } \text{rank}_p M_1 \leq n,$$

and the simultaneous equations $M_1 \mathbf{x} = \mathbf{0}$ have the linearly independent solution $\mathbf{a}_1^{1\mathbb{T}}$ on \mathbb{Z} , and two linearly independent solutions $\mathbf{a}_1^{1\mathbb{T}}$ and \mathbf{a}_1 on \mathbb{Z}_p .

We put $M_1 = (\mathbf{m}_1 \cdots \mathbf{m}_{n+2})$ and $\mathbf{x} = {}^t(x_1 \cdots x_{n+2})$. We transform M_1 , \mathbf{x} , \mathbf{a}_1 and $\mathbf{a}_1^{1\mathbb{T}}$ as follows. Suppose that $i, j \in \{1, \dots, n+2\}$ satisfy $i < j$ and $C(x_i) = C(x_j)$. For M_1 , we add \mathbf{m}_j to \mathbf{m}_i and delete \mathbf{m}_j . For \mathbf{x} , \mathbf{a}_1 and $\mathbf{a}_1^{1\mathbb{T}}$, we delete x_j , $C(x_j)$ and $C^{1\mathbb{T}}(x_j)$, respectively. We then have the new equation

$$(\mathbf{m}_1 \cdots \mathbf{m}_i + \mathbf{m}_j \cdots \widehat{\mathbf{m}_j} \cdots \mathbf{m}_{n+2}) {}^t(x_1 \cdots x_i \cdots \widehat{x_j} \cdots x_{n+2}) = \mathbf{0},$$

and the solutions

$${}^t(C(x_1) \cdots C(x_i) \cdots \widehat{C(x_j)} \cdots C(x_{n+2}))$$

on \mathbb{Z}_p , and

$${}^t(C^{1\mathbb{T}}(x_1) \cdots C^{1\mathbb{T}}(x_i) \cdots \widehat{C^{1\mathbb{T}}(x_j)} \cdots C^{1\mathbb{T}}(x_{n+2}))$$

on \mathbb{Z} and \mathbb{Z}_p . For the resultant matrix obtained from M_1 and the resultant column vectors obtained from \mathbf{x} , \mathbf{a}_1 and $\mathbf{a}_1^{1\mathbb{T}}$, we repeat the same transformations unless the number of column vectors for the resultant matrix originally from M_1 is ℓ . Then, as a consequence, we have the $n \times \ell$ -matrix M_2 , the column vectors $\mathbf{y} = {}^t(y_1, \dots, y_\ell)$, $\mathbf{a}_2 = {}^t(C(y_1), \dots, C(y_\ell))$ and $\mathbf{a}_2^{1\mathbb{T}} = {}^t(C^{1\mathbb{T}}(y_1), \dots, C^{1\mathbb{T}}(y_\ell))$, where \mathbf{y} is the column vector originally from \mathbf{x} .

We can easily see that $\text{rank}_{\mathbb{Z}} M_2 = \ell - 1$ and $\text{rank}_p M_2 \leq \ell - 2$, and $\mathbf{a}_2^{1\mathbb{T}}$ is a linearly independent solution of $M_2 \mathbf{y} = \mathbf{0}$ on \mathbb{Z} , and \mathbf{a}_2 and $\mathbf{a}_2^{1\mathbb{T}}$ are linearly independent solutions on \mathbb{Z}_p .

Hence, there exists an $(\ell - 1) \times (\ell - 1)$ -submatrix M_3 of M_2 such that $\det M_3 \neq 0 \in \mathbb{Z}$ and $\det M_3 = 0 \in \mathbb{Z}_p$, which implies that $p \leq |\det M_3|$. Furthermore, since we have $|\det M_3| \leq 2^{\ell-1}$ by Lemma 3.1, it holds that $p \leq 2^{\ell-1}$. This gives the conclusion $\#\mathcal{C}(D, C) = \ell \geq \log_2 p + 1$. \square

Proof of Theorem 1.5. From Lemmas 3.2 and 3.3, we have

$$\#\mathcal{C}(D, C) \geq \log_2 p + 1$$

for any nontrivially Dehn p -colored diagram (D, C) of a knot. Therefore, we have

$$\text{mincol}_p^{\text{Dehn}}(K) \geq \log_2 p + 1$$

for any Dehn p -colorable knot K . \square

4. \mathcal{R} -PALETTE GRAPHS

Let p be an odd prime number, and let $S \subset \mathbb{Z}_p$ with $S \neq \emptyset$. Set $\mu(S) = \{\{a_1, a_2\} \mid a_1, a_2 \in S\}$, where $\{a_1, a_2\}$ is regarded as the multiset $\{a_1, a_1\}$ when $a_1 = a_2$.

For $\{a_1, a_2\}, \{a_3, a_4\} \in \mu(S)$, we set

$$\{a_1, a_2\} \sim \{a_3, a_4\} \text{ if } a_1 + a_2 = a_3 + a_4 \text{ in } \mathbb{Z}_p.$$

We note that this relation \sim on $\mu(S)$ is an equivalence relation. We denote by $\overline{a_1 + a_2}$ the equivalence class of $\{a_1, a_2\} \in \mu(S)$.

Definition 4.1. The \mathcal{R} -palette graph of S is the simple graph $G_S = (V_S, E_S)$ composed of the vertex set

$$V_S = \mu(S) / \sim = \{\overline{a_1 + a_2} \mid \{a_1, a_2\} \in \mu(S)\}$$

and the edge set E_S satisfying that

$$e = \overline{b_1 b_2} \in E_S \iff \begin{array}{l} \text{there exist } \{a_1, a_2\} \in \overline{b_1} \text{ and } \{a_3, a_4\} \in \overline{b_2} \text{ such} \\ \text{that } a_1 + a_3 = a_2 + a_4 \text{ or } a_1 + a_4 = a_2 + a_3 \text{ in } \mathbb{Z}_p, \end{array}$$

where $e = \overline{b_1 b_2}$ means that e is an edge connecting the vertices $\overline{b_1}$ and $\overline{b_2}$. We attach the label $\overline{2^{-1}(b_1 + b_2)} \in \mathbb{Z}_p$ to the edge e between $\overline{b_1}$ and $\overline{b_2}$ (see Figure 4 for example).




FIGURE 4. \mathcal{R} -palette graphs

Let G_S be the \mathcal{R} -palette graph of S . A graph $G = (V, E)$ is an \mathcal{R} -subgraph of G_S if G is a subgraph of G_S , and

$$e \in E \implies \overline{b_e} \in V$$

$$\text{(i.e., } e = \overline{b_1 b_2} \in E \implies \overline{2^{-1}(b_1 + b_2)} \in V \text{)}$$

holds, where $\overline{b_e}$ is the label of e (see Figure 5 for example).




FIGURE 5. \mathcal{R} -palette graphs

Let (D, C) be a nontrivially Dehn p -colored diagram of a knot. The \mathcal{R} -palette graph of (D, C) is an \mathcal{R} -subgraph $G_{(D,C)} = (V_{(D,C)}, E_{(D,C)})$ of $G_{\mathcal{C}(D,C)}$ composed of the vertex set

$$V_{(D,C)} = \{\overline{b} \mid \text{there exists a } \overline{b}\text{-arc on } (D, C)\}$$

and the edge set $E_{(D,C)}$ satisfying that

$$e = \overline{b_1 b_2} \in E_{(D,C)} \iff \text{there exists a crossing on } (D, C) \text{ consisting of both the } \overline{b_1}\text{- and } \overline{b_2}\text{-under-arcs.}$$

As in the case of G_S , we attach the label $\overline{2^{-1}(b_1 + b_2)} \in \mathbb{Z}_p$ to the edge e between $\overline{b_1}$ and $\overline{b_2}$ (see Figure 6 for example).




FIGURE 6. The \mathcal{R} -palette graph of (D, C)

5. EVALUATIONS OF $\text{mincol}_p^{\text{Dehn}}(K)$ BY \mathcal{R} -PALETTE GRAPHS

Let p be an odd prime number.

The next theorem can be used to evaluate the minimum number of colors of knots or select some sets of colors, as candidates, each of which might be the set of colors for a nontrivial Dehn p -coloring of a knot diagram.

Theorem 5.1. *Let $S \subset \mathbb{Z}_p$. If $S = \mathcal{C}(D, C)$ for some nontrivially Dehn p -colored diagram (D, C) of a knot, the \mathcal{R} -palette graph G_S includes a connected \mathcal{R} -subgraph with at least three vertices.*

Proof. From the assumption, the \mathcal{R} -palette graph G_S includes the \mathcal{R} -palette graph $G_{(D,C)} = (V_{(D,C)}, E_{(D,C)})$ as an \mathcal{R} -subgraph. Hence we will show that $G_{(D,C)}$ is connected, and $G_{(D,C)}$ includes at least three vertices.

First, let us discuss the connectedness of $G_{(D,C)}$. For any $\bar{b}, \bar{b}' \in V_{(D,C)}$ with $\bar{b} \neq \bar{b}'$, there exist two arcs u and u' of D such that u and u' are a \bar{b} -arc and a \bar{b}' -arc, respectively. Since D is a knot diagram, there exists a finite sequence

$$u = u_1, c_1, u_2, c_2, \dots, c_{n-1}, u_n = u'$$

of arcs u_1, \dots, u_n and crossings c_1, \dots, c_{n-1} of D such that u_i and u_{i+1} are the under-arcs of c_i for each $i \in \{1, \dots, n-1\}$. This implies that there exists a finite sequence

$$\bar{b} = \bar{b}_1, e_1, \bar{b}_2, e_2, \dots, e_{n-1}, \bar{b}_n = \bar{b}'$$

of $\bar{b}_1, \bar{b}_2, \dots, \bar{b}_n \in V_{(D,C)}$ and $e_1, e_2, \dots, e_{n-1} \in E_{(D,C)}$ such that u_i is a \bar{b}_i -arc for each $i \in \{1, \dots, n\}$, and e_i is the edge between \bar{b}_i and \bar{b}_{i+1} for each $i \in \{1, \dots, n-1\}$, where if $\bar{b}_i = \bar{b}_{i+1}$, then $e_i = \emptyset$. Thus, \bar{b} and \bar{b}' can be connected.

Next, let us discuss the reason that $G_{(D,C)}$ includes at least three vertices. Since C is nontrivial, (D, C) includes at least one nontrivially colored crossing c as depicted in the left, the center or the right of Figure 7, where a_1, \dots, a_4 are distinct elements of \mathbb{Z}_p . For the case of the crossing in the left of Figure 7, we have the $\overline{2a_1}$ -, $\overline{a_1 + a_2}$ -, and $\overline{2a_2}$ -arcs whose labels are all distinct. For the case of the crossing in the center of Figure 7, we have the $\overline{a_1 + a_2}$ -, $\overline{2a_1}$ -, and $\overline{a_1 + a_3}$ -arcs whose labels are all distinct. For the case of the crossing in the right of Figure 7, we have the $\overline{a_1 + a_2}$ -, $\overline{a_1 + a_3}$ -, and $\overline{a_3 + a_4}$ -arcs whose labels are all distinct. Thus we have at least three vertices of $G_{(D,C)}$. This completes the proof. \square




FIGURE 7.

Example 5.2. For any subset $S \subset \mathbb{Z}_7$ such that $\#S \leq 3$, the \mathcal{R} -palette graph G_S includes no connected \mathcal{R} -subgraph with at least three vertices. This and Theorem 5.1 imply that $\text{mincol}_7^{\text{Dehn}}(K) \geq 4$ for any Dehn 7-colorable knot K .

Any subset of \mathbb{Z}_7 with four elements is affine equivalent to either $\{0, 1, 2, 3\}$ or $\{0, 1, 2, 4\}$. We note that by Lemma 1.2 and Remark 1.3, when we select candidates of sets of four colors each of which might be the set of colors for a nontrivially Dehn p -colored diagram of a knot, it suffices to discuss only the representatives of the affine equivalence classes, that is $\{0, 1, 2, 3\}$ and $\{0, 1, 2, 4\}$. As shown in the right of Figure 4, the \mathcal{R} -palette graph $G_{\{0,1,2,4\}}$ clearly has a connected \mathcal{R} -subgraph with at least three vertices, which is $G_{\{0,1,2,4\}}$ itself for example. On the other hand, the \mathcal{R} -palette graph $G_{\{0,1,2,3\}}$ depicted in the left of Figure 4 does not include a connected \mathcal{R} -subgraph with at least three vertices. Therefore, if there exists a nontrivially Dehn 7-colored diagram (D, C) for a knot with $\#\mathcal{C}(D, C) = 4 (= \lfloor \log_2 7 \rfloor + 2)$, then $\mathcal{C}(D, C)$ is affine equivalent to $\{0, 1, 2, 4\}$, and moreover, there exists a Dehn 7-coloring C' of D with $C \sim C'$ and $\mathcal{C}(D, C') = \{0, 1, 2, 4\}$.

Similarly as Example 5.2, for each odd prime p with $p < 2^5$, we can give the candidates of $\lfloor \log_2 p \rfloor + 2$ colors by using \mathcal{R} -palette graphs, where $\lfloor \log_2 p \rfloor + 2$ is the lower bound of the evaluation formula of Theorem 1.5. The next theorem gives the results.

Theorem 5.3. *Let p be an odd prime number with $p < 2^5$.*



- (1) *There is no subset $S \subset \mathbb{Z}_p$ with at most $\lfloor \log_2 p \rfloor + 1$ elements such that the \mathcal{R} -palette graph G_S includes a connected \mathcal{R} -subgraph with at least three vertices. This implies that*

$$\text{mincol}_p^{\text{Dehn}}(K) \geq \lfloor \log_2 p \rfloor + 2$$


for any Dehn p -colorable knot K .

- (2) *If there exists a nontrivially Dehn p -colored diagram (D, C) of a knot such that $\#\mathcal{C}(D, C) = \lfloor \log_2 p \rfloor + 2$, then*
- (i) $\mathcal{C}(D, C) \sim \{0, 1, 2\}$ when $p = 3$,
 - (ii) $\mathcal{C}(D, C) \sim \{0, 1, 2, 3\}$ when $p = 5$,
 - (iii) $\mathcal{C}(D, C) \sim \{0, 1, 2, 4\}$ when $p = 7$,
 - (iv) $\mathcal{C}(D, C) \sim \{0, 1, 2, 3, 6\}$ or $\{0, 1, 2, 4, 7\}$ when $p = 11$,
 - (v) $\mathcal{C}(D, C) \sim \{0, 1, 2, 4, 7\}$ when $p = 13$,
 - (vi) $\mathcal{C}(D, C) \sim \{0, 1, 2, 3, 5, 9\}, \{0, 1, 2, 3, 5, 10\}, \{0, 1, 2, 3, 5, 12\}, \{0, 1, 2, 3, 6, 9\},$
 $\{0, 1, 2, 3, 6, 10\}, \{0, 1, 2, 3, 6, 11\}, \{0, 1, 2, 3, 6, 13\}, \{0, 1, 2, 3, 7, 11\}, \{0, 1, 2, 4, 5, 9\},$
 $\{0, 1, 2, 4, 5, 10\}, \{0, 1, 2, 4, 5, 12\},$ or $\{0, 1, 2, 4, 10, 13\}$ when $p = 17$,
 - (vii) $\mathcal{C}(D, C) \sim \{0, 1, 2, 3, 5, 10\}, \{0, 1, 2, 3, 6, 10\}, \{0, 1, 2, 3, 6, 11\}, \{0, 1, 2, 3, 6, 12\},$
 $\{0, 1, 2, 3, 6, 13\}, \{0, 1, 2, 3, 6, 14\}, \{0, 1, 2, 3, 7, 12\}, \{0, 1, 2, 4, 5, 10\}, \{0, 1, 2, 4, 5, 14\},$
 $\{0, 1, 2, 4, 7, 12\},$ or $\{0, 1, 2, 4, 7, 15\}$ when $p = 19$,
 - (viii) $\mathcal{C}(D, C) \sim \{0, 1, 2, 3, 6, 12\}, \{0, 1, 2, 4, 7, 12\}, \{0, 1, 2, 4, 7, 13\}, \{0, 1, 2, 4, 7, 14\},$
 $\{0, 1, 2, 4, 9, 14\},$ or $\{0, 1, 2, 4, 10, 19\}$ when $p = 23$,
 - (ix) $\mathcal{C}(D, C) \sim \{0, 1, 2, 4, 8, 15\}$ when $p = 29$, and
 - (x) $\mathcal{C}(D, C) \sim \{0, 1, 2, 4, 8, 16\}$ when $p = 31$.


Proof. This can be shown by Mathematica computations. In particular, we found a connected \mathcal{R} -subgraph with at least three vertices for the \mathcal{R} -palette graph of each candidate of the sets of colors given in (2) of this theorem as depicted in Figures 8–12. \square

FIGURE 8. Connected \mathcal{R} -subgraphs for $p \in \{3, 5, 7, 11, 13\}$ FIGURE 9. Connected \mathcal{R} -subgraphs for $p \in \{17\}$


$p=17 \{0,1,2,3,6,13\}$




$p=17 \{0,1,2,3,7,11\}$




$p=17 \{0,1,2,4,5,9\}$



$p=17 \{0,1,2,4,5,10\}$



$p=17 \{0,1,2,4,5,12\}$



$p=17 \{0,1,2,4,10,13\}$






FIGURE 10. Connected \mathcal{R} -subgraphs for $p \in \{17\}$

FIGURE 11. Connected \mathcal{R} -subgraphs for $p \in \{19\}$

FIGURE 12. Connected \mathcal{R} -subgraphs for $p \in \{23, 29, 31\}$

We have the following proposition.

Proposition 5.4. *For each odd prime number p with $p < 2^5$ and $p \notin \{13, 29\}$, there exists a Dehn p -colorable knot K with $\text{mincol}_p^{\text{Dehn}}(K) = \lfloor \log_2 p \rfloor + 2$.*

Proof. For $p = 3$ (resp. 5, 7, 11, 17, 19, 23, 31), the diagram illustrated in Figure 13 (resp. 14, 15, 16, 17, 18, 19, 20) admits the Dehn p -coloring with $\lfloor \log_2 p \rfloor + 2$ colors. Hence, from these properties and Theorem 1.5, we can see that there exists a knot K with $\text{mincol}_p^{\text{Dehn}}(K) = \lfloor \log_2 p \rfloor + 2$ for each $p \in \{3, 5, 7, 11, 17, 19, 23, 31\}$.




FIGURE 13. A Dehn 3-colored diagram




FIGURE 14. A Dehn 5-colored diagram




FIGURE 15. A Dehn 7-colored diagram




FIGURE 16. A Dehn 11-colored diagram




FIGURE 17. A Dehn 17-colored diagram




FIGURE 18. A Dehn 19-colored diagram




FIGURE 19. A Dehn 23-colored diagram




FIGURE 20. A Dehn 31-colored diagram

□

Remark 5.5. For $p = 13$ or 29 , there is no Dehn p -colorable knot K with $\text{mincol}_p^{\text{Dehn}}(K) = \lfloor \log_2 p \rfloor + 2$. This result will be shown in our next paper [7].

Remark 5.6. The following properties hold.

- (1) When $p = 3$, any p -colorable knot K has $\text{mincol}_p^{\text{Dehn}}(K) = 3$.
- (2) When $p = 5$, any p -colorable knot K has $\text{mincol}_p^{\text{Dehn}}(K) = 4$.

The proof of (1) is left as an exercise to the reader. We give the proof of (2) in Appendix A.

APPENDIX A.

Proof of Remark 5.6 (2). In this proof, an n -gon means a region bounded by n semiarcs.

Let (D, C) be any nontrivially Dehn 5-colored diagram of a knot. We may construct a Dehn 5-colored diagram (D', C') with four colors by removing regions with the color 4 from (D, C) in the following steps.

- (Step 1) Remove crossings that have four or two regions colored by 4.
- (Step 2) Remove n -gons colored by 4 for $n \geq 4$.
- (Step 3) Remove 3-, and 2-gons colored by 4.

In this proof, suppose that a, b, c are colors with $a \neq b$, $a \neq c$, $a, b, c \neq 4$.

(Step 1) First, we remove crossings that have only the color 4 by the move depicted by Figure 21, where we note that there exists a region colored by some a .




FIGURE 21.

Next, we remove $\{4, 4\}$ -semiarcs by the moves as depicted in Figures 22–26, where the situation around each $\{4, 4\}$ -semiarc is represented by one of Figures 22–26.




FIGURE 22.




FIGURE 23.




FIGURE 24.




FIGURE 25.




FIGURE 26.

Here, we note that

- $2a - 4 \neq 4$ for Figure 22,
- $a - b + 4 \neq 4$ for Figure 23,
- $3 - b, a - b + 4 \neq 4$ for Figure 24,
- $3 - a, 2a + 1, -2a + 2 \neq 4$ for Figure 25,
- $3 - a, 3 - b, a - b + 4, -a + b + 4 \neq 4$ for Figure 26

from $a \neq b$ and $a, b \neq 4$.

Next we delete crossings that have two regions colored by 4s diagonally as the following move depicted in Figure 27. Here we note that there exists a crossing which has four regions colored by a, b, c and 4 near an $\{a, 4\}$ -semiarc.




FIGURE 27.

Here we obtain the color $-a + b + 4$ that is not equal to 4 from $a \neq b$. Thus, for the resultant Dehn 5-colored diagram, there exist no crossings that have four or two regions colored by 4.

(Step 2) For any n -gon, say x , colored by 4 with $n \geq 4$, we focus on an $\{a, 4\}$ -semiarc for some $a \neq 4$, say u , on the boundary of x . As in Figure 28, we move the semiarc u by RII-moves with each semiarc on the boundary of x which is not an $\{a, 4\}$ -semiarc, where we perform the RII-moves as u goes under each semiarc. Here, the 2-gons with no label in the right of Figure 28 are not colored by 4. Thus, we can delete the n -gon with the color 4.




FIGURE 28.

We note that the obtained new regions colored by 4 are 4-, 3-, 2-gons such that the color 4 is not appeared in the neighborhood of the regions. We also note that each 4-gon colored by 4 is as depicted in the left of Figure 29.

Next we delete 4-gons colored by 4 as depicted in Figure 29 if $b \neq c$, Figure 30 if $b = c, a \neq 2b + 1$ and Figure 31 if $b = c, a = 2b + 1$.




FIGURE 29.




FIGURE 30.




FIGURE 31.

Here, we note that

$$b - c + 4 \neq 4 \text{ for Figure 29,}$$

$$-a + b + 4, -a + 2b, -2a + 2b + 4 \neq 4 \text{ for Figure 30, and}$$

$$-b + 3, 2b + 1, 3b + 2 \neq 4 \text{ for Figure 31}$$

from $b \neq c, a \neq b; a \neq 2b + 1, \text{ and } b \neq 4$, respectively, and the 2-gons with no label in the right of Figures 30 and 31 are not colored by 4.

Thus, for the resultant Dehn 5-colored diagram, each region colored by 4 is a 3- or 2-gon around which the color 4 is not appeared.

(Step 3) Now we consider 3-gons colored by 4, where we note that such 3-gons are as depicted in the left of Figure 32, the left of Figure 33 or the mirror images for some distinct a, b, c .

We first remove 3-gons as depicted in the left of Figure 32 by the move shown in Figure 32.




FIGURE 32.

Here, we note that

$$-a + b + 4, -b + c + 4, a - c + 4, -a + c + 4 \neq 4$$

because a, b, c are distinct. Although the new 3-gon labeled by $a - b + c$ might be colored by 4, by considering the case of 3-gons in the left of Figure 33 or the mirror image, the 3-gon with the color 4 can be removed. As for the mirror images of the 3-gons in the left of Figure 32, they are also removed by taking the mirror of the same move.

Next we remove 3-gons as depicted in the left of Figure 33 by the move shown in Figure 33 if $a - b + c \neq 4$ and Figure 34 if $a - b + c = 4$.




FIGURE 33.




FIGURE 34.

Here, we note

$$\begin{aligned} & -a + b + 4, -b + c + 4, -a + c + 4 \neq 4 \text{ for Figure 33,} \\ & -a + c + 4, a + c + 1, -a + 3, 2a + 1 \neq 4 \text{ for Figure 34} \end{aligned}$$

because a, b, c are distinct and $a, b, c \neq 4$. As for the mirror images of the 3-gons in the left of Figure 33, they are also removed by taking the mirror of the same moves.

Next let us consider the case of 2-gons colored by 4, where we note that such 2-gons are as depicted in the left of Figure 35, the left of Figure 36 or the mirror image for some a, b .

The 2-gons as depicted in the left of Figure 35 can be deleted by the move in Figure 35.




FIGURE 35.

Here, note that

$$-a + b + 4 \neq 4$$

from $a \neq b$. As for the 2-gons as depicted in the left of Figure 36, we remove them by the move as shown in Figure 36 if $-a + 2b \neq 4$ and $-2a + 3b \neq 4$, Figure 37 if $-a + 2b = 4$ (and $-2a + 3b \neq 4$), and Figure 38 if $-2a + 3b = 4$ (and $-a + 2b \neq 4$).




FIGURE 36.




FIGURE 37.




FIGURE 38.

Here, we note that

$$\begin{aligned}
 & -a + b + 4, a - b + 4, -2a + 2b + 4 \neq 4 \text{ for Figure 36,} \\
 & 2a + 1, 3a + 2, 4a + 3 \neq 4 \text{ for Figure 37, and} \\
 & 3a + 2, 4a + 3, 2a + 1 \neq 4 \text{ for Figure 38}
 \end{aligned}$$

because a, b, c are distinct and $a, b, c \neq 4$. As for the mirror images of the 2-gons in the left of Figure 36, they are also removed by taking the mirror of the same moves.

The resultant diagram has no region colored by 4. This completes the proof. \square

ACKNOWLEDGMENTS

The authors wish to express their thanks to Professor Shin Satoh for several helpful comments on the property $\text{mincol}_5^{\text{Dehn}}(K) = 4$. The second author was supported by JSPS KAKENHI Grant Number 21K03233.

REFERENCES

- [1] H. Abchir, M. Elhamdadi and S. Lamsifer, *On the minimum number of Fox colorings of knots*, Grad. J. Math. **5** (2020), no. 2, 122–137.
- [2] F. Bento and P. Lopes, *The minimum number of Fox colors modulo 13 is 5*, Topology Appl. **216** (2017), 85–115.
- [3] R. H. Crowell and R. H. Fox, *Introduction to knot theory*, Graduate Texts in Mathematics, No. 57. Springer-Verlag, New York-Heidelberg, 1977. x+182 pp.
- [4] Y. Han and B. Zhou, *The minimum number of coloring of knots*, J. Knot Theory Ramifications **31** (2022), no. 2, Paper No. 2250013, 55 pp.
- [5] F. Harary and L. H. Kauffman, *Knots and graphs. I. Arc graphs and colorings*, Adv. Appl. Math. **22**(3) (1999), 312–337.
- [6] K. Ichihara and E. Matsudo, *A lower bound on minimal number of colors for links*, Kobe J. Math. **33** (2016), no. 1-2, 53–60.
- [7] E. Matsudo, K. Oshiro and G. Yamagishi, *Minimum numbers of Dehn colors of knots and symmetric local biquandle cocycle invariants*, preprint.
- [8] T. Nakamura, Y. Nakanishi, and S. Satoh, *The pallet graph of a Fox coloring*, Yokohama Math. J. **59** (2013), 91–97.
- [9] T. Nakamura, Y. Nakanishi, and S. Satoh, *11-colored knot diagram with five colors*, J. Knot Theory Ramifications **25** (2016), no. 4, Paper No. 1650017, 22 pp.
- [10] K. Oshiro, *Any 7-colorable knot can be colored by four colors*, J. Math. Soc. Japan **62**(3) (2010) 963–973.
- [11] S. Satoh, *5-colored knot diagram with four colors*, Osaka J. Math. **46** (2009), no. 4, 939–948.

THE INSTITUTE OF NATURAL SCIENCES, NIHON UNIVERSITY, 3-25-40 SAKURAJOSUI, SETAGAYA-KU, TOKYO 156-8550, JAPAN

Email address: matsudo.eri@nihon-u.ac.jp

DEPARTMENT OF INFORMATION AND COMMUNICATION SCIENCES, SOPHIA UNIVERSITY, TOKYO 102-8554, JAPAN

Email address: oshirok@sophia.ac.jp

Email address: g-yamagishi-3c9@eagle.sophia.ac.jp