

Fully viable DHOST bounce with extra scalar

Ok Song An, Jin U Kang, Yong Jin Kim, Ui Ri Mun and Un Gyong Ri

Department of Physics, Kim Il Sung University, Ryongnam Dong, Taesong District, Pyongyang, Democratic People's Republic of Korea

E-mail: os.an@ryongnamsan.edu.kp, ju.kang0718@ryongnamsan.edu.kp,
cioc6@ryongnamsan.edu.kp, ur.mun0826@ryongnamsan.edu.kp,
uk.ri0107@ryongnamsan.edu.kp

ABSTRACT: In this paper we construct a class of Degenerate Higher-Order Scalar-Tensor (DHOST) theories with an extra scalar field, which admits viable solutions of bouncing universe satisfying the following requirements: (i) absence of Belinski-Khalatnikov-Lifshitz (BKL) instability, ghost and gradient instability, (ii) absence of superluminality, (iii) generation of nearly scale-invariant curvature perturbations and very small tensor-to-scalar ratio, and (iv) conventional asymptotics in the distant past and future, where gravity sector is described by General Relativity and the DHOST scalar has a canonical form of Lagrangian. We also expect our models to have sufficiently small non-Gaussianities of primordial curvature perturbations to be compatible with observations. As such, this work exemplifies for the first time the fully viable two-field DHOST bouncing cosmology, which is free of instability and superluminality problems as well as compatible with observations.

Contents

1	Introduction	1
2	Exceptional Subclass of DHOST Ia with Extra Scalar	4
2.1	DHOST Ia with extra scalar	4
2.2	Quadratic action for perturbations	5
2.3	Stabilities and non-superluminality conditions: an exceptional subclass	7
2.4	Sufficient conditions for non-superluminality	8
3	DHOST Sector	9
3.1	Asymptotics and ansatz	10
3.2	Stability and non-superluminality	12
4	Extra Scalar and Comparison with Observations	16
4.1	Setup	16
4.2	Scalar spectral index for entropic perturbations	17
4.3	Conversion of entropic perturbations into curvature fluctuations	18
4.4	The amplitude of scalar power spectrum and characteristic energy scale	23
5	Summary and Conclusion	25
A	Background equations of motion	29
B	Exceptional Subclass	29

1 Introduction

The inflationary scenario [1–3] is the mainstream in studies of the early universe. Although it can solve many of the theoretical and phenomenological problems of the standard Big Bang cosmology, the initial singularity problem still remains unsolved [4]. Therefore, it deserves to study alternatives to the inflation.

The most promising alternative that can solve the initial singularity problem is a non-singular bouncing scenario (see [5–7] for a review), where the universe is supposed to undergo a smooth transition from the initial contracting phase to the currently expanding phase. In the non-singular bouncing scenario the Hubble parameter has to grow during the bounce period, which within the framework of General Relativity (GR) is possible only if there exists an exotic matter component that does not respect the Null Energy Condition (NEC). Here the problem is that it is challenging to realize the NEC violation without pathologies such as ghost and gradient instabilities and superluminality [8, 9].

The infamous no-go theorems [10, 11] tell us that in Horndeski theories [12] it is impossible for non-singular bouncing cosmologies not only to avoid ghost and gradient instabilities but also to have conventional asymptotics (so that strong coupling problem does not arise in the distant past and future). However, it turns out that beyond Horndeski [13, 14] and Degenerate Higher-Order Scalar-Tensor (DHOST) theories [15–17] (see [18, 19] for a review) admit non-singular bounce cosmology without ghost and gradient instabilities and with conventional asymptotics (see e.g. [20–25]).¹

An important yet unresolved issue is *superluminality*, which is another undesirable feature of low energy effective field theory, since it indicates that such a theory can not be UV-completed in a Lorentz-covariant way [26]. Beyond-Horndeski theories admit non-singular bouncing scenario without ghost and gradient instabilities and superluminality and with conventional asymptotics [27]. It was, however, shown in [28, 29] that superluminality issue reappears once we add to the theory an extra scalar field minimally coupled to gravity which has the non-zero kinetic energy and the luminal sound speed. This superluminality problem can be resolved by either assuming *strong gravity* in the asymptotic past and realizing the violation of NEC within Horndeski theories [30–35] or considering an exceptional subclass of DHOST Ia theories without the assumption of strong gravity as argued in [36, 37]. In general, strong gravity in the past causes a fast growth of quantum fluctuations, which implies that classical description of the model is no longer consistent. Although there exists a region in parameter space where this strong coupling problem does not occur [34], the bounce scenario with strong gravity in the past generates primordial scalar perturbations with large non-Gaussianities exceeding the observational constraints [38]. On the other hand, to our best knowledge, no explicit model without superluminality and consistent with observations in DHOST Ia theories has been constructed so far.

The aim of the this paper is to build a *novel* explicit bouncing scenario within the exceptional subclass of DHOST Ia theories such that

- it is completely free from the aforementioned issues, i.e., ghost and gradient instabilities, superluminality without assuming strong gravity in the past and future,
- it is observationally viable in the sense that it predicts the power spectrum of perturbations compatible with observations.

In order to seek such a healthy bouncing model within the exceptional subclass of DHOST Ia theories which does not suffer from any pathologies, we employ the inverse method used in [10, 21, 27, 39–41]. In addition, to make the model observationally viable, we should introduce an extra scalar. The reasoning for this deserves a detailed discussion below.

One serious issue that has to be addressed in constructing a bounce scenario is the so-called Belinsky-Khalatnikov-Lifshitz (BKL) instability [42], that is, the contracting phase is unstable due to the growth of anisotropies. In order to suppress the BKL instability within GR one needs to introduce a matter component with a steep equation of state that drives an ekpyrotic contraction phase (also known as a slow contraction phase) [43–45],

¹Although they contain higher-derivative terms, these general scalar-tensor theories do not possess Ostrogradski instability by construction.

which turns out to be a robust and rapid smoother even for initial conditions far from being spatially-flat, homogeneous and isotropic [46–49]. In a word, the bounce scenario should begin with the ekpyrotic contraction phase in order for the BKL instability to be absent.

The problem is that the ekpyrotic contraction makes all perturbations blue-tilted, which is incompatible with the observational data that the curvature perturbations have nearly scale-invariant power spectrum [50]. A typical resolution to this issue is to utilize the entropic mechanism, which consists of two steps: (i) adding an extra scalar field χ that generates nearly scale-invariant perturbations during the ekpyrotic contraction phase and (ii) converting the entropic perturbations into curvature fluctuations around the bounce phase. It is now manifest that the healthy and phenomenologically viable bouncing scenario with conventional asymptotics should consist of two fields, where the adiabatic field ϕ is described by the exceptional subclass of DHOST Ia theories.

As for the entropic field χ , we incorporate the entropic mechanism successfully with the exceptional subclass of DHOST Ia theories. Regarding the first step of the entropic mechanism, two classes of the scalar field χ have been studied in the literature: the entropic field χ has a canonical kinetic term and an unstable potential [51–53], or the entropic field χ has a non-minimal kinetic coupling to the ekpyrotic scalar field [54–57]. In either case, the sound speed of χ is equal to unity. The first class of models has a drawback that the potential has a tachyonic instability and thus the ekpyrotic contraction phase would not last long enough unless a special fine-tuned initial conditions are provided [58]. On the other hand, the second class of models is stable and does not suffer from fine-tuning of initial conditions. For this reason, we choose the second class for generating a nearly scale-invariant power spectrum.

In the second step of the entropic mechanism the conversion can be done either by a repulsive potential [53, 59–62] or by a higher-order kinetic coupling of χ , where the sound speed of χ becomes less than unity during the conversion process [63, 64]. In any case, the conversion is induced by a bending of the background trajectory in the field space. Here, the key requisite is that the conversion process should be efficient and smooth; otherwise the resulting curvature perturbations have a small amplitude and may have very large non-Gaussianities exceeding the current observational bounds, see [56, 61, 62] and references therein.

It is well-known that it is rather non-trivial to obtain an efficient conversion process [61, 62], which becomes more challenging in our case due to the stability constraint. This is because the efficient conversion requires the total bending almost unity (see the argument of [56, 61]), which can make the background trajectory get out of the stable region in the field space. In this paper we realize an efficient conversion process in terms of the entropic field with luminal sound speed. It must be noted, though, that all prior works related to conversion process have focused only on efficiency and non-Gaussianity but have not been concerned with the total background trajectory in the field space. In this paper we pay a careful attention to this point, namely we obtain an efficient conversion process maintaining the stability and non-superluminality throughout the whole background trajectory. By construction, our model is expected to have sufficiently small non-Gaussianities of primor-

dial curvature perturbations. We again emphasize that this work exemplifies for the first time the fully viable two-field DHOST bouncing cosmology, which is free of instability and superluminality problems as well as compatible with observations.

The rest of this paper is organized as follows. In Section 2 we briefly review the DHOST Ia theories with an extra scalar field, in relation to the no-go theorem [36, 37], and set up the theoretical framework and strategy employed in this paper. In Section 3 we consider the decoupled case (i.e., with unrolling extra scalar) and construct a stable and non-superluminal model in the DHOST sector. In Section 4 we consider the coupled case (i.e., with rolling χ) and construct the extra scalar sector such that the stability and non-superluminality are not spoiled and the conversion from entropy to curvature perturbations is efficient enough to be compatible with observations. We summarize the result and conclude in Section 5. Throughout this paper we use the $(+, -, -, -)$ signature of the metric and natural units with $c = \hbar = 1$. We define the reduced Planck mass as $M_{pl} \equiv 1/\sqrt{8\pi G_N} = 2.4 \times 10^{18} \text{GeV}$.

2 Exceptional Subclass of DHOST Ia with Extra Scalar

In this section we briefly describe an exceptional subclass of DHOST Ia theories with an extra scalar following the exposition in [36], with a new ingredient related to the superluminality issue which motivates our work.

2.1 DHOST Ia with extra scalar

The action of DHOST Ia theories with an extra scalar field χ is given by

$$\mathcal{S} = \mathcal{S}_\phi + \mathcal{S}_\chi \quad (2.1)$$

where \mathcal{S}_ϕ is the quadratic action of the DHOST theory [15, 17]

$$\mathcal{S}_\phi = \int d^4x \sqrt{-g} \left(F(\phi, X) + K(\phi, X) \square \phi + F_2(\phi, X) R + \sum_{I=1}^5 A_I(\phi, X) L_I \right), \quad (2.2a)$$

and \mathcal{S}_χ has the k-essence type [65]

$$\mathcal{S}_\chi = \int d^4x \sqrt{-g} P(\chi, Y, \phi). \quad (2.2b)$$

Here $\phi_{,\mu} \equiv \partial_\mu \phi$, $X \equiv g^{\mu\nu} \phi_{,\mu} \phi_{,\nu}$, $\chi_{,\mu} \equiv \partial_\mu \chi$, $Y \equiv g^{\mu\nu} \chi_{,\mu} \chi_{,\nu}$, and L_I ($I = 1, 2, 3, 4, 5$) are

$$\begin{aligned} L_1 &= \phi_{;\mu\nu} \phi^{;\mu\nu}, & L_2 &= (\square \phi)^2, & L_3 &= \phi^{;\mu} \phi_{;\mu\nu} \phi^{;\nu} \square \phi, \\ L_4 &= \phi^{;\mu} \phi_{;\mu\nu} \phi^{;\nu\rho} \phi_{,\rho}, & L_5 &= (\phi^{;\mu} \phi_{;\mu\nu} \phi^{;\nu})^2, \end{aligned}$$

where $\phi_{;\mu\nu} \equiv \nabla_\mu \nabla_\nu \phi$. The DHOST Ia theories are characterized by

$$\begin{aligned} A_2 &= -A_1, \\ A_4 &= \frac{1}{8(F_2 - X A_1)^2} [-16X A_1^3 + 4(3F_2 + 16X F_{2X}) A_1^2 - X^2 F_2 A_3^2] \end{aligned} \quad (2.3a)$$

$$\begin{aligned}
& - (16X^2 F_{2X} - 12X F_2) A_3 A_1 - 16F_{2X} (3F_2 + 4X F_{2X}) A_1 \\
& + 8F_2 (X F_{2X} - F_2) A_3 + 48F_2 F_{2X}^2 \Big]. \tag{2.3b}
\end{aligned}$$

$$A_5 = \frac{(4F_{2X} - 2A_1 + X A_3)(-2A_1^2 - 3X A_1 A_3 + 4F_{2X} A_1 + 4F_2 A_3)}{8(F_2 - X A_1)^2}, \tag{2.3c}$$

where the subscript X denotes the derivative with respect to X . We emphasize that the coupling of χ to the DHOST theory does not spoil the degeneracy, as the scalar field χ does not feature higher derivatives and the coupling terms do not involve derivatives of ϕ [37, 66]. Hence, the whole system does not suffer from the Ostrogradsky ghost.

2.2 Quadratic action for perturbations

In order to deal with the instability and superluminality issue we go to the perturbation level. The perturbed spatially-flat Friedmann-Lemaître-Robertson-Walker (FLRW) metric is given by

$$ds^2 = (1 + \alpha)^2 dt^2 - a^2 e^{2\zeta} \left(\delta_{ij} + h_{ij}^T + \frac{1}{2} h_{ik}^T h_j^k{}^T \right) (dx^i + \partial^i \beta dt)(dx^j + \partial^j \beta dt), \tag{2.4}$$

where α, β and ζ denote lapse, shift and curvature perturbations, respectively, and h_{ij}^T is traceless and transverse perturbation. Throughout this paper we work in the unitary gauge $\delta\phi = 0$. The lapse and shift perturbations α and β appear in the quadratic action as Lagrange multipliers, so they can be integrated out. After all, we obtain a quadratic action $S^{(2)} = S_{\text{tensor}}^{(2)} + S_{\text{scalar}}^{(2)}$ for the perturbations h_{ij}^T, ζ and $\delta\chi$, where

$$S_{\text{tensor}}^{(2)} = \int dt d^3x a^3 \left(\frac{\mathcal{G}_{\mathcal{T}}}{8} (\dot{h}_{ij}^T)^2 - \frac{\mathcal{F}_{\mathcal{T}}}{8a^2} (\partial_k h_{ij}^T)^2 \right), \tag{2.5}$$

with

$$\mathcal{G}_{\mathcal{T}} = -2F_2 + 2A_1 X, \tag{2.6a}$$

$$\mathcal{F}_{\mathcal{T}} = -2F_2. \tag{2.6b}$$

Here the dot notation denotes the derivative with respect to time t . Note that perturbation $\delta\chi$ of the extra scalar χ does not appear in the quadratic action for tensor perturbations, but appears only in the quadratic action $S_{\text{scalar}}^{(2)}$ for the scalar perturbations ζ and $\delta\chi$ given by [36, 37]

$$S_{\text{scalar}}^{(2)} = \int dt d^3x a^3 \left(G_{AB} \dot{v}^A \dot{v}^B - \frac{1}{a^2} F_{AB} \delta^{ij} \partial_i v^A \partial_j v^B + \Psi_1 \dot{\zeta} \delta\chi + \Psi_2 \delta\chi^2 \right). \tag{2.7}$$

Here $A, B = 1, 2$, and v^1 and v^2 denote $\tilde{\zeta} \equiv \zeta + \frac{\Gamma}{2\mathcal{G}_{\mathcal{T}}} \alpha$ and $\delta\chi$, respectively, where

$$\Gamma = X(-2A_1 + 4F_{2X} + A_3 X). \tag{2.8}$$

The functions $\Psi_{1,2}$ and the matrices G_{AB} and F_{AB} are given by respectively

$$\Psi_1 = \frac{\mathcal{G}_{\mathcal{T}}}{\Theta^2} \left[2(YQ + \Sigma) \dot{\chi} P_Y + \Theta \left(P_\chi - 2Y P_{\chi Y} + \frac{6\Delta}{a^3} \frac{d}{dt} (a^3 \dot{\chi} P_Y) \right) \right], \tag{2.9a}$$

$$\Psi_2 = \Omega + \frac{1}{a^3} \frac{d}{dt} \left(\frac{a^3 Y P_Y Q \left(1 - 3 \frac{P_Y}{Q} \Delta\right)}{\Theta} \right) + \frac{\dot{\chi} P_Y}{\Theta^2} \left[\dot{\chi} P_Y (Y Q + \Sigma) + \Theta \left(P_\chi - 2Y P_{\chi Y} + \frac{6\Delta}{a^3} \frac{d}{dt} (a^3 \dot{\chi} P_Y) \right) \right], \quad (2.9b)$$

$$G_{AB} = \begin{pmatrix} \mathcal{G}_S + \frac{g_T^2}{\Theta^2} Y Q & \dot{\chi} Q g \\ \dot{\chi} Q g & Q \end{pmatrix}, \quad F_{AB} = \begin{pmatrix} \mathcal{F}_S & \dot{\chi} P_Y f \\ \dot{\chi} P_Y f & P_Y \end{pmatrix}, \quad (2.9c)$$

where the subscripts χ and Y denote derivative with respect to χ and Y , respectively. Notice that in the case of $\dot{\chi} = 0$ the matrix G_{AB} and F_{AB} become diagonal so that the DHOST sector decouples from χ sector. We also introduced the new functions ²

$$\Omega = \frac{1}{2} P_{\chi\chi} - \frac{1}{a^3} \frac{d}{dt} (a^3 \dot{\chi} P_{\chi Y}), \quad (2.10a)$$

$$f = - \frac{\mathcal{G}_T + \mathcal{D}\dot{\phi} + \mathcal{F}_T \Delta}{\Theta}, \quad (2.10b)$$

$$g = - \frac{\mathcal{G}_T}{\Theta} \left(1 - 3 \frac{P_Y}{Q} \Delta \right), \quad (2.10c)$$

$$\mathcal{G}_S = \frac{\Sigma \mathcal{G}_T^2}{\Theta^2} + 3\mathcal{G}_T, \quad (2.10d)$$

$$\mathcal{F}_S = \frac{1}{a} \frac{d}{dt} \left(\frac{a \mathcal{G}_T (\mathcal{G}_T + \mathcal{D}\dot{\phi} + \mathcal{F}_T \Delta)}{\Theta} \right) - \mathcal{F}_T, \quad (2.10e)$$

$$\mathcal{D} = 2\dot{\phi} (F_{2X} - A_1), \quad (2.10f)$$

$$\Delta = \frac{X}{2\mathcal{G}_T} (2A_1 - 4F_{2X} - A_3 X), \quad (2.10g)$$

$$\begin{aligned} \Theta = & -\mathcal{G}_T \dot{\Delta} - 2F_2 H + \dot{\phi} \left(3\ddot{\phi} (A_1 - 2F_{2X}) - F_{2\phi} \right) + H \dot{\phi}^2 (3A_1 + 2F_{2X}) \\ & + \dot{\phi}^3 \left(\frac{1}{2} \ddot{\phi} (4A_{1X} - 3A_3 + 2A_4 - 8F_{2XX}) - 2F_{2\phi X} - K_X \right) \\ & + \frac{1}{2} H \dot{\phi}^4 (8A_{1X} + 3A_3) + \dot{\phi}^5 \ddot{\phi} (A_5 - A_{3X}), \end{aligned} \quad (2.10h)$$

$$\begin{aligned} \Sigma = & 3\mathcal{G}_T \dot{\Delta}^2 + 6\Theta \dot{\Delta} - \frac{3}{a^3} \frac{d}{dt} \left[a^3 (\Theta + \mathcal{G}_T \dot{\Delta}) \Delta \right] \\ & + 6F_2 H^2 + 6H \dot{\phi} \left(F_{2\phi} - 2(A_1 - 2F_{2X}) \ddot{\phi} \right) \\ & + \dot{\phi}^2 \left(12A_1 \dot{H} - 3A_3 \ddot{\phi}^2 - 3A_4 \ddot{\phi}^2 + F_X - 24F_{2X} \dot{H} - 42F_{2X} H^2 - K_\phi \right) \\ & - 3\dot{\phi}^3 \left(H \left(2A_{1X} \ddot{\phi} - 4A_{1\phi} + 3A_3 \ddot{\phi} + 6A_4 \ddot{\phi} - 4F_{2XX} \ddot{\phi} - 2F_{2\phi X} - 4K_X \right) + 2(A_3 + A_4) \ddot{\phi} \right) \\ & + \dot{\phi}^4 \left(-3H^2 (12A_{1X} + 9A_3 + 8F_{2XX}) + 6A_{1X} \dot{H} - 9A_3 \dot{H} - 9A_{3X} \ddot{\phi}^2 - 6A_{3\phi} \ddot{\phi} - 9A_{4X} \ddot{\phi}^2 \right. \\ & \quad \left. - 6A_{4\phi} \ddot{\phi} - 12A_5 \ddot{\phi}^2 + 2F_{XX} - 12F_{2XX} \dot{H} - K_{\phi X} \right) \\ & + \dot{\phi}^5 \left(3H \left(2A_{1\phi X} - A_{3X} \ddot{\phi} - 3A_{3\phi} - 2A_{4X} \ddot{\phi} - 8A_5 \ddot{\phi} + 2K_{XX} \right) - 2\ddot{\phi} (A_{3X} + A_{4X} + 4A_5) \right) \end{aligned}$$

²We note that the expression for Σ given in [36] has some typos; e.g., the prefactor of $\dot{\phi}^3$ should not contain $\ddot{\phi}$. In this paper Σ and Θ correspond to $\tilde{\Sigma}$ and $\tilde{\Theta}$ in [36, 37], respectively.

$$\begin{aligned}
& + \dot{\phi}^6 \left[-3H^2 (4A_{1XX} + 3A_{3X}) - 3A_{3X} \dot{H} \right. \\
& \quad \left. - \ddot{\phi} \left(2A_{3XX} \ddot{\phi} + 2A_{3\phi X} + 2A_{4XX} \ddot{\phi} + 2A_{4\phi X} + 13A_{5X} \ddot{\phi} + 8A_{5\phi} \right) \right] \\
& + \dot{\phi}^7 \left(-3H \left(A_{3\phi X} + 2A_{5X} \ddot{\phi} \right) - 2A_{5X} \ddot{\phi} \right) - 2\dot{\phi}^8 \ddot{\phi} \left(A_{5XX} \ddot{\phi} + A_{5\phi X} \right). \tag{2.10i}
\end{aligned}$$

Notice that terms with ζ^2 do not appear in the quadratic action $S_{\text{scalar}}^{(2)}$ since they cancel out upon using background equations of motion.

2.3 Stabilities and non-superluminality conditions: an exceptional subclass

The stability and non-superluminality conditions for the tensor sector immediately follow from the action (2.5) that

$$\mathcal{G}_{\mathcal{T}} \geq \mathcal{F}_{\mathcal{T}} > 0, \tag{2.11}$$

while those for the scalar sector require some involved algebra. First, the stability conditions amount to the positive definiteness of kinetic matrices (2.9c), i.e.

$$Q > 0, \quad \Pi \equiv \Theta^2 \mathcal{G}_S + (\mathcal{G}_{\mathcal{T}}^2 - \Theta^2 g^2) Y Q > 0, \tag{2.12a}$$

$$P_Y > 0, \quad \Xi \equiv \mathcal{F}_S - Y P_Y f^2 > 0. \tag{2.12b}$$

Second, to find the non-superluminality condition we compute the squared sound speeds, which are the eigenvalues of the matrix $G^{-1}F$

$$c_{s,\pm}^2 = \frac{1}{2\Pi} \left(\mathcal{A} \pm \sqrt{\mathcal{A}^2 - 4\Pi\Theta^2 c_m^2} \right), \tag{2.13}$$

where

$$\mathcal{A} = \Pi c_m^2 + \Theta^2 Q Y c_m^2 (f - g)^2 + \Theta^2 \Xi, \tag{2.14}$$

and $c_m^2 = P_Y/Q$ is the squared sound speed of the extra scalar field χ . Since $c_{s,+}^2 \geq c_{s,-}^2$ from (2.13), it is enough to consider only $c_{s,+}^2$ as long as we are concerned with the superluminality issue. Rewriting (2.13) as

$$\begin{aligned}
c_{s,+}^2 &= \frac{1}{2\Pi} \left(\Pi c_m^2 + \Theta^2 Q Y c_m^2 (f - g)^2 + \Theta^2 \Xi \right. \\
& \quad \left. + \sqrt{(\Pi c_m^2 + \Theta^2 Q Y c_m^2 (f - g)^2 - \Theta^2 \Xi)^2 + 4\Theta^4 Q Y c_m^2 \Xi (f - g)^2} \right), \tag{2.15}
\end{aligned}$$

it follows from the stability condition (2.12) that

$$c_{s,+}^2 \geq c_m^2 + \frac{\Theta^2 Q Y c_m^2 (f - g)^2}{\Pi} \geq c_m^2. \tag{2.16}$$

In the case that the extra scalar field has luminal sound speed, i.e. $c_m^2 = 1$, the non-superluminality condition $c_{s,+}^2 \leq 1$ combined with the inequality (2.16) implies

$$f = g \quad \text{or} \quad A_3 = \frac{2(XA_1 - 2F_2)(A_1 - 2F_{2X})}{X(3XA_1 - 4F_2)}. \tag{2.17}$$

The model that satisfies the above condition is dubbed *exceptional subclass* of DHOST Ia theories in [36, 37].³ We emphasize that DHOST Ia theories other than the exceptional subclass necessarily lead to superluminality, when combined with a luminal extra scalar.

2.4 Sufficient conditions for non-superluminality

The exceptional subclass may avoid the superluminality problem, but it is still not completely free of superluminality, since the condition (2.17) is necessary, but not sufficient for non-superluminality. In other words, we may have $c_{s,+}^2 > 1$ even though the condition (2.17) is satisfied.

Indeed, when $f = g$, the squared sound speed $c_{s,+}^2$ is reduced to

$$c_{s,+}^2 = \frac{1}{2\Pi} (\Pi c_m^2 + \Theta^2 \Xi + |\Pi c_m^2 - \Theta^2 \Xi|), \quad (2.18)$$

which becomes for $\Pi c_m^2 < \Theta^2 \Xi$ that

$$c_{s,+}^2 = \frac{\Theta^2 \Xi}{\Pi} = \frac{\Theta^2 (\mathcal{F}_S - Y P_Y g^2)}{\Theta^2 \mathcal{G}_S + (\mathcal{G}_T^2 - \Theta^2 g^2) Y Q}. \quad (2.19)$$

In the case of $\Theta^2 \Xi > \Pi$ one can readily see that $c_{s,+}^2 > 1$ and thus the superluminality appears in the scalar sector. Therefore, we find that the sufficient condition for the non-superluminality should be augmented by $\Theta^2 \Xi < \Pi$, which is achieved once $c_m^2 = 1$ and

$$\mathcal{F}_S > Y P_Y g^2. \quad (2.20)$$

To sum up, the sufficient conditions to have the non-superluminal sound speed in the exceptional subclass of DHOST Ia theories are: (i) the extra scalar field has the luminal sound speed, (ii) the stability conditions (2.11) and (2.12) holds, (iii) the scalar sector of the original DHOST action is non-superluminal (i.e., $\mathcal{G}_S \geq \mathcal{F}_S$), (iv) the newly augmented sufficient condition (2.20) is satisfied. All of these can be combined into the simple conditions

$$\mathcal{G}_T \geq \mathcal{F}_T > 0, \quad \mathcal{G}_S \geq \mathcal{F}_S > Y P_Y g^2. \quad (2.21)$$

In the subsequent sections we will construct an explicit bouncing scenario free of pathologies (that is, satisfying the constraint (2.21) together with (2.11)) as well as phenomenologically viable. The roadmap for this is as follows:

1. (Section 3) assuming that χ does not roll (i.e., $Y = 0$) so that the scalar perturbations $\tilde{\zeta}$ and $\delta\chi$ decouple from each other, we construct a completely healthy action \mathcal{S}_ϕ in the exceptional subclass of DHOST Ia theories, in the sense that it is free from

- BKL instability,

³As shown here, the condition (2.17) holds for any value of Y . In [36, 37] the emergent superluminality was analyzed for small Y .

- strong gravity assumption, in other words described both in the distant past and future by Einstein gravity and a conventional scalar field with luminal sound speed,
- superluminality as well as ghost and gradient instability, i.e. obeys the conditions (2.11) and (2.21) with $Y = 0$

$$\mathcal{G}_T \geq \mathcal{F}_T > 0, \quad \mathcal{G}_S \geq \mathcal{F}_S > 0, \quad (2.22)$$

2. (Section 4) allowing that the extra scalar χ rolls and interacts with ϕ , we construct the action \mathcal{S}_χ such that

- χ has luminal sound speed,
- during the ekpyrotic contraction phase a nearly scale-invariant power spectrum for $\delta\chi$ is generated,
- the perturbation $\delta\chi$ is converted into the curvature perturbation $\tilde{\zeta}$ after the bounce,
- while the original stability and non-superluminality of \mathcal{S}_ϕ are not spoiled, i.e.

$$\mathcal{F}_S > Y P_Y g^2. \quad (2.23)$$

3 DHOST Sector

In this section we consider the decoupled case (i.e., $\dot{\chi} = 0$) to construct a fully healthy model in the exceptional subclass of DHOST Ia, in which the independent Lagrangian functions are $F(\phi, X)$, $F_2(\phi, X)$, $K(\phi, X)$ and $A_1(\phi, X)$. To this end, we adopt the “inverse method” used in [10, 21, 27, 39–41] to choose a specific form of the Hubble parameter $H(t)$ and reconstruct the Lagrangian functions $F(\phi(t), X(t))$, $F_2(\phi(t), X(t))$, $K(\phi(t), X(t))$ and $A_1(\phi(t), X(t))$ that obey the background equations of motion and satisfy the constraint (2.22).

Let us suppose that the DHOST scalar field ϕ is a monotonic function of cosmic time t . Then, without loss of generality we can choose the form of ϕ as follows

$$\phi(t) = \mu^2 t, \quad (3.1)$$

where μ is a characteristic energy scale. This form of ϕ will greatly simplify the DHOST-related equations. Note that above form of ϕ is related to the scalar field φ with asymptotically canonical kinetic term (i.e. the coefficient of the kinetic term $\varphi_\mu \varphi^\mu$ approaching to 1/2 asymptotically) by $\phi \sim \exp(\varphi)$ [27].

It is convenient to work on the rescaled quantities:

$$t = \frac{M_{pl}}{\mu^2} \bar{t}, \quad H = \frac{\mu^2}{M_{pl}} \bar{H}, \quad \phi = M_{pl} \bar{\phi}, \quad X = \mu^4 \bar{X}, \quad (3.2a)$$

$$F = \mu^4 \bar{F}, \quad F_2 = M_{pl}^2 \bar{F}_2, \quad K = \mu \bar{K}, \quad A_1 = \frac{M_{pl}^2}{\mu^4} \bar{A}_1, \quad (3.2b)$$

$$\Theta = \mu^2 M_{pl} \bar{\Theta}, \quad \Sigma = \mu^4 \bar{\Sigma}, \quad \mathcal{D} = \frac{M_{pl}^2}{\mu^2} \bar{\mathcal{D}}, \quad (3.2c)$$

$$\mathcal{G}_{\mathcal{T}} = M_{pl}^2 \bar{\mathcal{G}}_{\mathcal{T}}, \quad \mathcal{F}_{\mathcal{T}} = M_{pl}^2 \bar{\mathcal{F}}_{\mathcal{T}}, \quad \mathcal{G}_{\mathcal{S}} = M_{pl}^2 \bar{\mathcal{G}}_{\mathcal{S}}, \quad \mathcal{F}_{\mathcal{S}} = M_{pl}^2 \bar{\mathcal{F}}_{\mathcal{S}}, \quad (3.2d)$$

where the quantities with an overbar are dimensionless. Note that the background solution (3.1) corresponds to $\bar{\phi}(\bar{t}) = \bar{t}$ from (3.2a), so that \bar{t} will be replaced by $\bar{\phi}$ at the end of the reconstruction procedure to obtain the Lagrangian functions as functions of the field ϕ . From now on, the dot notation denotes the derivative with respect to the dimensionless time \bar{t} .

3.1 Asymptotics and ansatz

Since in our model gravity and the scalar field ϕ become conventional in the asymptotic limit, the model functions $\bar{F}_2(\bar{\phi}, \bar{X})$, $\bar{K}(\bar{\phi}, \bar{X})$ and $\bar{A}_I(\bar{\phi}, \bar{X})$ ($I = 1, 2, 3, 4, 5$) should have the following asymptotic behavior:

$$\bar{F}_2(\bar{\phi}, \bar{X}) \sim -\frac{1}{2}, \quad \bar{K}(\bar{\phi}, \bar{X}), \bar{A}_I(\bar{\phi}, \bar{X}) \sim 0, \quad \text{for } |\bar{t}| \rightarrow +\infty. \quad (3.3)$$

The asymptotic behavior of the rescaled Hubble parameter \bar{H} and the rescaled model function $\bar{F}(\bar{\phi}, \bar{X})$ can be different in the far past and in the far future. We choose the contracting phase to be ekpyrotic in order to suppress the BKL instability [42]; otherwise anisotropies can grow and dominate, eventually spoiling the isotropy of FLRW cosmology and inducing a chaotic mixmaster phenomenon. Moreover, the ekpyrotic contraction is a rapid and robust smoother, meaning that homogeneity, isotropy and flatness are achieved within a few e-folds even in the case that the initial state is far from homogeneous, isotropic and flat universe [46, 47].

During the ekpyrotic contraction phase, the dimensionless Hubble parameter \bar{H} and the model function $\bar{F}(\bar{\phi}, \bar{X})$ behave as $\bar{H} \sim \frac{1}{\epsilon \bar{t}}$, $\bar{F}(\bar{\phi}, \bar{X}) \sim -2\dot{\bar{H}} - 3\bar{H}^2 \sim \frac{2\epsilon-3}{\epsilon^2 \bar{t}^2}$ with $\epsilon > 3$ as $\bar{t} \rightarrow -\infty$. On the other hand, we let the rescaled Hubble parameter $\bar{H} \sim \frac{1}{3\bar{t}}$ and the model function $\bar{F}(\bar{\phi}, \bar{X}) \sim \frac{1}{3\bar{t}^2}$ as $\bar{t} \rightarrow +\infty$ in the expanding phase as in [27]. This solution corresponds to the kinetic-dominated expanding universe, which is in fact a scaling attractor when the universe is filled only with a scalar field with sufficiently steep exponential potential [67]. In total, we let the scale factor $a(\bar{t})$, the Hubble parameter $\bar{H}(\bar{t}) \equiv \dot{a}/a$ and $\bar{F}(\bar{\phi}, \bar{X})$ behave asymptotically as follows:

$$a \propto \begin{cases} \bar{t}^{\frac{1}{3}}, & \bar{t} \rightarrow +\infty \\ (-\bar{t})^{\frac{1}{\epsilon}}, & \bar{t} \rightarrow -\infty \end{cases}, \quad \bar{H} \sim \begin{cases} \frac{1}{3\bar{t}}, & \bar{t} \rightarrow +\infty \\ \frac{1}{\epsilon \bar{t}}, & \bar{t} \rightarrow -\infty \end{cases}, \quad \bar{F}(\bar{\phi}, \bar{X}) \sim \begin{cases} \frac{1}{3\bar{t}^2}, & \bar{t} \rightarrow +\infty \\ \frac{2\epsilon-3}{\epsilon^2 \bar{t}^2}, & \bar{t} \rightarrow -\infty \end{cases}. \quad (3.4)$$

Our choice for $a(\bar{t})$ with the above asymptotic behavior is

$$a(\bar{t}) = \sigma\left(\frac{\bar{t}}{\tau}\right) \left[\left(\frac{\bar{t}}{\tau}\right)^2 + 1 \right]^{\frac{1}{6}} + \sigma\left(-\frac{\bar{t}}{\tau}\right) \left[\left(\frac{\bar{t}}{\tau}\right)^2 + 1 \right]^{\frac{1}{2\epsilon}}, \quad (3.5)$$

where $\sigma(\bar{t}) \equiv 1/(1 + \exp(-\bar{t}))$ is the logistic sigmoid function, and $\tau > 0$ is a parameter that controls the duration of the bounce phase. For this choice, the bounce occurs exactly




Figure 1. Plot of the scale factor a and the rescaled Hubble parameter \bar{H} . The parameters τ and ϵ are set to be $\tau = 10$ and $\epsilon = 10$.

at $\bar{t} = 0$, see Fig. 1. Note that ignoring the exponentially suppressed terms, the rescaled Hubble parameter \bar{H} behaves as

$$\bar{H}(\bar{t}) \sim \frac{\bar{t}}{\gamma(\bar{t}^2 + \tau^2)} \quad \text{for } |\bar{t}| \rightarrow +\infty, \quad (3.6)$$

where $\gamma = 3$ for $\bar{t} > 0$ and $\gamma = \epsilon$ for $\bar{t} < 0$.

We make an ansatz for the independent Lagrangian functions $\bar{F}(\bar{\phi}, \bar{X})$, $\bar{K}(\bar{\phi}, \bar{X})$, $\bar{F}_2(\bar{\phi}, \bar{X})$ and $\bar{A}_1(\bar{\phi}, \bar{X})$ as follows:

$$\bar{F}(\bar{\phi}, \bar{X}) = f_0(\bar{\phi}) + f_1(\bar{\phi})\bar{X} + f_2(\bar{\phi})\bar{X}^2, \quad (3.7a)$$

$$\bar{F}_2(\bar{\phi}, \bar{X}) = -\frac{1}{2} - g_0(\bar{\phi}) - g_1(\bar{\phi})\bar{X}, \quad (3.7b)$$

$$\bar{A}_1(\bar{\phi}, \bar{X}) = a_0(\bar{\phi}) + a_1(\bar{\phi})\bar{X}, \quad (3.7c)$$

$$\bar{K}(\bar{\phi}, \bar{X}) = 0. \quad (3.7d)$$

We let the functions $g_{0,1}(\bar{t})$ and $a_{0,1}(\bar{t})$ be exponentially suppressed as $|t| \rightarrow +\infty$, i.e.⁴

$$g_{0,1}(\bar{t}), a_{0,1}(\bar{t}) \propto e^{-\frac{2|\bar{t}|}{\tau}} \quad \text{for } |\bar{t}| \rightarrow +\infty. \quad (3.8)$$

During the ekpyrotic contraction phase ($\bar{t} < 0$), $f_0(\bar{t})$ and $f_1(\bar{t})$ behave as

$$f_0(\bar{t}) \sim \frac{\epsilon - 3}{\epsilon^2 \bar{t}^2}, \quad f_1(\bar{t}) \sim \frac{1}{\epsilon \bar{t}^2} \quad \text{for } \bar{t} \rightarrow -\infty, \quad (3.9)$$

while during the (kinetic-dominated) expanding phase ($\bar{t} > 0$) the potential term f_0 becomes subdominant, namely

$$f_0(\bar{t}) \propto \frac{1}{\bar{t}^4}, \quad f_1(\bar{t}) \sim \frac{1}{3\bar{t}^2} \quad \text{for } \bar{t} \rightarrow +\infty. \quad (3.10)$$

The function $f_2(\bar{t})$ should be subdominant compared to $f_1(\bar{t})$, and therefore $f_2(\bar{t})$ should behave asymptotically as follows:

$$f_2(\bar{t}) \propto \frac{1}{\bar{t}^4} \quad \text{for } |\bar{t}| \rightarrow +\infty. \quad (3.11)$$

⁴The functions $a_{0,1}(\bar{t})$ should not be confused with the scale factor $a(\bar{t})$.

3.2 Stability and non-superluminality

Now we choose functions $f_{0,1,2}(\bar{t})$, $g_{0,1}(\bar{t})$ and $a_{0,1}(\bar{t})$ such that not only the background equations of motion but also the constraint (2.22) is satisfied.

First, the constraint (2.11) for the tensor sector is easy to be satisfied. Indeed, for the ansatz (3.7), the coefficients of the quadratic action in the tensor sector become

$$\bar{\mathcal{G}}_{\mathcal{T}} = 1 + 2[a_0(\bar{t}) + a_1(\bar{t}) + g_0(\bar{t}) + g_1(\bar{t})], \quad \bar{\mathcal{F}}_{\mathcal{T}} = 1 + 2[g_0(\bar{t}) + g_1(\bar{t})]. \quad (3.12)$$

We select g_0 and a_0 such that $\bar{\mathcal{G}}_{\mathcal{T}} = \bar{\mathcal{F}}_{\mathcal{T}} = 1$, namely

$$g_0(\bar{t}) = -g_1(\bar{t}), \quad a_0(\bar{t}) = -a_1(\bar{t}). \quad (3.13)$$

That is, the speed of gravitational wave is always equal to unity and all the pathologies are absent in the tensor sector.

Now we determine the functions $f_{0,1,2}(\bar{t})$, $a_1(\bar{t})$ and $g_1(\bar{t})$ such that none of the pathologies appear in the scalar sector, i.e. $\bar{\mathcal{G}}_{\mathcal{S}} \geq \bar{\mathcal{F}}_{\mathcal{S}} > 0$. Notice that only $a_1(\bar{t})$ and $g_1(\bar{t})$ among these five Lagrangian functions appear in $\bar{\mathcal{F}}_{\mathcal{S}}$, namely

$$\bar{\mathcal{F}}_{\mathcal{S}} = \frac{1}{a} \frac{d\xi}{d\bar{t}} - 1, \quad (3.14)$$

where

$$\xi \equiv \frac{a\Lambda}{\bar{\Theta}}, \quad (3.15)$$

with

$$\Lambda \equiv \bar{\mathcal{G}}_{\mathcal{T}} + \bar{\mathcal{D}}\dot{\phi} + \bar{\mathcal{F}}_{\mathcal{T}}\Delta = 1 - 3g_1(\bar{t}), \quad (3.16)$$

$$\bar{\Theta} = \bar{H}(4a_1(\bar{t}) + g_1(\bar{t}) + 1) + \dot{g}_1(\bar{t}). \quad (3.17)$$

We therefore take the first step to choose $a_1(\bar{t})$ and $g_1(\bar{t})$ such that $\bar{\mathcal{F}}_{\mathcal{S}} > 0$.

Since $\dot{\xi} = a(1 + \bar{\mathcal{F}}_{\mathcal{S}}) > 0$, one can naively conclude that ξ should be a monotonically increasing function. However, one must be careful since $\bar{\Theta} \sim \bar{H}$ as $|\bar{t}| \rightarrow +\infty$ and thus $\bar{\Theta}$ should cross zero at least once. Indeed, both $\bar{\mathcal{G}}_{\mathcal{S}}$ and $\bar{\mathcal{F}}_{\mathcal{S}}$ become divergent when $\bar{\Theta} = 0$. As shown in [27], this is entirely harmless, because the equation of motion for $\tilde{\zeta}$ has regular solutions. A similar situation was discussed in [68].

We assume that $\bar{\Theta}$ crosses zero only once for simplicity and denote the time when $\bar{\Theta}$ vanishes by \bar{t}_0 . Then, the function $\xi(\bar{t})$ should increase monotonically except for the moment $\bar{t} = \bar{t}_0$. In particular, $\xi(\bar{t})$ should behave in the vicinity of $\bar{t} = \bar{t}_0$ as

$$\lim_{\bar{t} \rightarrow \bar{t}_0^-} \xi(\bar{t}) = +\infty, \quad \lim_{\bar{t} \rightarrow \bar{t}_0^+} \xi(\bar{t}) = -\infty. \quad (3.18)$$

On the other hand, it follows from the form of \bar{H} and the asymptotics (3.8) that $\xi(-\infty) < 0$ and $\xi(+\infty) > 0$. Therefore, Λ crosses zero once in the interval $(-\infty, \bar{t}_0)$ and $(\bar{t}_0, +\infty)$, respectively.




Figure 2. Plot of a_1 , $\bar{\Theta}$ and \bar{H} with the parameters given by (3.22).

To achieve this, we choose the form of $g_1(\bar{t})$ as follows:

$$g_1(\bar{t}) = \frac{1}{3}w \cosh^{-2}\left(\frac{\bar{t}}{\tau} + u\right), \quad (3.19)$$

where the parameter τ controls the duration of the bounce phase. The parameters $w > 1$ and u are introduced to avoid the fine-tuning problem that \bar{H} , $\bar{\Theta}$ and ξ become zero at the same time [27]. We also choose $a_1(\bar{t})$ such that $\bar{\Theta}$ is sufficiently close to \bar{H} for all times (see Fig. 2).

$$a_1(\bar{t}) = \frac{w}{12} \operatorname{sech}^2\left(\frac{\bar{t}}{\tau} + u\right) \left(\frac{2\left(\bar{t}^2 \tanh\left(\frac{\bar{t}}{\tau} + u\right) + \tau^2 \tanh\left(\frac{\bar{t}}{\tau}\right)\right)}{\bar{H}\tau(\bar{t}^2 + \tau^2)} - 1 \right). \quad (3.20)$$

Then, $\bar{\Theta}$ becomes

$$\bar{\Theta} = \frac{1}{3\tau\epsilon} \left[\frac{3\bar{t}\tau}{\bar{t}^2 + \tau^2} + \frac{e^{\bar{t}/\tau} (3\bar{t}^2\epsilon + \tau\bar{t}(\epsilon - 3) + 3\tau^2\epsilon)}{e^{\bar{t}/\tau} (\bar{t}^2 + \tau^2) + \tau^2 \left(\frac{\bar{t}^2}{\tau^2} + 1\right)^{\frac{1}{2\epsilon} + \frac{5}{6}}} + 3\epsilon \left(\frac{1}{e^{\bar{t}/\tau} + 1} - 1 \right) \right] - \frac{2\tau w}{3(\bar{t}^2 + \tau^2)} \sinh(u) \operatorname{sech}\left(\frac{\bar{t}}{\tau}\right) \operatorname{sech}^3\left(\frac{\bar{t}}{\tau} + u\right). \quad (3.21)$$

Fig. 3 confirms that we made an appropriate choice (3.19) and (3.20) for $g_1(\bar{t})$ and $a_1(\bar{t})$. Fig. 4 shows that $\bar{\mathcal{F}}_S$ becomes positive for all times, where the parameters are

$$\tau = 10, \quad w = 2, \quad u = \frac{1}{10}, \quad \epsilon = 5. \quad (3.22)$$

The next step is to determine the Lagrangian functions $f_{0,1,2}(\bar{t})$ such that the constraint $\bar{\mathcal{G}}_S \geq \bar{\mathcal{F}}_S$, which reads

$$\bar{\Sigma} \geq \bar{\Theta}^2 \left[\frac{d}{dt} \left(\frac{\Lambda}{\bar{\Theta}} \right) + \bar{H} \frac{\Lambda}{\bar{\Theta}} - 4 \right], \quad (3.23)$$

where the function $\bar{\Sigma}$ given in (2.10i) is reduced to

$$\bar{\Sigma} = f_1(\bar{t}) + 6f_2(\bar{t}) - 2g_1(\bar{t})\ddot{g}_1(\bar{t}) + 3\dot{g}_1(\bar{t})^2$$




Figure 3. Plot of Λ , $\bar{\Theta}$ and ξ with the parameters given by (3.22).

$$\begin{aligned}
& + \bar{H}(\bar{t}) [(15a_1(\bar{t}) - 4g_1(\bar{t}) - 15)g_1(\bar{t}) - (g_1(\bar{t}) - 3)\dot{a}_1(\bar{t})] \\
& - [a_1(\bar{t})(g_1(\bar{t}) - 3) + (g_1(\bar{t}) - 11)g_1(\bar{t})] \dot{\bar{H}}(\bar{t}) \\
& - 3\bar{H}^2(\bar{t}) [a_1(\bar{t})(g_1(\bar{t}) + 15) + (g_1(\bar{t}) - 1)g_1(\bar{t}) + 1]. \tag{3.24}
\end{aligned}$$

The form of $\bar{\Sigma}$ should be chosen such that not only the constraint (3.23) is satisfied but also it has the correct asymptotics. Once done, (3.24) will give a relationship between the Lagrangian functions $f_1(\bar{t})$ and $f_2(\bar{t})$.

It follows from (3.4) and (3.8)–(3.11) that $\bar{\Sigma}$ behaves asymptotically as

$$\bar{\Sigma} \sim \frac{\epsilon - 3}{\epsilon^2 \bar{t}^2} \text{ for } \bar{t} \rightarrow -\infty, \quad \Sigma \propto \frac{1}{\bar{t}^4} \text{ for } \bar{t} \rightarrow +\infty. \tag{3.25}$$

In the limiting case $|\bar{t}| \rightarrow +\infty$, the exponentially suppressed terms can be ignored, and thus the constraint (3.23) becomes

$$\bar{\Sigma} \geq -\dot{\bar{H}} - 3\bar{H}^2 \sim \frac{\bar{t}^2(\gamma - 3) - \gamma\tau^2}{\gamma^2(\bar{t}^2 + \tau^2)^2}, \tag{3.26}$$

where $\gamma = 3$ for $\bar{t} > 0$ and $\gamma = \epsilon$ for $\bar{t} < 0$. Our choice for $\bar{\Sigma}$ with the asymptotics (3.25) that satisfies the condition (3.26) is

$$\bar{\Sigma}(\bar{t}) = \frac{\tau^4}{(\bar{t}^2 + \tau^2)^2} + \frac{(\epsilon - 3) \left(1 - \tanh\left(\frac{\bar{t}}{\tau}\right)\right)}{2\epsilon^2 (\bar{t}^2 + \tau^2)}. \tag{3.27}$$

We numerically checked that $\bar{\mathcal{G}}_S > \bar{\mathcal{F}}_S > 0$ and thus $0 < c_s^2 < 1$ for all times, see Fig. 4 and 5.




Figure 4. Plot of $\bar{\mathcal{G}}_S$ and $\bar{\mathcal{F}}_S$. The parameters w , u , τ and ϵ are given by (3.22).




Figure 5. Plot of sound speed squared c_s^2 , which lies between 0 and 1 for all times. The main panel shows that c_s^2 tends to 1 as $|\bar{t}| \rightarrow +\infty$. The inset shows sound speed squared in the vicinity of bounce at $\bar{t} = 0$. The parameters w , u , τ and ϵ are given by (3.22).

It follows from (3.24) and (3.27) that

$$f_1(\bar{t}) + 6f_2(\bar{t}) = \frac{\tau^4}{(\bar{t}^2 + \tau^2)^2} + \frac{(\epsilon - 3) \left(1 - \tanh\left(\frac{\bar{t}}{\tau}\right)\right)}{2\epsilon^2 (\bar{t}^2 + \tau^2)} + 2g_1(\bar{t})\ddot{g}_1(\bar{t}) - 3\dot{g}_1(\bar{t})^2$$

$$\begin{aligned}
& -\bar{H}(\bar{t}) [(15a_1(\bar{t}) - 4g_1(\bar{t}) - 15)\dot{g}_1(\bar{t}) - (g_1(\bar{t}) - 3)\dot{a}_1(\bar{t})] \\
& + [a_1(\bar{t})(g_1(\bar{t}) - 3) + (g_1(\bar{t}) - 11)g_1(\bar{t})] \dot{\bar{H}}(\bar{t}) \\
& + 3\bar{H}^2(\bar{t}) [a_1(\bar{t})(g_1(\bar{t}) + 15) + (g_1(\bar{t}) - 1)g_1(\bar{t}) + 1]. \tag{3.28}
\end{aligned}$$

Here we do not give the explicit form of the Lagrangian functions $f_{0,1,2}$, which will be determined in the next section, since they are affected by the extra scalar field χ through the background equations of motion, though their asymptotic behavior remains unchanged.

4 Extra Scalar and Comparison with Observations

In this section we consider the coupled case, that is, allow χ to have non-zero kinetic energy and interact with ϕ so that the perturbations $\delta\chi$ can be converted into the curvature fluctuations.

4.1 Setup

As mentioned in the Introduction we focus on the typical case in which $P(\chi, Y, \phi)$ has the conventional form such that the perturbation $\delta\chi$ has luminal sound speed, namely

$$P(\chi, Y, \phi) = P_1(\chi, \phi)Y + P_0(\chi, \phi), \tag{4.1}$$

with $P_1(\chi, \phi) > 0$. As in the previous section, it is convenient to work on the rescaled quantities:

$$\chi = M_{pl}\bar{\chi}, \quad Y = \mu^4\bar{Y}, \quad P(\chi, Y, \phi) = \mu^4\bar{P}(\bar{\chi}, \bar{Y}, \bar{\phi}) = \mu^4(\bar{P}_0 + \bar{Y}P_1). \tag{4.2}$$

Let us first make an ansatz for $\bar{P}_0(\bar{\chi}, \bar{\phi})$ as follows:

$$\bar{P}_0(\bar{\chi}, \bar{\phi}) = -\bar{P}_{02}(\bar{\phi})\bar{\chi}^2 - \bar{P}_{04}(\bar{\phi})\bar{\chi}^4. \tag{4.3}$$

The presence of the quartic term is only for stability of the scalar potential. Of course, it is possible to make the scalar potential bounded from below only in terms of the quadratic term. It, however, turns out to be highly non-trivial in this case to make the conversion from $\delta\chi$ into $\tilde{\zeta}$ sufficiently efficient. Note that in order for interaction energy $-P_0(\bar{\chi}, \bar{\phi})$ to be bounded from below, $\bar{P}_{04}(\bar{\phi})$ should be positive and $-\bar{P}_{02}^2(\bar{\phi})/\bar{P}_{04}(\bar{\phi})$ should be bounded from below. We also assume that P_1 does not depend on χ and the scalar field χ has canonical kinetic term in the distant future. As we see below, the behavior of P_1 in the distant past is not canonical.

The background equations of motion (A.1), (A.2) and (A.3) become

$$\begin{aligned}
& (f_0(\bar{t}) - \bar{P}_{02}(\bar{t})\bar{\chi}^2 - \bar{P}_{04}(\bar{t})\bar{\chi}^4) - (f_1(\bar{t}) + \bar{Y}P_1(\bar{t})) - 3f_2(\bar{t}) \\
& + 3\bar{H}\bar{\Theta} + 3\bar{H}\dot{g}_1 - 3g_1(2\dot{\bar{H}} + 3\bar{H}^2) = 0, \tag{4.4a}
\end{aligned}$$

$$(f_0(\bar{t}) - \bar{P}_{02}(\bar{t})\bar{\chi}^2 - \bar{P}_{04}(\bar{t})\bar{\chi}^4) + (f_1(\bar{t}) + \bar{Y}P_1(\bar{t})) + f_2(\bar{t}) + 2\dot{\bar{H}} + 3\bar{H}^2 = 0, \tag{4.4b}$$

$$\frac{1}{a^3} \frac{d}{d\bar{t}} (a^3 \dot{\bar{\chi}} P_1(\bar{t})) + \bar{P}_{02}(\bar{t})\bar{\chi} + 2\bar{P}_{04}(\bar{t})\bar{\chi}^3 = 0. \tag{4.4c}$$

For simplicity, we let $\dot{\chi}$ decay exponentially in the asymptotic limit so that $\tilde{\zeta}$ and $\delta\chi$ decouple both in the distant past and in the distant future. It then follows from (4.4c) that $\bar{P}_{02}(\bar{t})$ and $\bar{P}_{04}(\bar{t})$ are also suppressed exponentially. This offers two advantages. The first one is that the asymptotic behavior of $f_{0,1,2}(\bar{t})$ derived in the previous subsection is not affected through the background equations of motion (4.4a) and (4.4b). The second one is that it is guaranteed that $-\bar{P}_{02}^2(\bar{\phi})/\bar{P}_{04}(\bar{\phi})$ is bounded from below as long as \bar{P}_{04} does not touch zero.

Our goal is to choose $\bar{\chi}(\bar{t})$ and the Lagrangian functions P_1 , \bar{P}_{02} and $\bar{P}_{04} > 0$ such that

- a nearly scale-invariant power spectrum of the extra scalar field χ is generated during the ekpyrotic phase,
- the perturbations $\delta\chi$ are converted into the curvature fluctuations $\tilde{\zeta}$ with high efficiency by allowing χ to have non-zero kinetic energy and interact with $\bar{\phi}$,
- the background equations of motion (4.4) and the constraint (2.23) are satisfied,
- $\dot{\chi}(\bar{t})$, $\bar{P}_{02}(\bar{t})$ and $\bar{P}_{04}(\bar{t})$ decay exponentially with the same exponent as $|\bar{t}| \rightarrow +\infty$.

4.2 Scalar spectral index for entropic perturbations

First of all, the nearly scale-invariant perturbations can be generated by taking the form of $P_1(\chi, \phi)$ as $\phi^{-2(1+b)}$ with $b \ll 1$ [54–56, 62]. To be concrete, in the decoupled case the equation for the Fourier mode $\delta\chi_k$ of the perturbation $\delta\chi$ becomes

$$\frac{1}{a^3} \frac{d}{dt} \left(a^3 P_1 \frac{d}{dt} \delta\chi_k \right) + \frac{k^2}{a^2} P_1 \delta\chi_k = 0, \quad (4.5)$$

or

$$\frac{d^2}{d\eta^2} v_k + \left(k^2 - \frac{1}{z} \frac{d^2 z}{d\eta^2} \right) v_k = 0, \quad (4.6)$$

where $v_k \equiv z \delta\chi_k$ with $z \equiv a\sqrt{P_1}$ and η denotes the conformal time defined by $dt = a d\eta$. Note that during the ekpyrotic contraction phase, the conformal time η is related to t by

$$-\eta \sim (-t)^{\frac{\epsilon-1}{\epsilon}}. \quad (4.7)$$

It follows from $P_1 \sim (-t)^{-2(1+b)}$ that $z \sim (-\eta)^{-b\epsilon/(\epsilon-1)-1}$. Since in the distant past the mode functions obey the equation for a free field in Minkowski space, we can impose the Bunch-Davies vacuum initial condition for the mode functions, namely

$$\lim_{k\eta \rightarrow -\infty} v_k = \frac{1}{\sqrt{2k}} e^{-ik\eta}. \quad (4.8)$$

The solution of (4.6) is then

$$v_k = \frac{\sqrt{-\pi\eta}}{2} H_\nu^{(1)}(-k\eta) \quad \text{with } \nu = \frac{3}{2} + \frac{b\epsilon}{\epsilon-1}, \quad (4.9)$$

where $H_\nu^{(1)}$ is a Hankel function of the first kind. In the super-horizon limit, i.e. $|k\eta| \ll 1$, we obtain

$$v_k \sim \frac{(-\eta)^{\frac{1}{2}-\nu}}{k^\nu}, \quad (4.10)$$

and thus we find the spectral index n_s related to b by

$$n_s = 1 - \frac{2b\epsilon}{\epsilon - 1}. \quad (4.11)$$

Since $\epsilon \gg 1$, for the value of b around 0.0158 the spectral tilt becomes $n_s \simeq 0.965$, which is in good agreement with the Planck data [50]. Eventually, once $P_1(\bar{\phi}) = P_1(\bar{t})$ behaves in the far past as

$$P_1(\bar{t}) \propto \frac{1}{(-\bar{t})^{2(1+b)}} \quad \text{for } \bar{t} \rightarrow -\infty, \quad (4.12)$$

the power spectrum of $\delta\bar{\chi}$ is nearly scale-invariant.

4.3 Conversion of entropic perturbations into curvature fluctuations

Note that during the ekpyrotic phase the curvature fluctuation $\tilde{\zeta}$ is highly blue-shifted, namely

$$a\sqrt{\mathcal{G}_S}M_{pl}\tilde{\zeta}_k = \frac{\sqrt{-\pi\eta}}{2}H_{\nu_1}^{(1)}(-k\eta) \quad \text{with} \quad \nu_1 = \frac{\epsilon - 3}{2(\epsilon - 1)}, \quad (4.13)$$

where $0 < \nu_1 < \frac{1}{2}$, given $\epsilon > 3$. It follows that $a\sqrt{\mathcal{G}_S}M_{pl}\tilde{\zeta}_k \sim \frac{(-\eta)^{\frac{1}{2}-\nu_1}}{k^{\nu_1}}$ for $|k\eta| \ll 1$, and hence the amplitude of $\tilde{\zeta}$ is much smaller than the one of $\delta\bar{\chi}$ during the ekpyrotic phase. We therefore need to introduce a mechanism to convert $\delta\bar{\chi}$ into $\tilde{\zeta}$ with high efficiency.

We choose the following form of $P_1(\bar{t})$ (see Fig. 6)

$$P_1(\bar{t}) = \frac{1}{2} \frac{\sigma\left(-\frac{\bar{t}}{\tau}\right)}{\left(\frac{\bar{t}}{\tau}\right)^{2(1+b)} + 1} + \frac{1}{2}\sigma\left(\frac{\bar{t}}{\tau}\right), \quad (4.14)$$

so that the scalar field χ has canonical kinetic energy in the far future.

Now we need to choose $\bar{\chi}$, \bar{P}_{02} and \bar{P}_{04} such that conversion from the perturbation $\delta\bar{\chi}$ into the curvature fluctuation $\tilde{\zeta}$ occurs efficiently. We remind that \bar{P}_{04} is introduced solely for stability of the potential and \bar{P}_{02} is determined by the background equation of motion (4.4c) once $\bar{\chi}$ and \bar{P}_{04} are specified. It is therefore crucial to choose the form of $\bar{\chi}$ appropriately.

Let us first investigate what is the main factor in conversion from entropy to curvature perturbations. It follows from (2.7) that the scalar perturbations obey the following equations:

$$\frac{1}{a^3} \frac{d}{dt} \left[a^3 \left(\left(\bar{\mathcal{G}}_S + \frac{\bar{Y}P_1}{\Theta^2} \right) \dot{\zeta}_k + \dot{\bar{\chi}} P_1 \bar{g} \delta\bar{\chi}_k + \frac{1}{2} \bar{\Psi}_1 \delta\bar{\chi}_k \right) \right] +$$




Figure 6. Plot of P_1 . The parameters are set to be $\tau = 10$ and $b = 0.0158$, respectively.

$$+\frac{\bar{k}^2}{a^2} \left(\bar{\mathcal{F}}_S \tilde{\zeta}_k + \dot{\chi} P_1 \bar{g} \delta \bar{\chi}_k \right) = 0, \quad (4.15a)$$

$$\frac{1}{a^3} \frac{d}{d\bar{t}} \left[a^3 \left(P_1 \delta \dot{\bar{\chi}}_k + \dot{\chi} P_1 \bar{g} \dot{\zeta}_k \right) \right] + \frac{\bar{k}^2}{a^2} \left(P_1 \delta \bar{\chi}_k + \dot{\chi} P_1 \bar{g} \tilde{\zeta}_k \right) - \frac{1}{2} \bar{\Psi}_1 \dot{\zeta}_k - \bar{\Psi}_2 \delta \bar{\chi}_k = 0, \quad (4.15b)$$

where $\bar{k} = \frac{M_{pl} k}{\mu^2}$ is the (dimensionless) rescaled wavenumber and

$$\bar{\Psi}_1 = \frac{2}{\bar{\Theta}^2} \left[(\bar{Y} P_1 + \bar{\Sigma}) \dot{\chi} P_1 - \bar{\Theta} (\bar{P}_{02} \bar{\chi} + 2 \bar{P}_{04} \bar{\chi}^3) (1 + 3\Delta) \right], \quad (4.15c)$$

$$\begin{aligned} \bar{\Psi}_2 = & -\bar{P}_{02} - 6 \bar{P}_{04} \bar{\chi}^2 + \frac{1}{a^3} \frac{d}{d\bar{t}} \left(\frac{a^3 \bar{Y} P_1^2 (1 - 3\Delta)}{\bar{\Theta}} \right) \\ & + \frac{\dot{\chi} P_1}{\bar{\Theta}^2} \left[\dot{\chi} P_1 (\bar{Y} P_1 + \bar{\Sigma}) + \bar{\Theta} P_{0\bar{\chi}} (1 + 3\Delta) \right], \end{aligned} \quad (4.15d)$$

$$\bar{g} = -\frac{1}{\bar{\Theta}} (1 - 3\Delta). \quad (4.15e)$$

In the long wavelength limit $\bar{k} \rightarrow 0$, which is of observational interest, the equations (4.15) become

$$\left(\bar{\mathcal{G}}_S + \frac{\bar{Y} P_1}{\bar{\Theta}^2} \right) \dot{\zeta}_k + \dot{\chi} P_1 \bar{g} \delta \dot{\bar{\chi}}_k + \frac{1}{2} \bar{\Psi}_1 \delta \bar{\chi}_k = \frac{C_0}{a^3}, \quad (4.16a)$$

$$\frac{1}{a^3} \frac{d}{d\bar{t}} \left[a^3 \left(P_1 \delta \dot{\bar{\chi}}_k + \dot{\chi} P_1 \bar{g} \dot{\zeta}_k \right) \right] - \frac{1}{2} \bar{\Psi}_1 \dot{\zeta}_k - \bar{\Psi}_2 \delta \bar{\chi}_k = 0, \quad (4.16b)$$

or

$$0 = \dot{\zeta}_k - \frac{1}{\bar{G}_{11}} \left(\frac{C_0}{a^3} - \bar{G}_{12} \delta \dot{\bar{\chi}}_k - \frac{1}{2} \bar{\Psi}_1 \delta \bar{\chi}_k \right), \quad (4.17a)$$

$$0 = \left(P_1 - \frac{\bar{G}_{12}^2}{\bar{G}_{11}} \right) \delta \ddot{\bar{\chi}}_k + \left[3 \bar{H} \left(P_1 - \frac{\bar{G}_{12}^2}{\bar{G}_{11}} \right) + \dot{P}_1 - \frac{d}{d\bar{t}} \left(\frac{\bar{G}_{12}^2}{\bar{G}_{11}} \right) \right] \delta \dot{\bar{\chi}}_k$$

$$\begin{aligned}
& + \left[\frac{\bar{\Psi}_1^2}{4\bar{G}_{11}} - 3\bar{H} \frac{\bar{G}_{12}}{2\bar{G}_{11}} \bar{\Psi}_1 - \frac{1}{2} \frac{d}{dt} \left(\frac{\bar{G}_{12} \bar{\Psi}_1}{\bar{G}_{11}} \right) - \bar{\Psi}_2 \right] \delta \bar{\chi}_k \\
& + \frac{C_0}{a^3} \left[\frac{d}{dt} \left(\frac{\bar{G}_{12}}{\bar{G}_{11}} \right) - \frac{1}{2} \frac{\dot{\bar{\Psi}}_1}{\bar{G}_{11}} \right], \tag{4.17b}
\end{aligned}$$

where C_0 is an integration constant, which is determined by the initial condition, and

$$\bar{G}_{11} = \bar{G}_S + \frac{\bar{Y} P_1}{\bar{\Theta}^2}, \quad \bar{G}_{12} = \dot{\bar{\chi}} P_1 \bar{g}. \tag{4.18}$$

It is now evident from (4.17a) that $\bar{\Psi}_1/\bar{G}_{11}$ is the main factor that controls the efficiency of conversion process. Since $\bar{\Psi}_1$ depends on $\dot{\bar{\chi}}$, one is tempted to make $|\dot{\bar{\chi}}|$ large. This approach has drawbacks, though, due to the constraint (2.23). Therefore, the only possible way to increase $\bar{\Psi}_1$ is to make $|\bar{P}_{02}\bar{\chi} + 2\bar{P}_{04}\bar{\chi}^3| = |\frac{d}{dt}(a\dot{\bar{\chi}}P_1)|$ large by increasing $\ddot{\bar{\chi}}$ instead of $\dot{\bar{\chi}}$. For this reason, we choose the following form of $\bar{\chi}$ (see Fig. 7)

$$\bar{\chi} = \bar{\chi}_0 + \lambda \operatorname{sech} \left(\frac{\bar{t} - \bar{t}_c}{2\tau} \right) \left[e^{-\left(\frac{\bar{t} - \bar{t}_c}{4\tau}\right)^2} - q \right], \tag{4.19}$$

where the parameter λ controls the overall rate of change of $\bar{\chi}$ and the parameter \bar{t}_c is related to the moment of conversion from $\delta\chi$ into $\tilde{\zeta}$. We set $\bar{t}_c \sim 10\tau$ so that the conversion occurs sufficiently long after the bounce phase. The bias $\bar{\chi}_0$ is introduced in order for $\bar{\chi}$ not to cross zero; otherwise \bar{P}_{02} and \bar{P}_{04} become singular at a certain point. The overall shape of the background trajectory around the conversion phase is determined by the function $\operatorname{sech} \left(\frac{\bar{t} - \bar{t}_c}{2\tau} \right)$ in (4.19). On the other hand, the factor in the square bracket plays two roles, the first one as enhancing the conversion efficiency and the second one as making \bar{P}_{02} positive in the asymptotic limit. The second role, which is actually played by the parameter q , is necessary since we do not want the tachyonic instability to appear in the asymptotic part of the background trajectory of interest. By adjusting the parameters $\bar{\chi}_0$, λ and q , we can make the conversion efficient and smooth, while not violating the constraint (2.23). Here we mean the conversion efficiency by [61, 62]

$$|\tilde{\zeta}_k(\bar{t}_{conv-end})/\delta\bar{\chi}_k(\bar{t}_{conv-begin})|, \tag{4.20}$$

where $\bar{t}_{conv-begin}$ and $\bar{t}_{conv-end}$ denote the beginning and end moment of the conversion process, respectively. And the conversion is said to be smooth when the comoving Hubble radius $(aH)^{-1}$ changes by order of one e-fold during conversion [61, 69].

We choose the following form of \bar{P}_{04}

$$\bar{P}_{04}(\bar{t}) = 10^{-10} \operatorname{sech} \left(\frac{\bar{t} - \bar{t}_c}{2\tau} \right), \tag{4.21}$$

so that it has the same asymptotics with $\dot{\bar{\chi}}$ and plays little role in the dynamics. The Lagrangian function \bar{P}_{02} is determined by combining (4.4c), (4.19) and (4.21), namely

$$\bar{P}_{02}(\bar{t}) = -2\bar{P}_{04}(\bar{t})\bar{\chi}^2 - \frac{1}{a^3\bar{\chi}} \frac{d}{dt} (a^3\dot{\bar{\chi}}P_1(\bar{t})). \tag{4.22}$$




Figure 7. Plot of $\bar{\chi}$ and $\dot{\bar{\chi}}$. The parameters $\bar{\chi}_0$, λ , τ and \bar{t}_c are given by (4.26).

One can readily see that the asymptotic behavior of \bar{P}_{02} and \bar{P}_{04} is given as follows:

$$\bar{P}_{02} \sim \frac{1}{\bar{t}^2} \exp\left(\frac{\bar{t}}{2\tau}\right) \text{ as } \bar{t} \rightarrow -\infty, \quad \bar{P}_{02} \sim \exp\left(\frac{-\bar{t}}{2\tau}\right) \text{ as } \bar{t} \rightarrow +\infty, \quad (4.23)$$

$$\bar{P}_{04} \sim \exp\left(-\frac{|\bar{t}|}{2\tau}\right) \text{ as } |\bar{t}| \rightarrow +\infty. \quad (4.24)$$

Fig. 8 confirms that \bar{P}_{02} and \bar{P}_{04} behave asymptotically as given in (4.23). It follows from (4.23) that $-\bar{P}_{02}^2/\bar{P}_{04}$ is bounded from below, see Fig. 9.

Now that all Lagrangian functions and the background evolution of the fields are specified, we solve the equations (4.17) numerically to confirm that $\delta\chi$ is converted into $\tilde{\zeta}$ with high efficiency. The initial condition is

$$\tilde{\zeta}_k(\bar{t}_i) = 10^{-5} \sqrt{\frac{P_1(\bar{t}_i)}{\bar{G}_S(\bar{t}_i)}}, \quad \delta\bar{\chi}_k(\bar{t}_i) = 10^{-5}, \quad \dot{\tilde{\zeta}}_k(\bar{t}_i) = 0, \quad \delta\dot{\bar{\chi}}_k(\bar{t}_i) = 0, \quad (4.25)$$

where \bar{t}_i is the initial time. The parameters are set to be

$$\tau = 10, \quad \epsilon = 10, \quad u = \frac{1}{10}, \quad w = 2, \quad (4.26a)$$

$$b = 0.0158, \quad \bar{t}_c = 10\tau = 100, \quad q = \frac{1}{12}, \quad \lambda = 0.35, \quad \bar{\chi}_0 = \frac{1}{10}. \quad (4.26b)$$

Fig. 10 shows the result of the numeric computation, where the initial time is $\bar{t}_i = -3 \times 10^3$. The conversion process takes place around $4\tau = \bar{t}_c - 6\tau \lesssim \bar{t} \lesssim \bar{t}_c + 6\tau = 16\tau$, during which about 98% of $\delta\bar{\chi}_k$ is converted into $\tilde{\zeta}_k$ and the comoving Hubble radius $(aH)^{-1}$ decreases by order of one e-fold. This implies that the conversion is smooth and highly efficient, cf. [62].

Fig. 11 shows that for the parameters given by (4.26) the constraint (2.23) is satisfied.

The remaining functions $f_{0,1,2}(\bar{t})$ are uniquely determined from (3.24) and the background equations of motion (4.4a) and (4.4b). We emphasize that non-trivial form of $\bar{\chi}$, \bar{P}_{02} and \bar{P}_{04} does not alter the asymptotics of $f_{0,1,2}(\bar{t})$ given in (3.11), (3.9) and (3.10).




Figure 8. The left panel shows plot of \bar{P}_{02} , while the right two panels show \bar{P}_{02} and \bar{P}_{04} behave asymptotically as given in (4.23). The parameters are given by (4.26).




Figure 9. Plot of $-\bar{P}_{02}^2 / \bar{P}_{04}$. The parameters are given by (4.26).




Figure 10. Plot of evolution of $\tilde{\zeta}_k$ and $\delta\bar{\chi}_k$. We set the initial time $\bar{t}_i = -3 \times 10^3$. The parameters are given by (4.26).

4.4 The amplitude of scalar power spectrum and characteristic energy scale

Now we determine the characteristic energy scale μ which is related to the amplitude of the perturbations $\delta\bar{\chi}$. Since $\delta\bar{\chi}$ is almost completely converted into $\tilde{\zeta}$, the power spectrum of $\delta\bar{\chi}$ should be

$$\Delta_{\delta\bar{\chi}}^2(k) \equiv k^3 \langle \delta\bar{\chi}_k^2 \rangle = A_s \left(\frac{k}{k_\star} \right)^{n_s-1}, \quad (4.27)$$

where k_\star denotes the pivot scale $0.05 \text{Mpc}^{-1} \simeq 10^{-59} M_{pl}$, and A_s denotes the measured amplitude at k_\star , namely [50]

$$A_s = (2.105 \pm 0.030) \times 10^{-9}. \quad (4.28)$$

After the mode v_k exits the horizon, its amplitude can be written as

$$a\sqrt{P_1}M_{pl}\delta\bar{\chi}_k = v_k \simeq \frac{\sqrt{-\pi\eta}}{2} \frac{1}{(-k\eta)^\nu}. \quad (4.29)$$

On the other hand, the conformal time η is related to the rescaled cosmic time \bar{t} by

$$\left(\frac{-\bar{t}}{\tau} \right)^{1-\frac{1}{\epsilon}} \sim \frac{\epsilon-1}{\epsilon} \frac{\mu^2}{M_{pl}} \left(\frac{-\eta}{\tau} \right). \quad (4.30)$$




Figure 11. Plot of $\bar{\mathcal{F}}_S$ and $\bar{Y}\bar{P}_Y\bar{g}^2$. The function $\bar{\mathcal{F}}_S$ always dominates over $\bar{Y}\bar{P}_Y\bar{g}^2$ throughout the whole time evolution, which shows the constraint (2.23) is satisfied. The parameters are given by (4.26).

It then follows that

$$a\sqrt{P_1} \sim \left[\frac{\epsilon-1}{\epsilon} \frac{\mu^2}{M_{pl}} \left(\frac{-\eta}{\tau} \right) \right]^{\frac{1}{2}-\nu}, \quad (4.31)$$

and thus

$$\Delta_{\delta\chi}^2(k) \sim \frac{\pi}{4} \left(\frac{\epsilon-1}{\epsilon\tau} \right)^2 \left(\frac{\mu}{M_{pl}} \right)^4 \left(\frac{\epsilon-1}{\epsilon\tau} \frac{\mu^2}{M_{pl}k_*} \right)^{1-n_s} \left(\frac{k}{k_*} \right)^{n_s-1}. \quad (4.32)$$

It follows from (4.32) that for $\mu \sim \mathcal{O}(10^{-2})M_{pl}$ the amplitude of the scalar power spectrum is in good agreement with the Planck data (4.28).

For completeness, we discuss the tensor-to-scalar ratio. Since $\bar{\mathcal{G}}_{\mathcal{T}} = \bar{\mathcal{F}}_{\mathcal{T}} = 1$ in our scenario, it is evident that the tensor perturbations carry a deep blue spectral index (actually equal to $4 - 2\nu_1$, where ν_1 is given by (4.13)). It then follows that at the pivot scale the tensor-to-scalar ratio is negligibly small, which is in fact a typical feature of bouncing scenarios that begin with the ekpyrotic contraction phase. Our scenario is therefore compatible with the observational data, regarding the tensor-to-scalar ratio as well.

5 Summary and Conclusion

In this paper we have constructed a class of DHOST theories with an extra scalar field, which admits viable bouncing solutions over which linear perturbations are free of all pathologies during entire evolution and compatible with observations. In particular, after finding the sufficient conditions for the scalar sector to be stable and non-superluminal (see (2.21)), we constructed for the first time a completely healthy two-field DHOST bounce model, in which the adiabatic field ϕ is described by the DHOST theories and the entropic field χ has luminal sound speed. Our model exhibits the conventional asymptotics for the DHOST sector without requiring strong gravity in the past employed in [30–35]. The overview of structure of the model is given in Table 1.

Table 1: Structure of the two-field model constructed, see (2.1), (5.1) and (5.2).

The exceptional subclass of DHOST Ia + an extra field									
DHOST sector						Extra field sector			
$F(\phi, X)$		$F_2(\phi, X)$		$A_1(\phi, X)$		$P(\chi, Y, \phi)$			
$f_0(\phi)$	$f_1(\phi)$	$f_2(\phi)$	$g_0(\phi)$	$g_1(\phi)$	$a_0(\phi)$	$a_1(\phi)$	$P_{02}(\phi)$	$P_{04}(\phi)$	$P_1(\phi)$

As the main calculations in the previous sections might look quite technical, in the flowchart in Fig. 12 we summarized the essentials/guideline of how the model has been constructed using the inverse method. The result of the found Lagrangian functions in the action (2.1) is given by

$$\mu^{-4}F(\phi, X) = \bar{F}(\bar{\phi}, \bar{X}) = f_0(\bar{\phi}) + f_1(\bar{\phi})\bar{X} + f_2(\bar{\phi})\bar{X}^2, \quad (5.1a)$$

$$M_{pl}^{-2}F_2(\phi, X) = \bar{F}_2(\bar{\phi}, \bar{X}) = -\frac{1}{2} + g_1(\bar{\phi}) - g_1(\bar{\phi})\bar{X}, \quad (5.1b)$$

$$M_{pl}^{-2}\mu^4A_1(\phi, X) = \bar{A}_1(\bar{\phi}, \bar{X}) = -a_1(\bar{\phi}) + a_1(\bar{\phi})\bar{X}, \quad (5.1c)$$

$$\mu^{-1}K(\phi, X) = \bar{K}(\bar{\phi}, \bar{X}) = 0, \quad (5.1d)$$

$$\mu^{-4}P(\chi, Y, \phi) = \bar{P}(\bar{\chi}, \bar{\phi}, \bar{Y}) = P_1(\bar{\phi})\bar{Y} - \bar{P}_{02}(\bar{\phi})\bar{\chi}^2 - \bar{P}_{04}(\bar{\phi})\bar{\chi}^4, \quad (5.1e)$$

where $\bar{\phi}$, \bar{X} , $\bar{\chi}$, \bar{Y} , \bar{F} , \bar{F}_2 , \bar{A}_1 , \bar{P} , \bar{P}_{02} and \bar{P}_{04} are rescaled quantities (see (3.2) and (4.2)), and

$$g_1(\bar{\phi}) = \frac{1}{3}w \cosh^{-2}\left(\frac{\bar{\phi}}{\tau} + u\right), \quad (5.2a)$$

$$a_1(\bar{\phi}) = \frac{w}{12} \operatorname{sech}^2\left(\frac{\bar{\phi}}{\tau} + u\right) \left(\frac{2\left(\bar{\phi}^2 \tanh\left(\frac{\bar{\phi}}{\tau} + u\right) + \tau^2 \tanh\left(\frac{\bar{\phi}}{\tau}\right)\right)}{\mathbf{H}(\bar{\phi})\tau(\bar{\phi}^2 + \tau^2)} - 1 \right), \quad (5.2b)$$

$$P_1(\bar{\phi}) = \frac{1}{2} \frac{\sigma\left(-\frac{\bar{\phi}}{\tau}\right)}{\left(\frac{\bar{\phi}}{\tau}\right)^{2(1+b)} + 1} + \frac{1}{2}\sigma\left(\frac{\bar{\phi}}{\tau}\right), \quad (5.2c)$$




Figure 12. Flowchart of the model construction using the inverse method. The left side shows the steps determining the Lagrangian functions, while the right side shows the corresponding constraints used. On the left side, the teal-colored rounded boxes are related to the DHOST sector, while those in orange correspond to the extra scalar sector.

$$\bar{P}_{04}(\bar{\phi}) = 10^{-10} \text{sech}\left(\frac{\bar{\phi} - \bar{t}_c}{2\tau}\right), \quad (5.2d)$$

$$\bar{P}_{02}(\bar{\phi}) = -2\bar{P}_{04}(\bar{\phi})\chi^2(\bar{\phi}) - \frac{1}{\mathbf{a}^3(\bar{\phi})\chi(\bar{\phi})} \frac{d}{d\bar{\phi}} \left(\mathbf{a}^3(\bar{\phi}) \frac{d\chi(\bar{\phi})}{d\bar{\phi}} P_1(\bar{\phi}) \right), \quad (5.2e)$$

$$\begin{aligned} f_2(\bar{\phi}) = & \frac{1}{8} \left(-30a_1(\bar{\phi}) \frac{dg_1}{d\bar{\phi}} \mathbf{H}(\bar{\phi}) + 2a_1(\bar{\phi}) \frac{d\mathbf{H}}{d\bar{\phi}} g_1(\bar{\phi}) - 6a_1(\bar{\phi}) \frac{d\mathbf{H}}{d\bar{\phi}} + 2 \frac{da_1}{d\bar{\phi}} g_1(\bar{\phi}) \mathbf{H}(\bar{\phi}) \right. \\ & - 6 \frac{da_1}{d\bar{\phi}} \mathbf{H}(\bar{\phi}) + 4 \frac{d^2 g_1}{d\bar{\phi}^2} g_1(\bar{\phi}) - 6 \left(\frac{dg_1}{d\bar{\phi}} \right)^2 + 8 \frac{dg_1}{d\bar{\phi}} g_1(\bar{\phi}) \mathbf{H}(\bar{\phi}) - 9 \frac{dg_1}{d\bar{\phi}} \mathbf{H}^2(\bar{\phi}) \theta(\bar{\phi}) \\ & \left. + 30 \frac{dg_1}{d\bar{\phi}} \mathbf{H}(\bar{\phi}) + 2 \frac{d\mathbf{H}}{d\bar{\phi}} g_1(\bar{\phi})^2 - 16 \frac{d\mathbf{H}}{d\bar{\phi}} g_1(\bar{\phi}) + 2 \frac{d\mathbf{H}}{d\bar{\phi}} + 9g_1(\bar{\phi}) \mathbf{H}^2(\bar{\phi}) \right) \end{aligned}$$

$$+3\mathbf{H}^2(\bar{\phi}) + 2P_1(\bar{\phi}) \left(\frac{d\chi}{d\bar{\phi}} \right)^2 + 2\Sigma(\bar{\phi}) \Big), \quad (5.2f)$$

$$f_1(\bar{\phi}) = \frac{3}{2}h^2(\bar{\phi}) \left(3\frac{dg_1}{d\bar{\phi}}\Theta(\bar{\phi}) - 3g_1(\bar{\phi}) - 1 \right) - 3\frac{d\mathbf{H}(\bar{\phi})}{d\bar{\phi}}g_1(\bar{\phi}) \\ - \frac{d\mathbf{H}(\bar{\phi})}{d\bar{\phi}} - 2f_2(\bar{\phi}) - P_1(\bar{\phi}) \left(\frac{d\chi}{d\bar{\phi}} \right)^2, \quad (5.2g)$$

$$f_0(\bar{\phi}) = \bar{P}_{02}(\bar{\phi})\chi^2(\bar{\phi}) + \bar{P}_{04}(\bar{\phi})\chi^4(\bar{\phi}) - f_1(\bar{\phi}) \\ - \left(\frac{d\chi}{d\bar{\phi}} \right)^2 P_1(\bar{\phi}) - f_2(\bar{\phi}) - 2\frac{dh}{d\bar{\phi}} - 3h^2(\bar{\phi}), \quad (5.2h)$$

with

$$\sigma(\bar{\phi}) = \frac{1}{1 + \exp(-\bar{\phi})}, \quad (5.3a)$$

$$\mathbf{H}(\bar{\phi}) = \frac{d}{d\bar{\phi}} \log \mathbf{a}(\bar{\phi}), \quad (5.3b)$$

$$\mathbf{a}(\bar{\phi}) = \sigma \left(\frac{\bar{\phi}}{\tau} \right) \left[\left(\frac{\bar{\phi}}{\tau} \right)^2 + 1 \right]^{\frac{1}{6}} + \sigma \left(-\frac{\bar{\phi}}{\tau} \right) \left[\left(\frac{\bar{\phi}}{\tau} \right)^2 + 1 \right]^{\frac{1}{2\epsilon}}, \quad (5.3c)$$

$$\Sigma(\bar{\phi}) = \frac{\tau^4}{(\bar{\phi}^2 + \tau^2)^2} + \frac{(\epsilon - 3) \left(1 - \tanh \left(\frac{\bar{\phi}}{\tau} \right) \right)}{2\epsilon^2 (\bar{\phi}^2 + \tau^2)}, \quad (5.3d)$$

$$\Theta(\bar{\phi}) = \mathbf{H}(\bar{\phi}) (4a_1(\bar{\phi}) + g_1(\bar{\phi}) + 1) + \frac{dg_1}{d\bar{\phi}}, \quad (5.3e)$$

$$\chi(\bar{\phi}) = \bar{\chi}_0 + \lambda \operatorname{sech} \left(\frac{\bar{\phi} - \bar{t}_c}{2\tau} \right) \left[e^{-\left(\frac{\bar{\phi} - \bar{t}_c}{4\tau} \right)^2} - q \right]. \quad (5.3f)$$

Here the parameter τ controls the duration of the bounce phase, ϵ characterizes the equation of state during the ekpyrotic contraction phase, b specifies the scalar spectral index, the characteristic energy scale μ determines the amplitude of the curvature perturbations, and \bar{t}_c characterizes the center of conversion period. The parameters u and w have been introduced to avoid the fine-tuning problem that H and Θ cross zero at the same time. The parameters q , $\bar{\chi}_0$ and λ are related to the efficiency of the conversion process. We numerically checked that there are no instabilities and superluminality for the parameters given by (4.26) and the characteristic energy scale $\mu \sim \mathcal{O}(10^{-2})M_{pl}$, see e.g. Figs. 4, 5 and 11.

In summary the model constructed in this work gives rise to a completely healthy and phenomenologically viable bounce scenario, which satisfies the followings:

- it is free of BKL, ghost and gradient instabilities,
- it has conventional asymptotics for DHOST sector,
- it is free of superluminality,
- it predicts nearly scale-invariant curvature perturbations, generated by an entropic mechanism, compatible with observations,

- it predicts a sufficiently low tensor-to-scalar ratio.

From the phenomenological point of view, it is important to make sure that the non-Gaussianities of the curvature perturbations do not exceed the observational bounds. It is actually obvious that the perturbations of the entropic field χ have negligible non-Gaussianities [56, 57]. Thus, the main contribution to the non-Gaussianities of the curvature perturbations should come from the non-linear conversion process from the entropic to curvature perturbations. Since our scenario has efficient (almost 100% of efficiency) and smooth conversion from entropy to curvature perturbations, we expect that the non-Gaussianities would not be generated too much [56, 61, 62]. In any case it would be interesting to confirm this point. We plan to pursue this direction in future work.

A Background equations of motion

This Appendix gives the background equations of motion for spatially-flat FLRW metric, which are obtained by varying the action (2.1) with respect to α , ζ and χ , respectively:

- α :

$$\begin{aligned}
& F + P - 2Y P_Y - 6F_2 H^2 - 6H \dot{\phi} \left(F_{2\phi} - (A_1 - 2F_{2X}) \ddot{\phi} \right) \\
& + \dot{\phi}^2 \left(-6A_1 \dot{H} + A_3 \ddot{\phi}^2 + A_4 \ddot{\phi}^2 - 2F_X + 12F_{2X} \dot{H} + 24F_{2X} H^2 + K_\phi \right) \\
& + \dot{\phi}^3 \left(H \left(-6A_{1\phi} + 3(A_3 + 2A_4) \ddot{\phi} - 6K_X \right) + 2(A_3 + A_4) \ddot{\phi} \right) \\
& + \dot{\phi}^4 \left(3H^2 (4A_{1X} + 3A_3) + \ddot{\phi} \left(\ddot{\phi} (2A_{3X} + 2A_{4X} + 3A_5) + 2A_{3\phi} + 2A_{4\phi} \right) + 3A_3 \dot{H} \right) \\
& + \dot{\phi}^5 \left(3H (A_{3\phi} + 2A_5 \ddot{\phi}) + 2A_5 \ddot{\phi} \right) + 2\dot{\phi}^6 \ddot{\phi} \left(A_{5X} \ddot{\phi} + A_{5\phi} \right) = 0 \tag{A.1}
\end{aligned}$$

- ζ :

$$\begin{aligned}
& 2A_1 \ddot{\phi}^2 + F + P - 4F_2 \dot{H} - 6F_2 H^2 - 4F_{2X} \ddot{\phi}^2 - 2F_{2\phi} \ddot{\phi} \\
& + 2\dot{\phi} \left[\ddot{\phi} (A_1 - 2F_{2X}) - 2H \left(2(F_{2X} - A_1) \ddot{\phi} + F_{2\phi} \right) \right] \\
& - \dot{\phi}^2 \left[-4A_1 \dot{H} - 6A_1 H^2 - 4A_{1X} \ddot{\phi}^2 - 2A_{1\phi} \ddot{\phi} + 2A_3 \ddot{\phi}^2 - A_4 \ddot{\phi}^2 \right. \\
& \quad \left. + 8F_{2XX} \ddot{\phi}^2 + 8F_{2\phi X} \ddot{\phi} + 2F_{2\phi\phi} + 2K_X \ddot{\phi} + K_\phi \right] \\
& + \dot{\phi}^3 \left[4H \left(2A_{1X} \ddot{\phi} + A_{1\phi} \right) - A_3 \ddot{\phi} \right] + \dot{\phi}^4 \ddot{\phi} \left[(A_5 - 2A_{3X}) \ddot{\phi} - A_{3\phi} \right] = 0 \tag{A.2}
\end{aligned}$$

- χ :

$$\frac{1}{a^3} \frac{d}{dt} (a^3 \dot{\chi} P_Y) - \frac{1}{2} P_\chi = 0. \tag{A.3}$$

One can readily check that in the case of single-field Beyond Horndeski theories where $P = 0$, $A_1 = -XF_4 - 2G_{4X}$, $A_3 = -2F_4$, $A_4 = 2F_4$ and $A_5 = 0$ the above background equations of motion coincide with those given in [27].

B Exceptional Subclass

For the exceptional subclass of DHOST Ia theories, the Lagrangian functions A_4 and A_5 become

$$A_4 = - \frac{2(A_1 - 2F_{2X}) (9A_1^2 X^2 - 17A_1 F_2 X + 2X F_{2X} (7F_2 - 6A_1 X) + 8F_2^2)}{X(3A_1 X - 4F_2)^2}, \tag{B.1a}$$

$$A_5 = \frac{2(A_1 - 2F_{2X})^2}{X(3A_1 X - 4F_2)}. \tag{B.1b}$$

Inserting the ansatz (3.7) into (2.17) and (B.1), we obtain

$$A_3 = \frac{2[a_0(\phi) + X a_1(\phi) + 2g_1(\phi)] [X(a_0(\phi) + X a_1(\phi) + 2g_1(\phi)) + 2g_0(\phi) + 1]}{X [X(3a_0(\phi) + 3X a_1(\phi) + 4g_1(\phi)) + 4g_0(\phi) + 2]}, \tag{B.2a}$$

$$A_4 = - \frac{(a_0(\phi) + Xa_1(\phi) + 2g_1(\phi))}{X(X(3a_0(\phi) + 3Xa_1(\phi) + 4g_1(\phi)) + 4g_0(\phi) + 2)^2} \times$$

$$\times [Xa_0(\phi) (36X^2a_1(\phi) + 34g_0(\phi) + 58Xg_1(\phi) + 17) + 18X^2a_0(\phi)^2$$

$$+ 18X^4a_1(\phi)^2 + X^2a_1(\phi)(34g_0(\phi) + 58Xg_1(\phi) + 17)$$

$$+ 2(2g_0(\phi) + 2Xg_1(\phi) + 1)(4g_0(\phi) + 11Xg_1(\phi) + 2)], \quad (\text{B.2b})$$

$$A_5 = \frac{2[a_0(\phi) + Xa_1(\phi) + 2g_1(\phi)]^2}{X[X(3a_0(\phi) + 3Xa_1(\phi) + 4g_1(\phi)) + 4g_0(\phi) + 2]}. \quad (\text{B.2c})$$

References

- [1] A. H. Guth, *The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems*, *Phys. Rev. D* **23** (1981) 347.
- [2] A. A. Starobinsky, *A New Type of Isotropic Cosmological Models Without Singularity*, *Phys. Lett. B* **91** (1980) 99.
- [3] A. D. Linde, *A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems*, *Phys. Lett. B* **108** (1982) 389.
- [4] A. Borde, A. H. Guth and A. Vilenkin, *Inflationary space-times are incomplete in past directions*, *Phys. Rev. Lett.* **90** (2003) 151301 [[gr-qc/0110012](#)].
- [5] M. Novello and S. E. P. Bergliaffa, *Bouncing Cosmologies*, *Phys. Rept.* **463** (2008) 127 [[0802.1634](#)].
- [6] D. Battfeld and P. Peter, *A Critical Review of Classical Bouncing Cosmologies*, *Phys. Rept.* **571** (2015) 1 [[1406.2790](#)].
- [7] R. Brandenberger and P. Peter, *Bouncing Cosmologies: Progress and Problems*, *Found. Phys.* **47** (2017) 797 [[1603.05834](#)].
- [8] V. A. Rubakov, *The Null Energy Condition and its violation*, *Phys. Usp.* **57** (2014) 128 [[1401.4024](#)].
- [9] S. Dubovsky, T. Gregoire, A. Nicolis and R. Rattazzi, *Null energy condition and superluminal propagation*, *JHEP* **03** (2006) 025 [[hep-th/0512260](#)].
- [10] M. Libanov, S. Mironov and V. Rubakov, *Generalized Galileons: instabilities of bouncing and Genesis cosmologies and modified Genesis*, *JCAP* **08** (2016) 037 [[1605.05992](#)].
- [11] T. Kobayashi, *Generic instabilities of nonsingular cosmologies in Horndeski theory: A no-go theorem*, *Phys. Rev. D* **94** (2016) 043511 [[1606.05831](#)].
- [12] G. W. Horndeski, *Second-order scalar-tensor field equations in a four-dimensional space*, *Int. J. Theor. Phys.* **10** (1974) 363.
- [13] M. Zumalacárregui and J. García-Bellido, *Transforming gravity: from derivative couplings to matter to second-order scalar-tensor theories beyond the Horndeski Lagrangian*, *Phys. Rev. D* **89** (2014) 064046 [[1308.4685](#)].
- [14] J. Gleyzes, D. Langlois, F. Piazza and F. Vernizzi, *Healthy theories beyond Horndeski*, *Phys. Rev. Lett.* **114** (2015) 211101 [[1404.6495](#)].
- [15] D. Langlois and K. Noui, *Degenerate higher derivative theories beyond Horndeski: evading the Ostrogradski instability*, *JCAP* **02** (2016) 034 [[1510.06930](#)].

- [16] J. Ben Achour, D. Langlois and K. Noui, *Degenerate higher order scalar-tensor theories beyond Horndeski and disformal transformations*, *Phys. Rev. D* **93** (2016) 124005 [[1602.08398](#)].
- [17] D. Langlois, M. Mancarella, K. Noui and F. Vernizzi, *Effective Description of Higher-Order Scalar-Tensor Theories*, *JCAP* **05** (2017) 033 [[1703.03797](#)].
- [18] D. Langlois, *Dark energy and modified gravity in degenerate higher-order scalar-tensor (DHOST) theories: A review*, *Int. J. Mod. Phys. D* **28** (2019) 1942006 [[1811.06271](#)].
- [19] T. Kobayashi, *Horndeski theory and beyond: a review*, *Rept. Prog. Phys.* **82** (2019) 086901 [[1901.07183](#)].
- [20] P. Creminelli, D. Pirtskhalava, L. Santoni and E. Trincherini, *Stability of Geodesically Complete Cosmologies*, *JCAP* **11** (2016) 047 [[1610.04207](#)].
- [21] R. Kolevatov, S. Mironov, N. Sukhov and V. Volkova, *Cosmological bounce and Genesis beyond Horndeski*, *JCAP* **08** (2017) 038 [[1705.06626](#)].
- [22] G. Ye and Y.-S. Piao, *Bounce in general relativity and higher-order derivative operators*, *Phys. Rev. D* **99** (2019) 084019 [[1901.08283](#)].
- [23] G. Ye and Y.-S. Piao, *Implication of GW170817 for cosmological bounces*, *Commun. Theor. Phys.* **71** (2019) 427 [[1901.02202](#)].
- [24] A. Ilyas, M. Zhu, Y. Zheng, Y.-F. Cai and E. N. Saridakis, *DHOST Bounce*, *JCAP* **09** (2020) 002 [[2002.08269](#)].
- [25] M. Zhu, A. Ilyas, Y. Zheng, Y.-F. Cai and E. N. Saridakis, *Scalar and tensor perturbations in DHOST bounce cosmology*, *JCAP* **11** (2021) 045 [[2108.01339](#)].
- [26] A. Adams, N. Arkani-Hamed, S. Dubovsky, A. Nicolis and R. Rattazzi, *Causality, analyticity and an IR obstruction to UV completion*, *JHEP* **10** (2006) 014 [[hep-th/0602178](#)].
- [27] S. Mironov, V. Rubakov and V. Volkova, *Bounce beyond Horndeski with GR asymptotics and γ -crossing*, *JCAP* **10** (2018) 050 [[1807.08361](#)].
- [28] S. Mironov, V. Rubakov and V. Volkova, *Subluminal cosmological bounce beyond Horndeski*, *JCAP* **05** (2020) 024 [[1910.07019](#)].
- [29] S. Mironov, V. Rubakov and V. Volkova, *Superluminality in beyond Horndeski theory with extra scalar field*, *Phys. Scripta* **95** (2020) 084002 [[2005.12626](#)].
- [30] Y. A. Ageeva, O. A. Evseev, O. I. Melichev and V. A. Rubakov, *Horndeski Genesis: strong coupling and absence thereof*, *EPJ Web Conf.* **191** (2018) 07010 [[1810.00465](#)].
- [31] Y. Ageeva, P. Petrov and V. Rubakov, *Horndeski genesis: consistency of classical theory*, *JHEP* **12** (2020) 107 [[2009.05071](#)].
- [32] Y. Ageeva, O. Evseev, O. Melichev and V. Rubakov, *Toward evading the strong coupling problem in Horndeski genesis*, *Phys. Rev. D* **102** (2020) 023519 [[2003.01202](#)].
- [33] Y. Ageeva, P. Petrov and V. Rubakov, *Nonsingular cosmological models with strong gravity in the past*, *Phys. Rev. D* **104** (2021) 063530 [[2104.13412](#)].
- [34] Y. Ageeva and P. Petrov, *On the strong coupling problem in cosmologies with "strong gravity in the past"*, *Mod. Phys. Lett. A* **37** (2022) 2250171 [[2206.10646](#)].
- [35] Y. Ageeva, P. Petrov and V. Rubakov, *Generating cosmological perturbations at Horndeski bounce*, *JHEP* **01** (2022) [[2207.04071](#)].

- [36] S. Mironov, V. Rubakov and V. Volkova, *Superluminality in DHOST theory with extra scalar*, *JHEP* **04** (2021) 035 [[2011.14912](#)].
- [37] S. Mironov and V. Volkova, *Non-singular cosmological scenarios in scalar-tensor theories and their stability: a review*, [2409.16108](#).
- [38] Y. Ageeva, M. Kottenko and P. Petrov, *The primordial non-Gaussianities for non-singular Horndeski cosmologies*, [2410.10742](#).
- [39] A. Ijjas and P. J. Steinhardt, *Classically stable nonsingular cosmological bounces*, *Phys. Rev. Lett.* **117** (2016) 121304 [[1606.08880](#)].
- [40] A. Ijjas and P. J. Steinhardt, *Fully stable cosmological solutions with a non-singular classical bounce*, *Phys. Lett. B* **764** (2017) 289 [[1609.01253](#)].
- [41] Y. Cai and Y.-S. Piao, *A covariant Lagrangian for stable nonsingular bounce*, *JHEP* **09** (2017) 027 [[1705.03401](#)].
- [42] V. Belinsky, I. Khalatnikov and E. Lifshitz, *Oscillatory approach to a singular point in the relativistic cosmology*, *Adv. Phys.* **19** (1970) 525.
- [43] J. Khoury, B. A. Ovrut, P. J. Steinhardt and N. Turok, *The Ekpyrotic universe: Colliding branes and the origin of the hot big bang*, *Phys. Rev. D* **64** (2001) 123522 [[hep-th/0103239](#)].
- [44] E. I. Buchbinder, J. Khoury and B. A. Ovrut, *New Ekpyrotic cosmology*, *Phys. Rev. D* **76** (2007) 123503 [[hep-th/0702154](#)].
- [45] A. M. Levy, A. Ijjas and P. J. Steinhardt, *Scale-invariant perturbations in ekpyrotic cosmologies without fine-tuning of initial conditions*, *Phys. Rev. D* **92** (2015) 063524 [[1506.01011](#)].
- [46] W. G. Cook, I. A. Glushchenko, A. Ijjas, F. Pretorius and P. J. Steinhardt, *Supersmoothing through Slow Contraction*, *Phys. Lett. B* **808** (2020) 135690 [[2006.01172](#)].
- [47] A. Ijjas, W. G. Cook, F. Pretorius, P. J. Steinhardt and E. Y. Davies, *Robustness of slow contraction to cosmic initial conditions*, *JCAP* **08** (2020) 030 [[2006.04999](#)].
- [48] A. Ijjas, A. P. Sullivan, F. Pretorius, P. J. Steinhardt and W. G. Cook, *Ultralocality and slow contraction*, *JCAP* **06** (2021) 013 [[2103.00584](#)].
- [49] A. Ijjas, P. J. Steinhardt, D. Garfinkle and W. G. Cook, *Smoothing and flattening the universe through slow contraction versus inflation*, [2404.00867](#).
- [50] PLANCK collaboration, *Planck 2018 results. VI. Cosmological parameters*, *Astron. Astrophys.* **641** (2020) A6 [[1807.06209](#)].
- [51] A. Notari and A. Riotto, *Isocurvature perturbations in the ekpyrotic universe*, *Nucl. Phys. B* **644** (2002) 371 [[hep-th/0205019](#)].
- [52] F. Finelli, *Assisted contraction*, *Phys. Lett. B* **545** (2002) 1 [[hep-th/0206112](#)].
- [53] J.-L. Lehners, P. McFadden, N. Turok and P. J. Steinhardt, *Generating ekpyrotic curvature perturbations before the big bang*, *Phys. Rev. D* **76** (2007) 103501 [[hep-th/0702153](#)].
- [54] T. Qiu, X. Gao and E. N. Saridakis, *Towards anisotropy-free and nonsingular bounce cosmology with scale-invariant perturbations*, *Phys. Rev. D* **88** (2013) 043525 [[1303.2372](#)].
- [55] M. Li, *Note on the production of scale-invariant entropy perturbation in the Ekpyrotic universe*, *Phys. Lett. B* **724** (2013) 192 [[1306.0191](#)].

- [56] A. Fertig, J.-L. Lehnert and E. Mallwitz, *Ekpyrotic Perturbations With Small Non-Gaussian Corrections*, *Phys. Rev. D* **89** (2014) 103537 [[1310.8133](#)].
- [57] A. Ijjas, J.-L. Lehnert and P. J. Steinhardt, *General mechanism for producing scale-invariant perturbations and small non-Gaussianity in ekpyrotic models*, *Phys. Rev. D* **89** (2014) 123520 [[1404.1265](#)].
- [58] E. I. Buchbinder, J. Khoury and B. A. Ovrut, *On the Initial Conditions in New Ekpyrotic Cosmology*, *JHEP* **11** (2007) 076 [[0706.3903](#)].
- [59] J.-L. Lehnert and P. J. Steinhardt, *Non-Gaussianity Generated by the Entropic Mechanism in Bouncing Cosmologies Made Simple*, *Phys. Rev. D* **80** (2009) 103520 [[0909.2558](#)].
- [60] J.-L. Lehnert, *Ekpyrotic Non-Gaussianity: A Review*, *Adv. Astron.* **2010** (2010) 903907 [[1001.3125](#)].
- [61] A. Fertig and J. L. Lehnert, *The Non-Minimal Ekpyrotic Trispectrum*, *JCAP* **01** (2016) 026 [[1510.03439](#)].
- [62] A. Fertig, J.-L. Lehnert, E. Mallwitz and E. Wilson-Ewing, *Converting entropy to curvature perturbations after a cosmic bounce*, *JCAP* **10** (2016) 005 [[1607.05663](#)].
- [63] A. Ijjas and R. Kolevatov, *Sourcing curvature modes with entropy perturbations in non-singular bouncing cosmologies*, *JCAP* **06** (2021) 012 [[2012.08249](#)].
- [64] A. Ijjas and R. Kolevatov, *Nearly scale-invariant curvature modes from entropy perturbations during the graceful exit phase*, *Phys. Rev. D* **103** (2021) L101302 [[2102.03818](#)].
- [65] C. Armendariz-Picon, V. F. Mukhanov and P. J. Steinhardt, *Essentials of k essence*, *Phys. Rev.* **D63** (2001) 103510 [[astro-ph/0006373](#)].
- [66] C. Deffayet and S. Garcia-Saenz, *Degeneracy, matter coupling, and disformal transformations in scalar-tensor theories*, *Phys. Rev. D* **102** (2020) 064037 [[2004.11619](#)].
- [67] I. P. C. Heard and D. Wands, *Cosmology with positive and negative exponential potentials*, *Class. Quant. Grav.* **19** (2002) 5435 [[gr-qc/0206085](#)].
- [68] M. Koehn, J.-L. Lehnert and B. Ovrut, *Nonsingular bouncing cosmology: Consistency of the effective description*, *Phys. Rev. D* **93** (2016) 103501 [[1512.03807](#)].
- [69] J.-L. Lehnert and P. J. Steinhardt, *Non-Gaussian density fluctuations from entropically generated curvature perturbations in Ekpyrotic models*, *Phys. Rev. D* **77** (2008) 063533 [[0712.3779](#)].