A Family of Controllable Momentum Coefficients for Forward-Backward Accelerated Algorithms

Mingwei $Fu¹$ and Bin Shi^{*} ^{2,3}

¹School of Mathematical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

²Shanghai Institute for Mathematics and Interdisciplinary Sciences, Shanghai 200433, China

³Center for Mathematics and Interdisciplinary Sciences, Fudan University, Shanghai 200433, China

January 20, 2025

Abstract

Nesterov's accelerated gradient method (NAG) marks a pivotal advancement in gradient-based optimization, achieving faster convergence compared to the vanilla gradient descent method for convex functions. However, its algorithmic complexity when applied to strongly convex functions remains unknown, as noted in the comprehensive review by [Chambolle and Pock](#page-20-0) [\[2016\]](#page-20-0). This issue, aside from the critical step size, was addressed by [Li et al.](#page-21-0) [\[2024b](#page-21-0)], with the monotonic case further explored by [Fu and Shi](#page-20-1) [\[2024\]](#page-20-1). In this paper, we introduce a family of controllable momentum coefficients for forward-backward accelerated methods, focusing on the critical step size $s = 1/L$. Unlike traditional linear forms, the proposed momentum coefficients follow an α -th power structure, where the parameter r is adaptively tuned to α . Using a Lyapunov function specifically designed for α , we establish a controllable $O(1/k^{2\alpha})$ convergence rate for the NAG- α method, provided that $r > 2\alpha$. At the critical step size, NAG- α achieves an inverse polynomial convergence rate of arbitrary degree by adjusting r according to $\alpha > 0$. We further simplify the Lyapunov function by expressing it in terms of the iterative sequences x_k and y_k . eliminating the need for phase-space representations. This simplification enables us to extend the controllable $O(1/k^{2\alpha})$ rate to the monotonic variant, M-NAG- α , thereby enhancing optimization efficiency. Finally, by leveraging the fundamental inequality for composite functions, we extended the controllable $O(1/k^{2\alpha})$ rate to proximal algorithms, including the fast iterative shrinkage-thresholding algorithm (FISTA- α) and its monotonic counterpart (M-FISTA- α).

1 Introduction

Over the past few decades, machine learning has experienced unprecedented growth, driving transformative advancements across a wide range of scientific disciplines and industrial applications. At

[∗]Corresponding author: <binshi@fudan.edu.cn>

the heart of this revolution lies the fundamental challenge of unconstrained optimization, particularly the task of minimizing a convex objective function without constraints. Mathematically, this problem can be expressed as:

$$
\min_{x \in \mathbb{R}^d} f(x).
$$

Addressing this problem has been a central focus of optimization research, leading to the development of gradient-based algorithms, which form the backbone of modern machine learning methodologies. These algorithms are celebrated for their computational efficiency and low memory requirements, making them particularly well-suited for large-scale problems, where computational and memory resources are often limited. Beyond their practical utility, gradient-based methods have also catalyzed profound theoretical advancements while enabled impactful real-world applications. As a result, these algorithms have now become indispensable tools in both academic research and industrial practice.

Among the family of gradient-based methods, the vanilla gradient descent method stands out as one of the simplest yet most fundamental approaches, which can be traced back to to the pioneering work of [Cauchy](#page-20-2) [\[1847\]](#page-20-2). Starting from an initial point $x_0 \in \mathbb{R}^d$, it follows a straightforward iterative update rule:

$$
x_k = x_{k-1} - s\nabla f(x_{k-1}),
$$

where $s > 0$ denotes the step size. For convex functions, the vanilla gradient descent method guarantees convergence with the following rate:

$$
f(x_k) - f(x^*) \le O\left(\frac{1}{sk}\right),\,
$$

where x^* denotes the global minimum point of $f(x)$. In the mid-1980s, [Nesterov](#page-21-1) [\[1983](#page-21-1)] introduced a groundbreaking innovation that revolutionized optimization theory: the accelerated gradient method, now widely known as Nesterov's accelerated gradient method (NAG). This method incorporates an elegant two-step mechanism to achieve faster convergence. Given an initial point $x_0 = y_0 \in \mathbb{R}^d$, NAG is implemented through the following iterative update rules:

$$
\begin{cases} x_k = y_{k-1} - s \nabla f(y_{k-1}), \\ y_k = x_k + \frac{k-1}{k+r} (x_k - x_{k-1}), \end{cases}
$$

where $s > 0$ is the step size, and $r \geq 2$ is the momentum coefficient. Compared to the vanilla gradient descent, NAG achieves an accelerated convergence rate:

$$
f(x_k) - f(x^*) \le O\left(\frac{1}{sk^2}\right).
$$

This acceleration represents a major leap in optimization efficiency, particularly for large-scale problems. To extend the applicability to composite optimization problems, especially those arising in image science, [Beck and Teboulle](#page-20-3) [\[2009\]](#page-20-3) introduced a proximal variant of NAG, known as the fast iterative shrinkage-thresholding algorithm (FISTA). This approach was later refined by [Beck](#page-20-4) [\[2017\]](#page-20-4), who introduced monotonic versions of both NAG and FISTA. Collectively, these algorithms are referred to as forward-backward accelerated gradient algorithms, highlighting their forwardbackward structure.

Despite their solid theoretical foundations, these forward-backward accelerated gradient algorithms have been primarily designed for convex functions. However, many real-world applications often involve strongly convex but ill-conditioned, characterized by Hessian eigenvalues bounded away from zero. For such strongly convex functions, the forward-backward accelerated gradient algorithms have yet to demonstrate superior performance over the vanilla gradient descent and its variants. Indeed, a critical question remains unsolved: whether these forward-backward algorithms can achieve linear convergence for strongly convex functions. This challenge was underscored in the seminal review by [Chambolle and Pock](#page-20-0) [\[2016](#page-20-0)]. Recent efforts have sought to address this gap. Building on the high-resolution differential equation framework [\[Shi et al.](#page-21-2), [2022\]](#page-21-2), [Li et al.](#page-20-5) [\[2024a](#page-20-5)] analyzed forward-backward accelerated method for step sizes $0 < s < 1/L$, demonstrating linear convergence.[1](#page-2-0) Additionally, [Fu and Shi](#page-20-1) [\[2024\]](#page-20-1) developed a novel Lyapunov function, which extends linear convergence to monotonic variants of these algorithms. Nevertheless, a critical limitation persists: for the critical step size $s = 1/L$, no theoretical guarantees currently exist for the forwardbackward accelerated gradient algorithms. This unresolved issue raises two fundamental questions in the context of strongly convex functions and the critical step size $s = 1/L$:

- How fast can forward-backward accelerated gradient algorithms converge?
- Can the forward-backward accelerated gradient algorithms be improved to achieve faster convergence?

Answering these questions represents a pivotal challenge in optimization theory, with the potential to further bridge the gap between theoretical developments and practical applications in machine learning.

1.1 Motivation: a family of controllable momentum coefficients

The classical NAG method leverages a momentum coefficient, which is typically expressed as:

$$
\frac{k-1}{k+r},
$$

where k denotes the iteration number, and r is a constant that governs the strength of the momentum term. This simple yet effective formulation has consistently demonstrated reliable performance across various convex optimization problems. However, its inherent simplicity imposes a notable limitation: the rigidity of this momentum coefficient restricts the algorithm's ability to adapt to problem-specific structures. This limitation becomes particularly pronounced in scenarios involving strongly convex or ill-conditioned objective functions, where a more flexible approach could better exploit the underlying structure for improved performance.

To address these limitations, we introduce a more generalized and flexible momentum coefficient. Specifically, we extend the classical linear form to an α -th power formulation:

$$
\frac{(k-1)^{\alpha}}{k^{\alpha}+rk^{\alpha-1}},\tag{1.1}
$$

where α is a tunable parameter that controls the degree of acceleration, and r is a controllable parameter that can be adjusted in accordance with the value of α . This generalized momentum

¹The parameter L is the Lipschitz constant, which is defined in detail in Section [2.](#page-5-0)

coefficient introduces enhanced adaptability, enabling the algorithm to dynamically tailor its momentum dynamics to the specific characteristics of the problem. By leveraging this flexibility, the proposed formulation has the potential to achieve faster convergence rates, particularly in challenging optimization landscapes.

Building upon this new momentum coefficient (1.1) , we propose an extended variant of the classical NAG method, termed NAG- α . The forward-backward update rules for this generalized algorithm, incorporating the new momentum coefficient from (1.1) , are as follows:

$$
\int x_k = y_{k-1} - s \nabla f(y_{k-1})
$$
\n(1.2a)

$$
\begin{cases} y_k = x_k + \frac{(k-1)^{\alpha}}{k^{\alpha} + rk^{\alpha-1}} (x_k - x_{k-1}) \end{cases}
$$
 (1.2b)

where $s > 0$ is the step size. This extension allows for greater control over the acceleration dynamics through the parameter α , making the algorithm better suited for a broader range of optimization problems. Inspired by prior works [\[Beck](#page-20-4), [2017](#page-20-4), [Fu and Shi,](#page-20-1) [2024](#page-20-1)], we further extend the momen-tum coefficient [\(1.1\)](#page-2-1) to construct a monotonic variant of NAG- α , which we denote as M-NAG- α . This variant incorporates an adaptive mechanism to ensure monotonicity in the progression of the function values, enhancing the robustness of the method. The update rules for $M-NAG-\alpha$ are defined as follows:

$$
(1.3a)
$$
\n
$$
z_{k-1} = y_{k-1} - s \nabla f(y_{k-1}),
$$
\n(1.3a)

$$
x_k = \begin{cases} z_{k-1}, & \text{if } f(z_{k-1}) \le f(x_{k-1}), \\ x_{k-1}, & \text{if } f(z_{k-1}) > f(x_{k-1}), \end{cases} \tag{1.3b}
$$

$$
\left(y_k = x_k + \frac{(k-1)^{\alpha}}{k^{\alpha} + rk^{\alpha-1}}(x_k - x_{k-1}) + \frac{(k-1)^{\alpha} + r(k-1)^{\alpha-1}}{k^{\alpha} + rk^{\alpha-1}}(z_{k-1} - x_k),\right)
$$
(1.3c)

where $s > 0$ is the step size. This monotonic extension integrates momentum-based acceleration with function-value monotonicity, thereby improving the algorithm's versatility and ensuring more stable convergence behavior.

To evaluate the practical performance of NAG- α and M-NAG- α , we conduct numerical experiments on a quadratic function under the critical step size. Figure [1](#page-4-0) illustrates the iterative progression of function values for both algorithms, highlighting the impact of varying α . The numerical results demonstrate that, as α increases and $r = 2\alpha + 1$ is appropriately adjusted, both NAG- α and M-NAG- α exhibit progressively faster convergence rates. These observations underscore the potential of the generalized momentum coefficient to significantly enhance algorithmic efficiency, leading to the following key conjectures:

- For strongly convex functions, the convergence rates of both NAG- α and M-NAG- α vary with the parameter α , provided that the control parameter r is appropriately tuned.
- At the critical step size $s = 1/L$, both NAG- α and M-NAG- α are expected to offer theoretical guarantees for faster convergence compared to the classical NAG method and its monotonic variant.

Figure 1: Iterative progression of function values for NAG- α and M-NAG- α applied to the quadratic function $f(x_1, x_2) = 5 \times 10^{-3} x_1^2 + x_2^2$. The experiments are performed with $s = 1/L = 0.5$.

1.2 Overview of contributions

In this paper, we introduce a novel family of momentum coefficients for forward-backward accelerated methods and thoroughly investigate their convergence behavior, with a particular focus on the critical step size $s = 1/L$. Instead of the conventional linear form, the momentum coefficient takes the α -th power form, where the parameter r is both controlled and adaptively tuned to the value of α .

- (I) We begin by constructing a novel Lyapunov function tailored to the momentum parameter α , which intentionally excludes the kinetic energy term. This formulation allows us to derive a controllable $O(1/k^{2\alpha})$ convergence rate for the NAG- α method, provided that the parameter r satisfies the condition $r > 2\alpha$. At the critical step size $s = 1/L$, the NAG- α method achieves an inverse polynomial convergence rate with arbitrary degree by adaptively tuning the parameter r in accordance with the parameter $\alpha > 0$.
- (II) The Lyapunov function that we construct for $NAG-\alpha$, by omitting the kinetic energy term, provides a simplified way to express the kinetic energy directly in terms of the iterative sequences x_k and y_k , avoiding the need for phase-space representations. This key observation enables us to extend the controllable $O(1/k^{2\alpha})$ convergence rate from NAG- α to its monotonic variant, M-NAG- α , offering an efficient method for accelerated optimization.
- (III) Building on the fundamental inequality for composite functions derived in [\[Li et al.](#page-21-0), [2024b\]](#page-21-0), which serves as the proximal counterpart to the strongly convex inequality, we extend the controllable $O(1/k^{2\alpha})$ convergence rate to the two proximal algorithms, the fast iterative shrinkage thresholding algorithm (FISTA- α) and its monotonic counterpart (M-FISTA- α).

1.3 Related works and organization

The introduction of NAG, originally proposed by [Nesterov](#page-21-1) [\[1983](#page-21-1)], represents a landmark moment in the evolution of gradient-based optimization algorithms. This two-step forward-backward algorithm fundamentally transformed optimization by incorporating acceleration into the traditional gradient descent method, resulting in significant improvements in convergence rates. Building upon this breakthrough, [Beck and Teboulle](#page-20-3) [\[2009](#page-20-3)] introduced a proximal version of the fundamental inequality, which paved the way for the development of FISTA. This new algorithm extended the power of NAG to handle composite objective functions, making it a more efficient tool for a wide range of applications, including signal processing and image reconstruction. [Beck](#page-20-4) [\[2017](#page-20-4)] further advanced this field by proposing a monotonically convergent version of the forward-backward accelerated gradient algorithms. This variant ensured that the performance of the algorithm would consistently improve over time. Simultaneously, adaptive restart strategies were explored to maintain monotonicity and enhance the robustness of forward-backward accelerated algorithms, with some contributions from the works of [Giselsson and Boyd](#page-20-6) [\[2014](#page-20-6)] and [O'Donoghue and Candes](#page-21-3) [\[2015\]](#page-21-3) .

Over the past decade, there has been growing interest in understanding the acceleration phenomenon in optimization algorithms through the lens of differential equations. This exploration began with works [\[Attouch et al.](#page-20-7), [2012](#page-20-7), [2014](#page-20-8)], which sparked a surge of research into the dynamics underlying accelerated methods. The topic gained further prominence with the development of a low-resolution differential equation for modeling NAG [\[Su et al.](#page-21-4), [2016](#page-21-4)], a variational perspective on acceleration [\[Wibisono et al.](#page-21-5), [2016\]](#page-21-5), and studies focusing on the faster convergence rate of function values [\[Attouch and Peypouquet,](#page-20-9) [2016](#page-20-9)]. The acceleration mechanism was ultimately clarified through comparisons between NAG method with Polyak's heavy-ball method, with pivotal insights emerging from the high-resolution differential equation framework introduced by [Shi et al.](#page-21-2) [\[2022\]](#page-21-2). This framework revealed that gradient correction is effectively achieved through an implicit velocity update, providing a more comprehensive understanding of how acceleration arises. Subsequent works [\[Chen et al.](#page-20-10), [2022a](#page-20-10), b] further demonstrated that this framework is particularly well-suited for NAG. Buildling on this insights, significant progress was made by extending the framework to address composite optimization problems, encompassing both convex and strongly convex functions [\[Li et al.](#page-20-12), [2022b](#page-20-12)[,a\]](#page-20-13). This extension was achieved by refining the proximal inequality and generalizing the framework to accommodate a wider range of optimization scenarios, including the overdamped case, as shown in [\[Chen et al.](#page-20-14), [2023](#page-20-14)]. These advancements have greatly enhanced our understanding of acceleration mechanisms and paved the way for more efficient, robust, and versatile optimization algorithms.

The remainder of this paper is structured as follows. Section [2](#page-5-0) introduces the definitions and fundamental inequalities for strongly convex functions and composite functions, laying the ground-work for the analysis. Section [3](#page-7-0) outlines the construction of the Lyapunov function and establishes the controllable $O(1/k^{2\alpha})$ convergence rate for both NAG- α and FISTA- α . Section [4](#page-14-0) extends he controllable $O(1/k^{2\alpha})$ convergence rate to the monotonically accelerated forward-backward algorithms, M-NAG- α and M-FISTA- α , leveraging the novel Lyapnov function that excludes the kinetic energy. Finally, Section [5](#page-19-0) concludes this papers and proposes potential avenues for future research.

2 Preliminaries

In this paper, we adopt notations that align closely with the conventions established in [\[Nesterov](#page-21-6), [2018,](#page-21-6) [Shi et al.](#page-21-2), [2022](#page-21-2), [Li et al.](#page-21-0), [2024b](#page-21-0)], while incorporating slight modifications tailored to the specific requirements of our analysis. Let $\mathcal{F}^0(\mathbb{R}^d)$ denote the class of continuous convex functions defined on \mathbb{R}^d . Specifically, a continuous function $g \in \mathcal{F}^0(\mathbb{R}^d)$ if it satisfies the convex condition:

$$
g(\alpha x + (1 - \alpha)y) \leq \alpha g(x) + (1 - \alpha)g(y), \quad \forall x, y \in \mathbb{R}^d, \ \alpha \in [0, 1].
$$

Within this class $\mathcal{F}^0(\mathbb{R}^d)$, we define the subclass $\mathcal{F}_L^1(\mathbb{R}^d)$, which consists of continuously differentiable functions with Lipschitz continuous gradients. Specifically, a function $f \in \mathcal{F}_L^1(\mathbb{R}^d)$ if $f \in \mathcal{F}^0(\mathbb{R}^d)$ is differentiable, and satisfies the Lipschitz gradient condition:

$$
\|\nabla f(x) - \nabla f(y)\| \le L\|x - y\|, \quad \forall x, y \in \mathbb{R}^d.
$$
\n(2.1)

We further narrow our focus to the class $\mathcal{S}_{\mu,L}^1(\mathbb{R}^d)$, a subclass of $\mathcal{F}_L^1(\mathbb{R}^d)$, where each function is μ -strongly convex for some $0 < \mu \leq L$. Specifically, a function $f \in \mathcal{S}_{\mu,L}^1(\mathbb{R}^d)$ if $f \in \mathcal{F}_L^1(\mathbb{R}^d)$ and satisfies the μ -strongly convex condition:

$$
f(y) \ge f(x) + \langle \nabla f(y), y - x \rangle + \frac{\mu}{2} \|y - x\|^2, \quad \forall x, y \in \mathbb{R}^d.
$$
 (2.2)

For any $f \in \mathcal{S}_{\mu,L}^1(\mathbb{R}^d)$, the following fundamental inequality holds for any step size $s \in (0,1/L]$:

$$
f(y - s\nabla f(y)) - f(x) \le \langle \nabla f(y), y - x \rangle - \underbrace{\frac{\mu}{2} \|y - x\|^2}_{\text{SC}} - \frac{s}{2} \|\nabla f(y)\|^2, \quad \forall x, y \in \mathbb{R}^d. \tag{2.3}
$$

We now turn our attention to composite functions of the form $\Phi = f + g$, where $f \in \mathcal{S}_{\mu,L}^1(\mathbb{R}^d)$ and $g \in \mathcal{F}^0(\mathbb{R}^d)$. Building upon the methodologies in [\[Beck and Teboulle](#page-20-3), [2009](#page-20-3), [Su et al.,](#page-21-4) [2016\]](#page-21-4), we introduce the concept of the s-proximal value, which serves as a cornerstone of our subsequent analysis.

Definition 2.1 (s-Proximal Value). Let the step size satisfy $s \in (0, 1/L]$. For any $f \in S^1_{\mu,L}(\mathbb{R}^d)$ and $g \in \mathcal{F}^0(\mathbb{R}^d)$, the s-proximal value is defined as:

$$
P_s(x) := \underset{y \in \mathbb{R}^d}{\arg \min} \left\{ \frac{1}{2s} \left\| y - (x - s \nabla f(x)) \right\|^2 + g(y) \right\}, \quad \forall x \in \mathbb{R}^d. \tag{2.4}
$$

The s-proximal value $P_s(x)$ minimizes a weighted sum of the squared Euclidean distance from the gradient-based update $x - s\nabla f(x)$ and the regularization term $g(y)$. This formulation facilitates efficient minimization of composite objectives, even when $g(x)$ is non-smooth but proximally simple. For instance, when $g(x) = \lambda ||x||_1$ (the ℓ_1 -norm), the s-proximal value admits a closed-form solution. Specifically, for any $x \in \mathbb{R}^d$, the *i*-th component of the *s*-proximal value $P_s(x)$ is given by:

$$
P_s(x)_i = (|(x - s\nabla f(x))_i| - \lambda s)_+ \text{sgn}((x - s\nabla f(x))_i),
$$

where $i = 1, \ldots, d$ and sgn(·) denotes the sign function.

Definition 2.2 (s-Proximal Subgradient). For any $x \in \mathbb{R}^d$, the s-proximal subgradient is defined as:

$$
G_s(x) := \frac{x - P_s(x)}{s},\tag{2.5}
$$

where the s-proximal value $P_s(x)$ is given by [\(2.4\)](#page-6-0). The subgradient $G_s(x)$ generalizes the classical gradient by incorporating the proximal update with respect to the regularizer g , capturing the direction of improvement for the composite function Φ.

Using the s-proximal subgradient from Definition [2.2,](#page-6-1) we extend the fundemental inequality [\(2.3\)](#page-6-2) to the composite setting. The proximal version is presented below, rigorously established in [\[Li et al.](#page-21-0), [2024b](#page-21-0)].

Lemma 2.3 (Lemma 4 in [\[Li et al.,](#page-21-0) [2024b](#page-21-0)]). Let $\Phi = f + g$ be a composite function with $f \in$ $\mathcal{S}_{\mu,L}^1(\mathbb{R}^d)$ and $g \in \mathcal{F}^0(\mathbb{R}^d)$. Then, the following inequality holds for any step size $s \in (0,1/L]$:

$$
\Phi(y - sG_s(y)) - \Phi(x) \le \langle G_s(y), y - x \rangle - \underbrace{\frac{\mu}{2} \|y - x\|^2}_{\text{SC}} - \frac{s}{2} \|G_s(y)\|^2, \quad \forall x, y \in \mathbb{R}^d. \tag{2.6}
$$

3 Lyapnov analysis for forward-backward accelerated algorithms

In this section, we first outline the process of constructing a novel Lyapunov function, following the principled way presented in [\[Chen et al.,](#page-20-11) [2022b](#page-20-11)]. Leveraging this approach, we then derive an $O(1/k^{2\alpha})$ convergence rate for the NAG- α method applied to strongly convex functions. A formal theorem is also provided to rigorously characterize this result.

3.1 Construction of a novel Lyapunov function

To analyze the NAG- α method, as given by $(1.2a)$ and $(1.2b)$, we first reformulate it in the phasespace representation. This reformulation leverages the implicit-velocity scheme, extending approaches previously applied to the classical NAG method [\[Chen et al.,](#page-20-11) [2022b](#page-20-11), [Li et al.](#page-21-0), [2024b](#page-21-0)]. To facilitate this process, we introduce a new iterative sequence $\sqrt{s}v_k = x_k - x_{k-1}$, which transforms the algorithm into the following phase-space representation:

$$
\int x_{k+1} - x_k = \sqrt{s}v_{k+1},
$$
\n(3.1a)

$$
\begin{cases} v_{k+1} - v_k = -\left(1 - \frac{(k-1)^{\alpha}}{k^{\alpha} + rk^{\alpha - 1}}\right) v_k - \sqrt{s} \nabla f(y_k), \end{cases}
$$
 (3.1b)

where the iterative sequence $\{y_k\}_{k=0}^{\infty}$ satisfies the following iterative relation:

$$
y_k = x_k + \frac{(k-1)^{\alpha}}{k^{\alpha} + rk^{\alpha - 1}} \cdot \sqrt{s}v_k.
$$
 (3.2)

(I) Construction of mixed energy In the context of Lyapunov analysis, the focus is often shifted from the iterative point x_k to the difference between the current iterate and the optimal solution x^* . The difference, $x_k - x^*$, provides deeper insights into the convergence behavior of the algorithm. Reformulating $(3.1a)$ in terms of this difference yields:

$$
(x_{k+1} - x^*) - (x_k - x^*) = \sqrt{s}v_{k+1}
$$
\n(3.3)

Substituting this expression (3.3) into $(3.1b)$, we derive the following iterative relation:

$$
\begin{aligned} \left[\sqrt{s}k^{\alpha}v_{k+1} + rk^{\alpha-1}(x_{k+1} - x^{\star})\right] - \left[\sqrt{s}(k-1)^{\alpha}v_{k} + r(k-1)^{\alpha-1}(x_{k} - x^{\star})\right] \\ = r\left[k^{\alpha-1} - (k-1)^{\alpha-1}\right](x_{k} - x^{\star}) - s\left(k^{\alpha} + rk^{\alpha-1}\right)\nabla f(y_{k}). \end{aligned} \tag{3.4}
$$

This iterative relation leads naturally to the mixed energy, which plays a central role in our Lyapunov analysis. The mixed energy is defined as:

$$
\mathcal{E}_{mix}(k) = \frac{1}{2} \left\| \sqrt{s}(k-1)^{\alpha} v_k + r(k-1)^{\alpha-1} (x_k - x^*) \right\|^2.
$$
 (3.5)

Next, we examine the change in mixed energy from one iteration to the next. Using the iterative relation [\(3.4\)](#page-7-4), the iterative difference in mixed energy is given by:

$$
\mathcal{E}_{mix}(k+1) - \mathcal{E}_{mix}(k)
$$
\n
$$
= \left\langle r \left[k^{\alpha-1} - (k-1)^{\alpha-1} \right] (x_k - x^\star) - s \left(k^\alpha + r k^{\alpha-1} \right) \nabla f(y_k), \right\rangle
$$
\n
$$
\sqrt{s}(k-1)^\alpha v_k + \frac{r}{2} \left[k^{\alpha-1} + (k-1)^{\alpha-1} \right] (x_k - x^\star) - \frac{s}{2} \left(k^\alpha + r k^{\alpha-1} \right) \nabla f(y_k) \right\rangle. \tag{3.6}
$$

Substituting the iterative relation (3.2) into the iterative difference (3.6) , we obtain:

$$
\mathcal{E}_{mix}(k+1) - \mathcal{E}_{mix}(k) \n= \left\langle -\frac{r \left[k^{\alpha-1} - (k-1)^{\alpha-1} \right]}{k^{\alpha} + rk^{\alpha-1}} \cdot \sqrt{s}(k-1)^{\alpha} v_k + r \left[k^{\alpha-1} - (k-1)^{\alpha-1} \right] (y_k - x^*) - s \left(k^{\alpha} + rk^{\alpha-1} \right) \nabla f(y_k), \n\left(1 - \frac{r}{2} \cdot \frac{k^{\alpha-1} + (k-1)^{\alpha-1}}{k^{\alpha} + rk^{\alpha-1}} \right) \sqrt{s}(k-1)^{\alpha} v_k + \frac{r}{2} \left[k^{\alpha-1} + (k-1)^{\alpha-1} \right] (y_k - x^*) - \frac{s \left(k^{\alpha} + rk^{\alpha-1} \right)}{2} \nabla f(y_k) \right\rangle.
$$
\n(3.7)

This difference involves several terms, which can be simplified further by introducing new notations:

$$
\int A_1(k, \alpha) = k^{\alpha - 1} - (k - 1)^{\alpha - 1},
$$
\n(3.8a)

$$
\begin{cases}\nA_1(\kappa, \alpha) = \kappa & -(\kappa - 1) \\
A_2(k, \alpha) = k^{\alpha - 1} + (k - 1)^{\alpha - 1},\n\end{cases}\n\tag{3.8b}
$$

$$
\left(B(k,\alpha,r) = k^{\alpha} + rk^{\alpha-1}.\right) \tag{3.8c}
$$

Using these new definitions, $(3.8a)$ — $(3.8c)$, the iterative difference (3.7) in mixed energy becomes:

$$
\mathcal{E}_{mix}(k+1) - \mathcal{E}_{mix}(k) = -\frac{s\sqrt{sk^{\alpha}(k-1)^{\alpha}}\langle \nabla f(y_k), v_k \rangle}{\Gamma}
$$

\n
$$
-rsk^{\alpha-1}B(k, \alpha, r) \langle \nabla f(y_k), y_k - x^{\star} \rangle
$$

\n
$$
+ \frac{s^2 B^2(k, \alpha, r)}{2} \|\nabla f(y_k)\|^2
$$

\n
$$
- \frac{rs(k-1)^{2\alpha} A_1(k, \alpha) [2k^{\alpha} + r A_1(k, \alpha)]}{2B^2(k, \alpha, r)} \|v_k\|^2
$$

\n
$$
+ \frac{r^2 A_1(k, \alpha) A_2(k, \alpha)}{2} \|y_k - x^{\star}\|^2
$$

\n
$$
+ r\sqrt{s}(k-1)^{\alpha} A_1(k, \alpha) \left(1 - \frac{r A_2(k, \alpha)}{B(k, \alpha, r)}\right) \langle y_k - x^{\star}, v_k \rangle.
$$
 (3.9)

This simplified representation enables us to bound and analyze the convergence properties of the algorithm effectively.

(II) Construction of potential energy We now proceed to define the potential energy, incorporating a dynamic coefficient $\tau(k)$, which is initially left undetermined:

$$
\mathcal{E}_{pot}(k) = s\tau(k)(f(x_k) - f(x^*)).\tag{3.10}
$$

Next, we compute the iterative difference in potential energy between consecutive iterations:

$$
\mathcal{E}_{pot}(k+1) - \mathcal{E}_{pot}(k) = s\tau(k) \left(f(x_{k+1}) - f(x_k) \right) + s \left(\tau(k+1) - \tau(k) \right) \left(f(x_{k+1}) - f(x^*) \right).
$$
 (3.11)

To bound this difference, we apply the fundamental inequality [\(2.3\)](#page-6-2) at two different points. First, for x_{k+1} and x_k , we obtain:

$$
f(x_{k+1}) - f(x_k) \le \langle \nabla f(y_k), y_k - x_k \rangle - \underbrace{\frac{\mu}{2} \| y_k - x_k \|^2}_{\text{SC}_1} - \frac{s}{2} \| \nabla f(y_k) \|^2. \tag{3.12}
$$

Similarly, applying the fundamental inequality (2.3) to x_{k+1} and x^* , we derive:

$$
f(x_{k+1}) - f(x^*) \le \langle \nabla f(y_k), y_k - x^* \rangle - \underbrace{\frac{\mu}{2} \| y_k - x^* \|^2}_{\mathbf{SC}_2} - \frac{s}{2} \| \nabla f(y_k) \|^2. \tag{3.13}
$$

Here, the terms denoted SC_1 in [\(3.12\)](#page-9-0) and SC_2 in [\(3.13\)](#page-9-1) are specific to the properties of strongly convex functions. These terms vanish in the case of convex functions, which aids in deriving convergence rates. Substituting the two inequalities, (3.12) and (3.13) , into the iterative difference (3.11) , we establish the following upper bound:

$$
\mathcal{E}_{pot}(k+1) - \mathcal{E}_{pot}(k) \le s\tau(k) \left(\langle \nabla f(y_k), y_k - x_k \rangle - \frac{\mu}{2} ||y_k - x_k||^2 \right)
$$

$$
+ s\left(\tau(k+1) - \tau(k)\right) \left(\langle \nabla f(y_k), y_k - x^* \rangle - \frac{\mu}{2} ||y_k - x^*||^2 \right)
$$

$$
- \frac{s^2 \tau(k+1)}{2} \cdot ||\nabla f(y_k)||^2
$$
(3.14)

To simplify further, we substitute the iterative relation [\(3.2\)](#page-7-5) into the right-hand side of [\(3.14\)](#page-9-3). This leads to the following expression for the iterative difference in potential energy:

$$
\mathcal{E}_{pot}(k+1) - \mathcal{E}_{pot}(k) \le \underbrace{s\sqrt{s}\tau(k) \cdot \frac{(k-1)^{\alpha}}{k^{\alpha} + rk^{\alpha-1}} \cdot \langle \nabla f(y_k), v_k \rangle}_{\text{T1}} - \underbrace{\frac{\mu s^2}{2} \cdot \frac{\tau(k)(k-1)^{2\alpha}}{B^2(k, \alpha, r)} \|v_k\|^2}_{\text{SC}_1} \times \left. f(\tau(k+1) - \tau(k)) \left(\langle \nabla f(y_k), y_k - x^{\star} \rangle - \frac{\mu}{2} \|y_k - x^{\star}\|^2 \right) \right)
$$

$$
-\frac{s^2\tau(k+1)}{2}\cdot\|\nabla f(y_k)\|^2.
$$
\n(3.15)

To ensure that the term $\mathbf{I} - \mathbf{II} = 0$, we choose the dynamic coefficient $\tau(k)$ as:

$$
\tau(k) = k^{\alpha} B(k, \alpha, r) = k^{\alpha} \left(k^{\alpha} + r k^{\alpha - 1} \right). \tag{3.16}
$$

Substituting the dynamic coefficient (3.16) into (3.15) , we simplify the upper bound, yielding the following expression:

$$
\mathcal{E}_{pot}(k+1) - \mathcal{E}_{pot}(k) \le \underbrace{s\sqrt{s}k^{\alpha}(k-1)^{\alpha} \cdot \langle \nabla f(y_{k}), v_{k} \rangle}_{\mathbf{H}} + s[(k+1)^{\alpha}B(k+1, \alpha, r) - k^{\alpha}B(k, \alpha, r)] \langle \nabla f(y_{k}), y_{k} - x^{\star} \rangle
$$

$$
- \underbrace{\frac{s^{2}(k+1)^{\alpha}B(k+1, \alpha, r)}{2} || \nabla f(y_{k}) ||^{2}}
$$

$$
- \underbrace{\frac{\mu s^{2}}{2} \cdot \frac{k^{\alpha}(k-1)^{2\alpha}}{B(k, \alpha, r)} ||v_{k}||^{2}}_{\mathbf{SC}_{1}}
$$

$$
- \underbrace{\frac{\mu s}{2} [(k+1)^{\alpha}B(k+1, \alpha, r) - k^{\alpha}B(k, \alpha, r)] ||y_{k} - x^{\star}||^{2}}_{\mathbf{SC}_{2}}.
$$
(3.17)

Finally, with the mixed energy defined in [\(3.5\)](#page-7-6) and the potential energy defined in [\(3.10\)](#page-9-4), we can now construct the Lyapunov function as:

$$
\mathcal{E}(k) = sk^{\alpha} \left(k^{\alpha} + rk^{\alpha - 1} \right) \left(f(x_k) - f(x^{\star}) \right) + \frac{1}{2} \left\| \sqrt{s} (k-1)^{\alpha} v_k + r(k-1)^{\alpha - 1} (x_k - x^{\star}) \right\|^2. \tag{3.18}
$$

This Lyapunov function encapsulates both the mixed and potential energies, serving as a key tool in deriving convergence rates for the $NAG-\alpha$ method, as described in $(1.2a)$ and $(1.2b)$.

3.2 Controllable $O(1/k^{2\alpha})$ convergence of NAG- α

By leveraging the Lyapunov function defined in [\(3.18\)](#page-10-2), we rigorously establish the convergence rate for the NAG- α method. This result is encapsulated in the following theorem:

Theorem 3.1. Let $f \in \mathcal{S}_{\mu,L}^1(\mathbb{R}^d)$, and suppose the step size satisfies $s \in (0,1/L]$. If the controllable parameter is chosen such that $r > 2\alpha$, then there exists a positive integer $K := K(\alpha, r)$ such that the iterative sequence ${x_k}_{k=0}^{\infty}$, generated by NAG- α under updates [\(1.2a\)](#page-3-0) and [\(1.2b\)](#page-3-1) with any initial $x_0 = y_0 \in \mathbb{R}^d$, converges at the following rate:

$$
f(x_k) - f(x^*) \le \frac{\mathcal{E}(K)}{sk^{\alpha} (k^{\alpha} + rk^{\alpha - 1})}, \quad \forall k \ge K.
$$
 (3.19)

Proof of Theorem [3.1.](#page-10-3) To establish the stated result, we analyze the iterative change in the Lyapunov function $\mathcal{E}(k)$ across iterations. By combining the inequalities derived for the iterative differences, (3.9) and (3.17) , we observe that the change in the Lyapunov function (3.18) satisfies:

$$
\mathcal{E}(k+1) - \mathcal{E}(k) \le \underbrace{s \left[(k+1)^{\alpha} B(k+1, \alpha, r) - B^2(k, \alpha, r) \right] \left(\langle \nabla f(y_k), y_k - x^{\star} \rangle - \frac{s}{2} \left\| \nabla f(y_k) \right\|^2 \right)}_{\text{III}}
$$

$$
-\frac{rs(k-1)^{2\alpha}A_{1}(k,\alpha)[2k^{\alpha}+rA_{1}(k,\alpha)]}{2B^{2}(k,\alpha,r)}||v_{k}||^{2} + \frac{r^{2}A_{1}(k,\alpha)A_{2}(k,\alpha)}{2}||y_{k}-x^{\star}||^{2} + r\sqrt{s}(k-1)^{\alpha}A_{1}(k,\alpha)\left(1-\frac{rA_{2}(k,\alpha)}{B(k,\alpha,r)}\right)\langle y_{k}-x^{\star},v_{k}\rangle + \frac{rs^{2}}{2}\cdot\frac{k^{\alpha}(k-1)^{2\alpha}}{B(k,\alpha,r)}||v_{k}||^{2} - \frac{\frac{\mu s^{2}}{2}\cdot\left[(k+1)^{\alpha}B(k+1,\alpha,r)-k^{\alpha}B(k,\alpha,r)\right]||y_{k}-x^{\star}||^{2}}{sC_{1}}.
$$
\n(3.20)

To guarantee $\mathcal{E}(k+1) - \mathcal{E}(k) \leq 0$, we systematically analyze the contributions of each term by grouping them into four major components:

(1) Analyzing Term III. The coefficient of Term III, after an asymptotic expansion, is given by:

$$
(k+1)^{\alpha} B(k+1,\alpha,r) - B^{2}(k,\alpha,r)
$$

= $(k+1)^{2\alpha} + r(k+1)^{2\alpha-1} - (k^{\alpha} + rk^{\alpha-1})^{2}$
= $k^{2\alpha} + (r+2\alpha) k^{2\alpha-1} + O(k^{2(\alpha-1)}) - k^{2\alpha} - 2rk^{2\alpha-1} - r^{2}k^{2(\alpha-1)}$
= $(2\alpha - r) k^{2\alpha-1} + O(k^{2(\alpha-1)}).$ (3.21)

It follows that when we control the parameter satisfing $r > 2\alpha$, the coefficient becomes asymptotically non-positive, ensuring a negative contribution to the iterative difference. On the other hand, it follows from the fact that

$$
\langle \nabla f(x) - \nabla f(y), x - y \rangle \ge \frac{1}{L} \|\nabla f(x) - \nabla f(y)\|^2, \quad \forall x, y \in \mathbb{R}^d
$$

for $f \in \mathcal{F}_L^1(\mathbb{R}^d)$ that

$$
\langle \nabla f(y_k), y_k - x^* \rangle - \frac{s}{2} \left\| \nabla f(y_k) \right\|^2 \ge \left(\frac{1}{L} - \frac{s}{2} \right) \left\| \nabla f(y_k) \right\|^2 \ge 0.
$$

Thus, the whole term III is negative in the iterative difference.

(2) Analyzing Term IV₁. The asymptotic expansion of the coefficient of Term IV₁ is:

$$
-\frac{rs(k-1)^{2\alpha}A_1(k,\alpha)[2k^{\alpha}+rA_1(k,\alpha)]}{2B^2(k,\alpha,r)}=-rs(\alpha-1)k^{2\alpha-2}+O(k^{2\alpha-3}).
$$

For $\alpha < 1$, this term is asymptotically non-negative. Incorporating the term $SC₁$ from the strongly convex inequality, we obtain:

$$
-\frac{rs(k-1)^{2\alpha}A_1(k,\alpha)\left[2k^{\alpha}+rA_1(k,\alpha)\right]}{2B^2(k,\alpha,r)}-\frac{\mu s^2}{2}\cdot\frac{k^{\alpha}(k-1)^{2\alpha}}{B(k,\alpha,r)}=-\frac{\mu s^2}{2}\cdot k^{2\alpha}+O(k^{2\alpha-1}).
$$
 (3.22)

Thus, the contribution of $||v_k||^2$ remains asymptotically negative.

(3) Analyzing Term IV₂. The asymptotic expansion of the coefficient of Term IV₂ is:

$$
\frac{r^2A_1(k,\alpha)A_2(k,\alpha)}{2} = r^2(\alpha-1)k^{2\alpha-3} + O(k^{2\alpha-4}).
$$

For $\alpha > 1$, this term is asymptotically non-negative. Incorporating the term SC_2 from the strongly convex inequality, we obtain:

$$
\frac{r^2 A_1(k, \alpha) A_2(k, \alpha)}{2} - \frac{\mu s}{2} \cdot [(k+1)^{\alpha} B(k+1, \alpha, r) - k^{\alpha} B(k, \alpha, r)]
$$

= $-\mu s \alpha k^{2\alpha - 1} + O(k^{2\alpha - 2})$. (3.23)

Thus, the contribution of $||y_k - x^*||^2$ also remains asymptotically negative.

(4) Analyzing Term IV₃. The coefficient of Term IV₃ is asymptotically expanded as:

$$
r\sqrt{s}(k-1)^{\alpha}A_1(k,\alpha)\left(1-\frac{rA_2(k,\alpha)}{B(k,\alpha,r)}\right) = r\sqrt{s}(\alpha-1)k^{2\alpha-2} + O(k^{2\alpha-3}).
$$
 (3.24)

 \Box

By combining the three asymptotic expansions, [\(3.22\)](#page-12-0), [\(3.23\)](#page-12-1), and [\(3.24\)](#page-12-2), the two strongly convex terms, SC_1 and SC_2 , reduce the combined sum $-IV_1 + IV_2 + IV_3$ to a sum of three non-positive perfect squares.

In summary, we conclude that when $r > 2\alpha$, there exists a positive integer $K = K(\alpha, r)$ such that for any $k \geq K$, the iterative difference $\mathcal{E}(k+1) - \mathcal{E}(k)$ remains non-positive. This ensures that the Lyapunov function (3.18) is non-increasing, thereby completing the proof.

From the above proof, it becomes clear that the fundamental inequality [\(2.3\)](#page-6-2) for strongly convex functions is pivotal in the analysis and cannot be simplified to its counterpart for convex functions. This distinction is further elaborated in the following remark:

Remark 3.2. In the inequality governing the iterative difference (3.20) , the coefficients of both terms, IV_1 and IV_2 , involve the expression

$$
A_1(k,\alpha) = k^{\alpha-1} - (k-1)^{\alpha-1} = (\alpha-1)k^{\alpha-2} + O(k^{\alpha-3}),
$$

which retains a consistent sign based on the value of α . Specifically, when $\alpha \geq 1$, the sign is positive, while for $\alpha < 1$, the sign is negative. As a result, in the inequality for the iterative difference (3.20) , the two terms, $-\mathbf{IV}_1$ and \mathbf{IV}_2 , always exhibits opposite signs. This observation reveals an important limitation: if the objective function under consideration is merely convex, rather than strongly convex, the terms SC_1 and SC_2 , which arise from the fundamental inequaliy for strongly convex functions, vanish. Without these additional terms, it is no longer possible to ensure that the iterative difference [\(3.20\)](#page-11-0) is non-positive. Consequently, the Lyapunov function [\(3.18\)](#page-10-2) cannot be guaranteed to decrease consistently. This underscores the indispensable role of strong convexity in achieving the desired convergence properties. The additional structure provided by strong convexity ensures that the iterative process maintains the non-positivity of the iterative difference, which is essential for guaranteeing a monotonic decrease in the Lyapunov function.

With Theorem [3.1,](#page-10-3) we can also derive the convergence rate of NAG- α at the critical step size $s = 1/L$, filling a gap left in prior analyses such as those in [\[Li et al.](#page-20-5), [2024a](#page-20-5), [Fu and Shi,](#page-20-1) [2024](#page-20-1)].

Corollary 3.3. Under the same assumptions of Theorem [3.1,](#page-10-3) if the step size satisfies $s = 1/L$, there exists a positive integer $K := K(\alpha, r)$ such that NAG- α converges at the following rate:

$$
f(x_k) - f(x^*) \le \frac{L\mathcal{E}(K)}{k^{\alpha} (k^{\alpha} + rk^{\alpha - 1})}, \quad \forall k \ge K.
$$
 (3.25)

This result confirms the theoretical guarantees of $NAG-\alpha$ for strongly convex functions and provides explicit conditions for achieving optimal convergence rates under the critical step size.

3.3 Generalization to FISTA- α

We now extend the controllable $O(1/k^{2\alpha})$ convergence of NAG- α , as established in Theorem [3.1](#page-10-3) and Corollary [3.3,](#page-13-0) to its proximal counterpart, FISTA- α . In accordance with Definition [2.1,](#page-6-3) FISTA- α employs the s-proximal value [\(2.4\)](#page-6-0) and adheres to the following iterative scheme, starting from any initial point $y_0 = x_0 \in \mathbb{R}^d$:

$$
(3.26a)
$$

$$
\begin{cases}\n x & (k-1)^{\alpha} \\
 y_k = x_k + \frac{(k-1)^{\alpha}}{k^{\alpha} + rk^{\alpha-1}} (x_k - x_{k-1})\n \end{cases}
$$
\n(3.26b)

where $s > 0$ is the step size. As specified in Definition [2.2,](#page-6-1) FISTA- α can alternatively be expressed using the s-proximal subgradient (2.5) , yielding the alternative iterative scheme:

$$
\int x_k = y_{k-1} - sG_s(y_{k-1})
$$
\n(3.27a)

$$
\begin{cases} y_k = x_k + \frac{(k-1)^{\alpha}}{k^{\alpha} + rk^{\alpha-1}} (x_k - x_{k-1}) \end{cases}
$$
 (3.27b)

where the s-proximal subgradient $G_s(y_k)$ generalizes the classical gradient $\nabla f(y_k)$ used in NAG- α . This formulation underscores the structural similarity between NAG- α and FISTA- α , while explicitly incorporating the non-smooth nature of the composite objective function Φ . By introducing the velocity iterative sequence $v_k = (x_k - x_{k-1})/\sqrt{s}$, the FISTA- α updates can be formulated in a phase-space representation akin to $NAG-\alpha$:

$$
\int x_{k+1} - x_k = \sqrt{5}v_{k+1},\tag{3.28a}
$$

$$
\left\{ v_{k+1} - v_k = -\left(1 - \frac{(k-1)^{\alpha}}{k^{\alpha} + rk^{\alpha-1}} \right) v_k - \sqrt{s} G_s(y_k), \right\}
$$
 (3.28b)

with the auxiliary sequence $\{y_k\}_{k=0}^{\infty}$ satisfying:

$$
y_k = x_k + \frac{(k-1)^{\alpha}}{k^{\alpha} + rk^{\alpha-1}} \cdot \sqrt{s}v_k.
$$
\n(3.29)

To establish the controllable $O(1/k^{2\alpha})$ convergence rate for FISTA- α , we construct a generalized Lyapunov function inspired by the smooth one (3.18) . This Lyapunov function is adapted to composite functions by substituting f with $\Phi = f + g$, yielding:

$$
\mathcal{E}(k) = sk^{\alpha} \left(k^{\alpha} + rk^{\alpha - 1} \right) \left(\Phi(x_k) - \Phi(x^{\star}) \right) + \frac{1}{2} \left\| \sqrt{s} (k-1)^{\alpha} v_k + r(k-1)^{\alpha - 1} (x_k - x^{\star}) \right\|^2. \tag{3.30}
$$

In Section [3.1,](#page-7-7) the $O(1/k^{2\alpha})$ rate for smooth functions hinges on two key inequalities, [\(3.12\)](#page-9-0) and [\(3.13\)](#page-9-1). For the composite setting, these are generalized to accommodate proximal components by leveraging Lemma [2.3,](#page-6-5) which extends the fundamental inequality for strongly convex functions to the proximal setting. These adaptations enable a rigorous analysis of $FISTA-\alpha$, effectively addressing the challenges posed by the non-smooth components in Φ . The convergence properties are formally encapsulated in the following theorem:

Theorem 3.4. Let $f \in \mathcal{S}_{\mu,L}^1(\mathbb{R}^d)$ and $g \in \mathcal{F}^0(\mathbb{R}^d)$, and suppose the step size satisfies $s \in (0,1/L]$. If the controllable parameter is chosen such that $r > 2\alpha$, then there exists a positive integer $K := K(\alpha, r)$ such that the iterative sequence $\{x_k\}_{k=0}^{\infty}$, generated by FISTA- α under updates [\(3.26a\)](#page-13-1) and [\(3.26b\)](#page-13-2) with any initial $x_0 = y_0 \in \mathbb{R}^d$, converges at the following rate:

$$
\Phi(x_k) - \Phi(x^\star) \le \frac{\mathcal{E}(K)}{sk^\alpha \left(k^\alpha + rk^{\alpha - 1}\right)}, \quad \forall k \ge K. \tag{3.31}
$$

Furthermore, if the step size is set critically as $s = 1/L$, the iterative sequence $\{x_k\}_{k=0}^{\infty}$ satisfies:

$$
\Phi(x_k) - \Phi(x^\star) \le \frac{L\mathcal{E}(K)}{k^{\alpha} (k^{\alpha} + rk^{\alpha - 1})}, \quad \forall k \ge K.
$$
\n(3.32)

4 Monotonically controllable $O\left(1/k^{2\alpha}\right)$ convergence

In this section, we extend the controllable $O(1/k^{2\alpha})$ convergence rate established for NAG- α in Sec-tion [3](#page-7-0) to its proximal variant, M-NAG- α . Following the methodology outlined by [Fu and Shi](#page-20-1) [\[2024\]](#page-20-1), the Lyapunov function [\(3.18\)](#page-10-2) is designed without incorporating the kinetic energy term. As a result, the derivation of the convergence rate relies solely on the iterative dynamics of $M-MAG-\alpha$, as specified in $(1.3a)$ — $(1.3c)$, bypassing the need for the complete set of NAG updates in $(1.2a)$ and $(1.2b)$. Building on this framework, we further generalize the controllable $O(1/k^{2\alpha})$ convergence to its proximal counterpart, $M-FISTA-_{\alpha}$.

4.1 Smooth optimization via M-NAG- α

According to the iterative relation [\(3.2\)](#page-7-5), the velocity term v_k can be expressed in terms of x_k and y_k . This substitution yields the following relation:

$$
\sqrt{s}(k-1)^{\alpha}v_k + r(k-1)^{\alpha-1}(x_k - x^*)
$$

=
$$
\left[k^{\alpha} + r(k^{\alpha-1} - (k-1)^{\alpha-1})\right](y_k - x_k) + r(k-1)^{\alpha-1}(y_k - x^*)
$$
. (4.1)

Leveraging this expression, we reformulate the Lyapunov function originally introduced in [\(3.18\)](#page-10-2) so that it depends exclusively on the variables, x_k and y_k . The reformulated Lyapunov function is given as:

$$
\mathcal{E}(k) = \underbrace{sk^{\alpha} (k^{\alpha} + rk^{\alpha-1}) (f(x_k) - f(x^*))}_{:= \mathcal{E}_{pot}(k)} + \underbrace{\frac{1}{2} || [k^{\alpha} + r (k^{\alpha-1} - (k-1)^{\alpha-1})] (y_k - x_k) + r(k-1)^{\alpha-1} (y_k - x^*) ||^2}_{:= \mathcal{E}_{mix}(k)} \tag{4.2}
$$

This reformulation, which decouples $\mathcal{E}(k)$ from v_k and depends only on x_k and y_k , is instrumental in deriving the controllable $O(1/k^{2\alpha})$ convergence rate for M-NAG- α . The formal result is encapsulated in the subsequent theorem.

Theorem 4.1. Let $f \in \mathcal{S}_{\mu,L}^1(\mathbb{R}^d)$, and suppose the step size satisfies $s \in (0,1/L]$. If the controllable parameter is chosen such that $r > 2\alpha$, then there exists a positive integer $K := K(\alpha, r)$ such that the iterative sequence $\{x_k\}_{k=0}^{\infty}$, generated by M-NAG- α under updates $(1.3a) - (1.3c)$ $(1.3a) - (1.3c)$ with any initial $x_0 = y_0 \in \mathbb{R}^d$, converges monotonically at the following rate:

$$
f(x_k) - f(x^*) \le \frac{\mathcal{E}(K)}{sk^{\alpha} (k^{\alpha} + rk^{\alpha - 1})}, \quad \forall k \ge K.
$$
 (4.3)

Furthermore, if the step size is set critically as $s = 1/L$, the iterative sequence $\{x_k\}_{k=0}^{\infty}$ satisfies:

$$
f(x_k) - f(x^*) \le \frac{L\mathcal{E}(K)}{k^{\alpha} (k^{\alpha} + rk^{\alpha - 1})}, \quad \forall k \ge K.
$$
 (4.4)

Proof of Theorem [4.1.](#page-15-0) To calculate the iterative difference of the Lyapunov function, we break the process into two distinct components: the potential energy and the mixed energy, as shown in [\(4.2\)](#page-15-1). To further streamline the calculation, we utilize the new definitions introduced in $(3.8a)$ — $(3.8c)$. These definitions provide convenient expressions that facilitate the management of interactions between the various terms in the Lyapunov function.

(1) For the potential energy $\mathcal{E}_{pot}(k)$, we calculate its iterative difference as

$$
\mathcal{E}_{pot}(k+1) - \mathcal{E}_{pot}(k) = sk^{\alpha} B(k, \alpha, r) (f(x_{k+1}) - f(x_k)) + s [(k+1)^{\alpha} B(k+1, \alpha, r) - k^{\alpha} B(k, \alpha, r)] (f(x_{k+1}) - f(x^*))
$$
. (4.5)

By combining $(1.3a)$ and $(1.3b)$, we can deduce that the iterative sequence of M-NAG- α satisfies:

$$
f(x_{k+1}) \le f(z_k) = f(y_k - s \nabla f(y_k)).
$$
\n(4.6)

Thus, we can apply the fundamental inequality (2.3) . First, substituting the points x_{k+1} and x_k , we obtain the following inequality:

$$
f(x_{k+1}) - f(x_k) \le f(y_k - s\nabla f(y_k)) - f(x_k)
$$

\n
$$
\le \langle \nabla f(y_k), y_k - x_k \rangle - \underbrace{\frac{\mu}{2} \| y_k - x_k \|^2}_{\text{SC}_1} - \frac{s}{2} \| \nabla f(y_k) \|^2.
$$
 (4.7)

Then, for the points x_{k+1} and x^* , we derive:

$$
f(x_{k+1}) - f(x^*) \le f(y_k - s\nabla f(y_k)) - f(x^*)
$$

\n
$$
\le \langle \nabla f(y_k), y_k - x^* \rangle - \underbrace{\frac{\mu}{2} \| y_k - x^* \|^2}_{\text{SC}_2} - \frac{s}{2} \| \nabla f(y_k) \|^2.
$$
 (4.8)

Substituting the two inequalities, (4.7) and (4.8) , into (4.5) , we establish the following upper bound for the iterative difference in the potential energy as:

$$
\mathcal{E}_{pot}(k+1) - \mathcal{E}_{pot}(k) = \underbrace{s k^{\alpha} B(k, \alpha, r) \langle \nabla f(y_k), y_k - x_k \rangle}_{\mathbf{I}} + s \left[(k+1)^{\alpha} B(k+1, \alpha, r) - k^{\alpha} B(k, \alpha, r) \right] \langle \nabla f(y_k), y_k - x^{\star} \rangle
$$

$$
- \frac{s^2 (k+1)^{\alpha} B(k+1, \alpha, r)}{2} \|\nabla f(y_k)\|^2
$$

$$
- \underbrace{\frac{\mu s}{2} \cdot k^{\alpha} B(k, \alpha, r) \|y_k - x_k\|^2}_{\mathbf{SC}_1}
$$

$$
- \underbrace{\frac{\mu s}{2} \cdot \left[(k+1)^{\alpha} B(k+1, \alpha, r) - k^{\alpha} B(k, \alpha, r) \right] \|y_k - x^{\star}\|^2}_{\mathbf{SC}_2}.
$$
(4.9)

(2) Next, we turn our attention to the mixed energy. By substituting [\(1.3a\)](#page-3-2) into [\(1.3c\)](#page-3-3) and re-expressing the differences between the iterative sequences and the optimal points, rather than using the sequences themselves, we arrive at the following iterative relation:

$$
\begin{aligned}\n\left[\left((k+1)^{\alpha} + r A_1(k+1,\alpha) \right) (y_{k+1} - x_{k+1}) + r k^{\alpha - 1} (y_{k+1} - x^{\star}) \right] \\
&- \left[(k^{\alpha} + r A_1(k,\alpha)) (y_k - x_k) + r(k-1)^{\alpha - 1} (y_k - x^{\star}) \right] \\
&= -r A_1(k,\alpha) (y_k - x_k) + r A_1(k,\alpha) (y_k - x^{\star}) - s B(k,\alpha,r) \nabla f(y_k).\n\end{aligned} \tag{4.10}
$$

Using the iterative relation [\(4.10\)](#page-16-1), we can compute the iterative difference in the mixed energy as follows:

$$
\mathcal{E}_{mix}(k+1) - \mathcal{E}_{mix}(k)
$$
\n
$$
= \left\langle -rA_1(k,\alpha) \left(y_k - x_k \right) + rA_1(k,\alpha) \left(y_k - x^{\star} \right) - sB(k,\alpha,r) \nabla f(y_k), \right\rangle
$$
\n
$$
\left(k^{\alpha} + \frac{rA_1(k,\alpha)}{2} \right) \left(y_k - x_k \right) + \frac{rA_2(k,\alpha)}{2} \left(y_k - x^{\star} \right) - \frac{sB(k,\alpha,r)}{2} \nabla f(y_k) \right\rangle \tag{4.11}
$$

Reformulating and simplifying the iterative difference [\(4.11\)](#page-16-2), we have:

$$
\mathcal{E}_{mix}(k+1) - \mathcal{E}_{mix}(k) = -\underbrace{\underbrace{s k^{\alpha} B(k, \alpha, r) \langle \nabla f(y_k), y_k - x_k \rangle}_{\text{IT}}}{\underbrace{\text{r} s k^{\alpha-1} B(k, \alpha, r) \langle \nabla f(y_k), y_k - x^* \rangle}_{2} + \frac{s^2 B^2(k, \alpha, r)}{2} \|\nabla f(y_k)\|^2}
$$

$$
-\frac{rA_1(k,\alpha) [2k^{\alpha} + rA_1(k,\alpha)]}{2} ||y_k - x_k||^2 + \frac{r^2 A_1(k,\alpha) A_2(k,\alpha)}{2} ||y_k - x^*||^2 + rA_1(k,\alpha) (B(k,\alpha,r) - rA_2(k,\alpha)) \langle y_k - x_k, y_k - x^* \rangle.
$$
 (4.12)

From the iterative differences of the potential and the mixed energies, as described in [\(4.9\)](#page-16-3) and (4.12) , we observe that the contributions from terms **I** and **II** are equal. This observation allows us to derive the following bound on the iterative difference:

$$
\mathcal{E}(k+1) - \mathcal{E}(k) \leq \underbrace{s\left[(k+1)^{\alpha} B(k+1, \alpha, r) - B^{2}(k, \alpha, r) \right] \left(\langle \nabla f(y_{k}), y_{k} - x^{\star} \rangle - \frac{s}{2} \|\nabla f(y_{k})\|^{2} \right)}_{\text{III}} \n- \underbrace{\frac{rA_{1}(k, \alpha)\left[2k^{\alpha} + rA_{1}(k, \alpha) \right]}{2} \|y_{k} - x_{k}\|^{2}}_{\text{IV}_{1}} \n+ \underbrace{\frac{r^{2}A_{1}(k, \alpha)A_{2}(k, \alpha)}{2} \|y_{k} - x^{\star}\|^{2}}_{\text{IV}_{2}} \n+ rA_{1}(k, \alpha)\left(B(k, \alpha, r) - rA_{2}(k, \alpha) \right) \langle y_{k} - x_{k}, y_{k} - x^{\star} \rangle_{\text{IV}_{3}} \n- \underbrace{\frac{\mu s}{2} \cdot k^{\alpha} B(k, \alpha, r) \|y_{k} - x_{k}\|^{2}}_{\text{SC}_{1}} \n- \underbrace{\frac{\mu s}{2} \cdot [(k+1)^{\alpha} B(k+1, \alpha, r) - k^{\alpha} B(k, \alpha, r)] \|y_{k} - x^{\star}\|^{2}}_{\text{SC}_{2}}.
$$
\n(4.13)

Analogously, we can estimate the asymptotic expansion for the iterative difference as follows: (1) Coefficient of Term III:

$$
(k+1)^{\alpha}B(k+1,\alpha,r) - B^{2}(k,\alpha,r) = (2\alpha - r)k^{2\alpha - 1} + O(k^{2(\alpha - 1)}).
$$
 (4.14)

(2) Coefficient of
$$
||y_k - x_k||^2
$$
 (Term IV₁ and Term SC₁):
\n
$$
-\frac{rA_1(k,\alpha)[2k^{\alpha} + rA_1(k,\alpha)]}{2} - \frac{\mu s}{2} \cdot k^{\alpha}B(k,\alpha,r) = -\frac{\mu s}{2} \cdot k^{2\alpha} + O(k^{2\alpha - 1}).
$$
\n(4.15)

(3) Coefficient of $||y_k - x^*||^2$ (Term **IV**₂ and Term **SC**₂):

$$
\frac{rA_1(k,\alpha)A_2(k,\alpha)}{2} - \frac{\mu s}{2} [(k+1)^{\alpha} B(k+1,\alpha,r) - k^{\alpha} B(k,\alpha,r)]
$$

= $-\mu s \alpha k^{2\alpha - 1} + O(k^{2\alpha - 2}).$ (4.16)

(4) Coefficient of $\langle y_k - x_k, y_k - x^* \rangle$ (Term IV₃): $rA_1(k, \alpha) (B(k, \alpha, r) - rA_2(k, \alpha)) = (\alpha - 1)rk^{2(\alpha - 1)} + O(k^{2\alpha - 3})$ (4.17)

By plugging (4.14) — (4.17) into the iterative inequality (4.13) , we can deduce that $\mathcal{E}(k+1)-\mathcal{E}(k) \le$ 0 asymptotically. This confirms that the Lyapunov function decreases over iterations, leading to the conclusion that the proof is complete. \Box

4.2 Composite optimization via M-FISTA- α

In this section, we extend the controllable $O(1/k^{2\alpha})$ convergence rate of M-NAG- α , as established in Theorem [4.1,](#page-15-0) to include its proximal variant, M-FISTA- α . As outlined in Definition [2.1,](#page-6-3) M-FISTA- α utilizes the s-proximal value defined in (2.4) and adheres to the following iterative scheme outlined below, starting from any initial point $y_0 = x_0 \in \mathbb{R}^d$:

$$
\begin{cases}\nz_{k-1} = P_s(y_{k-1}), \nx_k = \n\begin{cases}\nz_{k-1}, & \text{if } f(z_{k-1}) \le f(x_{k-1}), \\
\vdots \\
\vdots \\
\vdots \\
\vdots\n\end{cases} & \text{if } f(z_{k-1}) > f(z_{k-1}).\n\end{cases}\n\tag{4.18b}
$$

$$
x_k = \begin{cases} z_{k-1}, & \text{if } f(z_{k-1}) \le f(x_{k-1}), \\ x_{k-1}, & \text{if } f(z_{k-1}) > f(x_{k-1}), \end{cases} \tag{4.18b}
$$

$$
\left(y_k = x_k + \frac{(k-1)^{\alpha}}{k^{\alpha} + rk^{\alpha-1}}(x_k - x_{k-1}) + \frac{(k-1)^{\alpha} + r(k-1)^{\alpha-1}}{k^{\alpha} + rk^{\alpha-1}}(z_{k-1} - x_k),\right)
$$
(4.18c)

where $s > 0$ is the step size. To offer a more unified and efficient perspective, we reformulate M-FISTA- α , which mirrors the structure of M-NAG- α , with one key difference: the s-proximal value is replaced with the s-proximal subgradient as defined in (2.2) . This adjustment results in the following iterative scheme:

$$
(4.19a)
$$
\n
$$
z_{k-1} = y_{k-1} - sG_s(y_{k-1}),
$$
\n
$$
(4.19a)
$$

$$
x_k = \begin{cases} z_{k-1}, & \text{if } f(z_{k-1}) \le f(x_{k-1}), \\ x_{k-1}, & \text{if } f(z_{k-1}) > f(x_{k-1}), \end{cases} \tag{4.19b}
$$

$$
\left(y_k = x_k + \frac{(k-1)^{\alpha}}{k^{\alpha} + rk^{\alpha-1}}(x_k - x_{k-1}) + \frac{(k-1)^{\alpha} + r(k-1)^{\alpha-1}}{k^{\alpha} + rk^{\alpha-1}}(z_{k-1} - x_k),\right)
$$
(4.19c)

where the s-proximal subgradient $G_s(y_k)$ replaces the gradient $\nabla f(y_k)$ in NAG- α . This formulation explicitly accounts for the non-smooth nature of the composite objective function Φ , while maintaining a similar structure to $M-NAG-\alpha$.

To establish the controllable $O(1/k^{2\alpha})$ convergence rate for M-FISTA- α , we generalize the Lya-punov function [\(4.2\)](#page-15-1) used in M-NAG- α to accommodate the composite objective function Φ . The new Lyapunov function takes the following form:

$$
\mathcal{E}(k) = sk^{\alpha} (k^{\alpha} + rk^{\alpha-1}) (\Phi(x_k) - \Phi(x^*))
$$

+
$$
\frac{1}{2} \left\| \left[k^{\alpha} + r (k^{\alpha-1} - (k-1)^{\alpha-1}) \right] (y_k - x_k) + r(k-1)^{\alpha-1} (y_k - x^*) \right\|^2.
$$
 (4.20)

In Section [4.1,](#page-14-1) the derivation of the $O(1/k^{2\alpha})$ rate for smooth functions hinges on two key inequalities, [\(4.7\)](#page-15-2) and [\(4.8\)](#page-16-0). These inequalities are generalized in the composite setting to accommodate the proximal components, leveraging Lemma [2.3,](#page-6-5) which extends the fundamental inequality for strongly convex functions to the proximal case. These adaptations allow for a rigorous analysis of M-FISTA- α , effectively addressing the non-smooth components of Φ . The convergence properties of M-FISTA- α are formally encapsulated in the following theorem:

Theorem 4.2. Let $f \in \mathcal{S}_{\mu,L}^1(\mathbb{R}^d)$ and $g \in \mathcal{F}^0(\mathbb{R}^d)$, and suppose the step size satisfies $s \in (0,1/L]$. If the controllable parameter is chosen such that $r > 2\alpha$, then there exists a positive integer $K :=$ $K(\alpha, r)$ such that the iterative sequence $\{x_k\}_{k=0}^{\infty}$, generated by FISTA- α under updates, [\(4.18a\)](#page-18-0) — [\(4.18c\)](#page-18-1) with any initial $x_0 = y_0 \in \mathbb{R}^d$, converges at the following rate:

$$
\Phi(x_k) - \Phi(x^\star) \le \frac{\mathcal{E}(K)}{sk^\alpha \left(k^\alpha + rk^{\alpha - 1}\right)}, \quad \forall k \ge K. \tag{4.21}
$$

Furthermore, if the step size is set critically as $s = 1/L$, the iterative sequence $\{x_k\}_{k=0}^{\infty}$ satisfies:

$$
\Phi(x_k) - \Phi(x^\star) \le \frac{L\mathcal{E}(K)}{k^{\alpha} (k^{\alpha} + rk^{\alpha - 1})}, \quad \forall k \ge K. \tag{4.22}
$$

5 Conclusion and future work

In this paper, we introduce a family of controllable momentum coefficients for forward-backward accelerated methods, with a focus on the critical step size $s = 1/L$. Unlike traditional linear forms, the proposed momentum coefficients adopt an α -th power structure, with the parameter r adaptively tuned to α . A key contribution is the development of a Lyapunov function tailored to the parameter α , which excludes the kinetic energy term. This simplification enabled us to establish a controllable $O(1/k^{2\alpha})$ convergence rate for the NAG- α method, provided $r > 2\alpha$. At the critical step size, NAG- α achieves an inverse polynomial convergence rate of arbitrary degree by adjusting r according to $\alpha > 0$. By omitting the kinetic energy term, we further simplified its expression in terms of the iterative sequences x_k and y_k , thereby eliminating the need for phase-space representations. This insight allowed us to extend the controllable $O(1/k^{2\alpha})$ rate to the monotonic variant, M-NAG- α , further enhancing optimization efficiency. Finally, by leveraging the fundamental inequality for composite functions derived in [\[Li et al.,](#page-21-0) [2024b\]](#page-21-0), we extended the controllable $O(1/k^{2\alpha})$ rate to proximal algorithms, including FISTA- α and M-FISTA- α .

A notable reference point for first-order methods is the lower bound established in [\[Nemirovsky and Yudin,](#page-21-7) [1983,](#page-21-7) [Nesterov](#page-21-6), [2018\]](#page-21-6), which applies to a finite number of iterations $(k \le d/2)$. In the asymptotic sense, however, the convergence rate achieves $o(1/k^2)$, as shown in [\[Attouch and Peypouquet](#page-20-9), [2016\]](#page-20-9). From our analysis in Section [3,](#page-7-0) particularly [\(3.22\)](#page-12-0) and [\(3.23\)](#page-12-1), we observe that the strongly convex conditions used to bound the iterative difference is significantly larger than the corresponding terms, exceeding them by at least two orders $(O(k^2))$. If we consider a convex objective function augmented with a penalty term that asymptotically approaches zero, such as:

$$
\min_{x \in \mathbb{R}^x} g_k(x) = f(x) + \frac{1}{2k} ||x - x^*||^2,
$$

then by applying NAG- α with $r > 2\alpha$, it is possible to achieve the $O(1/k^{2\alpha})$ convergence rate for $\alpha \in (1, +\infty)$. However, in practical scenarios, constructing such a form is infeasible, as it presupposes prior knowledge of the optimal solution, which contradicts the essence of iterative optimization. This reveals a significant gap between theoretical results and practical applicability, as the strongly convex conditions used are overly restrictive and rarely hold in real-world problems. Therefore, exploring weaker conditions beyond strong convexity is a promising direction for future research. Such efforts could help bridge the gap between theory and practice, expanding the practical applicability of accelerated optimization methods.

Acknowledgements

We thanks Bowen Li for helpful discussions. Mingwei Fu was partially supported by the Hua Loo-Keng scholarship from CAS. This project was partially supported by the startup fund from SIMIS.

References

- H. Attouch and J. Peypouquet. The rate of convergence of nesterov's accelerated forward-backward method is actually faster than $1/k^2$. SIAM Journal on Optimization, 26(3):1824–1834, 2016.
- H. Attouch, P.-E. Maingé, and P. Redont. A second-order differential system with hessian-driven damping; application to non-elastic shock laws. Differential Equations and Applications, $4(1)$: 27–65, 2012.
- H. Attouch, J. Peypouquet, and P. Redont. A dynamical approach to an inertial forward-backward algorithm for convex minimization. SIAM Journal on Optimization, 24(1):232–256, 2014.
- A. Beck. First-order methods in optimization. SIAM, 2017.
- A. Beck and M. Teboulle. A fast iterative shrinkage-thresholding algorithm for linear inverse problems. SIAM journal on imaging sciences, 2(1):183–202, 2009.
- A. Cauchy. Méthode générale pour la résolution des systemes d'équations simultanées. Comp. Rend. Sci. Paris, 25(1847):536–538, 1847.
- A. Chambolle and T. Pock. An introduction to continuous optimization for imaging. Acta Numerica, 25:161–319, 2016.
- S. Chen, B. Shi, and Y.-x. Yuan. Gradient norm minimization of nesterov acceleration: $o(1/k^3)$. arXiv preprint arXiv:2209.08862, 2022a.
- S. Chen, B. Shi, and Y.-x. Yuan. Revisiting the acceleration phenomenon via high-resolution differential equations. *arXiv preprint arXiv:2212.05700*, 2022b.
- S. Chen, B. Shi, and Y.-x. Yuan. On underdamped nesterov's acceleration. arXiv preprint arXiv:2304.14642, 2023.
- M. Fu and B. Shi. Lyapunov analysis for monotonically forward-backward accelerated algorithms. arXiv preprint arXiv:2412.13527, 2024.
- P. Giselsson and S. Boyd. Monotonicity and restart in fast gradient methods. In 53rd IEEE Conference on Decision and Control, pages 5058–5063. IEEE, 2014.
- B. Li, B. Shi, and Y.-X. Yuan. Linear convergence of ISTA and FISTA. arXiv preprint arXiv:2212.06319, 2022a.
- B. Li, B. Shi, and Y.-x. Yuan. Proximal subgradient norm minimization of ista and fista. arXiv preprint arXiv:2211.01610, 2022b.
- B. Li, B. Shi, and Y.-x. Yuan. Linear convergence of forward-backward accelerated algorithms without knowledge of the modulus of strong convexity. $SIAM$ Journal on Optimization, $34(2)$: 2150–2168, 2024a.
- B. Li, B. Shi, and Y.-X. Yuan. Linear convergence of ISTA and FISTA. Journal of the Operations Research Society of China, pages 1–19, 2024b.
- A. S. Nemirovsky and D. B. Yudin. Problem complexity and method efficiency in optimization. John Wiley, New York, 1983.
- Y. Nesterov. A method of solving a convex programming problem with convergence rate $o(1/k^2)$. Soviet Mathematics-Doklady,, 27(2):372–376, 1983.
- Y. Nesterov. Lectures on Convex Optimization, volume 137. Springer, second edition, 2018.
- B. O'Donoghue and E. Candes. Adaptive restart for accelerated gradient schemes. Foundations of computational mathematics, 15:715–732, 2015.
- B. Shi, S. S. Du, M. I. Jordan, and W. J. Su. Understanding the acceleration phenomenon via high-resolution differential equations. Mathematical Programming, pages 79–148, 2022.
- W. Su, S. Boyd, and E. J. Candes. A differential equation for modeling nesterov's accelerated gradient method: Theory and insights. Journal of Machine Learning Research, 17(153):1–43, 2016.
- A. Wibisono, A. C. Wilson, and M. I. Jordan. A variational perspective on accelerated methods in optimization. proceedings of the National Academy of Sciences, 113(47):E7351–E7358, 2016.