THE LOCAL DIAMETER TWO PROPERTY AND THE DIAMETER TWO PROPERTY IN SPACES OF LIPSCHITZ FUNCTIONS

RAINIS HALLER, JAAN KRISTJAN KAASIK, AND ANDRE OSTRAK

Abstract. We separate the local diameter two property from the diameter two property and their weak-star counterparts from each other in spaces of Lipschitz functions. We characterise the w^* -LD2P, the w^* -D2P, the LD2P, and the SD2P in these spaces. We introduce a generalised version of cyclical monotonicity to study functionals on spaces of Lipschitz functions.

1. INTRODUCTION

Throughout the paper, let M be a pointed metric space, i.e. a metric space with a fixed point 0, with a distance function d. We denote by $\text{Lip}_0(M)$ the Banach space of all Lipschitz functions $f: M \to \mathbb{R}$ satisfying $f(0) = 0$ equipped with the norm

$$
||f|| = \sup \left\{ \frac{|f(x) - f(y)|}{d(x, y)} : x, y \in M, x \neq y \right\}.
$$

Let $\delta: M \to \mathrm{Lip}_0(M)^*$ be the canonical isometric embedding of M into $\mathrm{Lip}_0(M)^*$, which is given by $x \mapsto \delta_x$, where $\delta_x(f) = f(x)$. The norm closed linear span of $\delta(M)$ in $\text{Lip}_0(M)^*$ is called the *Lipschitz-free space* over M and is denoted by $\mathcal{F}(M)$. It is well known that $\mathcal{F}(M)^* = \text{Lip}_0(M)$.

Let X be a real Banach space with the closed unit ball B_X . Recall [\[2,](#page-15-0) [3\]](#page-15-1) that X has the

- (1) local diameter 2 property (briefly, $LD2P$) if every slice of B_X has diameter 2;
- (2) diameter 2 property (briefly, D2P) if every nonempty relatively weakly open subset of B_X has diameter 2;
- (3) strong diameter 2 property (briefly, SD2P) if every convex combination of slices of B_X has diameter 2, i.e. the diameter of $\sum_{i=1}^n \lambda_i S_i$ is 2 whenever $n \in \mathbb{N}, \lambda_1, \ldots, \lambda_n \geq 0$ with $\sum_{i=1}^n \lambda_i = 1$, and S_1, \ldots, S_n are slices of B_X ;
- (4) symmetric strong diameter 2 property (briefly, $SSD2P$) if for every $n \in \mathbb{N}$, every family $\{S_1, \ldots, S_n\}$ of slices of B_X , and every $\varepsilon > 0$, there exist $f_1 \in S_1, \ldots, f_n \in S_n$, and $g \in B_X$ with $||g|| > 1 - \varepsilon$ such that $f_i \pm g \in S_i$ for every $i \in \{1, \ldots, n\}.$

If X is a dual space, then the weak-star counterparts of these diameter two properties $(w^*$ -LD2P, w^* -D2P, w^* -SD2P, and w^* -SSD2P) are defined by replacing slices and weakly open subsets in the above definitions by w^* -slices and w^* -open subsets,

²⁰²⁰ Mathematics Subject Classification. Primary 46B04; Secondary 46B20.

Key words and phrases. Spaces of Lipschitz functions, Lipschitz-free spaces, diameter two properties, de Leeuw's map, finitely additive measures.

respectively. It is known that these properties are distinct and $(4) \Rightarrow (3) \Rightarrow (2) \Rightarrow$ (1) , and that these properties are distinct from their weak-star counterparts in dual Banach spaces.

In this paper, we study these diameter two properties in spaces of Lipschitz functions. The w^* -SD2P and the w^* -SSD2P in $\text{Lip}_0(M)$ have been characterised and shown to be different in [\[16,](#page-15-2) [17\]](#page-15-3). It has been shown in [\[13,](#page-15-4) [15\]](#page-15-5), that the D2P, the SD2P, and the SSD2P are all different and that all of the diameter two properties are different from their weak-star counterparts in $\text{Lip}_0(M)$ (in fact, it is known that the w ˚-SSD2P does not imply the LD2P in these spaces). It has remained unknown whether the $(w^*$ -)LD2P differs from the $(w^*$ -)D2P in $\text{Lip}_0(M)$ (see [\[13,](#page-15-4) Question 5.4] and [\[15,](#page-15-5) p. 3]). As far as we are aware, there are no known examples of dual Banach spaces with the LD2P but without the D2P (see $[1, 10, 11]$ $[1, 10, 11]$ $[1, 10, 11]$ for known examples of Banach spaces with the LD2P but without the D2P).

In Example [5.2,](#page-13-0) we construct a metric space M for which the space $\text{Lip}_0(M)$ has the LD2P, but lacks the w^* -D2P. Therefore, we answer [\[13,](#page-15-4) Question 5.4] (see also [\[15,](#page-15-5) p. 3]) and give an example of a dual Banach space with the LD2P that lacks the D2P. To achieve this, we introduce and make use of a representation theorem for $\text{Lip}_0(M)^*$, which might be of independent interest. We also characterise the w^* -LD2P, the w^* -D2P, the LD2P, and the SD2P in $\text{Lip}_0(M)$. Recall that the LD2P, the D2P, and the SD2P have dual formulations in terms of local octahedrality, weak octahedrality, and octahedrality (see [\[9\]](#page-15-9) and [\[14\]](#page-15-10)). Consequently, our results can also be restated in terms of octahedrality.

The paper is outlined as follows. In section 2, we introduce a representation theorem for the dual of the space of Lipschitz functions. In section 3, we present new conditions for the space $\text{Lip}_0(M)$ to have the LD2P and the SD2P. In section 4, we give characterisations of the w^* -LD2P and the w^* -D2P for the spaces of Lipschitz functions. Finally, in section 5, we provide an example of a metric space M , for which the space $\text{Lip}_0(M)$ has the LD2P but lacks the w^* -D2P.

In order to study the space $\text{Lip}_0(M)^*$, we make use of the following idea. Set

$$
\widetilde{M} = \big\{ (x, y) \colon x, y \in M, \ x \neq y \big\}.
$$

The de Leeuw's map $\Phi\colon \mathrm{Lip}_0(M) \to \ell_\infty(M)$ (see [\[20,](#page-15-11) Definition 2.31 and Theorem 2.35) is a linear isometry defined by $\Phi f = \hat{f}$, where

$$
\tilde{f}(x,y) = \frac{f(x) - f(y)}{d(x,y)}.
$$

Recall that $\ell_{\infty}(\widetilde{M})^* = ba(\widetilde{M})$, the Banach space of bounded and finitely additive signed measures on the power set of \widetilde{M} , with the norm $\|\mu\| = |\mu|(\widetilde{M})$. Thus, for every $F \in \text{Lip}_0(M)^*$, there exists $\mu \in ba(\tilde{M})$ with $\|\mu\| = \|F\|$ such that $\Phi^*\mu = F$, i.e.

$$
F(f) = \int_{\widetilde{M}} \widetilde{f} d\mu \quad \text{for all } f \in \text{Lip}_0(M).
$$

One may also regard the de Leeuw's map as embedding $\text{Lip}_0(M)$ into the Banach space $C(\beta M)$ of continuous functions on the Stone–Cech compactification of M (see [\[20,](#page-15-11) p. 102]). This approach has been used to associate elements of the dual space $\text{Lip}_0(M)^*$ with countably additive Radon measures on βM . See, e.g. [\[5,](#page-15-12) [6,](#page-15-13) [7,](#page-15-14) [18\]](#page-15-15) for recent developments using this method.

Notations. Given a real Banach space X , we denote the closed unit ball, the unit sphere, and the dual space of X by B_X , S_X , and X^* , respectively. A slice of B_X is a set of the form

$$
S(x^*, \alpha) = \{ x \in B_X : x^*(x) > 1 - \alpha \},
$$

where $x^* \in S_{X^*}$ and $\alpha > 0$. If X is a dual space, then slices whose defining functional comes from the predual of X are called w^* -slices.

For $x, y \in M$ with $x \neq y$, we define a *molecule* $m_{x,y}$ as a norm one element $\frac{\delta_x - \delta_y}{d(x,y)}$ in $\mathcal{F}(M)$.

Define $\pi \colon \widetilde{M} \to \mathcal{P}(M)$ by $\pi(x, y) = \{x, y\}$ for all $(x, y) \in \widetilde{M}$. Note, that for a subset A of \widetilde{M} , we have

 $\pi(A) = \{x \in M : \text{there exists } y \in M \text{ such that } (x, y) \in A \text{ or } (y, x) \in A\}.$

2. Dual of the space of Lipschitz functions

Every functional from the space $\text{Lip}_0(M)^*$ can be extended norm-preservingly to a finitely additive measure $\mu \in ba(\overline{M})$. However, given a measure $\mu \in ba(\overline{M})$, it is difficult to determine whether $\|\Phi^*\mu\| = \|\mu\|$. In this section, we provide a way of doing so.

Recall that a subset A of \tilde{M} is said to be *cyclically monotonic* (see, e.g. [\[8,](#page-15-16) [19\]](#page-15-17)) if for any finite sequence of pairs $(x_1, y_1), \ldots, (x_n, y_n) \in A$, we have

$$
\sum_{i=1}^{n} d(x_i, y_{i+1}) \geq \sum_{i=1}^{n} d(x_i, y_i),
$$

where $y_{n+1} = y_1$. We introduce the following generalisation of this notion.

Definition 2.1. Let $\gamma \in (0, 1]$ and let A be a subset of \widetilde{M} . We say that A is γ -cyclically monotonic if for any finite sequence of pairs $(x_1, y_1), \ldots, (x_n, y_n) \in A$, we have

$$
\sum_{i=1}^{n} \min \left\{ d(x_i, y_{i+1}) - \gamma d(x_i, y_i), d(y_i, y_{i+1}) \right\} \ge 0,
$$

where $y_{n+1} = y_1$.

Note that a subset A of \widetilde{M} is cyclically monotonic if and only if it is 1-cyclically monotonic. The following result generalises [\[8,](#page-15-16) Theorem 2.4 (i) \Leftrightarrow (iv)].

Proposition 2.2. Let A be a subset of \widetilde{M} and let $\gamma \in (0, 1]$. Then A is γ -cyclically monotonic if and only if there exists $f \in B_{\text{Lip}_0(M)}$ satisfying $f(m_{x,y}) \geq \gamma$ for all $(x, y) \in A$.

Proof. First, assume that A is γ -cyclically monotonic. Write $A = \{(x_i, y_i) \in M : i \in$ If for a set of indices I. For all $i, j \in I$, define

$$
\beta_{ij} = \min \left\{ d(x_i, y_j) - \gamma d(x_i, y_i), d(y_i, y_j) \right\}.
$$

It is clear that $\beta_{jj} = 0$ for all $j \in I$. Therefore, by [\[8,](#page-15-16) Lemma 2.2], there exist real numbers α_j , $j \in I$, such that for all $i, j \in I$,

$$
\alpha_i \leq \alpha_j + \beta_{ij}.
$$

(Note that [\[8,](#page-15-16) Lemma 2.2] is originally stated for the case $I = N$, but the result remains valid for any set I with an almost identical proof.)

Define $f \colon M \to \mathbb{R}$ by

$$
f(x) = \inf_{i \in I} (\alpha_i + d(x, y_i)).
$$

Note that for all $i, j \in I$,

$$
f(y_i) - f(y_j) = \alpha_i - \alpha_j \leq \beta_{ij} \leq d(y_i, y_j).
$$

Thus, f is a 1-Lipschitz McShane extension from $\{y_i : i \in I\}$ to the entire space M. For all $i, j \in I$, we have $\alpha_i \leq \alpha_j + \beta_{ij}$, and therefore,

$$
\alpha_i + \gamma d(x_i, y_i) \leq \alpha_j + d(x_i, y_j).
$$

Thus, for all $i \in I$, we have

$$
f(x_i) = \inf_{j \in I} (\alpha_j + d(x_i, y_j)) \geq \alpha_i + \gamma d(x_i, y_i) = f(y_i) + \gamma d(x_i, y_i).
$$

Consequently, $f(m_{x,y}) \geq \gamma$ for all $(x, y) \in A$.

Now assume that there exists $f \in B_{\text{Lip}_0(M)}$ satisfying $f(m_{x,y}) \geq \gamma$ for all $(x, y) \in$ A. For any finite sequence of pairs $(x_1, y_1), \ldots, (x_n, y_n) \in A$ and $y_{n+1} = y_1$, we get

$$
\sum_{i=1}^{n} \min \{ d(x_i, y_{i+1}) - \gamma d(x_i, y_i), d(y_i, y_{i+1}) \}
$$
\n
$$
\geqslant \sum_{i=1}^{n} \min \{ f(x_i) - f(y_{i+1}) - f(x_i) + f(y_i), f(y_i) - f(y_{i+1}) \}
$$
\n
$$
= \sum_{i=1}^{n} f(y_i) - f(y_{i+1}) = 0.
$$

 \Box

We are now prepared to provide criteria for determining whether a given $\mu \in$ $ba(\widetilde{M})$ satisfies $\|\Phi^*\mu\| = \|\mu\|$. Let $\mathfrak{r}: \widetilde{M} \to \widetilde{M}$ be the reflection mapping defined by $\mathfrak{r}(x, y) = (y, x).$

Theorem 2.3. Let $\mu = \mu^+ - \mu^- \in ba(\widetilde{M})$ be the Jordan decomposition of μ . The following statements are equivalent:

- (i) $\|\Phi^*\mu\| = \|\mu\|.$
- (ii) For every $\gamma \in (0, 1)$, there exists a subset A of M and an $f \in B_{\text{Lip}_0(M)}$ satisfying

$$
\mu^+(A) + \mu^-(\mathfrak{r}(A)) \geq \gamma |\mu|(\widetilde{M})
$$

and $f(m_{x,y}) \geq \gamma$ for all $(x, y) \in A$.

(iii) For every $\gamma \in (0, 1)$, there exists a γ -cyclically monotonic subset A of \widetilde{M} satisfying

$$
\mu^+(A) + \mu^-(\mathfrak{r}(A)) \geq \gamma|\mu|(\widetilde{M}).
$$

Proof. Throughout the proof, we assume without loss of generality that $\|\mu\| = 1$. Let $F = \Phi^* \mu$.

(i) \Rightarrow (ii). Let $\gamma \in (0, 1)$, let $f \in B_{\text{Lip}_0(M)}$ be such that $F(f) > 1 - (1 - \gamma)^2$, and let

$$
A = \big\{ (x, y) \in \widetilde{M} \colon f(m_{x, y}) \geqslant \gamma \big\}.
$$

Then $\mu^+(A) + \mu^-(\mathfrak{r}(A)) \ge \gamma$ because, by setting $B = \mathfrak{r}(A)$, we get

$$
1 - (1 - \gamma)^2 < F(f) = \int_A \tilde{f} d\mu + \int_B \tilde{f} d\mu + \int_{\widetilde{M} \setminus (A \cup B)} \tilde{f} d\mu
$$
\n
$$
\leq \mu^+(A) + \mu^-(B) + \gamma |\mu| \big(\widetilde{M} \setminus (A \cup B) \big)
$$
\n
$$
\leq \mu^+(A) + \mu^-(B) + \gamma \big(1 - \mu^+(A) - \mu^-(B) \big)
$$
\n
$$
= \gamma + (1 - \gamma) \big(\mu^+(A) + \mu^-(B) \big).
$$

 $(ii) \Leftrightarrow (iii)$ follows directly from Proposition [2.2.](#page-2-0)

(ii) \Rightarrow (i). It suffices to show that $||F|| \ge 1$. Fix $\gamma \in (0, 1)$, let a subset A of M and a function $f \in B_{\text{Lip}_0(M)}$ be as in (ii), and set $B = \mathfrak{r}(A)$. Then

$$
\int_A \tilde{f} d\mu \ge \gamma \mu^+(A) \quad \text{and} \quad \int_B \tilde{f} d\mu \ge \gamma \mu^-(B).
$$

Consequently,

$$
\|F\| \ge F(f) = \int_A \tilde{f} d\mu + \int_B \tilde{f} d\mu + \int_{\widetilde{M} \setminus (A \cup B)} \tilde{f} d\mu
$$

\n
$$
\ge \gamma \left(\mu^+(A) + \mu^-(B)\right) - (1 - \gamma)
$$

\n
$$
\ge \gamma^2 |\mu| (\widetilde{M}) - (1 - \gamma)
$$

\n
$$
= \gamma^2 - (1 - \gamma).
$$

Letting $\gamma \to 1$, we get $||F|| \ge 1$.

Given $F \in \text{Lip}_0(M)^*$, the measure $\mu \in ba(\widetilde{M})$ satisfying $\Phi^*\mu = F$ and $\|\mu\| = \|F\|$ is not unique. We note, similarly to [\[4,](#page-15-18) Proposition 3], that μ can be chosen to be positive.

Proposition 2.4. Let $\nu \in ba(\widetilde{M})$. Then there exists a positive $\mu \in ba(\widetilde{M})$ satisfying $\Phi^*\mu = \Phi^*\nu$ and $\|\mu\| = \|\nu\|$. Consequently, for every $F \in \text{Lip}_0(M)^*$, there exists a positive $\mu \in ba(\widetilde{M})$ satisfying $\Phi^*\mu = F$ and $\|\mu\| = \|F\|.$

Proof. Let $\nu = \nu^+ - \nu^- \in ba(\widetilde{M})$ be the Jordan decomposition of ν and let A, B be subsets of \widetilde{M} satisfying $A \cap B = \varnothing$, $A \cup B = \widetilde{M}$, and $\nu^+(B) = 0$ and $\nu^-(A) = 0$. Define $\mu \in ba(\widetilde{M})$ by

$$
\mu(C) = \nu(A \cap C) - \nu(B \cap \mathfrak{r}(C)).
$$

For any $f \in \text{Lip}_0(M)$, we have

$$
(\Phi^*\mu)(f) = \int_{\widetilde{M}} \widetilde{f} d\mu = \int_A \widetilde{f} d\mu + \int_{\mathfrak{r}(B)} \widetilde{f} d\mu
$$

=
$$
\int_A \widetilde{f} d\nu + \int_B \widetilde{f} d\nu = \int_{\widetilde{M}} \widetilde{f} d\nu = (\Phi^*\nu)(f).
$$

Furthermore,

$$
\|\mu\| = \mu(\widetilde{M}) = \nu(A) - \nu(B) = \nu^+(A) + \nu^-(B) = |\nu|(\widetilde{M}) = \|\nu\|.
$$

Definition 2.5. We call $\mu \in ba(\overline{M})$ *optimal* if it is positive and satisfies $\|\Phi^*\mu\|$ = $\|\mu\|.$

 \Box

Proposition [2.4](#page-4-0) thus establishes that for every $F \in \text{Lip}_0(M)^*$, there exists an optimal $\mu \in ba(\tilde{M})$ satisfying $\Phi^*\mu = F$.

Throughout the rest of the paper we work exclusively with positive measures. To this end, we state the following special case of Theorem [2.3.](#page-3-0)

Corollary 2.6. Let $\mu \in ba(\widetilde{M})$ be positive. The following statements are equivalent: (i) $\|\Phi^*\mu\| = \|\mu\|.$

- (ii) For every $\gamma \in (0, 1)$, there exist a subset A of M and an $f \in B_{\text{Lip}_0(M)}$ satisfying $\mu(A) \geq \gamma \mu(\widetilde{M})$ and $f(m_{x,y}) \geq \gamma$ for all $(x, y) \in A$.
- (iii) For every $\gamma \in (0, 1)$, there exists a γ -cyclically monotonic subset A of M satisfying $\mu(A) \geq \gamma \mu(M)$.

We conclude this section by showing that, for certain metric spaces, a strengthened version of Corollary [2.6](#page-5-0) holds. To this end, we rely on the following observation.

Lemma 2.7. Let $n \in \mathbb{N}$ and let M be a metric space with $d(x, y) \in \{0, 1, \ldots, n\}$ for all $x, y \in M$. Let $\mu \in ba(\overline{M})$ be positive, let $\gamma \in (0, 1)$, and let A be a γ -cyclically monotonic subset of \widetilde{M} . Then there exists a cyclically monotonic subset B of A with $\mu(B) \geq \mu(A) - 2n(1 - \gamma)\mu(\widetilde{M}).$

Proof. We assume without loss of generality, that $\mu(\widetilde{M}) = 1$ and $n(1 - \gamma) < 1$. Throughout the proof, we denote the floor, ceiling, and fractional part of $\alpha \in \mathbb{R}$ by $[\alpha]$, $[\alpha]$, and $\{\alpha\}$, respectively. Write $A = \{(x_i, y_i) \in M : i \in I\}$ for a set of indices I. For all $i, j \in I$, we define

$$
\beta_{ij} = d(x_i, y_j) - d(x_i, y_i)
$$

and

$$
\widetilde{\beta}_{ij} = \min\{d(x_i, y_j) - \gamma d(x_i, y_i), d(y_i, y_j)\},\
$$

and note that $\widetilde{\beta}_{ij} \leq \beta_{ij} + n(1 - \gamma)$.

By [\[8,](#page-15-16) Lemma 2.2], there exist $\tilde{\alpha}_i \in \mathbb{R}$, $i \in I$, so that $\tilde{\alpha}_i \leq \tilde{\alpha}_i + \tilde{\beta}_{ij}$ (as in the proof of Proposition [2.2,](#page-2-0) we note that [\[8,](#page-15-16) Lemma 2.2] remains valid for any set I). By $[8, \text{Lemma } 2.2]$, it suffices to show that there exist a subset J of I and elements $\alpha_i \in \mathbb{R}, i \in J$, so that $\alpha_i \leq \alpha_j + \beta_{ij}$ for all $i, j \in J$ and $\mu(B) \geq \mu(A) - 2n(1 - \gamma)$, where $B = \{(x_i, y_i): i \in J\}.$

Fix $K \in \mathbb{N}$ so that $\frac{1}{2K} \leq n(1-\gamma) \leq \frac{1}{K}$. For every $k \in \{1, ..., K\}$, define a subset I_k of I by

$$
I_k = \left\{ i \in I \colon \{ \widetilde{\alpha}_i \} \in \left[\frac{k-1}{K}, \frac{k}{K} \right) \right\}
$$

and a subset $A_k = \{(x_i, y_i): i \in I_k\}$ of A. Fix $k \in \{1, ..., K\}$ such that $\mu(A_k) \leq$ $1/K$. Let $J = I\backslash I_k$, let $B = \{(x_i, y_i) : i \in J\}$, and let $\alpha_i = [\tilde{\alpha}_i - \frac{k}{K}]$ for each $i \in J$. Note that $B = A \backslash A_k$ and that

$$
\mu(B) = \mu(A) - \mu(A_k) \ge \mu(A) - \frac{1}{K} \ge \mu(A) - 2n(1 - \gamma).
$$

Fix $i, j \in J$. It remains to verify that $\alpha_i \leq \alpha_j + \beta_{ij}$. Note that

$$
\alpha_i - \alpha_j = \begin{cases} \lfloor \widetilde{\alpha}_i - \widetilde{\alpha}_j \rfloor, & \text{if } \{ \widetilde{\alpha}_i - \frac{k}{K} \} \geqslant \{ \widetilde{\alpha}_j - \frac{k}{K} \}; \\ \lfloor \widetilde{\alpha}_i - \widetilde{\alpha}_j \rfloor + 1, & \text{if } \{ \widetilde{\alpha}_i - \frac{k}{K} \} < \{ \widetilde{\alpha}_j - \frac{k}{K} \}. \end{cases}
$$

If $\{\widetilde{\alpha}_i - \frac{k}{K}\} \geq \{\widetilde{\alpha}_j - \frac{k}{K}\}\text{, then}$

 $\alpha_i - \alpha_j = \left[\widetilde\alpha_i - \widetilde\alpha_j\right] \leqslant \left[\widetilde\beta_{ij}\right] \leqslant \left\lfloor\beta_{ij} + n(1-\gamma)\right\rfloor = \beta_{ij}.$

Now assume that $\{\widetilde{\alpha}_i - \frac{k}{K}\} < \{\widetilde{\alpha}_j - \frac{k}{K}\}\$. It suffices to show that $\widetilde{\alpha}_i - \widetilde{\alpha}_j < \beta_{ij}$ because then

$$
\alpha_i - \alpha_j = [\widetilde{\alpha}_i - \widetilde{\alpha}_j] + 1 \leq [\beta_{ij}] = \beta_{ij}.
$$

To obtain a contradiction, suppose that $\tilde{\alpha}_i - \tilde{\alpha}_j \ge \beta_{ij}$. Then

$$
\beta_{ij} \leq \tilde{\alpha}_i - \tilde{\alpha}_j \leq \tilde{\beta}_{ij} \leq \beta_{ij} + n(1 - \gamma),
$$

and therefore,

$$
\{\widetilde{\alpha}_i - \frac{k}{K}\} - \{\widetilde{\alpha}_j - \frac{k}{K}\} = \{\widetilde{\alpha}_i - \widetilde{\alpha}_j\} - 1 \le n(1 - \gamma) - 1.
$$

Consequently,

$$
\{\widetilde{\alpha}_j - \frac{k}{K}\} \geq 1 - n(1 - \gamma) + \{\widetilde{\alpha}_i - \frac{k}{K}\} \geq 1 - \frac{1}{K}.
$$

However, by the definition of J, we have $\{\widetilde{\alpha}_j - \frac{k}{K}\}\in [0, 1 - \frac{1}{K}]$ \Box

Proposition 2.8. Let $n \in \mathbb{N}$ and let M be a pointed metric space with $d(x, y) \in$ $\{0, 1, \ldots, n\}$ for all $x, y \in M$. Let $\mu \in ba(\widetilde{M})$ be positive. The following conditions are equivalent:

- (i) $\|\Phi^*\mu\| = \|\mu\|.$
- (ii) For every $\gamma \in (0, 1)$, there exist a subset B of M and an $f \in B_{\text{Lip}_0(M)}$ satisfying $\mu(B) \geq \gamma \mu(\widetilde{M})$ and $f(m_{x,y}) = 1$ for all $(x, y) \in B$.
- (iii) For every $\gamma \in (0, 1)$, there exists a cyclically monotonic set B of \widetilde{M} with $\mu(B) \geq \gamma \mu(M).$

Proof. (ii) \Leftrightarrow (iii) follows directly from Proposition [2.2.](#page-2-0)

(iii) \Rightarrow (i) is clear from Corollary [2.6](#page-5-0) (iii) \Rightarrow (i).

(i)⇒(iii). Let $\gamma \in (0, 1)$ and let $\tilde{\gamma} = \frac{2n+\gamma}{2n+1}$. By Corollary [2.6](#page-5-0) (i)⇒(iii), there exists a $\tilde{\gamma}$ -cyclically monotonic subset A of \widetilde{M} with $\mu(A) \geq \tilde{\gamma}\mu(\widetilde{M})$. By Lemma [2.7,](#page-5-1) there exists a cyclically monotonic subset B of A satisfying

$$
\mu(B) \ge \mu(A) - 2n(1 - \tilde{\gamma})\mu(\widetilde{M}) \ge (\tilde{\gamma} - 2n(1 - \tilde{\gamma}))\mu(\widetilde{M})
$$

$$
= ((2n + 1)\tilde{\gamma} - 2n)\mu(\widetilde{M}) = \gamma\mu(\widetilde{M}).
$$

 \Box

3. Characterisations of LD2P and SD2P for spaces of Lipschitz FUNCTIONS.

In this section we characterise the LD2P and the SD2P for the spaces of Lipschitz functions. The following lemma is useful for working with slices of such Banach spaces.

Lemma 3.1. Let $\mu \in S_{ba(\widetilde{M})}$ be optimal, let $\alpha > 0$, let $f \in S(\Phi^*\mu, \alpha^2)$, and let

$$
A = \{(x, y) \in \widetilde{M} : f(m_{x,y}) \geq 1 - \alpha\}.
$$

Then $\mu(A) \geq 1 - \alpha$.

Proof. It suffices to note that

$$
1 - \alpha^2 \leq (\Phi^* \mu)(f) = \int_A \tilde{f} d\mu + \int_{\widetilde{M} \setminus A} \tilde{f} d\mu \leq \mu(A) + (1 - \alpha)\mu(\widetilde{M} \setminus A)
$$

= $\mu(A) + (1 - \alpha)(1 - \mu(A)) = 1 - \alpha + \alpha \mu(A).$

We are now ready to characterise the LD2P for the spaces of Lipschitz functions.

Proposition 3.2. The space $\text{Lip}_0(M)$ has the LD2P if and only if for all optimal $\mu \in S_{ba(\widetilde{M})}$ and $\gamma \in (0, 1)$, there exist a subset A of M and $u, v \in M$ with $u \neq v$ such that $\mu(A) \geq \gamma$ and the sets $A \cup \{(u, v)\}\$ and $A \cup \{(v, u)\}\$ are both γ -cyclically monotonic.

Proof. (\Leftarrow) Let $F \in S_{\text{Lip}_0(M)}$ and let $\alpha > 0$. It suffices to show that for any $\gamma \in (0, 1)$, there exist $f, g \in S(F, \alpha)$ satisfying $||f - g|| \geq 2\gamma$. Let $\gamma \in (0, 1)$ be such that $\gamma^2 \geq 1 - \frac{\alpha}{2}$. By Proposition [2.4,](#page-4-0) there exists an optimal $\mu \in S_{ba(\widetilde{M})}$ with $\Phi^*\mu = F$. Therefore, there exist a subset A of \widetilde{M} and $u, v \in M$ with $u \neq v$ so that $\mu(A) \geq \gamma$ and the sets $A \cup \{(u, v)\}\$ and $A \cup \{(v, u)\}\$ are both γ -cyclically monotonic. By Proposition [2.2,](#page-2-0) there exist $f, g \in B_{\text{Lip}_0(M)}$ such that $f(m_{x,y}) \ge \gamma$ and $g(m_{x,y}) \geq \gamma$ for all $(x, y) \in A$, and $f(m_{u,v}) \geq \gamma$ and $g(m_{v,u}) \geq \gamma$. Therefore,

$$
||f - g|| \geq (f - g)(m_{u,v}) \geq 2\gamma.
$$

Note that $f \in S(F, \alpha)$ because

$$
F(f) = \int_{\widetilde{M}} \widetilde{f} d\mu = \int_A \widetilde{f} d\mu + \int_{\widetilde{M} \setminus A} \widetilde{f} d\mu \ge \gamma^2 - (1 - \gamma) > 2\gamma^2 - 1 \ge 1 - \alpha.
$$

Similarly, we get that $g \in S(F, \alpha)$.

 (\Rightarrow) Now assume that the space $\text{Lip}_0(M)$ has the LD2P. Let $\mu \in S_{ba(\widetilde{M})}$ be optimal and let $\gamma \in (0, 1)$. Set $\alpha = \frac{1-\gamma}{2}$. Since Lip₀(*M*) has the LD2P, there exist $f, g \in S(\Phi^*\mu, \alpha^2)$ with $||f - g|| > 2 - \alpha$. Therefore, there exist $u, v \in M$ so that

$$
f(m_{u,v}) + g(m_{v,u}) = (f - g)(m_{u,v}) \ge 2 - \alpha.
$$

Thus, $f(m_{u,v}) \geq 1 - \alpha$ and $g(m_{v,u}) \geq 1 - \alpha$. Set $A = B \cap C$, where

$$
B = \{(x, y) \in \widetilde{M} : f(m_{x,y}) \geq 1 - \alpha\} \quad \text{and} \quad C = \{(x, y) \in \widetilde{M} : g(m_{x,y}) \geq 1 - \alpha\}.
$$

By Proposition [2.2,](#page-2-0) B and C are γ -cyclically monotonic because $\gamma < 1 - \alpha$. Note that $A \cup \{(u, v)\}\$ and $A \cup \{(v, u)\}\$ are γ-cyclically monotonic because they are subsets of γ -cyclically monotonic sets B and C, respectively. By Lemma [3.1,](#page-6-0) we have $\mu(B) \geq 1 - \alpha$ and $\mu(C) \geq 1 - \alpha$, and hence,

$$
\mu(A) \geq 1 - 2\alpha = \gamma.
$$

The following lemma provides a tool to check whether the set $A \cup \{(u, v)\}\$ is γcyclically monotonic, given a γ -cyclically monotonic subset A of M and $(u, v) \in M$.

Lemma 3.3. Let $\gamma \in (0, 1]$, let A be a subset of M, and let $u, v \in M$ with $u \neq v$. The set $A \cup \{(u, v)\}\$ is γ -cyclically monotonic if and only if there exists $f \in B_{\text{Lip}_0(M)}$ satisfying for all $(x, y) \in A$, $f(m_{x,y}) \ge \gamma$, and for all $x, y \in \pi(A)$,

$$
f(y) - f(x) + \gamma d(u, v) \le d(x, u) + d(y, v).
$$

Proof. Assume first that $A \cup \{(u, v)\}$ is γ -cyclically monotonic. By Proposition [2.2,](#page-2-0) there exists $f \in B_{\text{Lip}_0(M)}$ satisfying $f(m_{x,y}) \geq \gamma$ for all $(x, y) \in A \cup \{(u, v)\}\$. For all $x, y \in \pi(A)$, we have

$$
\gamma d(u, v) \leqslant f(u) - f(v) \leqslant f(x) + d(x, u) - f(y) + d(y, v).
$$

Now assume that there exists $f \in B_{\text{Lip}_0(M)}$ satisfying $f(m_{x,y}) \geq \gamma$ for all $(x, y) \in$ A, and for all $x, y \in \pi(A)$,

$$
f(y) - f(x) + \gamma d(u, v) \le d(x, u) + d(y, v).
$$

We show that the set $A \cup \{(u, v)\}\$ is γ -cyclically monotonic. By Proposition [2.2,](#page-2-0) it suffices to find $g \in B_{\text{Lip}_0(M)}$ satisfying $g(m_{x,y}) \geq \gamma$ for all $(x, y) \in A \cup \{(u, v)\}.$ Without loss of generality, we may and do assume that $0 \in \pi(A)$. Define $g \in$ $B_{\text{Lip}_0(M)}$ by setting $g|_{\pi(A)} = f|_{\pi(A)}$ and $g(u) = \inf_{x \in \pi(A)} (f(x) + d(x, u))$, and then

$$
g(y) = \sup_{x \in \pi(A) \cup \{u\}} (g(x) - d(x, y)) \quad \text{for all } y \in M \setminus (\pi(A) \cup \{u\}).
$$

Clearly $g(m_{x,y}) = f(m_{x,y}) \geq \gamma$ for all $(x, y) \in A$. Note that $g(m_{u,v}) \geq \gamma$. Indeed, if $g(v) = g(u) - d(u, v)$, then this is clear; otherwise,

$$
g(u) - g(v) = \inf_{x, y \in \pi(A)} (f(x) + d(x, u) - f(y) + d(y, v)) \ge \gamma d(u, v).
$$

From Proposition [3.2](#page-7-0) and Lemma [3.3,](#page-7-1) we immediately get another characterisation of the LD2P for the spaces of Lipschitz functions.

Proposition 3.4. The space $\text{Lip}_0(M)$ has the LD2P if and only if for all optimal $\mu \in S_{ba(\widetilde{M})}$ and $\gamma \in (0, 1)$, there exist a subset A of M with $\mu(A) \geq \gamma$, functionals $f, g \in B_{\text{Lip}_0(M)}$, and elements $u, v \in M$ with $u \neq v$ satisfying for all $(x, y) \in A$, $f(m_{x,y}) \geq \gamma$ and $g(m_{x,y}) \geq \gamma$, and for all $x, y \in \pi(A)$,

$$
\max \{ f(x) - f(y), g(y) - g(x) \} + \gamma d(u, v) \le d(x, u) + d(y, v).
$$

We now characterise the SD2P for the spaces of Lipschitz functions in the spirit of Proposition [3.2.](#page-7-0) It was shown in [\[13,](#page-15-4) Example 3.1] that the w^* -SD2P and the SD2P are not equivalent properties for the spaces of Lipschitz functions. Therefore, our characterisation is not equivalent to the one given for the w^* -SD2P in [\[17,](#page-15-3) Theorem 3.1].

Proposition 3.5. The space $\text{Lip}_0(M)$ has the SD2P if and only if for all optimal $\mu_1, \ldots, \mu_n \in S_{ba(\widetilde{M})}$ and $\gamma \in (0, 1)$, there exist subsets A_1, \ldots, A_n of M and $u, v \in M$ with $u \neq v$ such that for all $i \in \{1, ..., n\}$, $\mu_i(A_i) \geq \gamma$ and the sets $A_i \cup \{(u, v)\}$ and $A_i \cup \{(v, u)\}\$ are γ -cyclically monotonic.

Proof. (\Leftarrow) Let $F_1, \ldots, F_n \in S_{\text{Lip}_0(M)}$, let $\alpha > 0$, let $\lambda_1, \ldots, \lambda_n \geq 0$ satisfy $\sum_{i=1}^{n} \lambda_i = 1$, and let $C = \sum_{i=1}^{n} \lambda_i S(F_i, \alpha)$. It suffices to show that for any $\gamma \in (0, 1)$, there exist $f, g \in C$ satisfying $||f - g|| \ge 2\gamma$. Let $\gamma \in (0, 1)$ be such that $\gamma^2 \ge 1 - \frac{\alpha}{2}$. By Proposition [2.4,](#page-4-0) there exist optimal $\mu_1, \ldots, \mu_n \in S_{ba(\widetilde{M})}$ with $\Phi^* \mu_i = F_i$ for every $i \in \{1, ..., n\}$. Therefore, there exist subsets $A_1, ..., A_n$ of M and $u, v \in M$ with $u \neq v$ such that for all $i \in \{1, ..., n\}$, $\mu_i(A_i) \geq \gamma$ and the sets $A_i \cup \{(u, v)\}$ and $A_i \cup \{(v, u)\}\$ are γ -cyclically monotonic.

Fix $i \in \{1, \ldots, n\}$. By Theorem [2.3,](#page-3-0) there exist $f_i, g_i \in B_{\text{Lip}_0(M)}$ such that $f_i(m_{x,y}) \geq \gamma$ and $g_i(m_{x,y}) \geq \gamma$ for all $(x, y) \in A_i$, and $f_i(m_{u,v}) \geq \gamma$ and $g_i(m_{v,u}) \geq$ γ . Note that $f_i \in S(F_i, \alpha)$ because

$$
F_i(f_i) = \int_{\widetilde{M}} \widetilde{f}_i d\mu_i = \int_{A_i} \widetilde{f}_i d\mu_i + \int_{\widetilde{M} \setminus A_i} \widetilde{f}_i d\mu_i \ge \gamma^2 - (1 - \gamma) > 2\gamma^2 - 1 \ge 1 - \alpha.
$$

Similarly, we get that $g_i \in S(F_i, \alpha)$.

Define $f = \sum_{i=1}^{n} \lambda_i f_i$ and $g = \sum_{i=1}^{n} \lambda_i g_i$. Then $f, g \in C$ and

$$
||f-g|| \geq \sum_{i=1}^{n} \lambda_i (f_i - g_i)(m_{u,v}) \geq 2\gamma.
$$

 (\Rightarrow) Now assume that $\mathrm{Lip}_0(M)$ has the SD2P. Let $\mu_1, \ldots, \mu_n \in S_{ba(\widetilde{M})}$ be optimal and let $\gamma \in (0, 1)$. Set $\alpha = \frac{1-\gamma}{2}$. Since Lip₀(M) has the SD2P, there exist $f_i, g_i \in$ $S(\Phi^*\mu_i, \alpha^2), i \in \{1, \ldots, n\}$, such that

$$
\left\|\frac{1}{n}\sum_{i=1}^n(f_i-g_i)\right\|>2-\frac{\alpha}{n}.
$$

Let $u, v \in M$ with $u \neq v$ be such that

$$
\frac{1}{n}\sum_{i=1}^{n}(f_i-g_i)(m_{u,v}) \geq 2-\frac{\alpha}{n}.
$$

Fix $i \in \{1, \ldots, n\}$. We have

$$
f_i(m_{u,v}) + g_i(m_{v,u}) = (f_i - g_i)(m_{u,v}) \ge n(2 - \frac{\alpha}{n}) - 2(n-1) = 2 - \alpha,
$$

and therefore, $f_i(m_{u,v}) \geq 1 - \alpha$ and $g_i(m_{v,u}) \geq 1 - \alpha$. Set $A_i = B_i \cap C_i$, where $B_i = \left\{ (x, y) \in \widetilde{M} : f_i(m_{x,y}) \geq 1 - \alpha \right\}$ and $C_i = \left\{ (x, y) \in \widetilde{M} : g_i(m_{x,y}) \geq 1 - \alpha \right\}$. By Proposition [2.2,](#page-2-0) B_i and C_i are γ -cyclically monotonic because $\gamma < 1 - \alpha$. Note that $A_i \cup \{(u, v)\}\$ and $A_i \cup \{(v, u)\}\$ are γ-cyclically monotonic because they are subsets of γ -cyclically monotonic sets B_i and C_i , respectively. By Lemma [3.1,](#page-6-0) we have $\mu_i(B_i) \geq 1 - \alpha$ and $\mu_i(C_i) \geq 1 - \alpha$, and hence,

$$
\mu_i(A_i) \geq 1 - 2\alpha = \gamma.
$$

From Lemma [3.3](#page-7-1) and Proposition [3.5,](#page-8-0) we get another characterisation of the SD2P for the spaces of Lipschitz functions, in the spirit of Proposition [3.4.](#page-8-1)

Proposition 3.6. The space $\text{Lip}_0(M)$ has the SD2P if and only if for all optimal $\mu_1, \ldots, \mu_n \in S_{ba(\widetilde{M})}$ and $\gamma \in (0, 1)$, there exist subsets A_1, \ldots, A_n of M with $\mu_1(A_1) \geqslant \gamma, \ldots, \mu_n(A_n) \geqslant \gamma$, functionals $f_1, g_1, \ldots, f_n, g_n \in B_{\mathrm{Lip}_0(M)}$, and elements $u, v \in M$ with $u \neq v$ satisfying for all $i \in \{1, ..., n\}$ and $(x, y) \in A_i$, $f_i(m_{x,y}) \ge \gamma$ and $g_i(m_{x,y}) \ge \gamma$, and for all $i \in \{1, \ldots, n\}$ and $x, y \in \pi(A_i)$,

$$
\max \{ f_i(x) - f_i(y), g_i(y) - g_i(x) \} + \gamma d(u, v) \le d(x, u) + d(y, v).
$$

Remark. In [\[13\]](#page-15-4), it was shown that if M has a property called the FLTP, then $\text{Lip}_0(M)$ has the SD2P. It remains unknown whether the FLTP is equivalent to the condition described in Proposition [3.6.](#page-9-0)

4. CHARACTERISATIONS OF w^* -D2P and w^* -LD2P for spaces of Lipschitz **FUNCTIONS**

Recall that the space $\text{Lip}_0(M)$ has the w^{*}-SD2P if and only if the space M has the LTP [\[17,](#page-15-3) Theorem 3.1], and that the space $\text{Lip}_0(M)$ has the w^{*}-SSD2P if and only if the space M has the SLTP [\[16,](#page-15-2) Theorem 2.1]. Inspired by these results, we characterise the w^* -D2P and the w^* -LD2P for the spaces of Lipschitz functions.

Definition 4.1. We say that M has the $Lip-LTP$ if, given a finite subset N of M, $\varepsilon > 0$, and $f \in B_{\text{Lip}_0(M)}$, there exist $u, v \in M$ with $u \neq v$ satisfying for all $x, y \in N$,

(4.1)
$$
(1 - \varepsilon)(|f(x) - f(y)| + d(u, v)) \le d(x, u) + d(y, v).
$$

Clearly the Lip-LTP follows from the LTP.

Proposition 4.2. The space $\text{Lip}_0(M)$ has the w^{*}-D2P if and only if M has the Lip-LTP.

Proof. If M is unbounded or not uniformly discrete, then M has the LTP and $\text{Lip}_0(M)$ has the w^{*}-SD2P by [\[12,](#page-15-19) Theorem 2.4] and [\[17,](#page-15-3) Theorem 3.1]. Therefore, we assume throughout the proof that M is bounded and uniformly discrete. i.e. there exist $r, R > 0$ such that for all $x, y \in M$ with $x \neq y$ we have $r < d(x, y) < R$.

Assume first that $\text{Lip}_0(M)$ has the w^{*}-D2P. Let N be a finite subset of M, let $\varepsilon > 0$, and let $f \in B_{\text{Lip}_0(M)}$. Let $\delta = \frac{\varepsilon r}{2R}$ and define

$$
U = \big\{ g \in B_{\text{Lip}_0(M)} \colon |(f - g)(m_{x,y})| < \delta \text{ for all } x, y \in N \text{ with } x \neq y \big\}.
$$

Note that U is not empty because $f \in U$. Since $\text{Lip}_0(M)$ has the w^{*}-D2P, there exist $g, h \in U$ with $||g - h|| > 2 - \varepsilon/2$. Consequently, there exist $u, v \in M$ with $u \neq v$ such that $g(m) + h(m) = (a + h)(m) > 0$

$$
g(m_{u,v}) + h(m_{v,u}) = (g - h)(m_{u,v}) \ge 2 - \varepsilon/2.
$$

\nThus, $g(m_{u,v}) \ge 1 - \varepsilon/2$ and $h(m_{v,u}) \ge 1 - \varepsilon/2$.
\nFix $x, y \in N$. If $f(x) - f(y) \ge 0$, then
\n
$$
(1 - \varepsilon)(|f(x) - f(y)| + d(u, v)) \le f(x) - f(y) + (1 - \varepsilon)d(u, v)
$$
\n
$$
\le h(x) - h(y) + \delta d(x, y) + (1 - \varepsilon)d(u, v)
$$
\n
$$
\le h(x) - h(y) + \frac{\varepsilon}{2}d(u, v) + (1 - \varepsilon)d(u, v)
$$
\n
$$
\le h(x) - h(y) + h(v) - h(u)
$$
\n
$$
\le d(x, u) + d(y, v).
$$

Similarly, if $f(x) - f(y) < 0$, then

$$
(1 - \varepsilon)(|f(x) - f(y)| + d(u, v)) \le f(y) - f(x) + (1 - \varepsilon)d(u, v)
$$

\n
$$
\le g(y) - g(x) + \delta d(x, y) + (1 - \varepsilon)d(u, v)
$$

\n
$$
\le g(y) - g(x) + \frac{\varepsilon}{2}d(u, v) + (1 - \varepsilon)d(u, v)
$$

\n
$$
\le g(y) - g(x) + g(u) - g(v)
$$

\n
$$
\le d(x, u) + d(y, v).
$$

Now assume that M has the Lip-LTP. Let $\mu_1, \ldots, \mu_n \in S_{\mathcal{F}(M)}$ be finitely supported and let $\alpha_1, \ldots, \alpha_n > 0$. Assume that $U = \bigcap_{i=1}^n S(\mu_i, \alpha_i)$ is not empty and fix $f \in U$. Let $N = \{0\} \cup (\bigcup_{i=1}^n \text{supp }\mu_i)$, let $\delta \in (0, 1)$, and let $\varepsilon = \frac{\delta r}{2R}$. Since M

has the Lip-LTP, there exist $u, v \in M$ with $u \neq v$ such that the inequality [\(4.1\)](#page-10-0) holds for all $x, y \in N$. Define a function $g \in B_{\text{Lip}_0(M)}$ by first setting $g|_N = f|_N$ and $g(u) = \min_{x \in N} (f(x) + d(x, u))$, and then

$$
g(y) = \max_{x \in N \cup \{u\}} (g(x) - d(x, y)) \quad \text{for all } y \in M \setminus (N \cup \{u\}).
$$

Note that $g(u) - g(v) \geq (1 - \delta) d(u, v)$. Indeed, if $g(v) = g(u) - d(u, v)$, then this is clear; otherwise, there exist $x, y \in N$ such that

$$
g(u) - g(v) = f(x) + d(x, u) - f(y) + d(y, v)
$$

\n
$$
\geq (1 - \varepsilon)d(u, v) + \varepsilon(f(x) - f(y))
$$

\n
$$
\geq (1 - \varepsilon)d(u, v) - \varepsilon d(x, y)
$$

\n
$$
\geq (1 - \frac{\delta r}{2R})d(u, v) - \frac{\delta}{2}d(u, v) \geq (1 - \delta)d(u, v).
$$

Similarly, define a function $h \in B_{\text{Lip}_0(M)}$ by first setting $h|_N = f|_N$ and $h(v) =$ $\min_{x \in N} (f(x) + d(x, v))$, and then

$$
h(y) = \max_{x \in N \cup \{v\}} (h(x) - d(x, y)) \quad \text{for all } y \in M \setminus (N \cup \{v\}),
$$

and verify that $h(v) - h(u) \geq (1 - \delta)d(u, v)$. It remains to note that $g, h \in U$ and that

$$
||g - h|| \geq (g - h)(m_{u,v}) \geq 2 - 2\delta.
$$

In Proposition [3.2](#page-7-0) and Proposition [3.4,](#page-8-1) we gave two characterisations of the LD2P for the spaces of Lipschitz functions. We now give similar characterisations of the w^* -LD2P. Note that the w^* -LD2P and the LD2P are not equivalent properties for the spaces of Lipschitz functions by [\[13,](#page-15-4) Example 3.1].

Proposition 4.3. The space $\text{Lip}_0(M)$ has the w^* -LD2P if and only if, given a finite cyclically monotonic subset A of \widetilde{M} and $\gamma \in (0, 1)$, there exist $u, v \in M$ with $u \neq v$ such that $A \cup \{(u, v)\}\$ and $A \cup \{(v, u)\}\$ are γ -cyclically monotonic.

Proof. (\Rightarrow) Assume that $\text{Lip}_0(M)$ has the w*-LD2P. Let $A = \{(x_i, y_i): i = 1, ..., n\}$ be a finite cyclically monotonic subset of \widetilde{M} and let $\gamma \in (0, 1)$. Set $\alpha = \frac{1-\gamma}{n}$ and $\mu = \sum_{i=1}^n \frac{1}{n} m_{x_i, y_i}$. By Proposition [2.2,](#page-2-0) we have $\mu \in S_{\mathcal{F}(M)}$. Since $\text{Lip}_0(M)$ has the w^{*}-LD2P, there exist $f, g \in S(\mu, \alpha)$ satisfying $||f - g|| > 1 + \gamma$. Therefore, there exist $u, v \in M$ with $u \neq v$ such that $(f - g)(m_{u,v}) \geq 1 + \gamma$. Thus, $f(m_{u,v}) \geq \gamma$ and $g(m_{v,u}) \geq \gamma$. By Proposition [2.2,](#page-2-0) $A \cup \{(u, v)\}\$ is γ-cyclically monotonic because from $f \in S(\mu, \alpha)$ we get for any $i \in \{1, \ldots, n\},\$

$$
f(m_{x_i,y_i}) = nf(\mu) - \sum_{\substack{j=1 \ j \neq i}}^n f(m_{x_j,y_j}) \geq n(1-\alpha) - (n-1) = 1 - n\alpha = \gamma.
$$

Analogously, we get that $A \cup \{(v, u)\}\$ is γ-cyclically monotonic because $g(m_{x_i, y_i}) \ge \gamma$ for all $i \in \{1, \ldots, n\}$.

(\Leftarrow) Let $\mu \in S_{\mathcal{F}(M)}$ be finitely supported, let $0 < \alpha \leq 1$, and let $\gamma \in (1 - \alpha, 1)$. It suffices to show that there exist $f, g \in S(\mu, \alpha)$ with $||f - g|| \geq 2\gamma$. By [\[20,](#page-15-11) Proposition 3.16], there exist $n \in \mathbb{N}$, $(x_1, y_1), \ldots, (x_n, y_n) \in \widetilde{M}$, and $\lambda_1, \ldots, \lambda_n > 0$ with $\sum_{i=1}^n \lambda_i = 1$ such that $\mu = \sum_{i=1}^n \lambda_i m_{x_i, y_i}$. By Proposition [2.2,](#page-2-0) $A = \{(x_i, y_i) : i =$ $1, \ldots, n\}$ is cyclically monotonic. Thus, there exist $u, v \in M$ with $u \neq v$ such that $A \cup \{(u, v)\}\$ and $A \cup \{(v, u)\}\$ are γ -cyclically monotonic. By Proposition [2.2,](#page-2-0) there exist $f, g \in B_{\text{Lip}_0(M)}$ satisfying $f(m_{x,y}) \geq \gamma$ and $g(m_{x,y}) \geq \gamma$ for all $(x, y) \in A$, and $f(m_{u,v}) \geq \gamma$ and $g(m_{v,u}) \geq \gamma$. Therefore, we have $f(\mu) \geq \gamma$ and $g(\mu) \geq \gamma$, which implies $f, g \in S(\mu, \alpha)$, and

$$
||f - g|| \ge (f - g)(m_{u,v}) \ge 2\gamma.
$$

From Lemma [3.3](#page-7-1) and Proposition [4.3,](#page-11-0) we immediately get another characterisation of the w ˚-LD2P for the spaces of Lipschitz functions.

Definition 4.4. We say that M has the 2-Lip-LTP if, given a finite cyclically monotonic subset A of M and $\varepsilon > 0$, there exist $f, g \in B_{\text{Lip}_0(M)}$ and $u, v \in M$ with $u \neq v$ satisfying for all $(x, y) \in A$, $f(m_{x,y}) \geq 1 - \varepsilon$ and $g(m_{x,y}) \geq 1 - \varepsilon$, and for all $x, y \in \pi(A),$

$$
\max \{ f(x) - f(y), g(y) - g(x) \} + (1 - \varepsilon) d(u, v) \le d(x, u) + d(y, v).
$$

Proposition 4.5. The space $\text{Lip}_0(M)$ has the w^{*}-LD2P if and only if M has the 2-Lip-LTP.

5. Separating LD2P and D2P in spaces of Lipschitz functions

In this section, we give an example of a metric space M for which the space $\text{Lip}_0(M)$ has the LD2P but not the D2P or even the w^{*}-D2P. To our knowledge, the only known examples of Banach spaces with the LD2P but without the D2P have appeared in [\[1,](#page-15-6) [10,](#page-15-7) [11\]](#page-15-8).

In [\[10\]](#page-15-7), it was shown that every Banach space containing an isomorphic copy of c_0 admits a renorming with the LD2P but without the D2P. Similar ideas were recently applied in [\[11\]](#page-15-8) to special renormings of $L_\infty[0, 1]$. It remains unclear whether any of these renormings is a dual Banach space.

In [\[1\]](#page-15-6), the authors constructed a Banach space with the LD2P but without the D₂P, using a binary tree with the root removed. This space is not a dual space as it is separable and contains (an isomorphic copy of) c_0 .

Lemma 5.1. Let M be a pointed metric space with $d(x, y) \in \mathbb{N} \cup \{0\}$ for all $x, y \in M$, and let A be a cyclically monotonic subset of M. Then there exists $f \in B_{\text{Lip}_0(M)}$ such that for all $x \in M$, $f(x) \in \mathbb{Z}$, and for all $(x, y) \in A$, $f(m_{x,y}) = 1$.

Proof. By Proposition [2.2,](#page-2-0) there exists $g \in B_{\text{Lip}_0(M)}$ satisfying $g(m_{x,y}) = 1$ for all $p(x, y) \in A$. Define a function $f : M \to \mathbb{R}$ by $f(x) = |g(x)|$, the floor of $g(x)$. Note that $f \in B_{\text{Lip}_0(M)}$ because for all $x, y \in M$,

$$
f(x) - f(y) = [g(x)] - [g(y)] \le [g(y) + d(x, y)] - [g(y)]
$$

= [g(y)] + d(x, y) - [g(y)] = d(x, y).

Furthermore, note that if $(x, y) \in A$, then

$$
f(x) - f(y) = [g(x)] - [g(y)] = [g(y) + d(x, y)] - [g(y)]
$$

= $[g(y)] + d(x, y) - [g(y)] = d(x, y).$

Example 5.2. Let $M = \{x_i, y_i, u_i^j, v_i^j : i \in \{1, 2, 3\}, j \in \mathbb{N}\}\)$ be a metric space where $d(y_1, x_2) = d(y_2, x_3) = d(y_3, x_1) = 1,$

for all $i \in \{1, 2, 3\}$ and $j \in \mathbb{N}$,

$$
d(x_i, u_i^j) = d(u_i^j, v_i^j) = d(v_i^j, y_i) = 1,
$$

and the distance between two different elements is equal to 2 in all other cases (see Figure [1\)](#page-13-1). Then the space $\text{Lip}_0(M)$ lacks the w^* -D2P but has the LD2P.

Figure 1. A representation of the metric space M in Example [5.2.](#page-13-0) The distances between points connected by a straight line segment are 1, the distances between other different points are 2.

Proof. We start by showing that the space $\text{Lip}_0(M)$ does not have the w*-D2P. By Proposition [4.2,](#page-10-1) it suffices to show that the space M lacks the Lip-LTP. Without loss of generality, we assume that $0 = x_1$. Let $N = \{x_1, x_2, x_3, y_1, y_2, y_3\}$ and let $\varepsilon = 1/14$. Define $f \in S_{\text{Lip}_0(M)}$ by

$$
f(x) = \begin{cases} 0, & \text{if } x \in \{x_1, y_3\}; \\ 1/2, & \text{if } x = y_2; \\ 3/2, & \text{if } x \in \{y_1, x_3\}; \\ 2, & \text{if } x = x_2; \\ 1, & \text{otherwise.} \end{cases}
$$

Fix $u, v \in M$ with $u \neq v$. We show that there exist $x, y \in N$ such that the inequality [\(4.1\)](#page-10-0) does not hold. First, assume that $u \in N$ or $v \in N$. Without loss of generality, let $u \in N$. Let $x = u$ and let $y \in N$ be such that $|f(x) - f(y)| = 3/2$. Then

$$
(1 - \varepsilon) (|f(x) - f(y)| + d(u, v)) \ge \frac{13}{14} \cdot \frac{5}{2} > 2 \ge d(x, u) + d(y, v).
$$

Now assume that $u, v \in M\backslash N$. If $d(u, v) = 2$, then there exist $x, y \in N$ satisfying $d(x, u) = 1$ and $|f(x) - f(y)| = 3/2$. Then

$$
(1 - \varepsilon) (|f(x) - f(y)| + d(u, v)) = \frac{13}{14} \cdot \frac{7}{2} > 3 \ge d(x, u) + d(y, v).
$$

If $d(u, v) = 1$, then there exist $i \in \{1, 2, 3\}$ and $j \in \mathbb{N}$ such that $u, v \in \{u_i^j, v_i^j\}$. Therefore, there exist $x, y \in \{x_i, y_i\}$ satisfying $d(x, u) = 1$, $d(y, v) = 1$, and $|f(x) - f(x)|$ $|f(y)| = 3/2$. Thus,

$$
(1 - \varepsilon) (|f(x) - f(y)| + d(u, v)) = \frac{13}{14} \cdot \frac{5}{2} > 2 = d(x, u) + d(y, v).
$$

Consequently, $\text{Lip}_0(M)$ does not have the w*-D2P.

Next, we show that the space $\text{Lip}_0(M)$ has the LD2P. We make use of Propo-sition [3.4.](#page-8-1) Let $\mu \in S_{ba(\widetilde{M})}$ be optimal and let $\gamma \in (0, 1)$. By Proposition [2.8,](#page-6-1) there exists a cyclically monotonic subset B of \widetilde{M} with $\mu(B) \geq \frac{1+\gamma}{2}$. Choose $j \in \mathbb{N}$ such that $C = \pi^{-1}(\{u_1^j, u_2^j, u_3^j, v_1^j, v_2^j, v_3^j\})$ satisfies $\mu(C) \leq \frac{1-\gamma}{2}$. Set $A = B\setminus C$ and note that $\mu(A) \geq \gamma$.

By Lemma [5.1,](#page-12-0) there exists $f \in B_{\text{Lip}_0(M)}$ such that for all $x \in M$, $f(x) \in \mathbb{Z}$, and for all $(x, y) \in A$, $f(m_{x,y}) = 1$. Without loss of generality, we may and do assume that $0 \in \pi(A)$ and that $f(0) = \min_{x \in \pi(A)} f(x)$. Then $f(x) \in \{0, 1, 2\}$ for all $x \in \pi(A)$.

It remains to show that there exist $u, v \in M$ with $u \neq v$ satisfying for all $x, y \in \pi(A),$

$$
|f(x) - f(y)| + d(u, v) \le d(x, u) + d(y, v).
$$

Indeed, if this is the case, then set $g = f$ and note that the subset A of \widetilde{M} , functionals $f, g \in B_{\text{Lip}_0(M)}$, and elements $u, v \in M$ satisfy the conditions of Proposition [3.4.](#page-8-1)

If there exists $i \in \{1, 2, 3\}$ such that $x_i \notin \pi(A)$ or $y_i \notin \pi(A)$, then letting $u = u_i^j$ and $v = v_i^j$, it follows that for all $x, y \in \pi(A)$, $d(x, u) \geq 2$ or $d(y, v) \geq 2$, and hence

$$
|f(x) - f(y)| + d(u, v) \le d(x, y) + 1 \le 3 \le d(x, u) + d(y, v).
$$

Now assume that $x_1, x_2, x_3, y_1, y_2, y_3 \in \pi(A)$. Note that $|f(x_i) - f(y_i)| \leq 1$ for some $i \in \{1, 2, 3\}$. Indeed, suppose for example that $f(x_1) - f(y_1) = 2$. Then $f(x_1) = 2$ and $f(y_1) = 0$, and therefore, $f(x_2) \le 1$ and $f(y_3) \ge 1$. Thus, if $|f(x_2) - f(y_2)| > 1$ and $|f(x_3) - f(y_3)| > 1$, then $f(x_2) = 0$, $f(y_2) = 2$, $f(y_3) = 2$, and $f(x_3) = 0$. But then $||f|| \ge 2$ because

$$
f(y_2) - f(x_3) = 2 = 2d(y_2, x_3).
$$

Let $i \in \{1, 2, 3\}$ be such that $|f(x_i) - f(y_i)| \leq 1$. Let $u = u_i^j$ and $v = v_i^j$, and fix $x, y \in \pi(A)$. Note that $d(u, v) = 1$, $d(x, u) \geq 1$, and $d(y, v) \geq 1$. If $x \notin \{x_i, y_i\}$ or $y \notin \{x_i, y_i\}$, then $d(x, u) \geq 2$ or $d(y, v) \geq 2$, and therefore,

$$
|f(x) - f(y)| + d(u, v) \le d(x, y) + 1 \le 3 \le d(x, u) + d(y, v).
$$

If $x, y \in \{x_i, y_i\}$, then $|f(x) - f(y)| \leq 1$, and therefore,

$$
|f(x) - f(y)| + d(u, v) \le 2 \le d(x, u) + d(y, v).
$$

 \Box

ACKNOWLEDGEMENTS

This work was supported by the Estonian Research Council grant (PRG1901).

The authors thank Trond A. Abrahamsen and Vegard Lima for helpful discussions on the content of this paper.

REFERENCES

- 1. Trond A. Abrahamsen, Vegard Lima, and André Martiny, *Delta-points in Banach spaces* generated by adequate families, Illinois J. Math. 66 (2022), no. 3, 421–434. MR 4477423
- 2. Trond A. Abrahamsen, Vegard Lima, and Olav Nygaard, Remarks on diameter 2 properties, J. Convex Anal. 20 (2013), no. 2, 439–452. MR 3098474
- 3. Trond A. Abrahamsen, Olav Nygaard, and Märt Põldvere, New applications of extremely regular function spaces, Pacific J. Math. **301** (2019), no. 2, 385–394. MR 4023351
- 4. Ramón J. Aliaga, Extreme points in Lipschitz-free spaces over compact metric spaces, Mediterr. J. Math. 19 (2022), no. 1, Paper No. 32, 12. MR 4369296
- 5. Ramón J. Aliaga, Eva Pernecká, and Richard J. Smith, Convex integrals of molecules in Lipschitz-free spaces, J. Funct. Anal. 287 (2024), no. 8, Paper No. 110560, 43. MR 4771935
- 6. \ldots , de Leeuw representations of functionals on Lipschitz spaces, arXiv e-prints (2024), arXiv:2403.09546.
- 7. $____$, A solution to the extreme point problem and other applications of Choquet theory to Lipschitz-free spaces, arXiv e-prints (2024), arXiv:2412.04312.
- 8. Ramón J. Aliaga and Abraham Rueda Zoca, Points of differentiability of the norm in Lipschitz-free spaces, J. Math. Anal. Appl. 489 (2020), no. 2, 124171, 17. MR 4095811
- 9. Julio Becerra Guerrero, Ginés López-Pérez, and Abraham Rueda Zoca, Octahedral norms and convex combination of slices in Banach spaces, J. Funct. Anal. 266 (2014), no. 4, 2424–2435. MR 3150166
- 10. $_____\$ g slices versus big relatively weakly open subsets in Banach spaces, J. Math. Anal. Appl. 428 (2015), no. 2, 855–865. MR 3334951
- 11. Christian Cobollo, Daniel Isert, Ginés López-Pérez, Miguel Martín, Yoël Perreau, Alicia Quero, Andrés Quilis, Daniel L. Rodríguez-Vidanes, and Abraham Rueda Zoca, Banach spaces with small weakly open subsets of the unit ball and massive sets of Daugavet and Δ -points, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM 118 (2024), no. 3, Paper No. 96, 17. MR 4735972
- 12. Julio Becerra Guerrero, Ginés López-Pérez, and Abraham Rueda Zoca, Octahedrality in Lipschitz-free Banach spaces, Proc. Roy. Soc. Edinburgh Sect. A 148 (2018), no. 3, 447–460. MR 3826487
- 13. Rainis Haller, Jaan Kristjan Kaasik, and Andre Ostrak, Separating diameter two properties from their weak-star counterparts in spaces of Lipschitz functions, Studia Math. 280 (2025), no. 1, 87–102. MR 4849357
- 14. Rainis Haller, Johann Langemets, and Märt Põldvere, On duality of diameter 2 properties, J. Convex Anal. 22 (2015), no. 2, 465–483. MR 3346197
- 15. Rainis Haller, Andre Ostrak, and Märt Põldvere, *Diameter two properties for spaces of Lip*schitz functions, J. Funct. Anal. 287 (2024), no. 7, Paper No. 110524, 20. MR 4757422
- 16. Andre Ostrak, Characterisation of the weak-star symmetric strong diameter 2 property in Lipschitz spaces, J. Math. Anal. Appl. 483 (2020), no. 2, 123630, 10. MR 4026495
- 17. Antonín Procházka and Abraham Rueda Zoca, A characterisation of octahedrality in Lipschitz-free spaces, Ann. Inst. Fourier (Grenoble) 68 (2018), no. 2, 569–588. MR 3803112
- 18. Richard J. Smith, A Choquet theory of Lipschitz-free spaces, arXiv e-prints (2024), arXiv:2412.05177.
- 19. Cédric Villani, *Optimal transport*, vol. 338, Springer-Verlag, Berlin, 2009, Old and new. MR 2459454
- 20. Nik Weaver, Lipschitz algebras, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2018, Second edition of [MR 1832645]. MR 3792558

(R. Haller) Institute of Mathematics and Statistics, University of Tartu, Narva mnt 18, 51009, Tartu, Estonia

 $Email \;address: \; \verb|rainis.haller@ut.ee|$

(J. K. Kaasik) Institute of Mathematics and Statistics, University of Tartu, Narva mnt 18, 51009, Tartu, Estonia

 $\emph{Email address: }$ jaan.kristjan.kaasik@ut.ee

(A. Ostrak) Department of Mathematics, University of Agder, Postbox 422 4604 Kristiansand, Norway.

Email address: andre.ostrak@uia.no