# GENERALIZATIONS OF TWO HYPERGEOMETRIC SUMS RELATED TO CONJECTURES OF GUO

ARIJIT JANA AND LITON KARMAKAR

ABSTRACT. In 2021, the first author and Kalita obtained two general hypergeometric formulas for sums involving certain rising factorials to prove some supercongruence conjectures of Guo related to (B.2) and (C.2). In this paper, we further generalize those formulas by using the WZ-method and the Zeilberger algorithm respectively.

### 1. INTRODUCTION AND STATEMENT OF RESULTS

In 1914, Ramanujan [13] developed 17 infinite series representations of  $\frac{1}{\pi}$ , including the following identity

$$\sum_{n=0}^{\infty} \frac{4n+1}{(-64)^n} {\binom{2n}{n}}^3 = \frac{2}{\pi}.$$
(1.1)

Throughout the paper, let p be an odd prime and  $r \ge 1$  be integers. In [14], Van Hamme listed a number of conjectural supercongruences relating those Ramanujan's formulas for  $\frac{1}{\pi}$ , like (1.1), has a nice p-adic analogue

(B.2) 
$$\sum_{n=0}^{\frac{p-1}{2}} \frac{4n+1}{(-64)^n} {\binom{2n}{n}}^3 \equiv (-1)^{\frac{p-1}{2}} p \pmod{p^3}.$$
(1.2)

The supercongruence (1.2) was first proved by Mortenson [11] using a  $_{6}F_{5}$  hypergeometric identity, and later reconfirmed by Zudilin [16] using the WZ-method [12], and by Long [10] through certain hypergeometric series identities and evaluations. In [5], Guo proved the following generalization of (1.2) by providing a q-analogue of (1.2) using q-WZ method

$$\sum_{n=0}^{\frac{p'-1}{2}} \frac{4n+1}{(-64)^n} {\binom{2n}{n}}^3 \equiv p^r (-1)^{\frac{(p-1)r}{2}} \pmod{p^{r+2}}.$$
(1.3)

Employing the powerful WZ-method, Guo further gave the generalization of (1.2) as follows

$$\sum_{n=0}^{\frac{p-1}{2}} \frac{(4n+1)^3}{(-64)^n} {2n \choose n}^3 \equiv -3p(-1)^{\frac{(p-1)}{2}} \pmod{p^2}.$$
(1.4)

Following this, first author and Kalita [9] deduced the further generalization of (1.4) as

$$\sum_{n=0}^{\frac{p^{r}-1}{2}} \frac{(4n+1)^{3}}{(-64)^{n}} {\binom{2n}{n}}^{3} \equiv -3p^{r}(-1)^{\frac{(p-1)r}{2}} \pmod{p^{3}}.$$
(1.5)

Since  $(4n+1)^3 + 3(4n+1) = 4(4n+1)(4n^2+2n+1)$ , first author and Kalita [7] remarked from (1.3) and (1.5) that

$$\sum_{n=0}^{\frac{p^{r}-1}{2}} \frac{(4n+1)(4n^2+2n+1)}{(-64)^n} {2n \choose n}^3 \equiv 0 \pmod{p^{r+2}}.$$
(1.6)

2010 Mathematics Subject Classification. 11A07, 11D88, 33B15, 33C20.

Key words and phrases. Rising factorial, WZ-Method, Zeilberger's Alogorithm.

In this context, Guo [5, Conjecture 4.2] posed a stronger supercongruence conjecture for (1.6) as follows:

$$\begin{cases} \sum_{\substack{n=0\\p^{r}-1\\n=0}}^{\frac{p^{r}-1}{2}} \frac{(4n+1)(4n^{2}+2n+1)}{(-64)^{n}} {\binom{2n}{n}}^{3} \equiv p^{3r} \pmod{p^{3r+1}}, \\ \sum_{n=0}^{p^{r}-1} \frac{(4n+1)(4n^{2}+2n+1)}{(-64)^{n}} {\binom{2n}{n}}^{3} \equiv p^{3r} \pmod{p^{3r+1}}. \end{cases}$$
(1.7)

To prove the above Conjecture, first author and Kalita [9] gave a more general result using telescopic method as follows

$$\sum_{n=0}^{M} (-1)^n (2\ell n + 1)(\ell^2 n^2 + \ell n + 1) \frac{(\frac{1}{\ell})_n^3}{(1)_n^3} = (-1)^M \frac{(1 + \frac{1}{\ell})_M^3}{(1)_M^3},$$
(1.8)

where the rising factorial  $(a)_n$  is defined by

$$(a)_0 = 1, \ (a)_n = a(a+1)\cdots(a+n-1) \text{ for } n \ge 1.$$

The rising factorial for negative index [1, (1.2.27)] is defined as

$$(a)_{-n} = \frac{1}{(a-1)(a-2)\cdots(a-n)} = \frac{1}{(a-n)_n} = \frac{(-1)^n}{(1-a)_n}.$$

Motivated by the above result, we here give a further generalization of (1.8) by giving a new WZ-pair in the following theorem.

**Theorem 1.1.** Let  $\ell \geq 1$ ,  $s \geq 0$ ,  $M \geq 0$  with  $M \geq s$  be integers. Then

$$\sum_{n=s}^{M} (-1)^{n} (2\ell n+1) \left(\ell^{2} n^{2} + \ell n + 1 - \ell^{2} s^{2}\right) \frac{\left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_{n}}{(1)_{n+s} (1)_{n-s} (1)_{n}}$$
$$= (-1)^{M} \frac{\left(1 + \frac{1}{\ell}\right)_{M+s} \left(1 + \frac{1}{\ell}\right)_{M-s} \left(1 + \frac{1}{\ell}\right)_{M}}{(1)_{M+s} (1)_{M-s} (1)_{M}}.$$

In 2011, using hypergeometric identities, Long [10] proved that for prime  $p \ge 5$ 

(C.2) 
$$\sum_{n=0}^{\frac{p-1}{2}} \frac{4n+1}{256^n} {\binom{2n}{n}}^4 \equiv p \pmod{p^4}.$$
 (1.9)

Then, Guo and Wang [6] gave the following generalization of (1.9) by establishing a q-analogue

$$\sum_{n=0}^{\frac{p^r-1}{2}} \frac{4n+1}{256^n} {2n \choose n}^4 \equiv p^r \pmod{p^{r+3}}.$$
(1.10)

Using hypergeometric series identities and evaluations, first author and Kalita [8] have also given a proof of (1.10). In the same paper, first author and Kalita proved the following generalization of (1.10) as

$$\sum_{n=0}^{\frac{p^{r}-1}{2}} \frac{(4n+1)^{3}}{256^{n}} {2n \choose n}^{4} \equiv -p^{r} \pmod{p^{r+3}}.$$
(1.11)

In this context, Guo [5, Conjecture 4.4] has provided another supercongruence conjecture related to (1.10) and (1.11) as follows:

$$\begin{cases} \sum_{\substack{n=0\\p^r-1\\p^r-1}}^{\frac{p^r-1}{2}} \frac{(4n+1)(8n^2+4n+1)}{256^n} {\binom{2n}{n}}^4 \equiv p^{4r} \pmod{p^{4r+1}},\\ \sum_{n=0}^{p^r-1} \frac{(4n+1)(8n^2+4n+1)}{256^n} {\binom{2n}{n}}^4 \equiv p^{4r} \pmod{p^{4r+1}}. \end{cases}$$
(1.12)

To prove the above Conjecture, first author and Kalita [9] deduced a more general result in the closed form via telescopic method as follows

$$\sum_{n=0}^{M} (2\ell n+1)(2\ell^2 n^2 + 2\ell n+1)\frac{(\frac{1}{\ell})_n^4}{(1)_n^4} = \frac{(1+\frac{1}{\ell})_M^4}{(1)_M^4}.$$
(1.13)

Motivated by the above result, we here give a further generalization of (1.13) as

**Theorem 1.2.** Let  $\ell \geq 1$ ,  $s, M \geq 0$  with  $M \geq s$  be integers. Then

$$\sum_{n=s}^{M} (2\ell n+1) \left( 2\ell^2 n^2 + 2\ell n + 1 - \ell^2 s^2 \right) \frac{\left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_n^2}{(1)_{n+s} (1)_{n-s} (1)_n^2} = \frac{\left(1 + \frac{1}{\ell}\right)_{M+s} \left(1 + \frac{1}{\ell}\right)_{M-s} \left(1 + \frac{1}{\ell}\right)_M^2}{(1)_{M+s} (1)_{M-s} (1)_M^2}.$$

The rising factorial  $\frac{(\frac{1}{t})_n}{(1)_n}$  appears in a number of hypergeometric closed-sum formulas. In the literature, we have not found any general identities that resemble the sum containing  $\frac{(\frac{1}{t})_{n+s}(\frac{1}{t})_{n-s}}{(1)_{n+s}(1)_{n-s}}$ . The paper is organised as follows. In Section 2, we shall first give a new WZ-pair via the WZ-method designed by Wilf and Zeilberger [12], and then using this we prove Theorem 1.1. In Section 3, we prove Theorem 1.2 by using Zeilberger algorithm.

## 2. WZ-METHOD AND PROOF OF THEOREM 1.1

Wilf-Zeilberger method is based on preliminary human guess to finding a WZ-pair. Motivated by the work of [2, 3, 4, 16], we here employ the below hypergeometric function to deduce the following lemma

$$F(n,k) = (-1)^{n+k} (2\ell n+1) \frac{\left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_{n+k}}{(1)_{n+s} (1)_{n-s} (1)_{n-k} \left(\frac{1}{\ell}\right)_{k+s} \left(\frac{1}{\ell}\right)_{k-s}}$$

**Lemma 2.1.** Let  $\ell \ge 1$ ,  $n, s, k \ge 0$  with  $n \ge s$  and  $n \ge k$  be integers. Suppose

$$F(n,k) = (-1)^{n+k} (2\ell n+1) \frac{\left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_{n+k}}{(1)_{n+s} (1)_{n-s} (1)_{n-k} \left(\frac{1}{\ell}\right)_{k+s} \left(\frac{1}{\ell}\right)_{k-s}}$$

and

$$G(n,k) = (-1)^{n+k} \ell \frac{\left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_{n+k-1}}{(1)_{n+s-1} (1)_{n-s-1} (1)_{n-k} \left(\frac{1}{\ell}\right)_{k+s} \left(\frac{1}{\ell}\right)_{k-s}}$$

where  $1/(1)_m = 0$  for m = -1, -2, .... Then

$$F(n,k-1) - F(n,k) = G(n+1,k) - G(n,k).$$
(2.1)

Proof. We have

$$\frac{F(n,k-1)}{F(n,k)} = -\frac{\left(k+s+\frac{1-\ell}{\ell}\right)\left(k-s+\frac{1-\ell}{\ell}\right)}{\left(n+k+\frac{1-\ell}{\ell}\right)\left(n-k+1\right)},$$
$$\frac{G(n+1,k)}{F(n,k)} = -\frac{\ell\cdot\left(n+\frac{1}{\ell}+s\right)\left(n+\frac{1}{\ell}-s\right)}{\left(2\ell n+1\right)\left(n-k+1\right)},$$

3

and

$$\frac{G(n,k)}{F(n,k)} = \frac{\ell \cdot \left(n^2 - s^2\right)}{\left(2\ell n + 1\right)\left(n + k + \frac{1-\ell}{\ell}\right)}.$$

Noting that

$$\frac{F(n,k-1)}{F(n,k)} - 1 = -\left\{\frac{k^2 + 2k\left(\frac{1-\ell}{\ell}\right) + \left(\frac{1-\ell}{\ell}\right)^2 - s^2 + \left(n+k+\frac{1-\ell}{\ell}\right)(n-k+1)}{\left(n+k+\frac{1-\ell}{\ell}\right)(n-k+1)}\right\},\$$

and

$$\frac{G(n+1,k)}{F(n,k)} - \frac{G(n,k)}{F(n,k)} = -\ell \cdot \left\{ \frac{\left(n^2 + \frac{2n}{\ell} + \frac{1}{\ell^2} - s^2\right)\left(n+k+\frac{1-\ell}{\ell}\right) + \left(n^2 - s^2\right)\left(n-k+1\right)}{(2\ell n+1)\left(n+k+\frac{1-\ell}{\ell}\right)\left(n-k+1\right)} \right\}.$$

It is easy to obtain that

$$\frac{F(n,k-1)}{F(n,k)} - 1 = \frac{G(n+1,k)}{F(n,k)} - \frac{G(n,k)}{F(n,k)},$$

and hence

$$F(n, k - 1) - F(n, k) = G(n + 1, k) - G(n, k).$$

Г		
L		
L		

Proof of theorem 1.1. Multiplying both sides of (2.1) by  $\ell^2 \left(\frac{1}{\ell}\right)_{k+s} \left(\frac{1}{\ell}\right)_{k-s}$ , we get

$$\ell^2 \left(\frac{1}{\ell}\right)_{k+s} \left(\frac{1}{\ell}\right)_{k-s} \left\{F(n,k-1) - F(n,k)\right\} = \ell^2 \left(\frac{1}{\ell}\right)_{k+s} \left(\frac{1}{\ell}\right)_{k-s} \left\{G(n+1,k) - G(n,k)\right\}.$$

Substituting k = 1 and then taking sum on both sides with n ranging from s to M, we obtain

$$\sum_{n=s}^{M} \ell^2 \left(\frac{1}{\ell}\right)_{1+s} \left(\frac{1}{\ell}\right)_{1-s} \left\{F(n,0) - F(n,1)\right\} = \ell^2 \left(\frac{1}{\ell}\right)_{1+s} \left(\frac{1}{\ell}\right)_{1-s} \left\{G(M+1,1) - G(s,1)\right\}.$$

We have

$$\begin{split} &\sum_{n=s}^{M} \ell^{2} \left(\frac{1}{\ell}\right)_{1+s} \left(\frac{1}{\ell}\right)_{1-s} \left\{F(n,0) - F(n,1)\right\} \\ &= \sum_{n=s}^{M} \ell^{2} \left(\frac{1}{\ell}\right)_{1+s} \left(\frac{1}{\ell}\right)_{1-s} \left\{\frac{(-1)^{n} (2\ell n+1) \left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_{n}}{(1)_{n+s} (1)_{n-s} (1)_{n} \left(\frac{1}{\ell}\right)_{s} \left(\frac{1}{\ell}\right)_{-s}} + \frac{(-1)^{n} (2\ell n+1) \left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_{n+1}}{(1)_{n+s} (1)_{n-s} (1)_{n-1} \left(\frac{1}{\ell}\right)_{n-s}}\right\} \\ &= \sum_{n=s}^{M} \frac{(-1)^{n} (2\ell n+1) \ell^{2} \left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_{n}}{(1)_{n+s} (1)_{n-s} (1)_{n}} \left\{\frac{\left(\frac{1}{\ell}\right)_{1+s} \left(\frac{1}{\ell}\right)_{1-s}}{\left(\frac{1}{\ell}\right)_{s} \left(\frac{1}{\ell}\right)_{-s}} + \frac{\left(\frac{1}{\ell}\right)_{n+1} (1)_{n}}{(1)_{n-1} \left(\frac{1}{\ell}\right)_{n}}\right\} \\ &= \sum_{n=s}^{M} \frac{(-1)^{n} (2\ell n+1) \ell^{2} \left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_{n}}{(1)_{n+s} (1)_{n-s} (1)_{n}} \left\{\left(\frac{1}{\ell}+s\right) \left(\frac{1}{\ell}-s\right) + n \left(\frac{1}{\ell}+n\right)\right\} \\ &= \sum_{n=s}^{M} (-1)^{n} (2\ell n+1) \left(\ell^{2} n^{2} + \ell n+1 - \ell^{2} s^{2}\right) \frac{\left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_{n}}{(1)_{n+s} (1)_{n-s} (1)_{n}}. \\ &\text{Since } \frac{1}{(1)_{-1}} = 0, \text{ we have } G(s,1) = 0. \text{ As a result,} \end{split}$$

$$\ell^{2} \left(\frac{1}{\ell}\right)_{1+s} \left(\frac{1}{\ell}\right)_{1-s} \left\{ G(M+1,1) - G(s,1) \right\} = (-1)^{M} \ell^{3} \frac{\left(\frac{1}{\ell}\right)_{M+s+1} \left(\frac{1}{\ell}\right)_{M-s+1} \left(\frac{1}{\ell}\right)_{M+1}}{(1)_{M+s} (1)_{M-s} (1)_{M}} = (-1)^{M} \frac{\left(1 + \frac{1}{\ell}\right)_{M+s} \left(1 + \frac{1}{\ell}\right)_{M-s} \left(1 + \frac{1}{\ell}\right)_{M}}{(1)_{M+s} (1)_{M-s} (1)_{M}}.$$

This completes the proof.

4

#### 3. Zeilberger Algorithm and Proof of Theorem 1.2

For integers  $n \ge k \ge 0$ , suppose F(n,k) is a hypergeometric function in n and k. Employing Zeilberger's algorithm, one can find another hypergeometric function G(n,k), as well as polynomials p(k) and q(k) such that

$$p(k)F(n, k-1) - q(k)F(n, k) = G(n+1, k) - G(n, k).$$

However the process is not so obvious always. We here use the Zeilberger's algorithm for the function

$$F(n,k) = (-1)^k (2\ell n + 1) \frac{\left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_n \left(\frac{1}{\ell}\right)_{n+k}}{(1)_{n+s} (1)_{n-s} (1)_n (1)_{n-k} \left(\frac{1}{\ell}\right)_{k+s} \left(\frac{1}{\ell}\right)_{k-s}}.$$

Our motivation of taking the above hypergeometric function is based on WZ-pairs of [2, 3, 4, 15].

**Lemma 3.1.** Let  $\ell \ge 1$ ,  $n, s, k \ge 0$  with  $n \ge s$  and  $n \ge k$  be integers. Suppose

$$F(n,k) = (-1)^k (2\ell n + 1) \frac{\left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_n \left(\frac{1}{\ell}\right)_{n+k}}{(1)_{n+s} (1)_{n-s} (1)_n (1)_{n-k} \left(\frac{1}{\ell}\right)_{k+s} \left(\frac{1}{\ell}\right)_{k-s}}$$

and

$$G(n,k) = (-1)^{k-1} \ell^2 \frac{\left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_n \left(\frac{1}{\ell}\right)_{n+k-1}}{(1)_{n+s-1} (1)_{n-s-1} (1)_{n-1} (1)_{n-k} \left(\frac{1}{\ell}\right)_{k+s} \left(\frac{1}{\ell}\right)_{k-s}}$$

where  $1/(1)_m = 0$  for  $m = -1, -2, \ldots$ . Then

$$(\ell k - \ell + 1) F(n, k - 1) - (\ell k - \ell + 2) F(n, k) = G(n + 1, k) - G(n, k).$$
(3.1)

Proof. We have

$$\frac{F(n,k-1)}{F(n,k)} = -\frac{\left(k+s+\frac{1-\ell}{\ell}\right)\left(k-s+\frac{1-\ell}{\ell}\right)}{\left(n+k+\frac{1-\ell}{\ell}\right)\left(n-k+1\right)},$$
$$\frac{G(n+1,k)}{F(n,k)} = -\frac{\ell^2 \cdot \left(n+\frac{1}{\ell}+s\right)\left(n+\frac{1}{\ell}-s\right)\left(\frac{1}{\ell}+n\right)}{(2\ell n+1)\left(n-k+1\right)}$$

and

$$\frac{G(n,k)}{F(n,k)} = -\frac{\ell^2 \cdot \left(n^3 - ns^2\right)}{\left(2\ell n + 1\right)\left(n + k + \frac{1-\ell}{\ell}\right)}$$

Noting that

$$\left(\ell k - \ell + 1\right) \frac{F(n,k-1)}{F(n,k)} - \left(\ell k - \ell + 2\right) = -\left\{\frac{\left(k + s + \frac{1-\ell}{\ell}\right)\left(k - s + \frac{1-\ell}{\ell}\right)\left(\ell k - \ell + 1\right) + \left(n + k + \frac{1-\ell}{\ell}\right)\left(n - k + 1\right)\left(\ell k - \ell + 2\right)}{\left(n + k + \frac{1-\ell}{\ell}\right)\left(n - k + 1\right)}\right\}$$

and

$$\frac{G(n+1,k)}{F(n,k)} - \frac{G(n,k)}{F(n,k)} = -\ell^2 \cdot \left\{ \frac{\left(n^2 + \frac{2n}{\ell} + \frac{1}{\ell^2} - s^2\right)\left(n+k+\frac{1-\ell}{\ell}\right)\left(\frac{1}{\ell} + n\right) - \left(n^3 - ns^2\right)\left(n-k+1\right)}{(2\ell n+1)\left(n+k+\frac{1-\ell}{\ell}\right)\left(n-k+1\right)} \right\}.$$

It is easy to obtain that

$$\left(\ell k - \ell + 1\right) \frac{F(n,k-1)}{F(n,k)} - \left(\ell k - \ell + 2\right) = \frac{G(n+1,k)}{F(n,k)} - \frac{G(n,k)}{F(n,k)}$$

and hence

$$(\ell k - \ell + 1) F(n, k - 1) - (\ell k - \ell + 2) F(n, k) = G(n + 1, k) - G(n, k).$$

5

Proof of theorem 1.2. Multiplying both sides of (3.1) by  $\ell^2 \left(\frac{1}{\ell}\right)_{k+s} \left(\frac{1}{\ell}\right)_{k-s}$ , we get

$$\ell^{2}\left(\frac{1}{\ell}\right)_{k+s}\left(\frac{1}{\ell}\right)_{k-s}\left\{\left(\ell k - \ell + 1\right)F(n, k-1) - \left(\ell k - \ell + 2\right)F(n, k)\right\}$$
$$= \ell^{2}\left(\frac{1}{\ell}\right)_{k+s}\left(\frac{1}{\ell}\right)_{k-s}\left\{G(n+1, k) - G(n, k)\right\}.$$

Substituting k = 1 and then taking sum on both sides with n ranging from s to M, we obtain

$$\sum_{n=s}^{M} \ell^2 \left(\frac{1}{\ell}\right)_{1+s} \left(\frac{1}{\ell}\right)_{1-s} \left\{F(n,0) - 2F(n,1)\right\} = \ell^2 \left(\frac{1}{\ell}\right)_{1+s} \left(\frac{1}{\ell}\right)_{1-s} \left\{G(M+1,1) - G(s,1)\right\}.$$

One can easily find that

$$\sum_{n=s}^{M} \ell^2 \left(\frac{1}{\ell}\right)_{1+s} \left(\frac{1}{\ell}\right)_{1-s} \left\{F(n,0) - 2F(n,1)\right\} = \sum_{n=s}^{M} \left(2\ell n + 1\right) \left(2\ell^2 n^2 + 2\ell n + 1 - \ell^2 s^2\right) \frac{\left(\frac{1}{\ell}\right)_{n+s} \left(\frac{1}{\ell}\right)_{n-s} \left(\frac{1}{\ell}\right)_n^2}{\left(1\right)_{n+s} \left(1\right)_{n-s} \left(1\right)_n^2}$$

Again,

. .

$$\ell^{2} \left(\frac{1}{\ell}\right)_{1+s} \left(\frac{1}{\ell}\right)_{1-s} \left\{ G(M+1,1) - G(s,1) \right\} = \ell^{4} \frac{\left(\frac{1}{\ell}\right)_{M+s+1} \left(\frac{1}{\ell}\right)_{M-s+1} \left(\frac{1}{\ell}\right)_{M+1}^{2}}{(1)_{M+s} (1)_{M-s} (1)_{M}^{2}} \\ = \frac{\left(1 + \frac{1}{\ell}\right)_{M+s} \left(1 + \frac{1}{\ell}\right)_{M-s} \left(1 + \frac{1}{\ell}\right)_{M}^{2}}{(1)_{M+s} (1)_{M-s} (1)_{M}^{2}}.$$

This completes the proof.

#### 4. Acknowledgement

The second author has been supported by the institute doctoral fellowship of National Institute of Technology, Silchar.

#### References

- G. Gasper and M. Rahman, *Basic Hypergeometric series*, Encyclopedia of Mathematics and its Applications. vol. 96, 2nd edn, Cambridge University Press, Cambridge (2004).
- [2] S. B. Ekhad and D. Zeilberger, A WZ proof of Ramanujan's formula for π, Geometry, Analysis, and Mechanics, J.M. Rassias (ed.), World Scientific, Singapore (1994), 107–108.
- [3] J. Guillera, Some binomial series obtained by the WZ-method, Adv. Appl. Math. 29 (2002), no. 4, 599–603.
- [4] J. Guillera, Generators of some Ramanujan formulas, Ramanujan J. 11 (2006), no. 1, 41–48.
- [5] V. J. W. Guo, Some generalizations of a supercongruence of van Hamme, Integral Transforms Spec. Funct. 28 (2017), no. 12, 888-899.
- [6] V. J. W. Guo and S.-D. Wang, Some congruences involving fourth powers of central q-binomial coefficients, Proc. Roy. Soc. Edinburgh Sect. A 150 (2020), 1127-1138.
- [7] A. Jana and G. Kalita, Supercongruences for sums involving rising factorial (<sup>1</sup>/<sub>ℓ</sub>)<sup>3</sup><sub>k</sub>, Integral Transforms Spec. Funct. 30 (2019), no. 9, 683–692.
- [8] A. Jana and G. Kalita, Supercongruences for sums involving fourth power of some rising factorials, Proc Math Sci 130 (2020), Art. 59, 13pp.
- [9] A. Jana and G. Kalita, On Some Conjectural Supercongruences for Sums Involving Certain Rising Factorials, Results Math. 76 (2021), Art. 155, 14pp.
- [10] L. Long, Hypergeometric evaluation identities and supercongruences, Pacific J. Math. 249 (2011), no. 2, 405-418.
- [11] E. Mortenson, A p-adic supercongruence conjecture of vanHamme, Proc. Amer. Math. Soc. 136 (2008), no. 12, 4321-4328.
- [12] M. Petkovŝek, H. S. Wilf, and D. Zeilberger, A = B, A. K. Peters, Ltd., Wellesley, Mass. (1996).
- [13] S. Ramanujan, Modular equations and approximations to π, Quart. J. Math. 45, (1914), 350-372. In Collected papers of Srinivasa Ramanujan, pages 23–39. AMS Chelsea Publ., Providence, RI, 2000.
- [14] L. VanHamme, Some conjectures concerning partial sums of generalized hypergeometric series, p-adic functional analysis (Nijmegen, 1996), Lecture Notes in Pure and Appl. Math. 192 (1997), 223–236.
- [15] S.-D. Wang, Some supercongruences involving  $\binom{2k}{k}^4$ , J. Differ. Equ. Appl., **24** (2018), no. 9, 1375–1383.
- [16] W. Zudilin, Ramanujan-type supercongruences, J. Number Theory 129 (2009), no. 8, 1848-1857.

7

DEPARTMENT OF MATHEMATICS, NATIONAL INSTITUTE OF TECHNOLOGY, SILCHAR, 788010, INDIA *Email address:* jana94arijit@gmail.com

DEPARTMENT OF MATHEMATICS, NATIONAL INSTITUTE OF TECHNOLOGY, SILCHAR, 788010, INDIA *Email address:* litonofficial8638@gmail.com