# 2D HQFTS AND FROBENIUS  $(\mathcal{G}, \mathcal{V})$ -CATEGORIES

PAUL GROSSKOPF

Abstract. Homotopy Quantum Field Theories as variants of Topological Quantum Field Theories are described by functors from some cobordism category, enriched with homotopical data, to a symmetric monoidal category  $\mathcal V$ . A new notion of HQFTs is introduced using target pairs of spaces  $(X, Y)$  acounting for basepoints being sent to points in Y. Such  $(X, Y)$ -HQFTs are classified in dimension 1 by dualizable representations of  $\mathcal{G} := \Pi_1(X, Y)$ , the relative fundamental groupoid. For dimension 2, the notion of crossed loop Frobenius  $(G, V)$ categories is introduced, generalizing crossed Frobenius G-algebras, where G is only a group. After stating generalities of these multi-object generalizations, a classification theorem of 2-dimensional  $(X, Y)$ -HQFTs via crossed loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -categories is proven.

### CONTENTS



### 1. INTRODUCTION

<span id="page-0-0"></span>Homotopy Quantum Field Theories were introduced by Turaev in [\[10\]](#page-30-1) and generalize the notion of Topological Quantum Field Theories. These can be seen as invariants of manifolds, that behave functorially with respect to gluings of manifolds along the boundary. One therefore considers the category  $\text{Cob}_{d+1}$ , which has d-dimensional closed oriented manifolds as objects and compact oriented  $(d+1)$ -dimensional cobordisms as morphisms between them. This category further is endowed with a symmetric monoidal structure, given by the disjoint union. A TQFT can now be described as a symmetric monoidal functor from  $\textsf{Cob}_{d+1}$  to the category Vect<sub>K</sub> of vector spaces over a field  $K$  or more generally any symmetric monoidal

target category  $V$ . In dimension 2, TQFTs are fully classified by commutative Frobenius monoids in  $V$  [\[1\]](#page-30-2).

For HQFTs we fix a pointed space  $X$  and endow all manifolds with maps to this target space considered up to homotopy. One again can define a category  $\text{Cob}_{d+1}^{X}$  and define X-HQFTs to be symmetric monoidal functors  $\textsf{Cob}^{X}_{d+1} \to \textsf{Vect}_{\mathbb{K}}$ . In dimension 1, such X-HQFTs with target space X are classified by finite dimensional representations of  $\pi_1(X)$ . Futhermore, for contractible spaces, the theory of  $X$ -HQFTs specifies to the theory of TQFTs. If X is homotopically more complex we get more involved algebraic structures. Particularly, if  $X = K(G, 1)$  is the Eilenberg-MacLane space of a group G, then 2D X-HQFTs are fully classified by so-called crossed Frobenius G-algebras [\[10\]](#page-30-1). For homotopy 2-types work has been conducted in the cases  $X = K(A, 2)$ , for A an abelian group, [\[2\]](#page-30-3) as well as the general case where both  $\pi_1(X)$  and  $\pi_2(X)$  are non trivial [\[7,](#page-30-4)[9\]](#page-30-5). Furthermore, it is known that  $(d+1)$ dimensional X-HQFTs only depend on the homotopy  $(d+1)$ -type of the target space [\[8\]](#page-30-6). However, in this article I want to explore a more general notion of HQFTs with target

 $X$  a homotopy 1-type. Classically we consider cobordisms whose bounday components are pointed and pointed maps to  $X$ . We can loosen these restrictions and consider some subset  $Y \subset X$ , whose elements are possible images of the base points. This leads to the notion of  $(X, Y)$ -HQFTs as symmetric monoidal functors

$$
\mathsf{Cob}^{X,Y}_{d+1}\to \mathcal{V}.
$$

We can classify these easily in dimension 1 by dualizable representations of the relative fundamental groupoid  $\mathcal{G} = \Pi_1(X, Y)$  in  $\mathcal{V}$ , assuming some connectedness condition on the pair  $(X, Y)$ .

**Theorem A.** Let V be a (strict) symmetric monoidal category. Further let X be a homotopy 1-type,  $Y \subset X$ , such that for any connected component C of X we have  $C \cap Y \neq \emptyset$ , and denote  $\mathcal{G} = \Pi_1(X, Y)$ . The 1-dimensional  $(X, Y)$ -HQFTs

$$
\mathcal{Z}:\mathsf{Cob}^{X,Y}_1\to\mathcal{V}
$$

are fully classified by functors

 $\mathcal{G} \rightarrow \mathcal{V}$ ,

such that all objects in the image are dualizable.

Moreover, the main result of this article is a classification of  $2D(X, Y)$ -HQFTs via so-called crossed loop Frobenius  $(G, V)$ -categories. I will give two equivalent definitions of Frobenius  $(\mathcal{G}, \mathcal{V})$ -categories in analogue to the characterisation of Frobenius algebras via comultiplication and counit or via a non-degenerate pairing. I further introduce crossings and proof the main classification theorem of this article.

**Theorem B.** Let V be a (strict) symmetric monoidal category. Further, let X be a homotopy 1-type,  $Y \subset X$ , such that for any connected component C of X we have  $C \cap Y \neq \emptyset$ , and denote  $\mathcal{G} = \Pi_1(X, Y)$ . Then the 2-dimensional  $(X, Y)$ -HQFTs

$$
\mathcal{Z}:{\sf Cob}_2^{X,Y}\to\mathcal{V}
$$

are fully classified by crossed loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -categories.

As indicated by the nomenclature this theorem can be thought of a groupoid-version of Turaev's original result. Particularly, choosing  $V = \text{Vect}_{K}$  and  $Y = \{*\}$  one retrieves the classification of X-HQFTs via crossed Frobenius G-algebras.

<span id="page-2-0"></span>Acknowledgements. I want to thank my PhD supervisor Joost Vercruysse at Université Libre de Bruxelles, who gave me the intial idea to work on this project. Further, I want to thank Alexis Virelizier, Nils Carqueville and André Henriques for valuable discussions and challenging me to be more precise and careful at times. I was supported by a FRIA fellowship of FNRS FNRS (Fonds National de la Recherche Scientifique), grant number FC41285, while working on this project. Furthermore, while finishing this article I am supported by the FWA (Foundation Wiener-Anspach), recieving a postdoctoral fellowship.

# 2. Prelimaries and Conventions

<span id="page-2-1"></span>In this section we discuss elementary notions and conventions used in the rest of the article.

## <span id="page-2-2"></span>2.1. Monoidal categories.

**Definition 2.1.** A monoidal category  $(\mathcal{V}, \otimes, I, \alpha, \lambda, \rho)$  consists of a category  $\mathcal{V}$ , an object I of  $V$  called **monoidal unit** and bifunctor

$$
\otimes: \mathcal{V} \times \mathcal{V} \rightarrow \mathcal{V}
$$

called **tensor product**. Further, there are natural isomorphisms

$$
\alpha: \otimes(\mathcal{V}\times \otimes) \rightarrow \otimes(\otimes\times\mathcal{V}),
$$

called the associator, and

$$
\lambda : \otimes (I \times \mathcal{V}) \to \mathcal{V}, \qquad \rho : \otimes (\mathcal{V} \times I) \to \mathcal{V},
$$

called the **left and right unitor**, satisfying the *Pentagram rule* for all objects  $A, B, C, D$ of  $\mathcal{V}$ :



and

$$
A \otimes (I \otimes C) \xrightarrow{\alpha_{A,I,C}} (A \otimes I) \otimes C
$$
  

$$
A \otimes \lambda_C \downarrow \qquad \qquad \downarrow \rho_A \otimes C
$$
  

$$
A \otimes C \longrightarrow A \otimes C.
$$

If  $\alpha, \lambda, \rho$  are identities, the monoidal category is called **strict**.

**Definition 2.2.** Let  $(\mathcal{V}, \otimes, I, \alpha, \lambda, \rho)$  and  $(\mathcal{V}', \boxtimes, I', \alpha', \lambda', \rho')$  be two monoidal categories. A *monoidal functor* between V and V' is a triple  $(F, \phi, \phi_0)$ , where  $F: V \to V'$  is a functor,  $\phi: F \boxtimes F \to F(\mathcal{V} \otimes \mathcal{V})$  a natural transformation and  $\phi_0: I' \to FI$  a morphism in  $\mathcal{V}'$  making

the following diagrams commute:

$$
FA \boxtimes (FB \boxtimes FC) \xrightarrow{\alpha'_{FA,FB,FC}} (FA \boxtimes FB) \boxtimes FC
$$
  
\n
$$
FA \boxtimes F(B \otimes C) \qquad F(A \otimes B) \boxtimes FC
$$
  
\n
$$
\phi_{A,B \otimes C} \downarrow \phi_{A,B \otimes FC}
$$
  
\n
$$
F(A \otimes (B \otimes C)) \xrightarrow{F_{\alpha_{A,B,C}}} F((A \otimes B) \otimes C),
$$
  
\n
$$
FB \boxtimes I' \xrightarrow{\rho'_{FB}} FB \qquad I' \boxtimes FB \xrightarrow{\lambda'_{FB}} FB
$$
  
\n
$$
FB \boxtimes FI \xrightarrow{\rho'_{FB}} FB \qquad \phi_0 \boxtimes FB \downarrow \qquad F \wedge B
$$
  
\n
$$
FB \boxtimes FI \xrightarrow{\phi_{B,I}} F(B \otimes I), \qquad FI \boxtimes FB \xrightarrow{\lambda'_{FB}} F(I \otimes B).
$$

If  $\phi$  and  $\phi_0$  are identities, the monoidal functor is called **strict**.

**Definition 2.3.** Let  $(\mathcal{V}, \otimes, I)$  and  $(\mathcal{V}', \boxtimes, I')$  be two monoidal categories and  $(F, \phi, \phi_0)$  and  $(G, \psi, \psi_0)$  two monoidal functors  $\mathcal{V} \to \mathcal{V}'$ . A natural transformation  $\rho : F \Rightarrow G$  is called monoidal, if

$$
FA \boxtimes FB \xrightarrow{\rho_A \boxtimes \rho_B} GA \boxtimes GB \qquad I' \xrightarrow{\psi_0} GI
$$
  
\n
$$
\phi_{A,B} \downarrow \qquad \qquad \downarrow \psi_{A,B} \qquad \qquad \phi_0 \downarrow \qquad \qquad \rho_I
$$
  
\n
$$
F(A \otimes B) \xrightarrow{\rho_{A \otimes B}} G(A \otimes B), \qquad FI
$$

are commutative diagrams.

We define *braided* and *symmetric* monoidal categories. These give us the right setting to talk for example about any commutative and cocommutative phenomena.

**Definition 2.4.** A *braiding* of a monoidal category  $V$  is a natural isomorphims  $\sigma_{A,B}$ :  $A \otimes B \to B \otimes A$  such that  $\lambda_A \sigma_{I,A} = \rho_A$  for all objects A of V and the diagrams

$$
(A \otimes B) \otimes C \xrightarrow{\sigma_{A \otimes B,C}} C \otimes (A \otimes B) \qquad A \otimes (B \otimes C) \xrightarrow{\sigma_{A,B \otimes C}} (B \otimes C) \otimes A
$$
  
\n
$$
\alpha_{A,B,C} \uparrow \qquad \qquad \downarrow \alpha_{C,A,B} \qquad \qquad \alpha_{A,B,C}^{-1} \uparrow \qquad \qquad \downarrow \alpha_{B,C,A}^{-1}
$$
  
\n
$$
A \otimes (B \otimes C) \qquad (C \otimes A) \otimes B \qquad (A \otimes B) \otimes C, \qquad B \otimes (C \otimes A)
$$
  
\n
$$
A \otimes \sigma_{B,C} \uparrow \qquad \qquad \uparrow \sigma_{A,C} \otimes B \qquad \qquad \sigma_{A,B} \otimes C \uparrow \qquad \qquad \uparrow B \otimes \sigma_{A,C}
$$
  
\n
$$
A \otimes (C \otimes B) \xrightarrow[\alpha_{A,C,B}]{\sigma_{A,C} \otimes B} (A \otimes C) \otimes B, \qquad (B \otimes A) \otimes C \xrightarrow[\alpha_{B,A,C}^{-1}]} B \otimes (A \otimes C),
$$

commute for all objects  $A, B, C$  in  $\mathcal V$ .

A monoidal category is called **braided**, if it has a braiding  $\sigma$  and **symmetric**, if  $\sigma_{A,B}^{-1} = \sigma_{B,A}$ for all objects  $A, B$  in  $\mathcal V$ .

A monoidal functor  $(F, \phi, \phi_0)$  between two braided monoidal categories  $(\mathcal{V}, \otimes, I, \alpha, \lambda, \rho, \sigma)$ and  $(\mathcal{V}', \boxtimes, I', \alpha', \lambda', \rho', \sigma')$  is called **braided** if

$$
FA \boxtimes FB \xrightarrow{\sigma'} FB \boxtimes FA
$$
  
\n
$$
\phi_{A,B} \downarrow \qquad \qquad \phi_{B,A}
$$
  
\n
$$
F(A \otimes B) \xrightarrow{F\sigma} F(B \otimes A)
$$

are commutative diagrams for all  $A, B$  in  $\mathcal V$ . A symmetric monoidal functor is a braided monoidal functor between symmetric monoidal categories.

**Definition 2.5.** Let  $(\mathcal{V}, \otimes, I)$  and  $(\mathcal{V}', \boxtimes, I')$  be two symmetric monoidal categories. They are symmetric monoidally equivalent, if there exist symmetric monoidal functors  $(F, \phi, \phi_0)$ :  $V \to V'$  and  $(G, \psi, \psi_0) : V' \to V$  and monoidal natural isomorphisms  $\epsilon : FG \Rightarrow \mathsf{Id}_{V'}$  and  $\eta$  : Id  $\nu \Rightarrow GF$ .

**Definition 2.6.** Let  $(\mathcal{V}, \otimes, I)$  and  $(\mathcal{V}', \boxtimes, I')$  be two monoidal categories. The **symmetric** *monoidal functor category* is defined as the full subcategory  $\text{Fun}^{\otimes,s}(\mathcal{V},\mathcal{V}') \subseteq \text{Fun}^{\otimes}(\mathcal{V},\mathcal{V}')$ with objects symmetric monoidal functors and morphisms monoidal natural transformations.

<span id="page-4-1"></span>The following proposition is well know and is therefore only stated without proof.

**Proposition 2.7.** Let  $C$  be a symmetric monoidal category and  $V$ ,  $V'$  two symmetric monoidally equivalent categories. Then the monoidal functor categories  $\text{Fun}^{\otimes,s}(\mathcal{V},\mathcal{C})$  and  $\text{Fun}^{\otimes,s}(\mathcal{V}',\mathcal{C})$  are equivalent as categories.

In order to avoid writing down associators and unitors, we might consider MacLane's coherence theorem for symmetric monoidal categories:

**Theorem 2.8** ( [\[6,](#page-30-7) Thm.1. in XI.3.] as well as [\[3,](#page-30-8) Thm.XI.5.3.]). Any symmetric monoidal category  $\mathcal V$  is symmetric monoidally equivalent to a strict symmetric monoidal category  $\mathcal S$ .

Henceforth, we assume every symmetric monoidal category to be strict.

## <span id="page-4-0"></span>2.2. Dualizability and Rigid categories.

**Definition 2.9.** Let  $(\mathcal{V}, \otimes, I)$  be a monoidal category and X an object of  $\mathcal{V}$ . A right dual of X is an object  $X^*$  together with two morphisms  $ev_X : X \otimes X^* \to I$  and  $coev_X : I \to X^* \otimes X$ in  $V$  called *evaluation* and *coevaluation* satisfying the *zig-zag equations*:

$$
(\text{ev}_X \otimes X)(X \otimes \text{coev}_X) = X,
$$
  

$$
(X^* \otimes \text{ev}_X)(\text{coev}_X \otimes X^*) = X^*.
$$

X is called right dualizable, if there exist a right dual of X and V is called right rigid, if all objects is right dualizable. Similarly, given an object X, its **left dual**  $*X$  can be defined via ev<sup>X</sup>: \* $X \otimes X \to I$ , coev<sup>X</sup>:  $I \to X \otimes^* X$  satisfying the analogous equations. X is called left dualizable, if there exist a left dual of X and V is called left rigid, if all objects are left dualizable.

Duals are not unique, however they are unique up to unique isomorphisms in the following sense:

<span id="page-5-2"></span>**Proposition 2.10.** Let  $V$  be a monoidal category and  $X$  an object in  $V$ . Let  $(X^*, \text{ev}_X, \text{coev}_X)$ and  $(X', \text{ev}_X', \text{coev}_X')$  be two right duals of X, then there exists a unique isomorphism  $\phi: X^* \to Y$ X' such that  $ev'_X(X \otimes \phi) = ev_X$  and  $coev'_X = (\phi \otimes X)coev_X$ .

Remark 2.11. Duals are also compatible with the tensor product: given  $X, Y \in V$  with right duals  $X^*, Y^*,$  then  $(X \otimes Y)$  is right dualizable with right dual  $(Y^* \otimes X^*)$  via

$$
\begin{array}{rcl}\n\mathsf{ev}_{X\otimes Y} & := & \mathsf{ev}_X(X\otimes \mathsf{ev}_Y\otimes X^*) : X\otimes Y\otimes Y^*\otimes X^*\to I, \\
\mathsf{coev}_{X\otimes Y} & := & (Y^*\otimes \mathsf{coev}_X\otimes Y)\mathsf{coev}_Y : I\to Y^*\otimes X^*\otimes X\otimes Y.\n\end{array}
$$

Furthermore, in symmetric monoidal categories one need not to differentiate between left and right duals, since any right dual  $(X^*, ev_X, cov_X)$  of X is also a left dual via  $ev_X \sigma$  and  $\sigma$ coev $_X$ .

**Definition 2.12.** Let V be a symmetric monoidal category. X is called *dualizable* if has a left/right dual.  $V$  is called **rigid**, if every object is dualizable.

Given a morphisms  $f : A \to B$  between dualizable objects A and B we consider the morphisms  $f^* : B^* \to A^*$  defined by

$$
f^* := (A^* \otimes \text{ev}_B)(A^* \otimes f \otimes B^*)(\text{coev}_A \otimes B^*).
$$
  

$$
\begin{array}{c}\nB^* \\
\hline\nf^* \\
A^* \\
A^* \\
\end{array} = \n\begin{array}{ccc}\nB^* \\
\hline\nf \\
A^* \\
\end{array}
$$

One easily checks that this morphism satisfies:

$$
(f^* \otimes B) \operatorname{coev}_B = (A^* \otimes f) \operatorname{coev}_A
$$
  
\n
$$
\operatorname{ev}_A(A \otimes f^*) = \operatorname{ev}_B(f \otimes B^*)
$$

<span id="page-5-1"></span>The following is again well known and therefore stated without proof.

**Proposition 2.13.** Let  $(V, \otimes, I)$ ,  $(W, \boxtimes, J)$  be two strict monoidal categories,  $(F, \phi, \phi_0)$ ,  $(G, \psi, \psi_0)$ :  $V \to W$  be two monoidal functors and  $\alpha : F \Rightarrow G$  a monoidal natural transformation. If V is right rigid, then  $\alpha$  is a natural isomorphism.

## <span id="page-5-0"></span>2.3. The relative fundamental groupoid.

**Definition 2.14.** Let  $(X, Y)$  be a pair of topological spaces. Define  $\Pi_1(X, Y)$  the **funda***mental groupoid* of  $(X, Y)$  as the following category:

- Objects are given by  $y \in Y$ .
- Given  $y, z \in Y$ , the Hom-sets are given by

$$
\Pi_1(X,Y)(y,z) := \{ [f] | f : [0,1] \to X, f(0) = y, f(1) = z \},
$$

where  $[f] = [q]$  iff they are f and g are homotopic relativ to the endpoints.

• The composition of  $[f]$  and  $[g]$  is given by  $[g] \circ [f] = [f \cdot g]$ , where

$$
(f \cdot g)(t) := \begin{cases} f(2t) & t \in [0, \frac{1}{2}] \\ g(2t - 1) & t \in [\frac{1}{2}, 1] \end{cases}
$$

- The units are given by the constant paths  $1_y$  for  $y \in Y$ .
- The inverse of  $[f]$  is given by  $[f]$ , where

$$
\bar{f}(t) := f(1-t)
$$

Denote the homsets  $\text{Hom}_{\Pi_1(X,Y)}(y, z) =: \Pi_1(y, z)$ , if there is no ambiguity about the pair  $(X, Y)$ .

One checks easily that the fundamental groupoid is in fact a groupoid, meaning that  $[f] \circ [\bar{f}] = 1_z$  and  $[\bar{f}] \circ [f] = 1_y$  for all  $f \in \Pi_1(y, z)$ . In the following we write  $\alpha : x \to y$ or  $\beta : y \to z$  for elements in the fundamental groupoid and  $\alpha\beta : x \to z$  for the composition  $\beta \circ \alpha \in \Pi_1(x,z)$ .

## 3. THE CATEGORY OF  $(X, Y)$ -COBORDISMS

<span id="page-6-0"></span>To provide a categorical definition of  $(X, Y)$ -HQFTs it is helpful to put homotopy conditions already in the definition of the source category and not as a condition on the functor. So throughout the following let X be a homotopy 1-type,  $Y \subset X$  and  $d \geq 0$  be a natural number. By manifold we will always mean a differentiable manifold and by closed we understand it to be compact and with empty boundary.

## <span id="page-6-1"></span>3.1.  $(X, Y)$ -cobordisms.

.

**Definition 3.1.** A topological space Z is **pointed** if every connected component of Z is provided with a base point. Denote the set of base points of  $Z$  by  $Z_{\bullet}$ .

**Definition 3.2.** A d-dimensional  $(X, Y)$ -manifold is a pair  $(M, [g])$ , where  $(M, M_{\bullet})$  is a pointed, oriented, d-dimensional manifold and  $[g] \in [(M, M_{\bullet}), (X, Y)]_{\bullet}$  a homotopy class of maps relative to the base points. Particularly,  $M$  is allowed to be the empty set and  $g$  the empty function. Denote it by  $\mathcal{O}_d$  to avoid ambiguities. g is called the *characteristic* map. An  $(X, Y)$ -homeomorphism of  $(X, Y)$ -manifolds between  $(M, [g])$  and  $(M', [g'])$  is an orientation preserving homeomorphism  $f : (M, M_{\bullet}) \to (M', M'_{\bullet})$  such that  $[g] = [g'f]$  in  $[(M, M_{\bullet}), (X, Y)]_{\bullet}.$ 

**Definition 3.3.** A  $(d+1)$ -dimensional  $(X, Y)$ -cobordism is a tuple  $(W, M_0, M_1, f)$ , together with a homotopy class of maps  $[g] \in [(W, \partial W), (X, Y)]_{\partial W}$  relative to the boundary  $\partial W$ . M<sub>0</sub> and  $M_1$  are disjoint, pointed, closed, oriented, d-dimensional  $(X, Y)$ -manifolds with characteristic maps  $[g_0] \in [(M_0, M_{0\bullet}), (X, Y)]_{\bullet}, [g_1] \in [(M_1, M_{1\bullet}), (X, Y)]_{\bullet}$  and W is a compact, oriented,  $(d+1)$ -dimensional manifold with  $f = (f_0 \coprod f_1) : \partial W \to -M_0 \coprod M_1$  an orientation preserving homeomorphisms, where  $-M_0$  denotes the manifold  $M_0$  with opposite orientation. Further for any representative  $g: M \to X$  we have that  $[g \circ f_0^{-1}] = [g_0]$  and  $[g \circ f_1^{-1}] = [g_1]$ in  $[(M_0, M_{0\bullet}), (X, Y)]_{\bullet}$  and  $[(M_0, M_{0\bullet}), (X, Y)]_{\bullet}$  respectively. An  $(X, Y)$ -homeomorphism of  $(X, Y)$ -cobordisms is a triple  $(F, \phi_0, \phi_1) : (W, M_0, M_1, f) \rightarrow (W', M'_0, M'_1, f')$ , where  $F: W \to W'$  is an orientation preserving homeomorphism and  $\phi_i: M_i \to M'_i$  are  $(X, Y)$ homeomorphisms of  $(X, Y)$ -manifolds for  $i \in \{0, 1\}$ , such that  $(\phi_0 \coprod \phi_1) \circ f \circ F|_{\partial W} = f'$  and  $[g] = [g'F]$  in  $[(W, \partial W), (X, Y)]$ .

<span id="page-7-0"></span>3.2. Glueing of  $(X, Y)$ -cobordisms. Since we work with a homotopy 1-type X glueing of  $(X, Y)$ -cobordisms poses no problems: Let  $(M_0, [g_0]), (M_1, [g_1])$  and  $(N, [h])$  be  $(X, Y)$ manifolds and fix representatives  $g_0 : M_0 \to X$  and  $g_1 : M_1 \to X$ . Let  $(W, M_0, N, f)$ and  $(W', N, M_1, f')$  be  $(X, Y)$ -cobordisms together with two representatives  $g: W \to X$ and  $g' : W' \to X$ . Since  $g \circ f_1^{-1}$  and  $g' \circ f'^{-1}$  are both representatives of the class  $[h] \in$  $[(N, N_{\bullet}), (X, Y)]_{\bullet}$  we have a homotopy  $T : N \times [0, 1] \rightarrow X$  such that  $T(n, 0) = (g \circ f_1^{-1})(n)$ ,  $T(n, 1) = (g' \circ (f'_0)^{-1})(n)$  for all  $n \in N$  and  $T(n_0, t) = y_0$  for base points  $n_0 \in N_\bullet$  and  $y_0 \in Y$ , hence we can glue W to  $N \times [0,1]$  from the left via  $\pi_0^{-1} f_1$  and glue W' from the right via  $f_0'^{-1}\pi_1$ , where  $\pi = \pi_0 \coprod \pi_1 : N \times \{0,1\} \to -N \coprod N$ . We define this new  $(X,Y)$ -cobordism by  $(W \circ W', M_0, M_1, (f_0, f'_1))$  and the homotopy class of  $g \circ g' := g \coprod_{\pi_0^{-1} f_1} T \coprod_{f_0'^{-1} \pi_1} g'.$  We call  $(W \circ W', [g \circ g'])$  the glueing of W and W'.

<span id="page-7-1"></span>3.3. The category of  $(X, Y)$ -cobordisms.  $(X, Y)$ -cobordisms form the following category  $\mathsf{Cob}^{X,Y}_{d+1}$ :

- Objects are oriented, *d*-dimensional compact  $(X, Y)$ -manifolds.
- Morphisms are homeomorphism classes of  $d+1$ -dimensional  $(X, Y)$ -cobordism.
- Composition is given by gluing of  $(X, Y)$ -cobordisms. Given  $(W, M_0, N, f), (W', N, M_1, f')$ their composition is defined to be  $(W \circ W', [g \circ g'])$ .
- Identities  $\mathsf{Id}_{(M,g)}$  on a given  $(M,g)$  an oriented, d-dimensional  $(X,Y)$ -manifold are given by the homeomorphism class of the  $(X, Y)$ -bordism  $(M \times [0, 1], M \times 0, M \times$  $1, \pi|_{M\times\{0,1\}}$  with homotopy class  $[g \circ \pi]$  the composition of g and projection on M. This might be called the *cylinder* over  $(M, g)$ .

Further, there is a monoidal structure on  $\mathsf{Cob}^{X,Y}_{d+1}$ :

- The Tensor product is given by the disjoint union. Given two  $(X, Y)$ -manifolds  $(M, [g]), (N, [g'])$  consider  $(M \coprod N, [g \coprod g'])$  and similarly for the disjoint union of  $(X, Y)$ -bordisms, satisfying associativity constraints.
- The Monoidal unit is the empty  $(X, Y)$ -manifold  $\emptyset_d$ , satisfying unitality constraints.
- With the braiding  $\sigma_{MN} : (M \coprod N, [g \coprod g']) \to (N \coprod M, [g' \coprod g])$ , with  $\sigma_{NM} \sigma_{MN} =$  $\mathsf{Id}_{M \coprod N}$ ,  $\mathsf{Cob}_{d+1}^{X,Y}$  satisfies the conditions of a symmetry of monoidal categories.

This symmetric monoidal category is called the *category of*  $(d+1)$ -dimensional  $(X, Y)$ cobordisms.

# <span id="page-7-2"></span>**Proposition 3.4.** The category  $\text{Cob}_{d+1}^{X,Y}$  is rigid.

This is easily seen by considering the cylinders  $M \times [0, 1]$  over oriented d-dimensional compact  $(X, Y)$ -manifolds  $(M, [g])$  and interpreting them as morphisms  $C : \emptyset \to -M \coprod M$ as well as  $E: M \coprod -M \to \emptyset$ , where  $-M$  is the manifold M with opposite orientation, by setting their homotopy class  $[q \circ \pi]$  to be the composition of q and projection on M. E and C statisfy the zig-zag-equations exhibiting  $(-M, [g])$  as the dual of M. Furthermore, for any bordism  $(W, M_0, M_1, f_0 \coprod f_1)$  the cobordism  $-W = (-W, -M_1, -M_0, f_1 \coprod f_0)$  with the same characteristic map is given by

$$
W^* = (-M_0 \otimes \text{ev}_{M_1})(-M_0 \otimes W \otimes -M_1)(\text{coev}_{M_0} \otimes -M_1)
$$

.

# 4.  $(X, Y)$ -HQFTS AND THEIR CLASSIFICATION IN DIMENSION 1

<span id="page-8-0"></span>In this formalism d-dimensional  $(X, Y)$ -HQFTs can now be defined quite easily with a variable target category. Please note, that there is some ambiguity in the term target! In the theory of TQFT it refers to the domain of the functor, whereas in [\[11\]](#page-30-9) it refers to the space X. We make this nomenclature precise. Through this section let  $d > 0$ .

**Definition 4.1.** Let  $(\mathcal{V}, \otimes, I, \sigma)$  be a (strict) symmetric, monoidal category. A d-dimensional **Homotopy Quantum Field Theory** with **target pair of spaces**  $(X, Y)$ , or shorter  $(X, Y)$ -HQFT is a symmetric monoidal functor

$$
\mathcal{Z}: \mathsf{Cob}^{X,Y}_d \to \mathcal{V}.
$$

 $V$  is called its **target category**.

**Definition 4.2.** The symmetric monoidal functor category  $\text{Fun}^{\otimes,\text{s}}(\text{Cob}_d^{\text{X},\text{Y}},\mathcal{V})$  of d-dimensional  $(X, Y)$ -HQFTs and target space pair  $(X, Y)$  and target category V is denoted by  $\mathcal{Q}_d(X, Y, V)$ . A morphism between two  $(X, Y)$ -HQFTs  $\mathcal Z$  and  $\mathcal Z'$  is hence a monoidal natural transformation  $\rho : \mathcal{Z} \Rightarrow \mathcal{Z}'$ .

Remark 4.3. Since  $\mathsf{Cob}^{X,Y}_{d}$  is rigid, all natural transformations between  $(X, Y)$ -HQFTs are isomorphisms due to [2.13.](#page-5-1) This means  $\mathcal{Q}_d(X, Y, V)$  is a groupoid.

A natural question to ask is how  $(X, Y)$ -HQFT's behave under change of target space. Given an  $(X, Y)$ -HQFT  $\mathcal Z$  and a map  $f: X \to X'$ , then the HQFT can be pushed forward to an  $(X', f(Y))$ -HQFT via  $f_* : \mathcal{Q}_d(X, Y, V) \to \mathcal{Q}_d(X', Y, V)$  by composing the characteristic map with f. Similarly if we are interested in changing the target category and can always compose with a symmetric monoidal functor  $F: \mathcal{V} \to \mathcal{V}'$  to get a push forward  $F_*: \mathcal{Q}_d(X, Y, \mathcal{V}) \to$  $\mathcal{Q}_d(X, Y, \mathcal{V}')$ .

Although the definition of  $\mathsf{Cob}^{X,Y}_d$  and d-dimensional  $(X,Y)$ -HQFTs makes sense for any target pair with X a homotopy 1-type, we impose some light connectivity condition on  $(X, Y)$ , namely that every connected component of  $X$  has non-empty intersection with  $Y$ . The reason for that is, that we otherwise might have morphisms in  $\text{Cob}_{d}^{X,Y}$  from the empty manifold to the empty manifold, whose characteristic map maps into a component without any point of Y in it. Such a morphism cannot factor through any object different from the empty manifold. Therefore we might not represent is by composition of simple generators. It is therefore reasonable to assume that any connected component of  $X$  has a point in  $Y$ .

**Definition 4.4.** A pair of topological spaces  $(X, Y)$  is called **admissible**, if X is a homotopy 1-type and for any connected component C of X we have  $C \cap Y \neq \emptyset$ .

**Definition 4.5.** Let  $\mathcal G$  be a category and  $\mathcal V$  a monoidal category. A **dualizable representation** of G in V is a functor  $F : \mathcal{G} \to \mathcal{V}$  such that  $Fx$  is dualizable in V for all objects x of G. A morphism of dualizable representations is a natural transformation  $\rho : F \to F'$ such that

$$
\begin{array}{rcl} {\sf ev}_{F'x}(\rho_x\otimes\rho_{x^*})&=&{\sf ev}_{Fx}\\ (\rho_{x^*}\otimes\rho_x){\sf coev}_{Fx}&=&{\sf coev}_{F'x}\end{array}
$$

Denote the category of dualizable representations and morphisms of dualizable representations as  $\mathsf{Fun}^*(\mathcal{G},\overline{\mathcal{V}}).$ 

**Theorem 4.6.** Let  $(V, \otimes, I, \sigma)$  be a (strict) symmetric monoidal category. Let  $(X, Y)$  be an admissible pair of spaces and denote the relative fundamental groupoid of the pair  $(X, Y)$  by  $\mathcal{G} = \Pi_1(X, Y)$ . The category  $\mathcal{Q}_1(X, Y, \mathcal{V})$  of 1-dimensional  $(X, Y)$ -HQFTs is equivalent to the category  $\text{Fun}^*(\mathcal{G}, \mathcal{V})$  of dualizable representations.

*Proof.* The objects in  $\text{Cob}_{1}^{X,Y}$  are oriented, 0-dimensional, closed manifolds, hence consist of a finite, disjoint union of oriented points, together with a homotopy class of maps to Y fixing basepoints. One therefore can identify them with points in  $Y$  together with an orientation  $\in \{+, -\}.$  The duality between  $+y$  and  $-y$  is exhibited by  $ev_y : +y \coprod -y \to \emptyset$ and  $\cos y : \emptyset \to -y \coprod +y$ , whose characteristic maps are the constant maps on y. Consider  $y, z \in Y$  and a paths  $g : [0, 1] \to X$  from y to z, then there exists a morphism from  $+y$  to  $+z$ given by the interval and characteristic class [g], that we might also denote by [g]. We claim that any morphisms in  $\text{Cob}_1^{X,Y}$  can be glued from  $\sigma, \text{ev}_y, \text{coev}_y$  for any  $y \in Y$  and [g] for any  $g: y \to z \in \mathcal{G}$ .



First of all, any morphisms  $-y \to -z$  with characteristic map [g] can be represented as  $[\bar{g}]^* = (\mathsf{Id}_{-z} \otimes \mathsf{ev}_{+y})(\mathsf{Id}_{-z} \otimes [\bar{g}] \otimes \mathsf{Id}_{-y})(\mathsf{coev}_{+z} \otimes \mathsf{Id}_{-y}).$ 



Furthermore, any morphism  $\emptyset \to y \otimes -z$  with characteristic class [g] can be represented as  $(+y \otimes [\bar{g}])$ coev<sub>z</sub> and similarly for  $-y \otimes +z \to \emptyset$ . Pre- or postcomposing with  $\sigma$  yield all morphisms  $\emptyset \to -y \otimes +z$  and  $+y \otimes -z \to \emptyset$ . Therefore any morphisms with underlying 1-manifold an interval is a composition our generators. For any morphisms with underlying 1-manifold a circle, we can assume by connectivity that the caracteristic maps hits at least on point in Y, say y. Any circle therefore factors through  $ev_y\sigma(\text{Id}_{-y}\otimes [g])$ coev<sub>y</sub>, where  $g: y \to y$ in G. Since morphisms in  $\text{Cob}_1^{X,Y}$  are finite unions of interval and circles, all morphisms are generated by  $\sigma$ ,  $ev_y$ ,  $coev_y$  for any  $y \in Y$  and  $[g]$  for any  $g : y \to z \in \mathcal{G}$ .

Now given a 1-dimensional  $(X, Y)$ -HQFT  $\mathcal{Z}$ , the functor determines objects  $\mathcal{Z}(+y)$  in V with choice of duals  $\mathcal{Z}(-y)$ ,  $\mathcal{Z}(\mathsf{coev}_y)$ ,  $\mathcal{Z}(\mathsf{ev}_y)$  together morphisms

$$
\mathcal{Z}([g]): \mathcal{Z}(+y) \to \mathcal{Z}(+z)
$$

in V for  $g: y \to z$ . One checks easily that these form an action of the groupoid G on the objects  $\mathcal{Z}(y)$ .

Conversely, any such data is sufficient to define a 1-dimensional  $(X, Y)$ -HQFTs, since  $\mathcal Z$  is determined by the value on the morphisms  $ev_y$ , coev<sub>y</sub> and [g].

Let now  $\rho : \mathcal{Z} \Rightarrow \mathcal{Z}'$  be a natural isomorphism between two 1-dimensional  $(X, Y)$ -HQFTs, hence we get isomorphisms  $\rho_{\pm y} : \mathcal{Z}(\pm y) \to \mathcal{Z}'(\pm y)$  such that  $ev'_y(\rho_{+y} \otimes \rho_{-y}) = ev_y$  and

<span id="page-10-0"></span> $(\rho_{-y}\otimes\rho_{+y})$ coev<sub>y</sub> = coev'<sub>y</sub> and  $\rho_z\mathcal{Z}([g]) = \mathcal{Z}([g])\rho_y$ , hence this is the same data than a morphism of dualizable representations.

## 5. CROSSED LOOP FROBENIUS  $(\mathcal{G}, \mathcal{V})$ -CATEGORIES

In this section the main notion in order to classify 2-dimensional  $(X, Y)$ -HQFTs is defined. These structures should specify to G-graded Frobenius algebras

$$
L=\bigoplus_{\alpha\in G}L_\alpha
$$

in the case where  $\mathcal{G} = G$  a group and  $\mathcal{V} = \mathsf{Vect}_{K}$ . However, since we do not assume the existence of (co)-products in  $\mathcal{V}$ , consider the  $L_{\alpha}$  themselves as the objects. As one can see in the proof of the main theorem, this is actually much more in the spirit of HQFTs.

**Definition 5.1.** Let G be a groupoid and V a (strict) symmetric monoidal category. A  $(\mathcal{G}, \mathcal{V})$ -category is a collection of objects  $L_{\alpha} \in \mathcal{V}$  for all  $\alpha : x \to y$  in the groupoid  $\mathcal{G},$ together with two families of morphisms in  $V$ 

$$
m_{\alpha,\beta}: L_{\alpha}\otimes L_{\beta}\to L_{\alpha\beta} \qquad \alpha:x\to y, \beta:y\to z, x,y,z\in\mathcal{G},
$$

called *composition/multiplication*, and

$$
j_x: I \to L_{1_x} \qquad x \in \mathcal{G},
$$

called  $unit$ , satisfying associativity:

$$
L_{\alpha} \otimes L_{\beta} \otimes L_{\gamma} \xrightarrow{m_{\alpha,\beta} \otimes L_{\gamma}} L_{\alpha\beta} \otimes L_{\gamma}
$$
  

$$
L_{\alpha} \otimes m_{\beta,\gamma} \downarrow \qquad \qquad \downarrow m_{\alpha\beta,\gamma}
$$
  

$$
L_{\alpha} \otimes L_{\beta\gamma} \xrightarrow{m_{\alpha,\beta\gamma}} L_{\alpha\beta\gamma}
$$

for all composable triple of morphisms  $\alpha, \beta, \gamma$  in G and unitality:



for  $\alpha: x \to y$  in  $\mathcal{G}$ .

A  $(G, V)$ -functor is a collection of morphisms  $F_{\alpha}: L_{\alpha} \to L'_{\alpha}$  in V commuting with composition and unit:

$$
L_{\alpha} \otimes L_{\beta} \xrightarrow{F_{\alpha} \otimes F_{\beta}} L'_{\alpha} \otimes L'_{\beta} \qquad I \xrightarrow{j_x} L_{1_x}
$$
  
\n
$$
m_{\alpha,\beta} \downarrow \qquad \qquad m'_{\alpha,\beta} \downarrow \qquad \qquad j'_{F_x} \searrow \qquad \qquad k
$$
  
\n
$$
L_{\alpha\beta} \xrightarrow{F_{\alpha\beta}} L'_{\alpha\beta}, \qquad \qquad L'_{1_{F_x}}.
$$

Denote the category of  $(\mathcal{G}, \mathcal{V})$ -categories as  $(\mathcal{G}, \mathcal{V})$  – Cat.

Remark 5.2. Let X be a class and  $\mathcal{G} = X \times X$  the groupoid with objects X, that has exactely one arrow  $x \to y$  for all  $x, y \in X$ . Such  $(X \times X, V)$ -categories are precisely categories enriched over  $\mathcal V$ , hence, this notion generalizes enriched categories. Note that generally, for given objects x, y in  $\mathcal{G}$ , there are multiple object  $L_{\alpha}$  in  $\mathcal{V}$ , namely for all  $\alpha: x \to y$  in  $\mathcal{G}$ , and not just a collection  $L_{xy} = \text{Hom}_L(x, y) \in \mathcal{V}$  as for enriched categories. Further, note that the objects  $x, y$  play only a supporting role by clarifying, which elements can be composed.

**Example 5.3.** Given a field K and a groupoid G. Define  $L_{\alpha} = \mathbb{K}\langle l_{\alpha} \rangle$  to be the 1-dimensional vector spaces generated by  $l_{\alpha}$  and  $m_{\alpha,\beta}(l_{\alpha} \otimes l_{\beta}) := l_{\alpha\beta}$  and  $j_x(1) := l_{1_x}$ .  $(L, m, j)$  is a  $(\mathcal{G}, \mathsf{Vect}_{\mathbb{K}})$ -category in analogue to the group algebra for a group G. We may therefore call it **groupoid algebra** of G and denote it by  $L = \mathbb{K}[\mathcal{G}]$ .

**Definition 5.4.** Let G be a groupoid and  $(\mathcal{V}, \otimes, I, \sigma)$  a (strict) symmetric monoidal category. A  $(\mathcal{G}, \mathcal{V})$ -opcategory is a  $(\mathcal{G}, \mathcal{V}^{op})$ -category L.

Explicitely, L consists of a collection of objects  $L_{\alpha} \in V$  for all  $\alpha : x \to y$  in the groupoid  $\mathcal{G}$ , together with two families of morphisms in  $\mathcal{V}$ :

$$
\Delta_{\alpha,\beta}: L_{\alpha\beta} \to L_{\alpha} \otimes L_{\beta} \qquad \alpha: x \to y, \beta: y \to z, x, y, z \in \mathcal{G},
$$

called *cocomposition*/*comultiplication*, and

$$
\nu_x: L_{1_x} \to I \qquad x \in \mathcal{G},
$$

called  $\textit{counti}$ , satisfying coassociativity:

$$
L_{\alpha} \otimes L_{\beta} \otimes L_{\gamma} \xleftarrow{\Delta_{\alpha,\beta} \otimes L_{\gamma}} L_{\alpha\beta} \otimes L_{\gamma}
$$
  

$$
L_{\alpha} \otimes \Delta_{\beta,\gamma} \qquad \qquad \Delta_{\alpha,\beta\gamma} \qquad \Delta_{\alpha\beta,\gamma}
$$
  

$$
L_{\alpha} \otimes L_{\beta\gamma} \xleftarrow{\Delta_{\alpha,\beta\gamma}} L_{\alpha\beta\gamma}
$$

for all composable triples of morphisms  $\alpha, \beta, \gamma$  in G and counitality

$$
L_{\alpha} \simeq I \otimes L_{\alpha} \xleftarrow{\nu_x \otimes L_{\alpha}} L_{1_x} \otimes L_{\alpha} \qquad L_{\alpha} \simeq L_{\alpha} \otimes I \xleftarrow{L_{\alpha} \otimes \nu_y} L_{\alpha} \otimes L_{1_y}
$$
  

$$
L_{\alpha} \qquad L_{\alpha}, \qquad L_{\alpha} \qquad L_{\alpha} \qquad L_{\alpha}.
$$

for  $\alpha: x \to y$  in  $\mathcal{G}$ .

A  $(G, V)$ -op-functor is a collection  $F_{\alpha}: L_{\alpha} \to L'_{\alpha}$  of maps in V commuting with cocomposition and counit:

$$
L_{\alpha} \otimes L_{\beta} \xrightarrow{F_{\alpha} \otimes F_{\beta}} L'_{\alpha} \otimes L'_{\beta} \qquad I \xleftarrow{\nu_x} L_{1_x}
$$
  
\n
$$
\Delta_{\alpha,\beta} \qquad \qquad \Delta'_{\alpha,\beta} \qquad \qquad \Delta'_{\alpha,\beta} \qquad \qquad \nu'_{Fx} \qquad \qquad \downarrow F_{1_x}
$$
  
\n
$$
L_{\alpha\beta} \xrightarrow{F_{\alpha\beta}} L'_{\alpha\beta}, \qquad \qquad L'_{1_{Fx}}.
$$

Denote the category of  $(\mathcal{G}, \mathcal{V})$ -opcategories as  $(\mathcal{G}, \mathcal{V})$  – opCat.

Frobenius algebras can be defined by two equivalent definitions the existence of an inner product as done in [\[11\]](#page-30-9) or the categorical definition stating the existence of a compatible comultiplication. In the following generalizations of both are introduced and it is shown, that these are equivalent.

<span id="page-12-0"></span>**Definition 5.5.** Let G be a groupoid and  $(\mathcal{V}, \otimes, I, \sigma)$  a (strict) symmetric monoidal category. A **Frobenius**  $(G, V)$ -category is a 5-tuple consisting of a family  $(L_{\alpha})_{\alpha \in \mathcal{G}}$  of objects in V and four families of morphisms  $m, j, \Delta, \nu$  in  $\mathcal V$  such that:

- (FC1)  $(L, m, i)$  is an  $(G, V)$ -category.
- (FC2)  $(L, \Delta, \nu)$  is a  $(\mathcal{G}, \mathcal{V})$ -opcategory.
- $(FC3)$  The **Frobenius conditions** are satisfied:



A **Frobenius**  $(\mathcal{G}, \mathcal{V})$ -functor is a  $(\mathcal{G}, \mathcal{V})$ -functor as well as a  $(\mathcal{G}, \mathcal{V})$ -opfunctor. Denote the category of Frobenius  $(G, V)$ -categories and -functors by Frob  $^{\mathcal{G}}(\mathcal{V})$ . We call L symmetric, if for all  $\alpha : x \to y$ 

$$
\nu_x m_{\alpha,\alpha^{-1}} = \nu_y m_{\alpha^{-1},\alpha} \sigma : L_\alpha \otimes L_{\alpha^{-1}} \to I.
$$

<span id="page-12-1"></span>Equivalently we can define Frobenius  $(\mathcal{G}, \mathcal{V})$ -categories by defining an inner product:

**Definition 5.6.** Let L be a  $(\mathcal{G}, \mathcal{V})$ -category. A family of morphisms  $\eta_{\alpha}: L_{\alpha} \otimes L_{\alpha^{-1}} \to I$  is called an *inner product* or *evaluation* on  $L$  if:

(IP1) it is **non-degenerate**, that is, if there exists a family  $\text{coev}_{\alpha}: I \to L_{\alpha^{-1}} \otimes L_{\alpha}$  called coevaluation, such that

$$
(\eta_{\alpha} \otimes L_{\alpha})(L_{\alpha} \otimes \mathsf{coev}_{\alpha}) = L_{\alpha},
$$
  

$$
(L_{\alpha^{-1}} \otimes \eta_{\alpha})(\mathsf{coev}_{\alpha} \otimes L_{\alpha^{-1}}) = L_{\alpha^{-1}}
$$

and

(IP2) it is compatibility with the multiplication:

$$
L_{\alpha} \otimes L_{\beta} \otimes L_{\gamma} \xrightarrow{m_{\alpha,\beta} \otimes L_{\gamma}} L_{\alpha\beta} \otimes L_{\gamma}
$$
  

$$
L_{\alpha} \otimes m_{\beta,\gamma} \downarrow \qquad \qquad \downarrow n_{\alpha\beta}
$$
  

$$
L_{\alpha} \otimes L_{\beta\gamma} \xrightarrow{n_{\alpha}} I,
$$

where  $\gamma = (\alpha \beta)^{-1}$ , hence  $\alpha^{-1} = \beta \gamma$ . The inner product is called *symmetric* if

$$
\eta_{\alpha} = \eta_{\alpha^{-1}} \sigma : L_{\alpha} \otimes L_{\alpha^{-1}} \to I.
$$

A **Frobenius**  $(\mathcal{G}, \mathcal{V})$ -category is a  $(\mathcal{G}, \mathcal{V})$ -category L with an inner product. If the inner product is symmetric, L is called a **symmetric** Frobenius  $(\mathcal{G}, \mathcal{V})$ -category.

The first conditions states that  $L_{\alpha^{-1}}$  it the (right) dual of  $L_{\alpha}$  in V via  $\eta_{\alpha}$  and coev<sub> $\alpha$ </sub>. The second connects this duality with the  $(G, V)$ -category structure on.

**Example 5.7.** (1) If  $\mathcal{G} = 1$  is the trivial group, with one morphism then every  $(1, \mathcal{V})$ category is a Frobenius monoid in  $\mathcal V$ .

(2) If  $\mathcal{G} = G$  is a group and  $\mathcal{V} = \mathsf{Vect}_{\mathbb{K}}$ , then any  $(G, \mathsf{Vect}_{\mathbb{K}})$ -category L is equivalent to a Frobenius  $G$ -algebra  $L'$  as defined in [\[11\]](#page-30-9) given by

$$
L'=\bigoplus_{\alpha\in G}L_{\alpha},
$$

with  $x \cdot y := m_{\alpha,\beta}(x \otimes y)$  for  $x \in L_{\alpha}, y \in L_{\beta}$  with  $\alpha, \beta \in G$  and  $1_{L'} = j_{\epsilon}(1) \in L_{\epsilon}$  for  $\epsilon \in G$  the neutral element.

(3) The groupoid algebra  $L = \mathbb{K}[\mathcal{G}]$  has a  $(\mathcal{G}, \text{Vect}_{\mathbb{K}})$ -opcategory structure via  $\Delta_{\alpha,\beta}(l_{\alpha\beta}) =$  $l_{\alpha} \otimes l_{\beta}$  and  $\nu(l_{1_x}) = 1$ . Furthermore, the Frobenius conditions are satisfied and therefore  $L = \mathbb{K}[\mathcal{G}]$  is a Frobenius  $(\mathcal{G}, \mathcal{V})$ -category, that is symmetric.

Lemma 5.8. Given an inner product, the corresponding coevaluation is unique. Further a Frobenius  $(G, V)$ -category is symmetric, if and only if

 $\sigma$ coev $\alpha$  = coev $\alpha$ <sup>-1</sup>.

Proof. The first statement was shown in [2.10.](#page-5-2) The second statement follows from the fact that  $\eta_X\sigma$  and  $\sigma$ coev<sub>X</sub> satisfy the zig-zag-equations. The uniqueness of  $\eta$  and coev shows that  $\sigma$ coev<sub> $\alpha$ </sub> = coev<sub> $\alpha$ </sub>-1, if and only iff  $\eta_{\alpha}\sigma = \eta_{\alpha^{-1}}$ .

**Lemma 5.9.** Let L be a Frobenius  $(G, V)$ -category then

<span id="page-13-0"></span>
$$
(1) \t m_{\alpha,\beta} = (L_{\alpha\beta}\otimes \eta_{(\alpha\beta)^{-1}})(L_{\alpha\beta}\otimes L_{(\alpha\beta)^{-1}}\otimes m_{\alpha,\beta})(\text{coev}_{(\alpha\beta)^{-1}}\otimes L_{\alpha}\otimes L_{\beta})
$$

$$
(2) \qquad = (L_{\alpha\beta}\otimes\eta_{\beta^{-1}})(L_{\alpha\beta}\otimes m_{(\alpha\beta)^{-1},\alpha}\otimes L_{\beta})(\mathsf{coev}_{(\alpha\beta)^{-1}}\otimes L_{\alpha}\otimes L_{\beta})
$$

(3) 
$$
= (\eta_{\alpha\beta}\otimes L_{\alpha\beta})(m_{\alpha,\beta}\otimes L_{(\alpha\beta)^{-1}}\otimes L_{\alpha\beta})(L_{\alpha}\otimes L_{\beta}\otimes \text{coev}_{\alpha\beta})
$$

(4) 
$$
= (\eta_{\alpha} \otimes L_{\alpha\beta})(L_{\alpha} \otimes m_{\beta,(\alpha\beta)^{-1}} \otimes L_{\alpha\beta})(L_{\alpha} \otimes L_{\beta} \otimes \text{coev}_{\alpha\beta})
$$

Proof. This proof is an easy deduction of the axiom for the coevaluation. By commutativity of tensor product and composition we can push coev and  $\eta$  together in [\(1\)](#page-13-0) to get

 $(L_{\alpha\beta} \otimes \eta_{(\alpha\beta)^{-1}})(\text{coev}_{(\alpha\beta)^{-1}} \otimes L_{\alpha\beta})m_{\alpha,\beta} = m_{\alpha,\beta}.$ 

[\(2\)](#page-13-0) follows from [\(1\)](#page-13-0) by compability of the multiplication and inner product. [\(3\)](#page-13-0), [\(4\)](#page-13-0) can be deduced anagolously.

**Proposition 5.10.** Both definitions [\(5.5](#page-12-0)  $\mathcal{O}(5.6)$  of (symmetric) Frobenius  $(\mathcal{G}, \mathcal{V})$ -categories are equivalent.

*Proof.* Let L be a Frobenius  $(G, V)$ -category in the sense of [5.5.](#page-12-0) We will show that there is an inner product on L. For  $\alpha: x \to y$  define  $\eta_{\alpha} := \nu_x m_{\alpha,\alpha^{-1}}$ . The proof works in analogue to the one-object case, but making sure we respect codomains and domains of arrows.

- $\eta_{\alpha}$  is symmetric if and only if L is a symmetric Frobenius  $(\mathcal{G}, \mathcal{V})$ -category.
- Non-degeneracy condition: Define  $\operatorname{coev}_{\alpha} := \Delta_{\alpha^{-1},\alpha} j_y$ . Then

$$
(L_{\alpha^{-1}} \otimes \eta_{\alpha})(\operatorname{coev}_{\alpha} \otimes L_{\alpha^{-1}}) = (L_{\alpha^{-1}} \otimes \nu_x)(L_{\alpha} \otimes m_{\alpha,\alpha-1})(\Delta_{\alpha^{-1},\alpha} \otimes L_{\alpha})(j_y \otimes L_{\alpha})
$$
  
\n
$$
= (L_{\alpha^{-1}} \otimes \nu_x)\Delta_{\alpha^{-1},1_x} m_{1_y,\alpha^{-1}}(j_y \otimes L_{\alpha^{-1}}) = L_{\alpha^{-1}},
$$
  
\n
$$
(\eta_{\alpha} \otimes L_{\alpha})(L_{\alpha} \otimes \operatorname{coev}_{\alpha}) = (\nu_x \otimes L_{\alpha})(m_{\alpha,\alpha^{-1}} \otimes L_{\alpha})(L_{\alpha} \otimes \Delta_{\alpha^{-1},\alpha})(L_{\alpha} \otimes j_y)
$$
  
\n
$$
= (\nu_x \otimes L_{\alpha})\Delta_{1_x,\alpha} m_{\alpha,1_y}(L_{\alpha} \otimes j_y) = L_{\alpha},
$$

where we used the Frobenius conditions as well as (co)-unitality.

• For the compatibility with the multiplication we can look at the following commuting diagram:



The square commutes by associativity of the multiplication and the two triagles commute per definitionem.

Therefore  $\eta$  is an inner product on the  $(\mathcal{G}, \mathcal{V})$ -category L, making it is Frobenius in terms of [5.6.](#page-12-1) Further L is symmetric if the inner product is symmetric.

Now let L be a Frobenius  $(G, V)$ -category in the sense of [5.6.](#page-12-1) We need to construct a comultiplication and a counit. We define

$$
\Delta_{\alpha,\beta} := (L_{\alpha} \otimes m_{\alpha^{-1},\alpha\beta})(\mathsf{coev}_{\alpha^{-1}} \otimes L_{\alpha\beta}) : L_{\alpha\beta} \to L_{\alpha} \otimes L_{\beta}, \nu_x := \eta_{1_x}(j_x \otimes L_{1_x}) : L_{1_x} \to I.
$$

First note that  $\Delta_{\alpha,\beta} = (m_{\alpha\beta,\beta^{-1}} \otimes L_{\beta})(L_{\alpha\beta} \otimes \text{coev}_{\beta})$  by using [\(2\)](#page-13-0) and [\(3\)](#page-13-0):

$$
\begin{array}{rcl}\n\Delta_{\alpha,\beta} & = & (L_{\alpha} \otimes m_{\alpha^{-1},\alpha\beta})(\operatorname{coev}_{\alpha^{-1}} \otimes L_{\alpha\beta}) \\
& = & (L_{\alpha} \otimes \eta_{\beta} \otimes L_{\beta})(L_{\alpha} \otimes m_{\alpha^{-1},\alpha\beta} \otimes L_{\beta^{-1}} \otimes L_{\beta})(L_{\alpha} \otimes L_{\alpha^{-1}} \otimes L_{\alpha\beta} \otimes \operatorname{coev}_{\beta})(\operatorname{coev}_{\alpha^{-1}} \otimes L_{\alpha\beta}) \\
& = & (L_{\alpha} \otimes \eta_{\beta} \otimes L_{\beta})(L_{\alpha} \otimes m_{\alpha^{-1},\alpha\beta} \otimes L_{\beta^{-1}} \otimes L_{\beta})(\operatorname{coev}_{\alpha} \otimes L_{\alpha\beta} \otimes L_{\beta^{-1}} \otimes L_{\beta})(L_{\alpha\beta} \otimes \operatorname{coev}_{\beta}) \\
& = & (m_{\alpha\beta,\beta^{-1}} \otimes L_{\beta})(L_{\alpha\beta} \otimes \operatorname{coev}_{\beta}).\n\end{array}
$$

Using this identity, we show coassociativity:

$$
\begin{array}{rcl}\n(\Delta_{\alpha,\beta}\otimes L_{\gamma})\Delta_{\alpha\beta,\gamma} & = & (L_{\alpha}\otimes m_{\alpha^{-1},\alpha\beta}\otimes L_{\gamma})(\text{coev}_{\alpha^{-1}}\otimes L_{\alpha\beta}\otimes L_{\gamma})(m_{\alpha\beta\gamma,\gamma^{-1}}\otimes L_{\gamma})(L_{\alpha\beta\gamma}\otimes \text{coev}_{\gamma}) \\
 & = & (L_{\alpha}\otimes m_{\alpha^{-1},\alpha\beta}\otimes L_{\gamma})(L_{\alpha}\otimes L_{\alpha^{-1}}\otimes m_{\alpha\beta\gamma,\gamma^{-1}}\otimes L_{\gamma})(\text{coev}_{\alpha^{-1}}\otimes L_{\alpha\beta\gamma}\otimes \text{coev}_{\gamma}) \\
 & = & (L_{\alpha}\otimes m_{\beta\gamma,\gamma^{-1}}\otimes L_{\gamma})(L_{\alpha}\otimes m_{\alpha^{-1},\alpha\beta\gamma}\otimes L_{\gamma}^{-1}\otimes L_{\gamma})(\text{coev}_{\alpha^{-1}}\otimes L_{\alpha\beta\gamma}\otimes \text{coev}_{\gamma}) \\
 & = & (L_{\alpha}\otimes m_{\beta\gamma,\gamma^{-1}}\otimes L_{\gamma})(L_{\alpha}\otimes L_{\beta\gamma}\otimes \text{coev}_{\gamma})(L_{\alpha}\otimes m_{\alpha^{-1},\alpha\beta\gamma})(\text{coev}_{\alpha^{-1}}\otimes L_{\alpha\beta\gamma}) \\
 & = & (L_{\alpha}\otimes \Delta_{\beta,\gamma})\Delta_{\alpha,\beta\gamma},\n\end{array}
$$

and counitality:

$$
(\nu_x \otimes L_{\alpha})\Delta_{1_x,\alpha} = (\eta_{1_x,1_x} \otimes L_{\alpha})(j_x \otimes L_{1_x} \otimes L_{\alpha})(L_{1_x} \otimes m_{1_x,\alpha})(\text{coev}_{1_x} \otimes L_{\alpha})
$$
  
\n
$$
= m_{1_x,\alpha}(\eta_{1_x,1_x} \otimes L_{1_x} \otimes L_{\alpha})(L_{1_x} \otimes \text{coev}_{1_x} \otimes L_{\alpha})(j_x \otimes L_{\alpha})
$$
  
\n
$$
= m_{1_x,\alpha}(j_x \otimes L_{\alpha}) = L_{\alpha},
$$

and analogously for the right counitality.  $(L, \Delta, \nu)$  therefore forms a  $(\mathcal{G}, \mathcal{V})$ -opcategory. We finish by checking the Frobenius conditions:

$$
(L_{\alpha} \otimes m_{\beta,\gamma})(\Delta_{\alpha,\beta} \otimes L_{\gamma}) = (L_{\alpha} \otimes m_{\beta,\gamma})(L_{\alpha} \otimes m_{\alpha^{-1},\alpha\beta} \otimes L_{\gamma})(\text{coev}_{\alpha^{-1}} \otimes L_{\alpha\beta} \otimes L_{\gamma})
$$
  
\n
$$
= (L_{\alpha} \otimes m_{\alpha^{-1},\alpha\beta\gamma})(L_{\alpha} \otimes L_{\alpha^{-1}} \otimes m_{\alpha\beta,\gamma})(\text{coev}_{\alpha^{-1}} \otimes L_{\alpha\beta} \otimes L_{\gamma})
$$
  
\n
$$
= (L_{\alpha} \otimes m_{\alpha^{-1},\alpha\beta\gamma})(\text{coev}_{\alpha^{-1}} \otimes m_{\alpha\beta,\gamma}) = \Delta_{\alpha,\beta\gamma}m_{\alpha\beta,\gamma}
$$

$$
\begin{array}{rcl}\n(m_{\alpha,\beta}\otimes L_{\gamma})(L_{\alpha}\otimes \Delta_{\beta,\gamma}) & = & (m_{\alpha,\beta}\otimes L_{\gamma})(L_{\alpha}\otimes m_{\beta\gamma,\gamma^{-1}}\otimes L_{\gamma})(L_{\alpha}\otimes L_{\beta\gamma}\otimes \mathsf{coev}_{\gamma}) \\
 & = & (m_{\alpha\beta\gamma,\gamma^{-1}}\otimes L_{\gamma})(m_{\alpha,\beta\gamma}\otimes L_{\gamma^{-1}}\otimes L_{\gamma})(L_{\alpha}\otimes L_{\beta\gamma}\otimes \mathsf{coev}_{\gamma}) \\
 & = & (m_{\alpha\beta\gamma,\gamma^{-1}}\otimes L_{\gamma})(m_{\alpha,\beta\gamma}\otimes \mathsf{coev}_{\gamma}) = \Delta_{\alpha\beta,\gamma}m_{\alpha,\beta\gamma},\n\end{array}
$$

where we used associativity. Therefore L forms a Frobenius  $(\mathcal{G}, \mathcal{V})$  –category in the sense of of [5.5.](#page-12-0) Let  $\alpha: x \to y$ , then

$$
\nu_x m_{\alpha,\alpha^{-1}} = \eta_{1_x}(j_x \otimes m_{\alpha,\alpha^{-1}}) = \eta_\alpha (m_{1_x,\alpha} \otimes L_{\alpha^{-1}})(j_x \otimes L_{\alpha} \otimes L_{\alpha^{-1}}) = \eta_\alpha,
$$

by compatibility of multiplication and inner product. Hence L is symmetric if and only if  $\eta$  $\overline{\phantom{a}}$  is.

We are finally able to define crossed loop Frobenius  $(G, V)$ -categories and therefore introduce the notion of crossings. By reasons explained in section [6](#page-16-0) these crossings are defined for all paths, but only act on loops  $\alpha : x \to x$  in G. This will be the main definition in classifying 2-dimensional  $(X, Y)$ -HQFTs.

<span id="page-15-0"></span>**Definition 5.11.** Given a Frobenius  $(\mathcal{G}, \mathcal{V})$ -category L and a morphism  $f : L_{\alpha} \otimes L_{\beta} \to L_{\beta}$ in V for any  $\alpha, \beta \in \mathcal{G}$ . Its **partial trace** is defined as

$$
\text{Trace}_{\beta}(f) := \eta_{\beta}(f \otimes L_{\beta^{-1}})(L_{\alpha} \otimes \text{coev}_{\beta^{-1}}) : L_{\alpha} \to I.
$$
\n
$$
L_{\alpha} \qquad L_{\alpha}
$$
\n
$$
\boxed{\text{Trace}_{\beta}(f)} = \boxed{f}
$$

**Definition 5.12.** Given a groupoid  $\mathcal{G}$ , consider the totally disconnected groupoid of loops  $\mathcal{G}_0$ . It has the same objects, but only endomorphisms  $\alpha : x \to x$ . A loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -category is a Frobenius  $(\mathcal{G}_0, \mathcal{V})$ -category. A loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -category is called crossed, if there is a family of morphisms called the crossing  $\phi_\beta^\alpha: L_\alpha \to L_{\beta\alpha\beta^{-1}}$  for all  $\alpha: x \to x \in \mathcal{G}_0$  and  $\beta: y \to x \in \mathcal{G}$  such that:

(LF1) They respect the  $(\mathcal{G}_0, \mathcal{V})$ -category and  $(\mathcal{G}_0, \mathcal{V})$ -opcategory structure:

$$
\begin{aligned}\n\phi_{\beta}^{\alpha\alpha'} m_{\alpha,\alpha'} &= m_{\beta\alpha\beta^{-1},\beta\alpha'\beta^{-1}} (\phi_{\beta}^{\alpha} \otimes \phi_{\beta}^{\alpha'}), & \phi_{\beta}^{1_x} j_x &= j_y, \\
(\phi_{\beta}^{\alpha} \otimes \phi_{\beta}^{\alpha'}) \Delta_{\alpha\alpha'} &= \Delta_{\beta\alpha\beta^{-1},\beta\alpha'\beta^{-1}} \phi_{\beta}^{\alpha\alpha'}, & \nu_y \phi_{\beta}^{1_x} &= \nu_x,\n\end{aligned}
$$

for all  $\alpha, \alpha': x \to x$  and  $\beta: y \to x$ .

(LF2) They are compatible composition:

$$
\phi^{\beta\alpha\beta^{-1}}_{\gamma}\phi^{\alpha}_{\beta}=\phi^{\alpha}_{\gamma\beta}, \quad \phi^{\alpha}_{1_x}=L_{\alpha},
$$

for all  $\alpha: x \to x$  and  $\beta: y \to x, \gamma: z \to y$ .  $(LF3)$  The *crossed conditions* are satisfied:

$$
m_{\beta,\alpha}\sigma = m_{\beta\alpha\beta^{-1},\beta}(\phi_{\beta}^{\alpha} \otimes L_{\beta}) : L_{\alpha} \otimes L_{\beta} \to L_{\beta\alpha} \qquad \forall \alpha : x \to x, \beta : x \to x,
$$
  

$$
\phi_{\alpha}^{\alpha} = L_{\alpha} : L_{\alpha} \to L_{\alpha} \qquad \forall \alpha : x \to x,
$$
  
Trace <sub>$\alpha$</sub>  $(m_{\alpha\beta(\beta\alpha)^{-1},\alpha}(L_{\alpha\beta\alpha^{-1}\beta^{-1}} \otimes \phi_{\beta}^{\alpha})) = \text{Trace}_{\beta}(\phi_{\alpha^{-1}}^{\alpha\beta\alpha^{-1}}m_{\alpha\beta(\beta\alpha)^{-1},\beta}(L_{\alpha\beta\alpha^{-1}\beta^{-1}} \otimes L_{\beta})),$   
for all  $\alpha, \beta : x \to x.$ 

A crossed loop Frobenius  $(G, V)$ -functor  $F: L \to L'$  is a Frobenius  $(\mathcal{G}_0, V)$ -functor, such that  $F(\phi_{\beta}^{\alpha}) = \phi_{\beta}'^{\alpha}$ . Denote the category of loop Frobenius  $(G, V)$ -categories and -functors by  $\mathsf{Frob}^{\mathcal{G}}_{\circ}(\mathcal{V}).$ 

From this definition general properties of crossed loop Frobenius  $(G, V)$ -categories can already be deduced.

**Corollary 5.13.** For  $\alpha : x \to x$  the operation  $m_{\alpha,\alpha} : L_{\alpha} \otimes L_{\alpha} \to L_{\alpha\alpha}$  is commutative. Particularly,  $L_{\alpha}$  is a commutative monoid for any idempotent  $\alpha = \alpha^2$ . For  $\alpha = 1_x$  and  $G := \mathcal{G}(x, x)$  there is a left action of G on  $L_{1_x}$  by  $\{\phi_{\beta}^{1_x}\}$  ${}_{\beta}^{\perp_x} \rbrace_{\beta \in G}$ 

**Proposition 5.14.** Any crossed loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -category is symmetric as a Frobenius  $(\mathcal{G}_0, \mathcal{V})$ -category.

Proof. First of all note that  $\phi_{\alpha}^{\alpha} = L_{\alpha} = \phi_{1_x}^{\alpha}$  and  $\phi_{\alpha^{-1}}^{\alpha} : L_{\alpha} \to L_{\alpha}$  satisfies  $\phi_{\alpha^{-1}}^{\alpha} \phi_{\alpha}^{\alpha} = \phi_{1_x}^{\alpha}$ , hence it is the identity on  $L_{\alpha}$ . Now

$$
m_{\alpha^{-1},\alpha}\sigma=m_{\alpha,\alpha^{-1}}(\phi_{\alpha^{-1}}^{\alpha}\otimes L_{\alpha})=m_{\alpha,\alpha^{-1}}.
$$

Particularly  $\nu_x m_{\alpha^{-1}.\alpha} \sigma = \nu_x m_{\alpha,\alpha^{-1}}$ .

## 6. CLASSIFICATION OF 2-DIMENSIONAL  $(X, Y)$ -HQFTs

<span id="page-16-0"></span>By the same reasoning as for the classification of 1-dimensional  $(X, Y)$ -HQFTs assume some connectedness condition on the pair  $(X, Y)$ . Fix therefore  $(X, Y)$  an admissible pair and  $V$  a strict symmetric monoidal target category. We classify HQFTs with target space pair  $(X, Y)$  by means of crossed loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -categories. This is done in two steps: determining a crossed loop Frobenius  $(G, V)$ -category for all 2-dimensional  $(X, Y)$ -HQFTs, called the *underlying* crossed loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -category. Secondly, showing that every crossed loop Frobenius  $(G, V)$ -category induces a 2-dimensional  $(X, Y)$ -HQFT.

By Proposition [2.7](#page-4-1) it is enough to consider a skeleton of the category  $\textsf{Cob}^{X,Y}_{2}$ . Since any connected 1-dimensional  $(X, Y)$ -manifold is  $(X, Y)$ -homeomorphic to the 1-sphere  $S^1$  together with a homotopy classes of pointed maps to X, identify them with elements  $\alpha \in \mathcal{G}(x, x)$ . Hence, a 2-dimensional  $(X, Y)$ -HQFT  $\mathcal Z$  maps a connected  $(X, Y)$ -manifold to an object  $L_{\alpha}$  in V. By Proposition [3.4](#page-7-2) and the fact, that the symmetric monoidal functor  $\mathcal Z$  preserves duals, identify  $L_{\alpha^{-1}} = L_{\alpha}^*$ . Further, for M an arbitrary object in  $\textsf{Cob}_{2}^{X,Y}$  with the components  $\{\alpha_p\}$  its image under Z is  $\mathcal{Z}(M) = \bigotimes L_{\alpha_p}$ .

Morphisms in  $\mathsf{Cob}^{X, Y}_2$  are 2-dimensional  $(X, Y)$ -cobordisms, called  $(X, Y)$ -surfaces.  $\mathcal Z$  maps such an  $(X, Y)$ -surface  $(W, M, N, f)$  with  $M = \coprod \alpha_p$  and  $N = \coprod \beta_q$  to a morphism  $\mathcal{Z}(W)$ :  $\bigotimes L_{\alpha_p} \to \bigotimes L_{\beta_q}$  in V. Given any  $(X, Y)$ -surface W label their boundary components with orientations  $\epsilon_i \in \{+, -\}$ , depending if the orientation on the boundary agrees with W or not. Hence, the negative compontents will be inputs of  $\mathcal{Z}(W)$  and the positive its output.

The easiest  $(X, Y)$ -surface is the closed disc  $B_{\epsilon}(\alpha)$ , with orientation on the boundary  $\epsilon$  and  $g|_{\partial B} = \alpha$  with  $\alpha \in \mathcal{G}(x, x)$ . Note, that given a map  $B_{\epsilon}(\alpha) \to X$  there is a contraction of the loop to the base point, hence  $\alpha$  is trivial and there are only two types of closed discs:  $B_+(1_x)$ 

and  $B_-(1_x)$ .



Another two elementary  $(X, Y)$ -surfaces are the annulus and the disc with two holes. Let  $C := \mathbb{S}^1 \times [0,1], C^0 := \mathbb{S}^1 \times \{0\}$  and  $C^1 := \mathbb{S}^1 \times \{1\}$  and provide them with base points  $(s,0)$ and  $(s,1)$  for  $s \in \mathbb{S}^1$ . Given two signs  $\epsilon, \mu = \pm$  we define  $C_{\epsilon,\mu}$  the oriented annulus C such that

$$
\partial C_{\epsilon,\mu} = \epsilon C_{\epsilon}^0 \coprod \mu C_{\mu}^1.
$$

The homotopy class of  $g: C_{\epsilon,\mu} \to X$  is determined by a loop  $\alpha = [g|_{C_{\epsilon}}]$  and a path  $\beta^{-1} =$  $[g|_{s\times[0,1]}]$ . Note that  $[g|_{C_{\mu}^1}] = \beta \alpha^{-\epsilon \mu} \beta^{-1} \in \mathcal{G}$ . Denote the annulus by  $C_{\epsilon,\mu}(\alpha;\beta)$ .



Let now D be a 2-disc with two holes, with boundary components  $S, T, U$  and base points s, t, u. Fix two arcs us, ut in D. Given three signs  $\epsilon, \mu, \nu$  we define analgously  $D_{\epsilon,\mu,\nu}$  such that

$$
\partial D_{\epsilon,\mu,\nu} = \epsilon S_{\epsilon} \coprod \mu T_{\mu} \coprod \nu U_{\nu}.
$$

The homotopy class of  $g: D_{\epsilon,\mu,\nu} \to X$  is given by loops  $\alpha := [g|_{S_{\epsilon}}]$  and  $\beta := [g|_{T_{\mu}}]$  and two paths  $\rho := [g_{\vert u s}]$  and  $\delta := [g_{\vert u t}]$ . Denote this  $(X, Y)$ -surface by  $D_{\epsilon,\mu,\nu}(\alpha,\beta;\rho,\delta)$ . Note that the loop  $\gamma := [g|_{U_{\nu}}]$  is given by  $(\rho \alpha^{-\epsilon} \rho^{-1} \delta \beta^{-\mu} \delta^{-1})^{\nu}$ . There are  $(X, Y)$ -homeomorphisms

$$
D_{\epsilon,\mu,\nu}(\alpha,\beta;\rho,\delta) \simeq D_{\mu,\nu,\epsilon}(\beta,\gamma;\rho^{-1}\delta,\rho^{-1}) \simeq D_{\nu,\epsilon,\mu,}(\gamma,\alpha;\delta^{-1},\delta^{-1}\rho).
$$



Analogously, denote  $(X, Y)$ -discs with *n*-holes by  $D_{\epsilon_1,\ldots,\epsilon_n,\mu}(\alpha_1, \ldots, \alpha_n; \rho_1, \ldots, \rho_n)$  where  $\epsilon_i =$  $\pm$  are the orientations on the inner boundary components and  $\mu = \pm$  the orientation on the outer circle. The loops determined by the inner boundary are denote by  $\alpha_i$  and the paths connecting the basepoint of the outer circle to the base point of the inner circles are denote by  $\rho_i$ .

The objects  $(L_{\alpha})_{\alpha \in \mathcal{G}}$  can be endowed with a crossed loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -category structure, using these elementary  $(X, Y)$ -surfaces. The two closed discs are considered cobordisms from or to the empty manifold, hence the  $(X, Y)$ -HQFT  $\mathcal Z$  maps them to

$$
\mathcal{Z}(B_+(1_x)) =: j_x : I \to L_{1_x}, \qquad \mathcal{Z}(B_-(1_x)) =: \nu_x : L_{1_x} \to I
$$

Let  $\alpha, \beta : x \to x$  be two loops, then  $D_{-+}(\alpha, \beta; 1_x, 1_x)$  is mapped to the morphism

$$
\mathcal{Z}(D_{--+}(\alpha,\beta;1_x,1_x)) =: m_{\alpha,\beta}: L_{\alpha} \otimes L_{\beta} \to L_{\alpha\beta},
$$

and similarly

$$
\mathcal{Z}(D_{++-}(\beta,\alpha;1_x,1_x)) =: \Delta_{\alpha,\beta}: L_{\alpha\beta} \to L_{\alpha} \otimes L_{\beta}.
$$

For  $\alpha \in \mathcal{G}(x,x)$  and  $\beta : y \to x$ , we can consider the  $(X,Y)$ -annulus  $C_{-+}(\alpha;\beta)$  as a cobordism between  $(C^0_-, \alpha)$  and  $(C^1_+, \beta \alpha \beta^{-1})$ . Therefore

$$
\mathcal{Z}(C_{-+}(\alpha;\beta)) =: \phi_{\beta}^{\alpha} : L_{\alpha} \to L_{\beta\alpha\beta^{-1}}.
$$

Particularly,  $C_{-+}(\alpha; 1_x)$  is  $(X, Y)$ -homeomorphic to the identity cylinder, hence

$$
\mathcal{Z}(C_{-+}(\alpha; 1_x)) = L_{\alpha}.
$$

Furthermore, we define

$$
\mathcal{Z}(C_{--}(\alpha; 1_x)) =: \eta_{\alpha}: L_{\alpha} \otimes L_{\alpha^{-1}} \to I.
$$
  

$$
\mathcal{Z}(C_{++}(\alpha; 1_x)) =: \operatorname{coev}_{\alpha}: I \to L_{\alpha^{-1}} \otimes L_{\alpha} \to I
$$

and we will see in the proof of Lemma [6.7,](#page-24-0) that this is consitent with both definitions of Frobenius  $(G, V)$ -categories. These operations form a crossed loop Frobenius  $(G, V)$ -category and determine the functor  $\mathcal Z$  uniquely.

**Theorem 6.1.** Let  $(V, \otimes, I, \sigma)$  be a (strict) symmetric monoidal category. Let  $(X, Y)$  be an admissble pair of spaces and denote the relative fundamental groupoid of the pair  $(X, Y)$  by  $\mathcal{G} = \Pi_1(X, Y)$ . The category  $\mathcal{Q}_2(X, Y, V)$  of 2-dimensional  $(X, Y)$ -HQFTs is equivalent to the category  $\mathsf{Frob}_{\circ}^{\mathcal{G}}(\mathcal{V})$  of crossed loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -categories.

The proof is done in multiple lemmas by adapting the proof of Turaev in [\[11\]](#page-30-9).

**Lemma 6.2.** The objects  $(L_{\alpha})_{\alpha \in \mathcal{G}_0}$  form a  $(\mathcal{G}_0, \mathcal{V})$ -category with multiplication  $m_{\alpha,\beta}$  and unit  $j_x$ .

*Proof.* We prove associativity and unitality. Let  $\alpha, \beta, \gamma \in \mathcal{G}(x, x)$  and

 $W = D_{---+}(\alpha, \beta, \gamma; 1_x, 1_x, 1_x)$ 

be a disc with 3 holes, labeled by  $\alpha$ ,  $\beta$ ,  $\gamma$  and the three paths connecting the basepoints labeled by identities.



W can be disected into elementary  $(X, Y)$ -surfaces in two different ways. Let

$$
W_0 := D_{-+}(\alpha, \beta; 1_x, 1_x) \coprod C_{-+}(\gamma; 1_x)
$$

and  $W_1 := D_{-+}(\alpha\beta, \gamma; 1_x, 1_x)$  and further

$$
W_2 := C_{-+}(\alpha; 1_x) \coprod D_{--+}(\beta, \gamma; 1_x, 1_x)
$$

and  $W_3 := D_{-+}(\alpha, \beta\gamma; 1_x, 1_x)$ . Gluing  $W_0$  to  $W_1$  and  $W_2$  to  $W_3$  yields discs with 3 holes  $(X, Y)$ -homeomorphic to W. Since  $\mathcal Z$  is functorial:

$$
m_{\alpha\beta,\gamma}(m_{\alpha,\beta} \otimes L_{\gamma}) = \mathcal{Z}(D_{--+}(\alpha\beta,\gamma;1_x,1_x))(\mathcal{Z}(D_{--+}(\alpha,\beta;1_x,1_x) \otimes \mathcal{Z}(C_{-+}(\gamma;1_x)))
$$
  
\n
$$
= \mathcal{Z}(W_1)\mathcal{Z}(W_0) = \mathcal{Z}(W) = \mathcal{Z}(W_3)\mathcal{Z}(W_2)
$$
  
\n
$$
= \mathcal{Z}(D_{--+}(\alpha,\beta\gamma;1_x,1_x))(\mathcal{Z}(C_{-+}(\alpha;1_x)) \otimes \mathcal{Z}(D_{--+}(\beta,\gamma;1_x,1_x)))
$$
  
\n
$$
= m_{\alpha,\beta\gamma}(L_{\alpha} \otimes m_{\beta,\gamma}).
$$

For unitality, look at  $C_{-+}(\alpha; 1_x)$  and the gluing of  $D_{-+}(\alpha, 1_x; 1_x, 1_x)$  with  $C_{-+}(\alpha; 1_x)$  and  $B_{+}(1_{x})$  glued to the first and second inner boundary respectively. These two  $(X, Y)$ -surfaces are  $(X, Y)$ -homeomorphic; hence

$$
m_{\alpha,1_x}(L_\alpha \otimes j_x) = \mathcal{Z}(D_{--+}(\alpha, 1_x; 1_x, 1_x))(\mathcal{Z}(C_{-+}(\alpha; 1_x)) \otimes \mathcal{Z}(B_{+}(1_x))) = \mathcal{Z}(C_{-+}(\alpha; 1_x)) = L_\alpha
$$
  
and analogously for the left unitality. Hence  $(L_\alpha)_{\alpha \in \mathcal{G}_0}$  form a  $(\mathcal{G}_0, \mathcal{V})$  – *category*.

**Lemma 6.3.** The objects  $(L_{\alpha})_{\alpha \in \mathcal{G}_0}$  form a  $(\mathcal{G}_0, \mathcal{V})$  – opcategory with comultiplication  $\Delta_{\alpha,\beta}$ and counit  $\nu_x$ .

This proof is dual to the previous, by reversing orientations on  $D_{-+}$  and  $B_{+}$ .

**Lemma 6.4.** The objects  $(L_{\alpha})_{\alpha \in \mathcal{G}_0}$  form a loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -category.

Proof. We need to prove the Frobenius conditions:

$$
\Delta_{\alpha,\beta\gamma} m_{\alpha\beta,\gamma} = (L_{\alpha} \otimes m_{\beta,\gamma}) (\Delta_{\alpha,\beta} \otimes L_{\gamma}), \Delta_{\alpha\beta,\gamma} m_{\alpha,\beta\gamma} = (m_{\alpha,\beta} \otimes L_{\gamma}) (L_{\alpha} \otimes \Delta_{\beta,\gamma})
$$

for  $\alpha, \beta, \gamma \in \mathcal{G}(x, x)$ . To prove the first condition we look at the  $(X, Y)$ -disc with three holes  $W = D_{-++}(\alpha, \beta\gamma, \gamma; 1_x, 1_x, 1_x)$ , which can be represented as gluing  $W_1 = D_{-++}(\alpha\beta\gamma, \gamma; 1_x, 1_x)$  $\simeq D_{++-}(\gamma, \alpha\beta; 1_x, 1_x)$  to  $W_2 = D_{--+}(\alpha, \beta\gamma; 1_x, 1_x)$ . Similarly, we can represent W as gluing  $W_3 = D_{-+}(\alpha, \beta; 1_x, 1_x) \coprod C_{-+}(\gamma, 1_x)$  to  $W_4 = C_{-+}(\alpha; 1_x) \coprod D_{-++}(\beta \gamma, \gamma; 1_x, 1_x)$ , where the last disc with two holes is  $(X, Y)$ -homeomorphic to  $D_{++-}(\gamma, \beta; 1_x, 1_x)$ .



Hence

$$
\Delta_{\alpha\beta,\gamma}m_{\alpha,\beta\gamma} = \mathcal{Z}(D_{+-+}(\alpha\beta\gamma,\gamma;1_x,1_x))\mathcal{Z}(D_{--+}(\alpha,\beta\gamma;1_x,1_x)) = \mathcal{Z}(W)
$$
  
\n
$$
= \mathcal{Z}\left(D_{--+}(\alpha,\beta;1_x,1_x)\coprod C_{-+}(\gamma,1_x)\right)\mathcal{Z}\left(C_{-+}(\alpha;1_x)\coprod D_{-++}(\beta\gamma,\gamma;1_x,1_x)\right)
$$
  
\n
$$
= (m_{\alpha,\beta}\otimes L_{\gamma})(L_{\alpha}\otimes \Delta_{\beta,\gamma}).
$$

Similar argumentation on  $D_{+--+}(\alpha, \alpha\beta, \gamma; 1_x, 1_x, 1_x)$  shows that

$$
\Delta_{\alpha,\beta\gamma}m_{\alpha\beta,\gamma}=(L_\alpha\otimes m_{\beta,\gamma})(\Delta_{\alpha,\beta}\otimes L_\gamma).
$$

**Lemma 6.5.** Given some 2-punctured  $(X, Y)$ -disc  $D_{-+}(\alpha, \beta; \rho, \delta)$ , such that the output label is  $\gamma = \rho \alpha \rho^{-1} \delta \beta \rho^{-1} = \beta$ , let P be the  $(X, Y)$ -surface obtained from gluing together the two boundary components labeled  $\beta$ . Then

$$
\mathcal{Z}(P) = \text{Trace}_{\beta}(\mathcal{Z}(D_{--+}(\alpha, \beta; \rho, \delta))).
$$

 $\Box$ 

*Proof.* The  $(X, Y)$ -surface is  $(X, Y)$ -homeomorphic to gluing  $D_{-+}(\alpha, \beta; \rho, \delta) \coprod C_{-+}(\beta^{-1}; 1_x)$ to  $C_{-+}(\alpha; 1_x) \coprod C_{--}(\beta; 1_x)$  and  $C_{--}(\beta; 1_x)$ .



This yields

$$
\mathcal{Z}(P)=\eta_{\beta}(\mathcal{Z}(D_{--+}(\alpha,\beta;\rho,\delta))\otimes L_{\beta^{-1}})(L_{\alpha}\otimes \mathsf{coev}_{\beta^{-1}}),
$$

hence it is the partial trace as define in [5.11.](#page-15-0)

**Lemma 6.6.** The loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -category  $(L, m, j, \Delta, \nu)$  is crossed via  $\mathcal{Z}(C_{-+}(\alpha; \beta))$  =  $\phi^{\alpha}_{\beta}$ .

*Proof.* We turn our attention to  $C_{-+}(\alpha;\beta)$  where  $\alpha: x \to x$  is a loop and  $\beta: y \to x$  a path in X with  $x, y \in Y$ . The family  $\phi^{\alpha}_{\beta}$  satisfies all axioms for a crossing of the loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -category  $(L_{\alpha})_{\alpha \in \mathcal{G}_0}$ .

We begin by showing compatibility with the  $(\mathcal{G}_0, \mathcal{V})$ -category and  $(\mathcal{G}_0, \mathcal{V})$ -opcategory structure: The X-disc  $B_+(1_y)$  is mapped to  $j_y: I \to L_{1_y}$  by  $\mathcal Z$ . It can also be obtained by gluing  $B_+(1_x)$  to  $C_{-+}(1_x;\beta)$ . Similarly  $B_-(1_y) \simeq B_-(1_x) \coprod_{\partial B_-(1_x)} C_{+-}(1_x;\beta)$ 



This shows  $\phi_{\beta}^{1_x} j_x = j_y$  and  $\nu_x \phi_{\beta^{-1}}^{1_y} = \nu_y$ , where we used that  $C_{+-}(1_x;\beta) \simeq C_{-+}(1_y;\beta^{-1})$ . For (co)multiplication consider  $D_{-+}(\alpha, \alpha'; \beta, \beta)$  and  $D_{++-}(\beta \alpha' \beta^{-1}, \beta \alpha \beta^{-1}; \beta^{-1}, \beta^{-1})$  respectively. We can represent  $D_{-+}(\alpha, \alpha'; \beta, \beta)$  either as the gluing of  $D_{-+}(\beta \alpha \beta^{-1}, \beta \alpha' \beta^{-1}; 1_y, 1_y)$ and  $C_{-+}(\alpha, \beta) \coprod C_{-+}(\alpha', \beta)$  along the inner circles or  $D_{--+}(\alpha, \alpha'; 1_x, 1_x)$  and  $C_{-+}(\alpha \alpha'; \beta)$ along the outer circle, hence  $\phi_{\beta}^{\alpha}m_{\alpha,\alpha'}=m_{\beta\alpha\beta^{-1},\beta\alpha'\beta^{-1}}(\phi_{\beta}^{\alpha}\otimes\phi_{\beta}^{\alpha'}$  $_{\beta}^{\alpha'}$  ).



Similarly, for  $D_{++-}(\beta \alpha' \beta^{-1}, \beta \alpha \beta^{-1}; \beta^{-1}, \beta^{-1})$  we have

 $(C_{+-}(\beta\alpha\beta^{-1},\beta^{-1})\coprod C_{+-}(\beta\alpha'\beta^{-1},\beta^{-1})\coprod$ ∂C ∐∂C  $D_{++-}(\alpha', \alpha; 1_x, 1_x)$  $= D_{++-}(\beta \alpha' \beta^{-1}, \beta \alpha \beta^{-1}, 1_y, 1_y)$ ∂C  $C_{+-}(\beta \alpha \alpha' \beta^{-1}, \beta^{-1}).$ 



Since  $C_{+-}(\beta \alpha \beta^{-1}, \beta^{-1}) \simeq C_{-+}(\alpha, \beta)$ , this results in

$$
(\phi^{\alpha}_{\beta} \otimes \phi^{\alpha'}_{\beta})\Delta_{\alpha\alpha'} = \Delta_{\beta\alpha\beta^{-1},\beta\alpha'\beta^{-1}}\phi^{\alpha\alpha'}_{\beta}.
$$

We have  $\phi_{1_x}^{\alpha} = \mathcal{Z}(C_{-+}(\alpha; 1_x)) = L_{\alpha}$  and regarding the composition

$$
\begin{array}{rcl}\n\phi_{\gamma\beta}^{\alpha} & = & \mathcal{Z}(C_{-+}(\alpha;\gamma\beta)) = \mathcal{Z}(C_{-+}(\beta\alpha\beta^{-1};\gamma))\mathcal{Z}(C_{-+}(\alpha;\beta)) \\
& = & \phi_{\gamma}^{\beta\alpha\beta^{-1}}\phi_{\beta}^{\alpha}.\n\end{array}
$$

A Dehn twist about the circle  $\mathbb{S} \times \{1/2\} \subset C_{-+}(\alpha; 1_x)$  defines an  $(X, Y)$ -homeomorphism  $C_{-+}(\alpha; 1_x) \simeq C_{-+}(\alpha, \alpha)$  and since Z is invariant under  $(X, Y)$ -homeomorphisms we have  $\phi_{\alpha}^{\alpha} = L_{\alpha}$  for all  $\alpha : x \to x$ .

Now for  $\alpha, \beta : x \to x$  look at  $D_{-+}(\alpha, \beta; 1_x, 1_x)$ , but  $(X, Y)$ -homeomorphically switch the positions of the inner circles:



This is a  $(X, Y)$ -homeomorphic to  $D_{-+}(\beta, \alpha; 1_x, \rho)\sigma$  for some  $\rho : x \to x$  and since the composition  $1_x \beta 1_x^{-1} \rho = 1_x$ , the homotopy class of  $\rho$  is the same as of  $\beta^{-1}$ . Hence

$$
m_{\alpha,\beta} = \mathcal{Z}(D_{--+}(\alpha,\beta,1_x,1_x)) = \mathcal{Z}(D_{--+}(\beta,\alpha;1_x,\beta^{-1}))\sigma =
$$
  
=  $\mathcal{Z}(D_{--+}(\beta,\beta^{-1}\alpha\beta,1_x,1_x))(\mathcal{Z}(C_{-+}(\beta,1_x)) \otimes \mathcal{Z}(C_{-+}(\alpha,\beta)))\sigma$   
=  $m_{\beta,\beta^{-1}\alpha\beta}(L_{\beta} \otimes \phi_{\beta^{-1}}^{\alpha})\sigma.$ 

We apply this to  $m_{\beta\alpha\beta^{-1},\beta}$  to get

$$
\begin{array}{lcl} m_{\beta\alpha\beta^{-1},\beta}(\phi^{\alpha}_{\beta}\otimes L_{\beta}) & = & m_{\beta,\alpha}\sigma(\phi^{\beta\alpha\beta^{-1}}_{\beta^{-1}}\otimes L_{\beta})(\phi^{\alpha}_{\beta}\otimes L_{\beta}) \\ \\ & = & m_{\beta,\alpha}\sigma(\phi^{\alpha}_{\beta^{-1}\beta}\otimes L_{\beta}) = m_{\beta,\alpha}\sigma, \end{array}
$$

as in the form of the definition of a crossing on loop Frobenius  $(G, V)$ -categories. We are only left to prove the crossed condition:

$$
\text{Trace}_{\,\alpha}\big(m_{\alpha\beta(\beta\alpha)^{-1},\alpha}\big(L_{\alpha\beta\alpha^{-1}\beta^{-1}}\otimes\phi^{\alpha}_{\beta}\big)\big)=\quad \text{Trace}_{\,\beta}\big(\phi^{\alpha\beta\alpha^{-1}}_{\alpha^{-1}}m_{\alpha\beta(\beta\alpha)^{-1},\beta}\big(L_{\alpha\beta\alpha^{-1}\beta^{-1}}\otimes L_{\beta}\big)\big).
$$

Consider the punctured torus  $P = \mathbb{S}^1 \times \mathbb{S}^1 \setminus B$ . Let  $(s, s)$  be the basepoint of the circle and deform the hole, such that it does not intersect  $\mathbb{S}^1 \times \{s\}$  and  $\{s\} \times \mathbb{S}^1$ . Further let  $g: P \to X$ with  $g(s, s) = x \in Y$ , then g is fully determined by  $\alpha, \beta : x \to x$  in G with  $\alpha = g|_{\mathbb{S} \times \{s\}}$  and  $\beta = g|_{\{s\}\times\mathbb{S}}$  and  $\partial P$  is represented by the loop  $\alpha\beta\alpha^{-1}\beta^{-1}$ .



Hence  $\mathcal{Z}(P): L_{\alpha\beta\alpha^{-1}\beta^{-1}} \to I$ . On one hand P can be constructed by taking  $D_{--+}(\alpha\beta\alpha^{-1}\beta^{-1},$  $\alpha$ ;  $1_x$ ,  $\beta$ ) and gluing the two circles labeled by  $\alpha$  together. Since

 $\mathcal{Z}(D_{--+}(\alpha\beta\alpha^{-1}\beta^{-1},\alpha;1_x,\beta))=m_{\alpha\beta\alpha^{-1}\beta^{-1},\beta\alpha\beta^{-1}}(L_{\alpha\beta\alpha^{-1}\beta^{-1}}\otimes\phi_{\beta}^{\alpha}):L_{\alpha\beta\alpha^{-1}\beta^{-1}}\otimes L_{\alpha}\rightarrow L_{\alpha},$ by the previous lemma

$$
\mathcal{Z}(P) = \mathrm{Trace}_{\alpha}(m_{\alpha\beta(\beta\alpha)^{-1},\alpha}(L_{\alpha\beta\alpha^{-1}\beta}\otimes \phi^{\alpha}_{\beta})).
$$

On the other hand, P can be obtained by taking  $D_{-+}(\alpha\beta\alpha^{-1}\beta^{-1},\beta;\alpha^{-1},\alpha^{-1})$  and gluing the two circles labeled by  $\beta$  together.

$$
\mathcal{Z}(D_{--+}(\alpha\beta\alpha^{-1}\beta^{-1},\beta;\alpha^{-1},\alpha^{-1})) = m_{\beta\alpha^{-1}\beta^{-1}\alpha^{-1},\alpha\beta\alpha^{-1}}(\phi_{\alpha^{-1}}^{\alpha\beta\alpha^{-1}\beta^{-1}}\otimes\phi_{\alpha^{-1}}^{\beta})
$$
  

$$
= \phi_{\alpha^{-1}}^{\alpha\beta\alpha^{-1}}m_{\alpha\beta(\beta\alpha)^{-1},\beta}:L_{\alpha\beta\alpha^{-1}\beta^{-1}}\otimes L_{\beta}\to L_{\beta},
$$

and therefore

$$
\mathcal{Z}(P) = \mathsf{Trace}_{\,\beta}(\phi_{\alpha^{-1}}^{\alpha\beta\alpha^{-1}}m_{\alpha\beta(\beta\alpha)^{-1},\beta}).
$$

<span id="page-24-0"></span>**Lemma 6.7.** The functor  $F : \mathcal{Q}_2(X, Y, V) \to \mathsf{Frob}_{\circ}^{\mathcal{G}}(V)$  assigning to every HQFT  $\mathcal{Z}$  :  $\mathsf{Cob}^{X,Y}_2 \to \mathcal{V}$  its underlying crossed loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -category is an equivalence of categories.

*Proof.* Firstly the assignement is functorial. Let  $\rho : \mathcal{Z} \Rightarrow \mathcal{Z}'$  be a morphism of HQFT, hence  $\rho$ is a monoidal natural isomorphism. Explicitely,  $\rho$  is given by a family of maps  $\rho_{\alpha}: L_{\alpha} \to L'_{\alpha}$ , given by  $\mathcal{Z}(\mathbb{S}, \alpha) =: L_{\alpha}, \mathcal{Z}'(\mathbb{S}, \alpha) =: L'_{\alpha}, \text{ satisfying:}$ 

$$
\bigotimes_{i=1}^{n} L_{\alpha_i} \xrightarrow{\bigotimes_{i=1}^{m} \rho_{\alpha_i}} \bigotimes_{i=1}^{n} L'_{\alpha}
$$
  

$$
\bigotimes_{j=1}^{m} L_{\beta_j} \xrightarrow{\bigotimes_{j=1}^{m} \rho_{\beta_j}} \bigotimes_{j=1}^{n} L_{\beta_j},
$$

for all  $(W, g)$  cobordant to  $\coprod_{i=1}^{n} (\mathbb{S}, \alpha_i)$  and  $\coprod_{j=1}^{n} (\mathbb{S}, \beta_j)$ . Particularly, that means  $\rho$  is a crossed loop Frobenius functor. Further the functor  $\overrightarrow{F}$  is faithful. To show that  $F$  is full, we show

 $\Box$ 

that any morphism  $\mathcal{Z}(W, g)$  can be represented by a composition of the structural morphisms  $(m, j, \Delta, \nu, \phi)$  (and  $\eta$ , coev respectively) in L.

One shows, that it is essentially surjective, by constructing an HQFT given an arbitrary crossed loop Frobenius  $(G, V)$ -category. We use the fact that any  $(X, Y)$ -surface can be obtained by gluing discs with  $\leq 2$  holes and by connectivity condition we can assume that such cutting along circles always go through some point sent to Y. Define  $\mathcal Z$  on  $B, C, D$  and show that this assignement is independent of the gluing, hence obtaining a full  $(X, Y)$ -HQFT. For the disc with no holes we have  $\mathcal{Z}(B_-(1_x)) := \nu_x$  and  $\mathcal{Z}(B_+(1_x)) := j_x$ .

 $(X, Y)$ -discs with one hole are  $(X, Y)$ -homeomorphic to one of the four  $C_{-}(\alpha; \beta), C_{-+}(\alpha; \beta),$  $C_{+-}(\alpha;\beta) \simeq C_{-+}(\beta\alpha\beta^{-1};\beta^{-1})$  or  $C_{++}(\alpha;\beta)$ . For  $\alpha \in \mathcal{G}(x,x)$  the  $(X,Y)$ -annulus  $C_{--}(\alpha;1_x)$ is an  $(X, Y)$ -cobordisms between  $(C^0_-, \alpha) \coprod (C^1_-, \alpha^{-1})$  and  $\emptyset_1$ . Therefore

$$
\mathcal{Z}(C_{--}(\alpha;1_x)) := \eta_\alpha : L_\alpha \otimes L_{\alpha^{-1}} \to I
$$

and similarly,

$$
\mathcal{Z}(C_{++}(\alpha;1_x)):=\mathsf{coev}_\alpha:I\to L_{\alpha^{-1}}\otimes L_\alpha.
$$

Let

$$
\mathcal{Z}(C_{-+}(\alpha;\beta)) := \phi^{\alpha}_{\beta}
$$

and since  $C_{--}(\alpha;\beta)$  and  $C_{++}(\alpha;\beta)$  can be optained by gluing  $C_{-+}(\alpha^{-1};\beta)$  to  $C_{--}(\alpha;1_x)$  and  $C_{++}(\alpha; 1_x)$  define

$$
\begin{array}{rcl}\mathcal{Z}(C_{--}(\alpha;\beta))&:=&\eta_{\alpha}(L_{\alpha}\otimes\phi_{\beta^{-1}}^{\beta\alpha^{-1}\beta^{-1}})=\eta_{\beta\alpha\beta^{-1}}(\phi_{\beta}^{\alpha}\otimes L_{\beta\alpha^{-1}\beta^{-1}})\\ \mathcal{Z}(C_{++}(\alpha;\beta))&:=&(\phi_{\beta}^{\alpha^{-1}}\otimes L_{\alpha})\text{coev}_{\alpha}=(L_{\beta\alpha^{-1}\beta^{-1}}\otimes\phi_{\beta^{-1}}^{\beta\alpha\beta^{-1}})\text{coev}_{\beta\alpha\beta^{-1}}.\end{array}
$$



The equality of the two expressions follows from the axioms of a crossed loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -category

$$
\eta_{\beta\alpha\beta^{-1}}(\phi_{\beta}^{\alpha}\otimes L_{\beta\alpha^{-1}\beta^{-1}}) = \nu_{y}m_{\beta\alpha\beta^{-1},\beta\alpha^{-1}\beta^{-1}}(\phi_{\beta}^{\alpha}\otimes L_{\beta\alpha^{-1}\beta^{-1}})
$$
  
\n
$$
= \nu_{y}\phi_{\beta}^{1_{x}}m_{\alpha,\alpha^{-1}}(\phi_{\beta^{-1}}^{\beta\alpha\beta^{-1}}\otimes\phi_{\beta^{-1}}^{\beta\alpha^{-1}\beta^{-1}})(\phi_{\beta}^{\alpha}\otimes L_{\beta\alpha^{-1}\beta^{-1}})
$$
  
\n
$$
= \nu_{x}m_{\alpha,\alpha^{-1}}(\phi_{\beta^{-1}\beta}^{\alpha}\otimes\phi_{\beta^{-1}}^{\beta\alpha^{-1}\beta^{-1}})
$$
  
\n
$$
= \eta_{\alpha}(L_{\alpha}\otimes\phi_{\beta^{-1}}^{\beta\alpha^{-1}\beta^{-1}})
$$

and dually using comultiplication and coevaluation.

Every  $(X, Y)$ -disc with two holes  $D_{\epsilon,\mu,\nu}$  can be brought to either one of the 4 forms:  $D_{--}(\alpha,\beta;$  $(\rho, \delta), D_{-+}(\alpha, \beta; \rho, \delta), D_{++-}(\alpha, \beta; \rho, \delta)$  or  $D_{+++}(\alpha, \beta; \rho, \delta)$  via the  $(X, Y)$ -homeomorphisms

$$
D_{\epsilon,\mu,\nu}(\alpha,\beta;\rho,\delta) \simeq D_{\mu,\nu,\epsilon}(\beta,\gamma;\rho^{-1}\delta,\rho^{-1}) \simeq D_{\nu,\epsilon,\mu,}(\gamma,\alpha;\delta^{-1},\delta^{-1}\rho)
$$

with  $\gamma = (\rho \alpha^{-\epsilon} \rho^{-1} \delta \beta^{-\mu} \delta^{-1})^{\nu}$ . We have

$$
\mathcal{Z}(D_{--}(\alpha,\beta;\rho,\delta)) = \eta_{\gamma^{-1}}(m_{\rho\alpha\rho^{-1},\delta\beta\delta^{-1}} \otimes L_{\gamma})(\phi_{\rho}^{\alpha} \otimes \phi_{\delta}^{\beta} \otimes L_{\gamma}),
$$
  
\n
$$
\mathcal{Z}(D_{--+}(\alpha,\beta;\rho,\delta)) = m_{\rho\alpha\rho^{-1},\delta\beta\delta^{-1}}(\phi_{\rho}^{\alpha} \otimes \phi_{\delta}^{\beta}),
$$
  
\n
$$
\mathcal{Z}(D_{++-}(\alpha,\beta;\rho,\delta)) = (\phi_{\delta^{-1}}^{\delta\beta\delta^{-1}} \otimes \phi_{\rho^{-1}}^{\rho\alpha\rho^{-1}})\Delta_{\delta\beta\delta^{-1},\rho\alpha\rho^{-1}},
$$
  
\n
$$
\mathcal{Z}(D_{+++}(\alpha,\beta;\rho,\delta)) = (\phi_{\delta^{-1}}^{\delta\beta\delta^{-1}} \otimes \phi_{\rho^{-1}}^{\rho\alpha\rho^{-1}} \otimes L_{\gamma})(\Delta_{\delta\beta\delta^{-1},\rho\alpha\rho^{-1}} \otimes L_{\gamma})\text{coev}_{\gamma}.
$$

This can be seen by gluing together  $D_{-+}(\alpha, \beta; 1_x, 1_x)$  and  $D_{++-}(\alpha, \beta; 1_x, 1_x)$  with cylinders  $C_{\epsilon,\mu}$  with corresponding signs. Since every  $(X, Y)$ -surface can be disected into  $(X, Y)$ -discs with  $\leq 2$  holes, this shows that any morphism in the image of  $\mathcal Z$  can be obtained by composition of structural morphisms and the functor  $F$  is fully faithful.

We show that this assignement is topologically invariant and since the mapping class group is generated by the Dehn twists along boundary parallel circles and the reflection, it suffices to show invariance under these two operations. The Dehn twist is an  $(X, Y)$ -homeomorphism  $C_{\epsilon,\mu}(\alpha;\beta) \to C_{\epsilon,\mu}(\alpha;\beta\alpha)$  hence both cylinders must be mapped to the same map:

$$
\mathcal{Z}(C_{-+}(\alpha;\alpha\beta)) = \phi^{\alpha}_{\beta\alpha} = \phi^{\alpha}_{\beta}\phi^{\alpha}_{\alpha} = \phi^{\alpha}_{\beta} = \mathcal{Z}(C_{-+}(\alpha;\beta))
$$

and since the rest of the cylinders are defined using  $C_{-+}(\alpha;\beta)$  this holds for all  $\epsilon,\mu =$  $\pm$ . The reflection is an  $(X, Y)$ -homeomorphism  $C_{\epsilon,\epsilon}(\alpha;\beta) \to C_{\epsilon,\epsilon}(\beta\alpha^{-1}\beta^{-1};\beta^{-1})$  given by a permutation of the boundary components and the product of the orientation reversing involutions  $\mathbb{S} \to \mathbb{S}, s \mapsto -\bar{s}$  and  $[0, 1] \to [0, 1], t \mapsto 1 - t$ . We get

$$
\mathcal{Z}(C_{--}(\beta\alpha^{-1}\beta^{-1};\beta^{-1}))\sigma = \eta_{\beta\alpha^{-1}\beta^{-1}}(L_{\beta\alpha^{-1}\beta^{-1}}\otimes \phi_{\beta}^{\alpha})\sigma \n= \eta_{\beta\alpha^{-1}\beta^{-1}}\sigma(\phi_{\beta}^{\alpha}\otimes L_{\beta\alpha^{-1}\beta^{-1}}) \n= \eta_{\beta\alpha\beta^{-1}}(\phi_{\beta}^{\alpha}\otimes L_{\beta\alpha^{-1}\beta^{-1}}) = \mathcal{Z}(C_{--}(\alpha;\beta)),
$$

which has been shown above and equivalently for  $C_{++}(\beta \alpha^{-1} \beta^{-1}, \beta^{-1})$ . The invariance of the  $(X, Y)$ -disc with two holes under the Dehn twists follows form the fact that we defined them by gluing  $C_{-+}(\alpha;\rho)$  and  $C_{-+}(\beta;\delta)$  at the inner circles, which satisfy the invariance. For the Dehn twist along the outer circle observe that multiplication and comultiplication preserve the action of  $\phi$  and  $\eta$ , coev are invariant under it. Hence, we can reduce this to

invariance of cylinders. The reflection of  $D_{---}(\alpha, \beta; \rho, \delta)$  is  $D_{---}(\beta, \gamma; \rho^{-1}\delta, \rho^{-1}\rho)$  and using  $\rho^{-1}\delta\beta\delta^{-1}\rho\rho^{-1}\gamma\rho = \rho^{-1}\delta\beta\delta^{-1}(\delta\beta^{-1}\delta^{-1}\rho\alpha^{-1}\rho^{-1})\rho = \alpha^{-1}$  we deduce:

$$
D_{---}(\beta, \gamma; \rho^{-1}\delta, \rho^{-1})\sigma_{(132)}
$$
\n
$$
= \eta_{\alpha^{-1}}(m_{\rho^{-1}\delta\beta\delta^{-1}\rho, \rho^{-1}\gamma\rho} \otimes L_{\alpha})(\phi_{\rho^{-1}\delta}^{\beta} \otimes \phi_{\rho^{-1}}^{\gamma} \otimes L_{\alpha})\sigma_{(132)}
$$
\n
$$
= \eta_{\alpha^{-1}}(m_{\alpha^{-1}\rho^{-1}\gamma^{-1}\rho, \rho^{-1}\gamma\rho} \otimes L_{\alpha})(\phi_{\rho^{-1}\delta}^{\beta} \otimes \phi_{\rho^{-1}}^{\gamma} \otimes L_{\alpha})\sigma_{(132)}
$$
\n
$$
= \eta_{\alpha^{-1}\rho^{-1}\gamma^{-1}\rho}(L_{\alpha^{-1}\rho^{-1}\gamma^{-1}\rho} \otimes m_{\rho^{-1}\gamma\rho,\alpha})\sigma_{(132)}(L_{\alpha} \otimes \phi_{\rho^{-1}\delta}^{\beta} \otimes \phi_{\rho^{-1}}^{\gamma})
$$
\n
$$
= \eta_{\alpha^{-1}\rho^{-1}\gamma^{-1}\rho}(L_{\alpha^{-1}\rho^{-1}\gamma^{-1}\rho} \otimes m_{\rho^{-1}\gamma\rho\alpha\rho^{-1}\gamma^{-1}\rho,\rho^{-1}\gamma\rho})(L_{\alpha^{-1}\rho\gamma^{-1}\rho} \otimes \phi_{\rho^{-1}\gamma\rho}^{\alpha})\sigma_{(\sigma^2\mu_{\rho^{-1}\gamma\rho})}
$$
\n
$$
(\sigma \otimes L_{\rho^{-1}\gamma\rho})(L_{\alpha} \otimes \phi_{\rho^{-1}\delta}^{\beta} \otimes \phi_{\rho^{-1}}^{\gamma})
$$
\n
$$
= \eta_{\rho^{-1}\gamma^{-1}\rho}(m_{\alpha^{-1}\rho^{-1}\gamma^{-1}\rho,\rho^{-1}\gamma\rho\alpha\rho^{-1}\gamma^{-1}\rho}} \otimes L_{\rho^{-1}\gamma\rho})(\sigma \otimes L_{\rho^{-1}\gamma\rho})
$$
\n
$$
(\phi_{\rho^{-1}\gamma\rho}^{\alpha} \otimes \phi_{\rho^{-1}\delta}^{\beta} \otimes \phi_{\rho^{-1}}^{\gamma})
$$
\n
$$
= \eta_{\rho^{-1}\gamma^{-1}\rho}(m_{\alpha,\alpha^{-1}\rho^{-1}\gamma^{-1}\rho} \otimes L_{\rho^{-1}\gamma\rho})(\phi_{\alpha^{-1}\rho^{-1}\gamma^{-1}\rho}^{\alpha}
$$

where we denote by  $\sigma_{(132)} = \sigma_{A,B\otimes C} : A \otimes B \otimes C \to B \otimes C \otimes A$  for all object  $A, B, C \in V$ . Similarly, the invariance of  $D_{+++}(\alpha, \beta; \rho, \delta)$  under the reflexion is shown.

We finish by showing this assignement is invariant under gluing. The gluing of  $C_{\epsilon,\mu}(\alpha;\beta)$  to  $C_{-\mu,\nu}(\beta\alpha^{-\epsilon\mu}\beta^{-1};\delta)$  yields  $C_{\epsilon,\nu}(\alpha,\delta\beta)$ . Without loss of generality assume that  $\mu = +$ , since both cases are topologically equivalent.

• 
$$
\epsilon = -, \nu = +:
$$
  
\n
$$
\mathcal{Z}\left(C_{-,+}(\beta \alpha \beta^{-1}; \delta)\coprod C_{-+}(\alpha; \beta)\right) = \phi_{\delta}^{\beta \alpha \beta^{-1}} \phi_{\beta}^{\alpha} = \phi_{\delta \beta}^{\alpha} = \mathcal{Z}(C_{-,+}(\alpha; \delta \beta)).
$$
\n•  $\epsilon = -, \nu = -:$   
\n
$$
\mathcal{Z}\left(C_{-, -}(\beta \alpha \beta^{-1}; \delta)\coprod C_{-+}(\alpha; \beta)\right) = \eta_{\delta \beta \alpha \beta^{-1} \delta^{-1}} (\phi_{\delta}^{\beta \alpha \beta^{-1}} \otimes L_{\delta \beta \alpha^{-1} \beta^{-1} \delta^{-1}}) (\phi_{\beta}^{\alpha} \otimes L_{\delta \beta \alpha^{-1} \beta^{-1} \delta^{-1}})
$$
\n
$$
= \eta_{\delta \beta \alpha \beta^{-1} \delta^{-1}} (\phi_{\delta \beta}^{\alpha} \otimes L_{\delta \beta \alpha^{-1} \beta^{-1} \delta^{-1}})
$$
\n
$$
= \mathcal{Z}(C_{-, -}(\alpha; \delta \beta)).
$$
\n•  $\epsilon = +, \nu = +:$ 

$$
\mathcal{Z}\left(C_{-,+}(\beta\alpha^{-1}\beta^{-1};\delta)\coprod C_{++}(\alpha;\beta)\right) = (\phi_{\delta}^{\beta\alpha^{-1}\beta^{-1}} \otimes L_{\alpha})(\phi_{\beta}^{\alpha^{-1}} \otimes L_{\alpha})\text{coev}_{\alpha}
$$

$$
= (\phi_{\delta\beta}^{\alpha^{-1}} \otimes L_{\alpha})\text{coev}_{\alpha}
$$

$$
= \mathcal{Z}(C_{+,+}(\alpha,\delta\beta)).
$$

$$
\bullet\;\epsilon=+,\nu=-:
$$

$$
\mathcal{Z}\left(C_{-,-}(\beta\alpha^{-1}\beta^{-1};\delta)\coprod C_{++}(\alpha;\beta)\right)
$$
\n
$$
= (\eta_{\beta\alpha^{-1}\beta^{-1}} \otimes L_{\alpha})(L_{\beta\alpha^{-1}\beta^{-1}} \otimes \phi_{\delta^{-1}}^{\delta\beta\alpha\beta^{-1}\delta^{-1}} \otimes L_{\alpha})(\sigma \otimes L_{\alpha})
$$
\n
$$
(L_{\delta\beta\alpha\beta^{-1}\delta^{-1}} \otimes L_{\beta\alpha^{-1}\beta} \otimes \phi_{\beta^{-1}}^{\beta\alpha\beta^{-1}})(L_{\delta\beta\alpha\beta^{-1}\delta^{-1}} \otimes \text{coev}_{\beta\alpha\beta^{-1}})
$$
\n
$$
= \phi_{\beta^{-1}}^{\beta\alpha\beta^{-1}}(\eta_{\beta\alpha\beta^{-1}} \otimes L_{\beta\alpha\beta^{-1}})(L_{\beta\alpha\beta^{-1}} \otimes \text{coev}_{\beta\alpha\beta^{-1}})\phi_{\delta^{-1}}^{\delta\beta\alpha\beta^{-1}\delta^{-1}}
$$
\n
$$
= \phi_{\beta^{-1}}^{\beta\alpha\beta^{-1}}\phi_{\delta^{-1}}^{\delta\beta\alpha\beta^{-1}\delta^{-1}} = \phi_{(\delta\beta)^{-1}}^{\delta\beta\alpha\beta^{-1}\delta^{-1}}
$$
\n
$$
= \mathcal{Z}(C_{-,+}(\delta\beta\alpha\beta^{-1}\delta^{-1};(\delta\beta)^{-1}))
$$
\n
$$
= \mathcal{Z}(C_{+,-}(\alpha;\delta\beta)).
$$

To complete the proof we need gluings  $D_{\epsilon,\mu,\nu}$  to  $C_{\pi,\omega}$ . If the cylinder is glued to any of the inner circles, the relation for composition of the crossings yields the equality. So let us only consider gluings along the outer circle. Further, if we consider  $C_{-+}$  or  $C_{+-}$  we can use compatibility of  $m, \Delta$  with the crossing or invariance of  $\eta$ , coev under the crossing to prove the equality of the two sides. Hence we concentrate our efforts on the case, where  $\pi = \omega$ 

• 
$$
\epsilon = \mu = \nu = -, \pi = \omega = +:
$$

$$
\mathcal{Z}\left(D_{-,-,-}(\alpha,\beta;\rho,\delta)\coprod C_{++}(\gamma;\xi)\right) =
$$
\n
$$
= (\eta_{\gamma^{-1}} \otimes L_{\xi\gamma^{-1}\xi^{-1}})(L_{\gamma^{-1}} \otimes \sigma)(L_{\gamma^{-1}} \otimes \phi_{\xi}^{\gamma^{-1}} \otimes L_{\gamma})(L_{\gamma^{-1}} \otimes \operatorname{coev}_{\gamma})m_{\rho\alpha\rho^{-1},\delta\beta\delta^{-1}}(\phi_{\rho}^{\alpha} \otimes \phi_{\delta}^{\beta})
$$
\n
$$
= \phi_{\xi}^{\gamma^{-1}}(\eta_{\gamma^{-1}} \otimes L_{\gamma})(L_{\gamma^{-1}} \otimes \operatorname{coev}_{\gamma^{-1}})m_{\rho\alpha\rho^{-1},\delta\beta\delta^{-1}}(\phi_{\rho}^{\alpha} \otimes \phi_{\delta}^{\beta})
$$
\n
$$
= \phi_{\xi}^{\gamma^{-1}}m_{\rho\alpha\rho^{-1},\delta\beta\delta^{-1}}(\phi_{\rho}^{\alpha} \otimes \phi_{\delta}^{\beta})
$$
\n
$$
= m_{\xi\rho\alpha\rho^{-1}\xi^{-1},\xi\delta\beta\delta^{-1}\xi^{-1}}(\phi_{\xi\rho}^{\alpha} \otimes \phi_{\xi\delta}^{\beta})
$$
\n
$$
= \mathcal{Z}(D_{-,-,+}(\alpha,\beta;\xi\rho,\xi\delta)).
$$

$$
\bullet \ \epsilon = \mu = -, \nu = +, \pi = \omega = -:
$$

$$
\mathcal{Z}\left(D_{-,-,+}(\alpha,\beta;\rho,\delta)\coprod C_{--}(\gamma;\xi)\right) =
$$
\n
$$
= \eta_{\gamma}(L_{\gamma}\otimes\phi_{\xi^{-1}}^{\xi\gamma^{-1}\xi^{-1}})(m_{\rho\alpha\rho^{-1},\delta\beta\delta^{-1}}\otimes L_{\xi\gamma^{-1}\xi^{-1}})(\phi_{\rho}^{\alpha}\otimes\phi_{\delta}^{\beta}\otimes L_{\xi\gamma^{-1}\xi^{-1}})
$$
\n
$$
= \eta_{\xi\gamma\xi^{-1}}(\phi_{\xi}^{\gamma}\otimes L_{\xi\gamma^{-1}\xi^{-1}})(m_{\rho\alpha\rho^{-1},\delta\beta\delta^{-1}}\otimes L_{\xi\gamma^{-1}\xi^{-1}})(\phi_{\rho}^{\alpha}\otimes\phi_{\delta}^{\beta}\otimes L_{\xi\gamma^{-1}\xi^{-1}})
$$
\n
$$
= \eta_{\xi\gamma\xi^{-1}}(m_{\xi\rho\alpha\rho^{-1}\xi^{-1},\xi\delta\beta\delta^{-1}\xi^{-1}}\otimes L_{\xi\gamma^{-1}\xi^{-1}})(\phi_{\xi\rho}^{\alpha}\otimes\phi_{\xi\delta}^{\beta}\otimes L_{\xi\gamma^{-1}\xi^{-1}})
$$
\n
$$
= \mathcal{Z}(D_{-,-,-}(\alpha,\beta;\xi\rho,\xi\delta)).
$$

$$
\epsilon = \mu = + = \nu = +, \pi = \omega = -:
$$
\n
$$
\mathcal{Z}\left(D_{+,+,+}(\alpha,\beta;\rho,\delta)\coprod C_{--}(\gamma;\xi)\right) =
$$
\n
$$
= (L_{\alpha} \otimes L_{\beta} \otimes \eta_{\gamma})(\phi_{\delta^{-1}}^{\beta\beta\delta^{-1}} \otimes \phi_{\rho^{-1}}^{\rho\alpha\rho^{-1}} \otimes L_{\gamma} \otimes \phi_{\xi^{-1}}^{\xi\gamma^{-1}\xi^{-1}})
$$
\n
$$
(\Delta_{\delta\beta\delta^{-1},\rho\alpha\rho^{-1}} \otimes L_{\gamma} \otimes L_{\xi\gamma^{-1}\xi^{-1}})(\operatorname{coev}_{\gamma} \otimes L_{\xi\gamma^{-1}\xi^{-1}})
$$
\n
$$
= (\phi_{\delta^{-1}}^{\delta\beta\delta^{-1}} \otimes \phi_{\rho^{-1}}^{\rho\alpha\rho^{-1}})\Delta_{\delta\beta\delta^{-1},\rho\alpha\rho^{-1}}(L_{\gamma^{-1}} \otimes \eta_{\gamma})(\operatorname{coev}_{\gamma} \otimes L_{\gamma^{-1}})\phi_{\xi^{-1}}^{\xi\gamma^{-1}\xi^{-1}}
$$
\n
$$
= (\phi_{\delta^{-1}}^{\delta\beta\delta^{-1}} \otimes \phi_{\rho^{-1}}^{\rho\alpha\rho^{-1}})\Delta_{\delta\beta\delta^{-1},\rho\alpha\rho^{-1}}\phi_{\xi^{-1}}^{\xi\gamma^{-1}\xi^{-1}}
$$
\n
$$
= (\phi_{(\xi\delta)^{-1}}^{\xi\delta\beta\delta^{-1}\xi^{-1}} \otimes \phi_{(\xi\rho)^{-1}}^{\xi\alpha\rho^{-1}\xi^{-1}})\Delta_{\xi\delta\beta\delta^{-1}\xi^{-1},\xi\rho\alpha\rho^{-1}\xi^{-1}}
$$
\n
$$
= \mathcal{Z}(D_{+,+,-}(\alpha,\beta;\xi\rho,\xi\delta)).
$$
\n
$$
\epsilon = \mu = + = \nu = -, \pi = \omega = +:
$$
\n
$$
\mathcal{Z}\left(D_{+,+,-}(\alpha,\beta;\rho,\delta)\coprod C_{++}(\gamma;\xi)\right) =
$$
\n
$$
= (L_{\xi\gamma^{-1}\xi^{-1}} \otimes \phi_{\delta^{-1}}^{\delta\beta\delta^{-1}} \otimes \phi_{\rho^{-1}}^{\rho\alpha\rho^{-1}})(L_{\xi\gamma^{-1}\xi} \otimes \Delta_{
$$

<span id="page-29-0"></span>For arbitrary  $(X, Y)$ -surfaces we conclude the proof analogously to the proof of Turaev [\[11,](#page-30-9) Thm 3.1. stating the independence of the splitting system of loops.  $\square$ 

### 7. OUTLOOK

The structure of a crossed loop Frobenius  $(\mathcal{G}, \mathcal{V})$ -category appears somehow incomplete, since it uses the full groupoid for the crossings, however only loops on the indices of the objects. An attempt to solve this asymmetry is to define a category of open/closed 2-dimensional  $(X, Y)$ -cobordisms, whose objects are given by open or closed strings together with appropriate homotopy classes of maps to  $Y$ . In order to define morphisms in this category, one needs to consider 2-manifolds with corners with additional data. Given such a symmetric monoidal category  $\mathsf{Cob}^{X,Y}_{2,oc}$ , one might define *open/closed HQFTs* as symmetric monoidal functors  $\mathcal{Z}$  :  $\mathsf{Cob}_{2,oc}^{X,Y} \to \mathcal{V}$ . For  $X = Y = \{*\}$  this theory should reduce to open/closed TQFTs, as defined by Lauda and Pfeiffer in [\[5\]](#page-30-10), who classified them by knowledgeable Frobenius algebras. For general target pairs of spaces, one therefore can expect the resulting algebraic structure to be some sort of *crossed knowledgable Frobenius*  $(\mathcal{G}, \mathcal{V})$ -category. The study of such a theory of open/closed HQFTs will be subject to an upcoming paper.

## <span id="page-30-0"></span>**REFERENCES**

- <span id="page-30-2"></span>[1] L. Abrams, Two-dimensional topological quantum field theories and Frobenius algebras, *Journal of Knot Theory and Its Ramifications* 5 (5), (1996), 569–587.
- <span id="page-30-8"></span><span id="page-30-3"></span>[2] M. Brightwell and P. Turner, Representations of the homotopy surface category of a simply connected space, *Journal of Knot theory and its Ramifications* 9 (7), (2000), 855–864.
- [3] C. Kassel, Quantum Groups, *Springer-Verlag*, New York 1995.
- <span id="page-30-10"></span>[4] J. Kock, Frobenius Algebras and 2D Topological Quantum Field Theories, *Cambridge University Press*, Cambridge (2003).
- <span id="page-30-7"></span>[5] A. Lauda and H. Pfeiffer, Open-closed strings: Two-dimensional extended TQFTs and Frobenius algebras, *Topology and its applications* 155 (7), (2008), 623–666.
- <span id="page-30-4"></span>[6] S. Mac Lane, Categories for the Working Mathematician, Graduate Texts in Mathematics 5, *Springer-Verlag*, New York 1998.
- <span id="page-30-6"></span>[7] T. Porter and V. Turaev, Formal Homotopy Quantum Field Theories, I: Formal Maps and Crossed C-algebras, *Journal of Homotopy and Related Strucutures* 3 (1), (2006).
- <span id="page-30-5"></span>[8] G. Rodrigues, Homotopy Quantum Field Theories and the Homotopy Cobordism Category in Dimension 1 + 1, *J. Knot Theory and its Ramifications* 12, (2003), 287–317.
- [9] M.D. Staic and V. Turaev, Remarks on 2-dimensional HQFT's, *Algebraic & Geometric Topology* 10, (2010), 1367–1393.
- <span id="page-30-1"></span>[10] V. Turaev, Homotopy field theory in dimension 2 and group-algebras, *<http://arxiv.org/abs/math/9910010>*.
- <span id="page-30-9"></span>[11] V. Turaev, Homotopy Quantum Field Theory, EMS Tracts in Mathematics, 2010.

Paul Großkopf, Mathematical Institute, University of Oxford, United Kingdom *Email address*: paul.grosskopf@gmx.at