# LEFSCHETZ DEFECT IN FAMILIES

MATTEO VERNI

ABSTRACT. We provide a cohomological characterization of the Lefschetz defect of smooth complex projective varieties. As a consequence, we deduce that the Lefschetz defect of a smooth Fano variety is invariant under smooth deformation. We also characterize the Lefschetz defect of an abelian variety in terms of its isogeny factors, and study it in families.

## 1. INTRODUCTION

In [Cas12], the author introduces the Lefschetz defect  $\delta_X$  (see Definition 2.1) for X a smooth Fano variety. It measures how many "special" steps occur in runs of the MMP of (X, -D) for all effective divisors D. This invariant has proven useful in the effort to classify Fano fourfolds: for the latest developments in this direction, see [Cas24].

Since there are finitely many deformation families of Fano manifolds in each dimension, to classify these varieties one is particularly interested in those invariants that are preserved under deformation. One is then naturally led to ask whether the Lefschetz defect is such an invariant. Previously, this had been established in the case  $\delta_X = 2$  and  $\rho_X = 3$  as an indirect consequence of the classification result of [Sec23].

In the present work, our main result is that the Lefschetz defect is indeed deformation invariant.

**Theorem 1.1.** Let  $X_1$  and  $X_2$  be two smooth Fano varieties which are deformation equivalent via a smooth deformation. Then  $\delta_{X_1} = \delta_{X_2}$ .

It follows that the strong characterization results of [Cas12], [CD15], [CRS22] and [Cas24] hold throughout each deformation family.

On the way to the proof we make the following remark of independent interest (see Section 2.1 for definitions).

**Proposition 1.2.** For any linearly equivalent divisors D and D' on a smooth projective variety, we have

$$\delta_X(D) = \delta_X(D').$$

In the second part of the paper, we begin to study the Lefschetz defect of smooth projective varieties other than Fano. The second main result precisely relates the Lefschetz defect of an abelian variety with its isogeny factors and their multiplicity.

### MATTEO VERNI

**Theorem 1.3.** Let A be an abelian variety of dimension n. Then  $\delta_A = 0$  if and only if A does not contain elliptic curves nor abelian surfaces of Picard number greater than 1. If instead  $\delta_A > 0$ , then exactly one of the following is verified:

•  $A \sim C \times E$  where E is an elliptic curve, and if  $k_E \geq 1$  is its multiplicity as an isogeny factor of A then

$$\delta_A = \begin{cases} k_E & \text{if } E \text{ does not have } CM \\ 2k_E - 1 & \text{if } E \text{ has } CM \end{cases}$$

•  $A \sim C \times S$  where S is a simple abelian surface of type II,  $\rho_S = 3$  and

$$\delta_A = 2.$$

Furthermore, any elliptic curve showing up as an isogeny factor of A does so with multiplicity at most one.

•  $A \sim C \times S'$  where S' is a simple abelian surface of type I or IV,  $\rho_{S'} = 2$ and

$$\delta_A = 1.$$

Furthermore, there are no elliptic curves contained in A.

This result can also be seen as one possible way of addressing the problem brought up in the introduction of [Cas23, Section 8], namely to find splitting results in the spirit of [Cas12, Th. 3.3.], outside the smooth Fano setting. This was done for mildly singular Fano varieties in [Noc12], and our result provides an instance completely outside the Fano context.

1.1. Outline of the article. In Section 2 we recall basic definitions and remarks about the Lefschetz defect. In Section 2.2 we prove a generalization (Proposition 2.4) of [Voi92, Lemme 1.5], from which we deduce that  $\delta_X(D)$  depends only on the cohomological class [D] and not on D itself. This provides a purely cohomological characterization of  $\delta_X$  to be used throughout, and it proves Proposition 1.2.

In Section 3 we collect the supporting results and give the proof of Theorem 1.1: the necessary ingredients are Corollary 2.5 and a result on deformation invariance of the cone of effective divisors, which was established at increasing levels of generality in the works [Siu02], [DH11] and [Tot12].

In Section 4 we make the link between Lefschetz defect and isogeny factors of an abelian variety, and prove Theorem 1.3. In subsection 4.1, we deduce from Theorem 1.3 that the Lefschetz defect is "stable" under deformations in the abelian setting too, however here "stable" should be taken with more care as in the presence of nontrivial Noether-Lefschetz loci on the moduli space, one can only hope for invariance in a more subtle sense.

Acknowledgements. I would first like to thank my advisors Claire Voisin and Emanuele Macrì, for sharing their knowledge and for their invaluable advice on how to write an article. In particular, I thank Claire Voisin for the suggestion of adapting her result of [Voi92] to the study of  $\delta_X$ . I would like to thank Saverio

Secci for explaining to me the previously known case of the invariance result, and also Angel Rios-Ortiz and Enrico Lampetti for helpful conversations.

This work was supported by the ERC Synergy Grant HyperK (Grant agreement No. 854361)

# 2. Lefschetz Defect

2.1. Basic definitions and remarks. In this section, X will always denote a smooth projective variety, if not specified otherwise. We first recall general definitions and remarks, and then establish the characterization (8).

For any algebraic variety Y,  $N_1(Y)_{\mathbb{Q}}$  denotes the  $\mathbb{Q}$ -vector space generated by all integral curves  $C \subset Y$  modulo numerical equivalence. Proper morphisms  $Y \to Z$  induce functorially linear maps  $N_1(Y)_{\mathbb{Q}} \to N_1(Z)_{\mathbb{Q}}$  in the obvious way.

**Definition 2.1.** Let  $i: D \hookrightarrow X$  be the embedding of an integral divisor. We define

$$\delta_X(D) := \operatorname{codim}_{N_1(X)_{\mathbb{Q}}} i_* N_1(D)_{\mathbb{Q}}.$$

We define the **Lefschetz defect of** X as the number

(1) 
$$\delta_X := \max_{\substack{D \subset X \\ \text{integral}}} \delta_X(D)$$

For a detailed survey on the Lefschetz defect and its applications, see [Cas23].

We observe that we could have taken the maximum in (1) over all (possibly nonreduced, possibly reducible) effective divisors. Firstly, for any nonreduced divisor D, one has  $\delta_X(D) = \delta_X(D_{red})$ . Indeed, when computing  $\dim_{\mathbb{Q}} i_*N_1(D)_{\mathbb{Q}}$ we are considering integral curves inside X: since the embedding of a reduced subscheme only depends on the topological embedding, the class of a curve  $C \subset$ D does not care about the nonreduced structure of D, the only thing that matters is the topological subspace  $|C| \subset |D| \subset |X|$ . Moreover, if  $D = D_1 \cup D_2$ , one sees that  $\delta_X(D) \leq \delta_X(D_1)$  since all curves lying on  $D_1$  are also lying on D.

We conclude that

(2) 
$$\delta_X = \max_{D \subset X} \delta_X(D).$$

Denoting the Picard rank of X by  $\rho_X$ , we remark that

$$(3) 0 \le \delta_X \le \rho_X - 1.$$

Indeed,  $\delta_X \neq \rho_X$  since any integral curve *C* on a divisor *D* is intersected nontrivially by hyperplane sections of *X*, so [*C*] represents a nontrivial class of  $i_*N_1(D)_{\mathbb{Q}}$ .

Another situation where  $\delta_X(D)$  is easily computed is the following. Let  $D \subset X$ be as above and suppose there exists a retraction  $X \to D$ , i.e., a morphism of varieties which is a left-inverse to the inclusion morphism  $D \hookrightarrow X$ . Then  $X \to D$ induces on numerical 1-cycles a retraction to the map  $N_1(D, \mathbb{Q}) \to N_1(X, \mathbb{Q})$ , which then must be injective. We conclude that

(4) 
$$\delta_X(D) = \rho_X - \rho_D.$$

### MATTEO VERNI

**Remark 2.2.** [Cas23, Remark 3] For any complex projective variety X,  $N_1(X)^{\vee}_{\mathbb{Q}}$  is naturally identified with  $N^1(X)_{\mathbb{Q}}$ . Moreover the latter injects into  $H^2(X, \mathbb{Q})$ , as homological equivalence of divisors is equivalent to numerical equivalence up to torsion. Thus we obtain a more cohomological formulation of  $\delta_X(D)$ , namely

(5) 
$$\delta_X(D) = \dim \operatorname{Ker} \left( \eta^* : N^1(X)_{\mathbb{Q}} \to H^2(D, \mathbb{Q}) \right)$$

Note also that in case  $H^2(X, \mathcal{O}_X) = 0$ , we have the diagram

which implies

$$\delta_X(D) = \dim \operatorname{Ker} \left( \eta^* : H^2(X, \mathbb{Q}) \to H^2(D, \mathbb{Q}) \right)$$

2.2. Another cohomological characterization of  $\delta_X(D)$ . For X a compact complex manifold, recall that there exists a cycle class map

$$CH^k(X) \to H^{2k}(X, \mathbb{Q})$$
  
 $Z \mapsto [Z]$ 

Recall that for any closed analytic subvariety  $\sigma: V \hookrightarrow X$ , [V] can be defined via any resolution of singularities  $\widetilde{V} \xrightarrow{\tau} V$  by pushing forward the fundamental class. More precisely, setting  $\widetilde{\sigma} := \sigma \circ \tau$ 

$$[V] = \sigma_*(\underline{1}_{\widetilde{Z}}) = PD_X^{-1} \circ (\widetilde{\sigma}^*)^{\vee} \circ PD_{\widetilde{Z}}(\underline{1}_{\widetilde{Z}})$$

where  $PD_X : H^k(X, \mathbb{Q}) \to H^{2n-k}(X, \mathbb{Q})^{\vee}$  are Poincaré duality morphisms and

$$\underline{1}_{\widetilde{Z}} \in H^0(\widetilde{Z}, \mathbb{Q}) \simeq \mathbb{Q}$$

is the generator associated to a given choice of orientation.

Remarkably, the subspace of cohomology classes on a Kähler manifold X that are trivial when restricted to a submanifold V depends only on the cohomology class [V]. More precisely, one has

**Proposition 2.3.** [Voi92, Lemme 1.5] Let Y be a Kähler manifold and W a submanifold of codimension c. Then

$$\operatorname{Ker}\left(H^{2}(Y,\mathbb{R})\xrightarrow{\sigma^{*}}H^{2}(W,\mathbb{R})\right) = \operatorname{Ker}\left(H^{2}(Y,\mathbb{R})\xrightarrow{\smile[W]}H^{2+2c}(Y,\mathbb{R})\right).$$

Note that this statement fails for cohomology of higher degrees, as explained by Voisin in Appendix B to [SY22, Remark B.3].

We can in fact prove a generalization of the above proposition which works for certain singular subvarieties. Its proof follows in part that of [Voi92], with adequate adjustments.

**Proposition 2.4.** Let X be a smooth projective variety over  $\mathbb{C}$  of dimension n and  $\sigma : V \hookrightarrow X$  a locally complete intersection subvariety of dimension  $d \ge 2$ and codimension c = n - d. Then

(6) 
$$\operatorname{Ker}\left(H^{2}(X,\mathbb{Q}) \xrightarrow{\sigma^{*}} H^{2}(V,\mathbb{Q})\right) = \operatorname{Ker}\left(H^{2}(X,\mathbb{Q}) \xrightarrow{\smile [V]} H^{2+2c}(X,\mathbb{Q})\right).$$

It follows immediately that we also have the equality

(7) 
$$\operatorname{Ker}\left(N^{1}(X,\mathbb{Q}) \xrightarrow{\sigma^{*}} H^{2}(V,\mathbb{Q})\right) = \operatorname{Ker}\left(N^{1}(X,\mathbb{Q}) \xrightarrow{\smile[V]} H^{2+2c}(X,\mathbb{Q})\right).$$

Using (5) we deduce from (7) the following.

**Corollary 2.5.** Let  $D \subset X$  be a divisor in X smooth complex projective variety. Then

(8) 
$$\delta_X(D) = \dim \operatorname{Ker}\left(N^1(X, \mathbb{Q}) \xrightarrow{\smile [D]} H^4(X, \mathbb{Q})\right).$$

In particular, Corollary 1.2 holds.

For the proof of Proposition 2.4 we will use the following results on the Betti cohomology of complex algebraic varieties:

**Theorem 2.6.** For Y a locally complete intersection subvariety of  $\mathbb{P}^N$ , and  $H \cap Y$  a general hyperplane section of Y, one has that

$$H^k(Y,\mathbb{Q}) \to H^k(H \cap Y,\mathbb{Q})$$

is an isomorphism for  $k < \dim(Y) - 1$  and injective for  $k = \dim(Y) - 1$ .

*Proof.* This is the theorem "LHT for singular spaces" at page 24 of [GM88], applied to the locally complete intersection case.  $\Box$ 

**Theorem 2.7.** [Del74, Proposition 8.2.7] Let  $\varphi : Y \to X$  be a proper morphism with X smooth projective. Let  $\tau : \widetilde{Y} \to Y$  be a desingularization and let  $\widetilde{\varphi} := \varphi \circ \tau : \widetilde{Y} \to X$ . Then for any k,

$$\operatorname{Ker}\left(\varphi^*: H^k(X, \mathbb{Q}) \to H^k(Y, \mathbb{Q})\right) = \operatorname{Ker}\left(\widetilde{\varphi}^*: H^k(X, \mathbb{Q}) \to H^k(\widetilde{Y}, \mathbb{Q})\right).$$

Proof of Proposition 2.4. Consider a resolution of singularities  $\widetilde{V} \xrightarrow{\tau} V$  and denote the induced composite morphism  $\widetilde{V} \to X$  by  $\widetilde{\sigma} := \sigma \circ \tau$ . One inclusion in (6) is easy: by the projection formula,

$$\sim [V] = \widetilde{\sigma}_* \circ \widetilde{\sigma}^* = \widetilde{\sigma}_* \tau^* \sigma^*,$$

thus

$$\operatorname{Ker} \sigma^* \subset \operatorname{Ker} (\smile [V]).$$


We are left to prove the opposite inclusion, which we break down into steps.

Step 1: We first reduce to the case where V is a surface. Let  $\alpha \in \text{Ker}(\smile [V])$ , and let h be the ample class associated to an ample divisor H. We denote by S

the surface we obtain from V by taking a general H-hyperplane section a total of d-2 times. Let us denote by j the embedding of S in V, and set  $\eta = \sigma \circ j$ . We have  $[S] = h^{d-2} \smile [V]$ , hence

$$\alpha \smile [S] = 0.$$

We have the following commutative diagram



where  $j^*$  is injective by Theorem 2.6. In order to conclude that  $\sigma^* \alpha = 0$  it thus suffices to prove the implication

$$\alpha \smile [S] = 0 \Rightarrow \eta^* \alpha = 0$$

which amounts exactly to the d = 2 case.

Step 2: Let  $\pi: \widetilde{S} \to S$  be a resolution of singularities and let  $\widetilde{\eta} := \eta \circ \pi$ . In this step we prove that

$$\alpha \smile [S] \Rightarrow \widetilde{\eta}^* \alpha = 0$$

The argument will follow very closely the one of [Voi92], which proves Proposition 2.3 for any d. However, note that it would not work here if we had not reduced to d = 2.

Consider the usual nondegenerate quadratic form on  $H^2(\widetilde{S}, \mathbb{Q})$ 

$$q(\gamma,\nu):=\int_{\widetilde{S}}\gamma\smile\nu$$

For any  $\alpha, \beta \in H^2(X, \mathbb{Q})$  we have

$$q(\widetilde{\eta}^*\alpha, \widetilde{\eta}^*\beta) := \int_X \alpha \smile \beta \smile [S],$$

from which we see that  $\alpha \smile [S] = 0$  implies  $\tilde{\eta}^* \alpha \in \operatorname{Ker} q_{|\operatorname{Im} \tilde{\eta}^*}$ . This means it suffices to prove that the restriction  $q_{|Im \tilde{\eta}^*}$  is nondegenerate.

The signature of q is described by the Hodge index theorem. The real cohomology  $H^2(\widetilde{S}, \mathbb{R})$  decomposes as

$$H^{2}(\widetilde{S},\mathbb{R}) = (H^{2,0}(\widetilde{S}) \oplus H^{0,2}(\widetilde{S}))_{\mathbb{R}} \oplus H^{1,1}(\widetilde{S})_{\mathbb{R}},$$

where the direct sum is orthogonal for q and the subscript  $\mathbb{R}$  designates the subspace of real cohomology classes. q is positive definite on the first subspace and it has signature  $(+, -, \ldots, -)$  on the second subspace. The real cohomology  $H^2(X, \mathbb{R})$  has a similar decomposition and the morphism  $\tilde{\eta}^*$ , being a morphism of Hodge structures, preserves such a decomposition. It follows that

$$\operatorname{Im} \widetilde{\eta}^* = \operatorname{Im} \widetilde{\eta}^{*(2,0)+(0,2)} \oplus \operatorname{Im} \widetilde{\eta}^{*(1,1)}.$$

Hence it suffices to prove that  $q_{|\text{Im }\tilde{\eta}^{*}(2,0)+(0,2)}$  and  $q_{|\text{Im }\tilde{\eta}^{*}(1,1)}$  are nondegenerate. For the first one this is obvious because  $q_{(H^{2,0}(\tilde{S})\oplus H^{0,2}(\tilde{S}))_{\mathbb{P}}}$  is positive definite. For the second one, this follows from the fact that the image contains the big and nef class  $\tilde{\eta}^* h$ , which thus has positive self intersection, and from the signature of q on  $H^{1,1}(\tilde{S})_{\mathbb{R}}$ .

Step 3: We are left to prove the implication

$$\widetilde{\eta}^* \alpha = 0 \Rightarrow \eta^* \alpha = 0$$

But this is exactly the content of Theorem 2.7, hence we are done.

## 3. Lefschetz defect in families of Fano Manifolds

In this section we focus on the Fano manifold case: it is restricted to this context that the Lefschetz defect has been introduced, because of its interpretation in terms of MMP runs ([Cas23, Section 3]). The advantage of the Fano hypothesis for our study of the behaviour in families is that the cohomology of these variety is more closely related to its Picard groups; the first varies well with deformations, while the second is the one to which the Lefschetz defect heavily depends on.

Throughout this section, X will denote a smooth Fano variety. First, note that  $H^2(X, \mathcal{O}_X) = 0$ , thus  $N^1(X)_{\mathbb{Q}} = H^2(X, \mathbb{Q})$ . This means (8) becomes

(9) 
$$\delta_X(D) = \dim \operatorname{Ker} \left( H^2(X, \mathbb{Q}) \xrightarrow{\sim [D]} H^4(X, \mathbb{Q}) \right)$$

We consider a **smooth deformation family of** X **over** B: this means a smooth morphism  $f: \mathcal{X} \to B$ , with B connected, for which the relative canonical is an anti-ample line bundle relatively over the base, making each fiber a smooth Fano variety, together with a chosen point  $0 \in B$  such that  $\mathcal{X}_t \simeq X$ . We denote such data as an arrow  $f: \mathcal{X} \to (B, 0)$ .

**Question 3.1.** How does  $\delta_{\mathcal{X}_t}$  vary with t?

For  $\rho_X = 1$  or dim  $X \leq 2$ ,  $\delta_X$  is trivially deformation invariant. The first is clear from the bounds  $0 \leq \delta_X \leq \rho_X - 1$  of (3), and the fact that  $\rho_X$  stays constant for smooth deformations of varieties with  $h^{2,0} = 0$ . This also covers the case dim X = 1 since then  $X \simeq \mathbb{P}^1$ . If instead X is a del Pezzo surface, then any integral curve C has  $N_1(C)_{\mathbb{R}} = \mathbb{R}[C]$  hence  $\delta_X = \delta_X(D) = \rho_X - 1$ 

3.1. Extending line bundles. Ideally, we want to spread effective divisors from one fiber to a whole family of effective divisors over the base. This in general is only possible up to basechanging the family: as long as the basechange is surjective, this does not pose any problem towards our final goal of comparing specific fibers  $X_t$  and  $X_s$ , as we do not care in which parameter space t and s live.

The first step is easy: it is well known that up to surjective basechange one can spread a line bundle on a fiber to a line bundle over the entire total space of the family, whenever the relative Picard scheme is smooth over its base.

**Theorem 3.2.** Let  $f: \mathcal{X} \to B$  be a smooth projective morphism with integral fibers with B smooth connected quasi-projective over a field k, and assume  $h^2(\mathcal{X}_t, \mathcal{O}_{\mathcal{X}_t}) = 0$  for all  $t \in B$ . Let  $0 \in B$  be a point and  $L \in \operatorname{Pic}_{\mathcal{X}_0}$ . Then there exists a projective, generically étale morphism  $(B', 0') \xrightarrow{\tau} (B, 0)$  with B' smooth connected and a line bundle  $\mathcal{L}$  on  $\mathcal{X}_{B'} = \mathcal{X} \times_B B'$  such that  $\mathcal{L}_{|\mathcal{X}_{0'}} \simeq O_{\mathcal{X}_0}(D)$ .

Proof. This is well known, following from the fact that by [FGI<sup>+</sup>05, Theorem 9.4.8] the relative Picard scheme  $\operatorname{Pic}_{\mathcal{X}/B} \xrightarrow{\pi} B$  exists and by [FGI<sup>+</sup>05, Proposition 9.5.19] it is smooth. Finding the line bundle  $\mathcal{L}$  as requested amounts to finding a basechange  $B' \to B$  such that  $\pi'_B \colon \operatorname{Pic}_{\mathcal{X}_{B'}/B'} \to B'$  has a section passing through the fiber of the projection  $\operatorname{Pic}_{\mathcal{X}_{B'}/B'} \to \operatorname{Pic}_{\mathcal{X}/B}$  above the point  $\mathcal{O}_{\mathcal{X}_0}(D) \in \operatorname{Pic}_{\mathcal{X}/B}$ . This can be done, for example by looking at the Henselianization of the local ring at  $0 \in B$ , from which one can lift morphisms along any smooth morphism. This lifting property then holds for a smaller, quasi-finite étale neighbourhood of  $U \to B$  of  $0 \in B$ , and one concludes by passing to a Nagata compactification  $U \subset B' \to B$ .

3.2. Effectivity of Cartier divisors in a family. By the previous section we can extend (up to basechange) a Cartier divisor D on one fiber  $\mathcal{X}_0$  to a Cartier divisor  $\mathcal{D}$  over the entire domain  $\mathcal{X}$ . What is not at all clear a priori is whether or not the other fibers  $\mathcal{D}_t$  will be effective in case  $D = \mathcal{D}_0$  is effective. Luckily, this happens to be true in the setting we are interested in, that of families of smooth Fano manifolds.

**Theorem 3.3.** Let  $\mathcal{X} \to B$  be smooth family of Fano varieties, with B a smooth curve. There exists a smooth curve B' and a surjective morphism  $B' \to B$  such that, setting  $\mathcal{X}' := \mathcal{X}_{B'}$ , for any  $\mathcal{L} \in \operatorname{Pic}(\mathcal{X}')$  one has that  $h^0(\mathcal{X}'_s, \mathcal{L}_{|\mathcal{X}'_s})$  is independent of s. In particular, if  $\mathcal{L}_{|\mathcal{X}_0}$  defines an effective divisor then so does  $\mathcal{L}_{|\mathcal{X}_t}$ .

*Proof.* This is an immediate application of the much more general theorem [Tot12, Theorem 6.1]. As the author himself mentions, the case of his result we need here is less general and was already present in the literature beforehand, see the works [DH11] and [Siu02].  $\Box$ 

3.3. Invariance of  $\delta_X$  for Fano families. In this section we prove the main Theorem 1.1, namely that the Lefschetz Defect of a smooth Fano variety is invariant under smooth deformation.

Before beginning the proof, recall that for any complex manifold X there exists a natural map

$$\operatorname{Pic}(X) \to H^2(X, \mathbb{Z}).$$

which sends L to its first chern class  $c_1(L)$ . When X compact, for any D effective Cartier divisor one can define [D] as discussed in Section 2.2: then one has that  $[D] = c_1(\mathcal{O}_X(D)).$ 

Proof of Theorem 1.1. It suffices to show  $\delta_{X_1} \leq \delta_{X_2}$ . Let  $D_1 \subset X_1$  be an effective Cartier divisor such that  $\delta_{X_1}(D) = \delta_{X_1}$ . Let  $\mathcal{X} \to B$  be a smooth family and

 $t_1, t_2 \in B$  such that  $\mathcal{X}_{t_i} \simeq X_i$ . Up to basechanging we may assume that the conclusions of Proposition 3.2 and Theorem 3.3 hold. Then there is some  $\mathcal{D}$  Cartier on  $\mathcal{X}$  such that  $\mathcal{D}_{|\mathcal{X}_{t_1}} \simeq D_1$ . Since  $D_1$  is effective then  $D_2 := \mathcal{D}_{|\mathcal{X}_{t_2}}$  is effective too, hence  $\delta_{X_2}(D_2) \leq \delta_{X_2}$ 

Fix some  $t \in B$ . For a small enough  $U \subset B$  contractible analytic neighbourhood of t, the submersion  $\mathcal{X}_U \to U$  is  $C^{\infty}$ -trivial. In particular the restriction morphism  $H^2(\mathcal{X}_U, \mathbb{Q}) \to H^2(\mathcal{X}_s, \mathbb{Q})$  is an isomorphism for all  $s \in U$ . By the functoriality of taking first chern classes of line bundles on complex manifolds,  $c_1(\mathcal{O}(\mathcal{D}))|_{\mathcal{X}_S} = c_1(\mathcal{O}(\mathcal{D}|_{\mathcal{X}_s})) = [\mathcal{D}|_{\mathcal{X}_s}]$  for all  $s \in U$ . In particular, for any  $s \in U$ we have the following commutative diagram

In particular, the kernel of the left and right vertical maps are identified via the isomorphism in cohomology induced by the diffeomorphism of the fibers  $\mathcal{X}_t \simeq \mathcal{X}_s$ . By (9), we can thus conclude that  $\delta_{\mathcal{X}_t}(D_1) = \delta_{\mathcal{X}_s}(\mathcal{D}_s)$ . This shows  $s \mapsto \delta_{\mathcal{X}_s}(\mathcal{D}_s)$  is locally constant, around any  $t \in B$ . Since B is connected, we obtain that  $\delta_{\mathcal{X}_{t_1}}(D_1) = \delta_{\mathcal{X}_{t_2}}(D_2)$ .

Putting everyting together, we conclude

$$\delta_{X_1} = \delta_{X_1}(D_1) = \delta_{X_2}(D_2) \le \delta_{X_2}.$$

# 4. Lefschetz defect on Abelian varieties

In this section, we study the number  $\delta_A$  in the case A is an abelian variety. In particular, for any effective divisor  $D \subset A$ , we observe that the number  $\delta_A(D)$  behaves in a way that heavily depends on the *Itaka dimension* of the line bundle  $\mathcal{O}_A(D)$ .

Recall the definition of Itaka dimension: for L a line bundle on a normal algebraic variety X,

$$\kappa(X,L) := \min\left\{ d \in \mathbb{N} \mid \frac{h^0(X,L^{\otimes m})}{m^d} \text{ bounded as } m \to \infty \right\}.$$

If we denote by  $\phi_{L^{\otimes m}} : X \dashrightarrow \mathbb{P}(H^0(X, L^{\otimes m}))$  the rational map induced by the linear system  $|L^{\otimes m}|$ , and by  $W_{L^{\otimes m}} \subset \mathbb{P}(H^0(X, L^{\otimes m}))$  the closure of  $\operatorname{Im} \phi_{L^{\otimes m}}$ , then

(10) 
$$\kappa(X,L) = \dim W_m \quad \text{for} \quad m >> 0.$$

Let us fix a complex abelian variety A of dimension n. Then the Itaka dimension of the line bundle  $\mathcal{O}_A(D)$  for  $D \subset A$  an effective divisor manifests in a very explicit geometric way, which we now recall. It is a remarkable basic fact about abelian varieties that for any effective divisor  $D \subset A$ , the linear system |2D| is basepoint-free. Moreover, if

$$A \xrightarrow{\varphi} B \xrightarrow{\psi} \mathbb{P}(|2D|)$$

is the Stein factorization of  $\phi_{2D}$ , then *B* itself is an abelian variety (by Proposition, page 88 of [Mum08]). Since  $\psi$  is finite, by (10) we have

$$\kappa(A, \mathcal{O}(D)) = \dim B.$$

We know discuss what is the link between  $\delta_A(D)$  and dim *B*. This analysis will also show that the finer datum of the isogeny factors of *B* and their multiplicity in *A* determines  $\delta_A(D)$  even more precisely.

**Proposition 4.1.** Let A be an abelian variety,  $D \subset A$  an effective divisor and  $A \rightarrow B$  constructed as above. Set  $b := \dim B$ . Then we have one of the following cases:

- If  $b \ge 3$ , then  $\delta_A(D) = 0$ .
- If b = 2, then  $\delta_A(D) = \rho_B 1 \in \{0, \dots, 3\}$ .
- If b = 1, meaning B is an elliptic curve which appears with multiplicity  $k \ge 1$  in the factorization up to isogeny of A, then

(11) 
$$\delta_A(D) = \begin{cases} k & \text{if } B \text{ does not have } CM \\ 2k-1 & \text{if } B \text{ has } CM \end{cases}$$

Proof. Note that by the discussion preceeding (2), or a posteriori by (8), one has  $\delta_A(D) = \delta_A(2D)$ , so up to substituting D with 2D, we may suppose that D is basepoint-free. Set  $C := \text{Ker } \varphi$ ;  $\varphi$  has connected fibers so C is a connected abelian subvariety of A. By the Poincaré splitting theorem, there exists another abelian subvariety  $T \subset A$  and an isogeny  $\mu: C \times T \to A$  fitting in the commutative diagram

$$\begin{array}{ccc} C \times T & \stackrel{\mu}{\longrightarrow} & A \\ & & & \downarrow^{\varphi} \\ T & \stackrel{}{\xrightarrow{}} & D & \stackrel{}{\longrightarrow} & B & \stackrel{}{\xrightarrow{}} & \psi \end{array} \mathbb{P}(|D|) \end{array}$$

In particular, the lower horizontal map is an isogeny as well, hence the pullback along finite maps  $N := (\varphi \circ \mu)_{T}^* \psi^* \mathcal{O}_{\mathbb{P}}(1)$  is ample, and

$$pr_T^*N = \mu^* \varphi^* \psi^* \mathcal{O}_{\mathbb{P}(|D|)}(1) = \mu^* L.$$

Looking at the diagram

where the horizontal arrows are isomorphisms since induced by an isogeny, we see that the two vertical arrows have isomorphic kernels, hence by (7) we get

$$\delta_A(L) = \dim K \text{ where } K = \operatorname{Ker}\left(N^1(C \times T)_{\mathbb{R}} \xrightarrow{\smile pr^*[N]} H^4(C \times T, \mathbb{R})\right)$$

In other words, we reduced the computation of  $\delta_A(D)$  to the case where A is a product and  $\varphi$  is the projection onto one factor.

By the Kunneth formula, we have the splitting

(12) 
$$\operatorname{NS}(C \times T) = pr_C^* \operatorname{NS}(C) \oplus \operatorname{Hom}(T, C) \oplus pr_T^* \operatorname{NS}(T).$$

where the middle summand is the subgroup of Hodge classes in  $H^1(C) \otimes H^1(T)$  via the identifications

$$T \simeq \operatorname{Alb}_T$$
,  $(\operatorname{Pic}^0_C)^{\vee} \simeq C$ .

Since the map  $\sim pr_T^*[N]$  preserves the Kunneth decompositions of domain and codomain, we can write K as

$$K = (K \cap pr_C^* N^1(C)_{\mathbb{R}}) \oplus (K \cap \operatorname{Hom}(T, C)_{\mathbb{R}}) \oplus (K \cap pr_T^* N^1(T)_{\mathbb{R}}).$$

But by Hard Lefschetz for the pair (T, [N]), we have some understanding of the dimension of each summand: since dim  $T = \dim B = b$ , and N is ample on T, the map

$$\sim [N]: H^2(T, \mathbb{R}) \to H^4(T, \mathbb{R})$$

is injective as soon as  $b \geq 3$ . Similarly,  $H^1(T, \mathbb{R}) \to H^3(T, \mathbb{R})$  and  $H^0(T, \mathbb{R}) \to H^2(T, \mathbb{R})$  are injective for  $b \geq 2$  and  $b \geq 1$  respectively. Since all morphisms  $pr_T^*$  are injective, we get immediately that

$$b \ge 3 \Rightarrow \dim_{\mathbb{R}} K = 0,$$
  
 $b = 2 \Rightarrow \dim K = \rho_T - 1.$ 

Now let us suppose b = 1, meaning T is an elliptic curve. If  $O \in T$  denotes the neutral element of T, we have  $[C] = \varphi^*[O]$  as codimension 1 cycles on A. The 0-cycle [O] is an ample generator of Pic T, and  $\psi^* \mathcal{O}_{\mathbb{P}}(1) = \mathcal{O}_T(m[O])$  where  $m = \deg \psi^* \mathcal{O}_{\mathbb{P}}(1) > 0$ . In conclusion,

$$\mathcal{O}_A(D) = L = \phi_L^* \mathcal{O}_{\mathbb{P}}(1) = \varphi^* \mathcal{O}_T(m[O]) = m\varphi^* \mathcal{O}_T([O]) = m\mathcal{O}_A(C),$$

hence D is linearly equivalent to m[C].

In particular,  $\delta_A(D) = \delta_{C \times T}(C)$ , hence by (4) and (12) we get

$$\delta_A(D) = \rho_A - \rho_C = 1 + \operatorname{rk}(\operatorname{Hom}(T, C)).$$

Let  $k_T$  be the multiplicity of the elliptic curve T in the factorization up to isogeny of A. Then

$$\operatorname{Hom}(T, C) = \operatorname{Hom}(T, T^{k_T - 1}) = (k_t - 1) \operatorname{End}(T),$$

and since clearly  $k_T = k_B$  we obtain the formula (11).

#### MATTEO VERNI

The case b = 0 cannot occur, since that would mean  $\delta_X(L) = \rho_A$ , which cannot happen as remarked in (3). Alternatively, note that  $C = \text{Ker } \varphi$  is by construction ([Mum08] page 88) a connected component of

$$H(D) = \{ z \in A | t_z^* D = D \},\$$

where  $t_z$  is the translation by z, but this subset is never the whole A since for any choice of  $x \in D$ ,  $y \notin D$ , then  $y - x \notin H(D)$ .

As the above proposition shows, it turns out that  $\delta_A$  is very often reached by  $\delta_A(D)$  where D is an abelian subvariety of codimension one, apart from certain cases where  $\delta_A(D)$  is low and determined by very specific simple abelian surfaces showing up as isogeny factors of A. We can now prove Theorem 1.3.

Proof of Theorem 1.3. The statement in the case  $\delta_A = 0$  follows immediately from Proposition 4.1. Suppose then  $\delta_A(D) = \delta_A > 0$  for some effective divisor  $D \subset A$ . Then  $b \leq 2$ : if b = 1, we fall in the first case of the present corollary, by the third point of Proposition 4.1. If instead b = 2, we have  $\delta_A = \delta_A(\varphi^* N) \in$  $\{1, 2, 3\}$  for some N ample on B. If B where not simple, meaning  $B \sim E \times E'$ , and if L denotes the pullback of an ample line bundle of E onto A, then  $\delta_A(L) \geq$  $\rho_B - 1 = \delta_A(D) = \delta_A$  so  $\delta_A(L) = \delta_A$  and we fall again in the first case of the corollary.

So let us suppose we are not in the first case of the corollary, which by the above argument means that b = 2 and B simple. The latter implies that  $\delta_A \in \{1, 2\}$ ; once again by  $\delta_A(\varphi^* N) = \rho_B - 1$ , these two possible values correspond to the second and third cases of the corollary. The type of the simple abelian surfaces follows from the restrictions on Picard rank imposed by the type, as in [BL04, Proposition 5.5.7]. The stated obstruction to the presence of elliptic curves is due to the working assumption of us not being in the first case of the corollary.  $\Box$ 

As an example, Theorem 1.3 says that if A is an abelian threefold, then surely  $\delta_A \in \{0, 1, 2, 3, 5\}$ , and we will have the following cases ( $E_{RM}$  and  $E_{CM}$  denote respectively real-multiplication and complex-multiplication elliptic curves):

- $\delta_A = 5$  and  $A \sim E_{CM}^3$ .
- $\delta_A = 3$  and  $A \sim E_{RM} \times E_{CM}^2$ .
- $\delta_A = 2$  and either  $A \sim S_{\text{II}} \times E$ , where  $S_{\text{II}}$  is a simple abelian surface of type II (hence  $\rho_{S_{\text{II}}} = 3$ ), or  $A \sim E_{RM}^2 \times E$  with  $E \not\sim E_{RM}$ .
- $\delta_A = 1$  and either  $A \sim E \times E' \times E''$  with all three curves non-isogenous, or  $A \sim S \times E$  with S simple not isogenous to  $S_{\text{II}}$  such as above.
- $\delta_A = 0$  and A simple.

4.1. Families of abelian varieties. Before looking at the behaviour of the Lefschetz defect in families of abelian varieties, we should point out that the families we must take into consideration cannot be the naive ones: this is because for abelian varieties we have Noether-Lefschetz phenomena complicating the picture, which were totally absent in the Fano case. More specifically, the Picard number

of Fano varieties stays constant under deformation, while for abelian varieties it jumps on a countable union of analytic subvarieties of the moduli space, so the cohomology classes we are trying to count are disappearing and reappearing in an uncontrolled manner as we deform. This is the same problem that appears when trying to parametrize, say, abelian varieties with complex multiplication: one cannot hope for a good coarse moduli space only of abelian varieties X such that  $\operatorname{End}_{\mathbb{Q}}(X) = F$  for some non-real number field F. Rather, one should impose the weaker requirement  $F \subset \operatorname{End}_{\mathbb{Q}}(X)$ : this condition does indeed yield (coarse) moduli spaces ([BL04, Chapter 9]). Analogously, to control NS(X) we consider the following families.

**Definition 4.2.** Let X be a compact complex manifold and  $\Lambda$  an abelian group. A  $\Lambda$ -polarization on X is an embedding of abelian groups

$$j: \Lambda \hookrightarrow \mathrm{NS}(A).$$

For example, a polarization in the usual sense is a  $\mathbb{Z}$ -polarization which lands  $\mathbb{Z}$  into the ample cone. Fix (X, j) a  $\Lambda$ -polarized manifold.

**Definition 4.3.** A  $\Lambda$ -polarized deformation of (X, j) is a smooth proper family  $\mathcal{X} \to B$  such that B is a connected complex manifold,  $0 \in B$  a point such that  $\mathcal{X}_0 \simeq X$  and for all  $b \in B$ , there is a chosen  $\Lambda$ -polarization

$$j_b \colon \Lambda \hookrightarrow \mathrm{NS}(\mathcal{X}_b)$$

which corresponds to the inclusion j under the identification

$$H^2(\mathcal{X}_b,\mathbb{Z})\simeq H^2(X,\mathbb{Z}).$$

Nontrivial deformations of this type are plentiful: their existence follows from the existence of deformations of abelian varieties X with  $F \subset \operatorname{End}_{\mathbb{Q}}(X)$  mentioned in the above discussion (see also [BL04, Proposition 5.5.7]).

**Definition 4.4.** Let (X, j) be a  $\Lambda$ -polarized complex manifold. For any effective divisor  $D \subset X$  we define the number

$$\delta_{(X,j)}(D) := \operatorname{Ker}(\Lambda \otimes_{\mathbb{Z}} \mathbb{Q} \xrightarrow{\sim c_1(D)} H^4(X, \mathbb{Q}))$$

and accordingly we define the  $\Lambda$ -defect of X as the number

$$\delta_{(X,j)} = \max_{D \subset X} \delta_{(X,j)}(D).$$

Note that for a smooth projective complex variety X and  $D \subset X$  a divisor, we have by definition that the Lefschetz defect is exactly the NS(X)-defect as above, i.e., we have

$$\delta_{(X,id)}(D) = \delta_X(D),$$

where on the left we see X as a NS(X)-polarized manifold with the obvious polarization.

**Proposition 4.5.** For any  $\Lambda$ -deformation  $\mathcal{A} \to S$  of abelian varieties and any two points  $t, t' \in S$ , one has

$$\delta_{(\mathcal{A}_t, j_t)} = \delta_{(\mathcal{A}_{t'}, j_{t'})}.$$

In particular, the Lefschetz defect of an abelian variety A can only increase for any NS(A)-polarized deformation of A, and it stays constant outside a countable union of proper analytic subsets of the base.

*Proof.* The second statement follows from the first, where the countably many analytic subsets mentioned are exactly the ones where the Picard number jumps up. Note that the Picard number cannot jump down as there is an inclusion  $NS(A) \hookrightarrow NS(A_t)$  by the definition of NS(A)-polarized deformation.

Let  $\delta_{(\mathcal{A}_t,j_t)} = \delta_{(\mathcal{A}_t,j_t)}(D)$ . Then we saw above that there exists a morphism of abelian varieties  $\varphi : \mathcal{A}_t \to B$  and  $\mathcal{O}_{\mathcal{A}_t}(D) = \varphi^* N$  for N ample on B. Once again we denote  $C := \text{Ker } \varphi$ , and we set  $b := \dim B = \kappa(\mathcal{A}_t, \mathcal{O}_{\mathcal{A}_t}(D))$ .

From the complex tori viewpoint, a deformation of A corresponds to a deformation of its defining lattice  $\Pi_A$  inside  $\mathbb{C}^g$ . The subtorus  $C \subset A$  corresponds to a sublattice  $\Pi_C \subset \Pi_A$ , and the deformation of  $\Pi_A$  induces a unique deformation on  $\Pi_C$ . By taking the quotients of this inclusion of lattices, we get a deformation of the lattice  $\Pi_B$ . This means there exist deformations  $\mathcal{B}$  and  $\mathcal{C}$  of B and Crespectively, and a morphism

$$\phi \colon \mathcal{A} \to \mathcal{B}$$

relative over S, whose fiber over s has kernel equal to  $C_s$ . Since all fibers of  $\mathcal{A}$  are abelian, then all subtori  $C_s$  and all quotient tori  $\mathcal{B}_s$  are abelian as well.

Suppose first that b = 2. By [BL04, Lemma 8.7.1] and the discussion preceding it, N on B extends to a line bundle  $\mathcal{N}$  on  $\mathcal{B}$  and  $\mathcal{N}_s$  is ample on each  $\mathcal{B}_s$ . We define  $\mathcal{L} := \phi^* \mathcal{N}$ ; we then have  $\mathcal{L}_t = \mathcal{O}_{\mathcal{A}_t}(D)$ , and  $\mathcal{L}_s$  is effective for all  $s \in S$ . If  $[D] \in \Lambda \subset H^2(\mathcal{A}_t, \mathbb{Z})$ , then  $[\mathcal{L}_s] \in \Lambda \subset H^2(\mathcal{A}_s, \mathbb{Z})$  for all  $s \in S$  and

$$\delta_{\Lambda}(\mathcal{L}_s) = \operatorname{rk}(j_s(\Lambda) \cap \phi_s^* NS(\mathcal{B}_s)) - 1$$

since  $\mathcal{N}_s$  being ample on each  $\mathcal{B}_s$  means  $\phi_s^*[\mathcal{N}_s] \in \Lambda_{\mathbb{Q}} \cap \phi_s^* NS(\mathcal{B}_s)$  is the only class coming from  $\mathcal{B}_s$  that does not vanish after  $\sim [\mathcal{L}_s]$ . Now note that, for all  $s \in S$ ,

$$j_s(\Lambda) \cap \phi_s^* \operatorname{NS}(\mathcal{B}_s) = j_s(\Lambda) \cap \phi_s^* H^2(\mathcal{B}_s, \mathbb{Z}).$$

The right side has a dimension which is invariant for  $s \in S$ , so  $\delta_{(\mathcal{A}_s, j_s)}(\mathcal{L}_s)$  stays constant as s varies.

Applying this to s = t, t' we conclude that

$$\delta_{(\mathcal{A}_t,j_t)} = \delta_{(\mathcal{A}_t,j_t)}(D) = \delta_{(\mathcal{A}_{t'},j_{t'})}(\mathcal{L}_{t'}) \le \delta_{(\mathcal{A}_{t'},j_{t'})}.$$

Now the case b = 1: we have  $\alpha \smile [D] = 0 \iff \alpha \smile [C] = 0$ , thus we can switch back to the more convenient characterization (5), writing

$$\delta_{(\mathcal{A}_s,j_s)}(\mathcal{L}_s) = \dim \operatorname{Ker}(\Lambda_{\mathbb{Q}} \xrightarrow{\imath_s} H^2(\mathcal{C}_s,\mathbb{Q})),$$

where  $i_s$  is the natural embedding  $\mathcal{C}_s \hookrightarrow \mathcal{A}_s$ . The above number does not depend on s, by definition of  $\Lambda$ -polarized family, which is what we were looking for.  $\Box$ 

#### LEFSCHETZ DEFECT IN FAMILIES

### References

- [BL04] Birkenhake C. and Lange H. : Complex Abelian Varieties. Grundlehren der mathematischen Wissenschaften. Springer Berlin Heidelberg, 2004.
- [Cas12] Casagrande, C. : On the Picard number of divisors in Fano manifolds. Ann. Sci. Éc. Norm. Supér. (4), 45(3):363–403, 2012.
- [Cas23] Casagrande C. : The Lefschetz defect of Fano varieties. Rend. Circ. Mat. Palermo (2), 72(6):3061–3075, 2023.
- [Cas24] Casagrande C. : Fano 4-folds with  $b_2 > 12$  are products of surfaces. *Invent. Math.*, **236**(1):1–16, 2024.
- [CD15] Casagrande C. and Druel S. : Locally unsplit families of rational curves of large anticanonical degree on Fano manifolds. *IMRN*, **2015**(21):10756–10800, 2015.
- [CRS22] Casagrande C., Romano E. A., and Secci S. A. : Fano manifolds with Lefschetz defect 3. J. Math. Pures Appl. (9), 163:625–653, 2022.
- [Del74] Deligne P. : Théorie de Hodge. III. Inst. Hautes Études Sci. Publ. Math., pages 5–77, 1974.
- [DH11] De Fernex T. and Hacon C. D. : Deformations of canonical pairs and Fano varieties. J. Reine Angew. Math., 651:97–126, 2011.
- [FGI<sup>+</sup>05] Fantechi B., Göttsche L., Illusie L., Kleiman S., Nitsure N., and Vistoli A. : Fundamental Algebraic Geometry: Grothendieck's FGA Explained. Mathematical surveys and monographs. American Mathematical Society, 2005.
- [GM88] Goresky M. and MacPherson R. : Stratified Morse theory, volume 14 of Ergeb. Math. Grenzgeb., 3. Folge. Berlin etc.: Springer-Verlag, 1988.
- [Mum08] Mumford D. : Abelian varieties. With appendices by C. P. Ramanujam and Yuri Manin. New Delhi: Hindustan Book Agency/distrib. by American Mathematical Society (AMS); Bombay: Tata Institute of Fundamental Research, corrected reprint of the 2nd ed. 1974 edition, 2008.
- [Noc12] Della Noce G. : On the Picard number of singular Fano varieties. *IMRN*, **2014**(4):955–990, 2012.
- [Sec23] Secci S.A. : Fano fourfolds having a prime divisor of Picard number 1. Adv. Geom., 23(2):267–280, 2023.
- [Siu02] Siu Y.-T. : Extension of twisted pluricanonical sections with plurisubharmonic weight and invariance of semipositively twisted plurigenera for manifolds not necessarily of general type. In *Complex geometry. Collection of papers dedicated to Hans Grauert on the occasion of his 70th birthday*, pages 223–277. Berlin: Springer, 2002.
- [SY22] Shen J. and Yin Q. : Topology of Lagrangian fibrations and Hodge theory of hyper-Kähler manifolds. *Duke Math. J.*, **171**(1):209–241, 2022.
- [Tot12] Totaro B. : Jumping of the nef cone for Fano varieties. J. Algebr. Geom., **21**(2):375–396, 2012.
- [Voi92] Voisin C. : Sur la stabilité des sous-variétés lagrangiennes des variétés symplectiques holomorphes. In Complex projective geometry. Selected papers from the conference on projective varieties, 19-24 June 1989 in Trieste, Italy, and the conference on vector bundles and special projective embeddings, 3-16 July 1989 in Bergen, Norway, pages 294–303. Cambridge: Cambridge University Press, 1992.

Sorbonne Université, Université Paris Cité, CNRS, IMJ-PRG, F-75005 Paris, France

Email address: matteo.verni@imj-prg.fr