# Binary superfluids: low-energy properties and dissipative processes from spontaneous emission of massive phonons

Silvia Trabucco,<sup>1,2</sup> Luca Lepori,<sup>3,4</sup> Maria Luisa Chiofalo,<sup>5,6</sup> and Massimo Mannarelli<sup>2,1</sup>

<sup>1</sup>Gran Sasso Science Institute, Viale Francesco Crispi, 7, 67100 L'Aquila.

<sup>2</sup>INFN, Laboratori Nazionali del Gran Sasso, Via G. Acitelli, 22, 67100 Assergi (AQ), Italy.

<sup>3</sup>Dipartimento di Scienze Matematiche, Fisiche e Informatiche,

Università di Parma, Parco Area delle Scienze, 53/A, I-43124 Parma, Italy.

<sup>4</sup>Gruppo Collegato di Parma, INFN-Sezione Milano-Bicocca, I-43124 Parma, Italy.

<sup>5</sup>Dipartimento di Fisica, Università di Pisa, Polo Fibonacci, Largo B. Pontecorvo 3.

<sup>6</sup>INFN Sezione di Pisa, Polo Fibonacci, Largo B. Pontecorvo 3, 56127 Pisa, Italy.

We discuss the low-energy properties of binary superfluids with density-dependent interactions. Adding an intra-species coupling that induces an explicit soft symmetry-breaking, we determine the background pressure and we show that the low-energy spectrum consists of a massless Nambu-Goldstone boson and a massive (pseudo) Nambu-Goldstone boson. When the background velocities of the two superfluids are transonic, the system is characterized by two distinct acoustic horizons: the hydrodynamic analogue of the black hole event horizon. The Hawking-like emission occurring at these horizons produces an effective friction on the fluids. We compute the viscosity-to-entropy ratios close to the two acoustic horizons, finding that the emission of pseudo Nambu-Goldstone bosons violates the bound conjectured by Kovtun, Son and Starinet.

#### I. INTRODUCTION

The current generation of quantum technologies opens up the possibility of investigating the physics of ultracompact objects, like neutron stars or black-holes [1-13], in analogue experiments involving quantum fluids with tunable degree of correlations. Concerning black holes, key aspects are the dissipative effects close to horizons, as characterized by the shear viscosity-to-entropy ratio  $\eta/s$  [14, 15]. This quantity is known to depend on the degree of correlations in the fluid and reaches a minimum in correspondence of a critical point [15]. Interestingly enough, this minimum is of the order of  $\hbar/k_B$  in strongly correlated quantum fluids,  $k_B$  being the Boltzmann constant, with a value that is similar in quite diverse systems, as for instance the quark-gluon plasma and ultracold Fermi gases, irrespective of their so different microscopic details. The lower bound  $\eta/s = 1/4\pi$  in units of  $\hbar/k_B$ , known from Kovtun, Son, Starinet (KSS) [16], has been derived in different frameworks, i.e. gravity-dual theories [17], analogue gravity [1], kinetic theories [18, 19], and more generally from Kubo relations [1, 15, 17]. While the KSS bound is seen to hold for a remarkably large class of theories, specific situations show that it is nonuniversal [15, 20]. These include translationally invariant non-Fermi liquids in two dimensions [21], the presence of higher curvature corrections in the dual gravitational theory and massive gravity in a higher derivative theory, such as the Gauss–Bonnet theory [22]. The latter introduces a lower bound  $\eta/s_{GB} = (1/4\pi)(16/25)$ , which in fact avoids violations of causality and of energy positivity [15]. Overall, lower bounds  $\eta/s$  establish that the low-shear viscosity systems, i.e. nearly perfect fluids, are also strongly correlated, bearing important information to fully exploit the analogy between black hole horizons and quantum fluids. Additionally, whether only one universal  $\eta/s$  bound does exist, comprising all possible theories, remains a relevant open question. It is therefore especially interesting to design a theoretical framework, amenable to realistic experimental implementations, to explore the degree of KSS-bound violation under the tuning of specific external parameters.

Multicomponent quantum gases offer a versatile platform for investigating a wide range of phenomena, for instance due to the possibility to manage multiple levels simultaneously, and to the high tunability of both intraspecies and interspecies interactions. These features have been shown to allow access to novel phases of matter [23–25], possibly supporting nontrivial topological defects [26-32], as well as self-bound quantum droplets [33, 34]. The first experimental realization of two overlapping quantum bosonic mixtures was achieved by sympathetic cooling of two integer spin states of <sup>87</sup>Rb atoms [35]. Subsequently, binary superfluids of fermions, of bosons as well as of a mixture of fermionic and bosonic gases have been explored to investigate various relevant phenomena for elementary physics [36-42]. Among them, we mention the Andreev-Bashkin effect [43], analog black-holes [44–47] and bubble formation in first-order phase transitions [48]. The possible phases of multicomponent quantum gases are also of relevance to characterize the interior of compact stars [8]. Inside them, superfluid neutrons should coexist with superconducting protons, moreover, if deconfined quark matter is present, baryonic superfluidity may be realized along with color superconductivity [49–54].

In this work, we focus on a binary mixture of weakly interacting bosons at vanishing temperature. In each component, superfluidity arises when the global symmetry associated with the conservation of number of particles is spontaneously broken [55, 56], resulting in the presence of massless Goldstone bosons in the low-energy spectrum, *i.e.* the phonons. When an intra-species Rabi coupling is added, such symmetry is explicitly broken [26, 57] and the low-energy spectrum of the binary mixture is modified. If the Rabi coupling has a subleading effect, the low-energy spectrum consists of both a massless and a massive phonon [58, 59]. These degrees of freedom may be interpreted as density waves [60], and they can be described as massless and massive scalar particles propagating on top of an emergent acoustic metric [1, 61-65]. This system is particularly interesting when the flow of one or both species is transonic, providing the hydrodynamic analog of black-holes [14, 44, 47]. Similarly to astrophysical black-holes, Hawking-like emission of phonons occurs at the acoustic horizon, imprinting a signature in the density-density correlation functions [47, 66, 67]. This spontaneous phonon emission is an irreversible process that occurs at the expense of the fluid kinetic energy, and can be described as effective shear and bulk viscosities of the superfluid [19].

Within this framework, we compute both the shearto-entropy density ratio and the bulk-to-entropy density ratio associated to the spontaneous emission at the acoustic horizon of binary transonic superfluid, extending the previous results for a single species [19]. We show how the saturation of the bound occurs for the spontaneous emission of massless phonons, while the bound is violated when a massive phonon is emitted. In fact, in the latter condition, the corresponding pressure and energy density decrease, reducing the effective shear and bulk viscosities with respect to the massless case. These findings are shown in the nonrelativistic limit and are directly applicable to homonuclear bosonic mixtures of alkali atoms with coupled hyperfine states [34, 35].

This paper is organized as follows. In Sec. II, we briefly review the analogue gravity set-up with a single-species boson gas. In Sec. III, we describe the general features of binary bosonic systems, while in Sec. IV we determine the ground-state pressure. The low-energy field theory of binary superfluids is developed in Sec. V. The Hawkinglike emission of massive phonons is studied in Sec. VI and in Sec. VII we evaluate the resulting effective viscosity coefficients. We draw our conclusions in Sec. VIII. Two distinct and general derivations of the effective lowenergy Lagrangian are given in the Appendix A.

Along all the work, we use natural units  $\hbar = c = k_B = 1$ .

### II. EFFECTIVE THEORY OF A SINGLE ULTRACOLD BOSON FLUID

As a first step, we briefly recap some useful facts regarding dilute single-species boson gas with a U(1) global symmetry. In particular, we show how the acoustic metric emerges in the low-energy effective theory.

#### A. General considerations

We consider a complex scalar field,  $\Phi$ , with nonvanishing chemical potential and local interactions that do not explicitly break the U(1) global symmetry associated to the conserved number of particles. We assume that the bulk temperature of the system is negligible. For definiteness, we examine a simple model with contact self-interaction; for a more general system see the Appendix A. The corresponding Lagrangian density, hereafter the Lagrangian, is given by

$$\mathcal{L} = (D_{\nu}\Phi)^* D^{\nu}\Phi - m^2 |\Phi|^2 - U|\Phi|^4 \,, \tag{1}$$

where m is the mass of the bosons,  $0 < U \ll 1$  a perturbative dimensionless interaction constant, and

$$D_{\nu} = \partial_{\nu} + i \frac{\mu}{\gamma} v_{\nu} , \qquad (2)$$

is the covariant derivative, depending on the medium properties, see for instance [68] for more details. We identify  $\mu$  with the boson chemical potential, while  $v_{\nu} =$  $\gamma(1, \boldsymbol{v})$  is the fluid four-velocity:  $\gamma$  is the Lorentz factor, such that  $v^{\mu}v^{\nu}\eta_{\mu\nu} = 1$ , with  $\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1)$ the Minkowski metric. The non-relativistic limit of Eq. (1) is the Gross-Pitaevskii (Hartree-Fock) Lagrangian describing a diluted and weakly interacting boson gas at vanishing temperature [56, 69]. The excitation spectrum over the superfluid ground-state can be calculated linearizing the Gross-Pitaevskii Lagrangian, this approach being equivalent to the Bogoliubov formalism [24, 70].

Expanding the covariant derivative in Eq. (2), we have

$$\mathcal{L} = \partial_{\nu} \Phi^* \partial^{\nu} \Phi - i \frac{\mu}{\gamma} v_{\nu} (\Phi^* \partial^{\nu} \Phi - \Phi \partial^{\nu} \Phi^*) - \left( m^2 - \frac{\mu^2}{\gamma^2} \right) |\Phi|^2 - U |\Phi|^4 , \qquad (3)$$

showing that the chemical potential gives two distinct contributions to the Lagrangian: an explicit Lorentz symmetry breaking, due to the term linear in the chemical potential, and the effective mass shift  $\mu^2/\gamma^2$ . When  $\mu = \gamma m$  the effective mass of the scalar field vanishes, signaling a spontaneous symmetry breaking. For  $\mu > \gamma m$ , the U(1) global symmetry is indeed spontaneously broken [71], and the minimum of the potential

 $V = \left(m^2 - \frac{\mu^2}{\gamma^2}\right)|\Phi|^2 + U|\Phi|^4, \qquad (4)$ 

$$|\Phi|^{2} = \frac{\mu^{2} - m^{2}\gamma^{2}}{2U\gamma^{2}}.$$
(5)

The fact that the potential depends only on  $|\Phi|$  and not on its phase, invites to use the Madelung representation

$$\Phi = \frac{\rho}{\sqrt{2}} e^{i\theta},\tag{6}$$

is

where the two real scalar fields  $\rho$  and  $\theta$  are called the radial and phase fields, respectively. In the Madelung representation the Lagrangian in Eq. (3) reads

$$\mathcal{L} = \frac{1}{2} \partial_{\nu} \rho \partial^{\nu} \rho + \frac{1}{2} \rho^{2} \partial_{\nu} \theta \partial^{\nu} \theta + \frac{\rho^{2} \mu}{\gamma} v_{\nu} \partial^{\nu} \theta - \frac{m^{2} \gamma^{2} - \mu^{2}}{2 \gamma^{2}} \rho^{2} - \frac{U}{4} \rho^{4} , \qquad (7)$$

where the radial field is massive and with quartic selfinteraction. On the other hand, the  $\theta$  field is a phase, thus it only appears in derivative terms. Since  $\theta$  is a cyclic variable, the current

$$J_{\nu} = \frac{\delta \mathcal{L}}{\delta \partial^{\nu} \theta} = \rho^2 \left( \frac{\mu}{\gamma} v_{\nu} + \partial_{\nu} \theta \right) , \qquad (8)$$

is conserved. In the ground state, the phase is homogeneous, therefore

$$J_{\nu} = \rho^2 \frac{\mu}{\gamma} v_{\nu} = n v_{\nu} \,, \tag{9}$$

where

$$n = \frac{\mu}{\gamma} \rho^2 \,, \tag{10}$$

is the number density. From Eq. (8), it is clear that we can introduce the chemical potential and the fluid velocity in an alternative way. If we replace in Eq. (1) the covariant derivative with the standard derivative, then in the expression of the current, see Eq. (8), the first term in the round brackets vanishes. Then, we can define a mean field value of the phase field in such a way that

$$\partial_{\nu}\theta = \frac{\mu}{\gamma}v_{\nu}\,,\tag{11}$$

and thus the mean field value of the current is equal to Eq. (9). This procedure implies that the mean field value of the phase field is equal to  $\mu v_{\nu} x^{\nu} / \gamma$ . One can see that this is equivalent to a gauge transformation: we can replace the covariant derivative in Eq. (1) with a standard derivative and then by the gauge transformation

$$\Phi \to \Phi \, e^{i\frac{\mu}{\gamma}v_{\nu}x^{\nu}} \,, \tag{12}$$

we obtain the same Lagrangian reported in Eq. (3). This procedure has the merit of treating the radial and the phase field on the same footing: the background medium properties are taken into account by appropriate mean field values of  $\rho$  and  $\theta$ . The former is related to the number density, while the latter to the chemical potential and the fluid four-velocity.

The radial field can be viewed as a variational parameter: its mean field value, see Eq. (5),

$$\rho^2 = \frac{\mu^2 - m^2 \gamma^2}{U\gamma^2} \quad \text{for } \mu \ge \gamma m \,, \tag{13}$$

corresponds to the solution of the stationary condition that minimizes the potential in Eq. (3). This is equivalent to maximize the ground state Lagrangian. Since the grand potential per unit volume is given by

$$\Omega = -\frac{1}{\mathcal{V}} \int d^4 x \mathcal{L} \,, \tag{14}$$

where V is the total volume, the configuration that maximizes the Lagrangian minimizes the thermodynamic potential. Given that  $P = -\Omega$ , substitution of Eq. (13) in Eq. (7) yields the quantum pressure of the system:

$$P = \frac{(\mu^2 - m^2 \gamma^2)^2}{4U\gamma^4} \,. \tag{15}$$

Notice that this includes only the contribution of condensation and neglects temperature as well as quantum fluctuations. The velocity dependence, determined by the Lorentz factor, gives the relativistic analogue of the Bernoulli equation. Indeed, we have that

$$P + \frac{\mu n v^2}{2} = \frac{(\mu^2 - m^2)^2}{4U} + \mathcal{O}(v^4), \qquad (16)$$

where the number density is defined in Eq. (10). We remark that  $\theta$  is not a variational parameter: its derivative determines the velocity of the fluid. If we were to take it as a variational parameter, we should have considered the value that maximizes the pressure, that is the value for which the velocity vanishes [9]. This approach is however incorrect, because the fluid current  $J_{\nu}$  in Eq. (8) is conserved. On the other hand, it indicates that the addition of dissipative interactions in the Lagrangian in Eq. (7) would have the effect of changing  $\theta$  so that the velocity of the fluid is reduced and thus the pressure maximized.

#### B. Effective low-energy theory

We now determine the effective Lagrangian for the lowenergy excitations of the superfluid. We separate both the radial and phase fluctuations,  $\tilde{\rho}$  and  $\tilde{\theta}$ , from their mean field solutions as

$$\rho \to \rho + \tilde{\rho} , \quad \theta \to \theta + \theta ,$$
 (17)

where  $\rho$  and  $\theta$  now indicate the solutions of the classical equations of motion, corresponding to a stationary point of the action.

Expanding around the mean field we have that

$$\mathcal{L} = \mathcal{L}_0(\rho, \partial_\mu \rho, \partial_\mu \theta) + \tilde{\rho} \left( \frac{\delta \mathcal{L}}{\delta \rho} - \partial_\mu \frac{\delta \mathcal{L}}{\delta \partial_\mu \rho} \right) \Big|_{\rho, \theta} + \delta \partial_\mu \tilde{\theta} \left. \frac{\delta \mathcal{L}}{\delta \partial_\mu \theta} \right|_{\rho, \theta} + \mathcal{L}_2 + \dots, \qquad (18)$$

where the dots indicate neglected cubic terms and higher, as well as surface terms. The term  $\mathcal{L}_0(\rho, \partial_\mu \rho, \partial_\mu \theta)$  is the background pressure previously evaluated in Eq. (15). The linear term vanishes, because we are evaluating it at the stationary point of the action. The quadratic part of the Lagrangian reads

$$\mathcal{L}_2 = \frac{1}{2} \partial_\nu \tilde{\rho} \partial^\nu \tilde{\rho} - \frac{\tilde{m}^2}{2} \tilde{\rho}^2 + V^\nu \tilde{\rho} \partial_\nu \tilde{\theta} + \frac{\rho^2}{2} \partial_\nu \tilde{\theta} \partial^\nu \tilde{\theta} \,, \quad (19)$$

where

$$\tilde{m} = \sqrt{2U\rho^2} \,, \tag{20}$$

is the radial field mass. The phase fluctuation is instead massless, and is the Nambu Goldstone boson (NGB) associated to the spontaneous breaking of the U(1) symmetry. Hereafter, we will call it phonon. The vector coupling between  $\tilde{\rho}$  and  $\partial_{\nu}\tilde{\theta}$ 

$$V^{\nu} = 2\rho \frac{\mu}{\gamma} v^{\nu} \,, \tag{21}$$

is determined by the background properties. Since the radial field is massive, we can integrate it out using its equation of motion

$$(\Box + \tilde{m}^2)\tilde{\rho} = V^{\mu}\partial_{\mu}\tilde{\theta}.$$
(22)

The formal inversion of the previous equation yields

$$\tilde{\rho} = \frac{1}{\Box + \tilde{m}^2} V^{\mu} \partial_{\mu} \tilde{\theta} \,. \tag{23}$$

For phonon momenta p smaller than the radial field mass, the expansion of  $(\Box + \tilde{m}^2)^{-1}$  gives terms suppressed as powers of  $(p/\tilde{m})^2$ . Therefore, the relevant scale for the momentum expansion is not m but  $\tilde{m}$ , that is the curvature of the potential at the minimum. Close to any second order phase transition point the curvature of the potential vanishes, and the energy and momentum range of validity of the low-energy effective Lagrangian shrinks.

Upon substituting Eq. (23) in Eq. (19), we obtain

$$\mathcal{L}_{\text{eff}} = \frac{1}{2} V^{\mu} \partial_{\mu} \tilde{\theta} \frac{1}{\Box + \tilde{m}^2} V^{\nu} \partial_{\nu} \tilde{\theta} + \frac{\rho^2 \eta^{\mu\nu}}{2} \partial_{\nu} \tilde{\theta} \partial_{\mu} \tilde{\theta} , \quad (24)$$

and the leading order in  $p/\tilde{m}$  consists in replacing

$$\frac{1}{\Box + \tilde{m}^2} \to \frac{1}{\tilde{m}^2} \,. \tag{25}$$

We can finally rewrite the effective low-energy Lagrangian as

$$\mathcal{L}_{\text{eff}} = \frac{\rho^2}{2} \left( \eta^{\mu\nu} + \frac{V^{\mu}V^{\nu}}{\rho^2 \tilde{m}^2} \right) \partial_{\nu} \tilde{\theta} \partial_{\mu} \tilde{\theta} \equiv \frac{\sqrt{-g}g^{\mu\nu}}{2} \partial_{\nu} \tilde{\theta} \partial_{\mu} \tilde{\theta} \,, \tag{26}$$

where the emergent acoustic metric for the phonon field is given by

$$\sqrt{-g}g^{\mu\nu} = \rho^2 \left(\eta^{\mu\nu} + \left(\frac{1}{c_s^2} - 1\right)v^{\mu}v^{\nu}\right),$$
 (27)

and

$$c_s = \sqrt{\frac{U\rho^2}{2m^2 + 3U\rho^2}},$$
 (28)

is the adiabatic speed of sound. This quantity vanishes for  $U\rho^2 \rightarrow 0$ , corresponding to the transition from the broken to the normal phase, while its upper bound is  $\sqrt{1/3}$ .

We observe that retaining the leading order in  $p/\tilde{m}$ in Eq. (26) is equivalent to neglect the kinetic term  $\partial_{\nu}\tilde{\rho}\partial^{\nu}\tilde{\rho}/2$  in Eq. (19), and treat  $\tilde{\rho}$  as a homogeneous field. This will be useful in the development of the low-energy theory of binary superfluids. More general approaches to the effective low-energy theory are reported in the Appendix **A**. Now, we move to consider binary mixture of superfluids.

# III. BINARY BOSON SUPERFLUIDS

In the previous section, we have seen how it is possible to define an emergent acoustic metric for the longwavelength excitations of a single superfluid. We now extend the same approach to a mixture composed of two *flavors A* and *B*. In experiments with homonuclear gases, they may correspond to two atomic hyperfine states. In astrophysics, they may indicate the superfluid components in the neutron star interior. We take a schemeindependent approach and, as in the previous section, we employ a relativistic framework. We assume vanishing temperature and that the two fluids move with velocities  $v_A$  and  $v_B$  with respect to an external observer.

If the number of particles of each flavor is separately conserved, we have a U(1) symmetry group associated to each flavor yielding to  $U_A(1) \otimes U_B(1)$  as a symmetry group. The spontaneous symmetry breaking of such group can produce one or two NGBs, depending on the interaction mechanism. We discuss in detail the case in which a U(1) subgroup of  $U_A(1) \otimes U_B(1)$  is explicitly broken by a soft coupling term. Then, we expect that the spontaneous symmetry breaking mechanism could produce one genuine NGB and one pseudo-NGB, with a mass proportional to the soft coupling.

#### A. Lagrangian of binary bosonic systems

The general form of the Lagrangian for two complex scalar fields,  $\Phi_{A/B}$ , is

$$\mathcal{L} = \mathcal{L}_K(\Phi_A, \Phi_B) - \frac{1}{2}m_A^2 \Phi_A^* \Phi_A - \frac{1}{2}m_B^2 \Phi_B^* \Phi_B - V(\Phi_A, \Phi_B)$$
(29)

where  $m_{A/B}$  are the masses of the two fields. The kinetic term is

$$\mathcal{L}_K = (D_{\mu,A} \mathbf{\Phi}_A)^* D_A^{\mu} \mathbf{\Phi}_A + (D_{\mu,B} \mathbf{\Phi}_B)^* D_B^{\mu} \mathbf{\Phi}_B \,, \quad (30)$$

with the covariant derivatives defined as in Eq. (2), that is

$$D_A^{\nu} = \partial^{\nu} + i \frac{\mu_A}{\gamma_A} v_A^{\nu}, \quad D_B^{\nu} = \partial^{\nu} + i \frac{\mu_B}{\gamma_B} v_B^{\nu}, \qquad (31)$$

as each component has distinct chemical potential and velocity. Here,  $\gamma_{A/B} = (1 - v_{A/B}^2)^{-1/2}$  are the Lorentz factors of each fluid component. We consider only quartic intra- and inter-species interactions,  $V_U$ , to which a Rabi term,  $V_{\lambda}$ , is conveniently added:

$$V = V_U + V_\lambda = \sum_{i=A,B} U_{ii} (\Phi_i^* \Phi_i)^2 + \sum_{i \neq j} \frac{1}{2} U_{ij} \Phi_i^* \Phi_i \Phi_j^* \Phi_j + \lambda (\Phi_A^* \Phi_B + \Phi_B^* \Phi_A),$$
(32)

with  $U_{ij}$  dimensionless coupling constants. The strength of the Rabi coupling,  $\lambda$ , has dimension of a mass squared and it is assumed to be so small that it does not sensibly modify the energy spectrum obtained for  $\lambda = 0$ . In this setting, the quartic interactions are the leading terms, while  $V_{\lambda}$  is assumed to be a subleading perturbation.

This aspect can be better clarified considering the nonrelativistic limit of Eq. (29), relevant to current and realistic experiments with ultracold atom platforms [38, 39]. The above description takes the form of a two-species interacting Hamiltonian [56, 69] given by

$$H = \sum_{l,i} \left[ b_{l,i}^{\dagger} \left( -\frac{\hbar^2}{2m_i} \frac{\partial^2}{\partial \mathbf{r}_l^2} + V_{\text{trap}}(\mathbf{r}_l) \right) b_{l,i} + \tilde{U}_{ii} \left( b_{l,i}^{\dagger} b_{l,i} \right)^2 \right]$$
$$+ \sum_{i \neq i} \frac{1}{2} \tilde{U}_{ij} \sum_{l,m} b_{l,i}^{\dagger} b_{l,i} b_{m,j}^{\dagger} b_{m,j}$$
$$+ \tilde{\lambda} \sum_{l} \left( b_{l,A}^{\dagger} b_{l,B} + b_{l,B}^{\dagger} b_{l,A} \right), \qquad (33)$$

where  $b_{l,i}$  are the ladder operators, the indices (i, j) label the two bosonic flavors, that in principle can have different numbers or fillings, while the indices (l, m) refers to the position in a lattice or in continuous space. The added one-particle potential,  $V_{\text{trap}}(\mathbf{r}_l)$ , is typically harmonic and is necessary to provide particle trapping.

The above Hamiltonian can be obtained from the relativistic Lagrangian by rescaling the interaction couplings as

$$\tilde{U}_{ij} = U_{ij} \sqrt{m_i m_j}, \quad \tilde{\lambda} = \frac{\lambda}{\sqrt{m_i m_j}}, \quad (34)$$

as a result all couplings have the same dimension and can be compared. For harmonic  $V_{\text{trap}}(\mathbf{r}_l)$ , with isotropic trapping frequency  $\Omega$ , the requirement that all the couplings are perturbative and that the Rabi term has a subleading effect with respect to the quartic ones implies that

$$\tilde{\lambda} \ll \tilde{U}_{ij} \ll \hbar \Omega \simeq 10^{-15} - 10^{-13} \,\mathrm{eV}\,, \qquad (35)$$

with  $\hbar\Omega$  being the scale of the energy difference between the ground state and the first excited state. The estimated value of  $\hbar\Omega$  is obtained considering that in current ultracold atom experiments, typical stable ranges to avoid gravity effects (at lower frequencies) and threebody losses (at higher frequencies) are  $\Omega \sim 10 - 10^3$  Hz, then  $\hbar\Omega \simeq 10^{-15} - 10^{-13}$  eV. This bound for the intensity of the Rabi coupling can drastically change on a real space lattice, where low values of  $\tilde{\lambda}$  are needed to avoid the population of excited bands [57].

Referring back to Eq. (32), the  $U_{ij}$  quartic couplings determine the strengths of the density-density boson interactions. Perturbation theory holds when  $|U_{ij}| \ll 1$ , moreover for

$$U_{AA} > 0$$
,  $U_{BB} > 0$ ,  $|U_{AB}| < 2\sqrt{U_{AA}U_{BB}}$ , (36)

the two species are miscible and are homogeneously distributed in space [72]. Without loss of generality, we take

$$0 < U_{BB} \le U_{AA} \ll 1, \qquad (37)$$

while the sign of the inter-flavor coupling,  $U_{AB}$ , is not fixed.

As in the single fluid model, the quartic couplings can produce the condensation of one or of both flavors. In other words, they are responsible of the corresponding spontaneous symmetry breaking. On the other hand, the Rabi term in Eq. (32) mixes the phases of the  $\Phi_A$  and  $\Phi_B$  fields, explicitly breaking the  $U_A(1) \otimes U_B(1)$  global symmetry. Given the relative richness of the set of possible symmetry breaking patterns, we characterize them in detail in the following section.

### **B.** Symmetries

In order to discuss the symmetries of the two-fluid system, we introduce the vector notation

$$\mathbf{\Phi} = \begin{pmatrix} \Phi_A \\ \Phi_B \end{pmatrix}, \tag{38}$$

that allows a more compact representation. We refer to the space spanned by the  $\Phi$  field as the flavor space. We begin with the discussion of the symmetries of the kinetic term in Eq. (30). In the vector basis, it can be generalized to

$$\mathcal{L}_K = (D_\nu \Phi)^{\dagger} (D^\nu \Phi) \,, \tag{39}$$

where the covariant derivative is

$$D_{\nu} = \mathcal{I}_2 \,\partial_{\nu} - iA_{\nu} \,, \tag{40}$$

with  $\mathcal{I}_2$  the 2 × 2 identity, and  $A_{\nu}$  the hermitian 2 × 2 matrix that represents a general external vector field. As in Eqs. (30) and (31), we assume that it has only time-like components that are diagonal in the flavor basis. Indicating with  $\sigma_i$  the Pauli matrices, the only nonvanishing terms in  $A_{\nu}$  can be cast as

$$A_{\nu} = A_{\nu}^{I} \mathcal{I}_{2} + A_{\nu}^{3} \sigma_{3} = \begin{pmatrix} \mu_{A}/\gamma_{A} & 0\\ 0 & \mu_{B}/\gamma_{B} \end{pmatrix} \delta_{\nu 0} \,. \tag{41}$$

Given the expression of these covariant derivatives, the kinetic term in Eq. (39) turns out to be invariant under the  $U_A(1) \otimes U_B(1)$  symmetry group, which we rewrite as

$$G = U(1)_{\mathcal{V}} \otimes U(1)_{\mathcal{A}}, \qquad (42)$$

where  $U(1)_{\mathcal{V}}$  is associated to a global phase change, generated by  $\mathcal{I}_2$ , while  $U(1)_{\mathcal{A}}$  is the *axial* group generated by  $\sigma_3$ . If the two species have the same chemical potentials and velocities, then  $A^3_{\nu}$  vanishes and  $\mathcal{L}_K$  is invariant under the symmetry group

$$G_T = U(1)_{\mathcal{V}} \otimes SU(2) \supset G, \qquad (43)$$

where SU(2) generates flavor rotations. The above relation simply states that for two fluids having the same chemical potentials and velocities, the kinetic term does not distinguish the two fluids, thus it is invariant under a global phase redefinition and flavor rotations.

In the vector notation, the mass –or quadratic– term contains also the Rabi interaction and reads

$$\frac{1}{2}m_{A}^{2}\Phi_{A}^{*}\Phi_{A} + \frac{1}{2}m_{B}^{2}\Phi_{B}^{*}\Phi_{B} + V_{\lambda}(\Phi_{A}, \Phi_{B}) = \frac{1}{2}\Phi^{\dagger}M\Phi,$$
(44)

with mass matrix

$$M = \begin{pmatrix} m_A^2 & 2\lambda \\ 2\lambda & m_B^2 \end{pmatrix} = m^2 \mathcal{I}_2 + \Delta m^2 \sigma_3 + 2\lambda \sigma_1 \,, \quad (45)$$

where  $m_{A/B}^2 = m^2 \pm \Delta m^2$ .

Let us scrutinize the symmetries of this term in various cases:

- if  $\lambda = 0$  and  $\Delta m^2 = 0$ , the mass term symmetry is  $G_T$ : the two fields are indistinguishable and invariant under global phase and flavor rotations;
- if  $\lambda \neq 0$  and  $\Delta m^2 = 0$ , the mass term is invariant under  $U(1)_{\mathcal{V}}$  times the U(1) group generated by  $\sigma_1$ . The mass eigenstates are distinguishable: they have mass squared,  $m^2 \pm 2\lambda^2$ .
- if  $\lambda = 0$  and  $\Delta m^2 \neq 0$ , the mass term symmetry is G: the two fields are distinguishable because they have different masses;
- if both λ ≠ 0 and Δm<sup>2</sup> ≠ 0, the mass term is only invariant under the U(1)<sub>ν</sub> symmetry.

In this work, we will focus on the last two cases. For sufficiently small values of  $\lambda$ , the perturbative Rabi term produces a soft breaking of the axial symmetry. As a consequence, in the spontaneously broken phase, we expect to have one true NGB, originating from the breaking of the  $U(1)_{\mathcal{V}}$  symmetry, plus a pseudo-NBG associated to the breaking of the non-exact  $U(1)_{\mathcal{A}}$  symmetry: a massless phonon and a massive phonon. In the remaining of this paper, we will consider the corrections induced by the Rabi term only at leading order  $\mathcal{O}(\lambda)$ , keeping in mind that one-loop corrections are of order  $\lambda \log(\lambda)$  [73]. We consider now the density-density interactions in Eq. (32): in the vector basis both the inter-species and intra-species interactions can be rewritten as

$$V_U = U_{AA} (\mathbf{\Phi}^{\dagger} \mathcal{P}_A \mathbf{\Phi})^2 + U_{BB} (\mathbf{\Phi}^{\dagger} \mathcal{P}_B \mathbf{\Phi})^2 + U_{AB} (\mathbf{\Phi}^{\dagger} \mathcal{P}_A \mathbf{\Phi}) (\mathbf{\Phi}^{\dagger} \mathcal{P}_B \mathbf{\Phi}), \qquad (46)$$

with

$$\mathcal{P}_A = \frac{\mathcal{I}_2 + \sigma_3}{2} \quad \text{and} \quad \mathcal{P}_B = \frac{\mathcal{I}_2 - \sigma_3}{2}, \qquad (47)$$

the two flavor projectors. Since the interaction terms are proportional to  $\mathcal{I}_2$  and  $\sigma_3$ , they are invariant under G but not under the enlarged group  $G_{\mathrm{T}}$ : they distinguish the two bosonic species. The terms proportional to  $\sigma_3$  vanish only when  $U_{AA} = U_{BB}$ , and  $U_{AA} = 2|U_{AB}|$ , however, this condition violates the stability requirement in Eq. (36).

Let us briefly recap the most important aspects of the above analysis. For vanishing Rabi coupling, the Lagrangian has the G symmetry given in Eq. (42). If this symmetry is spontaneously-broken the system is expected to have two genuine NGBs, that is two massless phonons. If  $\lambda \neq 0$ , the system has only the  $U(1)_{\mathcal{V}}$  symmetry and in the superfluid ground-state only one NGB can appear. Furthermore, if the Rabi term can be treated as a soft explicit symmetry breaking, the low-energy theory is expected to include a pseudo-NGB, with a square mass proportional to  $\lambda$ .

## IV. PRESSURE OF BINARY SUPERFLUIDS

The considered system consists of two interacting gases having both intra-species and inter-species interactions, as well as a perturbative Rabi coupling. In absence of both the inter-species and Rabi interactions, the two fluids would be non-interacting and the total pressure of the system would be equal to the sum of two partial ones associated to each flavor. By adding the interactions between the two components, we might naively expect that the total pressure be cannot be separated into two independent contributions. Nevertheless, if the Rabi coupling vanishes and the G symmetry holds, the ground-state Lagrangian can be cast as a sum of two terms, and the total pressure is the sum of two partial pressures. Furthermore, the same separation can be extended for perturbative Rabi coupling. In the following, we investigate which are the conditions for such separation.

#### A. Flavor rotation in the ground state

As for a single fluid in the broken phase, it is useful to adopt the Madelung representation for both  $\Phi_A$  and  $\Phi_B$ . Introducing two pairs of real scalar fields,  $\rho_i$ ,  $\theta_i$ , such that

$$\Phi_i = \frac{\rho_i}{\sqrt{2}} e^{i\theta_i} , \qquad i = A, B , \qquad (48)$$

the Lagrangian in Eq. (29) with the potential in Eq. (32) reads

$$\mathcal{L} = \sum_{i=A,B} \left[ \frac{1}{2} \partial_{\nu} \rho_i \partial^{\nu} \rho_i + \frac{1}{2} \rho_i^2 \partial_{\nu} \theta_i \partial^{\nu} \theta_i - \frac{1}{2} m_i^2 \rho_i^2 + \frac{1}{2} \frac{\mu_i^2}{\gamma_i^2} \rho_i^2 - \frac{1}{4} U_{ii} \rho_i^4 \right] - \frac{U_{AB}}{4} \rho_A^2 \rho_B^2 - \lambda \rho_A \rho_B \cos(\theta_B - \theta_A) \,, \tag{49}$$

where we have replaced the covariant derivatives with standard derivatives. As illustrated in Sec. II, chemical potentials and fluid velocities can be introduced as expectation values of the phase fields. In the last line, the Rabi term explicitly violates the conservation of the currents

$$J_i^{\nu} = \frac{\delta \mathcal{L}}{\delta \partial_{\nu} \theta_i}, \qquad i = A, B, \qquad (50)$$

while conserving the total one  $J_B^{\nu} + J_A^{\nu}$ . Their difference satisfies

$$\partial_{\nu}(J_B^{\nu} - J_A^{\nu}) = 2\lambda\rho_A\rho_B\sin(\theta_B - \theta_A), \qquad (51)$$

which, for  $\rho_A \simeq \rho_B$  both homogeneous, reduces to the sine-Gordon equation

$$\Box(\theta_B - \theta_A) - 2\lambda\sin(\theta_B - \theta_A) = 0, \qquad (52)$$

suggesting that solitonic solutions could be present [74, 75]. Unlike the phase of a single fluid, the relative phase  $(\theta_B - \theta_A)$  is an observable that is accessible in ultracold atom experiments [60].

In the present section, we focus on the background properties, while the effective action for the long-wavelength excitations will be discussed in Sec. V. Hereafter, we will indicate with  $\rho_A$ ,  $\rho_B$ ,  $\theta_A$  and  $\theta_B$  the mean field values. In analogy with Eq. (11), we write the gradients of the phase fields as

$$\partial^{\nu}\theta_{i} = \frac{\mu_{i}}{\gamma_{i}}v_{i}^{\nu}, \qquad i = A, B, \qquad (53)$$

where  $v_A^{\mu}$  and  $v_B^{\mu}$  are the two fluids four-velocities. Hence, from Eq. (49), we obtain the background Lagrangian

$$\mathcal{L}_{0} = \frac{\rho_{A}^{2}}{2\gamma_{A}^{2}} (\mu_{A}^{2} - m_{A}^{2}\gamma_{A}^{2}) + \frac{\rho_{B}^{2}}{2\gamma_{B}^{2}} (\mu_{B}^{2} - m_{B}^{2}\gamma_{B}^{2}) - \frac{1}{4}\rho_{A}^{4}U_{AA} - \frac{1}{4}\rho_{B}^{4}U_{BB} - \frac{1}{4}\rho_{B}^{2}\rho_{A}^{2}U_{AB} - \lambda \rho_{A}\rho_{B}\cos(\theta_{B} - \theta_{A}).$$
(54)

Before estimating the mean field values of the radial fields, we remark a subtle aspect about the phase fields. They are not variational parameters, as they are connected by Eq. (53) with the fluids' flow (the same holds in the single-fluid case). Since the Rabi coupling explicitly depends on the relative phase, it tends to fix such a difference. For instance, in the broken phase, the mean

field values  $\rho_A$  and  $\rho_B$  are positive, thus if  $\lambda < 0$  the background pressure is maximized for

$$\theta_A = \theta_B \,. \tag{55}$$

For this reason we shall refer to the last term in Eq. (54) as the *Rabi pressure*. If at each space-time point the two phases are equal, Eq. (53) implies that

$$\frac{\mu_A}{\gamma_A} v_A^{\nu} = \frac{\mu_B}{\gamma_B} v_B^{\nu} \,. \tag{56}$$

Since for each flavor  $v_i^{\nu} = \gamma_i(1, -v_i)$ , i = A, B, Eq. (55) in turn implies that  $\mu_A = \mu_B$  and  $v_A = v_B$ . Thus, the Rabi pressure tends to make the two fluids flow with the same velocity and to have equal chemical potentials. In the background Lagrangian of Eq. (54), we consider  $\theta_A \neq \theta_B$ , because both the chemical potentials and the fluids' velocities are assumed to be externally fixed.

Following the same procedure of SecIIA, we now determine the pressure by maximizing  $\mathcal{L}_0$  with respect to  $\rho_A$ and  $\rho_B$ , which are the two independent variational parameters. We introduce the flavor vector notation

$$\boldsymbol{\rho}_{2} = \begin{pmatrix} \rho_{A}^{2} \\ \rho_{B}^{2} \end{pmatrix}, \quad \boldsymbol{\mu}_{2} = \begin{pmatrix} \mu_{A}^{2} \\ \mu_{B}^{2} \end{pmatrix}, \quad \boldsymbol{m}_{2} = \begin{pmatrix} m_{A}^{2} \\ m_{B}^{2} \end{pmatrix}, \quad (57)$$

where the subscripts indicate that they contain squared quantities. For the sake of brevity, we define

$$\lambda_{AB} = \lambda \cos(\theta_B - \theta_A), \qquad (58)$$

and we redefine the chemical potentials

$$\frac{\mu_i}{\gamma_i} \to \mu_i \,.$$
 (59)

The Lagrangian in Eq. (54) now reads

$$\mathcal{L}_0 = \frac{1}{2} \boldsymbol{\rho}_2 \cdot (\boldsymbol{\mu}_2 - \boldsymbol{m}_2) - \frac{1}{4} \boldsymbol{\rho}_2^t \boldsymbol{U} \boldsymbol{\rho}_2 - \lambda_{AB} \rho_A \rho_B , \quad (60)$$

where

$$\boldsymbol{U} = \begin{pmatrix} U_{AA} & U_{AB}/2 \\ U_{AB}/2 & U_{BB} \end{pmatrix}, \qquad (61)$$

and we left the last term as a function of  $\rho_A$  and  $\rho_B$ . Since we treat the Rabi term as a small perturbation, we proceed first to diagonalize  $\mathcal{L}_0$  in the sum of two independent terms neglecting the contribution of the Rabi. Any transformation

$$\boldsymbol{\rho}_2' = R(\alpha)\boldsymbol{\rho}_2, \quad \boldsymbol{\mu}_2' = R(\alpha)\boldsymbol{\mu}_2, \quad \boldsymbol{m}_2' = R(\alpha)\boldsymbol{m}_2, \quad (62)$$

where

$$R(\alpha) = \begin{pmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & -\cos \alpha \end{pmatrix} = \sigma_3 \cos \alpha + \sigma_1 \sin \alpha \,, \quad (63)$$

is a matrix in flavor space depending on the mixing angle  $\alpha$ , leaves invariant the first term of the Lagrangian in Eq. (60). This transformation clearly mixes the different

flavors: the primed vectors are a mixing of the vector components in the not-primed basis. We refer to this transformation as *rotation* in flavor space.

Although the first term in Eq. (60) locks the flavor rotation angles of  $\rho_2$ ,  $\mu_2$  and  $m_2$  to be the same, the value of  $\alpha$  is not fixed. On the other hand, the diagonalization of

$$V_U = \frac{1}{4} \boldsymbol{\rho}_2^t \boldsymbol{U} \boldsymbol{\rho}_2 \,, \tag{64}$$

sets

$$\alpha = \pm \arcsin\left(\sqrt{\frac{1}{2} - \frac{\Delta U}{\sqrt{4\Delta U^2 + U_{AB}^2}}}\right), \qquad (65)$$

where

$$\Delta U = \frac{U_{AA} - U_{BB}}{2} \,. \tag{66}$$

For later convenience, we also define the following quantities

$$U = \frac{U_{AA} + U_{BB}}{2}, \qquad U_{\pm} = U \pm \sqrt{\Delta U^2 + \frac{U_{AB}^2}{4}}.$$
 (67)

Given the choice of ordering of the couplings in Eq. (37), we have  $\Delta U \geq 0$ ; moreover, in order to have positivedefined squared radial fields, chemical potentials and masses, we take the positive sign in the right-hand side of Eq. (65).

Indicating the components of the rotated vectors as

$$\boldsymbol{\rho}_{2}^{\prime} = \begin{pmatrix} \rho_{I}^{2} \\ \rho_{II}^{2} \end{pmatrix}, \quad \boldsymbol{\mu}_{2}^{\prime} = \begin{pmatrix} \mu_{I}^{2} \\ \mu_{II}^{2} \end{pmatrix}, \quad \boldsymbol{m}_{2}^{\prime} = \begin{pmatrix} m_{I}^{2} \\ m_{II}^{2} \end{pmatrix},$$
(68)

the rotated squared radial fields are given by

$$\rho_I^2 = \rho_A^2 \cos \alpha + \rho_B^2 \sin \alpha ,$$
  

$$\rho_{II}^2 = -\rho_B^2 \cos \alpha + \rho_A^2 \sin \alpha ,$$
(69)

and, with our choice of the interaction couplings, it follows that

$$\rho_I^2 \ge \rho_{II}^2 \,. \tag{70}$$

Similar expressions hold for the squared masses and chemical potentials.

Now, we consider some specific cases. For  $U_{AA} = U_{BB}$ , i.e.  $\Delta U = 0$ , the mixing angle is maximal  $\alpha = \pi/4 \pm n\pi$ , with *n* integer, and the rotation matrix is

$$R = \pm \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1\\ 1 & -1 \end{pmatrix} = \pm \frac{1}{\sqrt{2}} (\sigma_3 + \sigma_1) \,. \tag{71}$$

Close to maximal mixing,

$$\rho_I^2 \simeq \frac{1}{\sqrt{2}} (\rho_A^2 + \rho_B^2), \quad \rho_{II}^2 \simeq \frac{1}{\sqrt{2}} (\rho_A^2 - \rho_B^2), \quad (72)$$

and for  $\rho_A \simeq \rho_B$ , this implies that  $\rho_I^2 \gg \rho_{II}^2$ .

For general values of  $\rho_A$  and  $\rho_B$ , the rotation angle

$$\alpha = \arctan\left(\frac{\rho_B^2}{\rho_A^2}\right),\tag{73}$$

makes  $\rho_I$  maximal while  $\rho_{II}$  vanishes. Regarding the masses in the rotated basis, in analogy with Eqs. (70) and (73), the mass of the radial field II is smaller than the mass of the field I, and it vanishes when

$$\alpha = \arctan\left(\frac{m_B^2}{m_A^2}\right). \tag{74}$$

We remark that the vanishing of  $m_{II}$  is not associated to any specific phenomenon: it does not signal a phase transition. The phase transition to the broken phase occurs when the rotated chemical potentials exceed the corresponding masses. This can be observed by rewriting the Lagrangian in Eq. (60) in the I, II flavor basis

$$\mathcal{L}_{0} = \frac{\rho_{I}^{2}}{2} (\mu_{I}^{2} - m_{I}^{2}) - \frac{\rho_{I}^{4}}{4} U_{+} + \frac{\rho_{II}^{2}}{2} (\mu_{II}^{2} - m_{II}^{2}) - \frac{\rho_{II}^{4}}{4} U_{-} - V_{\lambda} , \qquad (75)$$

which, apart from the Rabi term, is the sum of two independent pieces. We now determine the quantum pressure of the system for vanishing Rabi coupling; we will include it in Sec. IV C.

### B. Vanishing Rabi coupling

For vanishing Rabi coupling, the Lagrangian in Eq. (75) can be cast as that of two non-interacting fluids

$$\mathcal{L}_{0} = \mathcal{L}_{I} + \mathcal{L}_{II} = \frac{\rho_{I}^{2}}{2} (\mu_{I}^{2} - m_{I}^{2}) - \frac{\rho_{I}^{4}}{4} U_{+} + \frac{\rho_{II}^{2}}{2} (\mu_{II}^{2} - m_{II}^{2}) - \frac{\rho_{II}^{4}}{4} U_{-}, \quad (76)$$

where the two Lagrangians,  $\mathcal{L}_I$  and  $\mathcal{L}_{II}$ , are independent. We determine the expectation values of the radial fields in the same manner as in the single-fluid case discussed in Sec. II. The values maximizing the Lagrangian are

$$\rho_I^2 = \begin{cases} \frac{\mu_I^2 - m_I^2}{U_+} & \text{if } \mu_I \ge m_I \\ 0 & \text{if } \mu_I \le m_I \end{cases}, \quad (77)$$

and

$$\rho_{II}^2 = \begin{cases} \frac{\mu_{II}^2 - m_{II}^2}{U_-} & \text{if } \mu_{II} \ge m_{II} \\ 0 & \text{if } \mu_{II} \le m_{II} \end{cases},$$
(78)

and are the analogous of the expression for the single scalar field in Eq. (13). In the I, II basis the conditions to have nonvanishing expectation values of the scalar

fields  $\rho_I$  and  $\rho_{II}$ , are  $\mu_I > m_I$  and  $\mu_{II} > m_{II}$ , respectively. When expressing these conditions in the A, B basis, we have that

$$(\mu_A^2 - m_A^2)\cos\alpha + (\mu_B^2 - m_B^2)\sin\alpha > 0, (\mu_A^2 - m_A^2)\sin\alpha - (\mu_B^2 - m_B^2)\cos\alpha > 0,$$
(79)

which can be cast as

$$\tan \alpha > -\frac{\mu_A^2 - m_A^2}{\mu_B^2 - m_B^2} \quad \text{and} \quad \tan \alpha > \frac{\mu_B^2 - m_B^2}{\mu_A^2 - m_A^2}, \quad (80)$$

respectively. Since

$$\frac{\mu_B^2 - m_B^2}{\mu_A^2 - m_A^2} > -\frac{\mu_A^2 - m_A^2}{\mu_B^2 - m_B^2},\tag{81}$$

it is possible that  $\rho_I > 0$ , while  $\rho_{II} = 0$ . In the following, we will assume for simplicity that the second condition in (80) is satisfied, therefore both  $\rho_I$  and  $\rho_{II}$  are positive.

The quantum pressure of the system is given by the value taken by the background Lagrangian at the stationary points and it is the sum of two partial pressures

$$P = P_I + P_{II} , \qquad (82)$$

with

$$P_I = \frac{(\mu_I^2 - m_I^2)^2}{4U_+}$$
 and  $P_{II} = \frac{(\mu_{II}^2 - m_{II}^2)^2}{4U_-}$ . (83)

Using standard thermodynamic relations, we obtain the number density for each rotated component

$$n_I = \mu_I \rho_I^2$$
,  $n_{II} = \mu_{II} \rho_{II}^2$ , (84)

showing that when  $|U_{AB}| = 2\sqrt{U_{AA}U_{BB}}$ , the number density  $n_{II}$  diverges. This signals the transition to a dilute liquidlike droplet state [33].

#### C. Including the Rabi coupling

We examine the contribution of the Rabi interaction to the background pressure. In the I, II basis, the Rabi term takes the following form:

$$V_{\lambda} = \frac{\lambda_{AB}}{\sqrt{2}} \sqrt{(\rho_I^4 - \rho_{II}^4) \sin(2\alpha) - 2\rho_I^2 \rho_{II}^2 \cos(2\alpha)}, \quad (85)$$

and is a non-trivial coupling between the  $\rho_I$  and  $\rho_{II}$ fields. If the explicit breaking is small, we can treat  $V_{\lambda}$ as a perturbation. In this case, we expand the mean field solutions at leading order in  $\lambda$  as follows

$$\rho_j \to \rho_j + \frac{\lambda_{AB}}{\rho_j^2} \delta \rho_j , \quad j = I, II$$
(86)

where

$$\frac{|\lambda_{AB}|}{\rho_I^2} \ll 1 \quad \text{and} \quad \frac{|\lambda_{AB}|}{\rho_{II}^2} \ll 1.$$
 (87)

while  $\delta \rho_I$  and  $\delta \rho_{II}$  are two variations to be determined. Upon substituting these expressions in the equations for the stationary point of the Lagrangian, we find that the leading order contributions are

$$\delta\rho_I = -\frac{\rho_I^3 \sin(2\alpha) - \rho_I \rho_{II}^2}{U_+ \sqrt{2(\rho_I^4 - \rho_{II}^4) \sin(2\alpha) - 4\rho_I^2 \rho_{II}^2 \cos(2\alpha)}},$$
(88)

and

$$\delta\rho_{II} = +\frac{\rho_{II}^3 \sin(2\alpha) + \rho_{II}\rho_I^2}{U_-\sqrt{2(\rho_I^4 - \rho_{II}^4)\sin(2\alpha) - 4\rho_I^2\rho_{II}^2\cos(2\alpha)}}.$$
(89)

Even when the Rabi coupling satisfies the conditions in Eq. (87), the subsystems I and II are coupled:  $\delta\rho_I$  and  $\delta\rho_{II}$  are functions of both  $\mu_I$  and  $\mu_{II}$ .

If we want to still write the pressure as the sum of two independent terms as in Eq. (82), we have to impose an additional constraint:

$$\rho_{II} \ll \rho_I \,. \tag{90}$$

This, in turn, implies that  $\rho_A \simeq \rho_B$  (see Eq. (73)) and thus

$$\delta \rho_I \simeq -\frac{\rho_I}{U_+} \sqrt{\frac{\sin(2\alpha)}{2}},$$
  
$$\delta \rho_{II} \simeq +\frac{\rho_{II} \cos(2\alpha)}{U_- \sqrt{2} \sin(2\alpha)},$$
 (91)

meaning that each radial field depends on the corresponding chemical potential, see Eqs. (77) and (78). Upon replacing these expressions in Eq. (86) we obtain the shifts of the radial fields due to the Rabi term

$$\rho_I \to \rho_I \left( 1 + \frac{\lambda_{AB}}{U_+ \rho_I^2} \sqrt{\frac{\sin(2\alpha)}{2}} \right),$$
(92)

$$\rho_{II} \to \rho_{II} \left( 1 + \frac{\lambda_{AB}}{U_- \rho_{II}^2} \frac{\cos(2\alpha)}{\sqrt{2\sin(2\alpha)}} \right) \,. \tag{93}$$

Comparing these expressions with the estimate in Eq. (35), we better comprehend the condition to treat perturbatively the Rabi term. In the broken phase, the actual dimensionless expansion parameter is the ratio between the Rabi coupling and the expectation values of the rotated radial fields. Since  $\rho_{I,II}$  are of the order of  $m_{A,B}$ , the condition  $\tilde{\lambda} \ll \tilde{U}_{ij}$  holds.

Upon substituting Eqs. (92) and (93) in Eq. (85), one obtains that the two subsystems are decoupled and the pressure can be expressed as the sum of two independent contributions:

$$P_{I} \to P_{I}(\lambda_{AB} = 0) + \lambda_{AB} \,\delta P_{I} ,$$
  

$$P_{II} \to P_{II}(\lambda_{AB} = 0) + \lambda_{AB} \,\delta P_{II} , \qquad (94)$$

where  $\lambda_{AB}$  is defined in Eq. (58) and  $\delta P_I$  and  $\delta P_{II}$  are the pressure variations depending only on  $\mu_I$  and  $\mu_{II}$ , respectively. In this case, each subsystem is characterized by its thermodynamic quantities that do not depend on the properties of the other subsystem.

Alternatively, a more straightforward approach can be used to derive the perturbative effect of the Rabi coupling on the pressure. Combining Eq. (85) with the decoupling condition  $\rho_{II} \ll \rho_{I}$ , we have that

$$V_{\lambda} \simeq \lambda_{AB} \rho_I^2 \sqrt{\sin(2\alpha)/2} - \lambda_{AB} \rho_{II}^2 \sqrt{\sin(2\alpha)/2} \cot(2\alpha) \,. \tag{95}$$

Consistently with the observations in Sec. III B, these terms can be interpreted as small shifts of the  $m_I$  and  $m_{II}$  masses, that is

$$m_I^2 \to m_I^2 + \lambda_{AB} \sqrt{\sin(\alpha) \cos(\alpha)} ,$$
  
$$m_{II} \to m_{II}^2 - \lambda_{AB} \sqrt{\sin(\alpha) \cos(\alpha)} \cot(2\alpha) .$$
(96)

In fact, the corresponding partial pressures can be readily obtained from those in Eq. (83), with the above shifted masses. In detail, we have that

$$P_I = \frac{(\mu_I^2 - m_I^2)^2}{4U_+} - \frac{\lambda_{AB}\sqrt{\sin(2\alpha)}}{2\sqrt{2}U_+}(\mu_I^2 - m_I^2), \quad (97)$$

and

$$P_{II} = \frac{(\mu_{II}^2 - m_{II}^2)^2}{4U_-} + \frac{\lambda_{AB}\sqrt{\sin(2\alpha)}\cot(2\alpha)}{2\sqrt{2}U_-}(\mu_{II}^2 - m_{II}^2).$$
(98)

We remark that for both non-vanishing  $U_{AB}$  and  $\lambda$ , in general, it is not possible to simultaneously diagonalize  $V_U$  and  $V_{\lambda}$ . They differ in the dependence in  $\rho_A$  and  $\rho_B$ : quadratic for the inter-species interaction, and linear for the Rabi term. Consequently, the diagonalization of  $V_U$  requires a non linear transformation of the fields, as in Eq. (69), leaving the Rabi term non diagonal. The specular situation occurs when the Rabi term dominates over the inter-species interaction. If we diagonalize the Rabi term, the transformed fields remain interacting due to a non-vanishing  $U_{AB}$  [57].

### V. QUANTUM FLUCTUATIONS IN BINARY SUPERFLUIDS

In the previous section, we characterized the mean field Lagrangian of the boson mixture. Following the same steps of the single-fluid model, we analyze the effective action for the longwavelength excitations. We replace

$$\rho_i \to \rho_i + \tilde{\rho}_i, \quad \theta_i \to \theta_i + \theta_i, \quad i = A, B,$$
(99)

and throughout the remaining on this work,  $\rho_i$  and  $\theta_i$ indicate the mean field solutions, while  $\tilde{\rho}_i$  and  $\tilde{\theta}_i$  are their corresponding fluctuations. We expand the Lagrangian as

$$\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_2 \,, \tag{100}$$

with  $\mathcal{L}_0$  the mean-field one in Eq. (54), while  $\mathcal{L}_2$  is quadratic in the fields' fluctuations. In the following sections, we will discuss the case for vanishing Rabi interaction ( $\lambda = 0$ ), and the non-vanishing case ( $\lambda \neq 0$ ). We will show that  $\mathcal{L}_2$  can be written as the one of two non-interacting scalar fields, propagating on an emergent acoustic metric. The Rabi term will induce an effective mass in one of the scalar fields.

#### A. Fluctuations with vanishing Rabi term

Given Eq. (99), we compute the Lagrangian,  $\mathcal{L}_2$ , quadratic in the fluctuations. We consider a homogeneous and stationary background, with both fluids moving at the same velocity  $v^{\nu}$ . For  $\lambda = 0$ , we have

$$\mathcal{L}_{2} = \frac{1}{2} \tilde{\rho}_{A}^{2} \left[ (\mu_{A}^{2} - m_{A}^{2}) - \frac{3}{2} U_{AA} \rho_{A}^{2} - \frac{U_{AB}}{4} \rho_{B}^{2} \right] + \frac{1}{2} \tilde{\rho}_{B}^{2} \left[ (\mu_{B}^{2} - m_{B}^{2}) - \frac{3}{2} U_{BB} \rho_{B}^{2} - \frac{U_{AB}}{4} \rho_{A}^{2} \right] - U_{AB} \rho_{A} \rho_{B} \tilde{\rho}_{A} \tilde{\rho}_{B} + \frac{1}{2} \rho_{A}^{2} \partial_{\nu} \tilde{\theta}_{A} \partial^{\nu} \tilde{\theta}_{A} + \frac{1}{2} \rho_{B}^{2} \partial_{\nu} \tilde{\theta}_{B} \partial^{\nu} \tilde{\theta}_{B} + 2 \rho_{A} \tilde{\rho}_{A} \mu_{A} v^{\nu} \partial_{\nu} \tilde{\theta}_{A} + 2 \rho_{B} \tilde{\rho}_{B} \mu_{B} v^{\nu} \partial_{\nu} \tilde{\theta}_{B} , \quad (101)$$

and we disregarded the kinetic terms of the radial fluctuations because they give subleading corrections to the low-energy effective theory (see Sec. II B). For later convenience, we introduce the rescaled fluctuations

$$\hat{\rho}_i = \mu_i \tilde{\rho}_i, \quad \hat{\theta}_i = \rho_i \tilde{\theta}_i, \quad i = A, B,$$
 (102)

and the corresponding flavor vectors

$$\hat{\boldsymbol{\rho}} = \begin{pmatrix} \hat{\rho}_A\\ \hat{\rho}_B \end{pmatrix}, \qquad \hat{\boldsymbol{\theta}} = \begin{pmatrix} \hat{\theta}_A\\ \hat{\theta}_B \end{pmatrix}.$$
 (103)

Thus, the quadratic Lagrangian in Eq. (101) can be cast as

$$\mathcal{L}_2 = \frac{1}{2} \hat{\boldsymbol{\rho}}^t H \hat{\boldsymbol{\rho}} + 2v^{\nu} \hat{\boldsymbol{\rho}} \cdot \partial_{\nu} \hat{\boldsymbol{\theta}} + \frac{1}{2} \partial_{\nu} \hat{\boldsymbol{\theta}} \cdot \partial^{\nu} \hat{\boldsymbol{\theta}} , \qquad (104)$$

where H is the  $2 \times 2$  symmetric matrix with elements

$$H_{11} = \frac{1}{\mu_A^2} \left[ \mu_A^2 - m_A^2 - \frac{3}{2} U_{AA} \rho_A^2 - \frac{U_{AB}}{4} \rho_B^2 \right] ,$$
  

$$H_{22} = H_{11} (A \leftrightarrow B) ,$$
  

$$H_{12} = -U_{AB} \frac{\rho_A \rho_B}{2\mu_A \mu_B} .$$
(105)

We diagonalize  $\mathcal{L}_2$  with a suitable transformation in flavor space of the fields  $\hat{\rho}$  and  $\hat{\theta}$ . The first and last term in the r.h.s. of Eq. (104) are separately invariant under any flavor mixing transformation of  $\hat{\rho}$  and  $\hat{\theta}$ , as the one in Eq. (63). However, the invariance of the second term locks the mixing angle to be the same for  $\hat{\rho}$  and  $\hat{\theta}$ . The locking implies that the vectors in the new basis are

$$\tilde{\rho}' = R(\beta)\hat{\rho} \text{ and } \tilde{\theta}' = R(\beta)\hat{\theta}, \quad (106)$$

where  $\beta$  is the common rotation angle. The flavor rotation angle is still arbitrary, and we can use such freedom to rotate the  $\hat{\rho}$  field to diagonalize *H*, similarly to the the procedure in Sec. IV A. This is achieved choosing

$$\beta = \pm \arcsin\left(\frac{1}{\sqrt{2}}\sqrt{1 - \frac{H_{11} - H_{22}}{\sqrt{(H_{11} - H_{22})^2 + 4H_{12}^2}}}\right).$$
(107)

We remark that the flavor rotation angle required to disentangle the radial fluctuations, differs from the one in Eq. (63) used to diagonalize the background. As a consequence, the components  $\tilde{\rho}_{a/b}$  and  $\tilde{\theta}_{a/b}$  of the rotated fields

$$\tilde{\boldsymbol{\rho}}' = \begin{pmatrix} \tilde{\rho}_a \\ \tilde{\rho}_b \end{pmatrix}, \quad \tilde{\boldsymbol{\theta}}' = \begin{pmatrix} \tilde{\theta}_a \\ \tilde{\theta}_b \end{pmatrix}, \quad (108)$$

are not the fluctuations of  $\rho_{I/II}$  and  $\theta_{I/II}$ . They can be written as a linear combination of  $\tilde{\rho}_{a/b}$  and  $\tilde{\theta}_{a/b}$ , respectively.

In the a, b basis the matrix H is diagonal, thus no mixing exists, in other words

$$\mathcal{L}_2 = \mathcal{L}_{2a} + \mathcal{L}_{2b} \,, \tag{109}$$

where

$$\mathcal{L}_{2a} = -\frac{1}{2}\tilde{m}_a^2\tilde{\rho}_a^2 + 2v^{\nu}\tilde{\rho}_a\partial_{\nu}\tilde{\theta}_a + \frac{1}{2}\partial_{\nu}\tilde{\theta}_a\partial^{\nu}\tilde{\theta}_a\,,\qquad(110)$$

and  $\mathcal{L}_{2b}$  is given by an analogous expression. They have the same formal expression obtained in the single-fluid case, see Eq. (19), when neglecting the kinetic term of the radial mode. Note that the radial modes have dimension 2, therefore their *masses* 

$$\tilde{m}_{a}^{2} = -\frac{H_{11} + H_{22}}{2} + \frac{1}{2}\sqrt{(H_{11} - H_{22})^{2} + 4H_{12}^{2}},$$
  
$$\tilde{m}_{b}^{2} = -\frac{H_{11} + H_{22}}{2} - \frac{1}{2}\sqrt{(H_{11} - H_{22})^{2} + 4H_{12}^{2}}, \quad (111)$$

are dimensionless. Upon integrating out the radial modes, the effective Lagrangian can be written as

$$\mathcal{L}_2 = \frac{1}{2} \left( \eta^{\mu\nu} + \left( \frac{1}{c_{sa}^2} - 1 \right) v^{\mu} v^{\nu} \right) \partial_{\mu} \tilde{\theta}_a \partial_{\nu} \tilde{\theta}_a + (a \to b) ,$$
(112)

where the two sound speeds are given by

$$c_{sa}^2 = \frac{2\tilde{m}_a^2}{1+\tilde{m}_a^2}, \qquad c_{sb}^2 = \frac{2\tilde{m}_b^2}{1+\tilde{m}_b^2}, \qquad (113)$$

and given the expression of the H-matrix components in (105), each sound mode depends on the intra- and interspecies couplings, as well as on the chemical potentials  $\mu_A$  and  $\mu_B$  and the masses  $m_A$  and  $m_B$ .

### B. Including the Rabi term in the fluctuations

Including fluctuations in the A, B flavor basis, the Rabi term is given by

$$V_{\lambda} = \lambda \left(\rho_A + \tilde{\rho}_A\right) \left(\rho_B + \tilde{\rho}_A\right) \cos(\theta_A - \theta_B + \tilde{\theta}_A - \tilde{\theta}_B), \quad (114)$$

and it produces a pletora of terms. The linear terms in the fluctuations vanish at the stationary point. There are two quadratic terms in the fluctuations, which have distinct effects. One term is proportional to  $\tilde{\rho}_A \tilde{\rho}_B$  and thus contributes to the  $H_{12}$  matrix element, see Eq. (105). This produces a trivial small shift of the effective masses of the radial fluctuations. The second quadratic term in the fluctuations is

$$\mathcal{L}_{\lambda\theta^2} = \frac{\lambda_{AB}}{2} \rho_A \rho_B (\tilde{\theta}_A - \tilde{\theta}_B)^2 , \qquad (115)$$

where  $\lambda_{AB}$  is defined in Eq. (58). This is an effective mass term for the mode corresponding to the relative phase fluctuations. However, this term is expressed in the *A*, *B* basis, but to diagonalize the Lagrangian in the fluctuations and write it as in Eq. (109), we performed a rotation in flavor space by the angle  $\beta$  in Eq. (107). Therefore, we need to express the Rabi quadratic term (115) in the rotated basis *a*, *b*.

Before doing that, we observe that to have the total pressure as the sum of two partial pressure in the background analysis, we had to assume that  $\rho_I \gg \rho_{II}$  and, correspondingly, a background flavor rotation angle close to maximal mixing, that is  $\alpha \simeq \pi/4$ . We expect that the same conditions would lead to an effective theory with two decoupled modes, one massless and one massive. To this end, we follow the same procedure discussed for vanishing Rabi coupling. In particular, we rescale the phases as in Eq. (102) and obtain that, close to maximal background mixing, the condition  $\rho_I \gg \rho_{II}$  implies  $\rho_B \simeq \rho_A$ , see Eq. (72). Then, Eq. (115) can be rewritten as

$$\mathcal{L}_{\lambda\theta^2} \simeq \lambda_{AB} \frac{\rho_I^2}{\rho_A^2} \sqrt{\frac{\sin 2\alpha}{2}} (\hat{\theta}_B - \hat{\theta}_A)^2 , \qquad (116)$$

thus it corresponds to the mass term for the relative phase mode in the rescaled basis, see Eq. (102). Now we perform a flavor rotation to the a, b basis as in Eq. (106). This is a necessary operation, because to integrate out the radial mode we have first to obtain a Lagrangian that is diagonal in flavor basis. Note that we are forced to do this flavor rotation for the following chain of reasons: we need to rotate the radial fluctuations to diagonalize the H matrix, and the radial and phase fluctuations are locked by the interaction term between radial and phase fluctuations. In the rotated basis

$$\hat{\theta}_B - \hat{\theta}_A = \tilde{\theta}_a(\sin\beta - \cos\beta) - \tilde{\theta}_b(\sin\beta + \cos\beta), \quad (117)$$

the approximations  $\rho_A \simeq \rho_B$  and of maximal background mixing lead to  $H_{11} \simeq H_{22}$ , thus the fluctuation flavor mixing angle  $\beta$  is close to maximal as well. In this case, we can neglect the mass of the  $\hat{\theta}_a$  field and rewrite the Rabi coupling as

$$\mathcal{L} \simeq 2 \,\lambda_{AB} \,\tilde{\theta}_b^2 \,. \tag{118}$$

Upon integrating out the radial fluctuations using the same procedure discussed for the single-fluid case in Sec. II B, we obtain the quadratic effective Lagrangian in the phases fluctuations

$$\mathcal{L}_{\rm eff} = \mathcal{L}_{\rm eff,a} + \mathcal{L}_{\rm eff,b} \,, \tag{119}$$

where

$$\mathcal{L}_{\text{eff,a}} = \frac{1}{2} \left( \eta^{\mu\nu} + \left( \frac{1}{c_{sa}^2} - 1 \right) v^{\mu} v^{\nu} \right) \partial_{\mu} \tilde{\theta}_a \partial_{\nu} \tilde{\theta}_a \,, \quad (120)$$

and

$$\mathcal{L}_{\text{eff,b}} = \frac{1}{2} \left( \eta^{\mu\nu} + \left( \frac{1}{c_{sb}^2} - 1 \right) v^{\mu} v^{\nu} \right) \partial_{\mu} \tilde{\theta}_b \partial_{\nu} \tilde{\theta}_b - \frac{1}{2} m_b^2 c_{sb}^2 \tilde{\theta}_b^2$$
(121)

with

$$m_b^2 = -\frac{4\lambda_{AB}}{c_{sb}^2} = -\frac{4\lambda}{c_{sb}^2}\cos(\theta_A - \theta_B), \qquad (122)$$

the mass squared of the  $\tilde{\theta}_b$  phonon. A similar expression was obtained in [14] with a different method. The above expression implies that  $\lambda \cos(\theta_A - \theta_B)$  must be negative, otherwise the  $\tilde{\theta}_b$  field would be tachyonic. This is the same condition needed to have a positive contribution of the Rabi term to the background pressure, see Eq. (54). Taking  $\lambda < 0$ , the Rabi pressure is maximal for  $\theta_A = \theta_B$ , see Eq. (55), which is also the condition that maximizes the mass of the  $\tilde{\theta}_b$  field. Alternatively, the favored state may be a crystal of sine-Gordon kinks, see Eq. (52), and the negative value of the squared mass would actually indicate the presence of phonon excitations of the crystal [76].

An interesting point is that, from Eq. (53), the difference ence of the background phases is related to a difference in the chemical potentials and fluid velocities. Since in the derivation we have assumed that the two velocities are equal (it was not a necessary condition but it helped to simplify the analysis), the condition to have a positive defined squared mass can be related to a condition on the chemical potentials. As we have previously discussed, it is possible to prepare the system in such a way that

$$\theta_A = \theta_B + \Theta(x) \,, \tag{123}$$

where  $\Theta(x)$  is determined by the chemical potential difference between the two fluids: from Eq. (53), we have that for equal velocities of the two fluids

$$\mu_A = \mu_B + \gamma \, v_\mu \partial^\mu \Theta(x) \,, \tag{124}$$

where  $\gamma$  is their common Lorentz factor. Considering for simplicity the nonrelativistic limit, we have that

$$\Theta = \Theta_0 + (\mu_A - \mu_B) t, \qquad (125)$$

where  $\Theta_0$  is some constant phase that can be reabsorbed in the definition of the background phases. Then, the Rabi coupling will tend to make the two chemical potentials equal; indeed any chemical potential difference would produce a mode that oscillates in time between a stable and a tachyonic configuration, signaling that the system is not in the ground-state.

### VI. SPONTANEOUS PHONON EMISSION

Having obtained the low-energy effective theory for binary superfluids, we now consider the effect of a background velocity profile. In particular, we are interested to determine the effect of the phonon mass on the spontaneous emission process at the acoustic horizon [14, 77]. Given the two Lagrangians in Eqs. (120) and (121) for the massless and massive phonons, respectively, the corresponding acoustic horizons are in the positions,  $x_a^H$  and  $x_b^H$ , such that

$$v(x_a^H) = c_{sa}$$
 and  $v(x_b^H) = c_{sb}$ , (126)

where we assumed that the flow is along the x-direction and thus the acoustic horizons are two planes centered at  $x_a^H$  and  $x_b^H$ . More precisely, the horizon positions can be determined with precision of the order of the healing length, that is the threshold scale length required for the validity of the hydrodynamic description of realistic superfluids [69]. Close to the acoustic horizons, we consider that the fluid velocity is given by  $\boldsymbol{v} = -v\hat{x}$  with velocity profile

$$v = c_s - Cx, \qquad (127)$$

with C a constant and  $c_s = (c_{sa} + c_{sb})/2$ . Then we have that

$$x_a^H = \frac{c_{sb} - c_{sa}}{2C} = -x_b^H, \qquad (128)$$

are the positions of the horizons of the two acoustic holes. Notice that in our three-dimensional setup, the horizon must be intended as a two-dimensional surface orthogonal to the fluid flow.

For each acoustic hole it is possible to define the corresponding Hawking temperature following the same reasoning used in one component superfluid, see for instance [1]. Since the two species a and b are independent, in principle they have distinct Hawking temperatures

$$T_{a,b} = \frac{1}{2\pi} \left( \frac{|c_{sa,b} - v|}{1 - |c_{sa,b}v|} \right)' \Big|_{H}, \qquad (129)$$

where the prime indicates the derivative with respect to the direction orthogonal to the horizon. However, in the nonrelativistic limit the two temperatures are equal and given by

$$T \simeq \frac{C}{2\pi} \,, \tag{130}$$

thus they only depend on the space gradient of the velocity field. Given the expression in Eq. (128), and considering Eqs. (113), in the nonrelativistic limit the relative distance between the two horizons is

$$x_a^H - x_b^H \simeq \frac{\tilde{m}_b - \tilde{m}_a}{\sqrt{2\pi}T} \,. \tag{131}$$

Since the discussion of the Hawking emission of massless phonons is equivalent to the one of a one component superfluid, hereafter we focus on the massive mode. For the sake of simplicity in the notation, in the following we drop the b index on the corresponding quantities.

#### A. Dispersion close to the acoustic horizon

The emission of massive particles at the acoustic horizon can be studied by one of the approaches used to explain the Hawking radiation, for instance one may determine how the tunneling probability depends on the mass of the emitted particles [19, 78].

Here we present a qualitative description of the phenomenon, based on semi-classical arguments. For this purpose, it is useful to analyze first the dispersion law of the phonons around the acoustic horizon. The equation of motion of phonons with mass  $m_{\rm ph}$  is

$$g_{\mu\nu}p^{\mu}p^{\nu} = m_{\rm ph}^2 c_s^2 \,, \qquad (132)$$

then, taking  $p_{\mu} = (E, -p, 0, 0)$  and the metric from Eqs. (27) and (121), we obtain the dispersion law

$$E_{\pm} = \frac{-(1-c_s^2) v p_x \pm \frac{1}{\gamma} \sqrt{c_s^2 p_x^2 + (1-c_s^2 v^2) m_{\rm ph}^2 c_s^4}}{1-c_s^2 v^2},$$
(133)

which is valid in the collinear regime, meaning that both the fluid velocity and the phonon momentum are along the x-axis. In the following, we restrict ourselves, to positive energy states in the nonrelativistic limit ( $v \ll 1$ and  $c_s \ll 1$ ). In this case the dispersion law of phonons simplifies:

$$E_{+} = -vp_{x} + \sqrt{c_{s}^{2}p_{x}^{2} + m_{\rm ph}^{2}c_{s}^{4}}, \qquad (134)$$

which has still a relativistic-like form, because we did not assume that  $|p_x| < m_{\rm ph}c_s$ . In this way we can take into account the case of an arbitrarily small phonon mass. The corresponding group velocity is given by

$$v_g = \frac{\mathrm{d}E_+}{\mathrm{d}p_x} = -v + \frac{c_s p_x}{\sqrt{p_x^2 + m_{\mathrm{ph}}^2 c_s^2}},$$
 (135)

indicating that at the acoustic horizon, corresponding to  $v = c_s$ , the group velocity of massive phonons is negative: they are trapped unless their momenta are larger than the threshold momentum

$$p_{\rm c} = \frac{m_{\rm ph} c_s v}{\sqrt{c_s^2 - v^2}} \simeq \frac{m_{\rm ph} c_s^{3/2}}{\sqrt{2Cx}} \simeq \frac{m_{\rm ph} c_s^{3/2}}{\sqrt{4\pi Tx}}, \qquad (136)$$

where in the last equation we exploited Eq. (130) (x being the distance form the horizon). Clearly, for  $m_{\rm ph} =$ 0 we have  $p_c = 0$  and this is independent of x. Instead, for nonvanishing phonon mass, the above expression implies that the largest threshold momentum is obtained at the shorter distance from the horizon, that is for  $x \simeq L_c$ , where  $L_c$  is a distance, already introduced after Eq. (130), of the order of the healing length [69] (see also [26] for a discussion of healing lengths in binary superfluids with Rabi coupling).

From the relation in Eq. (136), it follows that the threshold energy is

$$E_c = -p_c v + \sqrt{p_c^2 c_s^2 + m_{\rm ph}^2 c_s^4} \simeq m_{\rm ph} c_s^{3/2} \sqrt{4\pi T L_c} \,,$$
(137)

where we have considered the regime

$$c_s \ll \frac{1}{4\pi T L_c},\tag{138}$$

so that the phonon threshold momentum is much larger than  $m_{\rm ph} c_s$ . In other words, the phonon mass produces a small variation of the phonon energy.

#### B. Thermodynamics close to the horizon

At a distance from the horizon smaller than the phonon mean free path, phonons can be treated as an ideal Bose gas at the Hawking temperature [79]. Therefore, in this limit, the energy density of massive phonons is

$$\epsilon_{\rm ph} = \int Ef(p) \frac{d^3p}{(2\pi)^3} \simeq \frac{1}{2\pi L_c^2} \int_{p_c}^{\infty} \frac{E}{e^{E/T} - 1} dp_x \,, \ (139)$$

where the distribution function

$$f(p) = \frac{1}{e^{E/T} - 1} \frac{2\pi}{L_c} \delta(p_y) \frac{2\pi}{L_c} \delta(p_x) \theta(p_x - p_c), \quad (140)$$

takes into account that phonons are only emitted in the direction orthogonal to the horizon [19, 80], that is along the x-axis, and that the phonon momenta must be larger than the threshold momentum, given in Eq. (136). Upon changing variable,  $p_x \to E$ , and then defining z = E/T, this expression can be rewritten as

$$\epsilon_{\rm ph}(m_{\rm ph}) = \frac{T^2}{2\pi L_c^2 (c_s^2 - v^2)} \int_{z_c}^{\infty} \frac{vz + \frac{c_s z^2}{\sqrt{z^2 - z_c^2}}}{e^z - 1} dz \,, \ (141)$$

where  $z_c = E_c/T$ . Close to the acoustic horizon, that is at a distance  $x \sim L_c$ , the approximation  $c_s^2 - v^2 \simeq 4\pi T c_s L_c$  can be used, where we employed Eqs. (127) and (130). Then, at the leading order in  $E_c/T$  we find that

$$\epsilon_{\rm ph}(m_{\rm ph}) \simeq \epsilon_{\rm ph}(0) \left(1 - \frac{3E_c}{\pi^2 T}\right) ,$$
 (142)

where

$$\epsilon_{\rm ph}(0) = \frac{T}{24L_c^3},\qquad(143)$$

is the energy density of a massless phonon gas [79]. Naively, one may have expected a correction to the energy density of the order of  $m_{\rm ph}c_s^2/T$ , but it is instead given by

$$\frac{3E_c}{\pi^2 T} = \frac{6m_{\rm ph}c_s}{\pi^{3/2}} \sqrt{\frac{c_s L_c}{T}} \,, \tag{144}$$

then ruled not only by the Rabi coupling, entering the phonon mass, but also by the ratio between  $L_c$  and the Hawking temperature. Note that the correction is nonperturbative, in the sense that it is of order  $\lambda^{1/2}$ , see Eqs. (122) and (137). This result is similar to the one found in the calculation of the free-energy of a scalar field in thermal field theory, see for instance [81]. Such behavior reflects the breakdown of perturbation theory due to the presence of infrared divergences.

We can estimate the energy density reduction due to the Rabi coupling assuming that  $L_c = \xi$ , where  $\xi$  is the healing length. In the set-up of [82], where the experiment is performed with <sup>87</sup>Rb atoms, the healing length is  $\xi \simeq 2 \,\mu$ m, the speed of sound is  $c_s \simeq 0.3 \,\text{mm/s}$ , and the estimated Hawking temperature is  $T \simeq 0.1 \,mc_s^2 \simeq 0.1$ nK, where *m* is the mass of a <sup>87</sup>Rb atom. Then, considering that  $m_{\rm ph}c_s \simeq 2\sqrt{\lambda}$  and that  $\lambda = m\tilde{\lambda}$  (see Eq. (34)) with  $\tilde{\lambda} \sim 10^{-16} \,\text{eV}$  (see Eq. (35)), we have that

$$\frac{3E_c}{\pi^2 T} \simeq 0.2 \,, \tag{145}$$

therefore the effect is a reduction of the order of 20% with respect to the same value for vanishing Rabi coupling. Note that the effect is proportional to  $\tilde{\lambda}^{1/2}$  and it is independent of the mass of the used atoms. In a similar way, we can expand the phonon pressure

$$P_{\rm ph}(m_{\rm ph}) = \frac{1}{2\pi L_c^2} \int_{p_c}^{\infty} \frac{E}{e^{E/T} - 1} v_g^{-1} dp \,, \qquad (146)$$

obtaining

$$P_{\rm ph}(m_{\rm ph}) \simeq P_{\rm ph}(0) \left(1 - \frac{6E_c}{\pi^2 T}\right) ,\qquad(147)$$

where

$$P_{\rm ph}(0) = \epsilon_{\rm ph}(0), \qquad (148)$$

indeed for vanishing phonon masses the system is scale invariant in (1+1) dimensions: as already noticed phonons are mostly emitted in the direction orthogonal to the horizon. The effect of the phonon mass is to reduce both the energy density and the pressure with respect to the values obtained with massless phonons, in such a way that

$$P_{\rm ph}(m_{\rm ph}) < \epsilon_{\rm ph}(m_{\rm ph}), \qquad (149)$$

where the difference between these two quantities at the leading order in  $E_c/T$  is

$$\epsilon_{\rm ph}(m_{\rm ph}) - P_{\rm ph}(m_{\rm ph}) \simeq \epsilon_{\rm ph}(0) \frac{3E_c}{\pi^2 T} \,.$$
 (150)

Thus, the same difference is proportional to the square root of the Rabi coupling.

### VII. VISCOSITY AND THE CONFORMAL LIMIT CLOSE TO THE HORIZON

As a consequence of the spontaneous phonon emission close to the acoustic horizon, the transonic superfluid loses energy because a non-equilibrium (relaxation) dynamics sets in [19, 79, 80]: the superfluid kinetic energy is transformed in heat. At the effective level, this behavior can be characterized by effective bulk,  $\zeta_{\text{eff}}$ , and shear,  $\eta_{\text{eff}}$ , viscous coefficients, associated to the spontaneous phonon emission. As shown in [19], the shear-toentropy density ratio,  $\eta_{\text{eff}}/s_{\text{ph}}$ , of the phonon gas emitted at the acoustic horizon saturates the so-called KSS bound. First, we briefly recap the results of [19], obtained for a single superfluid and then move to the evaluation of the ratios  $\zeta_{\text{eff}}/s_{\text{ph}}$  and  $\eta_{\text{eff}}/s_{\text{ph}}$  in binary superfluids.

Let us first discuss the bulk viscosity for a fluid flowing with no shear. Close to the acoustic horizon, the viscous stress tensor is

$$\sigma'_{ik} = \zeta_{\text{eff}} \,\delta_{ix} \delta_{kx} \boldsymbol{\nabla} \cdot \boldsymbol{v} \simeq 2\pi \, T \, \zeta_{\text{eff}} \,\delta_{ix} \delta_{kx} \,, \qquad (151)$$

where we have taken into account that phonons are mostly emitted along the direction orthogonal to the horizon and on the r.h.s. we have used Eqs. (127) and (130). Equating  $\sigma'_{xx} = P_{\rm ph}$ , where  $P_{\rm ph}$  is given in Eq. (146), we have that

$$2\pi T \zeta_{\text{eff}} = P_{\text{ph}} \,, \tag{152}$$

which relates the bulk viscosity to the pressure of the phonon gas. This equation states that close to the acoustic horizon the only considered dissipative mechanism is the spontaneous phonon emission; any other viscous process is assumed negligible. However, it is well known that scattering processes are dissipative and contribute to the fluid viscosity [83]. Here we consider the phonon gas in a region close to the acoustic horizon of extension smaller that the typical phonon mean free path so that phonon scattering processes can be neglected. Then, we use the thermodynamic relation

$$\epsilon_{\rm ph} = -P_{\rm ph} + Ts_{\rm ph} \,, \tag{153}$$

combined with the relation in Eq. (148), valid for a gas of massless particles in 1+1 dimensions, to rewrite Eq. (152) as

$$2\pi T \zeta_{\text{eff}} = \frac{T s_{\text{ph}}}{2} \,. \tag{154}$$

and so it readily follows that

$$\frac{\zeta_{\text{eff}}}{s_{\text{ph}}} = \frac{1}{4\pi} \,. \tag{155}$$

In a similar way, we can obtain the ratio between the shear viscosity coefficient and the entropy density. In this case, we have to assume that the flow has shear, thus that the velocity, which is along the x-direction, depends on a transverse coordinate, for instance  $v_x$  depends on the y-coordinate. In this conditions, there is a "transverse" pressure [19], such that

$$\sigma'_{yx} = K \eta = -\frac{K}{C} P_{\rm ph} \,, \qquad (156)$$

where  $K = \partial_y v_x$  and C is given in Eq. (130). Using the same reasoning used above, we readily obtain that

$$\frac{\eta_{\text{eff}}}{s_{\text{ph}}} = \frac{1}{4\pi} \,, \tag{157}$$

corresponding to the KSS bound.

These results can be easily extended to binary superfluids with vanishing Rabi coupling. In the particular conditions considered in Sec. V A, the low-energy spectrum consists of two decoupled massless phonons that we called a and b. We have assumed that both fluids move with the same velocity v, however each phonon mode has its specific speed of sound, see Eqs. (113), meaning that in general there will be two distinct sonic horizons, corresponding to the conditions  $v = c_{sa}$ , and  $v = c_{sb}$  for the modes a and b, respectively. Then, as in the singlefluid case, one can determine the effective bulk and shear viscosity to entropy ratios for each fluid.

For nonvanishing Rabi coupling, one of the two phonons becomes massive and this is expected to change the viscosity to entropy density ratios. The reason is that for massive phonons the relation in Eq. (148) does not hold. The basic reason is that the phonon mass breaks the scale invariance. More in details, the needed ingredients to saturate the KSS bound in the presence of an acoustic horizon are the followings:

- i) close to the horizon the system is effectively (1+1)dimensional;
- ii) the relation  $\nabla \cdot \boldsymbol{v} = 2\pi T$ ;
- iii) the relation  $P_{\rm ph} = \epsilon_{\rm ph}$ .

While the conditions i) and ii) hold as well for a massive phonon gas, the condition iii) is violated by the phonon mass. As we have shown in the previous section, a mass term implies Eq. (149), hence

$$Ts_{\rm ph} = P_{\rm ph} + \epsilon_{\rm ph} \simeq 2P_{\rm ph} \left(1 + \frac{3E_c}{2\pi^2 T}\right),$$
 (158)

and then

$$\frac{\eta_{\text{eff}}}{s_{\text{ph}}} = \frac{\zeta_{\text{eff}}}{s_{\text{ph}}} \simeq \frac{1}{4\pi} \left( 1 - \frac{3E_c}{2\pi^2 T} \right) \,, \tag{159}$$

and using the estimate in Eq. (145), we have a reduction of about 10% with respect to the KSS bound. This is the main result of the present work.

Let us remark an important aspect of the obtained result: it holds because the phonon mass is not an interaction term, therefore it does not produce phonon scatterings. Its only effect is to break the scale invariance and thus the phonon pressure must be less than the phonon energy density. This means that the bulk and shear viscosities can still be written as in Eqs. (154) and (156), respectively, but now the phonon pressure is given by Eq. (147). Since the phonon mass reduces the phonon pressure, it follows that with increasing phonon mass the effective viscosities are reduced, as well.

Concerning the effect of the interactions on the KSS bound, we observe that they immediately violate the conditions i) and iii) (encoding scale invariance), therefore they are expected to induce deviations from the same bound. This aspect deserves future investigation.

### VIII. CONCLUSIONS

Binary ultracold gases are versatile playgrounds for producing nontrivial phases of matter [8]. In the present paper, we have determined the properties of the superfluid ground-state for a two-boson mixture coupled by intra- and inter-species interactions, as well as by a Rabi exchange term. Later on, the corresponding effective lowenergy excitation theory has been determined. Assuming a perturbative Rabi coupling, we have shown that when the number densities of the two fluids are similar, it is possible to write the total pressure of the system as the sum of two partial pressures. Moreover, we have obtained that in the same conditions, the low-energy excitations of the binary superfluid are two decoupled modes, corresponding to the relative and the total phase oscillations. The former is massive, with a squared mass proportional to the Rabi coupling, while the latter remains massless.

Separating the total system as the sum of two noninteracting subsystems is useful when discussing the spontaneous phonon emission close to an acoustic horizon. Indeed, we can treat the emission of the two phonon modes as independent processes. The spontaneous phonon emission is an irreversible process, that at the hydrodynamic level amounts to effective viscosities. In binary superfluids, the spontaneous emission of massless phonons is analogous to the phonon process in a single superfluid, thus the viscosity-to-entropy ratios saturates the KSS bound [16, 19]: it is equal to  $1/4\pi$ . However, the spontaneous emission of massive phonons at the acoustic horizon violates the same bound: in this case,  $\eta_{\rm eff}/s_{\rm ph} = \zeta_{\rm eff}/s_{\rm ph} < 1/4\pi$ . The violation is estimated of the order of 10% using the specific setup in [82]. In general, the magnitude of the same deviation scales with the square root of the Rabi frequency, and it is independent on the mass of the used atoms. Therefore, it seems that to enhance the deviation from the KSS bound one

needs to design systems with the highest possible Rabi frequency and with sufficiently tunable speeds of sound and/or fluid velocity as necessary ingredients to have an acoustic horizon. Our setup provides a versatile tool to explore the degree of KSS-bound violation and even advance the exciting quest for a universal  $\eta/s$  bound, under controllable external conditions that can be engineered in current quantum technology platforms.

In actual experiments, the measurements of the local values of the entropy density and viscosity are required and seem feasible. The former may be achieved by employing local-thermometry techniques, as in [84], which allow to measure extremely small values of the entropy by means of radio-frequency spectroscopy. Local values of the shear viscosity may be obtained by appropriate inversion of the cloud-averaged viscosity [85].

Acknowledgments – The authors are pleased to thank Stefano Carretta, Leonardo Fallani, Stefano

Liberati, Luca Salasnich, Paolo Santini, and Andrea Trombettoni for support and fruitful discussions.

L. L. acknowledges financial support by a project funded under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.3 - Call for tender No. 341 of 15/03/2022 of Italian Ministry of University and Research funded by the European Union – NextGenerationEU, award number PE0000023, Concession Decree No. 1564 of 11/10/2022 adopted by the Italian Ministry of University and Research, CUP D93C22000940001, Project title "National Quantum Science and Technology Institute" (NQSTI), spoke 2.

This study has been also carried out within the National Centre on HPC, Big Data and Quantum Computing - SPOKE 10 (Quantum Computing) and received funding from the European Union Next-GenerationEU - National Recovery and Resilience Plan (NRRP) – MIS-SION 4 COMPONENT 2, INVESTMENT N. 1.4 – CUP N. I53C22000690001.

#### Appendix A: General derivations of the acoustic metric and of the low-energy Lagrangian

We determine the effective low-energy Lagrangian for a superfluid, using two different methods. For simplicity, we consider a single-species boson gas, but the two approaches can be straightforwardly extended to binary superfluids.

## 1. Expansion around mean field of a generic Lagrangian

In this first method we employ the same procedure used in Sec. II, but we start from an unspecified Lagrangian for a complex scalar field in the Madelung representation,

$$\mathcal{L} \equiv \mathcal{L}(\rho, \partial_{\mu}\rho, \partial_{\mu}\theta), \qquad (A1)$$

where  $\rho$  is the radial field and  $\theta$  is the cyclic coordinate. We assume vanishing temperature and that the Lagrangian is invariant under a global U(1) transformation. In the ground-state, such group is assumed to be spontaneously broken.

We include fluctuations on top of the mean field by the as in Eq. (17), hence  $\rho$  and  $\theta$  now indicate the solutions of the classical equation of motion, while  $\tilde{\rho}$  and  $\tilde{\theta}$  are the quantum fluctuations. Expanding in powers of the fluctuations, and neglecting surface terms, we have that

$$\mathcal{L} = \mathcal{L}_0(\rho, \partial_\mu \rho, \partial_\mu \theta) + \tilde{\rho} \left( \frac{\delta \mathcal{L}}{\delta \rho} - \partial_\mu \frac{\delta \mathcal{L}}{\delta \partial_\mu \rho} \right) \Big|_{\rho, \theta} + \delta \partial_\mu \tilde{\theta} \left. \frac{\delta \mathcal{L}}{\delta \partial_\mu \theta} \right|_{\rho, \theta} + \mathcal{L}_2 + \dots$$
(A2)

where the dots indicate cubic and higher terms in the fluctuations. The first term on the right-hand side, when evaluated in the stationary point, is the background pressure: it does not include the effect of fluctuations; in the model with quartic interactions it is given in Eq. (15). Since  $\rho$  and  $\theta$  are solutions of the classical equations of motion, it follows that the terms linear in the perturbations vanish. Thus the first nontrivial term is the quadratic Lagrangian

$$\mathcal{L}_{2} = \frac{\tilde{\rho}^{2}}{2} \left. \frac{\delta^{2} \mathcal{L}}{\delta \rho^{2}} \right|_{\rho,\theta} + \tilde{\rho} \partial_{\mu} \tilde{\rho} \left. \frac{\delta^{2} \mathcal{L}}{\delta \rho \delta \partial_{\mu} \rho} \right|_{\rho,\theta} + \frac{\partial_{\mu} \tilde{\rho} \partial_{\nu} \tilde{\rho}}{2} \left. \frac{\delta^{2} \mathcal{L}}{\delta \partial_{\mu} \rho \delta \partial_{\nu} \rho} \right|_{\rho,\theta} + \tilde{\rho} \partial_{\mu} \tilde{\theta} \left. \frac{\delta^{2} \mathcal{L}}{\delta \rho \delta \partial_{\mu} \theta} \right|_{\rho,\theta} + \left. \frac{\partial_{\nu} \tilde{\theta} \partial_{\mu} \tilde{\theta}}{2} \left. \frac{\delta^{2} \mathcal{L}}{\delta \partial_{\nu} \theta \delta \partial_{\mu} \theta} \right|_{\rho,\theta} \right|_{\rho,\theta} \right.$$
(A3)

where we have taken into account that  $\theta$  has only derivative couplings.

We now discuss in some details the various terms in the right-hand side of Eq. (A3). The first term is proportional to minus the curvature of the potential at the minimum:

$$\frac{\tilde{\rho}^2}{2} \left. \frac{\delta^2 \mathcal{L}}{\delta \rho^2} \right|_{\rho,\theta} = -\frac{\tilde{m}^2}{2} \tilde{\rho}^2 \,, \tag{A4}$$

where  $\tilde{m}$  is the effective mass of the radial field. The second term is a total derivative and can be neglected. The third term is

$$\frac{\partial_{\mu}\tilde{\rho}\partial_{\nu}\tilde{\rho}}{2} \left. \frac{\delta^{2}\mathcal{L}}{\delta\partial_{\mu}\rho\delta\partial_{\nu}\rho} \right|_{\rho,\theta} = \frac{K^{\mu\nu}}{2} \partial_{\mu}\tilde{\rho}\partial_{\nu}\tilde{\rho}, \qquad (A5)$$

where  $K^{\mu\nu}$  is a matrix depending on the background; in flat spacetime  $K^{\mu\nu} = \eta^{\mu\nu}$ . The fourth term can be written as

$$\tilde{\rho}\partial_{\mu}\tilde{\theta} \left. \frac{\delta^{2}\mathcal{L}}{\delta\rho\delta\partial_{\mu}\theta} \right|_{\rho,\theta} = V^{\mu}\tilde{\rho}\,\partial_{\mu}\tilde{\theta}\,,\tag{A6}$$

which is a mixing term. Since  $V^{\mu}$  is by construction a four-vector, it explicitly breaks the Lorentz symmetry. By dimensional analysis it follows that it has dimension 2. The fifth term can be written as

$$\partial_{\nu}\tilde{\rho}\partial_{\mu}\tilde{\theta} \left. \frac{\delta^{2}\mathcal{L}}{\delta\partial_{\nu}\rho\delta\partial_{\mu}\theta} \right|_{\rho,\theta} = A \eta^{\mu\nu}\partial_{\nu}\tilde{\rho}\partial_{\mu}\tilde{\theta} , \qquad (A7)$$

where A is a constant of dimension 1. It vanishes if the interactions in  $\mathcal{L}$  stem from a potential term, thus the radial field has no derivative couplings. Finally, the last term can be written as

$$\frac{\partial_{\nu}\tilde{\theta}\partial_{\mu}\tilde{\theta}}{2} \left. \frac{\delta^{2}\mathcal{L}}{\delta\partial_{\nu}\theta\delta\partial_{\mu}\theta} \right|_{\rho,\theta} = \frac{L^{\mu\nu}}{2} \partial_{\nu}\tilde{\theta}\partial_{\mu}\tilde{\theta} , \qquad (A8)$$

where  $L^{\mu\nu}$  depends on the background; in flat spacetime  $L^{\mu\nu} = B^2 \eta^{\mu\nu}$ , where B has dimension 1. In the end, the quadratic Lagrangian reads

$$\mathcal{L}_2 = \frac{1}{2} \partial_\nu \tilde{\rho} \partial^\nu \tilde{\rho} - \frac{\tilde{m}^2}{2} \tilde{\rho}^2 + V^\mu \tilde{\rho} \partial_\mu \tilde{\theta} + \frac{B^2}{2} \partial_\nu \tilde{\theta} \partial^\nu \tilde{\theta} + A \partial^\mu \tilde{\rho} \partial_\mu \tilde{\theta} , \qquad (A9)$$

where  $\hat{\theta}$  is clearly the NGB associated to the spontaneous symmetry breaking of the U(1) global symmetry. Since  $\hat{\rho}$  field is massive, we can integrate it out. Its equation of motion reads,

$$(\Box + \tilde{m}^2)\tilde{\rho} = V^{\mu}\partial_{\mu}\tilde{\theta} - A\Box\tilde{\theta}, \qquad (A10)$$

which we can formally invert to obtain

$$\tilde{\rho} = \frac{1}{\Box + \tilde{m}^2} \left( V^{\mu} \partial_{\mu} \tilde{\theta} - A \Box \tilde{\theta} \right) \,, \tag{A11}$$

showing a generic feature of effective-field theories: the nonlocality. However, higher derivative terms are suppressed as powers of  $p^2/\tilde{m}^2$ , where p is the momentum of the NGB. Note that the relevant scale for the expansion is  $\tilde{m}$  and not m. Since  $\tilde{m}$  vanishes close to the second order phase transition point, it follows that the energy and momentum range of validity of the low-energy effective Lagrangian shrinks close to the transition to the normal phase.

Upon substituting Eq. (A11) in Eq. (A9), we obtain

$$\mathcal{L}_{2} = \frac{1}{2} \left( V^{\mu} \partial_{\mu} \tilde{\theta} - A \Box \tilde{\theta} \right) \frac{1}{\Box + \tilde{m}^{2}} \left( V^{\mu} \partial_{\mu} \tilde{\theta} - A \Box \tilde{\theta} \right) + \frac{B^{2} \eta^{\mu\nu}}{2} \partial_{\nu} \tilde{\theta} \partial_{\mu} \tilde{\theta} , \qquad (A12)$$

which is the most general form of the low-energy effective Lagrangian. At the leading order in momenta, we replace

$$\frac{1}{\Box + \tilde{m}^2} \to \frac{1}{\tilde{m}^2}, \quad V^{\mu} \partial_{\mu} \tilde{\theta} - A \Box \tilde{\theta} \to V^{\mu} \partial_{\mu} \tilde{\theta} , \qquad (A13)$$

so that we can rewrite the effective low-energy Lagrangian as

$$\mathcal{L}_2 \simeq \frac{1}{2} \left( B^2 \eta^{\mu\nu} + \frac{V^{\mu}V^{\nu}}{\tilde{m}^2} \right) \partial_{\nu} \tilde{\theta} \partial_{\mu} \tilde{\theta} \equiv \frac{\sqrt{-g} g^{\mu\nu}}{2} \partial_{\nu} \tilde{\theta} \partial_{\mu} \tilde{\theta} \,, \tag{A14}$$

where  $g_{\mu\nu}$  is the acoustic metric. The obtained low-energy Lagrangian is valid for momenta  $p \ll \tilde{m}$ ; alternatively, it can be obtained neglecting the kinetic term  $\frac{1}{2}\partial_{\nu}\tilde{\rho}\partial^{\nu}\tilde{\rho}$  and taking as well A = 0 in Eq. (A9).

#### 2. Derivative expansion of the pressure

We now present a different method to derive the low-energy theory and the acoustic metric, based on the observation that the low-energy Lagrangian at the leading order in momenta of any system with a U(1) broken symmetry can be determined from the knowledge of the pressure [18, 68]. This method has the advantage of bypassing the integration of the radial field: we start directly with the effective Lagrangian of the  $\tilde{\theta}$  field. However, it has the disadvantage to include only the leading order terms in the phonon momentum. Taking the global symmetry as a local one, the low-energy Lagrangian including leading order derivatives is given by

$$\mathcal{L} = P\left[\sqrt{D_{\mu}\tilde{\theta}D^{\mu}\tilde{\theta}}\right],\tag{A15}$$

where  $D_{\mu}\tilde{\theta} = \partial_{\mu}\tilde{\theta} - A_{\mu}$  and P is a functional having the same algebraic expression of the pressure. As in Eq. (2), we take the external field given by

$$A_{\nu} = \frac{\mu}{\gamma} v_{\nu} , \qquad (A16)$$

thus we assume a background flow with four-velocity  $v_{\nu}$ . Note that the present definition of the covariant derivative slightly differs from the expression in Eq. (2), because here it acts on the phase field. For convenience we replace  $\mu/\gamma \rightarrow \mu$ . Upon expanding the right hand side of Eq. (A15) around  $\mu$ , we obtain that

$$\mathcal{L} = P(\mu) - \frac{\partial P}{\partial \mu} v_{\mu} \partial^{\mu} \tilde{\theta} + \frac{1}{2} \left[ \eta^{\mu\nu} \frac{1}{\mu} \frac{\partial P}{\partial \mu} + v^{\mu} v^{\nu} \left( \frac{\partial^2 P}{\partial \mu^2} - \frac{1}{\mu} \frac{\partial P}{\partial \mu} \right) \right] \partial_{\nu} \tilde{\theta} \partial_{\mu} \tilde{\theta} , \qquad (A17)$$

thus the quadratic term is

$$\mathcal{L}_2 = \frac{n}{2\mu} \left( \eta^{\mu\nu} + \left( \frac{1}{c_s^2} - 1 \right) v^{\mu} v^{\nu} \right) \partial_{\nu} \tilde{\theta} \partial_{\mu} \tilde{\theta} \,, \tag{A18}$$

where we used the thermodynamic relations

$$n = \frac{\partial P}{\partial \mu}$$
 and  $c_s^2 = \frac{n}{\mu \frac{\partial^2 P}{\partial \mu^2}}$ , (A19)

the second expression stemming from  $c_s^2 = \frac{\partial P}{\partial \rho}$ . The low-energy Lagrangian can be further expanded including higher orders of  $\partial_\mu \tilde{\theta}$ ; this generates interaction terms at the leading order in the momenta, see the discussion in [18].

- [1] C. Barcelo, S. Liberati, and M. Visser, Liv. Rev. Rel. 8, 12 (2005), arXiv:gr-qc/0505065.
- [2] E. Altman et al., PRX Quantum 2, 017003 (2021), arXiv:1912.06938 [quant-ph].
- [3] L. Warszawski and A. Melatos, Mon. Not. Roy. Astron. Soc. 415, 1611 (2011), arXiv:1103.6090 [astro-ph.SR].
- [4] L. Warszawski, A. Melatos, and N. Berloff, Phys. Rev. B 85, 104503 (2012), arXiv:1203.5133 [cond-mat.other].
- [5] D. Faccio, F. Belgiorno, S. Cacciatori, V. Gorini, S. Liberati, and U. Moschella, eds., *Analog. Grav. Phenomen.*, Vol. 870 (2013).
- [6] E. Poli, T. Bland, S. J. M. White, M. J. Mark, F. Ferlaino, S. Trabucco, and M. Mannarelli, Phys. Rev. Lett. 131, 223401 (2023), arXiv:2306.09698 [cond-mat.quant-gas].
- [7] T. Bland, F. Ferlaino, M. Mannarelli, E. Poli, and S. Trabucco, Few Body Syst. 65, 81 (2024), arXiv:2407.03212 [cond-mat.quant-gas].
- [8] J. Yago Malo, L. Lepori, L. Gentini, and M. L. Chiofalo, Technologies 12 (2024), 10.3390/technologies12050064.
- [9] L. Cipriani, M. Mannarelli, F. Nesti, and S. Trabucco, Phys. Rev. D 110, L021301 (2024), arXiv:2403.03833 [astro-ph.CO].
- [10] C. Coviello, M. L. Chiofalo, D. Grasso, S. Liberati, M. Mannarelli, and S. Trabucco, (2024), arXiv:2410.00264 [gr-qc].
- [11] P. Magierski, A. Barresi, A. Makowski, D. Pcak, and G. Wlazłowski, Eur. Phys. J. A 60, 186 (2024), arXiv:2406.14158 [cond-mat.quant-gas].
- [12] Z.-H. Tu and A. Li, (2024), arXiv:2412.09219 [nucl-th].
- [13] R. Schützhold, (2025), arXiv:2501.03785 [quant-ph].
- [14] S. Liberati, M. Visser, and S. Weinfurtner, Phys. Rev. Lett. 96, 151301 (2006), arXiv:gr-qc/0512139.
- [15] A. Adams, L. D. Carr, T. Schäfer, P. Steinberg, and J. E. Thomas, New J. Phys. 14, 115009 (2012).

- [16] P. Kovtun, D. T. Son, and A. O. Starinets, Phys. Rev. Lett. 94, 111601 (2005), arXiv:hep-th/0405231.
- [17] S. Cremonini, Mod. Phys. Lett B 25, 1867 (2011), https://doi.org/10.1142/S0217984911027315.
- [18] M. Mannarelli and C. Manuel, Phys. Rev. D 81, 043002 (2010).
- [19] M. L. Chiofalo, D. Grasso, M. Mannarelli, and S. Trabucco, New J. Phys. 26, 053021 (2024), arXiv:2202.13790 [gr-qc].
- [20] A. Cherman, T. D. Cohen, and P. M. Hohler, JHEP 02, 026 (2008), arXiv:0708.4201 [hep-th].
- [21] X.-H. Ge, S.-K. Jian, Y.-L. Wang, Z.-Y. Xian, and H. Yao, Phys. Rev. Res. 2, 023366 (2020).
- [22] M. Sadeghi and S. Parvizi, Class. Quant. Gravity **33**, 035005 (2016).
- [23] E. A. Cornell and C. E. Wieman, Rev. Mod. Phys. 74, 875 (2002).
- [24] A. Recati and S. Stringari, Ann. Rev. Cond. Matt. Phys. 13, 407 (2022).
- [25] C. Baroni, G. Lamporesi, and M. Zaccanti, Nat. Rev. Phys. (2024), 10.1038/s42254-024-00773-6.
- [26] D. T. Son and M. A. Stephanov, Phys. Rev. A 65, 063621 (2002), arXiv:cond-mat/0103451.
- [27] P. G. Kevrekidis, H. E. Nistazakis, D. J. Frantzeskakis, B. A. Malomed, and R. Carretero-Gonzalez, Eur. Phys. J. D 28, 181–185 (2004).
- [28] B. Bakkali-Hassani, C. Maury, Y.-Q. Zou, E. Le Cerf, R. Saint-Jalm, P. Castilho, S. Nascimbene, J. Dalibard, and J. Beugnon, Phys. Rev. Lett. 127 (2021), 10.1103/physrevlett.127.023603.
- [29] A. Romero-Ros, G. C. Katsimiga, S. I. Mistakidis, S. Mossman, G. Biondini, P. Schmelcher, P. Engels, and P. G. Kevrekidis, Phys. Rev. Lett. 132, 033402 (2024), arXiv:2304.05951 [nlin.PS].
- [30] C. Hamner, J. J. Chang, P. Engels, and M. A. Hoefer, Phys. Rev. Lett. 106 (2011), 10.1103/physrevlett.106.065302.
- [31] A. Farolfi, D. Trypogeorgos, C. Mordini, G. Lamporesi, and G. Ferrari, Phys. Rev. Lett. **125** (2020), 10.1103/physrevlett.125.030401.
- [32] A. Richaud, G. Lamporesi, M. Capone, and A. Recati, Phys. Rev. A 107 (2023), 10.1103/physreva.107.053317.
- [33] D. S. Petrov, Phys. Rev. Lett. 115, 155302 (2015).
- [34] G. Semeghini, G. Ferioli, L. Masi, C. Mazzinghi, L. Wolswijk, F. Minardi, M. Modugno, G. Modugno, M. Inguscio, and M. Fattori, Phys. Rev. Lett. 120 (2018), 10.1103/physrevlett.120.235301.
- [35] C. J. Myatt, E. A. Burt, R. W. Ghrist, E. A. Cornell, and C. E. Wieman, Phys. Rev. Lett. 78, 586 (1997).
- [36] J. Tuoriniemi, J. Martikainen, E. Pentti, A. Sebedash, S. Boldarev, and G. Pickett, J. Low Temp. Phys. 129, 531 (2002).
- [37] J. Rysti, J. Tuoriniemi, and A. Salmela, Phys. Rev. B 85, 134529 (2012).
- [38] M. Lewenstein, A. Sanpera, and V. Ahufinger, Ultracold Atoms in Optical Lattices: Simulating quantum many-body systems (OUP Oxford, 2012).
- [39] M. Inguscio and L. Fallani, Atomic Physics: Precise Measurements and Ultracold Matter (OUP Oxford, 2013).
- [40] I. Ferrier-Barbut, M. Delehaye, S. Laurent, A. T. Grier, M. Pierce, B. S. Rem, F. Chevy, and C. Salomon, Science 345, 1035 (2014), arXiv:1404.2548 [cond-mat.quant-gas].
- [41] L. Lepori, A. Trombettoni, and W. Vinci, Europhys. Lett. **109**, 50002 (2015).
- [42] L. Lepori and M. Mannarelli, Phys. Rev. D 99, 096011 (2019), arXiv:1901.07488 [hep-ph].
- [43] J. Nespolo, G. E. Astrakharchik, and A. Recati, New J. Phys. 19, 125005 (2017).
- [44] U. R. Fischer and R. Schutzhold, Phys. Rev. A 70, 063615 (2004), arXiv:cond-mat/0406470.
- [45] M. Visser and S. Weinfurtner, Phys. Rev. D 72, 044020 (2005), arXiv:gr-qc/0506029.
- [46] S. Liberati, M. Visser, and S. Weinfurtner, Class. Quant. Grav. 23, 3129–3154 (2006).
- [47] A. Berti, L. Fernandes, S. G. Butera, A. Recati, M. Wouters, and I. Carusotto, (2024), arXiv:2408.17292 [cond-mat.quant-gas].
- [48] A. Zenesini, A. Berti, R. Cominotti, C. Rogora, I. G. Moss, T. P. Billam, I. Carusotto, G. Lamporesi, A. Recati, and G. Ferrari, Nat. Phys. 20, 558 (2024), arXiv:2305.05225 [hep-ph].
- [49] S. L. Shapiro and S. A. Teukolsky, Black holes, white dwarfs, and neutron stars: The physics of compact objects (1983).
- [50] M. G. Alford, K. Rajagopal, and F. Wilczek, Nucl. Phys. B 537, 443 (1999), arXiv:hep-ph/9804403 [hep-ph].
- [51] P. F. Bedaque and T. Schafer, Nucl. Phys. A 697, 802 (2002), arXiv:hep-ph/0105150 [hep-ph].
- [52] D. Kaplan and S. Reddy, Phys. Rev. D 65, 054042 (2002), arXiv:hep-ph/0107265 [hep-ph].
- [53] M. G. Alford, A. Schmitt, K. Rajagopal, and T. Schafer, Rev. Mod. Phys. 80, 1455 (2008), arXiv:0709.4635 [hep-ph].
- [54] R. Anglani, R. Casalbuoni, M. Ciminale, N. Ippolito, R. Gatto, et al., Rev. Mod. Phys. 86, 509 (2014), arXiv:1302.4264 [hep-ph].
- [55] J. Annett, Superconductivity, Superfluids and Condensates, Oxford Master Series in Physics (OUP Oxford, 2004).
- [56] C. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases* (Cambridge University Press, 2008).
- [57] L. Lepori, A. Maraga, A. Celi, L. Dell'Anna, and A. Trombettoni, Cond. Matt. 3 (2018), 10.3390/condmat3020014.
- [58] H. Pu and N. P. Bigelow, Phys. Rev. Lett. 80, 1130 (1998).
- [59] H. Pu and N. P. Bigelow, Phys. Rev. Lett. 80, 1134 (1998).
- [60] D. S. Hall, M. R. Matthews, C. E. Wieman, and E. A. Cornell, Phys. Rev. Lett. 81, 1543 (1998), arXiv:cond-mat/9805327.
- [61] W. Unruh, Phys.Rev.Lett. 46, 1351 (1981).
- [62] N. Bilic, Class. Quant. Grav. 16, 3953 (1999), arXiv:gr-qc/9908002.
- [63] M. Visser and C. Molina-Paris, New J. Phys. 12, 095014 (2010), arXiv:1001.1310 [gr-qc].
- [64] R. Brout, S. Massar, R. Parentani, and P. Spindel, Phys. Rept. 260, 329 (1995), arXiv:0710.4345 [gr-qc].
- [65] G. E. Volovik, Found. Phys. **33**, 349 (2003), arXiv:gr-qc/0301043.
- [66] I. Carusotto, S. Fagnocchi, A. Recati, R. Balbinot, and A. Fabbri, New J. Phys. 10, 103001 (2008), 0803.0507.
- [67] J. Steinhauer, Nat. Phys. **12**, 959–965 (2016).
- [68] D. T. Son, (2002), arXiv:hep-ph/0204199 [hep-ph].
- [69] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, Rev. Mod. Phys. 71, 463 (1999), arXiv:cond-mat/9806038.

- [70] L. Pitaevskii and S. Stringari, *Bose-Einstein Condensation*, International Series of Monographs on Physics (Clarendon Press, 2003).
- [71] This is what occurs in the framework of the Bogoliubov theory for superfluid bosons, see for instance [86].
- [72] E. Timmermans, Phys. Rev. Lett. 81, 5718 (1998).
- [73] L.-F. Li and H. Pagels, Phys. Rev. Lett. 26, 1204 (1971).
- [74] D. T. Son and M. A. Stephanov, Phys. Rev. A 65, 063621 (2002).
- [75] C. Qu, M. Tylutki, S. Stringari, and L. P. Pitaevskii, Phys. Rev. A 95, 033614 (2017).
- [76] K. Takayama and M. Oka, Nucl. Phys. A 551, 637 (1993).
- [77] S. Weinfurtner, S. Liberati, and M. Visser, Lect. Notes Phys. 718, 115 (2007), arXiv:gr-qc/0605121.
- [78] M. K. Parikh and F. Wilczek, Phys. Rev. Lett. 85, 5042 (2000), arXiv:hep-th/9907001.
- [79] M. Mannarelli, D. Grasso, S. Trabucco, and M. L. Chiofalo, Phys. Rev. D 103, 076001 (2021), arXiv:2011.00019 [gr-qc].
- [80] M. Mannarelli, D. Grasso, S. Trabucco, and M. L. Chiofalo, (2021), arXiv:2109.11831 [gr-qc].
- [81] M. Le Bellac, *Thermal Field Theory*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, 2000).
- [82] J. R. Muñoz de Nova, K. Golubkov, V. I. Kolobov, and J. Steinhauer, Nature 569, 688 (2019), arXiv:1809.00913 [gr-qc].
- [83] I. M. Khalatnikov, Introduction to the theory of superfluidity (W. A. Benjamin, New York, 1965).
- [84] T. Hartke, B. Oreg, N. Jia, and M. Zwierlein, Phys. Rev. Lett. 125, 113601 (2020).
- [85] J. A. Joseph, E. Elliott, and J. E. Thomas, Phys. Rev. Lett. 115, 020401 (2015).
- [86] G. E. Volovik, The Universe in a Helium Droplet (Oxford University Press, 2009).