Vector-Valued Gaussian Processes and their Kernels on a Class of Metric Graphs

Tobia Filosi^A, Emilio Porcu^{*B}, Xavier Emery^C, Claudio Agostinelli^A, and Alfredo Alegría^D

^ADepartment of Mathematics, University of Trento ^BDepartment of Mathematics, Khalifa University & ADIA LAB

^CDepartment of Mining Engineering, Universidad de Chile ^DDepartment of Mathematics, Universidad Técnica Federico Santa María

January 20, 2025

Abstract

Despite the increasing importance of stochastic processes on linear networks and graphs, current literature on multivariate (vectorvalued) Gaussian random fields on metric graphs is elusive. This paper challenges several aspects related to the construction of proper matrixvalued kernels structures. We start by considering matrix-valued metrics that can be composed with scalar- or matrix-valued functions to implement valid kernels associated with vector-valued Gaussian fields. We then provide conditions for certain classes of matrix-valued functions to be composed with the univariate resistance metric and ensure positive semidefiniteness. Special attention is then devoted to Euclidean trees, where a substantial effort is required given the absence of literature related to multivariate kernels depending on the ℓ_1 metric. Hence, we provide a foundational contribution to certain classes of matrix-valued positive semidefinite functions depending on the ℓ_1 metric. This fact is then used to characterise kernels on Euclidean trees with a finite number of leaves. Amongst those, we provide classes of matrix-valued covariance functions that are compactly supported.

1 Introduction

1.1 Context: Random Fields on Networks

Networks analysis has become ubiquitous in several branches of theoretical and applied sciences, including mathematics, statistics, machine learning, artificial intelligence, with applications connected to the data planet. Networks can be used, either, to represent connections between random variables, or to provide a topological structure where a given process (called a random field) is observed. Our paper deals with the second situation, under the condition that the observed process is a vector-valued Gaussian random field that is second-order isotropic in the sense that the matrix-valued covariance kernel depends on distance between the points in the network.

Random fields can be continuously [Anderes et al., 2020] or discretely indexed over a network; for the second case, the common nomenclature is that of point processes on networks [Moradi and Mateu, 2020, Baddeley et al., 2021, Rakshit et al., 2017]. For both cases, defining a random field over a network is a challenging task.

The electrical engineering and machine learning communities have been very active on the subject of networks, and the reader is referred to Ortega et al. [2018] and Chami et al. [2022] for a comprehensive review.

For this manuscript, the network defines a topological structure representing the domain of definition of a given random field. This fact is exploited by Anderes et al. [2020]: endowing the network with a metric provides a metric space that is then (quasi) embeddable over Hilbert spaces. This is the crux of the argument that allows to build isotropic covariance functions over networks, where isotropy is understood with respect to two alternative metrics (the geodesic and the resistance metrics). A relevant remark is that networks endowed with metrics can be suitably represented through metric graphs. This approach is adopted by Bolin and Lindgren [2011], Anderes et al. [2020] and Bolin et al. [2022]. We shall make use of this fact subsequently. Scalar random fields that evolve temporally over graphs have been challenged by Porcu et al. [2023], Filosi et al. [2023] and Tang and Zimmerman [2020]. The way a metric graph can be specified is certainly not unique. Anderes et al. [2020] work over graphs with Euclidean edges — called generalised networks in Porcu et al. [2023] — which extend linear networks to nonlinear edges. Further, the random field defined over such structures can have realisations over any point over the edges, and not only in the nodes. Roughly, these are graphs where each edge is associated with an abstract set in bijective correspondence with a segment of the real line. This provides each edge with a Cartesian coordinate system to measure distances between any two points on that edge.

The fact that linear networks are overly limited to provide a suitable topological structure to random fields is well understood in the machine learning literature [Alsheikh et al., 2014, Hamilton et al., 2017, Borovitskiy et al., 2022], as well as in spatial statistics [Cressie et al., 2006, Ver Hoef et al., 2006, Peterson et al., 2013, 2007, Montembeault et al., 2012, Xiao et al., 2017, Perry and Wolfe, 2013, Deng et al., 2014, Baddeley et al., 2017].

1.2 Challenge

Our paper is unique according to the state of the art. The literature on vector-valued random fields on (linear or generalised) networks is elusive. This is not surprising: kernels on metric graphs are a very recent subject, and so far the main efforts have been focused to the case of scalar-valued random fields.

A subset of the graphs used in this paper is represented by the so-called Euclidean trees with a given number of leaves, which are substantially linear networks. For such a case, since a linear network is embedded into the two-dimensional Euclidean space, any matrix-valued covariance functions depending on the ℓ_1 norm (the so-called Manhattan distance) can be used as a covariance function for a vector-valued field on a linear network. The reader is referred to Zastavnyi [2000] for relevant results in terms of isometric embeddings. Unfortunately, there are two problems related to this fact:

- 1. To the knowledge of the authors, there are no models for matrix-valued covariances depending on the ℓ_1 distance;
- 2. Such constructions are no longer valid for the case of generalised networks.

Another relevant aspect is related to what we term *cross metrics* in the paper. To illustrate the principle, consider two random fields — call them Z_1 and Z_2 . Since the network does not necessarily represent a geographic space, there is no reason to believe that the distance between $Z_1(u)$ and $Z_2(v)$, for two different points u, v on the network, should be the same as the distance between $Z_1(u)$ and $Z_1(v)$ for instance. We take this aspect into account and devote attention to matrix-valued metrics.

1.3 Our Constructions: Road Map

Our constructions are based on the following steps. We provide here a concise description. Careful notation and terminology are established in Section 2. The Road Map

- 1. The purpose of the work is to construct a *p*-variate Gaussian random field, denoted $\mathbf{Z} := \{\mathbf{Z}(u) = (Z_1(u), \ldots, Z_p(u))^\top, u \in \mathcal{G}\}$ throughout and defined over a metric graph, \mathcal{G} , with a given matrix-valued covariance function that is isotropic, that is it depends on the *distance* between the points of the graph. Hence, the topological space \mathcal{G} , endowed with a proper matrix-valued (semi) metric $\mathbf{D} : \mathcal{G} \times \mathcal{G} \to [0, +\infty)^{p \times p}$, becomes a (semi) metric space $(\mathcal{G}, \mathbf{D})$.
- 2. Such a (semi) metric space can be (quasi) embedded on a suitable Hilbert space, which in turns allows to resort classical machinery for covariance functions on metric spaces [Schoenberg, 1942].
- Hence, a substantial part of the job is to build the multivariate metric D.
- 4. We then take advantage of isometric embedding arguments as in Anderes et al. [2020] to claim that, for a certain class of continuous mappings $\psi : [0, +\infty) \to (0, +\infty)$, the element-wise composition

$$\mathbf{K}(u,v) := \psi\left(\mathbf{D}(u,v)\right), \qquad u, v \in \mathcal{G},\tag{1}$$

provides a valid matrix-valued covariance function.

5. We then generalise this construction by considering a matrix-valued class of continuous mappings $\Psi : [0, +\infty)^{p \times p} \to [0, +\infty)^{p \times p}$ having elements Ψ_{ij} , such that the element-wise composition

$$\mathbf{K}(u,v) := \mathbf{\Psi}\left(\mathbf{D}(u,v)\right) = \left[\Psi_{ij}(D_{ij}(u,v))\right]_{i,j=1}^{p}, \qquad u,v \in \mathcal{G}, \qquad (2)$$

is a valid matrix-valued covariance function.

In turn, the construction of the multivariate metric \mathbf{D} requires a collection of steps, which entails several technical challenges:

- (a) Following the construction in Anderes et al. [2020], the vector-valued random field Z is defined as the sum of two random fields, Z_V and Z_E , where the subscripts V and E stand, respectively, for the sets containing vertices and the edges associated with the metric graph, \mathcal{G} . Both random fields are continuously indexed over the graph;
- (b) The random field Z_V at any point of an edge, $e \in E$, is constructed through interpolation of the random field defined at the extremes of each edge (which are obviously vertices in V);
- (c) The multivariate metric is then checked for desiderata. Specifically, that the marginal metrics (those in the diagonal of **D**) respect the ingenious univariate condition provided by Anderes et al. [2020]. Further, the cross-elements in **D** are required to be homogeneous, in the sense that $D_{ij} = D_{i'j'}$ for $i \neq j$ and $i' \neq j'$, for D_{ij} a generic element of **D**.

For a subclass termed *Euclidean trees*, we provide multivariate kernels with compact support depending on the so-called Manhattan distance.

The structure of the paper is the following. Section 2 provides the necessary background and notation. Results are provided in Sections 3 and 4. Section 5 concludes the paper. As proofs are rather technical and lengthy, they are deferred to Appendix A. Further, Appendix B contains a worked example that show some relevant property of the proposed construction.

Finally, to guide the reader into the structure of the results of this manuscript, we provide a road-map in Figure 1.

2 Mathematical Background

2.1 Notation handout

Throughout, p is an integer greater than 1 (the case p = 1 has been treated by Anderes et al. [2020]). Bold lowercase letters denote (deterministic) vectors (*e.g.* $\boldsymbol{\delta}$). Bold uppercase letters denote (deterministic) matrices (*e.g.* \mathbf{A}). Bold uppercase italic letters denote random vectors (*e.g.* \mathbf{Z}). $\mathbf{A}^{\top}, \boldsymbol{\delta}^{\top}$ and \mathbf{Z}^{\top} are the transposes of $\mathbf{A}, \boldsymbol{\delta}$ and \mathbf{Z} , while \mathbf{A}^{-} stands for the Moore-Penrose inverse of \mathbf{A} . Non-bold letters represent scalar (real-valued) quantities or



Figure 1: Road-map of the main results in this manuscript. L, P, T and C stand for Lemma, Proposition, Theorem and Corollary, respectively. The suffix indicates that the result is stated in Appendix A. The missing results have either no direct connection with the above results (L4.1, P6) or are essential for many others (LA.1).

non-numerical objects (sets, graphs, vertices, edges, or points). The (i, j)-th entry of **A** is denoted A_{ij} , while the *i*-th components of $\boldsymbol{\delta}$ and \boldsymbol{Z} are δ_i and Z_i , respectively. The all-ones vector of size n, all-ones matrix of size $n \times n$, and identity matrix of size $n \times n$ are denoted as $\mathbf{1}_n, \mathbf{1}_{n \times n}$ and \mathbf{I}_n , respectively. Finally, we define $\mathbf{J}_n := \text{diag} [0, 0, \dots, 0, 1]^\top$ $(n \times n \text{ diagonal matrix with only one non-zero entry}).$

Continuity, differentiation, integration and scalar- or matrix-valued functions (except power functions) involving matrices are understood as being performed element-wise. Concerning the power functions, \mathbf{A}^2 is \mathbf{A} multiplied by itself, $\sqrt{\mathbf{A}} = \mathbf{A}^{1/2}$ is the principal square root of \mathbf{A} , and \mathbf{A}^{-1} is the inverse of \mathbf{A} in the matrix sense. Products (*e.g.* \mathbf{AB} or $\mathbf{A\delta}$) and Kronecker products (*e.g.* $\mathbf{A} \otimes \mathbf{B}$) are also understood in the matrix sense, not as element-wise operations.

2.2 Gaussian random fields on graphs with Euclidean edges

The paper deals with a vector-valued Gaussian random field Z defined on a metric graph as exposed below. We assume a random field with zero-mean

vector and with a matrix-valued covariance matrix $\mathbf{K} : \mathcal{G} \times \mathcal{G} \to \mathbb{R}^{p \times p}$ that is isotropic in the sense there exists a pair (Ψ, \mathbf{D}) , with $\Psi : [0, +\infty)^{p \times p} \to \mathbb{R}^{p \times p}$ and $\mathbf{D} : \mathcal{G} \times \mathcal{G} \to [0, +\infty)^{p \times p}$, both continuous, such that Equation (2) is satisfied. We work with a specific class of metric graphs, termed graphs with Euclidean edges as being proposed by Anderes et al. [2020] to generalise linear networks. The main idea underlying this construction is to associate every edge of the graph to a segment of the real line. As a consequence, it is possible to consider both the vertices and the points over the edges as actual points of the graph, where distances can be computed and random fields can be defined. As this is natural on linear networks as well, we stress that graphs with Euclidean edges do not have any restriction on the topology, *i.e.* the edges' lengths are free to vary and so are the connections between the vertices. Here we rephrase their original definition.

Definition 2.1 (Graph with Euclidean edges). The type of topology considered by Anderes et al. [2020] (to which the reader is referred for additional details and motivation) is called graph with Euclidean edges, denoted with a triple $\mathcal{G} = (V, E, \{\varphi_e\}_{e \in E})$ throughout, where the elements are blended in the following way:

- (a) (V, E) has a graph structure, *i.e.* V is the set of vertices and $E \subset V \times V$ accounts for the edges. We assume that this graph is simple and connected, *i.e.* V is finite, the graph has not repeated edges or edges that join a vertex to itself and every pair of vertices is connected by a path.
- (b) Each edge $e \in E$ is provided with a *length* $\ell(e) > 0$ and a *weight* $w(e) := 1/\ell(e)$; furthermore, it is associated with a unique abstract set, also denoted e, such that V and all the edge sets are mutually disjoint.
- (c) Let v_1 and v_2 be vertices connected by $e \in E$. Then, φ_e is a continuous and bijective mapping defined on $e \cup \{v_1, v_2\}$, such that φ_e maps e onto an open interval $(0, \ell(e)) \subset \mathbb{R}$ and $\{v_1, v_2\}$ onto $\{0, \ell(e)\}$.

Henceforth, we shall assume the existence of a total order relation on the set of vertices V and that every edge is represented through the ordered pair (v_1, v_2) , where $v_1 < v_2$. By abuse of notation, we write $u \in \mathcal{G}$ to denote a point on the graph: it can be either a vertex or a point on an edge. Each point $u \in \mathcal{G}$ will be identified with the triple $(\underline{u}, \overline{u}, \delta)$, where: $\underline{u} < \overline{u}$ are the



Figure 2: Left: a graph with Euclidean edges, where the bijections φ_{e_1} and φ_{e_2} have been highlighted. Right: a Euclidean tree with 5 leaves where the role of \underline{u} , \overline{u} and $\delta_{e_3}(u)$ have been stressed. Adaptation of Filosi et al. [2023, Figure 1].

endpoints of the edge e containing u and $\delta \in [0, 1]$ is the relative distance of u from \underline{u} ; formally: $\delta = \delta(u) := w(e)\varphi_e(u)$. Notice that, whenever u is a vertex, it is always possible to write $u = (\underline{u}, \overline{u}, \delta)$ for $\delta \in \{0, 1\}$, though the choice of \underline{u} and \overline{u} may not be unique. Finally, we use the notation e(u), where $e : \mathcal{G} \to E$, to denote the edge that contains u. If $u \in V$, then e(u)is any of the edges that have an endpoint in u. It is in order to note that our definition slightly deviates from the original one in Anderes et al. [2020]: we do not require any distance consistency property, as we do not use the geodesic metric.

A graph with Euclidean edges is called an *Euclidean tree* if it has a tree structure, *i.e.* given any two points $u_1, u_2 \in \mathcal{G}$, there is exactly one path that connects u_1 to u_2 . A vertex of a Euclidean tree is called a *leaf* if it is connected to only one other vertex. Figure 2 aims to clarify the notions introduced so far.

2.3 Classes of matrices used in the paper

An $n \times n$ real matrix **A** is said to be *positive semidefinite* if and only if $\mathbf{c}^{\top} \mathbf{A} \mathbf{c} \geq 0$ for all $\mathbf{c} \in \mathbb{R}^n$, and *conditionally negative semidefinite* if and only if $\mathbf{c}^{\top} \mathbf{A} \mathbf{c} \leq 0$ for all $\mathbf{c} \in \mathbb{R}^n$ such that $\mathbf{1}_n^{\top} \mathbf{c} = 0$.

An $n \times n$ matrix **L** is called *quasi-Laplacian* if and only if it is symmetric, positive semidefinite, and has exactly one null eigenvalue, with corresponding eigenvector $\mathbf{1}_n$. A quasi-Laplacian matrix **L** is called *Laplacian* if and only if it has non-positive off-diagonal entries. Notice that Laplacian matrices arise naturally from weighted graphs: there is a bijection between the set of (positively) weighted, simple and undirected graphs and the set of Laplacian matrices [Devriendt, 2022, Subsection 2.1]. Henceforth, we will consider the Laplacian matrix of a given graph with Euclidean edges, with entries defined as Devriendt [2022, Equation (1)]:

$$L_{ij} = \begin{cases} \sum_{v \in V} w((v, v_i)) & \text{if } i = j \\ -w((v_i, v_j)) & \text{if } i \neq j, \end{cases}$$
(3)

with $w((v_i, v_j))$ the weight of the edge joining vertices v_i and v_j if these vertices are connected, 0 otherwise.

Quasi-Laplacian matrices have been introduced as a generalisation of Laplacian matrices, as they will often appear in this manuscript. Their interpretation is slightly blurred, since, thinking about the bijection between Laplacian matrices and weighted graphs, it seems natural to interpret quasi-Laplacian matrices as Laplacian matrices of graphs with possibly negative weights. Nevertheless, as one may expect, quasi-Laplacian matrices share several properties of Laplacian matrices formally stated below.

Proposition 1. Let L be a quasi-Laplacian matrix partitioned as follows:

$$\mathbf{L} = \begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{B}^\top & \mathbf{C} \end{bmatrix},\tag{4}$$

where A and C are square matrices. Then the following statements hold.

- 1. $\mathbf{B} \neq \mathbf{0}$.
- 2. A and C are strictly positive definite.

- 3. Both $\mathbf{L}/\mathbf{C} := \mathbf{A} \mathbf{B}\mathbf{C}^{-1}\mathbf{B}^{\top}$ and $\mathbf{L}/\mathbf{A} := \mathbf{C} \mathbf{B}^{\top}\mathbf{A}^{-1}\mathbf{B}$ are quasi-Laplacian matrices, i.e., the set of quasi-Laplacian matrices is closed under the Schur complement operation.
- 4. \mathbf{L}^- is a quasi-Laplacian matrix.

2.4 Metrics used in the paper

The work by Anderes et al. [2020] shows that there is no unique way to define a metric over a graph with Euclidean edges. While the geodesic distance is physically intuitive, Anderes et al. [2020] prove that it has very limited use in terms of available covariance functions. Further, when considering the geodesic distance not all graphs with Euclidean edges become *permissible* [Anderes et al., 2020], and a collection of technical restrictions is required. Alternatively, one can use the resistance metric, being a generalisation of the electric distance used for electric circuits [Klein and Randic, 1993]. To provide a description of the resistance metric, some further background is needed. Let X be a set and $Z : X \to \mathbb{R}$ a square-integrable random field. The *variogram* of Z is defined via

$$\gamma_Z : X \times X \to \mathbb{R}$$

(x₁, x₂) $\mapsto \gamma_Z(x_1, x_2) := \mathbb{V}\mathrm{ar} \left(Z(x_1) - Z(x_2) \right).$

Clearly, a variogram is always symmetric and non-negative valued. In addition, it is conditionally negative semidefinite, *i.e.* it satisfies [Chilès and Delfiner, 2012]

$$\sum_{i,j=1}^{n} c_i c_j \gamma(x_i, x_j) \le 0 \tag{5}$$

whenever $\sum_{i=1}^{n} c_i = 0$, for any $x_1, \ldots, x_n \in X$.

Anderes et al. [2020] define the resistance metric, denoting d_R throughout, as the variogram of a random field Z on the graph with Euclidean edges \mathcal{G} , that is constructed *ad hoc* in order to be as much as possibly parenthetical to a Brownian bridge on the graph. Technicalities are deferred to their paper.

The concept of variogram can be generalised to the case of a zero-mean square-integrable vector-valued random field \mathbf{Z} on a non-empty set X, through the following lines. The *pseudo-variogram* of \mathbf{Z} is defined through the iden-

tity

$$\Gamma_{\mathbf{Z}} : X \times X \to [0, +\infty)^{p \times p}$$

$$(x_1, x_2) \mapsto \Gamma_{\mathbf{Z}}(x_1, x_2) := \left[\operatorname{Var} \left(Z_i(x_1) - Z_j(x_2) \right) \right]_{i,j=1}^p.$$
(6)

Straightforward calculations allow to rewrite the above as

$$\boldsymbol{\Gamma}_{\boldsymbol{Z}}(x_1, x_2) = (\operatorname{diag} \operatorname{\mathbb{V}ar} \boldsymbol{Z}(x_1)) \ \boldsymbol{1}_p^{\top} + \boldsymbol{1}_p \ (\operatorname{diag} \operatorname{\mathbb{V}ar} \boldsymbol{Z}(x_2))^{\top} - 2 \operatorname{\mathbb{C}ov} \left(\boldsymbol{Z}(x_1), \boldsymbol{Z}(x_2)\right),$$
(7)

where \mathbb{V} ar $\mathbf{Z}(x_1)$ is the $p \times p$ collocated variance-covariance matrix of $\mathbf{Z}(x_1)$, diag \mathbb{V} ar $\mathbf{Z}(x_1)$ is the $p \times 1$ vector containing the main diagonal of \mathbb{V} ar $\mathbf{Z}(x_1)$ and \mathbb{C} ov $(\mathbf{Z}(x_1), \mathbf{Z}(x_2)) := \mathbb{E}(\mathbf{Z}(x_1)\mathbf{Z}(x_2)^{\top})$. The pseudo-variogram satisfies the following properties [Dörr and Schlather, 2023]:

• for any $x_1, x_2 \in X$,

$$\boldsymbol{\Gamma}_{\boldsymbol{Z}}(x_1, x_2) = \boldsymbol{\Gamma}_{\boldsymbol{Z}}(x_2, x_1)^{\top}; \tag{8}$$

- for any $x \in X$, the diagonal entries of $\Gamma_{\mathbf{Z}}(x, x)$ are zero;
- $\Gamma_{\mathbf{Z}}$ is $\mathbf{1}_p$ -conditionally negative semidefinite, *i.e.* for all $n \in \mathbb{N}^+$, $x_1, \ldots, x_n \in X$ and $\mathbf{c}_1, \ldots, \mathbf{c}_n \in \mathbb{R}^p$ such that $\mathbf{1}_p^\top \sum_{i=1}^n \mathbf{c}_i = 0$, it holds

$$\sum_{i,j=1}^{n} \boldsymbol{c}_{i}^{\mathsf{T}} \boldsymbol{\Gamma}_{\boldsymbol{Z}}(x_{i}, x_{j}) \boldsymbol{c}_{j} \leq 0.$$
(9)

In the following, we define the multivariate distance $\mathbf{D} : X \times X \rightarrow [0, +\infty)^{p \times p}$ via $\mathbf{D}(x_1, x_2) := \mathbf{\Gamma}_{\mathbf{L}}(x_1, x_2)$. It is in order to notice that at this point the matrix-valued distance \mathbf{D} is *not* symmetric, *i.e.* in general $D_{ij}(x_1, x_2) \neq D_{ji}(x_1, x_2)$. This may pave the way for new asymmetric metrics.

2.5 Classes of functions and Schoenberg characterisations

This section follows closely Zastavnyi [2023]. A function $k : X \times X \to \mathbb{R}$ is positive semidefinite if, for all $n \in \{1, 2, ...\}, x_1, ..., x_n \in X$ and $c_1, ..., c_n \in \mathbb{R}$, one has

$$\sum_{i,j=1}^{n} c_i c_j k(x_i, x_j) \ge 0.$$
(10)

In addition, k is a strictly positive definite function if it is positive semidefinite and

$$\sum_{i,j=1}^{n} c_i c_j k(x_i, x_j) = 0 \implies c_1 = \dots = c_n = 0.$$
(11)

The extension to the matrix-valued case reads as follows. A matrix-valued function $\mathbf{K}: X \times X \to \mathbb{R}^{p \times p}$ is positive semidefinite if, for all $n \in \{1, 2, \ldots\}$, $c_1, \ldots, c_n \in \mathbb{R}^p$ and $x_1, \ldots, x_n \in X$, it holds:

$$\sum_{i,j=1}^{n} \boldsymbol{c}_{i}^{\top} \mathbf{K}(x_{i}, x_{j}) \boldsymbol{c}_{j} \ge 0.$$
(12)

In addition, \mathbf{K} is strictly positive definite if it is positive semidefinite and the condition

$$\sum_{i,j=1}^{n} \boldsymbol{c}_{i}^{\mathsf{T}} \mathbf{K}(x_{i}, x_{j}) \boldsymbol{c}_{j} = 0$$
(13)

implies that there exists a pair $i \neq j$ such that $x_i = x_j$ or $c_1 = \cdots = c_n = 0$. A relevant result for our developments is coming from Proposition 1 in Zastavnyi [2023], which is formally stated for a neater exposition.

Proposition 2. Let $\mathbf{K} : X \times X \to \mathbb{R}^{p \times p}$. Then, the following are equivalent:

- K is positive semidefinite;
- for all $n \in \{1, 2, \ldots\}$ and for all $x_1, \ldots, x_n \in X$, the $np \times np$ block matrix $\mathbf{C} := [C_{ij}]_{i,j=1}^{np} := [\mathbf{K}(x_i, x_j)]_{i,j=1}^n$ is positive semidefinite, i.e., $\mathbf{c}^{\top} \mathbf{C} \mathbf{c} \geq 0$ for any $\mathbf{c} \in \mathbb{R}^{np}$.

A function $\psi : [0, +\infty) \to [0, +\infty)$ is called *completely monotone* if it is continuous on $[0, +\infty)$, infinitely differentiable on $(0, +\infty)$ and for each $i \in \mathbb{N}$ it holds

$$(-1)^{i}\psi^{(i)}(x) \ge 0,$$

where $\psi^{(i)}$ denotes the *i*th derivative of ψ and $\psi^{(0)} := \psi$. A function ψ : $[0, +\infty) \rightarrow [0, +\infty)$ is completely monotone if and only if it is the Laplace transform of a (unique) finite Borel measure μ on $[0, +\infty)$, *i.e.*

$$\psi(x) = \int_0^{+\infty} \exp(-tx) \,\mu(\mathrm{d}t), \quad x \in [0, +\infty).$$
 (14)

A function $g: [0, +\infty) \to [0, +\infty)$ is called a *Bernstein function* if g in infinitely differentiable on $[0, +\infty)$ and $g^{(1)}$ is a completely monotone function.

Our expository material is now completed by reporting Theorem 5 of Zastavnyi [2023].

Theorem 2.2. Let $\psi : [0, +\infty) \to (0, +\infty)$ be a non-constant completely monotone function, $g : [0, +\infty) \to [0, +\infty)$ be a Bernstein function and let $\Gamma_{\mathbf{Z}} : X \times X \to [0, +\infty)^{p \times p}$, for a non-empty set X. If, for all $x_1, x_2 \in X$, $\Gamma_{\mathbf{Z}}^{\top}(x_1, x_2) = \Gamma_{\mathbf{Z}}(x_2, x_1)$ and $\Gamma_{\mathbf{Z}}$ is $\mathbf{1}_p$ -conditionally negative semidefinite, then, for all $\xi > 0$,

$$(x_1, x_2) \mapsto \psi \Big(\xi g \left(\Gamma_{\boldsymbol{Z}}(x_1, x_2) \right), \qquad x_1, x_2 \in X,$$

where ψ and g are applied element-wise, is a positive semidefinite function on X.

3 General Results

The following notation will ease the exposition throughout:

- n is the number of vertices of the graph \mathcal{G} (cardinality of V),
- L is the Laplacian matrix of \mathcal{G} as per Equation (3),
- $\Sigma := L^{-}$ is the Moore-Penrose inverse of L,
- Λ and \mathbf{W} are the eigenvalues and eigenvectors matrices of Σ , *i.e.* $\Sigma = \mathbf{W} \Lambda \mathbf{W}^{\top}$,
- for two matrices A, B of the same size and for a positive integer p, we define

$$\mathcal{M}_{p}\left(\mathbf{A},\mathbf{B}\right) := \mathcal{M}_{p}\left(\begin{array}{cc}\mathbf{A}\\\mathbf{B}\end{array}\right) := \mathbf{I}_{p} \otimes \mathbf{A} + (\mathbf{1}_{p \times p} - \mathbf{I}_{p}) \otimes \mathbf{B} = \begin{bmatrix} \mathbf{A} & \mathbf{B} & \dots & \mathbf{B}\\ \mathbf{B} & \mathbf{A} & \dots & \mathbf{B}\\ \vdots & \vdots & \ddots & \vdots\\ \mathbf{B} & \mathbf{B} & \dots & \mathbf{A} \end{bmatrix}$$
(15)

The construction provided in this section extends, to the matrix-valued case, that in Anderes et al. [2020]. Specifically, we shall consider a vector-valued random field, Z, that is obtained through the identity

$$\mathbf{Z}(u) := \mathbf{Z}_V(u) + \mathbf{Z}_E(u), \qquad u \in \mathcal{G}.$$
 (16)

In turn, the random fields Z_V and Z_E obeys to different construction principles. Specifically, Z_V is a multivariate Gaussian random vector on the vertices and is linearly interpolated on the edges, whilst Z_E is zero on the vertices and adds some variability on the edges. We note that Anderes et al. [2020] consider this construction for the case p = 1 and this entails the use of a specific class of variograms that are then used to build the resistance metric, d_R . It is not surprising that our construction will rely on the pseudo-variogram Γ_Z for the generalisation to the *p*-variate setting with p > 1. While these constructions are mathematically involved, a simple description is provided here, with technicalities and proofs deferred to Appendix A.

3.1 Construction for the Vertices

This section provides a construction for the random field \mathbf{Z}_V as per the identity (16). While the search for metrics is probably unlimited in terms of alternatives, our spectrum is suitably restricted by providing two *desiderata*, denoted \mathbb{D} throughout.

- D1: The construction (16) needs to provide a pseudo-variogram Γ_Z for which the diagonal entries should obey to the variogram construction as in Anderes et al. [2020].
- D2: The multivariate random field Z_V on V should enjoy a homogeneous conditional independence structure. More precisely, we set the block-precision matrix Θ of Z_V to have the following structure:

$$\boldsymbol{\Theta} = \mathcal{M}_p \begin{pmatrix} \mathbf{Q} \\ -\alpha \mathbf{I}_n \end{pmatrix}, \qquad (17)$$

where α is a positive real value and \mathbf{Q} is a suitable $n \times n$ matrix that will be defined next. This means that, for $i \neq j$ and $v_1 \neq v_2$, $Z_{V,i}(v_1)$ and $Z_{V,j}(v_2)$ are conditionally independent given everything else. Define the $n \times n$ matrices:

$$\mathbf{Q} := \frac{1}{2} \left(\mathbf{L} + \alpha(p-2)\mathbf{I}_n + \sqrt{\mathbf{L}^2 - 2\alpha(p-2)\mathbf{L} + \alpha^2 p^2 \mathbf{I}_n} \right)$$
(18)
$$= \frac{1}{2} \mathbf{W} \left(\mathbf{\Lambda}^- + \alpha(p-2)\mathbf{I}_n + \sqrt{(\mathbf{\Lambda}^-)^2 - 2\alpha(p-2)\mathbf{\Lambda}^- + \alpha^2 p^2 \mathbf{I}_n} \right) \mathbf{W}^\top,$$
(19)

$$\mathbf{X} := \frac{1}{\alpha(p-1)} \left(\boldsymbol{\Sigma} \mathbf{Q} - \mathbf{I}_n \right) \tag{20}$$

$$=\frac{1}{2\alpha(p-1)}\mathbf{W}\left(\alpha(p-2)\mathbf{\Lambda}+\sqrt{\mathbf{I}_n-\mathbf{J}_n-2\alpha(p-2)\mathbf{\Lambda}+\alpha^2p^2\mathbf{\Lambda}^2}-\mathbf{I}_n-\mathbf{J}_n\right)\mathbf{W}^{\top}$$
(21)

Notice that in Equations (19) and (21) the principal square root coincides with the element-wise square root, being its arguments diagonal matrices. These equations have been established from Equations (18) and (20) by using the fact that \mathbf{W} is an orthogonal matrix and that $\mathbf{\Lambda}\mathbf{\Lambda}^- = \mathbf{I}_n - \mathbf{J}_n$ insofar as the first p - 1 diagonal entries of $\mathbf{\Lambda}$ are positive and the last diagonal entry is zero. In addition, we define the two positive constants

$$k_1 := \frac{p-1}{\alpha n p^2}$$
 $k_2 := \frac{p^2 - p + 1}{\alpha n p^2 (p-1)},$

and the $n \times n$ matrices

$$\tilde{\mathbf{\Sigma}} := \mathbf{\Sigma} + k_1 \mathbf{1}_{n imes n}$$
 $ilde{\mathbf{X}} := \mathbf{X} + k_2 \mathbf{1}_{n imes n}$

Although these definitions may appear peculiar, they are the only solution to the *desiderata* \mathbb{D} previously stated and are crucial to define the random field \mathbf{Z}_V . We start with the vertices, where \mathbf{Z}_V is assumed a zero-mean *p*-variate Gaussian random vector $\mathbf{Z}_V|_V : V \to \mathbb{R}^p$ having covariance matrix-valued function

$$\mathbb{C}\mathrm{ov}\left(\boldsymbol{Z}_{V}(v_{1}), \boldsymbol{Z}_{V}(v_{2})\right) := \mathcal{M}_{p}\left(\begin{array}{c} \boldsymbol{\tilde{\Sigma}}[v_{1}, v_{2}]\\ \boldsymbol{\tilde{X}}[v_{1}, v_{2}] \end{array}\right) \in \mathbb{R}^{p \times p}.$$
(22)

Checking the positive semidefiniteness of (22) relies on a result of independent interest which implies that both Θ and Θ^- are positive semidefinite matrices. We state this formally below.

Proposition 3. The matrix Θ defined in Equation (17) and its Moore-Penrose inverse Θ^- are quasi-Laplacian. **Proposition 4.** The function defined in (22) is positive semidefinite.

Once \mathbf{Z}_V is defined over the vertices, the corresponding values over the edges are attained through a sheer linear interpolation. Namely, for $u = (\underline{u}, \overline{u}, \delta(u)) \in \mathcal{G}$, we set

$$\mathbf{Z}_{V}(u) := (1 - \delta(u))\mathbf{Z}_{V}(\underline{u}) + \delta(u)\mathbf{Z}_{V}(\overline{u}).$$
⁽²³⁾

3.2 Construction for the Edges

We define Z_E as a *p*-variate zero-mean Gaussian random field independent of Z_V whose covariance matrix-valued function is:

$$\mathbb{C}\operatorname{ov}\left(\boldsymbol{Z}_{E}(u_{1}), \boldsymbol{Z}_{E}(u_{2})\right) := \mathbb{1}\left(e_{1} = e_{2}\right)\ell(e_{1})\left(\delta_{1} \wedge \delta_{2} - \delta_{1}\delta_{2}\right)\mathcal{M}_{p}\begin{pmatrix}1\\\beta\end{pmatrix} \in [0, +\infty)^{p \times p},$$
(24)

for $e_i := e(u_i)$, $\delta_i := \delta(u_i)$, $i \in \{1, 2\}$ and where $\beta \in [0, 1]$ is a fixed parameter. Notice that $\mathbf{Z}_E\Big|_{e_1}$ is independent from $\mathbf{Z}_E\Big|_{e_2}$ whenever $e_1 \neq e_2$, in addition $\ell(e_1)$ ($\delta_1 \wedge \delta_2 - \delta_1 \delta_2$) is the covariance function of a standard Brownian bridge on $[0, \ell(e_1)]$, whilst $\mathcal{M}_p(1, \beta)$ is positive semidefinite for $\beta \in [0, 1]$. This ensures that (24) is positive semidefinite, as the element-wise product of positive semidefinite functions is itself positive semidefinite.

3.3 Compendium

The following result comes directly from the above definitions.

Proposition 5. Let $\mathbf{K}_{\mathbf{Z}_V}$, $\mathbf{K}_{\mathbf{Z}_E}$ be the covariance mappings associated respectively with the random fields \mathbf{Z}_V and \mathbf{Z}_E . Then, it is true that

$$\mathbf{K}_{\boldsymbol{Z}_{V}}(u_{1}, u_{2}) = \mathcal{M}_{p} \begin{pmatrix} \boldsymbol{\delta}_{1,V}^{\top} \tilde{\boldsymbol{\Sigma}} \boldsymbol{\delta}_{2,V} \\ \boldsymbol{\delta}_{1,V}^{\top} \tilde{\mathbf{X}} \boldsymbol{\delta}_{2,V} \end{pmatrix} \in \mathbb{R}^{p \times p},$$
(25)

for $u_i = (\underline{u}_i, \overline{u}_i, \delta_i) \in \mathcal{G}$ $(i \in \{1, 2\})$, and where $\delta_{i,V}$ is the n-dimensional vector whose entries are

$$\delta_{i,V} := \begin{cases} 1 - \delta_i & \text{if } v = \underline{u}_i \\ \delta_i & \text{if } v = \overline{u}_i \\ 0 & \text{otherwise} \end{cases}, \quad v \in V.$$
(26)

Further, it is true that

$$\mathbf{K}_{\mathbf{Z}_{E}}(u_{1}, u_{2}) = \mathcal{M}_{p} \begin{pmatrix} \mathbb{1} (e_{1} = e_{2}) \ell(e_{1}) (\delta_{1} \wedge \delta_{2} - \delta_{1} \delta_{2}) \\ \beta \mathbb{1} (e_{1} = e_{2}) \ell(e_{1}) (\delta_{1} \wedge \delta_{2} - \delta_{1} \delta_{2}) \end{pmatrix}.$$

As a direct implication of Proposition 5 in concert with the fact that \mathbf{Z}_V and \mathbf{Z}_E are independent, we obtain the following.

Corollary 1. It is true that

$$\mathbf{K}_{\mathbf{Z}}(u_{1}, u_{2}) = \mathbf{K}_{\mathbf{Z}_{V}}(u_{1}, u_{2}) + \mathbf{K}_{\mathbf{Z}_{E}}(u_{1}, u_{2})$$

$$= \mathcal{M}_{p} \begin{pmatrix} \boldsymbol{\delta}_{1,V}^{\top} \tilde{\boldsymbol{\Sigma}} \boldsymbol{\delta}_{2,V} + \mathbb{1} (e_{1} = e_{2}) \ell(e_{1}) (\delta_{1} \wedge \delta_{2} - \delta_{1} \delta_{2}) \\ \boldsymbol{\delta}_{1,V}^{\top} \tilde{\mathbf{X}} \boldsymbol{\delta}_{2,V} + \beta \mathbb{1} (e_{1} = e_{2}) \ell(e_{1}) (\delta_{1} \wedge \delta_{2} - \delta_{1} \delta_{2}) \end{pmatrix}.$$
(27)

We are now able to derive the matrix-valued metric associated with the random field \mathbf{Z} in (16). To do so, we let $\mathbf{D} : \mathcal{G} \times \mathcal{G} \to [0, +\infty)^{p \times p}$ be the pseudo-variogram of \mathbf{Z} , *i.e.*

$$\mathbf{D}(u_1, u_2) := \mathbf{\Gamma}_{\mathbf{Z}}(u_1, u_2) = \left[\mathbb{V}\mathrm{ar} \left(Z_i(u_1) - Z_j(u_2) \right) \right]_{i,j=1}^p.$$
(28)

A straightforward application of Equation (7) in concert with tedious calculations provides the following fact.

Proposition 6. It is true that

$$\mathbf{D}(u_{1}, u_{2}) = \mathcal{M}_{p} \begin{pmatrix} (\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V})^{\top} \tilde{\boldsymbol{\Sigma}} (\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}) \\ \boldsymbol{\delta}_{1,V}^{\top} \tilde{\boldsymbol{\Sigma}} \tilde{\boldsymbol{\delta}}_{1,V} + \boldsymbol{\delta}_{2,V}^{\top} \tilde{\boldsymbol{\Sigma}} \tilde{\boldsymbol{\delta}}_{2,V} - 2\boldsymbol{\delta}_{1,V}^{\top} \tilde{\boldsymbol{X}} \boldsymbol{\delta}_{2,V} \end{pmatrix} + \mathcal{M}_{p} \begin{pmatrix} \ell(e_{1})\delta_{1}(1 - \delta_{1}) + \ell(e_{2})\delta_{2}(1 - \delta_{2}) - 2\mathbb{1} (e_{1} = e_{2}) \ell(e_{1}) (\delta_{1} \wedge \delta_{2} - \delta_{1}\delta_{2}) \\ \ell(e_{1})\delta_{1}(1 - \delta_{1}) + \ell(e_{2})\delta_{2}(1 - \delta_{2}) - 2\beta\mathbb{1} (e_{1} = e_{2}) \ell(e_{1}) (\delta_{1} \wedge \delta_{2} - \delta_{1}\delta_{2}) \end{pmatrix},$$
(29)

for $u_1, u_2 \in \mathcal{G}$, with notation as in Proposition 5.

Notice that the former line coincides with the contribution of the random field \mathbf{Z}_V , whilst the latter is the contribution of \mathbf{Z}_E .

We describe below some properties of the metric \mathbf{D} provided through our construction.

Proposition 7. The matrix-valued expression (28) satisfies properties similar to the (real-valued) quasi-metrics, i.e. it satisfies:

- 1. $D_{ij}(u_1, u_2) \ge 0;$
- 2. $D_{ij}(u_1, u_2) = 0 \iff u_1 = u_2 \text{ and } i = j;$
- 3. $\mathbf{D}(u_1, u_2) = \mathbf{D}(u_1, u_2)^\top = \mathbf{D}(u_2, u_1) = \mathbf{D}(u_2, u_1)^\top$ for any $u_1, u_2 \in \mathcal{G}$;
- 4. The mapping $(u_1, u_2) \mapsto \mathbf{D}(u_1, u_0) \mathbf{A} + \mathbf{A}^\top \mathbf{D}(u_0, u_2) \mathbf{D}(u_1, u_2) \mathbf{A}^\top \mathbf{D}(u_0, u_0) \mathbf{A}$ is positive semidefinite for any $u_0 \in \mathcal{G}$ and any $p \times p$ matrix \mathbf{A} such that $\mathbf{A}^\top \mathbf{1}_p = \mathbf{1}_p$;
- 5. The matrices $\mathbf{D}(u_1, u_2) \mathbf{D}(u_0, u_0)$ and $2\mathbf{D}(u_1, u_0) + 2\mathbf{D}(u_2, u_0) \mathbf{D}(u_1, u_2) 3\mathbf{D}(u_0, u_0)$ are symmetric and positive semidefinite for any $u_0, u_1, u_2 \in \mathcal{G}$.

Note the resemblance between the fourth property in Proposition 7 and the second statement in Zastavnyi [2023, Proposition 4].

Proposition 8. Let \mathcal{G} be a graph with Euclidean edges and let $\mathbf{D} : \mathcal{G} \times \mathcal{G} \rightarrow [0, +\infty)^{p \times p}$ as defined in (28) and having elements D_{ij} , $i, j = 1, \ldots, p$. Then, the following properties hold:

- 1. for all $i \in \{1, ..., p\}$, the distance D_{ii} coincides with the resistance distance defined by Anderes et al. [2020], i.e. $D_{ii}(u_1, u_2) = d_R(u_1, u_2)$;
- 2. the distance **D** is matrix-homogeneous, i.e. for every $i \neq j$ and $i' \neq j'$, we have $D_{ij}(u_1, u_2) = D_{i'j'}(u_1, u_2)$ and $D_{ii}(u_1, u_2) = D_{i'i'}(u_1, u_2)$.

Albeit counter-intuitive, the distance within the same variable is not always less than the same distance among variables. In formulae, $D_{ii}(u_1, u_2) \not\leq D_{ij}(u_1, u_2)$, in general. Appendix B shows this fact through a worked example.

The following result illustrates some asymptotic properties of the proposed metric.

Proposition 9. Let \mathcal{G} be a graph with Euclidean edges and let \mathbf{Q} and \mathbf{X} as defined in Equations (18)-(21). Then, the asymptotic results indicated in Table 1 hold.

Theorem 3.1. Let \mathcal{G} be a graph with Euclidean edges and $\mathbf{D} : \mathcal{G} \times \mathcal{G} \rightarrow [0, +\infty)^{p \times p}$ the distance defined at (28). In addition, let $\psi : [0, +\infty) \rightarrow (0, +\infty)$ be a non-constant completely monotone function and $g : [0, +\infty) \rightarrow (0, +\infty)$

Table 1: Asymptotic results with respect to p and α . In the 4th row, $0 < \tau \leq \lambda_{n-1}$, with λ_{n-1} being the smallest positive eigenvalue of Σ , while in the last row, $\tau \geq \lambda_1$. In the last column, the results involving $d_R(u_1, u_2)$ hold only for $\beta \to 1$ (the other result holds for every $\beta \in [0, 1]$). Emilio, per essere formali qui dovremmo modificare le prime due righe della colonna Q e metterci il limite: sei d'accordo o lasciamo così?

p	α	Q	X	$D_{ij}(u_1, u_2)$ for $i \neq j$
$+\infty$	fixed	$\alpha(p-1)\mathbf{I}_n$	Σ	$d_R(u_1, u_2)$
fixed	$+\infty$	$\alpha(p-1)\mathbf{I}_n$	Σ	$d_R(u_1, u_2)$
fixed	0+	\mathbf{L}	$-rac{1}{lpha n(p-1)}1_{n imes n}$	$\frac{2}{\alpha np}$
$+\infty$	$\frac{1}{p\tau}$	$\frac{\mathbf{I}_n}{\tau}$	$\mathbf{\Sigma} - \mathbf{\tau} \mathbf{I}_n$	$d_R(u_1,u_2) + 2 au oldsymbol{\delta}_{1,V}^ op oldsymbol{\delta}_{2,V}$
$+\infty$	$\frac{1}{p\tau}$	$\mathbf{L} + rac{1}{n au} 1_{n imes n}$	$-rac{ au}{n}1_{n imes n}$	$d_R(u_1, u_2) + \frac{2\tau}{n} + 2\boldsymbol{\delta}_{1,V}^{\top} \mathbf{X} \boldsymbol{\delta}_{2,V}$

 $[0, +\infty)$ a Bernstein function. Then, for every $\xi > 0$, the mapping **K** defined as

$$(u_1, u_2) \mapsto \mathbf{K}(u_1, u_2) = \psi\left(\xi g\left(\mathbf{D}(u_1, u_2)\right)\right),\tag{30}$$

where ψ and g are applied element-wise, is a valid covariance function. If, in addition, $\psi(0) = 1$, then (30) is a valid correlation function.

One of the main drawbacks of this construction is that the distance is homogeneous (see Proposition 8). Consequently, not only do the resulting matrix-valued covariance function (30) assigns the same marginal covariance for all the variables, *i.e.* $\psi(\xi g(D_{ii}(u_1, u_2))) = \psi(\xi g(d_R(u_1, u_2)))$ for any $u_1, u_2 \in \mathcal{G}$ and any $i \in \{1, \ldots, p\}$, but also the cross-covariance between two different variables is independent from the variables themselves, *i.e.* $\psi(\xi g(D_{ij}(u_1, u_2)))$ does not depend on i, j as long as $i \neq j$. Such a restriction can be easily circumvented. Let $[\rho_{ij}]_{i,j=1}^p$ be a collocated correlation matrix, *e.g.*, $\rho_{ii} = 1$ and $-1 \leq \rho_{ij} \leq 1$ when $i \neq j$, and the matrix being symmetric and positive semidefinite. Let $\boldsymbol{\sigma} = (\sigma_1, \ldots, \sigma_p)^{\top}$ with $\sigma_i > 0$ for $i = 1, \ldots, p$. Let **K** be the mapping in (30) and define the mapping $\widetilde{\mathbf{K}} : \mathcal{G} \times \mathcal{G} \to \mathbb{R}^{p \times p}$ with entries

$$\bar{K}_{ij}(u_1, u_2) := \sigma_i \sigma_j \rho_{ij} K_{ij}(u_1, u_2), \quad i, j = 1, \dots, p, \quad u_1, u_2 \in \mathcal{G}$$

Then, by straight application of the Schur product theorem, one gets that $\widetilde{\mathbf{K}}$ is a positive semidefinite matrix-valued function. However, all the marginal covariances (diagonal entries of $\widetilde{\mathbf{K}}$) are proportional, and the same happens for all the cross-covariances (off-diagonal entries of $\widetilde{\mathbf{K}}$), which is still restrictive. The results following throughout allow to circumvent such a limitation.

3.4 Generalisations

An interesting generalisation is provided below.

Theorem 3.2. Let $\theta \in (0,1]$. Let $\mathbf{F} : [0, +\infty) \to \mathbb{R}^{p \times p}$ be a Borel measure such that, for any fixed $\xi \geq 0$, the matrix $\mathbf{F}(d\xi)$ is symmetric and positive semidefinite. Let \mathbf{D} as in Equation (28) and let $\Psi(\cdot) : [0, +\infty) \to \mathbb{R}^{p \times p}$ be a continuous mapping determined through

$$\Psi(\mathbf{D}) := \left[\int_0^{+\infty} \exp(-\xi D_{ij}^{\theta}) F_{ij}(\mathrm{d}\xi) \right]_{i,j=1}^p,$$
(31)

where $F_{ij}(\mathbf{d}\cdot)$ and D_{ij} are the (i, j)-th entries of $\mathbf{F}(\mathbf{d}\cdot)$ and \mathbf{D} , respectively. Then, Ψ is the covariance function of a Gaussian random field \mathbf{Z} constructed according to the addition principle (16).

As a first application of this result, we adapt a construction that has been originally provided by Porcu et al. [2018] in Euclidean spaces and working with the classical Euclidean norm. The Shkarofsky-Gneiting family of functions $SG_{\alpha,\beta,\nu}(\cdot)$ is defined on $[0, +\infty)$ through

$$\mathcal{SG}_{\alpha,\beta,\nu}(t) = \left(1 + \frac{t}{\beta}\right)^{-\nu/2} \frac{\mathcal{K}_{\nu}\left(\frac{1}{\sqrt{\alpha}}\sqrt{\beta + t}\right)}{\mathcal{K}_{\nu}\left(\sqrt{\frac{\beta}{\alpha}}\right)}, \qquad t \ge 0,$$
(32)

where \mathcal{K}_{ν} stands for the modified Bessel function of the second kind. Arguments therein show that \mathcal{SG} is the Laplace transform of a positive and bounded function. The family is very interesting as it admits the special limit cases

$$\mathcal{SG}_{\alpha,0,-\nu}(t) = \mathcal{M}_{\alpha,\nu}(t) := \frac{2^{1-\nu}}{\Gamma(\nu)} \left(\frac{t}{\alpha}\right)^{\nu/2} \mathcal{K}_{\nu}\left(\sqrt{\frac{t}{\alpha}}\right), \qquad t \ge 0,$$

for $\alpha, \nu > 0$, and

$$\mathcal{SG}_{0,\beta,\nu}(t) = \mathcal{C}_{\beta,\nu}(t) := \left(1 + \frac{t}{\beta}\right)^{-\nu}, \qquad t \ge 0,$$

for $\beta, \nu > 0$, where $\mathcal{M}_{\alpha,\nu}(t) = \frac{2^{1+\nu}}{\Gamma(-\nu)} \left(\frac{\sqrt{t}}{\sqrt{\alpha}}\right)^{-\nu} \mathcal{K}_{-\nu} \left(\frac{\sqrt{t}}{\sqrt{\alpha}}\right), \alpha > 0, \nu > 0$ and where $\mathcal{C}_{\alpha,a,\nu}(t) = \left(1 + \frac{t^a}{\alpha}\right)^{-\nu}, \alpha > 0, \nu > 0$. We consider the case p = 2 and define the bivariate Shkarofsky-Gneiting mapping $\mathcal{SG}_2 : [0, +\infty)^{2\times 2} \to \mathbb{R}^{2\times 2}$ through

$$\boldsymbol{\mathcal{SG}}_{2}(\mathbf{D}) = \begin{bmatrix} \sigma_{1}^{2} \, \mathcal{SG}_{\alpha_{1},\beta_{1},\nu_{1}}(D_{11}^{\theta}) & \sigma_{1}\sigma_{2}\rho \, \mathcal{SG}_{\alpha_{12},\beta_{12},\nu_{12}}(D_{12}^{\theta}) \\ \\ \sigma_{1}\sigma_{2}\rho \, \mathcal{SG}_{\alpha_{12},\beta_{12},\nu_{12}}(D_{12}^{\theta}) & \sigma_{2}^{2} \, \mathcal{SG}_{\alpha_{2},\beta_{2},\nu_{2}}(D_{22}^{\theta}) \end{bmatrix}, \quad (33)$$

where $\theta \in (0, 1]$, $(\sigma_1, \sigma_2, \alpha_1, \alpha_2, \alpha_{12}, \beta_1, \beta_2, \beta_{12}) \in (0, +\infty)^8$ and $(\rho, \nu_1, \nu_2, \nu_{12}) \in \mathbb{R}^4$. The following result is a straightforward combination of Theorem 3.2 with the arguments in Theorem 1 of Porcu et al. [2018], so that the proof is omitted.

Theorem 3.3. Let SG_2 be the matrix-valued function defined by Equation (33). If either

A. PARSIMONIOUS \mathcal{SG}_2 : $\alpha_{12} = \frac{1}{2} \frac{\alpha_1 \alpha_2}{(\alpha_1 + \alpha_2)}, \ \beta_{12} = \frac{1}{2} (\beta_1 + \beta_2), \ \nu_{12} = \frac{1}{2} (\nu_1 + \nu_2),$ and

$$|\rho| \le \sqrt{\frac{A(\alpha_1, \beta_1, \nu_1)A(\alpha_2, \beta_2, \nu_2)}{A^2(\alpha_{12}, \beta_{12}, \nu_{12})}},$$

with $A(\cdot, \cdot, \cdot)$ defined as

$$A(\alpha,\beta,\nu) = \frac{2^{\nu-1} (\alpha\beta)^{\nu/2}}{\mathcal{K}_{\nu} \left(\sqrt{\frac{\beta}{\alpha}}\right)},$$

or

B. FULL \mathcal{SG}_2 : $\alpha_{12} < \frac{1}{2} \frac{\alpha_1 \alpha_2}{(\alpha_1 + \alpha_2)}, \ \beta_{12} > \frac{1}{2} (\beta_1 + \beta_2), \ \nu_{12} \neq \frac{1}{2} (\nu_1 + \nu_2), \ and$

$$|\rho| \le \sqrt{\frac{A(\alpha_1, \beta_1, \nu_1)A(\alpha_2, \beta_2, \nu_2)}{A^2(\alpha_{12}, \beta_{12}, \nu_{12})}} B\left(\frac{1}{\alpha_1} + \frac{1}{\alpha_2} - \frac{1}{2\alpha_{12}}, \beta_1 + \beta_2 - 2\beta_{12}, \nu_1 + \nu_2 - 2\nu_{12}\right)$$

with B defined as

$$B(\alpha,\beta,\nu) = \left(\frac{\left(\nu - \sqrt{\alpha\beta + \nu^2}\right)}{2\beta}\right)^{\nu} \exp\left(\frac{\alpha\beta + \nu(-\nu - \sqrt{\alpha\beta + \nu^2})}{-\nu + \sqrt{\alpha\beta + \nu^2}}\right),$$

then \mathcal{SG}_2 is positive semidefinite on any graph with Euclidean edges.

As a second application to a full multivariate setting $(p \ge 2)$, consider the mappings \mathcal{M}_p , \mathcal{C}_p and \mathcal{SG}_p defined on $[0, +\infty)^{p \times p}$ by

$$\mathcal{M}_{p}(\mathbf{D}) = \left[\sigma_{ij} \,\mathcal{M}_{\alpha_{ij},\nu_{ij}} \left(D_{ij}^{\theta}\right)\right]_{i,j=1}^{p},\tag{34}$$

$$\boldsymbol{\mathcal{C}}_{p}(\mathbf{D}) = \left[\sigma_{ij} \, \mathcal{C}_{\alpha_{ij},\nu} \left(D_{ij}^{\theta}\right)\right]_{i,j=1}^{p},\tag{35}$$

$$\boldsymbol{\mathcal{SG}}_{p}(\mathbf{D}) = \left[\sigma_{ij}\,\mathcal{SG}_{\alpha_{ij},\beta_{ij},\nu}\left(D_{ij}^{\theta}\right)\right]_{i,j=1}^{p},\tag{36}$$

with $[\sigma_{ij}]_{i,j=1}^p$, $[\nu_{ij}]_{i,j=1}^p$, $[\alpha_{ij}]_{i,j=1}^p$ and $[\beta_{ij}]_{i,j=1}^p$ being real symmetric matrices, ν a real number and $\theta \in (0, 1]$. Then the following statement holds.

Theorem 3.4. Let \mathcal{M}_p defined by Equation (34). If either

- A. $[\nu_{ij}]_{i,j=1}^p$ and $[\nu_{ij}\alpha_{ij}]_{i,j=1}^p$ are conditionally negative semidefinite and $\left[\frac{\sigma_{ij}\nu_{ij}^{\nu_{ij}}\exp(-\nu_{ij})}{\Gamma(\nu_{ij})}\right]_{i,j=1}^p$ is positive semidefinite or
- **B.** there exists $\beta > 0$ such that $[\nu_{ij}]_{i,j=1}^p$ and $[\alpha_{ij}^{-1} \beta \nu_{ij}]_{i,j=1}^p$ are conditionally negative semidefinite and $[\sigma_{ij}(\beta \alpha_{ij})^{-\nu_{ij}} \exp(-\nu_{ij})]_{i,j=1}^p$ is positive semidefinite,

then \mathcal{M}_p is positive semidefinite on any graph with Euclidean edges.

Theorem 3.5. Let C_p defined by Equation (35) with $\nu > 0$. If $[\beta_{ij}]_{i,j=1}^p$ is conditionally negative semidefinite and $[\sigma_{ij}\beta_{ij}^{\nu}]_{i,j=1}^p$ is positive semidefinite, then C_p is positive semidefinite on any graph with Euclidean edges.

Theorem 3.6. Let \mathcal{SG}_p defined by Equation (36). If $[\alpha_{ij}^{-1}]_{i,j=1}^p$ and $[\beta_{ij}]_{i,j=1}^p$ are conditionally negative semidefinite and $\left[\sigma_{ij}(\alpha_{ij}\beta_{ij})^{\nu/2}/\mathcal{K}_{\nu}\left(\sqrt{\frac{\alpha_{ij}}{\beta_{ij}}}\right)\right]_{i,j=1}^p$ is positive semidefinite, then \mathcal{SG}_p is positive semidefinite on any graph with Euclidean edges.

4 Euclidean Trees

Euclidean trees with a number m of leaves can be embedded on the m'dimensional Euclidean space with $m' := \lceil m/2 \rceil$ [Anderes et al., 2020] endowed with the ℓ_1 distance: for a positive integer m' and two points $x, y \in \mathbb{R}^{m'}$, the ℓ_1 or Manhattan-city block distance d_M can be defined as $d_M(x, y) = \sum_{k=1}^{m'} |x_k - y_k|$. According to this embedding, for every $m' \ge 2$, every matrixvalued covariance function depending on d_M can be used as an isotropic covariance function for a Euclidean tree The surprising fact is that, to the best of our knowledge, the literature has no models of multivariate covariance functions depending on such a metric. There is a fact that is even more surprising. Call Φ_m^p the class of continuous mappings $\mathbf{K} : [0, +\infty) \to \mathbb{R}^{p \times p}$ such that the composition $\mathbf{K}(d_M(\cdot, \cdot))$ is positive semidefinite on a Euclidean tree with m leaves. A characterisation of this class is missing in the literature and we provide it below.

Proposition 10. Let m, p be two positive integers. Let $m' := \lceil m/2 \rceil$. Let $\mathbf{K} : [0, +\infty) \to \mathbb{R}^{p \times p}$ be continuous with $\mathbf{K}(0) = \mathbf{1}_{p \times p}$. Then, \mathbf{K} belongs to the class Φ_m^p if and only if

$$\mathbf{K}(t) = \int_0^{+\infty} \omega_{m'}(rt) \,\mathbf{F}(\mathrm{d}r), \qquad t \ge 0, \tag{37}$$

where \mathbf{F} is a bounded symmetric matrix-valued measure such that $\mathbf{F}(dr)$ is positive semidefinite for all r > 0, and where

$$\omega_{m'}(t) = \frac{\Gamma(m'/2)}{\sqrt{\pi}\Gamma((m'-1)/2)} \int_{1}^{+\infty} \Omega_{m'}(\sqrt{v}t) v^{-m'/2} (v-1)^{(m'-3)/2} \mathrm{d}v, \quad (38)$$

with $\Omega_{m'}$ being the characteristic function of a random vector that is uniformly distributed over the unit sphere $\mathbb{S}^{m'-1}$ embedded in $\mathbb{R}^{m'}$.

Some comments are in order. For p = 1, the result has been proved by Cambanis et al. [1983]. The following strict inclusion relations hold:

$$\Phi_1^p \supset \Phi_2^p \supset \cdots \supset \bigcap_m \Phi_m^p =: \Phi_\infty^p.$$

A convergence argument from Schoenberg [1942] applies mutatis mutandis to assert (no proof needed) that $\mathbf{K} \in \Phi^p_{\infty}$ if and only if

$$\mathbf{K}(t) = \int_0^{+\infty} \exp(-rt) \mathbf{F}(\mathrm{d}r), \qquad t \ge 0, \tag{39}$$

with **F** as in Proposition 10. Closed-form expressions for the inner kernel $\omega_{m'}$ have been available thanks to Gneiting [1998]. In particular, we have

$$\omega_2(t) = -\frac{2}{\pi}\operatorname{si}(t) \quad \text{and} \quad \omega_3(t) = \frac{1}{2}\left(\frac{\sin t}{t} + \cos t + t\,\operatorname{si}(t)\right), \quad t \ge 0,$$

where si denotes the sine integral function.

There exists a wealth of multivariate covariance models that depend on the Euclidean distance, denoted $\|\cdot\|_m$, in \mathbb{R}^m . Call Ψ^p_m the class of continuous mappings $\mathbf{H} : [0, +\infty) \to \mathbb{R}^{p \times p}$ with $\mathbf{H}(0) = \mathbf{1}_{p \times p}$ and $\mathbf{H}(\|\cdot\|_m)$ being positive semidefinite in \mathbb{R}^m . The following result proves something extremely useful even for the scalar-case, that apparently was overlooked by Anderes et al. [2020].

Proposition 11. Let m, p be positive integers. Let $m' := \lceil m/2 \rceil$. Let **H** be a member of the class $\Psi_{2m'-1}^p$. Let the mapping **K** be defined through

$$\mathbf{K}(t) = I^{(m'-1)}\mathbf{H}(t), \qquad t \ge 0, \tag{40}$$

where $I^{(m'-1)}$ is the (m'-1) iterated application of the operator I, defined as

$$(If)(t) := \frac{\int_t^{+\infty} f(u) \mathrm{d}u}{\int_0^{+\infty} f(u) \mathrm{d}u}.$$

Then, **K** belongs to the class Φ_m^p .

Propositions 10 and 11 provide a recipe to build new members of the class Φ_m^p . We start by univariate models, as described in Table 2, where we use the shortcuts Ψ_m and Φ_m for Ψ_m^1 and Φ_m^1 , respectively.

We can now use Proposition 11 in concert with Table 2 and the following Lemma, for which a proof is not provided as it follows the same path as Theorem 3.2.

Lemma 4.1. Let m, p be positive integers and $\phi : [0, +\infty) \to \mathbb{R}$ be a member of the class Φ_m . Let \mathbf{F} be a matrix-valued measure as in Proposition 10. Then, the scale mixture

$$\mathbf{K}(t) = \int_0^{+\infty} \phi(rt) \, \mathbf{F}(\mathrm{d}r), \qquad t \ge 0, \tag{41}$$

provides an element of the class Φ_m^p .

Parameter restriction	Member of $\Psi_{2m'-1}$	Member of Φ_m
$\nu \ge m'$	$\psi_{\nu}(t) = (1-t)_{+}^{\nu}$	$\phi_{\nu}(t) = (1-t)_{+}^{\nu+m'-1}$
$\nu \geq m'+1$	$\psi_{\nu}(t) = (1-t)_{+}^{\nu} (1+\nu t)$	$\phi_{\nu}(t) = \frac{1}{m'} \left(1 - t \right)_{+}^{\nu + m' - 1} \left(m' + \nu t \right)$

Table 2: Elements of the class $\Psi_{2m'-1}$ and corresponding element of the class Φ_m after iterative application of the operator I as defined through Proposition 11. The first column provides the parametric restriction ensuring the corresponding element in the second column to be in $\Psi_{2m'-1}$.

Using the arguments in Theorem 1 of Daley et al. [2015] in concert with Lemma 4.1, one can prove that the multivariate model $\mathbf{K}(t) = [K_{ij}(t)]_{i,j=1}^{p}$ with

$$K_{ij}(t) = \sigma_i \sigma_{ij} \rho_{ij} \phi_{\nu_{ij}} \left(\frac{t}{b_{ij}} \right), \qquad t \ge 0,$$

is a member of the class Φ_m^p provided the constraints in Theorem 1 of Daley et al. [2015] hold. Here, ϕ_{ν} can be any of the entries in Table 2.

We finish this section by providing a direct construction that is based on the kernel ω_m as being defined through Equation (38).

Proposition 12. Let m, p be positive integers. Let $m' := \lceil m/2 \rceil$. Let $[\rho_{ij}]_{i,j=1}^p$, $[b_{ij}]_{i,j=1}^p$ and $[\nu_{ij}]_{i,j=1}^p$ be real symmetric matrices such that $b_{ij} > 0$ and $\nu_{ij} > 2m' - 1$ for i, j = 1, ..., p. For r > 0, define $\mathbf{A}(r)$ with entries

$$A_{ij}(r) = \frac{\rho_{ij}\Gamma(\frac{\nu_{ij}}{2})}{\Gamma(\frac{\nu_{ij}}{2} - m' + \frac{1}{2})\Gamma(m' - \frac{1}{2})} \left(1 - b_{ij}^2 r^2\right)_+^{\frac{\nu_{ij}}{2} - m' - \frac{1}{2}}, \quad i, j = 1, \dots, p.$$
(42)

Then, the mapping $\mathbf{K}: [0, +\infty) \to \mathbb{R}^{p \times p}$ with entries

$$K_{ij}(x) = \frac{\rho_{ij}}{b_{ij}^{2m'-1}} \,\omega_{\nu_{ij}}\left(\frac{x}{b_{ij}}\right), \qquad x \ge 0, \quad i, j = 1, \dots, p, \tag{43}$$

belongs to Φ_m^p provided the matrix $\mathbf{A}(r)$ as defined in (42) is positive semidefinite for any r > 0.

Sufficient conditions ensuring $\mathbf{A}(r)$ to be positive semidefinite for every fixed $r \geq 0$ are provided by Emery and Porcu [2023, Proposition 1]. The kernel (43) is especially flexible as it can be used to generate new families of multivariate kernels by using the scale mixture approaches highlighted above.



Figure 3: Marginal and cross covariance functions in terms of the resistance metric, over a tree with m = 4 leaves, measured from a reference location (black circle).



Figure 4: Realisation of a positively correlated bivariate random field on a tree with m = 4 leaves, with a covariance function of Askey type.

We conclude this section with an illustration to the bivariate setting (p = 2) for a Euclidean tree with m = 4 leaves. The entries of the covariance function are given by $K_{11}(d_R(u,v)) = (1 - d_R(u,v)/4)^3_+, K_{22}(d_R(u,v)) = (1 - d_R(u,v)/4)^3_+, \text{ and } K_{12}(d_R(u,v)) = 0.6 \times (1 - d_R(u,v)/2.5)^3_+, \text{ for all } u, v \in \mathcal{G}.$ Here, the collocated correlation coefficient between the two random field components is 0.6.

Figure 3 displays the decay of this covariance function in terms of the resistance metric measured from a reference location. It is evident that the first component of the random field has a larger correlation range, as reflected in the left panel: high (low) values of this variable tend to be surrounded by a wider extent of high (low) values. Figure 4 shows a realisation of a bivariate Gaussian random field over 750 sites on the tree. The Cholesky decomposition of the covariance matrix was used to perform this simulation.

5 Conclusions

This paper has shown the intricacies related to the construction to multivariate kernels over generalised frameworks that are represented through a broad class of metric graphs. One might argue that other classes of metric graphs are preferable in certain situations. However, to our knowledge, the only alternative has been proposed by Bolin et al. [2022] and coupled with the substantially different approach of stochastic partial differential equations. It is extremely challenging to attempt for a comparison between these approaches. One important advantage provided by Bolin et al. [2022] for scalar-valued random fields is that an example of a once mean-square differentiable random field is provided. Yet, this is limited to a specific example, while the approach by Anderes et al. [2020] — hence, our approach — allows to embrace a wealth of examples. Being the first contribution related to vector-valued random fields on metric graphs, this paper does not have a competitor to compare with.

A next intuitive step to this research is represented by vector-valued space-time random fields. The machinery provided in this paper allows for a fairly general building block to start with. Space-time models might also be the building blocks to nonstationary models for networks.

The impact of this research is apparent. The works by Baddeley et al. [2017] and Moradi and Mateu [2020] are a clear indication of the importance of this work to modelling point processes over networks. There is a fervent activity related to vector-valued processes in the machine learning community, with the reader referred to Borovitskiy et al. [2022] for details.

Acknowledgements

The authors are grateful to Valeria Simoncini for her support regarding the solutions of matrix quadratic equations. X. Emery acknowledges the support of the National Agency for Research and Development of Chile, through grants ANID Fondecyt 1210050 and ANID PIA AFB230001. Emilio Porcu is grateful to Horst Simon for enlightening discussion about multivariate isometric embeddings.

References

- Ethan Anderes, Jesper Møller, and Jakob G Rasmussen. Isotropic covariance functions on graphs and their edges. *The Annals of Statistics*, 48(4):2478– 2503, 2020.
- M.M. Moradi and J. Mateu. First-and second-order characteristics of spatiotemporal point processes on linear networks. *Journal of Computational* and Graphical Statistics, 29(3):432–443, 2020.
- Adrian Baddeley, Gopalan Nair, Suman Rakshit, Greg McSwiggan, and Tilman M. Davies. Analysing point patterns on networks – a review. *Spatial Statistics*, 42:100435, 2021. ISSN 2211-6753.
- Suman Rakshit, Gopalan Nair, and Adrian Baddeley. Second-order analysis of point patterns on a network using any distance metric. *Spatial Statistics*, 22:129–154, 2017.
- A. Ortega, P. Frossard, J. Kovacevic, J.M. Moura, and P. Vandergheynst. Graph signal processing: Overview, challenges, and applications. *Proceed*ings of the IEEE, 106(5):808–828, 2018.
- Ines Chami, Sami Abu-El-Haija, Bryan Perozzi, Christopher Ré, and Kevin Murphy. Machine learning on graphs: A model and comprehensive taxonomy. The Journal of Machine Learning Research, 23(1):3840–3903, 2022.
- D. Bolin and F. Lindgren. Spatial models generated by nested stochastic partial differential equations, with an application to global ozone mapping. *The Annals of Applied Statistics*, 5(1):523–550, 2011.
- David Bolin, Alexandre B Simas, and Jonas Wallin. Gaussian Whittle-Matérn fields on metric graphs. arXiv preprint arXiv:2205.06163, 2022.
- Emilio Porcu, Philip A White, and Marc G Genton. Stationary nonseparable space-time covariance functions on networks. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 85(5):1417–1440, 2023.
- Tobia Filosi, Claudio Agostinelli, and Emilio Porcu. Temporally-evolving generalised networks and their reproducing kernels. *arXiv preprint arXiv:2309.15855*, 2023.

- Jun Tang and Dale Zimmerman. Space-time covariance models on networks with an application on streams. arXiv:2009.14745, 2020.
- Mohammad Abu Alsheikh, Shaowei Lin, Dusit Niyato, and Hwee-Pink Tan. Machine learning in wireless sensor networks: Algorithms, strategies, and applications. *IEEE Communications Surveys & Tutorials*, 16(4):1996– 2018, 2014.
- William L Hamilton, Rex Ying, and Jure Leskovec. Representation learning on graphs: Methods and applications. arXiv preprint arXiv:1709.05584, 2017.
- Viacheslav Borovitskiy, Mohammad Reza Karimi, Vignesh Ram Somnath, and Andreas Krause. Isotropic gaussian processes on finite spaces of graphs. arXiv preprint arXiv:2211.01689, 2022.
- Noel Cressie, Jesse Frey, Bronwyn Harch, and Mick Smith. Spatial prediction on a river network. *Journal of Agricultural, Biological, and Environmental Statistics*, 11(2):127, 2006.
- Jay M Ver Hoef, Erin Peterson, and David Theobald. Spatial statistical models that use flow and stream distance. *Environmental and Ecological Statistics*, 13(4):449–464, 2006.
- Erin E Peterson, Jay M Ver Hoef, Dan J Isaak, Jeffrey A Falke, Marie-Josée Fortin, Chris E Jordan, Kristina McNyset, Pascal Monestiez, Aaron S Ruesch, Aritra Sengupta, et al. Modelling dendritic ecological networks in space: An integrated network perspective. *Ecology Letters*, 16(5):707–719, 2013.
- Erin E Peterson, David M Theobald, and Jay M ver Hoef. Geostatistical modelling on stream networks: Developing valid covariance matrices based on hydrologic distance and stream flow. *Freshwater Biology*, 52(2):267–279, 2007.
- Maxime Montembeault, Sven Joubert, Julien Doyon, Julie Carrier, Jean-François Gagnon, Oury Monchi, Ovidiu Lungu, Sylvie Belleville, and Simona Maria Brambati. The impact of aging on gray matter structural covariance networks. *Neuroimage*, 63(2):754–759, 2012.

- Shuai Xiao, Junchi Yan, Xiaokang Yang, Hongyuan Zha, and Stephen Chu. Modeling the intensity function of point process via recurrent neural networks. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 31. Association for the Advancement of Artificial Intelligence, 2017.
- Patrick O Perry and Patrick J Wolfe. Point process modelling for directed interaction networks. *Journal of the Royal Statistical Society: Series B*, 75 (5):821–849, 2013.
- Na Deng, Wuyang Zhou, and Martin Haenggi. The ginibre point process as a model for wireless networks with repulsion. *IEEE Transactions on Wireless Communications*, 14(1):107–121, 2014.
- Adrian Baddeley, Gopalan Nair, Suman Rakshit, and Greg McSwiggan. Stationary point processes are uncommon on linear networks. *Stat*, 6(1):68–78, 2017.
- Victor P Zastavnyi. On positive definiteness of some functions. Journal of Multivariate Analysis, 73(1):55–81, 2000. ISSN 0047-259X.
- I. J. Schoenberg. Positive definite functions on spheres. Duke Mathematical Journal, 9:96–108, 1942.
- Karel Devriendt. Effective resistance is more than distance: Laplacians, simplices and the Schur complement. *Linear Algebra and its Applications*, 639:24–49, 2022. ISSN 00243795. URL http://arxiv.org/abs/2010. 04521.
- D.J. Klein and M. Randic. Resistance distance. *Journal of Mathematical Chemistry*, 12:81–95, 1993.
- Jean-Paul Chilès and Pierre Delfiner. *Geostatistics: Modeling Spatial Uncertainty*, volume 713. John Wiley & Sons, 2012.
- Christopher Dörr and Martin Schlather. Characterization theorems for pseudo cross-variograms. *Journal of Applied Probability*, 60(4):1219–1231, 2023.
- V. P. Zastavnyi. Analog of Schoenberg's theorem for a-conditionally negative definite matrix-valued kernels. *Mathematical Notes*, 114(1):66–76, 2023. ISSN 0001-4346, 1573-8876.

- Emilio Porcu, Moreno Bevilacqua, and Amanda S Hering. The shkarofskygneiting class of covariance models for bivariate gaussian random fields. *Stat*, 7(1):e207, 2018.
- Stamatis Cambanis, Robert Keener, and Gordon Simons. On-symmetric multivariate distributions. Journal of Multivariate Analysis, 13(2):213– 233, 1983.
- Tilmann Gneiting. On α-symmetric multivariate characteristic functions. Journal of Multivariate Analysis, 64(2):131–147, 1998.
- Daryl J Daley, Emilio Porcu, and Moreno Bevilacqua. Classes of compactly supported covariance functions for multivariate random fields. *Stochastic Environmental Research and Risk Assessment*, 29:1249–1263, 2015.
- Xavier Emery and Emilio Porcu. The schoenberg kernel and more flexible multivariate covariance models in euclidean spaces. Computational and Applied Mathematics, 42(4):148, 2023.
- F. Zhang. The schur complement and its applications. In *The Schur complement and its applications*, volume 4 of *Numerical Methods and Algorithms*. Springer-Verlag, 2005. ISBN 978-0-387-24271-2. URL http://link.springer.com/10.1007/b105056.
- Carl D. Meyer. Generalized inversion of modified matrices. SIAM Journal on Applied Mathematics, 24(3):315–323, 1973.
- Xavier Emery, Emilio Porcu, and Philip White. New validity conditions for the multivariate Matérn coregionalization model, with an application to exploration geochemistry. *Mathematical Geosciences*, 54(6):1043–1068, 2022.
- I. S. Gradshteyn and I. M. Ryzhik. *Tables of Integrals, Series, and Products.* Academic Press, Amsterdam, seventh edition, 2007.
- Robert Reams. Hadamard inverses, square roots and products of almost semidefinite matrices. *Linear Algebra and its Applications*, 288:35–43, 1999.
- S. Bochner. Harmonic Analysis and the Theory of Probability. California Monographs in Mathematical Sciences. University of California Press, 1955. URL https://books.google.cl/books?id=7_1QAAAAMAAJ.

Harald Cramer. On the theory of stationary random processes. Annals of Mathematics, 41(1):215-230, 1940. ISSN 0003486X. URL http://www.jstor.org/stable/1968827.

A Technical Results and Proofs

This part of the Appendix contains proofs and technical results. To help the reader, Figure 1 provides the road-map of the main results of this manuscript.

Lemma A.1. Let $\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}$ be $n \times n$ matrices and let $p \in \{1, 2, \ldots\}$. Then

$$\mathcal{M}_p\begin{pmatrix}\mathbf{A}\\\mathbf{B}\end{pmatrix}\mathcal{M}_p\begin{pmatrix}\mathbf{C}\\\mathbf{D}\end{pmatrix} = \mathcal{M}_p\begin{pmatrix}\mathbf{A}\mathbf{C} + (p-1)\mathbf{B}\mathbf{D}\\\mathbf{A}\mathbf{D} + \mathbf{B}\mathbf{C} + (p-2)\mathbf{B}\mathbf{D}\end{pmatrix}.$$
 (44)

Proof of Lemma A.1. By the standard properties of the Kronecker product,

$$\begin{split} (\mathbf{I}_p \otimes \mathbf{A} + (\mathbf{1}_{p \times p} - \mathbf{I}_p) \otimes \mathbf{B}) & (\mathbf{I}_p \otimes \mathbf{C} + (\mathbf{1}_{p \times p} - \mathbf{I}_p) \otimes \mathbf{D}) = \\ &= (\mathbf{I}_p \otimes \mathbf{A}) (\mathbf{I}_p \otimes \mathbf{C}) + (\mathbf{I}_p \otimes \mathbf{A}) ((\mathbf{1}_{p \times p} - \mathbf{I}_p) \otimes \mathbf{D}) \\ &+ ((\mathbf{1}_{p \times p} - \mathbf{I}_p) \otimes \mathbf{B}) (\mathbf{I}_p \otimes \mathbf{C}) + ((\mathbf{1}_{p \times p} - \mathbf{I}_p) \otimes \mathbf{B}) ((\mathbf{1}_{p \times p} - \mathbf{I}_p) \otimes \mathbf{D}) \\ &= (\mathbf{I}_p \mathbf{I}_p) \otimes (\mathbf{A}\mathbf{C}) + (\mathbf{I}_p (\mathbf{1}_{p \times p} - \mathbf{I}_p)) \otimes (\mathbf{A}\mathbf{D}) \\ &+ ((\mathbf{1}_{p \times p} - \mathbf{I}_p)\mathbf{I}_p) \otimes (\mathbf{B}\mathbf{C}) + ((\mathbf{1}_{p \times p} - \mathbf{I}_p)(\mathbf{1}_{p \times p} - \mathbf{I}_p)) \otimes (\mathbf{B}\mathbf{D}) \\ &= \mathbf{I}_p \otimes (\mathbf{A}\mathbf{C}) + (\mathbf{1}_{p \times p} - \mathbf{I}_p) \otimes (\mathbf{A}\mathbf{D} + \mathbf{B}\mathbf{C}) + (p\mathbf{1}_{p \times p} - 2 \cdot \mathbf{1}_{p \times p} + \mathbf{I}_p) \otimes (\mathbf{B}\mathbf{D}) \\ &= \mathbf{I}_p \otimes (\mathbf{A}\mathbf{C}) + (\mathbf{1}_{p \times p} - \mathbf{I}_p) \otimes (\mathbf{A}\mathbf{D} + \mathbf{B}\mathbf{C}) \\ &+ ((p-2)(\mathbf{1}_{p \times p} - \mathbf{I}_p) + (p-1)\mathbf{I}_p) \otimes (\mathbf{B}\mathbf{D}) \\ &= \mathbf{I}_p \otimes (\mathbf{A}\mathbf{C} + (p-1)\mathbf{B}\mathbf{D}) + (\mathbf{1}_{p \times p} - \mathbf{I}_p) \otimes (\mathbf{A}\mathbf{D} + \mathbf{B}\mathbf{C} + (p-2)\mathbf{B}\mathbf{D}). \end{split}$$

Proof of Proposition 1. Throughout this proof, we assume that $\mathbf{L} \in \mathbb{R}^{n \times n}$, $\mathbf{A} \in \mathbb{R}^{n_1 \times n_1}$ and $\mathbf{C} \in \mathbb{R}^{n_2 \times n_2}$, with $n_1, n_2 > 0$ and $n = n_1 + n_2$.

- 1. Assume $\mathbf{B} = 0$. Then $n 1 = \operatorname{rank}(\mathbf{L}) = \operatorname{rank}(\mathbf{A}) + \operatorname{rank}(\mathbf{C})$. But since $\mathbf{L}\mathbf{1}_n = \mathbf{0}$, it follows that $\mathbf{A}\mathbf{1}_{n_1} = \mathbf{0}$ and $\mathbf{C}\mathbf{1}_{n_2} = \mathbf{0}$. This implies that $\operatorname{rank}(\mathbf{A}) \leq n_1 1$ and $\operatorname{rank}(\mathbf{C}) \leq n_2 1$. Contradiction.
- 2. We will show only that **C** strictly positive definite, as the proof for the other case relies on the same argument. Since **C** is a principal submatrix of **L**, it is positive semidefinite. As a consequence, it is sufficient to show that it is invertible. Assume, by contradiction, that **C** is singular. This is equivalent to assume that $\exists y \neq 0$ such that

 $\mathbf{C}\boldsymbol{y} = \mathbf{0}$. Since \mathbf{L} is positive semidefinite we have, for each $\boldsymbol{x} \in \mathbb{R}^{n_1}$ and for each $\gamma \in \mathbb{R}$:

$$0 \leq \begin{bmatrix} \boldsymbol{x} \\ \gamma \boldsymbol{y} \end{bmatrix}^{\top} \begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{B}^{\top} & \mathbf{C} \end{bmatrix} \begin{bmatrix} \boldsymbol{x} \\ \gamma \boldsymbol{y} \end{bmatrix}$$

= $\boldsymbol{x}^{\top} \mathbf{A} \boldsymbol{x} + \boldsymbol{x}^{\top} \mathbf{B} (\gamma \boldsymbol{y}) + (\gamma \boldsymbol{y})^{\top} \mathbf{B}^{\top} \boldsymbol{x} + (\gamma \boldsymbol{y})^{\top} \mathbf{C} (\gamma \boldsymbol{y}) = \boldsymbol{x}^{\top} \mathbf{A} \boldsymbol{x} + 2\gamma \boldsymbol{x}^{\top} \mathbf{B} \boldsymbol{y}.$

If $\mathbf{x}^{\top} \mathbf{B} \mathbf{y} \neq 0$, we can find a suitable γ that contradicts the previous inequality. Hence $\mathbf{x}^{\top} \mathbf{B} \mathbf{y} = 0$ and therefore, being \mathbf{x} arbitrary, $\mathbf{B} \mathbf{y} = \mathbf{0}$. Consider now the vector $\mathbf{z} := (0, ..., 0, \mathbf{y}^{\top})^{\top}$: we have $\mathbf{L} \mathbf{z} = \mathbf{0}$, but clearly \mathbf{z} does not lie in the null space of \mathbf{L} , which is $\{\lambda \mathbf{1}_n : \lambda \in \mathbb{R}\}$. Contradiction.

3. Clearly \mathbf{L}/\mathbf{C} is symmetric. Furthermore, the Guttman rank additivity formula [Zhang, 2005, Equation 0.9.2] states that rank(\mathbf{L}) = rank(\mathbf{C})+rank(\mathbf{L}/\mathbf{C}). Here we have rank(\mathbf{L}) = n-1 and rank(\mathbf{C}) = n_2 , as a consequence rank(\mathbf{L}/\mathbf{C}) = $n_1 - 1$. Furthermore, from Zhang [2005, Theorem 1.12], being \mathbf{C} strictly positive definite and L positive semidefinite, we have that \mathbf{L}/\mathbf{C} is positive semidefinite as well. The last thing that is left to be shown is that the null eigenvector of \mathbf{L}/\mathbf{C} is $\mathbf{1}_{n_1}$. Since $\mathbf{1}_n$ is the null eigenvector of \mathbf{L} , we get

$$\begin{bmatrix} \mathbf{A} & \mathbf{B} \end{bmatrix} \mathbf{1}_n = \mathbf{0} \qquad \text{and} \qquad \begin{bmatrix} \mathbf{B}^\top & \mathbf{C} \end{bmatrix} \mathbf{1}_n = \mathbf{0},$$

videlicet $\mathbf{A1}_{n_1} + \mathbf{B1}_{n_2} = \mathbf{0}$ and $\mathbf{B}^\top \mathbf{1}_{n_1} + \mathbf{C1}_{n_2} = \mathbf{0}$. Using these relations, the following chain of equivalences concludes the proof:

$$(\mathbf{A} - \mathbf{B}\mathbf{C}^{-1}\mathbf{B}^{\top})\mathbf{1}_{n_1} = \mathbf{0} \iff \mathbf{A}\mathbf{1}_{n_1} = \mathbf{B}\mathbf{C}^{-1}\mathbf{B}^{\top}\mathbf{1}_{n_1}$$

 $\iff -\mathbf{B}\mathbf{1}_{n_2} = \mathbf{B}\mathbf{C}^{-1}(-\mathbf{C}\mathbf{1}_{n_2}) \iff -\mathbf{B}\mathbf{1}_{n_2} = -\mathbf{B}\mathbf{1}_{n_2}.$

4. Consider the eigendecomposition of L:

$$\mathbf{L} = \mathbf{W} \begin{bmatrix} \mathbf{\Delta} & \mathbf{0} \\ \mathbf{0} & 0 \end{bmatrix} \mathbf{W}^{\top}, \tag{45}$$

with Δ a $(n-1) \times (n-1)$ diagonal matrix with positive diagonal entries. Since **W** is an orthogonal matrix, we have

$$\mathbf{L}^{-} = \left(\mathbf{W} \begin{bmatrix} \boldsymbol{\Delta} & \mathbf{0} \\ \mathbf{0} & 0 \end{bmatrix} \mathbf{W}^{\top} \right)^{-} = (\mathbf{W}^{\top})^{-} \begin{bmatrix} \boldsymbol{\Delta} & \mathbf{0} \\ \mathbf{0} & 0 \end{bmatrix}^{-} \mathbf{W}^{-} = \mathbf{W} \begin{bmatrix} \boldsymbol{\Delta}^{-1} & \mathbf{0} \\ \mathbf{0} & 0 \end{bmatrix} \mathbf{W}^{\top}$$
(46)

Therefore \mathbf{L}^- is symmetric, is positive semidefinite and with exactly one null eigenvalue, corresponding to the same null-space eigenvector of \mathbf{L} .

Lemma A.2. Let \mathbf{L} be an $n \times n$ quasi-Laplacian matrix and $\Sigma = \mathbf{L}^-$. Let $\Sigma =: \mathbf{W} \mathbf{\Lambda} \mathbf{W}^\top$ be the eigendecomposition of Σ , with $\mathbf{\Lambda} = \operatorname{diag} \left(\begin{bmatrix} \lambda_1, \ldots, \lambda_n \end{bmatrix} \right)$ and $\lambda_1 \geq \cdots \geq \lambda_{n-1} > \lambda_n = 0$. Let \mathbf{Q} be defined as in Subsection 3.1. Then, the eigenvalues $\lambda_n(\mathbf{Q}) < \lambda_1(\mathbf{Q}) \leq \cdots \leq \lambda_{n-1}(\mathbf{Q})$ of \mathbf{Q} are:

$$\lambda_i(\mathbf{Q}) = \begin{cases} \frac{1}{2} \left(\lambda_i^{-1} + \sqrt{\lambda_i^{-2} - 2\alpha(p-2)\lambda_i^{-1} + \alpha^2 p^2} + \alpha(p-2) \right) & i \neq n \\ \alpha(p-1) & i = n. \end{cases}$$

In particular the smallest eigenvalue of \mathbf{Q} , $\lambda_n(\mathbf{Q})$, is strictly positive, thus \mathbf{Q} is strictly positive definite. In addition, $\mathbf{Q}\mathbf{1}_n = \alpha(p-1)\mathbf{1}_n$.

Proof of Lemma A.2. Recall the eigendecomposition of matrix \mathbf{Q} given in Equation (19): it is clear that the eigenvalues of \mathbf{Q} are the elements on the diagonal of the matrix

$$\frac{1}{2} \left(\mathbf{\Lambda}^{-} + \sqrt{(\mathbf{\Lambda}^{-})^2 - 2\alpha(p-2)\mathbf{\Lambda}^{-} + \alpha^2 p^2 \mathbf{I}_n} + \alpha(p-2)\mathbf{I}_n \right) + \frac{1}{2} \left(\mathbf{\Lambda}^{-} + \sqrt{(\mathbf{\Lambda}^{-})^2 - 2\alpha(p-2)\mathbf{\Lambda}^{-} + \alpha^2 p^2 \mathbf{I}_n} + \alpha(p-2)\mathbf{I}_n \right) + \frac{1}{2} \left(\mathbf{\Lambda}^{-} + \sqrt{(\mathbf{\Lambda}^{-})^2 - 2\alpha(p-2)\mathbf{\Lambda}^{-} + \alpha^2 p^2 \mathbf{I}_n} + \alpha(p-2)\mathbf{I}_n \right) + \frac{1}{2} \left(\mathbf{\Lambda}^{-} + \sqrt{(\mathbf{\Lambda}^{-})^2 - 2\alpha(p-2)\mathbf{\Lambda}^{-} + \alpha^2 p^2 \mathbf{I}_n} + \alpha(p-2)\mathbf{I}_n \right) + \frac{1}{2} \left(\mathbf{\Lambda}^{-} + \sqrt{(\mathbf{\Lambda}^{-})^2 - 2\alpha(p-2)\mathbf{\Lambda}^{-} + \alpha^2 p^2 \mathbf{I}_n} + \alpha(p-2)\mathbf{I}_n \right) + \frac{1}{2} \left(\mathbf{\Lambda}^{-} + \sqrt{(\mathbf{\Lambda}^{-})^2 - 2\alpha(p-2)\mathbf{\Lambda}^{-} + \alpha^2 p^2 \mathbf{I}_n} + \alpha(p-2)\mathbf{I}_n \right) + \frac{1}{2} \left(\mathbf{\Lambda}^{-} + \sqrt{(\mathbf{\Lambda}^{-})^2 - 2\alpha(p-2)\mathbf{\Lambda}^{-} + \alpha^2 p^2 \mathbf{I}_n} + \alpha(p-2)\mathbf{I}_n \right) + \frac{1}{2} \left(\mathbf{\Lambda}^{-} + \mathbf{\Lambda}^{-$$

which are $\{f(\lambda^{-}) : \lambda \in \{\lambda_1, \ldots, \lambda_n\}\}$, where $\lambda_1, \ldots, \lambda_n$ are the diagonal entries of Λ (positive, except for λ_n that is null) and

$$f(\lambda^{-}) := \frac{1}{2} \left(\lambda^{-} + \sqrt{(\lambda^{-})^{2} - 2\alpha(p-2)\lambda^{-} + \alpha^{2}p^{2}} + \alpha(p-2) \right).$$

Here, we recall that, for a scalar $\lambda \in \mathbb{R}$, $\lambda^- := \lambda^{-1}$ if $\lambda \neq 0$ and $\lambda^- := 0$ if $\lambda = 0$. It is straightforward to show that $\lambda^- \mapsto f(\lambda^-)$ is a strictly increasing function on $\lambda^- \geq 0$. As a consequence, since $\lambda \mapsto \lambda^-$ is strictly decreasing on $\lambda > 0$, $\lambda \mapsto f(\lambda^-)$ is strictly decreasing on $\lambda > 0$. Therefore, $\lambda_1 \geq \cdots \geq \lambda_{n-1}$ are mapped in $\lambda_1(\mathbf{Q}) = f(\lambda_1^{-1}) \leq \cdots \leq \lambda_{n-1}(\mathbf{Q}) = f(\lambda_{n-1}^{-1})$. For the case $\lambda_n = 0$, we have $\lambda_n(\mathbf{Q}) = f(\lambda_n^-) = f(0) = \frac{1}{2}(\alpha p + \alpha(p-2)) = \alpha(p-1)$. In addition, since $\lambda_n = 0 < \lambda_1(\mathbf{Q})$, we have $\alpha(p-1) = \lambda_n(\mathbf{Q}) < \lambda_1(\mathbf{Q})$.

Finally, to show that $\mathbf{Q}\mathbf{1}_n = \alpha(p-1)\mathbf{1}_n$, it is sufficient to notice from the eigendecomposition of \mathbf{Q} given in Equation (19) that $\mathbf{1}_n$, being (proportional to) the last column of \mathbf{W} , is an eigenvector of \mathbf{Q} . And by the above result it is clear that the corresponding eigenvalue is $\lambda_n(\mathbf{Q}) = \alpha(p-1)$. This concludes the proof.

Lemma A.3. Let **L** be an $n \times n$ quasi-Laplacian matrix, $\Sigma = \mathbf{L}^-$ and $\alpha > 0$. Furthermore, let **Q** and **X** as defined in Subsection 3.1. Then the following holds:

1. Σ , L, Q and X commute;

2.
$$\Sigma \mathbf{L} = \mathbf{L} \Sigma = \mathbf{I}_n - \frac{1}{n} \mathbf{1}_{n \times n};$$

3. $\mathbf{Q}^2 - \alpha^2 (p-1)\mathbf{I}_n - \mathbf{Q}\mathbf{L} - \alpha(p-2)\mathbf{Q} + \alpha(p-2)\mathbf{L} = \mathbf{0};$

4.
$$\Sigma \mathbf{Q}^2 - \alpha^2 (p-1)\Sigma - \mathbf{Q} - \alpha (p-2)\Sigma \mathbf{Q} + \alpha (p-2)\mathbf{I}_n + \frac{\alpha}{n} \mathbf{1}_{n \times n} = \mathbf{0}.$$

Proof of Lemma A.3. 1. It is sufficient to notice that all the matrices share the same eigenvector matrix \mathbf{W} , *i.e.* for any $\mathbf{A}, \mathbf{B} \in \{\Sigma, \mathbf{L}, \mathbf{Q}, \mathbf{X}\}$, there are two diagonal matrices $\Lambda_{\mathbf{A}}, \Lambda_{\mathbf{B}}$ such that $\mathbf{A} = \mathbf{W} \Lambda_{\mathbf{A}} \mathbf{W}^{\top}$ and $\mathbf{B} = \mathbf{W} \Lambda_{\mathbf{B}} \mathbf{W}^{\top}$. Therefore

$$\begin{split} \mathbf{A}\mathbf{B} &= \mathbf{W}\boldsymbol{\Lambda}_{\mathbf{A}}\mathbf{W}^{\top}\mathbf{W}\boldsymbol{\Lambda}_{\mathbf{B}}\mathbf{W}^{\top} = \mathbf{W}\boldsymbol{\Lambda}_{\mathbf{A}}\boldsymbol{\Lambda}_{\mathbf{B}}\mathbf{W}^{\top} = \mathbf{W}\boldsymbol{\Lambda}_{\mathbf{B}}\boldsymbol{\Lambda}_{\mathbf{A}}\mathbf{W}^{\top} \\ &= \mathbf{W}\boldsymbol{\Lambda}_{\mathbf{B}}\mathbf{W}^{\top}\mathbf{W}\boldsymbol{\Lambda}_{\mathbf{A}}\mathbf{W}^{\top} = \mathbf{B}\mathbf{A}. \end{split}$$

2. By exploiting the diagonalisations $\Sigma = \mathbf{W} \mathbf{\Lambda} \mathbf{W}^{\top}$ (Lemma A.2) and $\mathbf{L} = \mathbf{W} \mathbf{\Lambda}^{-} \mathbf{W}^{\top}$ (Eqs. (45) and (46)), we get:

$$\mathbf{\Sigma}\mathbf{L} = \mathbf{W}\mathbf{\Lambda}\mathbf{\Lambda}^{-}\mathbf{W}^{ op} = \mathbf{W}(\mathbf{I}_n - \mathbf{J}_n)\mathbf{W}^{ op} = \mathbf{I}_n - \mathbf{W}\mathbf{J}_n\mathbf{W}^{ op}$$

Given that \mathbf{L} is a quasi-Laplacian matrix, its eigenvector matrix \mathbf{W} can be written as

$$\mathbf{W} = \begin{bmatrix} \mathbf{U} & c\mathbf{1}_n \end{bmatrix},$$

where $\mathbf{U} \in \mathbb{R}^{n \times (n-1)}$ and $nc^2 = 1$. Therefore

$$\mathbf{W}\mathbf{J}_{n}\mathbf{W}^{\top} = \begin{bmatrix} \mathbf{U} & c\mathbf{1}_{n} \end{bmatrix} \begin{bmatrix} \mathbf{0}_{(n-1)\times(n-1)} & \mathbf{0}_{(n-1)\times1} \\ \mathbf{0}_{1\times(n-1)} & 1 \end{bmatrix} \begin{bmatrix} \mathbf{U}^{\top} \\ c\mathbf{1}_{n}^{\top} \end{bmatrix} = \frac{1}{n}\mathbf{1}_{n}\mathbf{1}_{n}^{\top},$$
(47)

and this concludes the proof.

3. From the definition (18) of \mathbf{Q} , one has:

$$2\mathbf{Q} - \mathbf{L} - \alpha(p-2)\mathbf{I}_n = \sqrt{\mathbf{L}^2 - 2\alpha(p-2)\mathbf{L} + \alpha^2 p^2 \mathbf{I}_n}.$$

Now, taking the square of both sides (recall that **Q** and **L** commute):

$$4\mathbf{Q}^{2} + \mathbf{L}^{2} + \alpha^{2}(p-2)^{2}\mathbf{I}_{n} - 4\mathbf{Q}\mathbf{L} - 4\alpha(p-2)\mathbf{Q} + 2\alpha(p-2)\mathbf{L}$$
$$= \mathbf{L}^{2} - 2\alpha(p-2)\mathbf{L} + \alpha^{2}p^{2}\mathbf{I}_{n}.$$

By grouping the common terms and dividing everything by 4 we get the desired result.

4. To show the second assertion, it is sufficient to multiply the first equation by $\Sigma = L^{-}$:

$$\begin{split} \mathbf{\Sigma}\mathbf{Q}^2 &- \alpha^2(p-1)\mathbf{\Sigma} - \mathbf{Q}(\mathbf{\Sigma}\mathbf{L}) - \alpha(p-2)\mathbf{\Sigma}\mathbf{Q} + \alpha(p-2)(\mathbf{\Sigma}\mathbf{L}) = \mathbf{0} \\ \mathbf{\Sigma}\mathbf{Q}^2 &- \alpha^2(p-1)\mathbf{\Sigma} - \mathbf{Q} + \frac{1}{n}\lambda_n(\mathbf{Q})\mathbf{1}_{n\times n} - \alpha(p-2)\mathbf{\Sigma}\mathbf{Q} + \alpha(p-2)\mathbf{I}_n \\ &- \frac{\alpha(p-2)}{n}\mathbf{1}_{n\times n} = \mathbf{0}, \end{split}$$

where we have used $\Sigma \mathbf{L} = \mathbf{I}_n - \frac{1}{n} \mathbf{1}_{n \times n}$ and $\mathbf{Q} \mathbf{1}_n = \lambda_n(\mathbf{Q}) \mathbf{1}_n = \alpha(p-1) \mathbf{1}_n$ (Lemma A.2).

Lemma A.4. Let $\mathbf{X}, \mathbf{Y} \in \mathbb{R}^{n \times n}$ be symmetric matrices. For $i \in \{1, \ldots, n\}$, let $\lambda_i(\mathbf{X})$ and $\lambda_i(\mathbf{Y})$ denote the *i*th eigenvalues of \mathbf{X} and \mathbf{Y} , respectively. Let $c \in \mathbb{R}$ such that $\lambda_i(\mathbf{X}) \geq c$ and $\lambda_i(\mathbf{Y}) \geq -c$ for all $i \in \{1, \ldots, n\}$. Then $\mathbf{X} + \mathbf{Y}$ is positive semidefinite.

Proof of Lemma A.4. Let $\mathbf{X} = \mathbf{W}_1 \boldsymbol{\Delta}_1 \mathbf{W}_1^{\top}$ and $\mathbf{Y} = \mathbf{W}_2 \boldsymbol{\Delta}_2 \mathbf{W}_2^{\top}$ be the eigendecompositions of \mathbf{X} and \mathbf{Y} and let $\boldsymbol{v} \in \mathbb{R}^n$. In the following lines, we have defined $\boldsymbol{a} := \mathbf{W}_1^{\top} \boldsymbol{v}$ and $\boldsymbol{b} := \mathbf{W}_2^{\top} \boldsymbol{v}$.

$$\boldsymbol{v}^{\top}(\mathbf{X} + \mathbf{Y})\boldsymbol{v} = \boldsymbol{v}^{\top}\mathbf{W}_{1}\Delta_{1}\mathbf{W}_{1}^{\top}\boldsymbol{v} + \boldsymbol{v}\mathbf{W}_{2}\boldsymbol{\Delta}_{2}\mathbf{W}_{2}^{\top}\boldsymbol{v} = \boldsymbol{a}^{\top}\boldsymbol{\Delta}_{1}\boldsymbol{a} + \boldsymbol{b}^{\top}\boldsymbol{\Delta}_{2}\boldsymbol{b}$$
$$= \sum_{i=1}^{n}\lambda_{i}(\mathbf{X})a_{i}^{2} + \sum_{i=1}^{n}\lambda_{i}(\mathbf{Y})b_{i}^{2} \ge c\sum_{i=1}^{n}a_{i}^{2} - c\sum_{i=1}^{n}b_{i}^{2}$$
$$= c(\boldsymbol{a}^{\top}\boldsymbol{a} - \boldsymbol{b}^{\top}\boldsymbol{b}) = c(\boldsymbol{v}^{\top}\boldsymbol{v} - \boldsymbol{v}^{\top}\boldsymbol{v}) = 0.$$

Videlicet, $\mathbf{X} + \mathbf{Y}$ is positive semidefinite.

Lemma A.5. Let \mathbf{L} an $n \times n$ Laplacian matrix, let $\Sigma := \mathbf{L}^-$ its Moore-Penrose inverse and let $\alpha > 0$. Let Θ , \mathbf{Q} , \mathbf{X} , k_1 , k_2 , $\tilde{\Sigma}$ and $\tilde{\mathbf{X}}$ as described in Subsection 3.1. Finally let:

$$\tilde{\mathbf{K}} := \mathcal{M}_p \begin{pmatrix} \tilde{\boldsymbol{\Sigma}} \\ \tilde{\mathbf{X}} \end{pmatrix}.$$
(48)

Then $\tilde{\mathbf{K}} = \Theta^{-}$.

Proof of Lemma A.5. In order to show that $\tilde{\mathbf{K}} = \Theta^{-}$, by definition of the Moore-Penrose inverse, we need to prove

- 1. $\Theta \tilde{\mathbf{K}}$ and $\tilde{\mathbf{K}} \Theta$ are symmetric,
- 2. $\Theta \tilde{\mathbf{K}} \Theta = \Theta$,
- 3. $\tilde{\mathbf{K}}\Theta\tilde{\mathbf{K}} = \tilde{\mathbf{K}}.$

To show symmetry, recall from Lemma A.3 that all the matrices $\mathbf{Q}, \alpha \mathbf{I}_n, \mathbf{\Sigma}, \mathbf{X}, \mathbf{1}_{n \times n}$ commute. As a consequence, $\forall \mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D} \in {\mathbf{Q}, \alpha \mathbf{I}_n, \mathbf{\Sigma}, \mathbf{X}, \mathbf{1}_{n \times n}}$, it holds (Lemma A.1):

$$\mathcal{M}_{p}\begin{pmatrix}\mathbf{A}\\\mathbf{B}\end{pmatrix}\mathcal{M}_{p}\begin{pmatrix}\mathbf{C}\\\mathbf{D}\end{pmatrix}=\mathcal{M}_{p}\begin{pmatrix}\mathbf{A}\mathbf{C}+(p-1)\mathbf{B}\mathbf{D}\\\mathbf{A}\mathbf{D}+\mathbf{B}\mathbf{C}+(p-2)\mathbf{B}\mathbf{D}\end{pmatrix}$$
$$=\mathcal{M}_{p}\begin{pmatrix}\mathbf{C}\mathbf{A}+(p-1)\mathbf{D}\mathbf{B}\\\mathbf{D}\mathbf{A}+\mathbf{C}\mathbf{B}+(p-2)\mathbf{D}\mathbf{B}\end{pmatrix}=\mathcal{M}_{p}\begin{pmatrix}\mathbf{C}\\\mathbf{D}\end{pmatrix}\mathcal{M}_{p}\begin{pmatrix}\mathbf{A}\\\mathbf{B}\end{pmatrix}$$

In addition, $\mathcal{M}_p(\mathbf{A}, \mathbf{B})^{\top} = \mathcal{M}_p(\mathbf{A}^{\top}, \mathbf{B}^{\top}) = \mathcal{M}_p(\mathbf{A}, \mathbf{B})$, as \mathbf{A}, \mathbf{B} are symmetric. Therefore,

$$\begin{split} \left(\boldsymbol{\Theta}\tilde{\mathbf{K}}\right)^{\top} &= \tilde{\mathbf{K}}^{\top}\boldsymbol{\Theta}^{\top} = \mathcal{M}_{p} \begin{pmatrix} \boldsymbol{\Sigma} + k_{1}\boldsymbol{1}_{n\times n} \\ \mathbf{X} + k_{2}\boldsymbol{1}_{n\times n} \end{pmatrix}^{\top} \mathcal{M}_{p} \begin{pmatrix} \mathbf{Q} \\ -\alpha \mathbf{I}_{n} \end{pmatrix}^{\top} \\ &= \mathcal{M}_{p} \begin{pmatrix} \boldsymbol{\Sigma}^{\top} + k_{1}\boldsymbol{1}_{n\times n}^{\top} \\ \mathbf{X}^{\top} + k_{2}\boldsymbol{1}_{n\times n}^{\top} \end{pmatrix} \mathcal{M}_{p} \begin{pmatrix} \mathbf{Q}^{\top} \\ -\alpha \mathbf{I}_{n} \end{pmatrix} \\ &= \mathcal{M}_{p} \begin{pmatrix} \boldsymbol{\Sigma} \\ \mathbf{X} \end{pmatrix} \mathcal{M}_{p} \begin{pmatrix} \mathbf{Q} \\ -\alpha \mathbf{I}_{n} \end{pmatrix} + \mathcal{M}_{p} \begin{pmatrix} k_{1}\boldsymbol{1}_{n\times n} \\ k_{2}\boldsymbol{1}_{n\times n} \end{pmatrix} \mathcal{M}_{p} \begin{pmatrix} \mathbf{Q} \\ -\alpha \mathbf{I}_{n} \end{pmatrix} \\ &= \mathcal{M}_{p} \begin{pmatrix} \mathbf{Q} \\ -\alpha \mathbf{I}_{n} \end{pmatrix} \mathcal{M}_{p} \begin{pmatrix} \boldsymbol{\Sigma} \\ \mathbf{X} \end{pmatrix} + \mathcal{M}_{p} \begin{pmatrix} \mathbf{Q} \\ -\alpha \mathbf{I}_{n} \end{pmatrix} \mathcal{M}_{p} \begin{pmatrix} k_{1}\boldsymbol{1}_{n\times n} \\ k_{2}\boldsymbol{1}_{n\times n} \end{pmatrix} \\ &= \mathcal{M}_{p} \begin{pmatrix} \mathbf{Q} \\ -\alpha \mathbf{I}_{n} \end{pmatrix} \mathcal{M}_{p} \begin{pmatrix} \boldsymbol{\Sigma} + k_{1}\boldsymbol{1}_{n\times n} \\ \mathbf{X} + k_{2}\boldsymbol{1}_{n\times n} \end{pmatrix} = \boldsymbol{\Theta}\tilde{\mathbf{K}}. \end{split}$$

The same arguments show that $(\tilde{\mathbf{K}}\Theta)^{\top} = \tilde{\mathbf{K}}\Theta$. To prove the other two relations, we start with following equalities. In the next lines, we used Lemmas

A.1, A.2 and A.3.

$$\begin{split} \Theta \mathbf{K} &= \mathcal{M}_p \begin{pmatrix} \mathbf{Q} \\ -\alpha \mathbf{I}_n \end{pmatrix} \mathcal{M}_p \begin{pmatrix} \boldsymbol{\Sigma} \\ \mathbf{X} \end{pmatrix} = \mathcal{M}_p \begin{pmatrix} \mathbf{Q} \boldsymbol{\Sigma} - \alpha(p-1) \mathbf{X} \\ \mathbf{Q} \mathbf{X} - \alpha \boldsymbol{\Sigma} - \alpha(p-2) \mathbf{X} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} \mathbf{I}_n \\ \frac{1}{-\alpha(p-1)} (\mathbf{Q} - \boldsymbol{\Sigma} \mathbf{Q}^2) - \alpha \boldsymbol{\Sigma} - \alpha(p-2) \mathbf{X} \end{pmatrix} = \mathcal{M}_p \begin{pmatrix} \mathbf{I}_n \\ -\frac{1}{n(p-1)} \mathbf{1}_{n \times n} \end{pmatrix} \\ \Theta \mathbf{K} \Theta &= \mathcal{M}_p \begin{pmatrix} \mathbf{I}_n \\ -\frac{1}{n(p-1)} \mathbf{1}_{n \times n} \end{pmatrix} \mathcal{M}_p \begin{pmatrix} \mathbf{Q} \\ -\alpha \mathbf{I}_n \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} \mathbf{Q} + \alpha(p-1) \frac{1}{n(p-1)} \mathbf{1}_{n \times n} \\ -\alpha \mathbf{I}_n - \frac{1}{n(p-1)} \mathbf{Q} \mathbf{1}_{n \times n} + \alpha(p-2) \frac{1}{n(p-1)} \mathbf{1}_{n \times n} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} \mathbf{Q} + \frac{\alpha}{n} \mathbf{1}_{n \times n} \\ -\alpha \mathbf{I}_n - \frac{\alpha}{n(p-1)} \mathbf{Q} \mathbf{1}_{n \times n} + \alpha(p-2) \frac{1}{n(p-1)} \mathbf{1}_{n \times n} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} \mathbf{Q} + \frac{\alpha}{n} \mathbf{1}_{n \times n} \\ k_2 \mathbf{1}_{n \times n} \end{pmatrix} = \mathcal{M}_p \begin{pmatrix} \mathbf{k}_1 \mathbf{Q} \mathbf{1}_{n \times n} - \alpha(p-1) k_2 \mathbf{1}_{n \times n} \\ k_2 \mathbf{Q} \mathbf{1}_{n \times n} - \alpha k_1 \mathbf{1}_{n \times n} - \alpha k_2 (p-2) \mathbf{1}_{n \times n} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} -\alpha(p-1)(k_2 - k_1) \mathbf{1}_{n \times n} \\ -\alpha(k_1 - k_2) \mathbf{1}_{n \times n} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} \alpha(p-1)(k_2 - k_1) \mathbf{1}_{n \times n} \\ -\alpha(k_1 - k_2) \mathbf{1}_{n \times n} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} \alpha(p-1)(k_2 - k_1) \mathbf{Q} \mathbf{1}_{n \times n} + \alpha^2(p-1)(k_1 - k_2) \mathbf{1}_{n \times n} \\ -\alpha(p-1)(k_2 - k_1) \mathbf{Q} \mathbf{1}_{n \times n} + \alpha^2(p-2)(k_1 - k_2) \mathbf{1}_{n \times n} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} \alpha^2 p(p-1)(k_2 - k_1) \mathbf{1}_{n \times n} \\ \alpha^2 (p-1)(k_2 - k_1) \mathbf{1}_{n \times n} \end{pmatrix} \end{pmatrix} \end{pmatrix}$$

As a consequence:

$$\begin{split} \boldsymbol{\Theta} \tilde{\mathbf{K}} \boldsymbol{\Theta} &= \boldsymbol{\Theta} \left(\mathbf{K} + \mathcal{M}_p \begin{pmatrix} k_1 \mathbf{1}_{n \times n} \\ k_2 \mathbf{1}_{n \times n} \end{pmatrix} \right) \boldsymbol{\Theta} \\ &= \mathcal{M}_p \begin{pmatrix} \mathbf{Q} + \frac{\alpha}{n} \mathbf{1}_{n \times n} \\ -\alpha \mathbf{I}_n - \frac{\alpha}{n(p-1)} \mathbf{1}_{n \times n} \end{pmatrix} + \mathcal{M}_p \begin{pmatrix} \alpha^2 p(p-1)(k_1 - k_2) \mathbf{1}_{n \times n} \\ -\alpha^2 p(k_1 - k_2) \mathbf{1}_{n \times n} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} \mathbf{Q} + \frac{\alpha}{n} \mathbf{1}_{n \times n} + \alpha^2 p(p-1)(k_1 - k_2) \mathbf{1}_{n \times n} \\ -\alpha \mathbf{I}_n - \frac{\alpha}{n(p-1)} \mathbf{1}_{n \times n} - \alpha^2 p(k_1 - k_2) \mathbf{1}_{n \times n} \end{pmatrix} = \mathcal{M}_p \begin{pmatrix} \mathbf{Q} \\ -\alpha \mathbf{I}_n \end{pmatrix} = \boldsymbol{\Theta}. \end{split}$$

Similarly we prove that $\tilde{\mathbf{K}} \Theta \tilde{\mathbf{K}} = \tilde{\mathbf{K}}$:

$$\begin{split} \mathbf{K} \Theta \mathbf{K} &= \mathcal{M}_p \begin{pmatrix} \mathbf{\Sigma} \\ \mathbf{X} \end{pmatrix} \mathcal{M}_p \begin{pmatrix} \mathbf{I}_n \\ -\frac{1}{n(p-1)} \mathbf{1}_{n \times n} \end{pmatrix} = \mathcal{M}_p \begin{pmatrix} \mathbf{\Sigma} - \frac{1}{n} \mathbf{X} \mathbf{1}_{n \times n} \\ -\frac{1}{n(p-1)} \mathbf{\Sigma} \mathbf{1}_{n \times n} + \mathbf{X} - \frac{p-2}{n(p-2)} \mathbf{X} \mathbf{1}_{n \times n} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} \mathbf{\Sigma} + \frac{1}{\alpha n(p-1)} \mathbf{1}_{n \times n} \\ \mathbf{X} + \frac{p-2}{\alpha n(p-1)^2} \mathbf{1}_{n \times n} \end{pmatrix} \\ \mathcal{M}_p \begin{pmatrix} k_1 \mathbf{1}_{n \times n} \\ k_2 \mathbf{1}_{n \times n} \end{pmatrix} \Theta \mathbf{K} = \mathbf{K} \Theta \mathcal{M}_p \begin{pmatrix} k_1 \mathbf{1}_{n \times n} \\ k_2 \mathbf{1}_{n \times n} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} k_1 \mathbf{1}_{n \times n} \\ k_2 \mathbf{1}_{n \times n} \end{pmatrix} \mathcal{M}_p \begin{pmatrix} \mathbf{I}_n \\ -\frac{1}{n(p-1)} \mathbf{1}_{n \times n} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} k_1 \mathbf{1}_{n \times n} \\ -\frac{k_1}{n(p-1)} \mathbf{1}_{n \times n} \mathbf{1}_{n \times n} + k_2 \mathbf{1}_{n \times n} \mathbf{1}_{n \times n} \\ -\frac{k_1}{n(p-1)} \mathbf{1}_{n \times n} \mathbf{1}_{n \times n} + k_2 \mathbf{1}_{n \times n} - \frac{k_2(p-2)}{n(p-1)} \mathbf{1}_{n \times n} \mathbf{1}_{n \times n} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} (k_1 - k_2) \mathbf{1}_{n \times n} \\ -\frac{1}{p-1} (k_1 - k_2) \mathbf{1}_{n \times n} \\ k_2 \mathbf{1}_{n \times n} \end{pmatrix} = \mathcal{M}_p \begin{pmatrix} k_1 \mathbf{1}_{n \times n} \\ k_2 \mathbf{1}_{n \times n} \end{pmatrix} \mathcal{M}_p \begin{pmatrix} -\alpha (p-1)(k_2 - k_1) \mathbf{1}_{n \times n} \\ -\alpha (k_1 - k_2) \mathbf{1}_{n \times n} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} \alpha (p-1)(k_1 - k_2)^2 \mathbf{1}_{n \times n} \\ (-\alpha k_1 (k_1 - k_2) n - \alpha (p-1)k_2 (k_2 - k_1)n - \alpha (p-2)k_2 (k_1 - k_2)n \end{pmatrix} \mathbf{1}_{n \times n} \end{pmatrix} \\ &= \mathcal{M}_p \begin{pmatrix} \alpha n(p-1)(k_1 - k_2)^2 \mathbf{1}_{n \times n} \\ -\alpha n(k_1 - k_2)^2 \mathbf{1}_{n \times n} \end{pmatrix} \end{pmatrix}$$

Therefore,

$$\tilde{\mathbf{K}}\boldsymbol{\Theta}\tilde{\mathbf{K}} = \mathbf{K}\boldsymbol{\Theta}\mathbf{K} + 2\,\mathcal{M}_p\begin{pmatrix}k_1\mathbf{1}_{n\times n}\\k_2\mathbf{1}_{n\times n}\end{pmatrix}\boldsymbol{\Theta}\mathbf{K} + \mathcal{M}_p\begin{pmatrix}k_1\mathbf{1}_{n\times n}\\k_2\mathbf{1}_{n\times n}\end{pmatrix}\boldsymbol{\Theta}\mathcal{M}_p\begin{pmatrix}k_1\mathbf{1}_{n\times n}\\k_2\mathbf{1}_{n\times n}\end{pmatrix} = \tilde{\mathbf{K}}.$$

Proof of Proposition 3. We first show that Θ is quasi-Laplacian. To this end, we need to show that Θ is symmetric, positive semidefinite, $\Theta \mathbf{1}_{np} = \mathbf{0}$, and Θ has only one null eigenvalue.

- 1. Since **Q** and $\alpha \mathbf{I}_n$ are symmetric and $\Theta := \mathcal{M}_p(\mathbf{Q}, -\alpha \mathbf{I}_n), \Theta$ is symmetric.
- 2. First recall that $\Theta = \mathbf{I}_p \otimes (\mathbf{Q} + \alpha \mathbf{I}_n) + \mathbf{1}_{p \times p} \otimes (-\alpha \mathbf{I}_n)$: we aim to apply Lemma A.4 to the matrix Θ with $\mathbf{X} := \mathbf{I}_p \otimes (\mathbf{Q} + \alpha \mathbf{I}_n)$ and $\mathbf{Y} :=$

 $\mathbf{1}_{p \times p} \otimes (-\alpha \mathbf{I}_n)$. In the following, we write $\lambda(\mathbf{A})$ to denote the vector of eigenvalues of \mathbf{A} . In addition, the inequalities apply to all elements: we write $\lambda(\mathbf{A}) \geq c$ as a shortcut of $\forall i \in \{1, \ldots, n\}$, $\lambda_i(\mathbf{A}) \geq c$, for an $n \times n$ matrix \mathbf{A} .

$$\boldsymbol{\lambda}(\mathbf{X}) = \boldsymbol{\lambda}(\mathbf{I}_p \otimes (\mathbf{Q} + \alpha \mathbf{I}_n)) = \boldsymbol{\lambda}(\mathbf{I}_p) \otimes \boldsymbol{\lambda}(\mathbf{Q} + \alpha \mathbf{I}_n) = \mathbf{1}_p \otimes (\boldsymbol{\lambda}(\mathbf{Q}) + \alpha)$$

$$\geq \lambda_n(\mathbf{Q}) + \alpha = \alpha(p-1) + \alpha = \alpha p$$

$$\boldsymbol{\lambda}(\mathbf{Y}) = \boldsymbol{\lambda}(\mathbf{1}_{p \times p} \otimes (-\alpha \mathbf{I}_n)) = \begin{bmatrix} p \\ 0 \\ \vdots \\ 0 \end{bmatrix} \otimes \begin{bmatrix} -\alpha \\ \vdots \\ -\alpha \end{bmatrix} \geq -\alpha p.$$

Therefore Θ is positive semidefinite.

3. To show $\Theta \mathbf{1}_{np} = \mathbf{0}$, given the special structure of Θ , it is sufficient to show that

$$\begin{bmatrix} \mathbf{Q} & -\alpha \mathbf{I}_n & \dots & -\alpha \mathbf{I}_n \end{bmatrix} \mathbf{1}_{np} = \mathbf{0},$$

i.e. $\mathbf{Q1}_n = \alpha(p-1)\mathbf{1}_n$, which has already been proved in Lemma A.2.

4. We now need to show that Θ has no other null eigenvalues, *i.e.* the only solution to

$$\begin{bmatrix} \mathbf{Q} & -\alpha \mathbf{I}_n & \dots & -\alpha \mathbf{I}_n \\ -\alpha \mathbf{I}_n & \mathbf{Q} & \dots & -\alpha \mathbf{I}_n \\ \vdots & \vdots & \ddots & \vdots \\ -\alpha \mathbf{I}_n & -\alpha \mathbf{I}_n & \dots & \mathbf{Q} \end{bmatrix} \begin{bmatrix} \boldsymbol{v}_1 \\ \boldsymbol{v}_2 \\ \vdots \\ \boldsymbol{v}_n \end{bmatrix} = \begin{bmatrix} \mathbf{0} \\ \mathbf{0} \\ \vdots \\ \mathbf{0} \end{bmatrix}$$
(49)

is (up to a multiplicative constant) $v_1 = \cdots = v_n = \mathbf{1}_n$. By reading Equation (49) by blocks, we get the following: $\forall i \in \{1, \ldots, n\}$

$$\mathbf{Q} \boldsymbol{v}_i - \alpha \sum_{k \neq i} \boldsymbol{v}_k = \mathbf{0}.$$

Taking now the difference for i = i and i = j, we get

$$(\mathbf{Q} + \alpha \mathbf{I}_n)(\boldsymbol{v}_i - \boldsymbol{v}_j) = \mathbf{0}$$

and, since $\mathbf{Q} + \alpha \mathbf{I}_n$ is strictly positive definite directly by Lemma A.2, we obtain that $\mathbf{v}_i = \mathbf{v}_j =: \mathbf{v}$ for all i, j. The problem thus becomes

$$(\mathbf{Q} - \alpha(p-1)\mathbf{I}_n)\boldsymbol{v} = \boldsymbol{0}.$$

To conclude it is sufficient to notice that the eigenvalues of $\mathbf{Q} - \alpha(p - 1)\mathbf{I}_n$ are $\lambda_i(\mathbf{Q}) - \alpha(p - 1)$ and, by Lemma A.2, all the eigenvalues are not less than $\alpha(p - 1)$ and exactly one achieve such a value. As a consequence, the kernel of $\mathbf{Q} - \alpha(p - 1)\mathbf{I}_n$ has dimension one and, since we already know that $(\mathbf{Q} - \alpha(p - 1)\mathbf{I}_n)\mathbf{1}_n = \mathbf{0}$, the only possibility is $\mathbf{v} = \mathbf{1}_n$.

That Θ^- is also a quasi-Laplacian matrix comes from Lemma A.5 and the closure of the set of quasi-Laplacian matrices under Moore-Penrose inversion (Proposition 1).

Proof of Proposition 4. It is sufficient to apply Proposition 2 with X := V. We show that the $np \times np$ matrix $[\mathbf{K}(v_i, v_j)]_{i,j=1}^{np}$ is positive semidefinite. First, notice that $[\mathbf{K}(v_i, v_j)]_{i,j=1}^{np} = \mathcal{M}_p(\tilde{\Sigma}, \tilde{\mathbf{X}}) = \tilde{\mathbf{K}}$. Since $\tilde{\mathbf{K}} \in \mathbb{R}^{np \times np}$ is positive semidefinite (see Lemma A.5 and Proposition 3), each matrix \mathbf{C} as defined in Proposition 2, being a principal submatrix of $\tilde{\mathbf{K}}$, is positive semidefinite as well. This proves the Proposition.

- *Proof of Proposition 7.* 1. The proof of this point is straightforward, as the pseudo variogram only takes non-negative values.
 - 2. The condition $u_1 = u_2$ and i = j is sufficient to have $D_{ij}(u_1, u_2) = 0$ due to the fact that the diagonal entries of $\Gamma_{\mathbf{Z}}(u_1, u_1)$ are zero. Let us show that it is also necessary. From the definition of the pseudo-variogram as per Equation (6), $D_{ij}(u_1, u_2) = 0$ if and only if $Z_i(u_1) = Z_j(u_2)$. Since $\mathbf{Z} = \mathbf{Z}_V + \mathbf{Z}_E$ with \mathbf{Z}_V and \mathbf{Z}_E being mutually independent, a necessary condition is that $Z_{V,i}(u_1) = Z_{V,j}(u_2)$. Given the correlation structure (22) at the vertices, the variance-covariance matrix $\tilde{\mathbf{K}}$ of $\mathbf{Z}_V|_V$ is quasi-Laplacian (Lemma A.5 and Proposition 3) and thus it has only one null eigenvalue associated with the vector $\mathbf{1}_{np}$. That is, the only non-trivial linear combination of $\mathbf{Z}_V|_V$ having a zero variance is, up to a constant factor, $\mathbf{1}_{np}^{\top} \mathbf{Z}_V|_V$. Therefore, $Z_{V,i}(u_1) = Z_{V,j}(u_2)$ for $u_1, u_2 \in V$ only if i = j and $u_1 = u_2$. Given the linear interpolation on the edges (Equation (23)), the same holds true for any two points u_1 and u_2 belonging to \mathcal{G} .
 - 3. The claim stems from Equations (8) and (29).

4. Let $u_0, u_1, \ldots, u_n \in \mathcal{G}$ and $c_1, \ldots, c_n \in \mathbb{R}^p$. Let $c_0 = -\mathbf{A} \sum_{i=1}^n c_i$, where **A** is a $p \times p$ matrix such that $\mathbf{A}^{\top} \mathbf{1}_p = \mathbf{1}_p$. Then, from Equation (9), one has:

$$\begin{split} 0 &\geq \sum_{i,j=0}^{n} \boldsymbol{c}_{i}^{\top} \boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_{i}, u_{j}) \boldsymbol{c}_{j} \\ &= \sum_{i,j=1}^{n} \boldsymbol{c}_{i}^{\top} \boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_{i}, u_{j}) \boldsymbol{c}_{j} + \sum_{i=1}^{n} \boldsymbol{c}_{i}^{\top} \boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_{i}, u_{0}) \boldsymbol{c}_{0} + \sum_{j=1}^{n} \boldsymbol{c}_{0}^{\top} \boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_{0}, u_{j}) \boldsymbol{c}_{j} + \boldsymbol{c}_{0}^{\top} \boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_{0}, u_{0}) \boldsymbol{c}_{0} \\ &= \sum_{i,j=1}^{n} \boldsymbol{c}_{i}^{\top} \left[\boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_{i}, u_{j}) - \boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_{i}, u_{0}) \mathbf{A} - \mathbf{A}^{\top} \boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_{0}, u_{j}) + \mathbf{A}^{\top} \boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_{0}, u_{0}) \mathbf{A} \right] \boldsymbol{c}_{j}, \end{split}$$

i.e., the mapping $(u, v) \mapsto -[\Gamma_{\mathbf{Z}}(u, v) - \Gamma_{\mathbf{Z}}(u, u_0)\mathbf{A} - \mathbf{A}^{\top}\Gamma_{\mathbf{Z}}(u_0, v) + \mathbf{A}^{\top}\Gamma_{\mathbf{Z}}(u_0, u_0)\mathbf{A}]$ is positive semidefinite (Equation 12).

5. Taking $c_1 = \ldots = c_n := c$ and $\mathbf{A} = \mathbf{I}_p$, one gets

$$0 \ge \boldsymbol{c}^{\top} \sum_{i,j=1}^{n} \left[\boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_i, u_j) - \boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_i, u_0) - \boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_0, u_j) + \boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_0, u_0) \right] \boldsymbol{c}.$$

Accounting for the fact that $\Gamma_{\mathbf{Z}} = \Gamma_{\mathbf{Z}}^{\top}$ and $\Gamma_{\mathbf{Z}}(u_0, u_0) = \Gamma_{\mathbf{Z}}(u_1, u_1) = \Gamma_{\mathbf{Z}}(u_2, u_2)$, one obtains, for n = 1 and $u_0 = u_2$:

$$0 \ge \boldsymbol{c}^{\top} \left[\boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_0, u_0) - \boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_1, u_2) \right] \boldsymbol{c},$$

and for n = 2:

$$0 \ge \boldsymbol{c}^{\top} \left[\boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_1, u_2) + 3\boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_0, u_0) - 2\boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_1, u_0) - 2\boldsymbol{\Gamma}_{\boldsymbol{Z}}(u_2, u_0) \right] \boldsymbol{c},$$

i.e., $\Gamma_{\mathbf{Z}}(u_1, u_2) - \Gamma_{\mathbf{Z}}(u_0, u_0)$ and $2\Gamma_{\mathbf{Z}}(u_1, u_0) + 2\Gamma_{\mathbf{Z}}(u_2, u_0) - \Gamma_{\mathbf{Z}}(u_1, u_2) - 3\Gamma_{\mathbf{Z}}(u_0, u_0)$ are positive semidefinite matrices.

Proof of Proposition 8. 1. We split the computation of the distances (the one proposed here, **D**, and the one defined by Anderes et al. [2020], d_R) in two: the contribution (*i.e.* the variogram) of the \mathbf{Z}_V process and the one of \mathbf{Z}_E , denoted via the superscripts V and E respectively. We show that $\mathbf{D}_{ii}^V(u_1, u_2) = d_R^V(u_1, u_2)$ and $\mathbf{D}_{ii}^E(u_1, u_2) = d_R^E(u_1, u_2)$. The latter comes straightforwardly by comparing the diagonal entries

of Equation (24), *i.e.* $\mathbb{1}(e_1 = e_2) \ell(e_1) (\delta_1 \wedge \delta_2 - \delta_1 \delta_2)$, with the kernel R_e in Anderes et al. [2020, Equation 15].

Thus, let us show that $\mathbf{D}_{ii}^{V}(u_1, u_2) = d_R^{V}(u_1, u_2)$. Let $u_1, u_2 \in \mathcal{G}$ and $k_{V,ii}$ be the *i*-th diagonal entry of $\mathbf{K}_{\mathbf{Z}_V}$. Then, on account of Proposition 5:

$$\begin{split} \mathbf{D}_{ii}^{V}(u_{1}, u_{2}) &= k_{V,ii}(u_{1}, u_{1}) + k_{V,ii}(u_{2}, u_{2}) - 2k_{V,ii}(u_{1}, u_{2}) \\ &= \boldsymbol{\delta}_{1,V}^{\top} \tilde{\boldsymbol{\Sigma}} \boldsymbol{\delta}_{1,V} + \boldsymbol{\delta}_{2,V}^{\top} \tilde{\boldsymbol{\Sigma}} \boldsymbol{\delta}_{2,V} - 2\boldsymbol{\delta}_{1,V}^{\top} \tilde{\boldsymbol{\Sigma}} \boldsymbol{\delta}_{2,V} \\ &= \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right)^{\top} \left(\boldsymbol{\Sigma} + k_{1} \mathbf{1}_{n \times n}\right) \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right) \\ &= \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right)^{\top} \boldsymbol{\Sigma} \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right) + k_{1} \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right)^{\top} \mathbf{1}_{n \times n} \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right) \\ &= \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right)^{\top} \boldsymbol{\Sigma} \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right) + k_{1} \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right)^{\top} \mathbf{1}_{n \times n} \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right) \end{split}$$

where in the last step we have used $\delta_{1,V}^{\top} \mathbf{1}_n = \delta_{2,V}^{\top} \mathbf{1}_n = 1$ owing to Equation (26).

It is left to show that this quantity coincides with the resistance counterpart of Anderes et al. [2020]. They define their covariance matrix, here denoted by $\Sigma_{\mathbf{A}}$, as the (proper) inverse of a modified Laplacian matrix \mathbf{L} , where a one has been added on an arbitrary diagonal entry. In formulae: $\Sigma_{\mathbf{A}} := (\mathbf{L} + \mathbf{e}_j \mathbf{e}_j^{\top})^{-1}$, where \mathbf{e}_j is the j^{th} vector of the canonical basis of \mathbb{R}^n . It follows directly from Anderes et al. [2020, Equation 13] that $d_R^V = (\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V})^{\top} \Sigma_{\mathbf{A}} (\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V})$.

Let us show that, for a given quasi-Laplacian matrix **L** and for any $\boldsymbol{x} \in \mathbb{R}^n$ such that $\mathbf{1}_n^{\top} \boldsymbol{x} \neq 0$, it holds:

$$(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V})^{\top} \left(\mathbf{L} + \boldsymbol{x} \, \boldsymbol{x}^{\top} \right)^{-} (\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}) = (\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V})^{\top} \, \mathbf{L}^{-} \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V} \right).$$

$$(50)$$

In order to express the Moore-Penrose inverse of the rank-1 update $\mathbf{L} + \boldsymbol{x} \boldsymbol{x}^{\mathsf{T}}$, we exploit Theorem 1 of Meyer [1973] with $\mathbf{A} := \mathbf{L}$ and $\boldsymbol{c} := \boldsymbol{d} := \boldsymbol{x}$. Firstly, let us show that the hypotheses of the theorem are satisfied: we have to prove that $\boldsymbol{x} \notin R(\mathbf{L})$, where R denotes the range or column space. $R(\mathbf{L})$ is perpendicular to the kernel of \mathbf{L} , that is (recall that \mathbf{L} is a quasi-Laplacian matrix) ker $(\mathbf{L}) = \{h \mathbf{1}_n : h \in \mathbb{R}\}$. Now, $\boldsymbol{x} \in R(\mathbf{L}) \iff \boldsymbol{x} \perp \text{ker}(\mathbf{L}) \iff \mathbf{1}_n^{\mathsf{T}} \boldsymbol{x} = 0$, yet $\mathbf{1}_n^{\mathsf{T}} \boldsymbol{x} \neq 0$ by hypothesis. Therefore $\boldsymbol{x} \notin R(\mathbf{L})$, *i.e.* we can apply the above-mentioned theorem. In the following computations, we have adopted

the notation used by Meyer [1973, Section 2], i.e. :

$$egin{aligned} m{k} &:= \mathbf{L}^{-}m{x} & m{h} &:= m{x}^{ op} \mathbf{L}^{-} \ m{u} &:= ig(\mathbf{I}_n - \mathbf{L}\mathbf{L}^{-}ig)m{x} = rac{1}{n}m{1}_{n imes n}m{x} & m{v} &:= m{x}^{ op}ig(\mathbf{I}_n - \mathbf{L}^{-}\mathbf{L}ig) = rac{1}{n}m{x}^{ op}m{1}_{n imes n} \ m{eta} &:= 1 + m{x}^{ op} \mathbf{L}^{-}m{x}. \end{aligned}$$

In addition, we have $\|\boldsymbol{u}\|^2 = \|\boldsymbol{v}\|^2 = \boldsymbol{u}^\top \boldsymbol{u} = \frac{1}{n^2} \mathbf{1}_n^\top (\mathbf{1}_n^\top \boldsymbol{x}) (\mathbf{1}_n^\top \boldsymbol{x}) \mathbf{1}_n = \frac{1}{n} (\mathbf{1}_n^\top \boldsymbol{x})^2 \neq 0$. Let us finally apply Equation (3.1) of Meyer [1973]:

$$egin{aligned} \left(\mathbf{L}+oldsymbol{x}oldsymbol{x}^{ op}
ight)^{ op} &= \mathbf{L}^{ op}-oldsymbol{k}oldsymbol{u}^{ op}-oldsymbol{v}^{ op}oldsymbol{h}+etaoldsymbol{v}^{ op}oldsymbol{u}^{ op} &= \mathbf{L}^{ op}-oldsymbol{k}oldsymbol{u}^{ op}oldsymbol{h}^{ op}oldsymbol{u}^{ op}ol$$

To show (50), it is now sufficient to prove that

$$\left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right)^{\top} \left(\left(1 + \boldsymbol{x}^{\top} \mathbf{L}^{-} \boldsymbol{x}\right) \frac{\mathbf{1}_{n \times n}}{\left(\mathbf{1}_{n}^{\top} \boldsymbol{x}\right)^{2}} - \frac{\mathbf{L}^{-} \boldsymbol{x} \mathbf{1}_{n}^{\top}}{\mathbf{1}_{n}^{\top} \boldsymbol{x}} - \frac{\mathbf{1}_{n} \boldsymbol{x}^{\top} \mathbf{L}^{-}}{\mathbf{1}_{n}^{\top} \boldsymbol{x}} \right) \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right) = 0.$$

It is possible to see this by noticing that $\mathbf{1}_n^{\top} (\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}) = \mathbf{1}_n^{\top} \boldsymbol{\delta}_{1,V} - \mathbf{1}_n^{\top} \boldsymbol{\delta}_{2,V} = 1 - 1 = 0$ and similarly $(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V})^{\top} \mathbf{1}_n = 0$, therefore each term of the above equation equals 0.

As a consequence,

$$\begin{split} d_R^V(u_1, u_2) &= \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right)^\top \boldsymbol{\Sigma}_{\mathbf{A}} \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right) \\ &= \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right)^\top \left(\mathbf{L} + \boldsymbol{e}_j \boldsymbol{e}_j^\top\right)^- \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right) \\ &= \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right)^\top \mathbf{L}^- \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right) = D_{ii}^V(u_1, u_2). \end{split}$$

2. The proof of this point is straightforward from Equation (29): notice that $D_{ij}(u_1, u_2)$, for $i \neq j$, is the off-diagonal element of $\mathbf{D}(u_1, u_2)$, whilst $D_{ii}(u_1, u_2)$ is the main diagonal one.

Proof of Proposition 9. We prove each row of the table, in turn. In the following computations, \sim_a stands for "asymptotically equivalent to".

1. Fix $\alpha > 0$ and let $p \to +\infty$. Considering Equations (19) and (21), we have:

$$\mathbf{\Lambda}^{-} + \alpha(p-2)\mathbf{I}_{n} + \sqrt{(\mathbf{\Lambda}^{-})^{2} - 2\alpha(p-2)\mathbf{\Lambda}^{-} + \alpha^{2}p^{2}\mathbf{I}_{n}}$$

$$\sim_{a} \alpha(p-2)\mathbf{I}_{n} + \alpha p\mathbf{I}_{n} = 2\alpha(p-1)\mathbf{I}_{n}$$
(51)

$$\alpha(p-2)\mathbf{\Lambda} + \sqrt{\mathbf{I}_n - \mathbf{J}_n - 2\alpha(p-2)\mathbf{\Lambda} + \alpha^2 p^2 \mathbf{\Lambda}^2} - \mathbf{I}_n - \mathbf{J}_n \qquad (52)$$

$$\sim_a \alpha(p-2)\mathbf{\Lambda} + \alpha p \mathbf{\Lambda} = 2\alpha(p-1)\mathbf{\Lambda}.$$

As a consequence, $\mathbf{Q} \sim_a \alpha(p-1)\mathbf{I}_n$ and $\mathbf{X} \to \mathbf{\Sigma}$. Furthermore, $k_1, k_2 \to 0$ and thus $\tilde{\mathbf{\Sigma}}, \tilde{\mathbf{X}} \to \mathbf{\Sigma}$ as $p \to +\infty$. From this last relation, we obtain that the process \mathbf{Z}_V satisfies \mathbb{C} or $(Z_{V,i}, Z_{V,j}) \to 1$ for all $i, j \in \{1, \ldots, p\}$. Throughout the whole proof, the expression for the distance changes only via the first line of Equation (29), reported below for an neat exposition.

$$\boldsymbol{\delta}_{1,V}^{\top} \tilde{\boldsymbol{\Sigma}} \boldsymbol{\delta}_{1,V} + \boldsymbol{\delta}_{2,V}^{\top} \tilde{\boldsymbol{\Sigma}} \boldsymbol{\delta}_{2,V} - 2 \boldsymbol{\delta}_{1,V}^{\top} \tilde{\mathbf{X}} \boldsymbol{\delta}_{2,V}$$

$$= \boldsymbol{\delta}_{1,V}^{\top} \boldsymbol{\Sigma} \boldsymbol{\delta}_{1,V} + \boldsymbol{\delta}_{2,V}^{\top} \boldsymbol{\Sigma} \boldsymbol{\delta}_{2,V} - 2 \boldsymbol{\delta}_{1,V}^{\top} \mathbf{X} \boldsymbol{\delta}_{2,V} + 2k_1 - 2k_2.$$
(53)

In this case,

$$(53) \to (\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V})^{\top} \, \tilde{\boldsymbol{\Sigma}} \, (\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V})$$

If we add the hypothesis $\beta \to 1$, \mathbb{C} or $(Z_{E,i}, Z_{E,j}) \to 1$ as well. This implies that all the $p \times p$ entries of $\mathbf{D}(u_1, u_2)$ converge to the same term, that is $d_R(u_1, u_2)$.

- 2. Fix $p \ge 2$ and let $\alpha \to +\infty$. The same asymptotic properties (51) and (52) hold (notice that p and α always appear with the same exponent in each monomial). Therefore, the same reasoning applies.
- 3. Fix $p \ge 2$ and let $\alpha \to 0^+$. From Equation (18), it is patent that $\mathbf{Q} \to \frac{1}{2}(\mathbf{L} + \sqrt{\mathbf{L}^2}) = \mathbf{L}$; whilst from Equation (21), we have:

$$\mathbf{X} = \frac{1}{2(p-1)} \mathbf{W} \left((p-2)\mathbf{\Lambda} + \frac{1}{\alpha} \sqrt{\mathbf{I}_n - \mathbf{J}_n - 2\alpha(p-2)\mathbf{\Lambda} + \alpha^2 p^2 \mathbf{\Lambda}} - \frac{1}{\alpha} \mathbf{I}_n - \frac{1}{\alpha} \mathbf{J}_n \right) \mathbf{W}^{\top}$$

Consider the diagonal matrix in the big brackets: it is possible to expand the square root term by means of the Taylor expansion $\sqrt{a - b \cdot \alpha + c \cdot \alpha^2} =$

 $\sqrt{a} - \frac{b}{2\sqrt{a}}\alpha + o(\alpha)$ as $\alpha \to 0^+$: the *i*th element on the main diagonal of $\sqrt{\mathbf{I}_n - \mathbf{J}_n - 2\alpha(p-2)\mathbf{\Lambda} + \alpha^2 p^2 \mathbf{\Lambda}}$ is therefore:

$$\begin{cases} \sqrt{1} - \frac{2(p-2)\lambda_i}{2\sqrt{1}}\alpha + o(\alpha) = 1 - (p-2)\lambda_i \alpha + o(\alpha) & \text{if } i \neq n \\ 0 & \text{if } i = n. \end{cases}$$

As a consequence, we have:

$$(p-2)\mathbf{\Lambda} + \frac{1}{\alpha}\sqrt{\mathbf{I}_n - \mathbf{J}_n - 2\alpha(p-2)\mathbf{\Lambda} + \alpha^2 p^2 \mathbf{\Lambda}} - \frac{1}{\alpha}\mathbf{I}_n - \frac{1}{\alpha}\mathbf{J}_n$$

$$\sim_a (p-2)\mathbf{\Lambda} + \frac{1}{\alpha}(\mathbf{I}_n - \mathbf{J}_n) - (p-2)\mathbf{\Lambda} - \frac{1}{\alpha}\mathbf{I}_n - \frac{1}{\alpha}\mathbf{J}_n$$

$$= -\frac{2}{\alpha}\mathbf{J}_n,$$

that is:

$$\mathbf{X} \sim \frac{1}{2(p-1)} \mathbf{W} \left(-\frac{2}{\alpha} \mathbf{J}_n \right) \mathbf{W}^{\top} = -\frac{1}{\alpha(p-1)} \frac{1}{n} \mathbf{1}_{n \times n},$$

where in the last step we have used from the proof of Lemma A.3 that $\mathbf{W}\mathbf{J}_n\mathbf{W}^{\top} = \frac{1}{n}\mathbf{1}_{n\times n}$ (Equation (47)). In this case

(53)
$$\sim_a \boldsymbol{\delta}_{1,V}^{\top} \boldsymbol{\Sigma} \boldsymbol{\delta}_{1,V} + \boldsymbol{\delta}_{2,V}^{\top} \boldsymbol{\Sigma} \boldsymbol{\delta}_{2,V} - 2 \boldsymbol{\delta}_{1,V}^{\top} \left(-\frac{1}{\alpha n(p-1)} \mathbf{1}_{n \times n} \right) \boldsymbol{\delta}_{2,V} - 2 \frac{1}{\alpha n p(p-1)} = \boldsymbol{\delta}_{1,V}^{\top} \boldsymbol{\Sigma} \boldsymbol{\delta}_{1,V} + \boldsymbol{\delta}_{2,V}^{\top} \boldsymbol{\Sigma} \boldsymbol{\delta}_{2,V} + \frac{2}{\alpha n p}.$$

Clearly, the contribution of the process \mathbf{Z}_E in Equation (29) and the above term $\boldsymbol{\delta}_{1,V}^{\top} \Sigma \boldsymbol{\delta}_{1,V} + \boldsymbol{\delta}_{2,V}^{\top} \Sigma \boldsymbol{\delta}_{2,V}$ do not depend on α . As a consequence, $D_{ij}(u_1, u_2) \sim_a \frac{2}{\alpha n p}$ for $i \neq j$ as $\alpha \to 0^+$.

4. Let $\alpha \to 0^+$ and $p \to +\infty$ such that $\alpha p \tau \to 1$ with $0 < \tau \leq \lambda_{n-1}$ being fixed. Then, from Equations (19) and (46), one has $\mathbf{Q} \to \frac{\mathbf{I}_n}{\tau}$. Likewise, from Equation (21), it comes $\mathbf{X} \to \mathbf{\Sigma} - \tau \mathbf{I}_n$. The computation of the distances $D_{ij}(u_1, u_2)$ is straightforward: following the same reasoning of the previous derivation, we have

$$(53) = \boldsymbol{\delta}_{1,V}^{\top} \boldsymbol{\Sigma} \boldsymbol{\delta}_{1,V} + \boldsymbol{\delta}_{2,V}^{\top} \boldsymbol{\Sigma} \boldsymbol{\delta}_{2,V} - 2 \boldsymbol{\delta}_{1,V}^{\top} \left(\boldsymbol{\Sigma} - \tau \mathbf{I}_n\right) \boldsymbol{\delta}_{2,V} - 2 \frac{1}{\alpha n p(p-1)} \rightarrow \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right)^{\top} \boldsymbol{\Sigma} \left(\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{2,V}\right) + 2\tau \boldsymbol{\delta}_{1,V}^{\top} \boldsymbol{\delta}_{2,V}.$$

The first addendum, together with the contribution of the process Z_E (if $\beta \to 1$), gives the distance $d_R(u_1, u_2)$.

5. Let $\alpha \to 0^+$ and $p \to +\infty$, with $\alpha p \tau \to 1$ with $\tau \ge \lambda_1$ fixed. Equation (19) gives

$$\mathbf{Q} \to \mathbf{W} \operatorname{diag} \left[\lambda_1^{-1}, \dots, \lambda_{n-1}^{-1}, \frac{1}{\tau}\right]^\top \mathbf{W}^\top = \mathbf{W} \left(\mathbf{\Lambda}^- + \frac{1}{\tau} \mathbf{J}_n\right) \mathbf{W}^\top = \mathbf{L} + \frac{1}{n\tau} \mathbf{1}_{n \times n}.$$

Therefore, from Equation (20):

$$\mathbf{X} = \frac{p\tau}{p-1} \left(\mathbf{\Sigma} \left(\mathbf{L} + \frac{1}{n\tau} \mathbf{1}_{n \times n} \right) - \mathbf{I}_n \right) \to -\frac{\tau}{n} \mathbf{1}_{n \times n}, \quad \text{as } p \to +\infty.$$

Hence,

$$(53) \to \boldsymbol{\delta}_{1,V}^{\top} \boldsymbol{\Sigma} \boldsymbol{\delta}_{1,V} + \boldsymbol{\delta}_{2,V}^{\top} \boldsymbol{\Sigma} \boldsymbol{\delta}_{2,V} - 2 \boldsymbol{\delta}_{1,V}^{\top} \left(-\frac{\tau}{n} \mathbf{1}_{n \times n} \right) \boldsymbol{\delta}_{2,V} - 2 \frac{1}{\alpha n p (p-1)} \to \boldsymbol{\delta}_{1,V}^{\top} \boldsymbol{\Sigma} \boldsymbol{\delta}_{1,V} + \boldsymbol{\delta}_{2,V}^{\top} \boldsymbol{\Sigma} \boldsymbol{\delta}_{2,V} + \frac{2\tau}{n}.$$

The result for the distance $D_{ij}(u_1, u_2)$ is obtained from the last equation by adding and subtracting $2\delta_{1,V}^{\top} \mathbf{X} \delta_{2,V}$ and by using the fact that $\beta \to 1$.

Proof of Theorem 3.1. The result stems from Theorem 2.2 and the fact that the pseudo-variogram \mathbf{D} satisfies $\mathbf{D}^{\top}(u_1, u_2) = \mathbf{D}(u_2, u_1)$ and is $\mathbf{1}_p$ -conditionally negative semidefinite.

Proof of Theorem 3.2. The result stems from Theorem 3.1 with $\psi(t) = \exp(-t)$ and $g(t) = t^{\theta}$ and the fact that the set of positive semidefinite matrix-valued functions is closed under sums, point-wise limits and element-wise products with symmetric positive semidefinite matrices.

Proof of Theorem 3.4. From Emery et al. [2022, Equations (28)-(30) and Theorem 3], \mathcal{M}_p has a representation of the form (31) with $dF_{ij}(\xi) = R_{ij}(\xi)d\xi$, where $[R_{ij}(\xi)]_{i,j=1}^p$ is positive semidefinite under the stated conditions (A) or (B). The theorem thus results from the application of Theorem 3.2. *Proof of Theorem 3.5.* Using formula 3.381.4 of Gradshteyn and Ryzhik [2007], one can write

$$\boldsymbol{\mathcal{C}}_{p}(\mathbf{D}) = \left[\frac{\sigma_{ij}\beta_{ij}^{\nu}}{\Gamma(\nu)}\int_{0}^{+\infty}\xi^{\nu-1}\exp(-\beta_{ij}\xi)\exp(-\xi D_{ij}^{\theta})\mathrm{d}\xi\right]_{i,j=1}^{p}.$$

The conditional negative semidefiniteness of $[\beta_{ij}]_{i,j=1}^p$ implies that $\left[\frac{\xi^{\nu-1}}{\Gamma(\nu)}\exp(-\beta_{ij}\xi)\right]_{i,j=1}^p$ is positive semidefinite for any $\xi \ge 0$ [Reams, 1999, Lemma 2.5]. The theorem results from the application of the Schur product theorem and Theorem 3.2.

Proof of Theorem 3.6. One can write [Porcu et al., 2018]

$$\mathcal{SG}_{p}(\mathbf{D}) = \left[\frac{\sigma_{ij}2^{\nu-1}(\alpha_{ij}\beta_{ij})^{\nu/2}}{\mathcal{K}_{\nu}\left(\sqrt{\frac{\beta_{ij}}{\alpha_{ij}}}\right)}\int_{0}^{+\infty}\xi^{\nu-1}\exp\left(-\frac{1}{4\alpha_{ij}\xi} - \beta_{ij}\xi\right)\exp(-\xi D_{ij}^{\theta})\mathrm{d}\xi\right]_{i,j=1}^{p}$$

The fact that $[\alpha_{ij}^{-1}]_{i,j=1}^p$ and $[\beta_{ij}]_{i,j=1}^p$ are conditionally negative semidefinite implies that

$$\left[\exp\left(-\frac{1}{4\alpha_{ij}\xi}-\beta_{ij}\xi\right)\right]_{i,j=1}^{p}$$

is positive semidefinite for any $\xi \ge 0$. The theorem results from the application of the Schur product theorem and Theorem 3.2.

Proof of Proposition 10. We start with the case p = 1 and observe that, by Bochner's theorem [Bochner, 1955], a continuous mapping $C : \mathbb{R}^{m'} \to \mathbb{R}$ is positive semidefinite if and only if

$$C(\boldsymbol{x}) = \int_{\mathbb{R}^{m'}} \exp(i\boldsymbol{x}^{\top}\boldsymbol{\omega}) G(\mathrm{d}\boldsymbol{\omega}), \qquad \boldsymbol{x} \in \mathbb{R}^{m'},$$

for G being a non-decreasing and bounded measure in \mathbb{R}^m . Here, i denotes the unit complex number. Using the above representation, Cambanis et al. [1983] show that a mapping $K : [0, +\infty) \to \mathbb{R}$ belongs to the class Φ_m^1 if and only if Equation (37) holds for p = 1 and for F being non-decreasing and bounded. In particular, choosing a Dirac measure for F, one finds that $t \mapsto \omega_{m'}(rt)$ belongs to Φ_m^1 for any $r \ge 0$, that is, $\omega_{m'}(rd_M(\cdot, \cdot))$ is positive semidefinite on $\mathbb{R}^{m'}$.

We now address the general case with $p \geq 1$. First, consider a matrixvalued mapping **K** belonging to the class Φ_m^p . We invoke the extension of Bochner's theorem to the *p*-variate case [Cramer, 1940], for which a continuous mapping $\mathbf{C} : \mathbb{R}^{m'} \to \mathbb{R}^{p \times p}$ is positive semidefinite if and only if

$$\mathbf{C}(\boldsymbol{x}) = \int_{\mathbb{R}^{m'}} \exp(\mathrm{i} \boldsymbol{x}^{\top} \boldsymbol{\omega}) \mathbf{G}(\mathrm{d} \boldsymbol{\omega}), \qquad \boldsymbol{x} \in \mathbb{R}^{m'},$$

for **G** a bounded complex Hermitian matrix-valued measure. Hence, the same calculations as much as in Cambanis et al. [1983] show that the representation (37) holds for a bounded symmetric real matrix-valued measure **F** (that **F** is bounded stems from the fact that $\mathbf{K}(0)$ is finite). To finish this part of the proof we need to show that $d\mathbf{F}(r)$ is positive semidefinite for any $r \geq 0$, which happens if and only if the mapping

$$H_{\boldsymbol{c}}(r) := \boldsymbol{c}^{\top} \mathbf{F}(\mathrm{d}r) \boldsymbol{c}, \quad r \in [0, +\infty),$$

is non-negative for any arbitrary $\boldsymbol{c} \in \mathbb{R}^p$. To this end, consider a *p*-variate random field \boldsymbol{Z} in $\mathbb{R}^{m'}$ with isotropic covariance \mathbf{K} , and define $Z_{\boldsymbol{c}} := \boldsymbol{c}^{\top} \boldsymbol{Z}$, which is a univariate random field with isotropic covariance $K_{\boldsymbol{c}} = \boldsymbol{c}^{\top} \mathbf{K} \boldsymbol{c}$. Hence, $K_{\boldsymbol{c}}$ obeys to the representation provided by Cambanis et al. [1983], with a measure $F_{\boldsymbol{c}}$ such that

$$F_{\boldsymbol{c}} = \boldsymbol{c}^{\top} \mathbf{F} \boldsymbol{c}.$$

This proves the sufficiency part of the theorem by noting that F_c is non-decreasing.

Reciprocally, assume that the representation (37) holds with a bounded symmetric matrix-valued measure \mathbf{F} such that $d\mathbf{F}(r)$ is positive semidefinite for any $r \geq 0$. Being positive semidefinite on $\mathbb{R}^{m'}$ for any $r \geq 0$, $\omega_{m'}(rd_M(\cdot, \cdot))$ is the covariance function of a univariate Gaussian random field Z_r on $\mathbb{R}^{m'}$. Accordingly, the matrix-valued mapping $\omega_{m'}(rd_M(\cdot, \cdot))\mathbf{1}_{p\times p}$ belongs to Φ_m^p as the covariance function of $Z_r\mathbf{1}_p$, and so do the mappings $\omega_{m'}(rd_M(\cdot, \cdot))d\mathbf{F}(r)$ and $\mathbf{K}(d_M(\cdot, \cdot))$ owing to the fact that the set of positive semidefinite functions is closed under sums, element-wise products, and pointwise limits.

Proof of Proposition 11. We provide a proof by direct construction. Let m' be a positive integer. Arguments in Theorem 2.1 of Gneiting [1998] provide the identity

$$\omega_{m'}(t) = \kappa_{m'} I^{(m'-1)} \Omega_{2m'-1}(t), \qquad t \ge 0, \tag{54}$$

for $\kappa_{m'}$ a strictly positive constant. Hence, the proof comes from the integral representation (37) in concert with a straight use of Fubini's theorem. \Box

Proof of Proposition 12. The results comes from a straight application of Proposition 2 in Emery and Porcu [2023] in concert with Lemma 4.1, Proposition 11 and Fubini's theorem. $\hfill \Box$



Figure 5: An example of graph for which the distance $D_{11}(v_1, v_3) \not\leq D_{12}(v_1, v_3)$, for $p \geq 3$ and α wisely chosen.

B A Worked Example

Here we provide a simple example of the multivariate distance properties.

Example. Let $p \geq 3$ and $\alpha > 0$. Consider the graph depicted in Figure 5 with $V := \{v_1, v_2, v_3\}$ and whose Laplacian matrix **L** and $\Sigma = \mathbf{L}^-$ are

$$\mathbf{L} = \begin{bmatrix} 1 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 1 \end{bmatrix}, \qquad \mathbf{\Sigma} = \frac{1}{9} \begin{bmatrix} 5 & -1 & -4 \\ -1 & 2 & -1 \\ -4 & -1 & 5 \end{bmatrix}.$$

In addition, the eigendecomposition $\boldsymbol{\Sigma} = \mathbf{W} \boldsymbol{\Lambda} \mathbf{W}^{\top}$ is

$$\mathbf{W} = \begin{bmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ 0 & -\frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \end{bmatrix}, \qquad \mathbf{\Lambda} = \begin{bmatrix} 1 & & \\ & \frac{1}{3} & \\ & & 0 \end{bmatrix}.$$

Let us consider the distance between v_1 and v_3 for the same variable and for different variables, *i.e.* $D_{11}(v_1, v_3)$ and $D_{12}(v_1, v_3)$. Since both v_1 and v_3 are vertices, the contribution of Z_E is null, therefore, by Equation (29), we have:

$$D_{11}(v_1, v_3) = (\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{3,V})^{\top} \tilde{\boldsymbol{\Sigma}} (\boldsymbol{\delta}_{1,V} - \boldsymbol{\delta}_{3,V})$$

= $[1, 0, -1] \tilde{\boldsymbol{\Sigma}} \begin{bmatrix} 1\\0\\-1 \end{bmatrix}$
= $\tilde{\Sigma}_{11} + \tilde{\Sigma}_{33} - 2\tilde{\Sigma}_{13}$
= $\Sigma_{11} + \Sigma_{33} - 2\Sigma_{13} = 2,$
$$D_{12}(v_1, v_3) = \boldsymbol{\delta}_{1,V}^{\top} \tilde{\boldsymbol{\Sigma}} \boldsymbol{\delta}_{1,V} + \boldsymbol{\delta}_{3,V}^{\top} \tilde{\boldsymbol{\Sigma}} \boldsymbol{\delta}_{3,V} - 2\boldsymbol{\delta}_{1,V}^{\top} \tilde{\boldsymbol{X}} \boldsymbol{\delta}_{3,V}$$

= $\tilde{\Sigma}_{11} + \tilde{\Sigma}_{33} - 2\tilde{X}_{13}$
= $\Sigma_{11} + \Sigma_{33} + 2k_1 - 2X_{13} - 2k_2$
= $\frac{10}{9} - 2 \cdot \frac{1}{3\alpha p(p-1)} - 2X_{13}.$

Now it is possible to compute X_{13} by means of Equation (20):

$$\begin{split} X_{13} &= \frac{1}{2\alpha(p-1)} W_{1\star} \left(\alpha(p-2)\mathbf{\Lambda} + \sqrt{\mathbf{I}_n - \mathbf{J}_n - 2\alpha(p-2)\mathbf{\Lambda} + \alpha^2 p^2 \mathbf{\Lambda}^2} - \mathbf{I}_n - \mathbf{J}_n \right) W_{\star 3}^{\top} \\ &= \frac{1}{2\alpha(p-1)} \sum_{i=1}^3 W_{1i} W_{3i} \left(\alpha(p-2)\mathbf{\Lambda} + \sqrt{\mathbf{I}_n - \mathbf{J}_n - 2\alpha(p-2)\mathbf{\Lambda} + \alpha^2 p^2 \mathbf{\Lambda}^2} - \mathbf{I}_n - \mathbf{J}_n \right)_{ii} \\ &= \frac{1}{2\alpha(p-1)} \left(-\frac{1}{2} \left(\alpha(p-2) \cdot 1 + \sqrt{1 - 2\alpha(p-2) \cdot 1 + \alpha^2 p^2 \cdot 1^2} - 1 \right) \right. \\ &+ \frac{1}{6} \left(\alpha(p-2) \cdot \frac{1}{3} + \sqrt{1 - 2\alpha(p-2) \cdot \frac{1}{3} + \alpha^2 p^2 \cdot \frac{1}{3^2}} - 1 \right) \\ &+ \frac{1}{3} \left(\alpha(p-2) \cdot 0 + \sqrt{0 - 2\alpha(p-2) \cdot 0 + \alpha^2 p^2 \cdot 0^2} - 2 \right) \right) \\ &= \frac{1}{2\alpha(p-1)} \left(-\frac{1}{2} \sqrt{1 - 2\alpha(p-2) + \alpha^2 p^2} \\ &+ \frac{1}{6} \sqrt{1 - \frac{2}{3}\alpha(p-2) + \frac{\alpha^2 p^2}{9}} - \frac{4}{9} \alpha(p-2) - \frac{1}{3} \right). \end{split}$$

With p = 3, the maximum discrepancy $d(\alpha) := D_{11}(v_1, v_3) - D_{12}(v_1, v_3)$ is

reached at $\alpha^{\star} = 1.32092$. In such a case, $X_{13} = -0.48598$ and thus:

$$D_{12}(v_1, v_3) = \frac{10}{9} - 2 \cdot \frac{1}{18 \cdot 1.32092} - 2 \cdot (-0.48598) = 1.99896 < 2 = D_{11}(v_1, v_3),$$

that is, $d(\alpha^*) = 0.00103$. Notice that for $\alpha := \hat{\alpha} = 0.97222$ and for $\alpha \to +\infty$, we achieve $d(\hat{\alpha}) = 0$, and for $0 < \alpha < \hat{\alpha}$ we have $d(\alpha) < 0$. One can obtain larger values for $d(\alpha)$ by taking $\alpha := \frac{1}{p}$ and letting $p \to +\infty$. In this setting, we have $X_{13} \to -\frac{1}{3}$ and $d(\alpha) \to \frac{2}{9}$.