On Explicit Estimations for the Bloch Eigenvalues of the One-dimensional Schrödinger Operator and the Kronig-Penney Model

Cemile Nur^{*} and Oktay Veliev^{**} Department of Basic Sciences, Yalova University Yalova, Turkey. e-mail: cnur@yalova.edu.tr^{*} Department of Mechanical Engineering, Dogus University Istanbul, Turkey. e-mail: oveliev@dogus.edu.tr^{**}

Abstract

In this paper, we consider the small and large eigenvalues of the one-dimensional Schrödinger operator L(q) with a periodic, real and locally integrable potential q. First we explicitly write out the first and second terms of the asymptotic formulas for the large periodic and antiperiodic eigenvalues and illustrate these formulas for the Kronig-Penney model. Then we give estimates for the small periodic and antiperiodic eigenvalues and for the length of the first gaps in the case of the Kronig-Penney model. Moreover, we give error estimations and present a numerical example.

Key Words: One-dimensional Schrödinger operator, Periodic potential, Kronig-Penney model. AMS Mathematics Subject Classification: 34L20, 47E05.

1 Introduction and Preliminary Facts

In the present paper, we investigate the spectrum of the operator L(q) generated in $L_2(-\infty,\infty)$ by the differential expression

$$l(y) = -y'' + qy, \tag{1}$$

where q is 1-periodic integrable on [0, 1] and a real-valued potential. Without loss of generality, it is assumed that

$$\int_{0}^{1} q(x)dx = 0.$$
 (2)

It is well known that (see for example [1, 3, 4, 5, 8, 10, 12]) the spectrum $\sigma(L(q))$ of L(q) is the union of the spectra $\sigma(L_t(q))$ of the operators $L_t(q)$ for $t \in (-\pi, \pi]$ generated in $L_2[0, 1]$ by (1) and the boundary conditions

$$y(1) = e^{it}y(0), \qquad y'(1) = e^{it}y'(0).$$
 (3)

Moreover, $\sigma(L(q))$ consists of the closed intervals whose end points are the eigenvalues of $L_t(q)$ for $t = 0, \pi$ (see (3)). Therefore, to study the spectrum of the self adjoint operator L(q) it is enough to investigate the eigenvalues of $L_0(q)$ and $L_{\pi}(q)$, which are called the periodic and antiperiodic eigenvalues, respectively. The first periodic eigenvalue is denoted by λ_0 . The other eigenvalues of $L_0(q)$ and the eigenvalues of $L_{\pi}(q)$ are denoted by $\lambda_{n,j}$ and $\mu_{n,j}$, respectively, for $n \in \mathbb{N}$, j = 1, 2, where \mathbb{N} is the set of positive integers. Without loss of generality, it is assumed that $\lambda_{n,1} \leq \lambda_{n,2}$ and $\mu_{n,1} \leq \mu_{n,2}$, for $n \in \mathbb{N}$. It is known that (see [4]), the spectrum of the Schrödinger operator L(q) consists of the real intervals

$$\Gamma_1 := [\lambda_0, \mu_{1,1}], \quad \Gamma_2 := [\mu_{1,2}, \lambda_{1,1}], \quad \Gamma_3 := [\lambda_{1,2}, \mu_{2,1}], \quad \Gamma_4 := [\mu_{2,2}, \lambda_{2,1}], \dots$$

The bands $\Gamma_1, \Gamma_2, \ldots$ of the spectrum $\sigma(L(q))$ of L(q) are separated by the gaps

$$\Delta_1 := (\mu_{1,1}, \mu_{1,2}), \quad \Delta_2 := (\lambda_{1,1}, \lambda_{1,2}), \quad \Delta_3 := (\mu_{2,1}, \mu_{2,2}), \quad \Delta_4 := (\lambda_{2,1}, \lambda_{2,2}), \dots$$

For this reason, the investigation of the periodic and antiperiodic eigenvalues is of great importance. In

[2] and [17], Veliev investigated the asymptotic behavior of large periodic and antiperiodic eigenvalues and obtained asymptotic formulas for the lengths of the gaps in the spectrum.

In this paper, we obtain more explicit formulas for the large eigenvalues of $L_0(q)$ and $L_{\pi}(q)$ and estimate the small eigenvalues of these operators. Moreover, we take the Kronig-Penney model as an example and illustrate asymptotic formulas and estimations with this example. For this we use some formulas from [2, 13, 15, 16, 17]. For the independent reading of this paper, first of all, we list the formulas of these papers that are essentially used here.

In [2], we established the following equalities

$$(\lambda_{n,j} - (2\pi n)^2 - A_m(\lambda_{n,j}))(\Psi_{n,j}(x), e^{i2\pi nx}) - (q_{2n} + B_m(\lambda_{n,j}))(\Psi_{n,j}(x), e^{-i2\pi nx}) = R_m(\lambda_{n,j}),$$
(4)

where

$$A_{m}(\lambda_{n,j}) = \sum_{k=1}^{m} a_{k}(\lambda_{n,j}), \quad B_{m}(\lambda_{n,j}) = \sum_{k=1}^{m} b_{k}(\lambda_{n,j}),$$

$$A_{m}(\lambda_{n,j}) = \sum_{k=1}^{m} a_{k}(\lambda_{n,j}), \quad B_{m}(\lambda_{n,j}) = \sum_{k=1}^{m} b_{k}(\lambda_{n,j}), \quad B_{m}(\lambda_{n,j}) = \sum_{k=1$$

$$a_k(\lambda_{n,j}) = \sum_{n_1, n_2, \dots, n_k} \frac{1}{\prod_{s=1}^k (\lambda_{n,j} - (2\pi(n - n_1 - n_2 - \dots - n_s))^2)},$$
(5)

$$b_k(\lambda_{n,j}) = \sum_{n_1, n_2, \dots, n_k} \frac{q_{n_1} q_{n_2} \dots q_{n_k} q_{2n-n_1-n_2-\dots-n_k}}{\prod_{s=1}^k (\lambda_{n,j} - (2\pi (n-n_1-n_2-\dots-n_s))^2)},$$
(6)

$$R_m(\lambda_{n,j}) = \sum_{n_1, n_2, \dots, n_{m+1}} \frac{q_{n_1} q_{n_2} \dots q_{n_m} q_{n_{m+1}}(q(x) \Psi_{n,j}(x), e^{i2\pi(n-n_1-\dots-n_{m+1})x})}{\prod_{s=1}^{m+1} (\lambda_{n,j} - (2\pi(n-n_1-n_2-\dots-n_s))^2)}.$$
(7)

The summations in these formulas are taken over the indices satisfying the conditions

$$n_s \neq 0, \quad n_1 + n_2 + \dots + n_s \neq 0, 2n,$$

for s = 1, 2, ..., m + 1. Moreover, it was proved that

$$a_k(\lambda_{n,j}) = O((\frac{\ln|n|}{n})^k), \qquad b_k(\lambda_{n,j}) = O((\frac{\ln|n|}{n})^k)$$

for $k \geq 1$ and

$$R_m(\lambda_{n,j}) = O((\frac{\ln |n|}{n})^{m+1}).$$

In [17], the terms $a_1(\lambda_{n,j})$ and $a_2(\lambda_{n,j})$ were investigated in detail and the following estimations were proved:

$$a_1(\lambda_{n,j}) = a_1((2\pi n)^2) + o(n^{-2}) = o(n^{-1})$$
(8)

and

$$a_2(\lambda_{n,j}) = a_2((2\pi n)^2) + o(n^{-3}\ln n),$$

where $a_1((2\pi n)^2)$ and $a_2((2\pi n)^2)$ are obtained from $a_1(\lambda_{n,j})$ and $a_2(\lambda_{n,j})$ by changing $\lambda_{n,j}$ to $(2\pi n)^2$ in the corresponding formulas and $q \in L_1[0, 1]$.

Similarly, in [13], we proved that the following relations are valid:

$$b_1(\lambda_{n,j}) = b_{1,n}((2\pi n)^2) + o(n^{-2}), \qquad b_2(\lambda_{n,j}) = o(n^{-2}), b_{1,n}((2\pi n)^2) = 2Q_0Q_{2n} - S_{2n},$$
(9)

where $b_1((2\pi n)^2)$ is obtained from $b_1(\lambda_{n,j})$ by changing $\lambda_{n,j}$ to $(2\pi n)^2$ in (6), $Q_k = (Q, e^{2\pi i kx})$ and $S_k = (S, e^{2\pi i kx})$ are the Fourier coefficients of the following functions:

$$Q(x) = \int_0^x q(t) dt, \qquad S(x) = Q^2(x)$$
(10)

(see Lemma 6 of [13]).

Using (8) and these relations, in [17], the following asymptotic formulas were obtained. If there exists a positive number ε such that

$$|q_k| \ge \frac{\varepsilon}{k},\tag{11}$$

and

$$|q_k - S_k + 2Q_0 Q_k| \ge \varepsilon k^{-2},\tag{12}$$

for k = 2n > N, where N is a sufficiently large positive integer, then the periodic eigenvalues $\lambda_{n,j}$, for n > N and j = 1, 2, are simple and satisfy the asymptotic formulas

$$\lambda_{n,j} = (2\pi n)^2 + (-1)^j |q_{2n}| + o(n^{-1})$$
(13)

and

$$\lambda_{n,j} = (2\pi n)^2 + a_1((2\pi n)^2) + a_2((2\pi n)^2) + (-1)^j |q_{2n} - S_{2n} + 2Q_0Q_{2n}| + o(n^{-2}), \tag{14}$$

respectively (see pages 66 and 70 in [17]). Similarly, the antiperiodic eigenvalues $\mu_{n,j}$ satisfy the formulas obtained from (13) and (14) by replacing 2n with 2n-1 if (11) and (12) are satisfied for k = 2n-1 > N. Moreover, from these formulas we obtained the following formulas for the lengths $|\Delta_k|$ of the gaps Δ_k :

$$|\Delta_k| = 2 |q_k| + o(k^{-1}) \tag{15}$$

and

$$\Delta_k | = 2 |q_k - S_k + 2Q_0 Q_k| + o(k^{-2})$$
(16)

without using the conditions (11) and (12) (see page 71 in [17]).

The superiority of formula (15) over the well-known Titchmarsh formula

$$|\Delta_{2n}| = 2|q_{2n}| + O(n^{-1}) \tag{17}$$

(see [14, Chapter 21], [1, Chapter 3], and their references) for any potential $q \in L_1[0, 1]$ can be explained as follows. Generally speaking, certain Fourier coefficients q_{2n} of many functions in $L_1[0, 1]$ can be of order n^{-1} in the sense that $q_{2n} = O(n^{-1})$ and $n^{-1} = O(q_{2n})$ when n belongs to some subset of N (e.g., when the potential q exhibits a jump discontinuity, as occurs in the well known Kronig-Penney model). In this context, equation (15) with the error term $o(n^{-1})$ explicitly isolates the term $2|q_{2n}|$ from the error $o(n^{-1})$, whereas equation (17) does not. In this sense, $2|q_{2n}|$ can be seen as the first term of the asymptotic formula for the gaps in $\sigma(L(q))$ with the potential $q \in L_1[0, 1]$. Similarly, (16) gives us the second term $2|q_{2n} - S_{2n} + 2Q_0Q_{2n}| - 2|q_{2n}|$ of the asymptotic formula for the gaps in $\sigma(L(q))$.

Note that, in [17], we obtained asymptotic formulas with the error term $O\left(\left(\frac{\ln n}{n}\right)^k\right)$ for the *n*th periodic and antiperiodic eigenvalues and the *n*th gaps in the spectrum of the operator L(q) with $q \in L_1[0, 1]$, where $k = 3, 4, \ldots$ Many studies have been devoted to such asymptotic formulas for differentiable potentials (see [1, Chap. 3], [9, Chap. 1] and [11, Chap. 2] and their references), whereas we obtain asymptotic formulas of arbitrary order for the potential q from $L_1[0, 1]$. In Section 2, we improve the asymptotic formulas obtained in [17] and give explicit asymptotic formulas for the large eigenvalues.

In Section 3, we consider the small periodic and antiperiodic eigenvalues of the Schrödinger operator L(q) and estimate the length of the gaps Δ_n for the small values of n in the Kronig-Penney model. We note that, asymptotic formulas obtained in Section 2 cannot be used for estimating the small eigenvalues. To give estimates for the small eigenvalues, we use completely different methods than those of Section 2. In Section 3, we find conditions on the Kronig-Penney potential for which the iteration formula (4) is also valid for the small eigenvalues.

Finally, note that the Kronig-Penney model is a simplified approach to understanding the band structure of materials, it provides a foundational understanding of band theory and is a stepping stone to more complex models in solid-state physics. This model is particularly important in explaining the quantum mechanical basis for electrical conduction in solids and the properties of semiconductors.

2 Asymptotic Formulas for the Large Eigenvalues

To obtain explicit formulas for the large eigenvalues from (14), we use the following lemmas.

Lemma 1 If $q \in L_1[0,1]$, then the following formula holds:

$$a_1((2\pi n)^2) = \frac{|q_{2n}|^2}{-16\pi^2 n^2} + D(2n),$$
(18)

where

$$D(k) =: \frac{i}{2\pi k} \int_0^1 q(x)(Q(x,k) - Q_{k,0}))e^{-i2\pi kx} dx,$$
$$Q(x,k) = \int_0^x q(t)e^{i2\pi kt} dt - q_{-k}x, \ Q_{k,0} = \int_0^1 Q(x,k)dx.$$

1

Proof. Since $q_{-k} = \overline{q_k}$, $q_0 = 0$, from the definition of $a_1((2\pi n)^2)$ (see introduction), we obtain

$$a_1((2\pi n)^2) = \sum_{\substack{k=-\infty\\k\neq 0,2n}}^{\infty} \frac{|q_k|^2}{2\pi k(4\pi n - 2\pi k)}.$$

Using the equality

$$\frac{1}{2\pi k(4\pi n - 2\pi k)} = \frac{1}{4\pi n} \left(\frac{1}{2\pi k} + \frac{1}{4\pi n - 2\pi k} \right),\tag{19}$$

we see that

$$a_1((2\pi n)^2) = \frac{1}{4\pi n} \sum_{\substack{k=-\infty\\k\neq 0,2n}}^{\infty} \frac{|q_k|^2}{2\pi k} + \frac{1}{4\pi n} \sum_{\substack{k=-\infty\\k\neq 0,2n}}^{\infty} \frac{|q_k|^2}{(4\pi n - 2\pi k)}.$$
 (20)

Since

$$\frac{|q_k|^2}{2\pi k} + \frac{|q_{-k}|^2}{2\pi(-k)} = 0$$

we have the following equality for the first term on the right side of (20):

$$\frac{1}{4\pi n} \sum_{\substack{k=-\infty\\k\neq 0,2n}}^{\infty} \frac{|q_k|^2}{2\pi k} = \frac{|q_{2n}|^2}{-16\pi^2 n^2}.$$
(21)

Now let us estimate the second term on the right side of (20). For this, we consider the Fourier coefficients of Q(x, 2n). It is clear that

$$Q(1,2n) = Q(0,2n) = 0, \quad Q'(x,2n) = q(x)e^{i4\pi nx} - q_{-2n}.$$

Therefore, using the integration by parts, we see that the Fourier coefficients $Q_{2n,2n-k} = (Q(x,2n), e^{i2\pi(2n-k)x})$ of Q(x,2n) with respect to the orthonormal basis $\{e^{2\pi ikx} : k \in \mathbb{Z}\}$, for $2n - k \neq 0$, are

$$Q_{2n,2n-k} = \frac{q_{-k}}{i(4\pi n - 2\pi k)}.$$

Moreover, $q_0 = 0$ according to (2). Thus, the Fourier decomposition of Q(x, 2n) has the form

$$Q(x,2n) = Q_{2n,0} + \sum_{\substack{k=-\infty\\k\neq 0,2n}}^{\infty} \frac{q_{-k}e^{i2\pi(2n-k)x}}{i(4\pi n - 2\pi k)}.$$
(22)

Thus, (18) follows from (20)-(22). The lemma is proved. \blacksquare

Now we consider $a_2((2\pi n)^2)$. It is clear that

$$a_2((2\pi n)^2) = \sum_{k,k+l\neq 0,2n} \frac{q_k q_l q_{-k-l}}{[(2\pi n)^2 - (2\pi (n-k))^2][(2\pi n)^2 - (2\pi (n-k-l))^2]}$$
$$= \sum_{k,l} \frac{q_k q_l q_{-k-l}}{2\pi k (4\pi n - 2\pi k)(2\pi k + 2\pi l)(4\pi n - 2\pi k - 2\pi l)}.$$

Using (19) and

$$\frac{1}{(k+l)(2n-k-l)} = \frac{1}{2n} \left(\frac{1}{k+l} + \frac{1}{2n-k-l} \right),$$

we obtain that

$$a_2((2\pi n)^2) = \frac{1}{16\pi^2 n^2} (I_1 + I_2 + I_3 + I_4),$$
(23)

where

$$I_1(q) = \sum_{k,l} \frac{q_k q_l q_{-k-l}}{2\pi k (2\pi k + 2\pi l)},$$

$$I_2(n,q) = \sum_{k,l} \frac{q_k q_l q_{-k-l}}{(4\pi n - 2\pi k) (2\pi k + 2\pi l)},$$

$$I_3(n,q) = \sum_{k,l} \frac{q_k q_l q_{-k-l}}{2\pi k (4\pi n - 2\pi k - 2\pi l)},$$

and

$$I_4(n,q) = \sum_{k,l} \frac{q_k q_l q_{-k-l}}{(4\pi n - 2\pi k)(4\pi n - 2\pi k - 2\pi l)}$$

Now we estimate $I_1(q)$ and $I_j(n,q)$, for j = 2, 3, 4, in the following lemmas.

Lemma 2 The equality $I_1(q) = 0$ holds.

Proof. We prove this lemma in two ways. First grouping the terms

$$\frac{q_k q_l q_{-k-l}}{4\pi^2 k(k+l)}, \quad \frac{q_l q_k q_{-k-l}}{4\pi^2 l(k+l)}, \quad \frac{q_k q_{-k-l} q_l}{4\pi^2 k(-l)}, \\ \frac{q_l q_{-k-l} q_k}{4\pi^2 l(-k)}, \quad \frac{q_{-k-l} q_l q_k}{4\pi^2 (-k-l)(-k)}, \quad \frac{q_{-k-l} q_k q_l}{4\pi^2 (-k-l)(-l)}$$

with the equal multiplicands,

$$q_k q_l q_{-k-l}, \quad q_l q_k q_{-k-l}, \quad q_k q_{-k-l} q_l, \\ q_l q_{-k-l} q_k, \quad q_{-k-l} q_l q_k, \quad q_{-k-l} q_k q_l$$

and using the equality

$$\frac{1}{k(k+l)} + \frac{1}{l(k+l)} = \frac{1}{kl}$$

we obtain the proof of the lemma.

The second way is the following. Using the integration by parts one can easily verify that the Fourier coefficients $Q_k = (Q, e^{2\pi i k x})$ of Q (see (10)) with respect to the orthonormal basis $\{e^{2\pi i k x} : k \in \mathbb{Z}\}$, for $k \neq 0$, are $\frac{q_k}{-2\pi k i}$. Therefore, we have

$$Q(x) - Q_0 = \sum_{k \neq 0} \frac{q_k}{-2\pi ki} e^{2\pi i kx}.$$
(24)

Using this and (2), it is easy to verify that

$$I_1 = \int_0^1 q(x)(Q(x) - Q_0)^2 \, dx.$$

On the other hand, $(Q(x) - Q_0)' = q(x)$, Q(1) = Q(0) = 0, and hence

$$\int_0^1 q(x)(Q(x) - Q_0)^k \, dx = \frac{1}{k+1}((Q(1) - Q_0)^{k+1} - (Q(0) - Q_0)^{k+1}) = 0, \tag{25}$$

for $k = 1, 2, \dots$ The lemma is proved.

Lemma 3 The equalities $I_j(n,q) = o(n^2)$ for j = 2, 3, 4 hold.

Proof. Using (22) and (24), we obtain

$$I_2(n,q) = \int_0^1 q(x) \left(Q(x) - Q_0\right) \left(Q(x,2n) - Q_{2n,0}\right) \, dx. \tag{26}$$

Formula (22) can be written in the form

$$Q(x,2n) - Q_{2n,0} = \sum_{\substack{l=-\infty\\l+k\neq 0,2n}}^{\infty} \frac{q_{-k-l}e^{i2\pi(2n-k-l)x}}{i2\pi(2n-k-l)}.$$
(27)

Using these notations, we obtain

$$I_3(n,q) = \int_0^1 q(x) \left(Q(x) - Q_0 \right) \left(Q(x,2n) - Q_{2n,0} \right) \, dx = I_2(n,q). \tag{28}$$

In the same way, we establish that

$$I_4(n,q) = \int_0^1 q(x) \left(Q(x,2n) - Q_{2n,0} \right)^2 \, dx.$$
⁽²⁹⁾

Now the proof of the lemma follows from (2), (25)-(29) and the following obvious statements: Q(x, 2n) converges uniformly, for $x \in [0, 1]$, to zero as $n \to \infty$ (see page 57 of [17]). The lemma is proved.

Now from (14), (18), (23) and Lemmas 1-3, we obtain the following asymptotic formulas for the periodic eigenvalues:

Theorem 1 If (12), for k = 2n, holds, then the eigenvalues $\lambda_{n,j}$, for n > N and j = 1, 2, are simple and satisfy the asymptotic formula

$$\lambda_{n,j} = (2\pi n)^2 + D(2n) + (-1)^j |q_{2n} - S_{2n} + 2Q_0 Q_{2n}| + o(n^{-2})$$
(30)

as $n \to \infty$, where D(2n) is defined in Lemma 1.

The proof of the corresponding results for the antiperiodic problem can be carried out in a similar way.

Theorem 2 If (12), for k = 2n - 1, holds, then the eigenvalues $\mu_{n,j}$ of the operator $L_{\pi}(q)$, for n > N and j = 1, 2, are simple and satisfy the asymptotic formula

$$\mu_{n,j} = (2\pi n - \pi)^2 + D(2n - 1) + (-1)^j |q_{2n-1} - S_{2n-1} + 2Q_0 Q_{2n-1}| + o(n^{-2}), \tag{31}$$

as $n \to \infty$.

Now, we take the Kronig-Penney model as an example and use this example to illustrate the obtained asymptotic formulas (30) and (31). The Kronig-Penney model has been studied in many works (see, for example, [7], [1, Chapter 3] and [14, Chapter 21]). In this case, the potential q(x) has the form

$$q(x) = \begin{cases} a & \text{if } x \in [0, c] \\ b & \text{if } x \in (c, d] \end{cases}$$

and q(x+d) = q(x), where $c \in (0, d)$. For simplicity of notation and without loss of generality, we assume that d = 1, a < b and (2) is satisfied. Then, we have

$$q(x) = \begin{cases} a & \text{if } x \in [0, c] \\ b & \text{if } x \in (c, 1], \end{cases}$$

$$(32)$$

where a < 0 < b and

$$ac + (1 - c)b = 0. (33)$$

To estimate the second term of the asymptotic formulas for the periodic eigenvalues when q is defined by formula (32), we need to calculate the term D(2n) in (30), since the terms q_k , Q_k , S_k were calculated in [17] (see pages 76-78 in [17]) and the following equalities were obtained:

$$q_k = \frac{a-b}{2\pi ki} (1 - e^{-2\pi kic}), \tag{34}$$

$$Q_k = \frac{q_k}{2\pi ki} = \frac{a-b}{(2\pi k)^2} (e^{-2\pi kic} - 1), \qquad Q_0 = \frac{1}{2}b(c-1), \tag{35}$$

$$S_{k} = \frac{a^{2}}{\pi ki} \left(\frac{e^{-2\pi ikc} - 1}{(2\pi k)^{2}} - \frac{ce^{-2\pi ikc}}{2\pi ik} \right) + \frac{b^{2}}{\pi ki} \left(\frac{1 - e^{-2\pi ikc}}{(2\pi k)^{2}} + \frac{ce^{-2\pi ikc} - 1}{2\pi ik} \right) - \frac{b^{2}}{\pi ki} \left(\frac{e^{-2\pi ikc} - 1}{2\pi ik} \right).$$
(36)

Using the definition of Q(x,k) (see Lemma 1), by direct calculations we obtain

$$Q(x,k) = \begin{cases} \frac{a}{i2\pi k} (e^{i2\pi kx} - 1) - q_{-k}x & \text{if } x \in [0,c] \\ \frac{b}{i2\pi k} (e^{i2\pi kx} - 1) - q_{-k}x + q_{-k} & \text{if } x \in (c,1]. \end{cases}$$
(37)

Therefore, we have

$$Q_{k,0} = \int_{0}^{c} \left(\frac{a}{i2\pi k} (e^{i2\pi kx} - 1) - q_{-k}x\right) dx + \int_{c}^{1} \left(\frac{b}{i2\pi k} (e^{i2\pi kx} - 1) - q_{-k}x + q_{-k}\right) dx$$

$$= \frac{(b-a)(e^{i2\pi kc} - 1)}{4\pi^{2}k^{2}} + q_{-k}(\frac{1}{2} - c) = \frac{(b-a)(e^{i2\pi kc} - 1)}{4\pi^{2}k^{2}} + \frac{(a-b)(c-\frac{1}{2})}{2\pi ki}(1 - e^{2\pi kic})$$

$$= \frac{(b-a)(e^{i2\pi kc} - 1)}{4\pi^{2}k^{2}} + \frac{(a+b)(e^{i2\pi kc} - 1)}{4\pi ki}$$
(38)

and

$$D(k) = \frac{i}{2\pi k} \int_0^1 q(x)Q(x,k)e^{-i2\pi kx} dx - \frac{iQ_{k,0}}{2\pi k} \int_0^1 q(x)e^{-i2\pi kx} dx$$
$$= \frac{i}{2\pi k} \left(\int_0^c q(x)Q(x,k)e^{-i2\pi kx} dx + \int_c^1 q(x)Q(x,k)e^{-i2\pi kx} dx \right) - \frac{iQ_{k,0}q_k}{2\pi k}.$$
(39)

Using (32), (37), and the integration by parts in the last integrals give

$$\int_{0}^{c} q(x)Q(x,k)e^{-i2\pi kx} dx = \int_{0}^{c} a \left(\frac{a}{i2\pi k}(e^{i2\pi kx}-1)-q_{-k}x\right)e^{-i2\pi kx} dx$$
$$= \frac{a^{2}}{i2\pi k} \int_{0}^{c} (1-e^{-i2\pi kx}) dx - aq_{-k} \int_{0}^{c} xe^{-i2\pi kx} dx$$
$$= \frac{a^{2}}{i2\pi k} \left(c + \frac{e^{-i2\pi kc}-1}{i2\pi k}\right) + \frac{aq_{-k}}{i2\pi k} \left(ce^{-i2\pi kc} + \frac{e^{-i2\pi kc}-1}{i2\pi k}\right)$$
(40)

and

$$\int_{c}^{1} q(x)Q(x,k)e^{-i2\pi kx} dx = \int_{c}^{1} b\left(\frac{b}{i2\pi k}(e^{i2\pi kx}-1)-q_{-k}x+q_{-k}\right)e^{-i2\pi kx} dx$$
$$= \frac{b^{2}}{i2\pi k}\int_{c}^{1}(1-e^{-i2\pi kx}) dx - bq_{-k}\int_{c}^{1}xe^{-i2\pi kx} dx + bq_{-k}\int_{c}^{1}e^{-i2\pi kx} dx$$
$$= \frac{b^{2}}{i2\pi k}\left(1-c-\frac{e^{-i2\pi kc}-1}{i2\pi k}\right) + \frac{bq_{-k}}{i2\pi k}\left(1-ce^{-i2\pi kc}-\frac{e^{-i2\pi kc}-1}{i2\pi k}\right) + \frac{bq_{-k}}{i2\pi k}(e^{-i2\pi kc}-1). \quad (41)$$

Using (38), (40), (41) and (33) in (39) and doing some simplification, we obtain

$$D(k) = \frac{-ab}{4\pi^2 k^2} + O(\frac{1}{k^3}).$$

Now using (34)-(36) and (33), we obtain

$$q_k - S_k + 2Q_0Q_k = \frac{a-b}{2\pi ki}(1 - e^{-2\pi kic}) + \frac{ba}{(2\pi k)^2}(e^{-2\pi ikc} + 1) + O(k^{-3})$$
$$= e^{-\pi kic} \left(\frac{a-b}{\pi k}\sin(\pi kc) + \frac{ba}{2(\pi k)^2}\cos(\pi kc)\right) + O(k^{-3}).$$
(42)

In Theorem 2.1.11 of [17], we proved that if $c \in \mathbb{Q}$, then $|q_k - S_k + 2Q_0Q_k| > c_0k^{-2}$, for some positive constant c_0 , where \mathbb{Q} is the set of rational numbers. Therefore the following theorems follow from Theorem 1 and Theorem 2, respectively.

Theorem 3 If $c \in \mathbb{Q}$, then the periodic eigenvalues $\lambda_{n,j}$, for n > N and j = 1, 2, are simple and satisfy the asymptotic formula

$$\lambda_{n,j} = (2\pi n)^2 + \frac{-ab}{16\pi^2 n^2} + (-1)^j |q_{2n} - S_{2n} + 2Q_0 Q_{2n}| + o(n^{-2})$$
(43)

as $n \to \infty$.

and

Theorem 4 If $c \in \mathbb{Q}$, then the antiperiodic eigenvalues $\mu_{n,j}$ of the operator $L_{\pi}(q)$, for n > N and j = 1, 2, are simple and satisfy the asymptotic formula

$$\mu_{n,j} = (2\pi n - \pi)^2 + \frac{-ab}{4\pi^2(2n-1)^2} + (-1)^j |q_{2n-1} - S_{2n-1} + 2Q_0Q_{2n-1}| + o(n^{-2})$$
(44)

as $n \to \infty$.

Now, let us consider the general case $c \in \mathbb{R}$. By (42), we have

$$|q_k - S_k + 2Q_0Q_k| = |\alpha\sin(\pi kc) + \beta\cos(\pi kc)| + O(\frac{1}{k^3}),$$

where $\alpha = \frac{a-b}{\pi k}$ and $\beta = \frac{ab}{2\pi^2 k^2}$. There exists $\theta \in [0, 2\pi)$ such that

$$\cos\theta = \frac{\alpha}{\sqrt{\alpha^2 + \beta^2}}, \qquad \sin\theta = \frac{\beta}{\sqrt{\alpha^2 + \beta^2}},$$

where $\sqrt{\alpha^2 + \beta^2} = \frac{b-a}{\pi k} + O(\frac{1}{k^3})$. Then, we have

$$|q_k - S_k + 2Q_0Q_k| = \frac{b-a}{\pi k} |\sin(\pi kc + \theta)| + O(\frac{1}{k^3}),$$

Therefore, if there exists $\varepsilon > 0$ such that the inequality

$$\sin(\pi kc + \theta)| > \frac{\varepsilon}{k} \tag{45}$$

is satisfied, then (12) holds. Thus the following theorems follow from Theorem 1 and Theorem 2, respectively.

Theorem 5 If (45) is satisfied for k = 2n, then the periodic eigenvalues $\lambda_{n,j}$, for n > N and j = 1, 2, are simple and satisfy asymptotic formula (43).

and

Theorem 6 If (45) is satisfied for k = 2n - 1, then the antiperiodic eigenvalues $\mu_{n,j}$ of the operator $L_{\pi}(q)$, for n > N and j = 1, 2, are simple and satisfy asymptotic formula (44).

3 On the Small Eigenvalues

In this section, we consider the small periodic and antiperiodic eigenvalues of the Schrödinger operator L(q) with potential (32). We shall focus on the operator $L_0(q)$ with potential (32). The investigation of $L_{\pi}(q)$ is similar.

It is known that [6]

$$(2\pi n)^2 - M \le \lambda_{n,j} \le (2\pi n)^2 + M,$$

for $n \ge 1$, where $M = \max\{|a|, b\}$. Besides, $|\lambda_0| \le M$. If $k \ne \pm n$, then

$$\begin{aligned} |\lambda_{n,j} - (2\pi k)^2| &\ge |(2\pi n)^2 - (2\pi k)^2| - M = 4\pi^2 |n-k| |n+k| - M \\ &\ge 4\pi^2 (2n-1) - M, \end{aligned}$$
(46)

for $n \ge 1$ and under the assumption $M \le \frac{4\pi^2(2n-1)}{3}$. If n = 1, we have $|\lambda_{1,j}| \le 4\pi^2 + M$ and

$$|\lambda_{1,j} - (2\pi k)^2| \ge ||\lambda_{1,j}| - (2\pi k)^2| \ge 16\pi^2 - |\lambda_{1,j}| \ge 12\pi^2 - M,$$

for $|k| \ge 2$. Moreover, if $n \ge 2$, we have $|\lambda_{n,j}| \ge |\lambda_{2,j}| \ge 16\pi^2 - M$ and

$$|\lambda_{n,j} - (2\pi k)^2| \ge ||\lambda_{2,j}| - (2\pi k)^2| \ge |\lambda_{2,j}| - 4\pi^2 \ge 12\pi^2 - M,$$

for $k \neq \pm n$. The analogous inequalities can be written for the antiperiodic eigenvalues, from the inequalities

$$(2n-1)^2 \pi^2 - M \le \mu_{n,j} \le (2n-1)^2 \pi^2 + M$$

for $n \ge 2$. If $k \ne \pm n$, then

$$|\mu_{n,j} - (2k-1)^2 \pi^2| \ge |(2n-1)^2 \pi^2 - (2k-1)^2 \pi^2| - M = 4\pi^2 |n-k| |n+k-1| - M$$

$$\ge 4\pi^2 (2n-2) - M, \tag{47}$$

for $n \ge 2$, under the assumption $M \le \frac{8\pi^2(n-1)}{3}$. If n = 1, we have $|\mu_{1,j}| \le \pi^2 + M$ and

$$|\mu_{1,j} - (2k-1)^2 \pi^2| \ge ||\mu_{1,j}| - (2k-1)^2 \pi^2| \ge 9\pi^2 - |\mu_{1,j}| \ge 8\pi^2 - M,$$

for $|k| \ge 2$, and we assume $M \le 8\pi^2/3$, for n = 1. Besides, if $n \ge 2$, we have $|\mu_{n,j}| \ge |\mu_{2,j}| \ge 9\pi^2 - M$ and

$$|\mu_{n,j} - (2k-1)^2 \pi^2| \ge ||\mu_{2,j}| - (2k-1)^2 \pi^2| \ge |\mu_{2,j}| - \pi^2 \ge 8\pi^2 - M,$$

for $k \neq \pm n$.

We note that, the iteration formula (4) was obtained in [2] and has also been used in the previous section for the large eigenvalues to obtain subtle asymptotic formulas. In this section, we find conditions

on potential (32) for which the iteration formula (4) is also valid for the small eigenvalues, as m tends to infinity. We also note that, it is not easy to give such conditions, the calculations are very technical and long. Since the potential q is real, we have the following:

$$q_{-k} = \overline{q_k}, \qquad v_{n,j} = \overline{u_{n,j}},\tag{48}$$

where $u_{n,j} = (\Psi_{n,j}, e^{i2\pi nx})$ and $v_{n,j} = (\Psi_{n,j}, e^{-i2\pi nx})$ (see (2.1.83) in [17]). Now, in order to give the main results of this section, we prove the following lemmas. Without loss of generality, we assume that $\Psi_{n,j}(x)$ is a normalized eigenfunction corresponding to the eigenvalue $\lambda_{n,j}$. First, we consider the case $n \geq 1.$

Lemma 4 If $M \leq \frac{4\pi^2(2n-1)}{3}$, where $M = \max\{|a|, b\}$, then the statements (a) $\lim_{m \to \infty} R_m(\lambda_{n,j}) = 0$ (b) $|u_{n,j}| > 0$ are valid, where $R_m(\lambda_{n,j})$ is defined by (7).

Proof. (a) Considering the greatest summands of $R_{2m+1}(\lambda_{n,j})$ in absolute value and by (46), we obtain for n = 1,

$$\begin{aligned} |R_{2m+1}(\lambda_{1,j})| &< \frac{4^m |q_1| |q_{-2}|^m |q_2|^m M}{|\lambda_{1,j}|^{m+1} |\lambda_{1,j} - 16\pi^2|^m} + \frac{4^m |q_{-1}| |q_{-2}|^m |q_2|^m M}{|\lambda_{1,j}|^m |\lambda_{1,j} - 16\pi^2|^{m+1}} + \frac{4^{2m} |q_{-1}|^{m+1} |q_1|^m M}{|\lambda_{1,j} - 16\pi^2|^{m+1} |\lambda_{1,j} - 36\pi^2|^m} \\ &< \frac{4^m (b-a)^{2m+1} M}{2^{2m} \pi^{2m+1} (4\pi^2 - M)^{m+1} (12\pi^2 - M)^m} + \frac{4^m (b-a)^{2m+1} M}{2^{2m} \pi^{2m+1} (4\pi^2 - M)^m (12\pi^2 - M)^{m+1}} \\ &+ \frac{4^{2m} (b-a)^{2m+1} M}{\pi^{2m+1} (12\pi^2 - M)^{m+1} (32\pi^2 - M)^m} < \frac{5\pi}{3} (\frac{1}{4\pi^2})^m + \frac{\pi}{3} (\frac{4}{23\pi^2})^m, \end{aligned}$$

for n = 2,

$$\begin{split} |R_{2m+1}(\lambda_{2,j})| &< \frac{4^m |q_1| |q_{-2}|^m |q_2|^m M}{|\lambda_{2,j} - 4\pi^2|^{2m+1}} + \frac{4^m |q_2| |q_{-1}|^m |q_1|^m M}{|\lambda_{2,j} - 4\pi^2|^m |\lambda_{2,j}|^{m+1}} + \frac{4^{2m} |q_{-1}| |q_{-2}|^m |q_2|^m M}{|\lambda_{2,j} - 4\pi^2|^{m+1} |\lambda_{2,j} - 4\pi^2|^{m+1} |\lambda_{2,j} - 36\pi^2|^m} \\ &< \frac{4^m (b-a)^{2m+1} M}{2^{2m} \pi^{2m+1} (12\pi^2 - M)^{2m+1}} + \frac{4^m (b-a)^{2m+1} M}{2^{2m} \pi^{2m+1} (12\pi^2 - M)^m (16\pi^2 - M)^{m+1}} \\ &+ \frac{4^{2m} (b-a)^{2m+1} M}{\pi^{2m+1} (12\pi^2 - M)^{m+1} (20\pi^2 - M)^m} < 4\pi (\frac{1}{\pi^2})^m + \frac{8\pi}{3} (\frac{2}{3\pi^2})^m + 4\pi (\frac{8}{\pi^2})^m, \end{split}$$

and in general, we have $|R_{2m+1}(\lambda_{n,j})| < dr^m$, for $n \geq 3$, for some constant d > 0 and 0 < r < 1.

Therefore, $\lim_{m\to\infty} R_m(\lambda_{n,j}) = 0$. Similarly, we prove that $\lim_{m\to\infty} R_m^*(\lambda_{n,j}) = 0$. (b) Suppose the contrary, $u_{n,j} = 0$. Since the system of root functions $\{e^{2\pi i k x} : k \in \mathbb{Z}\}$ of $L_0(0)$ forms an orthonormal basis for $L_2[0,1]$ and by (48), we have the decomposition

$$\Psi_{n,j} = u_{n,j}e^{i2\pi nx} + \overline{u_{n,j}}e^{-i2\pi nx} + \sum_{k\in\mathbb{Z}, k\neq\pm n} \left(\Psi_{n,j}, e^{i2\pi kx}\right)e^{i2\pi kx}$$

for the normalized eigenfunction $\Psi_{n,j}$ corresponding to the eigenvalue $\lambda_{n,j}$ of $L_0(q)$. By Parseval's equality, we obtain

$$\sum_{k \in \mathbb{Z}, k \neq \pm n} |(\Psi_{n,j}, e^{i2\pi kx})|^2 = 1.$$
(49)

First, we consider the case n = 1. Using the relation

$$(\lambda_{N,j} - (2\pi n)^2)(\Psi_{N,j}, e^{i2\pi nx}) = (q\Psi_{N,j}, e^{i2\pi nx}),$$

(46), and the Bessel inequality, we obtain

$$\sum_{k \in \mathbb{Z}, k \neq \pm 1} |(\Psi_{1,j}, e^{i2\pi kx})|^2 = \sum_{k \in \mathbb{Z}, k \neq \pm 1} \frac{|(q\Psi_{1,j}, e^{i2\pi kx})|^2}{|\lambda_{1,j} - (2\pi k)^2|^2} \le \frac{1}{(4\pi^2 - M)^2} \sum_{k \in \mathbb{Z}, k \neq \pm 1} |(q\Psi_{1,j}, e^{i2\pi kx})|^2 < \frac{M^2}{(4\pi^2 - M)^2} < 1,$$

and in the case $n \geq 2$, we have

$$\sum_{k \in \mathbb{Z}, k \neq \pm n} |(\Psi_{n,j}, e^{i2\pi kx})|^2 = \sum_{k \in \mathbb{Z}, k \neq \pm n} \frac{|(q\Psi_{n,j}, e^{i2\pi kx})|^2}{|\lambda_{n,j} - (2\pi k)^2|^2} \le \frac{1}{(4\pi^2(2n-1) - M)^2} \sum_{k \in \mathbb{Z}, k \neq \pm n} |(q\Psi_{n,j}, e^{i2\pi kx})|^2 < \frac{M^2}{(4\pi^2(2n-1) - M)^2} < 1,$$

which contradicts (49) and completes the proof. \blacksquare

Now, we consider the case n = 0.

Lemma 5 If $M \le 4\pi^2/3$, then the statements (a) $\lim_{m\to\infty} R_m(\lambda_0) = 0$, (b) $|(\Psi_0, 1)| > 0$ are valid. **Proof.** The proof is similar to the proof of Lemma 1.

Before stating the main results, we write the following relations:

$$a_1((2\pi n)^2) = \frac{-ab}{16\pi^2 n^2} + \frac{(b^2 - a^2)\sin(4\pi nc)}{64\pi^3 n^3} + \frac{3(b-a)^2(\cos(4\pi nc) - 1)}{128\pi^4 n^4}$$
(50)

and

$$q_{2n} + b_1((2\pi n)^2) = q_{2n} + 2Q_0Q_{2n} - S_{2n}$$

= $\frac{(a-b)(1-e^{-i4\pi nc})}{4\pi ni} + \frac{ab(1+e^{-i4\pi nc})}{16\pi^2 n^2} + \frac{(a^2-b^2)(1-e^{-i4\pi nc})}{32\pi^3 n^3 i},$ (51)

which follow from (18), (39), (33) and (9), (34)-(36), (33), respectively, by direct calculations. Now we state the following interesting remark:

Remark 1 We have the formulas $q_{-k} = e^{i2\pi kc}q_k$ and $q_{-k} = \overline{q_k}$. These give, respectively,

$$\arg q_{-k} = \arg q_k + 2\pi kc \ and \ \arg q_{-k} = -\arg q_k$$

and hence $\arg q_k = -\pi kc$, $q_k = e^{-i\pi kc} |q_k|$, from which we obtain

$$q_{2n} = e^{-i2\pi nc} \, |q_{2n}|$$

and

$$q_{n_1}q_{n_2}...q_{n_k}q_{2n-n_1-n_2-...-n_k} = |q_{n_1}q_{n_2}...q_{n_k}q_{2n-n_1-n_2-...-n_k}| e^{-i2\pi nc}.$$

It means that $e^{i2\pi nc}\left(q_{2n} + \sum_{k=1}^{\infty} b_k(\lambda)\right)$ is a real number.

Using this remark and letting m tend to infinity in equation (4), we obtain the following main results. First, we consider the case $n \ge 1$:

Theorem 7 (a) If $M \leq \frac{4\pi^2(2n-1)}{3}$, then $\lambda_{n,j}$ is an eigenvalue of $L_0(q)$ if and only if it is either the root of the equation

$$\lambda - (2\pi n)^2 - \sum_{k=1}^{\infty} a_k(\lambda) - e^{i2\pi nc} \left(q_{2n} + \sum_{k=1}^{\infty} b_k(\lambda) \right) = 0$$
(52)

or the root of

$$\lambda - (2\pi n)^2 - \sum_{k=1}^{\infty} a_k(\lambda) + e^{i2\pi nc} \left(q_{2n} + \sum_{k=1}^{\infty} b_k(\lambda) \right) = 0$$
(53)

in the set $D_n := [(2\pi n)^2 - M, (2\pi n)^2 + M]$, where $n = 1, 2, \ldots$ Moreover, the roots of (52) and (53) in D_n , coincide with the (2n)th and (2n + 1)st periodic eigenvalues $\lambda_{n,1}$ and $\lambda_{n,2}$.

Proof. (a) By Lemma 4, letting m tend to infinity in equation (4), we obtain

$$\left(\lambda_{n,j} - (2\pi n)^2 - \sum_{k=1}^{\infty} a_k(\lambda_{n,j})\right) u_{n,j} = \left(q_{2n} + \sum_{k=1}^{\infty} b_k(\lambda_{n,j})\right) \overline{u_{n,j}},\tag{54}$$

where $\left|\frac{u_{n,j}}{\overline{u_{n,j}}}\right| = 1$. Therefore, we have

$$\left|\lambda_{n,j} - (2\pi n)^2 - \sum_{k=1}^{\infty} a_k(\lambda_{n,j})\right| = \left|q_{2n} + \sum_{k=1}^{\infty} b_k(\lambda_{n,j})\right|.$$

Now taking into account that

$$\lambda_{n,j} - (2\pi n)^2 - \sum_{k=1}^{\infty} a_k(\lambda_{n,j})$$

and

$$e^{i2\pi nc}\left(q_{2n}+\sum_{k=1}^{\infty}b_k(\lambda)\right)$$

are real numbers (see Lemma 2.1.1 (b) of [17] and Remark 1), we obtain that $\lambda_{n,j}$ is either the root of equation (52) or the root of (53).

Now, we prove that the roots of (52) and (53) lying in the interval D_n are the eigenvalues of L_0 . To do this, we first give the following relations:

$$a_1(\lambda) = a_1((2\pi n)^2) - \sum_{\substack{n_1 = -\infty\\n_1 \neq 0, 2n}}^{\infty} \frac{|q_{n_1}|^2 [\lambda - (2\pi n)^2]}{4\pi^2 n_1 (2n - n_1) [\lambda - (2\pi (n - n_1))^2]}$$

and

$$b_1(\lambda) = b_1((2\pi n)^2) - \sum_{\substack{n_1 = -\infty\\n_1 \neq 0, 2n}}^{\infty} \frac{q_{n_1}q_{2n-n_1}[\lambda - (2\pi n)^2]}{4\pi^2 n_1(2n-n_1)[\lambda - (2\pi (n-n_1))^2]}.$$

The equation ??(function)

$$F_{n,j}(\lambda) := \lambda - (2\pi n)^2 - a_1((2\pi n)^2) + (-1)^j e^{i2\pi nc} \left(q_{2n} + b_1((2\pi n)^2) \right)$$

has one root, for each j, in the interval $D_n = [(2\pi n)^2 - M, (2\pi n)^2 + M]$ and

$$|F_{n,j}(\lambda)| \ge |\lambda - (2\pi n)^2| - |a_1((2\pi n)^2)| - |q_{2n} + b_1((2\pi n)^2)|.$$

By Remark 1 and (51), we write

$$e^{i2\pi nc} (q_{2n} + b_1((2\pi n)^2)) = e^{i2\pi nc} (q_{2n} + 2Q_0 Q_{2n} - S_{2n})$$

= $\frac{(a-b)\sin(2\pi nc)}{2\pi n} + \frac{ab\cos(2\pi nc)}{8\pi^2 n^2} + \frac{(a^2-b^2)\sin(2\pi nc)}{16\pi^3 n^3}.$ (55)

Estimating $|a_1((2\pi n)^2)|$ and $|q_{2n} + b_1((2\pi n)^2)|$ by (50) and (55), we have

$$|F_{n,j}(\lambda)| > \frac{9\pi^2}{20},$$

$$G_{n,j}(\lambda) := \lambda - (2\pi n)^2 - \sum_{k=1}^{\infty} a_k(\lambda) + (-1)^j e^{i2\pi nc} \left(q_{2n} + \sum_{k=1}^{\infty} b_k(\lambda) \right),$$
(56)

for j = 1, 2. Estimating the summands of $|a_1(\lambda) - a_1((2\pi n)^2)|$, $|a_k(\lambda)|$, $|b_1(\lambda) - b_1((2\pi n)^2)|$ and $|b_k(\lambda)|$ by considering the greatest summands of them, for n = 1, we obtain

$$|a_1(\lambda) - a_1((2\pi n)^2)| < 1, \qquad |a_k(\lambda)| < \frac{2}{\pi^{k-1}},$$

$$|b_1(\lambda) - b_1((2\pi n)^2)| < 1, \qquad |b_k(\lambda)| < \frac{2}{\pi^{k-1}},$$

for $k \geq 2$. Therefore, it follows by the geometric series formula that

$$|a_1(\lambda) - a_1((2\pi n)^2)| + \sum_{k=2}^{\infty} |a_k(\lambda)| < \frac{\pi + 1}{\pi - 1}$$
$$|b_1(\lambda) - b_1((2\pi n)^2)| + \sum_{k=2}^{\infty} |b_k(\lambda)| < \frac{\pi + 1}{\pi - 1}$$

for n = 1. Hence by (56), we obtain

$$\begin{aligned} |G_{n,j} - F_{n,j}| &= \left| a_1(\lambda) - a_1((2\pi n)^2) + \sum_{k=2}^{\infty} a_k(\lambda) + (-1)^j e^{i2\pi nc} \left(b_1(\lambda) - b_1((2\pi n)^2) + \sum_{k=2}^{\infty} b_k(\lambda) \right) \right| \\ &\leq |a_1(\lambda) - a_1((2\pi n)^2)| + \sum_{k=2}^{\infty} |a_k(\lambda)| + |b_1(\lambda) - b_1((2\pi n)^2)| + \sum_{k=2}^{\infty} |b_k(\lambda)| \\ &< \frac{2(\pi + 1)}{\pi - 1}, \end{aligned}$$

for all λ from the boundary of D_n , for n = 1. Similar estimations can be obtained also for $n \geq 2$. Therefore $|G_{n,j}(\lambda) - F_{n,j}(\lambda)| < |F_{n,j}(\lambda)|$ holds for all λ from the boundary of D_n and by Rouche's theorem, $G_{n,j}(\lambda)$ has one root in the set D_n , for j = 1 and j = 2. Hence, L_0 has one eigenvalue (counting multiplicity) in D_n , which is the root of (52) and it has one eigenvalue (counting multiplicity) in D_n , which is the root of the equations (52) and (53) has exactly one root (counting multiplicity) in D_n . Thus, $\lambda \in D_n$ is an eigenvalue of L_0 if and only if, it is either the root of (52) or the root of (53) and the roots of (52) and (53) coincide with the eigenvalues $\lambda_{n,1}$ and $\lambda_{n,2}$ of L_0 .

Now, we consider the case n = 0.

Theorem 8 If $M \leq 4\pi^2/3$, then the first periodic eigenvalue λ_0 is the root of the equation

$$\lambda - \sum_{k=1}^{\infty} a_k(\lambda) = 0, \tag{57}$$

in the set $D_0 = [-M, M]$. Moreover, (57) has exactly one root (counting multiplicity) in the set D_0 and this root coincides with the first eigenvalue λ_0 of L_0 .

Proof. (a) Iterating the equation $\lambda_N(\Psi_N, 1) = (q\Psi_N, 1)$ m times, for N = 0, by isolating the terms containing $(\Psi_0, 1)$ gives

$$\left(\lambda_0 - \sum_{k=1}^m a_k(\lambda_0)\right)(\Psi_0, 1) = R_m(\lambda_0),$$

where a_k and R_m are defined by (5) and (7). Letting m tend to infinity in the last equation, by Lemma 5,

we obtain (57). Let $F_0(\lambda) := \lambda - a_1(0)$ and

$$G_0(\lambda) := \lambda - \sum_{k=1}^{\infty} a_1(\lambda).$$

Then, using the relations $Q_0 = ac/2$,

$$a_1(0) = \sum_{\substack{k=-\infty\\k\neq 0}}^{\infty} \frac{|q_k|^2}{-(2\pi(-k))^2} = -\sum_{\substack{k=-\infty\\k\neq 0}}^{\infty} \frac{|q_k|^2}{4\pi^2 k^2} = Q_0^2 - \sum_{\substack{k=-\infty\\k\neq 0}}^{\infty} Q_k Q_{-k}$$
$$= Q_0^2 - \int_0^1 \left(\sum_{\substack{k=-\infty\\k=-\infty}}^{\infty} Q_k e^{i2\pi kx}\right)^2 dx = Q_0^2 - \int_0^1 Q^2(x) dx = Q_0^2 - S_0$$

and

$$S_0 = \int_0^1 S(x)dx = \int_0^1 Q^2(x)dx = \int_0^c a^2 x^2 dx + \int_c^1 (bx - b)^2 dx = \frac{a^2 c^2}{3},$$

by direct calculation we obtain

$$-a_1(0) = S_0 - Q_0^2 = \frac{a^2c^2}{3} - \frac{a^2c^2}{4} = \frac{a^2c^2}{12}.$$

If $c \in (0, 1/2)$ and $M < 4\pi^2/3$, then $|ac| \leq (1/2)M < 2\pi^2/3$ since in this case b < |a| and |a| = M. If $c \in (1/2, 1)$, then $|ac| = (1 - c)b \leq (1 - 1/2)M < 2\pi^2/3$. In this case b > |a| and b = M. Therefore, in any case $|ac| < 2\pi^2/3$ if $M < 4\pi^2/3$ and we have $A_{1,0}(0) < \pi^4/27$. It is easy to verify that the inequality $|G_0(\lambda) - F_0(\lambda)| < |F_0(\lambda)|$ holds for all λ from the boundary of D_0 and by Rouche's theorem, $G_0(\lambda)$ has one root in the set D_0 . Hence, L_0 has one eigenvalue (counting multiplicity) in D_0 , which is the root of (57). On the other hand, equation (57) has exactly one root (counting multiplicity) in D_0 . Thus, $\lambda \in D_0$ is an eigenvalue of L_0 if and only if, it is the root of (57) and the root of (57) in D_0 coincides with the first eigenvalue λ_0 of L_0 .

Now, we can use numerical methods by taking finite sums instead of the infinite series in (52), (53) and (57). In this case, we write

$$\lambda - (2\pi n)^2 - \sum_{k=1}^r a_{s,k,n}(\lambda) + (-1)^j e^{i2\pi nc} \left(q_{2n} + \sum_{k=1}^r b_{s,k,n}(\lambda) \right) = 0,$$

for j = 1 and j = 2, and

$$\lambda - \sum_{k=1}^{\prime} a_{s,k,0}(\lambda) = 0,$$

respectively, where

$$a_{s,k,n}(\lambda) = \sum_{n_1,n_2,\dots,n_k=-s}^{s} \frac{q_{n_1}q_{n_2}\cdots q_{n_k}q_{-n_1-n_2-\dots-n_k}}{[\lambda - (2\pi(n-n_1))^2]\cdots [\lambda - (2\pi(n-n_1-\dots-n_k))^2]},$$

$$b_{s,k,n}(\lambda) = \sum_{n_1,n_2,\dots,n_k=-s}^{s} \frac{q_{n_1}q_{n_2}\cdots q_{n_k}q_{2n-n_1-n_2-\dots-n_k}}{[\lambda - (2\pi(n-n_1))^2]\cdots [\lambda - (2\pi(n-n_1-\dots-n_k))^2]}.$$

We define the functions

$$K_{n,j}(\lambda) := \lambda - (2\pi n)^2 - g_{n,j}(\lambda)$$
(58)

and

$$K_0(\lambda) := \lambda - g_0(\lambda), \tag{59}$$

where

$$g_{n,j}(\lambda) = \sum_{k=1}^{r} a_{s,k,n}(\lambda) - (-1)^{j} e^{i2\pi nc} \left(q_{2n} + \sum_{k=1}^{r} b_{s,k,n}(\lambda) \right)$$
(60)

15

and

$$g_0(\lambda) = \sum_{k=1}^r a_{s,k,0}(\lambda).$$
 (61)

Then,

$$\lambda = (2\pi n)^2 + g_{n,j}(\lambda), \tag{62}$$

for j = 1 and j = 2, and $n \ge 1$.

Now we state another main result.

Theorem 9 If $M \leq \frac{4\pi^2(2n-1)}{3}$, then for all x and y from the interval $D_n = [(2\pi n)^2 - M, (2\pi n)^2 + M]$, the relations

$$|g_{n,j}(x) - g_{n,j}(y)| \le C_n |x - y|,$$

$$C_n = \frac{4(b - a)^2}{\pi (4\pi^2 (2n - 1) - M) [\pi (4\pi^2 (2n - 1) - M) - (b - a)]} \le \frac{4}{\pi (\pi - 1)} < 1,$$
(63)

hold for j = 1, 2, and equation (62) has a unique solution $\rho_{n,j}$ in D_n , for each j, where n = 1, 2, ...Moreover

$$\begin{aligned} |\lambda_{n,j} - \rho_{n,j}| &< \frac{3(b-a)^{r+2}}{2\pi^{r+1}(4\pi^2(2n-1) - M)^r [\pi(4\pi^2(2n-1) - M) - (b-a)](1-C_n)} \\ &+ \frac{6(b-a)^2}{\pi^2(s+1)^2 [4\pi^2(s+1)|s+1-2n|-M](1-C_n)}, \end{aligned}$$
(64)

for j = 1, 2.

Proof. First we prove (63) by using the mean-value theorem. To do this, we estimate $|g'_{n,j}(\lambda)| = |\frac{d}{d\lambda}g_{n,j}(\lambda)|$. By (60) we have

$$|g_{n,j}'(\lambda)| = \left|\sum_{k=1}^{r} \frac{d}{d\lambda} a_{s,k,n}(\lambda) + (-1)^{j} e^{i2\pi nc} \sum_{k=1}^{r} \frac{d}{d\lambda} b_{s,k,n}(\lambda)\right|$$
$$\leq \sum_{k=1}^{r} \left|\frac{d}{d\lambda} a_{s,k,n}(\lambda)\right| + \sum_{k=1}^{r} \left|\frac{d}{d\lambda} b_{s,k,n}(\lambda)\right|.$$

First let us estimate the summands of the first term $\left|\frac{d}{d\lambda}(A_{s,k,n}(\lambda))\right|$:

$$\begin{aligned} \left| \frac{d}{d\lambda} (a_{s,k,n}(\lambda)) \right| &= \left| \sum_{n_1, n_2, \dots, n_k = -s}^{s} \frac{d}{d\lambda} \frac{q_{n_1} q_{n_2} \cdots q_{n_k} q_{-n_1 - n_2 - \dots - n_k}}{[\lambda - (2\pi(n - n_1))^2] \cdots [\lambda - (2\pi(n - n_1 - \dots - n_k))^2]} \right| \\ &< \frac{2(b-a)^{k+1}}{\pi^{k+1} (4\pi^2 (2n-1) - M)^{k+1}} \le \frac{2}{\pi^{k+1}}, \end{aligned}$$

for $k \geq 1$. Thus, by the geometric series formula, we obtain

$$\sum_{k=1}^{r} \left| \frac{d}{d\lambda} a_{s,k,n}(\lambda) \right| < \frac{2}{\pi(\pi-1)}.$$

In the same way, one can prove that

$$\sum_{k=1}^{r} \left| \frac{d}{d\lambda} b_{s,k,n}(\lambda) \right| < \frac{2}{\pi(\pi-1)}.$$

Hence,

$$|g_{n,j}'(\lambda)| < \frac{4(b-a)^2}{\pi(4\pi^2(2n-1)-M)[\pi(4\pi^2(2n-1)-M)-(b-a)]} = C_n \le \frac{4}{\pi(\pi-1)} < 1.$$

Since the inequalities

$$|g_j'(\lambda)| < C_n < 1 \tag{65}$$

hold for all $x, y \in D_n$, (63) holds by the mean value theorem and equation (62) has a unique solution

 $\rho_{n,j} \text{ in } D_n$, for each j (j = 1, 2), by the contraction mapping theorem. Now let us prove (64). By (58), we have $K_{n,j}(x) = x - (2\pi n)^2 - g_{n,j}(x)$ and by the definition of $\rho_{n,j}$, we write $K_{n,j}(\rho_{n,j}) = 0$, for j = 1, 2. Therefore by (52), (53) and (60), we obtain

$$|K_{n,j}(\lambda_{n,j}) - K_{n,j}(\rho_{n,j})| = |K_{n,j}(\lambda_{n,j})|$$

$$= \left|\lambda_{n,j} - (2\pi n)^2 - \sum_{k=1}^r a_{s,k,n}(\lambda_{n,j}) - (-1)^j e^{i2\pi nc} \left(q_{2n} + \sum_{k=1}^r b_{s,k,n}(\lambda_{n,j})\right)\right|$$

$$\leq \left|\sum_{k=1}^\infty a_{k,n}(\lambda_{n,j}) - \sum_{k=1}^r a_{s,k,n}(\lambda_{n,j})\right| + \left|\sum_{k=1}^\infty b_{k,n}(\lambda_{n,j}) - \sum_{k=1}^r b_{s,k,n}(\lambda_{n,j})\right|.$$
(66)

For the estimation of the first term of the right-hand side of (66), we obtain

$$\begin{split} \sum_{k=1}^{\infty} a_{k,n}(\lambda_{n,j}) &- \sum_{k=1}^{r} a_{s,k,n}(\lambda_{n,j}) \bigg| \\ &\leq \left| \sum_{k=1}^{\infty} a_{k,n}(\lambda_{n,j}) - \sum_{k=1}^{r} a_{k,n}(\lambda_{n,j}) \right| + \left| \sum_{k=1}^{r} a_{k,n}(\lambda_{n,j}) - \sum_{k=1}^{r} a_{s,k,n}(\lambda_{n,j}) \right| \\ &\leq \sum_{k=r+1}^{\infty} |a_{k,n}(\lambda_{n,j})| + \sum_{k=1}^{r} |a_{k,n}(\lambda_{n,j}) - a_{s,k,n}(\lambda_{n,j})|. \end{split}$$

Using the estimations for $\sum_{k=r+1}^{\infty} |a_{k,n}(\lambda_{n,j})|$ and $\sum_{k=1}^{r} |a_{k,n}(\lambda_{n,j}) - a_{s,k,n}(\lambda_{n,j})|$, by considering the greatest summands of them, we obtain

$$\left|\sum_{k=1}^{\infty} a_{k,n}(\lambda_{n,j}) - \sum_{k=1}^{r} a_{s,k,n}(\lambda_{n,j})\right| < \sum_{k=r+1}^{\infty} \frac{3(b-a)^{k+1}}{4\pi^{k+1}(4\pi^{2}(2n-1)-M)^{k}} + \frac{3(b-a)^{2}}{\pi^{2}(s+1)^{2}[4\pi^{2}(s+1)|s+1-2n|-M]}$$

and

$$\left|\sum_{k=1}^{\infty} a_{k,n}(\lambda_{n,j}) - \sum_{k=1}^{r} a_{s,k,n}(\lambda_{n,j})\right| < \frac{3(b-a)^{r+2}}{4\pi^{r+1}(4\pi^{2}(2n-1)-M)^{r}[\pi(4\pi^{2}(2n-1)-M)-(b-a)]} + \frac{3(b-a)^{2}}{\pi^{2}(s+1)^{2}[4\pi^{2}(s+1)|s+1-2n|-M]}.$$
(67)

In the same way, one can obtain

$$\sum_{k=1}^{\infty} b_{k,n}(\lambda_{n,j}) - \sum_{k=1}^{r} b_{s,k,n}(\lambda_{n,j}) \bigg| < \frac{3(b-a)^{r+2}}{4\pi^{r+1}(4\pi^{2}(2n-1)-M)^{r}[\pi(4\pi^{2}(2n-1)-M)-(b-a)]} + \frac{3(b-a)^{2}}{\pi^{2}(s+1)^{2}[4\pi^{2}(s+1)|s+1-2n|-M]}.$$
(68)

Thus, by (66)–(68) we have

$$|K_{n,j}(\lambda_{n,j}) - K_{n,j}(\rho_{n,j})| < \frac{3(b-a)^{r+2}}{2\pi^{r+1}(4\pi^2(2n-1)-M)^r[\pi(4\pi^2(2n-1)-M)-(b-a)]} + \frac{6(b-a)^2}{\pi^2(s+1)^2[4\pi^2(s+1)|s+1-2n|-M]}.$$
(69)

To apply the mean value theorem, we estimate $|K'_{n,j}(\lambda)|$:

$$|K'_{n,j}(\lambda)| = |1 - g'_{n,j}(\lambda)| \ge |1 - |g'_{n,j}(\lambda)|| \ge 1 - C_n.$$
(70)

By the mean value formula, (69) and (70), we obtain

$$|K_{n,j}(\lambda_{n,j}) - K_{n,j}(\rho_{n,j})| = |K'_{n,j}(\xi)||\lambda_{n,j} - \rho_{n,j}|, \qquad \xi \in [(2\pi n)^2 - M, (2\pi n)^2 + M]$$

and

$$\begin{aligned} |\lambda_{n,j} - \rho_{n,j}| &= \frac{|K_{n,j}(\lambda_{n,j}) - K_{n,j}(\rho_{n,j})|}{|K'_{n,j}(\xi)|} \\ &< \frac{3(b-a)^{r+2}}{2\pi^{r+1}(4\pi^2(2n-1) - M)^r[\pi(4\pi^2(2n-1) - M) - (b-a)](1-C_n)} \\ &+ \frac{6(b-a)^2}{\pi^2(s+1)^2[4\pi^2(s+1)|s+1-2n|-M](1-C_n)}. \end{aligned}$$

Now, we give an analogous theorem to Theorem 9 for the case n = 0.

Theorem 10 If $M \leq 4\pi^2/3$, then for all x and y from the interval $D_0 = [-M, M]$ the relations

$$|g_0(x) - g_0(y)| \le C_0 |x - y|,$$

$$C_0 = \frac{3(b - a)^2}{4\pi (2\pi^2 - M) [\pi (4\pi^2 - M) - (b - a)]} \le \frac{3}{\pi (\pi - 1)} < 1,$$

hold and the equation $% \left(f_{i} \right) = \left(f_{i} \right) \left($

 $\lambda = g_0(\lambda)$

has a unique solution ρ_0 in D_0 , where $g_0(\lambda)$ is defined by (61). Moreover

$$\begin{aligned} |\lambda_0 - \rho_0| &< \frac{9(b-a)^{r+2}}{16\pi^{r+1}(4\pi^2 - M)^{r-1}(2\pi^2 - M)[\pi(4\pi^2 - M) - (b-a)](1 - C_0)} \\ &+ \frac{3(b-a)^2}{\pi^2(s+1)^2[4\pi^2(s+1)^2 - M](1 - C_0)}. \end{aligned}$$
(71)

Proof. The proof is similar to the proof of Theorem 9 by the following estimates:

$$\left| \frac{d}{d\lambda} (a_{s,k,0}(\lambda)) \right| = \left| \sum_{n_1,n_2,\dots,n_k=-s}^{s} \frac{d}{d\lambda} \frac{q_{n_1}q_{n_2}\cdots q_{n_k}q_{-n_1-n_2-\dots-n_k}}{[\lambda - (2\pi(-n_1))^2]\cdots [\lambda - (2\pi(-n_1-\dots-n_k))^2]} \right| < \frac{3(b-a)^{k+1}}{4\pi^{k+1}(4\pi^2 - M)^k(2\pi^2 - M)} \le \frac{3}{\pi^{k+1}},$$

for $k \geq 1$,

$$|g_0'(\lambda)| = \left|\sum_{k=1}^r \frac{d}{d\lambda} a_{s,k,0}(\lambda)\right| < \frac{3(b-a)^2}{4\pi (2\pi^2 - M)[\pi (4\pi^2 - M) - (b-a)]} = C_0 \le \frac{3}{\pi (\pi - 1)} < 1,$$
$$|a_{k,0}(\lambda_0)| < \frac{9(b-a)^{k+1}}{16\pi^{k+1} (4\pi^2 - M)^{k-1} (2\pi^2 - M)} \le \frac{6}{\pi^{k-1}},$$

for $k \geq 1$, and

$$\begin{aligned} |K_0(\lambda_0) - K_0(\rho_0)| &\leq \sum_{k=r+1}^{\infty} |a_{k,0}(\lambda_0)| + \sum_{k=1}^r |a_{k,0}(\lambda_0) - \sum_{k=1}^r a_{s,k,0}(\lambda_0)| \\ &< \frac{9(b-a)^{r+2}}{16\pi^{r+1}(4\pi^2 - M)^{r-1}(2\pi^2 - M)[\pi(4\pi^2 - M) - (b-a)]} + \frac{3(b-a)^2}{\pi^2(s+1)^2[4\pi^2(s+1)^2 - M]}, \end{aligned}$$

where $K_0(\lambda)$ is defined by (59).

Now let us approximate $\rho_{n,j}$ by the fixed point iterations:

$$x_{n,i+1} = (2\pi n)^2 + g_{n,1}(x_{n,i}), \tag{72}$$

and

$$y_{n,i+1} = (2\pi n)^2 + g_{n,2}(y_{n,i}), \tag{73}$$

where $g_{n,j}(x)$ (j = 1, 2) is defined by (60). Since

$$|g_{n,j}(\lambda_{n,j})| = \left|\sum_{k=1}^{r} a_{s,k,n}(\lambda_{n,j}) - (-1)^{j} e^{i2\pi nc} \left(q_{2n} + \sum_{k=1}^{r} b_{s,k,n}(\lambda_{n,j}) \right) \right|$$

$$\leq \left|\sum_{k=1}^{r} a_{s,k,n}(\lambda_{n,j})\right| + \left|q_{2n} + \sum_{k=1}^{r} b_{s,k,n}(\lambda_{n,j})\right|$$

$$\leq \sum_{k=1}^{r} |a_{s,k,n}(\lambda_{n,j})| + |q_{2n}| + \sum_{k=1}^{r} |b_{s,k,n}(\lambda_{n,j})|,$$

first let us estimate $|a_{s,k,n}(\lambda)|$. The estimation of $|b_{s,k,n}(\lambda)|$ is similar.

$$\sum_{k=1}^{r} |a_{s,k,n}(\lambda)| < \sum_{k=1}^{r} \frac{3(b-a)^{k+1}}{4\pi^{k+1}(4\pi^2(2n-1)-M)^k} \le \sum_{k=1}^{r} \frac{2}{\pi^{k-1}} < \frac{2\pi}{\pi-1}.$$
(74)

Similarly, we obtain

$$|q_{2n}| + \sum_{k=1}^{r} |b_{s,k,n}(\lambda)| < \frac{(b-a)}{2\pi n} + \frac{2\pi}{\pi - 1}$$

Hence, we have

$$|g_{n,j}(\lambda_{n,j})| < \frac{(b-a)}{2\pi n} + \sum_{k=1}^{r} \frac{3(b-a)^{k+1}}{2\pi^{k+1}(4\pi^2(2n-1)-M)^k} < \frac{(b-a)}{2\pi n} + \frac{4\pi}{\pi - 1}.$$
 (75)

On the other hand, writing $4\pi^2(2n-1)$ instead of $4\pi^2(2n-1) - M$ in (74) and (75), we obtain

$$|g_{n,j}((2\pi n)^{2})| \leq \sum_{k=1}^{r} |a_{s,k,n}((2\pi n)^{2})| + |q_{2n}| + \sum_{k=1}^{r} |b_{s,k,n}((2\pi n)^{2})| < \frac{(b-a)}{2\pi n} + \sum_{k=1}^{r} \frac{3(b-a)^{k+1}}{2\pi^{k+1}(4\pi^{2}(2n-1))^{k}} < \frac{(b-a)}{2\pi n} + \frac{3(b-a)^{2}}{2\pi [4\pi^{3}(2n-1)-(b-a)]},$$
(76)

since $|(2\pi n)^2 - (2\pi k)^2| \ge 4\pi^2(2n-1)$, for $n = 1, 2, \dots$ Now we state the following result.

Theorem 11 If $M \leq \frac{4\pi^2(2n-1)}{3}$, then the following estimations hold for the sequences $\{x_{n,i}\}$ and

 $\{y_{n,i}\}\ defined\ by\ (72)\ and\ (73):$

$$|x_{n,i} - \rho_{n,1}| < (C_n)^i \left(\frac{(b-a)}{2\pi n(1-C_n)} + \frac{3(b-a)^2}{2\pi [4\pi^3(2n-1) - (b-a)](1-C_n)}\right),\tag{77}$$

$$|y_{n,i} - \rho_{n,2}| < (C_n)^i \left(\frac{(b-a)}{2\pi n(1-C_n)} + \frac{3(b-a)^2}{2\pi [4\pi^3(2n-1) - (b-a)](1-C_n)}\right),\tag{78}$$

for i = 1, 2, 3, ..., where C_n is defined in (63) and n = 1, 2, ...

Proof. Without loss of generality, we can take $x_{n,0} = (2\pi n)^2$. By (62), (63), and (72) we have

$$\begin{aligned} |x_{n,i} - \rho_{n,1}| &= |(2\pi n)^2 + g_{n,1}(x_{n,i-1}) - ((2\pi n)^2 + g_{n,1}(\rho_{n,1}))| \\ &= |g_{n,1}(x_{n,i-1}) - g_{n,1}(\rho_{n,1})| < C_n |x_{n,i-1} - \rho_{n,1}| < (C_n)^i |x_{n,0} - \rho_{n,1}|. \end{aligned}$$

Therefore it is enough to estimate $|x_{n,0} - \rho_{n,1}|$. By definitions of $\rho_{n,j}$ and $x_{n,0}$ we obtain

$$\rho_{n,1} - x_{n,0} = g_{n,1}(\rho_{n,1}) + (2\pi n)^2 - x_{n,0} = g_{n,1}(\rho_{n,1}) - g_{n,1}(x_{n,0}) + g_{n,1}((2\pi n)^2)$$

and by the mean value theorem there exists $x \in [(2\pi n)^2 - M, (2\pi n)^2 + M]$ such that

$$g_{n,1}(\rho_{n,1}) - g_{n,1}(x_{n,0}) = g'_{n,1}(x)(\rho_{n,1} - x_{n,0}).$$

The last two equalities imply that

$$(\rho_{n,1} - x_{n,0})(1 - g'_{n,1}(x)) = g_{n,1}((2\pi n)^2).$$

Hence by (65) and (76) we have

$$|\rho_{n,1} - x_{n,0}| \le \frac{|g_{n,1}((2\pi n)^2)|}{1 - C_n} < \frac{(b-a)}{2\pi n(1 - C_n)} + \frac{3(b-a)^2}{2\pi [4\pi^3(2n-1) - (b-a)](1 - C_n)}$$

and

$$|x_{n,i} - \rho_{n,1}| < (C_n)^i \left(\frac{(b-a)}{2\pi n(1-C_n)} + \frac{3(b-a)^2}{2\pi [4\pi^3(2n-1) - (b-a)](1-C_n)}\right).$$

One can easily show by the same way that

$$|y_{n,i} - \rho_{n,2}| < (C_n)^i \left(\frac{(b-a)}{2\pi n(1-C_n)} + \frac{3(b-a)^2}{2\pi [4\pi^3(2n-1) - (b-a)](1-C_n)}\right)$$

holds.

An analogous theorem to Theorem 11 can be stated for the case n = 0.

Theorem 12 If $M \leq 4\pi^2/3$, then the following estimation holds for the sequence $\{x_{0,i}\}$ defined by $x_{0,i} = g_0(x_{0,i})$, where $g_0(\lambda)$ is defined by (61):

$$|x_{0,i} - \rho_0| \le (C_0)^i \frac{(b-a)^2}{2\pi [2\pi^3 - (b-a)](1-C_0)},\tag{79}$$

where C_0 is defined in Theorem 10.

Proof. The proof is similar to the proof of Theorem 11 by the following estimates:

$$|g_0(0)| \le \sum_{k=1}^r |A_{s,k,0}(0)| < \sum_{k=1}^r \frac{(b-a)^{k+1}}{2^{k+1}\pi^{3k+1}} < \frac{(b-a)^2}{2\pi[2\pi^3 - (b-a)]}$$

Thus by (64), (71), (77)-(79), we have the approximations $x_{0,i}$, $x_{n,i}$, and $y_{n,i}$ for λ_0 , $\lambda_{n,1}$, and $\lambda_{n,2}$,

$$\begin{aligned} |\lambda_0 - x_{0,i}| &< \frac{9(b-a)^{r+2}}{16\pi^{r+1}(4\pi^2 - M)^{r-1}(2\pi^2 - M)[\pi(4\pi^2 - M) - (b-a)](1-C_0)} \\ &+ \frac{3(b-a)^2}{\pi^2(s+1)^2[4\pi^2(s+1)^2 - M](1-C_0)} + (C_0)^i \frac{(b-a)^2}{2\pi[2\pi^3 - (b-a)](1-C_0)}, \end{aligned}$$

$$\begin{aligned} |\lambda_{n,1} - x_{n,i}| &< \frac{3(b-a)^{r+2}}{2\pi^{r+1}(4\pi^2(2n-1) - M)^r [\pi(4\pi^2(2n-1) - M) - (b-a)](1-C_n)} \\ &+ \frac{6(b-a)^2}{\pi^2(s+1)^2 [4\pi^2(s+1)|s+1-2n|-M](1-C_n)} \\ &+ (C_n)^i \bigg(\frac{(b-a)}{2\pi n(1-C_n)} + \frac{3(b-a)^2}{2\pi [4\pi^3(2n-1) - (b-a)](1-C_n)}\bigg), \end{aligned}$$

and

$$\begin{aligned} |\lambda_{n,2} - y_{n,i}| &< \frac{3(b-a)^{r+2}}{2\pi^{r+1}(4\pi^2(2n-1)-M)^r[\pi(4\pi^2(2n-1)-M)-(b-a)](1-C_n)} \\ &+ \frac{6(b-a)^2}{\pi^2(s+1)^2[4\pi^2(s+1)|s+1-2n|-M](1-C_n)} \\ &+ (C_n)^i \bigg(\frac{(b-a)}{2\pi n(1-C_n)} + \frac{3(b-a)^2}{2\pi [4\pi^3(2n-1)-(b-a)](1-C_n)}\bigg). \end{aligned}$$

By these error formulas it is clear that the error gets smaller as r and s increase.

Now, we consider the operator $L_{\pi}(q)$ which is associated with the antiperiodic boundary conditions. The analogous formula to (4) is

$$\left(\mu_{n,j} - (2n-1)^2 \pi^2 - \sum_{k=1}^m \eta_k(\mu_{n,j})\right) (\Phi_{n,j}, e^{i(2n-1)\pi x}) - \left(q_{2n-1} + \sum_{k=1}^m \nu_k(\mu_{n,j})\right) (\Phi_{n,j}, e^{-i(2n-1)\pi x}) = \delta_m(\mu_{n,j}),$$
(80)

where $\Phi_{n,j}$ is an eigenfunction corresponding to the antiperiodic eigenvalue $\mu_{n,j}$ and

$$\eta_{k}(\mu) = \sum_{n_{1},n_{2},\dots,n_{k}} \frac{q_{n_{1}}q_{n_{2}}\cdots q_{n_{k}}q_{-n_{1}-n_{2}-\dots-n_{k}}}{[\mu - (2(n-n_{1})-1)^{2}\pi^{2}]\cdots [\mu - (2(n-n_{1}-\dots-n_{k})-1)^{2}\pi^{2}]},$$

$$\nu_{k}(\mu) = \sum_{n_{1},n_{2},\dots,n_{k}} \frac{q_{n_{1}}q_{n_{2}}\cdots q_{n_{k}}q_{2n-1-n_{1}-n_{2}-\dots-n_{k}}}{[\mu - (2(n-n_{1})-1)^{2}\pi^{2}]\cdots [\mu - (2(n-n_{1}-\dots-n_{k})-1)^{2}\pi^{2}]},$$

$$\delta_{m}(\mu) = \sum_{n_{1},n_{2},\dots,n_{m+1}} \frac{q_{n_{1}}q_{n_{2}}\cdots q_{n_{m}}q_{n_{m+1}}(q\Phi_{n,j}, e^{i(2(n-n_{1}-\dots-n_{m+1})-1)\pi x})}{[\mu - (2(n-n_{1})-1)^{2}\pi^{2}]\cdots [\mu - (2(n-n_{1}-\dots-n_{m+1})-1)^{2}\pi^{2}]}.$$

Here, the sums are taken under the conditions $\sum_{i=1}^{l} n_i \neq 0, 2n-1$ for l = 1, 2, ..., m+1. Therefore, the analogous formula to (54) is

$$\left(\mu_{n,j} - (2n-1)^2 \pi^2 - \sum_{k=1}^m \eta_k(\mu_{n,j})\right) (\Phi_{n,j}, e^{i(2n-1)\pi x}) = \left(q_{2n-1} + \sum_{k=1}^m \nu_k(\mu_{n,j})\right) (\Phi_{n,j}, e^{-i(2n-1)\pi x}).$$
(81)

Using (47), (80), (81) and similar estimations to those for the periodic eigenvalues, one can obtain analogous theorems to Theorem 7, Theorem 9, and Theorem 11 for the operator $L_{\pi}(q)$.

Now, we present a numerical example. From the numerical results, we conclude that, we can impose the conditions $M < 2\pi^2$, for n = 0, and $M < 2\pi^2(2n - 1)$, for n = 1, 2, ..., where $M = \max\{|a|, b\}$,

for the periodic eigenvalues, instead of the conditions $M \leq \frac{4\pi^2}{3}$, for n = 0, and $M \leq \frac{4\pi^2(2n-1)}{3}$, for $n = 1, 2, \ldots$, for some specific values of $c \in (0, 1)$. Similarly, we can impose the conditions $M < 4\pi^2$, for n = 1, and $M < 4\pi^2(n-1)$, for $n = 2, 3, \ldots$, for the antiperiodic eigenvalues, instead of the conditions $M \leq \frac{8\pi^2}{3}$, for n = 1, and $M \leq \frac{8\pi^2(n-1)}{3}$, for $n = 2, 3, \ldots$, for some specific values of $c \in (0, 1)$.

Example 1 For $a = -\pi^2$, $b = \pi^2$, and c = 1/2, we have the following approximations for the first periodic eigenvalues λ_0 , $\lambda_{1,1}$, $\lambda_{1,2}$, $\lambda_{2,1}$, $\lambda_{2,2}$ and antiperiodic eigenvalues $\mu_{1,1}$, $\mu_{1,2}$, $\mu_{2,1}$, $\mu_{2,2}$:

$$\begin{split} \lambda_0 &= -0.100720167503\pi^2, \\ \lambda_{1,1} &= 3.953707280198\pi^2, \\ \lambda_{2,1} &= 15.974913551204\pi^2, \\ \mu_{1,1} &= 0.317539742073\pi^2, \\ \mu_{2,1} &= 8.768711027230\pi^2, \\ \mu_{2,2} &= 9.180457181326\pi^2. \end{split}$$

In our calculations, we take r = s = 5. Usually it takes 8 - 10 iterations with the tolerance 1e - 18 by the fixed point iteration method, even if we choose an initial value that is not too close to the exact value, which means that convergence is quite fast.

Moreover we have the following estimations for the first 4 gaps:

$$\begin{split} |\Delta_1| &:= \mu_{1,2} - \mu_{1,1} = 1.260523373896\pi^2 = 12.440867038680 \\ |\Delta_2| &:= \lambda_{1,2} - \lambda_{1,1} = 0,023186881638\pi^2 = 0.228845349062 \\ |\Delta_3| &:= \mu_{2,2} - \mu_{2,1} = 0.411746154096\pi^2 = 4.063771654597 \\ |\Delta_4| &:= \lambda_{2,2} - \lambda_{2,1} = 0.008508819037\pi^2 = 0.083978677816. \end{split}$$

References

- B. M. Brown, M. S. P. Eastham, K. M. Schmidt, Periodic Differential Operators, Springer Basel (2013).
- [2] N. Dernek, O. A. Veliev, On the Riesz basisness of the root functions of the non-selfadjoint Sturm-Liouville operator. Isr. J. Math. 145, 113–123 (2005).
- [3] N. Dunford, J. T. Schwartz, Linear Operators, Part 2. Spectral Theory, Self Adjoint Operators in Hilbert Space. John Wiley & Sons, New York (1988).
- [4] M. S. P. Eastham, The Spectral Theory of Periodic Differential Equations. Scotting Academic Press, Edinburgh (1973).
- [5] I. M. Gelfand, Expansions in series of eigenfunctions of an equation with periodic coefficients, Sov. Math. Dokl. 73 (1950), 1117–1120.
- [6] T. Kato, Perturbation Theory for Linear Operators, Berlin. Springer-Verlag, Germany (1980).
- [7] R. de L. Kronig, W. G. Penney, Quantum mechanics in cristal lattices, Proc. Roy. Soc. 130 (1931), 499-513.
- [8] W. Magnus, S. Winkler, Hill's Equation. Interscience Publishers, New York (1966).
- [9] V. A. Marchenko, Sturm-LiouvilleOperators and Applications. BirkhauserVerlag, Basel (1986).
- [10] D. C. McGarvey, Differential operators with periodic coefficients in $L_p(-\infty, \infty)$. Journal of Mathematical Analysis and Applications 11 (1965), 564-596.
- [11] M. A. Naimark, Linear Differential Operators. George G. Harap & Company, London (1967)

- [12] F. S. Rofe-Beketov, The spectrum of non-self-adjoint differential operators with periodic coefficients, Soviet Math. Dokl. 4 (1963), 1563-1566.
- [13] A. A. Shkalikov, O. A. Veliev, On the Riesz basis property of the eigen- and associated functions of periodic and antiperiodic Sturm-Liouville Problems, Mathematical Notes, Vol 85, No. 5, pp.647-660, (2009).
- [14] E. C. Titchmarsh, Eigenfunction Expansion (Part II). Oxford University Press, London (1958).
- [15] O. A. Veliev, Toppamuk Duman, The spectral expansion for the non-self-adjoint Hill operators with a locally integrable potential. Journal of Mathematical Analysis and Applications, 265, no. 1, pp. 76-90 (2002).
- [16] O. A. Veliev, Non-self-adjoint Schrödinger Operator with a Periodic Potential, Springer, Switzerland 2021.
- [17] O. A. Veliev, Multidimensional Periodic Schrödinger Operator. Springer Tracts in Modern Physics, vol 291. Springer, Switzerland 2024.