# THE WORST-CASE ROOT-CONVERGENCE FACTOR OF GMRES(1)

YUNHUI HE˚

Abstract. In this work, we analyze the asymptotic convergence factor of minimal residual iteration (MRI) (or GMRES(1)) for solving linear systems  $Ax = b$  based on vector-dependent nonlinear eigenvalue problems. The worst-case root-convergence factor is derived for linear systems with A being symmetric or  $I - A$  being skew-symmetric. When A is symmetric, the asymptotic convergence factor highly depends on the initial guess. While  $M = I - A$  is skew-symmetric, GMRES(1) converges unconditionally and the worst-case root-convergence factor relies solely on the spectral radius of M. We also derive the q-linear convergence factor, which is the same as the worst-case root-convergence factor. Numerical experiments are presented to validate our theoretical results.

Key words. Root-convergence factor, minimal residual iteration, GMRES(1), restarted AA(1), vector-dependent eigenvalue problems

AMS subject classifications. 65F10, 15A18

<span id="page-0-0"></span>1. Introduction. In this work, we consider solving the following linear system

$$
(1.1) \t\t Ax = b,
$$

where  $A \in \mathbb{R}^{n \times n}$  is invertible. One of the many iterative methods for solving  $(1.1)$ is the Generalized Minimal Residual Method (GMRES) [\[29\]](#page-20-0). In the literature, there is a substantial amount of work on the convergence analysis of GMRES, see [\[15,](#page-20-1) [8,](#page-19-0) [33,](#page-20-2) [35,](#page-20-3) [26,](#page-20-4) [19,](#page-20-5) [18,](#page-20-6) [32,](#page-20-7) [20,](#page-20-8) [3\]](#page-19-1). The convergence behavior of GMRES is quite complex. For example, [\[14\]](#page-19-2) shows that any nonincreasing convergence curve can be obtained with GMRES applied to a matrix having any desired eigenvalues. Many variants of GMRES have been developed, such as GMRES with deflated restarting [\[23,](#page-20-9) [13,](#page-19-3) [21,](#page-20-10) [1\]](#page-19-4), restarted GMRES (GMRES $(m)$ ) [\[22,](#page-20-11) [4,](#page-19-5) [12,](#page-19-6) [37\]](#page-20-12), nested GMRES methods [\[36\]](#page-20-13), hybrid GMRES [\[24,](#page-20-14) [30\]](#page-20-15), and weighted GMRES [\[16,](#page-20-16) [9\]](#page-19-7).

There have been many efforts on the convergence analysis of restarted GMRES  $[17, 11, 41, 40]$  $[17, 11, 41, 40]$  $[17, 11, 41, 40]$  $[17, 11, 41, 40]$  $[17, 11, 41, 40]$  $[17, 11, 41, 40]$  $[17, 11, 41, 40]$ , which selects a relatively large restart parameter m to imitate the full GMRES process. However, there still lacks better understanding of restarted GMRES, even for GMRES(1)(or Minimal Residual Iteration), see Algorithm [1.1.](#page-1-0) Some numerical results show that using small values of the restart parameter yields convergence in fewer iterations compared to larger values. For example, numerical experiments in [\[7\]](#page-19-9) show that GMRES(1) for some linear systems with dimension three converges exactly in three iterations, while GMRES(2) stagnates. Moreover, the convergence of GMRES(1) can be extremely sensitive to the initial guess. There is still much to understand about the convergence behavior of  $\mathrm{GMRES}(m)$ , especially regarding how the asymptotic convergence factor is sensitive to the initial guess. Improved GM-RES convergence theory is needed to offer practical guidance on using this algorithm. We limit ourselves to investigating GMRES(1). In [\[27,](#page-20-20) Proposition 2.1], the author derived the following upper bound for  $GMRES(1)$  when A is symmetric definite:

(1.2) 
$$
\|r_{k+1}\| \leq \left|\frac{\lambda_{\max}(A) - \lambda_{\min}(A)}{\lambda_{\max}(A) + \lambda_{\min}(A)}\right| \|r_{k}\|.
$$

However, there is not much investigation on the above bound. For different initial errors  $r_0$ , the upper bound might not be tight. What is the asymptotic convergence

<sup>˚</sup>Department of Mathematics, University of Houston, 3551 Cullen Blvd, Room 641, Houston, Texas 77204-3008, USA [\(yhe43@central.uh.edu\)](mailto:yhe43@central.uh.edu).

factor for GMRES(1)? Is it possible that there exists  $r_0$  such that the above bound is achieved for every  $k$ ? In this work, we address these questions and provide additional results on how the initial guess or initial error affects the asymptotic convergence factor.

<span id="page-1-0"></span>

DEFINITION 1.1 (Root convergence). Assume that a sequence  $\{y_k\}$  generated by an iterative method converges to  $y^*$ . We define the root-convergence factor of the iterative method as

<span id="page-1-1"></span>(1.3) 
$$
\varrho(y_0) = \lim \sup_{k \to \infty} \varrho_k(y_0), \quad \text{where} \quad \varrho_k(y_0) = \|y_k - y^*\|^{1/k},
$$

where  $y_0$  is the initial guess for the iterative method. Further more, we call

<span id="page-1-2"></span>
$$
\varrho^* = \max_{y_0} \varrho(y_0),
$$

the worst-case root-convergence factor of the iterative method.

DEFINITION 1.2 (q-linear convergence, [\[34\]](#page-20-21)). We call a sequence  $\{y_k\}$  converges q-linearly with q-factor  $\sigma \in (0, 1)$  to  $y^*$  if

<span id="page-1-3"></span>(1.5) 
$$
||y_k - y^*|| \le \sigma ||y_{k-1} - y^*||, \quad k \ge 1.
$$

Motivated by  $[5]$  and  $[6]$ , we analyze the convergence behavior of  $GMRES(1)$ based on eigenvector-dependent eigenvalue problems. The main goal of this work is to offer theoretical results of  $(1.3)$ ,  $(1.4)$ , and  $(1.5)$  for GMRES $(1)$ . The main contributions of this work are summarized as follows. We provide a clear picture of the eigenpairs of vector-dependent eigenvalue problems associated with the GMRES(1) iterations. We derive the worst-case root-convergence factor of GMRES(1) with A being symmetric and  $M = I - A$  being skew-symmetric. Our results also show that how the initial guess can affect the asymptotic convergence factor. For some special initial guesses for GMRES(1), we present the corresponding root-convergence factors. We add some theoretical results for  $\text{rAA}(1)$ . We provide counterexamples for a conjecture of restarted Anderson acceleration with window size one, denoted as rAA(1), presented in  $[6]$ . A comparison between GMRES $(1)$  and rAA $(1)$  is discussed. We see that for  $M$  being skew-symmetric,  $\text{GMRES}(1)$  always converges with the worst-case convergence factor depending only on the spectral radius of M and has a faster convergence factor compared to  $\text{rAA}(1)$ . For A being symmetric definite,  $\text{GMRES}(1)$ always converges, and the worse-case root-convergence factor is  $\frac{\lambda_{\text{max}}(A) - \lambda_{\text{min}}(A)}{\lambda_{\text{max}}(A) + \lambda_{\text{min}}(A)}$  $\lambda_{\max}(A)+\lambda_{\min}(A)$  $\int$  and is one for indefinite system. We also derive the q-linear convergence factor, which is the same as the worst-case root-convergence factor. These results can help us better understand  $GMRES(1)$  and  $rAA(1)$  methods. They have the potential to offer guidance on designing faster algorithms, such as preconditioning.

The rest of this work is organized as follows. In section [2,](#page-2-0) we analyze GMRES(1) for A being symmetric and derive the worst-case root-convergence factor. Some new results of rAA(1) have been presented and we provide counterexamples of a conjecture from [\[6\]](#page-19-11). In section [3,](#page-14-0) we consider  $M = I - A$  being skew-symmetric, where the worstcase convergence factor is proposed. GMRES(1) always converges for this case. Then, we briefly discuss the q-linear convergence of GMRES(1). Numerical experiments are presented in section [4](#page-16-0) to validate our theoretical results. Finally, we draw conclusions in section [5.](#page-18-0)

<span id="page-2-0"></span>**2. Symmetric case for GMRES(1).** From Algorithm [1.1,](#page-1-0) we have

(2.1) 
$$
r_{k+1} = (I - \alpha_k(r_k)A)r_k.
$$

Let us consider the following mapping Φ:

<span id="page-2-1"></span>
$$
\Phi(v) = (I - \alpha(v)A)v,
$$

where

$$
\alpha(v) = \frac{\langle v, Av \rangle}{\langle Av, Av \rangle} = \frac{\langle Av, S(Av) \rangle}{\langle Av, Av \rangle},
$$

where  $S = \frac{1}{2}(A^{-1} + A^{-T})$ . Then [\(2.1\)](#page-2-1) can be written as  $r_{k+1} = \Phi(r_k)$ .

It is known that for an iterative function  $\mathcal F$  if it is differentiable at  $x^*$  (a fixed point of  $\mathcal F$ ), then the worst-case root-convergence factor is given by the spectral radius of  $\mathcal{F}'(x^*)$ , see [\[25\]](#page-20-22). Here, we cannot employ this approach to analyze GMRES(1).

2.1. One step of GMRES(1). Define the following vector-dependent matrix

$$
\mathcal{I}(v) = I - \alpha(v)A, \quad v \neq 0.
$$

It can be seen that  $\mathcal{I}(v)v = \Phi(v)$ . Based on this, every one step of the residuals of GMRES(1) satisfy

$$
r_{k+1} = \mathcal{I}(r_k)r_k, \quad k = 0, 1, \cdots
$$

To understand the worst-case root-convergence factor of GMRES(1), we first investigate

$$
\max_{v \in \mathbb{R}^n} \frac{\|\mathcal{I}(v)v\|}{\|v\|}.
$$

We consider the following eigenvector-dependent nonlinear eigenvalue problem:

<span id="page-2-2"></span>(2.2) 
$$
\mathcal{I}_2(u)u = \lambda u, \text{ where } \mathcal{I}_2(u) = (I - \alpha(u)A^T)(I - \alpha(u)A).
$$

For simplicity, when considering A as symmetric, we assume that the following condition is satisfied.

<span id="page-2-3"></span>ASSUMPTION 2.1. Suppose that  $A \in \mathbb{R}^{n \times n}$  is symmetric with p distinguish eigenvalues, denoted as  ${a_i}_{i=1}^p$ . For each i, there are  $n_i$  linearly independent orthonormal values, denoted as  $\{a_i\}_{i=1}^r$ . For each  $i$ , the eigenvectors  $\{u_{i,j}\}_{j=1}^{n_i}$ , where  $\sum_{i=1}^p n_i = n$ .

<span id="page-3-0"></span>Lemma 2.1.

(2.3) 
$$
\max_{v \neq 0} \frac{\|\mathcal{I}(v)v\|}{\|v\|} = \Gamma^*,
$$

where  $\Gamma^* = \max_{\lambda}$  $|\lambda|$  with  $\lambda$  being the eigenvalues of  $(2.2)$ .

*Proof.* Since  $||\mathcal{I}(v)|| = \min_{c \in \mathbb{R}} ||(I - cA)v||$ , [\(2.3\)](#page-3-0) is a direct conclusion from [\[11,](#page-19-8) Theorems 3.5, 3.6].  $\Box$ 

In the following, we focus on the case where  $A$  is symmetric.

LEMMA 2.2. Let  $z = \sum_{i=1}^{n}$  $\sum_{i=1}^n y_i$ , where  $y_i$  are orthogonal eigenvectors of the symmetric matrix  $A \in \mathbb{R}^{n \times n}$  associated with eigenvalues  $s_i$ . Then,

<span id="page-3-2"></span>(2.4) 
$$
\alpha(z) = \frac{\langle z, Az \rangle}{\langle Az, Az \rangle} = \frac{\sum_{i=1}^{n} s_i \|y_i\|^2}{\sum_{i=1}^{n} s_i^2 \|y_i\|^2}.
$$

For convenience, we consider the orthonormal eigenvectors of A in the following.

<span id="page-3-3"></span>LEMMA 2.3. Let  $u_i, u_j$  be two orthonormal eigenvectors of the symmetric matrix A with eigenvalues  $a_i$  and  $a_j \neq a_i$ , respectively. Consider  $z_{ij} = c_i u_i + c_j u_j$  with  $\epsilon = \frac{c_j}{c_j}$  $\frac{c_j}{c_i}$ . Then,  $z_{ij}$  is an eigenvector of  $(2.2)$  associated with non-unit eigenvalue if and only if  $a_i a_j > 0$  and  $\epsilon^2 = \frac{a_i}{a_j}$ , and the corresponding eigenvalue is

<span id="page-3-1"></span>
$$
\lambda_{ij} = \frac{(a_j - a_i)^2}{(a_j + a_i)^2}.
$$

Proof. By standard calculation, we have

(2.5) 
$$
\alpha(z_{ij}) = \frac{a_i + a_j \epsilon^2}{a_i^2 + a_j^2 \epsilon^2}
$$

Assume that

$$
\mathcal{I}_2(z_{ij})(z_{ij}) = (1 - \alpha(z_{ij})a_i)^2 c_i u_i + (1 - \alpha(z_{ij})a_j)^2 c_j u_j = \lambda_{ij} z_{ij}.
$$

.

It follows that

$$
(1 - \alpha(z_{ij})a_i)^2 c_i u_i = \lambda_{ij} c_i u_i \quad \text{and} \quad (1 - \alpha(z_{ij})a_j)^2 c_j u_j = \lambda_{ij} c_j u_j.
$$

It is equivalent to  $(1 - \alpha(z_{ij})a_i)^2 = (1 - \alpha(z_{ij})a_j)^2$ , that is,  $2 = (a_j + a_i)\alpha(z_{ij})$ . Using  $\alpha(z_{ij})$  defined in [\(2.5\)](#page-3-1), we obtain  $\epsilon^2 = \frac{a_i}{a_j}$ . Next, we compute the eigenvalue

$$
\lambda_{ij} = (1 - \alpha(z_{ij})a_i)^2 = \left(\frac{a_j(a_j - a_i)\epsilon^2}{a_i^2 + a_j^2 \epsilon^2}\right)^2 = \frac{(a_j - a_i)^2}{(a_j + a_i)^2},
$$

which completes the proof.

<span id="page-3-4"></span>THEOREM 2.4. Assume that A is symmetric. Let  $i_1, i_2 \in \{1, 2, \dots, p\}$  and  $i_1 \neq i_2$ . Consider

$$
u^* = \sum_{s=1}^{n_{i_1}} c_{i_1,s} u_{i_1,s} + \sum_{r=1}^{n_{i_2}} c_{i_2,r} u_{i_2,r}, \quad c_{i_1,1} c_{i_2,1} \neq 0,
$$

and  $u_{i_1 i_2} = c_{i_1} u_{i_1,1} + c_{i_2} u_{i_2,1}$ , where

<span id="page-4-1"></span>(2.6) 
$$
c_{i_1}^2 = \sum_{s=1}^{n_{i_1}} c_{i_1,s}^2, \quad c_{i_2}^2 = \sum_{r=1}^{n_{i_1}} c_{i_1,r}^2,
$$

Then,

<span id="page-4-0"></span>
$$
\alpha(u^*) = \alpha(u_{i_1 i_2}).
$$

Moreover, if  $\left(\frac{c_{i_2}}{c}\right)$  $c_{i_1}$  $\sqrt{2}$  $=\frac{a_{i_1}}{a_{i_2}}$  $\frac{a_{i_1}}{a_{j_1}}>0$ , then  $u^*$  and  $u_{i_1i_2}$  are eigenvectors of  $(2.14)$  associated with the same eigenvalue, i.e.,  $\mathcal{I}_2(u^*)u^* = \lambda_{i_1i_2}u^*$  and  $\mathcal{I}_2(u_{i_1i_2})u_{i_1i_2} = \lambda_{i_1i_2}u_{i_1i_2}$ , where

(2.8) 
$$
\lambda_{i_1 i_2}(\epsilon_{i_1 i_2}) = \frac{(a_{i_2} - a_{i_1})^2}{(a_{i_1} + a_{i_2})^2}.
$$

*Proof.* When  $c_{i_1} = 0$  or  $c_{i_2} = 0$ , it is obvious that  $u^*$  and  $u_{i_1 i_2}$  are eigenvectors of [\(2.14\)](#page-6-0) and  $\lambda = 0$ . Next, we prove [\(2.7\)](#page-4-0). From [\(2.4\)](#page-3-2), we have

$$
\alpha(u^*)=\frac{\sum_{s=1}^{n_{i_1}}a_{i_1}c_{i_1,s}^2+\sum_{r=1}^{n_{i_2}}a_{i_2}c_{i_1,r}^2}{\sum_{s=1}^{n_{i_1}}a_{i_1}^2c_{i_1,s}^2+\sum_{r=1}^{n_{i_2}}a_{i_2}^2c_{i_1,r}^2}=\frac{a_{i_1}\sum_{s=1}^{n_{i_1}}c_{i_1,s}^2+a_{i_2}\sum_{r=1}^{n_{i_2}}c_{i_1,r}^2}{a_{i_2}^2\sum_{s=1}^{n_{i_1}}c_{i_1,s}^2+a_{i_2}^2\sum_{r=1}^{n_{i_2}}c_{i_1,r}^2},
$$

and

$$
\alpha(u_{i_1i_2})=\frac{a_{i_1}c_{i_1}^2+a_{i_2}c_{i_2}^2}{a_{i_1}^2c_{i_1}^2+a_{i_2}^2c_{i_2}^2}.
$$

Using conditions in [\(2.6\)](#page-4-1) gives  $\alpha(u^*) = \alpha(u_{i_1 i_2})$ .

From Lemma [2.3,](#page-3-3) we know that  $(\lambda_{i_1i_2}(\epsilon_{i_1i_2}), u_{i_1i_2})$  is eigenpair of [\(2.2\)](#page-2-2). For simplicity, let  $\beta = \alpha(u^*)$ . We have

$$
(1 - \beta a_{i_1})^2 c_{i_1} u_{i_1,1} = \lambda_{i_1 i_2} c_{i_1} u_{i_1,1} \text{ and } (1 - \beta a_{i_2})^2 c_{i_2} u_{i_2,1} = \lambda_{i_1 i_2} c_{i_2} u_{i_2,1}.
$$

It means  $(1 - \beta a_{i_1})^2 = \lambda_{i_1 i_2}$  and  $(1 - \beta a_{i_2})^2 = \lambda_{i_1 i_2}$ . Thus,

$$
\mathcal{I}_2(u^*)u^* = \sum_{s=1}^{n_{i_1}} (1 - \beta a_{i_1})^2 c_{i_1,s} u_{i_1,s} + \sum_{r=1}^{n_{i_2}} (1 - \beta a_{i_2})^2 c_{i_2,r} u_{i_2,r}
$$
  
\n
$$
= \sum_{s=1}^{n_{i_1}} \lambda_{i_1 i_2} c_{i_1,s} u_{i_1,s} + \sum_{r=1}^{n_{i_2}} \lambda_{i_1 i_2} c_{i_2,r} u_{i_2,r}
$$
  
\n
$$
= \lambda_{i_1 i_2} \left( \sum_{s=1}^{n_{i_1}} c_{i_1,s} u_{i_1,s} + \sum_{r=1}^{n_{i_2}} c_{i_2,r} u_{i_2,r} \right)
$$
  
\n
$$
= \lambda_{i_1 i_2} u^*,
$$

which completes the proof.

THEOREM 2.5. Suppose that A is symmetric. If A is definite, then all the eigenvectors of [\(2.14\)](#page-6-0) associated with nonzero eigenvalue are given by

<span id="page-4-2"></span>
$$
(2.9) \qquad u^* = \sum_{s=1}^{n_{i_1}} c_{i_1,s} u_{i_1,s} + \sum_{r=1}^{n_{i_2}} c_{i_2,r} u_{i_2,r}, \quad \text{where} \quad i_1 \neq i_2, i_1, i_2 \in \{1, 2, \cdots, p\},
$$

where at least one term in  ${c_{i_1,s}}_{s=1}^{n_{i_1}}$  is nonzero, at least one term in  ${c_{i_2,r}}_{r=0}^{n_{i_2}}$  is nonzero, and  $\frac{c_{i_2}^2}{c_{i_1}^2} = \frac{a_{i_1}}{a_{i_2}}$  $\frac{a_{i_1}}{a_{i_2}} > 0$  with  $\sum_{s=1}^{n_{i_1}} c_{i_1,s}^2$  and  $c_{i_2}^2 = \sum_{r=1}^{n_{i_1}} c_{i_1,r}^2$ .

If  $A$  is indefinite, then all the eigenvectors of  $(2.14)$  associated with non-unit eigenvalues are given by  $(2.9)$ , and the eigenvectors associated with eigenvalue one are given by  $u^*$ , where  $\langle Au^*, u^* \rangle = 0$ .

Formula [\(2.9\)](#page-4-2) means that the eigenvectors associated with nonzero eigenvalue of [\(2.14\)](#page-6-0) are any linear combination of eigenvectors of A associated with exactly any two distinguish eigenvalues of A.

Proof. For any  $i \in \{1, 2, \dots, p\}$ , let  $u_* = \sum_s^{n_i} c_{i,s} u_{i,s} \neq 0$ . Then,  $(I - \alpha(u_*)A)u_* =$ 0, which means  $\mathcal{I}_2(u_*)u_* = 0$  and  $u_*$  is an eigenvector corresponding to zero eigenvalue.

<span id="page-5-0"></span>Assume that the eigenvector of  $(2.2)$  is given by

(2.10) 
$$
u^* = \sum_{i=1}^p \sum_{s=1}^{n_i} c_{i,s} u_{i,s},
$$

which contains at least two different integers  $i, j$  such that  $c_{i,s}c_{j,t} \neq 0$ .

When A is indefinite, there exists a vector v such that  $\langle Av, v \rangle = 0$ . Then  $\alpha(v) = 0$ and  $\mathcal{I}_2(v)(v) = v$ . In this situation,  $\lambda = 1$  is an eigenvalue of [\(2.2\)](#page-2-2).

Next, we prove that in  $(2.10)$ , there are only two different integers i, j such that  $c_{i,s}c_{j,t} \neq 0$ . For simplicity, let  $\beta = \alpha(u^*)$ . Assume  $\mathcal{I}_2(u^*)u^* = \lambda u^*, \lambda \neq 0$ . Then, we have

<span id="page-5-1"></span>
$$
(2.11) \t c_{i,s}(1-\beta a_i)^2 u_{i,s} = \lambda c_{i,s} u_{i,s}, \t (i,s) \in \{1,2,\cdots,p\} \times \{1,2,\cdots,n_i\}.
$$

If  $\beta = 0$ , then  $\langle Au^*, u^* \rangle$  and  $\lambda = 1$ . Consider  $\beta \neq 0$ . Define the quadratic function  $g(b) = (1 - \beta b)^2$ . For a given value  $g^*$ , there are at most two different values  $b_1, b_2$ such that  $g(b_1) = g(b_2) = g^*$ . To guarantee [\(2.11\)](#page-5-1), there are at most two different integers i, j such that  $c_{i,s}c_{j,t} \neq 0$ , where  $s \in \{1, 2, \dots, n_i\}$  and  $t \in \{1, 2, \dots, n_j\}$ , and the coefficients conditions are given in Theorem [2.4.](#page-3-4) Thus, all the eigenvectors of [\(2.14\)](#page-6-0) associated with non-unit eigenvalues are any linear combination of eigenvectors of A with exactly two distinguish eigenvalues of A.  $\Box$ 

From the above discussion, we know that

<span id="page-5-3"></span>(2.12) 
$$
\Gamma^* = \begin{cases} \max_{i \neq j, i, j \in \{1, 2, \cdots, p\}} \frac{|a_j - a_i|}{|a_j| + |a_i|}, & \text{if } A \text{ is definite} \\ 1, & \text{if } A \text{ is indefinite} \end{cases}
$$

<span id="page-5-4"></span>THEOREM 2.6. For GMRES(1) applied to  $Ax = b$ , where A is symmetric, the worst-case root-convergence factor is #ˇ

<span id="page-5-2"></span>(2.13) 
$$
\varrho_{GMRES(1)}^* \leq \begin{cases} \left| \frac{\lambda_{\max}(A) - \lambda_{\min}(A)}{\lambda_{\max}(A) + \lambda_{\min}(A)} \right|, & \text{if } A \text{ is definite} \\ 1, & \text{if } A \text{ is indefinite} \end{cases}
$$

*Proof.* From Lemma [2.1,](#page-2-3) we have  $||r_k|| \leq \Gamma^* ||r_{k-1}||$ . Thus,  $||r_k|| \leq (\Gamma^*)^k ||r_0||$  and  $\rho^* \leqslant \Gamma^*.$  $\Box$ 

From the above result, we know that when  $A$  is symmetric definite, then the worst-From the disorder result, we have that when TI is symmetric corresponding to the convergence factor is not worse than  $\frac{\lambda_{\text{max}}(A) - \lambda_{\text{min}}(A)}{\lambda_{\text{min}}(A) + \lambda_{\text{min}}(A)}$  $\lambda_{\max}(A)+\lambda_{\min}(A)$  $\vert$ . When A is symmetric indefinite,  $\rho^* = 1$ . In the following, we will show that there exists  $r_0$  such that the upper bound can be achieved.

2.2. Two steps of GMRES(1). To better understand how the initial guess affects the root-convergence factor of GMRES(1), we consider two steps of GMRES(1), and define the following vector-dependent matrix

$$
\Pi(v) = (I - \alpha(\Phi(v))A)(I - \alpha(v)A), \quad v \neq 0.
$$

It can be seen that  $\Pi(v)v = \Phi(\Phi(v))$ . Based on this, every two steps of the residuals of GMRES(1) satisfy

<span id="page-6-0"></span>
$$
r_{2(k+1)} = \Pi(r_{2k})r_{2k}, \quad k = 0, 1, \cdots
$$

The above iteration is related to the following eigenvector-dependent nonlinear eigenvalue problem

$$
(2.14) \t \Pi(u)u = \lambda u.
$$

<span id="page-6-1"></span>THEOREM 2.7. Let  $u_i, u_j$  be two orthonormal eigenvectors of the symmetric matrix A with eigenvalues  $a_i$  and  $a_j \neq a_i$ , respectively. Define

$$
z_{ij} = c_i u_i + c_j u_j = c_i (u_i + \epsilon u_j), \quad \epsilon = \frac{c_j}{c_i}, \quad c_i c_j \neq 0.
$$

Then

$$
\Pi(z_{ij})z_{ij} = \lambda_{ij}(\epsilon)z_{ij},
$$

where

$$
\lambda_{ij}(\epsilon) = \frac{(a_j - a_i)^2}{a_i^2 + \epsilon^2 a_j^2} \frac{1}{1 + \frac{1}{\epsilon^2}}.
$$

Proof. This can be easily derived following [\[6\]](#page-19-11). Thus, we omit it here.

<span id="page-6-3"></span><span id="page-6-2"></span>LEMMA 2.8. The maximum value of  $\lambda_{ij}(\epsilon)$  is given by

(2.15) 
$$
\max_{\epsilon \in \mathbb{R}} \lambda_{ij}(\epsilon) = \lambda_{ij} \left( \pm \sqrt{\frac{|a_i|}{|a_j|}} \right) = \frac{(a_j - a_i)^2}{(|a_j| + |a_i|)^2} \leq 1.
$$

*Proof.* To find the maximum of  $\lambda_{ij}(\epsilon)$ , we only need to find the minimum of

$$
g(\tau) = (a_i^2 + \tau a_j^2)(1 + \frac{1}{\tau}),
$$

where  $\tau = \epsilon^2 > 0$ . In fact,  $g'(\tau) = a_j^2(1 + \frac{1}{\tau}) + (a_i^2 + \tau a_j^2)(-\frac{1}{\tau^2})$ . Let  $g'(\tau) = 0$ . We where  $\tau = \epsilon^2 > 0$ . In fact,  $g'(\tau) = a_{\tilde{j}}(1 + \frac{\tilde{\tau}}{\tau}) + (a_{\tilde{i}}^2 + \tau a_{\tilde{j}})(-\frac{\tilde{\tau}}{\tau^2})$ . Let  $g'(\tau) = 0$ . We obtain  $\tau^2 = \epsilon^4 = \frac{a_{\tilde{i}}^2}{a_{\tilde{j}}^2}$ . We notice that when  $\epsilon \to 0$ ,  $g(\tau) \to \infty$ . Thus, when  $\epsilon = \pm \sqrt{\$  $g(\tau)$  obtains the minimum.

From Theorem [2.7](#page-6-1) and Lemma [2.8,](#page-6-2) we see that the largest eigenvalue [\(2.15\)](#page-6-3) is the same as that of  $(2.2)$ . In fact, we have the following result.

COROLLARY 2.9. If  $(\lambda, u)$  is an eigenpair of [\(2.2\)](#page-2-2), then  $(\lambda, u)$  is an eigenpair of  $(2.14).$  $(2.14).$ 

We explain the above corollary in more details. If  $(\lambda, z_{ij})$  is an eigenpair of [\(2.2\)](#page-2-2), where  $z_{ij} = c_i u_i + c_j u_j$  with  $\epsilon = \frac{c_j}{c_i}$  $\frac{c_j}{c_i}$ ,  $a_i a_j > 0$  and  $\epsilon^2 = \frac{a_i}{a_j}$ , one can show that  $\alpha(\Phi(z_{ij})) = \alpha(z_{ij})$ . Then, it follows that  $\Pi(z_{ij})z_{ij} = (I - \alpha(\check{\Phi}(z_{ij}))A)z_{ij}$   $(I \alpha(z_{ij})A|z_{ij} = \lambda z_{ij}$ . For general eigenpairs of [\(2.2\)](#page-2-2), we can use the properties in the proof of Theorem [2.4](#page-3-4) to process the proof here.

THEOREM 2.10. Suppose that  $A \in \mathbb{R}^{n \times n}$  is symmetric. Let  $i_1, i_2 \in \{1, 2, \cdots, p\}$ and  $i_1 \neq i_2$ . Consider

$$
u^* = \sum_{s=1}^{n_{i_1}} c_{i_1,s} u_{i_1,s} + \sum_{r=1}^{n_{i_2}} c_{i_2,r} u_{i_2,r}, \quad c_{i_1,1} c_{i_2,1} \neq 0,
$$

and  $u_{i_1 i_2} = c_{i_1} u_{i_1,1} + c_{i_2} u_{i_2,1}$ , where

(2.16) 
$$
c_{i_1}^2 = \sum_{s=1}^{n_{i_1}} c_{i_1,s}^2, \quad c_{i_2}^2 = \sum_{r=1}^{n_{i_1}} c_{i_1,r}^2.
$$

Then,

<span id="page-7-1"></span><span id="page-7-0"></span>
$$
\Pi(u^*)u^* = \lambda_{i_1i_2}u^*, \quad \Pi(u_{i_1i_2})u_{i_1i_2} = \lambda_{i_1i_2}u_{i_1i_2},
$$

where

(2.17) 
$$
\lambda_{i_1 i_2}(\epsilon_{i_1 i_2}) = \frac{(a_{i_2} - a_{i_1})^2}{a_{i_1}^2 + \epsilon_{i_1, i_2}^2 a_{i_2}^2} \frac{1}{1 + \frac{1}{\epsilon_{i_1 i_2}^2}}, \quad \epsilon_{i_1 i_2} = \frac{c_{i_2}}{c_{i_1}}.
$$

*Proof.* From Theorem [2.4,](#page-3-4) we have  $\alpha(u^*) = \alpha(u_{i_1 i_2})$ . Consider

$$
u_1^* = (I - \alpha(u^*)A)u^* = \sum_{s=1}^{n_{i_1}} (1 - \alpha(u^*)a_{i_1})c_{i_1,s}u_{i_1,s} + \sum_{r=1}^{n_{i_2}} (1 - \alpha(u^*)a_{i_2})c_{i_2,r}u_{i_2,r},
$$

and

$$
u_{i_1i_2,1} = (I - \alpha(u_{i_1i_2})A)u_{i_1i_2} = (1 - \alpha(u^*)a_{i_1})c_{i_1}u_{i_1,1} + (1 - \alpha(u^*)a_{i_2})c_{i_2}u_{i_2,1}.
$$

We have

$$
\sum_{s=1}^{n_{i_1}}(1-\alpha(u^*)a_{i_1})^2c_{i_1,s}^2+\sum_{r=1}^{n_{i_2}}(1-\alpha(u^*)a_{i_2})^2c_{i_2,r}^2=(1-\alpha(u^*)a_{i_1})^2c_{i_1}^2+(1-\alpha(u^*)a_{i_2})^2c_{i_2}^2.
$$

It follows that  $\alpha((I - \alpha(u^*)A)u^*) = \alpha((I - \alpha(u_{i_1i_2})A)u_{i_1i_2}).$ 

From Theorem [2.7,](#page-6-1) we know that  $(\lambda_{i_1 i_2}(\epsilon_{i_1 i_2}), u_{i_1 i_2})$  is eigenpair of [\(2.14\)](#page-6-0). For simplicity, let  $\beta_2 = \alpha(u_1^*)$  and  $\beta_1 = \alpha(u^*)$ . We have

$$
(1 - \beta_2 a_{i_1})(1 - \beta_1 a_{i_1})c_{i_1}u_{i_1,1} = \lambda_{i_1 i_2}c_{i_1}u_{i_1,1},
$$
  

$$
(1 - \beta_2 a_{i_2})(1 - \beta_1 a_{i_2})c_{i_2}u_{i_2,1} = \lambda_{i_1 i_2}c_{i_2}u_{i_2,1}.
$$

It means

$$
(1 - \beta_2 a_{i_1})(1 - \beta_1 a_{i_1}) = \lambda_{i_1 i_2}
$$
 and  $(1 - \beta_2 a_{i_2})(1 - \beta_1 a_{i_2}) = \lambda_{i_1 i_2}$ .

Thus,

$$
\Pi(u^*)u^* = \sum_{s=1}^{n_{i_1}} (1 - \beta_2 a_{i_1})(1 - \beta_1 a_{i_1})c_{i_1,s}u_{i_1,s} + \sum_{r=1}^{n_{i_2}} (1 - \beta_2 a_{i_2})(1 - \beta_1 a_{i_2})c_{i_2,r}u_{i_2,r}
$$
  
\n
$$
= \sum_{s=1}^{n_{i_1}} \lambda_{i_1 i_2} c_{i_1,s}u_{i_1,s} + \sum_{r=1}^{n_{i_2}} \lambda_{i_1 i_2} c_{i_2,r}u_{i_2,r}
$$
  
\n
$$
= \lambda_{i_1 i_2} \left( \sum_{s=1}^{n_{i_1}} c_{i_1,s}u_{i_1,s} + \sum_{r=1}^{n_{i_2}} c_{i_2,r}u_{i_2,r} \right)
$$
  
\n
$$
= \lambda_{i_1 i_2} u^*,
$$

which completes the proof.

<span id="page-8-2"></span>THEOREM 2.11. Suppose that A is symmetric. If A is definite, then all the eigenvectors of  $(2.14)$  associated with nonzero eigenvalue are given by

<span id="page-8-0"></span>
$$
(2.18) \t u^* = \sum_{s=1}^{n_{i_1}} c_{i_1,s} u_{i_1,s} + \sum_{r=1}^{n_{i_2}} c_{i_2,r} u_{i_2,r}, \quad \text{where} \quad i_1 \neq i_2, i_1, i_2 \in \{1, 2, \cdots, p\},
$$

where at least one term in  ${c_{i_1,s}}_{s=1}^{n_{i_1}}$  is nonzero and at least one term in  ${c_{i_2,r}}_{r=0}^{n_{i_2}}$  is nonzero. Moreover, all the eigenvalues are less than one and are given by  $(2.17)$ .

If A is indefinite, then all the eigenvectors of  $(2.14)$  associated with non-unit eigenvalues are given by  $(2.18)$ , and the eigenvectors associated with eigenvalue one are given by  $u^*$ , where  $\langle Au^*, u^* \rangle = 0$ .

Proof. For any  $i \in \{1, 2, \dots, p\}$ , let  $u_* = \sum_s^{n_i} c_{i,s} u_{i,s} \neq 0$ . Then,  $(I - \alpha(u_*)A)u_* =$ 0. So, such  $u_*$  cannot be eigenvectors of  $(2.14)$   $(\alpha(0))$  is not defined).

o, such  $u_*$  cannot be eigenvectors of [\(2.14\)](#page-6-0)  $(\alpha(0)$  is not defined).<br>Assume that the eigenvector of (2.14) is given by  $u^* = \sum_{i=1}^p \sum_{s=1}^{n_i} c_{i,s} u_{i,s}$ , which contains at least two different integers  $i, j$  such that  $c_{i,s}c_{j,t} \neq 0$ .

When A is indefinite, there exists a vector v such that  $\langle Av, v \rangle = 0$ . Then  $\alpha(v) = 0$ and  $\Pi(v) = v$ . In this situation,  $\lambda = 1$  is an eigenvalue of [\(2.14\)](#page-6-0).

Next, we prove that in  $u^*$ , there are at most two different integers  $i, j$  such that  $c_{i,s}c_{j,t} \neq 0$ . For simplicity, let  $\beta_1 = \alpha(u^*)$  and  $\beta_2 = \alpha(\Phi(u^*)) = \alpha((I - \alpha(u^*)A)u^*)$ . Using  $\Pi(u^*)u^* = \lambda u^*, \lambda \neq 0$ , we have

<span id="page-8-1"></span>
$$
(2.19) \ c_{i,s}(1-\beta_2 a_i)(1-\beta_1 a_i)u_{i,s} = \lambda c_{i,s}u_{i,s}, \quad (i,s) \in \{1,2,\cdots,p\} \times \{1,2,\cdots,n_i\}.
$$

If  $\beta_1 = 0$ , then  $\beta_2 = 0$ . If  $\beta_1 \beta_2 \neq 0$ , let us consider the quadratic function  $g(b) =$  $(1-\beta_2b)(1-\beta_1b)$ . For a given value  $g^*$ , there are at most two different values  $b_1, b_2$  such that  $g(b_1) = g(b_2) = g^*$ . To guarantee [\(2.19\)](#page-8-1), there are at most two different integers i, j such that  $c_{i,s}c_{j,t} \neq 0$ , where  $s \in \{1, 2, \dots, n_i\}$  and  $t \in \{1, 2, \dots, n_j\}$ . If  $\beta_1 = 0$ and  $\beta_2 \neq 0$ , to guarantee [\(2.19\)](#page-8-1), there is only at most one integer  $i \in \{1, 2, \dots, p\}$ such that  $c_{i,s} \neq 0$ , which means that  $u^* = u_p$ . However, we know for such  $u_p$ ,  $(I - \alpha(u_p)A)u_p = 0$  and  $\Pi(u_p)$  is not defined. Thus, all the eigenvectors of [\(2.14\)](#page-6-0) associated with non-unit eigenvalues are any linear combination of eigenvectors of A associated with exactly two distinguish eigenvalues of A.  $\Box$ 

COROLLARY 2.12. If the initial residual  $r_0 = u^*$  is given by [\(2.18\)](#page-8-0), then the corresponding root-convergence factor is

$$
\varrho_k(r_0) = \left( \frac{(a_{i_2} - a_{i_1})^2}{a_{i_1}^2 + \epsilon^2 a_{i_2}^2} \frac{1}{1 + \frac{1}{\epsilon_{i_1 i_2}^2}} \right)^{1/2},
$$

where  $\epsilon_{i_1 i_2}^2 =$  $\frac{c_{i_2}}{2}$  $c_{i_1}$  $\sqrt{2}$ with  $c_{i_1}$  and  $c_{i_2}$  given by  $(2.16)$ .

THEOREM 2.13. Suppose that  $A$  is symmetric. If  $A$  is indefinite,

(2.20) 
$$
\max_{v \neq 0} \frac{\|\Pi(v)v\|}{\|v\|} = 1.
$$

If A is definite,

$$
\max_{v \neq 0} \frac{\|\Pi(v)v\|}{\|v\|} = (\varrho_{GMRES(1)}^*)^2,
$$

where  $\rho_{GMRES(1)}^*$  is given by [\(2.13\)](#page-5-2).

*Proof.* Let  $(I - \alpha(v)A)v = v - \frac{\langle Av, v \rangle}{\langle Av, Av \rangle}Av = \Phi(v)$ . If  $\Phi(v) = 0$ ,  $\Pi(v)v$  is not defined. Using [\(2.3\)](#page-3-0), we have

$$
\max_{v \neq 0} \frac{\|\Pi(v)v\|}{\|v\|} = \max_{v \neq 0} \frac{\|(I - \alpha(\Phi(v))A)\Phi(v)\|}{\|v\|} \n= \max_{v \neq 0} \frac{\|(I - \alpha(\Phi(v))A)\Phi(v)\|}{\|\Phi(v)\|} \frac{\|\Phi(v)\|}{\|v\|} \n\leq \max_{y \neq 0} \frac{\|T(y)y\|}{\|y\|} \max_{v \neq 0} \frac{\|T(v)v\|}{\|v\|} \n= \varrho_{GMRES(1)}^* \varrho_{GMRES(1)}^* = (\varrho_{GMRES(1)}^*)^2.
$$

Using  $(2.14)$  and Lemma [2.8,](#page-6-2) we have

$$
\max_{v \neq 0} \frac{\|\Pi(v)v\|}{\|v\|} \ge \max_{\lambda} \|\lambda\| = (\varrho_{GMRES(1)}^*)^2.
$$

Thus, we obtain the desired result.

THEOREM 2.14. Consider GMRES(1) applied to  $Ax = b$ , where  $A \in \mathbb{R}^{2 \times 2}$  is symmetric. Let  $r_0 = c_1u_1 + c_2u_2$  be the initial residual. Then, if  $c_1c_2 = 0$ ,

 $\rho(r_0) = 0.$ 

If  $c_1c_2 \neq 0$ , let  $\epsilon = \frac{c_2}{c_1}$ . Then

(2.21) 
$$
\varrho(r_0) = \sqrt{\lambda_{12}(\epsilon)} = \left(\frac{(a_2 - a_1)^2}{(a_1^2 + \epsilon^2 a_2^2)(1 + \frac{1}{\epsilon^2})}\right)^{1/2}
$$

THEOREM 2.15. For GMRES(1) applied to  $Ax = b$ , where A is symmetric, the worst-case root-convergence factor is

.

 $\Box$ 

<span id="page-9-0"></span>(2.22) 
$$
\varrho_{GMRES(1)}^* = \max_{i \neq j, i, j \in \{1, 2, \cdots, p\}} \frac{|a_j - a_i|}{|a_j| + |a_i|} = \left| \frac{\lambda_{\max}(A) - \lambda_{\min}(A)}{\lambda_{\max}(A) + \lambda_{\min}(A)} \right|,
$$

if A is definite, and  $\varrho_{GMRES(1)}^* = 1$  if A is indefinite.

*Proof.* Consider  $r_0$  such that  $\Pi(r_0)r_0 = (\Gamma^*)^2 r_0$ , where  $\Gamma^*$  is defined in [\(2.12\)](#page-5-3). Then, we have  $r_{2k} = (\Gamma^*)^{2k} r_0$ . It follows that  $\varrho(r_0) = \limsup_{k \to \infty} ||r_{2k}||^{1/(2k)} = \Gamma^*$ .

When  $A$  is definite, if the initial residual is only associated with the first  $s$  eigenvalues  ${a_i}_{i=1}^s$ , then the root-convergence factor is given by  $(2.22)$  with  $i, j \in \{1, 2, \dots, s\}$ . Although when A is indefinite, the worst-case root-convergence factor is one, there may exist some good initial guesses such that the corresponding root-convergence can be less than one.

**2.3.** Some results of  $rAA(1)$ . Anderson acceleration is a widely used method to speed up the convergence of fixed-point iterations and it has garnered a lot of attention for its convergence analysis [\[38,](#page-20-23) [34,](#page-20-21) [2,](#page-19-12) [10,](#page-19-13) [31\]](#page-20-24).

In [\[6\]](#page-19-11), the authors study the asymptotic root-convergence factor of restarted  $AA(1)$ , denoted as r $AA(1)$ . For completeness, we give r $AA(1)$  iterations as follows:

$$
x_{k+1} = Mx_k + b, \quad k = 0, 2, 4, \cdots
$$
  
\n
$$
x_{k+1} = Mx_k + \frac{\langle Ar_k, r_k \rangle}{\langle Ar_k, Ar_k \rangle} M(x_k - x_{k-1}) + b, \quad \text{where } r_k = Ax_k - b, \ k = 1, 3, 5, \cdots
$$

It can be shown that rAA(1) residuals satisfy  $r_{k+2} = M(I - \alpha(r_k)A)r_k$ . To analyze its asymptotic convergence factor, the authors of [\[6\]](#page-19-11) consider the following vectordependent matrix

$$
\Upsilon(v) = M(I - \alpha(\Psi(v))A)M(I - \alpha(v)A),
$$

where

<span id="page-10-0"></span>
$$
\Psi(v) = M(I - \alpha(v)A)v, \quad v \neq 0,
$$

and the corresponding nonlinear eigenvalue problem is

$$
\Upsilon(u)u = \mu u.
$$

The residuals of rAA(1) satisfy  $r_{4(k+1)} = \Upsilon(r_{4k})r_{4k}$ . There remain questions about the eigenpairs of [\(2.23\)](#page-10-0). [\[6\]](#page-19-11) proposes some conjectures. We will derive more results of  $rAA(1)$  when A is symmetric and address one of the conjectures from [\[6\]](#page-19-11).

We first consider the following nonlinear eigenvalue problem:

(2.24) 
$$
\Psi(u)u = M(I - \alpha(u)A)u = \mu u.
$$

We are interested in eigenpairs ( $\mu \neq 0$ ) of [\(2.24\)](#page-10-1). We have the following results.

<span id="page-10-2"></span>THEOREM 2.16. Suppose that A is symmetric. Let  $i_1, i_2 \in \{1, 2, \dots, p\}$  with  $i_1 \neq i_2$  $i_2$  and  $\frac{-(a_{i_2}-1)a_{i_1}}{(a_{i_1}-1)a_{i_2}} > 0$ . Consider

<span id="page-10-1"></span>
$$
u^* = \sum_{s=1}^{n_{i_1}} c_{i_1,s} u_{i_1,s} + \sum_{r=1}^{n_{i_2}} c_{i_2,r} u_{i_2,r}, \quad c_{i_1,1} c_{i_2,1} \neq 0,
$$

and

$$
u_{i_1i_2} = c_{i_1}u_{i_1,s_1} + c_{i_2}u_{i_2,r_1} = c_{i_1}(u_{i_1,s_1} + \epsilon u_{i_2,r_1}),
$$

where 
$$
\epsilon = \frac{c_{i_2}}{c_{i_1}}, \epsilon^2 = \frac{-(a_{i_2}-1)a_{i_1}}{(a_{i_1}-1)a_{i_2}},
$$
 and  
\n
$$
c_{i_1}^2 = \sum_{s=1}^{n_{i_1}} c_{i_1,s}^2, \quad c_{i_2}^2 = \sum_{r=1}^{n_{i_1}} c_{i_1,r}^2, \quad s_1 \in \{1, 2, \cdots, n_{i_1}\}, \quad s_2 \in \{1, 2, \cdots, n_{i_2}\}.
$$

Then,  $u^*$  and  $u_{i_1i_2}$  are eigenvectors of  $(2.24)$  associated with the same eigenvalue, i.e.,

<span id="page-11-0"></span>(2.25) 
$$
\Psi(u^*)u^* = \mu_{i_1i_2}u^*, \quad \Psi(u_{i_1i_2})u_{i_1i_2} = \mu_{i_1i_2}u_{i_1i_2},
$$

where

(2.26) 
$$
\mu_{i_1 i_2}(\epsilon_{i_1 i_2}) = \frac{(1 - a_{i_1})(1 - a_{i_2})(a_{i_1} - a_{i_2})}{(a_{i_1} - 1)a_{i_1} - (a_{i_2} - 1)a_{i_2}}.
$$

Moreover, all the eigenpairs  $(\mu \neq 0)$  of  $(2.24)$  are given by  $(\mu_{i_1,i_2}, u^*)$ .

*Proof.* From Theorem [2.4,](#page-3-4) we have  $\alpha(u^*) = \alpha(u_{i_1 i_2}) := \beta$ . Using the second equation in  $(2.25)$  gives us

$$
(1 - a_{i_1})(1 - \beta a_{i_1})c_{i_1}u_{i_1,1} = \mu_{i_1i_2}c_{i_1}u_{i_1,1} \text{ and } (1 - a_{i_2})(1 - \beta a_{i_2})c_{i_2}u_{i_2,1} = \mu_{i_1i_2}c_{i_2}u_{i_2,1}.
$$

Since  $c_{i_1}c_{i_2}\neq 0$ ,

<span id="page-11-1"></span>(2.27) 
$$
(1 - a_{i_1})(1 - \beta a_{i_1}) = (1 - a_{i_2})(1 - \beta a_{i_2}).
$$

Solving the above equation for  $\beta$ , we obtain  $\beta = \frac{1}{a_{i_2} + a_{i_1} - 1}$ . Recall  $\beta = \frac{a_{i_1} + a_{i_2} \epsilon^2}{a_{i_1}^2 + a_{i_2}^2 \epsilon^2}$  $\frac{a_{i_1} + a_{i_2}^2}{a_{i_1}^2 + a_{i_2}^2 \epsilon^2}$ . We further obtain  $\epsilon^2 = \frac{-(a_{i_2}-1)a_{i_1}}{(a_{i_2}-1)a_{i_2}}$  $\frac{-(a_{i_2}-1)a_{i_1}}{(a_{i_1}-1)a_{i_2}}$ .

Thus,

$$
\Psi(u^*)u^* = \sum_{s=1}^{n_{i_1}} (1 - a_{i_1})(1 - \beta a_{i_1})c_{i_1,s}u_{i_1,s} + \sum_{r=1}^{n_{i_2}} (1 - a_{i_2})(1 - \beta a_{i_2})c_{i_2,r}u_{i_2,r}
$$
  
\n
$$
= \sum_{s=1}^{n_{i_1}} \mu_{i_1i_2}c_{i_1,s}u_{i_1,s} + \sum_{r=1}^{n_{i_2}} \mu_{i_1i_2}c_{i_2,r}u_{i_2,r}
$$
  
\n
$$
= \mu_{i_1i_2} \left( \sum_{s=1}^{n_{i_1}} c_{i_1,s}u_{i_1,s} + \sum_{r=1}^{n_{i_2}} c_{i_2,r}u_{i_2,r} \right)
$$
  
\n
$$
= \mu_{i_1i_2}u^*.
$$

To satisfy  $(2.27)$ , there are at most two different  $a_{i_1}, a_{i_2}$  existing. Thus, all eigenpairs with  $\mu \neq 0$  of [\(2.24\)](#page-10-1) are given by  $(\mu_{i_1,i_2}, u^*)$ .  $\Box$ 

In the above theorem, we require that  $\frac{-(a_{i_2}-1)a_{i_1}}{(a_{i_1}-1)a_{i_2}} > 0$ . However, if the eigenvalues of  $A$  do not satisfy this condition, then it means that  $(2.24)$  does not have nonzero eigenvalue. This motivates us to examine the eigenpairs of [\(2.23\)](#page-10-0) below.

<span id="page-11-2"></span>THEOREM 2.17. Suppose that A is symmetric. Let  $i_1, i_2 \in \{1, 2, \dots, p\}$  with  $i_1 \neq i_2$ i2, and consider

$$
u^* = \sum_{s=1}^{n_{i_1}} c_{i_1,s} u_{i_1,s} + \sum_{r=1}^{n_{i_2}} c_{i_2,r} u_{i_2,r}, \quad c_{i_1,1} c_{i_2,1} \neq 0,
$$

and

$$
u_{i_1i_2} = c_{i_1}u_{i_1,s_1} + c_{i_2}u_{i_2,r_1},
$$

where

<span id="page-12-0"></span>
$$
(2.28) \quad c_{i_1}^2 = \sum_{s=1}^{n_{i_1}} c_{i_1,s}^2, \quad c_{i_2}^2 = \sum_{r=1}^{n_{i_1}} c_{i_1,r}^2, \quad s_1 \in \{1, 2, \cdots, n_{i_1}\}, \quad s_2 \in \{1, 2, \cdots, n_{i_2}\}.
$$

Then,  $u^*$  and  $u_{i_1i_2}$  are eigenvectors of  $(2.23)$  associated with the same eigenvalue, i.e.,

<span id="page-12-1"></span>(2.29) 
$$
\Upsilon(u_{i_1i_2})u_{i_1i_2} = \mu_{i_1i_2}u_{i_1i_2},
$$

and

<span id="page-12-4"></span>(2.30) 
$$
\Upsilon(u^*)u^* = \mu_{i_1i_2}u^*,
$$

where

<span id="page-12-2"></span>(2.31) 
$$
\mu_{i_1 i_2}(\epsilon_{i_1 i_2}) = \frac{(a_{i_2} - a_{i_1})^2}{a_{i_1}^2 + \epsilon_{i_1 i_2}^2 a_{i_2}^2} \frac{((1 - a_{i_1})(1 - a_{i_2}))^2}{(1 - a_{i_1})^2 + \frac{(1 - a_{i_2})^2}{\epsilon^2}}, \quad \epsilon_{i_1 i_2} = \frac{c_{i_1}}{c_{i_2}},
$$

and (2.32)

<span id="page-12-3"></span>2.32)  
\n
$$
\max_{i_1 i_2} \mu_{i_1 i_2}(\epsilon_{i_1 i_2}) = \mu_{i_1 i_2} \left( \pm \sqrt{\left| \frac{(1 - a_{i_2}) a_{i_1}}{(1 - a_{i_1}) a_{i_2}} \right|} \right) = \left( \frac{(1 - a_{i_1})(1 - a_{i_2})(a_{i_1} - a_{i_2})}{|(1 - a_{i_1}) a_{i_1}| + |(1 - a_{i_2}) a_{i_2}|} \right)^2.
$$

*Proof.* From Theorem [2.4,](#page-3-4) we have  $\alpha(u^*) = \alpha(u_{i_1 i_2})$ . Consider

$$
u_1^* = \Psi(u^*) = \sum_{s=1}^{n_{i_1}} (1 - a_{i_1})(1 - \alpha(u^*)a_{i_1})c_{i_1,s}u_{i_1,s} + \sum_{r=1}^{n_{i_2}} (1 - a_{i_2})(1 - \alpha(u^*)a_{i_2})c_{i_2,r}u_{i_2,r},
$$

and

$$
u_{i_1i_2,1} = \Psi(u_{i_1i_2}) = (1 - a_{i_1})(1 - \alpha(u^*)a_{i_1})c_{i_1}u_{i_1,1} + (1 - a_{i_2})(1 - \alpha(u^*)a_{i_2})c_{i_2}u_{i_2,1}.
$$

Using condition  $(2.28)$ , we have

$$
\sum_{s=1}^{n_{i_1}} (1 - a_{i_1})^2 (1 - \alpha(u^*)a_{i_1})^2 c_{i_1, s}^2 + \sum_{r=1}^{n_{i_2}} (1 - a_{i_2})^2 (1 - \alpha(u^*)a_{i_2})^2 c_{i_2, r}^2
$$
  
= 
$$
(1 - a_{i_1})^2 (1 - \alpha(u^*)a_{i_1})^2 c_{i_1}^2 + (1 - a_{i_2})^2 (1 - \alpha(u^*)a_{i_2})^2 c_{i_2}^2.
$$

It follows that  $\alpha(u_1^*) = \alpha(u_{i_1 i_2,1})$ . For simplicity, let  $\beta_2 = \alpha(u_1^*)$  and  $\beta_1 = \alpha(u^*)$ . The results  $(2.29)$ ,  $(2.31)$ , and  $(2.32)$  are from [\[6\]](#page-19-11). Using  $(2.29)$  gives us

$$
(1 - a_{i_1})^2 (1 - \beta_2 a_{i_1}) (1 - \beta_1 a_{i_1}) c_{i_1} u_{i_1,1} = \mu_{i_1 i_2} c_{i_1} u_{i_1,1},
$$
  

$$
(1 - a_{i_2})^2 (1 - \beta_2 a_{i_2}) (1 - \beta_1 a_{i_2}) c_{i_2} u_{i_2,1} = \mu_{i_1 i_2} c_{i_2} u_{i_2,1}.
$$

Since  $c_{i_1}c_{i_2}\neq 0$ ,

$$
(1 - a_{i_1})^2 (1 - \beta_2 a_{i_1}) (1 - \beta_1 a_{i_1}) = \mu_{i_1 i_2}, \quad (1 - a_{i_2})^2 (1 - \beta_2 a_{i_2}) (1 - \beta_1 a_{i_2}) = \mu_{i_1 i_2}.
$$

Similar to the proof of Theorem [2.16,](#page-10-2) we can obtain  $(2.30)$ .

From Theorems [2.16](#page-10-2) and [2.17,](#page-11-2) we see that the eigenvectors of  $(2.24)$  are also eigenvectors of  $(2.23)$ .

Remark 2.18. It can be shown that when the maximum of  $\mu_{i_1,i_2}(\epsilon_{i_1i_2})$  is achieved, see [\(2.32\)](#page-12-3),  $\beta_1 = \beta_2$ .

<span id="page-13-2"></span>Theorem 2.19. Suppose that A is symmetric. Then, all the eigenvectors of [\(2.23\)](#page-10-0) associated with nonzero eigenvalues are given by

<span id="page-13-0"></span>(2.33) 
$$
u = \sum_{i \in \{1, 2, \cdots, p\}} \sum_{s=1}^{n_i} c_{i,s} u_{i,s},
$$

where  $2 \leq \#\{i\} \leq 4$ , and at least one term in  ${c_{i_1,s}}_{s=1}^{n_{i_1}}$  is nonzero and at least one term in  ${c_{i_2,r}}_{r=0}^{n_{i_2}}$  is nonzero.

The formula [\(2.33\)](#page-13-0) means that the eigenvectors of [\(2.23\)](#page-10-0) are any linear combination of eigenvectors of A associated with at least two and at most four distinguish eigenvalues of A.

Proof. As shown in Theorem [2.11,](#page-8-2)  $u_* = \sum_{s=1}^{n_i} c_{i,s} u_{i,s} \neq 0$ , where  $i \in \{1, 2, \dots, p\}$ cannot be eigenvectors of  $(2.23)$   $(\alpha(0)$  is not defined). Assume that the eigenvector of [\(2.23\)](#page-10-0) is given by

<span id="page-13-1"></span>(2.34) 
$$
u^* = \sum_{i=1}^p \sum_{s=1}^{n_i} c_{i,s} u_{i,s},
$$

which contains at least two different integers i, j such that  $c_{i,s}c_{j,t} \neq 0$ . Next, we prove that in [\(2.34\)](#page-13-1), there are at most four different integers *i*. For simplicity, let  $\beta_1 = \alpha(u^*)$ and  $\beta_2 = \alpha(M(I - \alpha(u^*)A))u^*$ ). We know that  $\beta_1\beta_2 \neq 0$ . Using  $\Upsilon(u^*)u^* = \lambda u^*$ with  $\lambda \neq 0$ , we have  $(2.35)$ 

$$
c_{i,s}(1-a_i)^2(1-\beta_2a_i)(1-\beta_1a_i)u_{i,s}=\lambda c_{i,s}u_{i,s}, \forall (i,s)\in\{1,2,\cdots,p\}\times\{1,2,\cdots,n_i\}.
$$

Consider the quartic function  $g(b) = (1 - \beta_2 b)(1 - \beta_1 b)(1 - b)^2$ . For a given value  $g^*$ , there are at most four different values  $b_1$ ,  $b_2$  such that  $g(b_1) = g(b_2) = g(b_3) = g(b_4)$  =  $g^*$ . To guarantee [\(2.19\)](#page-8-1), there are at most four different integers  $i, j, k, \ell$  such that  $c_{i,s_1}c_{j,t_1}c_{k,s_2}c_{\ell,t_2} \neq 0$ , where  $s \in \{1, 2, \cdots, n_i\}$  and  $t \in \{1, 2, \cdots, n_j\}$ .

Thus, all the eigenvectors of [\(2.23\)](#page-10-0) associated with nonzero eigenvalues are any linear combination of eigenvectors of A associated with at least two and at most four distinguish eigenvalues of A.  $\Box$ 

We comment that in Theorem  $2.19$  we do not know whether  $(2.33)$  can contain three or four eigenvectors associated with three or four distinguish eigenvalues of A. If there exists, then what the corresponding eigenvalues of  $\Upsilon$  are. We leave these questions for the future.

Define

<span id="page-13-3"></span>
$$
\Lambda^* = \max_{i,j \in \{1,2,\cdots,p\}} \left( \frac{(1-a_i)(1-a_j)(a_i-a_j)}{|(1-a_i)a_i| + |(1-a_j)a_j|} \right)^2.
$$

In  $[6]$ , it has shown that worst-case convergence factor of  $\text{rAA}(1)$  has a lower bound  $(\Lambda^*)^{1/4}$ . The numerical results in that work seem to indicate that  $(\Lambda^*)^{1/4}$  is the worst-case convergence factor. Moreover, it includes a conjecture stating that

(2.36) 
$$
\max_{v \neq 0} \frac{\|\Upsilon(v)v\|}{\|v\|} = \Lambda^*.
$$

In this following, we provide three counterexamples for  $(2.36)$ . We consider A being  $A_i$  as follows.

EXAMPLE 2.1 (counterexamples).

$$
\mathcal{A}_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad \mathcal{A}_2 = \begin{pmatrix} -1/2 & 0 & 0 \\ 0 & -4 & 0 \\ 0 & 0 & 2 \end{pmatrix}, \quad \mathcal{A}_3 = \begin{pmatrix} 1/2 & 0 & 0 & 0 \\ 0 & 3/2 & 0 & 0 \\ 0 & 0 & 1/3 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix}.
$$

First, it can be shown that  $\Lambda^*(\mathcal{A}_1) = \frac{1}{16} = 0.0625$ . Consider  $v_1 = \begin{bmatrix} 15, & 5, & 1 \end{bmatrix}^T$  and  $M_1 = I - A_1$ . Then, "

$$
\alpha(v_1) = \frac{v_1^T A_1 v_1}{v_1^T A_1^T A_1 v_1} = \frac{139}{167}, \quad u = M_1 (I - \alpha(v_1) A_2) v_1 = \begin{bmatrix} 0, & \frac{555}{167}, & \frac{500}{167} \end{bmatrix}^T,
$$
  
\n
$$
\alpha(u) = \frac{u^T A_1 u}{u^T A_1^T A_1 u} = \frac{\frac{1366050}{27889}}{\frac{3482100}{27889}} = \frac{1366050}{3482100} = \frac{27321}{69642},
$$
  
\n
$$
w = M_1 (I - \alpha(u) A_1) u = \begin{bmatrix} 0, & \frac{-1387500}{1938369}, & \frac{205350}{193837} \end{bmatrix} \approx \begin{bmatrix} 0 & -0.7158 & 1.0594 \end{bmatrix}^T.
$$

Thus,  $\frac{\|\Upsilon(v_1)v_1\|}{\|v_1\|} \approx \frac{1.2786}{15.8430} \approx 0.0807 > \Lambda^*(\mathcal{A}_1) = 0.0625$ . For more accuracy, one may verify if  $\|\Upsilon(v_1)v_1\|^2 = \|v_1\|^2 (\Lambda^*)^2$ , that is,  $\|w\|^2 16^2 = \|v_1\|^2$ , which gives

 $16^2((1387500\times193837)^2+(205350\times1838369)^2)=(15^2+5^2+1^2)(1938369\times193837)^2$ .

We notice that the left-hand side is even, while the right-hand side is odd; therefore, they cannot be equal.

Consider  $v_2 = [38, 1, 45]^T$  for  $\mathcal{A}_2$  and  $v_3 = [23, 60, 77, 1]^T$  for  $\mathcal{A}_3$ . It can be shown that

$$
\Lambda^*(\mathcal{A}_2) = 225/121 \approx 1.8595, \quad \frac{\|\Upsilon(v_2)v_2\|}{\|v_2\|} \approx 2.1598 > \Lambda^*(\mathcal{A}_2),
$$
  

$$
\Lambda^*(\mathcal{A}_3) = \frac{49}{81} \approx 0.6049, \quad \frac{\|\Upsilon(v_3)v_3\|}{\|v_3\|} \approx 0.6388 > \Lambda^*(\mathcal{A}_3).
$$

Although  $\max_{v \neq 0}$  $\|\Upsilon(v)v\|$  $\frac{\partial^2 v}{\partial x^2} > \Lambda^*$ , this does not indicate that the worst-case root- $\sqrt{1/4}$ 

convergence of rAA(1) is larger or equals  $\left(\max_{v\neq 0} \frac{\|\Upsilon(v)v\|}{\|v\|}\right)$  $\|v\|$ . Here, we cannot have a similar Lemma [2.1](#page-2-3) for rAA(1) since  $\alpha(v)$  for rAA(1) does not minimize the residual. Note that for  $A_1$ ,  $w = \Upsilon(v_1)v_1$  is an eigenvector of [\(2.23\)](#page-10-0). From Theorem [2.17,](#page-11-2) we have  $\Upsilon(w)w = \mu w$ . This means that for  $\mathcal{A}_1^{-1}v_1$  as an initial guess for solving  $A_1x = 0$ , the corresponding root-convergence factor of rAA(1) is  $\rho = (\mu)^{1/4} < \left( \max_{v \neq 0} \frac{\|\Upsilon(v)v\|}{\|v\|} \right)$  $\|v\|$  $\frac{1}{4}$ .

<span id="page-14-0"></span>3. Skew-symmetric case of  $M$  for  $GMRES(1)$ . In this section, we discuss the situation that  $M = I - A \in \mathbb{R}^{n \times n}$  is skew-symmetric, i.e.,  $M = -M^T$ . For simplicity, we assume that  $n$  is even. Then, the eigenvalues of  $M$  are imaginary. Denote the eigenvalues of M as  $\{m_j, \bar{m}_j\}_{j=0}^{n/2}$ , where  $\bar{m}_j$  is the conjugate of  $m_j$ . We define

$$
m_* = \min_{j=1,2,\cdots,n/2} \{|m_j|\}, \quad m^* = \max_{j=1,2,\cdots,n/2} \{|m_j|\}.
$$

Let  $M = QHQ^T$  be the real Schur decomposition of M, where  $H \in \mathbb{R}^{n \times n}$  is a block diagonal matrix with the diagonal blocks of size  $2 \times 2$ . The eigenvalues of the  $2 \times 2$ diagonal blocks are the eigenvalues of  $M$ .  $Q$  is an orthogonal matrix with

<span id="page-15-5"></span>(3.1) 
$$
Q = [Q_1, \cdots, Q_{n/2}], \quad Q_j = [q_j, \tilde{q}_j].
$$

From [\[6\]](#page-19-11), we know that M is invariant on the subspaces span $\{Q_j\}$  for  $j = 1, 2, \cdots, n/2$ and the eigenvalues of  $S = \frac{1}{2}(A^{-1}+A^{-T})$  are  $\{(1+|m_j|^2)^{-1}\}_{j=1}^{n/2}$ . We cite some results from  $[6]$ , which will be used for our analysis.

LEMMA 3.1. [\[6\]](#page-19-11) Suppose that M is skew-symmetric and  $q \in \text{span}\{Q_j\}$ . Then LEMMA 3.1. [6] Suppose that M is skew-symmetric and  $q \in \text{span}\{Q_j\}$ . Then<br>  $\alpha(q) = \frac{1}{1+|m_j|^2}$ , and for general  $0 \neq v \in \mathbb{R}^n$ , we have  $\alpha(v) \in \left[\frac{1}{1+(m^*)^2}, \frac{1}{1+(m^*)^2}\right] \subseteq$  $(0, 1].$ 

<span id="page-15-0"></span>THEOREM 3.2. Suppose that M is skew-symmetric. If  $0 \neq q \in \text{span}\{Q_j\}$ , then

<span id="page-15-3"></span>
$$
\Phi(q) \in \text{span}\{Q_j\} \quad and \quad \frac{\|\Phi(q)\|}{\|q\|} = \frac{|m_j|}{\sqrt{1+|m_j|^2}}.
$$

Moreover,

(3.2) 
$$
\max_{v \neq 0} \frac{\|\Phi(v)\|}{\|v\|} \leq \frac{m^*}{\sqrt{1 + (m^*)^2}}.
$$

Proof. The proof can be derived following [\[6\]](#page-19-11). Thus, we omit it here.

<span id="page-15-1"></span>COROLLARY 3.3. Consider GMRES(1) applied to linear system  $Ax = b$  with  $M = I - A$  being skew-symmetric. Then, the worst-case root-convergence factor of GMRES(1) is

(3.3) 
$$
\varrho_{GMRES(1),ss}^* = \frac{m^*}{\sqrt{1 + (m^*)^2}} < 1.
$$

*Proof.* Recall that  $r_k = \Phi(r_{k-1})$ . From Theorem [3.2,](#page-15-0) we have

<span id="page-15-2"></span>
$$
||r_k|| \le \frac{m^*}{\sqrt{1 + (m^*)^2}} ||r_{k-1}|| \le \cdots \le \left(\frac{m^*}{\sqrt{1 + (m^*)^2}}\right)^k ||r_0||.
$$

From the first result in Theorem [3.2,](#page-15-0) when  $r_0 \in \text{span}\{Q_j\}$ , we have  $r_k = \Phi(r_{k-1}) \in$ span $\{Q_j\}$ . Thus,

$$
\varrho_{GMRES(1),ss}^{*} = \max_{r_0}(\limsup_{k \to \infty} \Gamma_k(r_0)) = \frac{m^*}{\sqrt{1 + (m^*)^2}},
$$

which completes the proof.

The above result tells us that  $GMRES(1)$  always converges if  $M$  is skew-symmetric. In  $[6]$ , the authors consider rAA(1), and show that for the case where M is skewsymmetric, the worst-case root-convergence factor of rAA(1) is  $\varrho_{rAA(1)}^* = \frac{m^*}{(1 + (m^*)^2)^2}$ se root-convergence factor of rAA(1) is  $\rho_{rAA(1)}^* = \frac{m^*}{(1 + (m^*)^2)^{1/4}},$ which requires  $m^* < \sqrt{\frac{1}{2}(1+\sqrt{5})} \approx 1.272$  to guarantee that rAA(1) converges for arbitrary initial guess. Notice that no matter what  $m^*$  is, we always have  $\varrho_{GMRES(1),ss}^* < \varrho_{rAA(1)}^*$ .

From the above discussion, we can conclude that the asymptotic convergence factor of  $GMRES(1)$  for M being skew-symmetric depends solely on the dominant eigenvalue (largest magnitude) with which the initial residual  $r_0$  is associated.

<span id="page-15-4"></span> $\Box$ 

COROLLARY 3.4. Suppose M is skew-symmetric, and the initial residual  $r_0 =$  $\nabla j^*$  $j_{j=1}^{\infty} u_j$ , where  $u_j \in \text{span}\{Q_j\}$ ,  $1 \leqslant j^* \leqslant n/2$ , and  $|m_{j^*}|$  is the largest magnitude of the eigenvalues. Then,

$$
\varrho(r_0) = \frac{|m_j^*|}{\sqrt{1+|m_j^*|^2}}.
$$

*Remark* 3.5. In [\[28\]](#page-20-25), it has shown that when A is a real positive definite matrix, then

$$
\|r_k\| \leqslant \sqrt{1-\lambda_{\min}\left(\frac{A^{-1}+A^{-T}}{2}\right)\lambda_{\min}\left(\frac{A+A^{T}}{2}\right)}\|r_{k-1}\|.
$$

In fact, when M is skew-symmetric,  $\frac{A+A^{T}}{2} = \frac{I-M+I-M^{T}}{2} = I$ , and  $\frac{A^{-1}+A^{-T}}{2} =$ In fact, when M is skew-symmetric,<br> $(I + MM^T)^{-1}$ . Then,  $\lambda_{\min} \left( \frac{A^{-1} + A^{-T}}{2} \right)$ 2 ¯  $= (1 + (m^*)^2))^{-1}$ . Thus,

$$
\sqrt{1-\lambda_{\min}\left(\frac{A^{-1}+A^{-T}}{2}\right)\lambda_{\min}\left(\frac{A+A^{T}}{2}\right)} = \sqrt{1-\frac{1}{1+(m^*)^2}},
$$

which is the same result as we obtained in Corollary [3.3.](#page-15-1)

3.1. q-linear convergence. Based on the previous discussion, we are able to determine the q-linear convergence of GMRES(1). The following results have been obtained.

THEOREM 3.6. For GMRES(1) applied to  $Ax = b$ , where A is symmetric, the residuals converge q-linearly with  $q$ -factor  $\sigma$ :

$$
(3.4) \t\t\t  $\|r_k\| \leq \sigma \|r_{k-1}\|,$
$$

where  $\sigma = \rho_{GMRES(1)}^*$  is given by [\(2.22\)](#page-9-0) if A is symmetric and  $\sigma = \rho_{GMRES(1),ss}^*$  is qiven by  $(3.3)$  if A is skew-symmetric.

Proof. These results can be derived directly from Lemma [2.1](#page-2-3) and [\(3.2\)](#page-15-3).  $\Box$ 

We point out that the q-linear convergence with q-factor  $\sigma$  actually is the worst-case rate. For different initial guesses/errors,  $\sigma$  might be different, but it must be less than  $\rho_{GMRES(1)}^*$  ( $\rho_{GMRES(1),ss}^*$ ). However, we do not further discuss this. In [\[37\]](#page-20-12), it has  $\rho_{GMRES(1)}$  ( $\rho_{GMRES(1),ss}$ ). now<br>been shown the sequence  $\left\{\frac{\|r_k\|}{\|r\|_{M}}\right\}$  $\frac{\|r_k\|}{\|r_{k-1}\|}$  is increasing. We know that  $\frac{\|r_k\|}{\|r_{k-1}\|}$  has an upper bound  $\sigma$ , so the sequence must be convergent.

<span id="page-16-0"></span>4. Numerical experiments. In this section, we present some numerical experiments to validate our theoretical results of worst-case root-convergence factor of  $GMRES(1)$ . For symmetric  $A$ , we consider three choices, which correspond to three cases of M: M is not invertible,  $\rho(M) < 1$ , and  $\rho(M) > 1$ .

EXAMPLE 4.1 (A is symmetric). We consider solving  $Ax = b = 0$ , where A is  $A_i$ .

$$
A_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix}, A_2 = \begin{pmatrix} 1/2 & 0 & 0 & 0 & 0 \\ 0 & 1/4 & 0 & 0 & 0 \\ 0 & 0 & 1/8 & 0 & 0 \\ 0 & 0 & 0 & 1/16 & 0 \\ 0 & 0 & 0 & 0 & 1/32 \end{pmatrix}, A_3 = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 4 \end{pmatrix}.
$$

Using [\(2.13\)](#page-5-2), we obtain the worst-case root-convergence factor for  $A_1, A_2, A_3$ :  $\varrho_1^* =$  $\frac{1}{2} = 0.5, \varrho_2^* = \frac{15}{17} \approx 0.8824, \varrho_3^* = 1$  for  $A_1, A_2$  and  $A_3$ , respectively. For our tests, we run 1000 random initial guesses, whose elements are uniform distribution in the interval  $(-1, 1)$ , and we require  $||r_k|| \le 1e - 30$  to make k large enough to capture the behavior of  $\varrho_k(r_0) = ||r_k||^{1/k}$  $\varrho_k(r_0) = ||r_k||^{1/k}$  $\varrho_k(r_0) = ||r_k||^{1/k}$ . Figure 1 reports the root-convergence factor as a function of the iteration index k for different initial guesses for  $A_1$  and  $A_2$ . The dashed line is our theoretical worst-case root-convergence factor, denoted as  $\varrho^*$ . We see that the numerical root-convergence factor is bounded by the worst-case root-convergence factor, and the root-convergence factor depends on initial guesses.

<span id="page-17-0"></span>

FIG. 1. Root-convergence factor  $\varrho_k(r_0) = ||r_k||^{1/k}$  (solid lines) as a function of the iteration index k with 1000 random initial guesses for  $A_1$  (left) and  $A_2$  (right). The dashed line is the theoretical worst-case root convergence factor, denoted as  $\varrho^*$ .

For  $A_3$ , the indefinite case, the root-convergence factor is displayed on the left side of Figure [2.](#page-17-1) To verify our result in Lemma [2.3,](#page-3-3) which states the convergence factor depends on the initial guess, we also consider  $A_3$  and choose initial guesses where the first component is zero. Then, the worst-case root-convergence factor depends only on the eigenvalues 2, 3, 4. Using Theorem [2.6,](#page-5-4) we have  $\rho^* = \frac{4-2}{4+2} = \frac{1}{3} \approx 0.3333$ . The result on the right side of Figure [2](#page-17-1) validates this.

<span id="page-17-1"></span>

FIG. 2. Root-convergence factor  $\varrho_k(r_0) = ||r_k||^{1/k}$  (solid lines) as a function of the iteration index k with 1000 random initial guesses for  $A_3$  (left) and for  $A_3$  (right) with the first elements of the initial guesses being zero. The dashed line is the theoretical worst-case root-convergence factor, denoted as  $\varrho^*$ .

To investigate how the initial guess affects the root-convergence factor, see The-orem [2.7,](#page-6-1) we consider two different initial guesses for  $A_2$  (denoted its eigenvalues orem 2.1, we consider two different initial guesses for  $A_2$  (denoted its eigenvalues<br>as  $a_1 = 1/2, a_2 = 1/4, a_3 = 1/16$ ). One initial guess is  $x_0 = [1, 2\sqrt{2}, 0, 0, 0]^T$ corresponding to  $r_0 = [1/2, \sqrt{2}/2, 0, 0, 0]^T$ . Then,  $\epsilon = \frac{c_2}{c_1}$  $\sqrt{2} = \sqrt{\frac{a_1}{a_2}}$ . We have  $\varrho(r_0) = \frac{a_1 - a_2}{a_1 + a_2} = \frac{1}{3}$ . The another is  $x_0 = [1, 0, 8, 0, 0]^T$  corresponding to

<span id="page-18-1"></span>

FIG. 3. Root-convergence factor  $\varrho_k(r_0) = ||r_k||^{1/k}$  (solid lines) as a function of the iteration index k with two different initial guesses for  $A_2$ . The dashed line is the theoretical worst-case rootconvergence factor. Cross markers corresponds to  $x_0 = [1, 2\sqrt{2}, 0, 0, 0]^T$ , where  $\varrho^* = 0.3333$ . Circle convergence factor. markers are for  $x_0 = [1, 0, 8, 0, 0]^T$ , where  $\varrho^* = 0.6000$ .

 $r_0 = [1/2, 0, 1, 0, 0]^T$ . Then,  $\epsilon = \frac{c_3}{c_1} = 2$  $\frac{a_1}{a_3}$  and  $\rho^* = \frac{a_1 - a_3}{a_1 + a_3} = 0.6000$ . The root-convergence factors are reported in Figure [3.](#page-18-1) The dashed line is the theoretical worst-case root-convergence factor. We observe that the numerical root-convergence factors align with our theoretical results.

In the following example, we consider the case that  $M$  is symmetric.

EXAMPLE 4.2 (M is skew-symmetric). Consider  $A_4 = I - M$ , where

$$
M = \frac{1}{8} \begin{pmatrix} 0 & 1 & 0 & -5 & 0 & 0 & 0 & 2 \\ -1 & 0 & 0 & 0 & 5 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 & -2 & -1 & 5 & 0 \\ 5 & 0 & 0 & 0 & -1 & -2 & 0 & 0 \\ 0 & -5 & 2 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 2 & 0 & 0 & 0 & -5 \\ 0 & 2 & -5 & 0 & 0 & 0 & 0 & 1 \\ -2 & 0 & 0 & 0 & 0 & 5 & -1 & 0 \end{pmatrix}
$$

,

which is taken from [\[39\]](#page-20-26) with some modification. The eigenvalues of M are  $\pm \iota_{\frac{1}{4}}^1$ ,  $\iota \pm \frac{2}{4}$ ,  $\pm \iota_{\frac{3}{4}}^2$ ,  $\pm 1$ . Thus,  $m^* = 1$  and  $\varrho^* = \frac{1}{\sqrt{1+1^2}} \approx 0.7071$ .  $\frac{1}{1+1^2} \approx 0.7071.$ 

On the left side of Figure [4,](#page-19-14) we run 1000 random initial guesses, whose elements are uniform distribution in the interval  $(-1, 1)$ , and report the root-convergence factor  $\varrho_k(r_0) = ||r_k||^{1/k}$  as a function of the iteration index k. The dashed line is our theoretical worst-case root-convergence factor, denoted as  $\varrho^*$ . We see that when  $k \to \infty$ , the root-convergence factor converges to the worst-case root-convergence factor  $\varrho^* = 0.7071$ . To validate Corollary [3.4,](#page-15-4) we consider  $x_0 = Q(:, 3) + Q(:, 5) + Q(:, 7)$ , see [\(3.1\)](#page-15-5), which is a vector in the space spanned by the columns of  $Q_2, Q_3, Q_4$  corresponding to eigenvalues  $\pm \iota \frac{1}{4}$ ,  $\iota \pm \frac{2}{4}$ ,  $\pm \iota \frac{3}{4}$  of M. Thus, the expected root-convergence factor is  $\frac{3/4}{\sqrt{1-(2)}}$  $\frac{3}{4(1+(3/4)^2)} = \frac{3}{5} = 0.6$  $\frac{3}{4(1+(3/4)^2)} = \frac{3}{5} = 0.6$  $\frac{3}{4(1+(3/4)^2)} = \frac{3}{5} = 0.6$ . Our numerical results on the right side of Figure 4 confirm this.

<span id="page-18-0"></span>5. Conclusion. We study the root-convergence factor of GMRES(1) in solving linear systems. We derive theoretical results on the worst-case root-convergence factor based on eigenvector-dependent nonlinear eigenvalue problems. When the coefficient matrix A of the underlying linear system is symmetric, the root-convergence factor highly depends on the initial guess. For some special initial guesses, we are able

<span id="page-19-14"></span>

FIG. 4. Root-convergence factor  $\varrho_k(r_0) = ||r_k||^{1/k}$  (solid lines) as a function of the iteration index k with 1000 random initial  $x_0$  for  $A_4$  (left). The dashed line is the theoretical worst-case root convergence factor, denoted as  $\varrho^*$ . Cross markers are  $\varrho_k(r_0) = ||r_k||^{1/k}$  as a function of the iteration index k with  $x_0 = Q(:, 3) + Q(:, 5) + Q(:, 7)$  for  $A_4$  (right).

to derive their root-convergence factors. When the matrix  $M = I - A$  is skewsymmetric, our theoretical results reveal that the worst-case root-convergence factor depends solely on the spectral radius of  $M$ , and  $\text{GMRES}(1)$  converges unconditionally. The q-linear convergence factor is derived too, which is the same as the worst-case root-convergence factor. Additionally, we present new results for restarted Anderson acceleration with window size one.

Acknowledgments. The author thanks Microsoft Copilot for spelling and grammar check.

### REFERENCES

- <span id="page-19-4"></span>[1] E. Agullo, L. Giraud, and Y.-F. Jing, Block GMRES method with inexact breakdowns and deflated restarting, SIAM Journal on Matrix Analysis and Applications, 35 (2014), pp. 1625–1651.
- <span id="page-19-12"></span>[2] D. G. ANDERSON, *Iterative procedures for nonlinear integral equations*, Journal of the ACM (JACM), 12 (1965), pp. 547–560.
- <span id="page-19-1"></span>[3] M. ARIOLI, An analysis of GMRES worst case convergence, Unpublished report, (2009), p. 240.
- <span id="page-19-5"></span>[4] A. H. BAKER, E. R. JESSUP, AND T. MANTEUFFEL, A technique for accelerating the convergence of restarted GMRES, SIAM Journal on Matrix Analysis and Applications, 26 (2005), pp. 962–984.
- <span id="page-19-10"></span>[5] J. W. BOTH, K. KUMAR, J. M. NORDBOTTEN, AND F. A. RADU, Anderson accelerated fixedstress splitting schemes for consolidation of unsaturated porous media, Computers & Mathematics with Applications, 77 (2019), pp. 1479–1502.
- <span id="page-19-11"></span>[6] H. DE STERCK, O. A. KRZYSIK, AND A. SMITH, Asymptotic convergence of restarted anderson acceleration for certain normal linear systems, arXiv preprint arXiv:2312.04776v3, (2024).
- <span id="page-19-9"></span><span id="page-19-0"></span>M. EMBREE, The tortoise and the hare restart GMRES, SIAM review, 45 (2003), pp. 259–266. [8] - Row descriptive are GMRES convergence bounds?, arXiv preprint arXiv:2209.01231, (2022).
- <span id="page-19-7"></span>[9] A. Essai, Weighted FOM and GMRES for solving nonsymmetric linear systems, Numerical Algorithms, 18 (1998), pp. 277–292.
- <span id="page-19-13"></span>[10] C. EVANS, S. POLLOCK, L. G. REBHOLZ, AND M. XIAO, A proof that Anderson acceleration improves the convergence rate in linearly converging fixed-point methods (but not in those converging quadratically), SIAM Journal on Numerical Analysis, 58 (2020), pp. 788–810.
- <span id="page-19-8"></span>[11] V. Faber, J. Liesen, and P. Tichy, Properties of worst-case GMRES, SIAM Journal on Matrix Analysis and Applications, 34 (2013), pp. 1500–1519.
- <span id="page-19-6"></span>[12] A. FROMMER AND U. GLÄSSNER, Restarted GMRES for shifted linear systems, SIAM Journal on Scientific Computing, 19 (1998), pp. 15–26.
- <span id="page-19-3"></span>[13] L. GIRAUD, S. GRATTON, X. PINEL, AND X. VASSEUR, Flexible GMRES with deflated restarting, SIAM Journal on Scientific Computing, 32 (2010), pp. 1858–1878.
- <span id="page-19-2"></span>[14] A. GREENBAUM, V. PTÁK, AND Z. E. K. STRAKOŠ, Any nonincreasing convergence curve is pos-

sible for GMRES, SIAM Journal on Matrix Analysis and Applications, 17 (1996), pp. 465– 469.

- <span id="page-20-1"></span>[15] A. GREENBAUM AND L. N. TREFETHEN, GMRES/CR and Arnoldi/Lanczos as matrix approximation problems, SIAM Journal on Scientific Computing, 15 (1994), pp. 359–368.
- <span id="page-20-16"></span>[16] S. GÜTTEL AND J. PESTANA, Some observations on weighted GMRES, Numerical Algorithms, 67 (2014), pp. 733–752.
- <span id="page-20-17"></span>[17] W. JOUBERT, On the convergence behavior of the restarted GMRES algorithm for solving nonsymmetric linear systems, Numerical Linear Algebra with Applications, 1 (1994), pp. 427– 447.
- <span id="page-20-6"></span>[18] J. Liesen, Computable convergence bounds for GMRES, SIAM Journal on Matrix Analysis and Applications, 21 (2000), pp. 882–903.
- <span id="page-20-5"></span>[19] J. Liesen and Z. Strakos, Convergence of GMRES for tridiagonal Toeplitz matrices, SIAM Journal on Matrix Analysis and Applications, 26 (2004), pp. 233–251.
- <span id="page-20-8"></span>[20] J. LIESEN AND P. TICHY, The worst-case GMRES for normal matrices, BIT Numerical mathematics, 44 (2004), pp. 79–98.
- <span id="page-20-10"></span>[21] Y. Lin, L. Bao, and Q. Wu, Simpler GMRES with deflated restarting, Mathematics and Computers in Simulation, 82 (2012), pp. 2238–2252.
- <span id="page-20-11"></span>[22] R. B. Morgan, A restarted GMRES method augmented with eigenvectors, SIAM Journal on Matrix Analysis and Applications, 16 (1995), pp. 1154–1171.
- <span id="page-20-9"></span>[23]  $\_\_\_\_\_$  GMRES with deflated restarting, SIAM Journal on Scientific Computing, 24 (2002), pp. 20–37.
- <span id="page-20-14"></span>[24] N. M. NACHTIGAL, L. REICHEL, AND L. N. TREFETHEN, A hybrid GMRES algorithm for nonsymmetric linear systems, SIAM Journal on Matrix Analysis and Applications, 13 (1992), pp. 796–825.
- <span id="page-20-22"></span>[25] J. M. ORTEGA AND W. C. RHEINBOLDT, *Iterative solution of nonlinear equations in several* variables, SIAM, 2000.
- <span id="page-20-4"></span>[26] M. ROBBÉ AND M. SADKANE, A convergence analysis of GMRES and FOM methods for Sylvester equations, Numerical Algorithms, 30 (2002), pp. 71–89.
- <span id="page-20-20"></span>[27] Y. SAAD, Further analysis of minimum residual iterations, Numerical Linear Algebra with Applications, 7 (2000), pp. 67–93.
- <span id="page-20-25"></span>[28]  $\longrightarrow$ , Iterative methods for sparse linear systems, SIAM, 2003.
- <span id="page-20-0"></span>[29] Y. SAAD AND M. H. SCHULTZ, GMRES: A generalized minimal residual algorithm for solving nonsymmetric linear systems, SIAM Journal on Scientific and Statistical Computing, 7 (1986), pp. 856–869.
- <span id="page-20-15"></span>[30] V. Simoncini and E. Gallopoulos, A hybrid block GMRES method for nonsymmetric systems with multiple right-hand sides, Journal of Computational and Applied Mathematics, 66 (1996), pp. 457–469.
- <span id="page-20-24"></span>[31] H. D. STERCK AND Y. HE, On the asymptotic linear convergence speed of Anderson acceleration, Nesterov acceleration, and nonlinear GMRES, SIAM Journal on Scientific Computing, 43 (2021), pp. S21–S46.
- <span id="page-20-7"></span>[32] D. TITLEY-PELOQUIN, J. PESTANA, AND A. J. WATHEN, GMRES convergence bounds that depend on the right-hand-side vector, IMA Journal of Numerical Analysis, 34 (2014), pp. 462– 479.
- <span id="page-20-2"></span>[33] K.-C. Toh, GMRES vs. ideal GMRES, SIAM Journal on Matrix Analysis and Applications, 18 (1997), pp. 30–36.
- <span id="page-20-21"></span>[34] A. TOTH AND C. T. KELLEY, Convergence analysis for Anderson acceleration, SIAM Journal on Numerical Analysis, 53 (2015), pp. 805–819.
- <span id="page-20-3"></span>[35] H. A. VAN DER VORST AND C. VUIK, The superlinear convergence behaviour of GMRES, Journal of Computational and Applied Mathematics, 48 (1993), pp. 327–341.
- <span id="page-20-13"></span>[36] H. A. VAN DER VORST AND C. VUIK, GMRESR: a family of nested GMRES methods, Numerical Linear Algebra with Applications, 1 (1994), pp. 369–386.
- <span id="page-20-12"></span>[37] E. Vecharynski and J. Langou, The cycle-convergence of restarted GMRES for normal matrices is sublinear, SIAM Journal on Scientific Computing, 32 (2010), pp. 186–196.
- <span id="page-20-23"></span>[38] H. F. Walker and P. Ni, Anderson acceleration for fixed-point iterations, SIAM Journal on Numerical Analysis, 49 (2011), pp. 1715–1735.
- <span id="page-20-26"></span>[39] R. WARD AND L. GRAY, *Eigensystem computation for skew-symmetric matrices and a class of* symmetric matrices.[subroutines trizd, imzd, and tbakzd in fortran for ibm 360/91 computer], tech. rep., Oak Ridge National Lab.(ORNL), Oak Ridge, TN (United States), 1976.
- <span id="page-20-19"></span>[40] B. Zhong and R. B. Morgan, Complementary cycles of restarted GMRES, Numerical Linear Algebra with Applications, 15 (2008), pp. 559–571.
- <span id="page-20-18"></span>[41] J. ZÍTKO, Generalization of convergence conditions for a restarted GMRES, Numerical Linear Algebra with Applications, 7 (2000), pp. 117–131.