DADA: Dual Averaging with Distance Adaptation

Mohammad Moshtaghifar[∗] Anton Rodomanov† Daniil Vankov‡

Sebastian U. Stich§

January 17, 2025

Abstract

We present a novel universal gradient method for solving convex optimization problems. Our algorithm—Dual Averaging with Distance Adaptation (DADA)—is based on the classical scheme of dual averaging and dynamically adjusts its coefficients based on observed gradients and the distance between iterates and the starting point, eliminating the need for problem-specific parameters. DADA is a universal algorithm that simultaneously works for a broad spectrum of problem classes, provided the local growth of the objective function around its minimizer can be bounded. Particular examples of such problem classes are nonsmooth Lipschitz functions, Lipschitz-smooth functions, Hölder-smooth functions, functions with high-order Lipschitz derivative, quasi-self-concordant functions, and (L_0, L_1) -smooth functions. Crucially, DADA is applicable to both unconstrained and constrained problems, even when the domain is unbounded, without requiring prior knowledge of the number of iterations or desired accuracy.

Keywords: Convex Optimization, Gradient Methods, Adaptive Algorithms, Dual Averaging, Distance Adaption, Universal Methods, Worst-Case Complexity Guarantees

1 Introduction

Gradient methods are among the most popular and efficient algorithms for solving optimization problems arising in machine learning, as they are highly adaptable and scalable across various settings [\[2\]](#page-11-0). Despite their popularity, these methods face a significant challenge of selecting appropriate hyperparameters, particularly stepsizes, which are critical to the performance of the algorithm. Hyperparameter tuning is one of the standard approaches to address this issue but is a time-consuming and resource-intensive process, especially as models become larger and more complex. Consequently, the cost of training these models has become a significant concern [\[19,](#page-12-0) [22\]](#page-12-1).

Typically, line-search techniques have been used to select stepsizes for optimization methods, and they are provably efficient for certain function classes, such as Höldersmooth problems [\[15\]](#page-12-2). However, in recent years, several so-called parameter-free algorithms have been developed which do not utilize line search [\[3,](#page-11-1) [4,](#page-11-2) [9,](#page-11-3) [10,](#page-11-4) [12,](#page-11-5) [18\]](#page-12-3). Notably,

[∗]Sharif University of Technology. E-mail: m.moshtaghi@sharif.edu.

[†]CISPA Helmholtz Center for Information Security. E-mail: anton.rodomanov@cispa.de.

[‡]Arizona State University. E-mail: dvankov@asu.edu.

[§]CISPA Helmholtz Center for Information Security. E-mail: stich@cispa.de.

one strategy involves dynamically adjusting stepsizes based on estimates of the initial distance to the optimal solution [\[3,](#page-11-1) [9,](#page-11-3) [10\]](#page-11-4). Another approach leverages lower bounds on the initial distance combined with the Dual Averaging (DA) scheme [\[5,](#page-11-6) [12\]](#page-11-5). However, these methods primarily focus on nonsmooth Lipschitz or, in some cases, Lipschitz-smooth functions. Some of these methods also come with additional limitations, such as requiring bounded domain assumptions [\[10\]](#page-11-4) or failing to extend to constrained optimization problems [\[5,](#page-11-6) [12\]](#page-11-5).

To formalize the discussion, we consider the following optimization problem:

$$
f^* := \min_{x \in Q} f(x),\tag{1.1}
$$

where $Q \subseteq \mathbb{R}^d$ is a nonempty closed convex set, and $f: \mathbb{R}^d \to \mathbb{R}$ is a convex function over Q . We assume that Q is a simple set, meaning that it is possible to efficiently compute the projection onto Q . We also assume problem (1.1) has a solution which we denote by x^* . The starting point in our methods is denoted by x_0 .

Contributions. In this paper, we introduce Dual Averaging with Distance Adaptation (DADA), a novel universal gradient method for solving [\(1.1\)](#page-1-0). Building on the classical framework of weighted DA [\[14\]](#page-12-4), DADA incorporates a dynamically adjusted estimate of $D_0 := ||x_0 - x^*||$, inspired by recent techniques from [\[3,](#page-11-1) [9\]](#page-11-3) and further developed in [\[10\]](#page-11-4), without requiring prior knowledge of problem-specific parameters. Furthermore, our approach applies to both unconstrained problems and those with simple constraints, possibly with unbounded domains. This makes DADA a powerful tool across a wide range of applications.

We start, in Section [2,](#page-2-0) by presenting our method and outline its foundational structure based on the DA scheme [\[14\]](#page-12-4). Our main theoretical result, Theorem [2.1,](#page-3-0) establishes convergence guarantees for a broad range of function classes.

To demonstrate the versatility and effectiveness of DADA, in Section [3,](#page-5-0) we provide complexity estimates across several interesting function classes: nonsmooth Lipschitz functions, Lipschitz-smooth functions, Hölder-smooth functions, quasi-self-concordant (QSC) functions, functions with Lipschitz high-order derivative, and (L_0, L_1) -smooth functions. These results underscore DADA's ability to deliver competitive performance without knowledge of class-specific parameters.

Notation. In this text, we work in the space \mathbb{R}^d equipped with the standard inner product $\langle \cdot, \cdot \rangle$ and the general Euclidean norm:

$$
||x|| \coloneqq \langle Bx, x \rangle^{1/2}, \qquad x \in \mathbb{R}^d,
$$

where B is a fixed symmetric positive definite matrix. The corresponding dual norm is defined in the standard way:

$$
||s||_* := \max_{||x||=1} \langle s, x \rangle = \langle s, B^{-1}s \rangle^{1/2}, \quad s \in \mathbb{R}^d.
$$

Thus, for any $s, x \in \mathbb{R}^d$, we have the Cauchy-Schwarz inequality $|\langle s, x \rangle| \leq ||s||_* ||x||$. For a convex function $f: \mathbb{R}^d \to \mathbb{R}$, we denote its subdifferential at a point $x \in \mathbb{R}^d$ by $\partial f(x)$. We also use $\nabla f(x) \in \partial f(x)$ to denote a subgradient of f at the point x.

Algorithm 2.1 General Scheme of DA

Input: $x_0 \in Q$, number of iterations $T \ge 1$, coefficients $(a_k)_{k=0}^{T-1}$, $(\beta_k)_{k=1}^T$ with nondecreasing β_k for $k = 1, \ldots, T$ do Compute arbitrary $g_k \in \partial f(x_k)$ $x_k = \operatorname{argmin}_{x \in Q} \left\{ \sum_{i=0}^{k-1} a_i \langle g_i, x - x_i \rangle + \frac{\beta_k}{2} ||x - x_0||^2 \right\}$ Output: $x_T^* = \operatorname{argmin}_{x \in \{x_0, ..., x_{T-1}\}} f(x)$

2 DADA Method

Measuring the quality of solution. Given an approximate solution $x \in Q$ to prob-lem [\(1.1\)](#page-1-0) and an arbitrary subgradient $\nabla f(x) \in \partial f(x)$, we measure the suboptimality of x by the distance from x^* to the hyperplane $\{y : \langle \nabla f(x), x - y \rangle = 0\}$:

$$
v(x) \coloneqq \frac{\langle \nabla f(x), x - x^* \rangle}{\|\nabla f(x)\|_{*}} \quad (\ge 0). \tag{2.1}
$$

This objective is meaningful because minimizing $v(x)$ also reduces the corresponding function residual $f(x) - f^*$. Indeed, there exists the following simple relationship between $v(x)$ and the function residual [\[16,](#page-12-5) Section 3.2.2] (see also [A](#page-13-0)ppendix A for the short proof):

$$
f(x) - f^* \le \omega(v(x)),\tag{2.2}
$$

where

$$
\omega(t) := \max_{x} \{ f(x) - f^* : ||x - x^*|| \le t \}
$$
\n(2.3)

measures the local growth of f around the solution x^* . By bounding $\omega(t)$, we can derive convergence-rate estimates that simultaneously apply to a broad range of problem classes (we discuss several examples in Section [3\)](#page-5-0).

The method. Our algorithm is based on the general scheme of DA [\[14\]](#page-12-4) shown in Algorithm [2.1.](#page-2-1) Using a standard (sub)gradient method with time-varying coefficients is also possible but requires either short steps by fixing the number of iterations in advance, or paying an extra logarithmic factor in the convergence rate [\[16,](#page-12-5) Section 3.2.3].

The classical method of Weighted DA (WDA) selects the coefficients $a_k = \frac{\hat{D}_0}{\|a_k\|}$ $\frac{D_0}{\|g_k\|_*}$ and $\beta_k = \Theta(\sqrt{k})$, where \hat{D}_0 is a user-defined estimate of D_0 . The convergence is guaranteed for any value of \hat{D}_0 but one must pay a multiplicative cost of ρ^2 , where $\rho \coloneqq \max\{\frac{\hat{D}_0}{D_0}\}$ $\frac{\bar{D}_0}{\bar{D}_0}, \frac{\bar{D}_0}{\hat{D}_0}\},$ if the parameter D_0 is unknown. This cost can be significantly high if D_0 is not known almost exactly. To address this issue, we propose DADA, which reduces the cost to a logarithmic term, $\log^2 \rho$, offering a substantial improvement.

Specifically, our approach utilizes the following coefficients:

$$
a_k = \frac{\bar{r}_k}{\|g_k\|_*}, \quad \beta_k = c\sqrt{k+1}, \quad \bar{r}_k := \max\{\max_{1 \le t \le k} r_t, \bar{r}\}, \quad r_t := \|x_0 - x_t\|, \tag{2.4}
$$

where $\bar{r} > 0$ is a parameter and c is a certain constant to be specified later. In what follows, we assume w.l.o.g. that $g_k \neq 0$ for all $0 \leq k \leq T-1$ since otherwise the exact solution has been found, and the method could be successfully terminated before making T iterations.

As we can see, the main difference between WDA and DADA, is that the latter dynamically adjusts its estimate of D_0 by exploiting r_t , the distance between x_t and the initial point x_0 . This idea has been explored in recent works [\[3,](#page-11-1) [9\]](#page-11-3), which similarly utilize r_t in various ways. Other methods also attempt to estimate this quantity using alternative strategies, based on DA and the similar principle of employing an increasing sequence of lower bounds for D_0 [\[5,](#page-11-6) [12\]](#page-11-5).

The convergence guarantees for our method are provided in the result below:

Theorem [2.1](#page-2-1). Consider Algorithm 2.1 for solving problem (1.1) using the coefficients **Theorem 2.1.** Consular Algorium 2.1 for solving problem (1.1) using the coefficient.
from [\(2.4\)](#page-2-2) with $c > \sqrt{2}$. Then, for any $T \ge 1$ and $v_T^* \coloneqq \min_{0 \le k \le T-1} v(x_k)$, it holds that

$$
f(x_T^*) - f^* \le \omega(v_T^*),
$$

and

$$
v_T^* \le \frac{eD}{\sqrt{T}} \log \frac{e\bar{D}}{\bar{r}},\tag{2.5}
$$

where $\bar{D} \coloneqq \max\{\bar{r}, \frac{2c}{\bar{r}}\}$ $\frac{2c}{c-\sqrt{2}}D_0$ } and D := √ $\overline{2}(cD_0 + \frac{1}{c}\overline{D})$. Consequently, for a given $\delta > 0$, it holds that $v^*_T \leq \delta$ whenever $T \geq T_v(\delta)$, where

$$
T_v(\delta) \coloneqq \frac{e^2 D^2}{\delta^2} \log^2 \frac{e \bar{D}}{\bar{r}}
$$

Let us provide a proof sketch for Theorem [2.1](#page-3-0) here and defer the detailed proof to Appendix [B.](#page-14-0) We begin by applying the standard result for DA (Lemma [B.1\)](#page-14-1), which holds for any choice of coefficients a_k and β_k :

$$
\sum_{i=0}^{k-1} a_i v_i \|g_i\|_{*} + \frac{\beta_k}{2} D_k^2 \le \frac{\beta_k}{2} D_0^2 + \sum_{i=0}^{k-1} \frac{a_i^2}{2\beta_i} \|g_i\|_{*}^2,
$$

where $D_i = ||x_i - x^*||$ and $v_i = v(x_i)$ for all $i \geq 0$. Use the specific choices for a_k and β_k as defined in [\(2.4\)](#page-2-2), we obtain (see Lemma [B.2\)](#page-15-0):

$$
\sum_{i=0}^{k-1} \bar{r}_i v_i + \frac{c\sqrt{k+1}}{2} D_k^2 \le \frac{c\sqrt{k+1}}{2} D_0^2 + \frac{\sqrt{k}}{c} \bar{r}_{k-1}^2.
$$
 (2.6)

.

Dropping the nonnegative $\bar{r}_i v_i$ from the left-hand side, we can show by induction that \bar{r}_k is uniformly bounded (see Lemma [B.3\)](#page-15-1):

$$
\bar{r}_k \leq \bar{D},
$$

where \bar{D} is the constant from Theorem [2.1.](#page-3-0) This bound is crucial to our analysis, as we need to eliminate \bar{r}_{k-1} from the right-hand side of [\(2.6\)](#page-3-1). Achieving this requires selecting the coefficients precisely as defined in [\(2.4\)](#page-2-2), which is the primary difference compared to the standard DA method [\[14\]](#page-12-4). Next, using the inequality $D_0^2 - D_k^2 \leq 2r_k D_0$, we get

$$
\sum_{i=0}^{k-1} \bar{r}_i v_i \le c\sqrt{k+1} r_k D_0 + \frac{\sqrt{k}}{c} \bar{r}_{k-1}^2 \le \left(cD_0 + \frac{1}{c}\bar{D}\right) \bar{r}_k \sqrt{k+1}.
$$

After establishing this, the rest of the proof follows straightforwardly by dividing both sides by $\sum_{i=0}^{k-1} \bar{r}_i$ and applying the following inequality (valid for any nondecreasing sequence \bar{r}_k , see Lemma [A.1\)](#page-13-1):

$$
\min_{1\leq k\leq T}\frac{\bar{r}_k}{\sum_{i=0}^{k-1}\bar{r}_i}\leq \frac{(\frac{\bar{r}_T}{\bar{r}_0})^{\frac{1}{T}}\log\frac{e\bar{r}_T}{\bar{r}_0}}{T}.
$$

This gives us

$$
v_T^* \leq \frac{D}{\sqrt{T}} \bigg(\frac{\bar{D}}{\bar{r}} \bigg)^{\frac{1}{T}} \log \frac{e\bar{D}}{\bar{r}},
$$

which is almost [\(2.5\)](#page-3-2) except for the extra factor of $(\frac{\bar{D}}{\bar{r}})^{\frac{1}{T}}$. This extra factor, however, is rather weak as it can be upper bounded by a constant (say, $e \equiv \exp(1)$) whenever $T \geq \log \frac{\bar{D}}{\bar{r}}$. The case of $T \leq \log \frac{\bar{D}}{\bar{r}}$ is not interesting since then [\(2.5\)](#page-3-2) holds trivially because, for any $k \geq 0$, in view of [\(2.1\)](#page-2-3) and Lemma [B.3,](#page-15-1) we have $v_k \leq D_k \leq D$.

How to choose the constant c. According to Theorem [2.1,](#page-3-0) our method converges for **How to choose the constant** c. According to Theorem 2.1, our method converges for any $c > \sqrt{2}$. However, the choice of c can influence the constant factor in the complexity of DADA. Hence, our goal here is to find the optimal constant c that minimizes $T_v(\delta)$. To determine this c, let \bar{r} be sufficiently small, so that

$$
\bar{D} \equiv \max\left\{\bar{r}, \frac{2c}{c - \sqrt{2}}D_0\right\} = \frac{2c}{c - \sqrt{2}}D_0.
$$

Then, disregarding the logarithmic factors, due to their minimal impact on the complexity of our method, we can determine the optimal constant c that minimizes

$$
D \equiv \sqrt{2} \Big(cD_0 + \frac{1}{c} \bar{D} \Big) = \sqrt{2} \Big(c + \frac{2}{c - \sqrt{2}} \Big) D_0.
$$

This is the value

$$
c = 2\sqrt{2}.\tag{2.7}
$$

For this optimal choice of c, we get $\overline{D} = \max{\lbrace \overline{r}, 4D_0 \rbrace}$ and $D = 4D_0 + \frac{1}{2}\overline{D}$, so the complexity of our method given by Theorem [2.1](#page-3-0) is

$$
T_v(\delta) = \frac{e^2 (4D_0 + \frac{1}{2}\bar{D})^2}{\delta^2} \log^2 \frac{e\bar{D}}{\bar{r}}.
$$

Comparison with recent distance-adaptation methods. Let us briefly compare our method with several recently proposed parameter-free algorithms, namely, DoG [\[9\]](#page-11-3), DoWG [\[10\]](#page-11-4), D-Adaptation [\[5\]](#page-11-6) and Prodigy [\[12\]](#page-11-5).

To begin, we clarify the key differences between our method and approaches like DoG. One immediate difference is that we use DA instead of the classical (sub)gradient method employed by DoG. However, the most significant difference lies in how the sequence of gradients is handled. DoG normalizes the current gradient g_k by the accumulated norms of the previous gradients, an idea inspired by AdaGrad [\[8\]](#page-11-7). In contrast, our method simply normalizes g_k by its own norm. This modification makes our method universal,

ensuring the convergence of v_T^* to zero, which is not known to be the case for DoG, even for deterministic problems.

Both DoG and DoWG employ a similar approach to estimate D_0 and achieve comparable convergence rates for Lipschitz-smooth and nonsmooth functions. Similarly to our approach, DoWG considers only the deterministic case, but with an additional assumption of the bounded feasible set. They have a different definition of universality, considering only smooth and nonsmooth settings.

On the other hand, D-Adaptation and Prodigy are similar to our method in their use of DA. However, their approaches cannot be extended to the constrained optimization setting and are limited to Lipschitz functions. Nonetheless, their methods yield notable results in experiments, demonstrating strong empirical performance.

In conclusion, the main limitation of recent distance-adaptation methods is their inability to automatically adapt to diverse problem classes. Specifically, these methods require using different hyperparameters, such as an estimate of the maximal gradient norm, to adjust to the specific problem class. Addressing this broader adaptability has been the central focus of this paper.

3 Universality of DADA: Examples of Applications

Let us demonstrate that our method is *universal* in the sense that it simultaneously works for multiple problem classes without the need for choosing different parameters for each of these function classes. For simplicity, we assume that $\nabla f(x^*) = 0$ (this happens, in particular, when our problem (1.1) is unconstrained) and measure the ϵ -accuracy in terms of the function residual. To simplify the notation, we also denote $\log_+ t := 1 + \log t$ and $\bar{D}_0 \coloneqq \max\{\bar{r}, \|x_0 - x^*\|\},\$ where \bar{r} is the parameter of our method.

Nonsmooth Lipschitz functions. This function class is defined by the inequality

$$
|f(x) - f(y)| \le L_0 \|x - y\|
$$

for all $x, y \in Q$. For this problem class, DADA requires at most (see Corollary [C.2\)](#page-18-0)

$$
O\left(\frac{L_0^2\bar{D}_0^2}{\epsilon^2}\log^2_+\frac{\bar{D}_0}{\bar{r}}\right)
$$

oracle calls to reach ϵ -accuracy, which coincides with the standard complexity of (sub)gradient methods [\[14,](#page-12-4) [16\]](#page-12-5), up to an extra logarithmic factor. Note that this logarithmic factor is common for all distance-adaptation methods [\[5,](#page-11-6) [9,](#page-11-3) [10,](#page-11-4) [12\]](#page-11-5).

Lipschitz-smooth functions. Another important class of functions are those with Lipschitz gradient:

$$
\|\nabla f(x) - \nabla f(y)\|_{*} \le L_1 \|x - y\|
$$

for all $x, y \in Q$. In this case, the complexity of our method is (see Corollary [C.4\)](#page-19-0)

$$
O\left(\frac{L_1\bar{D}_0^2}{\epsilon}\log_+^2\frac{\bar{D}_0}{\bar{r}}\right).
$$

This coincides with the standard complexity of the (nonaccelerated) gradient method on Lipschitz-smooth functions [\[16,](#page-12-5) Section 3] up to an extra logarithmic factor.

Note that the complexity of DADA is slightly worse than that of the classical gradient method with line search [\[15\]](#page-12-2), which achieves a complexity bound of $O(\frac{L_1D_0^2}{\epsilon} + \log|\frac{L_1}{\hat{L}_1}|)$, where \hat{L}_1 is the initial guess for L_1 . The difference is that the logarithmic factor in the latter estimate appears in an additive way instead of multiplicative.

Hölder-smooth functions. The previous two examples are subclasses of the more general class of Hölder-smooth functions. It is defined by the following inequality:

$$
\|\nabla f(x) - \nabla f(y)\|_{*} \le H_{\nu} \|x - y\|^{\nu}
$$

for all $x, y \in Q$, where $\nu \in [0, 1]$ and $H_{\nu} \geq 0$. Therefore, for $\nu = 0$, we get functions with bounded variation of subgradients (which contains all Lipschitz functions) and for $\nu = 1$ we get Lipschitz-smooth functions.

The complexity of DADA on this problem class is (see Corollary [C.6\)](#page-20-0)

$$
O\bigg(\bigg[\frac{H_{\nu}}{\epsilon}\bigg]^\frac{2}{1+\nu}\bar{D}_0^2\log_+^2\frac{\bar{D}_0}{\bar{r}}\bigg).
$$

This is similar to the $O\left(\left[\frac{H_{\nu}}{\epsilon}\right]^{\frac{2}{1+\nu}} D_0^2 + \log\left|\frac{H_{\nu}^{\frac{2}{1+\nu}}}{\hat{I} + \frac{1-\nu}{1+\nu}}\right|\right)$ $\frac{H_{\nu}^{1+\nu}}{\hat{L}\epsilon^{\frac{1-\nu}{1+\nu}}}$) complexity of the universal (nonaccel-erated) gradient method with line search (GM-LS) [\[15\]](#page-12-2), where \tilde{L} is the parameter of the method. Again, the complexity of GM-LS is slightly better since the logarithmic factor is additive (and not multiplicative). However, GM-LS is not guaranteed to work (well) on other problem classes such as those we consider next.

Functions with Lipschitz high-order derivative. This class generalizes the Lipschitzsmooth class. Functions in this class have the property that their pth derivative $(p \geq 2)$ is Lipschitz, i.e., for all $x, y \in Q$, we have

$$
\|\nabla^p f(x) - \nabla^p f(y)\| \le L_p \|x - y\|,
$$

where the ∥·∥ norm in the left-hand side is the usual operator norm of a symmetric p-linear opeator: $||A|| = \max_{h \in Q: ||h|| = 1} ||A[h]^p||$. For example, the pth power of the Euclidean norm is an example of a function in this class (see [\[21\]](#page-12-6)). The complexity of DADA on this problem class is (see Corollary [C.8\)](#page-21-0)

$$
O\bigg(\bigg[\max_{2\leq i\leq p}\bigg[\frac{p}{i!}\frac{\|\nabla^i f(x^*)\|_{*}}{\epsilon}\bigg]^{\frac{2}{i}}+\bigg[\frac{L_p}{p!\,\epsilon}\bigg]^{\frac{2}{p+1}}\bigg]\bar{D}_0^2\log_+^2\frac{\bar{D}_0}{\bar{r}}\bigg).
$$

Although line-search gradient methods might be better for Hölder-smooth problems, to our knowledge, they are not known to attain comparable bounds on this function class.

Quasi-self-concordant (QSC) functions [\[1\]](#page-10-0). A function f is called QSC with parameter $M > 0$ over the set Q if it is three times continuously differentiable and the following inequality holds for any $x \in Q$ and any $u, v \in \mathbb{R}^d$:

$$
\nabla^3 f(x)[u, u, v] \le M \langle \nabla^2 f(x)u, u \rangle ||v||. \tag{3.1}
$$

For example, the exponential, logistic, and softmax functions are QSC; for more details and other examples, see [\[6\]](#page-11-8). When applied to a QSC function, our method has the following complexity (Corollary [C.11\)](#page-22-0):

$$
O\biggl(\biggl[M^2\bar{D}_0^2+\frac{\|\nabla^2 f(x^*)\|\bar{D}_0^2}{\epsilon}\biggr]\log_+^2\frac{\bar{D}_0}{\bar{r}}\biggr).
$$

In terms of comparisons, second-order methods, such as those explored in [\[6\]](#page-11-8), are more powerful for minimizing QSC functions, as they leverage additional curvature information. Their complexity bound, in terms of queries to the second-order oracle, is $O(M\hat{D}_0 \log \frac{F_0}{\epsilon} +$ $\log \frac{\hat{D}_{0}g_{0}}{\epsilon F_{0}}$, where $F_{0} = f(x_{0}) - f^{*}$, \hat{D}_{0} is the diameter of the initial sublevel set, and $g_0 = ||\nabla f(x_0)||_*$ (see [\[6,](#page-11-8) Corollary 3.4]). However, each iteration of these methods is significantly more expensive.

To our knowledge, the QSC class has not been previously studied in the context of first-order methods. The only other first-order methods for which one can prove similar bounds are the nonadaptive variants of our scheme, namely the normalized gradient method (NGM) from [\[16,](#page-12-5) Section 5] and the recent improvement of this algorithm for constrained problems [\[17\]](#page-12-7).

 (L_0, L_1) -smooth functions. As introduced in [\[24\]](#page-12-8), a function f is said to be (L_0, L_1) smooth if for all $x \in \mathbb{R}^d$, we have

$$
\|\nabla^2 f(x)\| \le L_0 + L_1 \|\nabla f(x)\|_*
$$

The complexity of DADA on this class is (see Corollary [C.14\)](#page-23-0)

$$
O\bigg(\bigg[L_1^2\bar{D}_0^2 + \frac{L_0\bar{D}_0^2}{\epsilon}\bigg] \log_+^2 \frac{\bar{D}_0}{\bar{r}}\bigg).
$$

Up to the extra logarithmic factor, this matches the complexity of NGM from [\[23\]](#page-12-9), with the distinction that their approach is less robust to the initial guess of D_0 . Specifically, the penalty for underestimating it in the latter method is a multiplicative factor of $\rho^2 := \frac{D_0^2}{\bar{r}^2}$ while in our method this factor is logarithmic: $\log_+^2 \rho$.

4 Experiments

To evaluate the efficiency of our proposed method, DADA, we conduct a series of experiments on convex optimization problems. Our goal is to demonstrate the effectiveness of DADA in achieving competitive performance across various function classes without any hyperparameter tuning.

Figure 4.1: Comparison of different methods on the Softmax function.

We compare DADA against state-of-the-art distance-adaptation algorithms, namely, DoG [\[9\]](#page-11-3) and Prodigy [\[12\]](#page-11-5), using their official implementations without any modifications. We also consider the Universal Gradient Method (UGM) from [\[15\]](#page-12-2) and the classical Weighted Dual Averaging (WDA) method [\[14\]](#page-12-4). For UGM, we choose the initial value of the line-search parameter $L_0 = 1$ and set the target accuracy to $\epsilon = 10^{-6}$. For WDA, we use the coefficients $a_k = \frac{D_0}{\|a_k\|}$ $\frac{D_0}{\|g_k\|_*}$ and $\beta_k =$ √ \overline{k} , where $D_0 = ||x_0 - x^*||$.

For each method, we plot the best function value among all the test points generated by the algorithm against the number of first-order oracle calls. We set the starting point to $x_0 = (1, \ldots, 1)$ and select the initial guess for the distance to the solution as $\bar{r} = 10^{-6} (1 + ||x_0||).$

Softmax function. Our first test problem is

$$
\min_{x \in \mathbb{R}^d} \left\{ f(x) := \mu \log \left(\sum_{i=1}^n \exp \left[\frac{\langle a_i, x \rangle - b_i}{\mu} \right] \right) \right\},\
$$

where $a_i \in \mathbb{R}^d$, and $b_i \in \mathbb{R}$ for all $1 \leq i \leq n$, and $\mu > 0$. This function can be viewed as a smooth approximation of $\max_{1 \leq i \leq n} [\langle a_i, x \rangle - b_i]$ [\[13\]](#page-11-9).

To generate the data for our problem, we proceed as follows. First, we generate i.i.d. vectors \hat{a}_i with components uniformly distributed in the interval $[-1, 1]$ for $i = 1, \ldots, n$, and similarly for the scalar values b_i . Using this data, we form the preliminary version of our function, \hat{f} . We then compute $a_i = \hat{a}_i - \nabla \hat{f}(0)$ and use the obtained (a_i, b_i) to define our function f. This way of generating the data ensures that $x^* = 0$ is a solution of our problem.

The results are shown in Fig. [4.1,](#page-8-0) where we fix $n = 10^3$ and $d = 2n$, and consider different values of $\mu \in \{0.1, 0.01, 0.005\}$. As we can see, most methods exhibit similar performance for $\mu = 0.1$ except for Prodigy which stops converging after a few initial iterations. This issue, along with a decline in performance for UGM, persists as μ decreases, whereas DADA, DoG, and WDA remain largely unaffected. Notably, DoG performs very similarly to DADA, which we hypothesize is primarily due to the similarity in estimating D_0 .

Additionally, Fig. [4.2](#page-9-0) illustrates the ratio between D_0 and \bar{r} , showing the estimation error of Prodigy, DoG, and DADA throughout the optimization process. For Prodigy, we use $\frac{D_0}{d_{\text{max}}}$ to generate the plot. The figure demonstrates that DADA and DoG exhibit

Figure 4.2: The ratio $\frac{D}{\bar{r}_t}$ for the Softmax function with different optimal points x^* .

Figure 4.3: Comparison of different methods on the polyhedron feasibility problem.

similar behavior in estimating D_0 , despite employing different update methods—Dual Averaging and Gradient Descent, respectively. However, Prodigy appears to encounter challenges in estimating D_0 as its estimation stabilizes at a relatively large value.

Hölder-smooth function. Let us consider the following *polyhedron feasibility problem*:

$$
f^* := \min_{x \in \mathbb{R}^d} \left\{ f(x) := \frac{1}{n} \sum_{i=1}^n [\langle a_i, x \rangle - b_i]_+^q \right\},\tag{4.1}
$$

where $a_i, b_i \in \mathbb{R}^d$, $q \in [1, 2]$, and $[\tau]_+ = \max(0, \tau)$. This problem can be interpreted as finding a point $x^* \in \mathbb{R}^d$ lying inside the polyhedron $P = \{x : \langle a_i, x \rangle \leq b_i, i = 1, \ldots, n\}.$ Such a point exists if and only if $f^* = 0$.

Observe that f in problem [\(4.1\)](#page-9-1) is Hölder-smooth with parameter $\nu = q-1$. Therefore, by varying $q \in [1, 2]$, we can check the robustness of different methods to the smoothness level of the objective function.

The data for our problem is generated randomly, following the procedure in [\[20\]](#page-12-10). First, we sample x^* uniformly from the sphere of radius $0.95R$ centered at the origin. Next, we generate i.i.d. vectors a_i with components uniformly distributed in $[-1, 1]$. To ensure that $\langle a_n, x^* \rangle < 0$, we invert the sign of a_n if necessary. We then sample positive reals s_i uniformly from $[0, -0.1c_{\min}]$, where $c_{\min} \coloneqq \min_i \langle a_i, x^* \rangle < 0$, and set $b_i = \langle a_i, x^* \rangle + s_i$. By construction, x^* is a solution to the problem with $f^* = 0$.

We select $n = 10^4$, $d = 10^3$, $R = 10^3$ and consider different values of $q \in \{1, 1.5, 2\}$. As shown in Fig. [4.3,](#page-9-2) as q increases and approaches 2, the performance of DoG signifi-

Figure 4.4: Comparison of different methods on the worst-case function.

Figure 4.5: The ratio $\frac{D}{\bar{r}_t}$ for the worst-case function with different optimal points x^* .

cantly declines. However, DADA, Prodigy, and UGM demonstrate similar performance regardless of the choice of q.

Worst-case function. As an example of a function with Lipschitz high-order derivative, we consider the following worst-case problem from [\[7\]](#page-11-10):

$$
\min_{x \in \mathbb{R}^d} \left\{ f(x) := \frac{1}{p} \sum_{i=1}^{d-1} |x^{(i)} - x^{(i+1)}|^p + \frac{1}{p} |x^{(d)}|^p \right\},\tag{4.2}
$$

where $p \ge 2$, and $x^{(i)}$ is the *i*th element of x. The optimal point in this problem is $x^* = 0$.

As illustrated in Fig. [4.4,](#page-10-1) as in the Softmax case, nearly all methods exhibit similar performance when $p = 2$, except for Prodigy whose convergence becomes slow after a few initial iterations. Interestingly, this issue no longer persists as p increases. Instead, a noticeable decline in the performance of DoG is observed. This trend is further highlighted in Fig. [4.5,](#page-10-2) where the estimate of D_0 by Prodigy, despite showing some improvements, is still quite inaccurate. Furthermore, for DoG , the estimate degrades as p increases.

In contrast, both DADA and UGM demonstrate stable and consistent performance across different values of p, with DADA performing slightly better than UGM.

References

[1] F. Bach. Self-concordant analysis for logistic regression. Electronic Journal of Statis-tics, 4:384 -414, 2010. DOI: [10.1214/09-EJS521](https://doi.org/10.1214/09-EJS521). URL: [https://doi.org/10.1214/](https://doi.org/10.1214/09-EJS521) [09-EJS521](https://doi.org/10.1214/09-EJS521).

- [2] L. Bottou, F. E. Curtis, and J. Nocedal. Optimization methods for large-scale machine learning. $SIAM$ Review, $60(2):223-311$, 2018. URL: [https://doi.org/10.](https://doi.org/10.1137/16M1080173) [1137/16M1080173](https://doi.org/10.1137/16M1080173).
- [3] Y. Carmon and O. Hinder. Making SGD parameter-free. In Proceedings of Thirty Fifth Conference on Learning Theory, volume 178, pages 2360–2389, 2022. URL: <https://proceedings.mlr.press/v178/carmon22a.html>.
- [4] A. Cutkosky and F. Orabona. Black-box reductions for parameter-free online learning in Banach spaces. In Annual Conference Computational Learning Theory, 2018. url: <https://api.semanticscholar.org/CorpusID:3346292>.
- [5] A. Defazio and K. Mishchenko. Learning-rate-free learning by D-adaptation. In Proceedings of the 40th International Conference on Machine Learning, volume 202, pages 7449–7479, 2023. url: [https://proceedings.mlr.press/v202/defazio23a.](https://proceedings.mlr.press/v202/defazio23a.html) [html](https://proceedings.mlr.press/v202/defazio23a.html).
- [6] N. Doikov. Minimizing quasi-self-concordant functions by gradient regularization of Newton method, 2023. eprint: <2308.14742>. url: [https://arxiv.org/abs/2308.](https://arxiv.org/abs/2308.14742) [14742](https://arxiv.org/abs/2308.14742).
- [7] N. Doikov, K. Mishchenko, and Y. Nesterov. Super-universal regularized Newton method. *SIAM Journal on Optimization*, $34(1):27-56$, 2024 . DOI: 10.1137/ [22M1519444](https://doi.org/10.1137/22M1519444). eprint: <https://doi.org/10.1137/22M1519444>. url: [https://doi.](https://doi.org/10.1137/22M1519444) [org/10.1137/22M1519444](https://doi.org/10.1137/22M1519444).
- [8] J. Duchi, E. Hazan, and Y. Singer. Adaptive subgradient methods for online learning and stochastic optimization. Journal of Machine Learning Research, 12(61):2121– 2159, 2011. url: <http://jmlr.org/papers/v12/duchi11a.html>.
- [9] M. Ivgi, O. Hinder, and Y. Carmon. DoG is SGD's best friend: a parameter-free dynamic step size schedule. In Proceedings of the 40th International Conference on Machine Learning, pages 14465–14499, 2023. url: [https://proceedings.mlr.](https://proceedings.mlr.press/v202/ivgi23a.html) [press/v202/ivgi23a.html](https://proceedings.mlr.press/v202/ivgi23a.html).
- [10] A. Khaled, K. Mishchenko, and C. Jin. DoWG unleashed: an efficient universal parameter-free gradient descent method. In Advances in Neural Information Processing Systems, volume 36, pages 6748–6769, 2023. url: [https://proceedings.](https://proceedings.neurips.cc/paper_files/paper/2023/file/15ce36d35622f126f38e90167de1a350-Paper-Conference.pdf) [neurips.cc/paper_files/paper/2023/file/15ce36d35622f126f38e90167de1a35](https://proceedings.neurips.cc/paper_files/paper/2023/file/15ce36d35622f126f38e90167de1a350-Paper-Conference.pdf)0- [Paper-Conference.pdf](https://proceedings.neurips.cc/paper_files/paper/2023/file/15ce36d35622f126f38e90167de1a350-Paper-Conference.pdf).
- [11] Z. Liu and Z. Zhou. Stochastic nonsmooth convex optimization with heavy-tailed noises. ArXiv, abs/2303.12277, 2023. url: [https://api.semanticscholar.org/](https://api.semanticscholar.org/CorpusID:257663403) [CorpusID:257663403](https://api.semanticscholar.org/CorpusID:257663403).
- [12] K. Mishchenko and A. Defazio. Prodigy: an expeditiously adaptive parameter-free learner. In Proceedings of the 41st International Conference on Machine Learning, volume 235, pages 35779–35804, 2024. url: [https://proceedings.mlr.press/](https://proceedings.mlr.press/v235/mishchenko24a.html) [v235/mishchenko24a.html](https://proceedings.mlr.press/v235/mishchenko24a.html).
- [13] Y. Nesterov. Smooth minimization of non-smooth functions. Mathematical programming, 103:127–152, 2005.
- [14] Y. Nesterov. Primal-dual subgradient methods for convex problems. Mathematical Programming, 120:221–259, 2005. url: [https : / / api . semanticscholar . org /](https://api.semanticscholar.org/CorpusID:14935076) [CorpusID:14935076](https://api.semanticscholar.org/CorpusID:14935076).
- [15] Y. Nesterov. Universal gradient methods for convex optimization problems. Mathematical Programming, 152:381-404, 2015. URL: [https://api.semanticscholar.](https://api.semanticscholar.org/CorpusID:18062781) [org/CorpusID:18062781](https://api.semanticscholar.org/CorpusID:18062781).
- [16] Y. Nesterov. Lectures on Convex Optimization. Springer Publishing Company, Incorporated, 2nd edition, 2018, pages 221–259. isbn: 3319915770. url: [https://](https://api.semanticscholar.org/CorpusID:14935076) api.semanticscholar.org/CorpusID:14935076.
- [17] Y. Nesterov. Primal subgradient methods with predefined step sizes. Journal of Optimization Theory and Applications, 2024. DOI: [10.1007/s10957-024-02456-9](https://doi.org/10.1007/s10957-024-02456-9). url: <https://arxiv.org/abs/2308.14742>.
- [18] F. Orabona and T. Tommasi. Training deep networks without learning rates through coin betting. In Neural Information Processing Systems, 2017. URL: [https://api.](https://api.semanticscholar.org/CorpusID:6762437) [semanticscholar.org/CorpusID:6762437](https://api.semanticscholar.org/CorpusID:6762437).
- [19] D. Patterson, J. Gonzalez, Q. Le, C. Liang, L.-M. Munguia, D. Rothchild, D. So, M. Texier, and J. Dean. Carbon emissions and large neural network training, 2021. eprint: <2104.10350>. url: <https://arxiv.org/abs/2104.10350>.
- [20] A. Rodomanov, X. Jiang, and S. U. Stich. Universality of AdaGrad stepsizes for stochastic optimization: inexact oracle, acceleration and variance reduction. In The Thirty-eighth Annual Conference on Neural Information Processing Systems, 2024. url: <https://openreview.net/forum?id=rniiAVjHi5>.
- [21] A. Rodomanov and Y. Nesterov. Smoothness parameter of power of Euclidean norm. Journal of Optimization Theory and Applications, 185:303-326, 2019. URL: [https:](https://api.semanticscholar.org/CorpusID:198968030) [//api.semanticscholar.org/CorpusID:198968030](https://api.semanticscholar.org/CorpusID:198968030).
- [22] O. Sharir, B. Peleg, and Y. Shoham. The cost of training NLP models: a concise overview, 2020. eprint: <2004.08900>. url: <https://arxiv.org/abs/2004.08900>.
- [23] D. Vankov, A. Rodomanov, A. Nedich, L. Sankar, and S. U. Stich. Optimizing (L_0, L_1) -smooth functions by gradient methods, 2024. eprint: 2410.10800. URL: <https://arxiv.org/abs/2410.10800>.
- [24] J. Zhang, T. He, S. Sra, and A. Jadbabaie. Why gradient clipping accelerates training: a theoretical justification for adaptivity. In International Conference on Learning Representations, 2020. url: <https://openreview.net/forum?id=BJgnXpVYwS>.

A Auxiliary Results

The following result has been established in prior works such as [\[11,](#page-11-11) Lemma 30]. We include the proof here for the reader's convenience.

Lemma A.1. Let $(d_i)_{i=0}^{\infty}$ be a positive nondecreasing sequence. Then for any $T \geq 1$,

$$
\min_{1\leq k\leq T}\frac{d_k}{\sum_{i=0}^{k-1}d_i}\leq \frac{(\frac{d_T}{d_0})^{\frac{1}{T}}\log\frac{ed_T}{d_0}}{T}.
$$

Proof. Let $A_k \coloneqq \frac{1}{d_k}$ $\frac{1}{d_k} \sum_{i=0}^{k-1} d_i$ for each $k \geq 0$ (so that $A_0 = 0$). Then, for each $k \geq 0$, we have

$$
d_{k+1}A_{k+1} - d_kA_k = d_k,
$$

which implies that

$$
\frac{d_k}{d_{k+1}} = A_{k+1} - \frac{d_k}{d_{k+1}} A_k = A_{k+1} - A_k + \left(1 - \frac{d_k}{d_{k+1}}\right) A_k.
$$

Summing up these identities for all $0 \leq k \leq T-1$, we get

$$
S_T \coloneqq \sum_{k=0}^{T-1} \frac{d_k}{d_{k+1}} = A_T + \sum_{k=0}^{T-1} \left(1 - \frac{d_k}{d_{k+1}} \right) A_k \le A_T^*(1 + T - S_T),
$$

where $A_T^* = \max_{0 \le k \le T} A_k \equiv \max_{1 \le k \le T} A_k$ and we have used the fact that $(d_i)_{i=0}^{\infty}$ is nondecreasing. Hence,

$$
A_T^* \ge \frac{S_T}{1+T-S_T}.
$$

Applying now the AM-GM inequality and denoting $\gamma_T = (\frac{d_0}{d_T})^{\frac{1}{T}}$ ($\in (0, 1]$), we can further estimate $S_T \geq T \gamma_T$, giving us

$$
A_T^* \ge \frac{T\gamma_T}{1 + T(1 - \gamma_T)}.
$$

Thus,

$$
\min_{1 \le k \le T} \frac{d_k}{\sum_{i=0}^{k-1} d_i} = \frac{1}{A_T^*} \le \frac{\frac{1}{\gamma_T} \left(1 + T(1 - \gamma_T)\right)}{T}.
$$

Estimating further $T(1-\gamma_T) \leq -T \log \gamma_T \equiv \log \frac{1}{\gamma_T T}$ and substituting the definition of γ_T , we get the claim. \Box

The following lemma is a classical result from [\[16,](#page-12-5) Lemma 3.2.1].

Lemma A.2. For any $x \in Q$ we have $f(x) - f^* \leq \omega(v(x))$, where $\omega(\cdot)$ and $v(\cdot)$ are defined as in [\(2.1\)](#page-2-3) and [\(2.3\)](#page-2-4) (with $\nabla f(x)$ being an arbitrary subgradient from $\partial f(x)$).

Proof. Let \bar{x} denote the orthogonal projection of x^* onto the supporting hyperplane ${y : \langle \nabla f(x), x - y \rangle = 0}$:

$$
\bar{x} = x^* + v(x) \frac{\nabla f(x)}{\|\nabla f(x)\|_*}.
$$

Then, $\langle \nabla f(x), \bar{x} - x \rangle = 0$, and $\|\bar{x} - x^*\| = v(x)$. Therefore,

$$
f(x) \le f(\bar{x}) + \langle \nabla f(x), \bar{x} - x \rangle = f(\bar{x}),
$$

and hence,

$$
f(x) - f^* \le f(\bar{x}) - f^* \le \omega(\|\bar{x} - x^*\|) = \omega(v(x)).
$$

Lemma A.3. Consider the nonnegative sequence $(d_k)_{k=0}^{\infty}$ that satisfies, for each $k \geq 0$,

$$
d_{k+1} \le \max\{d_k, R + \gamma d_k\},\
$$

where $0 \leq \gamma < 1$ and $R \geq 0$ are certain constants. Then, for any $k \geq 0$, we have

$$
d_k \le \max\left\{\frac{1}{1-\gamma}R, d_0\right\}.
$$

Proof. We use induction to prove that $d_k \leq D$ for a certain constant D to be determined later. To ensure that this relation holds for $k = 0$, we need to choose $D \geq d_0$. Let us now suppose that our relation has already been proved for some $k \geq 0$ and let us prove it for the next index $k + 1$. Using the induction hypothesis and the given inequality, we obtain

$$
d_{k+1} \le \max\{d_k, R + \gamma d_k\} \le \max\{D, R + \gamma D\}.
$$

To prove that the right-hand side is $\leq D$, we need to ensure that $R + \gamma D \leq D$, which means that we need to choose $D \geq \frac{1}{1-\gamma}R$. Combining this requirement with that from the base of induction, we see that we can choose $D = \max\{\frac{1}{1-\gamma}R, d_0\}.$ \Box

B Proof of Theorem [2.1](#page-3-0)

Lemma B.1. In Algorithm [2.1,](#page-2-1) for any $1 \leq k \leq T$, it holds that

$$
\sum_{i=0}^{k-1} a_i \langle g_i, x_i - x^* \rangle + \frac{\beta_k}{2} \| x_k - x^* \|^2 \le \frac{\beta_k}{2} \| x_0 - x^* \|^2 + \sum_{i=0}^{k-1} \frac{a_i^2}{2\beta_i} \| g_i \|_*^2,
$$

where β_0 is an arbitrary coefficient in $(0, \beta_1]$.

Proof. For any $0 \leq k \leq T$, define the function $\psi_k(x)$ as follows:

$$
\psi_k(x) := \sum_{i=0}^{k-1} a_i \langle g_i, x - x_i \rangle + \frac{\beta_k}{2} ||x - x_0||^2,
$$

so that $\psi_0(x) = \frac{\beta_0}{2} ||x - x_0||^2$ (with β_0 as defined in the statement). Note that ψ_k is a β_k -strongly convex function and x_k is its minimizer. Hence, for any $x \in Q$ and $0 \leq k \leq T$, we have

$$
\psi_k(x) \ge \psi_k^* + \frac{\beta_k}{2} \|x - x_k\|^2,\tag{B.1}
$$

where $\psi_k^* \coloneqq \psi_k(x_k)$. Consequently,

$$
\psi_{k+1}^* = \psi_{k+1}(x_{k+1}) = \psi_k(x_{k+1}) + a_k \langle g_k, x_{k+1} - x_k \rangle + \frac{\beta_{k+1} - \beta_k}{2} ||x_{k+1} - x_0||^2
$$

\n
$$
\geq \psi_k^* + \frac{\beta_k}{2} ||x_{k+1} - x_k||^2 + a_k \langle g_k, x_{k+1} - x_k \rangle + \frac{\beta_{k+1} - \beta_k}{2} ||x_{k+1} - x_0||^2
$$

\n
$$
\geq \psi_k^* + \frac{\beta_k}{2} ||x_{k+1} - x_k||^2 + a_k \langle g_k, x_{k+1} - x_k \rangle \geq \psi_k^* - \frac{a_k^2}{2\beta_k} ||g_k||_*^2.
$$

Telescoping these inequalities and using the fact that $\psi_0^* = 0$, we obtain

$$
\psi_k^* \ge -\sum_{i=0}^{k-1} \frac{a_i^2}{2\beta_i} \|g_i\|_*^2.
$$

Combining this inequality with the definition of ψ_k and [\(B.1\)](#page-15-2), we thus obtain

$$
\sum_{i=0}^{k-1} a_i \langle g_i, x^* - x_i \rangle + \frac{\beta_k}{2} \| x_0 - x^* \|^2 = \psi_k(x^*) \ge \psi_k^* + \frac{\beta_k}{2} \| x_k - x^* \|^2
$$

$$
\ge -\sum_{i=0}^{k-1} \frac{a_i^2}{2\beta_i} \| g_i \|_*^2 + \frac{\beta_k}{2} \| x_k - x^* \|^2.
$$

Rearranging, we get the claim.

Lemma B.2. Consider Algorithm [2.1](#page-2-1) using the coefficients defined in [\(2.4\)](#page-2-2). Then, the following inequality holds for all $1 \leq k \leq T$:

$$
\sum_{i=0}^{k-1} \bar{r}_i v_i + \frac{c\sqrt{k+1}}{2} D_k^2 \le \frac{c\sqrt{k+1}}{2} D_0^2 + \frac{\sqrt{k}}{c} \bar{r}_{k-1}^2,
$$

where $D_k = ||x_k - x^*||$ and $v_i := v(x_i)$.

Proof. Applying Lemma [B.1](#page-14-1) and the definition of v_i , we obtain

$$
\sum_{i=0}^{k-1} a_i v_i \|g_i\|_{*} + \frac{\beta_k}{2} D_k^2 \le \frac{\beta_k}{2} D_0^2 + \sum_{i=0}^{k-1} \frac{a_i^2}{2\beta_i} \|g_i\|_{*}^2.
$$

Substituting our choice of the coefficients given by [\(2.4\)](#page-2-2), we get

$$
\sum_{i=0}^{k-1} \bar{r}_i v_i + \frac{c\sqrt{k+1}}{2} D_k^2 \leq \frac{c\sqrt{k+1}}{2} D_0^2 + \frac{1}{2c} \sum_{i=0}^{k-1} \frac{\bar{r}_i^2}{\sqrt{i+1}} \leq \frac{c\sqrt{k+1}}{2} D_0^2 + \frac{\sqrt{k}}{c} \bar{r}_{k-1}^2,
$$

where we have used the fact that \bar{r}_k is nondecreasing and $\sum_{i=0}^{k-1} \frac{1}{\sqrt{i+1}} \leq 2$ √ k.

Lemma B.3. Consider Algorithm [2.1](#page-2-1) using the coefficients defined in [\(2.4\)](#page-2-2) and assume **Lemma B.3.** Consuder Algorithm 2.1 using the coefficients defined in $\langle \text{that } c \rangle \sqrt{2}$. Then, we have the following inequalities for all $0 \leq k \leq T$.

$$
\bar{r}_k \leq \bar{D}, \qquad D_k \leq D_0 + \frac{\sqrt{2}}{c} \bar{D},
$$

where $\bar{D} \coloneqq \max\{\bar{r}, \frac{2c}{\bar{r}}\}$ $\frac{2c}{c-\sqrt{2}}D_0$ } and $D_k := ||x_k - x^*||$.

Proof. Both bounds are clearly valid for $k = 0$, so it suffices to consider only the case when $1 \leq k \leq T$.

Applying Lemma [B.2,](#page-15-0) dropping the nonnegative $\bar{r}_i v_i$ from the left-hand side and rearranging, we obtain

$$
D_k^2 \le D_0^2 + \frac{2\sqrt{k}}{c^2\sqrt{k+1}} \bar{r}_{k-1}^2 \le D_0^2 + \frac{2}{c^2} \bar{r}_{k-1}^2.
$$

Consequently,

$$
D_k \le D_0 + \frac{\sqrt{2}}{c} \bar{r}_{k-1}.
$$
 (B.2)

Therefore,

$$
r_k \equiv \|x_k - x_0\| \le D_k + D_0 \le 2D_0 + \frac{\sqrt{2}}{c}\bar{r}_{k-1}.
$$

Hence,

$$
\bar{r}_k \equiv \max{\{\bar{r}_{k-1}, r_k\}} \le \max{\{\bar{r}_{k-1}, 2D_0 + \frac{\sqrt{2}}{c}\bar{r}_{k-1}\}}.
$$

Since $k \geq 1$ was allowed to be arbitrary, we can apply Lemma [A.3](#page-14-2) to conclude that

$$
\bar{r}_k \le \max\left\{\bar{r}, \frac{2}{1 - \frac{\sqrt{2}}{c}} D_0\right\} = \max\left\{\bar{r}, \frac{2c}{c - \sqrt{2}} D_0\right\} \equiv \bar{D}.
$$

This proves the first part of the claim.

Substituting the already proved bound on \bar{r}_k into [\(B.2\)](#page-16-0), we obtain the claimed upper bound on D_k . \Box

We are now ready to prove the main result.

Proof of Theorem [2.1.](#page-3-0) Let $T \geq 1$ be arbitrary. According to Lemma [A.2](#page-13-2) and the fact that $\omega(\cdot)$ is nondecreasing, we can write

$$
f(x_T^*) - f^* = \min_{0 \le k \le T-1} [f(x_k) - f^*] \le \min_{0 \le k \le T-1} \omega(v_k) = \omega(v_T^*),
$$

where $v_k := v(x_k)$ and $v^*_T := \min_{0 \leq k \leq T-1} v_k$. This proves the first part of the claim.

Let us now estimate the rate of convergence of v_T^* . To that end, let us fix an arbitrary $1 \leq k \leq T$. In view of Lemma [B.2,](#page-15-0) we have

$$
\sum_{i=0}^{k-1} \bar{r}_i v_i \le \frac{c\sqrt{k+1}}{2}(D_0^2 - D_k^2) + \frac{\sqrt{k}}{c}\bar{r}_{k-1}^2,
$$

where $D_k = ||x_k - x^*||$. Note that

$$
D_0^2 - D_k^2 \equiv ||x_0 - x^*||^2 - ||x_k - x^*||^2 = (||x_0 - x^*|| - ||x_k - x^*||)(||x_0 - x^*|| + ||x_k - x^*||)
$$

\n
$$
\leq 2||x_k - x_0|| ||x_0 - x^*|| \equiv 2r_k D_0.
$$

Therefore, we can continue as follows:

$$
\sum_{i=0}^{k-1} \bar{r}_i v_i \le c\sqrt{k+1} r_k D_0 + \frac{\sqrt{k}}{c} \bar{r}_{k-1}^2 \le (cD_0 + \frac{1}{c} \bar{r}_{k-1})\sqrt{k+1} \bar{r}_k
$$

$$
\le (cD_0 + \frac{1}{c}\bar{D})\sqrt{k+1} \bar{r}_k = D\sqrt{\frac{k+1}{2}} \bar{r}_k,
$$

where the second inequality is due to the fact that $\bar{r}_k = \max{\{\bar{r}_{k-1}, r_k\}}$, the final inequality is due to Lemma [B.3,](#page-15-1) and the constants \bar{D} and D are as defined in the statement. Hence,

$$
v_k^* \equiv \min_{0 \le i \le k-1} v_i \le \frac{\sum_{i=0}^{k-1} \bar{r}_i v_i}{\sum_{i=0}^{k-1} \bar{r}_i} \le \frac{\bar{r}_k}{\sum_{i=0}^{k-1} \bar{r}_i} D \sqrt{\frac{k+1}{2}}
$$

.

Letting now $k^* = \operatorname{argmin}_{1 \le k \le T} \frac{\bar{r}_k}{\sum_{i=0}^{k-1} \bar{r}_i}$ and using Lemma [A.1,](#page-13-1) we obtain

$$
v_T^* \le v_{k^*}^* \le \frac{D\sqrt{\frac{k^*+1}{2}}}{T} \left(\frac{\bar{r}_T}{\bar{r}}\right)^{\frac{1}{T}} \log \frac{e\bar{r}_T}{\bar{r}} \le \frac{D}{\sqrt{T}} \left(\frac{\bar{D}}{\bar{r}}\right)^{\frac{1}{T}} \log \frac{e\bar{D}}{\bar{r}},
$$

where we have used the fact that $k^* + 1 \leq T + 1 \leq 2T$ (since $1 \leq k^* \leq T$) and that $\bar{r}_T \leq \bar{D}$ (see Lemma [B.3\)](#page-15-1). This proves [\(2.5\)](#page-3-2) in the case when $T \geq \log \frac{\bar{D}}{\bar{r}}$ since then we can further bound $(\frac{\bar{D}}{\bar{r}})^{\frac{1}{T}} \equiv \exp(\frac{1}{T} \log \frac{\bar{D}}{\bar{r}}) \leq e$.

On the other hand, by the definition of v_k and Lemma [B.3,](#page-15-1) we always have the following trivial inequality for any $0 \leq k \leq T-1$:

$$
v_k \equiv \frac{\langle \nabla f(x_k), x_k - x^* \rangle}{\|\nabla f(x_k)\|_{*}} \le D_k \le D_0 + \frac{\sqrt{2}}{c} \bar{D} \le D.
$$

This means that [\(2.5\)](#page-3-2) is also satisfied in the case when $T \leq \log \frac{\bar{D}}{\bar{r}}$ since then $\frac{eD}{\sqrt{T}}$ $\frac{D}{\overline{T}} \log \frac{e\overline{D}}{\overline{r}} \geq$ √ $\frac{\overline{D}}{\overline{T}}\log \frac{\overline{D}}{\overline{r}}\geq D$ $\frac{D}{\epsilon}$ $T \geq D$ (we still consider $T \geq 1$). The proof of (2.5) is now finished. The final part of the claim readily follows from [\(2.5\)](#page-3-2). \Box

C Convergence of DADA on Various Problem Classes

In this section, we analyze the complexity of DADA across different problem classes. To achieve this, we first establish bounds on the growth function:

$$
\omega(t) = \max_{x} \{ f(x) - f^* : ||x - x^*|| \le t \},\
$$

and determine the threshold t such that $\omega(t) \leq \epsilon$ for a given ϵ . Subsequently, we combine these results with the complexity bound $T(\delta)$ derived in Theorem [2.1,](#page-3-0) enabling us to estimate the oracle complexity of DADA for finding an ϵ -solution in terms of the function residual.

C.1 Nonsmooth Lipschitz Functions

In this section, we assume that the function f in problem (1.1) is Lipschitz: for all $x, y \in Q$, it holds that

$$
|f(x) - f(y)| \le L_0 \|x - y\|,
$$

where $L_0 > 0$ is a fixed constant.

Lemma C.1. Assume that f is an L_0 -Lipschitz function. Then, $\omega(t) \leq \epsilon$ for any given $\epsilon > 0$ whenever $t \leq \delta(\epsilon)$, where

$$
\delta(\epsilon) \coloneqq \frac{\epsilon}{L_0}.
$$

Proof. Indeed, for any $x \in Q$, we have

$$
f(x) - f^* \le L_0 ||x - x^*||,
$$

Therefore, for any $t \geq 0$, we have

 $\omega(t) \leq L_0 t$.

Making the right-hand side $\leq \epsilon$, we get the claim.

Combining Theorem [2.1](#page-3-0) and Lemma [C.1,](#page-18-1) we get the following complexity result.

Corollary C.2. Consider problem [\(1.1\)](#page-1-0) under the assumption that f is an L_0 -Lipschitz function. Let Algorithm [2.1](#page-2-1) with coefficients [\(2.4\)](#page-2-2) be applied for solving this problem. Then, $f(x_T^*) - f^* \leq \epsilon$ for any given $\epsilon > 0$ whenever $T \geq T(\epsilon)$, where

$$
T(\epsilon) = \frac{e^2 L_0^2 D^2}{\epsilon^2} \log^2 \frac{e \bar{D}}{\bar{r}},
$$

and the constants D and \bar{D} are as defined in Theorem [2.1.](#page-3-0)

C.2 Lipschitz-Smooth Functions

Let us now consider the case when f is Lipschitz-smooth, meaning that for any $x, y \in Q$, the following inequality holds:

$$
f(y) \le f(x) + \langle \nabla f(x), y - x \rangle + \frac{L_1}{2} ||y - x||^2,
$$

where $L_1 > 0$ is a fixed constant.

Lemma C.3. Assume that f is Lipschitz-smooth with constant L_1 . Then, $\omega(t) \leq \epsilon$ for any given $\epsilon > 0$ whenever $t \leq \delta(\epsilon)$, where

$$
\delta(\epsilon) \coloneqq \min \left\{ \sqrt{\frac{\epsilon}{L_1}}, \frac{\epsilon}{2\|\nabla f(x^*)\|_{*}} \right\}.
$$

Proof. Indeed, for any $x \in Q$, we have

$$
f(x) - f^* \le \langle \nabla f(x^*), x - x^* \rangle + \frac{L_1}{2} ||x - x^*||^2
$$

$$
\le ||\nabla f(x^*)||_* ||x - x^*|| + \frac{L_1}{2} ||x - x^*||^2.
$$

Hence, for any $t \geq 0$,

$$
\omega(t) \le \frac{L_1}{2}t^2 + \|\nabla f(x^*)\|_* t.
$$

To make the right-hand side $\leq \epsilon$, it suffices to ensure that each of the two terms is $\leq \frac{\epsilon}{2}$ $\frac{\epsilon}{2}$:

$$
\frac{L_1}{2}t^2 \le \frac{\epsilon}{2}, \qquad \|\nabla f(x^*)\|_* t \le \frac{\epsilon}{2}.
$$

Solving this system of inequalities, we get the claim.

Combining Theorem [2.1](#page-3-0) and Lemma [C.3,](#page-18-2) we get the following complexity result.

Corollary C.4. Consider problem (1.1) under the assumption that f is Lipschitz-smooth with constant L_1 . Let Algorithm [2.1](#page-2-1) with coefficients [\(2.4\)](#page-2-2) be applied for solving this problem. Then, $f(x_T^*) - f^* \leq \epsilon$ for any given $\epsilon > 0$ whenever $T \geq T(\epsilon)$, where

$$
T(\epsilon) = \max\left\{\frac{L_1}{\epsilon}, \frac{4\|\nabla f(x^*)\|_*^2}{\epsilon^2}\right\} e^2 D^2 \log^2 \frac{e\bar{D}}{\bar{r}},
$$

and the constants D and \overline{D} are as defined in Theorem [2.1.](#page-3-0)

C.3 Hölder-Smooth Functions

Let us now consider a more general case, when f is Hölder-smooth, meaning that for any $x, y \in Q$, it holds that

$$
f(y) \le f(x) + \langle \nabla f(x), y - x \rangle + \frac{H_{\nu}}{1 + \nu} ||y - x||^{1 + \nu},
$$

where $\nu \in [0, 1]$ and $H_{\nu} > 0$.

Lemma C.5. Assume that f is a Hölder-smooth function with constants ν and H_{ν} . Then, $\omega(t) \leq \epsilon$ for any given $\epsilon > 0$ whenever $t \leq \delta(\epsilon)$, where

$$
\delta(\epsilon) := \min \left\{ \left[\frac{(1+\nu)\epsilon}{2H_{\nu}} \right]^{\frac{1}{1+\nu}}, \frac{\epsilon}{2\|\nabla f(x^*)\|_{*}} \right\}.
$$

Proof. Indeed, for any $x \in Q$, we have

$$
f(x) - f(x^*) \le \langle \nabla f(x^*), x - x^* \rangle + \frac{H_\nu}{1 + \nu} \|x - x^*\|^{1 + \nu}
$$

$$
\le \|\nabla f(x^*)\|_{*} \|x - x^*\| + \frac{H_\nu}{1 + \nu} \|x - x^*\|^{1 + \nu}.
$$

Hence, for any $t \geq 0$,

$$
\omega(t)\leq \|\nabla f(x^*)\|_*t+\frac{H_\nu}{1+\nu}t^{1+\nu}.
$$

To make the right-hand side of the last inequality $\leq \epsilon$, it suffices to ensure that each of the two terms is $\leq \frac{\epsilon}{2}$ $\frac{\epsilon}{2}$:

$$
\|\nabla f(x^*)\|_* t \le \frac{\epsilon}{2}, \qquad \frac{H_\nu}{1+\nu} t^{1+\nu} \le \frac{\epsilon}{2}.
$$

Solving this system of inequalities, we get the claim.

Combining Theorem [2.1](#page-3-0) and Lemma [C.5,](#page-19-1) we get the following complexity result.

Corollary C.6. Consider problem (1.1) under the assumption that f is a Hölder-smooth function with constants ν and H_{ν} . Let Algorithm [2.1](#page-2-1) with coefficients [\(2.4\)](#page-2-2) be applied for solving this problem. Then, $f(x_T^*) - f^* \leq \epsilon$ for any given $\epsilon > 0$ whenever $T \geq T(\epsilon)$, where

$$
T(\epsilon) = \max \left\{ \left[\frac{2H_{\nu}}{(1+\nu)\epsilon} \right]^{\frac{2}{1+\nu}}, \frac{4\|\nabla f(x^*)\|_{*}^2}{\epsilon^2} \right\} e^2 D^2 \log^2 \frac{e\bar{D}}{\bar{r}},
$$

and the constants D and \overline{D} are as defined in Theorem [2.1.](#page-3-0)

C.4 Functions with Lipschitz High-Order Derivative

In this section, we assume that function f in problem [\(1.1\)](#page-1-0) has L_p -Lipschitz pth derivative, where $p \geq 2$. It means that for any $x, y \in Q$, the following inequality holds:

$$
\|\nabla^p f(x) - \nabla^p f(y)\| \le L_p \|x - y\|.
$$

This implies the following global upper bound on the function value:

$$
f(y) \le f(x) + \sum_{i=1}^p \frac{1}{i!} \nabla^i f(x) [y - x]^i + \frac{L_p}{(p+1)!} \|y - x\|^{p+1}.
$$

Lemma C.7. Assume that f has L_p -Lipschitz pth derivative. Then, $\omega(t) \leq \epsilon$ for any given $\epsilon > 0$ whenever $t \leq \delta(\epsilon)$, where

$$
\delta(\epsilon) \coloneqq \min \left\{ \min_{2 \leq i \leq p} \left[\frac{i! \epsilon}{(p+1) \|\nabla^i f(x^*)\|} \right]^{\frac{1}{i}}, \left[\frac{p! \epsilon}{L_p} \right]^{\frac{1}{p+1}}, \frac{\epsilon}{(p+1) \|\nabla f(x^*)\|_*} \right\}.
$$

Proof. Indeed, for any $x \in Q$, we have

$$
f(x) - f^* \le \langle \nabla f(x^*), x - x^* \rangle + \sum_{i=2}^p \frac{1}{i!} \nabla^i f(x^*) [x - x^*]^i + \frac{L_p}{(p+1)!} \|x - x^*\|^{p+1}
$$

$$
\le \|\nabla f(x^*)\|_{*} \|x - x^*\| + \sum_{i=2}^p \frac{1}{i!} \|\nabla^i f(x^*)\| \|x - x^*\|^i + \frac{L_p}{(p+1)!} \|x - x^*\|^{p+1}.
$$

Therefore, for any $t > 0$, we have

$$
\omega(t) \leq \|\nabla f(x^*)\|_* t + \sum_{i=2}^p \frac{1}{i!} \|\nabla^i f(x^*)\| t^i + \frac{L_p}{(p+1)!} t^{p+1}.
$$

To make the right-hand side $\leq \epsilon$, it suffices to ensure that each of the following inequalities holds:

$$
\|\nabla f(x^*)\|_* t \leq \frac{\epsilon}{p+1}, \quad \frac{1}{i!} \|\nabla^i f(x^*)\| t^i \leq \frac{\epsilon}{p+1}, \quad \frac{L_p}{(p+1)!} t^{p+1} \leq \frac{\epsilon}{p+1}, \quad i=2,\ldots,p.
$$

Solving this system of inequalities, we get the claim.

Combining Theorem [2.1](#page-3-0) and Lemma [C.7,](#page-20-1) we get the following complexity result.

Corollary C.8. Consider problem [\(1.1\)](#page-1-0) under the assumption that f has L_p -Lipschitz pth derivative. Let Algorithm [2.1](#page-2-1) with coefficients [\(2.4\)](#page-2-2) be applied for solving this problem. Then, $f(x_T^*) - f^* \leq \epsilon$ for any given $\epsilon > 0$ whenever $T \geq T(\epsilon)$, where

$$
T(\epsilon) = \max \left\{ \max_{2 \le i \le p} \left[\frac{(p+1) \|\nabla^i f(x^*)\|}{i! \epsilon} \right]^{\frac{2}{i}}, \left[\frac{L_p}{p! \epsilon} \right]^{\frac{2}{p+1}}, \frac{(p+1)^2 \|\nabla f(x^*)\|_{*}^2}{\epsilon^2} \right\} e^2 D^2 \log^2 \frac{e\bar{D}}{\bar{r}},
$$

and the constants D and \bar{D} are as defined in Theorem [2.1.](#page-3-0)

C.5 Quasi-Self-Concordant Functions

In this section, we assume that the function f in problem (1.1) is Quasi-Self-Concordant (QSC), meaning that it is three times continuously differentiable and for any $x \in Q$ and arbitrary directions $u, v \in \mathbb{R}^d$, it holds that

$$
\nabla^3 f(x)[u, u, v] \le M \langle \nabla^2 f(x)u, u \rangle ||v||,
$$

where $M \geq 0$ is a fixed parameter.

The following lemma provides an important global upper bound on the function value for QSC functions.

Lemma C.9. [\[6,](#page-11-8) Lemma 2.7] Let f be QSC with the parameter M. Then, for any $x, y \in Q$, the following inequality holds:

$$
f(y) \le f(x) + \langle \nabla f(x), y - x \rangle + \langle \nabla^2 f(x)(y - x), y - x \rangle \varphi(M||y - x||),
$$

where $\varphi(t) \coloneqq \frac{e^t - t - 1}{t^2}$ $\frac{-t-1}{t^2}$.

Lemma C.10. Assume that f is a QSC function with constant M. Then, $\omega(t) \leq \epsilon$ for any given $\epsilon > 0$ whenever $t \leq \delta(\epsilon)$, where

$$
\delta(\epsilon) \coloneqq \min \left\{ \frac{1}{M}, \sqrt{\frac{\epsilon}{2(e-2)\|\nabla^2 f(x^*)\|}}, \frac{\epsilon}{2\|\nabla f(x^*)\|_*} \right\}.
$$

$$
\overline{}
$$

Proof. According to Lemma [C.9,](#page-21-1) for any $x \in Q$, we have

$$
f(x) - f^* \le \langle \nabla f(x^*), x - x^* \rangle + \langle \nabla^2 f(x^*)(x - x^*), x - x^* \rangle \varphi(M || x - x^* ||)
$$

\n
$$
\le ||\nabla f(x^*)||_* ||x - x^*|| + ||\nabla^2 f(x^*)|| ||x - x^*||^2 \varphi(M || x - x^* ||).
$$

Therefore, for any $t \geq 0$, we get

$$
\omega(t) \leq \|\nabla f(x^*)\|_* t + \|\nabla^2 f(x^*)\| t^2 \varphi(Mt),\tag{C.1}
$$

where we have used the fact that $\varphi(\cdot)$ is an increasing function.

Note that, for any $0 \le t \le \frac{1}{M}$, we can estimate $\varphi(Mt) \le \varphi(1) = e - 2$. Substituting this bound into [\(C.1\)](#page-22-1), we obtain

$$
\omega(t) \le \|\nabla f(x^*)\|_* t + (e-2) \|\nabla^2 f(x^*)\| t^2.
$$

To make the right-hand side $\leq \epsilon$, it suffices to ensure that each of the two terms is $\leq \frac{\epsilon}{2}$ $\frac{\epsilon}{2}$:

$$
\|\nabla f(x^*)\|_* t \le \frac{\epsilon}{2}, \qquad (e-2) \|\nabla^2 f(x^*)\| t^2 \le \frac{\epsilon}{2}.
$$

Solving this system of inequalities, we get the claim.

Combining Theorem [2.1](#page-3-0) and Lemma [C.10,](#page-21-2) we get the following complexity result.

Corollary C.11. Consider problem (1.1) under the assumption that f is QSC with con-stant M. Let Algorithm [2.1](#page-2-1) with coefficients (2.4) be applied for solving this problem. Then, $f(x_T^*) - f^* \leq \epsilon$ for any given $\epsilon > 0$ whenever $T \geq T(\epsilon)$, where

$$
T(\epsilon)=\max\left\{M^2,\frac{2(e-2)\|\nabla^2 f(x^*)\|}{\epsilon},\frac{4\|\nabla f(x^*)\|_*^2}{\epsilon^2}\right\}e^2D^2\log^2\frac{e\bar{D}}{\bar{r}},
$$

and the constants D and \bar{D} are as defined in Theorem [2.1.](#page-3-0)

C.6 (L_0, L_1) -Smooth Functions

Let us now consider the case when f is (L_0, L_1) -smooth [\[24\]](#page-12-8), meaning that for any $x \in \mathbb{R}^d$,

$$
\|\nabla^2 f(x)\| \le L_0 + L_1 \|\nabla f(x)\|_*,
$$

where $L_0, L_1 \geq 0$ are fixed constants.

Lemma C.12. [\[23,](#page-12-9) Lemma 2.2] Let f be (L_0, L_1) -smooth. Then, for any $x, y \in \mathbb{R}^d$, it holds that

$$
f(y) \le f(x) + \langle \nabla f(x), y - x \rangle + \frac{L_0 + L_1 \|\nabla f(x)\|_{*}}{L_1^2} \xi(L_1 \|y - x\|),
$$

where $\xi(t) \coloneqq e^t - t - 1$.

Lemma C.13. Assume that f is an (L_0, L_1) -smooth function. Then, $\omega(t) \leq \epsilon$ for any given $\epsilon > 0$ whenever $t \leq \delta(\epsilon)$, where

$$
\delta(\epsilon) \coloneqq \min \left\{ \frac{1}{L_1}, \sqrt{\frac{2\epsilon}{3(L_0 + L_1 \|\nabla f(x^*)\|_*)}}, \frac{\epsilon}{2\|\nabla f(x^*)\|_*} \right\}.
$$

Proof. According to Lemma [C.12,](#page-22-2) for any $x \in \mathbb{R}^d$, we have

$$
f(x) - f^* \le \langle \nabla f(x^*), x - x^* \rangle + \frac{L_0 + L_1 \|\nabla f(x^*)\|_{*}}{L_1^2} \xi(L_1 \|x - x^*\|)
$$

$$
\le \|\nabla f(x^*)\|_{*} \|x - x^*\| + \frac{L_0 + L_1 \|\nabla f(x^*)\|_{*}}{L_1^2} \xi(L_1 \|x - x^*\|)
$$

Therefore, for any $t \geq 0$, we get

$$
\omega(t) \le \|\nabla f(x^*)\|_* t + \frac{L_0 + L_1 \|\nabla f(x^*)\|_*}{L_1^2} \xi(L_1 t),\tag{C.2}
$$

 \Box

where the second inequality uses the fact that $\xi(x)$ is an increasing function.

Note that, for any $0 \leq t \leq \frac{1}{L}$ $\frac{1}{L_1}$, we can estimate

$$
\xi(L_1t)\leq \frac{L_1^2t^2}{2(1-\frac{L_1t}{3})}\leq \frac{3}{4}L_1^2t^2.
$$

Substituting this bound into [\(C.2\)](#page-23-1), we obtain:

$$
\omega(t) \leq \|\nabla f(x^*)\|_* t + \frac{3(L_0 + L_1 \|\nabla f(x^*)\|_*)}{4} t^2.
$$

To make the right-hand side of the last inequality $\leq \epsilon$, it suffices to ensure that each of the two terms is $\leq \frac{\epsilon}{2}$ $\frac{\epsilon}{2}$:

$$
\|\nabla f(x^*)\|_* t \le \frac{\epsilon}{2}, \qquad \frac{3(L_0 + L_1 \|\nabla f(x^*)\|_*)}{4} t^2 \le \frac{\epsilon}{2}.
$$

Solving this system of inequalities, we get the claim.

Combining Theorem [2.1](#page-3-0) and Lemma [C.13,](#page-22-3) we get the following complexity result.

Corollary C.14. Consider problem [\(1.1\)](#page-1-0) under the assumption that f is an (L_0, L_1) smooth function. Let Algorithm [2.1](#page-2-1) with coefficients [\(2.4\)](#page-2-2) be applied for solving this problem. Then, $f(x_T^*) - f^* \leq \epsilon$ for any given $\epsilon > 0$ whenever $T \geq T(\epsilon)$, where

$$
T(\epsilon) = \max \left\{ L_1^2, \frac{3(L_0 + L_1 \|\nabla f(x^*)\|_*)}{2\epsilon}, \frac{4\|\nabla f(x^*)\|_*^2}{\epsilon^2} \right\} e^2 D^2 \log^2 \frac{e\bar{D}}{\bar{r}},
$$

and the constants D and \bar{D} are as defined in Theorem [2.1.](#page-3-0)