ON EPIC GROUPS OVER LANGUAGE CLASS C

RAAD AL KOHLI, COLLIN BLEAK, LUNA ELLIOTT

Abstract

We introduce a new framework linking group theory and formal language theory which generalizes a number of ways these topics have been linked in the past. For a language class C in the Chomsky hierarchy, we say a group is epiC if it admits a language L over a finite (monoidal) generating set $X \subseteq G$ in the class C such that the image of L under the evaluation map is $G \setminus \{1_G\}$. We provide some examples of epiC groups and prove that the property of being epiC is not dependent on the generating set chosen. We also prove that epiC groups are closed under passage to finite index overgroups, taking extensions, and taking graph products of finitely many groups. Furthermore, we prove that epiRegular groups are closed under passage to finite index subgroups. Finally, we provide a characterization of the property of having solvable word problem within the framework of epiC groups.

Contents

1. Introduction	1
1.1. Results	2
2. Languages and groups: some formal definitions	4
3. Independence of generating sets.	5
4. Closure Properties	6
4.1. Finite index overgroups	6
4.2. Extensions	6
4.3. Graph products of groups	7
4.4. Finite index subgroups	11
5. Autostackable groups	12
6. Non-examples	13
6.1. Word problems	13
6.2. Continuum many non-epiRE 2-generated groups	14
7. Questions	19
References	20

1. INTRODUCTION

We denote by \mathcal{C} the following set

$$\mathcal{C} := \{ \text{Regular}, \text{CF}, \text{CS}, \text{RE} \}.$$

MSC (2020): 20F10; 20M35; 20F05; 03D05; 03D40.

Keywords: group theory, automata, language theory, word problem

For the purpose of open access, the authors have applied a CC BY public copyright licence to any accepted manuscript version arising.

If $C \in C$, then C represents a standard formal language class from the Chomsky Hierarchy [8, 9], with CF representing **Context-Free**, CS representing **Context Sensitive**, and RE representing **Recursively Enumerable**.

In this article, for each $C \in C$, we introduce the class of groups epiC. A finitely generated group $G = \langle X \rangle$ is epiC if there is a language $L_G \subseteq X^+$ in class C so that the evaluation map $eval_G : L_C \to G$ has image equaling $G \setminus \{1_G\}$.

The classes of groups epiC nest in the obvious fashion, induced by the nesting of language classes

$Regular \subsetneq CF \subsetneq CS \subsetneq RE.$

By definition, the class epiRegular is a natural broadening of the class of groups with rational cross section, a notion which can be traced back to Eilenberg and Schützenberger from [14] and which was explored in depth by Gilman and also Bodart [16, 4]. As our results show, epiRegular groups also contain the autostackable groups of [5] (and hence, also the asynchronously automatic groups and particular groups of topical interest such as Thompson's group F [11]). Finally, those groups which admit a regular language of geodesics representing all group elements (see [10]) are also epiRegular, since the identity will be represented by the empty string which can be removed. Thus, e.g., extra-large type Artin groups and Garside groups are epiRegular. It follows that each of the classes epiC can be thought of as a broadening of these previously studied classes.

Note as well that it is natural to think of epiRegular as a generalisation of the class of groups with coword problem being a regular language (the finite groups by Anisimov's Theorem [2]), epiCF as a generalisation of the class $coC\mathcal{F}$ of groups (see [23, 26, 3]) and epiCS as a generalisation of Holt and Röver's co-indexed groups [24]. The class of epiRegular groups was initially proposed by James Belk as an interesting class of groups to consider, after some analysis of a construction of Ville Salo appearing in a preprint of [29]. That conversation led to an investigation of epiRegular groups in the first author's dissertation [1], which is the foundation of the work described here.

1.1. **Results.** Here we present our main results.

The first result is that for each language class C, the property of being epiC is a property of the groups themselves, and not of the choices of finite generating sets used to demonstrate membership in the class epiC.

Theorem A. Suppose $C \in \{Regular, CF, CS, RE\}$ and G is a group with finite subsets X and Y where both X and Y generate G as a monoid. If G is epiC with respect to X then G is epiC with respect to Y.

Our next result is that even the smallest class, epiC, is a fairly broad class of groups.

Theorem B. The class of epiRegular groups contains:

- (i) all Baumslag-Solitar groups BS(m, n),
- (ii) all automatic groups, and
- (iii) all autostackable groups.

The first two points above are already known through the theory of groups with rational cross section (see [4, 15]). Guba and Sapir show that Thompson's group F has rational cross section [18], so F is certainly epiRegular.

Meanwhile, Corwin, et al. show F is autostackable in [11]. We prove autostackable groups have rational cross section in Theorem 5.7, and hence they are epiRegular as well. This result about autostackable groups is apparently new. However, we are not highlighting this result here as it is not too surprising in the context of the existing work of Brittenham, et al., and of Bodart (see [4, 5]). Recall the class $coC\mathcal{F}$ of groups with context-free coword problem introduced in [23]. Theorem B implies that the Baumslag-Solitar groups BS(m, n) are in epiCF, but it is shown in [23] that BS(1, 2) is not a $coC\mathcal{F}$ group.

Corollary 2.1. There are groups in the class epiRegular (and thus in epiCF) that are not in the class coCF.

Recall that epiRegular groups contain the class of groups with rational cross section, and [4] states that all Baumslag-Solitar groups and all automatic groups have rational cross sections, as a consequence of groups with rational cross-section being closed under some Bass–Serre Theory constructions. Following a construction in [21] we show that the class of groups with rational cross section is closed under graph products. This closure property for graph products holds more widely, and in particular, for each of the classes epiC.

The following theorem is an amalgamation of Theorems 4.2 and 4.15, below.

Theorem C. Suppose $C \in \{Regular, CF, CS, RE\}$. The class of epiC groups is closed under graph products (of finitely many groups) and extensions.

In counterpoint to the closure of the epiC classes under extensions, the related results for finite index subgroups are not as clear. We ask in Section 7 if the part of Theorem D below relating to finite index subgroups can be extended to the other epiC classes.

Theorem D. The class epiRegular is closed under passage to finite index subgroup. Also, for $C \in \{Regular, CF, CS, RE\}$ the class epiC is closed under passage to finite index overgroups.

It therefore remains a question as to whether the class epiC is closed under passage to finite index subgroups for $C \in \{CF, CS, RE\}$. See Question 7.7.

There is an interplay between having a machine that accepts a representative product string for every non-trivial element of a finitely generated group G and the property of G having a solvable word problem. The following is shown in Subsection 6.1.

Theorem E. Let G be a group with finite generating set X. The following are equivalent:

- G has a solvable word problem; and
- G admits a recursively enumerable presentation $G \cong \langle X|R \rangle$ and is in the class epiRE.

We are unaware of any epiRE group which does not have a recursively enumerable presentation, even though it is plausible that such exists.

As mentioned earlier, we have

$epiRegular \subseteq epiCF \subseteq epiCS \subseteq epiRE.$

At this time, we have not yet discovered a proof that any of these classes are different from each other. We are also unaware of an example group which does not admit rational cross section but which is epiRegular. A number of potential examples provided by [4] include the Grigorchuk groups (as finitely generated infinite torsion groups cannot admit rational cross section) as well as the group $C_2 \wr (C_2 \wr \mathbb{Z})$. These groups all fail to admit rational cross sections but might be epiRegular. Note as well that Grigorchuk's "first" group Γ is co-indexed (see [24]), and hence is in epiCS. It follows that Γ shall either separate epiRegular from groups with rational cross section, or separate epiRegular from epiCS.

In another direction, there are fast algorithms to enumerate words in a regular language, our proof that recursively presented groups in epiRE have solvable word problem reduces, in the case of finitely presented groups in the class epiRegular, to an argument that resolves the word problem for these groups in time complexity Exp(Exp(n)) for any given initial word of length n under the assumption that group elements of G appear in some natural length-based ordering. As there are finitely presented groups in epiRE where the word problem has time complexity worse than any finite tower of exponentials [6, 7] our argument would then prove that $epiRegular \subsetneq epiRE$. However, the definition of epiC for any language class C does not give control on when group elements get represented in the relevant language, so we cannot use this argument to separate epiRegular from epiRE, for instance.

As it is plausible that there are groups in epiRE which do not admit recursively enumerable presentations, it may be the case that there are continuum many isomorphism types of groups in epiRE. In particular, the following theorem (proven in Subsection 6.2) does not immediately follow from Theorem E.

Theorem F. Up to isomorphism, there are continuum many finitely generated groups which are not epiRE.

Acknowledgements The authors wish to thank James Belk, Carl-Fredrik Nyberg Brodda, Gemma Crowe, Alex Levine, and Richard Thomas for various helpful comments and discussions around the contents of this article.

2. Languages and groups: some formal definitions

For $C \in \mathcal{C}$, to describe the class epiC of groups we need a few definitions.

Let X be a set. For positive natural k set X^k to be the set of all k-tuples with coordinates from X. We set

$$X^+ := \bigcup_{k=1}^{\infty} X^k,$$

and

$$X^* := X^+ \cup \{\varepsilon\}.$$

We view a k-tuple (x_1, x_2, \ldots, x_k) as the string $x_1x_2 \cdots x_k$ and ε as the empty string (the empty tuple). Allowing concatenation of strings, X^* becomes **the free monoid generated** by X with identity element ε , as is standard. By the phrase **a formal language over alphabet** X we mean any subset $L \subseteq X^*$. In this article we will only consider formal languages over finite alphabets.

Throughout, we shall never need to explore the grammars of these language classes. However, we will need to recall that languages are defined by computing machines that accept them, which we call **acceptor machines**. We shall write $\mathcal{L}(\mathcal{A}) = L$ to denote that L is the language accepted by an acceptor machine \mathcal{A} . Now recall that a language $\mathcal{L}(\mathcal{A})$ is in the class C if it is accepted by an acceptor machine \mathcal{A} where C is

- **Regular** if \mathcal{A} is a finite state automaton (FSA),
- Context-free if \mathcal{A} is a pushdown automaton (PDA),
- Context Sensitive if \mathcal{A} is a linearly bounded automaton (LBA), or
- Recursively Enumberable if \mathcal{A} is a Turing machine (TM).

For more details regarding these types of acceptor machines or the language classes they accept, we refer the reader to [25].

Definition 2.1 (Word and coword problem). If G is a group and X is a monoid generating set for G, then we define the word problem and coword problem of G with respect to X by

$$WP(G, X) := \{ w \in X^* \mid w =_G 1_G \},\$$

and

$$coWP(G, X) := X^* \setminus WP(G, X)$$

respectively.

q

Remark 2.2. Note that if $w \in X^*$ we will write $w = x_1x_2...x_j$ if w is precisely the string $x_1x_2...x_j$, while we write $w =_G g$ (for $g \in G$) to denote that the product described by the string w evaluates to g in G (here we are assuming that X is a subset of G). We extend this notation to its symmetric transitive closure so we may compare products of elements of G, taken as strings, to determine when such expressions evaluate to the same element of G under the group product.

Note further that we are denoting the identity of G by 1_G , and that we may refer to elements of WP(G, X) as words and elements of coWP(G, X) as cowords when the context seems clear.

We are now ready to formally define the classes epiC of groups which we examine in this article.

For a given language class C, the class of groups epiC is those groups G with a monoid generating set X so that the set of non-identity elements in G is the image of a language $\mathcal{L} \subseteq X^*$ in C under the natural evaluation map. In more detail:

Definition 2.3. Let G be a finitely generated group and $X \subseteq G$ be finite with X generating G as a monoid. Let C be a language class from C and let $L \subseteq X^*$ be a language over the alphabet X which is in the class C.

We say L demonstrates epiC for G if:

- $L \cap WP(G, X) = \emptyset$, and
- for every non-identity element $h \in G$, there exists $w_h \in coWP(G, X)$ such that $w_h \in L$ and $w_h =_G h$.

If \mathcal{A} is an acceptor machine of the appropriate type to accept a language in the class C (i.e., a finite state automaton for Regular, a pushdown automaton for CF, etc.), then we say that \mathcal{A} demonstrates epiC for G if $\mathcal{L}(\mathcal{A})$ demonstrates epiC for G.

The class epiC is the class of all groups H such that H admits a finite monoid generating set Y and a language $\mathcal{K} \subseteq Y^*$ which demonstrates epiC for H.

Remark 2.4. Let G be a group and let C be a language class from C and suppose further that $X \subseteq G$ is a finite set which generates G as a monoid. Extending the previous definition we also say G is epiC with respect to X if there is a language $\mathcal{L} \subseteq X^*$ which demonstrates epiC for G. We may also say G is epiC in this case.

Below are some examples of epiRegular groups. Finite groups represent an extreme case of epiRegular groups, and hence are in epiC for any of our language classes. Here are two arguments (demonstrating the flexibility of the definition of epiRegularity).

Firstly, by Anisimov's theorem [2], there exists a finite state automaton accepting exactly the entirety of the coword problem for any given set of non-trivial generators. One can also use the non-trivial elements of the group as the generators, and thus build an automaton that accepts exactly the length one words over this alphabet.

The group \mathbb{Z} generated by the symmetric generating set $\{a, a^{-1}\}$ is epiRegular with $\{a^n \mid n > 0\} \cup \{(a^{-1})^n \mid n > 0\}$ demonstrating epiRegularity.

3. INDEPENDENCE OF GENERATING SETS.

In this section we show our first claimed theorem from the Introduction. Specifically, that being in a class epiC does not depend on the choice of (finite) generating set.

Theorem A. Suppose $C \in \{Regular, CF, CS, RE\}$ and G is a group with finite subsets X and Y where both X and Y generate G as a monoid. If G is epiC with respect to X then G is epiC with respect to Y.

Proof. It is a standard result (see [30, 25]) that for each language class C amongst Regular, CF, CS, and RE, the class C is closed under monoid homomorphisms which do not map letters to the empty word. Specifically, if X and Y are finite sets, $\mathcal{L} \subseteq X^*$ with \mathcal{L} in the class C, and where $\phi_* : X^* \to Y^*$ is the unique monoid homomorphism extending a function $\phi : X \to Y^+$, then the set $\mathcal{L}\phi_* := \{v \in Y^* \mid \exists u \in \mathcal{L}, v = u\phi_*\}$ is also a language in the class C.

Let C be a chosen language class amongst Regular, CF, CS, or RE. Let G be a non-trivial group with finite monoid generating set X and let $\mathcal{L} \subseteq X^*$ which demonstrates G is epiC with respect to X (note that the trivial group is clearly epiC regardless of the generating set). Finally, let Y be some finite monoid generating set for G.

For each $x \in X$ there is $w_x \in Y^*$ so that $w_x =_G x$ and as G is non-trivial we can further choose $w_x \neq \varepsilon$. Let $\phi : X \to Y^*$ be the function defined by the rules that for all $x \in X$, we have $x \mapsto w_x$. Note that for all $u \in \mathcal{L}$ we have $u\phi_*$ has the property that $u =_G u\phi_*$.

The map ϕ_* now takes \mathcal{L} to a new language $\mathcal{K} := \mathcal{L}\phi_* \subseteq Y^*$ which is still in the language class C and which demonstrates that G is epiC with respect to Y.

Remark 3.1. As seen in the proof above, we only rely on the property of the language classes being closed under monoid homomorphisms that do not map letters to the empty word. Thus for any language class C with that property, a group G is epiC with respect to one finite monoid generating set X if and only if G is epiC with respect to another finite monoid generating set Y.

So one may wish to explore the class of epiC groups for a language class C with the above mentioned property even if C is not in the Chomsky Hierarchy.

4. Closure Properties

In this section we show that each class epiC is closed under finite index overgroups, extensions and graph products. We start with finite index overgroups.

4.1. Finite index overgroups. Our result is as follows. This is one half of Theorem D.

Theorem 4.1. The class epiC is closed under finite index overgroups.

Proof. For any language class C in $\{Regular, CF, RE\}$, suppose the group G is epiC with respect to X (with $\mathcal{L}_{\mathcal{G}} \subseteq X^*$ demonstrating epiC for G) and H is a finite index overgroup of G. The transversal set T together with X generates H. We may assume that $1_H \in T$, so $(T \setminus \{1_H\}) \cup X$ is a finite generating set for H. Observe that any element $h \in H$ can be written as gt for some $g \in G$ and $t \in T$, with $1_H = 1_G 1_H$. Thus any non-trivial element h of H is either a non-trivial element of G, or is in a coset of G with a non-trivial coset representative $t \neq 1_H$. In the latter case, h may be t itself and so can be represented as $1_G t$, otherwise h can be represented as gt for non-trivial g. Thus $\mathcal{L}_{\mathcal{G}} \cup (T \setminus \{1_H\}) \cup \mathcal{L}_{\mathcal{G}}(T \setminus \{1_H\})$ demonstrates epiC for H if C is known to be closed under finite unions and concatenation.

4.2. Extensions. This result forms one half of Theorem C.

Theorem 4.2. Suppose $C \in \{Regular, CF, CS, RE\}$ and let N and Q be epiC. If E is an extension of N by Q (so that N is normal in E) then E is epiC.

Proof. Suppose without meaningful loss of generality that Q = E/N. Let X be a finite monoid generating set for N and Y be a finite subset of E such that $\{Ny : y \in Y\}$ is a monoid generating set for Q (and these elements are distinct). Note that $X \cup Y$ is a generating set for E.

Every element of E can be written as a product nq for some $n \in N$ and $q \in \langle Y \rangle$. Therefore elements of E are either elements of N, elements of $\langle Y \rangle \backslash N$ or a product of both.

Let \mathcal{L}_1 be the language demonstrating N is epiC over the alphabet X. Let \mathcal{L}_2 be the language demonstrating Q is epiC over the alphabet Y (viewing the elements of Y as their image in the quotient). It is known (see for example [25]) that C is closed under concatenation and finite unions. Thus we see that $\mathcal{L}' := \mathcal{L}_1 \cup \mathcal{L}_2 \cup \mathcal{L}_1 \mathcal{L}_2$ is a language demonstrating E is epiC (note that every element represented by $\mathcal{L}_1 \mathcal{L}_2$ has a non-trivial image in Q).

Remark 4.3. The key ingredients in the proofs of Theorems 4.1 & 4.2 are that the language classes are closed under concatenations and finite unions. Thus, this result holds for other language classes as long as those properties are satisfied.

Corollary 4.4. Finitely generated polycyclic groups are epiRegular, and hence so are finitely generated nilpotent groups.

Theorem 4.5. Suppose $C \in \{Regular, CF, CS, RE\}$ and let H be epiC. If G is a finite index overgroup of H then G is epiC.

4.3. Graph products of groups. We now work to show the remaining part of Theorem C: for each language class C, the class of epiC groups is closed under graph products.

We introduce the definition of graph products of groups as in [19].

Definition 4.6. Let $\Gamma = (V(\Gamma), E(\Gamma))$ be a finite simple graph and associate to each vertex $v \in V(\Gamma)$ a group G_v with a fixed group presentation $G_v := \langle X_v | R_v \rangle$, where we further suppose the groups G_v are pairwise disjoint as sets. We define the graph product of groups associated with the structure $(\Gamma, \{\langle X_v | R_v \rangle | v \in V(\Gamma)\})$ as the group G_{Γ} given by the presentation

$\langle X \mid R \rangle$,

where $X = \bigsqcup_{v \in V(\Gamma)} X_v$, and $R \subseteq X^*$ is the set of relations

$$R = \left(\bigsqcup_{v \in V(\Gamma)} R_v\right) \bigsqcup \{[x_v, x_w] \mid x_v \in X_v, x_w \in X_w, \{v, w\} \in E(\Gamma)\}.$$

In what follows, we will only care about graph products of groups associated to structures $(\Gamma, \{\langle X_v \mid R_v \rangle \mid v \in V(\Gamma)\})$ where each set X_v is a finite monoid generating set for the corresponding vertex group G_v . Note in this case that the associated group G_{Γ} is finitely generated as a monoid by X. We will be less explicit in discussing the structure $(\Gamma, \{\langle X_v \mid R_v \rangle \mid v \in V(\Gamma)\})$ below and simply refer to a graph product of groups G_{Γ} associated to a graph Γ .

In particular, the graph product of groups associated to the structure $(\Gamma, \{\langle X_v \mid R_v \rangle \mid v \in V(\Gamma)\})$ is generated by all the groups G_v with the relations being a union of all the relations of the groups G_v together with relations that ensure $[G_v, G_w] =_{G_{\Gamma}} 1_{G_{\Gamma}}$ for any $\{v, w\} \in E(\Gamma)$. Consequently, direct products of finite sets of groups and free products of finite sets of groups are both types of products that can be described as particular graph products of groups.

Before proving Theorem 4.15, we shall define some terminology and state some results from [21] which we shall rely on in our proof. For this purpose, we follow the exposition of [21], and redirect the reader to [21] for more details. We note that the work in [21] is based on [17].

For the remainder of this subsection we shall fix a particular structure

$$(\Gamma, \{ \langle X_v \mid R_v \rangle \mid v \in V(\Gamma) \})$$

and its associated graph product $G_{\Gamma} = \langle X \mid R \rangle$ defined as in Definition 4.6. We shall also assume that there is a total order \preceq on the vertices $V(\Gamma)$ of Γ .

Definition 4.7 (Local string and type). Let $\omega \in X^*$. A non-empty contiguous substring ω' of ω is said to be a **local string** if it is written in letters from the generators of exactly one vertex group, and it is not contained in a longer contiguous substring containing ω' that is written in letters of the generators of exactly one vertex group. The **type of a local string** is the corresponding vertex.

Since any string in X^* is a product of local strings we can define the **type of a string**.

Definition 4.8 (Type of a string). Let $\omega \in X^*$ be a string. The concatenation of the types of the local strings of ω taken in order is the **type** of ω .

For example, suppose $\omega \in X^*$ factors as the concatenation of three non-empty local strings $\omega = \omega_u \omega_v \omega_w$ where ω_u is a local string of type u, ω_v is a local string of type v, and ω_w is a local string of type w, then the type of ω is uvw.

Note also that in Definition 4.8 we are allowing the empty string as a type (it is the type of the empty string from X^*).

Definition 4.9 (Global Length). The global length of a string is the number of local strings it contains.

That is, for $\omega \in X^*$ the global length of ω is the actual length of its type as a string over the alphabet of vertices of Γ .

We may **shuffle** local string ω_u and ω_v in a string if ω_u and ω_v are adjacent in that string, and $\{u, v\} \in E(\Gamma)$. That is, we may use the commuting relations defined by $[G_u, G_v] =_{G_{\Gamma}} 1_{G_{\Gamma}}$ to obtain an equivalent string in the group when the corresponding vertices are adjacent. If in the process of shuffling, two local strings of the same type become adjacent then they amalgamate into a single local string. If it is equal to the identity of the corresponding group, then it is removed.

The process we described above eventually stabilises global length. That is, it will no longer be possible to shuffle local strings of the same type to become adjacent.

We summarize the construction above (corresponding to the first two paragraphs of page 238 of Hermiller and Meier's [21], this is the 9th page of the article).

Lemma 4.10. The process outlined above of taking a string w and shuffling and amalgamating local strings in iterative fashion eventually produces a string z which has a smallest representative type string t with respect to ShortLex ordering on the types, and this type string t is unique.

Definition 4.11. In the statement of Lemma 4.10 we refer to z as a pruning of w or we say we have pruned w to z. We also say that t is the pruned-type of w.

The following is an equivalent formulation of Proposition 3.1 of [21].

Lemma 4.12 ([21],Proposition 3.1). Each group element in G_{Γ} uniquely determines the type of any pruned string representing it.

We note that the identity element is the only element with ε as the type of the pruned string representing it (which by definition is the empty string, as well).

In [21], the authors give a normal form for graph products of groups (assuming one already has a normal form for every vertex group) consisting of **proper** strings. They do this in their Proposition 3.2. We do not have such a normal form in general as our vertex groups need not have normal forms. However, our construction still relies on Hermiller and Meier's property (O), which is a sub-property of their definition of a proper string.

Definition 4.13. A string ω is said to be **proper** if satisfies the following conditions.

- (L) Each local string is in its prescribed normal form.
- (0) If $\omega = \cdots \omega_I \cdots \omega_J \cdots$ with $I \ge J$ and $\{v_I, v_J\} \in E(\Gamma)$ then there is a local string ω_K such that $\omega = \cdots \omega_I \cdots \omega_K \cdots \omega_J \cdots$ where $\{v_K, v_J\} \notin E(\Gamma)$.

The authors of [21] prove in Proposition 3.2 that the set of proper strings is a set of normal forms for G_{Γ} .

In the proof of Theorem B of [21], the authors construct a graph which they call the **admissible graph**. This graph is a finite state automaton $\mathcal{A}_{AG}(\Gamma)$ accepting the proper strings of a graph product of cyclic groups of order 2. Further, since cyclic groups of order 2 have only one non-trivial element, the map that sends a vertex group generator to its type induces a transformation taking any proper string precisely to its type string. Thus as in Theorem B of [21], \mathcal{A}_{AG} determines the types of strings that are allowed. That is, for every type of a proper string in a graph product of groups, the string can be read off by following a path in the admissible graph. Conversely, any path in the admissible graph reads the type of a proper string. In the proof below, we shall replace states of the admissible graph by the automata that demonstrate the vertex groups are epiC. First we shall state a few properties of the admissible graph so that our construction is clear. For the construction of the admissible graph itself, we redirect the reader to [21].

The admissible graph of a graph Γ , $A_{AG}(\Gamma)$, constructed in [21] is a finite state automaton with the following properties:

- (i) There is a unique initial state, this is not a final state.
- (ii) Every non-initial state is labelled by some vertex v_I of the graph Γ . (The construction of the admissible graph does not guarantee a unique non-initial state for every vertex v_I .)
- (iii) Every edge of the admissible graph from a state labelled by $v_J \in V(\Gamma)$ to a state labelled by $v_I \in V(\Gamma)$ has as a label the generator of G_I . (Recall the admissible graph is based on the graph product of cyclic groups of order 2 over Γ , hence these groups are generated by a single element.)
- (iv) Every non-initial state is an accept state.
- (v) There are no ε -transitions in the admissible graph.
- (vi) The language accepted by the admissible graph is the set of strings of types of proper strings of G_{Γ} .

Further, we extract the following statement from [21]. We do not prove this here as this is covered in Proposition 3.2 as well as Theorem B of [21].

Lemma 4.14. Let $g \in G_{\Gamma}$ be a non-trivial element of G_{Γ} . Then there is a string ω in the generators of G_{Γ} such that $\omega =_{G_{\Gamma}} g$ and the type string of ω is the type string of some proper string of the graph product of cyclic groups of order 2 over Γ . Further, every local string of ω is non-trivial in its corresponding vertex group.

We are now ready to prove the following theorem.

In the proof below we envisage changing between machines of the same type, each to do the local calculations over each vertex group: specifically we replace the graph above by a large machine of the appropriate type consisting of smaller machines replacing the vertices of the graph above, and which carry out the calculations local to each of those vertex groups individually. We only allow transitions between local machines when we are currently in an accept state on a local machine and the next input is a generator from a next valid group (validity being determined by the connectivity of the graph Γ). Each valid transition effectively flushes the previous memory structure (either by using a new alphabet and ignoring old alphabet letters or treating them as boundaries in the new local memory calculations (e.g., in the LBA case)), each local calculation starts if it entered the local machine at an initial state and read the one letter of input which triggered the transition to this machine.

Theorem 4.15. Let Γ be a finite graph as above and C a language class in

{Regular, CF, CS, RE}. For every $v_I \in V(\Gamma)$, let G_I be an epiC group with finite monoid generating set X_I . Let G_{Γ} be the graph product of groups $\{G_I \mid v_I \in V(\Gamma)\}$ over the graph Γ . Then G_{Γ} is epiC.

Proof. We will construct a machine (FSA, PDA, LBA, or TM) \mathcal{A} demonstrating that G_{Γ} is epiC.

For every $v_I \in V(\Gamma)$, let \mathcal{A}_I be a machine (FSA, PDA, LBA, or TM) demonstrating that G_I is epiC.

(We shall call these \mathcal{A}_I local machines.) We will construct \mathcal{A} by "gluing" the \mathcal{A}_I onto every vertex of the admissible graph $\mathcal{A}_{AG}(\Gamma)$ that is labelled by v_I as follows.

For all states p and q of $\mathcal{A}_{AG}(\Gamma)$ that are labeled by v_I and v_J , respectively with an edge from p to q in $\mathcal{A}_{AG}(\Gamma)$ we do the following:

Remove that edge, and insert an ε transition from each final state of \mathcal{A}_I to the new state t_q which is a transitory state for entering the machine \mathcal{A}_J as if no processing has occurred. How the state t_q functions depends on the type of machine we're working with, there are four cases:

- (i) FSA: The initial state of \mathcal{A}_J shall be identified with t_q (we assume by the powerset construction that \mathcal{A}_J has a unique initial state) and processing will carry on as normal for this machine.
- (ii) PDA: Each transition to t_q from the local machine \mathcal{A}_I shall be an ε -transition that ignores the stack. The state t_q will have loop transitions that are ε -transitions which delete any letters on the current stack one letter at a time, until the bottom-of-stack symbol \perp of \mathcal{A} appears. We shall use this same bottom-of-stack symbol for each of the local machines. With the stack cleared we now can employ any ε -transition to any initial state of the local machine \mathcal{A}_J (where we have one such transition for each of it's initial states). From this point, processing continues as normal for that local machine.
- (iii) LBA: For a reference to the descriptions of LBA's which we use, the reader can consult Section 10.5 of [28]. Unlike the three other machine types, the machine A for LBA's shall have a global accept state which is different from the local accept states which appear in the local machines. The essential idea for LBA's is that calculations in an LBA take place on a tape with delineated left and right boundary symbols ("[" and "]" for discussion) which bracket the input word. The LBA may write on this tape between the boundaries as part of its calculation. In our context, as we are calculating a sequence of accepted (local) words, when the machine sees a letter from a generator of a different local group, that letter shall be perceived as

the right relevant boundary. Left boundaries will be updated with a new "[" symbol replacing whatever symbol appears at the final position for allowed calculations in the previous local machine. When a local machine achieves an accept state, it shall read an ε -transition loop to this state which has the impact of moving the read head to the right until it is over the right boundary, which is either a letter from the generating set of the next group, or it is the right boundary symbol "]". In the latter case, there is a transition to the global accept state for \mathcal{A} . In the other case, there is an ε -transition to the appropriate t_q state for the next local machine. From the state t_q initial processing shall be through a collection of ε -transitions supporting the following. First the read head should move left one space and write a "[" over whatever was there so that the new processing for the machine \mathcal{A}_J has a good left boundary. Then it shall transition to one of the initial states of \mathcal{A}_J and processing begins as normal.

On processing of the final local word, the "]" appears as the final boundary, so if an accept state is achieved for the local machine, we can follow with an ε -transition to an accept state for the global machine \mathcal{A} .

(iv) TM: Move to any initial state of \mathcal{A}_I .

The initial state of this new automaton \mathcal{A} will be the initial state of $\mathcal{A}_{AG}(\Gamma)$. The final states consist of all the final states of every copy of \mathcal{A}_I that we have glued onto the admissible graph by the above process. (If there is an edge from a sub-automaton to another, then it is from a final state of one sub-automaton to the initial state of the other.)

We note that any path in \mathcal{A} projects onto a path in $\mathcal{A}_{AG}(\Gamma)$ by identifying all the edges in a sub-automaton \mathcal{A}_I , and edges between a final state of a sub-automaton and the initial state of another will be edges between the vertices of $\mathcal{A}_{AG}(\Gamma)$ that the sub-automata were glued onto. Therefore, any path in \mathcal{A} must yield the same type string as a path in $\mathcal{A}_{AG}(\Gamma)$.

Since the initial state is not an accept state and the identity element of G_{Γ} has type ε , by Lemma 4.12 the identity element of G_{Γ} is not represented by any accepted string of the above automaton. By Lemma 4.14, we know that every non-trivial element of G_{Γ} can be expressed as a string whose type is the type of a proper string of the graph product of cyclic groups of order 2 over Γ , with every local string ω_I being non-trivial in G_I . For each such local string, there is a string ω'_I in the language accepted by \mathcal{A}_I such that $\omega_I =_{G_I} \omega'_I$. Thus the automaton \mathcal{A} simply replaces the types of proper strings of the graph product of cyclic groups with these local strings ω'_I while preserving the type string. Therefore there is a string representing every non-trivial element of G_{Γ} in $\mathcal{L}(\mathcal{A})$, and there is no string in $\mathcal{L}(\mathcal{A})$ representing the identity element of G_{Γ} . Therefore G_{Γ} is epiC.

This completes the proof of Theorem C. We note that it follows from Theorem 4.15 that the class of groups epiC is closed under direct and free products.

4.4. Finite index subgroups. Epiregular is the only of our classes which we know to be closed under finite index subgroups, and hence we focus on that class here (see Question 7.7). We restate Lemma 2.4 of [27] with t = 1 below as this will be used in the proof of Theorem 4.17.

Lemma 4.16. Let G be a group with finite generating set Σ , and H be finite index in G with generating set Δ . If $R \subseteq G$ is rational over Σ , and $R \subseteq H$, then R is rational over Δ .

While Theorem 4.1 is one half of Theorem D, the result below is the other half.

Theorem 4.17. If G is epiregular and $H \leq G$ has finite index, then H is epiregular.

Proof. Let X_G and X_H be finite generating sets for G and H respectively.

By Poincaré lemma we can assume that H is normal as the class epiregular is closed under extensions.

Let $L \subseteq X_G^*$ be a language demonstrating that G is epiregular. Let $L_H \subseteq X_G^*$ consisting of all words which evaluate to an element of H. So an element $g \in G$ is represented by an element of $L_H \cap L$ if and only if $g \in H \setminus \{1_G\}$.

Moreover the language L_H is regular as the cayley graph of G/H with start/termianl state H is a FSA demonstrating this.

Thus $L_H \cap L$ is regular as well.

Finally Lemma 4.16 implies there is a new regular language L' over X_H with the same evaluations as $L_H \cap L$ as required.

5. Autostackable groups

In this section, we shall refer to generating sets as being inverse closed as this is standard in the literature regarding autostackable groups. Recall that a generating set X is inverse closed if for all $x \in X$ we have $x^{-1} \in X$. In particular, an inverse closed generating set generates its group as a monoid as well.

Definition 5.1. For a finitely generated group G and a finite inverse closed generating set X, we define $\Gamma(G, X)$ to be the associated Cayley graph. That is the directed graph with vertex set G, and directed edge set \vec{E} ; where for each $g \in G$ and $a \in X$, there is a directed edge $e_{g,a}$ with initial vertex g, terminal vertex ga, and label a.

Definition 5.2. Let G be a finitely generated group with a finite inverse closed generating set X. We say that $N \subseteq X^*$ is a set of normal forms for G if for all $g \in G$ there is a unique string which we denote by $y_g \in N$ such that $y_g =_G g$.

Definition 5.3. Let G be a finitely generated group with a finite inverse closed generating set X. Let N be a set of normal forms for G over X. Whenever we have an equality of strings $y_{ga} = y_{ga}$ or $y_g = y_{ga}a^{-1}$, we call the edge $e_{g,a}$ of $\Gamma(G, X)$ degenerate.

Let $\vec{E_{N,d}} = \vec{E_d}$ be the set of all degenerate directed edges of $\Gamma(G, X)$, and let $\vec{E}_{N,r} = \vec{E_r} := \vec{E} \setminus \vec{E_d}$; we refer to elements of $\vec{E_r}$ as **recursive** edges.

Definition 5.4 (Padding). Let X be an alphabet and assume # is a symbol not in X. Set $\widehat{X} := X \cup \{\#\}$ and call the set \widehat{X} the **padded alphabet associated with** X and call # the **padding symbol (for** X).

Let $u, v, w \in X^*$, where $u = a_1 a_2 \cdots a_r$, $v = b_1 b_2 \cdots b_s$, and $w = c_1 c_2 \cdots c_t$ with each $a_i, b_i, c_i \in X$. Set k := max(r, s, t) and $\overline{X} := \widehat{X} \times \widehat{X} \times \widehat{X} \setminus \{(\#, \#, \#)\}$. We call \overline{X} the trinary extension of X. We define $(u, v, w)^t \in \overline{X}^*$ to be the string

$$(\alpha_1, \beta_1, \gamma_1)(\alpha_2, \beta_2, \gamma_2) \cdots (\alpha_k, \beta_k, \gamma_k)$$

where

(i) $\alpha_i = a_i \text{ for } 1 \leq i \leq r \text{ and } \alpha_i = \# \text{ for } r < i \leq k;$

(ii) $\beta_i = b_i$ for $1 \leq i \leq s$ and $\beta_i = \#$ for $s < i \leq k$;

(iii) $\gamma_i = c_i \text{ for } 1 \leq i \leq t \text{ and } \gamma_i = \# \text{ for } t < i \leq k.$

We call $(u, v, w)^t$ the trinary projection corresponding to (u, v, w).

Ą

For example, for the strings hi, bye, hello the trinary projection corresponding to (hi, bye, hello) is

(h, b, h)(i, y, e)(#, e, l)(#, #, l)(#, #, o).

Definition 5.5 (Synchronously Regular). Let X be a set and let \overline{X} be the trinary extension of X. A subset $L \subseteq (X^*)^3$ is called **synchronously regular** if the set L^t of trinary projections of L forms a regular language over the alphabet \overline{X} .

Definition 5.6 (Autostackable). Let G be a finitely generated group, and let X be a finite generating set for G which is inverse closed. We say G is **autostackable** if there is a set N of normal forms for G over X containing the empty word ε , a constant k, and a function $\phi: N \times X \to X^*$ such that the following hold.

(i) The graph of the function ϕ

$$graph(\phi) := \{(y_q, x, \phi(y_q, x)) \mid q \in G, x \in X\}$$

is a synchronously regular language.

- (ii) For each $g \in G$ and $x \in X$, the word $\phi(y_g, x)$ has length at most k and represents the element x of G, and:
 - (2d) If $e_{g,x} \in \vec{E}_{N,d}$, then the equality of words $\phi(y_g, x) = x$ holds.
 - (2r) The transitive closure $<_{\phi}$ of the relation < on $\vec{E}_{N,r}$, defined by $e' < e_{g,x}$ whenever $e_{g,x}, e' \in \vec{E}_{N,r}$ and e' is on the directed path $\phi(y_g, x)$ starting at the vertex g, is a strict well-founded partial ordering.

Theorem 5.7. If G is an autostackable group with inverse closed generating set X then G has rational cross-section.

Proof. Let G, X, N and ϕ be as in Definition 5.6, and \overline{X} be the trinary extension of X (as in Definition 5.4). Denote by g_{ϕ} the set $graph(\phi)$.

By definition, as g_{ϕ} is a synchronously regular language we have that $(g_{\phi})^t$ is a regular language over the alphabet \overline{X} . The semigroup homomorphism which extends the projection map to the first coordinate from \overline{X} takes the regular language $(g_{\phi})^t$ to a regular language in $(X \cup \{\#\})^*$. We may further compose by the semigroup homomorphism from $(X \cup \{\#\})^*$ to X^* which replaces all appearances of # by the empty string ε to find a regular language in X^* . This is precisely the set N of normal forms for G. Thus showing that G has rational cross-section.

This completes the proof of Theorem **B**.

6. Non-examples

In this section we shall demonstrate two theorems. The first theorem is that there is an equivalence for finitely generated groups between having solvable word problem, and having a recursively enumerable presentation and being in the class epiRE. It follows (see Corollary 6.1) that there are finitely presented groups that are not epiRE. The second theorem is that there are continuum many pairwise distinct isomorphism classes of 2-generated groups which are solvable with derived length three, and which are not epiRE.

6.1. Word problems. For a set X (and formal set of "inverses" X^{-1} for elements of X) and $R \subseteq (X \cup X^{-1})^*$ we say a group G is presented by $\langle X | R \rangle$ if the normal closure of the images of R in F_X is a subgroup of F_X with quotient isomorphic to G, and we write in this case $G \cong \langle X | R \rangle$.

In this section, we shall recall that Turing machines can be used in other ways apart from accepting languages, such as enumerating and outputting sets.

Translating Higman [22] for the case of a finitely generated group, we say a group G with finite generating set X has a **recursively enumerable presentation** $G \cong \langle X | R \rangle$ if there

is a Turing machine which will output the family R of strings over $(X \cup X^{-1})^*$. We say G has a **computable presentation** $G \cong \langle X | R \rangle$ if there is a Turing machine which determines in finite time whether or not a given string in $(X \cup X^{-1})^*$ is in the set R.

We shall now prove the final theorem mentioned in the Introduction.

Theorem E. Let G be a group with finite generating set X. The following are equivalent:

- G has a solvable word problem; and
- G admits a recursively enumerable presentation $G \cong \langle X|R \rangle$ and is in the class epiRE.

Proof. (\Rightarrow) :

One can check for each string in X^* whether or not it is a coword, and accept just the cowords. Outputting the set of words shows that G admits a recursively enumerable presentation.

 (\Leftarrow) : Let $f : \mathbb{N} \to (X \cup X^{-1})^*$ be some recursive enumeration of a language demonstrating that G is epiRE.

Let T be the Turing machine which enumerates the language R.

From T we can build another Turing machine N which will output W, the set of all the words over $(X \cup X^{-1})^*$ which evaluate as elements of the normal closure.

Let $g: \mathbb{N} \to W$ be the corresponding enumeration of the output of N.

The algorithm for checking if a string $w \in (X \cup X^{-1})^*$ belongs to W is as follows:

- (i) Loop over all elements i of \mathbb{N}
 - (a) Check if (i)g = w, if yes then w is in the word problem.
 - (b) Check all $j, k \leq i$ so see if $(j)fw^{-1} = (k)g$. If the answer is ever yes then we declare that w is not in the word problem.

Corollary 6.1. There are finitely presented groups which are not epiRE.

Corollary 6.2. For $C \in \{Regular, CF, CS, RE\}$, every finitely presented epiC group has a solvable word problem.

6.2. Continuum many non-epiRE 2-generated groups. One might think that as there are continuum many two-generated groups with unsolvable word problem that it should be immediately clear that there are continuum many non-epiRE groups with unsolvable word problem. However, Theorem E only applies for groups with recursive presentations, and of course, there are only countably many such groups up to isomorphism. Sadly, we also do not have an idea of how to specify an epiRE group which does not admit a recursively enumerable presentation.

In this subsection we closely follow a general construction due to Phillip Hall (see [20]), but as given in de la Harpe's [12]. The discussion can be found in discussion points 40-43 Section III.C. We give more details than appear in that source, and along the way we also correct a minor error in the text (in [12], the author asserts that a set of group elements freely generate a free abelian group when they do not, but, that is not important for the overall construction). In doing so we give a slightly modified generating set and show freeness. We will use aspects of that given construction in our proof of Theorem F.

We aim to show that the cardinality of the set of isomorphism classes of 2-generated groups is of size continuum. We will actually aim for more than this statement, with an eye towards applications to epiRE groups.

The outline of what is to come is as follows:

- (i) demonstrate that the free group $F_{\{a,b\}}$ freely generated by $\{a,b\}$, has continuum many normal subgroups; and in particular we can find a quotient which is solvable of length 3 and whose center contains $\bigoplus_{i \in \mathbb{N}} \mathbb{Z}$.
- (ii) demonstrate that taking quotients by these normal subgroups will generate continuum many pairwise non-isomorphic groups; and moreover we can choose these normal subgroups to be in a chain.
- (iii) We then show that any two comparable normal subgroups of $F_{\{a,b\}}$ give rise to quotients which cannot be shown to be epiRE by a common recursively enumerable language.

Lemma 6.3 ([12], III.C., 40). The set of normal subgroups of $F_{\{a,b\}}$ is of size continuum.

Proof. We follow de la Harpe's offered proof, inserting some commentary along the way. Set $S = \{s_i \mid i \in \mathbb{Z}\}$ as a set of generators indexed by \mathbb{Z} . Define R a set of relators as

$$R := \{ [[s_i, s_j], s_k] \mid i, j, k \in \mathbb{Z} \} \cup \{ [s_i, s_j] [s_{j+k}, s_{i+k}] \mid i, j, k \in \mathbb{Z} \}.$$

We will initially be concerned with the group $\Gamma_0 := \langle S \mid R \rangle$. Set $u_j = [s_0, s_j] \in \Gamma_0$ for $j \in \mathbb{Z} \setminus \{0\}$, and $U := \{u_i \mid i \in \mathbb{Z} \setminus \{0\}\} \subseteq \Gamma_0$.

We see immediately from the relations of the form $[[s_i, s_j], s_k]$ that each element of U lies in the centre of Γ_0 . Secondly, the relators of the form $[s_i, s_j][s_{j+k}, s_{i+k}]$ force $[s_i, s_j] = [s_{i+k}, s_{j+k}]$ for all $i, j, k \in \mathbb{Z}$, so each commutator of the form $[s_i, s_j]$ is an element of U.

In particular, for positive natural i we have $[s_{-i}, s_0] = [s_0, s_i]$ from which it follows that $[s_0, s_{-i}] = [s_0, s_i]^{-1}$.

Claim (a): The set U is a generating set for the commutator subgroup $[\Gamma_0, \Gamma_0]$ of Γ_0 , which is contained in the center of Γ_0 .

Proof of claim (a):

For notational purposes in calculations, for $i \in \mathbb{Z} \setminus \{0\}$, we will use *i* to represent s_i and \overline{i} to represent s_i^{-1} . For a word $w \in (S \cup S^{-1})^*$, we use \overline{w} for the inverse word so that $w\overline{w} = 1_{\Gamma_0}$.

The first observation we make below is that if we have a commutator c of two words each of length one, then this can be re-expressed as a commutator of positive generators from S, so such a commutator is in the centre of Γ_0 and in fact there is a positive integer j so that $c = u_j^{\pm 1}$.

To verify this, see the following two calculations using our notational convention mentioned earlier (remembering that U is contained in the center of Γ_0).

$$[\overline{i}, j] = i\overline{j}\,\overline{i}\,j = i\overline{j}\,\overline{i}\,ji\overline{i} = i[j, i]\overline{i} = [j, i]i\overline{i} = [j, i]$$

$$[\overline{i},\overline{j}] = ij\overline{i}\,\overline{j} = ij\overline{i}\,\overline{j}\,ij\overline{j}\,\overline{i} = ij[i,j]\overline{j}\,\overline{i} = [i,j]$$

One can now use standard commutator identities to express any commutator of two words from $(S \cup S^{-1})^*$ as products of conjugates of commutators involving subwords of the two words, where each commutator is shorter in length than the original. To that end, let $x, y, z \in (S \cup S^{-1})^*$ be non-trivial. We have the standard commutator equivalences

$$[x,zy] = [x,y] \cdot [x,z]^y$$

and

$$[zy, x] = [z, x]^y \cdot [y, x].$$

(where we use the convention $a^b = b^{-1}ab$ for conjugation).

Using these and our process of removing inverse generators from short commutators, we may inductively break any commutator in Γ_0 down into products of conjugates of commutators of pairs of generators. By our relations we may ignore conjugation of commutators of pairs of generators, thus it follows that the set U generates the commutator subgroup of Γ_0 , which is also in the center of Γ_0 (as U is contained in the center of Γ_0).

This completes the proof of Claim (a).

Claim (b): The commutator subgroup $[\Gamma_0, \Gamma_0]$ is free abelian¹ with free generating set $U_+ := \{u_i \mid i > 0\}.$

 \diamond

Proof of claim (b):

We have already shown that this group is abelian and generated by U_+ . We show that there are no more relations by constructing an explicit group homomorphism from Γ_0 to a group such that the elements in the image of U_+ satisfy no more relations. Consider the set

$$(\oplus_{i\in\mathbb{Z}}\mathbb{Z})\times(\oplus_{i\in\mathbb{Z}_{>0}}\mathbb{Z})$$

(where \oplus means take all tuples with only finitely many non-zero coordinates) with operation

$$((a_i)_{i \in \mathbb{Z}}, (b_i)_{i > 0}) \cdot ((c_i)_{i \in \mathbb{Z}}, (d_i)_{i > 0}) = \left((a_i + c_i)_{i \in \mathbb{Z}}, \left(b_i + d_i + \sum_{\substack{k,l \\ l - k = i}} a_l c_k \right)_{i > 0} \right).$$

This operation is associative as

$$\left(\left((a_{i,0})_{i\in\mathbb{Z}},(b_{i,0})_{i>0}\right)\cdot\left((a_{i,1})_{i\in\mathbb{Z}},(b_{i,1})_{i>0}\right)\right)\cdot\left((a_{i,2})_{i\in\mathbb{Z}},(b_{i,2})_{i>0}\right)$$

$$= \left((a_{i,0} + a_{i,1})_{i \in \mathbb{Z}}, \left(b_{i,0} + b_{i,1} + \sum_{\substack{k,l \\ l-k=i}} a_{l,0} a_{k,1} \right)_{i>0} \right) \cdot ((a_{i,2})_{i \in \mathbb{Z}}, (b_{i,2})_{i>0})$$

$$= \left((a_{i,0} + a_{i,1} + a_{i,2})_{i \in \mathbb{Z}}, \left(b_{i,0} + b_{i,1} + b_{i,2} + \sum_{\substack{k,l \\ l-k=i}} a_{l,0} a_{k,1} + \sum_{\substack{k,l \\ l-k=i}} (a_{l,0} + a_{l,1}) a_{k,2} \right)_{i>0} \right)$$

$$= \left((a_{i,0} + a_{i,1} + a_{i,2})_{i \in \mathbb{Z}}, \left(b_{i,0} + b_{i,1} + b_{i,2} + \sum_{\substack{k,l \\ l-k=i}} a_{l,c} a_{k,d} \right)_{i>0} \right).$$

Thus we have a semigroup which we will denote by Γ'_0 . It then follows immediately from the definition that Γ'_0 is a group with the inverse operation given by:

$$((a_i)_{i \in \mathbb{Z}}, (b_i)_{i>0})^{-1} = \left((-a_i)_{i \in \mathbb{Z}}, \left(\sum_{\substack{k,l \\ l-k=i}} a_l a_k - b_i \right)_{i>0} \right).$$

¹It is claimed in point (a) of the proof of the lemma in [12], that the commutator subgroup $[\Gamma_0, \Gamma_0] < \Gamma_0$ is the free abelian group on the set $\{u_i \mid i \in \mathbb{Z}, i \neq 0\}$, but this is false because of the identities $[s_0, s_{-i}] = [s_0, s_i]^{-1}$.

It follows from von Dyck's Theorem that the map ϕ from S to Γ'_0 defined by $(s_i)\phi = ((a_j)_{j\in\mathbb{Z}}, (0)_{i>0})$ where $a_j = 0$ for $j \neq i$ and $a_i = 1$ extends to a group homomorphism over Γ_0 if and only if the required relations R are satisfied. Note that for all $i, j \in \mathbb{Z}$ with j > i we have

$$((s_i)\phi)^{-1}((s_j)\phi)^{-1}(s_i)\phi(s_j)\phi = ((s_i)\phi)^{-1}((s_j)\phi)^{-1}((\dots 00100\dots 00100\dots), (00\dots)) i^{th} \text{ position}$$

$$= ((s_i)\phi)^{-1}((\dots 00100\dots), (00\dots 00(-1)00\dots)) i^{th} \text{ position}$$

$$= ((\dots 00\dots), (00\dots 00(-1)00\dots)),$$

$$((j-i)^{th} \text{ position})$$

and for j < i we have

$$((s_{i})\phi)^{-1}((s_{j})\phi)^{-1}(s_{i})\phi(s_{j})\phi = ((s_{i})\phi)^{-1}((s_{j})\phi)^{-1}((\ldots 00100 \ldots 00100 \ldots)), (00 \ldots 001000 \ldots)))$$

$$\stackrel{i^{th} \text{ position}}{= ((s_{i})\phi)^{-1}((\ldots 00100 \ldots), (00 \ldots 001000 \ldots)))}$$

$$\stackrel{i^{th} \text{ position}}{= ((\ldots 00 \ldots), (00 \ldots 001000 \ldots)).}$$

$$\stackrel{i^{th} \text{ position}}{= ((\ldots 00 \ldots), (00 \ldots 001000 \ldots)).}$$

In both of the above cases the only non-zero entry of $[(s_i)\phi, (s_j)\phi]$ is in the |i-j| coordinate of the second coordinate and its value is determined by which is bigger. Thus the required relations defined by the set R hold. Moreover, this map sends the set U to a free basis for the free abelian group $\{(0)_{i\in\mathbb{Z}}\}\times \bigoplus_{i>0}\mathbb{Z} \leq \Gamma'_0$, hence U does indeed generate a free abelian group.

This completes the proof of Claim (b).

Now returning to the proof in [12], let $\phi : \Gamma_0 \to \Gamma_0$ be the automorphism which maps s_i to s_{i+1} for all indices *i*. Because of the second part of the relations *R*, the automorphism ϕ fixes the set *U* pointwise, and so, fixes the commutator subgroup [Γ_0, Γ_0] pointwise. Also observe that the action of $\langle \phi \rangle$ on the set *S* is transitive.

We define a group

$$\Gamma := \langle S \cup \{t\} \mid R \cup \{s_i^t s_{i+1}^{-1} \mid i \in \mathbb{Z}\} \rangle.$$

It follows from the construction that $\Gamma \cong \Gamma_0 \rtimes_{\phi} \mathbb{Z}$, with conjugation by t acting as the automorphism ϕ , and t generating the \mathbb{Z} factor in the quotient of Γ by Γ_0 . We consider in the background the homomorphism $\psi : F_{\{a,b\}} \to \Gamma$ determined by $a \mapsto s_0$ and $b \mapsto t$. Distinct normal subgroups of Γ are then the images of distinct normal subgroups of $F_{\{a,b\}}$, so we reduce the problem to finding continuum many normal subgroups of Γ .²

 \diamond

²From the first claim Γ_0 is nilpotent of class 2. Further, it is immediate that Γ is solvable with derived length three, but not nilpotent as $[s_i, t] = s_i^{-1}s_{i+1}, [s_i^{-1}s_{i+1}, t] = s_{i+1}^{-1}s_is_{i+1}^{-1}s_{i+2}$ and inductively, if we keep commutating the result of the previous calculation by t, the rightmost term of the resulting product will be a sole occurrence of s_{i+k} (where we have carried out k commutations).

We now complete the proof of the lemma. Let X be any subset of $\mathbb{N}\setminus\{0\}$,³ and let N_X denote the subgroup of $[\Gamma_0, \Gamma_0]$ generated by $U_X := \{u_x \mid x \in X\}$. As $[\Gamma_0, \Gamma_0]$ is central in Γ , it follows that each N_X is normal, and that Γ therefore admits continuum many normal subgroups.

There is now an observation in [12] that Lemma 6.3 in effect shows $F_{\{a,b\}}$ admits continuum many normal subgroups containing $D^3F_{\{a,b\}}$ (the third derived subgroup of $F_{\{a,b\}}$). This is a result of Hall from 1954 (see [20]) where he actually shows that there is a group G satisfying $[D^2G, G] = 1$ so that there are continuum many normal subgroups of a group G. Hall in that paper also shows that the number of normal subgroups of $F_{\{a,b\}}$ containing $D^2F_{\{a,b\}}$ is countable.

A version of the lemma in Point 42 of [12] is as follows:

Lemma 6.4 ([12], III.C., 42). For a finitely-generated group Γ , TFAE:

- (i) Γ has continuum many normal subgroups; and
- (ii) Γ has continuum many pairwise non-isomorphic quotients.

Proof. That (ii) implies (i) is immediate just by considering the kernels of the quotient maps. Thus we need only show that (i) implies (ii).

Let us now assume Γ is generated by k generators for some natural k, has continuum many normal subgroups, and only countably many quotients up to isomorphism. In this case there must be a countable group Q and continuum many distinct quotient maps from Γ to Q. However, this is impossible since any homomorphism from Γ to Q is determined by where the generators of Γ are sent, and there are only countably many sequences of length k in Q.

We have seen from Lemma 6.3 and Lemma 6.4 that the free group $F_{\{a,b\}}$ admits continuum many non-isomorphic quotients. In fact, the group Γ , which is a solvable quotient of $F_{\{a,b\}}$ with derived length three, has continuum many non-isomorphic quotients. We are now in a position to prove the final theorem mentioned in the Introduction: specifically, we argue that Γ has quotients which are not epiRE (and in fact we distinguish continuum many such isomorphism classes).

Theorem F. Up to isomorphism, there are continuum many finitely generated groups which are not epiRE.

Proof. The first point to observe is that the set of subsets of $\mathbb{N}\setminus\{0\}$ admits chains of size continuum under the containment relation. This can be seen for example by noting that the set of downwards closed subsets of (\mathbb{Q}, \leq) is of size continuum and totally ordered by $\subseteq (\mathbb{Q}$ has the same cardinality as $\mathbb{N}\setminus\{0\}$).

Our main result now follows easily from our construction. For every subset $X \subseteq \mathbb{N} \setminus \{0\}$, let N_X denote the normal subgroup of Γ generated by the set $\{u_i \mid i \in X\}$. For each normal subgroup N_X , the quotient group $Q_X = \Gamma/N_X$ is determined, and for each Q_X let us re-use $Y := \{s, s^{-1}, t, t^{-1}\}$ as symbols for the induced generating set of Q_X (that is, e.g., $s \in Q_X$ is formally the coset sN_X of N_X in Γ , and the four cosets induced from Y under the natural quotient map form a generating set for Q_X).

Suppose for a contradiction that only countably many of the quotients Q_X are not epiRE up to isomorphism. By the same argument given in Lemma 6.4, there are only countably many sets X (up to equality) for which Q_X is not epiRE. From this point on we only consider

³[12] uses $\mathbb{Z}\setminus\{0\}$ here, not recognizing e.g. that $u_{-3} = u_3^{-1}$.

the subsets $X \subseteq \mathbb{N} \setminus \{0\}$ which are contained in some fixed continuum sized chain of subsets of $\mathbb{N} \setminus \{0\}$. We may also choose this chain such that all of the corresponding groups Q_X are epiRE (by removing the countably many problematic sets).

As there are (up to relabelling) only countably many Turing machines which use the alphabet Y, we must have that one of these machines must determine a language \mathcal{L}_Y which demonstrates that two distinct groups Q_X and $Q_{X'}$ arising in our construction are epiRE and we may assume $X' \subseteq X$.

As the sets X and X' are distinct, there is some u_i a generator of N_X which is not in the group $N_{X'}$. As \mathcal{L}_Y demonstrates that $Q_{X'}$ is epiRE, there is a word $w \in \mathcal{L}_Y$ representing the non-trivial image of u_i in $Q_{X'}$. As \mathcal{L}_Y demonstrates that Q_X is epiRE and $w \in \mathcal{L}_Y$, the word w represents a non-trivial element of Q_X . But as w represents $u_i N_{X'}$ in $Q_{X'}$ and $u_i N_{X'} \subseteq u_i N_X$, w represents the element $u_i N_X$ of Q_X . This is a contradiction as $u_i N_X = N_X$ is the identity of Q_X .

7. Questions

In this section we will explore some questions which are natural given the explorations above.

In Theorem E, we explored the relation between the solvability of the word problem and epiC groups. This leads to the following natural question.

Question 7.1. For any language class C (regular, context-free, context sensitive, recursively enumerable) are there epiC groups which do not admit recursively enumerable presentations?

Having introduced the hierarchy of epiC groups, we wish to understand the relationship between these classes of groups and whether the classes can be separated in a way that is analogous to the Chomsky hierarchy. More precisely we have the following question.

Question 7.2. Let C_1 and C_2 be distinct language classes in the Chomsky hierarchy such that C_1 is a subclass of C_2 . Does there exist a group in $epiC_2$ but not $epiC_1$?

If the answer to the previous question is affirmative, then we are also interested in the following related question.

Question 7.3. Let C_1 and C_2 be distinct language classes such that C_1 is a subclass of C_2 . Does there exist a reasonable group theoretic property \mathcal{P} such that a group $G \in epiC_1$ is equivalent to G has \mathcal{P} and $G \in epiC_2$?

It is known that finitely generated infinite torsion groups do not admit rational cross sections [16, 13], so there is a fair chance that these groups are not epiRegular. However, Grigorchuk's first group Γ is known to be co-indexed [24], and so $\Gamma \in$ epiCS. Thus, there is a chance that Grigorchuk's group can be used to separate the classes epiRegular and epiCS. Specifically, these considerations lead us to ask the following questions.

Question 7.4. Does there exist an epiRegular group which does not admit a rational cross section?

We would be happy even with stronger versions of the above question, for instance if we assume the group is recursively presented or finitely presented.

Question 7.5. Is Grigorchuk's first group Γ in epiRegular?

If Grigorchuk's group is epiRegular then, being epiRegular is not equivalent to admitting a rational cross section. On the other hand, if Grigorchuk's group is not epiRegular then, being epiRegular is not equivalent to being epiCS. Further, it is shown in [4] that $C_2 \wr (C_2 \wr \mathbb{Z})$ does not have rational cross-section. We currently do not know whether or not $C_2 \wr (C_2 \wr \mathbb{Z})$ is epiRegular. Understanding this may also prove useful in answering Question 7.4.

Question 7.6. Is $C_2 \wr (C_2 \wr \mathbb{Z})$, the standard restricted wreath product of C_2 with the Lamplighter group, an epiRegular group?

Finally, while we have understood that for any language class $C \in \{Regular, CF, CS, RE\}$ the class epiC is closed under passage to finite index overgroups, and is also closed under extensions, we have not been able to verify that epiC is closed under passage to finite index subgroup, except for the class epiRegular.

Question 7.7. For $C \in \{CF, CS, RE\}$ is the class epiC closed under passage to finite index subgroups?

References

- R. Al Kohli, Groups defined by language theoretic classes, Ph.D. thesis, University of St Andrews, St Andrews, Scotland, 2024.
- [2] A. V. Anīsīmov, The group languages, Kibernetika (Kiev) (1971), no. 4, 18–24. MR 301981
- [3] Collin Bleak, Francesco Matucci, and Max Neunhöffer, Embeddings into Thompson's group V and coCF groups, J. Lond. Math. Soc. (2) 94 (2016), no. 2, 583–597. MR 3556455
- [4] C. Bodart, Rational cross-sections, bounded generation, and orders on groups, Journal of the London Mathematical Society 109 (2024), no. 6, e12920.
- M. Brittenham, S. Hermiller, and D. Holt, Algorithms and topology of Cayley graphs for groups, J. Algebra 415 (2014), 112–136. MR 3229510
- [6] F. B. Cannonito, Hierarchies of computable groups and the word problem, J. Symbolic Logic 31 (1966), 376–392.
- [7] F. B. Cannonito and R. W. Gatterdam, The word problem and power problem in 1-relator groups are primitive recursive, Pacific J. Math. 61 (1975), no. 2, 351–359.
- [8] N. Chomsky, Three models for the description of language, IRE Transactions on Information Theory 3 (1956), no. 2, 113–124.
- [9] N. Chomsky and M. P. Schützenberger, The algebraic theory of context-free languages, Computer programming and formal systems, North-Holland, Amsterdam, 1963, pp. 118–161. MR 152391
- [10] L. Ciobanu, S. Hermiller, D. Holt, and S. Rees, Conjugacy languages in groups, Israel J. Math. 211 (2016), no. 1, 311–347. MR 3474966
- [11] N. Corwin, G. Golan, S. Hermiller, A. Johnson, and Z. Šunić, Autostackability of Thompson's group F, J. Algebra 545 (2020), 111–134. MR 4044692
- [12] P. de la Harpe, Topics in geometric group theory, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 2000. MR 1786869
- [13] P. de La Harpe and E. Ghys, Sur les groupes hyperboliques d'après mikhael gromov, Birkhäuser, 1990.
- [14] S. Eilenberg and M. P. Schützenberger, Rational sets in commutative monoids, J. Algebra 13 (1969), 173–191. MR 246985
- [15] D. B. A. Epstein, J. W. Cannon, D. F. Holt, S. V. F. Levy, M. S. Paterson, and W. P. Thurston, Word processing in groups, Jones and Bartlett Publishers, Boston, MA, 1992. MR 1161694
- [16] R. H. Gilman, Groups with a rational cross-section, Combinatorial group theory and topology (Alta, Utah, 1984), Ann. of Math. Stud., vol. 111, Princeton Univ. Press, Princeton, NJ, 1987, pp. 175– 183. MR 895616
- [17] E. R. Green, Graph products of groups, Ph.D. thesis, University of Leeds, 1990.
- [18] V. S. Guba, Polynomial upper bounds for the Dehn function of R. Thompson's group F, J. Group Theory 1 (1998), no. 2, 203–211. MR MR1614265 (99j:20037)
- [19] V. S. Guba and M. V. Sapir, Diagram groups are totally orderable, J. Pure Appl. Algebra 205 (2006), no. 1, 48–73. MR 2193191
- [20] P. Hall, Finiteness conditions for soluble groups, Proc. London Math. Soc. (3) 4 (1954), 419–436. MR 72873
- [21] S. Hermiller and J. Meier, Algorithms and geometry for graph products of groups, J. Algebra 171 (1995), no. 1, 230–257. MR 1314099

- [22] G. Higman, Subgroups of finitely presented groups, Proc. Roy. Soc. London Ser. A 262 (1961), 455–475. MR 130286
- [23] D. F. Holt, S. Rees, C. E. Röver, and R. M. Thomas, Groups with context-free co-word problem, J. London Math. Soc. (2) 71 (2005), no. 3, 643–657. MR 2132375 (2006a:20065)
- [24] D. F. Holt and C. E. Röver, Groups with indexed co-word problem, Internat. J. Algebra Comput. 16 (2006), no. 5, 985–1014. MR 2274726
- [25] J. E. Hopcroft and J. D. Ullman, Introduction to automata theory, languages, and computation, Addison-Wesley Series in Computer Science, Addison-Wesley Publishing Co., Reading, Mass., 1979. MR 645539
- [26] Joerg Lehnert and Pascal Schweitzer, Context-Freeness of Higman-Thompson group's co-word problem, preprint.
- [27] Alex Levine, EDT0L solutions to equations in group extensions, J. Algebra 619 (2023), 860–899. MR 4534696
- [28] P. Linz and S. H. Rodger, An introduction to formal languages and automata, Jones & Bartlett Learning, 2022.
- [29] V. Salo, Graph and wreath products in topological full groups of full shifts, 2021.
- [30] M. Sipser, Introduction to the theory of computation, Course Technology Cengage Learning, 2012.

Raad Al Kohli, rsasak@st-andrews.ac.uk School of Mathematics & Statistics, University of St Andrews, St Andrews, UK

Collin Bleak, collin.bleak@st-andrews.ac.uk School of Mathematics & Statistics, University of St Andrews, St Andrews, UK

Luna Elliott, luna.elliott@manchester.ac.uk

Department of Mathematics, The University of Manchester, Manchester, UK