Pauli principle forbids $\Omega_{QQQ}\Omega_{QQQ}\Omega_{QQQ}$ bound states

H. Garcilazo^{1,*} and A. Valcarce^{2,[†](#page-0-1)}

 1 Escuela Superior de Física y Matemáticas, Instituto Politécnico Nacional, Edificio 9, 07738 Mexico D.F., Mexico 2 Departamento de Física Fundamental,

Universidad de Salamanca, E-37008 Salamanca, Spain

(Dated: January 20, 2025)

Abstract

Lattice QCD studies have shown the attractive character of the ¹S₀ $\Omega_{QQQ}\Omega_{QQQ}$, $Q = s, c, b$, interaction, predicting deeply bound states as the mass of the heavy quark increases. This has led to the question of the possible existence of bound states of more than two Ω_{QQQ} baryons, in particular $\Omega_{QQQ}\Omega_{QQQ}\Omega_{QQQ}$ bound states. We discuss how these states might not exist in nature in any flavor sector. The reason would be due to the simultaneous action of two consequences of the Pauli principle in the different partial waves: one at the baryon level, affecting the ${}^{1}S_{0}$ partial wave, and the other due to the quark substructure, affecting the ${}^{5}S_{2}$ partial wave.

[∗]Electronic address: hgarcilazos@ipn.mx

[†]Electronic address: valcarce@usal.es

I. INTRODUCTION.

Experimental findings since 2003 with the discovery of the $X(3872)$ |1| have opened a new era for hadron spectroscopy. As a general conclusion, the idea has emerged that the heavy hadron spectra shows the contribution of states that do not belong to the simplest quark-antiquark (meson) or three-quark (baryon) structures proposed by Gell-Mann [\[2\]](#page-13-1). Experimental evidence of more complex structures is steadily and continuously being found [\[3\]](#page-13-2).

One of the most striking predictions of the quark model was the possible existence of bound states in double heavy tetraquark configurations, $QQ\bar{q}\bar{q}$ [\[4\]](#page-13-3). Stability was shown to be a function of the mass ratio $M_Q/m_q.$ The mechanism that stabilizes $QQ\bar q\bar q$ at large M_Q/m_q is the same that makes the hydrogen molecule much more stable than the positronium one in atomic physics [\[5](#page-13-4)]. Such prediction was confirmed by the LHCb Collaboration that announced the discovery of a very sharp peak in the $DD\pi$ spectrum that was dubbed T_{cc} [\[6](#page-13-5), [7\]](#page-13-6). It corresponds to a minimal quark content $cc\bar{u}\bar{d} = T_{cc}^+$. The T_{cc}^+ misses binding by a very small amount. So, it is almost certain that the heavier partners, $T_{bc} = bc\bar{u}\bar{d}$ and $T_{bb} = bb\bar{u}\bar{d}$, could be stable with respect to the strong and electromagnetic interactions [\[6,](#page-13-5) [8\]](#page-13-7).

Another area that has recently attracted a great deal of interest is the case of dibaryons in the heavy flavor sectors. Only one stable state made of two baryons is known, the deuteron. This is a loosely bound state of a proton and a neutron with spin 1. Lattice QCD studies have opened the door to the existence of stable states made of heavy flavor baryons. Thus, the HAL QCD Collaboration [\[9](#page-13-8)] has studied the $\Omega\Omega$ system in the 1S_0 channel through $(2 + 1)$ −flavor lattice QCD simulations with a large volume and nearly physical pion mass $m_{\pi} \simeq 146$ MeV. Their results indicate that the ${}^{1}S_{0}$ $\Omega\Omega$ state has an overall attraction and is located near to the unitary regime with a binding energy $B_{\Omega\Omega}^{\text{QCD}} = 1.6(6) \left(\substack{+0.7 \\ -0.6} \right)$ MeV. The Coulomb repulsion reduces the binding energy by a factor of two, $B_{\Omega\Omega}^{\text{QCD+Coulomb}} = 0.75(5)(5)$ MeV, the system being still bound.

A few years later, the $\Omega_{ccc} \Omega_{ccc}$ system was studied on the basis of the HAL QCD method [\[10](#page-13-9)]. Without the Coulomb interaction, a $(2 + 1)$ −flavor lattice QCD study of the ${}^{1}S_{0}$ channel with nearly physical light-quark masses and the relativistic heavy-quark action with the physical charm quark mass, shows a weak repulsion at short distances surrounded by a relatively strong attractive well, which leads to a bound state with a binding energy $B_{\Omega_{ccc}\Omega_{ccc}}^{\text{QCD}} = 5.68(0.77) \left(^{+0.46}_{-1.02}\right)$ MeV. Taking into account the Coulomb repulsion be-

tween the Ω_{ccc}^{++} 's, with their charge form factor obtained from lattice QCD, a scattering length of $a_0^C = -19(7) \begin{pmatrix} +7 \\ -6 \end{pmatrix}$ fm was obtained. The ratio $r_{\text{eff}}^C/a_0^C = -0.024$ is considerably smaller than that of the dineutron (-0.149) , which indicates that $\Omega_{ccc}\Omega_{ccc}$ is located in the unitary regime.

More recently, the first lattice QCD investigation of heavy dibaryons $\Omega_{bbb}\Omega_{bbb}$ in the ${}^{1}S_0$ channel has been carried out [\[11](#page-13-10)]. It was reported a binding energy $B_{\Omega_{\rm tot}}^{\rm QCD}$ $\frac{\text{QCD}}{\Omega_{bbb}\Omega_{bbb}} = 81 \left(\frac{+16}{-14} \right)$ MeV. With such a deep binding, Coulomb repulsion serves only as a perturbation on the ground state wave function of the parametrized strong potential and may shift the strong binding only by a few percent. In particular, the associated maximum change in binding energy is found to be between 5 and 10 MeV.

On the other hand, it is also well known from nuclear physics that when a two-baryon interaction is attractive, if such a system is merged with additional nuclear matter and there are no severe Pauli principle constraints, the attraction can be strengthened. Simple examples can be given of the effect of a third or a fourth baryon in two-baryon systems with attractive character interactions. The deuteron, $(I)J^P = (0)1^+$, is bound by 2.225 MeV, while the triton, $(I)J^P = (1/2)I/2^+$, is bound by 8.480 MeV, and the α particle, $(I)J^P =$ $(0)0^+$, is bound by 28.295 MeV. The binding per nucleon B/A increases as 1 : 3 : 7. A similar argument is found in systems with strangeness -1 . The hypertriton ${}_{\Lambda}^{3}H$, $(I)J^{P} = (0)1/2^{+}$, is bound with a separation energy of 130 ± 50 keV $[12]^1$, and the $^{4}_{\Lambda}H$, $(I)J^P = (0)0^+$, is bound with a separation energy of 2.12 ± 0.01 (stat) ± 0.09 (syst) MeV [\[14\]](#page-13-12).

The template that has been observed for tetraquarks, that the binding energy increases for heavy quarks [\[4](#page-13-3), [8\]](#page-13-7), it therefore appears to be reproduced in the case of other multiquark hadrons involving more than one bottom quark [\[11](#page-13-10)]. Considering all the above together, a common interesting pattern might be emerging that the presence of more than one bottom quark enhances the binding in multihadron systems [\[15,](#page-13-13) [16\]](#page-13-14). Given the bound-state nature of two– Ω systems in the different heavy flavor sectors predicted by lattice QCD calculations $[9-11]$ $[9-11]$, the question immediately arises whether systems containing more than two such Ω like baryons might also be bound in nature [\[17](#page-13-15), [18\]](#page-13-16). This issue is particularly relevant in the bottom sector given the very large binding energy predicted for the two-body system

¹ Note that the above canonical value of the hypertriton binding energy has been challenged recently by the STAR Collaboration [\[13\]](#page-13-17), claiming a much more tightly bound hypertriton with $B = 0.41 \pm 0.12$ (stat) \pm 0.11 (syst) MeV, which does not affect our reasoning.

by lattice QCD calculations [\[11\]](#page-13-10). This is the question we analyze in this work making use of the quark substructure of the Ω^i baryons ², similar to what was done in the case of tetraquarks [\[4\]](#page-13-3) and other two-baryon systems [\[19,](#page-13-18) [20](#page-14-0)].

II. Ω $^i\Omega^i\Omega^i$ STATES.

Due to the Fermi-Dirac statistic the $\Omega^i \Omega^i$ system can only exist in two different $L = 0$ partial waves: ${}^{1}S_{0}$ and ${}^{5}S_{2}$. Moreover, there is only one fully antisymmetric $\Omega^{i}\Omega^{i}\Omega^{i}$ state with three Ω^{i} 's in relative S wave, the $J^{P} = 3/2^{+}$. Both two-body partial waves, $^{1}S_{0}$ and ⁵ S_2 , are basic to construct the fully antisymmetric $J^P = 3/2^+ \Omega^i \Omega^i \Omega^i$ state.

Let's first study the three-body problem considering only the contribution of the attractive interaction in the ${}^{1}S_{0} \Omega^{i} \Omega^{i}$ channels, which are known in detail from the lattice QCD calculations [\[9](#page-13-8)[–11\]](#page-13-10). The objective of this first analysis will be to understand if the attractive two-body interactions, especially in the $\Omega^b \Omega^b$ case, are reflected in an attractive character of the three-body system or if, on the contrary, there are Pauli effects that mask such attraction as it occurs in other three-body systems [\[21](#page-14-1)].

We have solved the $\Omega^i \Omega^i \Omega^i$ three-body problem by the Faddeev method as described in detail, for example, in Ref. [\[19\]](#page-13-18). We have used as input the ${}^{1}S_{0} \Omega^{i} \Omega^{i}$ potentials parametrized in Refs. [\[9](#page-13-8), [10\]](#page-13-9) for $i = s$ and c. In the case of the $i = b$ potential shown in Fig. 4 of Ref. [\[11\]](#page-13-10) we have parametrized the interaction by means of three Gaussians as done in Refs. [\[9,](#page-13-8) [10\]](#page-13-9). In spite of the attractive character of the ${}^{1}S_{0}$ two-body interaction, we have not found any $J^P = 3/2^+$ three-body bound state in any flavor sector considering only the ${}^1S_0 \Omega^i \Omega^i$ channels.

It is important to realize that this result is a direct consequence of the Pauli principle at the baryon level. It is due to the fact that the recoupling coefficient between the two spin-0 Faddeev amplitudes is a negative number, see Table [I,](#page-4-0) so that it effectively changes the nature of the two-body interaction from attractive to repulsive such that no bound state can be obtained in a one-channel calculation [\[21](#page-14-1)]. In fact, increasing the attraction in the ${}^{1}S_{0}$ two-body channel results in an increase of the repulsion in the three-body system. This result applies for all systems with three identical fermions, in particular for the three-neutron

² Since our arguments will be valid for any of the flavor sectors considered, henceforth we will denote by Ω^i the Ω for $i = s$, the Ω_{ccc} for $i = c$ and the Ω_{bbb} for $i = b$.

		$ s_{12}, s_3; S\rangle$		
		$s_{12}=0$	$s_{12}=2$	
	$s_{23}=0$			
$\big s_1,s_{23};S\rangle$	$s_{23}=2$			

TABLE I: Recoupling coefficients between the different Jacobi coordinate systems of the $\Omega^i \Omega^i \Omega^i$ $J^P = 3/2^+$ state.

case. However, unlike the case of neutrons that have spin $1/2$, the Ω^{i} 's are spin $3/2$ fermions and therefore, as explained above, for $L = 0$, in addition to the spin-0 two-body amplitudes also contribute spin-2 two-body amplitudes.

If we choose an arbitrary three-body Jacobi coordinate system, $|s_{12}, s_3; S\rangle$, the fully antisymmetric three-body $J^P = 3/2^+$ wave function has two different components, $s_{12} = 0$ and $s_{12} = 2$, that has to be expressed in the different Jacobi couplings, $|s_1, s_{23}; S\rangle$. The recoupling coefficients between the different Jacobi coordinate systems are given in Table [I.](#page-4-0) This table illustrates on the one hand the relevance of the 5S_2 partial wave to the $\Omega^i\Omega^i\Omega^i$ $J^P = 3/2⁺$ state, much higher than that of the attractive ${}^{1}S_{0}$ partial wave, and on the other hand the strong coupling between the ¹S₀ and the ⁵S₂ partial waves in the $J^P = 3/2^+$ threebody state. Moreover, the recoupling coefficient between the spin-2 amplitudes is positive and therefore free of the Pauli principle effect at the baryon level discussed above, that effectively changes the nature of the spin-0 two-body interaction from attractive to repulsive in the three-body system.

Therefore, a definitive conclusion about the existence of bound states in the only $\Omega^i \Omega^i \Omega^i$ S-wave state, $J^P = 3/2^+$, cannot be reached without a full fledged calculation considering the contribution of both two-body partial waves: ${}^{1}S_{0}$ and ${}^{5}S_{2}$.

III. ${}^5S_2 \Omega^i \Omega^i$ STATE.

We have explained in the introduction that lattice QCD simulations have derived the ${}^{1}S_{0}$ interaction for the different $\Omega^i \Omega^i$ systems, but so far the interaction in the 5S_2 channel has not been derived. A naive ansatz of spin independence of the $\Omega^i \Omega^i$ interaction [\[18\]](#page-13-16), as it could be deduced from a Fermi-Breit like potential at baryonic level, leads to bound states of hundreds of MeV for the $i = b$ case using the ¹S₀ lattice QCD interaction of Ref. [\[11\]](#page-13-10). This would be a surprising result with binding energies even higher than expected for the beautiful partners of the T_{cc}^+ tetraquark $[15, 16, 22]$ $[15, 16, 22]$ $[15, 16, 22]$ even more so considering that if the quark substructure is considered and the six-particle system is treated rigorously no bound states are found for very heavy flavor dibaryons [\[23\]](#page-14-3) nor for fully-heavy tetraquarks [\[24](#page-14-4)]. Thus, as it has been done for multiquark states [\[4](#page-13-3), [22](#page-14-2)[–24](#page-14-4)] and other two-baryon systems [\[19,](#page-13-18) [20](#page-14-0), [25\]](#page-14-5), one can use the quark substructure of the Ω^i baryons to gain some clues about the most significant features of the $\Omega^i \Omega^i$ interaction in the ⁵ S_2 partial wave.

Consequently, it is important to keep in mind that we are working with identical baryons, Ω^i , that are formed by identical quarks, which is not always the case for identical hadrons. Therefore, the antisymmetry of the two-baryon wave function to the exchange of identical quarks between the two baryons must be also guaranteed in the overlap region of the two baryons. If there were any consequences due to the identity of the baryon constituents, it would be reflected in the normalization of the wave function that appears in the calculation of any observable, in particular the potential as we will discuss in more detail in the next section. Thus, we describe the baryon-baryon system by means of a constituent quark cluster model, i.e., baryons are described as clusters of three constituent quarks. Assuming a twocenter shell model the wave function of an arbitrary baryon-baryon system can be written as [\[20](#page-14-0)]:

$$
\Psi_{B^{\alpha}B^{\beta}}^{LSI}(\vec{R}) = \frac{\mathcal{A}}{\sqrt{1+\delta_{B^{\alpha}B^{\beta}}}}\sqrt{\frac{1}{2}}\left\{ \left[\Phi_{B^{\alpha}}\left(123; -\frac{\vec{R}}{2}\right) \Phi_{B^{\beta}}\left(456; \frac{\vec{R}}{2}\right) \right]_{LSI} + (-1)^{f} \left[\Phi_{B^{\beta}}\left(123; -\frac{\vec{R}}{2}\right) \Phi_{B^{\alpha}}\left(456; \frac{\vec{R}}{2}\right) \right]_{LSI} \right\},
$$
\n(1)

where A is the antisymmetrization operator accounting for the possible existence of identical quarks inside the hadrons. The symmetry factor f satisfies $L+S_1+S_2-S+I_1+I_2-I+f$ = odd. For non-identical baryons f indicates the symmetry associated to a given set of values LSI. The non-possible symmetries correspond to forbidden states. For identical baryons, $B^{\alpha} = B^{\beta}$, f has to be even in order to have a non-vanishing wave function, recovering the well-known selection rule $L + S + I = \text{odd}$. In the case we are interested in, two baryons made of identical quarks with $I = 0$, the antisymmetrization operator comes given by,

$$
\mathcal{A} = \left(1 - \sum_{i=1}^{3} \sum_{j=4}^{6} P_{ij}^{LSC}\right) (1 - \mathcal{P}), \qquad (2)
$$

FIG. 1: Normalization diagrams of the two-baryon wave function.

where P_{ij}^{LSC} exchanges a pair of identical quarks i and j in the configuration, spin and color spaces, and $\mathcal P$ exchanges identical baryons. Assuming a Gaussian form for the wave functions of the quarks inside the hadrons, the normalization of the two–baryon wave function $\Psi_{B\alpha}^{LS(I=0)}(\vec{R}) \equiv \Psi_{B\alpha}^{LS}(\vec{R})$ of Eq. [\(1\)](#page-5-0) can be expressed as [\[20,](#page-14-0) [25,](#page-14-5) [26\]](#page-14-6),

$$
\mathcal{N}_{B^{\alpha}B^{\beta}}^{LS}(R) = \left\langle \Psi_{B^{\alpha}B^{\beta}}^{LS}(\vec{R}) \mid \Psi_{B^{\alpha}B^{\beta}}^{LS}(\vec{R}) \right\rangle = N_{\text{di}}^{L}(R) - C(S) N_{\text{ex}}^{L}(R),\tag{3}
$$

where $N_{\text{di}}^{L}(R)$ and $N_{\text{ex}}^{L}(R)$ are the direct and exchange radial normalization contributions shown in Fig. [1,](#page-6-0) whose explicit expressions are,

$$
N_{\text{di}}^{L}(R) = 4\pi \exp\left(-\frac{3}{4} \frac{R^{2}}{\alpha^{2}}\right) i_{L+1/2} \left(\frac{3}{4} \frac{R^{2}}{\alpha^{2}}\right),
$$

\n
$$
N_{\text{ex}}^{L}(R) = 4\pi \exp\left(-\frac{3}{4} \frac{R^{2}}{\alpha^{2}}\right) i_{L+1/2} \left(\frac{1}{4} \frac{R^{2}}{\alpha^{2}}\right),
$$
\n(4)

where α is the Gaussian parameter of the quark wave function. In the limit where the two baryons overlap $(R \to 0)$, the Pauli principle may impose antisymmetry requirements, due to the existence of identical quarks in the interacting baryons, that were not present in a hadronic description. The second diagram in Fig. [1](#page-6-0) takes into account the exchange of identical quarks between the two baryons and would not exist if the quark substructure of the baryons had not been considered, the normalization in this case being a mere multiplicative factor. These effects would be prominent in S-waves, when the two baryons can approach without centrifugal barrier constraints. Using the asymptotic form of the Bessel functions, $i_{L+1/2}$, in the overlapping region the normalization kernel of Eq. [\(3\)](#page-6-1) can be expressed as,

$$
\mathcal{N}_{\Omega^{i}\Omega^{i}}^{LS}(R) \longrightarrow_{R \to 0} 4\pi \left[1 - \frac{3R^{2}}{4\alpha^{2}} \right] \frac{1}{1 \cdot 3 \cdots (2L+1)} \left[\frac{R^{2}}{4\alpha^{2}} \right]^{L} \times \left\{ \left[3^{L} - 3 C(S) \right] + \frac{\left[3^{L+2} - 3 C(S) \right]}{2(2L+3)} \left[\frac{R^{2}}{4\alpha^{2}} \right]^{2} + \cdots \right\}.
$$
\n(5)

 $C(S)$ is a flavor-independent spin coefficient given by,

$$
C(S) = \langle \Omega^{i}(123), \Omega^{i}(456); S \, \big| \, P_{36}^{S} \, \big| \Omega^{i}(123), \Omega^{i}(456); S \rangle \; . \tag{6}
$$

The spin coefficients for the different $\Omega^i \Omega^i$ states are given in Table [II.](#page-7-0)

∼ $\tilde{}$		

TABLE II: $C(S)$ spin coefficients for the $\Omega^i \Omega^i$ states.

It can be seen from Eq. [\(5\)](#page-7-1) how in S-waves, the closer $C(S)$ is to $1/3$ the greater the suppression of the normalization of the two-baryon wave function at short distances, generating Pauli repulsion due to the presence of a quasi-forbidden state [\[26\]](#page-14-6). For S-waves, if $C(S) = \frac{1}{3}$ the normalization of the two-baryon wave function behaves as $R⁴$ instead of being a constant, indicating that Pauli blocking occurs or, in other words, a node appears in the relative wave function due to a forbidden state [\[27](#page-14-7)]. The repulsion due to the presence of a quasiforbidden state is much stronger than in other channels where the antisymmetrization alone produces the repulsion [\[26](#page-14-6)]. This is because the existence of a forbidden state is a saturation phenomenon that produces a repulsive hard-core independent of the dynamics [\[28,](#page-14-8) [29\]](#page-14-9).

There are examples in the literature about the existence of quasi-forbibben and forbidden states. Thus there are, for example, quasi-forbidden channels in the ΣN [\[30\]](#page-14-10) and $\Sigma_c N$ [\[31\]](#page-14-11) interactions with $(S, I) = (0, 1/2)$ and $(S, I) = (1, 3/2)$, where $C(S) = 8/27$ and $7/27$, respectively, giving rise to strong repulsive potentials. Forbidden states, Pauli blocking, appear, for example, in the $(S, I) = (1, 1) \Delta N$ interaction [\[28\]](#page-14-8). It is worth noting that the study of πd elastic scattering results in a configuration space potential representing the ${}^{3}S_{1}(I = 1)$ ΔN phase shifts which presents a strong repulsive core of large radius, greater than 1 fm [\[32\]](#page-14-12). Such repulsion reaches to the limit of the wave function, much further than in the case of quasi-forbidden states. There is also experimental evidence about such a repulsive ³S₁(I = 1) ΔN potential from other processes, such as the π-nucleus inelastic scattering [\[33\]](#page-14-13). The same is true for partial waves of the $\Delta\Delta$ interaction, see, for example, Fig. 5 of Ref. [\[25\]](#page-14-5). The effect of Pauli blocking is quite clear, it generates strongly repulsive phase shifts that leave no room for the existence of bound states, see, for example, Fig. 3 of Ref. [\[29\]](#page-14-9).

IV. $J^P = 3/2^+ \Omega^i \Omega^i \Omega^i$ STATE.

We can derive the ⁵ S_2 $\Omega^i \Omega^i$ interaction from a basic interaction between the quarks making use of the Born-Oppenheimer (BO) approximation. The BO method, also known as adiabatic approximation, has been frequently employed for the study of the nuclear force in terms of the microscopic degrees of freedom [\[20,](#page-14-0) [34,](#page-14-14) [35](#page-14-15)]. It is based on the assumption that quarks move inside the clusters much faster than the clusters themselves. Then, one can integrate out the fast degrees of freedom assuming a fixed position for the center of each cluster, obtaining in this way a local potential depending on the distance between the centers of mass of the clusters. Explicitly, the potential is defined as follows,

$$
V_{B^{\alpha}B^{\beta}(LS)\to B^{\alpha}B^{\beta}(L'S')}(R) = \xi_{LS}^{L'S'}(R) - \xi_{LS}^{L'S'}(\infty),\tag{7}
$$

where

$$
\xi_{LS}^{L'S'}(R) = \frac{\left\langle \Psi_{B^{\alpha}B^{\beta}}^{L'S'}(\vec{R}) \mid \sum_{i < j = 1}^{6} V_{q_i q_j}(\vec{r}_{ij}) \mid \Psi_{B^{\alpha}B^{\beta}}^{L'S'}(\vec{R}) \right\rangle}{\sqrt{\left\langle \Psi_{B^{\alpha}B^{\beta}}^{L'S'}(\vec{R}) \mid \Psi_{B^{\alpha}B^{\beta}}^{L'S'}(\vec{R}) \right\rangle} \sqrt{\left\langle \Psi_{B^{\alpha}B^{\beta}}^{L'S'}(\vec{R}) \mid \Psi_{B^{\alpha}B^{\beta}}^{L'S'}(\vec{R}) \right\rangle}}.
$$
\n
$$
(8)
$$

In the last expression the quark coordinates are integrated out keeping R fixed, the resulting interaction being a function of the two-baryon relative distance.

We derive the interaction making use of the chiral constituent quark model of Ref. [\[20\]](#page-14-0). This model was proposed in the early 90's in an attempt to obtain a simultaneous description of the nucleon-nucleon interaction and the light baryon spectra. It was later on generalized to all flavor sectors [\[36](#page-14-16), [37](#page-14-17)]. In this model hadrons are described as clusters of three interacting massive (constituent) quarks. The masses of the quarks are generated by the dynamical breaking of the original $SU(2)_L \otimes SU(2)_R$ chiral symmetry of the QCD Lagrangian at a momentum scale of the order of $\Lambda_{\text{CSB}} = 4\pi f_\pi \sim 1$ GeV, where f_π is the pion electroweak decay constant. For momenta typically below that scale, when using the linear realization of chiral symmetry, light quarks interact through potentials generated by the exchange of Goldstone bosons. Notice that for the particular case of heavy quarks (c or b) chiral symmetry is explicitly broken and therefore boson exchanges associated to the dynamical breaking of chiral symmetry do not contribute. Perturbative QCD effects are taken into account through the one-gluon-exchange (OGE) potential [\[38\]](#page-14-18). Finally, any model imitating QCD should incorporate confinement. Although it is a very important term from the spectroscopic point of view, it is negligible for the hadron−hadron interaction. Lattice QCD calculations suggest a screening effect on the potential when increasing the interquark distance [\[39\]](#page-14-19). We refer the reader to Refs. [\[20,](#page-14-0) [36\]](#page-14-16) for a detailed description of the model.

Fig. 3 of Ref. [\[20](#page-14-0)] shows the different quark diagrams contributing to the baryon-baryon interaction. The kernel of the different diagrams, in particular the Coulomb-like term of the OGE, is different from zero at short distances and independent of the mass of the interacting quarks. According to Eq. [\(8\)](#page-8-0), the kernels of the different diagrams must be divided by the normalization of the two-baryon wave function in Eq. [\(3\)](#page-6-1). Inserting the coefficients from Table [II](#page-7-0) into the normalization of the two-baryon wave function in Eq. [\(5\)](#page-7-1), the 5S_2 Ω ^{*i*} Ω ^{*i*} partial wave satisfies the condition $C(S) = \frac{1}{3}$, which translates into the suppression in the normalization of the two-baryon wave function at short-range not expected at baryonic level. ³ Consequently, a strong repulsive hard core, independent of the dynamics, is expected in the 5S_2 Ω ^{*i*} Ω ^{*i*} interaction for any flavor due to the underlying quark substructure [\[28](#page-14-8), [29\]](#page-14-9). It is important to highlight that the ¹ $S_0 \Omega^i \Omega^i$ partial-wave does not satisfy the S–wave Pauli blocking condition $C(S) = \frac{1}{3}$, see Table [II,](#page-7-0) so that the short-range normalization behaves as a constant and does not induce any repulsion beyond that obtained from the dynamics considered. This result is in complete agreement with the conclusions deduced from lattice QCD [\[9](#page-13-8)[–11](#page-13-10)].

The 5S_2 Ω ^{*i*} Ω ^{*i*} repulsive interaction is shown in Fig. [2](#page-10-0) using the parameters and the interacting potential of Ref. [\[31\]](#page-14-11). Since the dominant term is the Coulomb-like contribution of the OGE potential, which is flavor independent, the range of the repulsive core depends on the Gaussian parameter of the quark wave function, α . We show the potential for three different values of α , representing the typical size of the different Ω^i baryons [\[40](#page-14-20)]. As can

³ Let us also note the presence of the symmetry repulsive hard-core for $L \neq 0$ in the ${}^{7}P_{2,3,4}$ Ω ^{*i*} Ω ^{*i*} partial wave, what is also a feature of the $\Delta\Delta$ interaction in the ${}^{7}P_{2,3,4}(I=3)$ partial wave [\[25\]](#page-14-5). However, in these cases, the effects of Pauli blocking may be masked by the centrifugal barrier.

FIG. 2: ${}^{5}S_{2}$ $\Omega^{i}\Omega^{i}$ interaction for different values of the Gaussian parameter of the quark wave function, α .

be seen the potential is repulsive, the repulsive core going to the limit of the relative wave function. This result for the ${}^5S_2 \Omega^i \Omega^i$ interaction was to some extent obvious and expected, because it is impossible to accommodate six identical fermions of spin 1/2 of the same flavor in the ground state with three colors and total spin 2: two of them must necessarily be in the same quantum state. ⁴ As mentioned above, this result has already been reported in the literature for other two-baryon systems showing Pauli blocking [\[29](#page-14-9)], obtaining repulsive cores starting about $0.75-0.78$ fm in the case of baryons made of lights quarks, i.e., $\alpha \simeq 0.5$ fm. What is really important is that this effect is independent of the dynamical model considered as long as interactions at the quark level are considered.

It is also important to emphasize that such a repulsive hard-core would not appear if antisymmetry effects among the baryon constituent would have not been considered. This would be the case if the interactions were derived without taking into account the quark substructure of the baryons, i.e., by extrapolating them either from interactions of other two-body systems or from the same system with quantum numbers that were not affected

⁴ Note that, as in the case of the tetraquarks, the similarities with atomic physics are maintained because a dynamics-independent effect due to the Fermi-Dirac statistic is responsible, for example, for reducing the energy of parahelium compared to that of orthohelium.

by Pauli blocking. This has been clearly illustrated in Ref. [\[18](#page-13-16)], showing that a naive ansatz of spin independence of the $\Omega^i \Omega^i$ interaction leads to bound states of hundreds of MeV for the $i = b$ case using the lattice QCD interactions of Ref. [\[11](#page-13-10)]. Even if one assumes that the ⁵ S_2 Ω ^{*i*} Ω ^{*i*} interaction has a strong repulsive core, but of the same range as that of the ¹ S_0 $\Omega^i \Omega^i$ interaction, which would be equivalent to disregarding the Pauli blocking effects on the ${}^{5}S_{2}$ channel, the binding energy changes very slowly leading to bound states of between 250 and 350 MeV for $i = b$ systems.

In addition to the evidence discussed in the previous section, it is important to highlight that there are preliminary studies of lattice QCD about the S-wave scattering of strangeness -3 baryons [\[41\]](#page-14-21) concluding a strongly repulsive interaction for $S = 2$. Using lattice configurations with pion mass $m_{\pi} \sim 390$ MeV and two different volumes, the $\Omega \Omega^{-1} S_0$ channel was found to be most likely weakly repulsive or attractive, leading to an extrapolated scattering length $a_{S=0}^{\Omega\Omega} = 0.16 \pm 0.22$ fm. However, the states achieved for $S=2$ are at a significantly higher energy level than for the $S = 0$ case, implying a strongly repulsive channel. In fact, it is suggested that none of the extracted states were actually the $S = 2$ ground state, pointing to further studies at different pion masses approaching the physical point to gain a better understanding.

Finally, we have solved the full $\Omega^i \Omega^i \Omega^i$ three-body problem by the Faddeev method [\[19\]](#page-13-18) considering the ${}^{1}S_{0}$ and ${}^{5}S_{2}$ partial waves. We have used as input the ${}^{1}S_{0}$ $\Omega^{i}\Omega^{i}$ potentials derived from the lattice QCD calculations [\[9](#page-13-8)[–11](#page-13-10)] and the ${}^5S_0 \Omega^i \Omega^i$ potentials of Fig. [2.](#page-10-0) The repulsive character of the single channel calculation due to the Pauli effects in the ${}^{1}S_{0}$ partial wave are reinforced due to the strong repulsion of the 5S_2 Ω ^{*i*} Ω ^{*i*} interactions due to Pauli blocking effects. In spite of the attractive character of the ${}^{1}S_{0}$ two-body interactions, we have not found $J^P = 3/2^+$ three-body bound states in any flavor sector.

Thus, two different consequences of the Pauli principle conspire against the possible existence of $\Omega^{i} \Omega^{i}$ bound states. On the one hand, at the baryon level, the negative recoupling coefficient of the spin-0 two-body amplitudes that effectively changes the nature of the ${}^{1}S_{0}$ $\Omega^{i}\Omega^{i}$ two-body interactions from attractive to repulsive. On the other hand, since the Ω^{i} 's are spin 3/2 fermions, they also contribute spin-2 amplitudes that are free of Pauli effects at baryonic level. The contribution of the spin-2 amplitudes to the unique S-wave fully antisymmetric state made of three Ω^i , the $J^P = 3/2^+$, is greater than that of the ${}^{1}S_{0}$ partial wave. However, due to the underlying quark substructure, spin-2 amplitudes present Pauli blocking which translates into a strong repulsive interaction. The cooperative combination of these two consequences of the Pauli principle, at the baryon level and at the quark level, points to the difficulty of the existence of S-wave $\Omega^i \Omega^i \Omega^i$ bound states for any flavor sector.

V. SUMMARY AND OUTLOOK.

In brief, we have analyzed the current question regarding the possible existence of S-wave $\Omega^i \Omega^i \Omega^i$ bound states for any flavor $i = s, c, b$. This possibility is suggested by the attractive character of the ¹ $S_0 \Omega^i \Omega^i$ interactions derived by lattice QCD simulations, particularly strong for the heavier flavor $i = b$. We have justified the importance of performing a full fledged study of the three-body system, including the 5S_2 Ω ^{*i*} Ω ^{*i*} partial wave, which has not yet been studied by lattice QCD. We have shown that, as in the three-neutron case, the three-body recoupling coefficients effectively change the nature of the ${}^{1}S_{0} \Omega^{i} \Omega^{i}$ two-body interaction from attractive to repulsive in the three-body system. We have also found that the dominant 5S_2 partial wave shows Pauli blocking, a saturation phenomenon independent of the dynamics that translates into a strong repulsive interaction. It is important to highlight how the ¹S₀ $\Omega^i \Omega^i$ partial-wave does not satisfy the S–wave Pauli blocking condition. As stated above, the cooperative combination of these two consequences of the Pauli principle, at the baryon level and at the quark level, points to the difficulty of the existence of $\Omega^i \Omega^i \Omega^i$ bound states despite the attractive interaction – rather strong in some flavor sectors – predicted for the $\Omega^i \Omega^{i-1} S_0$ partial wave by lattice QCD calculations.

The results of this study suggest an effort by lattice QCD to derive the interaction in the ⁵ $S_2 \Omega^i \Omega^i$ channel and reveal the existence of a strong dynamics-independent repulsion, thus demonstrating the observable effects of Pauli blocking due to quark substructure. In addition, it would also highlight the caution to be taken when extrapolating interactions at the baryon level between different systems or quantum numbers without taking into account the observable effects of the quark substructure, the so-called Pauli blocking.

VI. ACKNOWLEDGMENTS.

The authors acknowledge enlightening correspondence with N. Mathur and M. Padmanath. This work has been partially funded by COFAA-IPN (México) and by the Spanish Ministerio de Ciencia e Innovación (MICINN) and the European Regional Development Fund (ERDF) under contracts PID2022-141910NB-I00 and RED2022-134204-E.

- [1] S. K. Choi et al. (Belle Collaboration), Phys. Rev. Lett. 91, 262001 (2003).
- [2] M. Gell-Mann, Phys. Lett. 8, 214 (1964).
- [3] A. Hayrapetyan *et al.* (CMS Collaboration), Phys. Rev. Lett. **132**, 111901 (2024).
- [4] J. -P. Ader, J. -M. Richard, and P. Taxil, Phys. Rev. D 25, 2370 (1982).
- [5] J. -M. Richard, J. Fröhlich, G. -M. Graf, and M. Seifert, Phys. Rev. Lett. **71**, 1332 (1993).
- [6] R. Aaij *et al.* (LHCb Collaboration), Nat. Phys. **18**, 751 (2022).
- [7] R. Aaij et al. (LHCb Collaboration), Nat. Commun. 13, 3351 (2022).
- [8] B. Colquhoun, A. Francis, R. J. Hudspith, R. Lewis, K. Maltman, and W. G. Parrott, Phys. Rev. D 110, 094503 (2024).
- [9] S. Gongyo et al. (HAL QCD Collaboration), Phys. Rev. Lett. 120, 212001 (2018).
- [10] Y. Lyu, H. Tong, T. Sugiura, S. Aoki, T. Doi, T. Hatsuda, J. Meng, and T. Miyamoto, Phys. Rev. Lett. 127, 072003 (2021).
- [11] N. Mathur, M. Padmanath, and D. Chakraborty, Phys. Rev. Lett. 130, 111901 (2023).
- [12] M. Jurič *et al.*, Nucl. Phys. B 52 , 1 (1973).
- [13] J. Adam et al. (STAR Collaboration), Nat. Phys. 16, 409 (2020).
- [14] A. Esser, S. Nagao, F. Schulz, P. Achenbach, C. Ayerbe Gayoso, R. Böhm et al. (A1 Collaboration), Phys. Rev. Lett. 114, 232501 (2015).
- [15] M. Karliner and J. L. Rosner, Phys. Rev. Lett. 119, 202001 (2017).
- [16] E. J. Eichten and C. Quigg, Phys. Rev. Lett. 119, 202002 (2017).
- [17] T. -W. Wu, S. -Q. Luo, M. -Z. Liu, L. -S. Geng, and X. Liu, Phys. Rev. D 108, L091506 (2023).
- [18] H. Garcilazo and A. Valcarce, Rev. Mex. Fis. 70, 041202 (2024).
- [19] H. Garcilazo, F. Fernández, A. Valcarce, and R. D. Mota, Phys. Rev. C 56, 84 (1997).
- [20] A. Valcarce, H. Garcilazo, F. Fernández, and P. González, Rep. Prog. Phys. 68, 965 (2005).
- [21] H. Garcilazo, J. Phys. G **13**, L63 (1987).
- [22] E. Hernández, J. Vijande, A. Valcarce, and J. -M. Richard, Phys. Lett. B 800, 135073 (2020).
- [23] J. -M. Richard, A. Valcarce, and J. Vijande, Phys. Rev. Lett. 124, 212001 (2020).
- [24] J. -M. Richard, A. Valcarce, and J. Vijande, Phys. Rev. D 95, 054019 (2017).
- [25] A. Valcarce, F. Fernández, and P. González, Phys. Rev. C 56, 3026 (1997).
- [26] M. Oka, K. Shimizu, and K. Yazaki, Nucl. Phys. A 464, 700 (1987).
- [27] S. Saito, Prog. Theor. Phys. 41, 705 (1969).
- [28] M. Oka and K. Yazaki, Prog. Theor. Phys. 66, 556 (1981).
- [29] M. Oka and K. Yazaki, Prog. Theor. Phys. 66, 572 (1981).
- [30] H. Garcilazo, T. Fernández-Caramés, and A. Valcarce, Phys. Rev. C 75 , 034002 (2007); H. Garcilazo, A. Valcarce, and T. Fernández-Caramés, Phys. Rev. C 76, 034001 (2007).
- [31] H. Garcilazo, A. Valcarce, and T. F. Caramés, Eur. Phys. J. C 79, 598 (2019); H. Garcilazo, A. Valcarce, and T. F. Caramés, Phys. Rev. C **92**, 024006 (2015).
- [32] E. Ferreira and H. G. Dosch, Phys. Rev. C 40, 1750 (1989).
- [33] T. Takaki, Ann. Phys. (N.Y.) 166, 1 (1986).
- [34] D. A. Liberman, Phys. Rev. D **16**, 1542 (1977).
- [35] M. Oka and K. Yazaki, Int. Rev. Nucl. Phys. 1, 489 (1984).
- [36] J. Vijande, F. Fernández, and A. Valcarce, J. Phys. G **31**, 481 (2005).
- [37] A. Valcarce, H. Garcilazo, and J. Vijande, Eur. Phys. J. A 37, 217 (2008).
- [38] A. de Rújula, H. Georgi, and S. L. Glashow, Phys. Rev. D 12, 147 (1975).
- [39] G. S. Bali, Phys. Rep. **343**, 1 (2001).
- [40] A. Gal, H. Garcilazo, A. Valcarce, and T. Fernández-Caramés, Phys. Rev. D 90, 014019 (2014).
- [41] M. I. Buchoff, T. C. Luu, and J. Wasem, Phys. Rev. D 85, 094511 (2012).