# $\Omega_{bbb}\Omega_{bbb}\Omega_{bbb}$ tribaryons

H.  $Garcilazo^{1,*}$  and A.  $Valcarce^{2,\dagger}$ 

<sup>1</sup>Escuela Superior de Física y Matemáticas Instituto Politécnico Nacional, Edificio 9, 07738 CDMX, Mexico <sup>2</sup>Departamento de Física Fundamental, Universidad de Salamanca, E-37008 Salamanca, Spain

(Dated: January 20, 2025)

## Abstract

We study the possible existence of bound states of three  $\Omega_{bbb}$  baryons. We consider only S wave interactions and we start from recent lattice QCD results which give a strongly attractive potential between two  $\Omega_{bbb}$  baryons in the  $^1S_0$  channel. We analyze different scenarios. At baryonic level, the  $\Omega_{bbb}\Omega_{bbb}$  interaction could be understood to be basically spin-independent, so that the two contributing channels,  $^1S_0$  and  $^5S_2$ , would have a very similar interaction. This baryonic analysis leads to the existence of bound states in the three-body system. At the quark level, repulsive effects would appear in the  $^5S_2$  channel, making it more repulsive than the  $^1S_0$  channel. We study the effect of such repulsion in terms of its range.

<sup>\*</sup>Electronic address: hgarcilazos@ipn.mx

<sup>†</sup>Electronic address: valcarce@usal.es

#### I. INTRODUCTION

Recently, there have been several interesting developements on the possible existence of bound states of two and three  $\Omega$  baryons. For example, Ref. [1], using a one-boson-exchange (OBE) model found bound states of the systems  $\Omega_{ccc}\Omega_{ccc}$  and  $\Omega_{bbb}\Omega_{bbb}$ . Ref. [2] derived a  $\Omega\Omega$  interaction based on lattice QCD. Similarly, Ref. [3], a lattice QCD calculation with nearly physical light-quark masses, derived the  $\Omega_{ccc}\Omega_{ccc}$  interaction in the  $^1S_0$  channel. They obtain a bound state with a binding energy of 5.68 MeV. More recently, Ref. [4] performed a lattice QCD calculation of the  $\Omega_{bbb}\Omega_{bbb}$  system finding a very deep bound state in the  $^1S_0$  channel, with a binding energy of 81 MeV. The energy of the bound states of the two-body systems would be the threshold of any possible three-body bound state. Finally, Ref. [5], using the existing lattice QCD interactions for the different  $\Omega\Omega$  systems, investigated the three-body systems  $\Omega\Omega\Omega$ ,  $\Omega_{ccc}\Omega_{ccc}\Omega_{ccc}$ , and  $\Omega_{bbb}\Omega_{bbb}\Omega_{bbb}$ . They found that none of the three-body systems binds. However, making use of the OBE interactions of Ref. [1] the  $\Omega\Omega\Omega$  system develops a bound state.

In this work, we investigate whether the  $\Omega_{bbb}\Omega_{bbb}\Omega_{bbb}$  system is bound. The  $\Omega_{bbb}$  baryon has spin 3/2 and no isospin, so that the two-body system can have total spin  $S_i = 0, 1, 2$ , and 3. However, the states  $S_i = 1$  and  $S_i = 3$  are not allowed in S-wave by the Pauli principle, so that one is left with only the states  $S_i = 0$  and  $S_i = 2$ . In Ref. [4], they obtained the  $\Omega_{bbb}\Omega_{bbb}$  interaction only for the channel  $S_i = 0$ , so that we will have to discuss the situation of the channel  $S_i = 2$ . We will use some hypotheses deduces either at the baryon level or at the quark level about the  $\Omega_{bbb}\Omega_{bbb}$   $^5S_2$  interaction.

We carry out our study within the formalism of the nonrelativistic Faddeev equations for three identical particles considering only S waves. We start our discussion of the three-body system considering only the  $S_i = 0$  two-body channel and afterwards we analyze the effect of the  $S_i = 2$  two-body channel for the three-body bound state.

## II. SINGLE CHANNEL FADDEEV PROBLEM

The Faddeev equations for three identical particles are

$$T = 2t_i G_0 T (1)$$

where  $t_i$  is the t-matrix of the two-body system,

$$t_i = V + VG_0t_i (2)$$

where V the two-body interaction in the  ${}^{1}S_{0}$  channel and  $G_{0}$  is the propagator for three free particles.

We use the complete set of basis states  $|i\rangle$ ,

$$|i\rangle = |p_i q_i((s_i, s_k) S_i, s_i) J\rangle , \qquad (3)$$

with  $p_i$  and  $q_i$  the standard Jacobi momenta,  $s_i$ ,  $s_j$ , and  $s_k$  the spins of the three particles,  $S_i$  the total spin of the pair jk, and J the total spin of the three-body system. In this basis the Faddeev equation (1) becomes,

$$\langle i | T | \phi_0 \rangle = 2 \langle i | t_i | i' \rangle \langle i' | j \rangle G_0 \rangle j | T | \phi_0 \rangle , \qquad (4)$$

where the explicit form of the integral equation in momentum space is given in the Appendix.

The recoupling coefficient,

$$\langle i'|j\rangle = \langle p_i'q_i'|p_iq_i\rangle \langle ((s_i, s_k)S_i, s_i)J|((s_k, s_i)S_j, s_j)J\rangle , \qquad (5)$$

is of great interest. The space part  $\langle p_i'q_i'|p_jq_j\rangle$  is positive definite [6]; however, the spin part is

$$\langle ((s_j, s_k)S_i, s_i)J | ((s_k, s_i)S_j, s_j)J \rangle = (-)^{S_j + s_j - J} \sqrt{(2S_i + 1)(2S_j + 1)} W(s_j s_k J s_i; S_i S_j) .$$
 (6)

Since  $S_i = S_j = 0$  and  $s_i = s_j = s_k = J = 3/2$  one gets,

$$\langle ((s_j, s_k)S_i, s_i)J| ((s_k, s_i)S_j, s_j)J\rangle = (-)^{2s_j} \frac{1}{2s_j + 1} = -\frac{1}{4} , \qquad (7)$$

which is a negative number, so that it effectively changes the nature of the two-body interaction from attractive to repulsive such that no bound state can be obtained in a one-channel calculation. <sup>1</sup>

<sup>&</sup>lt;sup>1</sup> On the opposite side, a repulsive potential together with a negative spin recoupling coefficient does not change the nature of the interaction so as to lead to a three-body bound state, as discussed in Ref. [7] for the case of the  $J = 3/2 \Sigma^- nn$  system.

The result of Eq. (7) is a direct consequence of the Pauli principle and it applies for all systems with three identical fermions, i.e., particles with spin half-integer, like the case of three neutrons where,

$$\langle ((s_j, s_k)S_i, s_i)J | ((s_k, s_i)S_j, s_j)J \rangle = (-)^{2s_j} \frac{1}{2s_j + 1} = -\frac{1}{2} .$$
 (8)

It is worth noting that in the three-neutron case there is only a two-body channel,  $S_i = 0$ , if one includes only S waves, so that there is no possibility for a three-neutron bound state with any interaction. However, in the case of the three omegas besides the  $S_i = 0$  channel one also has the  $S_i = 2$  two-body channel.

# III. BARYONIC LEVEL ${}^5S_2$ $\Omega\Omega$ INTERACTION

A two-body interaction acting in S-waves contains only central and spin-spin terms, since terms like spin-orbit, tensor, etc., act only for nonzero orbital angular momentum.

The phenomenological description of the spectra of mesons and baryons in a nonrelativistic approach is based in a two-body potential between quarks [8, 9]. In Ref. [8] such potential was taken to be,

$$V(r) = -\frac{\kappa}{r} + \lambda r - \Lambda + \frac{\kappa}{m_i m_i} \frac{exp^{-r/r_0}}{rr_0} \vec{\sigma}_i \cdot \vec{\sigma}_j$$
 (9)

and similarly in Ref. [9]. The four terms in the r.h.s. of Eq. (9) are, respectively, the Coulomb term, the linear confinement term, the constant term, and the spin-spin term. This interaction is able to reproduce reasonably well the masses and other properties of all the existing mesons and baryons [9].

The Yukawa function in the spin-spin term is an extended delta function which becomes a delta function if  $r_0 \to 0$ . This form of the spin-spin interaction is suggested by the non-relativistic reduction of the one-gluon-exchange diagram [10],

$$H = \frac{2\alpha_s}{3m_i m_j} \frac{8\pi}{3} \delta^3(\vec{r}) \vec{S}_i \cdot \vec{S}_j, \tag{10}$$

where  $m_i$  and  $\vec{S}_i$  are the mass and spin operator of particle i. As one can see in Eqs. (9) and (10), the spin-spin term is inversely proportional to  $m_i m_j$  so that in the case of baryons with identical quarks it goes as  $1/m_i^2$ . Thus, using the masses for the light quarks  $n \approx 0.3$  GeV) and for the heavy quark  $b \approx 5$  GeV from Refs. [8, 9], one gets that the spin-spin term

in the case of the bb interaction is about 250 times smaller that that of the nn interaction and therefore it is negligible. This means that the interaction between two b quarks is basically independent of the spin.

Let us now consider the spin-spin interaction in the case of two  $\Omega_{bbb}$  baryons. In a quark model description, the baryon-baryon interaction can be deduced from the quark-quark interaction following a well-known procedure [11, 12]. In the case where the interactions are restricted to S waves, if the quark-quark interaction is almost independent of the spin, the baryon-baryon interaction will also be almost independent of the spin [12]. This would imply a purely central  $\Omega_{bbb}\Omega_{bbb}$  interaction, so that  $V_{\Omega_{bbb}\Omega_{bbb}}(^1S_0) \sim V_{\Omega_{bbb}\Omega_{bbb}}(^5S_2)$  and, thus, one can use the  $S_i = 0$  interaction of Ref. [4] also for  $S_i = 2$ .

### IV. TWO-CHANNEL FADDEEV PROBLEM

The explicit form of the Faddeev equations for the two-channel problem in momentum space is also given in the Appendix. If one includes the two channels  $S_i = 0$  and  $S_i = 2$ , the spin recoupling coefficients of Eq. (6) are,

$$\langle 0|0\rangle = -\frac{1}{4} \qquad \langle 0|2\rangle = -\frac{\sqrt{5}}{4}$$

$$\langle 2|0\rangle = -\frac{\sqrt{5}}{4} \qquad \langle 2|2\rangle = +\frac{3}{4}.$$
(11)

The recoupling coefficient  $\langle 2|2\rangle$  is positive and much larger than the coefficient  $\langle 0|0\rangle$ , so that if the interaction in the channel  $S_i = 2$  is equal to that of the channel  $S_i = 0$ , one would expect to get a bound state.

Following the conclusion of the previous section we start by taking  $V_{\Omega_{bbb}\Omega_{bbb}}(^5S_2) = V_{\Omega_{bbb}\Omega_{bbb}}(^1S_0)$ . We find in this case that the ground state and one excited state of the three-body  $\Omega_{bbb}\Omega_{bbb}\Omega_{bbb}$  system, that lie at -393.8 MeV and -94.9 MeV, respectively (-363.9 MeV and -81.0 MeV, respectively, if one includes the Coulomb interaction).

## V. QUARK LEVEL ${}^5S_2$ $\Omega\Omega$ INTERACTION

As it has been discussed in the literature [11, 14–16] there appears quark Pauli blocking for particular channels of some two-baryon systems. This has been discussed in detail, for example, for the  $\Sigma N$  system [16] or for the  $\Delta N$  and  $\Delta \Delta$  systems [11, 14, 15]. Pauli blocking




FIG. 1: The  $\Omega_{bbb}\Omega_{bbb}$  interaction  $V_2$  for A=0,1, and 2.

translates into repulsive cores of of long-range ( $\sim 1$  fm). See, for example, Fig. 3 of Ref. [15]. The  $\Omega_{bbb}\Omega_{bbb}$  two-baryon belongs to the same flavor multiplet as the  $\Delta\Delta$  system and thus it also shows Pauli blocking for the  $^5S_2$  channel, which means a strong long-range repulsive hard core. In order to simulate these effects we will take the  $S_i=2$  interaction as,

$$V_2(r) = \begin{cases} V_0(r) & r > r_0 \\ V_0(r) + A[V_0(r) - V_0(r_0)] \left(1 - \frac{r}{r_0}\right)^2 & r \le r_0 \end{cases}$$
(12)

where A is a constant and  $r_0 = 0.104$  fm is the radius where  $V_0$  is minimum (see Fig. 1). The form (12) guarantees that  $V_2(r)$  and  $dV_2(r)/dr$  are continuous at  $r = r_0$ . This transformation has the effect of pushing upwards the short-range part of the interaction, thereby reducing the attraction. We show this behavior in Fig. 1 for the special cases A = 1 and A = 2 as compared with the A = 0 case, which corresponds to the model of Ref. [4]. Thus, our final model of the  $\Omega_{bbb}\Omega_{bbb}$  interaction contained in Eq. (12) and Fig. 1, has the  ${}^5S_2$  interaction less attractive than the  ${}^1S_0$  interaction which is similar to what is obtained in the OBE model of Ref. [1].

We show in Fig. 2 the evolution of the binding energy (with and without Coulomb) as a function of the parameter A. As one can see, the binding energy changes very slowly. We consider that values of A larger than 20 are not realistic so that we expect the binding energy to lie between 250 and 350 MeV.




FIG. 2: The  $\Omega_{bbb}\Omega_{bbb}\Omega_{bbb}$  binding energy as a function of the parameter A.

## VI. ACKNOWLEDGEMENTS

This work has been partially funded by COFAA-IPN (México) and by the Spanish Ministerio de Ciencia e Innovación (MICINN) and the European Regional Development Fund (ERDF) under contracts PID2019-105439GB-C22, PID2022-141910NB-I00, and RED2022-134204-E.

## VII. APPENDIX

In order to solve the single channel three-body problem we use the method of Ref. [13], where the two-body amplitudes are expanded in terms of Legendre polynomials. Thus, the Faddeev equations for the bound state problem take the simple form,

$$T^{n}(q_{i}) = \sum_{n} \int_{0}^{\infty} dq_{j} K^{nm}(q_{i}, q_{j}; E) T^{m}(q_{j}) , \qquad (13)$$

where

$$K^{nm}(q_i, q_j; E) = 2 \langle 0 | 0 \rangle \sum_{r} \tau_i^{nr} (E - q_i^2 / 2\nu_i) \frac{q_i^2}{2} \int_{-1}^{1} d\cos\theta \frac{P_r(x_i') P_m(x_j)}{E - p_j^2 / 2\eta_j - q_j^2 / 2\nu_j} , \qquad (14)$$

with  $\langle 0|0\rangle$  the spin recoupling coefficient (11),

$$\tau_i^{nr}(e) = \frac{2n+1}{2} \frac{2r+1}{2} \int_{-1}^1 dx_i \int_{-1}^1 dx_i' P_n(x_i) t_i(x_i, x_i'; e) P_r(x_i') , \qquad (15)$$

and

$$x_i = \frac{p_i - b}{p_i + b} \ . \tag{16}$$

 $p_j$  and  $q_j$  are the magnitudes of the Jacobi relative momenta while  $\eta_j$  and  $\nu_j$  are the corresponding reduced masses.

 $t_i(x_i, x_i'; e)$  corresponds to the off-shell two-body t-matrix  $t_i(p_i, p_i'; e)$  through the transformation (16), with b a scale parameter on which the solution does not depend. The off-shell two-body t-matrices are obtained by solving the Lippmann-Schwinger equation,

$$t_i(p_i, p_i'; e) = V_i(p_i, p_i') + \int_0^\infty p_i'' 2dp_i'' V_i(p_i, p_i'') \frac{1}{e - p_i''^2 / 2\eta_i + i\epsilon} t_i(p_i'', p_i'; e) , \qquad (17)$$

with  $e = E - q_i^2/2\nu_i$ . In order to solve the two-channel three-body problem, Eq. (13) becomes,

$$T_{\alpha}^{n}(q_i) = \sum_{m\beta} \int_0^\infty dq_j K_{\alpha\beta}^{nm}(q_i, q_j; E) T_{\beta}^{m}(q_j) , \qquad (18)$$

with

$$K_{\alpha\beta}^{nm}(q_i, q_j; E) = 2 \langle \alpha | \beta \rangle \sum_{r} \tau_{\alpha}^{nr} (E - q_i^2 / 2\nu_i) \frac{q_i^2}{2} \int_{-1}^{1} d\cos\theta \frac{P_r(x_i') P_m(x_j)}{E - p_j^2 / 2\eta_j - q_j^2 / 2\nu_j} . \tag{19}$$

- [1] M. -Z. Liu and L. -S. Geng, Prediction of an  $\Omega_{bbb}\Omega_{bbb}$  dibaryon in the extended one-boson exchange model, Chin. Phys. Lett. **38** 10, 101201 (2021).
- [2] S. Gongyo, K. Sasaki, S. Aoki, T. Doi, T. Hatsuda, Y. Ikeda, T. Inoue, T. Iritani, N. Ishii, T. Miyamoto, and H. Nemura (HAL QCD Collaboration), Most Strange Dibaryon from Lattice QCD, Phys. Rev. Lett. 120, 212001 (2018).
- [3] Y. Liu, H. Tong, T. Sugiura, S. Aoki, T. Doi, T. Hatsuda, J. Meng, and T. Miyamoto, Dibaryon with Highest Charm Number near Unitarity from Lattice QCD, Phys. Rev. Lett. 127, 072003 (2021).
- [4] N. Mathur, M. Padmanath, and D. Chakraborty, Strongly Bound Dibaryon with Maximal Beauty Flavor from Lattice QCD, Phys. Rev. Lett. 130, 111901 (2023).
- [5] T. -W. Wu, S. -Q. Luo, M. -Z. Liu, L. -S. Geng, and X. Liu, Tribaryons with Lattice QCD and one-boson exchange potentials, Phys. Rev. D 108, L091506 (2023).
- [6] R. Balian and E. Brezin, Angular-momentum reduction of the Faddeev equations, Nuov. Cim.61B, 403 (1969).

- [7] H. Garcilazo, Non-Existence of  $\Lambda NN$  and  $\Sigma^-NN$  Bound States, J. Phys. G 13, L63 (1987).
- [8] R. K. Bhaduri, L. E. Cohler, and Y. Nogami, A Unified Potential for Mesons and Baryons, Nuov. Cim. 65A, 376 (1981).
- [9] B. Silvestre-Brac, Spectrum and Static Properties of Heavy Baryons, Few-Body Syst. **20**, 1 (1996).
- [10] N. Isgur and G. Karl, P-Wave Baryons in the Quark Model, Phys. Rev. D 18, 4187 (1978).
- [11] A. Valcarce, H. Garcilazo, F. Fernández, and P. González, Quark Model Study of Few Baryon Systems, Rep. Prog. Phys. 68,965 (2005).
- [12] K. Shimizu, One-pion exchange potential based on a quark model, Phys. Lett. B **148**, 418 (1984).
- [13] H. Garcilazo and A. Valcarce, Deeply Bound  $\Xi$  tribaryon, Phys. Rev. C 93,034001 (2016).
- [14] M. Oka and K. Yazaki, Short Range Part of Baryon Baryon Interaction in a Quark Model. 1.
  Formulation, Prog. Theor. Phys. 66, 556 (1981).
- [15] M. Oka and K. Yazaki, Short Range Part of Baryon Baryon Interaction in a Quark Model. 2.
  Numerical Results for S-Wave, Prog. Theor. Phys. 66, 572 (1981).
- [16] M. Oka, K. Shimizu and K. Yazaki, Hyperon Nucleon and Hyperon-hyperon Interaction in a Quark Model, Nucl. Phys. A 464, 700 (1987).