ON THE NUMBER OF COFINALITIES OF CUTS IN ULTRAPRODUCTS OF LINEAR ORDERS

MOHAMMAD GOLSHANI

ABSTRACT. Suppose κ is a regular cardinal and $\bar{a} = \langle \mu_i : i < \kappa \rangle$ is a nondecreasing sequence of regular cardinals. We study the set of possible cofinalities of cuts $Pcut(\bar{a}) = \{(\lambda_1, \lambda_2) : \text{for some ultrafilter } D \text{ on } \kappa, (\lambda_1, \lambda_2) \text{ is the}$ cofinality of a cut of $\prod_{i < \kappa} \mu_i / D\}$.

§ 1. INTRODUCTION

In late 1980th, Shelah introduced the theory of possible cofinalities, abbreviated pcf and used it to prove many surprising and unexpected results in ZFC, showing that ZFC is much more powerful than what was expected before by many set theorists. Given a regular cardinal κ and an increasing sequence of regular cardinals $\bar{a} = \langle \mu_i : i < \kappa \rangle$, pcf(\bar{a}) is defined as

$$\operatorname{pcf}(\bar{a}) = \{\operatorname{cf}(\prod_{i < \kappa} / D) : D \text{ is an ultrafiler on } \kappa\}.$$

One of the main results in pcf theory is that under suitable assumptions on the sequence \bar{a} , $|pcf(\bar{a})|$ is bounded and indeed $|pcf(\bar{a})| < \min\{(2^{\kappa})^+, \kappa^{+4}\}$ (see [2]).

In this paper, we introduce and study a related concept, where instead of taking cofinalities of ultraproducts, we take cuts of such ultraproducts and their cofinalities. Let κ be a regular cardinal and let \bar{a} be a non-decreasing sequence of regular cardinals. The set of possible cuts of \bar{a} is defined as

 $\begin{aligned} \operatorname{Pcut}(\bar{a}) &= \{ (\lambda_1, \lambda_2) : \lambda_1, \lambda_2 \geq \aleph_0 \text{ and for some ultrafilter } D \text{ on } \kappa, \, (\lambda_1, \lambda_2) \text{ is the} \\ & \text{ cofinality of a cut of } \prod_{i < \kappa} \mu_i / D \}. \end{aligned}$

²⁰¹⁰ Mathematics Subject Classification. Primary: 03E05, 03E10.

Key words and phrases. Possible cuts, pcf, cardinal arithmetic.

The author's research has been supported by a grant from IPM (No. 1403030417). The author thanks Saharon Shelah for some useful discussions.

M. GOLSHANI

We find a bound on $Pcut(\bar{a})$ by showing that $|Pcut(\bar{a})| \leq 2^{\kappa}$. We also discuss the optimality of this result.

§ 2. BOUNDS ON Pcut

Let κ be a regular cardinal and let $\bar{a} = \langle \mu_i : i < \kappa \rangle$ be an increasing sequence of regular cardinals. In this section we discuss the cardinality of the set $Pcut(\bar{a})$. We first prove the following:

Theorem 2.1. Suppose κ and \bar{a} are as above. Then $|\operatorname{Pcut}(\bar{a})| \leq 2^{\kappa}$. Indeed if $2^{\kappa} = \aleph_{\alpha}$, then $|\operatorname{Pcut}(\bar{a})| \leq |\alpha + 1|^2$.

Proof. We prove the theorem by showing that if $(\lambda_1, \lambda_2) \in \text{Pcut}(\bar{a})$, then $\lambda_1, \lambda_2 \leq 2^{\kappa}$. Thus fix (λ_1, λ_2) as above and let D be an ultrafilter on κ such that (λ_1, λ_2) is the cofinality of a cut of $\prod_{i < \kappa} \mu_i / D$.

 $(*): \lambda_2 \le 2^{\kappa}.$

Proof. Suppose towards a contradiction that $\lambda_2 > 2^{\kappa}$ and let $\bar{g} = (g_{\xi} : \xi < (2^{\kappa})^+)$ be a $<_D$ -decreasing sequence in $\prod_{i < \kappa} \mu_i$. Define the function $F : [(2^{\kappa})^+]^2 \to \kappa$ by

$$F(\xi, \zeta) = \min\{i < \kappa : g_{\zeta}(i) < g_{\xi}(i)\}.$$

The function F is well-defined. By the Erdos-Rado partition theorem, we can find some set X of cardinality κ^+ and some $i_* < \kappa$ such that for all $\xi < \zeta$ in X, we have $g_{\zeta}(i_*) < g_{\xi}(i_*)$. Thus we get an infinite strictly decreasing sequence of ordinals which is absurd. Thus $\lambda_2 \leq 2^{\kappa}$, as requested.

 $(**): \lambda_1 \leq 2^{\kappa}.$

Proof. Suppose towards a contradiction that $\lambda_1 > 2^{\kappa}$. Let D and (\bar{f}, \bar{g}) witness that $(\lambda_1, \lambda_2) \in \text{Pcut}(\bar{a})$, where $\bar{f} = \langle f_{\alpha} : \alpha < \lambda_1 \rangle$ and $\bar{g} = \langle g_{\xi} : \xi < \lambda_2 \rangle$. For $i < \kappa$ set $U_i = \{g_{\xi}(i) : \xi < \lambda_2\}$. Note that $|U_i| \leq \lambda_2 \leq 2^{\kappa}$. Define the relation E_i on λ_1 by $\alpha E_i \beta$ iff the following hold:

(1) for all $\xi < \lambda_2$ and all $\delta \in U_i$,

(2) for all $\xi < \lambda_2$ and all $\delta \in U_i$,

$$f_{\alpha}(\xi) = \delta \iff f_{\beta}(\xi) = \delta.$$

The relation E_i is clearly an equivalence relation on λ_1 and it has at most $|U_i| \leq 2^{\kappa}$ many equivalence classes. Set $E = \bigcap_{i < \kappa} E_i$. Then E is also an equivalence relation and it has at most $(2^{\kappa})^{\kappa} = 2^{\kappa}$ many equivalence classes.

As $\lambda_1 > 2^{\kappa}$, there exists some equivalence class Y which is unbounded in λ_1 . Define $h \in \prod_{i < \kappa} \mu_i$ by

$$h(i) = \sup\{f_{\alpha}(i) : \alpha \in Y\}$$

Then h is easily seen to be the $<_D$ -least upper bound of \overline{f} and that for all $\xi < \lambda_2, h <_D g_{\xi}$. Thus

$$\forall \alpha < \lambda_1 \ \forall \xi < \lambda_2 \ (f_\alpha <_D h <_D g_\xi)$$

which contradicts the choice of (\bar{f}, \bar{g}) .

The theorem follows from (*) and (**).

Corollary 2.2. Suppose 2^{κ} is not a fixed point of the \aleph -function. Then $|\operatorname{Pcut}(\bar{a})| < 2^{\kappa}$.

Proof. Let α be such that $2^{\kappa} = \aleph_{\alpha}$. Then $\alpha < 2^{\kappa}$, and hence by Theorem 2.1,

$$|\operatorname{Pcut}(\bar{a})| \le |\alpha + 1|^2 < 2^{\kappa},$$

as required.

Remark 2.3. If in the definition of the set $Pcut(\bar{a})$, we allow λ_2 to be 1, then the set $Pcut(\bar{a})$ can be large. Indeed if D is an ultrafilter on κ and for all $i, \theta_i < \mu_i$ is a regular cardinal, then

$$(\operatorname{tcf}(\prod_{i<\kappa}\theta_i)/D,1)\in\operatorname{Pcut},$$

provided that $\operatorname{tcf}(\prod_{i < \kappa} \theta_i)/D$ exists. In particular if $\mu = \lim_D \langle \mu_i : i < \kappa \rangle = \aleph_{\alpha}$, then $|\alpha| \leq |\operatorname{Pcut}(\bar{a})|$.

The next lemma shows that Pcut can take its maximal possible number of elements. **Lemma 2.4.** Let $\bar{a} = \langle \mu_i : i < \kappa \rangle$ where $\kappa = \aleph_0$ and for $i < \omega$, $\mu_i = \omega$. Let $\theta > \aleph_0$ be an inaccessible cardinal and set $\mathbb{P} = \text{Add}(\omega, \theta)$. Then $V[\mathbf{G}_{\mathbb{P}}] \models \text{``}|\text{Pcut}(\bar{a})| = \theta = 2^{\aleph_0}$ ".

Proof. By [1],

 $V[\mathbf{G}_{\mathbb{P}}] \models \text{``Pcut}(\bar{a}) \supseteq \{(\lambda_1, \lambda_2) : \lambda_1, \lambda_2 \leq \kappa \text{ are uncountable regular cardinals}\}",$

from which the result follows.

References

 Canjar, Michael; Countable ultraproducts without CH. Ann. Pure Appl. Logic 37 (1988), no. 1, 1-79.

 Shelah, Saharon Cardinal arithmetic. Oxford Logic Guides, 29. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1994. xxxii+481 pp. ISBN: 0-19-853785-9

Mohammad Golshani, School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box: 19395–5746, Tehran, Iran.

 $Email \ address: \ {\tt golshani.m0gmail.com}$