# Excited s-wave 1<sup>--</sup> vector mesons, their leptonic decays and (in-)complete absence of abnormal states as seen from the constituent quark BSE

V. Šauli

DPT NPI Rez near Prague, Czech Academy of Sciences

January 22, 2025

#### Abstract

Within a lattice inspired interaction between quark and antiquark, we obtain hierarchy of Bethe-Salpeter equation (BSE) solutions for vector quarkonia excited states in the constituent quark mass approximation. As a toy model, we apply the similar to calculate ground and excited states of  $\phi$  and  $\omega/\rho$  meson. Performing detailed numerical search, our study provides the evidence that the quark propagator with single valued constant mass does not provide known meson spectra without the presence of abnormal (unphysical) solutions simultaneously. We classify normal and abnormal solutions and discuss necessary changes in calculation scheme preventing the spectrum from inconsistent solutions. While all experimentally narrow vector mesons are identified with a normal state of the BSE, the intriguing appearance of mutually canceled normal-abnormal states is reported. They appearance discriminate between heavy and the light mesons. In both cases they seems to be remainder of inconsistent use of the hmogeneous BSE for the description of broad resonances.

## 1 Introduction

The concept of constituent quark takes its historical place in development of our understanding to QCD. It is supposed to be approximately valid for description of phenomena where the quantum effects responsible for quark masses running can be ignored. Light quarks with constituent mass  $m_u \simeq m_d \simeq 300 MeV$  can be used to estimate nucleon properties, e.g. the mass and magnetic moments. This concept roughly works over entire baryonic sector, while a similar attempt to describe light pseudo-scalar mesons fails, due to their nambu-goldstone boson character.

For heavier mesons the Constituent Quark (CQ) approximation (CQA) turns to be more appropriate [1, 2] and model dependent constituent charm (bottom) quark mass in range 1.3 - 1.8(4.2 - 5.2)GeV is used as a successful quantum mechanical part of certain effective theories of QCD till these days [3, 4, 6, 5, 7, 8, 9, 10, 11, 12, 13, 14, 15].

An instantaneous Salpeter [16, 17, 18, 19] and quasipotential [20, 21, 22] approximations represent most widely used three dimensional reductions of BSE. Similarly to method based on the light cone hamiltonians, they often exploit the CQ concept as well. Further augmented by phenomenological confinement interaction, these approaches provide masses and leptonic decays with a good agreement with experiment. It is believed that proximity of sum of CQ masses to the physical mass of hadron M makes CQA particularly viable.

Here we provide, certainly not the first evidence, that a more detailed view offer a slightly more complicated picture. Using relativistic -quantum field theory- based description we will report on inherent inconnsistency when dealing with broad resonances in naive way we treat the excited states in quantum mechanics. We will show the homogeneous Bethe-Salpeter equation (BSE) provide more complicate description when applied naively to excited strangeonium and complex of  $\omega$  and  $\rho$  excitations. While CQA BSE works reasonably well for describing heavy quarkonia, however when trying to descrobe objects that we sometimes call excited light vector mesons i.e.  $\omega' \omega'', \rho' \dots$ , we get in adition to normal states the abnormal solutions as well. Within achieved numerical accuracy, we observe that the poles of abnormal states cancel against the normal solution. In this paper we explain, how the narrowness or broadness of assumed bound state or resonance is reflected or ignored by the BSE solutions. Furthermore, we explain under which condition the solitery abnormal states disappear from the solution in CQA BSE.

Before presenting the results we make a short overwiev of achievements in Dyson-Schwigner (DS) formalism. Within the DS/BS formalism the constituent mass concept has been questioned in [23], where BSE with constituent quark propagators has been compared with more elaborated Maris-Tandy Model (MTM) [24]. In the later model, the propagators were calculated from corresponding gap equations. The MTM is most studied phenomenological BSE model for mesons so far, it is based on ladder-rainbow truncation and originaly made to fit light pseudoscalr mesons. Note already their simple truncation respects Goldsteone character of pions, hence MTM is known working well in the sector of ground state pseudoscalars. Much lately, the parameters were used more freely and flawour dependent version of MTM has been boosted to the sector of heavy mesons [25, 26, 27, 28, 30, 29]. Spectra of heavy -light mesons have been further investigated in [30, 31] and their electromagnetic form factors were calculated [32]. All above mentioned 4d BSE MTM studies of mesons were obtained in the Landau gauge, which turns to be particularly useful for such purpose.

On the other side, MTM produces number of complex conjugated branch poles/points [33, 34] in the quark propagator functions. Originally attributed to the ladder approximation, this analytical structure is inherent to the functional form of MTM kernel instead. Further analyses [35, 36] of gap equations in complex plane reveal infinite tower of branch points generated by complex conjugated poles, i.e. by the quark propagators in MTM. Unavidably, in those models, there is an infinite ladder of branch cuts traversing complex plane transversely to imaginary axis of  $q^2$  variable. In practice, this structure is ignored and sorted in a simplified fit for the MTM quark propagator. Even doing so, it makes the solution of BSE particularly complicated for heavy states and heavier meson is considered in the MTM, more uncertainties one can expect for results. The size of estimated errors in masses and leptonic widths, as has been honestly estimated in [27, 28, 29] is a response on complicated complex structure the MTM actually has.

In attempt to make analytical properties of QCD vertices and propagators more simple in the LRA, new models were employed and suggested recently. They kernels are inspired by lattice data for which a simple analytical fits were made and they provide the first results in unlimited range of physical scales: from the pion consideration [37], till the heavy masses of excited bottomonia [38].

Presented study makes a step aside an we look what can be achieved within (UV improved) model [38] when considered in CQA. Thus instead of using quark gap equation, the single parameter 'classical' quark propagator is used. Up to short notion, we do not compare with dressed case yet, but extend to charmonium, strangeonium and light quark system as well. We discuss the presence (absence) of abnormal solution in CQA BSE. In case of vector mesons, these states correspond to negative norm and negative residuum BSE solution, while they keep the quantum numbers of the ordinary meson : C = -1, P =-1J = 1. The application of CQA to the light mesons should be regarded as a theoretical toy model, however the resulting spectroscopy is mesmerizing: We get the spectrum of narrow excited mesons, which is in nice agreement with PDG lists of experimentaly known broad resonances, however in a way that their poles are largely canceled by presence of opposite poles generated by ghost solutions. This happens regularly, suggesting there are only cuts in corresponding form factors used in therory for description of exclusive experimental data. In sections describing results we clarify under which condition an unphysical uncanceled ghost could disappear from the spectrum.

# 2 Quarkonia leptonic decays

The quarkonium meson decay constant  $F_V$  uniquely determine the leptonic partial decay rate  $\Gamma$  of meson carrying mass M such that

$$\Gamma_{V \to l^+ l^-} = \frac{\pi \alpha^2 e_q^2 F_V^2}{3M_V} (1 + 2r^2) \sqrt{1 - 4r^2}, \qquad (1)$$

where  $\alpha$  is the fine structure constant,  $e_q$  is the quark charge in units electron charge and r stands for the outgoing lepton and vector meson mass ratio  $r = m_l/M_V$ . The constant  $F_V$  is defined by the matrix element of the electromagnetic current

$$<0|j^{\mu}(0)|V(Q,\epsilon)>=F_V M\epsilon^{\mu}$$
<sup>(2)</sup>

with j being the electromagnetic current associated with quark content of the meson with polarization  $\epsilon$ .

For mesons where the mixing can be ignored or idealized, the electormagnetic charge factorize out like in the Eq.(1) and the leptonic decay can be calculated from a simple formula :

$$F_{Y(n)}M_{Y(n)} = \frac{Z_2}{3}Tr_{CD} \int \frac{d^4q}{(2\pi)^4} [\gamma_\mu S_f(q+Q/2)\Gamma_V^\mu(q,Q)S_f(q-Q/2)]$$
(3)

where  $S_f$  stands for given flawour quark propagator. The formula (3), as has been suggested in [39] and used in [24] for the first time, assumes that meson pole part does not enter in QED Ward identity,  $\Gamma_V^{\mu}$  is thus fully transverse Bethe-Salpeter meson vertex function satisfying  $\Gamma Q = 0$ .

The leptonic decays are hard to measure experimentaly and they have been determined for ground states and few narrow mesons [40]. No  $f_V$  has been measured and reported for excited light vectors.

# 3 The BSE for quarkonium

To obtain vertex functions for excited quarkonia we use the BSE form as elaborated in [38]. The heart of the interaction kernel is based on lattice and DSEs solutions in the Landau&Feynman gauge, however here we vary the parameters more freely in order to get more unified snapshot of vector  $1^{--}$  mesons. Like in [38] the BSE is solved in the Feynman gauge with infrared regulated gauge term.

In addition to existing functional form of interacting kernel [38] we introduce separable effective coupling behaving like  $\simeq 1/ln^2q^2$  at large loop momenta, which allows to put renormalization constant  $Z_2 = 1$  in the Eq. (3), as it should be for CQA. Since we solve the full two dimensional BSE without using expansion to Tchebychev polynomials (see the discussion in [24]), no UV regulator is required to subtract UV divergences. They are not presented here. For completeness we write down the BSE kernel at this place:

$$K(k,p,Q) = \frac{4}{3} \Gamma^{\mu}(k,Q) \times \gamma^{\nu} \left[ \left( -g_{\mu\nu} + \frac{q_{\mu}q_{\nu}}{q^2} \right) G(q^2) - \frac{q_{\mu}q_{\nu}}{(q^2 - m_r^2)^2} \right]$$
(4)

with q = k - p and Q is the total momentum of the quark-antiquark pair. The transverse part is taken as following

$$G(q^2) = \frac{r_1}{q^2 - m_{g1}^2} - \frac{r_2}{q^2 - m_{g2}^2}.$$
 (5)

These two real poles appearing in the kernel are an imperfect approximation of continuous gluonic spectral function [42, 41] and it makes the problem numerically manageable. It appears in the product with the following effective quark-gluon vertex  $\Gamma^{\mu} = \gamma^{\mu} \alpha(0) \frac{\ln^2(\zeta)}{\ln^2(\zeta+k^2/\Lambda^2)}(1+\delta)$ , where  $\delta$  is flawour independent for ground stares but flawour dependent part of the BSE kernel otherwise. This small fluctuation reads  $\delta = \delta_0 \sin\left(\left[\sqrt{Q^2} - M_V(1)\right]\frac{2\pi}{E_p}\right)$ .

$m_u$	$m_s$	$m_c$	$m_b$
0.29	0.522	1.49	4.709

Table 1: Constituent quark masses (in GeV) for excited vectors.  $m_u$  stands for up and down quark masses in the isospin limit.

Numerical values for parameters are taken as  $\zeta = 100$ ;  $\Lambda = 0.4 GeV$ ;  $\delta_0 = 0.15$ ;  $E_p = 1.13 GeV$  and M(1) stands for the ground state mass in given spin/flawour channel, i.e.  $M_{J\psi}$  and  $M_{Y(1)}$  for heavy quarkonia, and  $M_{\phi} = 1019 MeV$  $M_{\omega} = 782 MeV$  for the strange and the light quark sector.

The infrared regulator  $m_r = 0.113 GeV$  has been taken identical to the first pole position  $m_{g1}$ , defining thus the smallest scale allowed by our consideration. It prevents the gauge term from being too large, which would generate unphysical solutions otherwise (continuous and abnormal solutions). The position of the second pole is taken to be  $m_{g2} = E_p = 1.13 GeV$  for simplicity.

The constant mass  $m_f$  appears in the constituent quark propagator:

$$S_f(q) = [\not q - m_f]^{-1} \tag{6}$$

and is what constitutes the CQA for given flawour f. Note, beyond CQA the quark mass is scale changing, similarly like the coupling does. In addition, the CQA ignores momentum dependence of the renormalization wave function  $Z_f$ .

Numerical value of  $m_f$  is tuned in order to get nice description of excited states. The ground state is handled separately in order prevent unphysical states at each flawour channel.

The CQ masses we use for this purpose are listed in the Tab 1.

We use two component approximation  $F_1, F_5$ , not including the components which are responsible for D- wave orbital momentum. We anticipate already here, the P-wave admixture due to the  $F_5$  is small, reducing  $F_V$  about few percentage. The leptonic decays as well as the n.c. is calculated in  $F_1 + F_5$ and  $F_1$  approximation respectively, but shown for the later case assuming that  $f_V$  are more close to the case when complete set of components is taken into account.

#### **3.1** Abnormal states

When deriving the BSE and its normalization, one assumes there is a real pole in the S-matrix associated with a free move of the center mass of the meson. In meson case, we thus neglect its decays. It also ignores the fact, that the lifetime should be much longer then typical interaction time and (in addition to other shorcomings) it can lead to occurrence of abnormal state solutions in BSE treatment of unstable mesons. Obviously, the canonical Nakanishi normalization condition for BSE does not fit a broad resonances. As a consequence a negative norm solutions of BSE can be seen in addition to normal ones and they call for negative residuum as was first noticed in the scalar toy model [43] (without relating this phenomena to stability of the bound states).

We should mention that there are more ways to scrutinize the theory and generate unphysical states. To name it : too much strong renormalized coupling, when ignoring of dressing of some important Green's function (see [44]), etc. etc... However, we argue here that in QCD (and not only there), the abnormal solution of BSE can appear as a consequence inappropriate use of the homogeneous BSE for description of resonances.

In particular case of presented study, the appearance of abnormal states is slightly more intriguing since ghosts solutions can be further supported by the CQA itself. However as we will clarify in the next section, the appearance of single ghost solution can be prevented by the use of more - in fact scale dependent- values for constituent quark mass. While we do not fot oher spins, but for  $1^{--}$  neutral states the single ghost appearance seems to prevented by taking by two different masses: one for the ground state and the second one for the rest of excited states. Actually, we have found technically more easy to calculate all excited states in the first stage and then tune the CQ mass to get a ground state without presence of ghost. Of course, this is nothing else but the evidence that the dressing of constituent quark propagator is important and should not be neglected.

For purpose of presented CQA BSE study, we numerically identify all normal and perhaps all abnormal states at every flawour channel, and evaluate the leptonic decay  $F_V$  for both. We use canonical positive (and negative) residuum normalization condition to normalize BSE solution for mesons (ghosts). WE get rid of appearance of single ghost states by definining which CQ mass should be used.

However, doing so, there can be identified further semi-pathological or at least confusing structure. In some circumstances, the normal-abnormal (NA) states appear (nuemrically nearly) degenerated. The pole of (NA) pairs cancel each other. Interestingly, when comparing with experimental data, then they appearance suggests that these pairs are an artefact of ignored coupling to decay products of resonances, i.e. they seems to be a prelude of the presence of the complex cut in more elaborated approximations. To author best knowledge, the appearance of ghosts is also proper to MTM, however the observation of pairing of normal-abnormal states has been either overlooked or it does not happen in the MTM due to the very different analytical structure. Furthermore, there is no other known, already solved BSE solutions for vector meson available on the market.

At last but not at least we should stress, that all abnormal states which comes as a numerical solution for vector mesons in presented study can be normalized to -1 residuum, i.e. they would appear as a negative poles in the

name:	Y(1)	Y(2)	Y(3)	Y(4)	Y(5)
this work	9.460	$9.914 \pm 10$	10.36	$10.530\pm30$	$10.740\pm30$
Experiment	9.460	10.23	10.355	10.579	10.900
BSE [26]	9.490	10.089	10.327		
Salpeter [18]	9.461	10.020	10.363	10.622	10.835

Table 2: Bottomonium: Comparison of calculated masses with experiment.

four point quark-antiquark functions of the form

$$-\frac{\Gamma(p,P)\Gamma(p',P)}{P^2 - M^2 + i\epsilon}$$
(7)

keeping thus the normalization condition for BSE ghosts identical (up to the sign) to ordinary states. We will call these negative norm state abnormal or ghosts, although they have identical quantum numbers to normal states. These states are not an excitations in relative time (as those in the Wick-Cutkosky model [43]) and should not be confused with them.

To clarify motivations for solutions presented in the next section, the author regards a single located abnormal state as badness of the model approximation (say inside the BSE)and we try to prevent them. On contrary, pairwise appearance with meson states can regarded as consequence of not wise application of the homogeneous BSE, but does not need to be regarded as inherent pathology due the cancelation in the S-matrix. The position of these poles are not a random we list these cases in the next section for sake of completeness.

# 4 Results for heavy quarkonia

The spectroscopy and lepton decay constants as obtained from our BSE solution are shown in the tables 2 3 for bottomonia and 5 for charmonia respectively, where we compare with the PDG data and several selected theoretical approaches in special case of bottomonium.

To compare with further studies, the value for  $M_{Y(1)} = 9.488 GeV$  was obtained in [25] in MTM, in which model a similar calculation was extended to excited states in [26, 27]. Recall that MTM for heavy quarkonia used in [27] differs from [26] as well as the parameters of MTM for light mesons usually differ from those used to evaluate properties of heavy mesons. It was also shown in [30] that the a flawour non-universal LR interaction is needed to describe leptonic decay constant of heavy quarkonia in MTM. A large uncertainty of MTM [28] in determined  $f_V$  as shown in [28] are due to numerical extraction of the bottom propagator in MTM.

name:	Y(1)	Y(2)	Y(3)	Y(4)	Y(5)
this work	150	$210\pm10$	$120\pm5$	$110\pm3$	$100 \pm 3$
Experiment	715	497	430	341	
BSE [28]	707	393	9(5)	20(15))	
SE [18]	702	503	432	387	

Table 3: Bottomonium: Comparison of calculated leptonic decays  $F_Y$  (in MeV) with experiments

normal	9744	9990	10346
abnormal	9793	10001	10363
$f_V(\text{normal})$	131	166	
$f_V(abnormal)$	152	346	

Table 4: Self-excluded states. Shown are masses for normal (first line) and abnormal states (second line) and their leptonic decay constants. Normalization to negative unit residuum is used in order to get  $f_V$  for abnormal states.

Within 5 - 7MeV step we have carefully scan the masses above 9 GeV (and stopped at 11 GeV) and obtained all states of BSE under consideration (it is costly procedure, the inspection of single point takes nearly 3 hours). We briefly describe our founding in the following text: All narrow experimental states Y(1) - Y(3) match with some ordinary state given by the solution of our considered BSE (see table 2)

In adition we list NA pairs. Numerical values for masses of normal and abnormal state lie very close to each other, hence they degenerate within the numerical errors. Three pairs of such solutions we have found are listed in the Table 4. The first NA pair appears below open bottom threshold, hence it should be viewed as a mild imperfection of BSE kernel when applied to bottomonia.

Salpeter equation is the known approximation of the BSE, where various complications presented in more complex BSE solutions are circumvented. While the boosting to moving frames is complicated (if possible at all), 3d reduction is particularly meaningful to evaluate static properties for bottomonium system and we provide the comparison with the study [18] in the tables (10-20 % errors are not shown).

The masses for excited 1 - S-states charmonia as calculated and compared to PDG data at the table 5. We do not compare with others, the reader can compare with [18] for Salpeter equation, [21] for Gross equation as well as with number of BSE studies [26, 27, 28, 30].

name:	$J/\psi$	$\psi(2s)$	$\psi(3s)$	$\psi(4s)$	$\psi(5s)$
M (calc)	3097	$3650\pm10$	$4100\pm15$	$4210\pm20$	$4350\pm30$
M (Exp)	3097	3686	4039	4153	4421
$f_V$ (calc)	280	$120\pm2$	250	$290\pm20$	$226\pm10$
$f_V (\exp)$	416	308	187	146	_
Salpeter [18]	573	424	351	_	321

Table 5: Charmonia: Comparison of calculated masses (in MeV) and leptonic decays with experiment.

Several comments is suited at this place. There is a single normal state at 3650 MeV corresponding to the first excited state  $\psi(2s)$  to which the model was originally tuned. Then there is AN pair at 4100. Since these two states have different  $f_V$  we keep the normal state in the Table. All other states are normal, there is no other abnormal state seen above 4 GeV, however all states appear twice, i.e. there are two close poles- two very close solutions of BSE. The difference of masses is much smaller then width of experimentally observed counter-partners, thus the pairwise appearance should be regarded as the single solution and the difference as the systematical error related to our approach. These errors go hand by hand with the error  $\sigma^2 \simeq 10^{-7} - 10^{-4}$  achieving in our iteration procedure (the reduction of these errors would be painfully costly, note the same level of degeneracy is seen for AN pairs). For purpose of completeness we list more precise positions of all doublings observed to 5 GeV in the list in the Appendix. The ground state is evaluated for separatele since for different values of the quark mass quoted bellow.

# 5 Light vectors and their excitations- further rise and fall of abnormal states

The consideration and calculation methods can be extended to the light vector mesons, e.g. to  $\phi(1020)$  and its excitations modeled by strangeonium and  $\omega/\rho$  system for which we assume ideal mixing , i.e.  $\omega = \frac{1}{\sqrt{2}}[uu + dd]$ . For both systems we compare with collected PDG data as have been presented in decade 2010 to 2020. Since the LRA does not distinguish between isospin partner  $\rho$  and  $\omega$  and since  $\omega$  is the narrow state, we use the mass of omega in the phase factor in the g function.

Recall here, most information on light vector resonances come from meson production in  $e^+e^-$  anihilations into a few mesons. Related measured cross sections always involve peak of  $\omega$  and  $\phi$  ( $\rho$ ) for odd (even) g-parity channel, while several broad and overlapping resonances are used for for phenomenological description (and have been listed in the PDG) for moderately larger energy.

The BSE solutions for strangeonium are presented in in Tabules 6 and 7. The main feature is that the only state which is iniquely described is the ground

Exp. mass	$\phi(1020)$	X(1575)	$\phi(1680)$	$\phi(1750)$	$\phi(2170)$	
this work	1019	1588	1960	2190	2327	2481
$f_V(n)$	230	80	119	147	166	190
abnormal	_	1582	2056	2172	2302	2449
$f_V(a)$	_	84	135	151	173	204

Table 6: Strangeonium masses compared against PDG list of excited  $\phi$  (and one X ) mesons. Negative norm/residuum states and their decay constants are displayed in 4-th and 5-th line respectively. Ground state is solved separately as explained in the text.

Exp. mass	$\rho(770)$	$\rho(1450)$	$ \rho(1570) $	$ \rho(1700) $	$ \rho(2150) $
Exp. m. (singlet)	$\omega(782)$	$\omega(1420)$	$\omega(1650)$	X(1750)	$\omega(2205)$
this work	782	1400	1564 & 1650	1750 & 1881	2015
$f_V$	*	$109\pm20$	$107 \pm 30$	$135\pm20$	$160\pm30$

Table 7: Masses of  $q\bar{q}$  (third line) compared to masses of light 1<sup>--</sup> resonances as listed in PDG lists 2012-2020 (the first and the second line). Calculated decay constants values are shown in the forth line. Negative norm states are not displayed and the ground state is calculated for different constituent propagator as explained in the text.

state. All other states come with their abnormal brothers. We list AN pair only for strangeonium and do not list them for  $\omega/\rho$  system. There are two more pairs appearing at 1260 MeV and 1320 MeV energy, they disapper by construction when smaller value mass is allowed as will be done for the ground state. We do not write down all step functions that would prevent ghost, noting the CQA is not fully adequate for the light mesons due to other known reasonings.

The optimistic feature of BSE is that it provides (with aforementioned constrain) one to one correspondence to PDG listed meson mass values. Not appealing feature is that the pole of the normal state solution is canceled against. If they cancel completely (which we cannot confirm) then there will be nothing controversial left with the appearance of ghosts in the BSE solutions for vector mesons.

### 5.1 Getting rid of abnormal states

Up to one case, no single ghost appears among excited states. Hence to prevent them completely means to find different CQ mass value or perhaps a different constant  $Z_f$  which prevents ghost near the ground state. Varying just the mass we have found this is possible for bottomonia and for  $\phi$ , albeit the leptonic decay is not ideal. We do not fine tune the model to get best agreement with the experiment, leaving this comparison postponed after solving DSE and BSE simultaneously. We just mention that the CQ does exist and provide example. The leptonic width calculated in all tables represent results for our simple search. To get the  $\omega$  in given model and  $J\Psi$  without ghost nearby one needs to use  $Z_f$ as well. It is equivalent to slight softening of the effective coupling in this case.

Even in the case when the change in one constant is enough to prevent gost, the solution is not without ambiguity. Keeping the mass for the ground state fixed, we have found at least two possible values CQ masses for bottomonium 9460, the first value is  $m_b^{Y1} = 4.42 GeV$  and the second for which we get the bound state mass as well is  $m_b^{Y1} = 3.37 GeV$ . The solution with smaller mass offers the decay  $f_V = 150 MeV$  as a result, being still far from the experimental value, but nevertheless more suited for this.

Not the same happen to charmonium. While we get rid of ghost solution that we would face otherwise at 3150 MeV, then this is replaced by NA pair both components weighting 3097 MeV for a constituent quark mass  $m_c^{J/\Psi} = 1.263 GeV$  for normal state and  $m_c^{J/\Psi} = 1.266 GeV$  for the gost. To prevent the ghost one can suppres the coupling by taking  $Z_f$  into the game. We do not quote not unique value of our searches here, noting all we see has underestmated  $f_V \simeq 100 MeV$ .

In case of strangeonium, scanning the CQ mass for  $\phi$  meson , while keeping BSE interaction intact, we get more eigenvalues as well. The first is ghost which appears for CQ mass 395MeV then the pair of normal solutions appear for  $m_s = 342MeV$ , 343, 5MeV providing almost identical leptonic decay constant  $f_V = 232MeV$  and  $f_V = 234MeV$  respectively. The BSE kernel is enough to prevent the ghost, no change in renormalization function is needed.

To close this section we find the solution for physical mass of  $\omega$ , providing the quark mass should be taken slightly bellow 290 MeV. To get the stable result, change in  $Z_f$  is needed in this case and as there are more solutions providing more values for  $f_v$ , we leave this entry as unanswered in CQA.

# 6 Conclusion

We calculated mass spectrum and leptonic decays constant of flawour-less vector mesons by using CQ LRA BSE in the Feynman gauge. There is a striking difference between description of heavy quarkonia and the excited mesons made out of the light (anti)quarks. In later case, the solution for normal excited state comes regularly in pair with abnormal state. Obtaining a good (normal) solution for the narrow ground state and facing the cancelation between normal and abnormal state in these pairs, we argue this is a due to the inconsistent use of the BSE for description of broad resonances.

On the other side, admitting the fact that the NA pairs are an artifact of not a wise use rather then bad approximations inside the BSE, one gets remarkable description of excited heavy quarkonia.

Further shortcoming of single mass value CQ BSE is either bad agreement

with experiment of the appearance of uncanceled ghost at vicinity of the ground state. These ghosts have no good meanings, and they can be cured by using slightly different CQ mass in the propagators of the BSE. In case of heavy quarkonia, the values of CQ masses used for excited states and ground states calculation turns to be proportional to the ratio

$$\frac{m_c^{ground}}{m_c^{excited}} = \frac{V(1)}{V(2)} \tag{8}$$

which is prelude for running mass corrections.

Actually, we expect there will be no single ghost state in more consistent DSE/BSE treatment. On the other side, we expect an AN pair can appear in principle even in more advanced approximations.

On the other side using a single interaction kernel in BSE description of variety mesons is mesmerizing. We expect, after taking dressing of propagators into account we will achieve good description in other spin channels as well. Such and similar advanced approaches [45, 46] describe resonances in given exclusive process. They offer alternatives to popular (but more involved) coupled channel analyses (see for instance [47, 48]) without the use of effective models but within gluon and quarks degrees of freedom.

# A Normalization of BSE

Performing trivial integrations, making a trace, the normalization used for purpose of our numerics reads:

$$\pm 1 = \frac{N_c}{2\pi^3} \int_{-\infty}^{\infty} dq_4 \int_0^{\infty} dq_s q_s^2 I$$

$$I = \frac{q_4^2 - \Delta}{[(q_4^2 + \Delta)^2 + q_4^2 M_V^2]^2} [F_1^2 (3m_f^2 + \frac{3}{4}M_V^2 + q_E^2 + 2q_4^2)$$

$$+ q_s^2 F_5^2 (q_E^2 + \frac{M_V^2}{4} - m_f^2) - 4mq_s^2 F_1 F_5]$$

$$- \frac{1}{(q_4^2 + \Delta)^2 + q_4^2 M_V^2} (\frac{3}{2}F_1^2 + \frac{1}{2}F_5^2 q_s^2) + \dots$$

$$(9)$$

where ellipses stands for less then 1% contribution due to the derivative of the interacting kernel and where  $q_E^2=q_4^2+q_s^2$  and

$$\Delta = q_s^2 + m_f^2 - \frac{M_V^2}{4} \,. \tag{10}$$

where plus and minus stands for the normal and abnormal state respectively. In shallow bounded limit  $2m_c \to M$  the denominator is small but the  $q_4^2 - \Delta$  can abruptly change the sign. The result depends on a way how it is weighted by  $F_1^2$ . An interplay decides between normal and abnormal state.

М	$4084^g; 4100$	4200; 4226	4341; 4368	4495; 4527	4670; 4702	4865; 4899	5089; 5119
F	$350^g$ ; 250	309;270	230; 223	186; 181	162;160	140; 144	132;130

Table 8: Eigenvalues of the BSE and leptonic decays as obtained from the BSE for vector 1S charmonia in details. g stands for negative norm state.

A large number of integration points 288 \* 144 was used to integrate 2d BSE. Large number of mesh points is also needed to eliminate fake solutions, which do appear otherwise.

# B Detailed list of numerical results for charmonia

Observed pairs of solution for BSE for charmonia are listed in the table 8. Only the first one stand for NA pair, all others are ordinary states. Assuming exact degeneracy, the systematical error is determined form the mass difference for each pair, hence not worse that 1%. Leptonic width are more sensitive, mainly to the integral error represented by the norm. In few cases we have used 20GeV UV cut off to prevent from numerical noise at UV tails of vertex functions.

# References

- [1] C. Quigg, J. L. Rosner, Phys. Lett. B **71**, 153-157 (1977).
- [2] E. Eichten, K. Gottfried, T. Kinoshita, K. D. Lane, and T. M. Yan, Phys. Rev. D 21, 203 (1980).
- [3] N. Brambilla, A. Pineda, J. Soto, A. Vairo, Rev.Mod.Phys. 77 1423 (2005).
- [4] J. Segovia, P. G. Ortega, D. R. Entem, F. Fernández Phys. Rev. D 93, 074027 (2016).
- [5] N. R. Soni, A. N. Gadaria, J. J. Patel, J. N. Pandya, Phys.Rev.D 102 1, 016013 (2020).
- [6] B. Azhothkaran , V.K. Nilakanthan V. K., Int.J.Theor.Phys. 59 7, 2016-2028 (2020).
- [7] R. Chaturvedi, A. K. Rai, Eur.Phys.J. A 58 11, 228 (2022).
- [8] N. Brambilla, H. S. Chung, A. Vairo, X.-P. Wang, JHEP 03 (2023).
- [9] N. Brambilla, M. A. Escobedo, A. Islam, M. Strickland, A. Tiwari, A. Vairo, P. V. Griend, Phys.Rev.D 108 1, L011502 (2023).

- [10] B. Assi, M. L. Wagman Phys.Rev.D 108 9, 096004 (2023).
- [11] Christas Mony A., R. Dhir, J. Phys. G; Nucl. Part. Phys. 51 115004 (2024).
- [12] N. Kivel, JHEP **07** 065 (2024).
- [13] P.G. Ortega, D.R. Entem, F. Fernández, J. Segovia, Phys. Rev. D 109 (2024).
- [14] U.V.N. Kanago, A.A. Likéné, J.M. Ema'a Ema'a, P.Ele Abiama, G.H. Ben-Bolie, Eur. Phys.J. A 60 3, 47 (2024).
- [15] Z.Zhao et all, Phys.Rev.D **109** 1,016012 (2024).
- [16] J.-K. Chen, Eur. Phys. J. C78 3, 235 (2018).
- [17] E. Gebrehana, S. Bhatnagar, H. Negash, Phys. Rev. D 100, 054034 (2019).
- [18] G.Wang and Xing-Gang Wu, Chinese Phys. C 44 063104 (2020).
- [19] V. Guleria, E. Gebrehana, S. Bhatnagar Phys.Rev.D 104 9, 094045 (2021).
- [20] S. Leitão, Y. Li, P. Maris, M. T. Peña, A. Stadler, J. P. Vary, E. P. Biernat Eur.Phys.J.C 77 (2017) 10, 696
- [21] S. Leitão, A. Stadler, M. T. Peña, E. P. Biernat, Phys. Rev. D 96 7, 074007 (2017).
- [22] R.N. Faustov, V.O. Galkin, X.-W. Kang Phys. Rev. D 101 1, 013004 (2020).
- [23] N. Souchlas, Phys. Rev. D 81 114019 (2010).
- [24] P. Maris and P. C. Tandy, Phys. Rev. C 60, 055214 (1999) [nucl-th/9905056].
- [25] M. Blank and A. Krassnigg, Phys. Rev. D 84, 096014 (2011).
- [26] Ch. S. Fischer, S. Kubrak, R. Williams, Eur. Phys. J. A 51 10 (2015).
- [27] T. Hilger, C. Popopovici, M. Gomez-Rocha, A. Krassnigg, Phys.Rev.D 91 3, 034013 (2015).
- [28] A. Krassnigg, M. Gomez-Rocha, T. Hilger, J.Phys. Conf. Ser. 742, 1 012032 (2016).
- [29] A. Krassnigg, T. Hilger, J. Phys. Conf. Ser. 1024 1, 012010 (2018).
- [30] F. F. Mojica, C. E. Vera, E. Rojas, B. El-Bennich, Phys. Rev. D96, 1 014012 (2017).
- [31] Heavy-light mesons from a flavour-dependent interaction, F. Gao, A. S. Miramontes, J. Papavassiliou, J. M. Pawlowski, arXiv:2411.19680.

- [32] Yin-Zhen Xu, JHEP 07, 118 (2024).
- [33] Andreas Windisch, talk at EXCITEDQCD2017, 8.5. Sintra, Portugal: The analytic properties of the Landau gauge quark propagator Dyson-Schwinger equation at rainbow level.
- [34] S. M. Dorkin, L. P. Kaptari, and B. Kämpfer, Phys. Rev. C 91, 055201 (2015).
- [35] J. Horak, J. Papavasiliou, J.M. Pawlowski, N.Wink, Phys. Rev. D104, 074017 (2021), [arXiv:2103.16175[hep-th]].
- [36] J. Horak, J.M. Pawlowski, N.Wink, SciPost Phys. 15,149 (2023).
- [37] V. Sauli, Phys. Rev. D 102, 014049 (2020).
- [38] V. Sauli, arXiv:2410.10625 [hep-ph].
- [39] M. A. Ivanov , Yu. L. Kalinovsky , C.D. Roberts, Phys. Rev. D 60, 034018 (1999).
- [40] KLOE Collaboration, F. Ambrosino, Phys. Lett. 608, 199-205 (2005).
- [41] V. Sauli, Phys. Rev. D 106 9, 094022 (2022) [arXiv: 2011.00536[hep-lat]].
- [42] J. Horak, J. M. Pawlowski, J. Rodríguez-Quintero, J. Turnwald, J. M. Urban, Phys. Rev.D 105, 3 03601 (2022).
- [43] N. Nakanishi, Phys. Rev. **138** 5B, 1182 (1965).
- [44] S. Ahlig, R. Alkofer, Annals Phys. 275, 113-147 (1999).
- [45] V. Sauli, Phys. Rev. D 106 3, 034030 (2022).
- [46] A.S. Miramontes, A. Bashir, Phys. Rev. D 107 1, 014016 (2023).
- [47] N. Hüsken, R. E. Mitchell, E. S. Swanson Phys.Rev.D 106 (2022) 9, 094013 (2022).
- [48] W-H. Liang, N. Ikeno, E. Oset, Phys. Lett. B 803, 135340 (2020).