HOMOLOGY INCLUSION OF COMPLEX LINE ARRANGEMENTS

ADRIEN RODAU

ABSTRACT. We introduce a new topological invariant of complex line arrangements in \mathbb{CP}^2 , derived from the interaction between their complement and the boundary of a regular neighbourhood. The motivation is to identify Zariski pairs which have the same combinatorics but different embeddings. Building on ideas developed by B. Guerville-Ballé and W. Cadiegan-Schlieper, we consider the inclusion map of the boundary manifold to the exterior and its effect on homology classes. A careful study of the graph Waldhausen structure of the boundary manifold allows to identify specific generators of the homology. Their potential images are encoded in a group, the graph stabiliser, with a nice combinatorial presentation. The invariant related to the inclusion map is an element of this group. Using a computer implementation in **Sage**, we compute the invariant for some examples and exhibit new Zariski pairs.

1. INTRODUCTION

The study of the topology of plane algebraic curves was initiated by O. Zariski. The *combinat*orics of a curve is the topological type of the pair formed by a tubular neighbourhood of the curve and the curve itself. These data are equivalent to the topological type of the singularities and the incidence relations between the components. O. Zariski [Zar31; Zar37] and E. R. van Kampen [Kam33] have shown that there exists pairs of curves with the same combinatorics but different embeddings in \mathbb{CP}^2 , which were dubbed Zariski pairs by E. Artal in [Art94].

Line arrangements are finite collections of complex lines in \mathbb{CP}^2 , that is, plane algebraic curves whose irreducible components have degree one. The components are non-singular and the singularities all belong to a same class of simple type. The combinatorics of a line arrangement depend only on the incidence relations, which can be encoded in a graph Γ called the *incid*ence graph. The study of line arrangements provides a favourable setting to create topological methods and invariants that could then be extended to algebraic curves in general. It also offers some interesting questions in itself. The first Zariski pair of line arrangements was constructed by G. Rybnikov [Ryb11] and studied in detail by E. Artal, J. Carmona, J. I. Cogolludo and M. Á. Marco in [Art+06] using the fundamental groups of the complements. The existence of this pair showed that the combinatorics does not determine the topological type of a curve even in the simplest case of line arrangements. The search for more Zariski pairs and a finer comprehension of the relationship between combinatorics and topology of curves and line arrangements has been a very active topic since the 2000s. Notably, S. Nazir, M. Yoshinaga [NY12] and F. Ye [Ye13] have completely determined the isotopy classes of line arrangements up to 9 lines, and in particular that no Zariski pairs exist for these values. We also mention the works of B. Guerville-Ballé [Gue16] and J. Viu Sos [GV19] who discovered several new Zariski pairs with more than 11 lines. Readers can refer to [ACT08; Gue22] for a more detailed review of the subject.

A wide variety of common invariants from algebraic topology have been applied to the study of line arrangements and Zariski pairs. Consider the exterior $E_{\mathcal{A}}$ of the arrangement \mathcal{A} in \mathbb{CP}^2 . A direct comparison of the fundamental groups of the exteriors using the Zariskivan Kampen method [Kam33] can sometimes give a Zariski pair, as for the original example

²⁰²⁰ Mathematics Subject Classification. 32S22, 52C35, 57M05.

of G. Rybnikov. However, there are known examples of Zariski pairs where the fundamental groups of the exteriors are isomorphic (see [Shi09; Shi19; ACM19a; Gue20]). It has since been shown that not even the characteristic varieties determine the topology of line arrangements in general, including for the subcategory of arrangements with real equations.

A possible approach to build invariants of line arrangements is to consider the boundary manifold $B_{\mathcal{A}} := \partial E_{\mathcal{A}}$ of the arrangement. T. Jiang, S. S.-T. Yau [JY93] and E. R. Westlund [Wes97] have shown that this boundary manifold has the structure of a graph manifold as defined by F. Waldhausen [Wal67a; Wal67b] and W. D. Neumann [Neu81]. The incidence graph Γ provides a 'blueprint' to reconstruct the boundary manifold by gluing together circle bundles (Seifert pieces) corresponding to the boundary of local neighbourhoods around each line component and each singularity of the arrangement.

Our own interest lies in the study of the inclusion

$$i_{\mathcal{A}}: B_{\mathcal{A}} \hookrightarrow E_{\mathcal{A}}$$

of the boundary manifold inside the exterior of a line arrangement \mathcal{A} . E. Hironaka [Hir01] studied the morphism induced by the inclusion on the fundamental groups for the case of real line arrangements, using methods to compute the presentations of the groups due to W. Arvola [Arv92] for the exterior and E. R. Westlund [Wes97] for the boundary. This study was continued and generalised to complex line arrangements by V. Florens, B. Guerville-Ballé and M. Á. Marco in [FGM15]. Their results exposed that the study of the fundamental group 'inclusion' is made difficult by the algebraic complexity of the morphisms involved. Another approach is to consider the morphism induced by the inclusion $i_{\mathcal{A}}$ on the *first homology groups* instead:

$$i_{\mathcal{A}}^*: H_1(B_{\mathcal{A}}, \mathbb{Z}) \longrightarrow H_1(E_{\mathcal{A}}, \mathbb{Z})$$

Unlike what one might think, this is not a trivial matter. The study of that morphism was first considered by E. Artal, V. Florens and B. Guerville-Ballé in [AFG17] in the form of the \mathcal{I} -invariant, which they used to obtain new Zariski pairs. This invariant was later generalised as the *loop-linking number* of W. Cadegan-Schlieper [Cad18] and further developed by B. Guerville-Ballé in [Gue22].

We propose a new invariant that extends these constructions and fully exploits the homology 'inclusion' of the boundary manifold of any line arrangement inside its exterior. The main principle of the construction is as follows. The first homology groups of the boundary manifold and the exterior are both combinatorially determined. However, the induced morphism $i_{\mathcal{A}}^*$ still contains significant topological information. The main difficulty of describing $i_{\mathcal{A}}^*$ lies in the ambiguity of defining a set of generators on the boundary. Such a set can be obtained from a specific class of embeddings of the graph inside the boundary manifold. These graphed embeddings depend on a graph ordering Ω . We then define the graph stabiliser as a combinatorial group which computes the homological differences between all ordered graphed embeddings. This allows to remove the ambiguity on the generators and is the main novelty of our construction. The morphism $i_{\mathcal{A}}^*$ induces an element of the graph stabiliser called homology inclusion which is a topological invariant of ordered oriented line arrangements. Using a computer program written in **Sage** [Sag23] in collaboration with B. Guerville-Ballé and E. Artal, we have computed homology inclusion values and obtained a new Zariski quadruplet of oriented line arrangements with 11 lines. In particular, four pairs within the quadruplet are Zariski pairs.

Section 2 gives a presentation of line arrangements, with a specific focus on combinatorics and related concepts, in particular the orderings which are key in the construction. In Section 3 we recall the structure of the boundary manifold as a graph manifold. In Section 4 we present the *flat stars* which are the 'elementary bricks' used to build the graphed embeddings, which are themselves defined in Section 5. In Section 6, we define the graph stabiliser group and the homology inclusion invariant. Section 7 is dedicated to determining of a presentation of the graph stabiliser using results from Sections 4 and 5. This presentation is then used in Section 8 to establish that the homology inclusion extends the loop-linking number. Finally Section 9 summarises the results of the computations made with the invariant and presents the new oriented Zariski quadruplet.

The computation of the homology inclusion makes use of specific results and algorithms which are not immediately necessary for its definition. We therefore delegate the presentation of the details of this computation to another publication.

In addition, we mention that the homology inclusion invariant represents a preliminary step in the study of the *twisted* homology inclusion of complex line arrangements. Consider the morphism induced by the inclusion $i_{\mathcal{A}}$ between the *twisted* homology groups of the boundary and exterior manifolds. To compute the twisted homology group of $B_{\mathcal{A}}$ one needs to restrict abelian representations of the fundamental group of the exterior to the boundary. The homology inclusion precisely allows to compute these values. The twisted homology inclusion invariant could potentially be used to discriminate equality cases of the homology inclusion invariant and thus detect more Zariski pairs. This new invariant will be the object of a future publication.

The author would like to thank his former PhD advisors Enrique Artal and Vincent Florens for their support and reviewing of this research. Many thanks also to Benoît Guerville-Ballé for his invaluable contribution to the computations of the invariant.

2. Combinatorics of complex line arrangements

2.1. Generalities. A complex line arrangement \mathcal{A} is a union of complex lines on the projective complex plane \mathbb{CP}^2 . The intersections are called the *singular points*.

Definition 2.1. A *topological equivalence* between two complex line arrangements \mathcal{A} and \mathcal{A}' in \mathbb{CP}^2 is an homeomorphism

$$\Phi: (\mathbb{CP}^2, \mathcal{A}) \xrightarrow{\sim} (\mathbb{CP}^2, \mathcal{A}') \qquad \diamondsuit$$

If Φ respects the orientation of all the components of \mathcal{A} then the equivalence is *positive*, or *oriented*. A topological equivalence between two arrangements induces a bijection between the sets of lines and between the sets of singular points. This fact is made precise using the concept of *combinatorics*.

Definition 2.2. A *line combinatorics* is a triple $C = (\mathcal{L}, \mathcal{Q}, \in)$ where \mathcal{L} is a finite set, \mathcal{Q} is a finite subset of $\mathcal{P}(\mathcal{L})$ and \in is a relation from \mathcal{Q} to \mathcal{L} such that:

- (i) For every element $P \in \mathcal{Q}$, there exist at least two distinct elements $L, L' \in \mathcal{L}$ such that $P \in L$ and $P \in L'$.
- (ii) For every pair $L, L' \in \mathcal{L}$ with $L \neq L'$ there exists a unique $P \in \mathcal{Q}$ such that $P \in L$ and $P \in L'$.

For a line arrangement \mathcal{A} , there is a natural line combinatorics $C_{\mathcal{A}}$ associated with \mathcal{A} , given by the incidence of the lines. Combinatorics can also be encoded with graphs. Let $\mathcal{Q}^{>2}$ be the subset of singular points with multiplicity > 2.

Definition 2.3. The *incidence graph* $\Gamma(C)$ of the combinatorics C is defined by the following description:

Vertices: one vertex for each element of $\mathcal{L} \cup \mathcal{Q}^{>2}$.

Edges:

(i) $L \in \mathcal{L}$ and $P \in \mathcal{Q}^{>2}$ are linked by an edge $e_{P,L}$ if and only if $P \in L$.

(ii) $L, L' \in \mathcal{L}$ are linked by an edge $e_{L,L'}$ if and only if $L \cap L'$ has multiplicity 2.

The combinatorics C is equivalent to the data of the graph $\Gamma(C)$ up to every automorphism that respects the subsets \mathcal{L} and $\mathcal{Q}^{\geq 2}$.

Remark 2.4. We sometimes also consider the *full* incidence graph $\widehat{\Gamma}(C)$, with vertex set $\mathcal{L} \cup \mathcal{Q}$, and edge set given by the description: $L \in \mathcal{L}$ and $P \in \mathcal{Q}$ are linked by an edge $e_{P,L}$ if and only if $P \in L$.

Any topological equivalence $\Phi : (\mathbb{CP}^2, \mathcal{A}) \to (\mathbb{CP}^2, \mathcal{A}')$ induces an automorphism

(2.1)
$$G(\Phi): \Gamma(C_{\mathcal{A}}) \xrightarrow{\sim} \Gamma(C_{\mathcal{A}'})$$

see [JY93]. In particular, the combinatorics is a topological invariant of line arrangements. An arrangement equivalence such that $G(\Phi) = \text{Id}$ is called an *ordered* equivalence.

If two arrangements with the same combinatorics are such that there exists no equivalence between them, they form a *Zariski pair*. If there is no *ordered* (resp. *oriented*) equivalence, they form an *ordered* (resp. *oriented*) Zariski pair.

As an example, consider the two MacLane arrangements \mathcal{M}^+ and \mathcal{M}^- with 8 lines given by the equations:

$$L_{0}: 0 = z \qquad L_{3}: 0 = y \qquad L_{6}: 0 = -x - \omega^{2}y + z$$

$$L_{1}: 0 = -x + z \qquad L_{4}: 0 = \omega^{2}x + \omega y + z \qquad L_{7}: 0 = \omega y + z$$

$$L_{2}: 0 = x \qquad L_{5}: 0 = -x + y$$

where $\omega = e^{\frac{2i\pi}{3}}$ for \mathcal{M}^+ and $\omega = e^{-\frac{2i\pi}{3}}$ for \mathcal{M}^- . They share a common combinatorics whose incidence graph is given in Figure 1. The two MacLane arrangements form the smallest possible ordered oriented Zariski pair of line arrangements. However, there still exists an oriented equivalence between them that does not induce the identity on the graph, and in addition the complex conjugation in \mathbb{CP}^2 preserves the graph but not the orientation of the lines, see [Mac36]. Therefore the MacLane arrangements do not form neither an oriented nor an ordered Zariski pair, and in particular not a Zariski pair.



FIGURE 1. Incidence graph of the MacLane combinatorics

2.2. Orderings. Let W be a finite set with cardinal $m \ge 2$. A linear order on W is a bijection $\theta: W \to \{1, \ldots, m\}$. The set of all linear orders on W is denoted by $\mathfrak{L}(W)$. Consider the action of \mathfrak{S}_m on the set $\{1, \ldots, m\}$. The cyclic subgroup generated by the circular permutation $(1 \cdots m)$ induces a left free action on the same set. This in turn induces a left free action of $\langle (1 \cdots m) \rangle$ on $\mathfrak{L}(W)$ given by, for all $\theta \in \mathfrak{L}(W)$ and $r \in \mathbb{Z}$:

$$(1\cdots m)^r \cdot \theta = (1\cdots m)^r \circ \theta$$

The quotient set of $\mathfrak{L}(W)$ by this action is denoted by $\mathfrak{C}(W)$ and its elements are called *circular* orders on W. We denote the quotient map by

For every $\omega \in \mathfrak{C}(W)$, the *m* elements of $c^{-1}(\omega)$ are called the *linearisations* of the circular order ω . The set $\mathfrak{C}(W)$ has cardinal (m-1)!.

Now fix an element $\theta_0 \in \mathfrak{L}(W)$. For all $\theta \in \mathfrak{L}(W)$ there is a unique element σ of the permutation group \mathfrak{S}_m such that $\theta = \sigma \circ \theta_0$. There is a free transitive right action of \mathfrak{S}_m on $\mathfrak{L}(W)$ given by, for all $\tau \in \mathfrak{S}_m$:

$$\theta \cdot \tau = \sigma \circ \tau \circ \theta_0$$

Lemma 2.5. The right action of \mathfrak{S}_m on $\mathfrak{L}(W)$ respects the left quotient $c : \mathfrak{L}(W) \to \mathfrak{C}(W)$.

Proof. Let $\theta, \theta' \in \mathfrak{S}_m$ such that $\theta' = (1 \cdots m)^r \circ \theta$ for some $r \in \mathbb{Z}$. Then for every $\tau \in \mathfrak{S}_m$ one has again:

$$\theta' \cdot \tau = \sigma' \circ \tau \circ \theta_0 = (1 \cdots m)^r \circ \sigma \circ \tau \circ \theta_0 = (1 \cdots m)^r \circ \theta \cdot \tau \qquad \Box$$

There is therefore a transitive right action of \mathfrak{S}_m on $\mathfrak{C}(W)$. The stabiliser of an element $\omega \in \mathfrak{C}(W)$ is the subgroup \mathfrak{S}_ω generated by the permutation $(\sigma(1)\cdots\sigma(m))$ where $\sigma \circ \theta_0$ lies in $c^{-1}(\omega)$.

Consider now a graph Γ with vertex set $V(\Gamma)$. For any vertex $v \in V(\Gamma)$, the *neighbour set* W_v is the subset of vertices that are connected to v by an edge. The cardinal m_v of W_v is called the *multiplicity* of the vertex.

Definition 2.6. A graph ordering of Γ is a collection $\Omega = (\omega_v)_{v \in V(\Gamma)}$ where for every vertex v, $\omega_v \in \mathfrak{C}(W_v)$ is a circular order on the neighbour set W_v .

Lemma 2.7. Any graph ordering $\widehat{\Omega}$ of the full incidence graph $\widehat{\Gamma}(C_{\mathcal{A}})$ can naturally be restricted to a graph ordering Ω of the incidence graph $\Gamma(C_{\mathcal{A}})$.

Proof. The set of vertices of $\Gamma(C_A)$ is $\mathcal{L} \cup \mathcal{Q}^{>2}$. For every point-vertex $P \in \mathcal{Q}^{>2}$ one has directly $W_P = \widehat{W}_P$. For every line-vertex L, there is a natural bijection $\widehat{W}_L \to W_L$ that replaces any vertex $P = L \cap L' \in \mathcal{Q}^{>2}$ with L' itself. The local circular order of $\widehat{\Omega}$ on \widehat{W}_L thus extends to W_L .

3. Boundary manifold

Let \mathcal{A} be an arrangement in \mathbb{CP}^2 . Denote by $N_{\mathcal{A}}$ a closed tubular neighbourhood of \mathcal{A} . In [CS08], D. Cohen and A. Suciu give a geometrical construction of $N_{\mathcal{A}}$: write $\mathcal{A} = \{P_{\mathcal{A}} = 0\}$ where $P_{\mathcal{A}}$ is a defining homogenous polynomial of \mathcal{A} in \mathbb{CP}^2 . Let $\phi : \mathbb{CP}^2 \to \mathbb{R}$ defined by

$$\phi(\mathbf{x}) = \left[P_{\mathcal{A}}(\mathbf{x})\right]^2 / \|\mathbf{x}\|^{2(n+1)}$$

Then $N_{\mathcal{A}} := \phi^{-1}([0, \delta])$ does not depend on $\delta > 0$ for d sufficiently small.

The compact closed 3-manifold $B_{\mathcal{A}} := \partial N_{\mathcal{A}}$ is called the *boundary manifold* of \mathcal{A} and has the structure of a graph manifold, see [JY93]. The aim of this section is to describe this structure.

3.1. Circle bundles. Let Σ be an oriented surface of genus 0 and let W be a set of $m \geq 0$ marked points on Σ . Denote by Σ_W the surface obtained by removing from Σ an open neighbourhood around each point of W. A circle bundle (S, Σ_W, p) is a fibre bundle $p : S \to \Sigma_W$ with basis Σ_W , such that the fibres $p^{-1}(*)$ are homeomorphic to S^1 . For short, we often denote the circle bundle as simply S. An orientation of (S, Σ_W, p) is given by the data of two of the following three:

- (i) an orientation on S.
- (ii) an orientation on all fibres of p.
- (iii) an orientation of Σ_W .

Note that the choice of two of these fixes the third, while the choice of one leaves only two possibilities for the other two.

A homeomorphism $\phi : S \to S'$ is *fibrewise* if it commutes with p, and *fibre-positive* if it preserves the orientation of the fibres. A circle bundle is a special case of a *Seifert manifold*, namely one without singular fibres. The following results are standard, see for example [JS79; FM97].

Theorem 3.1. Let S, S' be two Seifert manifolds with more than 3 boundary components. Then any homeomorphism $S \to S'$ is isotopic to a fibrewise one.

Example 3.2 (Standard circle bundle). Let $\varepsilon \in \mathbb{Z}$ and let $D_W \subset \Sigma_W$ be a disc with m holes centred on the points of W. We denote by $\partial^{\infty} D_W$ the (m + 1)-th boundary component of D_W . Consider the oriented product bundle $T_W := D_W \times S^1$ and an oriented solid torus $T_{\infty} := D' \times S^1$.

Fix a section $\check{s}: D_W \hookrightarrow T_W$, which intersects the boundary ∂T_W on a collection $\check{\mu}$ of simple closed curves, with $\check{\mu}^w := \check{s}(\partial^w D_W)$. Consider a fibre $\check{\lambda}^{\infty}$ in T_W transverse to $\check{\mu}^{\infty}$. Similarly, let \check{s}' be a section of T_{∞} and consider a fibre $\check{\lambda}^{\infty'}$ transverse to $\check{\mu}^{\infty'} := \check{s}'(\partial D')$. Glue ∂T_{∞} to the toric boundary component $\partial^{\infty} D_W \times S^1$ of T_W using the gluing map:

$$g_{\varepsilon}: \begin{cases} \check{\mu}^{\infty} \longmapsto -\check{\mu}^{\infty'} - \varepsilon \cdot \check{\lambda}^{\infty} \\ \check{\lambda}^{\infty} \longmapsto \check{\lambda}^{\infty'} \end{cases}$$

where the gluing respects the orientations of the fibres of T_W and T_∞ . Then $S(W, \check{s}, \varepsilon) := T_W \cup_{g_\varepsilon} T_\infty$ is a circle bundle over Σ_W , and ε is called its *Euler number*. The definition obviously does not depend on the choice of \check{s}' .

Now let S be any circle bundle and fix a collection of closed curves $(\mu^w)_{w \in W}$ on ∂S such that $\mu^w \subset \partial^w S$ is transverse to the fibres of S. Such curves are called *horizontal*.

Theorem 3.3. Let (S, Σ_W, p) be a circle bundle with m boundary components and let $(\mu^w)_{w \in W}$ be a collection of horizontal curves on the boundary of S. Then there exists a unique number $\varepsilon \in \mathbb{Z}$ and a positive fibrewise homeomorphism $S(W, \check{s}, \varepsilon) \to S$ sending $\check{\mu}^w$ to μ^w for every $w \in W$.

By an abuse of language, we call the integer $\varepsilon \in \mathbb{Z}$ the *Euler number* of the oriented circle bundle S, since the usual Euler number vanishes for bundles with non-empty boundary.

3.2. Graph structure. Graph manifolds were first considered by F. Waldhausen [Wal67b; Wal67a] and were further developed by W. Neumann [Neu81]. The boundary manifold of a line arrangement is a graph manifold, see [JY93]. Detailed proofs can also be found in [Wes97; CS08].

A graph structure on a closed oriented 3-manifold M is a set Θ of pairwise disjoint joining tori such that $M \smallsetminus \Theta$ is a disjoint union of Seifert manifolds. A graph manifold is a closed oriented 3-manifold who admits a graph structure.

Theorem 3.4 ([Wal67b]). Any graph manifold admits a unique graph structure with a minimal number of tori, except for those listed in [Wal67b, Satz 8.1].

Exceptional classes of graph manifolds arise as boundary manifolds of only two specific families of complex line arrangements, listed in [JY93]. We leave these two families outside the scope of our study. From now on we only consider non-exceptional line arrangements.

One can naturally associate to a graph structure Θ of M a graph Γ as follows: each vertex v is associated to a Seifert manifold S_v and is decorated by its Euler number and singular fibres. Each edge $e_{v,w}$ corresponds to a gluing torus $T_{v,w} \in \Theta$ with $T_{v,w} = S_v \cap S_w$ and is decorated by the gluing map. The graph manifold M can be reconstructed from the graph by gluing together the Seifert pieces along the gluing tori. If M is a boundary manifold, then the graph has no deadends (i.e. every vertex has at least two neighbours), and the Seifert manifolds S_v are actually circle bundles of the form $S(W_v, \check{s}_v, \varepsilon_v)$ as described in Example 3.2. In particular they have no singular fibres. For every edge $e_{v,w}$, the Seifert pieces S_v and S_w are glued along the torus $T_{v,w}$ with a gluing map sending the horizontal curve $\check{\mu}_v^w \subset S_v$ to a fibre in S_w and the horizontal curve $\check{\mu}_w^v \subset S_v$ to a fibre in S_w and the horizontal curve $\check{\mu}_w^v \subset S_w$ to a fibre in S_v . This gluing map is the same for every edge, therefore the edges of the graph do not need to be decorated. By Theorem 3.4, M admits a unique such minimal graph, and we only consider this one from now on.

Theorem 3.5 ([JY93]). The boundary manifold $B_{\mathcal{A}}$ of a non-exceptional line arrangement \mathcal{A} is a graph manifold whose minimal graph coincides with the incidence graph $\Gamma(C_{\mathcal{A}})$ decorated with the following Euler numbers:

(i)
$$\varepsilon_L = 1 - b(L)$$
 for every line $L \in \mathcal{L}$.
(ii) $\varepsilon_P = -1$ for every singular point $P \in \mathcal{Q}^{>2}$.

where b(L) is the number of singular points of $\mathcal{Q}^{>2}$ meeting L.

Up to isotopy, homeomorphisms of a graph manifold always preserve the minimal graph structure.

Theorem 3.6 ([Wal67a]). Let M and N be two graph manifolds with respective minimal graph structures Θ_M and Θ_N . Let $\Phi : M \to N$ be a homeomorphism. Then Φ is isotopic to a homeomorphism $\Phi' : M \to N$ such that $\Phi'(\Theta_M) = \Phi'(\Theta_N)$.

According to Theorem 3.6, any homeomorphism of a graph manifold M acts on the graph structure Γ (eventually after isotopy). Therefore, there is a surjective morphism

(3.1)
$$G: \operatorname{Homeo}(M) \longrightarrow \operatorname{Aut}(\Gamma)$$

The kernel of G is called the group of graphed homeomorphisms of the graph manifold M and is denoted Homeo_{Γ}(M). In particular a graphed homeomorphism Ψ restricts to an homeomorphism $\Psi_v \in$ Homeo (S_v) for every bundle piece S_v of M, and Ψ_v fixes the boundary of S_v componentwise. Moreover, if every vertex of the incidence graph Γ has more than 3 neighbours (which is the case for boundary manifolds of non-exceptional line arrangements), then Theorem 3.1 applies and Ψ_v is fibrewise up to isotopy.

Definition 3.7. A graphed homeomorphism $\text{Homeo}_{\Gamma}(M)$ is *positive* if it respects the global orientation of M. It is *strongly positive* if its restriction on every piece S_v is fibre-positive. \diamond

We denote by $\operatorname{Homeo}_{\Gamma}^{++}(M)$ the group of isotopy classes of graphed strongly positive homeomorphisms of M.

Proposition 3.8. Let $\Phi : (\mathbb{CP}^2, \mathcal{A}) \to (\mathbb{CP}^2, \mathcal{A}')$ be a positive ordered equivalence of line arrangements. Then Φ induces an homeomorphism $B_{\mathcal{A}} \xrightarrow{\sim} B_{\mathcal{A}'}$ which is an element of $\operatorname{Homeo}_{\Gamma}^{++}(B_{\mathcal{A}})$, *i.e. graphed and strongly-positive.*

Proof. We write $B := B_A \simeq B_{A'}$. By (3.1), Φ induces a transformation $G(\Phi)$ on the graph Γ , which coincides with the transformation of (2.1). In particular since Φ is ordered then its restriction to the graph manifold B is graphed and positive. The gluing map between the bundle pieces of B respects their orientation. Therefore on every S_v the restriction Φ_v of Φ is also positive. However, Φ_v is not necessarily fibre-positive. If Φ_v changes the orientation of the fibres of some S_v , then Φ_v must also change the orientation of the basis Σ_{W_v} since it preserves the global orientation of S_v . These changes propagate through the gluings between the pieces, so much that if Φ is fibre-positive on one S_v then it must be on all of them, and vice versa. Let ν be the graphed positive homeomorphism of B that reverse the orientation of the fibres (and therefore of the bases) of all S_v 's. Then if Φ is not strongly-positive, $\nu \circ \Phi$ is.

4. FLAT STARS

A flat ordered star is an ordered set of non-intersecting arcs drawn on a disc with boundary and starting at a common point. They are used in Section 5 to define a way to embed all halfedges of the graph Γ with a common starting vertex $v \in V(\Gamma)$ inside the corresponding circle bundle component S_v of the boundary manifold B_A . Depending on the situation, we consider flat stars ordered with a linear or circular order, which are defined in Section 4.1. Sections 4.2 and 4.3 establish technical results which are then used in Section 4.4 to prove the main goals of this section, namely Propositions 4.12 and 4.13 that state that the pure mapping class group of surfaces acts transitively on flat stars.

4.1. Linear and circular stars.

Definition 4.1. Let M be a compact 3-dimensional manifold and let W be a subset of components of ∂M . A *star* on M is a collection of properly embedded simple arcs $\alpha = (\alpha^w)_{w \in W}$ in M such that:

- (i) for every $w \in W$, $\alpha^w([0,1[) \subset \mathring{M}$, and $\alpha^w(1) \in w$.
- (ii) there is a common point $b \in M$ such that for every $w \in W$, $\alpha^w(0) = b$.
- (iii) for every $w \neq w' \in W$, $\alpha^w \cap \alpha^{w'} = \{b\}$.

The point b is the *centre* of the star α and the arcs α^w are the *branches*.

Reusing notations from Section 3.1, let $D_W \subset \Sigma_W$ be a disc (resp. sphere) with *m* holes centred on the points of *W*. For each $w \in W$, the corresponding boundary component is denoted by $\partial^w D_W$ (resp. $\partial^w \Sigma_W$). The last boundary component of D_W is denoted by $\partial^\infty D_W$.

Definition 4.2. Let $\theta \in \mathfrak{L}(W)$ be a linear order on W. A *linear* θ -star drawn on D_W is a star $\bar{\alpha} = (\bar{\alpha}^w)_{w \in W}$ in D_W with centre b such that:

- (iv) $b \in \partial^{\infty} D_W$, and $\bar{\alpha}^w(1) \in \partial^w D_W$ for every $w \in W$.
- (v) there exists a neighbourhood U of b and a positive local chart $\phi : D_W \to \mathbb{R}^2$ that sends the pair $(D_W \cap U, \bar{\alpha} \cap U)$ to the model shown in Figure 2a ordered by θ .

The set of all isotopy classes of θ -linear stars on D_W is denoted by $\mathrm{LS}_{\theta}(\Delta_W)$.

Definition 4.3. Let $\omega \in \mathfrak{C}(W)$ be a circular order on W. A *circular* ω -star drawn on Σ_W is a star $\tilde{\alpha} = (\tilde{\alpha}^w)_{w \in W}$ in Σ_W with centre b such that:

(iv') $b \in \check{\Sigma}_m$, and $\check{\alpha}^w(1) \in \partial^w \Sigma_W$ for every $w \in W$.

(v') there exists a neighbourhood U of b and a positive local chart $\phi : \Sigma_W \to \mathbb{R}^2$ that sends the pair $(\Sigma_W \cap U, \tilde{\alpha} \cap U)$ to the model shown in Figure 2b ordered by ω .

The set of all isotopy classes of ω -circular stars on Σ_W is denoted by $CS_{\omega}(\Sigma_W)$.



FIGURE 2. Star orderings

 \diamond

 \Diamond



FIGURE 3. Examples of flat stars

Examples of a linear star and of a circular star are shown on Figures 3a and 3b respectively, with $W = \{x, y, z, v, w\}$ in this linear (and circular) order. We use the convention of denoting unordered stars as α , linear stars as $\bar{\alpha}$ and circular stars as $\tilde{\alpha}$.

We also make use of the set $CS(\Sigma_W)$ (resp. $LS(\Delta_W)$) of all isotopy classes of circular stars on Σ_W (resp. linear stars on D_W).

4.2. Mapping class groups. We recall some results about the mapping class groups of surfaces, see [Bir75; FM11] for details.

Let Σ be a compact planar oriented surface. The mapping class group $\mathcal{M}(\Sigma)$ is the group of isotopy classes of homeomorphisms of Σ . The pure mapping class group $\mathcal{P}(\Sigma)$ is the subgroup of homeomorphisms that restrict to the identity on $\partial \Sigma$.

Let Σ'_W be the surface obtained from D_W by filling in the boundary component $\partial^{\infty} D_W$ with a disk punctured with one point b_{∞} , and let $C : \mathcal{P}(D_W) \to \mathcal{P}(\Sigma'_W)$ be the map induced by the inclusion $f : D_W \hookrightarrow \Sigma'_W$.

Lemma 4.4 (Capping). There is a short exact sequence:

$$1 \longrightarrow \left\langle \Delta^2 \right\rangle \longrightarrow \mathcal{P}\left(D_W\right) \stackrel{C}{\longrightarrow} \mathcal{P}\left(\Sigma'_W\right) \longrightarrow 1$$

where Δ^2 is the full Dehn twist parallel to $\partial^{\infty} D_W$.

Let $F : \mathcal{P}(\Sigma'_W) \to \mathcal{P}(\Sigma_W)$ be the map induced by the inclusion $g : \Sigma'_W \to \Sigma_W$, where we fill in the puncture b_{∞} . There is a natural 'pushing' map $P : \pi_1(\Sigma_W, b_{\infty}) \to \mathcal{P}(\Sigma'_W)$ defined as such: for any simple closed curve δ in Σ_W based on b_{∞} , let δ_+ and δ_- be two curves parallel to δ , which enclose an annulus centred on δ . Then $P(\delta)$ is defined as the composition $T_{\delta_+} \circ T_{\delta_-}^{-1}$ of the Dehn twists along δ_+ and δ_- (see [FM11, Section 4.2.2] and Figure 6a).

Lemma 4.5 (Birman exact sequence). There is a short exact sequence:

$$1 \longrightarrow \pi_1(\Sigma_W, b_\infty) \xrightarrow{P} \mathcal{P}(\Sigma'_W) \xrightarrow{F} \mathcal{P}(\Sigma_W) \longrightarrow 1 \qquad \vartriangleleft$$

Lemmas 4.4 and 4.5 also stand for non-pure mapping class groups.

Let Δ_W be a disc with *m* punctures placed at the points of the set *W*. Suppose that $m \geq 2$. The braid group \mathbb{B}_m on *m* strands is generated by the braids $\sigma_{j,l}$ that performs a half twist on the strands *j* and *l*. The pure braid group \mathbb{P}_m is the subgroup generated by the braids $a_{j,l} := \sigma_{j,l}^2$ that performs a *full* twist on the strands *j* and *l*. Any element $\theta \in \mathfrak{L}(W)$ induces isomorphisms:

$$\mathbb{P}_m \xrightarrow{\sim} \mathcal{P}(\Delta_W) \qquad \mathbb{B}_m \xrightarrow{\sim} \mathcal{M}(\Delta_W)$$

The disc Δ_W can be obtained by capping every boundary components of D_W except $\partial^{\infty} D_W$. Applying Lemma 4.4 for each capping eventually gives the following result:

Proposition 4.6. Suppose that $m \ge 2$. Any element $\theta \in \mathfrak{L}(W)$ induces group isomorphisms

$$\rho_{\theta}: \mathbb{P}_m \times \mathbb{Z}^m \xrightarrow{\sim} \mathcal{P}\left(D_W\right) \qquad \rho_{\theta}: \mathbb{B}_m \times \mathbb{Z}^m \xrightarrow{\sim} \mathcal{M}\left(D_W\right) \qquad \lhd$$

The generator $a_{\theta(v),\theta(w)} \in \mathbb{P}_m$ is sent by ρ_{θ} to a full Dehn twist along a curve $\delta_{v,w}$ going around $\partial^v D_W$ and $\partial^w D_W$, as shown on Figure 8a. Similarly, $\sigma_{\theta(v),\theta(w)} \in \mathbb{B}_m$ is sent to a half Dehn twist along the same curve. The generator $d_{\theta(v)} = (0, \ldots, 1, \ldots, 0) \in \mathbb{Z}^m$ is sent to the full Dehn twist around a curve δ_v parallel to $\partial^v D_W$, as shown on Figure 9. Note that the subgroup \mathbb{Z}^m is isomorphic to the kernel of the capping map $C_W : \mathcal{P}(D_W) \to \mathcal{P}(\Delta_W)$.

To simplify notations, we often write $(\beta, d)_{\theta} := \rho_{\theta}(\beta, d)$. However, when using this notation the *left* action of the mapping class group becomes a *right* action of the braid group.

4.3. Model disc. Let Δ_m be the unit disc in the complex plane \mathbb{C} with $m \geq 2$ punctures x^1, \ldots, x^m aligned from right to left on the real axis, and let D_m be the disc obtained by removing a small open disc D^j around each puncture x^j in Δ_m . We define linear stars on D_m and on Δ_m similarly to Definition 4.2, replacing the boundary components with the punctures in the latter case. The base point is always -i and the linear order is always assumed to be the identity. The objective of this section is to establish Proposition 4.11 which states that the action of the pure mapping class group of D_m on these flat stars is transitive. This result on D_m will be used in Section 4.4 to generalise it to linear and circular stars on D_W and Σ_W respectively.

The proof of Proposition 4.11 uses a series of reductions to smaller structures: first we establish a similar result in the disc Δ_m in Proposition 4.10. The proof of this second result consists in turn in building a connection between linear stars on Δ_m and a construction due to [Fen+99] called *curve diagrams*. We prove with Proposition 4.9 that curve diagrams are in bijection with linear stars on Δ_m , and then we use Theorem 4.7 that establishes that $\mathcal{P}(\Delta_m)$ acts transitively on curve diagrams, and thus on linear stars on Δ_m . We then extend the result on Δ_m to D_m . Because of this entanglement of proofs, we present these results and their proofs by going from the smallest structure (curve diagrams) to the largest (linear stars on D_m).

A curve diagram is a curve embedded in the interior of Δ_m , starting from 1 and ending at -1, and going through all x^j 's in order. Let \mathcal{C}_m be the set of isotopy classes of curve diagrams in Δ_m . We denote by γ_0 the standard curve diagram shown on Figure 4.

Theorem 4.7 ([Fen+99]). The natural action of $\mathcal{P}(\Delta_m)$ on \mathcal{C}_m is free and transitive. In particular there is a bijection

$$\begin{array}{ccc} X: & \mathcal{C}_m \xrightarrow{\sim} \mathcal{P}\left(\Delta_m\right) \\ & \gamma \longmapsto \phi \text{ such that } \phi \cdot \gamma_0 = \gamma \end{array}$$

Denote by $LS(\Delta_m)$ the set of isotopy classes of linear stars on Δ_m (remember that they are all ordered by the identity).

Proposition 4.8. The natural action of $\mathcal{P}(\Delta_m)$ on Δ_m induces an action on $LS(\Delta_m)$ given by

$$\begin{array}{c} \mathcal{P}\left(\Delta_{m}\right) \times \mathrm{LS}(\Delta_{m}) \longrightarrow \mathrm{LS}(\Delta_{m}) \\ \left(\phi, \left(\bar{\alpha}^{j}\right)_{j}\right) \longmapsto \left(\phi \cdot \bar{\alpha}^{j}\right)_{j} \end{array} <$$

Proof. Remember from Section 4.2 that there is a group isomorphism $\mathbb{P}_m \to \mathcal{P}(\Delta_m)$. Take $\bar{\alpha} \in \mathrm{LS}(D_m)$ and $\beta \in \mathbb{P}_m$. The associated isomorphism is denoted by β_{Id} . It is obvious that β_{Id} preserves the properties of Definition 4.1, so $\bar{\alpha} \cdot \beta_{\mathrm{Id}}$ is still a star. The diffeomorphism β_{Id} is a product of Dehn twists along curves $\delta_{j,l}$ circling x^j and x^l . Up to isotopy, one can always

suppose that all $\delta_{j,l}$ do not intersect with the neighbourhood U where the ordering of $\bar{\alpha}$ is defined. Therefore $\bar{\alpha} \cdot \beta_{\text{Id}}$ still respects the Definition 4.2 of a linear star on Δ_m .



FIGURE 4. Standard linear star $\bar{\alpha}_0$ and curve diagram γ_0 on Δ_m

Proposition 4.9. There exists a bijection $LS(\Delta_m) \simeq C_m$ that respects the actions of $\mathcal{P}(\Delta_m)$ on each set.

Proof. We build a map

$$S: \mathcal{C}_m \longrightarrow \mathrm{LS}(\Delta_m)$$

as follows: Let $\gamma \in C_m$ be any curve diagram. By Theorem 4.7, there exists a diffeomorphism $X(\gamma) \in \mathcal{P}(\Delta_m)$ such that $X(\gamma) \cdot \gamma_0 = \gamma$. Let $\bar{\alpha}_0$ be the standard linear star shown on Figure 4. We then define

(4.1) $S(\gamma) := X(\gamma) \cdot \bar{\alpha}_0 \in \mathrm{LS}(\Delta_m)$

Conversely, we build another map

 $V: \mathrm{LS}(\Delta_m) \longrightarrow \mathcal{C}_m$

as follows: let $\bar{\alpha} \in \mathrm{LS}(\Delta_m)$ be a linear star and consider a regular neighbourhood N of $\bar{\alpha}$ inside the complex plane, see Figure 5a for an example. Define $N_{\bar{\alpha}} := N \cap \Delta_m \setminus \bar{\alpha}$ as the *stripe* of $\bar{\alpha}$. It can be obtained by cutting $N \cap \Delta_m$ along each branch of α . The boundary $\partial N_{\bar{\alpha}}$ can be separated in two parts: one corresponds to $\partial N \cap \Delta_m$, and the other corresponds to the cutting of the star $\bar{\alpha}$ with the m points x^j marked in order. Each time we cut along a branch of $\bar{\alpha}$ the base point -i is 'split'. There are thus m + 1 'copies' of -i on $\partial N_{\bar{\alpha}}$ intertwined with the x^j 's, as shown on Figure 5b. A wedge collection $(V^j)_j$ is a set of curves embedded in $N_{\bar{\alpha}}$ such that each V^j goes from $\partial N \cap \Delta_m$ to x^j and back to $\partial N \cap \Delta_m$, and such that $V^j \cap V^k = \emptyset$ if $j \neq k$. Then $V(\bar{\alpha})$ is defined as the blue curve shown on Figure 5b, embedded in Δ_m .

Let us prove that the map V is well-defined.

Let $\bar{\alpha}, \bar{\alpha}'$ be two linear stars in Δ_m such that there exists an isotopy I_t of Δ_m sending $\bar{\alpha}$ to $\bar{\alpha}'$. Then I_t also sends the pair $(N_{\bar{\alpha}}, \partial N_{\bar{\alpha}})$ to $(N_{\bar{\alpha}'}, \partial N_{\bar{\alpha}'})$. Since I_t fixes $\partial \Delta_m$, it also preserves the separation of $\partial N_{\bar{\alpha}}$ shown on Figure 5b, and in particular I_t fixes all the x^j 's. It is clear that one can always build an isotopy J_t of $N_{\bar{\alpha}}$ sending any wedge collection to any other. Extend J_t by the identity outside of $N_{\bar{\alpha}}$. Then the composition $J_t \circ I_t$ is an isotopy of Δ_m sending $V(\bar{\alpha})$ to $V(\bar{\alpha}')$.

We now prove that the maps S and V are inverse to each other.

Let $\gamma \in \mathcal{C}_m$ and let $\bar{\alpha} := S(\gamma)$. We want to show that $V(\bar{\alpha}) = \gamma$. It is clear from the definition of V that for the standard linear star we have $V(\bar{\alpha}_0) = \gamma_0$. Consider $N_{\bar{\alpha}_0}$ the stripe of $\bar{\alpha}_0$.



FIGURE 5. Stripe $N_{\bar{\alpha}}$ of a star $\bar{\alpha}$

By (4.1) we have $\bar{\alpha} = X(\gamma) \cdot \bar{\alpha}_0$. The diffeomorphism $X(\gamma)$ preserves the regular neighbourhoods up to isotopy and therefore sends the pair $(N_{\bar{\alpha}_0}, \partial N_{\bar{\alpha}_0})$ to $(N_{\bar{\alpha}}, \partial N_{\bar{\alpha}})$. In particular:

$$V(\bar{\alpha}) = X(\gamma) \cdot V(\bar{\alpha}_0) = X(\gamma) \cdot \gamma_0 = \gamma$$

Therefore $V \circ S = \operatorname{Id}_{\mathcal{C}_m}$. Reciprocally, let $\bar{\alpha} \in \operatorname{LS}(\Delta_m)$ and let $\gamma := V(\bar{\alpha})$. We want to show that $S(\gamma) = \bar{\alpha}$. By (4.1), we have $S(\gamma) = X(\gamma) \cdot \bar{\alpha}_0$. Let $\bar{\alpha}' = X(\gamma)^{-1} \cdot \bar{\alpha}$. We show that $\bar{\alpha}' = \bar{\alpha}_0$. The curve γ separates Δ_m in two halves and by construction $\bar{\alpha}$ is contained entirely within one half. The diffeomorphism $X(\gamma)^{-1}$ preserves that separation, which implies that $\bar{\alpha}'$ is contained entirely within the lower half $\Delta_m \cap \{\operatorname{Re}(z) \leq 0\}$, just like $\bar{\alpha}_0$. By Definition 4.2, the stars $\bar{\alpha}'$ and $\bar{\alpha}_0$ are isotopic inside a neighbourhood U of their common base point -i. We are now reduced to examining the 'rectangle' $R := \Delta_m \cap \{\operatorname{Re}(z) \leq 0\} \setminus U$. The boundary ∂R can be divided in four parts. The top part corresponds to γ_0 with the x^j 's marked. The bottom part corresponds to $\partial U \cap \Delta_m$ and has also m points marked, namely the $y^j := \partial U \cap \bar{\alpha}'^j = \partial U \cap \bar{\alpha}_0^j$. The points (x^j) and (y^j) are ordered oppositely on their respective parts of ∂R with respect to the the orientation of Δ_m . Inside R, the star $\bar{\alpha}'$ (and $\bar{\alpha}_0$) is a set of simple non-intersecting arcs joining each y^j to the corresponding x^j . It is clear that there is only one isotopy type of this construction. The stars $\bar{\alpha}'$ and $\bar{\alpha}_0$ are therefore isotopic inside both R and U, so $\bar{\alpha}' = \bar{\alpha}_0$. Therefore $S \circ V = \operatorname{Id}_{\operatorname{LS}(\Delta_m)}$, which completes the proof. \Box

Proposition 4.10. The action of *Proposition 4.8* is free and transitive.

Proof. This is immediate from Theorem 4.7 and the fact that the bijection of Proposition 4.9 commutes with the natural action of $\mathcal{P}(\Delta_m)$ on Δ_m by construction.

We can now go back to D_m .

Proposition 4.11. There is a natural action of $\mathcal{P}(D_m)$ on $\mathrm{LS}(D_m)$. This action is transitive.

Proof. The disc Δ_m can be obtained from D_m by 'capping' each boundary component of D_m (save for $\partial^{\infty} D_m$) with a disc D^j with one puncture at x^j . The capping operation induces a map

$$J: \mathrm{LS}(D_m) \longrightarrow \mathrm{LS}(\Delta_m)$$

described as follows: consider $\bar{\alpha}$ in $D_m \subset \Delta_m$. Extend each branch $\bar{\alpha}^j$ inside D^j by an arc joining the end point on ∂D^j to the puncture x^j . There is only one isotopy type of such an arc

inside D^j . This extended star is $J(\bar{\alpha})$. An isotopy between two stars in D_m can be extended to each D^j , giving isotopic extended stars in Δ_j . Therefore J is well-defined.

Define an equivalence relation on $LS(\Delta_m)$ by

$$\forall \,\bar{\alpha}, \bar{\alpha}' \in \mathrm{LS}(\Delta_m) : \quad \bar{\alpha} \simeq \bar{\alpha}' \Longleftrightarrow J(\bar{\alpha}) = J(\bar{\alpha}')$$

We first prove that the subgroup $\rho_{\mathrm{Id}}(\mathbb{P}_m)$ acts freely and transitively on the set of equivalence classes of $\mathrm{LS}(\Delta_m)$. For every $\phi \in \mathcal{P}(D_m)$ and $\bar{\alpha} \in \mathrm{LS}(\Delta_m)$, we have:

(4.2)
$$J(\phi \cdot \bar{\alpha}) = C(\phi) \cdot J(\bar{\alpha})$$

where $C : \mathcal{P}(D_m) \to \mathcal{P}(\Delta_m)$ is the capping map obtained from applying Lemma 4.4 on all boundary components $\partial^j D_m$. The map J is clearly surjective, and so is the map C by Lemma 4.4. The result then follows from the freeness and transitivity of the action of $\mathcal{P}(\Delta_m)$ on $\mathrm{LS}(\Delta_m)$ by Proposition 4.10.

We now prove that the orbits of the action of the subgroup $\rho_{\mathrm{Id}}(\mathbb{Z}^m)$ coincide with the equivalence classes of $\mathrm{LS}(\Delta_m)$.

Let $\bar{\alpha}, \bar{\alpha}' \in \mathrm{LS}(\Delta_m)$. Suppose that there exists $\phi \in \rho_{\mathrm{Id}}(\mathbb{Z}^m)$ such that $\bar{\alpha} = \phi \cdot \bar{\alpha}'$. Then by Lemma 4.4, $\phi \in \ker C$ so by (4.2), $J(\bar{\alpha}) = J(\bar{\alpha}')$. Reciprocally, suppose that $\bar{\alpha}$ is equivalent to $\bar{\alpha}'$. There is an isotopy of Δ_m sending $J(\bar{\alpha})$ to $J(\bar{\alpha}')$. By definition of J, this isotopy restricts to an homeomorphism ϕ of D_m sending $\bar{\alpha}$ to $\bar{\alpha}'$. We then have:

$$J(\bar{\alpha}') = C(\phi) \cdot J(\bar{\alpha}) = J(\bar{\alpha})$$

But the action of $\mathcal{P}(\Delta_m)$ on $\mathrm{LS}(\Delta_m)$ is free. Therefore, $\phi \in \ker C = \rho_{\mathrm{Id}}(\mathbb{Z}^m)$.

4.4. Action on flat stars. In this section we prove that the action of the pure mapping class group of surfaces on the sets of linear and circular stars is transitive in both cases. The case of linear stars on D_W follows directly from the results of Section 4.3. The case of circular stars on Σ_W can eventually be reduced to the previous one thanks to Lemma 4.14 which explains how to particularise a circular star into a linear one.

Proposition 4.12. Let θ be a linear order on W. The natural action of $\mathcal{P}(D_W)$ on D_W induces a well-defined action on $\mathrm{LS}_{\theta}(D_W)$ given by

$$\mathcal{P}(D_W) \times \mathrm{LS}_{\theta}(D_W) \longrightarrow \mathrm{LS}_{\theta}(D_W)$$
$$\left((\beta, d)_{\theta}, (\bar{\alpha}^w)_{w \in W} \right) \longmapsto (\bar{\alpha}^w \cdot (\beta, d)_{\theta})_{w \in W}$$

This action is transitive.

Proof. The choice of the ordering θ on W induces an homeomorphism $M_{\theta}: D_m \to D_W$, which in turn induces a bijection $\mathrm{LS}(D_m) \simeq \mathrm{LS}_{\theta}(D_W)$. The results of Proposition 4.11 are thus passed on to D_W and $\mathrm{LS}_{\theta}(D_W)$.

Proposition 4.13. Let ω be a circular order on W. The natural action of $\mathcal{P}(\Sigma_W)$ on Σ_W induces a well-defined action on $CS_{\omega}(\Sigma_W)$. This action is transitive.

Recall the circularisation map $c : \mathfrak{L}(W) \twoheadrightarrow \mathfrak{C}(W)$ from (2.2). It is clear from Definition 4.2 that the inclusion $i : D_W \hookrightarrow \Sigma_W$ induces a surjective map

$$i: \mathrm{LS}(D_W) \longrightarrow \mathrm{CS}(\Sigma_W)$$

In particular, for every $\theta \in \mathfrak{L}(W)$, the map restricts to:

$$i: \mathrm{LS}_{\theta}(D_W) \longrightarrow \mathrm{CS}_{c(\theta)}(\Sigma_W)$$

Lemma 4.14. Let $\theta, \theta' \in \mathfrak{L}(W)$ such that $c(\theta) = c(\theta') = \omega$. Then for every linear star $\bar{\alpha} \in \mathrm{LS}_{\theta}(D_W)$ there exists another linear star $\bar{\alpha}' \in \mathrm{LS}_{\theta'}(D_W)$ such that $i(\bar{\alpha}) = i(\bar{\alpha}') \in \mathrm{CS}_{\omega}(\Sigma_W)$.

Proof. It is enough to prove the result when $\theta' = (1 \cdots m) \circ \theta$. Fix $b_{\infty} \in \Sigma_W \setminus D_W$ and define $\Sigma'_W = \Sigma_W \setminus \{b_{\infty}\}$. We write

$$D_W \stackrel{f}{\longleftrightarrow} \Sigma'_W \stackrel{g}{\longleftrightarrow} \Sigma_W$$

with $i = g \circ f$. Let $\bar{\alpha} \in \mathrm{LS}_{\theta}(D_W)$ drawn on D_W and let $z \in W$ with $\theta(z) = m$. Consider a loop $\delta \subset \Sigma_W$ based on b_∞ that circles the entire star $i(\bar{\alpha})$ except the last branch (in linear order) $\bar{\alpha}^z$, which it crosses once. Such a loop can always be constructed as follows: starting from the centre b of the star, take a path parallel to the left of $\bar{\alpha}^z$, then turn around $\partial^z D_W$ and return to b with a path parallel to the right of $\bar{\alpha}^z$. Repeat the process for all other branches in reverse linear order. Then inside the neighbourhood U of b, shortcut the loop around $\partial^z D_W$. Finally, deform the loop in $\Sigma_W \setminus D_W$ to make it go around b_∞ . Now recall from Section 4.2 that δ induces a 'pushing' transformation $P(\delta) \in \mathcal{P}(\Sigma'_W)$. Its action on $f(\bar{\alpha})$ is shown on Figure 6a. By construction, only the last strand $\bar{\alpha}^z$ is affected. Let $\hat{\alpha}' = P(\gamma) \cdot f(\bar{\alpha})$. By dragging the modified branch $\hat{\alpha}'^z$ all across $\Sigma'_W \setminus D_W$, the star $\hat{\alpha}'$ can be isotoped inside Σ'_W to the star shown on Figure 6b, where $\hat{\alpha}'^z$ is now the first branch in linear order inside D_W . Thus $\bar{\alpha}' := f^{-1}(\hat{\alpha}') \in \mathrm{LS}_{\theta'}(\Delta_W)$. Finally, by Lemma 4.5, we have

$$f(\bar{\alpha}') = g(\hat{\alpha}') = g(P(\delta) \cdot f(\bar{\alpha})) = F(P(\delta)) \cdot i(\bar{\alpha}) = i(\bar{\alpha})$$

since $F \circ P =$ Id.

i



FIGURE 6. Pushing map action on Σ'_W

Proof of Proposition 4.13. Let $\omega \in \mathfrak{C}(W)$ and $\tilde{\alpha} \in \mathrm{CS}_{\omega}(\Sigma_W)$. Let D_{∞} be a disc on Σ_W that does not meet $\tilde{\alpha}$ except on its centre *b*. Removing \mathring{D}_{∞} from Σ_W , we obtain D_W . Consider the inclusion $i: D_W \hookrightarrow \Sigma_W$. There exists a linear star $\bar{\alpha} \in \mathrm{LS}(D_W)$ such that $i(\bar{\alpha}) = \tilde{\alpha}$. By Lemmas 4.4 and 4.5, the map *i* induces a surjective map $F \circ C : \mathcal{P}(D_W) \twoheadrightarrow \mathcal{P}(\Sigma_W)$. Let $\bar{\phi} \in \mathcal{P}(D_W)$ such that $\phi = (F \circ C)(\bar{\phi})$. Then:

$$\phi \cdot \tilde{\alpha} = \phi \cdot i(\bar{\alpha}) = (F \circ C)(\bar{\phi}) \cdot i(\bar{\alpha}) = i(\bar{\phi} \cdot \bar{\alpha})$$

By Proposition 4.6, there exists $(\beta, d) \in \mathbb{P}_m \times \mathbb{Z}^m$ such that $\overline{\phi} = \rho_\theta(\beta, d)$. Then by Proposition 4.12, $\overline{\phi}(\overline{\alpha})$ is again a θ -linear star. Its image by i is thus again a ω -circular star on Σ_W .

We now prove the transitivity of the action. Let $\tilde{\alpha}, \tilde{\alpha}' \in CS_{\omega}(\Sigma_W)$ be two ω -circular stars on Σ_W . Up to to isotopy we can suppose that the centre b of $\tilde{\alpha}$ does not lie on $\tilde{\alpha}'$ and vice versa. Let D_{∞} be a disc that does not meet neither $\tilde{\alpha}$ nor $\tilde{\alpha}'$. Up to isotopy one can always move the centres b and b' to ∂D_{∞} . Similarly to the proof of Proposition 4.13, by removing \mathring{D}_{∞} from Σ_W , we obtain D_W and two linear stars $\bar{\alpha}, \bar{\alpha}'$ ordered by two linearisations $\theta, \theta' \in c^{-1}(\omega)$

respectively. By Lemma 4.14, up to an isotopy of Σ_W we can actually suppose that $\theta = \theta'$. Then by Proposition 4.12 there exists $\bar{\phi} \in \mathcal{P}(D_W)$ such that $\bar{\phi} \cdot \bar{\alpha} = \bar{\alpha}'$. By Lemmas 4.4 and 4.5 we have

$$\tilde{\alpha}' = i(\bar{\alpha}') = i(\bar{\phi} \cdot \bar{\alpha}) = (F \circ C)(\bar{\phi}) \cdot \hat{c}$$

with $(F \circ C)(\overline{\phi}) \in \mathcal{P}(\Sigma_W)$.

5. Graphed embeddings

The first homology and fundamental group of the boundary manifold $B_{\mathcal{A}}$ have two types of generators: one corresponding to the meridians of the generators and the other to the cycles of the graph. To properly define the cycle generators, it is necessary to associate them with actual curves drawn on $B_{\mathcal{A}}$ itself. We therefore define a type of morphism

$$\gamma: \Gamma\left(C_{\mathcal{A}}\right) \longrightarrow B_{\mathcal{A}}$$

called graphed embeddings that respects the graph structure of $B_{\mathcal{A}}$. Graphed embeddings are built by tying together ordered stars, one for each circle bundle component of the graph manifold $B_{\mathcal{A}}$. Recall from Theorem 3.5 that $B_{\mathcal{A}}$ is a graph manifold over the graph $\Gamma = \Gamma(C_{\mathcal{A}})$.

Definition 5.1. Let $\Omega = (\omega_v)_{v \in V(\Gamma)}$ be a graph ordering of Γ . An ordered graphed embedding with respect to Ω is a map $\gamma : \Gamma \hookrightarrow B_{\mathcal{A}}$ described as such: for every vertex $v \in V(\Gamma), \gamma \cap S_v = \alpha_v$ is a star drawn on the circle bundle $p_v : S_v \to \Sigma_{W_v}$ such that $p_v(\alpha_v)$ is a circular star on Σ_{W_v} ordered by ω_v .

Note that the branches of the star α_v inside S_v can travel along the fibres but two branches can never meet the same fibre, except over the centre b_v .

The set of isotopy classes of ordered graphed embeddings with respect to a graph ordering Ω is denoted by $E_{\Gamma}(\Omega)$.

Proposition 5.2. There is a well-defined action of the group of graphed strongly positive homeomorphisms Homeo_{Γ^+} (B_A) on $E_{\Gamma}(\Omega)$.

Proof. Consider $\gamma \in E_{\Gamma}(\Omega)$. For every $v \in V(\Gamma)$, let $\alpha_v = \gamma \cap S_v$ be the corresponding star. Then $p_v(\alpha_v) \in CS_{\omega_v}(\Sigma_{W_v})$. Let $\Psi \in Homeo_{\Gamma}^{++}(M)$. Since Ψ is graphed then for every $v \in V(\Gamma)$ it restricts to an homeomorphism Ψ_v of S_v , which is fibre-positive and fixes the boundary of S_v componentwise. In particular Ψ_v induces an element $\widetilde{\Psi}_v \in \mathcal{P}(\Sigma_{W_v})$ on the basis Σ_{W_v} of the fibre bundle S_v , such that the following diagram commutes:

$$S_{v} \xrightarrow{\Psi_{v}} S_{v}$$

$$\downarrow^{p_{v}} \qquad \downarrow^{p_{v}}$$

$$\Sigma_{W_{v}} \xrightarrow{\widetilde{\Psi}_{v}} \Sigma_{W_{v}}$$

By Proposition 4.13, $\widetilde{\Psi}_v \cdot p_v(\alpha_v)$ is again a circular star on Σ_{W_v} for the same order ω_v . Since

$$p_v(\Psi_v \cdot \alpha_v) = \Psi_v \cdot p_v(\alpha_v)$$

then $\Psi_v \cdot \alpha_v$ respects the conditions of Definition 5.1. Since this is true for all $v \in V(\Gamma)$ then $\Psi \cdot \alpha \in \mathcal{E}_{\Gamma}(\Omega)$.

The following fundamental 'flattening' lemma allows to consider a graphed embedding as a union of linear stars rather than circular ones, which makes the action easier to handle. Recall from Theorem 3.4 that for every circle bundle S_v there is a model $\chi_v : S(m_v, \check{s}_v, \varepsilon_v) \to S_v$ with $S(m_v, \check{s}_v, \varepsilon_v) = T_{m_v} \cup T_{\infty}$, and where \check{s}_v is a fixed section of T_{m_v} .

Lemma 5.3. Let $\gamma = (\alpha_v)_{v \in V(\Gamma)} \in E_{\Gamma}(\Omega)$. Let $v \in V(\Gamma)$ be a fixed vertex. There exists a linear star $\bar{\alpha}_v \in LS_{\theta_v}(D_{W_v})$, where $\theta_v \in \mathfrak{L}(W_v)$ is a linearisation of ω_v , and such that α_v can be isotoped to $\check{s}_v(\bar{\alpha}_v)$ inside $B_{\mathcal{A}}$.

Proof. Up to isotopy, the star α_v can be moved entirely within the trivially fibred torus T_{m_v} . If no branch of α_v makes a full loop around a fibre of T_{m_v} then there exist a section s' of T_{m_v} such that $s' \cap \alpha_v = \emptyset$. Cut T_{m_v} along s' to get a cylinder $S_{m_v} \times I$, then retract the star α_v on the section \check{s}_v along the fibres to get the result. Now suppose that for some $w \in W_v$ the branch α_v^w makes at least one full loop around a fibre. Up to isotopy one can always move the loops at the extremity of the branch, i.e. on the boundary torus $\partial^w S_v$. Inside the graph manifold B_A , the fibres of $\partial^w S_v$ are isotopic to horizontal loops inside $T_{m_w} \subset S_w$. One can thus 'push' the loops to the other side of the gluing with an isotopy of B_A , so that the branch α_v^w has no more fibre loops. Finally, push the centre b of α_v to the boundary $\partial^\infty \check{s}_v$. This has the effect of particularising the circular order ω_v to give a flat θ_v -linear star on \check{s}_v .

Lemma 5.4. For any $v \in V(\Gamma)$, the isotopy of Σ_{W_v} described in the proof of Lemma 4.14 can be lifted to an isotopy of S_v .

Proof. Let $z = \theta_v^{-1}(m_v) \in W_v$ and let δ be a loop on Σ_{W_v} that separates $\partial^z \Sigma_W$ from the other boundary components, and let δ_+ and δ_- be two curves parallel to δ as shown on Figure 6. Then we have

$$\Sigma_{W_v} = E_{\delta_-} \cup_{\delta_-} D_{\delta_-}$$

with $E_{\delta_+} \simeq D_{W_v \smallsetminus \{z\}}$ and $D_{\delta_-} \simeq D_{\{z\}}$. The 'pushing' transformation $P(\delta)$ is the identity outside of D_{δ_-} . Using a trivialisation T_{δ} of the fibre bundle over D_{δ_-} , one can thus lift $P(\delta)$ to a partial isotopy I of S_v , which can then be extended by the identity to the entirety of S_v . By construction, $p_v \circ I = P(\delta)$ inside T_{δ} .

Proposition 5.5. The action of Proposition 5.2 is transitive.

Proof. Let $\gamma, \gamma' \in E_{\Gamma}(\Omega)$ be two ordered graphed embeddings. Fix $v \in W$ and write $\alpha_v = \gamma \cap S_v$ and $\alpha'_v = \gamma' \cap S_v$. By Lemma 5.3 up to isotopy one can suppose that $\alpha_v = \check{s}_v(\bar{\alpha}_v)$ (resp. $\alpha_v = \check{s}_v(\bar{\alpha}'_v)$) with $\bar{\alpha}_v \in \mathrm{LS}_{\theta_v}(D_{W_v})$ (resp. $\bar{\alpha}'_v \in \mathrm{LS}_{\theta'_v}(D_{W_v})$), where $\theta_v, \theta'_v \in c^{-1}(\omega_v)$. Projecting on the basis Σ_{W_v} of S_v , we have $p(\check{s}_v) \simeq D_{W_v}$. Write

$$\Sigma_{W_v} = p(\check{s}_v) \cup_{\partial^\infty} D_0$$

where $D_0 \simeq D^2$. The situation is similar to Figure 6 with $p(\check{s}_v)$ and D_0 as the upper and lower hemispheres respectively. By Lemma 5.4, we can suppose that $\theta'_v = \theta_v$. Then by Proposition 4.12 there exists $\psi_v \in \mathcal{P}(D_{W_v})$ such that $\psi_v \cdot \bar{\alpha}_v = \bar{\alpha}'_v$. Using the trivialisation of S_v over D_{W_v} and extending by the identity on the rest of S_v we can lift ψ_v to a fibre-positive homeomorphism Ψ_v of S_v . Then $\Psi_v \cdot \alpha_v = \alpha'_v$. Moreover Ψ_v restricts to the identity on ∂S_v . One can then define an homeomorphism Ψ of $B_{\mathcal{A}}$ as the reunion of all Ψ_v acting on their respective S_v for every $v \in V(\Gamma)$. By construction, $\Psi \in \text{Homeo}_{\Gamma}^{++}(B_{\mathcal{A}})$ and $\Psi \cdot \gamma = \gamma'$.

6. Homology inclusion

In this section we present the construction of our new ordered oriented line arrangements invariant. Consider the inclusion map

 $i_{\mathcal{A}}: B_{\mathcal{A}} \longrightarrow E_{\mathcal{A}}$

and the induced map on the first homology groups

$$i_{\mathcal{A}}^*: H_1(B_{\mathcal{A}}, \mathbb{Z}) \longrightarrow H_1(E_{\mathcal{A}}, \mathbb{Z})$$

In Section 6.1 we give a presentation of these homology groups. The group $H_1(E_A, \mathbb{Z})$ is generated by the meridians of the lines of \mathcal{A} (modulo the sum of all of them), corresponding to the fibres of the associated circle bundle components of B_A . Meridians of the singular points can also be added as redundant generators. On the other hand $H_1(B_A, \mathbb{Z})$ is generated by the same meridians plus the cycles of the graph. Theorem 6.5 presents how the choice of an ordered graphed embedding allows to define the cycle generators. With that setting, Section 6.2 gives the definition of the invariant. The restriction of i_A^* on the cycle generators depends on the choice of a graph ordering Ω and of an ordered graphed embedding $\gamma \in E_{\Gamma}(\Omega)$. In order to remove that last dependency, we define the graph stabiliser \mathcal{G}_{Γ} as the quotient of all potential images of the cycles as sums of meridians by the differences between every two graphed embeddings. Then the homology inclusion $\mathcal{J}_{\Omega}(\mathcal{A})$ is given in Definition 6.10 as the class of the restriction of i_A^* inside the graph stabiliser. Theorem 6.11 establishes that \mathcal{J}_{Ω} is an invariant of ordered oriented line arrangements.

6.1. **Presentations of the homology groups.** For the boundary manifold the presentation is well known since the works of Westlund [Wes97], see also [Mum61; ACM19b; KN14]. For the exterior the presentation is obtained by abelianising the fundamental group, which is itself computed using the Zariski-van Kampen method. We simply reformulate these results using our own notations and the new notion of *ordered* graphed embeddings.

For this entire section, $\gamma \in E_{\Gamma}(\Omega)$ is a fixed ordered graphed embedding. Two types of generators are involved: the *meridians* which correspond to the fibres of each circle bundle component of $B_{\mathcal{A}}$, and the cycles of the graph Γ .

We start with the exterior manifold $E_{\mathcal{A}}$. Let $C_0(\Gamma)$ be the free abelian group generated by the set $V(\Gamma)$ of vertices of the graph Γ . Recall also that every vertex $v \in V(\Gamma)$ is decorated with the Euler number $\varepsilon_v \in \mathbb{Z}$ of the corresponding S_v .

Definition 6.1. The meridian homology $M(\Gamma)$ of a graph Γ is the quotient of $C_0(\Gamma)$ by the normally generated subgroup

$$R(\Gamma) \mathrel{\mathop:}= \left\langle \varepsilon_v \cdot v + \sum_{w \in W_v} w, \ v \in V(\Gamma) \right\rangle$$

with the exact sequence:

$$0 \longrightarrow R(\Gamma) \longrightarrow C_0(\Gamma) \xrightarrow{\eta} M(\Gamma) \longrightarrow 0 \qquad \diamond$$

For every $x \in C_0(\Gamma)$ we note \bar{x} its image in $M(\Gamma)$.

Proposition 6.2. There is a natural isomorphism $M(\Gamma(C_{\mathcal{A}})) \xrightarrow{\sim} H_1(E_{\mathcal{A}})$ that sends the class \bar{v} to the homology class f_v of the fibres of the circle bundle S_v for every vertex $v \in V(\Gamma)$.

Proof. Is it known that:

$$H_1(E_{\mathcal{A}}) = \left\langle f_L, \ L \in \mathcal{L} \ \bigg| \ \sum_{L \in \mathcal{L}} f_L \right\rangle$$

where f_L is the line meridian of L, that is the homological class of the fibres of S_L . Now using Theorem 3.5, the relations of $M(\Gamma(C_A))$ become

$$\forall L \in \mathcal{L}: \quad (b(L) - 1) \cdot L = \sum_{P \in W_L} P + \sum_{L' \in W_L} L' \qquad \forall P \in \mathcal{Q}^{>2}: \quad P = \sum_{L \in W_P} L$$

Replacing every P in the first relation allows to remove them all as generators and yields

$$\forall L \in \mathcal{L}: \quad (b(L) - 1) \cdot L = \sum_{P \in W_L} \sum_{L'' \in W_P} L'' + \sum_{L' \in W_L} L'$$

There are exactly b(L) singular points in W_L and each has L once amongst their neighbours. We can thus simplify the relations to get:

$$\forall L \in \mathcal{L}: \quad 0 = L + \sum_{\substack{P \in W_L}} \sum_{\substack{L'' \in W_P \\ L'' \neq L}} L'' + \sum_{\substack{L' \in W_L}} L'$$

where L'' (resp. L') are all the lines meeting L at singular points with multiplicity > 2 (resp. 2). But since L meets every other line of \mathcal{L} exactly once, we have in fact

$$\forall L \in \mathcal{L} : \quad 0 = L + \sum_{\substack{L' \in \mathcal{L} \\ L' \neq L}} L' = \sum_{L' \in \mathcal{L}} L'$$

Identifying the formal generator L with f_L , this is exactly the defining relation of $H_1(E_A)$ repeated for every line.

Note that Proposition 6.2 implies that $H_1(E_A)$ is combinatorially determined. We use the name 'meridian homology' and notation $M(\Gamma)$ to designate the corresponding formal combinatorial group. This distinction is necessary because $M(\Gamma)$ also appears as an important subgroup of the first homology group of the boundary manifold $H_1(B_A)$, as we shall see now.

The fundamental group $\pi_1(B_A)$ has two types of generators: the meridians, which must be connected to a common base point, and the cycles of the graph. A graphed embedding is required to properly draw both types of generators on the boundary manifold B_A . One also needs additional combinatorial data to obtain a presentation of $\pi_1(B_A)$.

Fix a vertex $r \in V(\Gamma)$. Seeing the graph Γ as a topological space, one can define the fundamental group $\pi_1(\Gamma, r)$, which is freely generated by the cycles of the graph based in r. Note that there is no canonical set of generators. Let \mathcal{T} be a spanning tree on Γ , that is a connected acyclic subgraph of Γ containing all vertices. For every pair of vertices $v, w \in V(\Gamma)$ there always exist a unique path from v to w inside \mathcal{T} . Then for every edge $e_{v,w} \in \Gamma \setminus \mathcal{T}$, there is a unique cycle $c_{v,w}$ drawn on Γ going through $e_{v,w}$ and the root r. In other words, the choice of \mathcal{T} gives a cycle basis. We then write $\gamma_{v,w} := \gamma(c_{v,w})$.

Let \mathbb{F}_V be the group freely generated by the vertex set V. For every vertex $v \in V(\Gamma)$, let f_v be the fibre curve of the circle bundle S_v over the point $\gamma(v)$, and let c_v be the unique path from the root r to v in \mathcal{T} . Then define the *meridian* of v as the curve $\mu_v := \gamma(c_v) \cdot f_v \cdot \gamma(c_v)^{-1}$ on $B_{\mathcal{A}}$.

Finally, let also δ be an orientation on the graph Γ , that is a function assigning a sign $\delta_{v,w} = \pm 1$ to each edge $e_{v,w}$, with $\delta_{v,w} = -\delta_{w,v}$.

Theorem 6.3. Write $b := \gamma(r)$. Then there is an exact sequence

$$0 \longrightarrow P_{\Gamma}(\mathcal{T}, \gamma) \longrightarrow \mathbb{F}_{V} * \pi_{1}(\Gamma, r) \xrightarrow{\gamma_{\mathcal{T}}} \pi_{1}(B_{\mathcal{A}}, b) \longrightarrow 0$$

where $P_{\Gamma}(\mathcal{T},\gamma)$ is the subgroup of relations normally generated by

$$(i) \ \forall \ v \in V(\Gamma): \quad \mu_v^{\varepsilon_v} \cdot \prod_{k=1}^{m_v} \mu_{\theta_v^{-1}(k)}^{u(v,\theta_v^{-1}(k))}$$

$$(ii) \ \forall \ e_{v,w} \in \Gamma: \quad [\mu_v, \mu_w^{u(v,w)}]$$

$$where \ u(v,w) \coloneqq \begin{cases} (\gamma_{v,w})^{\delta_{v,w}} & \text{if } e_{v,w} \in \Gamma \smallsetminus \mathcal{T}. \\ 1 & \text{if } e_{v,w} \in \mathcal{T}. \end{cases}$$

For $x, y \in \pi_1(B_{\mathcal{A}}, b)$ the notations are $[x, y] = xyx^{-1}y^{-1}$ and $x^y = y^{-1}xy$.

Remark 6.4. This presentation can often be simplified, in particular when some Euler numbers ε_v are equal to 0, 1 or -1.

The first homology group $H_1(\Gamma)$ is a free abelian group generated by the cycles of the graph Γ . Note that again there is no canonical basis of cycle generators.

Theorem 6.5. Let $\gamma \in E_{\Gamma}(\Omega)$ be a graphed embedding. Then γ induces a group isomorphism

$$Y_*: M(\Gamma) \oplus H_1(\Gamma) \xrightarrow{\sim} H_1(B_\mathcal{A})$$

Proof. Abelianising the exact sequence of Theorem 6.3 yields a new exact sequence

which is exactly the defining exact sequence of $M(\Gamma) \oplus H_1(\Gamma)$. The morphism γ_* is defined as the morphism induced by $Ab(\gamma)$ on the quotient, which does not depend on the choice of the parameter \mathcal{T} .

Proposition 6.6. For every pair of graphed embeddings $\gamma, \gamma' \in E_{\Gamma}(\Omega)$, we have

$$\left(\gamma_*^{-1} \circ \gamma'_*\right)_{|M(\Gamma)} = \mathrm{Id}_{M(\Gamma)} \qquad \vartriangleleft$$

Proof. For every $v \in V(\Gamma)$, $\gamma_*(\bar{v})$ is equal to the homological class of the meridian μ_v in $H_1(B_A)$, which by construction is equal to the class of the fibre f_v over $\gamma(v)$ in $H_1(S_v)$. Since all fibres of S_v have the same homological class, the image $\gamma_*(\bar{v})$ does not depend on the choice of γ . \Box

6.2. Definition of the invariant. Let \mathcal{A} be a line arrangement and write $\Gamma := \widetilde{\Gamma}(C_{\mathcal{A}})$ its reduced incidence graph. Consider the inclusion map

$$i_{\mathcal{A}}: B_{\mathcal{A}} \longrightarrow E_{\mathcal{A}}$$

and the induced map on the first homology groups

$$i_{\mathcal{A}}^*: H_1(B_{\mathcal{A}}, \mathbb{Z}) \longrightarrow H_1(E_{\mathcal{A}}, \mathbb{Z})$$

Let Ω be an ordering of the graph Γ . By Theorem 6.5, every graphed embedding $\gamma \in E_{\Gamma}(\Omega)$ induces an isomorphism

$$\gamma_*: M(\Gamma) \oplus H_1(\Gamma) \xrightarrow{\sim} H_1(B_{\mathcal{A}}, \mathbb{Z})$$

By Proposition 6.2, there is a natural group isomorphism $H_1(E_A, \mathbb{Z}) \simeq M(\Gamma)$. Therefore, we have a map

$$i_{\mathcal{A}}^* \circ \gamma_* : M(\Gamma) \oplus H_1(\Gamma) \longrightarrow M(\Gamma)$$

Lemma 6.7. For every graphed embedding $\gamma \in E_{\Gamma}(\Omega)$, we have

$$(i_{\mathcal{A}}^* \circ \gamma_*)_{|M(\Gamma)} = \mathrm{Id}_{M(\Gamma)} \qquad \vartriangleleft$$

Proof. For every line L of \mathcal{A} , $i_{\mathcal{A}}^* \circ \gamma_*(\bar{L})$ is equal to the homology class of the fibre f_L over $\gamma(L)$ inside S_L , which is exactly the generator of $H_1(E_{\mathcal{A}})$ corresponding to \bar{L} by Proposition 6.2. \Box

Now consider the restriction

$$(i_{\mathcal{A}}^* \circ \gamma_*)_{|H_1(\Gamma)} \in \operatorname{Hom}(H_1(\Gamma), M(\Gamma))$$

which we simply write $i_{\mathcal{A}}^* \circ \gamma_*$ for short. This morphism obviously depends on the choice of the graphed embedding γ . The graph stabiliser is a quotient designed to express the morphism $i_{\mathcal{A}}^*$ in a way that no longer depends on the choice of $\gamma \in E_{\Gamma}(\Omega)$ but only on the choice of the graph ordering Ω itself.

Definition 6.8. The graph stabiliser $\mathcal{G}_{\Gamma}(\Omega)$ is defined as the quotient of $\operatorname{Hom}(H_1(\Gamma), M(\Gamma))$ by the subgroup

$$\left\langle \phi \circ (\gamma_* - \gamma'_*)_{|H_1(\Gamma)} \middle| \begin{array}{c} \gamma, \gamma' \in \mathcal{E}_{\Gamma}(\Omega), \phi \in \operatorname{Hom}(H_1(B_{\mathcal{A}}), M(\Gamma)) \\ (\phi \circ \gamma_*)_{|M(\Gamma)} = \operatorname{Id}_{M(\Gamma)} \end{array} \right\rangle \qquad \diamond$$

All objects involved in Definition 6.8, including $H_1(B_A)$ by Theorem 6.5, depend only on the the combinatorics Γ of A and the choice of a graph ordering Ω . However, this last dependency can actually be lifted.

Theorem 6.9. The group isomorphism type of the graph stabiliser $\mathcal{G}_{\Gamma}(\Omega)$ is the same for all graph orderings Ω on Γ .

The proof of Theorem 6.9 is obtained by establishing a presentation of the graph stabiliser, which is the object of Section 7. From now on we simply write \mathcal{G}_{Γ} .

We are now ready to define the homology inclusion invariant.

Definition 6.10. Let \mathcal{A} be an ordered line arrangement and let $\Gamma := \Gamma(C_{\mathcal{A}})$ be its incidence graph. Fix an ordering Ω of the graph. Then for any graphed embedding $\gamma \in E_{\Gamma}(\Omega)$ the class

$$\mathcal{J}_{\Omega}(\mathcal{A}) := [i_{\mathcal{A}}^* \circ \gamma_*] \in \mathcal{G}_{\Gamma}$$

is called the *homology inclusion* of the ordered line arrangement.

Lemma 6.7 ensures that $i_{\mathcal{A}}^*$ respects the condition of the function ϕ in Definition 6.8, so by construction of \mathcal{G}_{Γ} the class $[i_{\mathcal{A}}^* \circ \gamma_*]$ does not depend on the choice of $\gamma \in E_{\Gamma}(\Omega)$. Therefore $\mathcal{J}_{\Omega}(\mathcal{A})$ is well-defined. Our main result states that this class element of the graph stabiliser \mathcal{G}_{Γ} is a topological invariant of ordered oriented line arrangements.

Theorem 6.11. Let $\mathcal{A}, \mathcal{A}' \subset \mathbb{CP}^2$ be two non-exceptional line arrangements with the same combinatorics C. If there exists a positive ordered equivalence between \mathcal{A} and \mathcal{A}' then for any graph ordering Ω of the incidence graph Γ we have $\mathcal{J}_{\Omega}(\mathcal{A}) = \mathcal{J}_{\Omega}(\mathcal{A}')$ inside the graph stabiliser \mathcal{G}_{Γ} .

Proof. The topological equivalence between \mathcal{A} and \mathcal{A}' induces homeomorphisms Ψ and Φ such that the following diagram commutes:

$$\begin{array}{cccc}
B_{\mathcal{A}'} & \xrightarrow{\Psi} & B_{\mathcal{A}} \\
\downarrow^{i}_{\mathcal{A}'} & & & \downarrow^{i}_{\mathcal{A}} \\
E_{\mathcal{A}'} & \xrightarrow{\sim} & E_{\mathcal{A}}
\end{array}$$

Write $B = B_{\mathcal{A}'} \simeq B_{\mathcal{A}}$. By Proposition 3.8, we can always suppose that $\Psi \in \operatorname{Homeo}_{\Gamma}^{++}(B)$. Let $\gamma \in E_{\Gamma}(\Omega)$ be an ordered graph embedding. By Proposition 5.2, the image $\Psi(\gamma)$ is again an element of $E_{\Gamma}(\Omega)$. The map Ψ induces a group automorphism $\Psi_* : H_1(B, \mathbb{Z}) \to H_1(B, \mathbb{Z})$. By construction we have $\Psi(\gamma)_* = \Psi_* \circ \gamma_*$ and $(i_{\mathcal{A}} \circ \Psi)_* = i_{\mathcal{A}}^* \circ \Psi_*$. Similarly, the map Φ induces $\Phi_* : H_1(E_{\mathcal{A}'}, \mathbb{Z}) \to H_1(E_{\mathcal{A}}, \mathbb{Z})$. By Proposition 6.2 these last two groups are both naturally isomorphic to $M(\Gamma)$ generated by the line meridians. Since Φ is an ordered equivalence it induces the identity between the sets of lines of \mathcal{A} and \mathcal{A}' , therefore $\Phi_* = \operatorname{Id}_{M(\Gamma)}$. Then inside $\operatorname{Hom}(H_1(\Gamma), M(\Gamma))$ we have

$$^*_{\mathcal{A}'} \circ \gamma_* = \Phi^{-1}_* \circ i^*_{\mathcal{A}} \circ \Psi_* \circ \gamma_* = i^*_{\mathcal{A}} \circ \Psi(\gamma)_*$$

By Definition 6.8 of the graph stabiliser \mathcal{G}_{Γ} , this implies that inside the quotient:

i

$$0 = [i_{\mathcal{A}}^* \circ (\Psi(\gamma)_* - \gamma_*)] = [i_{\mathcal{A}'}^* \circ \gamma_*] - [i_{\mathcal{A}}^* \circ \gamma_*] = \mathcal{J}_{\Omega}(\mathcal{A}') - \mathcal{J}_{\Omega}(\mathcal{A}) \qquad \Box$$

Remark 6.12. By definition of the graph stabiliser, the class $[i_{\mathcal{A}}^* \circ \gamma_*] \in \mathcal{G}_{\Gamma}$ does not depend on the graphed embedding $\gamma_* \in E_{\Gamma}(\Omega)$. However, it does depend on the graph ordering Ω which is preserved only by ordered equivalences. The homology inclusion is therefore an *ordered*

 \diamond

line arrangement invariant. However, some combinatorics have trivial automorphism groups. The restriction of the homology inclusion to this subclass of line arrangements becomes an unordered topological invariant. \square

Remark 6.13. The graph stabiliser does not quotient the homology differences caused by the application of the complex conjugation inside \mathbb{CP}^2 , since it is not a positive homeomorphism of the boundary manifold. A line arrangement \mathcal{A} and its conjugate $\overline{\mathcal{A}}$ may thus have different homology inclusion values in general.

7. PRESENTATION OF THE GRAPH STABILISER

The graph stabiliser \mathcal{G}_{Γ} is the underlying group where the homology inclusion invariant is defined. In order to make this invariant effective, one needs to be able to not only compute the invariant itself, but also the graph stabiliser. The objective of this section is to establish Theorem 7.11 which gives an explicit finite presentation of the graph stabiliser. To obtain this result we analyse further the objects involved in Definition 6.8 of the graph stabiliser in order to actually compute the homological difference between every two graphed embeddings. The crucial step to achieve this is Proposition 5.5 which allows to reduce this difference computation to a finite number of cases.

In addition, the presentation of the graph stabiliser is the main tool used in Section 8 to establish the connection between the homology inclusion and the pre-existing loop-linking number invariant.

Since the graph stabiliser is a combinatorial object, in this section we fix a graph Γ . For any line arrangement \mathcal{A} with graph Γ , the boundary manifold $B_{\mathcal{A}}$ as a topological object only depends on the choice of Γ , see Theorem 3.5.

We reuse all notations introduced at the beginning of Section 6.1. Recall that Γ has a natural structure of a CW-complex generated by the vertices and the edges. The first homology group $H_1(\Gamma)$ is the kernel of the boundary map $\partial_1 : C_1(\Gamma) \to C_0(\Gamma)$. There is an inclusion homomorphism

$$\zeta: H_1(\Gamma) \longrightarrow C_1(\Gamma)$$

which decomposes every cycle into the sum of its oriented edges.

Let us denote by $\vec{e}_{v,w}$ the *half-edge* of Γ starting from v and going towards w (and reciprocally for $\vec{e}_{w,v}$). Let $\vec{C}_1(\Gamma)$ be the free abelian group generated by all half-edges. Fix an orientation δ of Γ . There is a natural map $\chi_{\delta} : C_1(\Gamma) \to \vec{C}_1(\Gamma)$ defined by $\chi_{\delta}(e_{v,w}) := \delta_{v,w} (\vec{e}_{v,w} - \vec{e}_{w,v})$. There is also a decomposition

$$\vec{C}_1(\Gamma) = \bigoplus_{v \in V(\Gamma)} \left\langle \vec{W}_v \right\rangle$$

where $\vec{W}_v := \{ \vec{e}_{v,w} \mid w \in W_v \} \simeq W_v.$

Finally, if A, B are two free abelian groups, we denote by A^{\dagger} the dual \mathbb{Z} -module of A. We make the natural identification $\operatorname{Hom}(A, B) \simeq A^{\dagger} \otimes B$, and for every homomorphism $\phi : A \to B$ we denote by $\phi^{\dagger} : B^{\dagger} \to A^{\dagger}$ its dual homomorphism. In addition, if we fix a basis \mathcal{B} of the free module A, then for every element $a \in A$ we denote by $a^{\ddagger} \in A^{\dagger}$ its dual element, which decomposes with the same coefficients in the dual basis of \mathcal{B} . In particular the free modules $C_0(\Gamma), C_1(\Gamma), \vec{C}_1(\Gamma)$ (and the modules of morphisms between them) all have obvious canonical bases which we use to define their dual elements.

7.1. Difference maps. The difference maps are used to compute the homological difference between two ordered graphed embeddings. This difference lies exclusively on the cycle generators of $H_1(B_A)$ since by Proposition 6.6 the homological values of two ordered graphed embeddings always coincide on the meridian generators. We build the difference maps step by step, by

decomposing both the graph manifold and the graphed embeddings into their elementary parts and then computing the homological difference at every level.

Two stars α, α' in a manifold M are called *joined* if their centres coincide, and if for every $w \in W$ the endpoint of their corresponding branches α^w and ${\alpha'}^w$ also coincide.

Definition 7.1. Let $\bar{\alpha}, \bar{\alpha}' \in \mathrm{LS}_{\theta}(D_W)$ be two *joined* linear stars on the disc D_W . The *star* difference map is the map

$$\nabla_W(\bar{\alpha}, \bar{\alpha}') : W \longrightarrow H_1(D_W)$$

that sends every $w \in W$ to the class of the closed curve

$$\bar{\lambda}^w := \bar{\alpha}^w \cdot \left(\bar{\alpha}'^w\right)^{-1} \qquad \diamond$$

Consider now two ordered graphed embeddings $\gamma, \gamma' \in \mathcal{E}_{\Gamma}(\Omega)$ and fix $v \in V(\Gamma)$. Write $\alpha_v = \gamma \cap S_v$ and $\alpha'_v = \gamma' \cap S_v$ the corresponding stars on S_v . By Lemmas 5.3 and 5.4, we have $\alpha_v = \check{s}_v(\bar{\alpha}_v)$ and $\alpha'_v = \check{s}_v(\bar{\alpha}'_v)$ with $\bar{\alpha}_v, \bar{\alpha}'_v \in \mathrm{LS}_{\theta_v}(D_{W_v})$.

Lemma 7.2. There exists an isotopy of B_A that makes α_v and α'_v joined inside S_v .

Proof. Fix $w \in W_v$. Since the gluing map between S_v and S_w permutes meridians and longitudes, $\check{s}_v \cap \check{s}_w \subset B_A$ is reduced to a single point $t_{v,w} \in T_{v,w}$. Applying Lemma 5.3 in S_v gives an isotopy of B_A that projects the two stars α_v, α'_v on \check{s}_v . Applying Lemma 5.3 now in S_w gives another isotopy of B_A that projects $\gamma \cap S_w$ and $\gamma' \cap S_w$ to \check{s}_w . This second isotopy acts on $T_{v,w}$ and therefore extends to S_v where it moves α_v and α'_v inside \check{s}_v to make them meet $t_{v,w}$. Note that any of these isotopies might introduce full loops around a fibre in S_v or S_w . These loops can always be 'pushed' to the other side of the gluing, where they will become loops in \check{s}_w or \check{s}_v . Repeat the process for all $w \in W_v$. Separately, the centres of α_v and α_w both lie on $\partial^{\infty} D_{W_v}$. Using an isotopy of S_v in the neighbourhood of $\partial^{\infty} D_{W_v}$ we can always bring one centre to the other.

Lemma 7.2 also implies that $\bar{\alpha}_v$ and $\bar{\alpha}'_v$ are joined inside $\check{s}_v \simeq D_{W_v}$.

Definition 7.3. Let $v \in V(\Gamma)$ be a vertex of the graph Γ . Let $\langle W_v \rangle$ be the submodule of $C_0(\Gamma)$ freely generated by the neighbour set W_v . The homological neighbour map is the group isomorphism

$$h_v: \langle W_v \rangle \longrightarrow H_1(D_{W_v})$$

that sends every neighbour vertex $w \in W_v$ to the class of the curve $\partial^w D_{W_v}$. \diamond **Definition 7.4.** The *half-edge difference map* is the morphism

$$\vec{\nabla} : \mathbf{E}_{\Gamma}(\Omega)^2 \longrightarrow \mathrm{Hom}\left(\vec{C}_1(\Gamma), C_0(\Gamma)\right)$$

defined on every generating subset \vec{W}_v of $\vec{C}_1(\Gamma)$ by:

$$\vec{\nabla}(\gamma,\gamma')_{|\vec{W}_v} \coloneqq h_v^{-1} \circ \nabla_{W_v}(\bar{\alpha}_v,\bar{\alpha}'_v) \qquad \diamond$$

Proposition 7.5. For every $\gamma, \gamma', \gamma'' \in E_{\Gamma}(\Omega)$, the half-edge difference map $\vec{\nabla}$ verifies the following properties:

(i) $\vec{\nabla}(\gamma, \gamma)$ is the trivial homomorphism.

(ii)
$$\vec{\nabla}(\gamma, \gamma') = -\vec{\nabla}(\gamma', \gamma).$$

iii)
$$\vec{\nabla}(\gamma, \gamma'') = \vec{\nabla}(\gamma, \gamma') + \vec{\nabla}(\gamma', \gamma'').$$

Definition 7.6. Let δ be an orientation of Γ . The cycle difference map is the map

$$\nabla : \mathrm{E}_{\Gamma}(\Omega)^2 \longrightarrow \mathrm{Hom}\left(H_1(\Gamma), M(\Gamma)\right)$$

defined by

$$\nabla(\gamma,\gamma') := ((\chi_{\delta} \circ \zeta)^{\dagger} \otimes \eta) \circ \vec{\nabla}(\gamma,\gamma')$$

where $\eta : C_0(\Gamma) \twoheadrightarrow M(\Gamma)$ is the projection from Definition 6.1.

Remark 7.7. The cycle difference map ∇ verifies properties similar to Proposition 7.5 which are induced by the properties of the half-edge difference map $\vec{\nabla}$.

As announced, the cycle difference map gives a first reformulation of the definition of the graph stabiliser.

Proposition 7.8. Let $\gamma, \gamma' \in E_{\Gamma}(\Omega)$. Then

$$\gamma_* \circ \nabla(\gamma, \gamma') = (\gamma_* - \gamma'_*)_{|H_1(\Gamma)} \in \operatorname{Hom}(H_1(\Gamma), H_1(B_{\mathcal{A}})) \qquad \vartriangleleft$$

Theorem 7.9. There is a natural identification

$$\mathcal{G}_{\Gamma}(\Omega) \simeq \operatorname{Hom}(H_1(\Gamma), M(\Gamma)) / \langle \nabla(\gamma, \gamma') \mid \gamma, \gamma' \in \mathcal{E}_{\Gamma}(\Omega) \rangle \qquad \vartriangleleft$$

Proof. Let $\gamma, \gamma' \in E_{\Gamma}(\Omega)$ and $\phi \in \text{Hom}(H_1(B_{\mathcal{A}}), M(\Gamma))$ such that $(\phi \circ \gamma_*)_{|M(\Gamma)} = \text{Id}_{M(\Gamma)}$ as in Definition 6.8 of $\mathcal{G}_{\Gamma}(\Omega)$. Then by Proposition 7.8:

$$\phi \circ (\gamma_* - \gamma'_*)_{H_1(\Gamma)} = \phi \circ \gamma_* \circ \nabla(\gamma, \gamma') = \nabla(\gamma, \gamma') \qquad \Box$$

The remainder of this subsection is dedicated to the proof of Proposition 7.8. Let $e_{v,w}$ be an edge. Reusing previous notations, we write:

$$\gamma(\vec{e}_{v,w}) = \check{s}_v(\bar{\alpha}_v^w) \qquad \qquad \gamma(\vec{e}_{w,v}) = \check{s}_w(\bar{\alpha}_w^v)$$

and similarly for γ' .

Lemma 7.10. Let $i_v: S_v \hookrightarrow B_{\mathcal{A}}$ be the inclusion. The following diagram commutes

$$\begin{array}{c} H_1\left(D_{W_v}\right) \xrightarrow{i_v^* \circ S_v^*} H_1(B_{\mathcal{A}}) \\ \\ h_v \uparrow & \uparrow^{\gamma_*} \\ \langle W_v \rangle \xrightarrow{\eta} M(\Gamma) \end{array} \end{array}$$

Proof. For any $w \in W_v$, the closed curve $\partial^w \check{s}_v$ inside S_v is sent by the gluing map along the torus $T_{v,w}$ to a fibre of S_w . Its homological class inside $H_1(B_A)$ is thus equal to f_w . This precisely means that $i_v^* \circ \check{s}_v^* \circ h_v(w) = f_w$. But by Theorem 6.5, we also have $\gamma_*(\overline{w}) = f_w$ by construction of γ_* .



FIGURE 7. Difference between two graphed embeddings

 \diamond

Proof of Proposition 7.8. Let $c \in H_1(\Gamma)$ and write $\zeta(c) = \sum_k e_{v_k, v_{k+1}}$. By construction of the embedding $\gamma \in E_{\Gamma}(\Omega)$ we have

$$\gamma(\zeta(c)) = \prod_k \gamma(e_{v_k, v_{k+1}})$$

Inside $H_1(M)$, $(\gamma_* - \gamma'_*)(c)$ can be seen as the class of the closed curve

$$\gamma(\zeta(c)) \cdot \gamma'(\zeta(c))^{-}$$

As shown on Figure 7, a 2-chain bordering that curve inside M can be decomposed into a sum of 2-chains bordering each of the closed curves

$$\lambda_{v_k}^{v_{k+1}} \coloneqq \alpha_{v_k}^{v_{k+1}} \cdot \left(\alpha_{v_k}'^{v_{k+1}}\right)^{-1} \qquad \qquad \lambda_{v_{k+1}}^{v_k} \coloneqq \alpha_{v_{k+1}}^{v_k} \cdot \left(\alpha_{v_{k+1}}'^{v_k}\right)^{-1}$$

lying in \check{s}_{v_k} and $\check{s}_{v_{k+1}}$ respectively. Given an orientation δ , this yields the equation inside $H_1(B_A)$:

$$(\gamma_* - \gamma'_*)(c) = \sum_k \delta_k \left([\lambda_{v_k}^{v_{k+1}}] - [\lambda_{v_{k+1}}^{v_k}] \right)$$

where $[\cdot]$ designates the class of the closed curve inside $H_1(B_A)$ and $\delta_k := \delta_{v_k, v_{k+1}}$. Applying successively Lemma 7.10 and Definition 7.4 of the half-edge difference map $\vec{\nabla}(\gamma, \gamma')$, we get the following equality:

$$\begin{split} [\lambda_{v_{k}+1}^{v_{k+1}}] &- [\lambda_{v_{k+1}}^{v_{k}}] = i_{v_{k}}^{*} \circ \check{s}_{v_{k}}^{*} \circ h_{v_{k}} \circ h_{v_{k}}^{-1} (\bar{\lambda}_{v_{k}+1}^{v_{k+1}}) - i_{v_{k+1}}^{*} \circ \check{s}_{v_{k+1}}^{*} \circ h_{v_{k+1}} \circ h_{v_{k+1}}^{-1} (\bar{\lambda}_{v_{k+1}}^{v_{k}}) \\ &= \gamma_{*} \circ \eta \circ \left(\vec{\nabla}(\gamma, \gamma') (\vec{e}_{v_{k}, v_{k+1}}) - \vec{\nabla}(\gamma, \gamma') (\vec{e}_{v_{k+1}, v_{k}}) \right) \end{split}$$

Replacing in the sum yields:

$$\begin{aligned} (\gamma_* - \gamma'_*)(c) &= \sum_k \delta_k \cdot \gamma_* \circ \eta \circ \left(\vec{\nabla}(\gamma, \gamma')(\vec{e}_{v_k, v_{k+1}}) - \vec{\nabla}(\gamma, \gamma')(\vec{e}_{v_{k+1}, v_k}) \right) \\ &= \gamma_* \circ \left((\chi_\delta \circ \zeta)^{\dagger} \otimes \eta \right) \circ \vec{\nabla}(\gamma, \gamma')(c) \\ &= \gamma_* \circ \nabla(\gamma, \gamma')(c) \end{aligned}$$

7.2. Relations of the graph stabiliser. The results obtained in Section 4.4 allows us to compute explicitly the image of the cycle difference map ∇ . Combining this with Theorem 7.9 gives an explicit combinatorial presentation of the graph stabiliser $\mathcal{G}_{\Gamma}(\Omega)$.

Theorem 7.11. The group $\mathcal{G}_{\Gamma}(\Omega)$ is finitely presented and admits the presentation:

Generators: $c \otimes \overline{v}$ for every $c \in H^1(\Gamma)$ and $v \in V(\Gamma)$.

Relations: the images by
$$\zeta^{\dagger} \otimes \eta$$
: Hom $(C_1(\Gamma), C_0(\Gamma)) \to$ Hom $(H_1(\Gamma), M(\Gamma))$ of

- (GS1) $e_{v,x}^{\ddagger} \otimes v \text{ and } e_{v,x}^{\ddagger} \otimes x \text{ for every edge } e_{v,x}.$
- (GS2) $e_{v,y}^{\ddagger} \otimes z + \delta_{v,y} \delta_{v,z} \cdot e_{v,z}^{\ddagger} \otimes y$ for every pair $(e_{v,y}, e_{v,z})$ of adjacent edges in Γ . \triangleleft Note that Theorem 7.11 directly implies Theorem 6.9.

Proof. We reuse the notations from Sections 4 and 7.1. By Theorem 7.9 we need to compute

$$\langle \nabla(\gamma, \gamma') \mid \gamma, \gamma' \in \mathcal{E}_{\Gamma}(\Omega) \rangle \subset H^{1}(\Gamma) \otimes M(\Gamma)$$

Applying Definition 7.6, this subgroup is isomorphic to the image by $(\chi_{\delta} \circ \zeta)^{\dagger} \otimes \eta$ of the subgroup

$$\left\langle \vec{\nabla}(\gamma,\gamma') \mid \gamma \in \mathcal{E}_{\Gamma}(\Omega), \Psi \in \operatorname{Homeo}_{\Gamma}^{++}(M) \right\rangle \subset \vec{C}^{1}(\Gamma) \otimes C_{0}(\Gamma)$$

By Proposition 5.5, this subgroup is in turn isomorphic to

 $\vec{G}_{\Gamma} := \left\langle \vec{\nabla}(\gamma, \Psi \cdot \gamma) \mid \gamma \in \mathcal{E}_{\Gamma}(\Omega), \Psi \in \operatorname{Homeo}_{\Gamma}^{++}(M) \right\rangle$

Therefore, we need to determine a generating set of \vec{G}_{Γ} as a submodule of $\operatorname{Hom}(\vec{C}_1(\Gamma), C_0(\Gamma))$.

Let $\Psi \in \text{Homeo}_{\Gamma}^{++}(M)$ and $\gamma \in E_{\Gamma}(\Omega)$. Fix $v \in V(\Gamma)$ and write $\alpha_v = \gamma \cap S_v$. Up to isotopy, Ψ restricts to a fibre-positive homeomorphism Ψ_v of S_v and we have $\Psi \cdot \gamma \cap S_v = \Psi_v \cdot \alpha_v$. By Lemma 5.3, we can also suppose that $\gamma \cap S_v = \check{s}_v(\bar{\alpha}_v)$ where $\bar{\alpha}_v \in \text{LS}_{\theta_v}(D_{W_v})$. Since Ψ_v is fibrewise and fixes ∂S_v componentwise, its action on \check{s}_v is completely determined by an element $\psi_v \in \mathcal{P}(D_{W_v})$, and we have

$$\Psi \cdot \gamma \cap S_v = \check{s}_v \left(\psi_v \cdot \bar{\alpha}_v \right)$$

We can then rewrite $\vec{G}_{\Gamma} = \bigoplus_{v \in V(\Gamma)} \vec{G}_{\Gamma}^{v}$ with

$$\vec{G}_{\Gamma}^{v} := \left\langle h_{v}^{-1} \circ \nabla_{W_{v}}(\bar{\alpha}_{v}, \psi_{v} \cdot \bar{\alpha}_{v}) \mid \bar{\alpha}_{v} \in \mathrm{LS}_{\theta_{v}}(D_{W_{v}}), \psi_{v} \in \mathcal{P}(D_{W_{v}}) \right\rangle$$

where the domain W_v of $\nabla_{W_v}(\bar{\alpha}_v, \psi_v \cdot \bar{\alpha}_v)$ is viewed as the subset W_v of $C_1(\Gamma)$.

To compute a presentation of \vec{G}_{Γ}^{v} we will reuse the model disc D_{m} from Section 4.3. Denote by ∇_{m} the star difference map on D_{m} , and let

$$G_m := \left\langle h^{-1} \circ \nabla_m(\bar{\alpha}, \psi \cdot \bar{\alpha}) \mid \bar{\alpha} \in \mathrm{LS}(D_m), \psi \in \mathcal{P}\left(D_m\right) \right\rangle$$

where $h : \langle x^1, \ldots, x^m \rangle \to H_1(D_m)$ is the natural group isomorphism. By Proposition 4.11, the action of $\mathcal{P}(D_m)$ on $\mathrm{LS}(D_m)$ is transitive. Using the standard star $\bar{\alpha}_0$ from Figure 4, we can therefore rewrite

$$G_{m} := \left\langle h^{-1} \circ \nabla_{m}(\bar{\alpha}_{0}, \psi \cdot \bar{\alpha}_{0}) \mid \psi \in \mathcal{P}\left(D_{m}\right) \right\rangle$$

Lemma 7.12. The group G_m is freely generated by the union $R_1 \sqcup R_2$, where

$$R_{1} := \left\{ (x^{j})^{\ddagger} \otimes x^{j} \mid 1 \le j \le m \right\}$$

$$R_{2} := \left\{ (x^{k})^{\ddagger} \otimes (x^{k} + x^{l}) + (x^{l})^{\ddagger} \otimes (x^{k} + x^{l}) \mid 1 \le k < l \le m \right\}$$

For every $v \in V(\Gamma)$, the bijection $M_{\theta_v}: D_m \to D_{W_v}$ induces a bijection $G_{m_v} \simeq \tilde{G}_{\Gamma}^v$ given by

(7.1)
$$(x^k)^{\ddagger} \otimes x^l \mapsto \vec{e}_{v,y}^{\ddagger} \otimes z$$
, where $\theta_v(y) = k$ and $\theta_v(z) = l$
From Lemma 7.12, we get that

From Lemma 7.12, we get that

$$\vec{G}_{\Gamma}^{v} = \left\langle \vec{e}_{v,x}^{\dagger} \otimes x, \quad \vec{e}_{v,y}^{\dagger} \otimes (y+z) + \vec{e}_{v,z}^{\dagger} \otimes (y+z) \mid x, y, z \in W_{v} \right\rangle$$

This set of generators can be further simplified by subtracting the first generator from the second when x = y and x = z. Therefore

$$\vec{G}_{\Gamma}^{v} = \left\langle \vec{e}_{v,x}^{\dagger} \otimes x, \quad \vec{e}_{v,y}^{\dagger} \otimes z + \vec{e}_{v,z}^{\dagger} \otimes y \mid x, y, z \in W_{v} \right\rangle$$

Repeating this for all vertices $v \in V(\Gamma)$ gives the following generating set of \vec{G}_{Γ} :

(7.2)
$$\vec{G}_{\Gamma} = \left\langle \vec{e}_{v,x}^{\dagger} \otimes x, \quad \vec{e}_{v,y}^{\dagger} \otimes z + \vec{e}_{v,z}^{\dagger} \otimes y \mid \vec{e}_{v,x}, \vec{e}_{v,y}, \vec{e}_{v,z} \in \vec{C}_{1}(\Gamma) \right\rangle$$

Finally, for every edge $e_{v,w}$, one has

$$\chi^{\dagger}_{\delta}(\vec{e}^{\,\ddagger}_{v,w}) = -\chi^{\dagger}_{\delta}(\vec{e}^{\,\ddagger}_{w,v}) = \delta_{v,w} \cdot e^{\ddagger}_{v,u}$$

Therefore taking the image of (7.2) by χ^{\dagger}_{δ} gives exactly the set of generators of Theorem 7.11, now lying in Hom $(C_1(\Gamma), C_0(\Gamma))$ as required.

Proof of Lemma 7.12. By Proposition 4.6, every $\psi \in \mathcal{P}(D_m)$ is of the form (β, d) , with

$$\beta = \prod_{m=1}^{r} a_{k_m, l_m} \qquad \qquad d = \sum_{m=1}^{s} d_{j_m}$$

Each generator $a_{k,l}$ (resp. d_j) correspond to a full Dehn twist inside a curve $\delta_{k,l}$ (resp. δ_j). We say that a star $\bar{\alpha} \in LS(m)$ is in *standard position* if:

- (S1) for all curves δ_j , the pair $(\bar{\alpha}, \delta_j)$ is isotopic to Figure 9a.
- (S2) for all curves $\delta_{k,l}$, the pair $(\bar{\alpha}, \delta_{k,l})$ is isotopic to Figure 8a with possibly several (or none) parallel 'middle lines', all of which are parts of $\bar{\alpha}$.

We will prove by double induction on $r\geq 0$ and $s\geq 0$ that

$$(\mathbf{P}_{r,s}) \qquad \begin{cases} h^{-1} \circ \nabla_m(\bar{\alpha}_0, \bar{\alpha}_0 \cdot (\beta, d)) \in R_1 \sqcup R_2 \\ \bar{\alpha}_0 \cdot (\beta, d) \text{ is in standard position.} \end{cases}$$



(B) Action of $a_{k,l}$

FIGURE 8. Action of \mathbb{P}_m on a linear star



FIGURE 9. Action of \mathbb{Z}^m on a linear star

The standard star $\bar{\alpha}_0$ is clearly in standard position, with no middle lines. Now fix $r, s \ge 0$ and suppose that $\bar{\alpha} := \bar{\alpha}_0 \cdot (\beta, d)$ verifies $(\mathbf{P}_{r,s})$.

Let $1 \leq j \leq m$. Since $\bar{\alpha}$ is in standard position, the action of d_j on $\bar{\alpha}$ is shown on Figure 9. By superimposing Figures 9a and 9b, one sees that

(7.3)
$$h^{-1} \circ \nabla_m(\bar{\alpha}, \bar{\alpha} \cdot (1, d_j)) = (x^j)^{\downarrow} \otimes x^j$$

and therefore

$$h^{-1} \circ \nabla_m(\bar{\alpha}_0, \bar{\alpha} \cdot (1, d_j)) = h^{-1} \left(\nabla_m(\bar{\alpha}_0, \bar{\alpha}) + \nabla_m(\bar{\alpha}, \bar{\alpha} \cdot (1, d_j)) \right) \in R_1 \sqcup R_2$$

By retracting the curve δ_j closer to $\partial^j D_m$ on Figure 9b, we can put $d_j \cdot \bar{\alpha}$ back on standard position with respect to all δ_j 's. By retracting the curves $\delta_{j,k}$ for every $k \neq j$, we can also put $d_j \cdot \bar{\alpha}$ in standard position with respect to them, with one middle line added.

Let $1 \leq k < l \leq m$. Since $\bar{\alpha}$ is in standard position, the action of $a_{k,l}$ on $\bar{\alpha}$ is shown on Figure 8. By superimposing Figures 8a and 8b, one sees that

(7.4)
$$h^{-1} \circ \nabla_m(\bar{\alpha}, \bar{\alpha} \cdot (a_{k,l}, 0)) = (x^k)^{\ddagger} \otimes (x^k + x^l) + (x^l)^{\ddagger} \otimes (x^k + x^l)$$

and therefore

$$h^{-1} \circ \nabla_m(\bar{\alpha}_0, \bar{\alpha} \cdot (a_{k,l}, 0)) = h^{-1} \circ (\nabla_m(\bar{\alpha}_0, \bar{\alpha}) + \nabla_m(\bar{\alpha}, \bar{\alpha} \cdot (a_{k,l}, 0))) \in R_1 \sqcup R_2$$

In particular the middle lines do not contribute to the homological difference. By retracting the curves δ_k and δ_l on Figure 8b, we can put $\bar{\alpha} \cdot a_{k,l}$ back on standard position with respect to all δ_j 's. Now let $1 \leq k' < l' \leq m$ be another pair and suppose that (say) l = k'. To put $\bar{\alpha} \cdot a_{k,l}$ back on standard position, we can retract $\delta_{l,l'}$ closer to $\partial^l D_m$ on Figure 8a. This creates some new middle lines for $\delta_{l,l'}$. Similarly, $\delta_{l,k}$ itself can be retracted closer to $\partial^l D_m$ and $\partial^k D_m$, and keep the same middle lines.

This achieves the induction, and proves that $G_m \subset \langle R_1 \sqcup R_2 \rangle$. Conversely, applying (7.3) and (7.4) to $\bar{\alpha}_0$ for all d_j 's and all $a_{k,l}$'s shows that $R_1 \sqcup R_2 \subset G_m$.

7.3. Change of graph ordering. By Theorem 6.9 the graph stabiliser \mathcal{G}_{Γ} does not depend as a group on the choice of a graph ordering Ω . However, the original presentation of \mathcal{G}_{Γ} given in Definition 6.8 does involve a choice of Ω , and it is this presentation that is used to define (and compute) the homology inclusion invariant \mathcal{J}_{Ω} . Proposition 7.13 defines an explicit transition function T that describes the effect of changing Ω in the original presentation of \mathcal{G}_{Γ} .

We reuse notations from Section 2.2. A graph ordering Ω on Γ is an element of the set

$$\mathcal{C}(\Gamma) := \prod_{v \in V(\Gamma)} \mathfrak{C}(W_v)$$

Consider the group of permutations on graph orderings given by:

$$\mathfrak{S}_{\Gamma} := \prod_{v \in V(\Gamma)} \mathfrak{S}_{m_v}$$

The group \mathfrak{S}_{Γ} has a transitive right action on $\mathcal{C}(\Gamma)$ by Lemma 2.5.

Fix a vertex $v \in V(\Gamma)$. For $\sigma \in \mathfrak{S}_{m_v}$ acting on W_v , let σ^v be the element of \mathfrak{S}_{Γ} defined as σ on W_v and as the identity on every other neighbour set. The group \mathfrak{S}_{m_v} can be generated only by the permutations of type $\tau_k := (k, k+1)$ with the relations:

(7.5)
$$\tau_k^2 = \text{Id}$$
 $\tau_k \tau_j = \tau_j \tau_k \text{ if } |k-j| > 1$ $(\tau_k \tau_{k+1})^3 = \text{Id}$

Therefore the group \mathfrak{S}_{Γ} can be generated by the τ_k^v for every $v \in V(\Gamma)$ and $1 \leq k \leq m_v$.

Proposition 7.13. The group homomorphism

$$T: \mathfrak{S}_{\Gamma} \longrightarrow \mathcal{G}_{\Gamma}$$

given by, for every $v \in V(\Gamma)$ and $1 \leq k \leq m_v$:

(7.6)
$$T(\tau_k^v) = (\zeta^{\dagger} \otimes \eta) \left(\delta_{v,y} \cdot e_{v,y}^{\dagger} \otimes x \right) \quad \text{where } \theta_v(x) = k, \ \theta_v(y) = k+1$$

verifies the following property: if $\Omega, \Omega' \in \mathcal{C}(\Gamma)$ are such that $\Omega' = \Omega \cdot \Sigma$ with $\Sigma \in \mathfrak{S}_{\Gamma}$, then for every $\phi \in \operatorname{Hom}(H_1(B_{\mathcal{A}}), M(\Gamma))$ as in Definition 6.8 and every $\gamma \in \mathcal{E}_{\Gamma}(\Omega), \gamma' \in \mathcal{E}_{\Gamma}(\Omega')$, one has:

$$\phi \circ \gamma'_*] = [\phi \circ \gamma_*] + T(\Sigma)$$

inside \mathcal{G}_{Γ} .

The kernel of T contains the subgroup $\{(1 \cdots m_v)^v \mid v \in V(\Gamma)\}$ generated by the circular permutations that leave any circular order Ω on Γ unchanged.

Remark 7.14. From a computing perspective, Proposition 7.13 makes it possible to compute two homology inclusion values $\mathcal{J}_{\Omega}(\mathcal{A})$ and $\mathcal{J}_{\Omega'}(\mathcal{A}')$ with different graph orderings and still compare them. If there is an ordered oriented equivalence between \mathcal{A} and \mathcal{A}' , then the difference $\mathcal{J}_{\Omega'}(\mathcal{A}') - \mathcal{J}_{\Omega}(\mathcal{A})$ would be equal to $T(\Sigma)$ with $\Omega' = \Omega \cdot \Sigma$.

Proof. We first check that T respects the relations of the presentation of \mathfrak{S}_{m_v} given in (7.5). The second one is obviously respected. Write $\sigma = \tau_k^2$. Then

$$T(\sigma^{v}) = \left(\zeta^{\dagger} \otimes \eta\right) \left(\delta_{v,y} \cdot e_{v,y}^{\ddagger} \otimes x + \delta_{v,x} \cdot e_{v,x}^{\ddagger} \otimes y\right) = 0$$

by the relation (GS2) of \mathcal{G}_{Γ} . With $\tau = (\tau_k \tau_{k+1})^3$, a similar elementary computation shows that $T(\tau^v) = 0$. Therefore the group homomorphism T is well defined.

It is enough to prove the proposition when $\Sigma = \tau_k^v$ for a fixed $v \in V(\Gamma)$ and $1 \leq k \leq m_v$. Let ω_v be the circular order corresponding to v in Ω and fix a linearisation θ_v . By Lemma 5.3, we can suppose that $\gamma \cap S_v = \bar{\alpha}_v \in \mathrm{LS}_{\theta_v}(W_v)$. By Definition 6.8 of the graph stabiliser, the class $[\phi \circ \gamma_*] \in \mathcal{G}_{\Gamma}$ does not depend on the choice of $\gamma \in \mathrm{E}_{\Gamma}(\Omega)$, and therefore on the choice of $\bar{\alpha}_v \in \mathrm{LS}_{\theta_v}(W_v)$. Let $M_{\theta_v}: D_m \to D_{W_v}$ be the homeomorphism that sends each boundary component ∂D^j of D_m to $\partial^w D_{W_v}$ with $\theta_v(w) = j$. Then we can always suppose that $\bar{\alpha}_v = M_{\theta_v}(\bar{\alpha}_0)$ where $\bar{\alpha}_0$ is the standard ordered star in D_m shown on Figure 4.

Consider now the braid $\sigma_{k,k+1} \in \mathbb{B}_m$ that permutes the strands k and k+1. By Proposition 4.6, there is an associated homeomorphism in $\mathcal{M}(D_m)$. Its action on $\bar{\alpha}_0$ is shown on Figure 10. We thus obtain a new star $\bar{\alpha}_1 := M_{\theta_k} (\bar{\alpha}_0 \cdot \sigma_{k,k+1})$ in D_{W_v} which is now ordered by $\theta_v \cdot (k, k+1)$,



FIGURE 10. Action of \mathbb{B}_m on the standard linear star $\bar{\alpha}_0$.

and a new graphed embedding γ_1 ordered by $\Omega_1 = \Omega \cdot (k, k+1)^v = \Omega'$. Therefore $[\phi \circ (\gamma_1)_*] = [\phi \circ \gamma'_*]$.

Remember the difference maps from Section 7.1. Reusing the same tools from the proof of Theorem 7.11, we write:

$$\begin{aligned} [\phi \circ \gamma'_*] - [\phi \circ \gamma_*] &= [\phi \circ ((\gamma_1)_* - \gamma_*)] \\ &= \left((\chi_\delta \circ \zeta)^{\dagger} \otimes \eta \right) \left(\vec{\nabla} (\gamma, \gamma_1) \right) \\ &= \left((\chi_\delta \circ \zeta)^{\dagger} \otimes \eta \right) \left(h_v^{-1} \circ \nabla_{W_v} (\bar{\alpha}, \bar{\alpha}_1) \right) \end{aligned}$$

Using M_{θ_v} to go back to the model disc D_{m_v} , $h_v^{-1} \circ \nabla_{W_v}(\bar{\alpha}, \bar{\alpha}_1)$ is the image by (7.1) of $h^{-1} \circ \nabla_m(\bar{\alpha}_0, \bar{\alpha}_0 \cdot \sigma_{k,k+1})$. By superimposing Figures 10a and 10b, we see that

$$h^{-1} \circ \nabla_m(\bar{\alpha}_0, \bar{\alpha}_0 \cdot \sigma_{k,k+1}) = (x^{k+1})^{\ddagger} \otimes x^k$$

Applying the (7.1) isomorphism as well as χ_{δ} we get exactly the value of $T(\tau_k^v)$ from (7.6).

8. LOOP-LINKING NUMBER

The first homology groups of the exterior and boundary manifold of a line arrangement are determined by its combinatorics. However, the map $i_{\mathcal{A}}^*: H_1(B_{\mathcal{A}}) \to H_1(E_{\mathcal{A}})$ is not necessarily combinatorial and contains topological information. One of the first invariants developed from $i_{\mathcal{A}}^*$ was the \mathcal{I} -invariant from [FGM15], which was then extended into the loop-linking number introduced in [Cad18] and further developed in [Gue22]. In this section we prove that the the homology inclusion extends again these two invariants. Since all three are derived from $i_{\mathcal{A}}^*$, the difference lies mainly on the involved algebraic structures which play a role similar to the graph stabiliser. The fact that the loop-linking number generalises the \mathcal{I} -invariant is already covered in [Gue22]. We briefly restate the definitions of the loop-linking number reusing our own notations from Section 7.

Let \mathcal{A} be a line arrangement. Recall from Section 2.1 that $\widehat{\Gamma} := \widehat{\Gamma}(C_{\mathcal{A}})$ is the *full* incidence graph of \mathcal{A} . As in Section 6.2, consider the map induced by the inclusion on the first homology groups

$$i_{\mathcal{A}}^*: H_1(B_{\mathcal{A}}, \mathbb{Z}) \longrightarrow H_1(E_{\mathcal{A}}, \mathbb{Z})$$

Let $\widehat{\Omega}$ be a graph ordering on $\widehat{\Gamma}$. The full incidence graph $\widehat{\Gamma}$ gives a non-minimal graph structure of the boundary manifold $B_{\mathcal{A}}$. One can nevertheless define an embedding of the graph $\gamma : \widehat{\Gamma} \hookrightarrow B_{\mathcal{A}}$ again as a union of stars, one for each bundle piece. Then all the results of Section 6.1 stand for $\widehat{\Gamma}$, or any other graph structure of $B_{\mathcal{A}}$ for that matter, see [Wes97]. Lemma 6.7 stands as well, so for every graphed embedding $\gamma \in E_{\widehat{\Gamma}}(\widehat{\Omega})$ we have a map

$$i_{\mathcal{A}}^* \circ \gamma_* : H_1(\widehat{\Gamma}) \longrightarrow M(\widehat{\Gamma})$$

with $M(\widehat{\Gamma}) \simeq H_1(E_{\mathcal{A}}, \mathbb{Z}).$

Let $T_{\widehat{\Gamma}}$ be the subgroup of $\operatorname{Hom}(C_0(\widehat{\Gamma}), C_1(\widehat{\Gamma}))$ generated by the following elements:

(TLG1) $L'^{\ddagger} \otimes e_{P,L}$ for every edge $e_{P,L}$ and $L' \in W_P$.

(TLG2) ${P'}^{\ddagger} \otimes e_{P,L}$ for for every edge $e_{P,L}$ and $P' \in W_L$.

For every free abelian group A, denote by $\pi : A^{\dagger} \otimes A \to \mathbb{Z}$ the natural pairing, which we write as $\pi(a', a)$ for every pure tensor element $a' \otimes a \in A^{\dagger} \otimes A$.

Definition 8.1. The *tensor linking group* $\text{TLG}(\widehat{\Gamma})$ is the subgroup of $\text{Hom}(M(\widehat{\Gamma}), H_1(\widehat{\Gamma}))$ given by all elements ϕ such that for every $\psi \in T_{\widehat{\Gamma}}$:

$$\pi \left(\psi^{\ddagger}, (\eta^{\dagger} \otimes \zeta)(\phi)\right) = 0$$
 \diamond

Note that ψ^{\ddagger} designates the *dual element* of ψ lying in the dual module of Hom $(C_0(\widehat{\Gamma}), C_1(\widehat{\Gamma}))$, which is different from the *dual homomorphism* $\psi^{\dagger} \in \text{Hom}(C^1(\widehat{\Gamma}), C^0(\widehat{\Gamma}))$.

Now define

$$\Psi := \mathrm{Id} \otimes (i_{\mathcal{A}}^* \circ \gamma_*) : \mathrm{Hom}(M(\widehat{\Gamma}), H_1(\widehat{\Gamma})) \to \mathrm{Hom}(M(\widehat{\Gamma}), M(\widehat{\Gamma}))$$

Definition 8.2. The *loop-linking number* of \mathcal{A} is the function

$$\mathcal{L}(\mathcal{A}) : \mathrm{TLG}(\widehat{\Gamma}) \longrightarrow \mathbb{Z}$$

defined as the restriction of $\pi \circ \Psi$ on $\text{TLG}(\widehat{\Gamma})$.

The algebraic properties of $\text{TLG}(\widehat{\Gamma})$ ensure that $\mathcal{L}(\mathcal{A})$ depends neither on the choice of the graphed embedding γ nor on the choice of the graph ordering Ω , see [Cad18, Section 3.3.1]. Just like the homology inclusion, the loop-linking number is an invariant of *ordered and oriented* line arrangements, although this is not obvious from the definition.

Remark 8.3. The tensor linking group and loop-linking number are defined in [Cad18; Gue22] using homology with coefficients in an abelian group G. Since the graph stabiliser uses homology over \mathbb{Z} , we only state the definitions for this case.

We now establish that the homology inclusion extends the loop-linking number.

Theorem 8.4. Let $\mathcal{A}, \mathcal{A}' \subset \mathbb{CP}^2$ be two non-exceptional line arrangements with the same combinatorics C and same graph Γ . If there exists a graph ordering Ω of Γ such that $\mathcal{J}_{\Omega}(\mathcal{A}) = \mathcal{J}_{\Omega}(\mathcal{A}')$ then necessarily $\mathcal{L}(\mathcal{A}) = \mathcal{L}(\mathcal{A}')$.

The tensor-linking group plays a very similar role for the loop-linking number than the graph stabiliser does for the homology inclusion. Indeed, both are used to express the map $i_{\mathcal{A}}^*$ in a way that does not depend on the choice of the graphed embedding. However, the tensor-linking group is defined using every possible graphed embeddings. On the other hand, the graph stabiliser quotients out the differences between the graphed embeddings with the same graph ordering only. This gives a 'bigger' quotient with more possible values for the invariant, as evidenced by Lemma 8.6, and is the main reason why the homology inclusion extracts a finer topological information.

The rest of the section is dedicated to the proof of Theorem 8.4.

The following lemma can be established through elementary computations on the presentation of the graph stabiliser given in Theorem 7.11, see [Rod23, Section 1.6.4].

Lemma 8.5. There is a natural group isomorphism $\mathcal{G}_{\widehat{\Gamma}} \simeq \mathcal{G}_{\Gamma}$. In particular if $\widehat{\Omega}$ is a graph ordering on $\widehat{\Gamma}$ and Ω is its restriction on Γ (see Lemma 2.7) then for every graphed embeddings $\gamma \in E_{\Gamma}(\Omega)$ and $\widehat{\gamma} \in E_{\widehat{\Gamma}}(\widehat{\Omega})$ one has:

$$[i_{\mathcal{A}}^* \circ \gamma_*] = [i_{\mathcal{A}}^* \circ \widehat{\gamma}_*] \in \mathcal{G}_{\widehat{\Gamma}} \qquad \vartriangleleft$$

Let A, B be two free abelian groups with fixed bases. For any submodule H of $\text{Hom}(A, B) \simeq A^{\dagger} \otimes B$, we write

$$\begin{aligned} H^{\ddagger} &= \{\phi^{\ddagger} \mid \phi \in H\} \subset \operatorname{Hom}(A, B)^{\intercal} \simeq A \otimes B^{\dagger} \\ H^{\dagger} &= \{\phi^{\dagger} \mid \phi \in H\} \subset \operatorname{Hom}(B^{\dagger}, A^{\dagger}) \simeq B \otimes A^{\dagger} \end{aligned}$$

Let $G_{\widehat{\Gamma}}$ be the subgroup of $\operatorname{Hom}(C_1(\widehat{\Gamma}), C_0(\widehat{\Gamma}))$ generated by the elements of type (GS1) and (GS2) from Theorem 7.11.

Lemma 8.6. $G_{\widehat{\Gamma}}^{\dagger} \subset T_{\widehat{\Gamma}}^{\ddagger}$

Proof. The submodule $G_{\widehat{\Gamma}}^{\dagger} \subset C_0(\widehat{\Gamma}) \otimes C^1(\widehat{\Gamma})$ is generated by the following elements:

 \diamond

- (GS1') $P \otimes e_{P,L}^{\ddagger}$ and $L \otimes e_{P,L}^{\ddagger}$ for every edge $e_{P,L}$.
- (GS2') $L \otimes e_{P,L'}^{\ddagger} + \delta_{P,L} \delta_{P,L'} \cdot L' \otimes e_{P,L}^{\ddagger}$ for every $P \in \mathcal{Q}$ and $L, L' \in W_P$. $P \otimes e_{L,P'}^{\ddagger} + \delta_{L,P} \delta_{L,P'} \cdot P' \otimes e_{L,P}^{\ddagger}$ for every $L \in \mathcal{L}$ and $P, P' \in W_L$.

Separately, the submodule $T^{\ddagger}_{\widehat{\Gamma}} \subset C_0(\widehat{\Gamma}) \otimes C^1(\widehat{\Gamma})$ is generated by the following elements:

(TLG1') $L' \otimes e_{P,L}^{\ddagger}$ for every edge $e_{P,L}$ and $L' \in W_P$.

(TLG2') $P' \otimes e_{P,L}^{\ddagger}$ for for every edge $e_{P,L}$ and $P' \in W_L$.

Particularising (TLG1') when L' = L gives the first part of (GS1'), and doing the same with (TLG2') when P' = P gives the second part. It is also clear that the first (resp. second) part of (GS2') is generated by adding two elements of the first part of (TLG1') (resp. (TLG2')). Therefore $G_{\widehat{\Gamma}}^{\dagger} \subset T_{\widehat{\Gamma}}^{\ddagger}$.

Proof of Theorem 8.4. Consider the submodule of $\operatorname{Hom}(H_1(\widehat{\Gamma}), M(\widehat{\Gamma}))$ defined by

$$D := \left\langle i_{\mathcal{A}}^* \circ \gamma_* - i_{\mathcal{A}'}^* \circ \gamma_*' \mid \gamma, \gamma' \in \mathcal{E}_{\widehat{\Gamma}}(\widehat{\Omega}) \right\rangle$$

By assumption and by Definition 6.10 of the homology inclusion, D projects to $\{0\}$ in $\mathcal{G}_{\widehat{\Gamma}}$. By Theorem 7.11, this means that $D \subset (\zeta^{\dagger} \otimes \eta)(G_{\widehat{\Gamma}})$. Taking the dual and applying Lemma 8.6 gives that $D^{\dagger} \subset (\eta \otimes \zeta^{\dagger})(G_{\widehat{\Gamma}}^{\dagger}) \subset (\eta \otimes \zeta^{\dagger})(T_{\widehat{\Gamma}}^{\dagger})$. Now fix $d \in D$ and $\phi \in \text{TLG}(\widehat{\Gamma})$. By Definition 8.2 of the loop-linking number, for every $x \in M(\widehat{\Gamma})$ we have:

$$\left(\mathcal{L}(\mathcal{A}) - \mathcal{L}(\mathcal{A}')\right)(x) = \pi\left(x^{\ddagger}, d \circ \phi(x)\right) = \pi\left(d^{\dagger}(x^{\ddagger}), \phi(x)\right)$$

There exists $\psi \in T_{\widehat{\Gamma}}$ such that $d^{\dagger} = (\eta \otimes \zeta^{\dagger})(\psi^{\ddagger})$. Therefore:

$$\mathcal{L}(\mathcal{A}) - \mathcal{L}(\mathcal{A}') = \pi \left((\eta \otimes \zeta^{\dagger})(\psi^{\ddagger}), \phi \right) = \pi \left(\psi^{\ddagger}, (\eta^{\dagger} \otimes \zeta)(\phi) \right)$$

which is equal to 0 by Definition 8.1.

9. Computations

The practical method for computing the homology inclusion of a line arrangement has been developed in collaboration with E. Artal and B. Guerville-Ballé and will be detailed in another publication. We give here a quick overview of this method before presenting its main results.

The computation is done in two main phases. The first is to compute an explicit presentation of the graph stabiliser \mathcal{G}_{Γ} , which is done in Section 7. The second is to compute the values of the morphism

$$i_{\mathcal{A}}^* \circ \gamma_* : H_1(\Gamma) \longrightarrow M(\Gamma)$$

By Definition 6.10 of the invariant, the values can be computed for any graphed embedding $\gamma \in E_{\Gamma}(\Omega)$, since the image of the morphism inside \mathcal{G}_{Γ} does not depend on this choice. In practice this requires to explicitly construct the following embeddings all at the same time:

$$\Gamma \xrightarrow{\gamma} B_{\mathcal{A}} \xrightarrow{i_{\mathcal{A}}} E_{\mathcal{A}} \longrightarrow \mathbb{CP}^2$$

The wiring diagram of W. Arvola [Arv92] and the braid monodromy, which is related to the Zariski-van Kampen method, are two slightly different but equivalent tools that allow to characterise the topology of a complex line arrangement in an abstract form. Both tools do in fact describe the embedding i_A up to isotopy in a way that allows to identify the graph structure of B_A . We devised methods to construct an explicit ordered graphed embedding using either one of these tools. These methods allow to keep control over the actual ordering of the embedding. This point is paramount since two homology inclusion values can be compared only if their respective graph ordering are known. We also devised an algorithm describing the computation of the value

of $i_{\mathcal{A}}^* \circ \gamma_*$ from the raw data of the wiring diagram or braid monodromy. The full computation process is implemented using the Sage [Sag23] language.

Since the graph stabiliser is an abelian group of finite type, it admits a Smith normal form which can easily be determined with a computer. The values of the homology inclusion are given inside this basis.

We now give several examples of Zariski pairs which are identified by the homology inclusion. **Example 9.1** (MacLane arrangements). The MacLane arrangements and their common combinatorics are presented at the end of Section 2.1. It is known (see [Bjö+99]) that this constitutes the smallest possible combinatorics that does not admit a realisation in \mathbb{RP}^2 . The automorphism group of the combinatorics is isomorphic to $\mathrm{GL}_2(\mathbb{F}_3)$. All automorphisms can be effectively realised as projective automorphisms of \mathbb{CP}^2 , and those lying in the subgroup $\mathrm{SL}_2(\mathbb{F}_3)$ preserve the order. We now give the results of the computations of the graph stabiliser and the homology inclusion of \mathcal{M}^+ and \mathcal{M}^- .

The Smith normal form of the graph stabiliser is

$$\mathcal{G}_{\Gamma} \simeq \mathbb{Z}_{3Z} \times \mathbb{Z}^{35}$$

The two values of the homology inclusion in the corresponding basis are given in Figure 11. The difference is a non-zero element of the torsion part of \mathcal{G}_{Γ} . The line arrangements $(\mathcal{M}^+, \mathcal{M}^-)$ therefore form an ordered oriented Zariski pair. \bigcirc

FIGURE 11. Homology inclusion values of the MacLane arrangements

Example 9.2 (Rybnikov quadruplet). This is the first Zariski pair of line arrangements identified by G. Rybnikov in [Ryb11]. It consists of four conjugated line arrangements with 13 lines R_1, R_2 and their complex conjugates $\overline{R_1}, \overline{R_2}$. Their common combinatorics has a non-trivial automorphism group. The Smith normal form of the graph stabiliser is:

$$\mathcal{G}_{\Gamma} \simeq \left(\mathbb{Z}_{3\mathbb{Z}}\right)^2 \times \mathbb{Z}^{220}$$

The homology inclusion values differ not only on the torsion part but also on the free abelian part:

$$R_1:(\bar{1},\bar{1},f_1) \qquad \overline{R_1}:(\bar{1},\bar{0},f_1) \qquad R_2:(\bar{0},\bar{0},f_2) \qquad \overline{R_2}:(\bar{0},\bar{1},f_2)$$

with $f_1, f_2 \in \mathbb{Z}^{220}$. This means that the four arrangements $(R_1, R_2, \overline{R_1}, \overline{R_2})$ form an ordered oriented Zariski quadruplet. Moreover, the two pairs (R_1, R_2) and $(R_1, \overline{R_2})$ and their respective conjugates all form unoriented ordered Zariski pairs.

Both the MacLane ordered pair and the Rybnikov quadruplet are known examples that were already detected by the loop-linking number [Gue22]. However, the next example is a new Zariski pair that in particular is *not* distinguished by the loop-linking number.

Example 9.3 (New Zariski quadruplet). Consider the polynomial

$$P = X^4 + 2X^3 + 4X^2 + 3X + 1$$

and the equations given by

where $\omega = -\frac{1}{2} \pm \frac{1}{2}i\sqrt{5 \pm 2\sqrt{5}}$ takes the values of the four roots of *P*. This defines four conjugated arrangements with 11 lines $B_1, B_2, \overline{B_1}, \overline{B_2}$ whose common ordered combinatorics is given by

$$\begin{split} & [[0,1,2],[0,3],[0,4,5],[0,6,7],[0,8,9,10],[1,3,6],[1,4,7],[1,5,8],\\ & [1,9],[1,10],[2,3,5],[2,4],[2,6,10],[2,7],[2,8],[2,9],[3,4,9],[3,7,10],\\ & [3,8],[4,6,8],[4,10],[5,6],[5,7,9],[5,10],[6,9],[7,8]] \end{split}$$

The Smith normal form of the graph stabiliser is:

$$\mathcal{G}_{\Gamma} \simeq \mathbb{Z}_{5\mathbb{Z}} \times \mathbb{Z}^{119}$$

The values of the homology inclusion of the four arrangements are identical on the free part but differ on the torsion part:

$$B_1:\overline{1} \qquad \qquad B_2:\overline{4} \qquad \qquad \overline{B_1}:\overline{3} \qquad \qquad \overline{B_2}:\overline{0}$$

The automorphism group of the combinatorics is trivial, which means that the four arrangements form an unordered oriented Zariski quadruplet. The two pairs (B_1, B_2) and $(B_1, \overline{B_2})$ and their respective conjugates are unordered unoriented (i.e. proper) Zariski pairs.

References

- [Art94] Enrique Artal Bartolo. 'Sur les couples de Zariski'. French. In: J. Algebraic Geom. 3.2 (1994), pp. 223–247.
- [Art+06] Enrique Artal Bartolo, Jorge Carmona Ruber, José Ignacio Cogolludo Agustín and Miguel Ángel Marco Buzunáriz. 'Invariants of combinatorial line arrangements and Rybnikov's example'. In: Singularity Theory and Its Applications. Adv. Stud. Pure Math. 43. Math. Soc. Japan, 2006, pp. 1–34.
- [ACM19a] Enrique Artal Bartolo, José Ignacio Cogolludo Agustín and Jorge Martín Morales. 'Triangular curves and cyclotomic Zariski tuples'. In: Collect. Math. 71.3 (2019), pp. 427–441.
- [ACM19b] Enrique Artal Bartolo, José Ignacio Cogolludo Agustín and Daniel Matei. 'Characteristic varieties of graph manifolds and quasi-projectivity of fundamental groups of algebraic links'. In: Eur. J. Math. 6.3 (2019), pp. 624–645.
- [ACT08] Enrique Artal Bartolo, José Ignacio Cogolludo Agustín and Hiro-o Tokunaga. 'A survey on Zariski pairs'. In: Algebraic Geometry in East Asia Hanoi 2005 (2005). Adv. Stud. Pure Math. 50. Math. Soc. Japan, 2008, pp. 1–100.
- [AFG17] Enrique Artal Bartolo, Vincent Florens and Benoît Guerville-Ballé. 'A topological invariant of line arrangements'. In: Ann. Sc. Norm. Super. Pisa Cl. Sci. 5th ser. 17.3 (2017), pp. 949–968.
- [Arv92] William A. Arvola. 'The fundamental group of the complement of an arrangement of complex hyperplanes'. In: *Topology* 31.4 (1992), pp. 757–765.

REFERENCES

- [Bir75] Joan S. Birman. Braids, Links, and Mapping Class Groups. Ann. of Math. Stud. 82. Princeton Univ. Press, 1975.
- [Bjö+99] Anders Björner, Michel Las Vergnas, Bernd Sturmfels, Neil White and Günter M. Ziegler. Oriented matroids. 2nd edition. Encyclopedia Math. Appl. 46. Cambridge Univ. Press, 1999.
- [Cad18] William Cadegan-Schlieper. 'On the geometry and topology of hyperplane complements associated to complex and quaternionic reflection groups'. PhD thesis. Los Angeles: University of California, 2018.
- [CS08] Daniel C. Cohen and Alexander I. Suciu. 'The boundary manifold of a complex line arrangement'. In: Groups, homotopy and configuration spaces. Geometry and Topology Monographs 13. Mathematical Sciences Publishers, 2008, pp. 105–146.
- [FM11] Benson Farb and Dan Margalit. A Primer on Mapping Class Groups. Ed. by Dan Margalit. Princeton Mathematical Series 49. Princeton Univ. Press, 2011.
- [Fen+99] Roger Fenn, Michael T. Greene, Dale Rolfsen, Colin Rourke and Bert Wiest. 'Ordering the braid groups'. In: Pac. J. Math. 191.1 (1999), pp. 49–74.
- [FGM15] Vincent Florens, Benoît Guerville-Ballé and Miguel Ángel Marco Buzunáriz. 'On complex line arrangements and their boundary manifolds'. In: Math. Proc. Cambridge Philos. Soc. 159.2 (2015), pp. 189–205.
- [FM97] Anatoly T. Fomenko and Sergei V. Matveev. Algorithmic and Computer Methods for Three-Manifolds. Ed. by S.V. Matveev. Mathematics and Its Applications 425. Springer, 1997.
- [Gue16] Benoît Guerville-Ballé. 'An arithmetic Zariski 4-tuple of twelve lines'. In: *Geom. Topol.* 20.1 (2016), pp. 537–553.
- [Gue20] Benoît Guerville-Ballé. 'Topology and homotopy of lattice isomorphic arrangements'. In: *Proc. Amer. Math. Soc.* 148.5 (2020), pp. 2193–2200.
- [Gue22] Benoît Guerville-Ballé. 'The loop-linking number of line arrangements'. In: *Math. Z.* 301.2 (2022), pp. 1821–1850.
- [GV19] Benoît Guerville-Ballé and Juan Viu-Sos. 'Configurations of points and topology of real line arrangements'. In: Math. Ann. 374.1-2 (2019), pp. 1–35.
- [Hir01] Eriko Hironaka. 'Boundary manifolds of line arrangements'. In: *Math. Ann.* 319.1 (2001), pp. 17–32.
- [JS79] William H. Jaco and Peter B. Shalen. Seifert fibered spaces in 3-manifolds. Mem. Amer. Math. Soc. 220. Amer. Math. Soc., 1979.
- [JY93] Tan Jiang and Stephen S.-T. Yau. 'Topological invariance of intersection lattices of arrangements in CP²'. In: Bulletin of the American Mathematical Society 29.1 (1993), pp. 88–93.
- [Kam33] Egbert R. van Kampen. 'On the fundamental group of an algebraic curve'. In: Am. J. Math. 55.1/4 (1933), p. 255.
- [KN14] János Kollár and András Némethi. 'Holomorphic arcs on singularities'. In: Invent. Math. 200.1 (2014), pp. 97–147.
- [Mac36] Saunders MacLane. 'Some Interpretations of Abstract Linear Dependence in Terms of Projective Geometry'. In: Am. J. Math. 58.1 (1936), p. 236.
- [Mum61] David Mumford. 'The topology of normal singularities of an algebraic surface and a criterion for simplicity'. In: *Publications mathématiques de l'IHÉS* 9.1 (1961), pp. 5–22.
- [NY12] Shaheen Nazir and Masahiko Yoshinaga. 'On the connectivity of the realization spaces of line arrangements'. In: Ann. Sc. Norm. Super. Pisa Cl. Sci. 5th ser. 11.4 (2012), pp. 921–937.

REFERENCES

- [Neu81] Walter D. Neumann. 'A calculus for plumbing applied to the topology of complex surface singularities and degenerating complex curves'. In: Trans. Amer. Math. Soc. 268.2 (1981), pp. 299–344.
- [Rod23] Adrien Rodau. 'Peripheral structures and topological invariants of knotted submanifolds'. PhD thesis. Université de Pau et des Pays de l'Adour ; Universidad de Zaragoza, 2023.
- [Ryb11] Grigory Rybnikov. 'On the fundamental group of the complement of a complex hyperplane arrangement'. In: *Funct. Anal. Appl.* 45.2 (2011). Original pre-print released in 1998. arXiv: math/9805056 [math.AG].
- [Sag23] The Sage Developers. SageMath. the Sage Mathematics Software System. Version 10.1. 2023. URL: http://www.sagemath.org.
- [Shi09] Ichiro Shimada. 'Non-homeomorphic conjugate complex varieties'. In: Singularities -Niigata-Toyama 2007. Adv. Stud. Pure Math. 56. Math. Soc. Japan, 2009, pp. 285– 301.
- [Shi19] Taketo Shirane. 'Galois covers of graphs and embedded topology of plane curves'. In: Topology Appl. 257 (2019), pp. 122–143.
- [Wal67a] Friedhelm Waldhausen. 'Eine Klasse von 3-dimensionalen Mannigfaltigkeiten, I'. German. In: *Invent. Math.* 3.4 (1967), pp. 308–333.
- [Wal67b] Friedhelm Waldhausen. 'Eine Klasse von 3-dimensionalen Mannigfaltigkeiten, II'. German. In: *Invent. Math.* 4.2 (1967), pp. 87–117.
- [Wes97] Eric Robert Westlund. 'The boundary manifold of an arrangement'. PhD thesis. Madison: University of Wisconsin, 1997.
- [Ye13] Fei Ye. 'Classification of moduli spaces of arrangements of nine projective lines'. In: *Pac. J. Math.* 265.1 (2013), pp. 243–256.
- [Zar31] Oscar Zariski. 'On the irregularity of cyclic multiple planes'. In: Ann. Math. 2nd ser. 32.3 (1931), pp. 485–511.
- [Zar37] Oscar Zariski. 'The topological discriminant group of a Riemann surface of genus p'. In: Am. J. Math. 59.2 (1937), pp. 335–358.

UNIV. LILLE, CNRS, UMR 8524 - LABORATOIRE PAUL PAINLEVÉ, F-59000 LILLE, FRANCE *Email address*: adrien.rodau@univ-lille.fr