

Gravitomagnetic tidal response of relativistic stars in partially screened scalar-tensor theories

Tsutomu Kobayashi^{1,*}

¹*Department of Physics, Rikkyo University, Toshima, Tokyo 171-8501, Japan*

In scalar-tensor theories beyond Horndeski, the Vainshtein screening mechanism is only partially effective inside astrophysical bodies. We investigate the potential to detect this partial breaking of Vainshtein screening through the tidal response of fluid bodies. Specifically, we calculate the gravitomagnetic tidal Love numbers and analyze how deviations from general relativity depend on parameters governing the breaking of Vainshtein screening in the weak-gravity regime. For fixed parameter values, the relative deviations increase with higher multipoles and larger compactness. However, we demonstrate that these parameters alone are insufficient to fully characterize the tidal response of relativistic bodies in scalar-tensor theories beyond Horndeski.

I. INTRODUCTION

The response of a self-gravitating body to an applied external field depends on its internal structure and the gravitational theory, which is characterized by a set of tidal Love numbers [1]. Love numbers were first introduced in Newtonian gravity [2]. The extension of Love numbers to general relativity (GR) was initiated by [3, 4] and their notion was made more precise in [5, 6]. For non-rotating bodies, the relativistic Love numbers are classified into two types based on parity: gravitoelectric Love numbers associated with an even-parity tidal field and gravitomagnetic Love numbers associated with an odd-parity tidal field, with the latter emerging as a purely relativistic effect. The relativistic tidal Love numbers lay the foundation for probing the equation of state of nuclear matter in neutron stars with gravitational waves from compact binaries [7, 8].

The tidal response is also sensitive to the underlying gravitational theory, potentially enabling us to test GR through the tidal deformability of neutron stars. Among various possibilities, scalar-tensor theories have the simplest field content consisting of one scalar and two tensor degrees of freedom and offer us a theoretically consistent and physically well-motivated framework to explore deviations from GR. Of particular interest are theories that incorporate a mechanism to screen the effects of the scalar degree of freedom near a matter source, thereby evading existing tests of gravity within the solar system. The tidal response of neutron stars in scalar-tensor theories has been discussed in Refs. [9–14]. However, to the best of our knowledge, no analysis in that direction has been carried out so far for scalar-tensor theories equipped with the Vainshtein screening mechanism. (See Ref. [15] for a review of the Vainshtein mechanism.) Scalar-tensor theories in the Horndeski family [16–18], though not all, typically feature this mechanism due to nonlinear derivative interactions of the scalar field [19–21].¹ This screening mechanism operates so efficiently in the Horndeski

family that practically no deviations from GR are expected in the Vainshtein regime, i.e. inside the Vainshtein radius. However, in scalar-tensor theories beyond Horndeski [22, 23], Vainshtein screening is only partially effective inside a region filled with matter, although it is complete outside [24], leading to astrophysical tests of gravity within this class through the modified internal structure of astrophysical bodies [25, 26]. This motivates us to study the tidal deformability of relativistic stars in partially screened scalar-tensor theories, given that the tidal response is sensitive to both the internal structure of objects and the underlying theory of gravity. See Ref. [27] for related work exploring a somewhat similar direction in the Newtonian limit.

Scalar-tensor theories beyond Horndeski have been systematically constructed and classified under the name of degenerate higher-order scalar-tensor (DHOST) theories [28–30]. See [31, 32] for reviews. The breaking of Vainshtein screening generically occurs in DHOST theories [33–35]. The Tolman-Oppenheimer-Volkoff (TOV) system for the relativistic stellar structure in DHOST theories has been considered in Refs. [36–39]. In this paper, we study the gravitomagnetic tidal response of relativistic stars in the Vainshtein regime of DHOST theories and calculate the associated tidal Love numbers, illustrating how they depend on the parameters of the underlying theory of modified gravity governing the partial breaking of Vainshtein screening. Enriching our understanding of the tidal response of relativistic stars in modified gravity would help us develop accurate waveform models and break the degeneracy between uncertainties in the nuclear equation of state and gravity beyond GR.

The paper is organized as follows. In the next section, we provide the basic equations that determine our unperturbed configuration, following closely the previous work [39]. We also see how the Newtonian limit can differ from the standard behavior inside a fluid body. Section III details the derivation of the equation for the odd-parity perturbations in the static limit and the basic

* Email: tsutomu@rikkyo.ac.jp

¹ The very recent work on the tidal deformability of neutron stars

in the Horndeski theory [14] has considered concrete models without the Vainshtein mechanism.

procedure to compute the gravitomagnetic Love numbers in DHOST theories. In Sec. IV, we give a family of concrete DHOST models that we use in calculating the Love numbers. Some discussions on the unperturbed configurations are also provided. Our main results are presented in Sec. V, clarifying how the gravitomagnetic Love numbers depend on the modified gravity parameters. Finally, we draw our conclusions in Sec VI.

II. TOV SYSTEM IN DHOST THEORIES

A. Quadratic DHOST theories

The action of quadratic DHOST theories is given by [28]

$$S = \int d^4x \sqrt{-g} \left[f(X) \mathcal{R} + \sum_{I=1}^5 A_I(X) L_I \right] + S_m, \quad (1)$$

where \mathcal{R} is the Ricci scalar, $X := -\phi^\mu \phi_\mu / 2$,

$$\begin{aligned} L_1 &:= \phi_{\mu\nu} \phi^{\mu\nu}, & L_2 &:= (\square\phi)^2, & L_3 &:= \square\phi \phi^\mu \phi_{\mu\nu} \phi^\nu, \\ L_4 &:= \phi^\mu \phi_{\mu\rho} \phi^{\rho\nu} \phi_\nu, & L_5 &:= (\phi^\mu \phi_{\mu\nu} \phi^\nu)^2, \end{aligned} \quad (2)$$

with the notations $\phi_\mu := \nabla_\mu \phi$ and $\phi_{\mu\nu} := \nabla_\mu \nabla_\nu \phi$, and S_m is the action for a perfect fluid minimally coupled to gravity. The functions f and A_I must obey the degeneracy conditions so that the system is described by two tensor and one scalar dynamical degrees of freedom. We assume the shift symmetry of ϕ , and hence f and A_I are dependent only on X . One may add the terms of the form $F_0(X) + F_1(X) \square\phi$ to the Lagrangian without changing the number of dynamical degrees of freedom. However, in this paper, we only study the Vainshtein regime of quadratic DHOST theories where these terms can be ignored [33–35].

In light of the joint observation of GW170817 and GRB 170817A [40–42], we focus on the subset of DHOST theories in which the speed of gravitational waves is equal to that of light. This subset satisfies $A_1 = 0$ [43]. The degeneracy conditions are then given by [28]

$$A_2 = -A_1 = 0, \quad (3)$$

$$A_4 = -\frac{1}{2f} (2fA_3 - 3f_X^2 - 2Xf_X A_3 + X^2 A_3^2), \quad (4)$$

$$A_5 = -\frac{A_3}{f} (f_X + X A_3), \quad (5)$$

while f and A_3 are free.

The field equations derived from the action (1) are of the form

$$\frac{2}{\sqrt{-g}} \frac{\delta S}{\delta g^{\mu\nu}} = \mathcal{E}_{\mu\nu} - T_{\mu\nu} = 0, \quad (6)$$

$$\nabla_\mu \mathcal{J}^\mu = 0, \quad (7)$$

where $T_{\mu\nu} := -(2/\sqrt{-g})\delta S_m/\delta g^{\mu\nu}$ is the energy-momentum tensor for the perfect fluid and $\mathcal{J}^\mu := (1/\sqrt{-g})\delta S/\delta\phi_\mu$ is the shift current. The energy-momentum tensor satisfies the conservation equations (the hydrodynamical equations)

$$\nabla_\nu T_\mu^\nu = 0. \quad (8)$$

The hydrodynamical equations are identical to those in GR because the fluid is assumed to be minimally coupled with gravity.

B. Governing equations for the TOV system

Let us consider a spherically symmetric solution in DHOST theories, which will be used as an unperturbed background configuration. The metric and the scalar field are taken to be

$$\bar{g}_{\mu\nu} dx^\mu dx^\nu = -e^{\nu(r)} dt^2 + e^{\lambda(r)} dr^2 + r^2 d\Omega^2, \quad (9)$$

$$\phi = t + \psi(r), \quad (10)$$

with $d\Omega^2 = d\theta^2 + \sin^2\theta d\varphi^2$. Although the metric is assumed to be static, ϕ can depend linearly on t thanks to the shift symmetry. Notice that ϕ has the dimension of time in our convention. The energy-momentum tensor is given by

$$\bar{T}_\mu^\nu = (\rho + p) \bar{u}_\mu \bar{u}^\nu + p \delta_\mu^\nu, \quad (11)$$

with the four-velocity

$$\bar{u}^\mu = \left(e^{-\nu/2}, 0, 0, 0 \right), \quad (12)$$

where $\rho = \rho(r)$ and $p = p(r)$ are the energy density and the pressure, respectively. They are related through the equation of state: $p = p(\rho)$.

Although the field equations appear to be of higher order in DHOST theories, the degeneracy of the system allows one to reduce the number of derivatives by combining different components of the field equations. In the present case, the procedure was elaborated in Ref. [39], which is reviewed in Appendix A for completeness. One arrives in the end at the following set of equations:

$$e^\lambda = \mathcal{F}_\lambda(\nu, \nu', X, X', p), \quad (13)$$

$$X' = \mathcal{F}_1(\nu, X, \rho, p) \nu' + \frac{\mathcal{F}_2(\nu, X, \rho, p)}{r}, \quad (14)$$

$$\nu' = \mathcal{F}_3(\nu, X, \rho, \rho', p), \quad (15)$$

supplemented with the radial component of the hydrodynamical equations (8),

$$p' = -\frac{\nu'}{2}(\rho + p), \quad (16)$$

where a prime denotes differentiation with respect to r . Here, it is more convenient to use $X = [e^{-\nu} - e^{-\lambda}(\psi')^2]/2$

rather than ψ' . Although the explicit expressions for \mathcal{F}_λ , \mathcal{F}_1 , \mathcal{F}_2 , and \mathcal{F}_3 are extremely messy, it is straightforward to reproduce them by following the steps outlined in Appendix A with the help of Mathematica. See Ref. [39] for the explicit expressions, but notice that our definition of X differs by a factor of $-1/2$ from that in Ref. [39]. Given the equation of state $p = p(\rho)$, we can remove p (or ρ) from the above equations.

C. External solution, boundary conditions at the center, and matching at the surface

Setting $p = \rho = 0$ in the external region, Eqs. (13)–(15) are simplified to

$$e^\lambda = 1 + r\nu', \quad X' = 0, \quad \nu' = -\frac{1}{r} + \frac{e^{-\nu}}{2rX}. \quad (17)$$

By requiring that $\psi'(r) \rightarrow 0$ and $\nu(r) \rightarrow 0$ as $r \rightarrow \infty$, we find [39]

$$e^\nu = e^{-\lambda} = 1 - \frac{2\mu}{r}, \quad (18)$$

$$X = \frac{1}{2}. \quad (19)$$

Here, μ is an integration constant that is determined by matching the external and internal solutions at the stellar surface $r = R$. The external solution is thus given by the Schwarzschild solution, which manifests complete Vainshtein screening in the external region.

In order to integrate the equations in the internal region, let us provide the boundary conditions imposed at the center. In the vicinity of the center, we have a series expansion of the form,

$$\nu = \nu_c + \frac{\nu_2}{2}r^2 + \dots, \quad (20)$$

$$X = \frac{e^{-\nu_c}}{2} \left(1 + \frac{X_2}{2}r^2 + \dots \right), \quad (21)$$

$$\rho = \rho_c + \frac{\rho_2}{2}r^2 + \dots, \quad (22)$$

$$p = p_c + \frac{p_2}{2}r^2 + \dots, \quad (23)$$

where $\psi'(0) = 0$ is assumed. Expanding Eq. (13) around $r = 0$, we see that

$$\lambda = \frac{\lambda_2}{2}r^2 + \dots, \quad (24)$$

where λ_2 can be expressed in terms of ρ_c , p_c , ν_c , ν_2 , and X_2 . Expanding then Eqs. (14), (15), and (16) around $r = 0$, one finds the expressions for ν_2 , X_2 , and p_2 in terms of ρ_c , p_c , and ν_c . The results are summarized as

follows:

$$\nu_2 = 8\pi G_c \left[\frac{\rho_c + 3p_c}{3} - \frac{(\tilde{\alpha}_H + \tilde{\beta}_1)^2}{\tilde{\alpha}_H + 2\tilde{\beta}_1} \rho_c - \frac{(2\tilde{\alpha}_H + \tilde{\beta}_1)(\tilde{\alpha}_H + 3\tilde{\beta}_1)}{\tilde{\alpha}_H + 2\tilde{\beta}_1} p_c \right], \quad (25)$$

$$X_2 = -\frac{8\pi G_c}{\tilde{\alpha}_H + 2\tilde{\beta}_1} \left[(2\tilde{\alpha}_H + 3\tilde{\beta}_1)\rho_c + 3(\tilde{\alpha}_H + 3\tilde{\beta}_1)p_c \right], \quad (26)$$

$$\lambda_2 = -\frac{p_c}{f(e^{-\nu_c}/2)} + 2\nu_2 - 2\tilde{\alpha}_H X_2, \quad (27)$$

$$p_2 = -\frac{\nu_2}{2}(\rho_c + p_c), \quad (28)$$

with

$$\tilde{\alpha}_H := -\frac{2Xf_X}{f} \Big|_{X=e^{-\nu_c}/2}, \quad (29)$$

$$\tilde{\beta}_1 := \frac{X}{f} (f_X + XA_3) \Big|_{X=e^{-\nu_c}/2}, \quad (30)$$

$$8\pi G_c := \frac{1}{2f(e^{-\nu_c}/2)(1 - \tilde{\alpha}_H - 3\tilde{\beta}_1)}. \quad (31)$$

It is interesting to compare these expressions with Eqs. (35) and (37).

For given (ρ_c, ν_c) , one can integrate Eqs. (14)–(16) from the center to the surface of the star, $r = R$, defined by $p(R) = 0$. At $r = R$, the internal solution is matched smoothly to the external solution described by Eqs. (18) and (19). We require that $\rho(R) = \rho'(R) = 0$ and exclude the case where ρ' is discontinuous across the surface. In order for X to satisfy $X(R) = 1/2$, ν_c must be adjusted to a suitable value. The value of the integration constant μ is then determined from $e^{\nu(R)} = 1 - 2\mu/R$. One can thus obtain a family of unperturbed configurations parametrized by the single parameter ρ_c . Note that suitable ν_c does not necessarily exist, depending on ρ_c .

D. Weak-field limit

In this paper, we will solve the field equations for the unperturbed background fully numerically to obtain results valid even in the strong-field regime. However, it is instructive to see analytically the case of weak gravitational fields sourced by a nonrelativistic fluid body, as our TOV system is greatly simplified. To do so, let us assume that

$$\nu = \delta\nu \ll 1, \quad X - \frac{1}{2} = \delta X \ll 1. \quad (32)$$

We also ignore the pressure p . To first order in $\delta\nu$, δX , and ρ , Eqs. (14) and (15) reduce to

$$\delta X' = \frac{\delta\nu' + (\delta\nu + 2\delta X)/r}{2(\alpha_H + \beta_1)} + \frac{r\rho}{4M^2(\alpha_H + 2\beta_1)}, \quad (33)$$

$$\delta\nu' = -\frac{\delta\nu + 2\delta X}{r} - \frac{r(\alpha_H + \beta_1)[(1 + 3\alpha_H + 3\beta_1)\rho + (\alpha_H + \beta_1)r\rho']}{2M^2(\alpha_H + 2\beta_1)(1 - \alpha_H - 3\beta_1)}. \quad (34)$$

Here we introduced the convenient parametrization [44],

$$M^2 := 2f(1/2), \quad \alpha_H := -\left.\frac{2Xf_X}{f}\right|_{X=1/2},$$

$$\beta_1 := \left.\frac{X}{f}(f_X + XA_3)\right|_{X=1/2}, \quad (35)$$

which is conventionally used in the context of the effective field theory of dark energy and modified gravity. Combining Eqs. (33) and (34), we obtain

$$(r^2\delta\nu')' = 2G_N [\mathcal{M} + \gamma_1 r^2 \mathcal{M}''], \quad (36)$$

where we defined

$$8\pi G_N := \frac{1}{M^2(1 - \alpha_H - 3\beta_1)}, \quad (37)$$

$$\gamma_1 := -\frac{(\alpha_H + \beta_1)^2}{2(\alpha_H + 2\beta_1)}, \quad (38)$$

and

$$\mathcal{M}(r) := 4\pi \int_0^r \rho(\tilde{r})\tilde{r}^2 d\tilde{r}. \quad (39)$$

This immediately leads to

$$\delta\nu' = 2G_N \left(\frac{\mathcal{M}}{r^2} + \gamma_1 \mathcal{M}'' \right), \quad (40)$$

which reproduces the Newtonian limit of DHOST theories [24, 33–35]. The second term in the right-hand side of Eq. (40) is nonvanishing only inside a fluid body, and hence it tells us how the breaking of Vainshtein screening occurs in the Newtonian regime. The deviation from the standard Newtonian result is parametrized by α_H and β_1 .

Using Eqs. (34) and (40), we obtain

$$\delta X = -\frac{\delta\nu}{2} - \frac{G_N \mathcal{M}}{r} - \frac{(\alpha_H + \beta_1)(1 + \alpha_H + \beta_1)}{2(\alpha_H + 2\beta_1)} G_N \mathcal{M}'. \quad (41)$$

To first order in the small quantities, Eq. (13) yields

$$e^\lambda = 1 + r(\delta\nu' - 2\alpha_H \delta X'). \quad (42)$$

Substituting Eqs. (40) and (41) to this, we obtain

$$e^\lambda = 1 + 2G_N \left(\frac{\mathcal{M}}{r} + \alpha_H \mathcal{M}' + \gamma_3 r \mathcal{M}'' \right), \quad (43)$$

where

$$\gamma_3 := -\frac{\beta_1(\alpha_H + \beta_1)}{2(\alpha_H + 2\beta_1)}. \quad (44)$$

This also reproduces the previous result for the Vainshtein regime in the weak-field approximation [24, 33–35]. The deviation from standard gravity is parametrized again by α_H and β_1 .

The observational constraints on α_H and β_1 are found to be [35]

$$-0.05 < \alpha_H < 0.26, \quad -0.08 < \beta_1 < 0.02, \quad (45)$$

which come from the combination of the Hulse-Taylor pulsar [45] and stellar structure physics [46].

III. ODD-PARITY PERTURBATIONS

For the calculation of the tidal Love numbers, we consider the linear metric perturbations:

$$g_{\mu\nu} = \bar{g}_{\mu\nu} + h_{\mu\nu}. \quad (46)$$

In this paper, we only study the gravitomagnetic Love numbers, which are obtained from the odd-parity sector of perturbations [5, 6, 47]. Note that there is no odd-parity perturbation in the scalar field. The nonvanishing components of the odd-parity metric perturbations in the Regge-Wheeler gauge are [48]

$$h_{t\theta} = -\frac{1}{\sin\theta} \sum_{\ell,m} h_0^{(\ell m)}(t,r) \partial_\varphi Y_{\ell m}, \quad (47)$$

$$h_{t\varphi} = \sin\theta \sum_{\ell,m} h_0^{(\ell m)}(t,r) \partial_\theta Y_{\ell m}, \quad (48)$$

$$h_{r\theta} = -\frac{1}{\sin\theta} \sum_{\ell,m} h_1^{(\ell m)}(t,r) \partial_\varphi Y_{\ell m}, \quad (49)$$

$$h_{r\varphi} = \sin\theta \sum_{\ell,m} h_1^{(\ell m)}(t,r) \partial_\theta Y_{\ell m}, \quad (50)$$

with $Y_{\ell m}(\theta, \varphi)$ being the spherical harmonics. It is straightforward to compute the nonvanishing components of the perturbed \mathcal{E}'_μ :

$$\delta\mathcal{E}_\varphi^t = \sum_{\ell,m} \left\{ \frac{e^{-\nu/2-\lambda/2}}{r^2} \left[r^2 e^{-\nu/2-\lambda/2} f \left(h'_0 - \frac{2}{r} h_0 - \dot{h}_1 \right) \right]' - \frac{(\ell-1)(\ell+2)}{r^2} e^{-\nu} f h_0 \right\} \sin\theta \partial_\theta Y_{\ell m}, \quad (51)$$

$$\delta\mathcal{E}_\varphi^r = \sum_{\ell,m} e^{-\nu-\lambda} f \left[\ddot{h}_1 - \dot{h}'_0 + \frac{2}{r} \dot{h}_0 + \frac{(\ell-1)(\ell+2)}{r^2} e^\nu h_1 \right] \sin\theta \partial_\theta Y_{\ell m}, \quad (52)$$

$$\delta\mathcal{E}_\varphi^\theta = \sum_{\ell,m} \left\{ -\frac{e^{-\nu/2-\lambda/2}}{r^2} \left[e^{\nu/2-\lambda/2} f h_1 \right]' + \frac{e^{-\nu}}{r^2} f \dot{h}_0 \right\} (\cos\theta \partial_\theta Y_{\ell m} - \sin\theta \partial_\theta \partial_\theta Y_{\ell m}), \quad (53)$$

where a dot stands for differentiation with respect to t . Here and hereafter, we omit the labels (ℓm) for notational simplicity.

The odd-parity sector of the fluid perturbations arises from the perturbed four-velocity,

$$\delta u^\mu = \sum_{\ell,m} \frac{e^{-\nu/2} U(t,r)}{r^2(\rho+p)} \left(0, 0, \frac{\partial_\varphi Y_{\ell m}}{\sin\theta}, \frac{\partial_\theta Y_{\ell m}}{\sin\theta} \right). \quad (54)$$

As clarified in Ref. [49], there are two approaches to the calculation of the gravitomagnetic Love numbers depending on the assumptions made on the fluid perturbations. In this paper, we follow the irrotational fluid approach of [5, 47], in which the zero-frequency limit of the Regge-Wheeler equation is taken at the end instead of setting $U = 0$ from the beginning. The perturbed energy-momentum tensor is then given by

$$\delta T_\varphi^t = \sum_{\ell,m} e^{-\nu} [(\rho+p)h_0 + U] \sin\theta \partial_\theta Y_{\ell m}, \quad (55)$$

$$\delta T_\varphi^r = \delta T_\varphi^\theta = 0. \quad (56)$$

The φ -component of the hydrodynamical equations gives

$$\delta(\nabla_\mu T_\varphi^\mu) = \partial_t \delta T_\varphi^t = 0. \quad (57)$$

We assume the time dependence of the perturbations as $h_0(t,r) = h_0(r)e^{-i\omega t}$, $h_1(t,r) = h_1(r)e^{-i\omega t}$, and $U(t,r) = U(r)e^{-i\omega t}$. It follows from Eqs. (55) and (57) that

$$-i\omega [(\rho+p)h_0 + U] = 0. \quad (58)$$

To derive a single master equation for the odd-parity perturbations, it is convenient to introduce [50, 51]

$$\chi(r) := f \left(h'_0 - \frac{2}{r} h_0 + i\omega h_1 \right). \quad (59)$$

Using this variable, one can write the perturbed gravitational-field equations $\delta\mathcal{E}_\mu^\nu = \delta T_\mu^\nu$ as

$$\begin{aligned} e^{\nu/2-\lambda/2} \left(r^2 e^{-\nu/2-\lambda/2} \chi \right)' - (\ell-1)(\ell+2) f h_0 \\ = r^2 [(\rho+p)h_0 + U] = 0, \end{aligned} \quad (60)$$

$$i\omega \chi + \frac{(\ell-1)(\ell+2)}{r^2} e^\nu f h_1 = 0, \quad (61)$$

$$e^{\nu/2-\lambda/2} \left[e^{\nu/2-\lambda/2} f h_1 \right]' + i\omega f h_0 = 0. \quad (62)$$

Using Eqs. (60) and (61), we can remove h_0 and h_1 from Eq. (59) to get

$$\begin{aligned} e^{-\nu}\omega^2 \chi + e^{-\lambda} \chi'' - e^{-\lambda} \frac{F'}{F} \chi' \\ - \left\{ \frac{\ell(\ell+1)-2}{r^2} + f \left[\frac{1}{F} \left(\frac{F}{f} \right)' \right]' \right\} \chi = 0, \end{aligned} \quad (63)$$

with $F(r) := e^{\nu/2+3\lambda/2} f/r^2$. This is the master equation for the odd-parity perturbations. One can reconstruct h_0 , h_1 , and U from χ with the help of Eqs. (58), (60), and (61).

We are interested in the solutions in the zero-frequency limit, $\omega \rightarrow 0$. Rather than solving the master equation (63), we derive a differential equation for h_0 , which is more directly related to the Love number calculation. Using Eqs. (60) and (63), it can be shown that h_0 in the zero-frequency limit obeys the equation

$$h_0'' - \frac{\mathcal{P}(r)}{r} h_0' - \frac{\mathcal{Q}(r)}{r^2} h_0 = 0, \quad (64)$$

where

$$\mathcal{P}(r) := r \left[\frac{\nu'}{2} + \frac{\lambda'}{2} - \frac{f_X X'}{f} \right], \quad (65)$$

$$\mathcal{Q}(r) := e^\lambda [\ell(\ell+1) - 2(1 - e^{-\lambda})] - 2\mathcal{P}(r). \quad (66)$$

We can see that the effects of modified gravity appear in two ways. First, the modification of the unperturbed configuration appears through ν and λ . Second, the new term $f_X X'/f$ modifies directly the form of the equation for h_0 as compared to the corresponding equation in GR.

In the external region where the unperturbed solution is given by Eqs. (18) and (19), Eq. (64) reduces to the same equation as in GR, leading to the same analytic solution [47]

$$h_0 = \frac{2}{3(\ell-1)} r^{\ell+1} \left[A_\ell(r) - 4 \frac{\ell+1}{\ell} \tilde{k}_\ell R^{2\ell} \mu \frac{B_\ell(r)}{r^{2\ell+1}} \right], \quad (67)$$

where

$$A_\ell(r) := {}_2F_1(-\ell+1, -\ell-2, -2\ell, 2\mu/r), \quad (68)$$

$$B_\ell(r) := {}_2F_1(\ell-1, \ell+2, 2\ell+2, 2\mu/r), \quad (69)$$

with ${}_2F_1$ being the hypergeometric function. Note that $A_\ell(r), B_\ell(r) = 1 + \mathcal{O}(\mu/r)$ for $r \gg \mu$. The dimensionless coefficient \tilde{k}_ℓ is the (rescaled) gravitomagnetic Love number [47]. By requiring that h_0 and h'_0 are continuous across the stellar surface $r = R$, we obtain

$$\tilde{k}_\ell = \frac{\ell}{4(\ell+1)} \frac{R}{\mu} \frac{R A'_\ell - (\kappa - \ell - 1) A_\ell}{R B'_\ell - (\kappa + \ell) B_\ell} \Big|_{r=R}, \quad (70)$$

where $\kappa(r) := r h'_0 / h_0$. For the calculation of \tilde{k}_ℓ , it is therefore convenient to rewrite Eq. (64) in terms of κ as

$$r \kappa' + \kappa^2 - (1 + \mathcal{P}) \kappa - \mathcal{Q} = 0. \quad (71)$$

In the vicinity of the center, ν , λ , and X are expanded as Eqs. (20), (21), and (24), and hence $\mathcal{P} = \mathcal{O}(r^2)$ and $\mathcal{Q} = \ell(\ell+1) + \mathcal{O}(r^2)$. This shows that the boundary condition is given by $h_0 \propto r^{\ell+1}$ at $r = 0$, which is translated to

$$\kappa(0) = \ell + 1. \quad (72)$$

Equation (71) can be integrated numerically from $r = 0$ to $r = R$ with the boundary condition (72). Since the internal solution for $\kappa(r)$ differs from that in GR due to the modified interior structure and nonvanishing X' , the Love numbers \tilde{k}_ℓ will also be different from the values in GR, signaling the effects of the partial breaking of Vainshtein screening.

IV. MODEL DESCRIPTION

A. Modified gravity model

So far we have derived the general equations governing the unperturbed TOV system and the odd-parity perturbations in DHOST theories. To proceed, we consider a family of concrete DHOST models with the functions of the form

$$f = f_0 + f_1 X^q, \quad A_3 = a_3, \quad (73)$$

where f_0, f_1, a_3 , and q are constant parameters. Using Eq. (35), one can express f_0, f_1 , and a_3 in terms of M^2, α_H , and β_1 to obtain the following useful expression:

$$f = \frac{M^2}{2} \left[1 + \frac{\alpha_H}{2q} - \frac{\alpha_H}{2q} (2X)^q \right], \quad (74)$$

$$A_3 = \frac{M^2}{4} (\alpha_H + 2\beta_1). \quad (75)$$

The deviations from the standard behavior of gravity in the weak-field regime are parametrized completely by α_H and β_1 , as we have seen in Sec. IID. However, they alone will be insufficient for the characterization of the strong-field regime and the effects of q are expected to emerge when gravity gets stronger. The model with $q = 2$ was studied in Refs. [39, 52–54]. By comparing the results with the same (α_H, β_1) but with different q , one would be able to disclose the effects that are not captured by the conventionally used parameters (α_H, β_1) .

We use the DHOST theory with the functions (74) and (75) to compute the tidal Love numbers.

B. Relativistic star model

We are not attempting to determine precisely the tidal Love numbers in realistic situations. We therefore compute the Love numbers for energy polytropes, which have the simple equation of state

$$p = K \rho^{1+1/n}. \quad (76)$$

The energy polytropes with $n \leq 1$ result in $\rho'(R) \neq 0$, which hinders a smooth matching of the internal and external metrics at the surface of the star in DHOST theories. Though the form of the equation of state is different, discontinuities essentially caused by this are found in the previous study [39]. To avoid such discontinuities, we restrict our investigation to the polytropic indices $n > 1$ (more specifically, $n = 2$ and $n = 3/2$).

It is convenient to introduce the parameter

$$b := K \rho_c^{1/n}, \quad (77)$$

which is a dimensionless measure of the central density. For a given set of the model parameters (α_H, β_1, q) and the polytropic index n , Eqs. (14)–(16) can be integrated to yield a sequence of unperturbed background solutions parametrized by b . We find that there is a maximum value $b = b_{\max}$ above which one cannot find ν_c that admits an appropriate matching of the internal and external solutions. The same absence of a solution above a certain central density in DHOST theories was reported in Ref. [39] using a different equation of state. A similar result was also found in Ref. [55] in a subset of the Horndeski theory with the coupling between the Einstein tensor and the first derivative of the scalar field. We also find the following qualitative results for the unperturbed configurations. For fixed q and β_1 , positive (negative) α_H makes the maximum compactness smaller (larger). Similarly, for fixed q and α_H , positive (negative) β_1 makes the maximum compactness smaller (larger). Here, the compactness is defined as

$$C := \frac{\mu}{R}. \quad (78)$$

V. LOVE NUMBERS

The gravitomagnetic Love numbers \tilde{k}_ℓ are computed numerically for different model parameters as functions of the compactness C . Our results are displayed in Figs. 1–5.

Figure 1 shows the Love numbers \tilde{k}_2, \tilde{k}_3 , and \tilde{k}_4 for different values of α_H . The other model parameters are fixed as $\beta_1 = 0$ and $q = 1$, and the polytropic index is given by $n = 2$. We present in Fig. 2 the relative difference, $(\tilde{k}_\ell - \tilde{k}_\ell^{\text{GR}}) / \tilde{k}_\ell^{\text{GR}}$, where $\tilde{k}_\ell^{\text{GR}}$ is the Love number in GR. Typically, we have

$$\frac{\tilde{k}_\ell - \tilde{k}_\ell^{\text{GR}}}{\tilde{k}_\ell^{\text{GR}}} \sim -\mathcal{O}(10) \times \alpha_H, \quad (79)$$




FIG. 1. Gravitomagnetic Love numbers \tilde{k}_2 (left), \tilde{k}_3 (upper right), and \tilde{k}_4 (lower right) for different values of α_H versus compactness $C = \mu/R$. The other model parameters are given by $\beta_1 = 0$ and $q = 1$. We use the polytropic index $n = 2$. The dashed lines represent the results for GR.




FIG. 2. Relative differences between the $\ell = 2, 3, 4$ gravitomagnetic Love numbers in DHOST theories and GR versus compactness. The left panel shows the results for $\alpha_H = \pm 10^{-2}$, while the right panel is for $\alpha_H = \pm 10^{-3}$. The other model parameters are given by $\beta_1 = 0$ and $q = 1$. We use the polytropic index $n = 2$.

and it is larger by some factor for higher multipoles. We also find that the difference increases with larger compactness.

For the different choice of the polytropic index, $n = 3/2$, the Love numbers and their relative differences compared to GR are plotted in Fig. 3. We find qualitatively the same results as those for $n = 2$.

To see the effects that are not encapsulated solely in α_H and β_1 , we present in Fig. 4 the Love numbers for different q , with α_H and β_1 being fixed as $\alpha_H = 10^{-2}$ and $\beta_1 = 0$. For small C , the Love numbers depend almost only on α_H and β_1 . However, the q dependence emerges as C gets larger and one goes away from the weak-field regime. This shows that α_H and β_1 alone are insufficient to capture the behavior of gravity in the strong-field

regime.

Figure 5 shows how the Love numbers depend on β_1 for selected values of α_H . We find that the two parameters α_H and β_1 have similar effects on the tidal response: positive (negative) α_H and β_1 contribute to decreasing (increasing) the value of $|\tilde{k}_\ell|$. The magnitude of the change in \tilde{k}_ℓ caused by β_1 is of the same order as that by α_H .

VI. CONCLUSIONS

We have studied the gravitomagnetic tidal deformability of relativistic stars in scalar-tensor theories beyond Horndeski, i.e. degenerate higher-order scalar-tensor (DHOST) theories. Though the Vainshtein mechanism




FIG. 3. Gravitomagnetic Love numbers in DHOST theories versus compactness (left) and their relative differences compared to GR (upper right and lower right). We use the polytropic index $n = 3/2$. The dashed lines in the left panel represent the results for GR.




FIG. 4. Gravitomagnetic Love numbers in DHOST theories with $\alpha_H = 10^{-2}$ and $\beta_1 = 0$ versus compactness. We compare the results for different values of q . We use the polytropic index $n = 2$. The dashed lines represent the results for GR.

operates completely outside, screening is only partially effective in a region filled with matter in DHOST theories [24, 33–35]. We have shown that this partial breaking of Vainshtein screening in astrophysical bodies can be probed by their tidal response.

We have derived the odd-parity perturbation equations for the Love number calculation in the zero-frequency limit, adopting the irrotational fluid approach of [5, 47]. The key equation is given by Eq. (64), in conjunction with the definitions (65) and (66), where Vainshtein-breaking effects arise from the nonvanishing gradient of the scalar-field kinetic term as well as from the modified internal structure of the fluid body. We numerically calculated the gravitomagnetic Love numbers, \tilde{k}_ℓ , for different parameters of the DHOST model defined by the functions (74) and (75). Two of the model parameters, α_H and β_1 , are conventionally used in the context of effective field theory of dark energy and modified gravity to characterize deviations from GR in the weak-field regime and linear cosmology. We have found that the relative differences of the Love numbers compared to GR are roughly $\sim \mathcal{O}(10) \times \alpha_H, \beta_1$. The differences increase by some factor with higher multipoles and larger compactness. Qualitatively, positive (negative) α_H and β_1 contribute to decreasing (increasing) the value of $|\tilde{k}_\ell|$. These two parameters are however insufficient to fully characterize the deviations from GR away from the weak-field limit, which we have demonstrated by changing the third parameter of the model, q . Indeed, the effect of this parameter is more clearly seen for larger compactness.

In this paper, we have focused on the gravitomagnetic (odd-parity) Love numbers and postponed the analysis of the gravitoelectric (even-parity) ones, even though the latter have a larger contribution than the former to the phase of the gravitational-wave signal from a binary inspiral. The reason for this limitation is mostly techni-




FIG. 5. $\ell = 2$ gravitomagnetic Love number \tilde{k}_2 for different values of β_1 versus compactness. The left panel shows the results for $\alpha_H = 5 \times 10^{-3}$, while the right panel is for $\alpha_H = -5 \times 10^{-3}$. We use $q = 1$ and the polytropic index $n = 2$. The dashed lines represent the results for GR.

cal. The even-parity sector of the linear perturbations of a spherically symmetric solution in DHOST theories is even more involved than that in the Horndeski theory. The reduction of the number of derivatives in the degenerate system and the derivation of master variables have been successfully achieved only for a specific background solution in vacuum DHOST theories [56], and at this moment it seems challenging to extend the procedure of [56] to the present setup for the Love number calculation. We hope to come back to this issue in the future.

ACKNOWLEDGMENTS

The work of TK was supported by JSPS KAKENHI Grant No. JP20K03936 and MEXT-JSPS Grant-in-Aid for Transformative Research Areas (A) “Extreme Universe”, No. JP21H05182 and No. JP21H05189.

Appendix A: Derivation of the equations in Sec. II B

In this appendix, we outline the derivation of the basic equation governing the TOV system in DHOST theories, following closely the previous work [39].

The first step is to notice that $\mathcal{J}^r \propto \mathcal{E}_{tr}$, and hence the gravitational field equation $\mathcal{E}_{tr} = 0$ implies

$$\mathcal{J}^r = 0. \quad (\text{A1})$$

The scalar-field equation (7) is then automatically satisfied.

Using $X = [e^{-\nu} - e^{-\lambda}(\psi')^2]/2$ in place of ψ' , we see, from explicit calculations, that the field equations are of

the form

$$\begin{aligned} \mathcal{E}_t^t &= b_1 \nu'' + b_2 X'' + b_3 \lambda' + \tilde{\mathcal{E}}_t(\nu, \nu', \lambda, X, X') \\ &= -\rho, \end{aligned} \quad (\text{A2})$$

$$\begin{aligned} \mathcal{E}_r^r &= c_1 \nu'' + c_2 X'' + c_3 \lambda' + \tilde{\mathcal{E}}_r(\nu, \nu', \lambda, X, X') \\ &= p, \end{aligned} \quad (\text{A3})$$

$$\begin{aligned} \psi' \mathcal{J}^r &= c_1 \nu'' + c_2 X'' + c_3 \lambda' + \tilde{\mathcal{J}}(\nu, \nu', \lambda, X, X') \\ &= 0, \end{aligned} \quad (\text{A4})$$

where b_1, b_2, c_1, c_2 are expressed in terms of ν, λ , and X , while b_3 and c_3 depend also on ν' and X' . The point is that \mathcal{E}_r^r and $\psi' \mathcal{J}^r$ share the same coefficients c_1, c_2 , and c_3 . This fact allows us to solve the equation $\mathcal{E}_r^r - \psi' \mathcal{J}^r = p$ for λ to obtain

$$e^\lambda = \mathcal{F}_\lambda(\nu, \nu', X, X', p). \quad (\text{A5})$$

Equation (A5) can be used to remove λ and λ' from Eq. (A4), yielding

$$\psi' \mathcal{J}^r = \eta_1 \nu'' + \eta_2 X'' + \tilde{\mathcal{E}}_1(\nu, \nu', X, X', \rho, p) = 0, \quad (\text{A6})$$

where $\eta_{1,2} = \eta_{1,2}(\nu, \nu', X, X', p)$. Here we used the hydrodynamical equation (16) to replace p' with ρ, p , and ν' . Similar manipulations show that Eq. (A2) can also be rewritten in the form

$$\eta_1 \nu'' + \eta_2 X'' + \tilde{\mathcal{E}}_2(\nu, \nu', X, X', \rho, p) = 0, \quad (\text{A7})$$

which shares the same coefficient η_1 and η_2 with Eq. (A6). We thus obtain $\tilde{\mathcal{E}}_1 = \tilde{\mathcal{E}}_2$, which can be rearranged to give the equation of the form

$$X' = \mathcal{F}_1(\nu, X, \rho, p) \nu' + \frac{\mathcal{F}_2(\nu, X, \rho, p)}{r}. \quad (\text{A8})$$

We then substitute Eq. (A8) to Eq. (A6) to eliminate X' and X'' . It turns out that this procedure also removes ν'' in Eq. (A6) (thanks to the degeneracy of the system), leading to the equation of the form

$$\nu' = \mathcal{F}_3(\nu, X, \rho, \rho', p). \quad (\text{A9})$$

It is not particularly illuminating to replicate here the explicit expressions for \mathcal{F}_λ , \mathcal{F}_1 , \mathcal{F}_2 , and \mathcal{F}_3 , which are extremely messy.

One can thus reduce the number of derivatives of the TOV system in DHOST theories. For given central val-

ues ρ_c and ν_c , one can integrate Eqs. (A8), (A9), and the hydrodynamical equation (16), equipped with the equation of state, to determine $\nu(r)$, $X(r)$, $\rho(r)$, and $p(r)$. The remaining metric function λ can then be determined from Eq. (A5). The resultant metric and X are required to be matched smoothly to the external solution at the surface $r = R$ of the fluid body, where $p(R) = 0$. In order for this to be possible, ν_c must be adjusted to a suitable value. As a result, one finds a sequence of solutions parameterized by ρ_c .

-
- [1] E. Poisson and C.M. Will, *Gravity: Newtonian, Post-Newtonian, Relativistic*, Cambridge University Press (2014), [10.1017/CBO9781139507486](https://doi.org/10.1017/CBO9781139507486). 1
- [2] A.E.H. Love, *The yielding of the earth to disturbing forces*, *Proc. Roy. Soc. Lond. A* **82** (1909) 73. 1
- [3] E.E. Flanagan and T. Hinderer, *Constraining neutron star tidal Love numbers with gravitational wave detectors*, *Phys. Rev. D* **77** (2008) 021502 [[0709.1915](https://arxiv.org/abs/0709.1915)]. 1
- [4] T. Hinderer, *Tidal Love numbers of neutron stars*, *Astrophys. J.* **677** (2008) 1216 [[0711.2420](https://arxiv.org/abs/0711.2420)]. 1
- [5] T. Damour and A. Nagar, *Relativistic tidal properties of neutron stars*, *Phys. Rev. D* **80** (2009) 084035 [[0906.0096](https://arxiv.org/abs/0906.0096)]. 1, 4, 5, 8
- [6] T. Binnington and E. Poisson, *Relativistic theory of tidal Love numbers*, *Phys. Rev. D* **80** (2009) 084018 [[0906.1366](https://arxiv.org/abs/0906.1366)]. 1, 4
- [7] T. Hinderer, B.D. Lackey, R.N. Lang and J.S. Read, *Tidal deformability of neutron stars with realistic equations of state and their gravitational wave signatures in binary inspiral*, *Phys. Rev. D* **81** (2010) 123016 [[0911.3535](https://arxiv.org/abs/0911.3535)]. 1
- [8] LIGO SCIENTIFIC, VIRGO collaboration, *GW170817: Measurements of neutron star radii and equation of state*, *Phys. Rev. Lett.* **121** (2018) 161101 [[1805.11581](https://arxiv.org/abs/1805.11581)]. 1
- [9] P. Pani and E. Berti, *Slowly rotating neutron stars in scalar-tensor theories*, *Phys. Rev. D* **90** (2014) 024025 [[1405.4547](https://arxiv.org/abs/1405.4547)]. 1
- [10] S.S. Yazadjiev, D.D. Doneva and K.D. Kokkotas, *Tidal Love numbers of neutron stars in $f(R)$ gravity*, *Eur. Phys. J. C* **78** (2018) 818 [[1803.09534](https://arxiv.org/abs/1803.09534)].
- [11] A. Saffer and K. Yagi, *Tidal deformabilities of neutron stars in scalar-Gauss-Bonnet gravity and their applications to multimessenger tests of gravity*, *Phys. Rev. D* **104** (2021) 124052 [[2110.02997](https://arxiv.org/abs/2110.02997)].
- [12] S.M. Brown, *Tidal Deformability of Neutron Stars in Scalar-tensor Theories of Gravity*, *Astrophys. J.* **958** (2023) 125 [[2210.14025](https://arxiv.org/abs/2210.14025)].
- [13] G. Creci, T. Hinderer and J. Steinhoff, *Tidal properties of neutron stars in scalar-tensor theories of gravity*, *Phys. Rev. D* **108** (2023) 124073 [[2308.11323](https://arxiv.org/abs/2308.11323)].
- [14] R.F. Diedrichs, S. Tsujikawa and K. Yagi, *Tidal Love Numbers of Neutron Stars in Horndeski Theories*, [2501.07998](https://arxiv.org/abs/2501.07998). 1
- [15] E. Babichev and C. Deffayet, *An introduction to the Vainshtein mechanism*, *Class. Quant. Grav.* **30** (2013) 184001 [[1304.7240](https://arxiv.org/abs/1304.7240)]. 1
- [16] G.W. Horndeski, *Second-order scalar-tensor field equations in a four-dimensional space*, *Int. J. Theor. Phys.* **10** (1974) 363. 1
- [17] C. Deffayet, X. Gao, D.A. Steer and G. Zahariade, *From k -essence to generalised Galileons*, *Phys. Rev. D* **84** (2011) 064039 [[1103.3260](https://arxiv.org/abs/1103.3260)].
- [18] T. Kobayashi, M. Yamaguchi and J. Yokoyama, *Generalized G -inflation: Inflation with the most general second-order field equations*, *Prog. Theor. Phys.* **126** (2011) 511 [[1105.5723](https://arxiv.org/abs/1105.5723)]. 1
- [19] R. Kimura, T. Kobayashi and K. Yamamoto, *Vainshtein screening in a cosmological background in the most general second-order scalar-tensor theory*, *Phys. Rev. D* **85** (2012) 024023 [[1111.6749](https://arxiv.org/abs/1111.6749)]. 1
- [20] T. Narikawa, T. Kobayashi, D. Yamauchi and R. Saito, *Testing general scalar-tensor gravity and massive gravity with cluster lensing*, *Phys. Rev. D* **87** (2013) 124006 [[1302.2311](https://arxiv.org/abs/1302.2311)].
- [21] K. Koyama, G. Niz and G. Tasinato, *Effective theory for the Vainshtein mechanism from the Horndeski action*, *Phys. Rev. D* **88** (2013) 021502 [[1305.0279](https://arxiv.org/abs/1305.0279)]. 1
- [22] M. Zumalacárregui and J. García-Bellido, *Transforming gravity: from derivative couplings to matter to second-order scalar-tensor theories beyond the Horndeski Lagrangian*, *Phys. Rev. D* **89** (2014) 064046 [[1308.4685](https://arxiv.org/abs/1308.4685)]. 1
- [23] J. Gleyzes, D. Langlois, F. Piazza and F. Vernizzi, *Healthy theories beyond Horndeski*, *Phys. Rev. Lett.* **114** (2015) 211101 [[1404.6495](https://arxiv.org/abs/1404.6495)]. 1
- [24] T. Kobayashi, Y. Watanabe and D. Yamauchi, *Breaking of Vainshtein screening in scalar-tensor theories beyond Horndeski*, *Phys. Rev. D* **91** (2015) 064013 [[1411.4130](https://arxiv.org/abs/1411.4130)]. 1, 4, 8
- [25] K. Koyama and J. Sakstein, *Astrophysical Probes of the Vainshtein Mechanism: Stars and Galaxies*, *Phys. Rev. D* **91** (2015) 124066 [[1502.06872](https://arxiv.org/abs/1502.06872)]. 1
- [26] R. Saito, D. Yamauchi, S. Mizuno, J. Gleyzes and D. Langlois, *Modified gravity inside astrophysical bodies*, *JCAP* **06** (2015) 008 [[1503.01448](https://arxiv.org/abs/1503.01448)]. 1
- [27] P. Banerjee, D. Garain, S. Paul, S.t. Rajibul and T. Sarkar, *Constraining modified gravity from tidal phenomena in binary stars*, *Astrophys. J.* **910** (2021) 23 [[2006.01646](https://arxiv.org/abs/2006.01646)]. 1
- [28] D. Langlois and K. Noui, *Degenerate higher derivative theories beyond Horndeski: evading the Ostrogradski instability*, *JCAP* **02** (2016) 034 [[1510.06930](https://arxiv.org/abs/1510.06930)]. 1, 2

- [29] M. Crisostomi, K. Koyama and G. Tasinato, *Extended Scalar-Tensor Theories of Gravity*, *JCAP* **04** (2016) 044 [1602.03119].
- [30] J. Ben Achour, M. Crisostomi, K. Koyama, D. Langlois, K. Noui and G. Tasinato, *Degenerate higher order scalar-tensor theories beyond Horndeski up to cubic order*, *JHEP* **12** (2016) 100 [1608.08135]. 1
- [31] D. Langlois, *Dark energy and modified gravity in degenerate higher-order scalar-tensor (DHOST) theories: A review*, *Int. J. Mod. Phys. D* **28** (2019) 1942006 [1811.06271]. 1
- [32] T. Kobayashi, *Horndeski theory and beyond: a review*, *Rept. Prog. Phys.* **82** (2019) 086901 [1901.07183]. 1
- [33] M. Crisostomi and K. Koyama, *Vainshtein mechanism after GW170817*, *Phys. Rev. D* **97** (2018) 021301 [1711.06661]. 1, 2, 4, 8
- [34] D. Langlois, R. Saito, D. Yamauchi and K. Noui, *Scalar-tensor theories and modified gravity in the wake of GW170817*, *Phys. Rev. D* **97** (2018) 061501 [1711.07403].
- [35] A. Dima and F. Vernizzi, *Vainshtein Screening in Scalar-Tensor Theories before and after GW170817: Constraints on Theories beyond Horndeski*, *Phys. Rev. D* **97** (2018) 101302 [1712.04731]. 1, 2, 4, 8
- [36] E. Babichev, K. Koyama, D. Langlois, R. Saito and J. Sakstein, *Relativistic Stars in Beyond Horndeski Theories*, *Class. Quant. Grav.* **33** (2016) 235014 [1606.06627]. 1
- [37] J. Sakstein, E. Babichev, K. Koyama, D. Langlois and R. Saito, *Towards Strong Field Tests of Beyond Horndeski Gravity Theories*, *Phys. Rev. D* **95** (2017) 064013 [1612.04263].
- [38] J. Chagoya and G. Tasinato, *Compact objects in scalar-tensor theories after GW170817*, *JCAP* **08** (2018) 006 [1803.07476].
- [39] T. Kobayashi and T. Hiramatsu, *Relativistic stars in degenerate higher-order scalar-tensor theories after GW170817*, *Phys. Rev. D* **97** (2018) 104012 [1803.10510]. 1, 2, 3, 6, 9
- [40] LIGO SCIENTIFIC, VIRGO collaboration, *GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral*, *Phys. Rev. Lett.* **119** (2017) 161101 [1710.05832]. 2
- [41] LIGO SCIENTIFIC, VIRGO, FERMI-GBM, INTEGRAL collaboration, *Gravitational Waves and Gamma-rays from a Binary Neutron Star Merger: GW170817 and GRB 170817A*, *Astrophys. J. Lett.* **848** (2017) L13 [1710.05834].
- [42] LIGO SCIENTIFIC, VIRGO, FERMI GBM, INTEGRAL, ICECUBE, ASTROSAT CADMIUM ZINC TELLURIDE IMAGER TEAM, IPN, INSIGHT-HXMT, ANTARES, SWIFT, AGILE TEAM, 1M2H TEAM, DARK ENERGY CAMERA GW-EM, DES, DLT40, GRAWITA, FERMI-LAT, ATCA, ASKAP, LAS CUMBRES OBSERVATORY GROUP, OzGRAV, DWF (DEEPER WIDER FASTER PROGRAM), AST3, CAASTRO, VINROUGE, MASTER, J-GEM, GROWTH, JAGWAR, CALTECHNRAO, TTU-NRAO, NuSTAR, PAN-STARRS, MAXI TEAM, TZAC CONSORTIUM, KU, NORDIC OPTICAL TELESCOPE, ePESSTO, GROND, TEXAS TECH UNIVERSITY, SALT GROUP, TOROS, BOOTES, MWA, CALET, IKI-GW FOLLOW-UP, H.E.S.S., LOFAR, LWA, HAWC, PIERRE AUGER, ALMA, EURO VLBI TEAM, PI OF SKY, CHANDRA TEAM AT MCGILL UNIVERSITY, DFN, ATLAS TELESCOPES, HIGH TIME RESOLUTION UNIVERSE SURVEY, RIMAS, RATIR, SKA SOUTH AFRICA/MEERKAT collaboration, *Multi-messenger Observations of a Binary Neutron Star Merger*, *Astrophys. J. Lett.* **848** (2017) L12 [1710.05833]. 2
- [43] C. de Rham and A. Matas, *Ostrogradsky in Theories with Multiple Fields*, *JCAP* **06** (2016) 041 [1604.08638]. 2
- [44] D. Langlois, M. Mancarella, K. Noui and F. Vernizzi, *Effective Description of Higher-Order Scalar-Tensor Theories*, *JCAP* **05** (2017) 033 [1703.03797]. 4
- [45] J. Beltran Jimenez, F. Piazza and H. Velten, *Evading the Vainshtein Mechanism with Anomalous Gravitational Wave Speed: Constraints on Modified Gravity from Binary Pulsars*, *Phys. Rev. Lett.* **116** (2016) 061101 [1507.05047]. 4
- [46] I.D. Saltas, I. Sawicki and I. Lopes, *White dwarfs and revelations*, *JCAP* **05** (2018) 028 [1803.00541]. 4
- [47] P. Landry and E. Poisson, *Gravitomagnetic response of an irrotational body to an applied tidal field*, *Phys. Rev. D* **91** (2015) 104026 [1504.06606]. 4, 5, 6, 8
- [48] T. Regge and J.A. Wheeler, *Stability of a Schwarzschild singularity*, *Phys. Rev.* **108** (1957) 1063. 4
- [49] P. Pani, L. Gualtieri, T. Abdelsalhin and X. Jiménez-Forteza, *Magnetic tidal Love numbers clarified*, *Phys. Rev. D* **98** (2018) 124023 [1810.01094]. 5
- [50] K. Takahashi, H. Motohashi and M. Minamitsuji, *Linear stability analysis of hairy black holes in quadratic degenerate higher-order scalar-tensor theories: Odd-parity perturbations*, *Phys. Rev. D* **100** (2019) 024041 [1904.03554]. 5
- [51] K. Tomikawa and T. Kobayashi, *Perturbations and quasinormal modes of black holes with time-dependent scalar hair in shift-symmetric scalar-tensor theories*, *Phys. Rev. D* **103** (2021) 084041 [2101.03790]. 5
- [52] M. Crisostomi and K. Koyama, *Self-accelerating universe in scalar-tensor theories after GW170817*, *Phys. Rev. D* **97** (2018) 084004 [1712.06556]. 6
- [53] T. Hiramatsu and D. Yamauchi, *Testing gravity theories with cosmic microwave background in the degenerate higher-order scalar-tensor theory*, *Phys. Rev. D* **102** (2020) 083525 [2004.09520].
- [54] T. Hiramatsu, *CMB constraints on DHOST theories*, *JCAP* **10** (2022) 035 [2205.11559]. 6
- [55] A. Cisterna, T. Delsate and M. Rinaldi, *Neutron stars in general second order scalar-tensor theory: The case of nonminimal derivative coupling*, *Phys. Rev. D* **92** (2015) 044050 [1504.05189]. 6
- [56] K. Takahashi and H. Motohashi, *Black hole perturbations in DHOST theories: master variables, gradient instability, and strong coupling*, *JCAP* **08** (2021) 013 [2106.07128]. 9