Continuity of the Revuz correspondence under the absolute continuity condition

Ryoichiro Noda[∗]

Abstract

In this paper, we consider symmetric Hunt processes that correspond to regular Dirichlet forms and satisfy the absolute continuity condition, i.e., processes possess transition densities. For such processes, the Revuz correspondence relates positive continuous additive functionals (PCAFs) to so-called smooth measures. We show the continuity of this correspondence. Specifically, we show that if the 1-potentials of smooth measures converge (locally) uniformly as functions, then the associated PCAFs converge. This result is derived by directly estimating the distance between the PCAFs using the distance between the 1-potentials of the associated smooth measures. Furthermore, in cases where the transition density is jointly continuous, we present sufficient conditions for the convergence of 1-potentials based on the weak or vague convergence of smooth measures.

1 Introduction

Positive continuous additive functionals (PCAFs) of Markov processes, such as local times, can be very useful for analyzing these processes. In [Rev70], Revuz discovered a one-to-one correspondence between PCAFs and measures known as smooth measures, which is now called the Revuz correspondence. This correspondence forms a core of the theory of PCAFs. However, little research has been conducted on the topological properties of the Revuz correspondence. Recently, Nishimori, Tomisaki, Tsuchida and Uemura [NTTU24] demonstrated a certain compactness property of the Revuz correspondence. In particular, they showed that for a symmetric Hunt process associated with a regular Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(S,m)$, where S is a locally compact, separable, metrizable topological space and m is a Radon measure on S with full support, if the 1-potentials of smooth measures of finite energy integrals converge with respect to \mathcal{E}_1 , where $\mathcal{E}_1(f,g) := \mathcal{E}(f,g) + \int fg \, dm$, then a subsequence of the associated PCAFs converges in the local uniform topology almost surely for quasi-everywhere starting point [NTTU24, Theorem 4.1].

We focus on the continuity of the Revuz correspondence, that is, convergence of PCAFs in terms of their associated smooth measures. The study of convergence of PCAFs is interesting in itself as the topological properties of the Revuz correspondence, but it is also important in applications such as the construction of time-changed processes, such as Liouville Brownian motions, and the discussion of their convergence [AK16, GRV16, CHK17, Ooi24]. We consider symmetric Hunt processes that correspond to regular Dirichlet forms and posses transition densities. In our first result, Theorem 1.3, we show that if the 1-potentials of smooth measures of finite energy integrals converge uniformly, then the associated PCAFs converge in the sense that the expected supremum of the squared differences of PCAFs over any compact interval converges to zero. In our second result, Theorem 1.4, the first result is extend to more general smooth measures, and it is shown that if the 1-potentials of smooth measures in the strict sense (see Definition 2.5 below) converge locally uniformly and the Hunt process is conservative, then the associated

[∗]Research Institute for Mathematical Sciences, Kyoto University, Kyoto, 606-8502, JAPAN. Email:sgrndr@kurims.kyoto-u.ac.jp

PCAFs converge uniformly on compacts in probability (often referred to as the ucp topology). Our approach differs from that of [NTTU24], where smooth measures are approximated by a sequence of measures that are absolutely continuous with respect to the reference measure m. Instead of such an approximation argument, we directly estimate the distance between two PCAFs by the distance between the 1-potentials of the corresponding smooth measures (see Theorem 1.6). Using this inequality, our convergence results are proven.

To state our main result, we introduce several pieces of notation. The details on our framework are presented in Section 2.1. We fix a locally compact separable metrizable topological space S and a Radon measure m on S with full support. Let $(\mathcal{E}, \mathcal{F})$ be a regular Dirichlet form on $L^2(S,m)$. We write $X = ((X_t)_{t\in [0,\infty)}, (P_x)_{x\in S})$ for the associated Hunt process. We suppose that X admits a transition density $p:(0,\infty)\times S\times S\to [0,\infty]$ with respect to m. We then define for each $\alpha \geq 0$ the α -potential density $(r_{\alpha}(x, y))_{x,y \in S}$ by setting

$$
r_{\alpha}(x,y) := \int_0^{\infty} e^{-\alpha t} p_t(x,y) dt.
$$

For a Borel measure μ on S, a Borel subset $E \subseteq S$ and $\alpha \geq 0$, we set

$$
\mu^E(dx) \coloneqq 1_E(x)\,\mu(dy), \quad R_\alpha\mu(x) \coloneqq \int r_\alpha(x,y)\,\mu(dy).
$$

We call $R_{\alpha}\mu$ the α -potential of μ . Following [FOT11], we define S_{00} to be the collection of finite Borel measures μ on S satisfying

$$
||R_1\mu||_{\infty} \coloneqq \sup\{|R_1\mu(x)| \mid x \in S\} < \infty.
$$

For Borel measures $\mu, \mu_1, \mu_2, \ldots$ on S, we consider the following conditions. Note that, for a Borel measure ν on S and a Borel measurable function f on S, we write $||f||_{L^{\infty}(S,\nu)}$ for the L^{∞} -norm of f with respect to ν , i.e.,

$$
||f||_{L^{\infty}(S,\nu)} := \inf\{C > 0 \mid |f(x)| \le C \text{ for } \nu\text{-a.e. } x \in S\},\tag{1.1}
$$

where we set inf $\emptyset := \infty$.

Assumption 1.1. It holds that $\mu_n \in \mathcal{S}_{00}$ for all $n \geq 1$, $\mu \in \mathcal{S}_{00}$, and

$$
\lim_{n \to \infty} ||R_1 \mu_n - R_1 \mu||_{L^{\infty}(S, \mu_n + \mu)} = 0.
$$

The following is a generalization of the above condition to measures not necessarily belonging to S_{00}

Assumption 1.2. There exists an increasing sequence $(V_k)_{k>1}$ of relatively compact open subsets of S with $\bigcup_{k\geq 1} V_k = S$ such that, for every $k \geq 1$, $\mu^{V_k} \in \mathcal{S}_{00}$, $\mu_n^{V_k} \in \mathcal{S}_{00}$ for all $n \geq 1$, and

$$
\lim_{n \to \infty} \|R_1 \mu_n^{V_k} - R_1 \mu^{V_k}\|_{L^{\infty}(S, \mu_n^{V_k} + \mu^{V_k})} = 0.
$$

Under Assumption 1.1 or 1.2, it is easy to see that μ_n and μ are smooth measures in the strict sense by definition (see Definition 2.5). We write $A_n = (A_n(t))_{t\geq 0}$ (resp. $A = (A(t))_{t\geq 0}$) for the PCAF in the strict sense associated with μ_n (resp. μ) by the Revuz correspondence. (The details regarding smooth measures and PCAFs in the strict sense are presented in Section 2.2.) Our main results are the following.

Theorem 1.3. Under Assumption 1.1, for any $T > 0$,

$$
\lim_{n \to \infty} \sup_{x \in S} E_x \left[\sup_{0 \le t \le T} |A_n(t) - A(t)|^2 \right] = 0.
$$

When Assumption 1.2 is satisfied, by additionally assuming the conservativeness of X , the following weaker convergence of PCAFs is deduced.

Theorem 1.4. Suppose that Assumption 1.2 is satisfied and moreover X is conservative, i.e.,

$$
P_x(X_t \in S, \forall t \in [0, \infty)) = 1, \quad \forall x \in S.
$$

Then, for any $\varepsilon, T > 0$ and $x \in S$.

$$
\lim_{n \to \infty} P_x \left(\sup_{0 \le t \le T} |A_n(t) - A(t)| > \varepsilon \right) = 0.
$$
\n(1.2)

As a consequence, $A_n \stackrel{P}{\rightarrow} A$ in the local uniform topology under P_x for any $x \in S$.

Remark 1.5.

- (i) Theorem 1.3 implies that the result of [NTTU24, Theorem 4.1] holds for every starting point. (Recall that their result holds for quasi-every starting point.) Specifically, under Assumption 1.1, there exists a subsequence $(n_k)_{k\geq 1}$ such that, for all $x \in S$, $A_{n_k} \to A$ in the local uniform topology almost surely with respect to P_x . This is proven similarly to the fact that convergence in probability implies existence of an subsequence that converges almost surely.
- (ii) In our approach, it is not possible to replace P_x of (1.2) with $\sup_{x \in S} P_x$. This is because the proof depends on that the conservativeness of X ensures that $P_x(\tau_{V_k} \leq T) \to 0$ as $k \to \infty$ for each $T \in (0,\infty)$ and $x \in S$, where τ_{V_k} denotes the exit time of V_k by X. Obviously, this convergence does not hold uniformly with respect to $x \in S$.
- (iii) In many examples of interest, the transition density p is jointly continuous. In Section 4, we provide sufficient conditions for Assumptions 1.1 and 1.2 that are tractable in applications.

The proofs of Theorems 1.3 and 1.4 are based on the following estimate for the distance between PCAFs in terms of the distance between the 1-potentials of the associated smooth measures. This result itself is new and can be useful for deducing rate of convergence of PCAFs.

Theorem 1.6. Fix $\mu, \nu \in S_{00}$. Let A and B be the associated PCAFs in the strict sense, respectively. It then holds that, for any $\alpha, T > 0$,

$$
\sup_{x \in S} E_x \left[\sup_{0 \le t \le T} |A_t - B_t|^2 \right] \le 18(\|R_\alpha \mu\|_\infty + \|R_\alpha \nu\|_\infty) \|R_\alpha \mu - R_\alpha \nu\|_\infty
$$

$$
+ 4e^{2T} (1 - e^{-\alpha T}) (\|R_1 \mu\|_\infty^2 + \|R_1 \nu\|_\infty^2).
$$

The remainder of the article is organized as follows. In Section 2, we clarify the framework for our main results and introduce PCAFs and smooth measures in the strict sense. We also study α -potentials of smooth measures. In Section 3, we prove the main results. In Section 4, we provide sufficient conditions Assumptions 1.1 and 1.2.

2 Preliminaries

This section is divided into three subsections. In Section 2.1, we set out the framework for the arguments of this article. In Section 2.2, we introduce PCAFs and smooth measures in the strict sense. Then, in Section 2.3, we study properties of α -potentials of smooth measures. Throughout this paper, we fix a locally compact separable metrizable topological space S and a Radon measure m on S with full support. For a subset E of S, we write \overline{E} for the closure of E

in S. We define $S_\Delta = S \cup \{\Delta\}$ to be the one-point compactification of S. (NB. If S is compact, then we add Δ to S as an isolated point.) Any function f defined on S is regarded as a function on S_Δ by setting $f(\Delta) := 0$. We define $||f||_{\infty} := \sup\{|f(x)| \mid x \in S\}$ and, for a Borel measure v on S, $||f||_{L^{\infty}(S,\nu)}$ to be the L^{∞} -norm of f with respect to v (recall this from (1.1)). We use the convention inf $\emptyset := \infty$. Given a topological space T, we denote the Borel σ -algebra on T by $\mathcal{B}(T)$. The space $[0,\infty]$ is the usual one-point compactification of $[0,\infty)$.

2.1 Dirichlet forms and Hunt processes

In this subsection, we clarify the setting for our arguments and recall some basics of the theory of symmetric Dirichlet forms and Hunt processes. For details of this theory, the reader is referred to [CF12, FOT11].

We first fix the setting that is assumed in the rest of this paper. We let $(\mathcal{E}, \mathcal{F})$ be a regular Dirichlet form on $L^2(S, m)$ (see [FOT11, Section 1.1] for the definition of regular Dirichlet forms), and let $X = (\Omega, \mathcal{M}, (X_t)_{t \in [0,\infty]}, (P_x)_{x \in S_\Delta}, (\theta_t)_{t \in [0,\infty]})$ be the Hunt process associated with $(\mathcal{E}, \mathcal{F})$ in the sense of [FOT11, Theorem 7.2.1]. Here, (Ω, M) denotes the measurable space, and θ_t denotes the shift operator, i.e., a map $\theta_t : \Omega \to \Omega$ satisfying $X_s \circ \theta_t = X_{s+t}$ for any $s \in [0, \infty]$. Note that $X_{\infty}(\omega) = \Delta$ for all $\omega \in \Omega$. We write $\mathcal{F}_{*} = (\mathcal{F}_{t})_{t \in [0,\infty]}$ for the minimum augmented admissible filtration (see [CF12, p. 397]) and $\zeta := \inf\{t \in [0,\infty] \mid X_t = \Delta\}$ for the lifetime of X. We assume that X satisfies the following absolute continuity condition.

(AC) For all $x \in S$ and $t > 0$, the measure $P_x(X_t \in dy)$ is absolutely continuous with respect to $m(dy)$.

Then, by [BCM, Proof of Theorem 3.8], there exists a Borel measurable function $p:(0,\infty)\times$ $S \times S \to [0,\infty]$ satisfying, for any $s,t > 0$ and $x, y \in S$, $P_x(X_t \in dz) = p_t(x, z) m(dz)$, $p_t(x, y) = p_t(y, x)$, and

$$
p_{t+s}(x,y) = \int_{S} p_t(x,z)p_s(z,y) \, m(dz). \tag{2.1}
$$

The function p , which is uniquely determined, is called the *transition density (or, heat kernel)* of X (with respect to m). By using (2.1), it is easy to see that, for any $x, y \in S$ and $\alpha \ge \beta > 0$,

$$
r_{\beta}(x,y) = r_{\alpha}(x,y) + (\alpha - \beta) \int_{S} r_{\beta}(x,z)r_{\alpha}(z,y) m(dz)
$$

= $r_{\alpha}(x,y) + (\alpha - \beta) \int_{S} r_{\alpha}(x,z)r_{\beta}(z,y) m(dz)$ (2.2)

(cf. [MR06, Lemma 3.3.4]). For every Borel subset E of S, we write τ_E for the first exit time of E by X , i.e.,

$$
\tau_E \coloneqq \inf \{ t \in [0, \infty) \mid X_t \notin E \}. \tag{2.3}
$$

We next recall some notions regarding the Dirichlet form and the Hunt process. A Borel measurable subset N is said to be properly exceptional if it satisfies $m(N) = 0$ and

$$
P_x(\{X_t, X_{t-} \mid t > 0\} \subseteq S \setminus N) = 1, \quad \forall x \in S \setminus N.
$$

For a subset A of S , we set

Cap₁(A) := inf{
$$
\mathcal{E}_1(f, f) | f \in \mathcal{F}, f \ge 1
$$
 m-a.e. on a neighborhood of A},

where we set $\mathcal{E}_1(f,g) \coloneqq \mathcal{E}(f,g) + \int fg \, dm$. We call $\text{Cap}_1(A)$ the *capacity* of A, and if $\text{Cap}_1(A) =$ 0, then we say that A is $\mathcal{E}\text{-}polar$ (or, simply, polar). For a statement $\mathscr{S}(x)$ regarding $x \in S$, we say that $\mathscr S$ holds *quasi-everywhere* (*q.e.* for short), if $\mathscr S(x)$ holds for any $x \in S \setminus N$ for some polar subset N. A function $u : S \to [-\infty, \infty]$ is said to be quasi-continuous if, for any $\varepsilon > 0$,

there exists an open subset G of S such that $\text{Cap}_1(G) < \varepsilon$ and $u|_{S \setminus G}$ is continuous with values in $(-\infty,\infty)$.

We introduce a transformation of X that is used in the proof of Proposition 2.14 below. Fix a non-empty open subset D of S. The part process $X^D = (\Omega, \mathcal{M}, (X_t^D)_{t \in [0,\infty]}, (P_x)_{x \in D_\Delta},$ $(\theta_t^D)_{t\in[0,\infty]}$ of X onto D is defined by setting, for each $\omega \in \Omega$ and $t \in [0,\infty]$,

$$
X_t^D(\omega) \coloneqq \begin{cases} X_t(\omega), & \text{if } t < \tau_D(\omega), \\ \Delta_D, & \text{if } t \ge \tau_D(\omega), \end{cases} \quad \theta_t^D(\omega) \coloneqq \begin{cases} \theta_t(\omega), & \text{if } t < \tau_D(\omega), \\ \theta_\infty(\omega), & \text{if } t \ge \tau_D(\omega), \end{cases}
$$

and $P_{\Delta_D} := P_{\Delta}$, where $D_{\Delta} = D \cup {\Delta_D}$ denotes the one-point compactification of D. By [FOT11, Lemma 4.1.3 and Theorem A.2.10], X^D is an m^D -symmetric Hunt process. Moreover, by [FOT11, Theorems 4.4.2 and 4.4.3], its associated Dirichlet form $(\mathcal{E}^D, \mathcal{F}^D)$ on $L^2(D, m^D)$ is regular, and is called the part Dirichlet form of $(\mathcal{E}, \mathcal{F})$ onto D. Since X satisfies (AC), so does X^D . We write $p^D : (0, \infty) \times D \times D \to [0, \infty]$ for the transition density of X^D with respect to m^D .

Lemma 2.1. For any $(t, x, y) \in (0, \infty) \times D \times D$, it holds that $p_t^D(x, y) \leq p_t(x, y)$.

Proof. Fix $x \in D$. For any $t > 0$ and non-negative Borel measurable function f on D, we have that

$$
\int_D p_t^D(x, y) f(y) m(dy) = E_x[f(X_t^D)] \le E_x[f(X_t) 1_D(X_t)] = \int_D p_t(x, y) f(y) m(dy).
$$

This implies that $p_t^D(x, \cdot) \leq p_t(x, \cdot) \ m^D$ -a.e. for every $t > 0$. Using (2.1), we deduce that, for any $t > 0$ and $y \in D$,

$$
p_t^D(x,y) = \int_D p_{t/2}^D(x,z) \, p_{t/2}^D(z,y) \, m(dz) \le \int_D p_{t/2}(x,z) \, p_{t/2}(z,y) \, m(dz) \le p_t(x,y),
$$

which completes the proof.

2.2 PCAFs and smooth measures in the strict sense

In this subsection, we introduce the main objects: PCAFs and smooth measures in the strict sense. The details can be found in [CF12, Chapter 4] and [FOT11, Section 5]. We proceed with the same setting as in Section 2.2.

Definition 2.2 (PCAF, [FOT11, p. 222]). Let $A = (A_t)_{t>0}$ be an \mathcal{F}_{*} -adapted non-negative stochastic process. It is called a *positive continuous additive functional (PCAF)* of X if there exist a set $\Lambda \in \mathcal{F}_{\infty}$, called a *defining set* of A, and an properly exceptional set $N \subseteq S$, called an exceptional set of A, satisfying the following.

- (i) It holds that $P_x(\Lambda) = 1$ for all $x \in S \setminus N$.
- (ii) For every $\omega \in \Lambda$, $A_0(\omega) = 0$, the function $t \mapsto A_t(\omega)$ from $[0,\infty)$ to $[0,\infty]$ is continuous, $A_t(\omega) < \infty$ for all $t < \zeta(\omega)$, $A_t(\omega) = A_{\zeta(\omega)}(\omega)$ for all $t \ge \zeta(\omega)$, and $A_{t+s}(\omega) = A_t(\omega) +$ $A_s(\theta_t \omega)$ for all $s, t \geq 0$.

We set $A_{\infty} := \lim_{t \to \infty} A_t$ on Λ and $A_{\infty} := 0$ otherwise.

Definition 2.3 (PCAF in the strict sense, [FOT11, p. 235]). A PCAF is a PCAF in the strict sense if it admits a defining set Λ with $P_x(\Lambda) = 1$ for all $x \in S$. In other words, it is a PCAF with exceptional set \emptyset . We say that two PCAFs $A = (A_t)_{t>0}$ and $B = (B_t)_{t>0}$ in the strict sense are equivalent if there exists a common defining set $\Lambda \in \mathcal{F}_{\infty}$ such that $P_x(\Lambda) = 1$ for all $x \in S$ and $A_t(\omega) = B_t(\omega)$ for all $t \geq 0$ and $\omega \in \Lambda$.

Definition 2.4 (Smooth measure, [FOT11, p. 83]). A Borel measure μ on S is said to be a smooth measure if it satisfies the following conditions:

- (i) μ charges no polar sets, i.e., $\mu(E) = 0$ for any $E \in \mathcal{B}(S)$ with $Cap_1(E) = 0$;
- (ii) there exists an increasing sequence $(F_n)_{n>1}$ of closed subsets such that $\mu(F_n) < \infty$ for each n and, for each compact subset K of S ,

$$
\lim_{n\to\infty} \operatorname{Cap}_1(K\setminus F_n)=0.
$$

Obviously, if μ is a Radon measure, then Definition 2.4(ii) is satisfied. (Take an increasing sequence of relatively compact open subsets $(D_n)_{n\geq 1}$ with $\bigcup_{n\geq 1} D_n = S$ and set $F_n \coloneqq \overline{D}_n$.

As in Section 1, for a Borel measure μ on S, we define the α -potential $R_{\alpha}\mu : S \to [0, \infty]$ of μ by setting

$$
R_{\alpha}\mu(x) := \int_{S} r_{\alpha}(x, y) \,\mu(dy). \tag{2.4}
$$

Definition 2.5 (Smooth measure in the strict sense, [FOT11, p. 238]). Define \mathcal{S}_{00} (resp. $\mathcal{S}_{00}^{(0)}$) to be the collection of finite Borel measures μ on S satisfying $||R_1\mu||_{\infty} < \infty$ (resp. $||R_0\mu||_{\infty} < \infty$). A Borel measure μ on S is called a *smooth measure in the strict sense* if there exists an increasing sequence $(S_n)_{n\geq 1}$ of Borel subsets such that $\mu|_{S_n} \in S_{00}$ for every n and $P_x(\lim_{n\to\infty} \tau_{S_n} \geq \zeta) = 1$ for each $x \in S$. We write S_1 for the collection of smooth measures in the strict sense.

It is known that there is a one-to-one correspondence between PCAFs (in the strict sense) and smooth measures (in the strict sense), which is often referred to as the Revuz correspondence due to Revuz [Rev70]. The following is the Revuz correspondence between PCAFs in the strict sense and smooth measures in the strict sense. For the correspondence between general PCAFs and smooth measures, see [FOT11, Theorem 5.1.4]

Theorem 2.6 ([FOT11, Theorem 5.1.7] and [KN, Theorem 2.5]). There is a one-to-one correspondence between equivalence classes of $PCAFs$ (A_t)_t >0 in the strict sense and smooth measures μ in the strict sense characterized by the following relation.

(RC) For any $\alpha > 0$ and non-negative Borel measurable function f on S,

$$
E_x \left[\int_0^\infty e^{-\alpha t} f(X_s) \, dA_s \right] = \int_S r_\alpha(x, y) f(y) \, \mu(dy), \quad \forall x \in S. \tag{2.5}
$$

For a PCAF A in the strict sense, we refer to the unique Borel measure $\mu \in \mathcal{S}_1$ satisfying (RC) as the measure associated with A. Similarly, for $\mu \in \mathcal{S}_1$, we refer to a PCAF A in the strict sense satisfying (RC), which is unique up to the equivalence relation, as the PCAF associated with μ . The following result will be used in the proof of our main results.

Lemma 2.7 ([CF12, Theorem A.3.5(iii)]). Fix $\mu \in S_1$ and a Borel subset $E \subseteq S$. Let $A =$ $(A_t)_{t\geq0}$ be the associated PCAF in the strict sense. Then, $(B_t)_{t\geq0}$ defined by setting $B_t \coloneqq$ $\int_0^t \mathbb{1}_E(X_s) dA_s$ is a PCAF in the strict sense, and the associated smooth measure is μ^E .

2.3 Properties of α -potentials

In this subsection, we study properties of the α -potentials introduced at (2.4). A general theory of α -potentials can be found in [FOT11, Section 2.2]. We first refine results in [FOT11, Section 2.2] in the case where the Hunt process satisfies (AC). We then provide a maximum principle on difference of potentials in Proposition 2.13 and a useful sufficient condition for a Radon measure to be smooth in Proposition 2.14. We proceed with the same setting as in Section 2.2.

Lemma 2.8. Fix a Borel measure μ on S. For any $\alpha \geq \beta > 0$, we have that

$$
R_{\beta}\mu(x) = R_{\alpha}\mu(x) + (\alpha - \beta) \int_{S} r_{\beta}(x, z) R_{\alpha}\mu(z) m(dz)
$$

= $R_{\alpha}\mu(x) + (\alpha - \beta) \int_{S} r_{\alpha}(x, z) R_{\beta}\mu(z) m(dz)$ (2.6)

In particular, if $\mu \in \mathcal{S}_{00}$, then the above equations hold for any $\alpha, \beta > 0$. As a consequence of (2.6), if $||R_{\alpha}\mu||_{\infty} < \infty$ for some $\alpha > 0$, then $||R_{\beta}\mu||_{\infty} < \infty$ for any $\beta > 0$.

Proof. Equation (2.6) follows from (2.2). If $\mu \in \mathcal{S}_{00}$, then the terms are all finite ans so we can consider subtraction, which yields that (2.6) holds for any $\alpha, \beta > 0$. Noting that

$$
\int_{S} r_{\beta}(x, z) m(dz) = \int_{0}^{\infty} \int_{S} e^{-\beta t} p_t(x, z) m(dz) dt \le \int_{0}^{\infty} e^{-\beta t} dt = \beta^{-1},
$$

we obtain the last assertion of the result.

To study α -potentials, it is convenient to introduce another class of Borel measures as follows.

$$
\mathcal{S}_0 := \Big\{ \mu \mid \mu \text{ is a Borel measure on } S \text{ such that } \int_S R_1 \mu(x) \, \mu(dx) < \infty \Big\}.
$$

Recalling that $\mu(S) < \infty$ and $||R_1\mu||_{\infty} < \infty$ for each $\mu \in \mathcal{S}_{00}$, we have that $\mathcal{S}_{00} \subseteq \mathcal{S}_0$. The α potentials of measures in S_0 are closely related to α -excessive functions. There appear two types of excessiveness in [FOT11, Equation (2.2.3) and p. 393]. To distinguish between the two, we refer to the property of [FOT11, Equation (2.2.3)] as quasi-excessiveness. Namely, we say that a non-negative Borel measurable function f on S is α -quasi-excessive if $e^{-\alpha t}E_x[f(X_t)] \leq f(x)$ for m-a.e. $x \in S$ for all $t > 0$. On the other hand, as usual, we say that a non-negative Borel measurable function f on S is α -excessive if $e^{-\alpha t}E_x[f(X_t)] \leq f(x)$ for all $t \geq 0$ and $x \in S$ and $e^{-\alpha t}E_x[f(X_t)] \uparrow f(x)$ as $t \downarrow 0$ for all $x \in S$. Below, we collect some fundamental results about excessive functions.

Lemma 2.9. Fix $\alpha > 0$ arbitrarily. The following statements hold.

- (i) Any non-negative constant function is α -excessive.
- (ii) Any α -excessive function is α -quasi-excessive.
- (iii) If f and g are α -quasi-excessive, then so are $f \wedge g$ and $f + g$.
- (iv) For any $\mu \in \mathcal{S}_0$, $R_{\alpha}\mu$ is α -excessive and quasi-continuous.

Proof. The first three assertions are straightforward. As for the last assertion, by using (2.1) , we deduce that

$$
e^{-\alpha t} E_x[R_\alpha \mu(X_t)] = \int_S \int_t^\infty e^{-\alpha s} p(s, x, z) \, ds \mu(dz),
$$

which yields that $R_{\alpha}\mu$ is α -excessive. Moreover, by [FOT11, Exercise 4.2.2], $R_{\alpha}\mu$ is quasicontinuous. \Box

The following result is useful for extending "almost-everywhere" statements regarding excessive functions to "pointwise" statements.

Lemma 2.10. Fix $\alpha > 0$ and α -excessive functions f and g. If $f \leq g$ m-a.e., then $f(x) \leq g(x)$ for every $x \in S$.

Proof. Fix $x \in S$. We have that, for any $t > 0$,

$$
e^{-\alpha t} E_x[f(X_t)] = e^{-\alpha t} \int_S p_t(x, y) f(y) m(dy) \le e^{-\alpha t} \int_S p_t(x, y) g(y) m(dy) = e^{-\alpha t} E_x[g(X_t)].
$$

itting $t \to 0$ in the above inequality, we obtain that $f(x) < q(x)$.

Letting $t \to 0$ in the above inequality, we obtain that $f(x) \leq g(x)$.

The following results are slight refinements of [FOT11, Theorem 2.2.1 and Lemma 2.2.4] in our setting. We say that a Borel measurable function f on S is square-integrable if $f \in L^2(S, m)$.

Lemma 2.11 (cf. [FOT11, Theorem 2.2.1]). Fix $\alpha > 0$ and a non-negative Borel measurable function f on S. The function f is square-integrable and α -quasi-excessive if and only if $R_{\alpha}\mu$ f, m-a.e., for some $\mu \in \mathcal{S}_0$.

Proof. This is an immediate consequence of [FOT11, Theorem 2.2.1 and Exercise 4.2.2]. \Box

Lemma 2.12 (cf. [FOT11, Lemma 2.2.4]). Fix $\alpha > 0$ and $\mu, \nu \in \mathcal{S}_0$.

(i) If $R_{\alpha}\mu \leq R_{\alpha}\nu$, μ -a.e., then $R_{\alpha}\mu(x) \leq R_{\alpha}\nu(x)$ for all $x \in S$.

(ii) If $R_{\alpha}\mu \leq C$, μ -a.e. for some constant $C > 0$, then $R_{\alpha}\mu(x) \leq C$ for all $x \in S$.

Proof. (i). By [FOT11, Lemma 2.2.4(i) and Exercise 4.2.2], if $R_{\alpha}\mu \leq R_{\alpha}\nu$, μ -a.e., then $R_{\alpha}\mu \leq$ $R_{\alpha} \nu$, *m*-a.e. So, the result follows from Lemma 2.10.

(ii). By [FOT11, Lemma 2.2.4(ii) and Exercise 4.2.2], if $R_{\alpha}\mu \leq C$, μ -a.e., then $R_{\alpha}\mu \leq C$, m-a.e. So, the result follows from Lemma 2.10.

Now, we are ready to prove a maximum principle on differences of potentials as follows.

Proposition 2.13. For any $\mu, \nu \in \mathcal{S}_{00}$, it holds that

$$
||R_{\alpha}\mu - R_{\alpha}\nu||_{\infty} = ||R_{\alpha}\mu - R_{\alpha}\nu||_{L^{\infty}(S,\mu+\nu)}.
$$

Proof. Fix $\mu, \nu \in \mathcal{S}_{00}$ and set $\varepsilon := ||R_{\alpha}\mu - R_{\alpha}\nu||_{L^{\infty}(S, \mu+\nu)}$. We then have that

$$
R_{\alpha}\mu \leq (\varepsilon \wedge R_{\alpha}\mu) + R_{\alpha}\nu, \quad \mu\text{-a.e.}
$$

By Lemmas 2.9 and 2.11, $f := (\varepsilon \wedge R_{\alpha}\mu) + R_{\alpha}\nu$ is square-integrable and α -quasi-excessive. Thus, by Lemma 2.11 again, there exists a measure $\nu' \in \mathcal{S}_{00}$ such that $f = R_{\alpha} \nu'$, m-a.e. We deduce from Lemma 2.9(iv) that both f and $R_{\alpha} \nu'$ are quasi-continuous. Hence, by [FOT11, Lemma 2.1.4], we obtain that $f = R_{\alpha} \nu'$, q.e. In particular, by [FOT11, Theorem 2.2.3] $f = R_{\alpha} \nu'$, μ -a.e. So, it follows that $R_{\alpha}\mu \leq R_{\alpha}\nu'$ μ -a.e. From Lemma 2.12(i), it holds that $R_{\alpha}\mu(x) \leq R_{\alpha}\nu'(x)$ for all $x \in S$, which implies that $R_{\alpha} \mu \leq f$, q.e. From the definition of f, we obtain that

$$
R_{\alpha}\mu - R_{\alpha}\nu \le \varepsilon, \quad \text{q.e.}
$$

Since any polar set is m-negligible by the definition of Cap_1 , the above inequality holds for m-a.e. $x \in S$. This, combined with (AC), immediately yields that, for any $x \in S$,

$$
e^{-\alpha t} E_x[R_\alpha \mu(X_t)] - e^{-\alpha t} E_x[R_\alpha \nu(X_t)] \le e^{-\alpha t} \varepsilon.
$$

Letting $t \to 0$ in the above inequality, we deduce from the α -excessiveness of $R_{\alpha}\mu$ and $R_{\alpha}\nu$ that

$$
R_{\alpha}\mu(x) - R_{\alpha}\nu(x) \le \varepsilon, \quad \forall x \in S. \tag{2.7}
$$

By the definition of ε , we have that

$$
R_{\alpha}\nu \leq (\varepsilon \wedge R_{\alpha}\nu) + R_{\alpha}\mu, \quad \nu\text{-a.e.}
$$

So, by the same argument, we deduce that

$$
R_{\alpha}\nu(x) - R_{\alpha}\mu(x) \le \varepsilon, \quad \forall x \in S. \tag{2.8}
$$

From (2.7) and (2.8), it follows that $||R_{\alpha}\mu - R_{\alpha}\nu||_{\infty} \leq \varepsilon$. The converse inequality is obvious and so we obtain the desired result. \Box We next provide a useful sufficient condition for a Radon measure being smooth.

Proposition 2.14. Let μ be a Radon measure on S. If, for any compact subset $K \subseteq S$,

$$
\lim_{\delta \to 0} \sup_{x \in K} \int_0^\delta \int_K p_t(x, y) \, dt \, \mu(dy) < \infty,\tag{2.9}
$$

then $\mu^D \in \mathcal{S}_{00}$ for any relatively compact Borel subset D of S. In particular, $\mu \in \mathcal{S}_1$.

To prove the above result, we use the theory of PCAFs and smooth measures that are not necessarily in the strict sense. See [FOT11, Section 5.1] for details. We also use the technique of part processes introduced at the end of Section 2.1.

Proof. Fix a relatively compact open subset D of S. By Lemma 2.1 and (2.9) , we have that

$$
\lim_{t \to 0} \sup_{x \in D} \int_0^t \int_D p_t^D(x, y) \, \mu(dy) \, ds < \infty.
$$

Thus, from [Mor21, Propositions 2.6 and 2.7], we deduce that μ^D is smooth in the strict sense with respect to $(\mathcal{E}^D, \mathcal{F}^D)$. In particular, μ^D charges no \mathcal{E}^D -polar sets (see [FOT11, Theorem 2.2.3]). Thus, by [FOT11, Theorem 4.4.3(ii)], μ^D charges no \mathcal{E} -polar sets. It is then the case that μ^D is a smooth measure with respect to $(\mathcal{E}, \mathcal{F})$. We write $A^D = (A_t^D)_{t \geq 0}$ for the PCAF associated with μ^D in the sense of [FOT11, Theorem 5.1.4]. Let $N \subseteq S$ be an exceptional set of A^D . We have from [FOT11, Theorem 5.1.3(iii)] that, for any $t > 0$,

$$
E_x[A_t^D] = \int_0^t \int_D p_s(x, y) \,\mu(dy) \, ds \quad \text{for } m\text{-a.e. } x \in S.
$$

By the same argument as in [KM24, Proposition 2.32], we deduce that, for any $t \geq 0$,

$$
E_x[A_t^D] = \int_0^t \int_D p_s(x, y) \,\mu(dy) \, ds, \quad \forall x \in S \setminus N. \tag{2.10}
$$

By (2.9), we can find $t_0 > 0$ satisfying

$$
c := \sup_{x \in \overline{D}} \int_0^{t_0} \int_{\overline{D}} p_s(x, y) \,\mu(dy) \, ds < \infty.
$$

By [FOT11, Lemma 5.1.1], we have that $\underline{A_t} = \int_0^t 1_{\overline{D}}(X_s) dA_s$ for all $t \geq 0$ P_x -a.s. for \underline{q} .e. $x \in S$. We write $\sigma_{\overline{D}}$ for the first hitting time of \overline{D} by X, i.e., $\sigma_{\overline{D}} := \inf\{t \in (0,\infty) \mid X_t \in \overline{D}\}$. Then, by using the strong Markov property, we deduce that, for q.e. $x \in S$,

$$
E_x[A_{t_0}^D] = E_x \left[\int_{\sigma_D}^{t_0} 1_{\overline{D}}(X_s) dA_s^D \cdot 1_{\{\sigma_{\overline{D}} \le t_0\}} \right]
$$

\n
$$
= E_x \left[\left(\int_0^{t_0 - \sigma_{\overline{D}}} 1_{\overline{D}}(X_s) dA_s^D \right) \circ \theta_{\sigma_{\overline{D}}} \cdot 1_{\{\sigma_{\overline{D}} \le t_0\}} \right]
$$

\n
$$
\le E_x \left[E_{X_{\sigma_{\overline{D}}}}[A_{t_0}^D] \cdot 1_{\{\sigma_{\overline{D}} \le t_0\}} \right]
$$

\n
$$
= E_x \left[\int_0^{t_0} \int_{\overline{D}} p_s(X_{\sigma_{\overline{D}}}, y) \mu(dy) ds \cdot 1_{\{\sigma_{\overline{D}} \le t_0\}} \right]
$$

\n
$$
\le c,
$$

where we use (2.10) to obtain the fourth equality. This, combined with (2.10) , yields that

$$
\int_0^{t_0} \int_D p_s(x, y) \,\mu(dy) \, ds \le c \quad \text{for q.e. } x \in S.
$$

In particular, the above equation holds m-a.e. Therefore, using (2.1) , we deduce that, for any $x \in S$ and $\varepsilon \in (0, t_0)$,

$$
\int_{\varepsilon}^{t_0} \int_{D} p_s(x, y) \,\mu(dy) \, ds = \int_{0}^{t_0 - \varepsilon} \int_{D} p_{s + \varepsilon}(x, y) \,\mu(dy) \, ds
$$

$$
= \int_{0}^{t_0 - \varepsilon} \int_{D} \int_{S} p_{\varepsilon}(x, z) p_s(z, y) \, m(dz) \,\mu(dy) \, ds
$$

$$
= \int_{S} p_{\varepsilon}(x, z) \int_{0}^{t_0 - \varepsilon} \int_{D} p_s(z, y) \,\mu(dy) \, ds \, m(dz)
$$

$$
\leq c.
$$

Letting $\varepsilon \to 0$, we obtain that

$$
\sup_{x \in S} \int_0^{t_0} \int_D p_s(x, y) \,\mu(dy) \, ds \le c.
$$

From [Mor21, Proposition 2.7], it follows that $||R_1\mu^D||_{\infty} < \infty$. By the local compactness of S, any relatively compact Borel subset D' is contained in some relatively compact open subset D'' . Since $R_1\mu^{D'} \leq R_1\mu^{D''}$, we obtain that $||R_1\mu^{D'}||_{\infty} < \infty$. In particular, $\mu^{D'} \in \mathcal{S}_{00}$, which proves the first assertion. Now, let $(D_k)_{k\geq 1}$ be an increasing sequence of relatively compact open subsets of S with $\bigcup_{k\geq 1} D_k = S$. Since $\mu^{D_k} \in \mathcal{S}_{00}$ and $\tau_{V_k} \to \zeta$ almost surely under P_x for every $x \in S$, we deduce that $\mu \in S_1$. \Box

3 Proof of main results

In this section, we prove the main results. We first show Theorem 1.6. Then, using it, we establish the convergence results, Theorems 1.3 and 1.4. We proceed in the same setting as in Section 2.1.

3.1 Proof of Theorem 1.6

In this subsection, we prove Theorem 1.6. We will use the following moment formula for the product of PCAFs.

Lemma 3.1 ([KN, Corollary 2.9(iii)]). Fix $\mu, \nu \in S_1$ and write A, B for the associated PCAFs in the strict sense. Then, for any $x \in S$,

$$
E_x[A_\infty B_\infty] = \int_S r_0(x, y) R_0 \mu(y) \nu(dy) + \int_S r_0(x, y) R_0 \nu(y) \mu(dy).
$$

We first show an estimate for PCAFs associated with measures in $\mathcal{S}_{00}^{(0)}$, where we recall the class $S_{00}^{(0)}$ from Definition 2.5.

Lemma 3.2. Fix $\mu, \nu \in S_{00}^{(0)}$. Let A and B be the associated PCAFs in the strict sense, respectively. Then it holds that

$$
\sup_{x \in S} E_x \left[\sup_{0 \le t \le \infty} |A_t - B_t|^2 \right] \le 18(\|R_0\mu\|_{\infty} + \|R_0\nu\|_{\infty}) \|R_0\mu - R_0\nu\|_{\infty}.
$$

Proof. Fix $x \in S$. We note that, by Lemma 3.1, A_{∞} and B_{∞} are square-integrable with respect to P_x . Define a \mathcal{F}_* -martingale $(M_t)_{t\in[0,\infty]}$ by setting $M_t := E_x[A_\infty - B_\infty | \mathcal{F}_t]$. Doob's martingale inequality yields that

$$
E_x \left[\sup_{0 \le t \le \infty} M_t^2 \right] \le 4 E_x [M_\infty^2] = 4 E_x \left[|A_\infty - B_\infty|^2 \right].
$$

Using Lemma 3.1, we obtain that

$$
E_x [|A_{\infty} - B_{\infty}|^2] = E_x [A_{\infty}^2] + E_x [B_{\infty}^2] - 2E_x [A_{\infty} B_{\infty}]
$$

= $2 \int_S r_0(x, y) R_0 \mu(y) \mu(dy) + 2 \int_S r_0(x, y) R_0 \nu(y) \nu(dy)$
 $- 2 \int_S r_0(x, y) R_0 \mu(y) \nu(dy) - 2 \int_S r_0(x, y) R_0 \nu(y) \mu(dy)$
 $\leq 2(||R_0 \mu||_{\infty} + ||R_0 \nu||_{\infty}) ||R_0 \mu - R_0 \nu||_{\infty}.$

Since we have that $A_{\infty} - B_{\infty} = (A_{\infty} - B_{\infty}) \circ \theta_t + (A_t - B_t)$ for every $t \geq 0$, the Markov property yields that

$$
M_t = A_t - B_t + E_{X_t}[A_{\infty} - B_{\infty}] = A_t - B_t + R_0\mu(X_t) - R_0\nu(X_t), \quad \forall t \ge 0.
$$

Thus, we deduce that

$$
E_x \left[\sup_{0 \le t \le \infty} |A_t - B_t|^2 \right]
$$

\n
$$
\le 2E_x \left[\sup_{0 \le t \le \infty} M_t^2 \right] + 2E_x \left[\sup_{0 \le t \le \infty} |R_0 \mu(X_t) - R_0 \nu(X_t)|^2 \right]
$$

\n
$$
\le 8E_x \left[|A_\infty - B_\infty|^2 \right] + 2||R_0 \mu - R_0 \nu||_\infty^2
$$

\n
$$
\le 16(||R_0 \mu||_\infty + ||R_0 \nu||_\infty) ||R_0 \mu - R_0 \nu||_\infty + 2(||R_0 \mu||_\infty + ||R_0 \nu||_\infty) ||R_0 \mu - R_0 \nu||_\infty^2
$$

\n
$$
\le 18(||R_0 \mu||_\infty + ||R_0 \nu||_\infty) ||R_0 \mu - R_0 \nu||_\infty
$$

which completes the proof.

To extend the above result to more general smooth measures, we use a killing technique. Let λ be the exponential distribution with mean 1, i.e., $\lambda((v,\infty)) = e^{-v}$ for all $v \ge 0$. We set $\tilde{\Omega} \coloneqq \Omega \times [0, \infty), \, \tilde{\mathcal{M}} \coloneqq \mathcal{M} \otimes \mathcal{B}([0, \infty)),$ and $\tilde{P}_x \coloneqq P_x \otimes \lambda$. Fix $\alpha > 0$. We define $T_\alpha : \Omega \times [0, \infty) \to$ $[0, \infty)$ by setting $T_\alpha(\omega, v) = \alpha^{-1}v$, and define, for each $t \in [0, \infty]$ and $(\omega, v) \in \tilde{\Omega}$,

$$
\tilde{X}_t^{(\alpha)}(\omega, v) \coloneqq \begin{cases} X_t(\omega), & t < T_\alpha(\omega, v), \\ \Delta, & t \ge T_\alpha(\omega, v), \end{cases} \quad \tilde{\theta}_t^{\alpha}(\omega, v) \coloneqq (\theta_t(\omega), (v - \alpha t) \vee 0).
$$

Then, $\tilde{X}^{(\alpha)} = (\tilde{\Omega}, \tilde{\mathcal{M}}, \{\tilde{X}_{t}^{(\alpha)}\})$ $\{\epsilon^{(\alpha)}_t\}_{t\in[0,\infty]}, \{\tilde{P}_x\}_{x\in S_\Delta},(\tilde{\theta}_t^\alpha)_{t\in[0,\infty]})$ is a Hunt process (see [FOT11, Theorem A.2.11]). (NB. In [FOT11, Theorem A.2.11], a more generalization transformation is considered and hence the measurability of $x \mapsto \tilde{P}_x(\cdot)$ becomes weak, so called universal measurability. However, in our setting, it is easy to check that the map is still $\mathcal{B}(S)$ -measurable.) It is also easy to check that the transition density of $\tilde{X}^{(\alpha)}$ is given by $\tilde{p}_t^{(\alpha)}$ $t^{(\alpha)}(x,y) = e^{-\alpha t} p_t(x,y)$. As a consequence, the β-potential density of $\tilde{X}^{(\alpha)}$ is $r_{\alpha+\beta}$. This implies that if μ is a smooth measure of X in the strict sense, then μ is also smooth in the strict sense with respect to $\tilde{X}^{(\alpha)}$. In the following arguments, any function Y defined on Ω is naturally identified with the function \tilde{Y} on Ω given by $Y(\omega, v) = Y(\omega)$.

Lemma 3.3. Let μ be a smooth measure of X in the strict sense and A be the associated PCAF of X in the strict sense. Fix $\alpha > 0$. We define, for each $t \in [0,\infty]$ and $(\omega, v) \in \Omega$,

$$
\tilde{A}_t^{(\alpha)}(\omega, v) \coloneqq \begin{cases} A_t(\omega), & t < T_\alpha(\omega, v), \\ A_{T_\alpha(\omega, v)}(\omega), & t \ge T_\alpha(\omega, v). \end{cases}
$$

Then, $(\tilde{A}_t^{(\alpha)})$ $\mathcal{L}_{t}^{(\alpha)}$ _t ≥ 0 is a PCAF of $\tilde{X}^{(\alpha)}$ in the strict sense and its associated smooth measure with respect to $\tilde{X}^{(\alpha)}$ is μ .

Proof. It is easy to see that $(\tilde{A}_t^{(\alpha)})$ $_{t}^{(\alpha)}$ _t₂₀ is a PCAF of $\tilde{X}^{(\alpha)}$ in the strict sense. We have that, for every non-negative measurable function f on S and $\beta > 0$,

$$
\tilde{E}_x \left[\int_0^\infty e^{-\beta t} f(X_t^{(\alpha)}) d\tilde{A}_t^{(\alpha)} \right] = \tilde{E}_x \left[\int_0^{T_\alpha} e^{-\beta t} f(X_t) dA_t \right]
$$
\n
$$
= \int_0^\infty e^{-\beta t} E_x \left[\int_0^s f(X_t) dA_t \right] \alpha e^{-\alpha s} ds
$$
\n
$$
= E_x \left[\int_0^\infty e^{-\beta t} f(X_t) \int_t^\infty \alpha e^{-\alpha s} ds dA_t \right]
$$
\n
$$
= E_x \left[\int_0^\infty e^{-(\alpha + \beta)t} f(X_t) dA_t \right]
$$
\n
$$
= \int_S r_{\alpha + \beta}(x, y) f(y) \mu(dy),
$$

where we use (2.5) to obtain the last equality. Hence, we deduce the desired result.

 \Box

Using the above lemma, we can prove Theorem 1.6 as follows.

Proof of Theorem 1.6. By regarding μ and ν as smooth measures of $\tilde{X}^{(\alpha)}$ in the strict sense, we define $\tilde{A}^{(\alpha)}$ and $\tilde{B}^{(\alpha)}$ to be the associated PCAFs in the strict sense. Using Lemma 3.3, we deduce that, for any $x \in S$,

$$
E_x \left[\sup_{0 \le t \le T} |A_t - B_t|^2 \right] \le \tilde{E}_x \left[\sup_{0 \le t \le T} |\tilde{A}_t^{(\alpha)} - \tilde{B}_t^{(\alpha)}|^2 \right] + \tilde{E}_x \left[\sup_{0 \le t \le T} |A_t - B_t|^2; T_\alpha < T \right]
$$
\n
$$
\le \tilde{E}_x \left[\sup_{0 \le t \le T} |\tilde{A}_t^{(\alpha)} - \tilde{B}_t^{(\alpha)}|^2 \right] + 2(1 - e^{-\alpha T})(E_x[A_T^2] + E_x[B_T^2]).
$$

We have that

$$
E_x[A_T^2] = E_x \left[\left(\int_0^T dA_t \right)^2 \right]
$$

\n
$$
\leq e^{2T} E_x \left[\left(\int_0^\infty e^{-t} dA_t \right)^2 \right]
$$

\n
$$
= 2e^{2T} E_x \left[\int_0^\infty e^{-2t} E_{X_t} \left[\int_0^\infty e^{-s} dA_s \right] dA_t \right]
$$

\n
$$
= 2e^{2T} \int_S r_2(x, y) \int_S r_1(y, z) \mu(dz) \mu(dy)
$$

\n
$$
\leq 2e^{2T} \|R_1 \mu\|_\infty^2,
$$

where we use [CF12, Exercise 4.1.7] and (2.5) to obtain the second and third equalities, respectively. Similarly, we deduce that $E_x[B_T^2] \le 2e^{2T} ||R_1\nu||_\infty^2$. Since r_α is the 0-potential density of $\tilde{X}^{(\alpha)}$, it follows from Lemma 3.2 that

$$
\sup_{x \in S} \tilde{E}_x \left[\sup_{0 \le t \le T} |\tilde{A}_t^{(\alpha)} - \tilde{B}_t^{(\alpha)}|^2 \right] \le 18 (\|R_\alpha \mu\|_\infty + \|R_\alpha \nu\|_\infty) \|R_\alpha \mu - R_\alpha \nu\|_\infty,
$$

which completes the proof.

3.2 Theorems 1.3 and 1.4

In this subsection, we prove Theorems 1.3 and 1.4. The key observation is that, by Proposition 2.13, Assumption 1.1 implies the convergence of the α -potentials with respect to the supremum norm. This is described precisely as follows.

Lemma 3.4. Under Assumption 1.1, for any $\alpha > 0$,

$$
\lim_{n \to \infty} ||R_{\alpha}\mu_n - R_{\alpha}\mu||_{\infty} = 0, \quad \sup_{n \ge 1} ||R_{\alpha}\mu_n||_{\infty} < \infty.
$$

Proof. Fix $\alpha > 0$. Using Lemma 2.8, we deduce that

$$
|R_{\alpha}\mu_n(x) - R_{\alpha}\mu(x)|
$$

\n
$$
\leq |R_1\mu_n(x) - R_1\mu(x)| + |1 - \alpha| \int_S r_{\alpha}(x, y)|R_1\mu_n(y) - R_1\mu(y)| m(dy)
$$

\n
$$
\leq ||R_1\mu_n - R_1\mu||_{\infty} + |1 - \alpha|\alpha^{-1}||R_1\mu_n - R_1\mu||_{\infty},
$$

where we use that $\int_S r_\alpha(x, y) m(dy) \leq \alpha^{-1}$ at the last inequality. Since we have from Assumption 1.2 and Proposition 2.13 that

$$
\lim_{n \to \infty} \|R_1 \mu_n - R_1 \mu\|_{\infty} = 0,
$$
\n(3.1)

we obtain the first result. Since $||R_1\mu||_{\infty} < \infty$, we have from Lemma 2.8 that $||R_{\alpha}\mu||_{\infty} < \infty$ for any $\alpha > 0$. Similarly, $||R_{\alpha}\mu_n||_{\infty} < \infty$ for any $n \ge 1$ and $\alpha > 0$. These, combined with the first result, yield the second result. 口

By Theorem 1.6 and Lemma 3.4, we can prove Theorem 1.3 as follows.

Proof of Theorem 1.3. Suppose that Assumption 1.1 is satisfied. It follows from Theorem 1.6 that, for any $\alpha > 0$,

$$
\sup_{x \in S} E_x \left[\sup_{0 \le t \le T} |A_n(t) - A(t)|^2 \right] \le 18(\|R_\alpha \mu_n\|_\infty + \|R_\alpha \mu\|_\infty) \|R_\alpha \mu_n - R_\alpha \mu\|_\infty
$$

$$
+ 4e^{2T} (1 - e^{-\alpha T}) (\|R_1 \mu_n\|_\infty^2 + \|R_1 \mu\|_\infty^2)
$$

Letting $n \to \infty$ and then $\alpha \to 0$ in the above inequality and using Lemma 3.4, we obtain the desired result. \Box

Finally, using Theorem 1.3, we obtain Theorem 1.4 as follows.

Proof of Theorem 1.4. Suppose that Assumption 1.2 is satisfied. By Proposition 2.13, similarly to (3.1), it holds that

$$
\lim_{n \to \infty} ||R_1 \mu_n^{V_k} - R_1 \mu^{V_k}||_{\infty} = 0, \quad \forall k \ge 1.
$$

Thus, the measures $\mu_n^{V_k}$ and μ^{V_k} satisfy Assumption 1.1 for each $k \geq 1$. Define $A_n^{V_k}(t) :=$ $\int_0^t 1_{V_k}(X_s) A_n(ds)$ and $A^{V_k}(t) := \int_0^t 1_{V_k}(X_s) A(ds)$. By Lemma 2.7, $A_n^{V_k}$ and A^{V_k} are the PCAFs in the strict sense associated with $\mu_n^{V_k}$ and μ^{V_k} , respectively. Therefore, Theorem 1.3 implies that, for any $\varepsilon, T > 0$ and $k \geq 1$,

$$
\lim_{n \to \infty} \sup_{x \in S} E_x \left[\sup_{0 \le t \le T} |A_n^{V_k}(t) - A^{V_k}(t)|^2 \right] = 0.
$$

Fix $x \in S$ and $\varepsilon, T > 0$. Recall from (2.3) that τ_{V_k} denotes the first exit time of V_k by X. If $\tau_{V_k} > T$, then $A_n^{V_k}(t) = A_n(t)$ and $A^{V_k}(t) = A(t)$ for all $t \in [0, T]$. Thus, we deduce that

$$
P_x \left(\sup_{0 \le t \le T} |A_n(t) - A(t)| > \varepsilon \right) \le P_x(\tau_{V_k} \le T) + P_x \left(\sup_{0 \le t \le T} |A_n^{V_k}(t) - A^{V_k}(t)| > \varepsilon \right)
$$

$$
\le P_x(\tau_{V_k} \le T) + \varepsilon^{-2} E_x \left[\sup_{0 \le t \le T} |A_n^{V_k}(t) - A^{V_k}(t)|^2 \right],
$$

where we use the Markov inequality to obtain the last inequality. Since the Hunt process X is conservative, $\tau_{V_k} \to \infty$ as $k \to \infty$ almost surely. Therefore, we can conclude that

$$
\lim_{n \to \infty} P_x \left(\sup_{0 \le t \le T} |A_n(t) - A(t)| > \varepsilon \right) = 0.
$$

 \Box

4 Sufficient conditions for Assumptions 1.1 and 1.2

In many examples of interest, the transition density p is jointly continuous. Below, we provide sufficient conditions for Assumptions 1.1 and 1.2 that are tractable in applications.

Theorem 4.1. Assume that the following conditions are satisfied.

- (A1) The transition density p is jointly continuous with values in $[0, \infty)$.
- (A2) All the measures $\mu, \mu_1, \mu_2, \ldots$ are finite Borel measures and $\mu_n \to \mu$ weakly.
- (A3) There exists a compact subset K_0 of S such that $\text{supp}(\mu_n) \subseteq K_0$ for all $n \geq 1$ and $supp(\mu) \subseteq K_0$.
- (A_4) For any compact subset K of S,

$$
\lim_{t \to 0} \sup_{n \ge 1} \sup_{x \in K} \int_0^t \int_S p_s(x, y) \,\mu_n(dy) \, ds = 0.
$$

Then Assumption 1.1 is satisfied.

Theorem 4.2. Assume that the following conditions are satisfied.

- (B1) The transition density p is jointly continuous with values in $[0, \infty)$.
- (B2) All the measures $\mu, \mu_1, \mu_2, \ldots$ are Radon measures and $\mu_n \to \mu$ vaguely, i.e., for all compactly supported functions $f : S \to \mathbb{R}$,

$$
\lim_{n \to \infty} \int_{S} f_n(x) \,\mu_n(dx) = \int_{S} f(x) \,\mu(dx).
$$

 $(B3)$ For any compact subset K of S, it holds that

$$
\lim_{t \to 0} \sup_{n \ge 1} \sup_{x \in K} \int_0^t \int_K p_s(x, y) \,\mu_n(dy) \, ds = 0.
$$

Then Assumption 1.2 is satisfied.

.

We first prove Theorem 4.1. By Proposition 2.14, (A4) yields that $\mu_n \in \mathcal{S}_{00}$ for each $n \geq 1$. The following result states that the limiting measure μ also belongs to \mathcal{S}_{00} .

Lemma 4.3. Under the conditions of Theorem 4.1, for any compact subset K

$$
\lim_{\delta \downarrow 0} \sup_{x \in K} \int_0^\delta \int_S p_t(x, y) \, \mu(dy) \, dt = 0.
$$

In particular, $\mu \in \mathcal{S}_{00}$.

Proof. By conditions (A2) and (A4), we have that, for each $t > 0$ and $x \in S$,

$$
\lim_{n \to \infty} \int_S p_t(x, y) \,\mu_n(dy) = \int_S p_t(x, y) \,\mu(dy).
$$

Thus, using Fatou's lemma and condition $(A4)$, we deduce that, for any compact subset K of S,

$$
\lim_{\delta \downarrow 0} \sup_{x \in K} \int_0^\delta \int_S p_t(x, y) \,\mu(dy) \, dt \le \lim_{\delta \downarrow 0} \sup_{x \in K} \lim_{n \to \infty} \int_0^\delta \int_S p_t(x, y) \,\mu_n(dy) \, dt
$$
\n
$$
\le \lim_{\delta \downarrow 0} \lim_{n \to \infty} \sup_{x \in K} \int_0^\delta \int_S p_t(x, y) \mu_n(dy) \, dt
$$
\n
$$
= 0.
$$

The last assertion follows from Proposition 2.14.

We recall some basic properties of transition densities.

Lemma 4.4. The following statements hold.

- (i) For any $t > 0$ and $x, y \in S$, we have that $p_t(x, y) \leq p_t(x, x)^{1/2} p_t(y, y)^{1/2}$.
- (ii) For any $x \in S$, the function $t \mapsto p_t(x, x)$ is non-increasing.

Proof. By (2.1) and the Cauchy-Schwarz inequality, we have that

$$
p_t(x,y) = \int p_{t/2}(x,z)p_{t/2}(z,y) m(dz)
$$

\n
$$
\leq \left(\int p_{t/2}(x,z)^2 m(dz)\right)^{1/2} \left(\int p_{t/2}(y,z)^2 m(dz)\right)^{1/2}
$$

\n
$$
= p_t(x,x)^{1/2} p_t(y,y)^{1/2},
$$

which shows (i). If we write $p_t^x(\cdot) := p_t(x, \cdot)$, then (2.1) implies that $p_{t/2}^x \in L^2(S, m)$ and $P_{t/2} p_{t/2}^x = p_t^x$, where P_t denotes the transition function of X, i.e., $P_t f(x) \coloneqq E_x[f(X_t)]$. It follows from [FOT11, Lemma 1.3.3(i)] that $p_t^x \in \mathcal{F}$. Hence, applying [FOT11, Lemma 1.3.3(i)] yields that, for any $s, t > 0$,

$$
(P_{t/2} p_{s/2}^x, P_{t/2} p_{s/2}^x) \le (p_{s/2}^x, p_{s/2}^x),
$$

where (\cdot, \cdot) denotes the inner product on $L^2(S,m)$. From the above inequality and (2.1), we deduce that, for any $s, t > 0$,

$$
p_{t+s}(x,x) \le p_s(x,x),
$$

which proves (ii).

Now, we are ready to show Theorem 4.1.

 \Box

Proof of Theorem 4.1. Suppose conditions $(A1)$, $(A2)$, $(A3)$, and $(A4)$ of Theorem 4.1 are satisfied. We have from (A4) and Proposition 2.14 that $\mu_n \in \mathcal{S}_{00}$ for each $n \geq 1$ and from Lemma 4.3 that $\mu \in \mathcal{S}_{00}$. By the weak convergence of μ_n to μ and the equicontinuity of $\{p_t(x, y)\}_{x \in K_0}$ as continuous functions of $(t, y) \in [\delta, T] \times K_0$, we deduce that, for any $\delta, T > 0$,

$$
\lim_{n \to \infty} \sup_{x \in K_0} \left| \int_S \int_{\delta}^T e^{-s} p_s(x, y) \, ds \, \mu_n(dy) - \int_S \int_{\delta}^T e^{-s} p_s(x, y) \, ds \, \mu(dy) \right| = 0 \tag{4.1}
$$

(see [Whi, Lemma 1.1]). We denote by M the maximum of $p_1(x, y)$ over $(x, y) \in K_0 \times K_0$. Then, by using Lemma 4.4, we obtain that, for any $T > 1$,

$$
\sup_{x \in K_0} \int_T^{\infty} \int_S e^{-t} p_t(x, y) dt \,\mu_n(dy) \le M\mu_n(S)e^{-T}.
$$

The weak convergence of μ_n to μ implies that $\mu_n(S)$ is uniformly bounded. Thus,

$$
\lim_{T \to \infty} \sup_{n \ge 1} \sup_{x \in K_0} \int_T^{\infty} \int_S e^{-t} p_t(x, y) dt \,\mu_n(dy) = 0. \tag{4.2}
$$

Similarly, it holds that

$$
\lim_{T \to \infty} \sup_{x \in K_0} \int_T^{\infty} \int_S e^{-t} p_t(x, y) dt \,\mu(dy) = 0. \tag{4.3}
$$

The triangle inequality yields that, for any $x \in S$ and $\delta, T > 0$ with $\delta < T$,

$$
|R_1\mu_n(x) - R_1\mu(x)| \le \left| \int_S \int_0^\infty e^{-t} p_t(x, y) \, dt \, \mu_n(dy) - \int_S \int_0^\infty e^{-t} p_t(x, y) \, dt \, \mu(dy) \right|
$$

\n
$$
\le \int_0^\delta \int_S p_t(x, y) \, dt \, \mu_n(dy) + \int_0^\delta \int_S p_t(x, y) \, dt \, \mu(dy)
$$

\n
$$
+ \left| \int_S \int_\delta^T e^{-t} p_t(x, y) \, dt \, \mu_n(dy) - \int_S \int_\delta^T e^{-t} p_t(x, y) \, dt \, \mu(dy) \right|
$$

\n
$$
+ \int_T^\infty \int_S e^{-t} p_t(x, y) \, dt \, \mu_n(dy) + \int_T^\infty \int_S e^{-t} p_t(x, y) \, dt \, \mu(dy).
$$

Taking the supremum over $x \in K_0$ in the above inequality, letting $n \to \infty$, $\delta \to 0$, and $T \to \infty$ in the above inequality, and using $(A4)$, Lemma 4.3, (4.1) , (4.2) , and (4.3) , we deduce that

$$
\lim_{n \to \infty} \sup_{x \in K_0} |R_1 \mu_n(x) - R_1 \mu(x)| = 0.
$$

Therefore, Assumption 1.1 is satisfied.

Next, we prove Theorem 4.2. By using the following result, Theorem 4.2 is easily deduced from Theorem 4.1.

Lemma 4.5. Under Assumption 4.1, there exists an increasing sequence $(V_k)_{k\geq 1}$ of relatively compact open subsets of S with $\bigcup_{k\geq 1} V_k = S$ such that $\mu_n^{V_k} \to \mu^{V_k}$ weakly as $n \to \infty$, for each $k \geq 1$.

Proof. By [WJ87, Theorem 1], there exists a metric d on S inducing the topology on S such that any bounded closed subset of S is compact. Fix $\rho \in S$. Let $(r_k)_{k>1}$ be an increasing sequence with $r_k \uparrow \infty$ such that $B_d(\rho, r_k) := \{x \in S \mid d(\rho, x) < r_k\}$ is relatively compact and $\mu(\partial B_d(\rho, r_k)) = 0$. Define $V_k := B_d(\rho, r_k)$. Condition (B2) of Theorem 4.2 implies that $\mu_n^{V_k} \to \mu^{V_k}$ weakly. Hence, we complete the proof. \Box

Proof of Theorem 4.2. Suppose that conditions $(B1)$, $(B2)$, and $(B3)$ of Theorem 4.2 are satisfied. Fix $(V_k)_{k\geq 1}$ appearing in Lemma 4.5. Then, for each $k\geq 1$, the measures $\mu_n^{V_k}$ and μ^{V_k} satisfy the conditions (A1), (A2), (A3), and (A4) of Theorem 4.1 with $K_0 = V_k$. Thus, we have from Theorem 4.1 that, for each $k \geq 1$, $\mu_n^{V_k} \in \mathcal{S}_{00}$ for all $n \geq 1$, $\mu^{V_k} \in \mathcal{S}_{00}$, and

$$
\lim_{n \to \infty} \|R_1 \mu_n^{V_k} - R_1 \mu^{V_k}\|_{L^{\infty}(S, \mu_n^{V_k} + \mu^{V_k})} = 0.
$$

Thus, Assumption 1.2 is verified.

Acknowledgement

I would like to thank my supervisor Dr. David Croydon for his support and fruitful discussions and Drs. Naotaka Kajino and Takahiro Mori for their valuable comments that helped me improve the earlier version of Proposition 2.14 to its current form. This work was supported by JSPS KAKENHI Grant Number JP 24KJ1447 and the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University.

References

- [AK16] S. Andres and N. Kajino. Continuity and estimates of the Liouville heat kernel with applications to spectral dimensions. Probab. Theory Related Fields, 166(3-4):713– 752, 2016.
- [BCM] M. T. Barlow, Z.-Q. Chen, and M. Murugan. Stability of EHI and regularity of MMD spaces. Preprint. Available at arXiv:2008.05152.
- [CF12] Z.-Q. Chen and M. Fukushima. Symmetric Markov processes, time change, and boundary theory, volume 35 of London Mathematical Society Monographs Series. Princeton University Press, Princeton, NJ, 2012.
- [CHK17] D. A. Croydon, B. M. Hambly, and T. Kumagai. Time-changes of stochastic processes associated with resistance forms. Electron. J. Probab., 22(82):41, 2017.
- [FOT11] M. Fukushima, Y. Oshima, and M. Takeda. Dirichlet forms and symmetric Markov processes, volume 19 of De Gruyter Studies in Mathematics. Walter de Gruyter & Co., Berlin, extended edition, 2011.
- [GRV16] C. Garban, R. Rhodes, and V. Vargas. Liouville Brownian motion. Ann. Probab., 44(4):3076–3110, 2016.
- [KM24] N. Kajino and M. Murugan. Heat kernel estimates for boundary traces of reflected diffusions on uniform domains, 2024. Preprint. Available at arXiv:2312.08546.
- [KN] N. Kajino and R. Noda. Generalized Kac's moment formula. In preparation.
- [Mor21] T. Mori. L^p-Kato class measures and their relations with Sobolev embedding theorems for Dirichlet spaces. J. Funct. Anal., 281(3):Paper No. 109034, 32, 2021.
- [MR06] M. B. Marcus and J. Rosen. *Markov processes, Gaussian processes, and local times*, volume 100 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2006.
- [NTTU24] Y. Nishimori, M. Tomisaki, K. Tsuchida, and T. Uemura. On a convergence of positive continuous additive functionals in terms of their smooth measures, 2024. Preprint. Available at arXiv:2405.03937.

- [Ooi24] T. Ooi. Convergence of processes time-changed by Gaussian multiplicative chaos, 2024. Preprint. Available at arXiv:2305.00734.
- [Rev70] D. Revuz. Mesures associées aux fonctionnelles additives de Markov. I. Trans. Amer. Math. Soc., 148:501–531, 1970.
- [Whi] W. Whitt. Internet supplement to stochastic-process limits. Available at https: //www.columbia.edu/~ww2040/supplementno.pdf.
- [WJ87] R. Williamson and L. Janos. Constructing metrics with the Heine-Borel property. Proc. Amer. Math. Soc., 100(3):567–573, 1987.