

BOUNDARY REPRESENTATIONS FROM CONSTRAINED INTERPOLATION

GAL BEN AYUN AND ELI SHAMOVICH

ABSTRACT. In this paper, we study C^* -envelopes of finite-dimensional operator algebras arising from constrained interpolation problems on the unit disc. In particular, we consider interpolation problems for the algebra H_{node}^∞ that consists of bounded analytic functions on the unit disk that satisfy $f(0) = f(\lambda)$ for some $0 \neq \lambda \in \mathbb{D}$. We show that there exist choices of four interpolation nodes that exclude both 0 and λ , such that if I is the ideal of functions that vanish at the interpolation nodes, then $C_e^*(H_{\text{node}}^\infty/I)$ is infinite-dimensional. This differs markedly from the behavior of the algebra corresponding to interpolation nodes that contain the constrained points studied in the literature. Additionally, we use the distance formula to provide a completely isometric embedding of $C_e^*(H_{\text{node}}^\infty/I)$ for any choice of n interpolation nodes that do not contain the constrained points into $M_n(G_{nc}^2)$, where G_{nc}^2 is Brown's noncommutative Grassmannian.

1. INTRODUCTION

Let $\{z_1, \dots, z_n\} \subset \mathbb{D} = \{z \in \mathbb{C} \mid |z| < 1\}$ be a finite set of points and $\{w_1, \dots, w_n\} \subset \mathbb{C}$. The Pick interpolation problem is the following - does there exist a holomorphic function $f : \mathbb{D} \rightarrow \mathbb{D}$ (called an *interpolating function*) such that $f(z_i) = w_i$? The solution to this problem is well known - a necessary and sufficient condition for the existence of an interpolating function is that the *Pick matrix*

$$\left[\frac{1 - w_i \overline{w_j}}{1 - z_i \overline{z_j}} \right]_{i,j=1}^n$$

is positive semi-definite. This question was solved independently by Pick and Nevanlinna (See for example [31]). More generally, one can formulate the matricial Pick interpolation problem, where the w_i 's are replaced by $s \times r$ matrices $\{W_1, \dots, W_n\}$. Analogously, an interpolating function f with $f(z_i) = W_i$ and $\|f\| \leq 1$ exists if and only if

$$\left[\frac{I - W_i W_j^*}{1 - z_i \overline{z_j}} \right] \geq 0.$$

The operator algebraic approach to the Pick interpolation problem was pioneered by Koranyi and Sz-Nagy [36], and a breakthrough was achieved by Sarason [34]. The key ingredient is the relationship between $H^2(\mathbb{D})$ and $H^\infty(\mathbb{D})$. Recall that $H^2(\mathbb{D})$ is the reproducing kernel Hilbert space (RKHS for short) of all analytic functions on \mathbb{D} with square summable MacLaurin coefficients. The multiplier algebra of $H^2(\mathbb{D})$ is $H^\infty(\mathbb{D})$, the algebra of all bounded analytic function on \mathbb{D} . One encodes the fact that one can test the existence of solutions by the positivity of the corresponding Pick matrix for all sizes of matrices by saying that $H^2(\mathbb{D})$ is a complete Pick space. Complete Pick spaces are well-studied and important objects in function theory. A celebrated theorem of Agler and McCarthy [3] tells us that under some mild assumptions, every complete Pick space is equivalent as an RKHS to a quotient of the Drury-Arveson space. Namely, the RKHS of analytic functions on the unit ball of \mathbb{C}^d (where we allow $d = \infty$) with the reproducing kernel $k(z, w) = \frac{1}{1 - \langle z, w \rangle}$.

However, even in the case of the annulus, the Pick interpolation problem does not have such a simple and elegant solution. Abrahamse [1] has discovered that to solve the scalar Pick interpolation problem on a multiply-connected domain of genus g , one requires a family of kernels parametrized by the g -dimensional torus. In [6], Ball has solved the matricial interpolation problem on multiply-connected domains. However, a sufficient family of kernels described by Ball is parametrized by g -tuples of unitaries of all sizes. In [2], Agler formulated a general approach to the Pick interpolation problem based on families of kernels (see also [24]).

E.S. was partially supported by BSF Grant no. 2022235.

Another generalization is the constrained Pick interpolation problem, where the interpolating function is to satisfy an additional condition. In [14], the authors solve an interpolation problem on \mathbb{D} with the additional constraint that the interpolating function f satisfies $f'(0) = 0$. The algebra of all such functions is denoted H_1^∞ . In this case, a family of kernel functions

$$\{k^{\alpha,\beta}\}_{|\alpha|^2+|\beta|^2=1}$$

is required. Ragupathi [32] generalized this result for subalgebras of H^∞ with *predual factorization*. One main family of examples of subalgebras with predual factorization consists of subalgebras of $H^\infty(\mathbb{D})$ of the form

$$H_B = \mathbb{C} \cdot 1 + BH^\infty,$$

where B is a Blaschke product. The constrained interpolation problem is analogous to the interpolation problem in multiply-connected domains. To be more precise, if one considers multiply-connected domains, the corresponding algebro-geometric object is a curve of genus g . In the constrained case (with a finite Blaschke product), the algebro-geometric object is a singular rational curve. One of the first examples of the interplay between dilation theory and singular rational curves is the Neal parabola [16]. The Neal parabola is the planar curve cut out by the equation $y^2 = x^3$. A parametrization of the Neal parabola is given by the map $z \mapsto (z^2, z^3)$. It is rather straightforward that the algebra H_1^∞ corresponds to the Neal parabola. More generally, H_B corresponds to a parametrization of a rational curve ramified at the zeroes of B with multiplicities prescribed by B . In [7], Ball, Bolotnikov, and ter Horst solved the matrix-valued constrained interpolation problem for H_1^∞ . An approach to the constrained interpolation problem through test functions was taken in [18, 19]. In [12], an approach was taken using the $\mathbb{A}_1(1)$ property of dual operator algebras, which was subsequently developed in [15] with the introduction of the conductor ideal.

The minimality of the corresponding family of kernels can be studied using boundary representations. Boundary representations were introduced by Arveson [4]. Given an operator algebra A one would like to find the smallest C^* -algebra generated by A . This C^* -algebra is called the C^* -envelope of A . Its existence was conjectured by Arveson and proved by Hamana [21]. In [28], Paulsen and McCullough calculate the C^* -envelopes of two operator algebras - the first one is $H^\infty(\mathbb{D}^2)/I$ with

$$I = \left\{ f \in H^\infty(\mathbb{D}^2) : f(0,0) = f\left(\frac{1}{\sqrt{2}},0\right) = f\left(0,\frac{1}{\sqrt{2}}\right) \right\}.$$

This operator algebra arises from an interpolation problem studied by Solazzo [35]. The second is $H^\infty(\mathbb{A})/I_F$, which arises from an interpolation problem in finitely connected domains, studied by Abrahamse [1]. Here, $F \subset \mathbb{A}$ is the finite set of interpolation nodes and I_F is the ideal of functions in $H^\infty(\mathbb{A})$ that vanish on F . It turns out that $C_e^*(H^\infty(\mathbb{A})/I_F) \cong M_n(C(\mathbb{T}))$, where $3 \leq n = |F|$ and $\mathbb{T} = \partial\mathbb{D}$ is the unit circle. Since, in the case of the annulus, the kernel family is parametrized by \mathbb{T} ; we see that this family is minimal and solves the complete Pick interpolation problem on the annulus. In [14], the authors show that if $0 \in F$, then $C_e^*(H_1^\infty/I_F) \cong M_{n+1}$. More generally, Ragupathi [32] calculated $C_e^*(H_B/I_F)$ in the special case that F contains some of the zeroes of B . The C^* -envelopes always turned out to be algebras of matrices.

In this paper, we study constrained interpolation algebras arising from nodal cubics. Namely, for $0 \neq \lambda \in \mathbb{D}$, we consider the algebra

$$H_{\text{node}}^\infty = \{f \in H^\infty : f(0) = f(\lambda)\} = \mathbb{C} \cdot 1 + \frac{z(z-\lambda)}{1-\bar{\lambda}z} H^\infty.$$

We describe the solution to the matricial Pick interpolation problem and the corresponding family of kernels in Section 2. The family is analogous to the one obtained by Ball, Bolotnikov, and ter Horst [7]. The first main result of the paper is in Section 3.

Theorem A (Theorem 3.11 and Theorem 3.18). *For $\lambda = \frac{1}{\sqrt{2}}$, there exists $F = \{z_1, z_2, z_3, z_4\} \subset \mathbb{D}$, such that $C_e^*(H_{\text{node}}^\infty/I_F)$ is infinite-dimensional.*

To this end, we produce a family of pairwise unitarily inequivalent dilation maximal irreducible representations parametrized by the circle without a finite set of points. This shows that the behavior of the C^* -envelope in the case when the set of interpolation nodes does not contain the ramification points is markedly different from the case studied in the literature so far. In Section 4, we provide a candidate for the C^* -envelope of $C_e^*(H_{\text{node}}^\infty/I_F)$ for F that does not contain 0 and λ .

Theorem B (Theorem 4.3). *Let $0, \lambda \notin F \subset \mathbb{D}$ be a finite set. Then, there exists a completely isometric embedding $\Gamma: H_{\text{node}}^\infty/I_F \rightarrow M_n(G_2^{nc})$.*

Here G_{nc}^2 is the noncommutative Grassmannian introduced by Brown [10]. The noncommutative Grassmannian G_{nc}^2 is the noncommutative analog of the Riemann sphere, which in turn parametrizes the family of kernels for the scalar Pick problem.

2. PICK FAMILY FOR H_{NODE}^∞

The content of this section is inspired by the work of Ball, Bolotnikov, and ter Horst [7]. Though some of the proofs are similar to [7], we include them for the sake of completeness. Our method differs from the approach in [7], and our goal is to obtain a distance formula.

For $k, l \in \mathbb{N}$, we denote by $M_{k \times l}(L^2)$ the Hilbert space of $k \times l$ matrices with entries in $L^2(\mathbb{T})$, with the inner product given by

$$\langle F, G \rangle = \frac{1}{2\pi} \int_{\mathbb{T}} \text{tr}[FG^*].$$

Similarly, $M_{k \times l}(L^1)$ is the Banach space of $k \times l$ matrices with entries in L^1 , with the norm given by

$$\|F\|_1 = \frac{1}{2\pi} \int_{\mathbb{T}} \text{tr}[(F^*F)^{\frac{1}{2}}],$$

and $M_{k \times l}(L^\infty)$ is the Banach space of $k \times l$ matrices with entries in L^∞ and the sup norm $\|\cdot\|_\infty$. We will also need the Hilbert space $M_{k \times l}(H^2)$ and the Banach spaces $M_{k \times l}(H^1)$ and $M_{k \times l}(H^\infty)$, which are the subspaces of analytic functions in $M_{k \times l}(L^2)$, $M_{k \times l}(L^1)$ and $M_{k \times l}(L^\infty)$, respectively. We recall that the Riesz representation theorem identifies $M_{k \times l}(L^1)^*$ with $M_{l \times k}(L^\infty)$ via:

$$\varphi_F(G) = \frac{1}{2\pi} \int_{\mathbb{T}} \text{tr}[FG]$$

for $F \in M_{k \times l}(L^1)$ and $G \in M_{l \times k}(L^\infty)$. Moreover, the preannihilator of $M_{k \times l}(H^\infty)$ is given by $M_{l \times k}(zH^1)$.

For a fixed $0 \neq \lambda \in \mathbb{D}$, consider the norm-closed, co-dimension 1 subalgebra of H^∞ defined by

$$H_{\text{node}}^\infty = \{f \in H^\infty : f(0) = f(\lambda)\} = \mathbb{C} \cdot 1 + B_\lambda H^\infty,$$

where $B_\lambda(z) = z \frac{z-\lambda}{1-\bar{\lambda}z}$ is the Blaschke product vanishing at 0 and λ . Moreover, for $z \in \mathbb{D}$ let $k_w(z) = \frac{1}{1-z\bar{w}}$ be the function in H^2 satisfying $\langle f, k_w \rangle = f(w)$ for all $f \in H^2$.

Lemma 2.1. *H_{node}^∞ separates points in $\mathbb{D} \setminus \{0, \lambda\}$.*

Proof. Let $z_1, \dots, z_n \in \mathbb{D} \setminus \{0, \lambda\}$. By Lagrange interpolation, there are polynomials $\{f_i\}_{i=1}^n \subset H^\infty$ satisfying

$$f_i(z_j) = \delta_{i,j}, f_i(0) = f_i(\lambda) = 0.$$

In particular, $f_i \in H_{\text{node}}^\infty$ for all $1 \leq i \leq n$. □

Lemma 2.2. *Let $(\alpha, \beta) \in \mathbb{C}^2$ such that $|\alpha|^2 + |\beta|^2 = 1$ and define*

$$H_{\alpha, \beta}^2 = \{f \in H^2 \mid \exists c \in \mathbb{C}, f(0) = c \cdot \alpha, f(\lambda) = c \cdot (\alpha + f_\lambda(\lambda)\beta)\}.$$

Here $f_\lambda(z) = \frac{\sqrt{1-|\lambda|^2}}{|\lambda|}(k_\lambda(z) - 1)$. Then, $H_{\alpha, \beta}^2$ is a reproducing kernel Hilbert space with kernel function

$$k^{\alpha, \beta}(z, w) = \overline{(\alpha + \beta f_\lambda(w))}(\alpha + \beta f_\lambda(z)) + \frac{B_\lambda(z)\overline{B_\lambda(w)}}{1 - z\bar{w}}.$$

Proof. Note that $f_\lambda(\lambda) = \frac{|\lambda|}{\sqrt{1-|\lambda|^2}}$. Hence, for every $g \in H_{\alpha, \beta}^2$,

$$\langle g, 1 \rangle = c\alpha, \langle g, f_\lambda \rangle = \frac{\sqrt{1-|\lambda|^2}}{|\lambda|}(g(\lambda) - g(0)) = c\beta.$$

In other words, $H_{\alpha,\beta}^2 = \text{span}\{\alpha + \beta f_\lambda(z)\} \oplus B_\lambda H^2$, where the sum is direct since $B_\lambda H^2$ is orthogonal to $\text{span}\{1, f_\lambda\} = \text{span}\{1, k_\lambda\}$, as $B_\lambda(0) = B_\lambda(\lambda) = 0$. Therefore, $H_{\alpha,\beta}^2 \subset H^2$ is a closed subspace. Hence, $H_{\alpha,\beta}^2$ is a RKHS with kernel functions $P_{H_{\alpha,\beta}^2} k_w$. Moreover,

$$k^{\alpha,\beta}(z, w) = \langle P_{H_{\alpha,\beta}^2} k_w, P_{H_{\alpha,\beta}^2} k_z \rangle = \langle k_w, \alpha + \beta f_\lambda \rangle \langle \alpha + \beta f_\lambda, \alpha + \beta f_\lambda \rangle + M_{B_\lambda} M_{B_\lambda}^* k_w = \overline{(\alpha + \beta f_\lambda(w))} (\alpha + \beta f_\lambda) + \frac{B_\lambda(z) \overline{B_\lambda(w)}}{1 - z\bar{w}}.$$

□

Lemma 2.3. For every $|\alpha|^2 + |\beta|^2 = 1$, $H_{\text{node}}^\infty \subseteq \text{Mult}(H_{\alpha,\beta}^2)$. Moreover, if $\alpha \neq 0, -f_\lambda(\lambda)\beta$, then we have equality. As a result, for every $|\alpha|^2 + |\beta|^2 = 1$, $M_f^* k_{z_i}^{\alpha,\beta} = \overline{f(z_i)} k_{z_i}^{\alpha,\beta}$.

Proof. Fix arbitrary (α, β) on the unit sphere. Let $f \in H_{\text{node}}^\infty$ and $f(0) = f(\lambda) = \tilde{c}$. Then, for every $g \in H_{\alpha,\beta}^2$,

$$(fg)(0) = cf(0)\alpha = c\tilde{c}\alpha, (fg)(\lambda) = c\tilde{c}(\alpha + f_\lambda(\lambda)\beta).$$

That is $fg \in H_{\alpha,\beta}^2$, hence $f \in \text{Mult}(H_{\alpha,\beta}^2)$. The last part then follows automatically by properties of $\text{Mult}(\mathcal{H})$ for a general RKHS.

On the other hand, for every $z_0 \in \mathbb{D} \setminus \{0, \lambda\}$,

$$k_{z_0}^{\alpha,\beta}(z_0) = \|k_{z_0}^{\alpha,\beta}\|^2 = |\alpha + f_\lambda(z_0)\beta|^2 + \frac{|B_\lambda(z_0)|^2}{1 - |z_0|^2} > 0.$$

Let $f \in \text{Mult}(H_{\alpha,\beta}^2)$. Since f is a multiplier, $\sup_{z \in \mathbb{D}} |f(z)| \leq \|M_f\|_{\text{mult}} < \infty$. Let $z_0 \in \mathbb{D} \setminus \{0, \lambda\}$. Since $k_{z_0}^{\alpha,\beta}(z_0) > 0$

and is analytic, $\frac{1}{k_{z_0}^{\alpha,\beta}(z)}$ is analytic in a neighborhood of z_0 . Moreover, $f k_{z_0}^{\alpha,\beta} \in H_{\alpha,\beta}^2$, and thus, $f = \frac{f k_{z_0}^{\alpha,\beta}}{k_{z_0}^{\alpha,\beta}}$ is analytic in a neighborhood of z_0 . By Riemann's removable singularity theorem, f is analytic in all of \mathbb{D} . Therefore, $f \in H^\infty$. Assume, in addition, that $\alpha \neq 0, -f_\lambda(\lambda)\beta$. Then, let $p \in H_{\alpha,\beta}^2$ be a polynomial satisfying

$$p(0) = \alpha, p(\lambda) = \alpha + f_\lambda(\lambda)\beta.$$

By assumption:

$$f(0)p(0) = c \cdot \alpha, f(\lambda)p(\lambda) = c \cdot (\alpha + f_\lambda(\lambda)\beta)$$

Therefore $f(0) = f(\lambda) = c$, that is $f \in H_{\text{node}}^\infty$.

□

For fixed $k, l' \geq l \in \mathbb{N}$, let $M_k(H_{\text{node}}^\infty)$ denote the space of $k \times k$ matrices with entries in H_{node}^∞ and define:

$$\mathbb{G}(l' \times l) = \{\alpha, \beta \in M_{l' \times l}(\mathbb{C}) : \alpha\alpha^* + \beta\beta^* = I_l\}$$

Furthermore, for $(\alpha, \beta) \in \mathbb{G}(l' \times l)$, define:

$$(H_{\alpha,\beta}^2)_{k,l,l'} = \{F \in M_{k \times l}(H^2) \mid \exists M \in M_{k \times l'}(\mathbb{C}) \mid F(0) = M\alpha, F(\lambda) = M(\alpha + f_\lambda(\lambda)\beta)\}$$

(if $k = l = l'$, we will shorten this by $(H_{\alpha,\beta}^2)_k$. If $k = l = l' = 1$, this is simply $H_{\alpha,\beta}^2$). By an argument similar to the one in the proof of Lemma 2.2,

$$(H_{\alpha,\beta}^2)_{k,l,l'} = M_{k \times l'}(\mathbb{C})(\alpha + \beta f_\lambda(z)) \oplus B_\lambda M_{k \times l}(H^2).$$

Proposition 2.4. $(H_{\alpha,\beta}^2)_{k,l,l'}$ is a matrix-valued RKHS with kernel function:

$$k^{\alpha,\beta}(z, w) = (\alpha^* + \overline{f_\lambda(w)}\beta^*)(\alpha + f_\lambda(z)\beta) + \frac{B_\lambda(z)\overline{B_\lambda(w)}}{1 - z\bar{w}} I_l.$$

To be more precise, for every $f \in (H_{\alpha,\beta}^2)_{k,l,l'}$, every $w \in \mathbb{D}$, and every $X \in M_{k \times l}(\mathbb{C})$,

$$\text{tr}(f(w)X^*) = \langle f, X k_w^{\alpha,\beta} \rangle.$$

Moreover, for every $F \in M_k(H_{\text{node}}^\infty)$,

$$M_F^* X k(\cdot, w) = F(w)^* X k(\cdot, w)$$

Proof. Let $f \in (H_{\alpha,\beta}^2)_{k,l,l'}$ and write $f = U(\alpha + \beta f_\lambda) + B_\lambda \tilde{f}$ for some $\tilde{f} \in M_k(H^2)$. Then for every $X \in M_{k \times l}(\mathbb{C})$,

$$\begin{aligned} \langle f, Xk^{\alpha,\beta}(\cdot, w) \rangle &= \langle U(\alpha + \beta f_\lambda(z)) + B_\lambda(z)\tilde{f}(z), Xk^{\alpha,\beta}(\cdot, w) \rangle = \\ &= \langle U(\alpha + \beta f_\lambda(z)), X(\alpha^* + \beta^* \overline{f_\lambda(w)})(\alpha + \beta f_\lambda(z)) \rangle + \left\langle B_\lambda(z)\tilde{f}(z), X \frac{B_\lambda(z)\overline{B_\lambda(w)}}{1 - z\overline{w}} \right\rangle = \\ &= \underbrace{\langle U(\alpha + \beta f_\lambda(z)), X(\alpha^* + \beta^* \overline{f_\lambda(w)})(\alpha + \beta f_\lambda(z)) \rangle}_{(1)} + \underbrace{\left\langle B_\lambda(z)\tilde{f}(z), X \frac{B_\lambda(z)\overline{B_\lambda(w)}}{1 - z\overline{w}} \right\rangle}_{(2)}. \end{aligned}$$

We treat each summand separately:

$$\begin{aligned} &\langle U(\alpha + \beta f_\lambda(z)), X(\alpha^* + \beta^* \overline{f_\lambda(w)})(\alpha + \beta f_\lambda(z)) \rangle = \\ &= \langle U\alpha, X(\alpha^* + \beta^* \overline{f_\lambda(w)})\alpha \rangle + \langle U\beta f_\lambda(z), X(\alpha^* + \beta^* \overline{f_\lambda(w)})\beta f_\lambda(z) \rangle = \\ &= \text{tr}[U\alpha\alpha^*(\alpha + \beta f_\lambda(w))X^*] + \text{tr}[U\beta\beta^*(\alpha + \beta f_\lambda(w))X^*] \\ &= \text{tr}[U(\alpha + \beta f_\lambda(w))X^*]. \end{aligned}$$

For the second summand, we observe that B_λ is an isometry. Therefore,

$$\left\langle B_\lambda(z)\tilde{f}(z), X \frac{B_\lambda(z)\overline{B_\lambda(w)}}{1 - z\overline{w}} \right\rangle = B_\lambda(w)\langle \tilde{f}(z), X \frac{1}{1 - z\overline{w}} \rangle = B_\lambda(w) \text{tr}(\tilde{f}(w)X^*).$$

Combining the two, we get our result. For the last claim of the lemma, note that for every $G \in (H_{\alpha,\beta}^2)_{k,l,l'}$,

$$\begin{aligned} \langle G, M_F^* Xk(\cdot, w) \rangle &= \langle FG, Xk(\cdot, w) \rangle = \text{tr}(X^*F(w)G(w)) = \text{tr}((F(w)^*X)^*G(w)) = \\ &= \langle G(w), F(w)^*Xk(\cdot, w) \rangle = \langle G, F(w)^*Xk(\cdot, w) \rangle. \end{aligned}$$

□

With this setup, we present the main theorem of this section:

Theorem 2.5. *For fixed $k \in \mathbb{N}$, let $\{z_1, \dots, z_n\} \subset \mathbb{D}$ and $\{W_1, \dots, W_n\} \in M_k(\mathbb{C})$. There exists a function $F \in M_k(H_{\text{node}}^\infty)$ satisfying:*

$$F(z_i) = W_i \quad \forall 1 \leq i \leq n, \quad \|F\|_\infty \leq 1$$

if and only if for every $(\alpha, \beta) \in \mathbb{G}(k \times k)$ and for every $(X_1, \dots, X_n) \in (M_k(\mathbb{C}))^n$,

$$(2.1) \quad \sum_{i,j=1}^n \text{tr}[X_j k^{\alpha,\beta}(z_i, z_j) X_i^* - W_j^* X_j k^{\alpha,\beta}(z_i, z_j) X_i^* W_i] \geq 0$$

Before proving this theorem, we need a few general results about $(H_{\alpha,\beta}^2)_{k,l,l'}$. To this end, fix some points $F = \{z_1, \dots, z_n\} \subset \mathbb{D}$, let B_F be the Blaschke product with simple zeroes at F and define the ideal:

$$\mathcal{I} = \{f \in H_{\text{node}}^\infty : \forall 1 \leq i \leq n, f(z_i) = 0\} = H_{\text{node}}^\infty \cap B_F H^\infty$$

Lemma 2.6. *For fixed $k \in \mathbb{N}$ set*

$$\mathcal{I} = \{G \in M_k(H_{\text{node}}^\infty) : \forall 1 \leq i \leq n, G(z_i) = 0\} = M_k(H_{\text{node}}^\infty) \cap B_F M_k(H^\infty).$$

Then

$$\mathcal{I}_\perp = M_k(\mathcal{I})_\perp \subset \overline{\varphi_\lambda(z) B_F(z)} M_k(H^1),$$

with $\varphi_\lambda(z) = \frac{z-\lambda}{1-\lambda z} = \frac{B_\lambda(z)}{z}$.

Proof. It is known that $(H^\infty)_\perp = H_0^1 = zH^1$. Since $B_\lambda B_F H^\infty \subset I$ and since taking pre-annihilators is inclusion reversing,

$$I_\perp \subset (B_\lambda B_F H^\infty)_\perp = \overline{B_\lambda B_F} z H_0^1 = \overline{\varphi_\lambda B_F} H^1.$$

Therefore

$$\mathcal{I}_\perp = M_k(I)_\perp = M_k(I_\perp) \subset \overline{\varphi_\lambda B_F} M_k(H^1).$$

□

Lemma 2.7.

$$S = \{h \in \mathcal{I}_\perp \mid \exists s \in \mathbb{D} : \det(h(s)) \neq 0\} \subset \mathcal{I}_\perp$$

is dense in \mathcal{I}_\perp .

Proof. Let $f \in \mathcal{I}_\perp$ be such that for every s , $\det(f(s)) = 0$ and fix $\varepsilon > 0$. Note that $f(\frac{1}{2})$ has a finite spectrum, hence there exists $0 < \delta$ such that $\frac{\delta}{2} \notin \sigma(f(\frac{1}{2}))$ and $\delta \cdot k < \varepsilon$. Now, $f - \delta z I_k$ is invertible at $\frac{1}{2}$ and $f - \delta z I_k \in S$ as $f, \delta z I_k \in \mathcal{I}_\perp$. Moreover,

$$\|f - (f - \delta z I_k)\|_1 = \delta \cdot \|I_k\|_1 = \delta \cdot k < \varepsilon.$$

□

Recall from [30] that a function $f \in M_{k \times \ell}(H^\infty)$ is called *inner* if $f(e^{i\theta})$ is an isometry almost everywhere on \mathbb{T} (in particular, $\ell \leq k$). Additionally, we will say that $g \in M_{s \times t}(H^2)$ is right outer (respectively, left outer) if $\overline{M_s(H^\infty)g} = M_{s \times t}(H^2)$ (respectively, $gM_t(H^\infty) = M_{s \times t}(H^2)$)

Lemma 2.8. *Let $g \in M_k(H^\infty)$ such that g^T is left outer. Then $\det(g(z)) \neq 0$ for all $z \in \mathbb{D}$.*

Proof. Denote by $K(z, w)$ the matricial Szego kernel. As g^T is left outer, there exists a sequence $\{f_n\} \subset M_k(H^\infty)$, such that $\lim_{n \rightarrow \infty} g^T f_n = I_k$ in $M_k(H^2)$. Hence, for every $z \in \mathbb{D}$ and $1 \leq i, j \leq k$:

$$(g^T(z) f_n(z))_{i,j} = \langle g^T(z) f_n(z), E_{i,j} \rangle = \langle g^T f_n, E_{i,j} K(\cdot, z) \rangle \rightarrow \langle I_k, E_{i,j} K(\cdot, z) \rangle = \langle I_k, E_{i,j} \rangle = (I_k)_{i,j}$$

i.e $g^T(z) f_n(z) \rightarrow I_k$ entry-wise. By continuity of the determinant:

$$\det(g^T(z) f_n(z)) \rightarrow \det(I_k) = 1$$

And therefore $\det(g^T(z)) = \det(g(z)) \neq 0$.

□

The following factorization result in $M_k(H^1)$ is, probably, well-known to experts. However, since we do not have a good reference, we include the proof here.

Lemma 2.9. *Every function $h \in M_k(H^1)$ can be factored as*

$$h = g \cdot f$$

with $f \in M_{l \times k}(H^2)$, $g \in M_{k \times l}(H^2)$, g^T left outer and $\|g\|_2^2 = \|f\|_2^2 = \|h\|_1$.

Proof. By [30, Propositions 4.1-4.3], h^T can be factored as $h^T = h_i^T h_o^T$, with h_i^T a $k \times l$ inner function and $h_o^T \in M_{l \times k}(H^2)$ a left outer function. Next, by [34, Theorem 4] we can factor h_o as:

$$h_o = g \tilde{f}$$

with $g \in M_{k \times l}(H^2)$, $\tilde{f} \in M_{l \times k}(H^2)$ and $\tilde{f}^* \tilde{f} = (h_o^* h_o)^{\frac{1}{2}}$, $g^* g = \tilde{f} \tilde{f}^*$ on \mathbb{T} . Lastly, define $f = \tilde{f} h_i$. Then the factorization:

$$h = h_o h_i = g \tilde{f} h_i = g \cdot f$$

satisfies the desired properties.

□

Theorem 2.10. *Every $h \in S$ can be factored as:*

$$h = g f^*$$

with $g \in (H_{\alpha, \beta}^2)_k$, $f \in M_k(L^2) \ominus ((H_{\alpha, \beta}^2)_k \cap B_F M_k(H^2))$, g^T left outer and $\|g\|_2^2 = \|f\|_2^2 = \|h\|_1$.

Proof. It is clear that if such factorization exists, then g and f^* (and, hence, f) are square matrices of size $k \times k$ (as $\det(h(s)) \neq 0$ for some s).

Write $h = \overline{\varphi_\lambda(z)B_F}h'$ with $h' \in M_k(H^1)$. By Lemma 2.9, h' can be factored as gf_0 with $f_0, g \in M_k(H^2)$, g^T outer and $\|g\|_2^2 = \|f_0\|_2^2 = \|h\|_1$. Now factor:

$$h = \overline{\varphi_\lambda(z)B_F}gf_0$$

Taking $f = \varphi_\lambda(z)B_Ff_0^*$ and observing that $\|f\|_2^2 = \|f_0\|_2^2$ (as both φ_λ and B_F are inner functions and $\|f_0\|_2^2 = \|f_0^*\|_2^2$) finishes the factorization.

It is left to prove the second part. Let

$$\tilde{\alpha} = I_k, \tilde{\beta} = \frac{1}{f_\lambda(\lambda)} (g(0)^{-1}g(\lambda) - I_k), \tilde{U} = g(0)$$

and define

$$\alpha = (I_k + \tilde{\beta}\tilde{\beta}^*)^{-\frac{1}{2}}, \beta = (I_k + \tilde{\beta}\tilde{\beta}^*)^{-\frac{1}{2}}\tilde{\beta}, U = \tilde{U} (I_k + \tilde{\beta}\tilde{\beta}^*)^{\frac{1}{2}}$$

Then $\alpha\alpha^* + \beta\beta^* = I_k$ and $U\alpha = g(0)$, $U(\alpha + f_\lambda(\lambda)\beta) = g(\lambda)$. Hence $g \in (H_{\alpha,\beta}^2)_k$. Note that U , α , and $\alpha + f_\lambda(\lambda)\beta$ are invertible.

Note that for every $\phi \in \mathcal{I}$,

$$\langle \phi g, f \rangle = \frac{1}{2\pi} \int_{\mathbb{T}} \phi g f^* = \frac{1}{2\pi} \int_{\mathbb{T}} \phi h = 0,$$

that is $f \in (\mathcal{I}g)^\perp$. So it's enough to prove that $\overline{\mathcal{I}g} = (H_{\alpha,\beta}^2)_k \cap B_F M_k(H^2)$. First, it is clear that

$$\mathcal{I}g \subset (H_{\alpha,\beta}^2)_k \cap B_F M_k(H^2).$$

For the other direction, let $\psi \in (H_{\alpha,\beta}^2)_k \cap B_F M_k(H^2)$, and write

$$\psi(\lambda) = V(\alpha + f_\lambda(\lambda)\beta), \psi(0) = V\alpha$$

As g^T is left-outer, g is right-outer, and therefore $\overline{M_k(H^\infty)g} = M_k(H^2)$. So there exists a sequence $\{\psi_m\}_m \subset M_k(H^\infty)$ such that $\psi_m g \rightarrow \psi$. Write:

$$\psi_m = \psi_m(0) + (\psi_m(\lambda) - \psi_m(0)) \frac{f_\lambda}{f_\lambda(\lambda)} + B_\lambda \tilde{\psi}_m$$

with $\tilde{\psi}_m \in M_k(H_{\text{node}}^\infty)$. Therefore

$$\varphi_m = \psi_m - (\psi_m(\lambda) - \psi_m(0)) \frac{f_\lambda}{f_\lambda(\lambda)} \in M_k(H_{\text{node}}^\infty).$$

Moreover,

$$\|\varphi_m g - \psi\| \leq \|\psi_m g - \psi\| + \left\| \frac{f_\lambda}{f_\lambda(\lambda)} \right\| \|\psi_m(\lambda) - \psi_m(0)\| \rightarrow 0,$$

where in the last step we used that

$$\lim_{m \rightarrow \infty} \psi_m(\lambda) - \psi_m(0) = \psi(\lambda)g(\lambda)^{-1} - \psi(0)g(0)^{-1} = VU^{-1} - VU^{-1} = 0$$

Lastly, we can always assume that the interpolation points don't contain both 0 and λ - otherwise, we can take one out and solve an equivalent interpolation problem with $n - 1$ points. Hence, by entry-wise application of Lagrange interpolation theorem, we have polynomials $\{P_i\} \subset H_{\text{node}}^\infty$ such that $P_i(z_j) = \delta_{i,j}I_k$. Now consider the sequence $\varphi_m - \sum_{i=1}^n \varphi_m(z_i)P_i$. Since $\varphi_m(z_i)g(z_i) \rightarrow \psi(z_i) = 0$ for every $1 \leq i \leq n$ and $g(z_i)$ is invertible,

$$\|(\varphi_m - \sum_{i=1}^n \varphi_m(z_i)P_i)g - \psi\| \leq \|\varphi_m g - \psi\| + \sum_{i=1}^n \|\varphi_m(z_i)\| \|P_i g\| \rightarrow 0.$$

Moreover, $(\varphi_m - \sum_{i=1}^n \varphi_m(z_i)P_i)g \in \mathcal{I}g$, concluding the proof. \square

Let $\mathcal{M}_{\alpha,\beta} = \text{span}\{Xk^{\alpha,\beta}(\cdot, z_i) \mid X \in M_k(\mathbb{C}) \text{ and } 1 \leq i \leq n\}$. It is an immediate consequence of the reproducing property that

$$(\mathcal{M}_{\alpha,\beta})_k = (H_{\alpha,\beta}^2)_k \ominus ((H_{\alpha,\beta}^2)_k \cap B_F M_k(H^2)).$$

Remark 2.11. Similarly, if $(\alpha, \beta) \in \mathbb{G}(l' \times l)$, then:

$$(\mathcal{M}_{\alpha,\beta})_{k,l',l} = \{Xk^{\alpha,\beta}(z_i, z_j) : X \in M_{k \times l}(\mathbb{C})\} = (H_{\alpha,\beta}^2)_{k,l',l} \ominus ((H_{\alpha,\beta}^2)_{k,l',l} \cap B_F M_{k \times l}(H^2))$$

Lastly, for Let $(\alpha, \beta) \in \mathbb{G}(k \times k)$, define:

$$(\mathcal{M}_{\alpha,\beta})_k = \text{span}\{Xk^{\alpha,\beta}(\cdot, z_j) : X \in M_k(\mathbb{C})\}$$

We are now ready to prove Theorem 2.5.

Proof of necessity in Theorem 2.5. Fix $k \in \mathbb{N}$ and assume there exists $F \in M_k(H_{\text{node}}^\infty)$ such that $F(z_i) = W_i$ and $\|F\|_\infty \leq 1$. Note that:

$$(2.2) \quad 1 \geq \|F\|_\infty = \|M_F\|_{op} \geq \|P_{(\mathcal{M}_{\alpha,\beta})_k} M_F|_{(\mathcal{M}_{\alpha,\beta})_k}\|_{op}$$

Therefore, $P_{(\mathcal{M}_{\alpha,\beta})_k} M_F|_{(\mathcal{M}_{\alpha,\beta})_k}$ is a contraction, i.e:

$$1 - P_{(\mathcal{M}_{\alpha,\beta})_k} M_F M_F^*|_{(\mathcal{M}_{\alpha,\beta})_k} \geq 0$$

where we used the fact that $(\mathcal{M}_{\alpha,\beta})_k$ is M_F^* invariant by properties of matrix valued RKHS. Therefore:

$$\begin{aligned} 0 &\leq \left\langle (1 - P_{(\mathcal{M}_{\alpha,\beta})_k} M_F M_F^*|_{(\mathcal{M}_{\alpha,\beta})_k}) \left(\sum_{j=1}^n X_j k^{\alpha,\beta}(\cdot, z_j) \right), \sum_{i=1}^n X_i k^{\alpha,\beta}(\cdot, z_i) \right\rangle = \\ &= \left\langle \sum_{j=1}^n X_j k^{\alpha,\beta}(\cdot, z_j), \sum_{i=1}^n X_i k^{\alpha,\beta}(\cdot, z_i) \right\rangle - \left\langle M_F^* \left(\sum_{j=1}^n X_j k^{\alpha,\beta}(\cdot, z_j) \right), M_F^* \left(\sum_{i=1}^n X_i k^{\alpha,\beta}(\cdot, z_i) \right) \right\rangle = \\ &= \sum_{i,j=1}^n \langle X_j k^{\alpha,\beta}(\cdot, z_j), X_i k^{\alpha,\beta}(\cdot, z_i) \rangle - \sum_{i,j=1}^n \langle F(z_j)^* X_j k^{\alpha,\beta}(\cdot, z_j), F(z_i)^* X_i k^{\alpha,\beta}(\cdot, z_i) \rangle = \\ &= \sum_{i,j=1}^n \text{tr} (X_i^* X_j k^{\alpha,\beta}(z_i, z_j) - X_i^* F(z_i) F(z_j)^* X_j k^{\alpha,\beta}(z_i, z_j)) = \\ &= \text{tr} \left(\sum_{i,j=1}^n X_j k^{\alpha,\beta}(z_i, z_j) X_i^* - F(z_j)^* X_j k^{\alpha,\beta}(z_i, z_j) X_i^* F(z_i) \right). \end{aligned}$$

Letting $F(z_j) = W_j$ concludes the proof of necessity. \square

Remark 2.12. In fact, the same proof of necessity works with $(\alpha, \beta) \in \mathbb{G}(l' \times l)$ and:

$$(X_1, \dots, X_n) \in (M_{k \times l}(\mathbb{C}))^n$$

Proof of sufficiency in Theorem 2.5. Let z_1, \dots, z_n be the interpolation nodes and $W_1, \dots, W_n \in M_k(\mathbb{C})$ be the interpolation data. Using Lagrange interpolation entry-wise, we obtain $Q \in M_k(H_{\text{node}}^\infty)$ such that $Q(z_i) = W_i$. Since, $(\mathcal{M}_{\alpha,\beta})_k$ is M_Q^* invariant,

$$\|P_{(\mathcal{M}_{\alpha,\beta})_k} M_Q|_{(\mathcal{M}_{\alpha,\beta})_k}\| = \|M_Q^*|_{(\mathcal{M}_{\alpha,\beta})_k}\| \leq 1,$$

where the last inequality is true by our assumption (2.1) and the calculation in the proof of necessity.

Define a linear functional on $(\mathcal{I})_\perp$ by:

$$L_Q(h) = \frac{1}{2\pi} \int_{\mathbb{T}} \text{tr}[Qh].$$

We show that $\|L_Q\| \leq 1$. By Theorem 2.10, for $h \in S$, we can convert the formula of L_Q to an L^2 -inner product as follows:

$$L_Q(h) = \frac{1}{2\pi} \int_{\mathbb{T}} \text{tr}[Qh] = \frac{1}{2\pi} \int_{\mathbb{T}} \text{tr}[Qg f^*] = \langle Qg, f \rangle_{M_k(L^2)} = \left\langle P_{H_{\alpha,\beta}^2} Qg, P_{(H_{\alpha,\beta}^2 \cap B_F M_k(H^2))^\perp} f \right\rangle_{M_k(L^2)}.$$

Since $(\mathcal{M}_{\alpha,\beta})_k = (H_{\alpha,\beta}^2)_k \ominus ((H_{\alpha,\beta}^2)_k \cap B_F M_k(H^2))$, we can rewrite the equality above as:

$$\begin{aligned}
L_Q(h) &= \langle P_{(\mathcal{M}_{\alpha,\beta})_k} Qg, P_{(\mathcal{M}_{\alpha,\beta})_k} f \rangle = \\
&= \left\langle P_{(\mathcal{M}_{\alpha,\beta})_k} M_Q (P_{(\mathcal{M}_{\alpha,\beta})_k} g + P_{H_{\alpha,\beta}^2 \cap B_F M_k(H^2)} g), P_{(\mathcal{M}_{\alpha,\beta})_k} f \right\rangle = \\
&= \langle P_{(\mathcal{M}_{\alpha,\beta})_k} M_Q P_{(\mathcal{M}_{\alpha,\beta})_k} g, P_{(\mathcal{M}_{\alpha,\beta})_k} f \rangle.
\end{aligned}$$

The last equality follows from the fact that $H_{\alpha,\beta}^2 \cap B_F M_k(H^2)$ is multiplier invariant. By density of S ,

$$\|L_Q\| = \sup_{h \in S, \|h\|=1} |L_Q(h)| \leq \sup_{(f,g) \in A} |\langle P_{(\mathcal{M}_{\alpha,\beta})_k} M_Q P_{(\mathcal{M}_{\alpha,\beta})_k} g, P_{(\mathcal{M}_{\alpha,\beta})_k} f \rangle|,$$

where $A = \{(f, g) : \|f\| = 1, \|g\| = 1, g \in (H_{\alpha,\beta}^2)_k, f \in ((H_{\alpha,\beta}^2)_k \cap B_F M_k(H^2))^\perp\}$. Lastly, by Cauchy-Schwarz:

$$\sup_{(f,g) \in A} |\langle P_{(\mathcal{M}_{\alpha,\beta})_k} M_Q P_{(\mathcal{M}_{\alpha,\beta})_k} g, P_{(\mathcal{M}_{\alpha,\beta})_k} f \rangle| \leq \|P_{(\mathcal{M}_{\alpha,\beta})_k} M_Q|_{(\mathcal{M}_{\alpha,\beta})_k}\| \leq 1.$$

By the Hahn-Banach Theorem, we can extend L_Q to a bounded functional on $M_k(L^1)$. By the Riesz representation theorem, such a functional has the form of L_S for some $S \in M_k(L^\infty)$, where

$$L_S(h) = \frac{1}{2\pi} \int_{\mathbb{T}} \text{tr}[S(z)h(z)]|dz|$$

and $1 \geq \|L_S\| = \|S\|_\infty$. As L_S is an extension of L_Q , $L_{S-Q}|_{(\mathcal{I})^\perp} = L_S|_{(\mathcal{I})^\perp} - L_Q = 0$, that is, $S - Q \in ((\mathcal{I})^\perp)^\perp = \mathcal{I}$. As $Q \in M_k(H_{\text{node}}^\infty)$ and $\mathcal{I} \subset M_k(H_{\text{node}}^\infty)$, $S \in M_k(H_{\text{node}}^\infty)$ and $S(z_i) = Q(z_i) = W_i$. We noted that $\|S\|_\infty \leq 1$, which means S is the desired solution. \square

Corollary 2.13. (Scalar case) Given $\{z_1, \dots, z_n\} \subset \mathbb{D}$, $\{w_1, \dots, w_n\} \subset \mathbb{C}$, there exists $f \in H_{\text{node}}^\infty$ such that $f(z_i) = w_i$ if and only if the Pick matrix

$$[(1 - w_i \bar{w}_j) k^{\alpha,\beta}(z_i, z_j)]_{i,j=1}^n$$

is positive semi-definite for all $\alpha, \beta \in \mathbb{G}(1 \times 1)$.

Proof. By what we proved, such a function exists if and only if for every $(\alpha, \beta) \in \mathbb{G}(1 \times 1)$ and for every choice of scalars $(x_1, \dots, x_n) \in \mathbb{C}^n$,

$$\sum_{i,j=1}^n (1 - w_i \bar{w}_j) k^{\alpha,\beta}(z_i, z_j) x_i \bar{x}_j \geq 0.$$

This condition is exactly the condition

$$\left\langle [(1 - w_i \bar{w}_j) k^{\alpha,\beta}(z_i, z_j)]_{i=1}^n \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}, \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \right\rangle \geq 0$$

for every $\begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{C}^n$, which is exactly the condition that

$$[(1 - w_i \bar{w}_j) k^{\alpha,\beta}(z_i, z_j)]_{i=1}^n \geq 0.$$

\square

We conclude this section with two important corollaries that connect the norm in H_{node}^∞/I and the norm of compression operators.

Corollary 2.14. (Distance formula) Let \mathcal{I} be as before. Then for every $F \in M_k(H_{\text{node}}^\infty)$,

$$\text{dist}(F, \mathcal{I}) = \|F + \mathcal{I}\| = \sup_{(\alpha,\beta) \in \mathbb{G}(k \times k)} \|M_F^*|_{\mathcal{M}_{\alpha,\beta}}\|.$$

Proof. By definition of the quotient norm,

$$\|F + \mathcal{I}\| = \inf\{\|G\| : G \in M_k(H_{\text{node}}^\infty), F(z_i) = G(z_i) = W_i, \forall i = 1, \dots, n\}.$$

From the contractive case, it follows that there exists a $G \in M_k(H_{\text{node}}^\infty)$, such that $G(z_i) = W_i$ and $\|G\| \leq M$ if and only if

$$(2.3) \quad \sum_{i,j=1}^n \text{tr} [M^2 X_j k^{\alpha,\beta}(z_i, z_j) X_i^* - W_j^* X_j k^{\alpha,\beta}(z_i, z_j) X_i^* W_i] \geq 0$$

for all n -tuples $(X_1, \dots, X_n) \in (M_k(\mathbb{C}))^n$ and $(\alpha, \beta) \in \mathbb{G}(k \times k)$. Therefore,

$$\begin{aligned} \text{dist}(F, \mathcal{I}) &= \inf\{M : (2.3) \text{ is satisfied } \forall (\alpha, \beta) \in \mathbb{G}(k \times k), (X_1, \dots, X_n) \in (M_k(\mathbb{C}))^n\} = \\ &= \inf\{M : \|M_F^*|_{\mathcal{M}_{\alpha,\beta}}\| \leq M, \forall (\alpha, \beta) \in \mathbb{G}(k \times k)\}. \end{aligned}$$

Clearly,

$$\sup_{(\alpha,\beta) \in \mathbb{G}(k \times k)} \|M_F^*|_{\mathcal{M}_{\alpha,\beta}}\| = \inf\{M : \|M_F^*|_{\mathcal{M}_{\alpha,\beta}}\| \leq M, \forall (\alpha, \beta) \in \mathbb{G}(k \times k)\}$$

□

For fixed $m, r \in \mathbb{N}$ and $(\alpha, \beta) \in \mathbb{G}(m \times m)$, define a map $\varphi_{\alpha,\beta} : M_r(H_{\text{node}}^\infty) \rightarrow B((\mathcal{M}_{\alpha,\beta})_{r,m})$ by

$$\varphi_{\alpha,\beta}(F) = P_{(\mathcal{M}_{\alpha,\beta})_{r,m}} M_F |_{(\mathcal{M}_{\alpha,\beta})_{r,m}}$$

By Sarason's Lemma [33, Lemma 0], this map is an algebra homomorphism, and it is straight-forward that $\ker(\varphi) = \mathcal{I}$.

Corollary 2.15. (Generalized distance formula) *Let $r \in \mathbb{N}$ and $F \in M_r(H_{\text{node}}^\infty)$. Denote $M_r(I) = \mathcal{I}$. Then:*

$$\|F + \mathcal{I}\| = \sup_m \sup \left\{ \|P_{(\mathcal{M}_{\alpha,\beta})_{r,m}} M_F |_{(\mathcal{M}_{\alpha,\beta})_{r,m}}\| \mid (\alpha, \beta) \in \mathbb{G}(m \times m) \right\}$$

Proof. The fact that the RHS is greater or equal to the LHS is clear by Corollary 2.14. Fix $m \in \mathbb{N}$ and $(\alpha, \beta) \in \mathbb{G}(m \times m)$, and consider φ as before. Clearly, $\varphi_{\alpha,\beta}$ is a contraction, as

$$\|\varphi_{\alpha,\beta}(F)\| = \|P_{(\mathcal{M}_{\alpha,\beta})_{r,m}} M_F |_{(\mathcal{M}_{\alpha,\beta})_{r,m}}\| \leq \|P_{(\mathcal{M}_{\alpha,\beta})_{r,m}}\| \|M_F|_{H_{\alpha,\beta}^2}\| \leq \|M_F\| = \|F\|_\infty,$$

where we used the fact that as a multiplier on $H_{\alpha,\beta}^2$, $M_F|_{H_{\alpha,\beta}^2}$ is the compression of $M_F \in \text{Mult}(M_k(H^2))$ to the invariant subspace $H_{\alpha,\beta}^2$, and that $\|M_F\|_{op} = \|F\|_\infty$ for every $F \in M_k(H^\infty)$. Lastly, note that for every $F \in M_r(H_{\text{node}}^\infty)$, $G \in \mathcal{I}$,

$$\|\varphi(F)\| = \|\varphi(F + G)\| \leq \|F + G\|.$$

Hence

$$\|P_{(\mathcal{M}_{\alpha,\beta})_{r,m}} M_F |_{(\mathcal{M}_{\alpha,\beta})_{r,m}}\| \leq \inf_{G \in \mathcal{I}} \|F + G\| = \|F + I\|.$$

As this is independent of m and (α, β) ,

$$\sup_m \sup \left\{ \|P_{(\mathcal{M}_{\alpha,\beta})_{r,m}} M_F |_{(\mathcal{M}_{\alpha,\beta})_{r,m}}\| \mid (\alpha, \beta) \in \mathbb{G}(m \times m) \right\} \leq \|F + I\|$$

as desired. □

3. BOUNDARY REPRESENTATIONS FROM SCALAR INTERPOLATION

We begin this section with some results on representations of H_{node}^∞/I for general $0 \neq \lambda \in \mathbb{D}$ and $\{z_1, \dots, z_n\} \subset \mathbb{D} \setminus \{0, \lambda\}$. Here I is again the ideal of function that vanish on the z_i . We view H_{node}^∞/I as an operator algebra with the natural structure. By a representation of an operator algebra on a Hilbert space \mathcal{H} , we mean a unital completely contractive homomorphism to $B(\mathcal{H})$. The spectral structure of H_{node}^∞/I is quite simple and is described in the following proposition:

Lemma 3.1. H_{node}^∞/I is generated as an algebra by $h + I$, with $h \in H_{\text{node}}^\infty$ being the degree $n + 1$ polynomial satisfying $h(z_i) = z_i$ for all $i = 1, \dots, n$ and $h(0) = h(\lambda) = 0$. The spectrum of H_{node}^∞/I consists of the characters of evaluation at z_1, \dots, z_n . Therefore, for every finite-dimensional representation $\varphi : H_{\text{node}}^\infty/I \rightarrow B(\mathcal{H})$ and every $f \in H_{\text{node}}^\infty$, $\varphi(f + I) = f(\varphi(h + I))$ is diagonalizable.

Proof. Let $f \in H_{\text{node}}^\infty$. There exists a polynomial $p(z) = \sum_{k=0}^{n+1} a_k z^k \in H_{\text{node}}^\infty$ satisfying $p(z_i) = f(z_i)$ for all $1 \leq i \leq n$. Define $\tilde{f}(z) = \sum_{k=0}^{n+1} a_k h^k \in H_{\text{node}}^\infty$. Then

$$\tilde{f}(z_i) = \sum_{k=0}^{n+1} a_k h^k(z_i) = \sum_{k=0}^{n+1} a_k z_i^k = p(z_i) = f(z_i).$$

So $f + I = \tilde{f} + I = \sum_{k=0}^{n+1} a_k h^k + I = \sum_{k=0}^{n+1} a_k (h + I)^k$. We conclude that every character of H_{node}^∞/I is determined on h . It is also easy to see that for every $\alpha \in \mathbb{C} \setminus \{z_1, \dots, z_n\}$, $h - \alpha + I$ is invertible in H_{node}^∞/I . The claim on the spectrum now follows.

For the claim that for every $f \in H_{\text{node}}^\infty$, $\varphi(f + I) = f(\varphi(h + I))$, we use a density argument. Firstly, since I is weak* closed, the quotient map

$$\pi : H_{\text{node}}^\infty \rightarrow H_{\text{node}}^\infty/I$$

is weak* continuous. Therefore $\varphi \circ \pi$ is weak* continuous. Since polynomials are weak* dense, it is enough to prove it for polynomials. However, this follows from the fact that φ is an algebra homomorphism. It is not hard to check that the minimal polynomial of $\varphi(h + I)$ factors into different linear factors, i.e, $\varphi(h + I)$ is diagonalizable. The general claim follows. \square

Lemma 3.2. For $(\alpha, \beta) \in \mathbb{G}(1 \times 1)$, the map

$$\rho_{\alpha, \beta} : H_{\text{node}}^\infty/I \rightarrow B(\mathcal{M}_{\alpha, \beta})$$

given by $f + I \rightarrow P_{\mathcal{M}_{\alpha, \beta}} M_f|_{\mathcal{M}_{\alpha, \beta}}$ is an operator algebra representation.

Proof. The claim follows at once from the Sarason Lemma [33, Lemma 0]. \square

Remark 3.3. For every (α, β) on the unit sphere of \mathbb{C}^2 , let $A = \rho_{\alpha, \beta}(h + I)$. Then, the minimal polynomial of A is precisely, $p(z) = \prod_{i=1}^n (z - z_i)$.

Now we need to recall some notations and terminology from the theory of C^* -covers and boundary representations of operator algebras. Recall, that a C^* -cover of an operator algebra \mathcal{A} is a C^* -algebra B together with a completely isometric homomorphisms $\iota : \mathcal{A} \rightarrow B$, such that $B = C^*(\iota(\mathcal{A}))$. For every operator algebra \mathcal{A} , there are two extremal C^* -covers. Let $C_{\text{max}}^*(\mathcal{A})$ denote its maximal C^* -cover. The maximal C^* -cover was constructed by Kirchberg and Wassermann [25]. The maximality of this C^* -cover is encoded in its universal property. For every C^* -algebra B and a completely contractive homomorphism $\varphi : \mathcal{A} \rightarrow B$, there exists a $*$ -homomorphism $\pi : C_{\text{max}}^*(\mathcal{A}) \rightarrow B$ extending φ . The minimal C^* -cover of \mathcal{A} is the C^* -envelope that we will denote by $C_e^*(\mathcal{A})$. The existence of the C^* -envelope was conjectured by Arveson [4] and proved by Hamana [21]. The C^* -envelope of \mathcal{A} has the universal property that for every C^* -cover $\iota : \mathcal{A} \rightarrow B$, there exists a surjective homomorphism $\pi : B \rightarrow C_e^*(\mathcal{A})$ that is completely isometric on $\iota(\mathcal{A})$.

Definition 3.4. Let $\mathcal{H} \subset \mathcal{K}$ be Hilbert spaces, \mathcal{A} an operator algebra and

$$\psi_1 : \mathcal{A} \rightarrow B(\mathcal{H}), \psi_2 : \mathcal{A} \rightarrow B(\mathcal{K})$$

be two representations. We say that ψ_2 is *dilation* of ψ_1 if $P_{\mathcal{H}} \psi_2|_{\mathcal{H}} = \psi_1$. Moreover, ψ_1 is called *dilation maximal* if every dilation of ψ_1 is trivial, i.e if ψ_2 dilates ψ_1 then $\psi_2 = \psi_1 \oplus \psi$ for some representation ψ .

It follows from Sarason's lemma that \mathcal{H} is semi-invariant for $\psi_2(A)$. The following definition is due to Arveson [4].

Definition 3.5. Let \mathcal{A} be an operator algebra. A representation

$$\pi : C^*(\mathcal{A}) \rightarrow B(\mathcal{H})$$

is said to have the unique extension property if, for every unital completely positive map (ucp for short) $\varphi : C^*(\mathcal{A}) \rightarrow B(\mathcal{H})$, if $\pi|_{\mathcal{A}} = \varphi|_{\mathcal{A}}$, then $\pi = \varphi$.

Definition 3.6. Let (B, ι) be a C^* -cover of an operator algebra \mathcal{A} . A representation

$$\psi : B \rightarrow B(\mathcal{H})$$

is said to be a boundary representation if it is irreducible and has the unique extension property.

It follows from results of Muhly and Solel [29] (see also [17] and [5]) that π has the unique extension property if and only if $\pi|_{\mathcal{A}}$ is dilation maximal. The following theorem is a particular case of [13, Proposition 6.5.1].

Theorem 3.7. *Let \mathcal{A} be an operator algebra and $\psi : C_{\max}^*(\mathcal{A}) \rightarrow B(\mathcal{H})$ be a boundary representation of \mathcal{A} . Let $\pi : C_{\max}^*(\mathcal{A}) \rightarrow C_e^*(\mathcal{A})$ be the surjective $*$ -homomorphism guaranteed by the universal properties. Then, there is a representation*

$$\sigma : C_e^*(\mathcal{A}) \rightarrow B(\mathcal{H})$$

such that $\sigma \circ \pi = \psi$ (that is, ψ factors through $C_e^*(\mathcal{A})$).

Before presenting the main result of this section, we need two technical lemmas.

Lemma 3.8. *Fix $0 \neq \lambda \in \mathbb{D}$ and let $\omega \in \mathbb{D}$.*

(1) *If $\exists \zeta \neq \omega$ such that $B_\lambda(\zeta) = B_\lambda(\omega)$ then*

$$B_{\omega, \zeta}(z) := \frac{z - \zeta}{1 - \bar{\zeta}z} \cdot \frac{z - \omega}{1 - \bar{\omega}z} \in H_{\text{node}}^\infty.$$

(2) *If $B'_\lambda(\omega) = 0$ then*

$$B_\omega^2(z) = \left(\frac{z - \omega}{1 - \bar{\omega}z} \right)^2 \in H_{\text{node}}^\infty.$$

Proof. Let $\varphi \in \text{Aut}(\mathbb{D})$ be such that $\varphi(B_\lambda(\omega)) = 0$.

(1) In the first case, $\varphi \circ B_\lambda$ is a Blaschke product with 2 zeroes that vanishes at ζ and ω , hence

$$\varphi \circ B_\lambda = e^{i\theta} \frac{z - \zeta}{1 - \bar{\zeta}z} \cdot \frac{z - \omega}{1 - \bar{\omega}z} \in H_{\text{node}}^\infty.$$

(2) In the second case, by the chain rule

$$(\varphi \circ B_\lambda)'(\omega) = \varphi'(B_\lambda(\omega)) B'_\lambda(\omega) = 0,$$

i.e $\varphi \circ B_\lambda$ is a Blaschke product with at most 2 zeroes with a double zero at ω . Therefore

$$\varphi \circ B_\lambda(z) = e^{i\theta} B_\omega^2 \in H_{\text{node}}^\infty.$$

□

Lemma 3.9. *Let $0 \neq \lambda \in \mathbb{D}$. Let $z_1, \dots, z_4 \in \mathbb{D} \setminus \{0, \lambda\}$ be such that $B_\lambda(z_1) = B_\lambda(z_4)$ and $B_\lambda(z_2) = B_\lambda(z_3)$. Assume, furthermore, that there exists $\omega \in \mathbb{D}$ such that $B_{z_1, z_2, \omega} \in H_{\text{node}}^\infty$. Then there exists some $\zeta \in \mathbb{D}$ such that*

$$B_{z_3, z_4, \zeta}(z) = \frac{z - z_3}{1 - \bar{z}_3 z} \cdot \frac{z - z_4}{1 - \bar{z}_4 z} \cdot \frac{z - \zeta}{1 - \bar{\zeta} z} \in H_{\text{node}}^\infty.$$

Moreover, if $z_1, \dots, z_4 \in \mathbb{R}$ and $\lambda \in \mathbb{R}$, then $\omega, \zeta \in \mathbb{R}$.

Proof. By Lemma 3.8 and by our assumption

$$B_{z_1, z_4}, B_{z_2, z_3} \in H_{\text{node}}^\infty.$$

If $B'_\lambda(\omega) = 0$, then $\left(\frac{z - \omega}{1 - \bar{\omega}z} \right)^2 \in H_{\text{node}}^\infty$. But then

$$B_{z_3, z_4, \omega}(0) = \frac{B_{z_1, z_2}(0) \cdot B_{z_3, z_4}(0) \cdot B_\omega^2(0)}{B_{z_1, z_2, \omega}(0)} = \frac{B_{z_1, z_2}(\lambda) \cdot B_{z_3, z_4}(\lambda) \cdot B_\omega^2(\lambda)}{B_{z_1, z_2, \omega}(\lambda)} = B_{z_3, z_4, \omega}(\lambda),$$

where we used the fact that

$$B_{z_1, z_2} \cdot B_{z_3, z_4} = B_{z_1, z_4} \cdot B_{z_2, z_3}.$$

Otherwise, since B_λ is 2 to 1, there exists some $\zeta \neq \omega$ such that $B_\lambda(\zeta) = B_\lambda(\omega)$. By Lemma 3.8, $B_{\zeta, \omega} \in H_{\text{node}}^\infty$. By similar arguments,

$$B_{z_3, z_4, \zeta}(0) = \frac{B_{z_1, z_2}(0) \cdot B_{z_3, z_4}(0) \cdot B_{\zeta, \omega}(0)}{B_{z_1, z_2, \omega}(0)} = \frac{B_{z_1, z_2}(\lambda) \cdot B_{z_3, z_4}(\lambda) \cdot B_{\zeta, \omega}(\lambda)}{B_{z_1, z_2, \omega}(\lambda)} = B_{z_3, z_4, \zeta}(\lambda).$$

Lastly, assume z_1, \dots, z_4, λ are real. By plugging 0 and λ into $B_{z_1, z_2, \omega}$ we obtain the following equality:

$$-z_1 z_2 \omega = \frac{\lambda - z_1}{1 - z_1 z} \cdot \frac{\lambda - z_2}{1 - z_2 z} \cdot \frac{\lambda - \omega}{1 - \bar{\omega} \lambda}$$

Therefore

$$\omega \cdot \frac{1 - \bar{\omega} \lambda}{\lambda - \omega} = \frac{\omega - |\omega|^2 \lambda}{\lambda - \omega} \in \mathbb{R}$$

Consequently, there exists $c \in \mathbb{R}$ such that

$$(c + 1)\omega = \lambda(c + |\omega|^2) \in \mathbb{R}.$$

If $c = -1$, $\lambda = 0$ or $|\omega| = 1$, contradiction. Hence $\omega \in \mathbb{R}$. A similar argument shows that $\zeta \in \mathbb{R}$. \square

From here on out, $0 \neq \lambda \in \mathbb{R} \cap \mathbb{D}$ is fixed. For every (α, β) on the unit sphere of \mathbb{C}^2 , we will denote The following definition encodes the technical properties that we need to ensure the abundance of boundary representations.

Definition 3.10. Four points $z_1, \dots, z_4 \in \mathbb{D} \cap \mathbb{R} \setminus \{0, \lambda\}$ will be called **good points** with respect to λ , if there exists an infinite family $A \subset \mathbb{T} \subset \mathbb{G}(1 \times 1)$ such that the following conditions hold:

- (1) $B_\lambda(z_1) = B_\lambda(z_4)$, $B_\lambda(z_2) = B_\lambda(z_3)$.
- (2) There exists $\omega \in \mathbb{D} \cap \mathbb{R}$ such that

$$B_{z_1, z_2, \omega}(z) := \frac{z - z_1}{1 - \bar{z}_1 z} \cdot \frac{z - z_2}{1 - \bar{z}_2 z} \cdot \frac{z - \omega}{1 - \bar{\omega} z} \in H_{\text{node}}^\infty.$$

- (3) For every $(\alpha, \beta) \in A$, the matrix

$$[k^{\alpha, \beta}(z_i, z_j)]_{i, j=1}^4$$

has a column (equivalently, row) with no zeroes.

- (4) For every $(\alpha, \beta) \neq (\alpha', \beta') \in A$ there exists a pair $z_i \neq z_j$ such that:

$$(3.1) \quad \left| \frac{k^{\alpha, \beta}(z_j, z_i)}{\sqrt{k^{\alpha, \beta}(z_i, z_i) \cdot k^{\alpha, \beta}(z_j, z_j)}} \right|^2 \neq \left| \frac{k^{\alpha', \beta'}(z_j, z_i)}{\sqrt{k^{\alpha', \beta'}(z_i, z_i) \cdot k^{\alpha', \beta'}(z_j, z_j)}} \right|^2$$

- (5) For every $(\alpha, \beta) \in A$,

$$\left\{ \begin{array}{l} k_{z_\ell} \notin \text{span}\{k_{z_3}^{\alpha, \beta}, k_{z_4}^{\alpha, \beta}, P_{\mathcal{M}_{\alpha, \beta}} k_\zeta^{\alpha, \beta}\} \quad \ell \in \{1, 2\} \\ k_{z_\ell} \notin \text{span}\{k_{z_1}^{\alpha, \beta}, k_{z_2}^{\alpha, \beta}, P_{\mathcal{M}_{\alpha, \beta}} k_\omega^{\alpha, \beta}\} \quad \ell \in \{3, 4\} \end{array} \right\}, \left\{ \begin{array}{l} f_\ell \notin \text{span}\{k_{z_1}^{\alpha, \beta}, k_{z_2}^{\alpha, \beta}, P_{\mathcal{M}_{\alpha, \beta}} k_\omega^{\alpha, \beta}\} \quad \ell \in \{1, 2\} \\ f_\ell \notin \text{span}\{k_{z_3}^{\alpha, \beta}, k_{z_4}^{\alpha, \beta}, P_{\mathcal{M}_{\alpha, \beta}} k_\zeta^{\alpha, \beta}\} \quad \ell \in \{3, 4\} \end{array} \right\},$$

where ζ is some point satisfying $B_{z_3, z_4, \zeta} \in H_{\text{node}}^\infty$ that exists by Lemma 3.9, and $\{f_1, \dots, f_4\} \subset \mathcal{M}_{\alpha, \beta}$ is the dual basis of $B = \{k_{z_i}^{\alpha, \beta} : 1 \leq i \leq 4\}$ (i.e. $f_i(z_j) = \delta_{i, j}$).

Theorem 3.11. Let $z_1, \dots, z_4 \in \mathbb{D} \setminus \{0, \lambda\}$ be good points and let $I \subset H_{\text{node}}^\infty$ be the ideal of functions that vanish on z_1, \dots, z_4 . Then, the representations $\sigma_{\alpha, \beta} : C_{\text{max}}^*(H_{\text{node}}^\infty/I) \rightarrow B(\mathcal{M}_{\alpha, \beta})$ that extend the representations $\rho_{\alpha, \beta}$ for $(\alpha, \beta) \in A$ form an infinite family of unitarily inequivalent boundary representations. In particular, $C_e^*(H_{\text{node}}^\infty/I)$ is infinite-dimensional.

We break the proof of Theorem 3.11 into pieces. First we show that for every $(\alpha, \beta), (\alpha', \beta') \in A$, the representations $\rho_{\alpha, \beta}$ and $\rho_{\alpha', \beta'}$ are unitarily inequivalent and irreducible. We then proceed to show that they are boundary representations for H_{node}^∞ .

Proposition 3.12. The family of representations $(\sigma_{\alpha, \beta})_{(\alpha, \beta) \in A}$ of $C_{\text{max}}^*(H_{\text{node}}^\infty/I)$ is a family of irreducible non-unitarily equivalent representations.

Proof. Fix $(\alpha, \beta) \in A$. Recall that in the scalar case, a basis for $\mathcal{M}_{\alpha, \beta}$ is

$$B = \{k_{z_1}^{\alpha, \beta}, k_{z_2}^{\alpha, \beta}, k_{z_3}^{\alpha, \beta}, k_{z_4}^{\alpha, \beta}\}.$$

For every $f \in H_{\text{node}}^\infty$, we saw that $M_f^* k_{z_i}^{\alpha, \beta} = \overline{f(z_i)} k_{z_i}^{\alpha, \beta}$. Therefore, M_f^* is diagonal with respect to B . Let G be the Gram matrix of B . Namely,

$$(G)_{i, j} = \langle k_{z_j}^{\alpha, \beta}, k_{z_i}^{\alpha, \beta} \rangle = k^{\alpha, \beta}(z_i, z_j).$$

Then

$$[P_{\mathcal{M}_{\alpha,\beta}} M_f |_{\mathcal{M}_{\alpha,\beta}}]_C = ([M_f^* |_{\mathcal{M}_{\alpha,\beta}}]_C)^* = G^{-1/2} \begin{pmatrix} f(z_1) & 0 & 0 & 0 \\ 0 & f(z_2) & 0 & 0 \\ 0 & 0 & f(z_3) & 0 \\ 0 & 0 & 0 & f(z_4) \end{pmatrix} G^{1/2} = G^{-\frac{1}{2}} D_f G^{\frac{1}{2}},$$

where C is the orthonormal basis $\left\{ \sum_{j=1}^4 (G^{-\frac{1}{2}})_{j,i} k_{z_j}^{\alpha,\beta} : 1 \leq i \leq 4 \right\}$.

Since $B(\mathcal{M}_{\alpha,\beta}) \cong M_4(\mathbb{C})$ as C^* -algebras via $T \rightarrow [T]_C$, it is enough to show that:

$$C^*(\rho_{\alpha,\beta}(H_{\text{node}}^\infty/I)) \cong C^*\left(\underbrace{\{[P_{\mathcal{M}_{\alpha,\beta}} M_f |_{\mathcal{M}_{\alpha,\beta}}]_C : f \in H_{\text{node}}^\infty\}}_{\mathcal{A}}\right) = M_4(\mathbb{C})$$

To see that, for $1 \leq i \leq 4$ let $f_i \in H_{\text{node}}^\infty$ be such that $f_i(z_j) = \delta_{i,j}$ and denote:

$$A_i = P_{\mathcal{M}_{\alpha,\beta}} M_{f_i} |_{\mathcal{M}_{\alpha,\beta}} = G^{-\frac{1}{2}} E_{ii} G^{\frac{1}{2}} \in C^*(\rho_{\alpha,\beta}(H_{\text{node}}^\infty/I)).$$

Then,

$$\sum_{i=1}^4 \frac{1}{k^{\alpha,\beta}(z_i, z_i)} A_i A_i^* = G^{-1} \in C^*(\mathcal{A})$$

As C^* algebras are closed under square-roots

$$G^{1/2} \in C^*(\mathcal{A})$$

Moreover, note that:

$$A_i A_j^* = G^{-\frac{1}{2}} D_{f_i} G D_{f_j} G^{-\frac{1}{2}} = k^{\alpha,\beta}(z_i, z_j) G^{-\frac{1}{2}} E_{i,j} G^{-\frac{1}{2}}$$

By (3) in Definition 3.10, we can assume without loss of generality that

$$k^{\alpha,\beta}(z_1, z_2), k^{\alpha,\beta}(z_1, z_3), k^{\alpha,\beta}(z_1, z_4) \neq 0$$

Hence:

$$\frac{1}{k^{\alpha,\beta}(z_i, z_j)} A_i A_j^* = G^{1/2} G^{-1/2} E_{i,j} G^{-1/2} G^{1/2} = E_{i,j} \in C^*(\rho_{\alpha,\beta}(H_{\text{node}}^\infty/I))$$

for $1 \leq i = j \leq 4$ and $i = 1, j = 2, 3, 4$. Consequently:

$$E_{2,3} = E_{2,1} E_{1,3}, E_{2,4} = E_{2,1} E_{1,4}, E_{3,4} = E_{3,1} E_{1,4} \in C^*(\rho_{\alpha,\beta}(H_{\text{node}}^\infty/I))$$

i.e. $\{E_{i,j} : 1 \leq i, j \leq 4\} \subset C^*(\rho_{\alpha,\beta}(H_{\text{node}}^\infty/I))$, concluding the proof of irreducibility. For the second part, recall that $\sigma_{\alpha,\beta}$ and $\sigma_{\alpha',\beta'}$ are unitarily equivalent if there exists a unitary $U : \mathcal{M}_{\alpha,\beta} \rightarrow \mathcal{M}_{\alpha',\beta'}$, such that for every $f + I \in H_{\text{node}}^\infty/I$:

$$U \rho_{\alpha,\beta}(f + I)^* = \rho_{\alpha',\beta'}(f + I)^* U$$

i.e, for all $f + I \in H_{\text{node}}^\infty/I$:

$$U M_f^* |_{\mathcal{M}_{\alpha,\beta}} = M_f^* |_{\mathcal{M}_{\alpha',\beta'}} U$$

In particular, if f_i is the polynomial satisfying $f_i(z_j) = \delta_{i,j}$, then for every $1 \leq i \leq 4$:

$$f_i(z_i) U k_{z_i}^{\alpha,\beta} = U k_{z_i}^{\alpha,\beta} = U M_{f_i}^* (k_{z_i}^{\alpha,\beta}) = M_{f_i}^* (U k_{z_i}^{\alpha,\beta})$$

Therefore $U k_{z_i}^{\alpha,\beta} \in \text{span} \left\{ k_{z_i}^{\alpha',\beta'} \right\}$ (one dimensional eigenspace), i.e $U k_{z_i}^{\alpha,\beta} = \gamma_i k_{z_i}^{\alpha',\beta'}$. Moreover, as U is unitary,

$$\|k_{z_i}^{\alpha,\beta}\| = |\gamma_i| \|k_{z_i}^{\alpha',\beta'}\|$$

and

$$\left\langle k_{z_i}^{\alpha,\beta}, k_{z_j}^{\alpha,\beta} \right\rangle = \gamma_i \overline{\gamma_j} \left\langle k_{z_i}^{\alpha',\beta'}, k_{z_j}^{\alpha',\beta'} \right\rangle.$$

Taking the absolute value on both sides:

$$\left| \left\langle k_{z_i}^{\alpha,\beta}, k_{z_j}^{\alpha,\beta} \right\rangle \right| = \frac{\|k_{z_i}^{\alpha,\beta}\|}{\|k_{z_i}^{\alpha',\beta'}\|} \frac{\|k_{z_j}^{\alpha,\beta}\|}{\|k_{z_j}^{\alpha',\beta'}\|} \left| \left\langle k_{z_i}^{\alpha',\beta'}, k_{z_j}^{\alpha',\beta'} \right\rangle \right|.$$

Equivalently,

$$(3.2) \quad \frac{|k^{\alpha,\beta}(z_j, z_i)|^2}{k^{\alpha,\beta}(z_i, z_i) \cdot k^{\alpha,\beta}(z_j, z_j)} = \frac{|k^{\alpha',\beta'}(z_j, z_i)|^2}{k^{\alpha',\beta'}(z_i, z_i) \cdot k^{\alpha',\beta'}(z_j, z_j)}$$

for every z_i, z_j , contradicting (4) in Definition 3.10. \square

Let $\{f_1, \dots, f_4\} \subset \mathcal{M}_{\alpha,\beta}$ be the dual basis of $\{k_{z_1}^{\alpha,\beta}, \dots, k_{z_4}^{\alpha,\beta}\}$ (i.e. $f_i(z_j) = \delta_{i,j}$). We will show that $\rho_{\alpha,\beta}$ are dilation maximal. Our strategy is to prove that every $(\alpha, \beta) \in A$, $\rho_{\alpha,\beta}$ has no non-trivial extensions and no non-trivial coextensions. This condition is known to be equivalent to dilation maximality. This approach is an adaptation of the approach taken by McCullough [27]. Due to the existence of the generator $h + I \in H_{\text{node}}^\infty/I$, for every representation, an invariant/coinvariant/reducing subspace for the image of $h + I$ is such for the image of the entire algebra. This phenomenon distinguishes this case from the case when the set of interpolation nodes contains the constraints.

Theorem 3.13. *For every $(\alpha, \beta) \in A$, $\rho_{\alpha,\beta}$ is co-extension maximal.*

Fix the permutation $\tau = (14)(23) \in S_4$. We begin by proving two propositions:

Proposition 3.14. *For every pair $(\alpha, \beta) \in A$, if $f \in \text{span}\{k_{z_\ell}^{\alpha,\beta}\} \subset \mathcal{M}_{\alpha,\beta}$ satisfies*

$$(3.3) \quad P_{\mathcal{M}_{\alpha,\beta}}(1 - M_\varphi M_\varphi^*)|_{\mathcal{M}_{\alpha,\beta}} \geq f f^*$$

for every Blaschke product $\varphi \in H_{\text{node}}^\infty$ vanishing on $z_{\tau(\ell)}$, then $f = 0$.

Proof. Let $\varphi \in H_{\text{node}}^\infty$ be a Blaschke product with zeroes $x_1, \dots, x_n \subset \mathbb{D} \setminus \{0, \lambda\}$. Since M_φ is an isometry on H^2 and $H_{\alpha,\beta}^2$ is invariant, M_φ is an isometry on $H_{\alpha,\beta}^2$. Moreover, clearly $K_\varphi = \text{span}\{k_{x_1}^{\alpha,\beta}, \dots, k_{x_n}^{\alpha,\beta}\} \subset H_{\alpha,\beta}^2 \ominus (\varphi H_{\alpha,\beta}^2)$. On the other hand, if $g \in H_{\alpha,\beta}^2 \ominus K_\varphi$, then g vanishes on the zeroes of φ . In particular, there exists $h \in H^2$, such that $\varphi h = g$. By our assumption, there exists $c, d \in \mathbb{C}$, such that $d \neq 0$, $\varphi(0) = \varphi(\lambda) = d$, $g(0) = c\alpha$, and $g(\lambda) = c(\alpha + f_\lambda(\lambda)\beta)$. Hence, $h \in H_{\alpha,\beta}^2$. Conclude that $K_\varphi = H_{\alpha,\beta}^2 \ominus (\varphi H_{\alpha,\beta}^2)$. Let $P_\varphi = 1 - M_\varphi M_\varphi^* \in B(H_{\alpha,\beta}^2)$ be the projection onto K_φ . Hence,

$$(3.4) \quad \text{Im}(P_{\mathcal{M}_{\alpha,\beta}} P_\varphi|_{\mathcal{M}_{\alpha,\beta}}) \subseteq P_{\mathcal{M}_{\alpha,\beta}} K_\varphi.$$

Let:

$$\varphi(z) = \begin{cases} B_{z_3, z_4, \zeta} & \ell \in \{1, 2\} \\ B_{z_1, z_2, \omega} & \ell \in \{3, 4\} \end{cases}.$$

By our assumptions $\varphi \in H_{\text{node}}^\infty$. Assume $\ell \in \{1, 2\}$. In that case, $\tau(\ell) \in \{3, 4\}$. By (3.4):

$$\text{Im}(1 - P_{\mathcal{M}_{\alpha,\beta}} M_\varphi M_\varphi^*|_{\mathcal{M}_{\alpha,\beta}}) \subset \text{span}\{k_{z_3}^{\alpha,\beta}, k_{z_4}^{\alpha,\beta}, P_{\mathcal{M}_{\alpha,\beta}} k_\zeta^{\alpha,\beta}\}$$

By (3.3), $f \in \text{span}\{k_{z_3}^{\alpha,\beta}, k_{z_4}^{\alpha,\beta}, P_{\mathcal{M}_{\alpha,\beta}} k_\zeta^{\alpha,\beta}\}$, hence by the assumption on f , either $f = 0$ or:

$$k_{z_\ell}^{\alpha,\beta} \in \text{span}\{k_{z_3}^{\alpha,\beta}, k_{z_4}^{\alpha,\beta}, P_{\mathcal{M}_{\alpha,\beta}} k_\zeta^{\alpha,\beta}\}$$

which contradicts (5) in Definition 3.10. Hence $f = 0$ for all $(\alpha, \beta) \in A$. The case of $\ell \in \{3, 4\}$ is similar. \square

Proposition 3.15. *Let $(\alpha, \beta) \in A$ and let $\psi_{\alpha,\beta}: H_{\text{node}}^\infty/I \rightarrow B(\mathcal{M}_{\alpha,\beta} \oplus \mathbb{C})$ be a coextension of $\rho_{\alpha,\beta}$. Write*

$$\psi_{\alpha,\beta}(h + I)^* = \begin{pmatrix} \rho_{\alpha,\beta}(h + I)^* & v \\ 0 & z_\ell \end{pmatrix}.$$

Then $v = 0$.

Proof. By Lemma 3.1, $\psi_{\alpha,\beta}(h + I)$ is diagonalizable, where $h + I$ is the analog of the coordinate function in the quotient. Taking $p(z) = \prod_{i=1}^4 (z - z_i)(z - \lambda)$ and abbreviating $A = \rho_{\alpha,\beta}(h + I)$, we have:

$$0 = p(\sigma_{\alpha,\beta}(h + I)^*) = \begin{pmatrix} 0 & \tilde{p}(A)v \\ 0 & 0 \end{pmatrix}$$

with $\tilde{p}(z) = \prod_{i \neq \ell}^4 (z - z_i)z(z - \lambda)$. Since $p(A) = 0$ and $\tilde{p}(A)v = 0$, by Remark 3.3 there exists some $f \in M_{\alpha, \beta}$ such that $(A^* - z_\ell I)f = v$. Note that if $f = \sum_{i=1}^4 \gamma_i k_{z_i}^{\alpha, \beta}$, then:

$$(A^* - z_\ell I) f^{g(z) = \sum_{i \neq \ell}^4 \gamma_i k_{z_i}^{\alpha, \beta}} (A^* - z_\ell I) \left(\sum_{i \neq \ell}^4 \gamma_i k_{z_i}^{\alpha, \beta} \right) = (A^* - z_\ell I) g$$

So we can assume $f \in \text{span}\{k_{z_i}^{\alpha, \beta} : 1 \leq i \neq \ell \leq 4\}$. For every Blaschke product $\varphi \in H_{\text{node}}^\infty$:

$$\varphi(\psi_{\alpha, \beta}(h + I))^* = \begin{pmatrix} \varphi(A)^* & (\varphi(A)^* - \overline{\varphi(z_\ell)})f \\ 0 & \varphi(z_\ell) \end{pmatrix}$$

Moreover, if $\varphi(\overline{z_\ell}) = \varphi(z_\ell) = 0$ (as $z_\ell \in \mathbb{R}$), then:

$$\varphi(\sigma_{\alpha, \beta}(h + I))^* = \begin{pmatrix} \varphi(A)^* & \varphi(A)^* f \\ 0 & 0 \end{pmatrix}.$$

Since $\varphi(\psi_{\alpha, \beta}(h + I)) = \psi_{\alpha, \beta}(\varphi)$,

$$1 - \varphi(A)^* \varphi(A) \geq 0$$

Doing the calculations, we see that:

$$(3.5) \quad \begin{pmatrix} 1 - \varphi(A)^*(I + ff^*)\varphi(A) & 0 \\ 0 & 1 \end{pmatrix} \geq 0.$$

Letting Q be the positive root of $I + ff^*$, $Q^{-2} = I - \frac{1}{1 + \|f\|^2} ff^*$, and by (3.5) $\varphi(A)^* Q$ is a contraction. Therefore:

$$Q^{-2} \geq \varphi(A)\varphi(A)^* \Rightarrow I - \varphi(A)\varphi(A)^* \geq \frac{1}{1 + \|f\|^2} ff^*.$$

This can be rewritten as, by definition of A^* :

$$I - P_{\mathcal{M}_{\alpha, \beta}} M_{\hat{\varphi}} M_{\hat{\varphi}}^* |_{\mathcal{M}_{\alpha, \beta}} \geq \frac{1}{1 + \|f\|^2} ff^*.$$

Letting $\varphi(z) = \frac{z - z_\ell}{1 - \overline{z_\ell} z} \cdot \frac{z - z_{\tau(\ell)}}{1 - \overline{z_{\tau(\ell)}} z} \in H_{\text{node}}^\infty$, by Proposition 3.14,

$$\sqrt{\frac{1}{1 + \|f\|^2}} f \in \text{span}\{k_{z_\ell}^{\alpha, \beta}, k_{z_{\tau(\ell)}}^{\alpha, \beta}\}.$$

Hence, by our assumption, $f \in \mathbb{C}k_{z_{\tau(\ell)}}$. Now the second part of Proposition 3.14, implies $f = 0$. Therefore, for every $(\alpha, \beta) \in A$, $v = 0$. \square

Now, we can prove that $\rho_{\alpha, \beta}$ has no non-trivial co-extensions.

Proof of Theorem 3.13. Assume that $\psi : H_{\text{node}}^\infty / I \rightarrow B(\mathcal{M}_{\alpha, \beta} \oplus \mathcal{H})$ is a co-extension of $\rho_{\alpha, \beta}$. Write

$$\psi_{\alpha, \beta}(h + I)^* = \begin{pmatrix} \rho_{\alpha, \beta}(h + I)^* & D \\ 0 & G \end{pmatrix}.$$

Let $m(x) = \prod_{k=1}^4 (z - z_k)$. As $m(\psi^*(h + I)) = \psi^*(m(h + I)) = 0$, $m(G) = 0$, that is $(G - z_1 I) \cdots (G - z_4 I) = 0$. Therefore, there is a non-orthogonal direct sum decomposition

$$\mathcal{H} = \sum_{i=1}^4 \ker(G - z_i I).$$

Assume that $D \neq 0$. Hence, there exists a vector $0 \neq v \in \ker(G - z_i)$ for some i such that $e = Dv \neq 0$. Consider $\mathcal{K} = \mathcal{M}_{\alpha, \beta} \oplus \text{span}\{v\} \subset \mathcal{M}_{\alpha, \beta} \oplus \mathcal{H}$. Then:

$$\psi(h + I)^* |_{\mathcal{K}} = \begin{pmatrix} A^* & e \\ 0 & z_i \end{pmatrix}$$

By Proposition 3.15, $e = 0$ and hence $D = 0$. \square

Theorem 3.16. For every $(\alpha, \beta) \in A$, $\rho_{\alpha, \beta}$ is extension maximal.

We start with a proposition similar to Proposition 3.15.

Proposition 3.17. *Let $(\alpha, \beta) \in A$ and let $\psi_{\alpha, \beta}: H_{\text{node}}^\infty/I \rightarrow B(\mathcal{M}_{\alpha, \beta} \oplus \mathbb{C})$ be an extension of $\rho_{\alpha, \beta}$. Write*

$$\psi_{\alpha, \beta}(h + I) = \begin{pmatrix} \rho_{\alpha, \beta}(h + I) & v \\ 0 & z_\ell \end{pmatrix}.$$

Then $v = 0$.

Proof. By the same reasoning as in the proof of Proposition 3.15, $v = (A - z_\ell I)f$ for some $f \in \mathcal{M}_{\alpha, \beta}$. Write $f = \sum_{i=1}^4 \gamma_i f_i$ with f_i being the function satisfying $f_i(z_j) = \delta_{i,j}$, and observe that:

$$(A - z_\ell I)f = (A - z_\ell I) \left(\sum_{i=1}^4 \gamma_i f_i \right) = (A - z_\ell I) \left(\sum_{i \neq \ell}^4 \gamma_i f_i \right).$$

since $f_\ell \in \ker(A - z_\ell I)$. Therefore we can assume that $f \in \text{span}\{f_i : 1 \leq i \neq \ell \leq 4\}$. Let

$$\varrho(z) = \frac{z - z_\ell}{1 - z_\ell z} \cdot \frac{z - z_{\tau(\ell)}}{1 - z_{\tau(\ell)} z}.$$

By Lemma 3.1:

$$\sigma_{\alpha, \beta}(\varrho + I) = \varrho(\sigma_{\alpha, \beta}(h + I)) = \begin{pmatrix} \varrho(A) & \varrho(A)f \\ 0 & 0 \end{pmatrix}.$$

As $\sigma_{\alpha, \beta}$ is a representation and ϱ is a Blaschke product and hence a contraction,

$$1 \geq \begin{pmatrix} \varrho(A)\varrho(A)^* + \varrho(A)ff^*\varrho(A)^* & 0 \\ 0 & 1 \end{pmatrix}.$$

In particular,

$$1 - P_{\mathcal{M}_{\alpha, \beta}} M_\varrho M_\varrho^* |_{\mathcal{M}_{\alpha, \beta}} \geq \varrho(A)ff^*\varrho(A)^*$$

Hence

$$\varrho(A)f = \sum_{\substack{i=1 \\ i \neq \ell, \neq \tau(\ell)}}^4 \gamma_i \varrho(z_i) f_i \in \text{Im}(1 - P_{\mathcal{M}_{\alpha, \beta}} M_\varrho M_\varrho^* |_{\mathcal{M}_{\alpha, \beta}}) \subset \text{span}\{k_{z_\ell}^{\alpha, \beta}, k_{z_{\tau(\ell)}}^{\alpha, \beta}\}$$

Note that for $i \neq j$, $\langle f_i, k_{z_j}^{\alpha, \beta} \rangle = 0$. Therefore

$$\text{span}\{f_i : 1 \leq i \leq 4, i \neq \ell, \tau(\ell)\} \cap \text{span}\{k_{z_\ell}^{\alpha, \beta}, k_{z_{\tau(\ell)}}^{\alpha, \beta}\} = \{0\}$$

hence $f \in \text{span}\{f_{\tau(\ell)}\}$, and we can write $f = \gamma \cdot f_{\tau(\ell)}$. Now let:

$$\tilde{\varphi}(z) = \begin{cases} B_{z_1, z_2, \omega} & \ell \in \{1, 2\} \\ B_{z_3, z_4, \zeta} & \ell \in \{3, 4\} \end{cases}$$

and assume again that $\ell \in \{1, 2\}$. By the same reasoning:

$$\tilde{\varphi}(A)f = \gamma \cdot \tilde{\varphi}(z_{\tau(\ell)}) f_{\tau(\ell)} \in \text{span}\{k_1^{\alpha, \beta}, k_{z_2}^{\alpha, \beta}, P_{\mathcal{M}_{\alpha, \beta}} k_\omega^{\alpha, \beta}\}.$$

By our choice of $\tilde{\varphi}$, $\tilde{\varphi}(z_{\tau(\ell)}) \neq 0$. Assume that $\gamma \neq 0$. Therefore $\tilde{\varphi}(A)f \neq 0$ and consequently:

$$f_{\tau(\ell)} \in \text{span}\{k_{z_1}^{\alpha, \beta}, k_{z_2}^{\alpha, \beta}, P_{\mathcal{M}_{\alpha, \beta}} k_\omega^{\alpha, \beta}\}.$$

contradiction Definition 3.10. Hence for all $(\alpha, \beta) \in A$, $f = 0$ and consequently $v = 0$. The case where $\ell \in \{3, 4\}$ is similar. \square

Proof of Theorem 3.16. Assume that $\psi : H_{\text{node}}^\infty/I \rightarrow B(\mathcal{M}_{\alpha, \beta} \oplus \mathcal{H})$ is an extension of $\rho_{\alpha, \beta}$. Write

$$\psi(h + I) = \begin{pmatrix} \rho_{\alpha, \beta}(h + I) & D \\ 0 & G \end{pmatrix}.$$

Let $m(x) = \prod_{k=1}^4 (z - z_k)$. As $m(\psi(h + I)) = \psi(m(h) + I) = 0$, $m(G) = 0$, that is $(G - z_1 I) \cdots (G - z_4 I) = 0$. Therefore, we have a non-orthogonal direct sum decomposition

$$\mathcal{H} = \sum_{i=1}^4 \ker(G - z_i I).$$

Assume that $D \neq 0$. Hence, there exists a vector $0 \neq v \in \ker(G - z_i I)$ for some i such that $e = Dv \neq 0$. Consider $\mathcal{K} = \mathcal{M}_{\alpha,\beta} \oplus \text{span}\{v\} \subset \mathcal{M}_{\alpha,\beta} \oplus \mathcal{H}$. By assumption, \mathcal{K} is invariant. Therefore:

$$\sigma_{\alpha,\beta}(h + I)|_{\mathcal{K}} = \begin{pmatrix} A & e \\ 0 & z_i \end{pmatrix}$$

By Proposition 3.17, $e = 0$. Hence, $D = 0$. □

For completeness we include the proof that for every $(\alpha, \beta) \in A$, $\rho_{\alpha,\beta}$ is dilation maximal.

Proof of Theorem 3.11. Let $(\alpha, \beta) \in A$. Assume that:

$$\psi : H_{\text{node}}^\infty / I \rightarrow B(\mathcal{K})$$

is a dilation of $\rho_{\alpha,\beta}$, i.e., $\mathcal{M}_{\alpha,\beta} \subset \mathcal{K}$ and

$$\rho_{\alpha,\beta} = P_{\mathcal{M}_{\alpha,\beta}} \psi|_{\mathcal{M}_{\alpha,\beta}}$$

Then, again by Sarason's Lemma

$$\psi(h + I) = \begin{pmatrix} A & B & C \\ 0 & \rho_{\alpha,\beta}(h + I) & D \\ 0 & 0 & E \end{pmatrix}.$$

As the 2×2 upper-left corner is a co-extension of $\rho_{\alpha,\beta}$, by Theorem 3.13. Similarly, since the 2×2 lower-right corner is an extension of $\rho_{\alpha,\beta}$, by Theorem 3.16, $D = 0$. Therefore,

$$\psi(h + I) = \begin{pmatrix} A & 0 & C \\ 0 & \rho_{\alpha,\beta} & 0 \\ 0 & 0 & E \end{pmatrix}$$

that is, $\mathcal{M}_{\alpha,\beta}$ is reducing for $\phi(h + I)$ and, thus, for all of $\psi(H_{\text{node}}^\infty / I)$, as desired. □

Now it remains to show that there are sets of good points for some $0 \neq \lambda \in \mathbb{D} \cap \mathbb{R}$. We will dedicate the rest of this section to an example of a good set of points. Fix $\lambda = \frac{1}{\sqrt{2}}$. We will show that for the choice of points:

$$z_1 = \frac{4}{3\sqrt{2}}, z_2 = \frac{1}{2\sqrt{2}}, z_3 = \frac{\sqrt{2}}{3}, z_4 = -\frac{1}{\sqrt{2}}$$

$C_e^*(H_{\text{node}}^\infty / I)$ is infinite-dimensional. By what we observed, it is enough to show that these points are good with respect to λ . Our goal is to prove the following theorem.

Theorem 3.18. z_1, \dots, z_4 are good points.

Due to the fact that the proof is quite computational, we again break it into pieces.

Proposition 3.19. z_1, \dots, z_4 satisfy conditions (1) – (4).

Proof. Under this setup:

$$B_\lambda(z) = \frac{\sqrt{2}z(\sqrt{2}z - 1)}{2 - \sqrt{2}z}, f_\lambda(z) = \frac{z}{\sqrt{2} - z}$$

Recall that for fixed $(\alpha, \beta) \in \mathbb{G}(1 \times 1)$, the kernel function $k^{\alpha,\beta}$ is given by

$$k^{\alpha,\beta}(z, w) = \overline{(\alpha + \beta f_\lambda(w))}(\alpha + \beta f_\lambda(z)) + \frac{B_\lambda(z) \cdot \overline{B_\lambda(w)}}{1 - z\bar{w}}.$$

To make calculations more manageable, we added two tables of values that will be useful to us, while using $|\alpha|^2 + |\beta|^2 = 1$ and that $\alpha, \beta \in \mathbb{R}$.

z	$B_\lambda(z)$	$f_\lambda(z)$
z_1	$\frac{2}{3}$	2
z_2	$-\frac{1}{6}$	$\frac{1}{3}$
z_3	$-\frac{1}{6}$	$\frac{1}{2}$
z_4	$\frac{2}{3}$	$-\frac{1}{3}$

TABLE 1. Values of $B_\lambda(z)$ and $f_\lambda(z)$ for different z .

(z, w)	$k^{\alpha, \beta}(z, w)$
(z_1, z_1)	$(\alpha + 2\beta)^2 + 4$
(z_1, z_2)	$\frac{1}{3}\alpha^2 + \frac{7}{3}\alpha\beta + \frac{1}{2}$
(z_1, z_3)	$\frac{5}{2}\alpha\beta + \frac{4}{5}$
(z_1, z_4)	$\frac{5}{3}\alpha^2 + \frac{5}{3}\alpha\beta - \frac{2}{5}$
(z_2, z_2)	$(\alpha + \frac{1}{3}\beta)^2 + \frac{2}{63}$
(z_2, z_3)	$\frac{5}{6}\alpha^2 + \frac{5}{6}\alpha\beta + \frac{1}{5}$
(z_2, z_4)	$\frac{10}{9}\alpha^2 - \frac{1}{5}$
(z_3, z_3)	$(\alpha + \frac{1}{2}\beta)^2 + \frac{1}{28}$
(z_3, z_4)	$\frac{7}{6}\alpha^2 + \frac{1}{6}\alpha\beta - \frac{1}{4}$
(z_4, z_4)	$(\alpha - \frac{1}{3}\beta)^2 + \frac{8}{9}$

TABLE 2. Values of $k^{\alpha, \beta}(z, w)$ for different pairs (z, w) .

In particular, we see that condition (1) of Definition 3.10 is satisfied. Secondly, it is straight-forward to check that:

$$B_{z_1, z_2, \sqrt{2}-1}, B_{z_3, z_4, \sqrt{2}-1} \in H_{\text{node}}^\infty.$$

That gives us condition (2). For condition (3), consider the polynomial $f(\alpha, \beta) = \alpha^2 + \beta^2 - 1$.

Clearly, the real zeroes of f are the circle. We will show that the first column/row is non-zero except for a finite subset of (α, β) in the circle. To see this, consider additionally $g_2(\alpha, \beta) = \alpha^2 + 7\alpha\beta + \frac{3}{2}$, $g_3(\alpha, \beta) = 5\alpha\beta + \frac{8}{5}$, and $g_4(\alpha, \beta) = 5\alpha^2 + 5\alpha\beta - \frac{6}{5}$. Since the vanishing of g_i is equivalent to the vanishing of $k^{\alpha, \beta}(z_1, z_i)$, we get that the set of points (α, β) , where the first column has zeroes in it is contained in

$$D = \bigcup_{i=2}^4 Z(f) \cap Z(g_i).$$

Here, $Z(f)$ stands for the zero locus of f and similarly for the g_i . Since these are algebraic planar curves each of degree 2, that do not have components in common, each of the intersection sets has at most 4 real points. Therefore, D is finite. Therefore, on the complement of D , condition (3) of Definition 3.10 holds.

As for condition (4), we first note that if $k^{\alpha, \beta} = k^{-\alpha, \beta}$. More generally, if we multiply (α, β) by a unimodular scalar, we do not change the kernel. Hence, we are considering, in fact, the real projective line. However, we will restrict our selves to a half-circle. Let $A' \subset \mathbb{T}$ be the upper open half-circle. We will show that for every $(\alpha, \beta) \neq (\alpha', \beta') \in A'$, either (3.1) is satisfied for (z_1, z_2) or for (z_2, z_3) . To this end, fix $(\alpha, \beta) \neq (\alpha', \beta') \in A'$ and let $\frac{\beta}{\alpha} = x$. By $\alpha^2 + \beta^2 = 1$:

$$x^2 = \left(\frac{\beta}{\alpha}\right)^2 = \frac{1 - \alpha^2}{\alpha^2} \Rightarrow \alpha^2 = \frac{1}{1 + x^2}$$

We can rewrite the kernels in terms of x :

$$\begin{aligned} k^{\alpha, \beta}(z_1, z_1) &= (\alpha + 2\beta)^2 + 4 = |\alpha|^2 \left(1 + 2\frac{\beta}{\alpha}\right) \left(1 + 2\frac{\beta}{\alpha}\right) + 4 = \\ &= \frac{1}{1 + x^2} (1 + 2x)^2 + 4 = \frac{8x^2 + 4x + 5}{1 + x^2} \end{aligned}$$

And similarly

$$k^{\alpha, \beta}(z_2, z_2) = \frac{1}{1 + x^2} \left(1 + \frac{x}{3}\right)^2 + \frac{2}{63} = \frac{9x^2 + 42x + 65}{63(1 + x^2)}$$

$$k^{\alpha, \beta}(z_3, z_3) = \frac{1}{1 + x^2} \left(1 + \frac{x}{2}\right)^2 + \frac{1}{28} = \frac{8x^2 + 28x + 29}{28(1 + x^2)}$$

$$k^{\alpha,\beta}(z_1, z_2) = \frac{1}{1+x^2} \left(1 + \frac{x}{3}\right) (1+2x) - \frac{1}{6} = \frac{3x^2 + 14x + 5}{6(1+x^2)}$$

$$k^{\alpha,\beta}(z_2, z_3) = \frac{1}{1+x^2} \left(1 + \frac{x}{2}\right) \left(1 + \frac{x}{3}\right) + \frac{1}{30} = \frac{6x^2 + 25x + 31}{30(1+x^2)}$$

By letting $y = \frac{\beta'}{\alpha'}$ and replacing x with y in the calculations above, we can write $k^{\alpha',\beta'}(z_i, z_j)$ in terms of y . By taking $j = 1, i = 2$, equation 3.1 becomes

$$(3.6) \quad \frac{\left(\frac{(3x^2+14x+5)}{6(1+x^2)}\right)^2}{\frac{(8x^2+4x+5)}{1+x^2} \cdot \frac{(9x^2+42x+65)}{1+x^2}} = \frac{\left(\frac{(3y^2+14y+5)}{6(1+y^2)}\right)^2}{\frac{(8y^2+4y+5)}{1+y^2} \cdot \frac{(9y^2+42y+65)}{1+y^2}}$$

Simplifying and canceling some terms on both sides, we obtain the equation

$$(3x^2 + 14x + 5)^2(8y^2 + 4y + 5)(9y^2 + 42y + 65) - (3y^2 + 14y + 5)^2(8x^2 + 4x + 5)(9x^2 + 42x + 65) = 0$$

As we are looking for solutions with $x \neq y$, and as $x = y$ is a solution, by dividing by $x - y$, refactoring, and dividing both sides by the constants, the equations become

$$(3.7) \quad (xy + x + y + 3) (12x^2y^2 + 31x^2y + 31xy^2 - 5x^2 + 28xy - 5y^2 - 65x - 65y - 50) = 0$$

We do the same analysis for $j = 2, i = 3$. For this choice of indices, 5.1 becomes

$$(3.8) \quad \frac{\left(\frac{(6x^2+25x+31)}{30(1+x^2)}\right)^2}{\frac{(9x^2+42x+65)}{1+x^2} \cdot \frac{(8x^2+28x+29)}{63(1+x^2)}} = \frac{\left(\frac{(6y^2+25y+31)}{30(1+y^2)}\right)^2}{\frac{(9y^2+42y+65)}{1+y^2} \cdot \frac{(8y^2+28y+29)}{63(1+y^2)}}$$

Again, canceling some terms on both sides, dividing by $x - y$, refactoring, and dividing both sides by the relevant constants, (3.8) becomes

$$(xy + x + y - 1) (12x^2y^2 + 25x^2y + 25xy^2 + 37x^2 + 37y^2 + 25x + 25y + 62) = 0$$

By our analysis, (x, y) with $x \neq y$ solves (3.7) if and only if

$$(3.9) \quad xy + x + y + 3 = 0$$

or

$$(3.10) \quad 12x^2y^2 + 31x^2y + 31xy^2 - 5x^2 + 28xy - 5y^2 - 65x - 65y - 50 = 0$$

and it solves (3.8) if and only if

$$(3.11) \quad xy + x + y - 1 = 0$$

or

$$(3.12) \quad 12x^2y^2 + 25x^2y + 25xy^2 + 37x^2 + 37y^2 + 25x + 25y + 62 = 0$$

Clearly, (3.9) and (3.11) can not be satisfied simultaneously. Moreover, notice that

$$\begin{aligned} & 12x^2y^2 + 25x^2y + 25xy^2 + 37x^2 + 37y^2 + 25x + 25y + 62 = \\ & = 6x^2 \left(y^2 + \frac{25}{6}y + \frac{30.75}{6} \right) + 6y^2 \left(x^2 + \frac{25}{6}x + \frac{30.75}{6} \right) + (2.5x + 5)^2 + (2.5y + 5)^2 + 12 = \\ & = 6x^2 \left(\left(y + \frac{25}{12} \right)^2 + \frac{113}{144} \right) + 6y^2 \left(\left(x + \frac{25}{12} \right)^2 + \frac{113}{144} \right) + (2.5x + 5)^2 + (2.5y + 5)^2 + 12 > 0, \end{aligned}$$

i.e (3.12) is never satisfied. Therefore, it is enough to check that (3.11) and (3.10) are not satisfied simultaneously. Assume that (x, y) with $x \neq y$ is a solution of (3.11). Plugging $xy = 1 - x - y$ into (3.10), we see that

$$\begin{aligned} & 12(1 - x - y)^2 + 31x(1 - x - y) + 31y(1 - x - y) - 5x^2 + 28(1 - x - y) - 5y^2 - 65x - 65y - 50 = \\ & = -24x^2 - 38xy - 24y^2 - 86x - 86y - 10 = -24x^2 - 38(1 - x - y) - 24y^2 - 86x - 86y - 10 = \\ & = -24(x^2 + y^2 + 2x + 2y + 2) = -24((x + 1)^2 + (y + 1)^2) < 0, \end{aligned}$$

i.e. (x, y) is not a solution of (3.10). Therefore, (3.1) can not be satisfied for both (z_1, z_2) and (z_2, z_3) , hence condition (4) is satisfied. \square

It is left to prove our points satisfy condition (5) of Definition 3.10. To this end, we first prove two technical lemmas.

Lemma 3.20. *Let $\{z_1, \dots, z_n\}, \{w_1, \dots, w_n\} \subset \mathbb{D} \setminus \{0, \lambda\}$ be two sets of points such that $z_i \neq z_j$ and $w_i \neq w_j$ for all $1 \leq i \neq j \leq n$. Then:*

$$P = \left[\frac{B_\lambda(z_i) \overline{B_\lambda(w_j)}}{1 - z_i \overline{w_j}} \right]_{i,j=1}^n$$

is invertible.

Proof. As $B_\lambda(z_i) \neq 0$ and $B_\lambda(w_j) \neq 0$ by assumption, and since:

$$P = \begin{pmatrix} B_\lambda(z_1) & 0 & \dots & 0 \\ 0 & B_\lambda(z_2) & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & B_\lambda(z_n) \end{pmatrix} \left[\frac{1}{1 - z_i \overline{w_j}} \right] \begin{pmatrix} \overline{B_\lambda(w_1)} & 0 & \dots & 0 \\ 0 & \overline{B_\lambda(w_2)} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \overline{B_\lambda(w_n)} \end{pmatrix}$$

it is enough to prove the middle matrix is invertible. We prove this by induction on n . For $n = 1$, the claim is clear. Assume this is true for n , and consider the sets $\{z_1, \dots, z_{n+1}\}, \{w_1, \dots, w_{n+1}\}$ such that they satisfy the assumption. Calculating the determinant of P as a function of z_1 gives us

$$\det(P) = \sum_{j=1}^{n+1} (-1)^{1+j} \det((P)_{1,j}) \cdot \frac{1}{1 - z_1 \overline{w_j}}.$$

Replacing z_1 with a variable z and thinking of $\det(P)$ as a function of z , $\det(P)(z)$ has poles at $\frac{1}{\overline{w_j}}$ for all $1 \leq j \leq n+1$ (by induction hypothesis, $\det((P)_{1,j}) \neq 0$), and:

$$\det(P)(z_2) = \dots = \det(P)(z_{n+1}) = \det(P)(\infty) = 0$$

i.e it has $n+1$ zeroes at z_2, \dots, z_n, ∞ . Therefore, $\det(P) = \det(P)(z_1) \neq 0$. \square

Lemma 3.21. *Let $\{z_1, \dots, z_n\}, \{w_1, \dots, w_n\} \subset (\mathbb{D} \cap \mathbb{R}) \setminus \{0, \lambda\}$ be two sets of points such that $z_i \neq z_j$ and $w_i \neq w_j$ for all $1 \leq i \neq j \leq n$. Then the set of all pairs $(\alpha, \beta) \in \mathbb{T}$ for which the matrix $[k^{\alpha, \beta}(z_i, w_j)]_{i,j=1}^n$ is singular is either \mathbb{T} or finite.*

Proof. Re-write the matrix as:

$$\underbrace{\begin{pmatrix} \alpha + f_\lambda(z_1)\beta \\ \vdots \\ \alpha + f_\lambda(z_n)\beta \end{pmatrix}}_{u_{\alpha, \beta}} \cdot \underbrace{\begin{pmatrix} \alpha + f_\lambda(w_1)\beta \\ \vdots \\ \alpha + f_\lambda(w_n)\beta \end{pmatrix}}_{v_{\alpha, \beta}^*} + \underbrace{\left[\frac{B_\lambda(z_i) \overline{B_\lambda(w_j)}}{1 - z_i \overline{w_j}} \right]}_P.$$

This is a rank one perturbation of the invertible matrix P . By the determinant lemma, its determinant is given by $\det(P)(1 + v_{\alpha, \beta}^* P^{-1} u_{\alpha, \beta})$. As everything is real-valued, the polynomial $1 + v_{\alpha, \beta}^* P^{-1} u_{\alpha, \beta}$ is a real polynomial of degree 2 in (α, β) . Since the unit circle is an irreducible real algebraic curve, the level-set

$$\{(\alpha, \beta) \in \mathbb{T} : 1 + v_{\alpha, \beta}^* P^{-1} u_{\alpha, \beta} = 0\}$$

is either finite or is equal to \mathbb{T} . \square

Proposition 3.22. *The points z_1, \dots, z_4 satisfy condition (5) of Definition 3.10.*

Proof. First of all, we note that for z_1, \dots, z_4 :

$$\omega = \zeta = \sqrt{2} - 1$$

Let $\sigma = (12)(34)$, $\tau = (14)(23) \in S_4$. Assume that $\ell \in \{1, 2\}$ and $(\alpha, \beta) \in A'$ satisfies:

$$k_{z_\ell}^{\alpha, \beta} \in \text{span}\{k_{z_{\tau(\ell)}}^{\alpha, \beta}, k_{z_{\sigma(\ell)}}^{\alpha, \beta}, P_{M_{\alpha, \beta}} k_{\sqrt{2}-1}^{\alpha, \beta}\}$$

Since $k_{z_{\tau(\ell)}}^{\alpha, \beta}$, $k_{z_{\sigma(\ell)}}^{\alpha, \beta}$, $k_{z_\ell}^{\alpha, \beta}$ are linearly independent, there are scalars $\gamma_1, \gamma_2, \gamma_3$ such that:

$$P_{M_{\alpha, \beta}} k_{\sqrt{2}-1}^{\alpha, \beta} = \gamma_1 k_{z_\ell}^{\alpha, \beta} + \gamma_2 k_{z_{\tau(\ell)}}^{\alpha, \beta} + \gamma_3 k_{z_{\sigma(\ell)}}^{\alpha, \beta}$$

By taking inner products with $k_{z_i}^{\alpha, \beta}$ on both sides for $1 \leq i \leq 4$, this implies that the matrix:

$$C_\ell(\alpha, \beta) = [k^{\alpha, \beta}(z_i, w_j)]$$

is singular, where:

$$w_j = \begin{cases} z_j & j \neq \sigma(\ell) \\ \sqrt{2} - 1 & j = \sigma(\ell) \end{cases}$$

Now assume $1 \leq \ell \leq 4$ and $(\alpha, \beta) \in A$ satisfies:

$$f_\ell \notin \text{span}\{k_{z_{\tau(\ell)}}^{\alpha, \beta}, k_{z_{\sigma(\ell)}}^{\alpha, \beta}, P_{M_{\alpha, \beta}} k_{\sqrt{2}-1}^{\alpha, \beta}\}$$

then there are scalars $\gamma_1, \gamma_2, \gamma_3$ such that:

$$f_\ell = \gamma_1 k_{z_{\tau(\ell)}}^{\alpha, \beta} + \gamma_2 k_{z_{\sigma(\ell)}}^{\alpha, \beta} + \gamma_3 P_{M_{\alpha, \beta}} k_{\sqrt{2}-1}^{\alpha, \beta}$$

by taking inner products with f_i for $1 \leq i \neq \ell \leq 4$, this implies that the 3×3 matrix:

$$(C_\ell(\alpha, \beta))_{\ell, \ell}$$

is singular. By taking $\alpha = 1$, $\beta = 0$ and calculating $k^{1,0}(z_\ell, \sqrt{2} - 1)$ for all $1 \leq \ell \leq 4$, we obtain the following four matrices:

$$C_1(1, 0) = \begin{pmatrix} 5 & \frac{17-8\sqrt{2}}{7} & \frac{4}{5} & \frac{19}{15} \\ 5 & \frac{13-7\sqrt{2}}{3} & \frac{31}{30} & \frac{41}{45} \\ \frac{6}{5} & \frac{5(\sqrt{2}-1)}{\sqrt{2}} & \frac{29}{28} & \frac{11}{12} \\ \frac{19}{15} & \frac{5(2\sqrt{2}+1)}{21} & \frac{11}{12} & \frac{7}{6} \end{pmatrix}, C_2(1, 0) = \begin{pmatrix} \frac{17-8\sqrt{2}}{7} & \frac{5}{6} & \frac{4}{5} & \frac{19}{15} \\ \frac{13-7\sqrt{2}}{3} & \frac{65}{63} & \frac{31}{30} & \frac{41}{45} \\ \frac{5(\sqrt{2}-1)}{\sqrt{2}} & \frac{31}{63} & \frac{29}{30} & \frac{11}{28} \\ \frac{5(2\sqrt{2}+1)}{21} & \frac{41}{45} & \frac{11}{12} & \frac{7}{6} \end{pmatrix}$$

$$C_3(1, 0) = \begin{pmatrix} 5 & \frac{5}{6} & \frac{4}{5} & \frac{17-8\sqrt{2}}{7} \\ \frac{5}{6} & \frac{65}{63} & \frac{31}{30} & \frac{13-7\sqrt{2}}{3} \\ \frac{4}{5} & \frac{31}{63} & \frac{29}{30} & \frac{5(\sqrt{2}-1)}{\sqrt{2}} \\ \frac{19}{15} & \frac{41}{45} & \frac{11}{12} & \frac{5(2\sqrt{2}+1)}{21} \end{pmatrix}, C_4(1, 0) = \begin{pmatrix} 5 & \frac{5}{6} & \frac{17-8\sqrt{2}}{7} & \frac{19}{15} \\ \frac{5}{6} & \frac{65}{63} & \frac{13-7\sqrt{2}}{3} & \frac{41}{45} \\ \frac{4}{5} & \frac{31}{63} & \frac{5(\sqrt{2}-1)}{\sqrt{2}} & \frac{11}{28} \\ \frac{19}{15} & \frac{41}{45} & \frac{5(2\sqrt{2}+1)}{21} & \frac{7}{6} \end{pmatrix}$$

The matrices $C_\ell(1, 0)$ and $(C_\ell(1, 0))_{\ell, \ell}$ are invertible for all $1 \leq \ell \leq 4$. By Lemmas 3.20 and 3.21, this implies that for every $1 \leq \ell \leq 4$, $C_\ell(\alpha, \beta)$ and $(C_\ell(\alpha, \beta))_{\ell, \ell}$ are invertible for all $(\alpha, \beta) \in A' \setminus S$ for some finite set $S \subset A'$. In particular, by taking $A = A' \setminus S$, A is the desired set, and condition (5) is satisfied. \square

Proof of Theorem 3.18. We saw that if we take A to be the complement of a finite set in the open upper-half circle, then the conditions of Definition 3.10 are satisfied. Hence, our points are good. \square

Remark 3.23. A comment on the choices made in the example is in order. The technicalities involved in proving the unitary inequivalence, had forced us to restrict our attention to real (α, β) and real points. We believe that every $[\alpha : \beta] \in \mathbb{P}^1(\mathbb{C})$ outside of a finite set, perhaps, gives rise to a boundary representation of H_{node}^∞/I . Therefore, $C(P^1(\mathbb{C}))$ is a quotient of $C_e^*(H_{\text{node}}^\infty/I)$. It is also challenging to find a higher-dimensional boundary representation. Though, it might be the case that most (α, β) in the Grassmannian will give such representations.

4. A C^* -COVER CANDIDATE

The non-commutative Grassmanian, G_n^{nc} , introduced by Brown [10], is the universal C^* -algebra generated by the identity element and $\{P_{i,j} : 1 \leq i, j \leq n\}$ such that the matrix:

$$P = [P_{i,j}]_{i,j=1}^n$$

is a projection. In [26], the author proves that G_2^{nc} is a C^* -subalgebra of the free product $M_2 *_{\mathbb{C}} \mathbb{C}^2$. Consequently, by a theorem of Exel and Loring [20, Theorem 3.2], G_2^{nc} is RFD. In this section, we present another characterization of G_2^{nc} , and define a completely isometric embedding of H_{node}^∞/I into $M_n(G_2^{nc})$, providing a candidate for the C^* -envelope of H_{node}^∞/I .

Let \mathfrak{A} be the universal C^* -algebra generated by two elements x, y such that $x^*x + y^*y = 1$. Note that such universal C^* algebra exists by [8, Section 2, 8.3.1], since the relation is of the form:

$$\|p(x, y, x^*, y^*)\| = 0$$

for the non-commutative polynomial $p(x, y, x^*, y^*) = x^*x + y^*y - 1$, this relation is realizable by bounded operators on a Hilbert space, and it places an upper bound on $\|x\|$ and $\|y\|$. This algebra is also known as the universal Pythagorean algebra (see for example [9] and [11]).

Theorem 4.1. *Let $\mathfrak{B} = C^*(1, xx^*, yy^*, xy^*) \subset \mathfrak{A}$. Then $G_2^{nc} \cong \mathfrak{B}$.*

Proof. As $\begin{pmatrix} x \\ y \end{pmatrix}$ is a column isometry, the matrix:

$$\begin{pmatrix} x \\ y \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}^* = \begin{pmatrix} xx^* & xy^* \\ yx^* & yy^* \end{pmatrix}$$

is a projection. By universality, if $1, a, b, c$ are the generators of G_2^{nc} , there exists a surjective $*$ -homomorphism:

$$\rho : G_2^{nc} \rightarrow \mathfrak{B}, 1 \rightarrow 1, a \rightarrow xx^*, b \rightarrow xy^*, c \rightarrow yy^*$$

It is enough to prove that ρ is injective. To this end, let $x \in \ker(\rho)$. As G_2^{nc} is RFD, to show $x = 0$ it is enough to prove that for every finite-dimensional representation π of G_2^{nc} , $\pi(x) = 0$. Fix a finite-dimensional representation:

$$\pi : G_2^{nc} \rightarrow M_n(\mathbb{C})$$

Then the matrix:

$$P = \begin{pmatrix} \pi(a) & \pi(b) \\ \pi(b)^* & \pi(c) \end{pmatrix}$$

is a projection. Let $m = \text{tr}(P)$. We have two cases:

(1) If $m \leq n$, let $\chi_1 : G_2^{nc} \rightarrow \mathbb{C}$ be the character satisfying:

$$\chi_1(a) = \chi_1(c) = 1, \chi_1(b) = 0$$

and consider the representation $\pi' = \pi \oplus \chi_1^{n-m}$. Then:

$$P' = \begin{pmatrix} \pi'(a) & \pi'(b) \\ \pi'(b)^* & \pi'(c) \end{pmatrix}$$

is a projection in $M_{2(2n-m)}(\mathbb{C})$ with $\text{tr}(P') = 2(n-m) + m = 2n - m$, hence there exists a column isometry:

$$V : \mathbb{C}^{2n-m} \rightarrow \mathbb{C}^{2(2n-m)}, V = \begin{pmatrix} \tilde{x} \\ \tilde{y} \end{pmatrix}$$

such that $VV^* = P'$. By universality of \mathfrak{A} , we have a representation:

$$\tilde{\pi} : \mathfrak{A} \rightarrow M_{2n-m}(\mathbb{C}), \tilde{\pi}(x) = \tilde{x}, \tilde{\pi}(y) = \tilde{y}$$

Therefore: $\pi' = \tilde{\pi}|_{\mathfrak{B}} \circ \rho$. As $x \in \ker(\rho)$, $\pi'(x) = 0$ and consequently $\pi(x) = 0$.

(2) If $n < m$, let $\chi_0 : G_2^{nc} \rightarrow \mathbb{C}$ be the character that send a, b , and c to zero, and consider the representation $\pi' = \pi \oplus \chi_0^{m-n}$. Then:

$$P' = \begin{pmatrix} \pi'(a) & \pi'(b) \\ \pi'(b)^* & \pi'(c) \end{pmatrix}$$

is a projection in $M_{2m}(\mathbb{C})$ with $\text{tr}(P') = m$. By similar arguments, again $\pi(x) = 0$.

In either case, $\pi(x) = 0$ and therefore $x = 0$. Hence, ρ is an isomorphism of C^* -algebras. \square

For every m and every $(\alpha, \beta) \in \mathbb{G}(m \times m)$, $\begin{pmatrix} \alpha^t \\ \beta^t \end{pmatrix}$ is a column isometry. By universality of \mathfrak{A} , there exists a representation

$$\pi_{\alpha, \beta} : \mathfrak{A} \rightarrow M_m(\mathbb{C})$$

given by $x \rightarrow \alpha^t, y \rightarrow \beta^t$. Moreover, note that every m dimensional representation of \mathfrak{A} is of this form for some pair $(\alpha, \beta) \in \mathbb{G}(m \times m)$.

Define $\Psi \in M_n(\mathfrak{A})$ by

$$\Psi = \sum_{i,j=1}^n \underbrace{\left((x + f_\lambda(z_i)y) (x^* + \overline{f_\lambda(z_j)}y^*) + \frac{B_\lambda(z_i)\overline{B_\lambda(z_j)}}{1 - z_i\overline{z_j}} \right)}_{i_{i,j}} F_{i,j}$$

where $F_{i,j}$ are the matrix units of $M_n(\mathbb{C})$. An important observation is the following.

Proposition 4.2. *Assuming $0, \lambda \notin \{z_1, \dots, z_n\}$, Ψ is positive definite.*

Proof. Let $L = \begin{pmatrix} x + f_\lambda(z_1)y \\ \vdots \\ x + f_\lambda(z_n)y \end{pmatrix}$. Then:

$$\Psi = LL^* + \underbrace{\left[\frac{B_\lambda(z_i)\overline{B_\lambda(z_j)}}{1 - z_i\overline{z_j}} \right]_{i,j=1}^n}_G.$$

Note that G is the Gram matrix of a linearly independent set, since the kernel functions of H^2 are linearly independent, our points do not contain the zeroes of B_λ , and, moreover, B_λ is an isometry. Hence, G is strictly positive and so is Ψ . \square

We are ready to present the main theorem of this section.

Theorem 4.3. *The homomorphism $\Gamma : H_{\text{node}}^\infty / I \mapsto M_n(\mathfrak{A})$ given by*

$$\Gamma(f + I) = \Psi^{-\frac{1}{2}} \underbrace{\begin{pmatrix} f(z_1) & 0 & \dots & 0 \\ 0 & f(z_2) & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & f(z_n) \end{pmatrix}}_{D_f} \Psi^{\frac{1}{2}}$$

is completely isometric. In fact, $\Gamma(H_{\text{node}}^\infty / I) \subset M_n(\mathfrak{B})$.

To prove Theorem 4.3, we begin by proving several lemmas to reformulate a generalized trace condition of the one presented in Section 2. For fixed $m, r \in \mathbb{N}$, $(\alpha, \beta) \in \mathbb{G}(m \times m)$ and $W_1, \dots, W_n \in M_r(\mathbb{C})$, define the generalized trace condition to be:

$$(4.1) \quad \sum_{i,j=1}^n \text{tr}[X_j k^{\alpha, \beta}(z_i, z_j) X_i^* - W_j^* X_j k^{\alpha, \beta}(z_i, z_j) X_i^* W_i] \geq 0$$

for all $X_1, \dots, X_n \in M_{r \times m}(\mathbb{C})$. Let $m, r \in \mathbb{N}$, and consider the space $M_{r, mn}(\mathbb{C})$, thought of as the space of row block matrices with n blocks of size $r \times m$. In other words, we identify $M_{r, mn} \cong M_{r, m} \otimes \mathbb{C}^n$. Fix some $W = (W_1, \dots, W_n) \in M_r(\mathbb{C})^{\oplus n}$ and define two linear operators:

$$\psi_{\alpha, \beta}, \varphi_W : M_{r \times mn}(\mathbb{C}) \rightarrow M_{r \times mn}(\mathbb{C})$$

by

$$\psi_{\alpha, \beta}(X_1 \ \dots \ X_n) = (X_1 \ \dots \ X_n) \cdot [k^{\alpha, \beta}(z_j, z_i)^t]_{i, j=1}^n$$

and

$$\varphi_W(X_1 \ \dots \ X_n) = (W_1 X_1 \ \dots \ W_n X_n)$$

Furthermore, define an inner product on $M_{r \times mn}(\mathbb{C})$ by

$$\left\langle \begin{pmatrix} X_1 \\ \vdots \\ X_n \end{pmatrix}^T, \begin{pmatrix} Y_1 \\ \vdots \\ Y_n \end{pmatrix}^T \right\rangle = \text{tr} \left(\sum_{i=1}^n X_i Y_i^* \right).$$

It is straightforward that with respect to this inner product:

$$\varphi_W^*(X_1 \ \dots \ X_n) = (W_1^* X_1 \ \dots \ W_n^* X_n)$$

Lemma 4.4. Let $E_{i, j}$ be the matrix units in $M_{r \times mn}$, and define the ordered basis

$$\varepsilon = \{E_{i, j} : 1 \leq i \leq r, 1 \leq j \leq mn\}$$

ordered by $E_{i, j} < E_{k, l}$ if and only if $j < l$ or $j = l$ and $i < k$. Then, in Kronecker's notation [23], for every $n \in \mathbb{N}$ and every $W = (W_1, \dots, W_n) \in M_r(\mathbb{C})^{\oplus n}$,

$$[\psi_{\alpha, \beta}]_\varepsilon = \sum_{i, j=1}^n F_{j, i} \otimes k^{\alpha, \beta}(z_j, z_i)^t \otimes I_r, \quad [\varphi_W]_\varepsilon = \sum_{i=1}^n F_{i, i} \otimes I_m \otimes W_i$$

with $F_{i, j}$ the matrix units of $M_n(\mathbb{C})$.

Proof. By definition, $\psi_{\alpha, \beta}$ is given by right multiplication by the matrix $\sum_{i, j=1}^n F_{i, j} \otimes k^{\alpha, \beta}(z_j, z_i)$. Hence by [23, Lemma 4.31], $[\psi_{\alpha, \beta}]_\varepsilon = \sum_{i, j=1}^n E_{i, j} \otimes k^{\alpha, \beta}(z_j, z_i)^t \otimes I_r$.

For the second part, by definition, φ_W acts on the i 'th block by multiplication on the left by W_i . Therefore, again by [23, Lemma 4.31], $[\varphi_W]_\varepsilon$ is a block diagonal matrix, with the i 'th block equal to $I_m \otimes W_i$. Therefore,

$$[\varphi_W]_\varepsilon = \sum_{i=1}^n F_{i, i} \otimes I_m \otimes W_i.$$

□

Remark 4.5. Immediately, ε is an orthonormal basis with respect to the inner product defined on $M_{r \times mn}(\mathbb{C})$.

Lemma 4.6. For every $(\alpha, \beta) \in \mathbb{G}(m \times m)$, $\pi_{\alpha^t, \beta^t}^{(n)}(\Psi) \otimes I_r = [\psi_{\alpha, \beta}]_\varepsilon$. In particular, $[\psi_{\alpha, \beta}]_\varepsilon$ is positive definite.

Proof.

$$\begin{aligned} (\pi_{\alpha^t, \beta^t}^{(n)}(\Psi))_{i, j} &= \pi_{\alpha^t, \beta^t}(\ell_{i, j}) = (\alpha^t + f_\lambda(z_i)\beta^t)(\bar{\alpha} + \overline{f_\lambda(z_j)\beta}) + \frac{B_\lambda(z_i)\overline{B_\lambda(z_j)}}{1 - z_i \bar{z}_j} I_m = \\ &= k^{\alpha, \beta}(z_i, z_j)^t \end{aligned}$$

Therefore

$$\pi_{\alpha^t, \beta^t}^{(n)}(\Psi) \otimes I_r = \sum_{i, j=1}^n F_{j, i} \otimes k^{\alpha, \beta}(z_j, z_i)^t \otimes I_r = [\psi_{\alpha, \beta}]_\varepsilon.$$

For the second part, by 4.2 and since $\pi^{(n)}$ is a homomorphism of unital C^* -algebras, $\pi_{\alpha^t, \beta^t}^{(n)}(\Psi)$ is positive definite. By properties of Kronecker product and since I_r is positive definite, $[\psi_{\alpha, \beta}]_\varepsilon$ is positive definite. Lastly, by remark 4.5, $\psi_{\alpha, \beta}$ is positive definite. □

Lemma 4.7. *The generalized trace condition for fixed $m, r \in \mathbb{N}$, $(\alpha, \beta) \in \mathbb{G}(m \times m)$ and $W_1, \dots, W_n \in M_r(\mathbb{C})$ is equivalent to*

$$\left\| \left[\psi_{\alpha, \beta}^{-\frac{1}{2}} \right]_{\varepsilon} \cdot [\varphi_W]_{\varepsilon} \cdot \left[\psi_{\alpha, \beta}^{\frac{1}{2}} \right]_{\varepsilon} \right\| \leq 1.$$

Proof. By our previous notation, the generalized trace condition is:

$$(4.2) \quad \begin{aligned} & \operatorname{tr} \left[\begin{array}{ccc} (X_1 & \dots & X_n) [k^{\alpha, \beta}(z_j, z_i)]_{i,j=1}^n \begin{pmatrix} X_1^* \\ \vdots \\ X_n^* \end{pmatrix} \\ - (W_1^* X_1 & \dots & W_n^* X_n) [k^{\alpha, \beta}(z_j, z_i)]_{i,j=1}^n \begin{pmatrix} X_1^* W_1 \\ \vdots \\ X_n^* W_n \end{pmatrix} \end{array} \right] \geq 0 \end{aligned}$$

for all $X_1, \dots, X_n \in M_{r \times m}(\mathbb{C})$. Denoting $X = (X_1 \ \dots \ X_n)$, this is the same as

$$\langle \psi_{\alpha, \beta}(X) - \varphi_W(\psi_{\alpha, \beta}(\varphi_W^*(X))), X \rangle \geq 0$$

for all $X_1, \dots, X_n \in M_{r \times m}(\mathbb{C})$. By definition, this is the same as

$$\psi_{\alpha, \beta} \geq \varphi_W \circ \psi_{\alpha, \beta} \circ \varphi_W^*.$$

Using Lemma 4.6, we can conjugate both sides by $\psi_{\alpha, \beta}^{-\frac{1}{2}}$. Consequently, the trace condition is equivalent to:

$$1 \geq \left(\psi_{\alpha, \beta}^{-\frac{1}{2}} \circ \varphi_W \circ \psi_{\alpha, \beta}^{\frac{1}{2}} \right) \left(\psi_{\alpha, \beta}^{-\frac{1}{2}} \circ \varphi_W \circ \psi_{\alpha, \beta}^{\frac{1}{2}} \right)^*$$

which is equivalent to:

$$\left\| \psi_{\alpha, \beta}^{-\frac{1}{2}} \circ \varphi_W \circ \psi_{\alpha, \beta}^{\frac{1}{2}} \right\| \leq 1$$

By Remark 4.5, we are done. □

We are now ready to prove Theorem 4.3.

Proof of Theorem 4.3. Throughout this proof, we will use the notation:

$$\widetilde{\psi}_{\alpha, \beta} := \sum_{i,j=1}^k F_{j,i} \otimes k^{\alpha, \beta}(z_j, z_i)^t$$

Let $r \in \mathbb{N}$ and consider the homomorphism $\Gamma^{(r)} : M_r(H_{\text{node}}^{\infty}/I) \rightarrow M_r(M_n(\mathfrak{A}))$ given by

$$\Gamma^{(r)}([f_{i,j} + I]) = \sum_{i,j=1}^r E_{i,j} \otimes \Gamma(f_{i,j} + I).$$

To show Γ^r is an isometry, it is enough to prove $\|[f_{i,j} + I]\| \leq 1 \iff \|\Gamma^{(r)}([f_{i,j} + I])\| \leq 1$. As \mathfrak{A} is RFD so is $M_r(M_n(\mathfrak{A}))$. Hence, our claim is equivalent to showing that $\|[f_{i,j} + I]\| \leq 1 \iff \|\tilde{\rho}(\Gamma^{(r)}([f_{i,j} + I]))\| \leq 1$ for every finite-dimensional representation $\tilde{\rho} : M_r(M_n(\mathfrak{A})) \rightarrow M_k(\mathbb{C})$. Fix a finite-dimensional representation $\tilde{\rho}$ of $M_r(M_n(\mathfrak{A}))$. It is well known that $\tilde{\rho}$ is unitarily equivalent to $\rho^{(r)}$ for some representation

$$\rho : M_n(\mathfrak{A}) \rightarrow M_s(\mathbb{C}).$$

Hence, $\|\rho^{(r)}(\Gamma^{(r)}([f_{i,j} + I])\| = \|\tilde{\rho}(\Gamma^{(r)}([f_{i,j} + I])\|$. Again, ρ is unitarily equivalent to $\hat{\rho}^{(n)}$ where $\hat{\rho}$ is a representation of \mathfrak{A} . As we noted, $\hat{\rho} = \pi_{\alpha, \beta}$ for some $(\alpha, \beta) \in \mathbb{G}(m \times m)$, hence ρ is unitarily equivalent to $\pi_{\alpha, \beta}^{(n)}$. Let U be the unitary satisfying $\rho([f_{p,q}]) = U\pi_{\alpha, \beta}^{(n)}([f_{p,q}])U^*$. Then

$$\begin{aligned} \rho^{(r)} \left(\Gamma^{(r)}([f_{i,j} + I]) \right) &= \sum_{i,j=1}^r E_{i,j} \otimes \rho(\Gamma(f_{i,j} + I)) = \sum_{i,j=1}^r E_{i,j} \otimes \left(U\pi_{\alpha, \beta}^{(n)}(\Gamma(f_{i,j} + I))U^* \right) = \\ &= \sum_{i,j=1}^r E_{i,j} \otimes \left(U\pi_{\alpha, \beta}^{(n)} \left(\Psi^{-\frac{1}{2}} D_{f_{i,j}} \Psi^{\frac{1}{2}} \right) U^* \right) \stackrel{\pi_{\alpha, \beta}^{(n)}(f_{i,j}(z_k))=f_{i,j}(z_k)I_m}{=} \end{aligned}$$

$$\begin{aligned}
&= \sum_{i,j=1}^n E_{i,j} \otimes \left(U \widetilde{\psi}_{\alpha,\beta}^{-\frac{1}{2}} \left(\sum_{k=1}^n F_{k,k} \otimes f_{i,j}(z_k) I_m \right) \widetilde{\psi}_{\alpha,\beta}^{\frac{1}{2}} U^* \right) = \\
&= (I_r \otimes U^*) \left(I_r \otimes \widetilde{\psi}_{\alpha,\beta}^{-\frac{1}{2}} \right) \left(\sum_{i,j=1}^r E_{i,j} \otimes \left(\sum_{k=1}^n F_{k,k} \otimes f_{i,j}(z_k) I_m \right) \right) \left(I_r \otimes \widetilde{\psi}_{\alpha,\beta}^{\frac{1}{2}} \right) (I_r \otimes U).
\end{aligned}$$

Since I_r and U are unitaries, $I_r \otimes U$ is unitary (with $(I_r \otimes U)^* = I_r \otimes U^*$) by properties of Kronecher product, that is $\rho^{(r)}(\Gamma^{(r)}([f_{i,j}]))$ is unitarily equivalent to

$$(4.3) \quad \left(I_r \otimes \widetilde{\psi}_{\alpha,\beta}^{-\frac{1}{2}} \right) \left(\sum_{i,j=1}^r E_{i,j} \otimes \left(\sum_{k=1}^n F_{k,k} \otimes f_{i,j}(z_k) I_m \right) \right) \left(I_r \otimes \widetilde{\psi}_{\alpha,\beta}^{\frac{1}{2}} \right).$$

Conjugating (4.3) by the unitary that interchanges the first two tensor products and then by the unitary that interchanges the last two tensor products, we get

$$\begin{aligned}
&\left(\widetilde{\psi}_{\alpha,\beta}^{-\frac{1}{2}} \otimes I_r \right) \left(\sum_{k=1}^n F_{k,k} \otimes I_m \otimes W_k \right) \left(\widetilde{\psi}_{\alpha,\beta}^{\frac{1}{2}} \otimes I_r \right) = [\psi_{\alpha,\beta}]_{\varepsilon}^{-\frac{1}{2}} \cdot [\varphi_W]_{\varepsilon} \cdot [\psi_{\alpha,\beta}]_{\varepsilon}^{-\frac{1}{2}} = \\
&= \left[\psi_{\alpha,\beta}^{-\frac{1}{2}} \circ \varphi_W \circ \psi_{\alpha,\beta}^{\frac{1}{2}} \right]_{\varepsilon}
\end{aligned}$$

with $W_k = [f_{i,j}](z_k)$. Now assume that $\|\tilde{\rho}(\Gamma^{(r)}(f))\| \leq 1$. By unitary equivalence, and since ε is an orthonormal basis,

$$\left\| \left[\psi_{\alpha,\beta}^{-\frac{1}{2}} \circ \varphi_W \circ \psi_{\alpha,\beta}^{\frac{1}{2}} \right]_{\varepsilon} \right\| = \left\| \psi_{\alpha,\beta}^{-\frac{1}{2}} \circ \varphi_W \circ \psi_{\alpha,\beta}^{\frac{1}{2}} \right\| \leq 1.$$

We observed this is equivalent to the generalized trace condition (4.1) for W_1, \dots, W_n being true for r and m . As $M_r(H_{\text{node}}^{\infty}/I)$ and $M_r(H_{\text{node}}^{\infty})/I$ are isometrically isomorphic, by the generalized distance formula this is equivalent to $\|[f_{i,j}] + I\| = \|[f_{i,j} + I]\| \leq 1$. Therefore, $\tilde{\rho}(\Gamma^{(r)})$ is an isometry, hence (again using the fact that \mathfrak{A} is RFD) $\Gamma^{(r)}$ is an isometry. Lastly, by examining the formula for Ψ , we see that $\Psi \in M_n(\mathfrak{B})$. Hence, the range of Γ is contained in $M_n(\mathfrak{B})$. \square

Question 4.8. Is $C^*(\Gamma(H_{\text{node}}^{\infty}/I))$ the C^* -envelope of $H_{\text{node}}^{\infty}/I$ at least for n big enough?

Remark 4.9. It follows from a general proposition in [14], that if we have only two points, then $C_e^*(H_{\text{node}}^{\infty}/I) \cong M_2(\mathbb{C})$. Hence, we only ask the above question for $n \geq 3$. However, maybe, a bigger n is necessary to ensure that $C_e^*(H_{\text{node}}^{\infty}/I) \cong C^*(\Gamma(H_{\text{node}}^{\infty}/I))$. It seems that a big obstruction in understanding the above question is a technical one. Namely, figuring out whether $\Psi \in C^*(\Gamma(H_{\text{node}}^{\infty}/I))$. If it is true, then one might be able to use [22, Theorem].

REFERENCES

- [1] M. B. Abrahamse. The Pick interpolation theorem for finitely connected domains. *Michigan Mathematical Journal*, 26(2):195–203, 1979.
- [2] J. Agler. Some interpolation theorems of Nevanlinna-Pick type. unpublished (1988).
- [3] J. Agler and J. E. McCarthy. Complete Nevanlinna-Pick kernels. *J. Funct. Anal.*, 175(1):111–124, 2000.
- [4] W. Arveson. Subalgebras of C^* -algebras. *Acta Math.*, 123:141–224, 1969.
- [5] W. Arveson. The noncommutative Choquet boundary. *J. Amer. Math. Soc.*, 21(4):1065–1084, 2008.
- [6] J. A. Ball. A lifting theorem for operator models of finite rank on multiply-connected domains. *J. Operator Theory*, 1(1):3–25, 1979.
- [7] J. A. Ball, V. Bolotnikov, and S. ter Horst. A constrained Nevanlinna-Pick interpolation problem for matrix-valued functions. *Indiana University mathematics journal*, pages 15–51, 2010.
- [8] B. Blackadar. *Operator algebras*, volume 122 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2006. Theory of C^* -algebras and von Neumann algebras, Operator Algebras and Non-commutative Geometry, III.
- [9] A. Brothier and V. F. R. Jones. Pythagorean representations of Thompson’s groups. *J. Funct. Anal.*, 277(7):2442–2469, 2019.
- [10] L. G. Brown. Ext of certain free product C^* -algebras. *Journal of Operator Theory*, pages 135–141, 1981.
- [11] K. Courtney and D. Sherman. The universal C^* -algebra of a contraction. *J. Operator Theory*, 84(1):153–184, 2020.
- [12] K. R. Davidson and R. Hamilton. Nevanlinna–Pick interpolation and factorization of linear functionals. *Integral Equations and Operator Theory*, 70(1):125–149, 2011.
- [13] K. R. Davidson and M. Kennedy. Noncommutative Choquet theory. *arXiv preprint arXiv:1905.08436*, 2019.

- [14] K. R. Davidson, V. Paulsen, M. Raghupathi, and D. Singh. A constrained Nevanlinna-Pick interpolation problem. *Indiana University Mathematics Journal*, pages 709–732, 2009.
- [15] K. R. Davidson and E. Shamovich. Nevanlinna-Pick families and singular rational varieties. In *Operator Theory, Operator Algebras and Their Interactions with Geometry and Topology: Ronald G. Douglas Memorial Volume*, pages 129–145. Springer, 2020.
- [16] M. A. Dritschel, M. T. Jury, and S. McCullough. Dilations and constrained algebras. *Oper. Matrices*, 10(4):829–861, 2016.
- [17] M. A. Dritschel and S. A. McCullough. Boundary representations for families of representations of operator algebras and spaces. *J. Operator Theory*, 53(1):159–167, 2005.
- [18] M. A. Dritschel and J. Pickering. Test functions in constrained interpolation. *Trans. Amer. Math. Soc.*, 364(11):5589–5604, 2012.
- [19] M. A. Dritschel and B. Undrakh. Rational dilation problems associated with constrained algebras. *J. Math. Anal. Appl.*, 467(1):95–131, 2018.
- [20] R. Exel and T. A. Loring. Finite-dimensional representations of free product C^* -algebras. *International Journal of Mathematics*, 3(04):469–476, 1992.
- [21] M. Hamana. Injective envelopes of C^* -algebras. *Journal of the Mathematical Society of Japan*, 31(1):181–197, 1979.
- [22] A. Hopenwasser. Boundary representations on C^* -algebras with matrix units. *Trans. Amer. Math. Soc.*, 177:483–490, 1973.
- [23] R. A. Horn and C. R. Johnson. *Topics in matrix analysis*. Cambridge university press, 1994.
- [24] M. T. Jury, G. Knese, and S. McCullough. Agler interpolation families of kernels. *Oper. Matrices*, 3(4):571–587, 2009.
- [25] E. Kirchberg and S. Wassermann. C^* -algebras generated by operator systems. *J. Funct. Anal.*, 155(2):324–351, 1998.
- [26] K. McClanahan. C^* -algebras generated by elements of a unitary matrix. *Journal of functional analysis*, 107(2):439–457, 1992.
- [27] S. McCullough. Isometric representations of some quotients of H^∞ of an annulus. *Integral Equations and Operator Theory*, 39(3):335–362, 2001.
- [28] S. McCullough and V. Paulsen. C^* -envelopes and interpolation theory. *Indiana University mathematics journal*, pages 479–505, 2002.
- [29] P. S. Muhly and B. Solel. An algebraic characterization of boundary representations. In *Nonselfadjoint operator algebras, operator theory, and related topics*, volume 104 of *Oper. Theory Adv. Appl.*, pages 189–196. Birkhäuser, Basel, 1998.
- [30] B. S. Nagy, C. Foias, H. Bercovici, and L. Kérchy. *Harmonic analysis of operators on Hilbert space*. Springer Science & Business Media, 2010.
- [31] G. Pick. Über die beschränkungen analytischer Funktionen, welche durch vorgegebene Funktionswerte bewirkt werden. *Mathematische Annalen*, 77(1):7–23, 1915.
- [32] M. Raghupathi. Nevanlinna-Pick interpolation for $\mathbb{C}+ BH^\infty$. *Integral Equa. Oper. Theory*, 63:103–125, 2009.
- [33] D. Sarason. On spectral sets having connected complement. *Acta Sci. Math. (Szeged)*, 26:289–299, 1965.
- [34] D. Sarason. Generalized interpolation in H^∞ . *Transactions of the American Mathematical Society*, 127(2):179–203, 1967.
- [35] J. P. Solazzo. *Interpolation and computability*. University of Houston, 2000.
- [36] B. Sz.-Nagy and A. Korányi. Operatortheoretische Behandlung und Verallgemeinerung eines Problemkreises in der komplexen Funktionentheorie. *Acta Math.*, 100:171–202, 1958.

DEPARTMENT OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, BEER-SHEVA 8410501, ISRAEL

DEPARTMENT OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, BEER-SHEVA 8410501, ISRAEL