Below the Schwinger critical magnetic field value, quantum vacuum and gamma-ray bursts delay

Iver H. Brevik¹, Moshe M. Chaichian² and Anca Tureanu²

¹Department of Energy and Process Engineering,

Norwegian University of Science and Technology, N-7491 Trondheim, Norway ²Department of Physics, University of Helsinki,

and Helsinki Institute of Physics, P. O. Box 64, FI-00014 Helsinki, Finland

Abstract

A magnetic field above the Schwinger critical value $B_{\rm crit} = 10^9$ Tesla is much higher than any magnetic field known by now in the interstellar bulk except in the vicinity of observed magnetars having magnetic fields between 10^9 and 10^{11} Tesla. Above the critical magnetic field limit, calculated by Schwinger in the lowest order perturbation in quantum electrodynamics (QED), one reaches the threshold for electron-positron pair creation (through the intermediate electric field, as known also from standard electrodynamics), which has interesting consequences. Therefore, finding out whether one could encounter some consequences of interest also for the values of the magnetic field below the Schwinger critical point, we invoke the next higher-order effect in QED, which is emerging from the *Quantum Vacuum Effect*. The latter is equivalent to the use of the Euler-Heisenberg effective theory in nonlinear electrodynamics, where the Lagrangian has now a term with a higher power, B^4 . In this case, in the region $B < B_{crit}$, we show that interesting effects appear, among them the Cherenkov radiation and the reduction in the speed of light. The latter effects appear because of the quantum vacuum mimicking a medium. We also present quantitative arguments for such a close analogy. As a rough estimate, we show that the time delay τ of gamma-ray bursts (GRB) having traveled through the entire cosmological distances in an average strong magnetic field such as 10^6 Tesla, reaches an experimentally considerable value of $\tau = 2.4$ hours. Of course in the vicinity of magnetars, the magnetic field is much stronger, of the order of $10^9 - 10^{11}$ Tesla. However, in this case the linear scale of GRB trajectory through such regions would be much smaller. For the latter, we also give a corresponding estimate for the number of the magnetars along the trajectory and also for the delay. Finally, we shall dwell on the recently raised issue in the literature, namely the Lorentz invariance violation (LIV).

I. INTRODUCTION

As is known, when an electromagnetic field in vacuum is exposed to an external strong static magnetic field \mathbf{B}_0 , the equations of motion derived from the generalized high-intensity action integral become formally analogous to the conventional Maxwell equations in a continuous medium [1]. Then it becomes natural to also introduce nondispersive dielectric constants ε_{ik} and μ_{ik} , which in turn lead to the refractive index components, greater than unity. The Schwinger critical limit $B_{\rm crit}$ [2] occurs when the magnetic field B_0 becomes so strong that electron-positron pair production occurs. The value $B_{\rm crit} = 10^9$ Tesla is far beyond what is achievable on Earth, while it is known to be surpassed in the vicinity of astrophysical objects such as magnetars, first announced in 2008 [3]. It is of interest to investigate also the case for the more accessible lower magnetic fields $B < B_{\rm crit}$, and to explore the implications of this kind of medium-like vacuum, appropriately called the *quantum vacuum effect*.

From a fundamental viewpoint, one notices that a constant magnetic field alone will not be able to produce electron-positron pairs. One needs an electric field as an intermediate factor. This is essentially the same kind of behaviour as encountered in standard electrodynamics – an electric field varying in space is needed so as to get a magnetic field developing in time, according to Maxwell's equations.

As mentioned above, the effect is related to the Euler-Heisenberg theory of nonlinear electrodynamics [4], derived from the higher-order corrections in QED. An extensive study of the effect from a quantum perspective was given by Adler [5]. The propagation of photons in an external field and photon splitting were considered in Refs. [6] and [7]. Reference [8] considered, from a geometric optics approximation, various nontrivial QED vacua. From an inflationary viewpoint, the possibility of generation of the origin of primordial magnetic field was discussed in Ref. [9]. As well, the study of the Quantum Magnetic Collapse in a strong magnetic field of the order of the critical magnetic fields has been performed in [10], with possible implications of the results for astroparticle physics and cosmology. Also there exist some works on the effect of homogeneous magnetic fields on the refractive index [11, 12]. In the latter works two different cases were considered: one in which high-energy photons propagate in fields weak compared with the critical field, and one where low-energy photons propagate in fields with arbitrary intensity.

In connection with the nonlinear electrodynamics effects, in [13] and [14], respectively the photon polarization and the vacuum birefringence have been considered in detail, while in [15] the polarization selection rules that take place in the photon splitting in a medium (i.e. in the presence of a chemical potential) were discussed.

There also exist studies of the supercritical limit i.e., when $B_0 > B_{\text{crit}}$ [16, 17]. An earlier work of the same group of authors had studied the case when there were both a constant electric and a constant magnetic field [18].

In the following we shall first give an overview of the electromagnetic properties of this

kind of magnetic-generated medium in the case of a subcritical magnetic field $B_0 < B_{crit}$, and will thereafter focus on some new aspects not considered previously in the literature. Specifically:

i) Considering that photons propagating in a medium are slowed down to the velocity c/n (here assuming that the medium is isotropic), one may ask: how large must B_0 be in order to make the effect observable? How does the photon velocity reduction, if calculated on the basis of the mean magnetic field in intergalactic space, compare with the dispersive slowing down of the velocity of photons in the electron plasma, as considered in [19, 20]?

ii) As a rough estimate, we can take the average magnetic field to be 10^6 Tesla, which although seems to be high, nevertheless is still well below the Schwinger critical limit, extending over a large cosmic distance. In such a case, the delay of GRB, τ , comes out to be large, estimated of the order of $\tau = 2.4$ hours. Therefore, when two species of particles, namely photons and neutrinos, are traveling through the interstellar distances, photons arrive later by a time delay τ compared with neutrinos, which have no interaction with the magnetic field. However, as mentioned in the Abstract, in the vicinity of magnetars magnetic field is much stronger, of the order of $10^9 - 10^{11}$ Tesla. With such values of the magnetic fields we also give an estimate for the delay, considering a realistic case with the number of magnetars along the trajectory passed by the GRB in the Subsection B of Section III. Actually several earlier observations such as [27, 28] and references therein led to the idea of a Lorentz invariance violation (LIV), assuming that the two different species of neutrino and GRB were emitted from the same source and at the same time. The important point, however, is that in those observations there was no assurance of any kind in their accuracy of measurements that the two species were emitted from the same source, not speaking of their simultaneous emission. In addition to all that, the measured delays were quite inaccurate. Still those observations gave boost to a possible LIV. Concerning the claimed observation of LIV based on different delays depending on the energy spectrum of emitted GRB, and the implication of the present work, we present a full detail in the Section IV, where we give arguments that there has been no evidence for the LIV extracted from the GRB spectrum. *iii*) Since the theoretical value of n is greater than one, the possibility of a Cherenkov radiation becomes of immediate interest. Under terrestrial conditions, such an effect appears to be far too small to be observed. But what can be said about the specific case of *atmospheric* constellation, and whether there is any situation, where such an effect as the Cherenkov radiation can be detected? Apparently, there are only a few works in the literature discussing such a possibility (see the recent paper [21] and references therein).

In the next section, we give an overview of the general theory describing in the subcritical case, where the magnetic field is lower than the critical value. In this work we use the Gaussian units, mostly with $\hbar = c = 1$, and follow the notation of Ref [1]. Finally, we present several new results and discuss the relation to their observability.

II. BASICS

Let the total magnetic field in a homogeneous region of outer space be denoted by **B**. It is composed of the strong static field \mathbf{B}_0 and a weak time-varying part **B'** associated with the traveling electromagnetic waves. No dielectric media are assumed present. The Schwinger critical limit is

$$B_{\rm crit} = \frac{m^2 c^3}{|e|\hbar} = 4.41 \times 10^9 \text{ Tesla} = 4.41 \times 10^{13} \text{ Gauss.}$$
(1)

In this formula, m is the electron mass and e is the electron charge. We re-emphasize that, while the critical field is far too strong to be producible on Earth, it is significantly exceeded in magnetars $(10^9 - 10^{11} \text{ Tesla})^1$.

It is in principle straightforward to consider a strong electric field \mathbf{E} in the same way. The critical value becomes then

$$E_{\rm crit} = \frac{m^2 c^3}{|e|\hbar} = 1,32 \times 10^{18} \text{ Volt/m.}$$
(2)

We will be concerned with low-order nonlinear theory, meaning that the fields are obtained from first order expansions in the parameter $(B_0/B_{\rm crit})^2$. We start from the perturbed Lagrangian density L', derived from the original (unperturbed) Maxwell Lagrangian density L. In the general case [1],

$$L' = \frac{e^4}{360\pi^2 m^4} [(E^2 - H^2)^2 + 7(\mathbf{E} \cdot \mathbf{H})^2].$$
 (3)

¹ If one would consider axion electrodynamics (cf., for instance, Ref. [22]), the form of $B_{\rm crit}$ would be different from Eq. (1) and it could happen that the value of $B_{\rm crit}$ would drastically change and even become much reduced. The case with axion electrodynamics is of interest to be investigated further. The latter can be of interest also due to an excess in the amount of axions around the magnetars as recently argued in [23].

In the following we shall assume that there is no external electric field, i.e. $\mathbf{E} = 0$. Thus,

$$L' = \frac{e^4}{360\pi^2 m^4} H^4,$$
(4)

which enables us to calculate the magnetization \mathbf{M} in the medium by using the general formula $\mathbf{M} = \partial L' / \partial \mathbf{H}$,

$$M = \frac{e^4}{90\pi^2 m^4} H^3.$$
 (5)

Now one has $\mathbf{B} = \mu \mathbf{H}$ where μ is the permeability. Introducing the susceptibility χ so that $\mathbf{M} = \chi \mathbf{H}$, we have in Gaussian units

$$\chi = \frac{\mu - 1}{4\pi},\tag{6}$$

leading to

$$\mu = 1 + \frac{2e^4}{45\pi m^4} H^2. \tag{7}$$

Along with the permeability μ , the general formalism introduces also a constant permittivity ε in the same way; cf. Ref. [1]. These two material parameters have thus the same fundamental basis. (The details of the derivation are given below, see formulas (10)-(17).) The corresponding expression for the permittivity ε is

$$\varepsilon = 1 + \frac{5e^4}{45\pi m^4} H^2. \tag{8}$$

Thus, the refractive index n becomes

$$n = \sqrt{\varepsilon\mu} = 1 + \frac{7e^4}{90\pi m^4} H^2.$$
(9)

Note: this simple calculation rests upon the assumption that it is physically meaningful to consider the medium as isotropic, possessing scalar expressions for ε and μ . This is not evident beforehand, but will be justified by a more detailed consideration below. The main aim of the calculation is to show that the two constants of the matter, μ and ε are quadratically dependent on the applied magnetic field².

We now consider the physical situation in more detail, focusing on the weak fields \mathbf{E}', \mathbf{B}' associated with propagating waves in the strong magnetic field background. With the decomposition

$$\mathbf{B} = \mathbf{B}_0 + \mathbf{B}', \quad \mathbf{E} = \mathbf{E}', \tag{10}$$

² There is an extra π in the denominators of the above formulas, compared with those given in Ref. [1].

assuming a common time factor $e^{-i\omega t}$, we then obtain from Maxwell's equations

$$\mathbf{k} \times \mathbf{H}' = -\omega \mathbf{D}', \quad \mathbf{k} \times \mathbf{E}' = \omega \mathbf{B}', \tag{11}$$

$$\mathbf{k} \cdot \mathbf{B}' = 0, \quad \mathbf{k} \cdot \mathbf{D}' = 0, \tag{12}$$

corresponding to the constitutive relations for an anisotropic medium,

$$D'_{i} = \varepsilon_{ik} E'_{k}, \quad B'_{i} = \mu_{ik} H'_{k}. \tag{13}$$

Explicitly, the matter's constants can be expressed as [1]

$$\varepsilon_{ik} = \delta_{ik} + \frac{2e^4}{45m^4} B_0^2 (-\delta_{ik} + \frac{7}{2}b_i b_k), \tag{14}$$

$$\mu_{ik} = \delta_{ik} + \frac{2e^4}{45m^4} B_0^2 (\delta_{ik} + 2b_i b_k), \tag{15}$$

where $\mathbf{b} = \mathbf{B}_0/B_0$ is the unit vector in the direction of the applied field. We can express these equations also in terms of the critical field,

$$\varepsilon_{ik} = \delta_{ik} + \frac{2e^2}{45} \frac{B_0^2}{B_{\text{crit}}^2} (-\delta_{ik} + \frac{7}{2} b_i b_k), \tag{16}$$

$$\mu_{ik} = \delta_{ik} + \frac{2e^2}{45} \frac{B_0^2}{B_{\text{crit}}^2} \left(\delta_{ik} + 2b_i b_k \right).$$
(17)

In Gaussian units, $e^2 = 1/137^3$.

As mentioned above, we limit ourselves to the case of weak nonlinearity, meaning that the condition for the use of Eqs. (16) and (17) is $B_0/B_{\rm crit} \ll 1$. The effective magnetic expansion parameter is [1]

$$eH/m^2 \ll 1,\tag{18}$$

with H the total magnetic field. In practice, H can here be replaced by B_0 .

We may distinguish between two cases of linear polarization of a propagating plane wave, taking into account the special nature of the **bk** plane. We shall however focus mainly on the perpendicular case, designated by a subscript \perp , where **B**' is perpendicular to the

³ There is a debate in the literature (see [24]) concerning the numerical factor in the second term within the brackets of Eq. (15), implying a numerical change compared with the corresponding values given in [1]. However, such minor numerical change in no way would alter the main results presented in our work.

We thank Yuri Obukhov for bringing this point to our attention.

mentioned plane. For simplicity we take \mathbf{B}_0 in the *x* direction, and the wave vector \mathbf{k} in the *z* direction. For later convenience we define the symbol *d* as

$$d = \frac{7e^4}{90m^4} B_0^2 = \frac{7e^2}{90} \frac{B_0^2}{B_{\rm crit}^2}.$$
 (19)

Therefore, from Eqs. (16) and (17), we obtain

$$\varepsilon_{xx} = 1 + \frac{10d}{7}, \quad \varepsilon_{yy} = \varepsilon_{zz} = 1 - \frac{4d}{7},$$
(20)

$$\mu_{xx} = 1 + \frac{12d}{7}, \quad \mu_{yy} = \mu_{zz} = 1 + \frac{4d}{7},$$
(21)

the nondiagonal components ε_{ik} and μ_{ik} being equal to zero.

Here the most significant constitutive relation is

$$B'_{y} = \mu_{yy}H'_{y} = \left(1 - \frac{4d}{7}\right)H'_{y},$$
(22)

since it corresponds to the real wave for B'_y . We see that μ_{yy} is equal to the permeability μ in the isotropic model above, Eq. (7), to the approximation considered. This justifies the introduction of the isotropic model. Once μ_{yy} is fixed, the other components ε_{ik} and μ_{ik} follow from Maxwell's equations.

With $k = n_{\perp}\omega$, n_{\perp} being the refractive index for perpendicular polarization we thus have, with $n_{\perp} = \sqrt{\varepsilon \mu}$,

$$n_{\perp} = 1 + d = 1 + \frac{7e^4}{90m^4}B_0^2 = 1 + \frac{7e^2}{90}\frac{B_0^2}{B_{\rm crit}^2}.$$
(23)

In the same way one can analyze the case of parallel polarization, subscript \parallel as mentioned. This case leads to a refractive index n_{\parallel} , which is smaller than n_{\perp} , and will not be further considered here.

We re-emphasize that the present dielectric theory, coming from the quantum vacuum, is nondispersive, i.e there is no frequency dependence in the matter's constants and thus in the refraction index too.

III. APPLICATIONS

We shall now illustrate the use of the above formalism, under various circumstances.

A. Reduction of the photon velocity in the medium (analogous to the works in [19, 20]). The weak magnetic field case

As is the case in all dielectric media, the propagation velocity is lower than in a vacuum. We exploit our earlier definition of the parameter d in Eq. (19) to write the velocity in the simple form

$$v = c/n = c(1 - d).$$
 (24)

Numerically, from Eq. (19) we get, inserting B_{crit} from Eq. (1),

$$d = 2.9 \times 10^{-31} \times B_0^2, \tag{25}$$

where B_0 is in gauss (G) units. Taking $B_0 = 100 \ \mu G$ to be representative for intergalactic space, we obtain

$$v = c(1 - 2.9 \times 10^{-39}). \tag{26}$$

This illustrates how extremely small is the reduction of the light velocity under these conditions. However, if an appreciable light reduction is to be achieved, one has to focus on local astrophysical regions where B_0 is very strong. It is to be noted that even with a field of $B_0 = 10^{10}$ G, very strong according to normal standards, we are still far below the critical limit (1).

Related to this point, it is of interest also to compare this case with the results one gets by looking at how much photons are slowed down because of the electron plasma in the interstellar space, i.e. without having a magnetic field. This means to compare with the velocity reduction of gamma-rays due to the conventional dispersive theory. We take the electrons with mass m and their mean density in the interstellar space to be N.

We start with the dispersive relation for photons in an electron(-positron) plasma,

$$\omega^2 = k^2 + \omega_p^2,\tag{27}$$

where ω_p is the plasma frequency determined by

$$\omega_p^2 = \frac{4\pi N e^2}{m}.\tag{28}$$

The system is thus dispersive, and we consider the group velocity

$$v_g = \frac{d\omega}{dk} = c \left(1 - \frac{\omega_p^2}{2\omega^2} \right).$$
⁽²⁹⁾

Here we insert the electron mass m = 0.511 MeV, and, similarly as in Ref. [20], focus attention on galaxy filaments at around z = 0.1, for which the electron density is about $N = 4 \times 10^{-4}$ cm⁻³. In view of the conversion factor 1 cm = 5.068×10^{13} GeV⁻¹ this means $N = 3.1 \times 10^{-45}$ GeV³ so that $\omega_p^2 = 5.5 \times 10^{-43}$ GeV² = 1.2×10^6 s⁻². Thus, for a gamma-ray with energy E = 100 GeV⁴ we obtain

$$v_q = c(1 - 3 \times 10^{-47}),\tag{31}$$

which shows an even smaller velocity reduction than the previous expression (26).

B. Reduction of the photon velocity in the medium in the presence of a strong magnetic field. The delay of photon detection time

Let us assume now an extreme case when the GRB photons travel over large intergalactic distances, experiencing strong magnetic fields all the time, before they each us, i.e. the detectors on the Earth. For definiteness, we choose an average value for the magnetic field as mentioned above, i.e.

$$B_0 = 10^{10} \text{ G}, \tag{32}$$

⁴ In order not to violate the parameter regime where the Heisenberg–Euler Lagrangian as the leading-order in the perturbative expansion can be reliably applied, besides the condition (18) $eB/m^2 \ll 1$, there is still another criterion [14], namely the condition

$$\frac{eB\omega}{m^3} \ll 1. \tag{30}$$

With a value of E = 100 GeV for the energy of GRB, one gets $eB\omega/m^3 = 44.4$, which is clearly not smaller than unity. However, with lower energies, such as E = 1 GeV, one can trust the leading term in the perturbative expansion as performed in the present work, and the qualitative results coming from them. As a result, for the GRB energy E = 1 GeV, the delay times calculated for E = 100 GeV, i.e. Eq. (35) and Eq. (36), will be reduced by the corresponding factor of 100.

Since the main result of the work is to show that there is no LIV derived from GRB delays, contrary to some opposite claims in the literature, the nondispersive character, emerging as a by-product of calculations and mentioned in the work, is not of importance for our arguments, which are based entirely on the emission of different frequencies of light from their source occurring NOT simultaneously. and make a crucial assumption that this magnetic field is constant over the *entire* photon trajectory. At the end of this Subsection we present a realistic case, where the trajectory of photon is considered to pass only through the vicinity of the magnetars, with an estimate on their number in the interstellar space. As compared with the value 100 μ G used in the previous subsection we are thus increasing B_0 by a factor of 10^{14} . We obtain now

$$v = c(1 - 2.9 \times 10^{-11}). \tag{33}$$

This number is actually of considerable physical interest: what is the *time delay* of a gammaray burst having traveled a large intergalactic distance in such a strong field before it reaches us? Denoting the effective distance by D we obtain for the delay time τ , omitting for simplicity the redshift correction,

$$\tau = \frac{D \times d}{c(1-d)} \approx \frac{D \times d}{c}.$$
(34)

As in Ref. [20] we choose for D the large value 3 Gpc = 9×10^{25} m, whereby

$$\tau = 8.7 \times 10^6 \text{ s} = 2.4 \text{ hours.}$$
 (35)

As mentioned before, at the point ii) above, the delay time τ in Eq. (35) is of considerable importance to explain the delay of the GRB compared with the neutrino bursts, without invoking the LIV.

The assumption made above on the choice of $B_0 = 10^6$ Tesla as an average cosmological magnetic field in the *entire* interstellar space, seems to be too strong. However, there are regions where still stronger magnetic fields, of the order of 10^9 to 10^{11} Tesla exist in the vicinity of magnetars, but the linear scale of such regions would be much smaller, around magnetars not exceeding some 10^9 cm or so. Assuming now that the mean separation between galaxies is about 1 Mpc = 3×10^{24} cm, then there could be about 10^4 galaxies along the line of sight, passing the entire interstellar distance of 3 Gpc = 10^{28} cm.

In such a case, instead of the time delay as $\tau = 8.7 \times 10^6$ s = 2.4 hours according to Eq. (35), we shall have a time delay to be anything between the values such as

$$\tau = 8.7 \times (10^2 - 10^{18})$$
 s (36)

according to the Eqs. (25) and (34). In the estimate (36) we have also taken into account that within the sphere around each magnetar, in average the magnetic field value is reduced

by a factor of 10^2 from $B_0 = 10^9 - 10^{11}$ Tesla. We are aware of the fact that the range of values given in (36) as compared with the neutrino bursts, are far from being realistic due to the unknown number of magnetars in the galactic space and as well the value of magnetic fields around them⁵. But the observations in comparing the neutrino with GRB bursts are also not accurate and conclusive by any standard all.

Let us also mention that the delay of neutrinos traveling through the entire galactic space is fully negligible, as shown in [19, 20] taking into account the masses of the three species and their oscillations.

C. Remarks on the Cherenkov effect

The fact that n > 1 in the nonlinear theory makes it natural, as mentioned above, to investigate the possibility of observing a Cherenkov effect although the smallness of the medium susceptibility necessarily imposes strong restrictions on the particle velocity. To our knowledge there are only a few earlier investigations along these lines, although we may mention Refs. [21, 25], and also the older review article Ref. [26]. It becomes now natural to consider protons instead of electrons as the fundamental particles. One may note that with the proton mass $m_p = 938$ MeV the critical magnetic field, Eq. (1), becomes enhanced by a factor $(m_p/m)^2$.

It is obvious that we have to look for extreme events in the intergalactic space. Cosmic radiation in the form of protons has been observed up to about 3×10^{20} eV. For definiteness, let us consider the "window" where the proton energy is

$$E > 10^{15} \text{ eV},$$
 (37)

and investigate under which conditions Cherenkov radiation in this energy interval can occur. A proton with energy $E = 10^{15}$ eV in rectilinear uniform motion has a velocity β equal to

$$\beta = c \left[1 - \frac{1}{2} \left(\frac{m_p}{E} \right)^2 \right] = c (1 - 4.4 \times 10^{-13}).$$
(38)

As before, we consider the case of perpendicular polarization, where the component B'_y propagates in the z direction in the nonlinear medium created by the x directed magnetic

⁵ For a celestial experiment, one might think of a Fizeau-type setup, in which the whole system is placed in a constant strong magnetic field to observe the reduction in the light velocity.

field B_0 . The threshold for the Cherenkov effect is

$$n_{\perp}\beta = 1. \tag{39}$$

Thus the threshold corresponds to the perpendicular refractive index

$$n_{\perp} = \frac{1}{\beta} = 1 + 4.4 \times 10^{-13},\tag{40}$$

where use is made of Eq. (38). We compare this with the equation (23), i.e. with

$$n_{\perp} = 1 + \frac{7e^4}{90m^4} B_0^2, \tag{41}$$

where the mass of the electrons m enters, because the medium, as before, is the quantum electrodynamics vacuum, which gives the dominant contribution as compared to the contributions coming from the other constituents present in the quantum vacuum. Then, from Eq. (41), we derive for the magnetic field B_0 the value:

$$B_0 = 4.5 \times 10^{-14} \text{ GeV}^2. \tag{42}$$

Using the conversion factors 1 GeV= 1.602×10^{-3} erg, 1 cm = 5.068×10^{13} GeV⁻¹, we have the following relationship between the energy densities in the natural and the Gaussian system of units,

$$\text{GeV}^4 = 2.08 \times 10^{38} \, \frac{\text{erg}}{\text{cm}^3}.$$
 (43)

Since the magnetic energy density in Gaussian units is $B_0^2/(8\pi)$, we obtain the magnetic threshold value to be

$$B_0 = 6.5 \times 10^5 \text{ G.} \tag{44}$$

This appears to be roughly as we might expect: this value is less than the critical field even for electrons (cf. Eq. (1)), and is much less than the magnetic field observed around the magnetars. So protons with energies in the mentioned energy "window" will in principle be able to initiate Cherenkov radiation if they pass through regions with so strong magnetic fields.

Finally, we give the Tamm-Frank seminal formula for the emitted Cherenkov radiation from a particle with charge e moving with velocity v along the x axis, in Gaussian units,

$$\frac{dE_{ch}}{dx} = \frac{e^2}{4\pi} \int_{\omega > c/n(\omega)} \mu(\omega)\omega\left(1 - \frac{c^2}{v^2 n^2(\omega)}\right) d\omega,\tag{45}$$

where $\mu(\omega)$ is the magnetic permeability of the medium, which is now given by the Eqs. (19) and (21). This formula can be used for a possible experimental detection of the Cherenkov radiation.

IV. DISCUSSION AND CONCLUSIONS

The quantum vacuum mimicking a medium carries over to all the characteristic properties of conventional electrodynamics of media, among which the character of the photon momentum and also the Cherenkov effect stand out as typical examples. As noted before, the theory becomes nondispersive in the present formulation. Physically, this is a consequence of the external magnetic field \mathbf{B}_0 being treated as a static one. More general theories, where the magnetic field \mathbf{B} (as well as an electric field \mathbf{E}) is time-dependent, would imply dispersive properties also in the magnetically-generated refractive index n.

Concerning the implication of the present work on Lorentz invariance violation (LIV), we would like to mention the following:

1. As we have shown, taking as an example the average magnetic field through which the gamma-ray burst has traveled through the cosmological distances to be 10^6 Tesla, then they delay compared with the neutrinos to reach the Earth by a time in the range of $\tau = 8.7 \times (10^2 - 10^{18})$ seconds. Although a value as 10^6 Tesla may look high, nevertheless it is well below the Schwinger critical limit. Such a delay is well compatible with the delays observed by ANTARES neutrino telescope [27] or by the IceCube data [28].

The observed time delays had originally been taken as an indication for possible LIV, interpreting the reduction in the speed of light to be the breaking of Special Relativity at high energies. Notice, however, that in the interpretation of those observations, it has been assumed that the GRB and neutrino bursts have been emitted simultaneously, though it is even not known whether the GRB and the neutrino bursts have come from the same source or not.

Therefore, the results presented in this Letter put doubt on the claim that the GRB delay compared with the neutrino bursts is a sign of LIV, since the delay can well be accounted for by the standard physics, namely by the dispersion of light in a strong magnetic field found in the interstellar space due to the existence of magnetars, as shown in the present work.

2. In the previous works [19] and [20], the dispersion of light and its delay were carefully studied by considering the effects of different media, electron-positron, CMB and the axion plasmas, existing within the cosmic distances. The results in [19] and [20] showed that, while the time delay of GRB by the CMB medium was the largest compared with the other two media, still it was by far too small to account for the delays of GRB observed in [27] and [28].

3. According to the dispersion relation in matter-filled media studied in [19] and [20], as well as in any dispersive medium, the more energetic the GRB, the less the delay - in other words, the more energetic GRB reach us earlier than the less energetic ones. Contrary to that, the strong magnetic field considered in the present Letter produces out of the quantum vacuum a nondispersive medium, which does not depend on the energy of light and thus the refraction index depends only on the value of the external magnetic field.

4. Irrespective of the experimental observations of delay in GRB and also inspired by the results such as in [27] and [28], there has been a remarkable activity, both on the theoretical and phenomenological possible occurrence of LIV at high energies. As a noncomprehesive list of references we mention [29–36], and references therein.

We notice that in all the so-far proposed theoretical models, such as in a stringy spacetime foam or in phenomenological models with some deformation of the usual dispersion relation $E^2 = \mathbf{p}^2$, and as well in the interpretation of the observations as arguments for LIV, either the space is not taken to be vacuum, but filled with some matter (i.e. the induced LIV), or the assumption of a simultaneous emission of particles with different energies in the gamma-ray bursts is used: For instance, typically in the LIV approach motivated by quantum gravity effects (which assumes the presence of gravity), the dispersion relation contains higher-powers in energy E, such as:

$$E^{2}\left[1 + \xi \frac{E}{E_{QG}} + O\left(\frac{E^{2}}{E_{QG}^{2}}\right)\right] = \mathbf{p}^{2}c^{2} + m^{2}c^{4},$$
(46)

where E_{QG} is an effective energy scale for quantum gravity, commonly taken to be of the order 10¹⁶ GeV, and ξ is an arbitrary parameter. Then, with a modified dispersion relation such as (46), in the leading order in E/E_{QG} , the group velocity becomes:

$$v_g = c \left(1 - \xi \frac{E}{E_{QG}} \right). \tag{47}$$

Other phenomenological models used for LIV based on the quantum deformation of the Lorentz or Poincaré groups lead to the same effect as (47) does, i.e. increasing the gamma-ray energy makes the velocity smaller.

According to (47), the more energetic the GRB, the later they reach the Earth. At the same time the analyses of GRB time delay as a function of their energy also show that the higher energy gamma-rays arrive later. Now, if one assumes that the GRB with all

different energies have been emitted by a single source simultaneously, then of course one can attribute such a observation to a LIV, which is also in accord with (47).

But could one justify such an assumption that all the GRB with different energies have been emitted by a single source *simultaneously*?

We do not have any clue about the dynamics of bursts and the emission of particles from their sources, as they are truly complicated. But what we do know from the other sources, such as from the 11-Year Solar Cycle and also from the atomic explosions, is that the lower energy particles are emitted first, while the higher energy ones later. The latter phenomenon can be naturally explained when one acknowledges that the temperatures in such explosive sources are much higher in their centre and also that the diameters of such sources are usually huge. This puts doubt on the assumption of simultaneous emission, and consequently on claiming the observation of Lorentz invariance violation based on the GRB time delays.

Our conclusion is that there is no evidence for the Lorentz invariance violation, i.e. for the basic assumption of the Special Relativity in the vacuum⁶ – all the data can be explained by the standard physics, unless one assumes the existence of some kind of matters in the space and invokes the use of an unjustified assumption.

ACKNOWLEDGEMENTS

We are most grateful to Efrain Ferrer, Kimball Milton, Viatcheslav Mukhanov, Olivier Piguet, Adam Schwimmer, Jenny Wagner and in particular to Felix Karbstein, Yuri Obukhov and Misao Sasaki for several illuminating discussions and suggestions.

 V. B. Berestetskii, E. M. Lifshitz and L. P. Pitaevskii, *Quantum Electrodynamics* (Pergamon Press, Oxford, 1982), Secs. 129,130.

⁶ We should keep in mind that, if the Lorentz invariance of the Special Relativity is violated, then it automatically implies that the diffeormorphism invariance of the General Relativity is also broken. Therefore, the Eintein gravitational theory and its modifications should be accordingly changed too, and only the latter theories with diffeormorphism symmetry broken should be used in all the applications.

- [2] J. Schwinger, On gauge invariance and vacuum polarization. Phys. Rev. 82, 664 (1951).
- [3] European Southern Observatory, 23 September 2008.
- [4] W. Heisenberg and H. Euler, Consequences of Dirac theory of the positron (translated). Zeitschr. Physik 98, 714 (1936).
- [5] S. L. Adler, Photon splitting and photon dispersion in a strong magnetic field. Annals of Physics 67, 599 (1971).
- [6] Z. Bialynicki-Birula and I. Bialynicki-Birula, Nonlinnear effects in quantum electrodynamics. Photon propagation and photon splitting in an external field. Phys. Rev. D 2, 2341 (1970).
- [7] J. E. Brezin and C. Itzykson, Polarization phenomena in vacuum nonlinear electrodynamics. Phys. Rev. D 3, 618 (1971).
- [8] W. Dittrich and H. Gies, Light propagation in nontrivial QED vacua. Phys. Rev. D 58, 025004 (1998).
- [9] V. Demozzi, V. Mukhanov and H. Rubinstein, Magnetic field from inflation? JCAP 08 (2009) 025.
- M. Chaichian, S. S. Masood, C. Montonen, A. Perez Martinez and H. Perez Rojas, Quantum magnetic and gravitational collapse, Phys. Rev. Lett. 84 (2000), 5261 [arXiv:hep-ph/9911218 [hep-ph]].
- [11] W.-Y.Tsai and T. Erber, Photon pair creation in intense magnetic fields. Phys. Rev. D 10, 492 (1974).
- [12] W.-Y.Tsai and T. Erber, Propagation of photons in homogeneous magnetic fields: Index of refraction. Phys. Rev. D 12, 1132 (1975).
- F. Karbstein, Photon polarization tensor in a homogeneous magnetic or electric field, Phys. Rev. D 88 (2013) 8, 085033, arXiv:1308.6184 [hep-th].
- [14] F. Karbstein, Vacuum Birefringence at the Gamma Factory, Annalen Phys. 534 (2022) 3, 2100137, arXiv: 2106.06359 [hep-ph].
- [15] E. J. Ferrer, V. de la Incera and A. F. Shabad, Covariant decomposition and polarizational selection rules for interaction among photons in a moving medium. Forthscr. Physik 32, 261 (1984).
- [16] A. E. Shabad, Photon propagation in a supercritical magnetic field. J. Exp. Theor. Phys. 98, 186 (2004).

- [17] F. Karbstein, All-Loop Result for the Strong Magnetic Field Limit of the Heisenberg-Euler Effective Lagrangian, Phys.Rev.Lett. 122 (2019) 21, 211602, arXiv:1903.06998 [hep-th]; Erratum Phys. Rev. Lett. 133, 199902 (2024).
- [18] I. A. Batalin and A. E. Shabad, Green's function of a photon in a constant homogeneous electromagnetic field of general form. Soviet Physics JETP 33, 483 (1971).
- [19] M. Chaichian, I. Brevik and M. Oksanen, Can the gamma-ray bursts traveling through the interstellar space be explained without invoking the drastic assumption of Lorentz invariance violation?, 40th Int. Conf. on High Energy Phys (ICHEP 2020). Proceedings of Science, Vol. 390, 2021, arXiv:2101.05758 [astro-ph.HE].
- [20] I. Brevik, M. Chaichian and M. Oksanen, Dispersion of light traveling through the interstellar space, induced and intrinsic Lorentz invariance violation, Eur. Phys. J. C 81, 926 (2021).
- [21] A. J. Macleod, A. Noble and D. A. Jaroszynski, Cherenkov radiation from the quantum vacuum. Phys. Rev. Lett. 122, 161601 (2019).
- [22] P. Sikivie, Invisible axion search methods, Rev. Mod. Phys. **93**, 015004 (2021).
- [23] D. Noordhuis, A. Prabhu, C. Weniger and S. J. Witte, Axion Clouds around Neutron Stars, Phys. Rev. X 14, 041015 (2024).
- [24] J. S. Heyl and L. Hernquist, Birefringence and dichroism of the QED vacuum, J. Phys. A: Math. Gen. 30 6485 (1997).
- [25] C.-Y. Lee, Cherenkov radiation in a strong magnetic field. Phys. Lett. B 810, 135794 (2020).
- [26] T. Erber, High-energy electromagnetic conversion processes in intense magnetic fields. Rev. Mod. Phys. 38, 626 (1966).
- [27] S. Adrian-Martinez et al. [ANTARES], Stacked search for time shifted high energy neutrinos from gamma ray bursts with the ANTARES neutrino telescope, Eur. Phys. J. C 77, 20 (2017) [arXiv:1608.08840 [astro-ph.HE]].
- [28] M. G. Aartsen et al. IceCube, Search for High-Energy Neutrino Emission from TeV Pulsar Wind Nebulae, Astrophys. J. 898, 117 (2020) [arXiv:2003.12071 [astro-ph.HE]]. Measurement of light-by-light scattering and search for axion-like particles with 2.2 nb⁻¹ of Pb+Pb data with the ATLAS detector, arXiv:2008.05355 [hep-ex].
- [29] G. Amelino-Camelia, J. R. Ellis, N. E. Mavromatos, D. V. Nanopoulos and S. Sarkar, Tests of quantum gravity from observations of gamma-ray bursts, Nature 393, 763 (1998) [astro-ph/9712103].

- [30] J. R. Ellis, K. Farakos, N. E. Mavromatos, V. A. Mitsou and D. V. Nanopoulos, A search in gamma-ray burst data for nonconstancy of the Astrophys. J. 535, 139 (2000) [astro-ph/9907340].
- [31] U. Jacob and T. Piran, Neutrinos from gamma-ray bursts as a tool to explore quantumgravity-induced Lorentz violation, Nature Phys. 3, 87 (2007) [hep-ph/0607145].
- [32] N. E. Mavromatos, String Quantum Gravity, Lorentz-Invariance Violation and Gamma-Ray Astronomy, Int. J. Mod. Phys. A 25, 5409 (2010) [arXiv:1010.5354 [hep-th]].
- [33] M. Rodriguez Martinez and T. Piran, Constraining Lorentz violations with gamma-ray bursts, JCAP 04, 006 (2006) [astro-ph/0601219].
- [34] Hanlin Song and Bo-Qiang Ma, Energy-dependent intrinsic time delay of gamma-ray bursts on testing Lorentz invariance violation, Phys.Lett.B 856 (2024) 138951, arXiv:2408.14719 [astro-ph.HE].
- [35] T. Piran and D. D. Ofengeim, Lorentz invariance violation limits from GRB 221009A, Phys. Rev. D 109 (2024) no.8, L081501 [arXiv:2308.03031 [astro-ph.HE]].
- [36] Z. Cao et al. [LHAASO], Stringent Tests of Lorentz Invariance Violation from LHAASO Observations of GRB 221009A, Phys. Rev. Lett. 133 (2024) no.7, 071501 [arXiv:2402.06009 [astro-ph.HE]].