# Multivariate Poisson approximation of joint subgraph counts in random graphs via size-biased couplings

Eulalia Nualart

Rui-Ray Zhang

Universitat Pompeu Fabra B Barcelona School of Economics Ramón Trias Fargas 25-27, Barcelona 08005, Spain eulalia.nualart@upf.edu

Barcelona School of Economics rui.zhang@bse.eu

#### Abstract

Using Chen-Stein method in combination with size-biased couplings, we obtain the multivariate Poisson approximation in terms of the Wasserstein distance. As applications, we study the multivariate Poisson approximation of the distribution of joint subgraph counts in an Erdős-Rényi random graph and the multivariate hypergeometric distribution giving explicit convergence rates.

*Keywords*: Poisson approximation; Chen-Stein method; size-biased couplings; random graphs; hypergeometric distribution.

AMS 2010 subject classification: 60F05; 60C05; 05C80.

# 1 Introduction

Let  $\mathcal{G}(n,p)$  denote the Erdős-Rényi random graph on n vertices with edge probability  $p = p(n) \in (0,1)$ . Let H be a strictly balanced graph with  $v_H > 0$  vertices and  $e_H$  edges, and let W denote the number of copies of H in  $\mathcal{G}(n,p)$ . It is well-known since the 80's (see e.g. Janson, Luczak, and Ruciński [5, Theorem 3.19]) that using the method of moments, if  $np^{d_H} \to c > 0$  as  $n \to \infty$ , where  $d_H = e_H/v_H$  denotes the density of H, then W converges in distribution to a Poisson random variable  $P_{\lambda}$  with parameter  $\lambda = c^{v_H}/a_H$ , where  $a_H$  denotes the size of the automorphism group of H. Furthermore, using the Chen-Stein method, one can obtain an explicit rate of convergence in terms of the total variation distance of both distributions, see Barbour, Holst, and Janson [2, Chapter 5] or [5, Example 6.26]. More specifically, if H has no isolated vertices, then as  $n \to \infty$ ,

$$d_{TV}(W, P_{\lambda}) = O(n^{-\gamma}),$$

where  $\gamma = \min \{ v_{H'} - e_{H'}/d_H : H' \subsetneq H, e_{H'} > 0 \} > 0$  and  $\lambda = \mathbb{E}(W) \to c^{v_H}/a_H$ . This is based on the notion of size-biased coupling that we recall in Section 2. See Ross [7, Section 4] for an extended survey of this topic.

One of the goals of this paper is to extend this rate of convergence to the joint convergence of several subgraph counts in terms of the Wasserstein distance, which is stronger than the total variation one. More specifically, let  $(H_1, \ldots, H_d)$  be a sequence of d distinct strictly balanced graphs with  $0 < v_{H_i} \leq n$  vertices,  $e_{H_i} > 0$  edges, same density  $\alpha = d_{H_i}$ , for  $i = 1, \ldots, d$ , and no isolated vertices. Let  $\mathbf{W} = (W_1, \ldots, W_d)$ , where for  $i = 1, \ldots, d$ ,  $W_i$  denotes the number of copies of  $H_i$  in  $\mathcal{G}(n, p)$ . Then, we show in Theorem 3.4, that if for some constant c > 0,  $p = cn^{-1/\alpha}(1 + o(1))$ , then as  $n \to \infty$ ,

$$d_{\mathrm{W}}(\mathbf{W}, \mathbf{P}_{\lambda}) = O(n^{-\gamma}),$$

where  $\mathbf{P}_{\lambda} = (P_1, \ldots, P_d)$  denotes a Poisson random vector with independent components and expectation  $\boldsymbol{\lambda} = (\lambda_1, \ldots, \lambda_d)$  with  $\lambda_i = \mathbb{E}(W_i) \to c^{\alpha v_{H_i}}/a_{H_i}$ , and

$$\gamma = \min_{i \in \{1, \dots, d\}} \min\{ v_{H'_i} - e_{H'_i} / \alpha : H'_i \subsetneq H_i, e_{H'_i} > 0 \} > 0$$

If the graphs do not have the same density, but there is a critical common density  $\alpha$  shared by a subset of them, our Theorem 3.4 also studies the behavior of  $W_i$  for the graphs that do not have density equal to  $\alpha$ . In the case that the graphs are not necessarily strictly balanced and  $p \to 0$  as  $n \to \infty$ , we obtain a bound for the Wasserstein distance of both distributions that can be applied to a wide class of examples in order to show multivariate Poisson convergence, see Corollary 3.2 and Remark 3.5.

In order to achieve this first aim, we use the multivariate Poisson approximation developed by Pianoforte and Turin [6], which differs from the multivariate extension given by Goldstein and Rinott [4] and that by Arratia, Goldstein, and Gordon [1]. In [6], the authors use Stein's equation for the Poisson distribution iteratively in order to derive a multivariate Poisson approximation bound in terms of the Wasserstein distance for general integer-valued random vectors. As applications, they obtain the multivariate Poisson approximation of dependent Bernoulli sums and Poisson process approximation of point processes of U-statistic structure. As the authors observe in their paper, a particular case of their bound is when one can find random vectors with an exact size-biased distribution in a multivariate setting that differs from previous multivariate extensions such as [4]. In our paper, we exploit this fact by formalizing the multivariate notion of size-biased coupling and studying the case of multivariate sums of indicators random variables. Then, as a consequence of the multivariate Poisson approximation theorem obtained in [6], we obtain explicit bounds in terms of the Wasserstein distance for increasing and decreasing multivariate size-biased couplings, extending the one-dimensional case (see for e.g. [7]). Although the multivariate Poisson distribution has independent components, our bounds take into account of the dependences of the random vector through its covariances. We then apply the increasing size-biased multivariate Poisson approximation bound to the joint convergence of several subgraph counts mentioned above. This needs a careful analysis of the covariance structure of subgraph counts, extending the covariance study done in Krokowski and Thäle [8]. In [8, Theorem 4.2], the rate of convergence to the multivariate Gaussian distribution of joint subgraph counts is achieved in the case that p is constant, using the fourth differentiable function distance and discrete Malliavin calculus. Finally, as an application of the decreasing size-biased multivariate Poisson approximation, we consider the multivariate hypergeometric distribution.

While we focused on Erdős-Rényi random graphs, the techniques developed here may potentially be extended to other random graph models, such as random geometric graphs, etc. The multivariate size-biased coupling framework could also be useful in other contexts such as pattern occurrences in random permutations, and local statistics in random discrete structures. It may be possible to obtain high-order asymptotic probability with smaller error, by considering clusters of subgraphs, as in Zhang [9, Chapter 5]. The paper is organized as follows. In Section 2 we develop the general theory of multivariate size-biased coupling and Poisson approximations. Then in Section 3 we consider the two applications, namely the distribution of subgraph counts in random graphs and the multivariate hypergeometric distribution.

**Notation:** We use the notation  $\mathbb{N} = \{1, 2, ...\}$  and  $\mathbb{N}_0 = \{0, 1, 2, ...\}$ . Throughout this paper,  $\mathbf{P}_{\lambda} = (P_1, ..., P_d)$  denotes a Poisson distributed random vector with independent components and expectation  $\boldsymbol{\lambda} = (\lambda_1, ..., \lambda_d)$ .

# 2 Size-biased coupling and Poisson approximation

Size-biased couplings first appeared in the context of Stein's method for normal approximation by Goldstein and Rinott [4], and they are also useful when used in conjunction with Stein's method to obtain Poisson approximation. Following the survey by Ross [7], we recall the notion of size-biased coupling.

**Definition 2.1** (Size-biased coupling). Let  $W \ge 0$  be a random variable with  $\mathbb{E}(W) = \lambda < \infty$ . We say that a random variable  $\widetilde{W}$  is a size-biased coupling of W if for all function f satisfying  $\mathbb{E}|Wf(W)| < \infty$ , we have

$$\mathbb{E}(Wf(W)) = \lambda \mathbb{E}(f(\widetilde{W})).$$

Observe that if  $W \ge 0$  is an integer-valued random variable with finite mean  $\lambda > 0$ , then  $\widetilde{W}$  is a size-biased coupling of W if and only if for all  $k \in \mathbb{N}_0$ ,

$$\mathbb{P}(\widetilde{W} = k) = \frac{k}{\lambda} \mathbb{P}(W = k).$$

Applying size-biased couplings gives Poisson approximation for real-valued random variables.

**Theorem 2.2.** [7, Theorem 4.13] Let  $W \ge 0$  be an integer-valued random variable with  $\mathbb{E}(W) = \lambda > 0$ , and let  $\widetilde{W}$  be a size-biased coupling of W. Then, if  $P_{\lambda}$  is a Poisson distributed random variable with parameter  $\lambda$ , we have

$$d_{\mathrm{TV}}(W, P_{\lambda}) \leq \min\{1, \lambda\} \mathbb{E}|\widetilde{W} - 1 - W|,$$

where the total variation (TV) distance between two distributions P and Q on some finite domain  $\mathcal{D}$  is defined by

$$d_{\rm TV}(P,Q) = \frac{1}{2} \sum_{x \in \mathcal{D}} |P(x) - Q(x)| = \max_{S \subseteq \mathcal{D}} |P(S) - Q(S)|.$$

### 2.1 Multivariate size-biased coupling

We extend the size-biased coupling to the multivariate case, following the result by Pianoforte and Turin [6].

**Definition 2.3** (Multivariate size-biased coupling). Let  $\mathbf{W} = (W_1, \ldots, W_d)$  be a random vector with  $\mathbb{E}(W_i) = \lambda_i < \infty$ , for  $i = 1, \ldots, d$ . Let  $\widetilde{\mathbf{W}} = (\widetilde{\mathbf{W}}^1, \ldots, \widetilde{\mathbf{W}}^d)$  be a d-dimensional triangular array such that  $\widetilde{\mathbf{W}}^i = (\widetilde{W}^i_1, \ldots, \widetilde{W}^i_i)$  is an i-dimensional random vector for each  $i = 1, \ldots, d$ . We say that  $\widetilde{\mathbf{W}}$  is a size-biased coupling of  $\mathbf{W}$  if

$$\mathbb{E}(W_i f(\mathbf{W}^i)) = \lambda_i \mathbb{E}(f(\widetilde{\mathbf{W}}^i)), \quad with \ \mathbf{W}^i = (W_1, \dots, W_i)$$
(2.1)

for all i = 1, ..., d, and all functions  $f : \mathbb{R}^i \to \mathbb{R}$  satisfying  $\mathbb{E}|W_i f(\mathbf{W}^i)| < \infty$ .

Observe that if  $\mathbf{W} = (W_1, \ldots, W_d)$  takes values in  $\mathbb{N}_0^d$ ,  $0 < \mathbb{E}(W_i) = \lambda_i < \infty$ , and the random vector  $\widetilde{\mathbf{W}}^i$  takes values in  $\mathbb{N}_0^i$  for all  $i = 1, \ldots, d$ , then condition (2.1) is equivalent of saying that for all  $i = 1, \ldots, d$  and  $\mathbf{k}^i = (k_1, \ldots, k_i) \in \mathbb{N}_0^i$ , we have

$$\mathbb{P}(\widetilde{\mathbf{W}}^{i} = \mathbf{k}^{i}) = \frac{k_{i}}{\lambda_{i}} \mathbb{P}(\mathbf{W}^{i} = \mathbf{k}^{i}).$$
(2.2)

The following extends [7, Corollary 4.14] to the multivariate setting.

**Lemma 2.4.** For all i = 1, ..., d, let  $(X_1^i, ..., X_{n_i}^i)$  be an  $n_i$ -dimensional random vector formed by indicator random variables such that for  $j = 1, ..., n_i$ ,  $\mathbb{P}(X_j^i = 1) = p_{i,j} \in (0,1)$ , and set  $\mathbf{W} = (W_1, ..., W_d)$ , with  $W_i = \sum_{j=1}^{n_i} X_j^i$ . For all i = 1, ..., d and  $\ell = 1, ..., n_i$ , consider an  $n_1 \cdots n_{i-1}(n_i - 1)$ -dimensional random vector

$$\mathbf{X}^{i,(i,\ell)} = \left( (X_j^{1,(i,\ell)})_{j=1}^{n_1}, \dots, (X_j^{i-1,(i,\ell)})_{j=1}^{n_{i-1}}, (X_j^{i,(i,\ell)})_{j=1,j\neq\ell}^{n_i} \right)$$

defined on the same probability space as the random vector

$$\mathbf{X}^{i,\ell} = \left( (X_j^1)_{j=1}^{n_1}, \dots, (X_j^{i-1})_{j=1}^{n_{i-1}}, (X_j^i)_{j=1, j \neq \ell}^{n_i} \right)$$

satisfying

$$\mathcal{L}\left(\mathbf{X}^{i,(i,\ell)}\right) = \mathcal{L}\left(\mathbf{X}^{i,\ell} \mid X^{i}_{\ell} = 1\right).$$
(2.3)

Let  $I_1, \ldots, I_d$  be a sequence of independent random variables, independent of all else, such that for all  $i = 1, \ldots, d$  and  $j = 1, \ldots, n_i$ ,

$$\mathbb{P}(I_i = j) = \frac{p_{i,j}}{\lambda_i}, \qquad \text{where } \lambda_i = \mathbb{E}(W_i) = \sum_{j=1}^{n_i} p_{i,j}.$$
(2.4)

Let  $\widetilde{\mathbf{W}} = (\widetilde{\mathbf{W}}^1, \dots, \widetilde{\mathbf{W}}^d)$  be the d-dimensional triangular array given by, for  $i = 1, \dots, d$ ,

$$\widetilde{\mathbf{W}}^{i} = \left(\sum_{j=1}^{n_{1}} X_{j}^{1,(i,I_{i})}, \dots, \sum_{j=1}^{n_{i-1}} X_{j}^{i-1,(i,I_{i})}, \sum_{j=1, j \neq I_{i}}^{n_{i}} X_{j}^{i,(i,I_{i})} + 1\right).$$
(2.5)

Then,  $\widetilde{\mathbf{W}}$  is a size-biased coupling of  $\mathbf{W}$ .

*Proof.* It suffices to prove (2.2). For i = 1, ..., d and  $\mathbf{k}^i = (k_1, ..., k_i) \in \mathbb{N}_0^i$ , in view of (2.4) and (2.5), we have, conditioning with respect to all the values of  $I_i$  and using independence together with the conditional probability law equality (2.3), that

$$\begin{split} \lambda_{i} \mathbb{P}(\widetilde{\mathbf{W}}^{i} = \mathbf{k}^{i}) &= \lambda_{i} \sum_{j=1}^{n_{i}} \mathbb{P}(\widetilde{\mathbf{W}}^{i} = \mathbf{k}^{i}, I_{i} = j) \\ &= \sum_{j=1}^{n_{i}} \mathbb{P}((W_{1}, \dots, W_{i-1}, W_{i} - X_{j}^{i} + 1) = \mathbf{k}^{i} \mid X_{j}^{i} = 1) p_{i,j} \\ &= \sum_{j=1}^{n_{i}} \frac{p_{i,j}}{\mathbb{P}(X_{j}^{i} = 1)} \mathbb{P}(\mathbf{W}^{i} = \mathbf{k}^{i}, X_{j}^{i} = 1) = \sum_{j=1}^{n_{i}} \mathbb{P}(\mathbf{W}^{i} = \mathbf{k}^{i}, X_{j}^{i} = 1) = k_{i} \mathbb{P}(\mathbf{W}^{i} = \mathbf{k}^{i}), \end{split}$$

recalling the definition of  $p_{i,j}$  in (2.4). This concludes the desired proof.

The following extends [7, Corollary 4.15] to the multivariate setting.

**Corollary 2.5.** For all i = 1, ..., d, let  $(X_1^i, ..., X_{n_i}^i)$  be an  $n_i$ -dimensional random vector formed by exchangeable indicator random variables and consider an  $n_1 \cdots n_{i-1}(n_i - 1)$ -dimensional random vector

$$\mathbf{X}^{i,(i,1)} = \left( (X_j^{1,(i,1)})_{j=1}^{n_1}, \dots, (X_j^{i-1,(i,1)})_{j=1}^{n_{i-1}}, (X_j^{i,(i,1)})_{j=2}^{n_i} \right)$$

defined on the same probability space as the random vector

$$\mathbf{X}^{i,1} = \left( (X_j^1)_{j=1}^{n_1}, \dots, (X_j^{i-1})_{j=1}^{n_{i-1}}, (X_j^i)_{j=2}^{n_i} \right)$$

satisfying

$$\mathcal{L}\left(\mathbf{X}^{i,(i,1)}\right) = \mathcal{L}\left(\mathbf{X}^{i,1} \mid X_1^i = 1\right)$$

Let  $\widetilde{\mathbf{W}} = (\widetilde{\mathbf{W}}^1, \dots, \widetilde{\mathbf{W}}^d)$  be the d-dimensional triangular array given by, for  $i = 1, \dots, d$ ,

$$\widetilde{\mathbf{W}}^{i} = \left(\sum_{j=1}^{n_{1}} X_{j}^{1,(i,1)}, \dots, \sum_{j=1}^{n_{i-1}} X_{j}^{i-1,(i,1)}, \sum_{j=2}^{n_{i}} X_{j}^{i,(i,1)} + 1\right).$$

Then,  $\widetilde{\mathbf{W}}$  is a size-biased coupling of  $\mathbf{W}$ .

*Proof.* The proof follows from Lemma 2.4 and the fact that exchangeability implies that the random variables  $I_i$ 's are uniformly distributed.

### 2.2 Multivariate Poisson approximation

Recall that the Wasserstein distance between random vectors  $\mathbf{X}$  and  $\mathbf{P}$  is defined by

$$d_{\mathrm{W}}(\mathbf{X}, \mathbf{P}) = \sup_{g \in \mathrm{Lip}^{d}(1)} |\mathbb{E}(g(\mathbf{X})) - \mathbb{E}(g(\mathbf{P}))|,$$

where  $\operatorname{Lip}^{d}(1)$  denotes the set of Lipschitz functions  $g: \mathbb{N}_{0}^{d} \to \mathbb{R}$  with Lipschitz constant bounded by 1 with respect to the 1-norm  $|\mathbf{x}|_{1} = \sum_{i=1}^{d} |x_{i}|$ , for a vector  $\mathbf{x} = (x_{1}, \ldots, x_{d}) \in \mathbb{R}^{d}$ . Since the indicator functions defined on  $\mathbb{N}_{0}^{d}$  are Lipschitz, for random vectors in  $\mathbb{N}_{0}^{d}$ , the Wasserstein distance dominates the total variation distance, and therefore, all bounds henceforth are also valid for total variation distance.

Our first result is the following Poisson approximation using multivariate size-biased coupling, which is a consequence of the result by Pianoforte and Turin [6].

**Theorem 2.6.** Let  $\mathbf{W} = (W_1, \ldots, W_d)$  be a *d*-dimensional random vector taking values in  $\mathbb{N}_0^d$ , with  $\mathbb{E}(W_i) = \lambda_i > 0$  for  $i = 1, \ldots, d$  and let  $\widetilde{\mathbf{W}}$  be a size-biased coupling of  $\mathbf{W}$ . Then,

$$d_{\mathbf{W}}(\mathbf{W}, \mathbf{P}_{\lambda}) \leqslant \sum_{i=1}^{d} \min\{1, \lambda_i\} \mathbb{E}|\widetilde{W}_i^i - 1 - W_i| + 2\sum_{i=2}^{d} \lambda_i \sum_{j=1}^{i-1} \mathbb{E}|\widetilde{W}_j^i - W_j|.$$

*Proof.* The proof is a direct application of [6, Theorem 1.1] by taking, for  $i = 1, \ldots, d$ ,  $Z_i^{(i)} = \widetilde{W}_i^i - 1 - W_i$ ,  $Z_j^{(i)} = \widetilde{W}_j^i - W_j$  for  $j = 1, \ldots, i - 1$ , and  $\mathbf{Y}^{(i)} = \widetilde{\mathbf{W}}^i$ . Then, we observe that (2.2) implies that all the  $q_{m_{1:i}}^{(i)}$ 's in Theorem 1.1 in [6] are zero, since in our notation, we have that for all  $i = 1, \ldots, d$  and  $\mathbf{k}^i = (k_1, \ldots, k_i) \in \mathbb{N}_0^i$  with  $k_i \neq 0$ ,

$$q_{m_{1:i}}^{(i)} = k_i \mathbb{P}(\mathbf{W}^i = \mathbf{k}^i) - \lambda_i \mathbb{P}(\widetilde{\mathbf{W}}^i = \mathbf{k}^i),$$

which concludes the proof.

**Remark 2.7.** We observe that [6, Theorem 1.1] is stated with  $\min\{1, \lambda_i\}$  replaced by  $\lambda_i$ . However, when going through the proof, we see that we can use a sharper bound given for e.g. in [7, Lemma 4.4], which allows to replace the  $\lambda_i$  by  $\min\{1, \lambda_i\}$ .

When the multi-dimensional size-biased coupling is increasing, we extend [7, Theorem 4.20].

**Theorem 2.8.** Under the assumptions of Lemma 2.4, suppose that for all i = 1, ..., d, j = 1, ..., i - 1,  $\ell = 1, ..., n_i$ , and  $k = 1, ..., n_j$ , we have  $X_k^{j,(i,\ell)} \ge X_k^j$ , and for j = i and  $k \neq \ell$  we have  $X_k^{i,(i,\ell)} \ge X_k^j$ . Then,

$$d_{W}(\mathbf{W}, \mathbf{P}_{\lambda}) \leqslant \sum_{i=1}^{d} \min\{1, \lambda_{i}^{-1}\} \left( \operatorname{Var}(W_{i}) - \lambda_{i} + 2\sum_{j=1}^{n_{i}} p_{i,j}^{2} \right) + 2\sum_{i=2}^{d} \lambda_{i}^{-1} \sum_{j=1}^{i-1} \operatorname{Cov}(W_{i}, W_{j}) + 2\sum_{i=1}^{d} \sum_{j=1}^{d} \sum_{i=1}^{d} \sum_{j=1}^{d} \sum_{j=1}^{d} \sum_{j=1}^{d} \sum_{i=1}^{d} \sum_{j=1}^{d} \sum_{i=1}^{d} \sum_{j=1}^{d} \sum_{i=1}^{d} \sum_{j=1}^{d} \sum_{j=1}^{d} \sum_{i=1}^{d} \sum_{j=1}^{d} \sum_{j=1}^{d} \sum_{i=1}^{d} \sum_{j=1}^{d} \sum_{j=1}^{d} \sum_{j=1}^{d} \sum_{i=1}^{d} \sum_{j=1}^{d} \sum_{i=1}^{d} \sum_{j=1}^{d} \sum_{j=1}^$$

*Proof.* Applying Theorem 2.6, we have, by using (2.4), that

$$\begin{aligned} d_{\mathrm{W}}(\mathbf{W}, \mathbf{P}_{\lambda}) \\ &\leqslant \sum_{i=1}^{d} \min\{1, \lambda_{i}^{-1}\} \sum_{j=1}^{n_{i}} p_{i,j} \mathbb{E}\left(\sum_{k=1, k \neq j}^{n_{i}} (X_{k}^{i,(i,j)} - X_{k}^{i}) + X_{j}^{i}\right) + 2 \sum_{i=2}^{d} \lambda_{i} \sum_{j=1}^{i-1} \mathbb{E}\left(\widetilde{W}_{j}^{i} - W_{j}\right) \\ &= \sum_{i=1}^{d} \min\{1, \lambda_{i}\} \mathbb{E}\left(\widetilde{W}_{i}^{i} - W_{i} - 1 + 2X_{I_{i}}^{i}\right) + 2 \sum_{i=2}^{d} \lambda_{i} \sum_{j=1}^{i-1} \mathbb{E}\left(\widetilde{W}_{j}^{i} - W_{j}\right). \end{aligned}$$

Now, using (2.1) and the fact that  $\lambda_i = \mathbb{E}(W_i)$ , we get that

$$\lambda_i \mathbb{E}(\widetilde{W}_i^i - W_i) = \mathbb{E}(W_i^2) - (\mathbb{E}(W_i))^2 = \operatorname{Var}(W_i),$$

and similarly,

$$\lambda_i \mathbb{E}(\widetilde{W}_j^i - W_j) = \mathbb{E}(W_i W_j) - \mathbb{E}(W_i) \mathbb{E}(W_j) = \operatorname{Cov}(W_i, W_j).$$

Finally, using (2.4) to rewrite  $\lambda_i$ , we have

$$\lambda_i \mathbb{E}(X_{I_i}^i) = \lambda_i \sum_{j=1}^{n_i} \mathbb{P}(I_i = j) \mathbb{E}(X_{I_i}^i \mid I_i = j) = \sum_{j=1}^{n_i} p_{i,j} \mathbb{E}(X_{I_i}^i \mid I_i = j) = \sum_{j=1}^{n_i} p_{i,j}^2,$$

where we use the definition of  $p_{i,j}$ . This completes the desired bound.

Next, we extend [7, Theorem 4.31] for the decreasing size-biased coupling.

**Theorem 2.9.** Under the assumptions of Lemma 2.4, suppose that for all i = 1, ..., d, j = 1, ..., i - 1,  $\ell = 1, ..., n_i$ , and  $k = 1, ..., n_j$ , we have  $X_k^{j,(i,\ell)} \leq X_k^j$  and for j = i and  $k \neq \ell$  we have  $X_k^{i,(i,\ell)} \leq X_k^j$ . Then,

$$d_{\mathrm{W}}(\mathbf{W}, \mathbf{P}_{\lambda}) \leqslant \sum_{i=1}^{d} \min\{1, \lambda_i^{-1}\} \left(\lambda_i - \operatorname{Var}(W_i)\right) - 2\sum_{i=2}^{d} \lambda_i^{-1} \sum_{j=1}^{i-1} \operatorname{Cov}(W_i, W_j).$$

*Proof.* Applying Theorem 2.6, we have that

$$d_{W}(\mathbf{W}, \mathbf{P}_{\lambda}) \leq \sum_{i=1}^{d} \min\{1, \lambda_{i}^{-1}\} \sum_{j=1}^{n_{i}} p_{i,j} \mathbb{E}\left(\sum_{k=2}^{n_{i}} (X_{k}^{i} - X_{k}^{i,(i,j)}) + X_{j}^{i}\right) + 2\sum_{i=2}^{d} \lambda_{i} \sum_{j=1}^{i-1} \mathbb{E}\left(W_{j} - \widetilde{W}_{j}^{i}\right) = \sum_{i=1}^{d} \min\{1, \lambda_{i}\} \mathbb{E}\left(W_{i} - \widetilde{W}_{i}^{i} + 1\right) + 2\sum_{i=2}^{d} \lambda_{i} \sum_{j=1}^{i-1} \mathbb{E}\left(W_{j} - \widetilde{W}_{j}^{i}\right).$$

Finally, proceedings as in the proof of Theorem 2.8, we obtain the desired bound.

# 3 Applications

### 3.1 Joint subgraph counts in random graphs

Recall that  $\mathcal{G}(n,p)$  denotes the Erdős-Rényi random graph on n vertices with edge probability  $p = p(n) \in (0,1)$ . Let  $(H_1,\ldots,H_d)$  be a sequence of distinct fixed graphs such that for  $i = 1,\ldots,d$ , each graph  $H_i$  has  $0 < v_{H_i} \leq n$  vertices,  $e_{H_i} > 0$  edges, and no isolated vertices. We analyze the joint distribution of subgraph counts of  $(H_1,\ldots,H_d)$  in  $\mathcal{G}(n,p)$ , that is, the number of copies of  $H_i$  in  $\mathcal{G}(n,p)$  for  $i = 1,\ldots,d$ .

Our main result in this section is the multivariate Poisson approximation of subgraph counts, as a consequence of the decreasing size-biased coupling in Theorem 2.8, as an extension of [7, Theorem 4.21]. Observe that in this application of Theorem 2.8 the  $p_{i,j}$ 's only depend on i.

**Theorem 3.1.** Let  $(H_1, \ldots, H_d)$  be a sequence of d distinct graphs with  $0 < v_{H_i} \leq n$  vertices,  $e_{H_i} > 0$  edges, and no isolated vertices. Let  $\mathbf{W} = (W_1, \ldots, W_d)$ , where  $W_i$  is the number of copies of  $H_i$  in  $\mathcal{G}(n, p)$  for  $i = 1, \ldots, d$ , and let  $\lambda_i = \mathbb{E}(W_i)$ . Then,

$$d_{\mathbf{W}}(\mathbf{W}, \mathbf{P}_{\lambda}) \leqslant \sum_{i=1}^{d} \min\{1, \lambda_i^{-1}\} \left( \operatorname{Var}(W_i) - \lambda_i + 2\lambda_i p^{e_{H_i}} \right) + 2\sum_{i=2}^{d} \lambda_i^{-1} \sum_{j=1}^{i-1} \operatorname{Cov}(W_i, W_j).$$

To obtain more explicit error bounds, we introduce some notation to analyse the covariances between subgraph counts. Let  $H_i$  and  $H_j$  be two graphs that can be isomorphic. For any fixed copies  $\alpha_i \in \Gamma_i$  and  $\beta_j \in \Gamma_j$  in the complete graph  $K_n$ , we denote by  $e_{\alpha_i,\beta_j}$  the number of edges shared between  $\alpha_i$  and  $\beta_j$ , and by  $v_{\alpha_i,\beta_j}$  the number of vertices incident to these shared edges. Define

$$M_{i,j} = \max\{e_{\alpha_i,\beta_j} : \alpha_i \in \Gamma_i, \beta_j \in \Gamma_j\},\$$

which represents the maximum possible number of shared edges between any copy of  $H_i$  and  $H_j$ . For each  $k \in \{1, \ldots, M_{i,j}\}$ , define

$$\ell_{k,i,j} = \begin{cases} \min\{v_{\alpha_i,\beta_j} : e_{\alpha_i,\beta_j} = k, \alpha_i \in \Gamma_i, \beta_j \in \Gamma_j\} & \text{if } \mathcal{S}_k(i,j) \neq \emptyset, \\ 0 & \text{if } \mathcal{S}_k(i,j) = \emptyset, \end{cases}$$

where  $S_k(i, j) = \{(\alpha_i, \beta_j) \in \Gamma_i \times \Gamma_j : e_{\alpha_i, \beta_j} = k\}$  is the set of pairs sharing exactly k edges. When  $\ell_{k,i,j} > 0$ , it represents the minimum number of vertices needed to realize k shared edges between

copies of  $H_i$  and  $H_j$ . Note that  $\binom{\ell_{k,i,j}}{2} \leq k \leq 2\ell_{k,i,j}$ , and  $\ell_{k,i,j} = 0$  when it is impossible for  $H_i$  and  $H_j$  to share exactly k edges. Let

$$\mathcal{K}_{i,j} = \{k \in \{1, \dots, M_{i,j}\} : \ell_{k,i,j} > 0\}$$

be the set of possible shared edge counts between  $H_i$  and  $H_j$ . Define

$$\gamma_i(t) = \min_{k \in \mathcal{K}_{i,i}} \left\{ \ell_{k,i,i} - \frac{k}{t} \right\}$$

As a consequence of Theorem 3.1, we obtain an explicit error bound in the case that  $p \to 0$  as  $n \to \infty$  using the above notation.

**Corollary 3.2.** Under the same conditions as in Theorem 3.1, assume that  $p \to 0$  as  $n \to \infty$ . Then, there exists a constant C > 0 such that for n sufficiently large, we have

$$d_{W}(\mathbf{W}, \mathbf{P}_{\lambda}) \leqslant C \left\{ \sum_{i=1}^{d} \min\{1, \lambda_{i}\} \left( p^{e_{H_{i}}} + n^{v_{H_{i}} - \gamma_{i}} p^{e_{H_{i}}} \right) + \sum_{i=2}^{d} \sum_{j=1}^{i-1} \lambda_{j} \sum_{k \in \mathcal{K}_{i,j}} p^{-k} n^{-\ell_{k,i,j}} \right\},$$

where for  $i = 1, \ldots, d$ ,

$$\gamma_i = \gamma_i(d_{H_i}) = \min\left\{ v_{H'_i} - \frac{e_{H'_i}}{d_{H_i}} : H'_i \subsetneq H_i, e_{H'_i} > 0 \right\},$$

and  $d_{H_i} = e_{H_i}/v_{H_i}$  is the density of the graph  $H_i$ .

#### 3.1.1 Proofs of Theorem 3.1 and Corollary 3.2

Proof of Theorem 3.1. For i = 1, ..., d, let  $\Gamma_i$  denote the set of all copies of  $H_i$  in the complete graph of n vertices  $K_n$ . Let  $X_{\alpha_i}^i$  the indicator that there is a copy of  $H_i$  in  $\mathcal{G}(n, p)$  at  $\alpha_i$ , and set  $W_i = \sum_{\alpha_i \in \Gamma_i} X_{\alpha_i}^i$ . We apply Theorem 2.8 to the vector  $\mathbf{W}$  by constructing a size-biased coupling of  $\mathbf{W}$  using Corollary 2.5 and the fact that each  $W_i$  is a sum of exchangeable indicators.

For all i = 1, ..., d, j = 1, ..., i, and  $\alpha_i \in \Gamma_i$ , let  $X_{\beta_j}^{j,(i,\alpha_i)}$  be the indicator that there is a copy of  $H_i$  in  $\mathcal{G}(n,p) \cup \{\alpha_i\}$  at  $\beta_j \in \Gamma_j$ , where  $\mathcal{G}(n,p) \cup \{\alpha_i\}$  denotes the graph obtained by adding the minimum edges necessary to  $\mathcal{G}(n,p)$  such that  $\mathcal{G}(n,p) \cup \{\alpha_i\}$  contains a copy of  $H_i$  at  $\alpha_i$ .

Then the following facts imply the theorem:

•  $\mathcal{L}(\mathbf{X}^{i,(i,\alpha_i)}) = \mathcal{L}(\mathbf{X}^{i,\alpha_i} \mid X^i_{\alpha_i} = 1)$ , where

$$\mathbf{X}^{i,(i,\alpha_i)} = \left( (X_{\beta}^{1,(i,\alpha_i)})_{\beta \in \Gamma_1}, \dots, (X_{\beta}^{i-1,(i,\alpha_i)})_{\beta \in \Gamma_{i-1}}, (X_{\beta}^{i,(i,\alpha_i)})_{\beta \in \Gamma_i, \beta \neq \alpha_i} \right)$$

and

$$\mathbf{X}^{i,\alpha_i} = \left( (X^1_\beta)_{\beta \in \Gamma_1}, \dots, (X^{i-1}_\beta)_{\beta \in \Gamma_{i-1}}, (X^i_\beta)_{\beta \in \Gamma_i, \beta \neq \alpha_i} \right)$$

- For all i = 1, ..., d, j = 1, ..., i 1,  $\alpha_i \in \Gamma_i$ , and  $\beta_j \in \Gamma_j$ , we have  $X_{\beta_j}^{j,(i,\alpha_i)} \ge X_{\beta_j}^j$ , and for  $j = i, \beta_i \neq \alpha_i$ , we have  $X_{\beta_i}^{i,(i,\alpha_i)} \ge X_{\beta_i}^j$ , by noting that we only add edges in the size-biased coupling construction.
- $\mathbb{E}(X_{\alpha_i}^i) = p^{e_{H_i}}$  for all  $\alpha \in \Gamma_i$ . This completes the proof.

Observe that applying the one-dimensional bound of [7, Corollary 4.22], we obtain the following bound for the first error term in Theorem 3.1. For  $i = 1, \ldots, d, \alpha_i \in \Gamma_i$ , and  $H'_i \subseteq H_i$ , let  $\Gamma_i^{H'_i,\alpha_i} \subseteq \Gamma_i$  be the set of subgraphs of  $K_n$  isomorphic to  $H_i$ , whose intersection with  $\alpha_i$  is  $H'_i$ . Then, for  $i = 1, \ldots, d$ ,

$$\operatorname{Var}(W_i) - \lambda_i + 2\lambda_i p^{e_{H_i}} \leqslant \lambda_i \left( p^{e_{H_i}} + \sum_{\substack{H_i' \subseteq H_i: e_{H_i'} > 0 \\ H_i' \subseteq H_i: e_{H_i'} > 0} |\Gamma_i^{H_i', \alpha_i}| p^{e_{H_i} - e_{H_i'}} \right),$$
(3.1)

where the terms  $|\Gamma_i^{H'_i,\alpha_i}|$  account for the number of covariance terms for different types of pair indicators.

We next give a more explicit bound on the second error term in Theorem 3.1. Recall that  $a_H$ denotes the size of the set of automorphisms of graph H. We observe that for  $i = 1, \ldots, d$ ,

$$\mathbb{E}(W_i) = |\Gamma_i| p^{e_{H_i}} = \binom{n}{v_{H_i}} \frac{v_{H_i}!}{a_{H_i}} p^{e_{H_i}}.$$

We have the following asymptotic result, which extends Lemma 4.1 in Krokowski and Thäle [8] to the case where p depends on n, and with errors therein corrected.

**Lemma 3.3.** Under the same conditions as in Corollary 3.2, for all  $i \in \{2, \ldots, d\}$  and  $j \in$  $\{1, \ldots, i-1\}$ , we have, for sufficiently large n,

$$\frac{\operatorname{Cov}(W_i, W_j)}{\mathbb{E}(W_i)\mathbb{E}(W_j)} = O\bigg(\sum_{k \in \mathcal{K}_{i,j}} p^{-k} n^{-\ell_{k,i,j}}\bigg).$$

*Proof.* The proof follows along the same lines as in [8, Lemma 4.1], but taking into account the fact that p depends on n. Let  $i \in \{2, \ldots, d\}$  and  $j \in \{1, \ldots, i-1\}$ . Then for sufficiently large n,

$$\operatorname{Cov}(W_i, W_j) = \sum_{\alpha_i \in \Gamma_i, \beta_j \in \Gamma_j : e_{\alpha_i, \beta_j} \ge 1} \left( \mathbb{E}(X_{\alpha_i}^i X_{\beta_j}^j) - \mathbb{E}(X_{\alpha_i}^i) \mathbb{E}(X_{\beta_j}^j) \right)$$
$$= \sum_{\alpha_i \in \Gamma_i, \beta_j \in \Gamma_j : e_{\alpha_i, \beta_j} \ge 1} \left( p^{e_{H_i} + e_{H_j} - e_{\alpha_i, \beta_j}} - p^{e_{H_i}} p^{e_{H_j}} \right)$$
$$= \sum_{k \in \mathcal{K}_{i,j}} \sum_{(\alpha_i, \beta_j) \in \mathcal{S}_k(i, j)} p^{e_{H_i} + e_{H_j} - k} (1 - p^k) \leqslant \sum_{k \in \mathcal{K}_{i, j}} \Delta_k,$$

where  $\Delta_k = p^{-k} \sum_{(\alpha_i, \beta_j) \in \mathcal{S}_k(i,j)} p^{e_{H_i} + e_{H_j}}$ . Next, we observe that since the  $v_{H_i} > 0$  and the graphs  $H_i$  have no isolated vertices, we have that  $1 \in \mathcal{K}_{i,j}$ . Let k = 1. We have  $\binom{n}{v_{H_i}} \frac{v_{H_i}!}{a_{H_i}} = \frac{\mathbb{E}(W_i)}{p^{e_{H_i}}}$  possible choices for  $\alpha_i \in \Gamma_i$ . Since k = 1, given  $\alpha_i \in \Gamma_i$ , we need to compute the number of  $\beta_j \in \Gamma_j$  that have exactly one edge in common with  $\alpha_i$ , by observing that we have  $e_{H_i}$  possibilities to choose this common edge in  $\alpha_i$ . Once the common edge in  $\alpha_i$  is fixed, we have  $\binom{n-v_{H_i}}{v_{H_j}-2}$  possibilities to choose the  $v_{H_j}-2$  remaining vertices of  $\beta_j$ ,  $e_{H_i}$  possibilities to choose the common edge in  $\beta_j$ , and  $2(v_{H_i}-2)!/a_{H_i}$  distinct possible arrangements of the vertices. Then

$$\Delta_1 = \mathbb{E}(W_i) e_{H_i} e_{H_j} \binom{n - v_{H_i}}{v_{H_j} - 2} \frac{2(v_{H_j} - 2)!}{a_{H_j}} p^{e_{H_j} - 1} = \mathbb{E}(W_i) \mathbb{E}(W_j) O(n^{-2} p^{-1}).$$

If k = 2, that is,  $\alpha_i$  and  $\beta_j$  have exactly two common edges, then we have two possibilities. The first one is that the edges share one common vertex, thus, there are only three distinct vertices. In this case, following as above we find that the contribution in the sum is  $\mathbb{E}(W_i)\mathbb{E}(W_j)O(n^{-3}p^{-2})$ . The second possibility is that the edges do not share any vertex, that is, we have 4 distinct vertices. In this case the contribution is  $\mathbb{E}(W_i)\mathbb{E}(W_j)O(n^{-4}p^{-2})$ . Therefore, for  $k \ge 2$ , we observe that the terms that will contribute in the bound will be those for which the k edges in common have a minimum number of distinct vertices, that is,  $\ell_{k,i,j}$ . This concludes the proof.  $\Box$ 

Proof of Corollary 3.2. Applying the bound (3.1) and Lemma 3.3, we get, from Theorem 3.1, that for sufficiently large n, the quantity  $d_{\rm W}(\mathbf{W}, \mathbf{P}_{\lambda})$  is bounded by

$$\begin{split} C \bigg\{ \sum_{i=1}^{d} \min\{1,\lambda_{i}\} \bigg( p^{e_{H_{i}}} + \sum_{H_{i}' \subsetneq H_{i}:e_{H_{i}'} > 0} |\Gamma_{i}^{H_{i}',\alpha_{i}}| p^{e_{H_{i}}-e_{H_{i}'}} \bigg) + \sum_{i=2}^{d} \sum_{j=1}^{i-1} \lambda_{j} \sum_{k \in \mathcal{K}_{i,j}} p^{-k} n^{-\ell_{k,i,j}} \bigg\} \\ &\leqslant C \bigg\{ \sum_{i=1}^{d} \min\{1,\lambda_{i}\} \bigg( p^{e_{H_{i}}} + n^{v_{H_{i}}} p^{e_{H_{i}}} \sum_{H_{i}' \subsetneq H_{i}:e_{H_{i}'} > 0} \bigg( n^{v_{H_{i}'}/e_{H_{i}'}-v_{H_{i}}/e_{H_{i}}} \bigg)^{-e_{H_{i}'}} \bigg) \\ &\quad + \sum_{i=2}^{d} \sum_{j=1}^{i-1} \lambda_{j} \sum_{k \in \mathcal{K}_{i,j}} p^{-k} n^{-\ell_{k,i,j}} \bigg\}, \end{split}$$

by noting that  $|\Gamma_i^{H'_i,\alpha_i}| \leq n^{v_{H_i}-v_{H'_i}}$ . This completes the proof.

#### 3.1.2 Strictly balanced graphs

We now apply Corollary 3.2 to a wide class of graphs called *strictly balanced* in order to obtain an exact convergence rate. A graph H is called *strictly balanced* if

$$d_{H'} < d_H$$
 whenever  $H' \subsetneq H$ , where  $d_H = \frac{e_H}{v_H}$ 

is the density of the graph H. It is well-known that trees, cycles, and complete graphs are all strictly balanced graphs. We have the following consequence Corollary 3.2 when all the graphs are strictly balanced.

**Theorem 3.4.** Let  $\mathcal{G}(n,p)$  be an Erdős-Rényi random graph with edge probability

$$p = cn^{-1/\alpha}(1 + o(1)),$$

where c > 0 and  $\alpha > 0$  are fixed constants. Let  $(H_1, \ldots, H_d)$  be a sequence of d strictly balanced distinct graphs with  $0 < v_{H_i} \leq n$  vertices,  $e_{H_i} > 0$  edges, for  $i = 1, \ldots, d$ , and no isolated vertices, Let  $W_i$  denote the number of copies of  $H_i$  in  $\mathcal{G}(n, p)$ . Define  $I = \{i : d_{H_i} = \alpha\}$  as the set of critical indices. Then as  $n \to \infty$ , we have the following.

1. For  $i \notin I$ : If  $d_{H_i} < \alpha$  then

$$\frac{W_i}{\mathbb{E}(W_i)} \xrightarrow{P} 1 \text{ at rate } n^{-\gamma_i},$$

where

$$\gamma_i = \gamma_i(\alpha) = \min_{k \in \mathcal{K}_{i,i}} \left\{ \ell_{k,i,i} - \frac{k}{\alpha} \right\} > 0.$$

If  $d_{H_i} > \alpha$  then

$$\mathbb{P}(W_i > 0) \leqslant \exp(-Cn^{\eta_i}),$$

where

$$\eta_i = \eta_i(\alpha) = v_{H_i}(d_{H_i}/\alpha - 1) > 0.$$

2. For the critical subsequence  $\mathbf{W}_I = (W_i)_{i \in I}$ , we have,

$$d_{\mathrm{W}}(\mathbf{W}_{I}, \mathbf{P}_{\lambda_{\mathbf{I}}}) \leqslant C n^{-\gamma},$$

where  $\boldsymbol{\lambda}_I = (\lambda_i)_{i \in I}$  with  $\lambda_i = \mathbb{E}(W_i) \to c^{\alpha v_{H_i}}/a_{H_i}$  and

$$\gamma = \min_{i \in I} \gamma_i(\alpha) = \min_{i \in I} \min\{ v_{H'_i} - e_{H'_i} / \alpha : H'_i \subsetneq H_i, e_{H'_i} > 0 \} > 0.$$
(3.2)

*Proof.* When  $d_{H_i} < \alpha$ , we use the second moment method. First note that  $\mathbb{E}[W_i] \to \infty$ . By Chebyshev's inequality and Lemma 3.3, for all  $\epsilon > 0$ ,

$$\mathbb{P}\left(\left|\frac{W_i}{\mathbb{E}[W_i]} - 1\right| \ge \epsilon\right) \leqslant \frac{\operatorname{Var}(W_i)}{\epsilon^2 \mathbb{E}[W_i]^2} = O\left(\sum_{k \in \mathcal{K}_{i,i}} n^{k/\alpha - \ell_{k,i,i}}\right) = O\left(n^{-\gamma_i}\right),$$

in view of the definition of  $\gamma_i$ . By the strictly balanced property, we have  $k/\ell_{k,i,i} < e_{H_i}/v_{H_i} = d_{H_i} < \alpha$ , and therefore  $\gamma_i > 0$ .

When  $d_{H_i} > \alpha$ , we have that

$$\mathbb{E}[W_i] \leqslant n^{v_{H_i}} (cn^{-1/\alpha})^{e_{H_i}} = c^{e_{H_i}} n^{v_{H_i}(1-d_{H_i}/\alpha)} \leqslant \exp(-Cn^{\eta_i}),$$

with  $\eta_i > 0$ . By Markov's inequality,

$$\mathbb{P}(W_i > 0) \leqslant \mathbb{E}[W_i] \leqslant \exp(-Cn^{\eta_i}).$$

We now consider the case  $d_{H_i} = \alpha$  for all  $i \in I$ . First observe that for all  $i \in I$ , as  $n \to \infty$ ,  $\mathbb{E}(W_i) \to c^{\alpha v_{H_i}}/a_{H_i}$ . We next bound the two terms in Corollary 3.2. For this step, in order to simplify the exposition, we assume, without loss of generality, that  $I = \{1, \ldots, d\}$ . We start bounding the first sum. We observe that  $\lambda_i = O(1)$  and that  $n^{v_{H_i}}p^{e_{H_i}} = O(1)$ . In particular,  $p^{e_{H_i}} = p^{e_{H_i}}n^{v_{H_i}}n^{-v_{H_i}} = O(n^{-v_{H_i}})$ . Then, recalling the definition of  $\gamma$  in (3.2) and observing that  $\gamma > 0$  since all the graphs are strictly balanced, we obtain that for large n,

$$\sum_{i=1}^{d} \min\{1, \lambda_i\} \left( p^{e_{H_i}} + n^{v_{H_i} - \gamma_i} p^{e_{H_i}} \right) = O(n^{-\gamma}).$$

We next bound the second sum. By the definition of p, we have that for some constant C > 0and large n,

$$\sum_{i=2}^{d} \sum_{j=1}^{i-1} \lambda_j \sum_{k \in \mathcal{K}_{i,j}} p^{-k} n^{-\ell_{k,i,j}} \leqslant C \sum_{i=2}^{d} \sum_{j=1}^{i-1} \sum_{k \in \mathcal{K}_{i,j}} n^{-(\ell_{k,i,j}-k/\alpha)}.$$

Now, observe that given (i, j) since  $\alpha = d_{H_i}$  and all the graphs have the same density and are strictly balanced, we have that  $\ell_{k,i,j} - k/\alpha \ge \gamma > 0$  since if a pair  $(k, \ell_{k,i,j})$  is in the sum, then the graph with k edges and  $\ell_{k,i,j}$  vertices is a subgraph of both  $H_i$  and  $H_j$ . Thus, the second sum is also  $O(n^{-\gamma})$  as n is large, which implies the desired result.

**Remark 3.5.** We note that it is possible to get results for subgraphs that are not strictly balanced, as Corollary 3.2 does not assume 'balancedness'. In particular, in the case where H has a unique densest subgraph, the desired result can be deduced immediately from our results. However, other cases are more delicate, with different subgraphs of H 'competing'. One would need to incorporate considerations similar to those in the determination the threshold of appearance of H, as was done by Bollobás [3, Section 4.2] to obtain approximation errors for particular graphs.

We end this section with an example of application of Theorem 3.4.

**Example 3.6** (Vector of cycles). Let  $d \ge 2$  and let  $H_1, \ldots, H_d$  be d distinct cycles, where each  $H_i$  is of length  $k_i \ge 3$ , with  $k_1 < k_2 < \cdots < k_d$ . Then we have

$$\mathbb{E}(W_i) = \frac{n(n-1)\cdots(n-k_i+1)}{2k_i}p^{k_i}.$$

For any two cycles  $H_i$  and  $H_j$  with i < j, the maximum shared edges is  $M_{i,j} = k_i - 1$ , and for  $k = 1, 2, ..., k_i - 1$ , the minimum vertices for k shared edges is  $\ell_{k,i,j} = k + 1$ . Since all cycles have density equal to one, taking  $p = cn^{-1}(1 + o(1))$ , by Theorem 3.4, we obtain, by noting that  $\gamma = 1$ , that for n sufficiently large,

$$d_{\mathrm{W}}(\mathbf{W}, \mathbf{P}_{\lambda}) \leqslant C n^{-1},$$

where  $\lambda_i = \mathbb{E}(W_i) \to c^{k_i}/(2k_i)$  for  $i = 1, \ldots, d$ .

### 3.2 Multivariate hypergeometric distribution

In this section we give Poisson approximation of the multivariate hypergeometric distribution, starting with its definition.

**Definition 3.7** (Multivariate hypergeometric distribution). Consider an urn containing N balls of d different colors, with  $n_i$  balls of color i (i = 1, ..., d) such that  $\sum_{i=1}^{d} n_i = N$ . When drawing m balls without replacement, let  $\mathbf{W} = (W_1, ..., W_d)$ , where  $W_i$  denotes the number of balls of color i in the sample. Then  $\mathbf{W}$  follows a multivariate hypergeometric distribution.

Next, as a consequence of the decreasing size-biased coupling in Theorem 2.9, we obtain the following Poisson approximation, which is the multivariate extension of [7, Example 4.32]. Observe that in this application of Theorem 2.9 the  $p_{i,j}$ 's do not depend on *i* nor *j*.

**Theorem 3.8.** Let  $\mathbf{W} = (W_1, \ldots, W_d)$  follow the multivariate hypergeometric distribution defined as above. Then

$$d_{W}(\mathbf{W}, \mathbf{P}_{\lambda}) \leqslant \sum_{i=1}^{d} \min\left\{1, \frac{mn_{i}}{N}\right\} \left(1 - \frac{(N-n_{i})(N-m)}{N(N-1)}\right) + \frac{2(N-m)}{N(N-1)} \sum_{i=2}^{d} \sum_{j=1}^{i-1} n_{j},$$

where  $\lambda_i = \mathbb{E}(W_i) = mn_i/N$  for each  $i = 1, \ldots, d$ .

**Remark 3.9.** As in the one-dimensional case, this approximation is useful when N is large and m/N and  $n_i/N$  are small for all i = 1, ..., d.

Proof. For each color  $i \in \{1, \ldots, d\}$ , note that  $W_i$  can be written as a sum of indicator random variables  $W_i = \sum_{j=1}^{n_i} X_j^i$ , where  $X_j^i$  is the indicator that ball j of color i is chosen in the m-sample for  $j = 1, \ldots, n_i$ . So  $\mathbb{P}(X_j^i = 1) = m/N$ . For  $\mathbf{W} = (W_1, \ldots, W_d)$ , we construct its size-biased coupling  $\widetilde{\mathbf{W}} = (\widetilde{\mathbf{W}}^1, \ldots, \widetilde{\mathbf{W}}^d)$  with the random vector  $\widetilde{\mathbf{W}}^i$  defined as follows for each  $i \in \{1, \ldots, d\}$ . Let  $I_i$  be uniformly distributed on  $\{1, \ldots, n_i\}$ , independent of all else. Then, if ball  $I_i$  of color i is already in the m-sample, let  $\widetilde{\mathbf{W}}^i = \mathbf{W}^i$ ; otherwise, we add ball  $I_i$  of color i to the sample, and remove a uniformly chosen ball from the current sample. Recall that the law of coupling from (2.3) specifies that the distribution of the coupled random vector matches the conditional distribution of the original vector given that ball  $\ell$  of color i was selected. Then we have, for all  $i = 1, \ldots, d$ , that we can write the  $\widetilde{\mathbf{W}}^i$  defined above as

$$\widetilde{\mathbf{W}}^{i} = \left(\sum_{j=1}^{n_{1}} X_{j}^{1,(i,I_{i})}, \dots, \sum_{j=1}^{n_{i-1}} X_{j}^{i-1,(i,I_{i})}, \sum_{j=1, j \neq I_{i}}^{n_{i}} X_{j}^{i,(i,I_{i})} + 1\right),$$

where for colors k < i,  $\sum_{j=1}^{n_k} X_j^{k,(i,I_i)}$  represents the count of color k balls after coupling, and for color i,  $\sum_{j=1, j\neq I_i}^{n_i} X_j^{i,(i,I_i)} + 1$  represents the count including the mandatory ball  $I_i$ . Then  $X_k^{j,(i,\ell)} \leq X_k^j$  for  $j = 1, \ldots, i-1$ ,  $\ell = 1, \ldots, n_i$ , and  $k = 1, \ldots, n_j$ , and for j = i and  $k \neq \ell$  we have  $X_k^{i,(i,\ell)} \leq X_k^j$ , and therefore,  $\widetilde{W}_j^i \leq W_j$  for all  $j \neq i$ , as no additional balls of any color other than i can be added during the coupling procedure. Thus, we can apply Theorem 2.9. Standard hypergeometric calculations yield

$$\mathbb{E}[W_i] = \frac{mn_i}{N} = \lambda_i, \text{ and } \operatorname{Var}(W_i) = \frac{mn_i(N - n_i)(N - m)}{N^2(N - 1)} = \lambda_i \frac{(N - n_i)(N - m)}{N(N - 1)}.$$
 (3.3)

For the covariance between counts with distinct colors  $i \neq j$ , similar calculations give

$$\operatorname{Cov}(W_i, W_j) = -\frac{mn_i n_j (N-m)}{N^2 (N-1)} = -\lambda_i \lambda_j \frac{N-m}{m(N-1)}.$$

Hence by Theorem 2.9, we have

$$d_{W}(\mathbf{W}, \mathbf{P}_{\lambda}) \leqslant \sum_{i=1}^{d} \min\{1, \lambda_{i}\} \left(1 - \frac{(N - n_{i})(N - m)}{N(N - 1)}\right) + \frac{2(N - m)}{m(N - 1)} \sum_{i=2}^{d} \sum_{j=1}^{i-1} \lambda_{j}.$$

Then the proof is complete in view of the expectation formula in (3.3).

### Acknowledgement

This project was initiated at the Probability and Combinatorics Workshop held at the Bellairs Research Institute in March 2024. The authors thank the organisers, Gábor Lugosi, Louigi Addario-Berry, and Nicolas Broutin. Special thanks go to Gábor Lugosi for useful comments, suggestions, and discussions. The first author acknowledges support from the Spanish MINECO grant PID2022-138268NB-100 and Ayudas Fundacion BBVA a Proyectos de Investigación Científica 2021.

# References

- [1] Richard Arratia, Larry Goldstein, and Louis Gordon. Two moments suffice for Poisson approximations: the Chen-Stein method, Annals of Probability, 17(1): 9–25, 1989.
- [2] Andrew Barbour, Lars Holst, and Svante Janson. *Poisson Approximation*, Oxford University Press, 1992.
- [3] Béla Bollobás. Random Graphs, 2nd ed. Cambridge University Press, 2001.
- [4] Larry Goldstein and Yosef Rinott. Multivariate normal approximations by Stein's method and size bias couplings, *Journal of Applied Probability*, 33(1):1–17, 1996.
- [5] Svante Janson, Tomasz Łuczak, and Andrzej Ruciński. Random Graphs, John Wiley & Sons, Inc., 2000.
- [6] Federico Pianoforte and Riccardo Turin. Multivariate Poisson and Poisson process approximations with applications to Bernoulli sums and U-statistics, Journal of Applied Probability, 60(1): 223–240, 2023.
- [7] Nathan Ross. Fundamentals of Stein's method, *Probability Surveys*, 8:210–293, 2011.
- [8] Krai Krokowski and Christoph Thäle. Multivariate central limit theorems for Rademacher functionals with applications, *Electronic Journal of Probability*, 22: 1–30, 2017.
- [9] Rui-Ray Zhang. The probability of non-existence of small substructures via clusters and cumulants. PhD thesis, Monash University, 2023.