Holomorphic geometric structures on Hopf manifolds

Matthieu Madera

Abstract

We construct integrable holomorphic G-structures and flat holomorphic Cartan geometries on every complex Hopf manifold, without using the normal forms given by the Poincaré-Dulac Theorem. We provide a new proof of the latter using charts adapted with the geometric structures.

Contents

| 1. | \mathbf{Intr} | oduction | 2 |
|----|-----------------|--|-----------|
| | 1.1. | Introduction to Hopf manifolds | 2 |
| | | Description of the results | 4 |
| 2. | Geo | ometric preliminaries | 6 |
| | 2.1. | Canonical forms on frame bundles | 7 |
| | 2.2. | Holomorphic G-structures on manifolds | 8 |
| | 2.3. | Holomorphic Cartan geometries | 10 |
| 3. | Res | onances of a contraction | 11 |
| 4. | Geo | metric structures of order 1 on Hopf manifolds | 12 |
| 5. | The | group of sub-resonant polynomial transformations | 23 |
| 6. | Geo | metric structures of higher order on Hopf manifolds | 27 |
| | 6.1. | High dimension | 28 |
| | | Hopf surfaces | 31 |
| | | 6.2.1. Mall theorem for Hopf surfaces | 32 |
| | | 6.2.2. Ornea and Verbitsky theorem for Hopf surfaces | 33 |
| | | 6.2.3. G-structures on resonant Hopf surfaces | 34 |

| 7. | Car | tan connection, integrability and Poincaré- Dulac theorem | 40 |
|----|------|---|----|
| | 7.1. | Another Lie group | 40 |
| | 7.2. | The Cartan geometry induced by the G-structures | 42 |
| | 7.3. | Integrability of the G-structure of order r+1 | 44 |
| | 7.4. | Poincaré-Dulac theorem | 47 |

1. Introduction

We study complex Hopf manifolds. This is a class of compact complex manifolds of any dimension $n \ge 2$ that do not admit kählerian metrics. As real manifolds of dimension 2n, they are in fact isomorphic to the product $\mathbb{S}^1 \times \mathbb{S}^{2n-1}$ and hence do not have the topology of symplectic manifolds.

The complex structures on $\mathbb{S}^1 \times \mathbb{S}^{2n-1}$, turning it into a Hopf manifold, are classified with the help of a famous dynamics theorem, the **Poincaré-Dulac Theorem** ([2], p.194). This theorem provides normal forms for Hopf manifolds.

The aim of this paper is to construct holomorphic geometric structures on Hopf manifolds of any dimension, without using the existence of normal forms. Throughout the paper, the term "geometric structure" can stand for several concepts. The main point of view is the one of G-structures of higher order (Section 2.2.), but we will also discuss about Gromov geometric structures (Section 2.2.) and Cartan geometries (Section 2.3.).

The reason why we do not want to take advantage of the Poincaré-Dulac Theorem is that one of the goals of the paper is to show the close relationship between the existence of a normal form for a given Hopf manifold and the existence of a holomorphic geometric structure on it, which can be seen as an integrable G-structure, or as a flat Cartan geometry.

In that sense, without any need of normal forms, we construct holomorphic Gstructures of higher order on every Hopf manifold and we show that they induce holomorphic Cartan geometries. Studying these structures, we prove that some G-structures are integrable (Definition 2.2.2) and that the induced Cartan geometry is flat. We infer that there exist special charts adapted with the integrable structure, providing a normal form for the Hopf manifold, and proving the Poincaré-Dulac Theorem.

1.1. Introduction to Hopf manifolds

A (primary) Hopf manifold is a compact complex manifold of dimension $n \geq 2$, having its universal covering space biholomorphic to $\mathbb{C}^n \setminus \{0\}$ and having an infinite cyclic fundamental group. As an example, Hopf [16] first considered actions of \mathbb{Z} on $\mathbb{C}^2 \setminus \{0\}$ of the type

$$\begin{array}{cccc} \mathbb{Z} \times \mathbb{C}^2 \setminus \{0\} & \longrightarrow & \mathbb{C}^2 \setminus \{0\} \\ (k, z) & \longmapsto & (\frac{1}{2})^k z \end{array}$$

This action of \mathbb{Z} preserves the standard torsion-free flat affine connection of $\mathbb{C}^2 \setminus \{0\}$, and therefore, the Hopf surface defined as the quotient of $\mathbb{C}^2 \setminus \{0\}$ under this action carries naturally a torsion-free flat affine connection. More generally, any Hopf manifold obtained as the quotient of $\mathbb{C}^n \setminus \{0\}$ by the action generated by an invertible linear map is naturally endowed with a torsion-free flat affine connection. Those Hopf manifolds are called **linear Hopf manifolds**.

A torsion-free affine connection can be seen from two points of view. First, we can see it as a Cartan geometry with model the affine group of \mathbb{C}^n and its closed subgroup the general linear group (see Section 2.3.). On the other hand, it can be seen as a $GL(n, \mathbb{C})$ -structure of order 2 (Example 2.2.2).

Borcea ([5], p.6) proved that the fundamental group of a Hopf manifold is generated by an element that has the topological properties of what is called a contraction of \mathbb{C}^n at 0 (Section 3.). Wehler ([33] for surfaces) and Haefliger ([15], Appendix) did work on the Kuranishi family of such a Hopf manifold. They took advantage of a wellknown theorem of holomorphic dynamics, the Poincaré-Dulac Theorem ([2], p.194). A contraction can be put, with a local holomorphic change of coordinates, under a normal form. This normal form is a resonant polynomial map ([2], p.194, Definition 3.0.1 for the definition of resonance). It appears that the space of normal forms of the germ of the contraction is closely related with the local deformation space of the complex structure of the corresponding Hopf manifold.

Using the Poincaré-Dulac theorem, McKay and Pokrovskiy ([23]) proved that any Hopf surface carries locally homogeneous structures. That is, the complex structure of a Hopf surface is always induced by an atlas of local charts with values in a suitable complex homogeneous space G/H. They classify all possible structures with model $\mathcal{O}(n)$ (the line bundle of degree n over the Riemann sphere $\mathbb{P}^1(\mathbb{C})$) on any Hopf surface, depending on the normal form of the contraction. They showed that every Hopf surface carries some flat Cartan geometries. These geometries generalize torsion-free flat affine connections that naturally exist on linear Hopf surfaces.

On the other hand, some authors did work on Hopf manifolds without using the Poincaré-Dulac Theorem, i.e. without assuming that the contraction defining the Hopf manifold is under a polynomial normal form. In this context, Mall ([21]) defined the notion of Mall bundles on Hopf manifolds. A Mall bundle on a Hopf manifold is a holomorphic vector bundle with a trivial pullback on the universal covering space $\mathbb{C}^n \setminus \{0\}$. Mall was able to give a full description of the cohomology of such a Mall bundle, using the fact that the linear operator given by the contraction on space of sections is compact. We will also use this property in Section 4., more precisely in the proof of Theorem 3.

Recently, keeping this idea of considering contractions under arbitrary forms, and using the link between the existence of an affine connection and cohomology of some tensor vector bundles, which are Mall bundles, Ornea and Verbitsky ([28]) proved that a Hopf manifold given by a non-resonant contraction (Definition 3.0.1) carries a unique affine connection, which is a posteriori torsion-free and flat. The contraction then preserves an affine atlas on \mathbb{C}^n , and an affine local system of coordinates around 0 linearises the contraction. This is a way, using the geometry of Hopf manifolds, to prove the non-resonant case of the Poincaré-Dulac Theorem, also known as the Poincaré linearization Theorem ([2], p.193). In [22] (p.21, lemma 14.2), McKay proved also that a Hopf manifold admitting a holomorphic affine connection (not necessarily torsion-free, nor flat) is isomorphic to a linear Hopf manifold, an isomorphism being given by the exponential chart at 0 of the pulled back affine connection extended to \mathbb{C}^n .

During the development of this work, we had the opportunity to discuss with Paul Boureau, who also presents in [6] new results on holomorphic geometric structures on Hopf manifolds, using a different approach. Indeed, he offers a re-examination of Poincaré-Dulac normal forms and the study of the group of sub-resonant polynomials, and uses these normal forms to exhibit locally homogeneous geometric structures on Hopf manifolds of any dimension. This result generalizes those of McKay and Pokrovskiy, who dealt in [23] with the case of Hopf surfaces.

1.2. Description of the results

We propose a generalization of the paper of Ornea and Verbitsky [28]. Given a contraction, we consider a group G_{β}^r of sub-resonant polynomial transformations of maximal degree r of \mathbb{C}^n (Section 5.). This group depends only on the eigenvalues of the differential at 0 of the contraction. The parameter β stands for the collection of eigenvalues and their multiplicity, and the parameter r is the maximal length of the resonances among the eigenvalues (Section 3.).

We construct a G_{β}^{1} -structure of order 1 on the Hopf manifold, where the group G_{β}^{1} is given by the jets of order 1 at 0 of the elements of the group G_{β}^{r} (Section 2.2. for the definition of *G*-structure, Section 4. Theorem 3 for the construction). More specifically, we prove that the contraction preserves an entanglement of vector sub-bundles of the holomorphic tangent bundle $T\mathbb{C}^{n}$. As a consequence, the Hopf manifold carries various sub-bundles of its holomorphic tangent bundle. For example, carrying a G_{β}^{1} -structure implies that there is a complete flag of vector sub-bundle of the holomorphic tangent bundle. This is the first step in the construction of *G*-structures of higher order on Hopf manifolds.

Given such a structure of order 1, using cohomology arguments on Mall bundles, we prove that the structure extends uniquely to a G_{β}^2 -structure of order 2, where the group G_{β}^2 is given by the jets of order 2 of elements of G_{β}^r (Section 6., Theorem 4 and Section 6.2.3. for surfaces).

This is a generalization of the cohomological technique used by Ornea and Verbitsky in [28]. Notice that the extension of our G^1_β -structure of order 1 is a generalization of an affine connection, but in the latter case, there is no need to reduce the group of the order 1 frame bundle. A torsion-free affine connection is indeed a $GL(n, \mathbb{C})$ - structure of order 2 ([7], p.68). In the non-resonant case, the holomorphic tangent bundle TM of the Hopf manifold admits a holomorphic affine connection because the space $H^1(M, TM^* \otimes TM^* \otimes TM)$ is trivial. A torsion-free affine connection is an equivariant way of extending all the jets of order 1 of charts to jets of order 2. In the resonant case, we have to consider cohomology in some vector bundles taking into account the resonances and we show that the first cohomology space of this vector bundle is trivial. With these considerations, we show that we can extend some jets of order 1 of charts, namely those given by a G^1_{β} -structure of order 1, into equivalent classes of jets of order 2. The latter equivalent classes are precisely given by what we call the sub-resonant monomials (Section 6.).

By induction, we can prove that every Hopf manifold admits special G-structures of every order (Section 6., Theorem 4), except for surfaces (Section 6.2.3.), where we prove that, a priori, the structure extends up to order r + 2.

Since the group G_{β}^{r} is a group of global polynomial transformations of degree at most r, the extension from the G_{β}^{r} -structure of order r to a structure of order r + 1 is realized by an equivariant way of extending jets of order r of the holomorphic charts given by the structure into jets of order r + 1 (see Section 7.2.). Using this equivariant section, we can pull back the canonical 1-form of the frame bundle of order r + 1 (see [18] and Section 2.1.) on the structure of order r, and we prove in Theorem 9, that it leads to a holomorphic Cartan geometry (see [31] and Section 2.3. for the definitions), with model defined in Section 7.1.. The strategy of the proof is the following. The pullback of the canonical form induces a horizontal distribution of the frame bundle of order r defined over the reduction (see Section 2. for the definition of the frame bundle of higher order). The defect of this distribution to be actually tangent to the reduction is measured by a holomorphic section of a Mall bundle. We prove that such a section, being invariant by the contraction and containing the data of the resonances, has to vanish.

The fact that the previous horizontal distribution is tangent to the reduction is a strong tool that will help us to prove that the G_{β}^{r} -structure of order r + 1 turns out to be integrable (see Definition 2.2.2). We will use a Frobenius-type theorem, given by Benoist in his paper bringing his point of view on the demonstration of Gromov's opendense orbit Theorem ([3], Theorem p.12, Section 2.2.). The existence of an integrability atlas is equivalent to finding solutions of a partial derivatives relation. Here, the completeness (Definition 2.2.5) of this relation is given by the equivariant section mentioned above. Roughly speaking, it corresponds to the rigidity in the sense of Gromov geometric structures (see Definition 2.2.4). The consistence is obtained by considering the extension of the structure of order r up to order r + 2 which, by the same argument as before, produces a Cartan geometry on the reduction of order r + 1.

We notice that, in the case of Hopf surfaces, the flat Cartan geometries we construct are part of McKay and Pokrovskiy's classification of $\mathcal{O}(n)$ -structures on Hopf surfaces in [23]. What is new here is that we construct those structures without using the normal form of the contraction.

The existence of this integrable structure, and *a fortiori*, a flat Cartan geometry (Theorem 7.3.1), is equivalent to the existence of special charts adapted to the geometric structure, which provide new coordinates where the contraction has a polynomial normal form. Therefore, we infer a proof of the Poincaré-Dulac Theorem in this context of global contractions, generalizing the proof given by Ornea and Verbitsky in [28] for the non-resonant case (see Theorem 11).

For a Hopf manifold given by a contraction in normal form, the geometric structures we consider arise naturally, as in the case of linear Hopf manifolds. What we prove in this paper is the converse. A contraction has to preserve some integrable G-structures, or equivalently some flat Cartan geometries, and special charts for those structures provide the change of coordinates putting the contraction under a normal form.

The main steps of the paper are the following. In Section 4., we prove the existence of structures of order 1. In Section 6., the induction step in the construction of geometric structures of higher order is explained. In Section 7. we explain how a Cartan geometry is induced by the *G*-structures we consider (Section 7.2.), we prove the integrability of the structure of order r + 1 and the flatness of the Cartan geometry (Section 7.3.) and we deduce a proof of the Poincaré-Dulac Theorem (Section 7.4.).

2. Geometric preliminaries

In this section, M is a complex manifold of dimension n and r is a positive integer.

We denote by $\mathcal{D}^r(\mathbb{C}^n)$ the set of equivalence classes of germs of biholomorphic maps $(\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ under the relation "having the same Taylor polynomial of order r at 0". Endowed with the truncated composition, $\mathcal{D}^r(\mathbb{C}^n)$ is a complex Lie group ([3]). We will denote by $\mathfrak{d}^r(\mathbb{C}^n)$ its Lie algebra. By convention, $\mathfrak{d}^0(\mathbb{C}^n)$ is the trivial Lie algebra. As an example, the group $\mathcal{D}^1(\mathbb{C}^n)$ is isomorphic to the general linear group $GL(n,\mathbb{C})$. For $r \geq s \geq 0$, we denote by $\rho_{r,s} : \mathcal{D}^r(\mathbb{C}^n) \to \mathcal{D}^s(\mathbb{C}^n)$ the standard projection, with the convention $\mathcal{D}^0(\mathbb{C}^n) = \{e\}$. This is a complex Lie groups homomorphism.

On the complex manifold M, we consider the set of equivalence classes of germs of biholomorphic maps $(\mathbb{C}^n, 0) \to (M, m)$ under the relation "having the same Taylor polynomial of order r at 0 in any fixed holomorphic chart in a neighborhood of $m \in M^{"}$. The collection of all those sets patches to form a principal $\mathcal{D}^r(\mathbb{C}^n)$ bundle called the order r frame bundle over M, with total space denoted by $\mathcal{R}^r(M)([3])$. Notice that $\mathcal{R}^1(M)$ is isomorphic to the bundle of linear frames of the holomorphic tangent bundle TM of M. For $r \geq s \geq 0$, we denote by $\pi_{r,s} : \mathcal{R}^r(M) \to \mathcal{R}^s(M)$ the standard projection, with the convention $\mathcal{R}^0(M) = M$. This is a homomorphism of principal bundles, with underlying Lie group homomorphism $\rho_{r,s}$.

2.1. Canonical forms on frame bundles

We recall here the definition and the main properties of the canonical forms on the frame bundles over a complex manifold M.

First, we notice that the translations of \mathbb{C}^n give rise to a global section of all the frame bundles of \mathbb{C}^n , and hence global trivializations

$$\begin{array}{rccc} \xi_r & : & \mathcal{R}^r(\mathbb{C}^n) & \longrightarrow & \mathbb{C}^n \times \mathcal{D}^r(\mathbb{C}^n) \\ & & j_0^r f & \longmapsto & (f(0), j_0^r(\theta_{-f(0)}f)) \end{array}$$

where θ_z is the translation of vector $z \in \mathbb{C}^n$.

We also notice that if U and V are complex manifolds, and if $f: U \to V$ is a holomorphic map, then it induces a holomorphic map

$$\begin{array}{cccc} j^r f & : & \mathcal{R}^r(U) & \longrightarrow & \mathcal{R}^r(V) \\ & & j_0^r g & \longmapsto & j_0^r(fg). \end{array}$$

We define ([24], p.14), for $r \ge 0$, the following holomorphic 1-form on $\mathcal{R}^{r+1}(M)$, called the canonical form of the r+1 order frame bundle

$$\begin{array}{rcl} \chi_{r+1} & : & T\mathcal{R}^{r+1}(M) & \longrightarrow & T_0\mathbb{C}^n \times \mathfrak{d}^r(\mathbb{C}^n) \\ & & (j_0^{r+1}f, X) & \longmapsto & \mathrm{d}_{j_0^r id}\xi_r((\mathrm{d}_{j_0^r id}j^r f)^{-1}(d_{j_0^{r+1}f}\pi_{r+1,r}(X))) \end{array}$$

This canonical form illustrates the following fact. For $r \geq 0$, an element $j_0^{r+1}f$ of $\mathcal{R}^{r+1}(M)$ can be seen as a horizontal plane for the bundle $\pi_{r,0} : \mathcal{R}^r(M) \to M$ at $j_0^r f$. Together with the standard trivialization of the vector bundle of vertical tangent vectors of $\mathcal{R}^r(M)$ ($\pi_{r,0}$ is a principal bundle) and the fact that $j_0^1 f$ gives a linear frame of $T_{f(0)}M$, $j_0^{r+1}f$ is naturally seen as a frame of order 1 of $\mathcal{R}^r(M)$ at $j_0^r f$. The canonical form at a vector X is obtained by reading the coordinates of the projection of X in this frame. Notice that for r = 0 the form on $\mathcal{R}^1(M)$ is given by reading the coordinates of a tangent vector of M in the given frame.

We will now give some properties of the canonical form. For $r \ge 0$, we consider the following group representation

$$\frac{\underline{Ad}^{r+1}}{j_0^{r+1}\phi} : \mathcal{D}^{r+1}(\mathbb{C}^n) \longrightarrow \underbrace{\underline{Ad}_{j_0^{r+1}\phi}^{r+1}}_{p_0^{r+1}\phi} := \mathrm{d}_{J_0^r id}(j_0^r f \mapsto j_0^r(\phi f \phi^{-1}))$$

Using ξ_r , we can consider that \underline{Ad}^{r+1} takes values in $GL(T_0\mathbb{C}^n \times \mathfrak{d}^r(\mathbb{C}^n))$. Using this representation, we have the following proposition.

Proposition 2.1.1. ([24], p.14) For $r \ge 0$, the canonical form χ_{r+1} satisfies the following properties:

- 1. $\forall v \in \mathfrak{d}^{r+1}(\mathbb{C}^n), \ \chi_{r+1}(X_v) \equiv (0, \mathrm{d}_e \rho_{r+1,r}(v)), \ where \ X_v \ is \ the \ fundamental \ vector field \ of \ \mathcal{R}^{r+1}(M) \ generated \ by \ v \ ;$
- 2. $\forall j_0^{r+1} \phi \in \mathcal{D}^{r+1}(\mathbb{C}^n), \ \mathcal{R}^*_{j_0^{r+1}\phi} \chi_{r+1} = \underline{Ad}^{r+1}_{j_0^{r+1}\phi^{-1}} \chi_{r+1}.$

2.2. Holomorphic G-structures on manifolds

Let G be a complex Lie subgroup of $\mathcal{D}^r(\mathbb{C}^n)$. We recall the definition of a G-structure of order r on the manifold M.

Definition 2.2.1. ([1], p.152 Example 5) A G-structure of order r on M is a principal G sub-bundle of the order r frame bundle of M. Equivalently, it is a global section of the quotient bundle $\mathcal{R}^r(M)/G$.

Roughly speaking, a G-structure of order r on a manifold is a system of local sections of the r order frame bundle (r-frames) such that, on the overlaps, the changes of frames are everywhere elements of G.

Example 2.2.1. ([32], p.315) On $M = \mathbb{C}^n$, the standard global coordinates define a trivialization of the frame bundle of any order over \mathbb{C}^n . For every r, this trivialization is a $\{e\}$ -structure of order r, where $\{e\}$ is the trivial group. Moreover, for every Lie subgroup G of $\mathcal{D}^r(\mathbb{C}^n)$, this $\{e\}$ -structure induces a G-structure of order r on \mathbb{C}^n . We call this G-structure the standard G-structure of \mathbb{C}^n .

Definition 2.2.2. ([1], p.187) A G-structure of order r over M is said to be integrable if it is locally isomorphic to the standard G-structure of \mathbb{C}^n , i.e. if there exists a holomorphic atlas on M such that on every open set of this atlas, the chart sends the standard local section of $\mathcal{R}^r(\mathbb{C}^n) \to \mathbb{C}^n$ to a local section of the G-structure over M.

- **Example 2.2.2.** 1. ([19], p.13 Example 2.11) An absolute parallelism on M is a trivialization of the tangent bundle of M. Equivalently, this is a global section of the order 1 frame bundle $\mathcal{R}^1(M)$. Therefore, it is a $\{e\}$ -structure of order 1. The parallelism is given by n holomorphic vector fields, pointwise linearly independents. The structure is therefore integrable if and only if the vector fields commute two by two.
 - 2. ([1], p.185 Example 2) A distribution of dimension d in the tangent bundle is a Gstructure of order 1 with the group G defined as the subgroup of $GL(n, \mathbb{C})$ composed of the elements preserving a d-dimensional vector subspace of \mathbb{C}^n . By the classical Frobenius Theorem, a distribution is integrable (and produces a foliation) if and only if it is involutive under the Lie bracket.
 - 3. ([7], p.68) The group $GL(n, \mathbb{C})$ is naturally a Lie subgroup of $\mathcal{D}^2(\mathbb{C}^n)$. A $Gl(n, \mathbb{C})$ structure of order 2 on M is a torsion-free holomorphic affine connection. The
 structure is integrable if and only if the curvature tensor vanishes.

Another notion is the notion of rigidity in the context of Gromov geometric structures. We recall here the definitions ([13], §0, [3], Section 1.2 p.4).

Let Z be a complex manifold endowed with a holomorphic action of $\mathcal{D}^r(\mathbb{C}^n)$.

Definition 2.2.3. ([13] §0, [3] p.4) A Gromov geometric structure of type Z and of order r on M is a holomorphic $\mathcal{D}^r(\mathbb{C}^n)$ -equivariant map $g: \mathcal{R}^r(M) \to Z$.

Remark 2.2.1. In [3], the manifold Z and its action of $\mathcal{D}^r(\mathbb{C}^n)$ are supposed to be algebraic. This assumption is fundamental for the open-dense orbit Theorem, but we will not use it here. Nevertheless, we notice that every Gromov geometric structure we consider in this paper is actually of algebraic type, because they are G-structures with G an algebraic subgroup of $\mathcal{D}^r(\mathbb{C}^n)$.

For G a subgroup of $\mathcal{D}^r(\mathbb{C}^n)$, a G-structure of order r is a Gromov geometric structure of type $\mathcal{D}^r(\mathbb{C}^n)/G$. Indeed, if $g: \mathcal{R}^r(M) \to \mathcal{D}^r(\mathbb{C}^n)/G$ is a $\mathcal{D}^r(\mathbb{C}^n)$ -equivariant map, then the subset $g^{-1}(\{id\})$ of $\mathcal{R}^r(M)$ is naturally endowed with a structure of holomorphic G-principal sub-bundle of $\mathcal{R}^r(M)$.

Definition 2.2.4. ([3], p.7) A Gromov geometric structure of type Z and of order r on M is said to be rigid at order $s \ge r-1$ if for every $x \in M$ and every t > s, a t-jet of biholomorphism of (M, x) that preserves the structure is completely determined by its s-jet.

- **Example 2.2.3.** 1. ([19], p.15) A jet of a germ of biholomorphism fixing a point of M and preserving an absolute parallelism is in fact the jet of the identity automorphism of M. Therefore, an absolute parallelism is rigid at order 0.
 - 2. A distribution in the tangent bundle is not rigid. As an example, we consider a germ of biholomorphism of \mathbb{C}^2 fixing 0 and preserving the foliation generated by the vector field $\frac{\partial}{\partial z_1}$ in the standard coordinates of \mathbb{C}^n . Then, in standard coordinates, the germ of biholomorphism reads

$$(z_1, z_2) \mapsto (f(z_1, z_2), g(z_2))$$

Conversely, any germ of this form preserves the foliation, showing that the latter is not rigid.

3. ([3], p.7) A torsion-free holomorphic affine connection is rigid at order 1, since a jet of biholomorphism preserving a point is conjugated by the exponential chart with its differential at the fixed point.

In his paper bringing his point of view on the demonstration of Gromov's open-dense orbit Theorem ([3], p.1), Benoist proved a Frobenius type theorem for existence and uniqueness of solutions of some partial derivatives relations.

If M and N are complex manifolds, we denote by $J^r(M, N)$ the bundle of r-jets of holomorphic maps from M to N. A partial derivatives relation is a complex submanifold R of $J^r(M, N)$ and a solution is a local holomorphic map $f : M \to N$ such that the section $j^r f : M \to J^r(M, N)$ induced by f takes values in R. **Definition 2.2.5.** ([3], p.12) (1) A partial derivatives relation $R \subset J^r(M, N)$ is said to be complete if the natural projection $J^r(M, N) \to J^{r-1}(M, N)$ induces an isomorphism between R and its image.

(2) A partial derivatives relation $R \subset J^r(M, N)$ is said to be consistent if, for every $p \in R$, there exists a section $M \to R$ through p and tangent at p to a holonomic section, *i.e.* a section of the form $j^r f$, for $f : M \to N$.

Using these two notions, Benoist proved the following.

Theorem 1. ([3], p.12) Let R be a complete and consistent partial derivatives relation. Then for every $p \in R$, there exists a unique germ of solution through p.

2.3. Holomorphic Cartan geometries

We will recall the concept of holomorphic Cartan geometry on a complex manifold M of dimension n. The main references here are [31] and [4].

Let G be a complex Lie group with Lie algebra \mathfrak{g} , and let H be a complex Lie subgroup with Lie algebra \mathfrak{h} .

Definition 2.3.1. ([31], p.184 Definition 3.1) A holomorphic Cartan geometry of type (G, H) on M is a holomorphic H-principal bundle $\pi : P \to M$ endowed with a holomorphic one form $\eta : TP \to \mathfrak{g}$ satisfying the following:

1. $\forall p \in P, \eta_p : T_p P \to \mathfrak{g} \text{ is an isomorphism (i.e. } \eta \text{ is an absolute parallelism on } P)$

2. $\forall v \in \mathfrak{h}, \eta(X_v) \equiv v$, where X_v is the fundamental vector field generated by v on P

3.
$$\forall h \in H, \ \mathcal{R}_h^* \eta = Ad_{h^{-1}}(\eta).$$

Example 2.3.1. A holomorphic affine connection on M is a holomorphic Cartan geometry of type $(Aff(n, \mathbb{C}), GL(n, \mathbb{C})))$. Indeed, it is an absolute parallelism of the order one frame bundle $\eta : T\mathcal{R}^1(\mathbb{C}^n) \to \mathfrak{aff}(n, \mathbb{C})$. The principal connection on $\mathcal{R}^1(M) \to M$ is given by the $Ad_{GL(n,\mathbb{C})}$ -invariant splitting $\mathfrak{aff}(n,\mathbb{C}) = T_0\mathbb{C}^n \oplus \mathfrak{gl}(n,\mathbb{C})$. On the other hand, given a principal connection on $\mathcal{R}^1(M)$, we obtain the Cartan geometry by adding the data of the canonical form (see Section 2.1.).

We recall the central notion of curvature and a remark on this tensor field.

Definition 2.3.2. ([31], p.184) Let $\eta : TP \to \mathfrak{g}$ be a holomorphic Cartan geometry on M. The holomorphic 2-form on P defined by $\Omega := \mathrm{d}\eta + \frac{1}{2}[\eta, \eta]$ is called the curvature form. The Cartan geometry η is said to be flat if the form Ω is identically zero.

Proposition 2.3.1. ([11], p.4) Let $\eta : TP \to \mathfrak{g}$ be a holomorphic Cartan geometry on M. The curvature form Ω can be seen as a holomorphic 2-form on M with value in the \mathfrak{g} -adjoint bundle $Ad_{\mathfrak{g}}(P)$ of P defined as the associated bundle

$$Ad_{\mathfrak{g}}(P) := P \times_H \mathfrak{g}$$

The interest of the curvature is the following Theorem that asserts that a flat Cartan geometry is locally modeled on the standard Cartan geometry.

Theorem 2. ([31], p.212 Theorem 5.1) Let η be a holomorphic Cartan geometry of type (G, H) on a bundle $P \to M$. Then η is flat if and only if at each x of M, there exists a neighborhood of x and a local isomorphism between the Cartan geometry η and the bundle $G \to G/H$ equipped with the Maurer-Cartan form of G ([31], p.98, definition 1.3).

3. Resonances of a contraction

Let $n \geq 2$ be an integer and $\gamma : \mathbb{C}^n \to \mathbb{C}^n$ be a biholomorphic contraction in 0, i.e. a map such that for any compact K and any neighborhood U of 0, a sufficiently high power of γ maps K in U. Denote by M the corresponding Hopf manifold, defined as the orbit space of the fixed point free and properly discontinuous action of the group \mathbb{Z} on $\mathbb{C}^n \setminus \{0\}$ generated by γ , with its compact complex manifold structure given by the quotient map ([15]). We will prove in Section 6. that M carries geometric structures, encoded in the contraction γ and, even more, encoded in the Jacobian of γ at 0.

Let $d_0\gamma: T_0\mathbb{C}^n \to T_0\mathbb{C}^n$ be the differential map of γ at 0. Haefliger ([15], Appendix A.2) proved that the eigenvalues of this linear map are of modulus smaller than 1. Up to a linear change of coordinates of \mathbb{C}^n , we can assume that the Jacobian matrix of γ at 0 has, in the standard coordinates, the form

$$J_{0}\gamma = \begin{pmatrix} \alpha_{1} & (0) & & & \\ * & & & (0) \\ * & * & \alpha_{1} & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \alpha_{m} & (0) \\ (0) & & & & \\ & & & & * & * & \alpha_{m} \end{pmatrix}$$

where

$$1 > |\alpha_1| \ge \dots \ge |\alpha_m| > 0$$

and the block corresponding to the eigenvalue α_i is of size n_i , for $j \in \{1, ..., m\}$.

We denote also by u_0 the order one Taylor jet of γ at 0. The map $u_0 : \mathbb{C}^n \to \mathbb{C}^n$ is complex linear, and its matrix in the standard basis of \mathbb{C}^n is also given by $J_0\gamma$. In

all that follows, α will be used to represent the data of the eigenvalues $(\alpha_1, ..., \alpha_m)$ and their multiplicity $(n_1, ..., n_m)$.

With this notations we will use the following central definition.

Definition 3.0.1. ([5], p.1) A α -resonance, or simply a resonance if there is no ambiguity, is a relation of the form

$$\alpha_j = \prod_{k=1}^m \alpha_k^{p_k}$$

where j is an element of $\{1, ..., m\}$ and $p = (p_1, ..., p_m)$ is a nonzero element of \mathbb{N}^m . We will say that α_i is dependent on α_k if there exists a resonance of the form

$$\alpha_j = \alpha_k \prod_{l=1}^m \alpha_l^{p_l}$$

and independent if there exist no resonance of this type.

Remark 3.0.1. Some authors do not consider the trivial relation $\alpha_j = \alpha_j$ as a resonance. In what follows, to be as general as possible, we will need to say that α_j is dependent of α_j .

4. Geometric structures of order 1 on Hopf manifolds

We will show that every Hopf manifold carries a G-structure of order 1, with G a linear complex Lie group induced by the set of eigenvalues α . Namely, let G^1_{α} be the complex Lie group of invertible triangular inferior matrices $A = (a_{i,j})_{i,j}$ such that for i > j, $a_{i,j} = 0$ if α_{k_i} is independent on α_{k_j} , where k_i is the element of $\{1, ..., m\}$ such that α_{k_i} is the diagonal coefficient of $J_0\gamma$ in the row i, and same for k_j (see Section 3.).

Example 4.0.1. Assume that n = 3 and that γ is such that

$$J_0\gamma = \begin{pmatrix} \frac{1}{2} & 0 & 0\\ 0 & \frac{1}{3} & 0\\ 0 & 0 & \frac{1}{4} \end{pmatrix}$$

then G^1_{α} is the group of invertible matrices of the form

$$\begin{pmatrix} * & 0 & 0 \\ 0 & * & 0 \\ * & 0 & * \end{pmatrix}.$$

If γ is such that

$$J_0 \gamma = \begin{pmatrix} \frac{1}{2} & 0 & 0\\ 0 & \frac{1}{3} & 0\\ 0 & 0 & \frac{1}{6} \end{pmatrix}$$

then G^1_{α} is the group of invertible matrices of the form

$$\begin{pmatrix} * & 0 & 0 \\ 0 & * & 0 \\ * & * & * \end{pmatrix}.$$

If γ is such that

$$J_0 \gamma = \begin{pmatrix} \frac{1}{2} & 0 & 0 & 0\\ 1 & \frac{1}{2} & 0 & 0\\ 0 & 0 & \frac{1}{4} & 0\\ 0 & 0 & 0 & \frac{1}{8} \end{pmatrix}$$

then G^1_{α} is the group of invertible matrices of the form

$$\begin{pmatrix} * & 0 & 0 & 0 \\ * & * & 0 & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix}.$$

In this section, we are interested in G^1_{α} -structures of order 1 (Section 2.2. for the definition of *G*-structures). To understand G^1_{α} -structures of order 1, we have to describe the group G^1_{α} as a subgroup of $GL(n, \mathbb{C})$. We will realize G^1_{α} as the subgroup of $GL(n, \mathbb{C})$ preserving an entanglement of sub-spaces of \mathbb{C}^n , under the standard action of $GL(n, \mathbb{C})$ on \mathbb{C}^n .

For every $j \in \{1, ..., m\}$, we define I_j the set of non-trivial dependence indices of α_j i.e.

 $I_j = \{0\} \cup \{l \in \{1, ..., j-1\} \mid \alpha_j \text{ depends on } \alpha_l\}.$

Using also the notation $V_{0,0} = \mathbb{C}^n$, the invertible linear maps that preserve the family of sub-spaces

$$V_{j,k} := \left(\bigcap_{l \in I_j} V_{l,0}\right) \cap \operatorname{Vect}\{e_i, i \in \{1, ..., n\} \setminus \{n_1 + ... + n_{j-1} + 1, ..., n_1 + ... + n_{j-1} + n_j - k\}\}$$

for every $j \in \{1, ..., m\}$ and $k \in \{0, ..., n_j - 1\}$, are exactly the elements of G^1_{α} . The vectors $e_1, ..., e_n$ denote the standard basis of \mathbb{C}^n .

Example 4.0.2. Assume that G^1_{α} is the group of invertible matrices of the form

$$\begin{pmatrix} * & 0 & 0 \\ 0 & * & 0 \\ * & 0 & * \end{pmatrix},$$

then it is the group of invertible matrices that preserve the spaces

$$Vect\{e_2, e_3\}$$
$$Vect\{e_1, e_3\}$$
$$Vect\{e_2\}.$$

Assume that G^1_{α} is the group of invertible matrices of the form

$$egin{pmatrix} * & 0 & 0 & 0 \ 0 & * & 0 & 0 \ 0 & * & * & 0 \ * & 0 & 0 & * \end{pmatrix},$$

then it is the group of invertible matrices that preserve the spaces

$$Vect\{e_{2}, e_{3}, e_{4}\} \\Vect\{e_{1}, e_{3}, e_{4}\} \\Vect\{e_{1}, e_{4}\} \\Vect\{e_{2}, e_{3}\}.$$

Therefore, a G^1_{α} -structure of order 1 on a manifold is an entanglement of sub-bundles of the tangent bundle. The following theorem asserts that every Hopf manifold carries such a structure, descending from an invariant structure on $\mathbb{C}^n \setminus \{0\}$ extending to \mathbb{C}^n .

Theorem 3. There exists a G^1_{α} -structure on \mathbb{C}^n invariant under the action of $\langle \gamma \rangle$.

Proof. Since γ is a contraction, there exists an open neighborhood D of 0 in \mathbb{C}^n such that $\gamma(D)$ is bounded and included in D. Then, the space of bounded holomorphic one forms on D (for a given hermitian metric) $H_b^0(D, \Omega_D^1)$ is a Banach space, and the pull-back operator

$$\gamma^*: H^0_b(D, \Omega^1_D) \longrightarrow H^0_b(D, \Omega^1_D)$$

is compact ([21], Proposition 4.1). The compactness of this operator allows to use the powerful Riesz-Schauder theorem ([10], Section 5.2, [25] and [28] for some use of the Riesz-Schauder theorem in the context of Hopf manifolds).

We will construct a G^1_{α} -structure by use of an induction argument. We start with the base case.

Looking at the form of the Jacobian matrix of $d_0\gamma$, we see that the 1-form dz_{n_1} is not in the image of the operator $\gamma^* - \alpha_1 Id$, because there can be no linear forms ω_0 on $T_0\mathbb{C}^n$ such that $\omega_0 \circ d_0\gamma - \alpha_1\omega_0 = d_0z_{n_1}$. Therefore, the operator $\gamma^* - \alpha_1 Id$ is not onto, and since α_1 is not equal to 0, we infer, using the Riesz-Schauder theorem, that α_1 is an eigenvalue of γ^* and that there exists a positive integer N_1 , that we can assume to be greater or equal than n_1 , such that the space ker $((\gamma^* - \alpha_1 Id)^{N_1})$ is finite dimensional and such that the following decomposition holds

$$H_b^0(D,\Omega_D^1) = \ker((\gamma^* - \alpha_1 Id)^{N_1}) \oplus \overline{\operatorname{im}((\gamma^* - \alpha_1 Id)^{N_1})}.$$

This allows us to consider first an arbitrary nonzero element ω of ker $(\gamma^* - \alpha_1 Id)$. We will show that ω does not vanish at 0. To do that, we write

$$\omega = \sum_{j=1}^{n} \sum_{k \ge 0} P_{j,k} dz_j$$

where $P_{j,k}$ is a homogeneous polynomial map of degree k. Since ω is nonzero, let k_0 be the smallest non negative integer such that one of the P_{j,k_0} is nonzero, and let j_0 be the largest integer in $\{1, ..., n\}$ such that P_{j_0,k_0} is nonzero. Then the equation

$$\gamma^*\omega = \alpha_1\omega$$

is equivalent to

$$\sum_{j=1}^{n} \sum_{k \ge k_0} (P_{j,k} \circ \gamma) d\gamma_j = \alpha_1 \sum_{j=1}^{n} \sum_{k \ge k_0} P_{j,k} dz_j$$

where $\gamma = (\gamma_1, ..., \gamma_n)$, which is itself equivalent to

$$\sum_{j=1}^{n} \left(\sum_{l=1}^{n} \left(\sum_{k \ge k_0} P_{l,k} \circ \gamma \right) \frac{\partial \gamma_l}{\partial z_j} \right) dz_j = \sum_{j=1}^{n} \left(\alpha_1 \sum_{k \ge k_0} P_{j,k} \right) dz_j.$$

This last equation is equivalent to the identification of two holomorphic functions, for every $j \in \{1, ...n\}$, namely

$$\sum_{l=1}^{n} \left(\sum_{k \ge k_0} P_{l,k} \circ \gamma \right) \frac{\partial \gamma_l}{\partial z_j} = \alpha_1 \sum_{k \ge k_0} P_{j,k}.$$

This implies, choosing $j = j_0$ and looking at the Taylor polynomial of order k_0 , the following equation

$$\sum_{l=1}^{j_0} (P_{l,k_0} \circ u_0) \frac{\partial \gamma_l}{\partial z_j}(0) = \alpha_1 P_{j_0,k_0}.$$

But looking at the form of the Jacobian matrix $J_0\gamma$ again, we obtain that

$$\alpha P_{j_0,k_0} \circ u_0 = \alpha_1 P_{j_0,k_0}$$

where α is the eigenvalue of $J_0 \gamma$ lying at row and column j_0 , say α_r with $r \in \{1, ..., m\}$. This proves that $\frac{\alpha_1}{\alpha_r}$ is an eigenvalue of the linear endomorphism

$$\begin{array}{cccc} \mathbb{C}^{\mathrm{Hom}}_{k_0}[Z_1,...,Z_n] &\longrightarrow & \mathbb{C}^{\mathrm{Hom}}_{k_0}[Z_1,...,Z_n] \\ P &\longmapsto & P \circ u_0 \end{array}$$

where the $\mathbb{C}_{k_0}^{\text{Hom}}[Z_1, ..., Z_n]$ is the space of homogeneous polynomials of degree k_0 with n variables. But the eigenvalues of this endomorphism are precisely all the possible products of k_0 eigenvalues of $J_0\gamma$. Therefore, if k_0 is positive, then there is a resonance

$$\alpha_1 = \alpha_r \prod_{j=1}^m \alpha_j^{p_j}$$

where the length of $p = (p_1, ..., p_m)$ is k_0 , therefore positive. But this kind of relation is impossible, since

$$1 > |\alpha_1| \ge \dots \ge |\alpha_m| > 0.$$

Hence, $k_0 = 0$ (and r = 1), which proves that ω does not vanish at 0.

Since the space ker($(\gamma^* - \alpha_1 Id)^{N_1}$) is finite dimensional, say of dimension r_1 , and stable by γ^* , let $(\omega_1, ..., \omega_{r_1})$ be a basis of ker $((\gamma^* - \alpha_1 Id)^{N_1})$ such that γ^* is triangular superior in this basis. We need to prove that the family $(\omega_1, ..., \omega_{r_1})$ is pointwise linearly independent. Let us prove it at 0. Assume that $\lambda_1, ..., \lambda_{r_1}$ are complex numbers such that

$$\sum_{j=1}^{r_1} \lambda_j(\omega_j)_0 = 0.$$

Then

$$\omega := (\gamma^* - \alpha_1 Id)^{N_1 - 1} \left(\sum_{j=1}^{r_1} \lambda_j \omega_j\right)$$

is an element of ker $(\gamma^* - \alpha_1 Id)$. On the other hand, we have

$$\omega = \sum_{j=1}^{r_1} \lambda_j (\gamma^* - \alpha_1 Id)^{N_1 - 1} \omega_j$$

= $\sum_{j=1}^{r_1} \lambda_j \sum_{k=0}^{N_1 - 1} {N_1 - 1 \choose k} (-\alpha_1)^{N_1 - 1 - k} (\gamma^*)^k \omega_j.$

Then

$$\begin{aligned}
\omega_0 &= \sum_{j=1}^{r_1} \lambda_j \sum_{k=0}^{N_1-1} \binom{N_1-1}{k} (-\alpha_1)^{N_1-1-k} (\omega_j)_0 \circ (d_0 \gamma^k) \\
&= \left(\sum_{j=1}^{r_1} \lambda_j (\omega_j)_0 \right) \circ \left(\sum_{k=0}^{N_1-1} \binom{N_1-1}{k} (-\alpha_1)^{N_1-1-k} (d_0 \gamma^k) \right) \\
&= 0.
\end{aligned}$$

We infer that ω is identically zero. But the operator $(\gamma^* - \alpha_1 Id)^{N_1-1}$ is identically zero on ker $((\gamma^* - \alpha_1 Id)^{N_1-1})$ and one-to-one onto its direct image on Vect $(\{\omega_j | \omega_j \notin \text{ker}((\gamma^* - \alpha_1 Id)^{N_1-1})\})$. Therefore, for every j such that $\omega_j \notin \text{ker}((\gamma^* - \alpha_1 Id)^{N_1-1})$, we get $\lambda_j = 0$. Doing the same argument and using smaller and smaller powers of $\gamma^* - \alpha_1 Id$, we get that $\lambda_j = 0$ for every j.

This proves that the family $((\omega_1)_0, ..., (\omega_{r_1})_0)$ is linearly independent. By a continuity argument, there exists an open neighborhood U of 0 in D such that for every $z \in U$, $((\omega_1)_z, ..., (\omega_{r_1})_z)$ is linearly independent.

We are now proving that $r_1 = n_1$ the multiplicity of α_1 as an eigenvalue of $d_0\gamma$. First, notice that $((\omega_1)_0, ..., (\omega_{r_1})_0)$ is a linearly independent family of elements of the characteristic space associated to α_1 of the endomorphism

$$\begin{array}{cccc} (T_0\mathbb{C}^n)^* & \longrightarrow & (T_0\mathbb{C}^n)^* \\ \phi & \longmapsto & \phi \circ (d_0\gamma) \end{array}$$

and that this characteristic space has dimension n_1 . Therefore, $r_1 \leq n_1$.

Now, for $j \in \{1, ..., n_1\}$, since we have the decomposition

$$H_b^0(D,\Omega_D^1) = \ker((\gamma^* - \alpha_1 Id)^{N_1}) \oplus \overline{\operatorname{im}((\gamma^* - \alpha_1 Id)^{N_1})}$$

there exists a unique $\eta_j \in \ker((\gamma^* - \alpha_1 Id)^{N_1})$ such that $dz_j - \eta_j$ is an element of $\overline{\operatorname{im}((\gamma^* - \alpha_1 Id)^{N_1})}$. Looking at the Jacobian $J_0\gamma$, for every $\xi \in \operatorname{im}((\gamma^* - \alpha_1 Id)^{N_1})$, the coefficient of ξ_0 in front of d_0z_l is zero for every $l \in \{1, ..., n_1\}$. This property extends easily to the space $\overline{\operatorname{im}((\gamma^* - \alpha_1 Id)^{N_1})}$. Moreover, $\eta_j \in \ker((\gamma^* - \alpha_1 Id)^{N_1})$, so $(\eta_j)_0 = d_0z_j$. Therefore, $((\eta_1)_0, ..., (\eta_{n_1})_0)$ is linearly independent, so is $(\eta_1, ..., \eta_{n_1})$. Since $\ker((\gamma^* - \alpha_1 Id)^{N_1})$ is of dimension r_1 , we have $n_1 \leq r_1$, achieving the proof that $r_1 = n_1$.

For every $j \in \{1, ..., n_1\}$ and every $z \in U$, we define

$$(H_{1,n_1-j})_z = \bigcap_{l=1}^j \ker((\omega_l)_z).$$

Then,

$$\{0\} \subset H_{1,0} \subset \ldots \subset H_{1,n_1-1} \subset TU$$

defines a γ -invariant (non complete) flag on U and since γ is a contraction in 0 and Uis an open neighborhood of 0, this flag extends to \mathbb{C}^n . We define \mathcal{H}_1 to be equal to $H_{1,0}$. Notice that the eigenvalues of $d_0\gamma$ on $(\mathcal{H}_1)_0$ are exactly $\alpha_2, ..., \alpha_m$ with respective multiplicity $n_2, ..., n_m$, since it is obvious that $d_0\gamma^*$ has a unique eigenvalue α_1 of multiplicity n_1 on $(\mathcal{H}_1)_0^*$.

This concludes the initial case. We will then deal with the induction step.

Assume that $j \in \{1, ..., m-1\}$ is such that for every $k \in \{1, ..., j\}$, we have a flag of γ -invariant sub-bundles

$$\mathcal{H}_k \subset H_{k,1} \subset ... \subset H_{k,n_k-1} \subset \bigcap_{l \in I_k} \mathcal{H}_l$$

with

$$\dim \mathcal{H}_k = \dim \left(\bigcap_{l \in I_k} \mathcal{H}_l\right) - n_k$$

and

$$\dim\left(\bigcap_{l\in I_k}\mathcal{H}_l\right) = n - \sum_{l\in I_k} n_l$$

as well as

$$\dim\left(\bigcap_{l\in I_{j+1}}\mathcal{H}_l\right)=n-\sum_{l\in I_{j+1}}n_l.$$

Finally, assume that the vector sub-bundles $\mathcal{H}_1, ..., \mathcal{H}_j$ are such that for every $k \in \{1, ..., j+1\}$, the eigenvalues of $d_0\gamma$ on

$$\left(igcap_{l\in I_k}\mathcal{H}_l
ight)_0$$

are the α_l such that $l \notin I_k$ and each α_l as multiplicity n_l .

The manifold

$$B := \bigcap_{k \in I_{j+1}} \mathcal{H}_k$$

is a holomorphic vector bundle of rank

$$d := n - \sum_{l \in I_{j+1}} n_l$$

over \mathbb{C}^n . By the Oka-Grauert principle ([12], p.144), it is holomorphically trivial. Let then $(s_1, ..., s_d)$ be a family of holomorphic sections trivializing B and let $(s_1^*, ..., s_d^*)$ be the dual family, trivializing the dual bundle B^* . Up to the action of $GL(d, \mathbb{C})$, we can assume that the following holds

where

$$\{k_1 < \dots < k_q\} = \{0, \dots, m\} \setminus I_{j+1}$$

and the block corresponding to the eigenvalue α_{k_l} is of size n_{k_l} , for $l \in \{1, ..., q\}$, as in $J_0\gamma$. Since j + 1 is an element of $\{0, ..., m\} \setminus I_{j+1}$, there exists a unique $l \in \{1, ..., q\}$ such that $k_l = j + 1$.

The space $H_b^0(D, B^*)$ is a Banach space, by the Montel Theorem, and the restriction $\tilde{\gamma}$ of γ^* to this space is itself a compact operator.

Looking at the previous matrix, we see that

 $(s_{n_{k_1}+...+n_{k_l}})^*$

is not in the image of $\tilde{\gamma} - \alpha_{j+1}Id$. Notice that $(s_{n_{k_1}+\ldots+n_{k_l}})^*$ is bounded since it is globally defined on \mathbb{C}^n and D is bounded, hence relatively compact. Using Riesz-Schauder Theorem, α_{j+1} is an eigenvalue of $\tilde{\gamma}$ and there exists a positive integer N_{j+1} , that we can assume to be greater or equal than n_{j+1} , such that the space ker $((\tilde{\gamma} - \alpha_{j+1}Id)^{N_{j+1}})$ is finite dimensional and such that the following decomposition holds

$$H_b^0(D, B^*) = \ker((\tilde{\gamma} - \alpha_{j+1}Id)^{N_{j+1}}) \oplus \overline{\operatorname{im}((\tilde{\gamma} - \alpha_{j+1}Id)^{N_{j+1}})}.$$


We take an arbitrary nonzero element φ of ker $(\tilde{\gamma} - \alpha_{j+1}Id)$ and we will show it does not vanish at 0. We write

$$\varphi = \sum_{\mu=1}^d \sum_{\nu \ge 0} P_{\mu,\nu} s_\mu^*$$

where $P_{\mu,\nu}$ is a homogeneous polynomial of degree ν . For every $\mu \in \{1, ..., d\}$, we write

$$\tilde{\gamma}(s^*_{\mu}) = \sum_{\tau=1}^d f_{\mu,\tau} s^*_{\tau}$$

where $f_{\mu,\tau}$ are holomorphic functions on D. Notice that the matrix $(f_{\mu,\tau}(0))_{\mu,\tau}$ is exactly the matrix



Since φ is nonzero, let ν_0 be the smallest non negative integer such that one of the P_{μ,ν_0} is nonzero, and let μ_0 be the largest integer in $\{1, ..., d\}$ such that P_{μ_0,ν_0} is nonzero. Then the equation

$$\tilde{\gamma}(\varphi) = \alpha_{j+1}\varphi$$

is equivalent to

$$\sum_{\mu=1}^d \sum_{\nu \ge \nu_0} (P_{\mu,\nu} \circ \gamma) \tilde{\gamma}(s_\mu^*) = \alpha_{j+1} \sum_{\mu=1}^d \sum_{\nu \ge \nu_0} P_{\mu,\nu} s_\mu^*$$

which is itself equivalent to

$$\sum_{\mu=1}^d \left(\sum_{\tau=1}^d \left(\sum_{\nu \ge \nu_0} P_{\tau,\nu} \circ \gamma \right) f_{\tau,\mu} \right) s_{\mu}^* = \sum_{\mu=1}^d \left(\alpha_{j+1} \sum_{\nu \ge \nu_0} P_{\mu,\nu} \right) s_{\mu}^*.$$

This last equation is equivalent to the identification of two holomorphic maps, for every $\mu \in \{1, ..., d\}$, namely

$$\sum_{\tau=1}^{d} \left(\sum_{\nu \ge \nu_0} P_{\tau,\nu} \circ \gamma \right) f_{\tau,\mu} = \alpha_{j+1} \sum_{\nu \ge \nu_0} P_{\mu,\nu}.$$

This implies, choosing $\mu = \mu_0$, and looking at the Taylor polynomial of order ν_0 , the following equation

$$\sum_{\tau=1}^{\mu_0} (P_{\tau,\nu_0} \circ u_0) f_{\tau,\mu_0}(0) = \alpha_{j+1} P_{\mu_0,\nu_0}.$$

But looking at the form of the matrix $(f_{\mu,\tau}(0))_{\mu,\tau}$, we obtain that

$$\alpha P_{\mu_0,\nu_0} \circ u_0 = \alpha_{j+1} P_{\mu_0,\nu_0}$$

where α is the eigenvalue of $(f_{\mu,\tau}(0))_{\mu,\tau}$ lying at row and column μ_0 , say α_r with $r \in \{0, ..., m\} \setminus I_{j+1}$, which exactly means that r = j + 1 or α_{j+1} does not depend on α_r . This proves that $\frac{\alpha_{j+1}}{\alpha_r}$ is an eigenvalue of the linear endomorphism

$$\begin{array}{ccc} \mathbb{C}^{\mathrm{Hom}}_{\nu_0}[Z_1,...,Z_n] &\longrightarrow & \mathbb{C}^{\mathrm{Hom}}_{\nu_0}[Z_1,...,Z_n] \\ P &\longmapsto & P \circ u_0. \end{array}$$

But the eigenvalues of this endomorphism are precisely all the possible products of ν_0 eigenvalues of $d_0\gamma$. Therefore, if ν_0 is positive, then there is a resonance

$$\alpha_{j+1} = \alpha_r \prod_{i=1}^m \alpha_i^{p_i}$$

where the length of $p = (p_1, ..., p_m)$ is ν_0 , therefore positive. But this kind of relation is impossible, since $r \in \{0, ..., m\} \setminus I_{j+1}$. Hence, $\nu_0 = 0$ (and r = j + 1), which proves that φ does not vanish at 0.

Since ker $((\tilde{\gamma} - \alpha_{j+1}Id)^{N_{j+1}})$ is finite dimensional, say of dimension r_{j+1} , and stable by $\tilde{\gamma}$, let $(\varphi_1, ..., \varphi_{r_{j+1}})$ be a basis of ker $((\tilde{\gamma} - \alpha_{j+1}Id)^{N_{j+1}})$ such that $\tilde{\gamma}$ is triangular superior in this basis. Then, as in the initial case, we can prove that $((\varphi_1)_0, ..., (\varphi_{r_{j+1}})_0)$ is linearly independent, and projecting the family

$$((s_{n_{k_1}+\ldots+n_{k_{l-1}}+1})^*, \ldots, (s_{n_{k_1}+\ldots+n_{k_l}})^*)$$

on ker $((\tilde{\gamma} - \alpha_{j+1}Id)^{N_{j+1}})$, as in the initial case, that $r_{j+1} = n_{j+1}$. We can then consider, for $k \in \{1, ..., n_{j+1}\}$

$$H_{j+1,n_{j+1}-k} = \bigcap_{l=1}^{\kappa} \ker(\varphi_l)$$

in a neighborhood of 0 and extend those γ -invariant vector bundles to \mathbb{C}^n , applying the contraction γ as much as we need. We denote by \mathcal{H}_{j+1} the vector bundle $H_{j+1,0}$. By construction, we have a flag

$$\mathcal{H}_{j+1} \subset H_{j+1,1} \subset \ldots \subset H_{j+1,n_{j+1}-1} \subset \bigcap_{k \in I_{j+1}} \mathcal{H}_k$$

with

$$\dim \mathcal{H}_{j+1} = \dim \bigcap_{k \in I_{j+1}} \mathcal{H}_k - n_{j+1}$$

To conclude the induction step, we need to show that, if j < m - 1, then

$$\dim \bigcap_{k \in I_{j+2}} \mathcal{H}_k = n - \sum_{k \in I_{j+2}} n_k.$$

We check first this dimension in $T_0\mathbb{C}^n$. The spectrum of $d_0\gamma$ on

$$\left(\bigcap_{k\in I_{j+2}}\mathcal{H}_k\right)_0$$

is exactly $\{\alpha_l, l \notin I_{j+2}\}$ and each α_l has multiplicity n_l . Therefore, at 0, the dimension is what we need. Let z be an element of \mathbb{C}^n and let d be the dimension of

$$\left(\bigcap_{k\in I_{j+2}}\mathcal{H}_k\right)_z.$$

Since the sequence $(\gamma^p(z))_{p\in\mathbb{N}}$ converges to 0, and since $T_0\mathbb{C}^n$ is trivial, the sequence of vector sub-spaces of \mathbb{C}^n

$$\left(d_{\gamma^{p-1}(z)}\gamma\left(\bigcap_{k\in I_{j+2}}\mathcal{H}_k\right)_z\right)_{p\in\mathbb{N}}$$

converges to

$$\left(\bigcap_{k\in I_{j+2}}\mathcal{H}_k\right)_0$$

so we have that $d = n - \sum_{k \in I_{j+2}} n_k$. Therefore,

$$\bigcap_{k\in I_{j+2}}\mathcal{H}_k$$

is a vector bundle of rank $n - \sum_{k \in I_{j+2}} n_k$. Hence we proved the induction step, and by induction, we proved the theorem.

This theorem implies that every Hopf manifold carries special geometries, namely a G-structure of order 1, with a group that is only determined by the eigenvalues of the differential map at 0 of the contraction. In the following section, we will define a group of polynomial transformations of \mathbb{C}^n . The group G^1_{α} can be identified as the group of jets of order 1 of elements of this group.

5. The group of sub-resonant polynomial transformations

From now on, we will use the following notation

$$J_0 \gamma = \begin{pmatrix} \beta_1 & & (0) \\ & \cdot & & \\ & & \cdot & \\ * & & & \beta_n \end{pmatrix}$$

so that

$$\beta_1 = \dots = \beta_{n_1} = \alpha_1$$

$$\vdots$$

$$\beta_{n_1 + \dots + n_{m-1} + 1} = \dots = \beta_n = \alpha_m$$

We define a positive integer r depending on the set of eigenvalues $\{\beta_1, ..., \beta_n\}$, namely the maximal length of the resonances, as follows

$$r = \max\{|p|, p \in \mathbb{N}^n \mid \exists j \in \{1, ..., n\} \mid \beta_j = \beta^p\}.$$

Notice that the fact that r is finite was proven by Poincaré ([2], Theorem 1 p.188).

We will define a group of polynomial transformations of \mathbb{C}^n of order r. This group depends on the set of resonances of $\beta = (\beta_1, ..., \beta_n)$. For $j \in \{1, ..., n\}$ we consider

$$r_{j} = \max\{|p|, p \in \mathbb{N}^{n} \mid \beta_{j} = \beta^{p}\}$$
$$\mathcal{P}_{j} = \{p \in \mathbb{N}^{n} \mid \beta_{j} = \beta^{p} \text{ and } \forall k \ge j, p \neq e_{k}\}$$
$$\mathcal{Q}_{j} = \{q \in \mathbb{N}^{n} \setminus \{0\} \mid \exists p \in \mathcal{P}_{j} \mid \forall l \in \{1, ..., n\}, q_{l} \le p_{l}\}$$

where e_k is the k-th element of the standard basis of \mathbb{Z}^n . The number r_j is the maximal length of resonance of β_j (and therefore we have $r = \max r_j$), the set \mathcal{P}_j is the set of resonances of β_j , where we do not take into account $\beta_j = \beta_{j+l}$ with $l \ge 0$, even if it is the case, and \mathcal{Q}_j is the set of nonzero *n*-tuples of natural integers that are component-wise not greater than some resonance of β_j . We refer to Example 5.0.1 for clarifications on those technical definitions.

Let $P : \mathbb{C}^n \to \mathbb{C}^n$. We say that P is a β sub-resonant polynomial transformation of \mathbb{C}^n if, in the standard coordinates,

$$P(z) = (a_1z_1, a_2z_2 + P_2(z_1), \dots, a_nz_n + P_n(z_1, \dots, z_{n-1}))$$

where

$$\forall j \in \{1, ..., n\}, \qquad a_j \neq 0 \\ \forall j \in \{2, ..., n\}, \quad P_j(z_1, ..., z_{j-1}) = \sum_{q \in \mathcal{Q}_j} c_{j,q} z^q.$$

Notice that such a polynomial map is at most of degree r.

Example 5.0.1. Assume that γ is such that

$$J_0 \gamma = \begin{pmatrix} \alpha & 0 & 0 \\ 0 & \alpha^3 & 0 \\ 0 & 0 & \alpha^5 \end{pmatrix} = \begin{pmatrix} \beta_1 & 0 & 0 \\ 0 & \beta_2 & 0 \\ 0 & 0 & \beta_3 \end{pmatrix}.$$

The non trivial resonances are

$$\begin{array}{rcl} \beta_2 & = & \beta_1^3 & = & \beta^{(3,0,0)} \\ \beta_3 & = & \beta_1^5 & = & \beta^{(5,0,0)} \\ \beta_3 & = & \beta_1^2 \beta_2 & = & \beta^{(2,1,0)} \end{array}$$

Therefore, we have

$$r = 5$$
 $r_1 = 1$ $r_2 = 3$ $r_3 = 5$

$$\begin{array}{c} \mathcal{P}_1 = \emptyset & \mathcal{Q}_1 = \emptyset \\ \mathcal{P}_2 = \{(3,0,0)\} & \mathcal{Q}_2 = \{(1,0,0), (2,0,0), (3,0,0)\} \\ \mathcal{P}_3 = \{(5,0,0), (2,1,0)\} & \mathcal{Q}_3 = \{(i,0,0), 1 \le i \le 5\} \cup \{(0,1,0), (1,1,0), (2,1,0)\} \end{array}$$

The polynomial map P given by the expression

$$P(z) = (z_1, z_2 + 2z_1 - z_1^3, z_3 + z_1 + z_1z_2 - 2z_1^2z_2 + z_1^4 + z_1^5)$$

is sub-resonant. Indeed, the monomials $z_1^3e_2$, $z_1^2z_2e_3$, $z_1^5e_3$ are resonant, i.e. they correspond to elements of the sets \mathcal{P}_j , and the monomials z_1e_2 , z_1e_3 , $z_1z_2e_3$, $z_1^4e_3$ are sub-resonant i.e. they correspond to elements of the sets \mathcal{Q}_j .

We denote by G_{β}^{r} the set of all β sub-resonant polynomial transformations of \mathbb{C}^{n} . This is a set of polynomial maps of degree at most r. We will show that it is a subgroup of the group of holomorphic automorphisms of \mathbb{C}^{n} .

Proposition 5.0.1. The set G^r_{β} , endowed with the composition of maps from \mathbb{C}^n to itself, is a subgroup of the group $Aut(\mathbb{C}^n)$ of all the holomorphic automorphisms of \mathbb{C}^n .

Proof. The following proof is technical, but we will give some intuitive ideas first. It is already known that the set of β resonant polynomial maps is a group for the composition of maps ([20], Section 1 Proposition 2). The idea is that since the sub-resonant monomials are by definition not greater than some resonant monomial and

since the composition of resonant monomials remains resonant, the composition of subresonant monomials remains sub-resonant.

We will show this result by induction on the dimension n. If n = 2, then let P, \tilde{P} be elements of G_{β}^{r} . We use the same notations as in the definition above, namely

$$P(z) = (a_1 z_1, a_2 z_2 + P_2(z_1))$$

$$\tilde{P}(z) = (b_1 z_1, b_2 z_2 + \tilde{P}_2(z_1)).$$

A monomial appearing in \tilde{P}_2 is a z^q with $q \in Q_2$, i.e. a $z_1^{q_1}$ with $1 \leq q_1 \leq d$ where $\beta_2 = \beta_1^d$. In case we don't have this type of resonances, then Q_2 is empty and G_{β}^r is obviously a group, since it is the group of linear diagonal transformations of \mathbb{C}^2 . In the resonant case, we have that

$$z_1^{q_1} \circ P(w) = a_1^{q_1} w_1^{q_1}$$

This identity proves that $\tilde{P} \circ P$ is an element of G_{β}^r . Moreover, P is obviously one-to-one, and

$$P^{-1}(w) = \left(\frac{w_1}{a_1}, \frac{w_2 - P_2(\frac{w_1}{a_1})}{a_2}\right)$$

which is also an element of G_{β}^r . Therefore, if n = 2, then G_{β}^r is a subgroup of $Aut(\mathbb{C}^2)$.

Let now $n \in \mathbb{N}$ be such that for every integer m satisfying $2 \leq m \leq n$ and every set of eigenvalues $(\eta_1, ..., \eta_m)$ satisfying

$$1 \ge |\eta_1| \ge \dots \ge |\eta_m| \ge 0$$

the set G_{η} is a group. Let $(\beta_1, ..., \beta_{n+1}) \in \mathbb{C}^{n+1}$ satisfying

$$1 \ge |\beta_1| \ge \dots \ge |\beta_{n+1}| \ge 0.$$

Let P, \tilde{P} be β sub-resonant. Looking at the composition $\tilde{P} \circ P$, the *n* first coordinates form a composition of two $(\beta_1, ..., \beta_n)$ sub-resonant polynomial maps. Therefore, using the induction hypothesis, we only focus on the (n + 1)-th coordinate. The polynomial $\tilde{P}_{n+1}(z_1, ..., z_n)$ contains monomials of the form z^q with $q \in Q_{n+1}$. By definition of Q_{n+1} , there exists an element p of \mathcal{P}_{n+1} such that $q \leq p$ (coordinate by coordinate). We use the notation

$$P(z) = (a_1 z_1 + P_1(z), ..., a_{n+1} z_{n+1} + P_{n+1}(z))$$

where $a_j \neq 0$ and

$$P_j(z) = \sum_{s \in \mathcal{Q}_j} c_{j,s} z^s.$$

Then we compute

$$z^{q} \circ P(w) = \prod_{\substack{j=1 \ j=1}}^{n+1} (a_{j}w_{j} + P_{j}(w))^{q_{j}}$$
$$= \prod_{j=1}^{n+1} (a_{j}w_{j} + \sum_{s \in \mathcal{Q}_{j}} c_{j,s}w^{s})^{q_{j}}$$

We denote by N_j the cardinality of \mathcal{Q}_j and $\mathcal{Q}_j = \{s_1^{(j)}, ..., s_{N_j}^{(j)}\}$. Then we have

$$z^{q} \circ P(w) = \prod_{j=1}^{n+1} \left(\sum_{i_{1}+\ldots+i_{N_{j}+1}=q_{j}} \frac{q_{j}!}{i_{1}!\ldots i_{N_{j}+1}!} \left(\prod_{k=1}^{N_{j}} \left(\left(c_{j,s_{k}^{(j)}} w^{s_{k}^{(j)}} \right)^{i_{k}} \right) \right) (a_{j}w_{j})^{i_{N_{j}+1}} \right) \\ = \prod_{j=1}^{n+1} \left(\sum_{i_{1}+\ldots+i_{N_{j}+1}=q_{j}} \frac{q_{j}!}{i_{1}!\ldots i_{N_{j}+1}!} \left(\prod_{k=1}^{N_{j}} c_{j,s_{k}^{(j)}}^{i_{k}} \right) a_{j}^{i_{N_{j}+1}} w^{\left(\sum_{k=1}^{N_{j}} i_{k}s_{k}^{(j)} \right) + i_{N_{j}+1}e_{j}} \right)$$

where, as usual, $(e_1, ..., e_{n+1})$ is the standard basis of \mathbb{Z}^{n+1} , so that the monomials appearing are w to the power of

$$\sum_{j=1}^{n+1} \left(\sum_{k=1}^{N_j} i_k^{(j)} s_k^{(j)} + i_{N_j+1}^{(j)} e_j \right)$$

where, for every $j \in \{1, ..., n+1\}$, we have

$$\sum_{k=1}^{N_j+1} i_k^{(j)} = q_j$$

We need to explain that for every

$$(i_1^{(1)}, ..., i_{N_1+1}^{(1)}), ..., (i_1^{(n+1)}, ..., i_{N_{n+1}+1}^{(n+1)})$$

satisfying

$$\sum_{k=1}^{N_j+1} i_k^{(j)} = q_j$$

the (n+1) tuple

$$\sum_{j=1}^{n+1} \left(\sum_{k=1}^{N_j} i_k^{(j)} s_k^{(j)} + i_{N_j+1}^{(j)} e_j \right)$$

is bounded above by an element of \mathcal{P}_{n+1} . First, we notice that $q_{n+1} = 0$, implying that, for every $k \in \{1, ..., N_{n+1} + 1\}$, we have $i_k^{(n+1)} = 0$. We can rewrite the (n+1) tuple as

$$\sum_{j=1}^{n} \left(\sum_{k=1}^{N_j} i_k^{(j)} s_k^{(j)} + i_{N_j+1}^{(j)} e_j \right).$$

By definition of the sets \mathcal{Q}_j , for every $j \in \{1, ..., n\}$, for every $k \in \{1, ..., N_j\}$, there exists an element $\underline{p}_k^{(j)}$ of \mathcal{P}_j such that $s_k^{(j)} \leq \underline{p}_k^{(j)}$. We recall that p is an element of \mathcal{P}_{n+1} such that $q \leq p$. We consider

$$\tilde{p} = p - \sum_{j=1}^{n} \left(\sum_{k=1}^{N_j} i_k^{(j)} \right) e_j + \sum_{j=1}^{n} \sum_{k=1}^{N_j} i_k^{(j)} \underline{p}_k^{(j)}$$

and we check that $\beta^{\tilde{p}} = \beta_{n+1}$, meaning that \tilde{p} is an element of \mathcal{P}_{n+1} . Since $p \ge q$ and $\underline{p}_k^{(j)} \ge s_k^{(j)}$, we have

$$\tilde{p} \ge q - \sum_{j=1}^{n} \left(\sum_{k=1}^{N_j} i_k^{(j)} \right) e_j + \sum_{j=1}^{n} \sum_{k=1}^{N_j} i_k^{(j)} s_k^{(j)},$$

but we have

$$q - \sum_{j=1}^{n} \left(\sum_{k=1}^{N_j} i_k^{(j)} \right) e_j = \sum_{j=1}^{n} i_{N_j+1}^{(j)} e_j.$$

Therefore, we deduce that

$$\tilde{p} \ge \sum_{j=1}^{n} \left(\sum_{k=1}^{N_j} i_k^{(j)} s_k^{(j)} + i_{N_j+1}^{(j)} e_j \right)$$

which is what we wanted. This shows that $\tilde{P} \circ P$ is an element of G_{β} . To conclude, we write

$$P^{-1}(w) = \left(\frac{w_1}{a_1}, \frac{w_2 - P_2(\frac{w_1}{a_1})}{a_2}, \dots, \frac{w_{n+1} - P_{n+1}(\dots)}{a_{n+1}}\right).$$

The induction hypothesis allows us to look only at the (n + 1)-th coordinate, that we treat exactly the same way as for the composition.

This concludes the induction and therefore, the proof that G_{β}^{r} is a subgroup of $Aut(\mathbb{C}^{n})$.

To conclude this section, we notice that the group G^r_β can be seen as a subgroup of the jets group $\mathcal{D}^r(\mathbb{C}^n)$, and that it is a closed subgroup. Therefore, it carries a natural structure of complex Lie subgroup of $\mathcal{D}^r(\mathbb{C}^n)$.

Remark 5.0.1. In relation to Remark 2.2.1, we notice also that G^r_β is an algebraic subgroup of $\mathcal{D}^r(\mathbb{C}^n)$.

6. Geometric structures of higher order on Hopf manifolds

For every $k \in \mathbb{N}^*$, we consider the group

$$G^k_\beta := \{j^k_0 P \mid P \in G^r_\beta\}$$

endowed with the order k truncated composition. Notice that since G_{β}^{r} is a group of polynomial maps of degree at most r, for every $k \geq r$, we have $G_{\beta}^{k} = G_{\beta}^{r}$. Notice also

that G^1_{β} coincides with the G^1_{α} of Section 4.. We will show that we can extend a given G^1_{β} -structure of order 1 step by step to get a uniquely determined G^k_{β} -structure of order k, for any positive integer k in the case of dimension $n \geq 3$, and for $k \in \{r+1, r+2\}$ for surfaces.

The case of Hopf surfaces has to be treated separately. The reason for this special treatment is a cohomological property that is different between $\mathbb{C}^2 \setminus \{0\}$ and $\mathbb{C}^n \setminus \{0\}$, for $n \geq 3$. Indeed, if \mathcal{O}_X stands for the the sheaf of holomorphic functions on a complex manifold X, then we have the following properties (see for example [9] or [20], Section 2.1)

$$\begin{cases} H^1(\mathbb{C}^n \setminus \{0\}, \mathcal{O}_{\mathbb{C}^n \setminus \{0\}}) = 0 \text{ for } n \ge 3\\ H^1(\mathbb{C}^2 \setminus \{0\}, \mathcal{O}_{\mathbb{C}^2 \setminus \{0\}}) \cong \left\{ \sum_{p \in (-\mathbb{N}^*)^2} a_p z^p \text{ convergent on } (\mathbb{C}^*)^2 \right\} \end{cases}$$

Because of this phenomenon, Mall's theorem fails in dimension two, as noticed in [21], Remark 4.4.. In Section 6.2., we will deal with the case of Hopf surfaces.

6.1. High dimension

This section is devoted to the proof of the following result.

Theorem 4. Assume that $n \geq 3$. Let k be a positive integer and let $s^{(k)} : M \to \mathcal{R}^k(M)/G^k_\beta$ be a reduction of the structure group of $\mathcal{R}^k(M)$ to G^k_β . Then there exists a unique reduction $s^{(k+1)} : M \to \mathcal{R}^{k+1}(M)/G^{k+1}_\beta$ of the structure group of $\mathcal{R}^{k+1}(M)$ to G^{k+1}_β such that

$$\overline{\pi}_{k+1,k} \circ s^{(k+1)} = s^{(k)}$$

where

$$\overline{\pi}_{k+1,k}: \mathcal{R}^{k+1}(M)/G_{\beta}^{k+1} \to \mathcal{R}^{k}(M)/G_{\beta}^{k}$$

is the natural projection.

Proof. Let $M = \bigcup_i U_i$ be a covering of M by open subsets and let (φ_i) local holomorphic charts on U_i such that

$$s^{(k)}|_{U_i} = [s_i^{(k)}]$$

where

$$\begin{array}{cccc} s_i^{(k)} & \colon & M & \longrightarrow & \mathcal{R}^k(M) \\ & & m & \longmapsto & j_0^k(\varphi_i^{-1}\theta_{\varphi_i(m)}f_{i,m}) \end{array}$$

where θ_z is the translation in \mathbb{C}^n of vector $z \in \mathbb{C}^n$ and $j_0^k(f_{i,m})$ is the jet of order k at 0 of a germ of biholomorphism $(\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$. We consider the following diagram

where the upper left object is the pullback of the diagram. We will denote by A the space

$$(s^{(k)})^* \left(\mathcal{R}^{k+1}(M) / G_{\beta}^{k+1} \right).$$

We claim that the map $p:A\to M$ is an affine bundle. The fiber of this bundle is isomorphic to the quotient

$$V^{k+1}/V_{\beta}^{k+1}$$

where V^{k+1} is the space of homogeneous polynomial maps of degree k+1 from \mathbb{C}^n to \mathbb{C}^n and V_{β}^{k+1} is the subspace generated by the monomials of the form

 $z^q e_j$

where $q \in Q_j$ is of length k + 1, i.e. q is a sub-resonance of β_j of length k + 1. To see that, we consider

$$\psi_i : A|_{U_i} \longrightarrow U_i \times V^{k+1} / V_{\beta}^{k+1} (\varphi_i^{-1} \theta_{\varphi_i(m)}(f_{i,m})_k (id+P))]) \longmapsto (m, [P])$$

where $(f_{i,m})_k$ is the truncating at order k of $f_{i,m}$. We compute

$$\psi_{j}\psi_{i}^{-1}(m,[P]) = \psi_{j}(m,[j_{0}^{k+1}(\varphi_{i}^{-1}\theta_{\varphi_{i}(m)}(f_{i,m})_{k}(id+P))])$$

$$= \psi_{j}(m,[j_{0}^{k+1}(\varphi_{j}^{-1}\theta_{\varphi_{j}(m)}(f_{j,m})_{k}(((f_{j,m})_{k})^{-1}\theta_{-\varphi_{j}(m)}\varphi_{j}$$

$$\varphi_{i}^{-1}\theta_{\varphi_{i}(m)}(f_{i,m})_{k})(id+P))]).$$

But, by construction,

$$j_0^k(((f_{j,m})_k)^{-1}\theta_{-\varphi_j(m)}\varphi_j\varphi_i^{-1}\theta_{\varphi_i(m)}(f_{i,m})_k)$$

is an element of G^k_{β} . Therefore, we see that

$$\psi_j \psi_i^{-1}(m, [P]) = (m, [\mathcal{L}_{j,i,m}(P) + Q_{j,i,m}])$$

where

$$\mathcal{L}_{j,i,m}(P) = \phi_{j,i,m} \circ P \circ \phi_{j,i,m}^{-1}$$

with

$$\phi = ((f_{j,m})_1)^{-1} (\theta_{-\varphi_j(m)} \varphi_j \varphi_i^{-1} \theta_{\varphi_i(m)})_1 (f_{i,m})_1$$

and $Q_{j,i,m} \in V^{k+1}$ is independent of P. This proves that p is an affine bundle with typical fiber V^{k+1}/V_{β}^{k+1} .

To prove the theorem is equivalent to prove that the affine bundle p admits a unique global section. We will use a cohomological argument in order to prove that. Namely, if ξ, ζ are two local sections of p, say on an open set U, then the difference $\xi - \zeta$ is well defined as a section of the vector bundle, denoted by B, associated with p. Since p is locally trivial, we can choose local sections on trivialization open sets and consider the difference of those sections on the intersection of the open sets of the covering. Therefore the existence of a global section of p is equivalent to the vanishing of a cohomology class in the first sheaf cohomology group of the associated vector bundle B. Moreover, a global section of the affine bundle is unique if and only if there exists a unique global section of B, namely the zero section. To rephrase, the sub-resonant monomials are exactly the indeterminacy to extend the structure to the superior order. Extending up to these monomials, i.e. considering the quotient V^{k+1}/V_{β}^{k+1} , determines completely how to extend the structure.

Let us then study this holomorphic vector bundle B on M. Using the above computation, we have a description of the cocycle of B. Namely, it is given by $\phi_{i,j,m}$. Therefore, we can see that the principal bundle associated with B is the first order frame bundle on M, i.e. $\mathcal{R}^1(M)$. Indeed, they have the same cocycle, up to the action of the cochain $\{(f_{i,m})_1\}$. This shows that if $\pi : \mathbb{C}^n \setminus \{0\} \to M$ is the universal covering of M, then the pulled back vector bundle $\overline{B} := \pi^* B$ is trivial over $\mathbb{C}^n \setminus \{0\}$ and it extends to \mathbb{C}^n . Therefore, B is a Mall bundle over the Hopf manifold M. Since we are in the case $n \geq 3$, the Mall theorem implies that

$$h^0(M,B) = h^1(M,B)$$

where $h^i(M, B)$ is the dimension of the *i*-th sheaf cohomology space of the vector bundle B. Using that, it is sufficient to prove that $H^0(M, B)$ is trivial to finish the proof of the theorem.

The fiber over 0 of the bundle \overline{B} , namely \overline{B}_0 is isomorphic to the space

$$V^{k+1}/V_{\beta}^{k+1}$$

and in a well chosen isomorphism, the action of γ on \overline{B}_0 reads

$$\gamma P = u_0^{-1} \circ P \circ u_0$$

where, as above, u_0 is the linear part of γ and where P is a homogeneous polynomial map of order k + 1 from \mathbb{C}^n to itself containing only non sub-resonant monomials, because P is an element of V^{k+1}/V_{β}^{k+1} . We deduce that the eigenvalues of the action of γ on \overline{B}_0 are exactly of the form

$$\beta_j^{-1}\beta^p$$

where the length of p is k + 1 and p is not j sub-resonant, i.e. $p \notin Q_j$ where this last set is defined above.

We choose $(\eta_1, ..., \eta_d)$ a family of global trivializing sections of \overline{B} . Suppose that \overline{B} carries a nonzero global section η . Then we can write

$$\eta = \sum_{i=1}^d \sum_{\nu \ge 0} P_{i,\nu} \eta_i$$

where $P_{i,\nu}$ is a homogeneous polynomial of degree ν . Since η is nonzero, let ν_0 the minimal degree where not all the $P_{i,\nu}$ are zero. Suppose also that η is γ invariant, then looking at the equation

$$\gamma.\eta=\eta$$

and looking at the order ν_0 , we deduce that the linear endomorphism

$$\begin{array}{ccc} \mathbb{C}^{\operatorname{Hom}}_{\nu_0}[Z_1,...,Z_n] \otimes V^{k+1}/V_{\beta}^{k+1} & \longrightarrow & \mathbb{C}^{\operatorname{Hom}}_{\nu_0}[Z_1,...,Z_n] \otimes V^{k+1}/V_{\beta}^{k+1} \\ Q \otimes P & \longmapsto & (Q \circ u_0) \otimes (u_0^{-1} \circ P \circ u_0) \end{array}$$

carries 1 as an eigenvalue. But this implies that there exists a resonance

$$\beta_i = \beta^p \beta^{p'}$$

where the length of p is k + 1 and p is not j sub-resonant, and where the length of p' is ν_0 . But this is impossible, because it would imply that p is sub-resonant.

This proves that

$$H^0(M,B) = \{0\}$$

which, as we said above, is enough to conclude the proof of the theorem.

This theorem combined with the existence of the G^1_β -structure of order 1 proved in Section 4. allow to write the following corollary, proved by induction.

Corollary 6.1.1. Let M be a Hopf manifold of dimension at least 3 and $\beta = (\beta_1, ..., \beta_n)$ be the set of eigenvalues associated with this Hopf manifold. Then, for every integer $k \ge 1$, M carries G^k_β -structure of order k. Moreover, every G^k_β -structure of order k is completely determined by the induced G^1_β -structure of order 1.

6.2. Hopf surfaces

In this subsection, we focus on Hopf surfaces. What fails in the proof of Theorem 4, is the moment when we use Mall's Theorem giving the equality

$$h^0(M,B) = h^1(M,B).$$

Notice that the proof of the fact that $h^0(M, B) = 0$ is still valid. Therefore, if we can extend a structure of order k at the order k + 1, this extension is unique. But the problem is that, a priori, the space $H^1(M, B)$ is not trivial.

In their paper [28], Ornea and Verbitsky have the same problem. As they mention, Theorem 6.6 is still true. We will explain why.

6.2.1. Mall theorem for Hopf surfaces

<

In [21], Mall shows a theorem that allows to compute the cohomology of any holomorphic vector bundle over a Hopf manifold of dimension $n \geq 3$ with a trivial pullback on $\mathbb{C}^n \setminus \{0\}$. Let us state his Theorem.

Theorem 5. ([21], Theorem 1.1) Let B be a Mall bundle on M a Hopf manifold of dimension $n \ge 3$. Then we have

$$\begin{cases} h^0(M,B) &= h^1(M,B) \\ h^p(M,B) &= 0 & \text{for } 2 \le p \le n-2 \\ h^{n-1}(M,B) &= h^n(M,B) \end{cases}$$

We want to give a version of this theorem for Hopf surfaces. As mentioned by Mall, by using the argument in the proof of Theorem 5, we only see that the Euler characteristic vanishes. The same result can be achieved by adding a hypothesis for Hopf surfaces.

Let $M = W / \langle \gamma \rangle$ be a Hopf surface, where $W := \mathbb{C}^2 \setminus \{0\}$ and $\gamma : \mathbb{C}^2 \to \mathbb{C}^2$ is a biholomorphic contraction in 0. We denote by $\pi : W \to M$ the projection.

Let B be a Mall bundle of rank r on M. Since $\pi^*(B)$ is trivial, there exists a group representation

$$\begin{array}{cccc} \rho_B & : & <\gamma > & \longrightarrow & GL(r, \Gamma(W, \mathcal{O})) \\ & \gamma & \longmapsto & A \end{array}$$

such that the bundle B is isomorphic to the quotient E_A of the trivial bundle $W \times \mathbb{C}^r$ by the action

$$\begin{array}{ccc} <\gamma > \times (W \times \mathbb{C}^r) & \longrightarrow & (W \times \mathbb{C}^r) \\ (\gamma, (z, v)) & \longmapsto & (\gamma(z), A(z)v). \end{array}$$

Mall uses the generalized Douady sequence (see [8], p.5). For Hopf surfaces, this exact sequence reads

$$0 \longrightarrow H^{0}(M, E_{A}) \xrightarrow{\pi^{*}} H^{0}(W, \mathcal{O}^{r}) \xrightarrow{A-\gamma^{*}} H^{0}(W, \mathcal{O}^{r}) \longrightarrow H^{1}(M, E_{A})$$
$$\xrightarrow{\pi^{*}} H^{1}(W, \mathcal{O}^{r}) \xrightarrow{A-\gamma^{*}} H^{1}(W, \mathcal{O}^{r}) \longrightarrow H^{2}(M, E_{A}) \longrightarrow 0.$$

From this exact sequence and with the work of Mall we infer the following result.

Theorem 6. Let E_A be a Mall bundle on M.

If the map

$$A - \gamma^* : H^1(W, \mathcal{O}^r) \longrightarrow H^1(W, \mathcal{O}^r)$$

is one-to-one, then we have $h^0(M, E_A) = h^1(M, E_A)$. If this map is onto, then we have $H^2(M, E_A) = 0$.

Proof. The second assertion is straightforward. The first assertion comes from the Corollary 4.2 in [21] ensuring that

$$A - \gamma^* : H^0(W, \mathcal{O}^r) \longrightarrow H^0(W, \mathcal{O}^r)$$

is a Fredholm operator of index 0, even for surfaces.

6.2.2. Ornea and Verbitsky theorem for Hopf surfaces

In [28], Ornea and Verbitsky proved their Theorem 6.6 for Hopf manifolds of dimension at least 3 and mentioned that the result was true for surfaces. We give here the proof for Hopf surfaces because we judge it interesting in order to understand what will come next in the resonant case.

Let γ be a non-resonant contraction of \mathbb{C}^2 at 0. In this section exceptionally, nonresonant means that the maximal length of resonance is 1. To be more specific, the linear map $d_0\gamma$ can have one eigenvalue of multiplicity 2. Here, this is the definition of "non-resonant" that Ornea and Verbitsky use in [28]. Let M be the corresponding Hopf manifold.

Theorem 7. ([28], p.26, Theorem 6.6) The Hopf surface M carries a unique flat torsion-free holomorphic connection. Moreover, there exists a biholomorphism $\varphi : \mathbb{C}^2 \to \mathbb{C}^2$ such that $\varphi \gamma \varphi^{-1}$ is linear.

Proof. The proof is the same as the one of Ornea and Verbitsky, except that we have to show that $h^0(M, TM^* \otimes TM^* \otimes TM) = h^1(M, TM^* \otimes TM^* \otimes TM)$. To do that, according to Theorem 6, we need to show that the map

$$H^{1}(\mathbb{C}^{2} \setminus \{0\}, (\mathcal{O}^{2})^{*} \otimes (\mathcal{O}^{2})^{*} \otimes \mathcal{O}^{2}) \longrightarrow H^{1}(\mathbb{C}^{2} \setminus \{0\}, (\mathcal{O}^{2})^{*} \otimes (\mathcal{O}^{2})^{*} \otimes \mathcal{O}^{2})$$

taking a convergent Laurent series on $(\mathbb{C}^*)^2$

$$s(z) = \sum_{j,k=1}^{2} \sum_{l=1}^{2} \sum_{p \in (-\mathbb{N}^*)^2} a_{j,l,m,p} z^p dz_j \otimes dz_k \otimes \frac{\partial}{\partial z_l}$$

to

$$\sum_{j,k=1}^{2} \sum_{l=1}^{2} \sum_{p \in (-\mathbb{N}^*)^2} a_{j,l,m,p} z^p dz_j \circ (d_z \gamma)^{-1} \otimes dz_k \circ (d_z \gamma)^{-1} \otimes d_z \gamma_* \frac{\partial}{\partial z_l} - s(\gamma(z))$$

is one-to-one.

Let s be such a convergent power series satisfying that its image is 0. Then assuming that s is nonzero, and looking at the lowest nonzero term in the Taylor expansion, we get a relation of the form

$$\alpha_1^j \alpha_2^{2-j} = \alpha_l \alpha_1^{-p_1} \alpha_2^{-p}$$

where α_1, α_2 are the eigenvalues of $d_0\gamma$, j is an element in $\{0, 1, 2\}$, l is an element in $\{1, 2\}$ and $p = (p_1, p_2)$ is an element of $(-\mathbb{N}^*)^2$. This relation automatically gives rise to a resonance of length greater than 1, which contradicts our assumptions. Therefore, the following equality holds

$$h^{0}(M, TM^{*} \otimes TM^{*} \otimes TM) = h^{1}(M, TM^{*} \otimes TM^{*} \otimes TM).$$

The rest of the proof is exactly the same as in [28].

6.2.3. G-structures on resonant Hopf surfaces

We could try to prove Theorem 4 the same way for Hopf surfaces. Nevertheless, once we reach the computation of

$$H^1(M,B)$$

where B has typical fiber isomorphic to V^{k+1}/V_{β}^{k+1} , the computation of the map

$$H^1(\mathbb{C}^2 \setminus \{0\}, \pi^*B) \longrightarrow H^1(\mathbb{C}^2 \setminus \{0\}, \pi^*B)$$

assuming that it is not one-to-one produces relations of the form

$$\alpha_1^j \alpha_2^{k+1-j} = \alpha_l \alpha_1^{-p_1} \alpha_2^{-p_2}$$

and, as soon as $k \geq 2$, the existence of this type of relation is not a contradiction. Therefore, the space $H^1(M, B)$ is not trivial, which means that the existence of an extension of our geometric structure is a class in $H^1(M, B)$ which is, a priori, non trivial.

Let $\gamma : \mathbb{C}^2 \to \mathbb{C}^2$ be a holomorphic contraction at 0, and let M be the corresponding Hopf manifold. Let α_1, α_2 be the eigenvalues of $d_0\gamma$, verifying

$$1 > |\alpha_1| \ge |\alpha_2| > 0.$$

According to Theorem 3, there exists a $\langle \gamma \rangle$ -invariant G^1_{α} -structure of order 1 on \mathbb{C}^2 . A consequence of the existence of this structure is that there exists a holomorphic distribution of lines in $T\mathbb{C}^2$ invariant under the action of $\langle \gamma \rangle$.

Lemma 6.2.1. The $\langle \gamma \rangle$ -invariant distribution is given by the kernel of a global holomorphic nowhere vanishing 1-form ω on \mathbb{C}^2 which verifies the following properties:

1. the form ω is closed,

2. the following equation holds: $\gamma^* \omega = \alpha_1 \omega$.

Proof. To prove this lemma, we need to go back to the proof of Theorem 3. According to that proof, there exist a bounded open neighborhood D of $0 \in \mathbb{C}^2$ verifying that $\gamma(D)$ is relatively compact in D, and a non-vanishing holomorphic one form ω_0 on D satisfying the equation $\gamma^*\omega_0 = \alpha_1\omega_0$. The distribution is given locally by ker(ω_0), and it extends to \mathbb{C}^2 because γ is a contraction.

Now, using again the fact that γ is a contraction, we have the following equality:

$$\mathbb{C}^2 = \bigcup_{n \in \mathbb{N}} \gamma^{-n}(D).$$

For every $n \in \mathbb{N}$, we define a holomorphic 1-form on $\gamma^{-n}(D)$ by the following formula:

$$(\omega_n)_z = \frac{1}{\alpha_1^n} (\omega_0)_{\gamma^n(z)} \circ d_z \gamma^n.$$

Then the family of nowhere vanishing holomorphic one forms $(\omega_n)_{n\in\mathbb{N}}$ verifies that

$$\begin{cases} \omega_{n+1}|_{\gamma^{-n}(D)} = \omega_n \\ \gamma^* \omega_n = \alpha_1 \omega_{n+1} \end{cases}$$

Therefore, all those forms patch together to produce a global nowhere vanishing holomorphic one form ω on \mathbb{C}^2 verifying $\gamma^* \omega = \alpha_1 \omega$.

To prove that ω is closed, we consider its exterior derivative $d\omega$. This is a holomorphic two form on \mathbb{C}^2 verifying $\gamma^* d\omega = \alpha_1 d\omega$. We assume that $d\omega$ is nonzero and we consider its power series expansion. Then, the equation $\gamma^* d\omega = \alpha_1 d\omega$ implies, looking at the lowest nonzero order of the Taylor expansion, a relation of the type

$$\alpha_1 = \alpha_j \alpha_k \alpha^p$$

where j, k are elements of $\{1, 2\}$ and p is an element of \mathbb{N}^2 . But the existence of such a relation is a contradiction because of the property

$$1 > |\alpha_1| \ge |\alpha_2| > 0.$$

Therefore, the form $d\omega$ is the zero form which implies that ω is closed.

Now, there are essentially three cases in our situation.

The first one is the non-resonant case (see Definition 3.0.1). In this situation, the proofs of Theorem 3 and Lemma 6.2.1 show that there exists another global holomorphic closed one form η on \mathbb{C}^2 satisfying $\gamma^* \eta = \alpha_2 \eta$. Moreover, ω and η are pointwise linearly independent. Since \mathbb{C}^2 is simply connected, we consider the holomorphic functions w_1

and w_2 on \mathbb{C}^2 , vanishing at 0, and such that $\omega = dw_1$ and $\eta = dw_2$. Then $\varphi = (w_1, w_2)$ is a local biholomorphism, and it verifies that

$$\varphi\gamma\varphi^{-1} = \begin{pmatrix} lpha_1 & 0 \\ 0 & lpha_2 \end{pmatrix}.$$

In other words, ω and η define a flat torsion-free affine connection on \mathbb{C}^2 invariant by $\langle \gamma \rangle$, and φ is the developing map of this affine connection. This gives a new proof of Theorem 6.6 in [28], p.27, in our non-resonant case.

The second one is the case where $\alpha_1 = \alpha_2$. In this case, according to the proof of Theorem 3, there exists a nowhere vanishing holomorphic one form η_0 on D satisfying $\gamma^*\eta_0 = \alpha_1\eta_0 + \varepsilon\omega_0$, where ε is 0 or 1, and such that ω_0 and η_0 are pointwise linearly independent.

Lemma 6.2.2. The form η_0 is closed.

Proof. Since we have the equality $\gamma^*\eta_0 = \alpha_1\eta_0 + \varepsilon\omega_0$, and since ω_0 is closed by Lemma 6.2.1 we infer that the exterior differential of η_0 satisfies the equation $\gamma^* d\eta_0 = \alpha_1 d\eta_0$. But then, by the same Taylor expansion argument as the one in the proof of Lemma 6.2.1, the 2-form $d\eta_0$ is identically 0. Therefore, η_0 is closed.

As in the non resonant case, we integrate ω and η_0 in a neighborhood of $0 \in \mathbb{C}^2$ and we produce local coordinates φ such that

$$\varphi\gamma\varphi^{-1} = \begin{pmatrix} \alpha_1 & 0\\ \varepsilon & \alpha_1 \end{pmatrix}.$$

This completes the new proof of Theorem 6.6 in [28], p.27 in the non resonant case.

Let us now deal with the third case, which is the remaining resonant case, namely, where there exists an integer $r \geq 2$ such that $\alpha_2 = \alpha_1^r$. In this case, we will denote by λ the eigenvalue α_1 , so that $\alpha_2 = \lambda^r$.

Lemma 6.2.3. The distribution ker(ω) is also given by a non-vanishing global holomorphic vector field X on \mathbb{C}^2 satisfying the equation $\gamma_* X = \lambda^r X$.

Proof. According to the proof of Theorem 3, there exists a section η_0 of the bundle $\ker(\omega)^*$ over D satisfying the equation $(\eta_0)_{\gamma(z)} \circ d_z \gamma|_{\ker(\omega)_z} = \lambda^r(\eta_0)_z$. Using the same argument as in the proof of the fact that ω_0 extends to \mathbb{C}^2 (see Lemma 6.2.1), η_0 extends to a global non-vanishing section of the bundle $\ker(\omega)$ verifying $\eta_{\gamma(z)} \circ d_z \gamma|_{\ker(\omega)_z} = \lambda^r \eta_z$. Now, since the bundle $\ker(\omega)$ is a line bundle (here we take advantage of the dimension 2), η produces a trivialization of the bundle $\ker(\omega)$. Namely, define X as the section of $\ker(\omega)$ satisfying $\eta(X) = 1$. Then, X can be seen as a non-vanishing holomorphic vector field on \mathbb{C}^2 generating the distribution of lines $\ker(\omega)$. We check that the equation $\eta_{\gamma(z)} \circ d_z \gamma|_{\ker(\omega)_z} = \lambda^r \eta_z$ implies that $\gamma_* X = \lambda^r X$, which completes the proof.

At this moment, we need to prove that the germ of the contraction at 0 can be transformed, under a local change of coordinates, into a simpler form. This simpler form is not the Poincaré-Dulac normal form that we will see down below (Section 7.). But this simpler form will help us to see geometric structures of higher order on our Hopf surface.

Lemma 6.2.4. There exists a local system of coordinates $\varphi : U \to V$ between two open neighborhoods of $0 \in \mathbb{C}^2$ such that $\varphi \gamma \varphi^{-1}$ has the form

$$(w_1, w_2) \longmapsto (\lambda w_1, \lambda^r w_2 + f(w_1))$$

where f is a local holomorphic function of one variable, verifying f(0) = 0.

Proof. According to Lemma 6.2.3, there exists a nowhere vanishing vector field X on \mathbb{C}^2 satisfying $\gamma_* X = \lambda^r X$. The flow-box theorem asserts that there exists a system of local coordinates (ξ_1, ξ_2) in a neighborhood of 0 such that $X = \frac{\partial}{\partial \xi_2}$. In these coordinates, the holomorphic 1-form ω reads

$$\omega = g(\xi_1, \xi_2) d\xi_1 + h(\xi_1, \xi_2) d\xi_2.$$

The condition that X belongs to the kernel of ω (see Lemma 6.2.3) implies that the map h is zero. Now, we use that the form ω is closed (see Lemma 6.2.1). This implies that the map g does not depend on ξ_2 . We therefore rewrite

$$\omega = g(\xi_1) d\xi_1.$$

Let G be the anti-derivative of g vanishing at 0. Then we have $\omega = dG$. We consider the map

$$(\xi_1, \xi_2) \mapsto (w_1, w_2) = (G(\xi_1), \xi_2).$$

Since ω does not vanish, G is a local biholomorphism, and the latter map is a change of coordinates in a neighborhood of 0. In this new system of coordinates, the vector field X reads $\frac{\partial}{\partial w_2}$ and the form ω reads dw_1 . Now, the equations $\gamma^*\omega = \lambda\omega$ and $\gamma_*X = \lambda^r X$ imply that, in the coordinates (w_1, w_2) , the contraction γ reads

$$(w_1, w_2) \longmapsto (\lambda w_1, \lambda^r w_2 + f(w_1))$$

which proves the lemma.

The consequence of this lemma is that we immediately see that the contraction γ preserves (integrable) *G*-structures of arbitrary order, namely, if we denote by β the couple of eigenvalues (λ, λ^r) , then γ obviously preserves a G^k_β -structure of order k for every $1 \leq k \leq r$. To see that, it suffices to notice that for any element (x_1, x_2) in \mathbb{C}^2 , the map

$$(w_1, w_2) \mapsto (\lambda(w_1 + x_1), \lambda^r(w_2 + x_2) + f(w_1 + x_1)) - (\lambda x_1, \lambda^r x_2 + f(x_1))$$

has its k-jet at 0 in the group G^k_{β} . In other words, at each point where the map

$$(w_1, w_2) \longmapsto (\lambda w_1, \lambda^r w_2 + f(w_1))$$

is defined, the translation of this map to build a map fixing the origin has its k-jet in the group G_{β}^{k} . Notice that all those structures are compatible. Indeed, they are all induced by the one of order r. We denote by R_{r} the principal G_{β}^{r} sub-bundle of $\mathcal{R}^{r}(\mathbb{C}^{2})$ given by the structure of order r.

What is natural to do is to try to extend the G_{β}^{r} -structure of order r in a G_{β}^{r} structure of order r + 1, using the technique of Theorem 4. The problem here is that the vector bundle we consider in the proof still have a non-trivial first cohomology set.

Let us introduce new groups. We denote by H_{β}^{r+1} (respectively H_{β}^{r+2}) the group $G_{\beta'}^{r+1}$ (respectively $G_{\beta'}^{r+1}$), where β' denotes the set of eigenvalues (λ, λ^{r+2}) . Notice that the introduction of this new set of eigenvalues is purely artificial. The reason to do that is to use what we already did to prove that H_{β}^{r+1} and H_{β}^{r+2} are complex Lie groups (see Section 5.). Using these definitions, Lemma 6.2.4 shows that γ preserves a H_{β}^{r+1} -structure of order r + 1, that will be denoted by \tilde{R}_{r+1} , and a H_{β}^{r+2} -structure of order r + 2, that will be denoted by \tilde{R}_{r+2} . Those structures are again compatible with the others, i.e. all the structures we consider are induced by \tilde{R}_{r+2} .

We consider now extensions of R_r at order r + 2 as sub-bundles of \overline{R}_{r+2} instead of sub-bundles of $\mathcal{R}^{r+2}(\mathbb{C}^2)$. This has the effect of changing the vector bundle that we consider and now we can argue as in Theorem 4 to prove the following.

Theorem 8. There exists a unique holomorphic section of the bundle $\widetilde{R}_{r+2} \to R_r$ which is G_{β}^r -equivariant and $\langle \gamma \rangle$ -equivariant.

Proof. We start with the problem of finding a G_{β}^{r} -equivariant and $\langle \gamma \rangle$ -equivariant section of $\widetilde{R}_{r+1} \to R_{r}$. The latter is an affine line bundle, with fiber isomorphic to $\mathbb{C}Z_{1}^{r+1}e_{2}$ where $Z_{1}^{r+1}e_{2}$ is considered as the homogeneous polynomial map $\mathbb{C}^{2} \to \mathbb{C}^{2}$ of order r + 1 given by $(z_{1}, z_{2}) \mapsto (0, z_{1}^{r+1})$. We infer that, by Hartogs' theorem, it is enough to work on $\mathbb{C}^{2} \setminus \{0\}$. To deal with the $\langle \gamma \rangle$ -equivariance, we work on M. We recall that all the geometric structures over \mathbb{C}^{2} that we consider are $\langle \gamma \rangle$ -invariant.

The difference between two local equivariant sections of $\tilde{R}_{r+1} \to R_r$ is a local section of the induced vector bundle B over M with fiber $\mathbb{C}Z_1^{r+1}e_2$.

As usual (see proofs of Theorems 4 and 7), we need to show, using Mall theorem on surfaces (Theorem 6), that $h^0(M, B) = h^1(M, B) = 0$.

We start by proving that $h^0(M, B) = 0$. We take a global section of B, we pull it back to $\mathbb{C}^2 \setminus \{0\}$ and extend it to \mathbb{C}^2 using Hartogs Theorem. Considering the Taylor expansion at 0 of the section and writing the γ -invariance, we infer that, if the section is nonzero, there exists a relation of the type

$$\alpha_2 = \alpha_1^{r+1} \alpha^p$$

where $p = (p_1, p_2)$ is an element of \mathbb{N}^2 . In our case, this relation reads

$$\lambda^r = \lambda^{r+1+p_1+rp_2}$$

and leads to a contradiction since $0 < |\lambda| < 1$.

Therefore, $h^0(M, B) = 0$. This implies that an eventual global equivariant section of $\widetilde{R}_{d+1} \to R_d$ would be unique, and that, using Theorem 6 and the same computations as in the proof of Theorem 7, we will have $h^1(M, B) = 0$ if a relation of the following type leads to a contradiction:

$$\alpha_1^{r+1} = \alpha_2 \alpha^{-p}$$

where $p = (p_1, p_2)$ is an element of $(-\mathbb{N}^*)^2$. However, such a relation reads

$$\lambda^{r+1} = \lambda^{r-p_1 - rp_2}$$

which is a contradiction. Therefore, we have that $h^1(M, B) = 0$.

We infer that there exists a unique G_{β}^{r} -equivariant section of $R_{r} \to \widetilde{R}_{r+1}$ over the manifold M. Pulling back everything on $\mathbb{C}^{2} \setminus \{0\}$ and extending everything to \mathbb{C}^{2} by Hartogs' Theorem, we get a unique G_{β}^{r} -equivariant section of $R_{r} \to \widetilde{R}_{r+1}$ over the manifold \mathbb{C}^{2} which is $\langle \gamma \rangle$ -equivariant. We denote by R_{r+1} the image of this section and this is a $\langle \gamma \rangle$ -invariant G_{β}^{r} -structure of order r + 1 over \mathbb{C}^{2} .

We apply the same argument to find an equivariant section of the bundle $R_{r+2} \rightarrow R_{r+1}$. The only thing that is different is that we have to consider relations of type

$$\lambda^{r+2} = \lambda^{r-p_1 - rp_2}$$

where $p = (p_1, p_2)$ is an element of $(-\mathbb{N}^*)^2$. Such a relation cannot exist because, by assumption, $r \geq 2$.

Therefore, there exists a unique equivariant section of $\widetilde{R}_{r+2} \to R_{r+1}$, and composing with the first section we computed, we get the result.

Reading the proof of Theorem 8, we can give two remarks. The first one is that the proof breaks if r = 1, i.e. if $d_0\gamma$ has a double eigenvalue. Fortunately, we discussed this case above. The second one is that we cannot produce with this proof for G^r_β -structures of any order, namely, relations of the form

$$\lambda^{2r+1} = \lambda^{r-p_1 - rp_2},$$

where $p = (p_1, p_2)$ is an element of $(-\mathbb{N}^*)^2$, can exist. Indeed, it suffices to choose $p_1 = p_2 = -1$. Since $r \ge 2$, Theorem 8 shows that we can a priori extend only twice the structure of order r. However, we will see in the next section that this is sufficient.

7. Cartan connection, integrability and Poincaré-Dulac theorem

7.1. Another Lie group

In this subsection, we introduce a new complex Lie group of interest. This Lie group will also depend on the data β of the eigenvalues of the contraction generating the Hopf manifold. We begin with a central lemma.

Lemma 7.1.1. For any element P in the complex Lie group G_{β}^{r} and for any z in \mathbb{C}^{n} , the automorphism of \mathbb{C}^{n} given by $\theta_{-P(z)}P\theta_{z}$, where θ_{w} is the translation of vector w, is an element of G_{β}^{r} .

Proof. Since P is a polynomial map of order r, the automorphism $\theta_{-P(z)}P\theta_z$, sending 0 to 0, is also a polynomial map of order r. Moreover, in the Taylor expansion at 0 of the latter polynomial map, the monomials appearing are "bounded above" by those of P, in the sense that if $Z^p e_j$ appears in the expansion of $\theta_{-P(z)}P\theta_z$, where $Z = (Z_1, ..., Z_n)$, $Z^p = Z_1^{p_1}...Z_n^{p_n}$ and e_j is as usual the *j*-th standard coordinate of \mathbb{C}^n , then there exists $q \in \mathbb{N}^n$ such that $p \leq q$ component by component, and such that $Z^q e_j$ appears in P. But then q is j sub resonant for the eigenvalues β , in the sense of Section 5., and therefore, p is also j sub resonant for the eigenvalues β . Therefore, the polynomial map $\theta_{-P(z)}P\theta_z$ is an element of G_{β}^r .

We consider the subgroup $\widetilde{G}_{\beta}^{r}$ of the group of automorphisms of \mathbb{C}^{n} generated by G_{β}^{r} and the translations. Then, as a corollary of Lemma 7.1.1, we get the following.

Corollary 7.1.1. The group $\widetilde{G}_{\beta}^{r}$ is a complex Lie group, biholomorphic to the complex manifold $\mathbb{C}^{n} \times G_{\beta}^{r}$. Through this biholomorphism, the group operation reads

$$(z, P)(w, Q) = (z + P(w), \theta_{-P(w)} P \theta_w Q).$$

Proof. By definition, we have

$$\widetilde{G}_{\beta}^{r} = \langle G_{\beta}^{r}, \{\theta_{z}, z \in \mathbb{C}^{n}\} \rangle_{Aut(\mathbb{C}^{n})}.$$

We consider the map

$$\begin{array}{rcccc} \rho & : & \mathbb{C}^n \times G^r_\beta & \longrightarrow & \widetilde{G}^r_\beta \\ & & (z,P) & \longmapsto & \theta_z P. \end{array}$$

This map is one-to-one, because if $\theta_z P = \theta_w Q$, then w and z are the respective images of 0 by both automorphisms, so z = w and then P = Q. Now the map ρ is onto because of Lemma 7.1.1. Indeed, an element f of \widetilde{G}_{β}^r is a finite composition of elements of G_{β}^r and translations. As an automorphism of \mathbb{C}^n , it sends the origin to a vector z = f(0). Now the composition $P = \theta_{-z} f$ is an element of $\widetilde{G}_{\beta}^{r}$ fixing the origin. But by Lemma 7.1.1, such an element belongs to G_{β}^{r} . Therefore, we have $f = \rho(z, P)$.

Computing the composition $\theta_z P \theta_w Q$, we get (notice that it illustrates the previous claims)

 $\begin{array}{llll} \theta_z P \theta_w Q & = & \theta_z \theta_{P(w)} \theta_{-P(w)} P \theta_w Q \\ & = & \theta_{z+P(w)} \theta_{-P(w)} P \theta_w Q \\ & = & \rho(z+P(w), \theta_{-P(w)} P \theta_w Q). \end{array}$

To finish the proof, we notice that the complex manifold $\mathbb{C}^n \times G_{\beta}^r$ endowed with the group operation given by ρ is naturally a complex Lie group, since every operation involved is holomorphic.

By definition, the complex Lie group G_{β}^r is a Lie subgroup of \widetilde{G}_{β}^r . In the next subsection, we will see that the geometric structures that we build in Section 6. induce a Cartan geometry on the Hopf manifold with model $(\widetilde{G}_{\beta}^r, G_{\beta}^r)$ (see Section 2.3.).

We recall from Section 2.1. the existence of the representation $\underline{Ad}^{r+1}: \mathcal{D}^{r+1}(\mathbb{C}^n) \to GL(T_0\mathbb{C}^n \times \mathfrak{d}^r(\mathbb{C}^n))$. Using Corollary 7.1.1, the Lie algebra of \widetilde{G}^r_{β} is isomorphic (as complex vector spaces, not as Lie algebras) with $T_0\mathbb{C}^n \times \mathfrak{g}^r_{\beta}$, where \mathfrak{g}^r_{β} is the Lie algebra of G^r_{β} . Therefore, it can be seen as a subspace of $T_0\mathbb{C}^n \times \mathfrak{d}^r(\mathbb{C}^n)$.

Lemma 7.1.2. Let P be an element of G_{β}^r . The space $T_0\mathbb{C}^n \times \mathfrak{g}_{\beta}^r$ is preserved by the linear map \underline{Ad}_P^{r+1} , where P is seen as an element of $\mathcal{D}^r(\mathbb{C}^n)$. Moreover, when restricted to $T_0\mathbb{C}^n \times \mathfrak{g}_{\beta}^r$, the linear transformation \underline{Ad}_P^{r+1} coincides with the adjoint representation of the element P in the group \widetilde{G}_{β}^r .

Proof. By definition of \underline{Ad}^{r+1} (see Section 2.1.), we have that

$$\underline{Ad}_{P}^{r+1} = d_{j_{0}^{r}id}\xi_{r} \circ d_{j_{0}^{r}id}(j_{0}^{r}f \mapsto j_{0}^{r}(PfP^{-1})) \circ (d_{j_{0}^{r}id}\xi_{r})^{-1} \\
= d_{(0,j_{0}^{r}id)}(\xi_{r} \circ j_{0}^{r}f \mapsto j_{0}^{r}(PfP^{-1}) \circ \xi_{r}^{-1}) \\
= d_{(0,j_{0}^{r}id)}((z,j_{0}^{r}\phi) \mapsto (P(z),j_{0}^{r}(\theta_{-P(z)}P\theta_{z}\phi P^{-1}))).$$

But if z is a vector in \mathbb{C}^n and ϕ is an element of G^r_β , then by Proposition 5.0.1 and by Lemma 7.1.1, then $(P(z), j^r_0(\theta_{-P(z)}P\theta_z\phi P^{-1}))$ is an element of $\mathbb{C}^n \times G^r_\beta$. Therefore, the space $T_0\mathbb{C}^n \times \mathfrak{g}^r_\beta$ is preserved by the linear map \underline{Ad}^{r+1}_P .

Moreover, computing the interior homomorphism Int_P in the group \widetilde{G}^r_{β} , we have

$$P\theta_w Q P^{-1} = \theta_{P(w)} \theta_{-P(w)} P\theta_w Q P^{-1},$$

so by Corollary 7.1.1, we have the equality

$$\underline{Ad}_P^{r+1}|_{T_0\mathbb{C}^n\times\mathfrak{g}_\beta^r} = Ad_P$$

where the adjoint representation of the right hand side is the one of the group G_{β}^{r} .

7.2. The Cartan geometry induced by the G-structures

We recall that r is the maximal length of the resonances of the set of eigenvalues $\beta = (\beta_1, ..., \beta_n)$. Using this definition, we deduce that for every $k \ge r$, we have

$$G^k_\beta = G^r_\beta$$

Moreover, since G_{β}^{r} is a Lie group for the usual composition (not the truncated composition), for every $k \geq r$, G_{β}^{r} can be seen as a Lie subgroup of $\mathcal{D}^{k}(\mathbb{C}^{n})$, the group of jets of order k at 0 of germs of biholomorphism $(\mathbb{C}^{n}, 0) \to (\mathbb{C}^{n}, 0)$.

If the dimension n is not lower than 3, then, Corollary 6.1.1 shows that there exists a G^r_{β} -structure of order r on M. If n = 2, this is proved in Section 6.2.3.. This structure can be written as a G^r_{β} -principal sub-bundle R_r of $\mathcal{R}^r(M)$. Using Theorem 4 if $n \geq 3$ and Theorem 8 for surfaces, and using the previous observations, there exists a unique G^r_{β} -structure of order r + 1 on M extending R_r .

This can be reformulated as follows. There exists a unique G_{β}^{r} -equivariant section

$$\Gamma_r: R_r \longrightarrow \mathcal{R}^{r+1}(M)$$

of the natural map

$$\mathcal{R}^{r+1}(M) \longrightarrow \mathcal{R}^r(M).$$

Notice that the uniqueness in the case of surfaces comes from the fact that in the proof of Theorem 4, the vanishing of the h^0 of the considered bundle is still true for surfaces.

We recall from Section 2.1. the existence of the canonical form $\chi_{r+1} : T\mathcal{R}^{r+1}(M) \to T_0\mathbb{C}^n \times \mathfrak{d}^r(\mathbb{C}^n)$ on $\mathcal{R}^{r+1}(M)$. We consider η_r the holomorphic 1-form on R_r with values in $T_0\mathbb{C}^n \times \mathfrak{d}^r(\mathbb{C}^n)$ defined by

$$\eta_r = \Gamma_r^* \chi_{r+1}.$$

As we saw in Section 2.1., this exhibits the fact that Γ_r is a $\mathcal{R}^r(M)$ -horizontal plane field on R_r , and it generates a field of frames of order 1 of $\mathcal{R}^r(M)$ over R_r . This implies that at each point $j_0^r f \in R_r$, the linear map $(\eta_r)_{j_0^r f}$ is one-to-one onto its image.

By the equivariance property of Γ_r , and by Proposition 2.1.1, the form η_r satisfies the following properties

- 1. $\forall v \in \mathfrak{g}_{\beta}^{r}, \eta_{r}(X_{v}) \equiv (0, v)$, where X_{v} is the fundamental vector field of R_{r} generated by v;
- 2. $\forall P \in G_{\beta}^r, \mathcal{R}_P^* \eta_r = \underline{Ad}_{P^{-1}}^{r+1} \eta_r.$

The main result of this subsection is the following.

Theorem 9. The form η_r takes values in $T_0\mathbb{C}^n \times \mathfrak{g}_{\beta}^r$ and defines a holomorphic Cartan geometry of type $(\widetilde{G}_{\beta}^r, G_{\beta}^r)$ on the principal bundle R_r .

Proof. We recall that we can consider every of our geometric objects as defined on \mathbb{C}^n and with the property of being $<\gamma>$ -equivariant.

According to the Lemma 7.1.2, the representation $\underline{Ad}^{r+1}|_{G^r_\beta}$ descends to a representation, denoted by \underline{Ad}

$$G^r_{\beta} \longrightarrow GL((T_0\mathbb{C}^n \times \mathfrak{d}^r(\mathbb{C}^n))/(T_0\mathbb{C}^n \times \mathfrak{g}^r_{\beta}))$$

and the form $\tilde{\eta}_r : TR_r \to (T_0\mathbb{C}^n \times \mathfrak{d}^r(\mathbb{C}^n))/(T_0\mathbb{C}^n \times \mathfrak{g}_\beta^r)$, obtained by applying the standard quotient map to η_r , satisfies the following properties

- 1. $\forall v \in \mathfrak{g}_{\beta}^{r}, \, \widetilde{\eta_{r}}(X_{v}) \equiv 0$, where X_{v} is the fundamental vector field of R_{r} generated by v;
- 2. $\forall P \in G_{\beta}^r, \mathcal{R}_P^* \widetilde{\eta_r} = \underline{Ad}_{P^{-1}}^{r+1} \widetilde{\eta_r}.$

Notice that we have an isomorphism

$$T_0\mathbb{C}^n \times \mathfrak{d}^r(\mathbb{C}^n))/(T_0\mathbb{C}^n \times \mathfrak{g}_\beta^r) \cong \mathfrak{d}^r(\mathbb{C}^n)/\mathfrak{g}_\beta^r$$

Therefore, $\tilde{\eta}_r$ can be seen as a holomorphic $\langle \gamma \rangle$ -invariant 1-form on \mathbb{C}^n with value in the associated vector bundle

$$R_r \times_{G^r_\beta} \mathfrak{d}^r(\mathbb{C}^n)/\mathfrak{g}^r_\beta$$

where G^r_{β} acts on $\mathfrak{d}^r(\mathbb{C}^n)/\mathfrak{g}^r_{\beta}$ via

$$\begin{array}{ccc} G^r_{\beta} & \longrightarrow & GL(\mathfrak{d}^r(\mathbb{C}^n)) \\ P & \longmapsto & \mathrm{d}_{j^r_0id}(j^r_0\phi \mapsto j^r_0(P\phi P^{-1})) \end{array}$$

We will show that such an invariant 1-form on \mathbb{C}^n vanishes.

Following the computations of Ornea and Verbitsky ([28], 5.2. Resonant equivariant bundles p.16), in the proof of Theorem 4 and the computations in the proofs of Section 6.2.3., we need to compute the eigenvalues of the action of γ on the fiber of the bundle $R_r \times_{G_{\beta}^r} \mathfrak{d}^r(\mathbb{C}^n)/\mathfrak{g}_{\beta}^r$ over the origin. The typical fiber of this bundle is the space $\mathfrak{d}^r(\mathbb{C}^n)/\mathfrak{g}_{\beta}^r$, and in a well chosen isomorphism, the action of γ is given by $\underline{Ad}_{j_0^{r+1}\gamma^{-1}}^{r+1}$.

The homomorphisms $\rho_{r,s} : \mathcal{D}^r(\mathbb{C}^n) \to \mathcal{D}^s(\mathbb{C}^n)$ for $0 \leq s \leq r$ gives a filtration by normal Lie subgroups of the group $\mathcal{D}^r(\mathbb{C}^n)$

$$\{id\}=N^r_r\subset N^r_{r-1}\subset \ldots \subset N^r_1\subset N^r_0=\mathcal{D}^r(\mathbb{C}^n)$$

where $N_s^r = \ker(\rho_{r,s})$. This filtration by normal subgroups gives a filtration by adjoint invariant Lie sub-algebras

$$\{0\} = \mathfrak{n}_r^r \subset \mathfrak{n}_{r-1}^r \subset \ldots \subset \mathfrak{n}_1^r \subset \mathfrak{n}_0^r = \mathfrak{d}^r(\mathbb{C}^n).$$

Therefore, as in the proof of Theorem 4, the eigenvalues of the linear map $\underline{Ad}_{j_0^{r+1}\gamma^{-1}}^{r+1}$ over the space $\mathfrak{d}^r(\mathbb{C}^n)/\mathfrak{g}_{\beta}^r$ are

$$\beta_i^{-1}\beta^p$$

with $j \in \{1, ..., n\}$, $p \in \mathbb{N}^n$ of length satisfying $1 \leq |p| \leq r$ and p is not j sub-resonant.

Hence, as usual, by taking Taylor expansion at 0 of a γ -invariant holomorphic 1-form with value in the bundle $R_r \times_{G^r_{\beta}} \mathfrak{d}^r(\mathbb{C}^n)/\mathfrak{g}^r_{\beta}$, we see that if the section is non identically zero, then there is a relation of the form

$$\beta^k \beta_l \beta_i^{-1} \beta^p = 1$$

where k is an element of \mathbb{N}^n , l is an element of $\{1, ..., n\}$ and j and p are as above. But the existence of such a relation is a contradiction, because it implies that p is j sub-resonant.

Therefore, our form $\tilde{\eta}_r$ is identically zero. We infer the first statement in the theorem, namely, that η_r takes values in $T_0\mathbb{C}^n \times \mathfrak{g}_{\beta}^r$.

According to our previous observations, at each point of R_r , the form η_r is oneto-one onto its image, so if we consider it as a form with value in $T_0\mathbb{C}^n \times \mathfrak{g}_{\beta}^r$, by an argument of dimensions, η_r is an absolute parallelism. Moreover, we already know that the fundamental vector fields of R_r are η_r constant. The equivariance property of η_r required in the definition of a Cartan geometry is given by Lemma 7.1.2.

This result is equivalent to the following fact. The map $\Gamma_r : R_r \to \mathcal{R}^{r+1}(M)$ is a holomorphic horizontal plane field of the bundle $R_r \to M$. Notice that this gives rise to a connection on the principal bundle $R_r \to M$ but that this connection is not principal (except if r = 1, in which case this connection induces the unique affine connection exhibited by Ornea and Verbitsky in [28], p.27 Theorem 6.6).

7.3. Integrability of the G-structure of order r+1

Let R_r be a G_{β}^r -structure of order r on M, and $\Gamma_r : R_r \to \mathcal{R}^{r+1}(M)$ be the G_{β}^r equivariant section given by Theorem 4 if $n \geq 3$, and Theorem 8 if n = 2.

Applying successively Theorem 4, or Theorem 8, we get also uniquely determined G_{β}^{r} equivariant sections

$$\Gamma_k : \operatorname{im}(\Gamma_{k-1}) \longrightarrow \mathcal{R}^{k+1}(M)$$

for every $k \ge r+1$ if $n \ge 3$, for every k = r+1 if n = 2. We denote by R_k the image of Γ_{k-1} . From this, we can prove the following.

Proposition 7.3.1. The G_{β}^{r} -structure of order r + 1 R_{r+1} is rigid of order r, in the sense of rigidity for Gromov geometric structures (see Section 2.2.4).

Proof. Let x be a point of the manifold M, and let $j_x^{r+1}f$ be a jet of order r+1 at x of a germ of biholomorphism of M at x, and fixing x. Assume, moreover, that this jet preserves the sub-bundle R_{r+1} . Then, by the equivariance property of a jet of a biholomorphism, $j_x^r f$ preserves R_r .

Let $j_0^{r+1}g$ be an element of $R_{r+1}|_x$. The jet $j_x^{r+1}f$ is completely determined by the jet $j_0^{r+1}(fg)$. Let P be the polynomial map of G_{β}^r such that $j_0^r(fg) = j_0^r(gP)$, then by the equivariance property of Γ_r , we have that

$$j_0^{r+1}(fg) = j_0^{r+1}(gP).$$

Therefore, $j_x^{r+1}f$ is completely determined by P, which is itself completely determined by $j_x^r f$. Hence, the structure R_{r+1} is rigid at order r.

We will now prove the integrability of the G_{β}^{r} -structure of order r+1. The strategy is to use the point of view of Benoist, and namely, to use Theorem 1.

Theorem 10. The G_{β}^{r} -structure of order r + 1 R_{r+1} is integrable.

Proof. We will prove that the reduction R_{r+1} is, roughly speaking, complete and consistent in the terminology of Benoist ([3], Section 3.2 p.12, see Definition 2.2.5). The Frobenius type theorem of Benoist ([3], Section 3.2 p.12, see Theorem 1) will provide a germ of solution of the partial derivative relation R_{r+1} at $0 \in \mathbb{C}^n$.

As we saw before, using Hartogs Theorem (the extensions of our geometric structures are sections of trivial affine bundles), every structure that we have can be seen as a geometric structure on \mathbb{C}^n , invariant under the action of the contraction γ , so we consider now R_k as a G^r_β -structure of order k on \mathbb{C}^n invariant by γ , for every $r \leq k \leq r+2$ and

$$\Gamma_k: R_k \longrightarrow R_{k+1}$$

for every $r \leq k \leq r+1$. Notice that we will only use the extension of order 2 or R_r , so we do not need to distinguish between $n \geq 3$ and n = 2.

For every integer $k \ge 1$, we consider $J_k := J^k(\mathbb{C}^n, \mathbb{C}^n)$ the space of jets of order k of germs of holomorphic maps from \mathbb{C}^n to itself. We also consider

$$E_{r+1} := \{ j_z^{r+1} f \in J_{r+1} \mid j_0^{r+1}(f\theta_z) \in R_{r+1} \}.$$

We will show that the partial derivative relation E_{r+1} is complete and consistent in the terminology of Benoist ([3] and Definition 2.2.5). We denote by p_{r+1} the standard projection

$$p_{r+1}: J_{r+1} \longrightarrow J_r.$$

Then we will show that p_{r+1} , when restricted to E_{r+1} , is a biholomorphism onto its image. This, by definition, proves that E_{r+1} is complete. Suppose that $j_z^{r+1}f$ and $j_w^{r+1}g$ are elements of E_{r+1} such that

$$j_z^r f = j_w^r g$$

Then z = w and

$$j_0^r(f\theta_z) = j_0^r(g\theta_z).$$

But since $j_z^{r+1}f$ and $j_w^{r+1}g$ are elements of E_{r+1} , then $j_0^r(f\theta_z)$ and $j_0^r(g\theta_z)$ are elements of R_r and

$$\begin{cases} \Gamma(j_0^r(f\theta_z)) &= j_0^{r+1}(f\theta_z) \\ \Gamma(j_0^r(g\theta_z)) &= j_0^{r+1}(g\theta_z) \end{cases}$$

Therefore, we have that

$$j_0^{r+1}(f\theta_z) = j_0^{r+1}(g\theta_z),$$

which implies that

$$j_z^{r+1}f = j_z^{r+1}g.$$

This proves that p_{r+1} restricted to E_{r+1} is one-to-one, hence a biholomorphism onto its image. We deduce that the partial derivative relation E_{r+1} is complete.

We will now show that E_{r+1} is consistent. By definition, for any element x of E_{r+1} , we need to find a local section of $E_{r+1} \to \mathbb{C}^n$ passing through x and tangent at x to an holonomic section, i.e. a section that is actually a jet of a holomorphic germ.

We consider the G_{β}^{r} equivariant section $\Gamma_{r+1} : R_{r+1} \to \mathcal{R}^{r+2}(M)$. Applying exactly the same proof as in Theorem 9, we can see the map Γ_{r+1} as a connection on the bundle $R_{r+1} \to M$.

We consider the map

$$\begin{array}{rcccc} H_{r+1} & : & E_{r+1} & \longrightarrow & J_{r+2} \\ & & j_z^{r+1} f & \longmapsto & j_z^{r+2} (\widehat{f\theta_z} \theta_{-z}) \end{array}$$

where $f\theta_z$ is a germ representing $\Gamma_{r+1}(j_0^{r+1}(f\theta_z))$.

Since the map Γ_{r+1} gives rise to a connection on the bundle $R_{r+1} \to M$, the map H_{r+1} is a horizontal plane field for the bundle $J^{r+1} \to \mathbb{C}^n$ over E_{r+1} which is tangent to E_{r+1} and holonomic. At each point of E_{r+1} , we consider a local section tangent at the point to the horizontal plane given by H_{r+1} . This is a section that is tangent to a holonomic plane. Therefore, the partial derivative relation E_{r+1} is consistent.

By Theorem 1, at each point of E_{r+1} , there exists a unique germ of solution passing through this point.

Let p be an element of R_{r+1} , and let f be a germ at 0 of solution of E_{r+1} such that $p = j_0^{r+1} f$. Then by definition, the biholomorphic germ f maps the translations of \mathbb{C}^n into R_{r+1} , and gives rise to a local isomorphism between the standard G_{β}^r -structure or order r+1 of \mathbb{C}^n and R_{r+1} .

This concludes the proof that R_{r+1} is integrable.

As an immediate consequence, R_r is also integrable. In the language of Cartan geometries, we have the following.

Corollary 7.3.1. The Cartan geometry η_r on the bundle R_r (see Theorem 9) is flat.

Proof. According to Theorem 10, the G_{β}^r -structure of order r + 1 R_{r+1} is integrable. Let U be an open set in M and V be an open set in \mathbb{C}^n such that there exists an integrability chart $f: V \to U$ for R_{r+1} . It implies that f is also an integrability chart for R_r and that the jet map

$$j^r f: V \times \mathcal{D}^r(\mathbb{C}^n) \longrightarrow \mathcal{R}^r(M)$$

induces a isomorphism of G_{β}^{r} -structures of order r, still denoted

$$j^r f: V \times G^r_\beta \longrightarrow R_r.$$

Notice that $V \times G_{\beta}^{r}$ can be seen as an open set of $\widetilde{G}_{\beta}^{r}$ (see Corollary 7.1.1). We compute the pullback of η_{r} by the map $j^{r}f$:

$$(j^r f)^* \eta_r = (j^r f)^* (\Gamma_r^* \chi_{r+1}) = (\Gamma_r \circ j^r f)^* \chi_{r+1} = (j^{r+1} f)^* \chi_{r+1}$$

where $j^{r+1}f$ is considered as a map $V \times G_{\beta}^r \longrightarrow R_{r+1}$ and the last equality holds because of the fact that f is an integrability chart for R_{r+1} .

The canonical form on $\mathcal{R}^{r+1}(M)$ is pulled back by $j^{r+1}f$ on the canonical form on $\mathcal{R}^{r+1}(\mathbb{C}^n)$. But now according to Corollary 7.1.1, the restriction of the canonical form of $\mathcal{R}^{r+1}(\mathbb{C}^n)$ to the group \widetilde{G}^r_{β} coincides with the Maurer Cartan form on the latter. Therefore, the form η_r is pulled back on the Maurer Cartan form of \widetilde{G}^r_{β} , meaning that $j^r f$ is an isomorphism of Cartan geometries. By Theorem 2, η_r is flat.

7.4. Poincaré-Dulac theorem

We will now give a new proof of the Poincaré-Dulac Theorem in the context of global holomorphic contractions of \mathbb{C}^n in 0.

Theorem 11. Let γ be an invertible contraction of \mathbb{C}^n in 0, where $n \geq 2$, let $\beta = (\beta_1, ..., \beta_n)$ be the collection of eigenvalues of $d_0\gamma$, and let r be the maximal length of the resonances of β . Then there exists a global biholomorphism $\phi : \mathbb{C}^n \to \mathbb{C}^n$ such that $\phi^{-1}\gamma\phi$ is an element of G^r_{β} .

Proof. According to Theorem 10, there exists an integrable G_{β}^{r} -structure on \mathbb{C}^{n} invariant under the action of γ , denoted by R_{r} . Let U, V be two neighborhoods of $0 \in \mathbb{C}^{n}$ and $\phi: U \to V$ be an integrable chart of R_{r} on V mapping 0 to 0. Now, since γ preserves $R_{r}, \gamma \circ \phi$ is also an integrable chart. Hence, for every z close enough to $0, j_{0}^{r}(\phi \circ \theta_{\phi^{-1}\gamma\phi(z)})$ and $j_{0}^{r}(\gamma \circ \phi \circ \theta_{z})$ are in the same G_{β}^{r} orbit. We deduce that for every z close enough to $0, j_{0}^{r}(\theta_{-\phi^{-1}\gamma\phi(z)} \circ \phi^{-1} \circ \gamma \circ \phi \circ \theta_{z})$ is an element of G_{β}^{r} . But this implies that the germ $\phi^{-1} \circ \gamma \circ \phi$ is itself an element of G_{β}^r . Therefore, it extends to a polynomial map P in G_{β}^r .

To conclude the proof, since γ is a contraction in 0, P is also a contraction in 0 and the germ of biholomorphism ϕ^{-1} gives a biholomorphism between the Hopf manifolds M and $(\mathbb{C}^n \setminus \{0\})/\langle P \rangle$. Therefore, ϕ^{-1} , and a fortiori ϕ are globally defined biholomorphic maps.

Notice that here, we produce new coordinates in which γ reads as a sub-resonant polynomial map. The Poincaré-Dulac theorem is slightly stronger, in the sense that we can find a system of coordinates in which γ reads as a resonant polynomial map. But, starting with Theorem 11 and following the algebraic part of the proof of the Poincaré-Dulac theorem, we can find a polynomial change of coordinates (actually an element of G_{β}^{r}) such that γ reads like a resonant polynomial map. In other words, in the conjugacy class of γ in G_{β}^{r} , there is a resonant polynomial contraction. One of these resonant polynomial contractions is what is called a Poincaré-Dulac normal form of the contraction.

If γ is a contraction of \mathbb{C}^n at 0, and γ reads in the standard coordinates of \mathbb{C}^n like an element of its associated group G_{β}^r , then γ obviously preserves all the standard G_{β}^k -structures of order k of \mathbb{C}^n . All of those structures descend to the Hopf manifold and are obviously integrable. We deduce the following corollary.

Corollary 7.4.1. Let M be a Hopf manifold given by a contraction γ that induces the set of eigenvalues $\beta = (\beta_1, ..., \beta_n)$ and the maximal length r. Then

- 1. there exist integrable G_{β}^{k} -structures of order k on M for every $k \geq 1$. Moreover, those structures can be chosen so that their pullback on $\mathbb{C}^{n} \setminus \{0\}$ extends to \mathbb{C}^{n} .
- 2. there exists a flat Cartan geometry on M with model $(\widetilde{G}_{\beta}^{r}, G_{\beta}^{r})$. Moreover, this structure can be chosen so that its pullback on $\mathbb{C}^{n} \setminus \{0\}$ extends to \mathbb{C}^{n} .

Acknowledgments

We are very grateful to Sorin Dumitrescu and Laurent Meersseman for their helpful support throughout the development of this work. We also thank Bertrand Deroin, Frank Loray and Paul Boureau for the interesting discussions around the subject.

References

- D. V. Alekseevskij, A. M. Vinogradov, and V. V. Lychagin. Basic ideas and concepts of differential geometry. In *Geometry*, I, volume 28 of *Encyclopaedia Math. Sci.*, pages 1–264. Springer, Berlin, 1991.
- [2] V. I. Arnold. Geometrical methods in the theory of ordinary differential equations, volume 250 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, New York, second edition, 1988. Translated from the Russian by Joseph Szücs [József M. Szűcs].
- [3] Yves Benoist. Orbites des structures rigides (d'après M. Gromov). In Integrable systems and foliations/Feuilletages et systèmes intégrables (Montpellier, 1995), volume 145 of Progr. Math., pages 1–17. Birkhäuser Boston, Boston, MA, 1997.
- [4] Indranil Biswas, Sorin Dumitrescu, and Georg Schumacher. Deformation theory of holomorphic Cartan geometries. *Indag. Math. (N.S.)*, 31(3):512–524, 2020.
- [5] Ciprian Borcea. Some remarks on deformations of Hopf manifolds. Rev. Roumaine Math. Pures Appl., 26(10):1287–1294, 1981.
- [6] Paul Boureau. Normal forms and geometric structures on hopf manifolds.
- [7] M. Brunella, S. Dumitrescu, P. Eyssidieux, A. Glutsyuk, L. Meersseman, and M. Nicolau. Complex manifolds, foliations and uniformization, volume 34/35 of Panoramas et Synthèses [Panoramas and Syntheses]. Société Mathématique de France, Paris, 2011. Courses from the Summer School "Uniformisation de Familles de Variétés Complexes" held in Dijon, August 31–September 11, 2009.
- [8] Adrien Douady. Déformations régulières. Séminaire Henri Cartan, 13(1):1–8, 1960.
- [9] Jean Frenkel. Cohomologie non abélienne et espaces fibrés. Bull. Soc. Math. France, 85:135–220, 1957.
- [10] Avner Friedman. Foundations of modern analysis. Holt, Rinehart and Winston, Inc., New York-Montreal, Que.-London, 1970.
- [11] Peter B Gothen. Representations of surface groups and higgs bundles. Moduli Spaces, LMS Lecture Notes Series, 411:151–178, 2014.
- [12] Hans Grauert and Reinhold Remmert. Theory of Stein spaces. Classics in Mathematics. Springer-Verlag, Berlin, 2004. Translated from the German by Alan Huckleberry, Reprint of the 1979 translation.
- [13] Michael Gromov. Rigid transformations groups. In Géométrie différentielle (Paris, 1986), volume 33 of Travaux en Cours, pages 65–139. Hermann, Paris, 1988.
- [14] Victor Guillemin. The integrability problem for g-structures. Transactions of the American Mathematical Society, 116:544–560, 1965.

- [15] A. Haefliger. Deformations of transversely holomorphic flows on spheres and deformations of Hopf manifolds. *Compositio Math.*, 55(2):241–251, 1985.
- [16] H. Hopf. Zur Topologie der komplexen Mannigfaltigkeiten. In Studies and Essays Presented to R. Courant on his 60th Birthday, January 8, 1948, pages 167–185. Interscience Publishers, New York, 1948.
- [17] Mikio Ise. On the geometry of Hopf manifolds. Osaka Math. J., 12:387–402, 1960.
- [18] Shoshichi Kobayashi. Canonical forms on frame bundles of higher order contact. In Proc. Sympos. Pure Math., Vol. III, pages 186–193. Amer. Math. Soc., Providence, RI, 1961.
- [19] Shoshichi Kobayashi. Transformation groups in differential geometry. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1972 edition.
- [20] Jean Jacques Loeb and Marcel Nicolau. Holomorphic flows and complex structures on products of odd-dimensional spheres. *Math. Ann.*, 306(4):781–817, 1996.
- [21] Daniel Mall. Contractions, Fredholm operators and the cohomology of vector bundles on Hopf manifolds. Arch. Math. (Basel), 66(1):71–76, 1996.
- [22] Benjamin McKay. The Hartogs extension problem for holomorphic parabolic and reductive geometries. Monatsh. Math., 181(3):689–713, 2016.
- [23] Benjamin McKay and Alexey Pokrovskiy. Locally homogeneous structures on Hopf surfaces. Indiana Univ. Math. J., 59(4):1491–1539, 2010.
- [24] Pierre Molino. Théorie des G-structures: le problème d'équivalence, volume 588. Springer, 2006.
- [25] Liviu Ornea and Misha Verbitsky. Embedding of LCK manifolds with potential into Hopf manifolds using Riesz-Schauder theorem. In *Complex and symplectic geometry*, volume 21 of *Springer INdAM Ser.*, pages 137–148. Springer, Cham, 2017.
- [26] Liviu Ornea and Misha Verbitsky. Embedding of lck manifolds with potential into hopf manifolds using riesz-schauder theorem. Complex and symplectic geometry, pages 137–148, 2017.
- [27] Liviu Ornea and Misha Verbitsky. Non-linear hopf manifolds are locally conformally kähler. The Journal of Geometric Analysis, 33(7):201, 2023.
- [28] Liviu Ornea and Misha Verbitsky. Mall bundles and flat connections. Online first, 2024.
- [29] Viktor Pavlovich Palamodov. Deformations of hopf manifolds and the poincaré—dulac theorem. Functional Analysis and Its Applications, 17(4):252–259, 1983.
- [30] L Reich. Das typenproblem bei formal-biholomorphen abbildungen mit anziehendem fixpunkt. Mathematische Annalen, 179(3):227–250, 1969.
- [31] R.W. Sharpe and S.S. Chern. Differential Geometry: Cartan's Generalization of Klein's Erlangen Program. Graduate Texts in Mathematics. Springer New York, 2000.
- [32] Shlomo Sternberg. Lectures on differential geometry. Prentice-Hall, Inc., Englewood Cliffs, NJ, 1964.
- [33] Joachim Wehler. Versal deformation of Hopf surfaces. J. Reine Angew. Math., 328:22-32, 1981.

UNIVERSITÉ COTE D'AZUR, LJAD, FRANCE Email address: matthieu.madera@univ-cotedazur.fr