# The distance problem via subadditivty

Renan Gross\*

#### Abstract

In a recent paper, Aldous, Blanc and Curien asked which distributions can be expressed as the distance between two independent random variables on some separable measured metric space. We show that every nonnegative discrete distribution whose support contains 0 arises in this way, as well as a class of finitely supported distributions with density.

### 1 Introduction

#### 1.1 The problem

The distance problem asks the following. Given a distribution  $\theta$  on  $[0, \infty)$ , is it possible to find a complete separable measured metric space  $(S, d, \mu)$ , so that when  $X, Y \sim \mu$  are independent, their distance satisfies  $d(X, Y) \sim \theta$ ? If so, we say that  $(S, d, \mu)$  achieves  $\theta$ , and that  $\theta$  is feasible. This very natural and quite inviting problem was first proposed by Aldous, Blanc and Curien in [ABC24], where they also give some statistical motivation.

#### 1.2 Examples

While any  $(S, d, \mu)$  gives some distribution  $\theta$  which can in principle be calculated, the inverse problem, of finding an explicit space which achieves a given distribution, is harder. The following examples show how to achieve some common distributions, and may give some intuition about the types of constructions one can use.

- 1. The uniform measure. Let  $S = S^1 = \mathbb{R} \mod \mathbb{Z}$  be the unit circle,  $\mu$  the uniform measure on  $S^1$ , and d(x, y) the arc length between x and y. Then  $d(X, Y) \sim U\left[0, \frac{1}{2}\right]$ .
- 2. The exponential distribution. Let  $S = \mathbb{R}$ , let d(x, y) = |x y|, and let  $\mu = \exp(1)$  be the exponential distribution. By the memoryless property, |X Y| is also exponentially distributed.
- 3. Bernoulli distribution. Let  $p_0 > 0$ . In order to achieve the Bernoulli distribution  $\theta = p_0 \delta_0 + (1-p_0) \delta_1$ , let  $m \ge 2$  be an integer and let  $q \in [0,1]$  to be chosen later. Let  $S = \{0, \ldots, m-1\}$ , equipped with the discrete metric

$$d(x,y) = \begin{cases} 0 & x = y \\ 1 & x \neq y \end{cases}$$

Let  $\mu$  be the distribution of the random variable on S whose value is 0 with probability q, and uniform among  $\{1, \ldots, m-1\}$  with probability 1-q. Then

$$\mathbb{P}\left[d\left(X,Y\right)=0\right] = q^2 + \left(\frac{1-q}{m-1}\right)^2.$$

<sup>\*</sup>University of Cambridge. Email: rg751@cam.ac.uk

Choosing m so that the quadratic equation  $p_0 = q^2 + \left(\frac{1-q}{m-1}\right)^2$  has a solution and setting

$$q = \frac{1 + \sqrt{1 + \left((m-1)^2 + 1\right)\left(p_0\left(m-1\right)^2 - 1\right)}}{\left(m-1\right)^2 + 1}$$
(1)

gives us the desired measured probability space.

- 4. The uniform measure, revisited. Let  $S = \{0, 1\}^{\mathbb{N}}$ , let  $\mu$  be the measure on S where all entries are i.i.d. 0-1 Bernoulli random variables with success probability 1/2, and let  $d(x, y) = \sum_{k=1}^{\infty} 2^{-k} |x_i y_i|$ . Then  $d(X, Y) \sim U[0, 1]$ . This is essentially a more explicit version of the first example.
- 5. The exponential distribution, revisited. Let S = (0, 1],  $\mu$  the uniform measure on S, and  $d(x, y) = \left|\int_x^y \frac{1}{t} dt\right| = \left|\log \frac{y}{x}\right|$ . A short calculation shows that  $\mathbb{P}\left[d(X, Y) \leq t\right] = 1 e^{-t}$ , and so  $d(X, Y) \sim \exp(1)$ .
- 6. Non-example. Let  $\theta$  be any distribution with density on  $[0, \infty)$ , and denote its cumulative distribution function by F. Let  $S = [0, \infty)$ , let  $\mu$  be the distribution whose CDF is  $\sqrt{F}$ , and let

$$d(x,y) = \begin{cases} 0 & x = y \\ \max(x,y) & x \neq y. \end{cases}$$

Since  $\theta$  has density, the event  $\{x = y\}$  has measure 0 under  $\mu$ , and so  $d(X,Y) = \max(X,Y)$  with probability 1. The CDF of the maximum of two i.i.d. random variables is the square of their individual CDF, and so  $d(X,Y) \sim \theta$ . However, (S,d) is not separable (for small enough  $\varepsilon$ , the only element  $\varepsilon$ -close to x is x itself).

#### 1.3 Results

In [ABC24], Aldous, Blanc and Curien showed some basic constraints on the support of the distribution – for example, it must contain 0 – and gave several constructions and approximations. In their main constructions, the metric space S was a weighted tree and the measure  $\mu$  a distribution on its leaves / rays. Choosing the weights and distribution in a clever way, they showed that every finite discrete distribution with 0 in its support is feasible.

Our first result resolves the question of discrete distributions.

**Theorem 1.** Every nonnegative discrete distribution  $\theta$  supported on 0 is feasible. (proof  $\nearrow$ )

Rather than building a metric space  $(S, d, \mu)$  which directly achieves  $\theta$  from scratch, our proof combines together several simpler spaces  $(S_n, d_n, \mu_n)$ , allowing us to sample from different  $\theta_n$ s with different probabilities. The main challenge is to decompose  $\theta$  into subdistributions so that combining the spaces preserves the triangle inequality. As will be described in the next section, this construction can also be interpreted as a metric on the rays of a tree, where two rays are close if they travel together for a long time, generalizing a commonly used metric on the space of ends of a tree.

Our second result concerns distributions with density.

**Theorem 2.** Let g be a probability density function on [0, 1], and suppose that there exist constants c, C > 0 such that

$$c < g\left(t\right) < C \tag{2}$$

for all  $t \in [0,1]$ . Then the distribution  $\theta$  with density g is feasible. (proof  $\nearrow$ )

The theorem is derived by altering the metric of a known-to-be-feasible distribution; here, subadditivity is used in order to preserve the metric. The theorem represents one step towards answering the open problem in [ABC24]:

**Problem 3.** Prove that for every probability density function f on  $[0, \infty)$  whose support contains 0, the distribution with density f is feasible.

### 2 Proof of Theorem 1

Our proof uses the following construction, which, given several feasible distributions, returns a mixture of these distributions. The resultant measure can be thought of as being computed by a "selection" mechanism: we linearly order the distributions, and iteratively flip a coin for each one to see whether or not it is the one that we sample from; if none of the distributions are chosen, the default 0 is returned.

More precisely, let  $I \subseteq \mathbb{Z}$ , and for each  $n \in I$ , let  $(S_n, d_n, \mu_n)$  be a complete countable measured metric space. Assume that the following two properties hold.

1. Let  $m, n \in I$  with m < n, and let  $x_1 \neq x_2 \in S_m$  and  $y_1 \neq y_2 \in S_n$ . Then

$$d_m(x_1, x_2) \le 2d_n(y_1, y_2).$$
(3)

2. For every  $n \in I \cap \mathbb{N}$  there exists a special  $s_n \in S_n$  such that

$$\sum_{n \in I \cap \mathbb{N}} 1 - \mu_n\left(\{s_n\}\right) < \infty. \tag{4}$$

We now define a new measured metric space out of these spaces.

1. The space. We say that a vector  $x \in \prod_{n \in I} S_n$  is eventually constant if there exists an N > 0 such that  $x_n = s_n$  for all  $n \ge N$ . Note that when  $\sup I < \infty$ , every vector is eventually constant. We set

$$S = \left\{ x \in \prod_{n \in I} S_n \mid x \text{ is eventually constant} \right\}.$$

2. The metric. For  $x, y \in S$ , let  $n(x, y) = \sup \{n \mid x_n \neq y_n\}$  if  $x \neq y$ , and  $n(x, y) = \infty$  when x = y. When  $x \neq y$ , this is always finite since x and y are eventually constant. We then define

$$d(x,y) = \begin{cases} 0 & x = y \\ d_{n(x,y)} \left( x_{n(x,y)}, y_{n(x,y)} \right) & x \neq y. \end{cases}$$

It is clear that d(x, y) = 0 if and only if x = y, so we only need to verify the triangle inequality. Let  $x, y, z \in S$  be distinct, and denote b = n(x, y).

(a) If either n(y,z) > b or n(x,z) > b, then since x and y agree on indices greater than b, we must have n(x,z) = n(y,z) > b, and also d(x,z) = d(y,z). Then by (3),

$$d(x,y) = d_b(x_b, y_b) \le 2d_{n(x,z)}(x_{n(x,z)}, z_{n(x,z)}) = d(x,z) + d(y,z).$$

- (b) If both n(y,z) = b and n(x,z) = b, then the triangle inequality is satisfied because  $d_b$  is a metric.
- (c) Otherwise, we have that at least one of n(y, z) and n(x, z) is strictly smaller than b. But it is impossible for both to be strictly smaller, since that would mean that  $x_b = z_b = y_b$ , contradicting the fact that  $x_b \neq y_b$ . Supposing w.l.o.g. that n(x, z) = b, we then trivially have

$$f(x, y) = d_b(x_b, y_b) = d_b(x_b, z_b) = d(x, z) \le d(x, z) + d(y, z)$$

3. The measure. Let  $\tilde{\mu} = \bigotimes_{n \in I} \mu_n$  be the product measure on  $\prod_{n \in I} S_n$ . By (4) the sum  $\sum_{n \in I \cap \mathbb{N}} 1 - \mu_n(\{s_n\})$  is finite, and the Borel-Cantelli lemma implies that  $X \sim \tilde{\mu}$  is eventually constant with probability 1 under  $\mu$ . Under the metric *d* defined above, an open ball around  $x \in S$  of radius *r* is given by

$$B(x,r) = \bigcup_{N \in I} \bigcup_{s \in S_N \mid d_N(x_N,s) < r, s \neq x_N} \left\{ y \in S \mid y_N = s, y_n = x_n \text{ for all } n > N \right\} \cup \left\{ x \right\}.$$

Each set of the form  $\{y \in S \mid y_N = s, y_n = x_n \text{ for all } n > N\}$  is measurable under  $\tilde{\mu}$ , and therefore so is their countable union. The measure  $\tilde{\mu}$  can therefore be restricted to a measure  $\mu$  defined on the  $\sigma$ -algebra generated by the open balls in S.

We call the space  $(S, d, \mu)$  the selection space of  $(S_n, d_n, \mu_n)$ . It is not necessarily separable, but in our applications it will be. Let us now find the distribution of d(X, Y) when  $X, Y \sim \mu$  are independent. For r = 0, we have

$$\mathbb{P}\left[d\left(X,Y\right)=0\right] = \prod_{n\in I} \mathbb{P}\left[X_n = Y_n\right] = \prod_{n\in I} \mathbb{P}\left[d_n\left(X_n,Y_n\right)=0\right],\tag{5}$$

while for r > 0,

$$\mathbb{P}\left[d\left(X,Y\right)=r\right] = \sum_{n\in I} \mathbb{P}\left[n\left(X,Y\right)=n\right] \cdot \mathbb{P}\left[d_{n}\left(X_{n},Y_{n}\right)=r \mid n\left(X,Y\right)=n\right]$$
$$= \sum_{n\in I} \left(\prod_{k\in I,k>n} \mathbb{P}\left[X_{k}=Y_{k}\right]\right) \mathbb{P}\left[X_{n}\neq Y_{n}\right] \cdot \mathbb{P}\left[d_{n}\left(X_{n},Y_{n}\right)=r \mid X_{n}\neq Y_{n}\right]$$
$$= \sum_{n\in I} \left(\prod_{k\in I,k>n} \mathbb{P}\left[d\left(X_{k},Y_{k}\right)=0\right]\right) \mathbb{P}\left[d_{n}\left(X_{n},Y_{n}\right)=r\right],$$
(6)

and we have indeed obtained a mixture of positive elements of the distributions achieved by  $(S_n, d_n, \mu_n)$ .

Remark 4. As a space of vectors indexed by  $I \subseteq \mathbb{Z}$ , the selection space can also be interpreted as the steps of a bi-directional walk, where at time-step n, the walker may go in the "direction"  $x_n \in S_n$ . When  $\sup I$  is finite and we view time as starting at  $\sup I$  and directed down towards  $\inf I$ , the index n(x, y) is the first time that two walkers x and y split up when starting at a common origin. In this case the elements of S can be seen as the space of ends on some rooted tree, and the metric d is a generalization of the metric  $e^{-h(x,y)}$ on the space of ends, where h(x, y) is the depth of the last common vertex of the rays x and y. See Figure 1. When  $\sup I = \infty$ , however, there is no such common root.



Figure 1: When  $\sup I = N < \infty$ , the metric d is a generalization of a class of metrics on the space of ends of a tree rooted, where the distance between two rays from the origin depends on the depth of their last common vertex.

**Proposition 5.** Let a > 0, let  $p_0 \in (0, 1)$  and let  $\lambda$  be a discrete probability distribution with supp  $(\lambda) \subseteq [a, 2a]$ . Then the distribution  $\theta = p_0 \delta_0 + (1 - p_0) \lambda$  is feasible.

*Proof.* Let  $a_0 = 0$ , and let  $(a_n)_{n \in I}$  be the positive atoms of  $\lambda$  arranged in some arbitrary order; here,  $I = \mathbb{N}$  if there is an infinite number of atoms and I = [N] for some N otherwise. For every  $n \in I$ , denote

 $p_n = \theta(\{a_n\})$ , and define  $q_n$  by

$$q_n = \frac{p_n}{p_0 + p_1 + \ldots + p_n}$$

For  $n \in I$ , let  $(S_n, d_n, \mu_n)$  be the metric space corresponding to the Bernoulli measure  $\theta_n = (1 - q_n) \delta_0 + q_n \delta_{a_n}$ , obtained by multiplying the metric described in the examples section by  $a_n$ . Then the two assumptions in the construction of the selection space hold:

1. A calculation shows that when setting  $p_0 = 1 - q_n$  in (1), the probability to obtain 0 under  $\mu_n$  is bounded by

$$\mu_n(\{0\}) \ge 1 - q_n. \tag{7}$$

Thus

$$\sum_{n \in I} 1 - \mu_n \left(\{0\}\right) \le \sum_{n \in I} q_n = \sum_{n \in I} \frac{p_n}{p_0 + p_1 + \ldots + p_n} \le \sum_{n \in I} \frac{p_n}{p_0} = \frac{1 - p_0}{p_0},\tag{8}$$

and the sum in (4) is finite.

2. Since  $a_n \in [a, 2a]$ , the requirement (3) is immediate: the left-hand side of (3) is at most 2a, while the right-hand side is at least 2a.

Let  $(S, d, \mu)$  be the selection space constructed from  $(S_n, d_n, \mu_n)$ . Since the positive atoms of the  $\mu_n$ s are all distinct, equations (5) and (6) imply that

$$\mathbb{P}[d(X,Y) = 0] = \prod_{n \in I} \mathbb{P}[d_n(X_n, Y_n) = 0] = \prod_{n \in I} (1 - q_n) = p_0$$

and

$$\mathbb{P}\left[d\left(X,Y\right) = a_{n}\right] = q_{n} \prod_{i \in I, i > n} \left(1 - q_{i}\right) = p_{n}$$

Since S is countable, it is also separable, and since the non-zero distances are bounded from below, it is complete, and the proposition is proven. We remark that by (7), the Weierstrass inequality, and (8),

$$\mathbb{P}\left[X_n = (0, 0, \ldots)\right] = \prod_{n \in I} \left(1 - \mu_n\left(\{0\}\right)\right) \ge \prod_{n \in I} \left(1 - q_n\right) \ge 1 - \sum_{n \in I} q_n \ge 1 - \frac{1 - p_0}{p_0}.$$
(9)

*Proof of Theorem 1.* The measure  $\theta$  can be written as

$$\theta = p_{\infty}\delta_0 + \sum_{n \in I} p_n \lambda_n,$$

where  $I \subseteq \mathbb{Z}$ , and for every  $n \in I$ ,  $p_n > 0$  and the measure  $\lambda_n$  is discrete and supported on  $(2^n, 2^{n+1}]$ . Note that if  $p_{\infty} = 0$ , then necessarily inf  $I = -\infty$ . Define the numbers

$$\beta_n = \frac{p_n}{1 - \sum_{i \in I, i > n} p_i},$$

and let  $(S_n, d_n, \mu_n)$  be the metric space defined in the proof of Proposition 5 corresponding to the measure  $\theta_n = (1 - \beta_n) \delta_0 + \beta_n \lambda_n$ . Then the two assumptions in the construction of the selection space hold for  $(S_n, d_n, \mu_n)$ :

1. If sup  $I < \infty$  then of course the sum in (4) is finite. Otherwise, by (9), if  $X_n \sim \mu_n$ , then

$$\mathbb{P}[X_n = (0, 0, \ldots)] \ge 1 - \frac{1 - (1 - \beta_n)}{1 - \beta_n} = 1 - \frac{\beta_n}{1 - \beta_n}.$$

Thus, since  $\beta_n \to 0$  as  $n \to \infty$ , there exists a constant C > 0 such that

$$\sum_{n \in I \cap \mathbb{N}} \mathbb{P}\left[X_n \neq (0, 0, \ldots)\right] \le \sum_{n \in I \cap \mathbb{N}} \frac{\beta_n}{1 - \beta_n} \le C \sum_{n \in I \cap \mathbb{N}} \beta_n = C \sum_{n \in I \cap \mathbb{N}} \frac{p_n}{1 - \sum_{i \in I, i > n} p_i} \le C \frac{1}{p_\infty + \sum_{i \in I, i < 1} p_i}$$

2. Since the range of  $d_n$  is  $(2^n, 2^{n+1}]$ , the requirement (3) is immediate; in fact, the left hand side of (3) is bounded by just half of the right hand side.

Let  $(S, d, \mu)$  be the selection space constructed from  $(S_n, d_n, \mu_n)$ . Since the supports of  $\lambda_n$  are disjoint, equations (5) and (6) imply that

$$\mathbb{P}\left[d\left(X,Y\right)=0\right] = \prod_{n\in I} \mathbb{P}\left[X_n = Y_n\right] = \prod_{n\in I} \left(1-\beta_n\right) = p_{\infty},$$

while the probability to sample from  $\lambda_n$  is exactly  $\beta_n \prod_{i \in I, i > n} (1 - \beta_i) = p_n$ .

If  $(x^{(k)})_{k\in\mathbb{N}}$  is a Cauchy sequence in S, then it is either constant from some k on, or for every  $N \in I$ , for all k large enough  $x_n^{(k)}$  is constant for  $n \geq N$ , and so  $x^{(k)}$  converges to an element in S, and S is complete.

All that remains is to show that S is separable. If  $\inf I > -\infty$ , then S is a countable union of countable sets, and so is countable itself. Otherwise, for each negative  $n \in I$ , let  $a^{(n)}, b^{(n)}$  be two distinct elements from  $S_n$ , and set

$$A_n = \left\{ x \in S \mid x_i = a^{(i)} \,\forall i \le n \right\} \cup \left\{ x \in S \mid x_i = b^{(i)} \,\forall i \le n \right\}.$$

Since all vectors in S are eventually constant, each  $A_n$  is a countable union of countable sets and is therefore countable, and so the union  $\bigcup_{n \in I} A$  is also countable. It is also dense in S: for any  $x \in S$  and any  $n \in I$ , either  $x_n \neq a^{(n)}$  or  $x_n \neq b^{(n)}$ , and so there exists  $y \in A_n$  such that n(x,y) = n, yielding  $d(x,y) \leq 2^{n+1}$ (recall that n is negative).

### 3 Proof of Theorem 2

Our proof first finds an easy-to-construct feasible "starter" distribution, and then changing its metric in order to obtain the distribution  $\theta$ . The idea is as follows. Let W be a continuous random variable on [0, 1] whose cumulative distribution function  $F_W$  is strictly increasing on [0, 1], and let  $\varphi : [0, 1] \to [0, \infty)$  be strictly increasing. The cumulative distribution function of the random variable  $\varphi(W)$  is given by

$$F_{\varphi(W)}(t) = \mathbb{P}\left[\varphi\left(W\right) \le t\right] = \mathbb{P}\left[W \le \varphi^{-1}\left(t\right)\right] = F_W\left(\varphi^{-1}\left(t\right)\right)$$

Denoting by G the cumulative distribution function of  $\theta$  and choosing  $\varphi(t) = G^{-1}(F_W(t))$  gives us, for  $t \in [0,1]$ ,  $F_W(\varphi^{-1}(t)) = G(t)$ , and so the random variable  $\varphi(W)$  has distribution  $\theta$ . If W is achievable by some  $(S, d, \mu)$ , then when  $X, Y \sim \mu$  are independent, we have  $\varphi(d(X, Y)) \sim \theta$ . However, this does not immediately mean that  $(S, \varphi \circ d, \mu)$  is a measured metric space that achieves  $\theta$ , since there is no guarantee that  $\varphi \circ d$  is still a metric (or that  $\mu$  is still Borel under the topology induced by  $\varphi \circ d$ ). The essence of the proof is to find a suitable W which allows  $\varphi$  to preserve the metric d.

**Definition 6.** A function  $f: [0, \infty) \to [0, \infty)$  is called *metric preserving* if for every metric space (S, d), the function  $f \circ d$  is also a metric on S. It is called *strongly metric preserving* if it is metric preserving and (S, d) is topologically equivalent to  $(S, f \circ d)$ .

Metric preserving functions have been well studied; see [Cor99] for a brief introduction. We will require only the following standard result.

**Theorem 7.** [Kel75, page 131] Let  $f : [0, \infty) \to [0, \infty)$  be continuous, nondecreasing, and satisfying the following two conditions:

- 1.  $f(x) = 0 \iff x = 0$ .
- 2. (Subadditivity)  $f(x+y) \leq f(x) + f(y)$  for all  $x, y \in [0, \infty)$ .

Then f is strongly metric preserving.

Theorem 7 implies that it suffices to find a continuous random variable W such that  $G^{-1}(F_W(t))$  is subadditive. If  $G^{-1}$  is itself already subadditive, then no modification by  $F_W$  is actually needed; in this case, we can take W to be the uniform measure on [0, 1], so that  $F_W(t) = t$  for  $t \in [0, 1]$  and  $G^{-1}(F_W(t)) =$  $G^{-1}(t)$ . Obtaining  $\theta$  is then just an instance of the inverse sampling theorem (as we saw in the examples section, the uniform measure is indeed feasible). However, when  $G^{-1}$  is not subadditive, the idea is that if  $F_W$  itself is in some sense very strongly subadditive, then its subadditivity is enough to counter the non-subadditivity of  $G^{-1}$ . This is made precise in the following simple proposition.

**Proposition 8.** Let  $\Psi : [0,1] \rightarrow [0,1]$  be an absolutely continuous nondecreasing function, and assume that there exist m, M > 0 such that

$$m < \Psi'(t) < M. \tag{10}$$

Let  $f: [0,\infty) \to [0,1]$  be such that for every  $x \leq y$ ,

$$f(x+y) \le \frac{m}{M}f(x) + f(y).$$
(11)

Then  $\Psi \circ f$  is subadditive.

*Proof.* By absolute continuity, for all  $x \leq y$ ,

$$\Psi(f(x+y)) = \Psi(f(y)) + \int_{f(y)}^{f(x+y)} \Psi'(t) dt$$

$$\stackrel{(10)}{\leq} \Psi(f(y)) + M(f(x+y) - f(y))$$

$$\stackrel{(11)}{\leq} \Psi(f(y)) + mf(x)$$

$$\stackrel{(10)}{\leq} \Psi(f(y)) + \Psi(f(x)).$$

The next proposition shows that it is indeed possible to obtain such f.

**Proposition 9.** For every  $\varepsilon \in (0, 1)$ , there exists a feasible continuous random variable W whose cumulative distribution function  $F_W$  satisfies

$$F_W(x+y) \le \varepsilon F_W(x) + F_W(y)$$

for all  $x \leq y$ .

Given the above, Theorem 2 quickly follows.

Proof of Theorem 2. By (2),  $G^{-1}$  is absolutely continuous, strictly increasing, and has upper and lower bounds on its derivative  $(G^{-1})(t)' = \frac{1}{g(G^{-1}(t))}$ . Then by Propositions 8 and 9, there is a feasible W achieved by  $(S, d, \mu)$  such that  $\varphi = G^{-1} \circ F_W$  is subadditive. Since both  $G^{-1}$  and  $F_W$  are continuous, by Theorem 7,  $\varphi$  is strongly metric preserving, and so  $(S, \varphi \circ d, \mu)$  achieves  $\theta$ .

Proof of Proposition 9. Let  $\alpha \in (0, \frac{1}{2})$  to be chosen later, define  $H : \mathbb{R} \to [0, 1]$  by

$$H(t) = \begin{cases} 0 & t < 0\\ t^{\alpha} & 0 \le t \le 1\\ 1 & t > 1, \end{cases}$$

- 1		
- 1		

and let  $\mu$  be the distribution whose cumulative distribution function is H. This distribution has density  $h(t) = \alpha t^{\alpha-1}$  for  $t \in [0,1]$ . Let W = |X - Y|, where  $X, Y \sim \mu$  are i.i.d. By definition, W is feasible. For  $t \in [0,1]$ , the cumulative distribution function  $F_W$  is given by

$$\begin{split} F_W(t) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} h(x) h(y) \, \mathbf{1}_{|x-y| \le t} \, dy dx \\ &= 2 \int_0^1 \int_x^{x+t} h(x) h(y) \, dy dx \\ &= 2 \int_0^1 h(x) \left( H(x+t) - H(x) \right) \, dx \\ &= 2 \int_0^{1-t} \alpha x^{2\alpha - 1} \left( \left( 1 + \frac{t}{x} \right)^{\alpha} - 1 \right) \, dx + 2 \int_{1-t}^1 \alpha x^{\alpha - 1} \left( 1 - x^{\alpha} \right) \, dx. \end{split}$$

The function  $F_W$  can readily be shown to be concave on  $[0,\infty)$  by calculating the second derivative of  $2\int_0^1 \int_x^{x+t} h(x) h(y) dy dx$ . Thus, for all  $z, w \in [0,\infty)$ ,

$$F_W(z) \le F_W(w) + (z - w) F'_W(w).$$

Let  $x, y \in (0, \infty)$  with  $x \leq y$ . Choosing z = x + y and w = y, we get

$$F_{W}(x+y) \leq F_{W}(y) + xF'_{W}(y)$$
  
$$\leq F_{W}(y) + xF'_{W}(x)$$
  
$$= F_{W}(y) + F_{W}(x) \cdot \frac{xF'_{W}(x)}{F_{W}(x)}$$

It therefore suffices to show that for all  $t \in [0, \infty)$ ,

$$\frac{tF_{W}^{\prime}\left(t\right)}{F_{W}\left(t\right)} \leq \varepsilon$$

As  $F_W$  is concave and increasing, it is enough to show a bound of  $\frac{1}{2}\varepsilon$  for all  $t \leq \frac{1}{2}$ , since for all  $t > \frac{1}{2}$ ,  $F'_W(t) \leq F'_W(\frac{1}{2})$  and  $F_W(t) \geq F_W(\frac{1}{2})$ . The denominator is bounded by

$$F_W(t) = 2\alpha \left( \int_0^{1-t} x^{2\alpha-1} \left( \left( 1 + \frac{t}{x} \right)^{\alpha} - 1 \right) dx + \int_{1-t}^1 x^{\alpha-1} (1 - x^{\alpha}) dx \right)$$
$$\geq 2\alpha \int_0^t x^{2\alpha-1} \left( \left( \frac{t}{x} \right)^{\alpha} - 1 \right) dx = 2\alpha \left( \frac{1}{\alpha} t^{2\alpha} - \frac{1}{2\alpha} t^{2\alpha} \right) = t^{2\alpha}.$$

Recalling that  $\alpha < \frac{1}{2}$ , the numerator is bounded by

$$\begin{split} F'_W(t) &= 2\int_0^1 h\left(x\right) h\left(x+t\right) \, dx \\ &= 2\alpha^2 \left(\int_0^t x^{2\alpha-2} \left(1+\frac{t}{x}\right)^{\alpha-1} \, dx + \int_t^{1-t} x^{2\alpha-2} \left(1+\frac{t}{x}\right)^{\alpha-1} \, dx\right) \\ &\leq 2\alpha^2 \left(\int_0^t x^{2\alpha-2} \left(\frac{t}{x}\right)^{\alpha-1} \, dx + \int_t^{1-t} x^{2\alpha-2} \, dx\right) \\ &= 2\alpha t^{2\alpha-1} - 2\alpha^2 \frac{1}{1-2\alpha} \left(\frac{1}{(1-t)^{1-2\alpha}} - \frac{1}{t^{1-2\alpha}}\right) \\ &\leq 2\alpha \frac{1}{t^{1-2\alpha}} \left(1 + \frac{\alpha}{1-2\alpha} \frac{1}{t^{1-2\alpha}}\right). \end{split}$$

Choosing  $\alpha \leq \frac{1}{4}$  then gives  $F'_W(t) \leq 4\alpha t^{2\alpha-1}$ . We thus have

$$\frac{tF'_W(t)}{F_W(t)} \le \frac{4\alpha t^{2\alpha}}{t^{2\alpha}} = 4\alpha,$$

and the proposition follows for  $\alpha \leq \frac{1}{4}\varepsilon$ .

Remark 10. The function  $F_W$  may be expressed in a slightly more closed form using hypergeometric functions; a short calculation reveals that for  $t \in [0, 1]$ ,

$$F_W(t) = 1 + 2(1-t)^{\alpha} (F(-\alpha, 1, 1+\alpha; 1-t) - 1),$$

where  $_{2}F_{1}(a,b;c;z)$  is the hypergeometric function

$${}_{2}F_{1}(a,b;c;z) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_{0}^{1} \frac{x^{b-1} (1-x)^{c-b-1}}{(1-xz)^{a}} dx.$$

A plot of  $F_W$  for some values of  $\alpha$  is given in Figure 2, where the asymptotic behavior of  $t^{2\alpha}$  can be seen near the origin.



Figure 2: The function  $F_W$  for various values of  $\alpha$ .

Remark 11. It might perhaps be possible to extend the proof method of Theorem 2 to distributions on  $[0, \infty)$  where the density is always positive. Indeed, if we compose  $G^{-1}$  with another function f, we still have

$$G^{-1}(f(x+y)) = G^{-1}(f(y)) + \int_{f(y)}^{f(x+y)} \frac{d}{dt} G^{-1}(t) dt$$
  
$$\leq G^{-1}(f(y)) + (f(x+y) - f(y)) \sup_{t \in [f(y), f(x+y)]} \frac{d}{dt} G^{-1}(t)$$

An analogue of Theorem 2 would follow if we could find, for each  $\varepsilon > 0$ , an unbounded feasible random variable W such that for all  $x \le y$ 

$$F_W(x+y) \le F_W(y) + \left(\varepsilon \sup_{t \in [F_W(y), F_W(x+y)]} \frac{d}{dt} G^{-1}(t)\right) F_W(x).$$

Unlike the random variable from Proposition 9, such a W would have to rely directly on G in some way. However, the method cannot be easily extended to distributions whose density is 0 in an interval: this would imply that  $G^{-1} \circ F_W$  is discontinuous, which would prevent it from preserving the metric.

# 4 Acknowledgments

We thank Wendelin Werner and Guillaume Blanc for useful comments.

## References

- [ABC24] David J. Aldous, Guillaume Blanc, and Nicolas Curien. The distance problem on measured metric spaces. 2024.
- [Cor99] Paul Corazza. Introduction to metric-preserving functions. Amer. Math. Monthly, 106(4):309–323, 1999.
- [Kel75] John L. Kelley. General topology, volume No. 27 of Graduate Texts in Mathematics. Springer-Verlag, New York-Berlin, 1975. Reprint of the 1955 edition [Van Nostrand, Toronto, Ont.].