

# Charged rotating black holes in scalar multipolar universes

Cristian Stelea,<sup>1</sup> Marina-Aura Dariescu,<sup>2</sup> Vitalie Lungu,<sup>3</sup>

<sup>1</sup> *Department of Exact and Natural Sciences, Institute of Interdisciplinary Research,  
“Alexandru Ioan Cuza” University of Iasi  
11 Bd. Carol I, Iasi, 700506, Romania*

<sup>2,3</sup> *Faculty of Physics, “Alexandru Ioan Cuza” University of Iasi  
11 Bd. Carol I, Iasi, 700506, Romania*

## Abstract

Recently, Cardoso and Natário [6] constructed an exact solution of Einstein-scalar field equations that describes a scalar counterpart of the Schwarzschild-Melvin Universe. In fact, this solution belongs to a more general class of solutions described by Herdeiro in [7]. In this work we show how to further generalize these solutions in presence of rotation and various charges. More specifically, we describe general charged rotating black holes with NUT charge and present some of their properties.

---

<sup>1</sup>Corresponding author e-mail: [cristian.stelea@uaic.ro](mailto:cristian.stelea@uaic.ro)

<sup>2</sup>E-mail: [marina@uaic.ro](mailto:marina@uaic.ro)

<sup>3</sup>E-mail: [vitalie.lungu@student.uaic.ro](mailto:vitalie.lungu@student.uaic.ro)

# 1 Introduction

Scalar fields belong to the simplest case of matter fields and they have been consistently used in various fields of modern physics, ranging from General Relativity (GR) to high energy particle physics. For example, in cosmology, scalar fields have been used to model dark energy and dark matter [1] (see also [2]). In the high energy particle physics context the first observational evidence of a scalar particle was obtained at the Large Hadron Collider, in the form of the long sought Higgs boson [3], [4]. Scalar fields also occur naturally in theories involving Kaluza-Klein dimensional reductions, such as string theory. Moreover, in the context of black hole physics in GR the role of scalar fields and their consequences on the no-scalar-hair theorems have been addressed in [5].

Quite recently, in [6] the authors found a new solution of the Einstein-scalar field equations that describes a black hole immersed into a scalar Universe for which the scalar field is asymptotically aligned with the  $z$ -axis. This solution has been called the scalar Schwarzschild-Melvin solution, by analogy with the well-known Schwarzschild-Melvin solution in the Einstein-Maxwell theory, for which the magnetic field is aligned asymptotically with the  $z$ -axis. Later, in [7] Herdeiro showed that the scalar Schwarzschild-Melvin solution is a particular case of a class of solutions of the Einstein-scalar field equations that describes black holes in multipolar scalar Universes. This class of solutions belongs to the well-known Weyl-Papapetrou metrics. The background on which the black holes reside is determined entirely by a massless and minimally coupled scalar field. Quite generically, black hole solutions on this scalar background contain naked curvature singularities that reside either on the black hole horizon or in the asymptotic regions. This is to be expected since for a massless and minimally coupled scalar field there are no asymptotically flat regular solutions with scalar fields [5]. However, in this class of solutions there do exist particular profiles of the scalar field that correspond to scalar black hole solutions which are regular on the event horizon and in the near horizon region, which means that the naked curvature singularities are in the asymptotic regions. These black hole solutions could still be physically interesting if one considers an appropriate cutoff to avoid the asymptotic regions.

The purpose of this work is to further generalize the Herdeiro solutions in presence of rotation and also electromagnetic and NUT charges. More precisely, in the next section one shows how to simply obtain such a solution in presence of a scalar field. Our approach is based on the symmetries of the dimensionally reduced theory down to three dimensions. Rotating black holes of this kind could also be constructed using the methods described in [8], however, in our approach one could easily include the effect of electric and magnetic charges as well.

The structure of this paper is as follows: in the next section one describes the solution generating technique. In section 3 one presents the scalar generalization of the Kerr-Newman black hole with NUT charge in scalar multipolar Universes. We focus on a special solution describing the rotating generalization of the Cardoso-Naváro solution and present some of its properties. The final section is dedicated to conclusions and avenues for further work.

## 2 The solution generating technique

As it is well-known by now, Einstein's field equations form a highly nonlinear system of partial differential equations. Solving them in full generality is a daunting task and in order to find new physically meaningful exact solutions one has to rely on various solution generating techniques. Usually, these solutions generating techniques are based on some kind of simplifying symmetries imposed in the ansatz. Of particular importance is the so-called Weyl-Papapetrou ansatz for the geometry, which is considered to be static and axisymmetric:

$$ds^2 = -e^{-2\psi} dt^2 + e^{2\psi} \left[ e^{2\gamma} (d\rho^2 + dz^2) + \rho^2 d\varphi^2 \right], \quad (1)$$

where the functions  $\psi$  and  $\gamma$  depend on the coordinates  $\rho$  and  $z$ . Performing now a dimensional reduction on the time direction one arrives at the three-geometry of the kind:

$$ds_3^2 = e^{2\gamma} (d\rho^2 + dz^2) + \rho^2 d\varphi^2. \quad (2)$$

The interesting thing about this geometry is that its Ricci tensor is linear in the function  $\gamma$  [9], since:

$$\begin{aligned} R_{\rho\rho}[\gamma] &= -(\partial_z^2 \gamma + \partial_\rho^2 \gamma) + \frac{1}{\rho} \partial_\rho \gamma, \\ R_{\rho z}[\gamma] &= \frac{1}{\rho} \partial_z \gamma, \\ R_{zz}[\gamma] &= -(\partial_z^2 \gamma + \partial_\rho^2 \gamma) - \frac{1}{\rho} \partial_\rho \gamma. \end{aligned} \quad (3)$$

One can see now that if one writes down the dimensionally reduced Einstein equations in the form  $R_{ij} = T_{ij} - Tg_{ij}$ , then for each source  $T_{ij} - Tg_{ij}$  one can associate a function  $\gamma$  in three dimensions, as long as the matter Lagrangeans of each source are decoupled.

To better understand this situation, let us consider first a static and axisymmetric vacuum solution of Einstein equations as in (1). After performing a dimensional reduction along the time direction one obtains the effective Lagrangian<sup>1</sup>:

$$\mathcal{L}_3 = \sqrt{g_3} [R - 2(\partial_i \psi)(\partial^i \psi)]. \quad (4)$$

Then Einstein's field equations can be written as  $R_{ij}[\gamma] = 2(\partial_i \psi)(\partial_j \psi)$  and they lead to the following equations for the  $\gamma$  function:

$$\begin{aligned} \partial_\rho \gamma &= \rho [(\partial_\rho \psi)^2 - (\partial_z \psi)^2], \\ \partial_z \gamma &= 2\rho (\partial_\rho \psi)(\partial_z \psi). \end{aligned} \quad (5)$$

On the other hand, the equation of motion for the scalar field  $\psi$  as derived from the Lagrangean (4) leads to:

$$\Delta \psi = e^{2\gamma} \left[ \partial_\rho^2 \psi + \partial_z^2 \psi + \frac{1}{\rho} \partial_\rho \psi \right] = 0. \quad (6)$$

---

<sup>1</sup>Here  $g_3 = e^{4\gamma} \rho^2$  is the metric determinant of the three geometry, which has Euclidean signature.

Since  $e^{2\gamma} \neq 0$  this means that the scalar  $\psi$  can be considered as a solution of the three-dimensional Laplace equation on an Euclidean geometry corresponding to  $\gamma = 0$  in (2) [9]. Therefore Einstein's equations in four dimensions for a static axisymmetric background are now essentially reduced to finding a solution of Laplace's equation on flat space.

Consider now a particular background of the form:

$$ds^2 = -dt^2 + e^{2\mu}[d\rho^2 + dz^2] + \rho^2 d\varphi^2, \quad (7)$$

solution of the Einstein-scalar field equations derived from the Lagrangean:

$$\mathcal{L}_4 = \sqrt{-g}[R - 2(\partial_\mu\phi)(\partial^\mu\phi)]. \quad (8)$$

After performing a dimensional reduction along the time direction one obtains the three-geometry of the form:

$$ds_3^2 = e^{2\mu}(d\rho^2 + dz^2) + \rho^2 d\varphi^2. \quad (9)$$

with the effective Lagrangian:

$$\mathcal{L}_3 = \sqrt{g_3}[R - 2(\partial_i\phi)(\partial^i\phi)]. \quad (10)$$

Then Einstein's field equations can be written as  $R_{ij}[\mu] = 2(\partial_i\phi)(\partial_j\phi)$  and they lead to the same functional equations for the  $\mu$  function as in (5):

$$\begin{aligned} \partial_\rho\mu &= \rho[(\partial_\rho\phi)^2 - (\partial_z\phi)^2], \\ \partial_z\mu &= 2\rho(\partial_\rho\phi)(\partial_z\phi). \end{aligned} \quad (11)$$

The equation of motion for the scalar field  $\phi$  shows that it is a harmonic function in the auxiliary flat background (9) (with  $\mu = 0$ ). The general solution for this equation can be expressed as [7] -[23]:

$$\phi = \frac{a_0}{R} + \sum_{l=1}^{\infty} \left( \frac{a_l}{R^{l+1}} + b_l R^l \right) P_l, \quad (12)$$

where  $R = \sqrt{\rho^2 + z^2}$  and  $P_l$  is the  $l$ -Legendre polynomial with argument  $\frac{z}{R}$ . Here  $a_0$ ,  $a_l$  and  $b_l$  are constants.

One can also integrate (11) to find the function  $\mu$  [7] -[23]:

$$\mu = \sum_{l,p=1}^{\infty} \left[ \frac{(l+1)(p+1)a_l a_p}{(l+p+2)R^{l+p+2}} (P_{l+1}P_{p+1} - P_l P_p) + \frac{l p b_l b_p R^{l+p}}{l+p} (P_l P_p - P_{l-1} P_{p-1}) \right]. \quad (13)$$

One can now combine the two solutions as follows: consider a four-dimensional geometry sourced by an Einstein-scalar Lagrangean of the form (8). If one assumes the geometry to belong to the Weyl-Papapetrou form:

$$ds^2 = -e^{-2\psi} dt^2 + e^{2\psi} \left[ e^{2\lambda}(d\rho^2 + dz^2) + \rho^2 d\varphi^2 \right], \quad (14)$$

then, after performing a dimensional reduction along the time direction one obtains a three-geometry of the form:

$$ds_3^2 = e^{2\lambda}(d\rho^2 + dz^2) + \rho^2 d\varphi^2, \quad (15)$$

sourced by a three-dimensional Lagrangean:

$$\mathcal{L}_3 = \sqrt{g_3} [R - 2(\partial_i \psi)(\partial^i \psi) - 2(\partial_i \phi)(\partial^i \phi)]. \quad (16)$$

The Einstein's field equations can now be written as:

$$R_{ij}[\lambda] = 2(\partial_i \psi)(\partial_j \psi) + 2(\partial_i \phi)(\partial_j \phi) = R_{ij}[\gamma] + R_{ij}[\mu] \quad (17)$$

and therefore, since the expression of the Ricci tensor in (3) is linear in the function  $\lambda$ , one can simply take  $\lambda = \gamma + \mu$  to satisfy the Einstein equations in three dimensions. The remaining equations of motion derived from the Lagrangean (16) show that the functions  $\psi$  and  $\phi$  are harmonic functions  $\Delta\psi = \Delta\phi = 0$ , while the functions  $\gamma$  and  $\mu$  can be found by integrating (5) and (11) (which are now equivalent to Einstein's equations (17)). In conclusion, for every static and axisymmetric vacuum solution of the form (1) one can find a scalar modification in the form:

$$ds^2 = -e^{-2\psi} dt^2 + e^{2\psi} \left[ e^{2\gamma+2\mu}(d\rho^2 + dz^2) + \rho^2 d\varphi^2 \right]. \quad (18)$$

This is the class of solutions considered by Herdeiro in [7]. In the followings we shall show that it is easy to generalize this construction in the presence of rotation and various other charges.

Consider first a charged and rotating generalization of the Weyl-Papapetrou ansatz of the form:

$$ds^2 = -e^{2\psi}(dt + \mathcal{A})^2 + e^{2\psi} \left[ e^{2\gamma}(d\rho^2 + dz^2) + \rho^2 d\varphi^2 \right], \quad (19)$$

where  $\mathcal{A} = \mathcal{A}_\varphi d\varphi$ . Let us assume that, in four dimensions, this geometry is a solution of the field equations derived from the Lagrangean:

$$\mathcal{L}_4 = \sqrt{-g} [R - F^2], \quad (20)$$

where one considers the Maxwell field  $F = dA$ , with  $A = \chi dt + A_\varphi d\varphi$ . Performing now a dimensional reduction on the time direction one obtains the three-dimensional Lagrangean:

$$\mathcal{L}_3 = \sqrt{g_3} \left[ R_3 - 2(\partial\psi)^2 + e^{-4\psi}(\mathcal{F})^2 - e^{-2\psi}F_{(2)}^2 + 2e^{2\psi}(\partial\chi)^2 \right], \quad (21)$$

while the three-geometry is given by (2). Here we defined  $F_{(2)} = dA_\varphi \wedge d\varphi - d\chi \wedge \mathcal{A}$  and  $\mathcal{F} = d\mathcal{A}$ . One can write the three-dimensional Einstein equations in the following form:

$$\begin{aligned} R_{ij}[\gamma] = T_{ij}^{(1)} &= 2(\partial_i \psi)(\partial_j \psi) - 2e^{-4\psi} \left[ \mathcal{F}_{ik} \mathcal{F}_j{}^k - \frac{1}{2} g_{ij} \mathcal{F}_{kl} \mathcal{F}^{kl} \right] - 2e^{2\psi} (\partial_i \chi)(\partial_j \chi) + \\ &+ 2e^{-2\psi} \left[ F_{(2)ik} F_{(2)j}{}^k - \frac{1}{2} g_{ij} F_{(2)kl} F_{(2)}^{kl} \right]. \end{aligned} \quad (22)$$

The remaining field equations in three dimensions are the field equations for the matter fields, as derived from the Lagrangean (21):

$$\begin{aligned}\nabla_i (e^{-4\psi} \mathcal{F}^{ij}) &= 0, & \nabla_i (e^{-2\psi} F_{(2)}^{ij}) &= 0, & \nabla_i (e^{2\psi} \nabla^i \chi) &= 0, \\ \Delta\psi &= -\frac{1}{2}e^{-2\psi} F_{(2)}^2 + e^{-4\psi} \mathcal{F}^2 + e^{2\psi} (\partial\chi)^2.\end{aligned}\quad (23)$$

Together, the system of equations given in (22) and (23) is equivalent to the system of equations derived from the Einstein-Maxwell Lagrangean (20).

Consider now a four-dimensional geometry that should be a solution of the Einstein-Maxwell-scalar system described by the Lagrangean:

$$\mathcal{L}_4 = \sqrt{-g} [R - 2(\partial_\mu \phi)(\partial^\mu \phi) - F^2]. \quad (24)$$

In general, one can write it in the Weyl-Papapetrou form:

$$ds^2 = -e^{2\psi} (dt + \mathcal{A})^2 + e^{2\psi} \left[ e^{2\lambda} (d\rho^2 + dz^2) + \rho^2 d\varphi^2 \right]. \quad (25)$$

Performing now a dimensional reduction on the timelike direction one is lead to the three-dimensional geometry of the form (15), which should be a solution of the field equations derived from the dimensionally reduced Lagrangean:

$$\mathcal{L}_3 = \sqrt{g_3} \left[ R_3 - 2(\partial\psi)^2 + e^{-4\psi} (\mathcal{F})^2 - e^{-2\psi} F_{(2)}^2 + 2e^{2\psi} (\partial\chi)^2 - 2(\partial_\mu \phi)(\partial^\mu \phi) \right]. \quad (26)$$

In this case Einstein's field equations can be cast in the form:

$$R_{ij}[\lambda] = T_{ij}^{(1)} + T_{ij}^{(2)}, \quad (27)$$

where  $T_{ij}^{(1)} = R_{ij}[\gamma]$  is listed in (22) while  $T_{ij}^{(2)} = 2(\partial_i \phi)(\partial_j \phi) = R_{ij}[\mu]$ . Thence one obtains  $R_{ij}[\lambda] = R_{ij}[\gamma] + R_{ij}[\mu]$  and therefore one can simply take  $\lambda = \gamma + \mu$  to simply satisfy Einstein's field equations in three dimensions.

In order to check that the remaining field equations for the various matter fields in (26) are satisfied, one should note that the scalar Lagrangean term for the scalar field  $\phi$  is decoupled from the rest of the matter fields and it only adds the field equation  $\Delta\phi = 0$  to them, while all the remaining field equations are the same as in (23) and they are satisfied as well if the initial solution (19) satisfies them.

In conclusion, starting with a known solution (19) of the Einstein-Maxwell equations then the new solution of the field equations derived from the Lagrangean (24) can be written in the form:

$$ds^2 = -e^{2\psi} (dt + \mathcal{A})^2 + e^{2\psi} \left[ e^{2\gamma+2\mu} (d\rho^2 + dz^2) + \rho^2 d\varphi^2 \right], \quad (28)$$

where the scalar field takes the form (12), while the function  $\mu$  is given by (13).

### 3 The Kerr-Newman-NUT solution with a scalar field

As an example of the solution generating technique described in the previous section, let us consider the Kerr-Newman-NUT solution in four dimensions (see for instance [10], [11])<sup>2</sup>:

$$ds^2 = -\frac{\Delta_r}{\Sigma} \left[ dt - (a \sin^2 \theta + 2l(a - \cos \theta)) d\varphi \right]^2 + \Sigma \left( \frac{dr^2}{\Delta_r} + d\theta \right)^2 + \frac{\sin^2 \theta}{\Sigma} [adt - (r^2 + (a + l)^2) d\varphi]^2, \quad (29)$$

where we defined:

$$\Delta_r = r^2 - 2mr + a^2 + e^2 + g^2 - l^2, \quad \Sigma = r^2 + (l + a \cos \theta)^2. \quad (30)$$

Here  $m$  is the mass parameter,  $a$  is the rotation parameter,  $l$  is the NUT charge, while  $e$  and  $g$  are the electric and magnetic charges. The components of the electromagnetic Maxwell field  $A = \chi dt + A_\varphi d\varphi$  are:

$$\chi = -\frac{er + g(l + a \cos \theta)}{\Sigma},$$

$$A_\varphi = \frac{er[(a + l)^2 - (l + a \cos \theta)^2] + g(1 + a \cos \theta)[r^2 + (a + l)^2]}{a\Sigma}. \quad (31)$$

As it is well-known, this geometry can be brought into the Weyl-Papapetrou form (19) by using the following coordinate transformations:

$$\rho = \sqrt{\Delta_r} \sin \theta, \quad z = (r - m) \cos \theta, \quad (32)$$

such that:

$$d\rho^2 + dz^2 = (\Delta_r \cos^2 \theta + (r - m) \sin^2 \theta) \left( \frac{dr^2}{\Delta_r} + d\theta^2 \right), \quad (33)$$

while:

$$e^{2\gamma} = \frac{\Delta_r - a^2 \sin^2 \theta}{\Delta_r \cos^2 \theta + (r - m) \sin^2 \theta}. \quad (34)$$

Then the general Kerr-Newman-NUT solution in a multipolar scalar universe will be given by:

$$ds^2 = -\frac{\Delta_r}{\Sigma} \left[ dt - (a \sin^2 \theta + 2l(a - \cos \theta)) d\varphi \right]^2 + \Sigma e^{2\mu} \left( \frac{dr^2}{\Delta_r} + d\theta \right)^2 + \frac{\sin^2 \theta}{\Sigma} [adt - (r^2 + (a + l)^2) d\varphi]^2, \quad (35)$$

---

<sup>2</sup>Note that we could also consider the most general solution with the acceleration parameter, however, for simplicity we set it to zero here.

where the Maxwell field components are given in (31), the scalar field  $\phi$  has the general expression (12), while the function  $\mu$  is given by (13), using (32).

Note that the horizon location in our new solution is again determined by the roots of  $\Delta_r = 0$ , while the ergosphere has the same properties as in the Kerr solution. The Ricci scalar for the geometry (35) is given by  $R = 2(\partial_\mu\phi)(\partial^\mu\phi)$  and, therefore, the curvature singularities in the metric are directly related to the divergencies of the scalar field. Moreover, the effects of the scalar field in the metric are encoded using the function  $e^{2\mu}$  and it will generically lead to distortions of the black hole horizon. For simplicity, let us set the NUT charge to zero  $l = 0$ . Then the metric induced on the black hole horizon  $r = r_H$  is:

$$ds_H^2 = \Sigma_H e^{2\mu_H} d\theta^2 + \frac{\Psi_H \sin^2 \theta}{\Sigma_H} d\varphi^2, \quad (36)$$

where we defined  $\Psi_H = (r_H^2 + a^2)^2$ ,  $\Sigma_H = r_H^2 + a^2 \cos^2 \theta$ , while  $\mu_H$  is the value of the function  $\mu$  on the black hole horizon. Then the area of the black hole horizon can be written as:

$$Area = 2\pi(r_H^2 + a^2) \int_0^\pi e^{\mu_H} \sin \theta d\theta. \quad (37)$$

As a particular example, let us consider the scalar profile used in [6] to construct the scalar-Schwarzschild-Melvin solution. In this case:

$$\phi = kz = k(r - m) \cos \theta, \quad e^{2\mu} = e^{-k^2 \rho^2} = e^{-k^2 \Delta_r \sin^2 \theta}. \quad (38)$$

If one sets to zero all the charges one obtains the rotating version of the scalar-Schwarzschild-Melvin [6] in the form:

$$ds^2 = -\frac{\Delta_r}{\Sigma} \left[ dt - a \sin^2 \theta d\varphi \right]^2 + \Sigma e^{2\mu} \left( \frac{dr^2}{\Delta_r} + d\theta \right)^2 + \frac{\sin^2 \theta}{\Sigma} \left[ a dt - (r^2 + a^2) d\varphi \right]^2, \quad (39)$$

where  $\Delta_r = r^2 - 2mr + a^2$  and  $\Sigma = r^2 + a^2 \cos^2 \theta$ . It is easy to check that it satisfies the Einstein-scalar equations  $R_{\mu\nu} = 2(\partial_\mu\phi)(\partial_\nu\phi)$ . If  $a = 0$  it reduces to the solution in [6]. Its Ricci scalar can be simply expressed as:

$$R = \frac{2k^2 e^{k^2 \Delta_r \sin^2 \theta} (\Delta_r + (m^2 - a^2) \sin^2 \theta)}{r^2 + a^2 \cos^2 \theta}. \quad (40)$$

Besides the usual curvature singularities presented in the Kerr metric one can see that the backreaction of the scalar field leads to another curvature singularity located in the asymptotic region as  $r \rightarrow \infty$ . Note that in this case the area of the black hole horizon is the same as the area computed for the Kerr black hole since  $\mu_H = 0$  on the black hole horizon.

## 4 Conclusions

Recently, Cardoso and Natário [6] constructed an exact solution of Einstein-scalar field equations that describes a scalar counterpart of the Schwarzschild-Melvin Universe. In fact, this

solution belongs to a more general class of solutions described by Herdeiro in [7]. These solutions describe black holes in the so-called scalar multipolar Universes, which are determined in terms of the multipolar terms appearing in the scalar field expansion (12). Generically, these black hole solutions present either singularities on the black hole horizon, or singularities in the asymptotic regions, rendering them non-asymptotically flat. However, for special profiles of the scalar fields the singularities can be pushed out to the asymptotic region, keeping the black hole horizon and the regions near it nonsingular.

In our work we further generalized this class of exact solutions of the Einstein-scalar system to include rotation and various charges. More specifically, in section 2 we presented our solution generating technique that allowed us to combine any stationary exact solution of the Einstein-Maxwell system with axial symmetry with the backreaction of a minimally coupled scalar field. Our technique is based on the symmetries of the dimensionally reduced Lagrangeans down to three dimensions along the timelike direction and it relies on the properties of the Weyl-Papapetrou ansatz for the metric.

As an example we derived the general Kerr-Newman-NUT solution in the scalar multipolar universes. Setting all the charges to zero and for a specific profile of the scalar field we showed how one can recover the rotating version of the Cardoso and Natário solution [6]. One should note that our solution can be further generalized by introducing an acceleration parameter in the original metric [10], [11]. We chose not to include this parameter to keep the final solution listed here in a simpler form.

One interesting application of our technique is to use the effect of the scalar field to balance multi-black hole systems. In absence of the scalar field such systems are not in equilibrium and they generically present various conical defects in order to accommodate the presence of other forces needed to balance the attractive gravity effects. For example, in four and higher dimensions the equilibrium conditions in presence of Maxwell fields have been considered in [12] - [19] (and references therein). These multi-black hole systems are usually constructed in Weyl-Papapetrou coordinates and therefore they can be used as seeds in our solution generating technique. By adding carefully chosen scalar profiles one can achieve equilibrium as in [22], [23]. The use of scalar fields to locally balance systems of multi-black holes is not new, as they have been used for instance in [20], [21] in order to achieve this goal.

Another straightforward generalization of our results is to obtain the corresponding solutions in five dimensions, such as the Kerr-Myers solution with scalar fields, or to study the effect of the scalar fields in balancing the static black rings in five dimensions.

Work on these matters is in progress and it will be presented elsewhere.

*Note added* After the completion of this work we learned about the paper [24] that presents similar rotating solutions in four and five dimensions for specific scalar profiles.

## References

- [1] I. Quiros, Int. J. Mod. Phys. D **28**, no.07, 1930012 (2019) doi:10.1142/S021827181930012X [arXiv:1901.08690 [gr-qc]].

- [2] V. Faraoni, doi:10.1007/978-1-4020-1989-0
- [3] G. Aad *et al.* [ATLAS], Phys. Lett. B **716**, 1-29 (2012) doi:10.1016/j.physletb.2012.08.020 [arXiv:1207.7214 [hep-ex]].
- [4] S. Chatrchyan *et al.* [CMS], Phys. Lett. B **716**, 30-61 (2012) doi:10.1016/j.physletb.2012.08.021 [arXiv:1207.7235 [hep-ex]].
- [5] C. A. R. Herdeiro and E. Radu, Int. J. Mod. Phys. D **24**, no.09, 1542014 (2015) doi:10.1142/S0218271815420146 [arXiv:1504.08209 [gr-qc]].
- [6] V. Cardoso and J. Natário, [arXiv:2410.02851 [gr-qc]].
- [7] C. A. R. Herdeiro, [arXiv:2410.12950 [gr-qc]].
- [8] M. Astorino, Phys. Rev. D **91**, 064066 (2015) doi:10.1103/PhysRevD.91.064066 [arXiv:1412.3539 [gr-qc]].
- [9] B. Chng, R. B. Mann and C. Stelea, Phys. Rev. D **74**, 084031 (2006) doi:10.1103/PhysRevD.74.084031 [arXiv:gr-qc/0608092 [gr-qc]].
- [10] M. Astorino, Phys. Rev. D **110**, no.10, 104054 (2024) doi:10.1103/PhysRevD.110.104054 [arXiv:2404.06551 [gr-qc]].
- [11] H. Ovcharenko, J. Podolsky and M. Astorino, [arXiv:2409.02308 [gr-qc]].
- [12] B. Chng, R. B. Mann, E. Radu and C. Stelea, JHEP **12**, 009 (2008) doi:10.1088/1126-6708/2008/12/009 [arXiv:0809.0154 [hep-th]].
- [13] C. Stelea, K. Schleich and D. Witt, Phys. Rev. D **83**, 084037 (2011) doi:10.1103/PhysRevD.83.084037 [arXiv:0909.3835 [hep-th]].
- [14] C. Stelea, C. Dariescu and M. A. Dariescu, Phys. Rev. D **84**, 044009 (2011) doi:10.1103/PhysRevD.84.044009 [arXiv:1107.3484 [gr-qc]].
- [15] C. Stelea, K. Schleich and D. Witt, Phys. Rev. D **91**, 024040 (2015) doi:10.1103/PhysRevD.91.024040 [arXiv:1108.5145 [gr-qc]].
- [16] C. Stelea and M. C. Ghilea, Phys. Lett. B **719**, 191-195 (2013) doi:10.1016/j.physletb.2013.01.009 [arXiv:1211.3725 [gr-qc]].
- [17] J. Kunz, P. G. Nedkova and C. Stelea, Nucl. Phys. B **874**, 773-791 (2013) doi:10.1016/j.nuclphysb.2013.06.013 [arXiv:1304.7020 [gr-qc]].
- [18] C. Stelea, C. Dariescu and M. A. Dariescu, Phys. Rev. D **87**, no.2, 024039 (2013) doi:10.1103/PhysRevD.87.024039 [arXiv:1211.3154 [gr-qc]].
- [19] C. Stelea, Phys. Rev. D **97**, no.2, 024044 (2018) doi:10.1103/PhysRevD.97.024044 [arXiv:1709.10309 [gr-qc]].

- [20] C. A. R. Herdeiro and E. Radu, Phys. Rev. Lett. **131**, no.12, 121401 (2023)  
doi:10.1103/PhysRevLett.131.121401 [arXiv:2305.15467 [gr-qc]].
- [21] C. A. R. Herdeiro and E. Radu, Phys. Rev. D **107**, no.6, 064044 (2023)  
doi:10.1103/PhysRevD.107.064044 [arXiv:2302.00016 [gr-qc]].
- [22] A. Viganò, [arXiv:2211.00436 [gr-qc]].
- [23] M. Astorino and A. Viganò, Eur. Phys. J. C **82**, no.9, 829 (2022)  
doi:10.1140/epjc/s10052-022-10787-y [arXiv:2105.02894 [gr-qc]].
- [24] J. Barrientos, C. Charmousis, A. Cisterna and M. Hassaine, [arXiv:2501.10223 [gr-qc]].