AVERAGING PRINCIPLES AND CENTRAL LIMIT THEOREMS FOR MULTISCALE MCKEAN-VLASOV STOCHASTIC SYSTEMS

JIE XIANG AND HUIJIE QIAO

Department of Mathematics, Southeast University, Nanjing, Jiangsu 211189, P.R.China

ABSTRACT. In this paper, we study a class of multiscale McKean-Vlasov stochastic systems where the entire system depends on the distribution of the fast component. First of all, by the Poisson equation method we prove that the slow component converges to the solution of the averaging equation in the L^p $(p \ge 2)$ space with the optimal convergence rate $\frac{1}{2}$. Then a central limit theorem is established by tightness.

1. Introduction

McKean-Vlasov stochastic differential equations (SDEs for short), also known as distribution-dependent or mean-field SDEs, are characterized by the dependence of their coefficients on the probability law of the solutions. The study of McKean-Vlasov SDEs dates back to the foundational work of McKean [19], which was inspired by Kac's program in Kinetic Theory. Over the years, extensive research on McKean-Vlasov SDEs has yielded significant results. For example, Ding and Qiao [5, 6] explored the well-posedness and stability of solutions for McKean-Vlasov SDEs with non-Lipschitz coefficients. Wang [29] established the exponential ergodicity of strong solutions for Landau-type McKean-Vlasov SDEs. For a broader and more detailed overview, references such as [1, 12, 13, 25] and the works cited therein provide valuable insights.

The averaging principle, first introduced by Khasminskii in the seminal work [15], has proven to be a highly effective and essential tool in the analysis of SDEs for modeling problems across various practical research fields. The core concept of the stochastic averaging principle is to simplify the study of complex stochastic equations by examining corresponding averaging stochastic equations, thereby offering a convenient and efficient approach for investigating many significant properties. In recent years, the stochastic averaging principle has been extended to encompass a broader range of SDEs, as demonstrated in works such as [7, 20, 26, 27].

In recent years, the averaging principle for multiscale McKean-Vlasov SDEs has attracted increasing attention ([11, 17, 21, 24, 33]). Let us mention some results related with our work. When the coefficients are independent of the distribution of the fast component, Röckner, Sun and Xie [24] achieved the optimal convergence order of $\frac{1}{2}$ in the L^2 space by leveraging the Poisson equation. Later, Qiao and Wei [21] considered a type of

AMS Subject Classification (2020): 60H10.

Keywords: Multiscale McKean-Vlasov stochastic systems, averaging principles, the Poisson equation, central limit theorems.

This work was supported by NSF of China (No.12071071) and the Jiangsu Provincial Scientific Research Center of Applied Mathematics (No. BK20233002).

Corresponding author: Huijie Qiao, hjqiaogean@seu.edu.cn.

multiscale McKean-Vlasov systems where the entire system depend on the distribution of the fast component and established the strong convergence rate $\frac{1}{4}$ in the L^2 space through a modified Khasminskii time discretization approach. Note that in [21] the strong convergence rate $\frac{1}{4}$ is not optimal. Thus, in this paper our first aim is to improve this strong convergence rate $\frac{1}{4}$ to the rate $\frac{1}{2}$. Concretely speaking, consider the following slow-fast system on $\mathbb{R}^n \times \mathbb{R}^m$:

$$\begin{cases}
dX_{t}^{\varepsilon} = b_{1}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}, Y_{t}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{t}^{\varepsilon, \xi}}) dt + \sigma_{1}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}) dB_{t}, \\
X_{0}^{\varepsilon} = \varrho, \quad 0 \leqslant t \leqslant T, \\
dY_{t}^{\varepsilon, \xi} = \frac{1}{\varepsilon} b_{2}(Y_{t}^{\varepsilon, \xi}, \mathcal{L}_{Y_{t}^{\varepsilon, \xi}}) dt + \frac{1}{\sqrt{\varepsilon}} \sigma_{2}(Y_{t}^{\varepsilon, \xi}, \mathcal{L}_{Y_{t}^{\varepsilon, \xi}}) dW_{t}, \\
Y_{0}^{\varepsilon, \xi} = \xi, \quad 0 \leqslant t \leqslant T, \\
dY_{t}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}} = \frac{1}{\varepsilon} b_{2}(Y_{t}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{t}^{\varepsilon, \xi}}) dt + \frac{1}{\sqrt{\varepsilon}} \sigma_{2}(Y_{t}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{t}^{\varepsilon, \xi}}) dW_{t}, \\
Y_{0}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}} = y_{0}, \quad 0 \leqslant t \leqslant T,
\end{cases} \tag{1}$$

where (B_t) , (W_t) are *n*-dimensional and *m*-dimensional standard Brownian motions, respectively, defined on the complete filtered probability space $(\Omega, \mathscr{F}, \{\mathscr{F}_t\}_{t\in[0,T]}, \mathbb{P})$ and are mutually independent. Moreover, these mappings $b_1: \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m) \to \mathbb{R}^n$, $\sigma_1: \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \to \mathbb{R}^{n \times n}$, $b_2: \mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m) \to \mathbb{R}^m$, $\sigma_2: \mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m) \to \mathbb{R}^{m \times m}$ are all Borel measurable, and ϱ, ξ are two random variables. We employ the method of the Poisson equation to demonstrate that for some $p \geqslant 2$

$$\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|X_t^{\varepsilon}-\bar{X}_t|^p\right)\leqslant C_T\varepsilon^{\frac{p}{2}},$$

where X^{ε} is the solution for the slow part of the system (1) and \bar{X} is the solution of the averaging equation (See Eq.(3)). Note that in [17], for a general multiscale McKean-Vlasov stochastic system depending on the distribution of the fast component, Li and Xie obtained that (See [17, Theorem 2.1])

$$\sup_{0 \le t \le T} \mathbb{E}|X_t^{\varepsilon} - \bar{X}_t|^p \leqslant C_T \varepsilon^p.$$

It is obvious that our result can not be covered by [17, Theorem 2.1]. Moreover, our result is stronger in some sense.

Next, it is natural to investigate whether $\frac{X^{\varepsilon}-\bar{X}}{\sqrt{\varepsilon}}$ converges and if it does, what is the limit. This type problem is usually solved by the central limit theorem. The study of the central limit theorem in the context of McKean-Vlasov SDEs has garnered significant attention in recent years ([8, 9, 28]). For a multiscale McKean-Vlasov stochastic system independent of the distribution of the fast component, Hong et al. [10] proved a central limit type theorem by the Poisson equation technique. In this paper, we follow the same line as that in [10] and establish a central limit theorem for the system (1). We mention that in [17] Li and Xie also observed the central limit theorem for a slow-fast McKean-Vlasov stochastic system, where the entire system depend on the distributions of both the slow motion and the fast component. Comparing our result with [17, Theorem 2.3], we find that our conditions are more general.

The novelty of this paper is reflected in two aspects. Firstly, we achieve the L^p convergence with the optimal strong convergence order of $\frac{1}{2}$. Secondly, we establish a central limit theorem for multiscale McKean-Vlasov SDEs where the entire system depends on the distribution of the fast component. It is worth stressing that due to the application of

Poisson equation techniques, it is necessary to estimate the moments of both the fast and slow processes to a high degree. Additionally, a time discretization technique is utilized to obtain the critical estimates required for establishing the central limit theorem.

The remainder of this paper is organized as follows. Section 2 introduces the notations and assumptions used throughout this work. Section 3 presents the main results. Two main theorems are proved in Sections 4 and 5, respectively. Lastly, the Appendix includes the proofs of several key estimates for clarity and completeness.

Throughout this paper, C with or without subscript T, denotes positive constants that may vary from one line to another. The subscript T is used specifically to indicate that the constant depends on the parameter T.

2. Notations and assumptions

In this section, we will recall some notations and list all assumptions.

2.1. **Notations.** In this subsection, we introduce some notations used in the sequel.

We use $|\cdot|$ and $|\cdot|$ for norms of vectors and matrices, respectively. Let $\langle\cdot,\cdot\rangle$ denote the scalar product in \mathbb{R}^d and A^* be the transpose of the matrix A.

Let $\mathcal{B}_b(\mathbb{R}^n)$ be the set of all bounded Borel measurable functions on \mathbb{R}^n . Let $C(\mathbb{R}^n)$ be the collection of all continuous functions on \mathbb{R}^n and $C^2(\mathbb{R}^n)$ be the set of all continuous functions on \mathbb{R}^n which have continuous partial derivatives of order up to 2.

Let $\mathscr{B}(\mathbb{R}^n)$ be the Borel σ -algebra on \mathbb{R}^n and $\mathcal{P}(\mathbb{R}^n)$ be the space of all probability measures defined on $\mathscr{B}(\mathbb{R}^n)$ carrying the usual topology of weak convergence. Let $\mathcal{P}_2(\mathbb{R}^n)$ be the collection of all the probability measures μ on $\mathscr{B}(\mathbb{R}^n)$ satisfying

$$\mu(|\cdot|^2) := \int_{\mathbb{P}^n} |x|^2 \mu(\mathrm{d}x) < \infty.$$

It is a Polish space under the L^2 -Wasserstein distance

$$W_{2}(\mu_{1}, \mu_{2}) := \inf_{\pi \in \mathscr{C}(\mu_{1}, \mu_{2})} \left(\int_{\mathbb{R}^{n} \times \mathbb{R}^{n}} |x - y|^{2} \pi(\mathrm{d}x, \mathrm{d}y) \right)^{\frac{1}{2}}, \mu_{1}, \mu_{2} \in \mathcal{P}_{2}(\mathbb{R}^{n}),$$

where $\mathscr{C}(\mu_1, \mu_2)$ is the set of all couplings π with marginals distributions μ_1 and μ_2 . Moreover, for any $x \in \mathbb{R}^n$, the Dirac measure δ_x belongs to $\mathcal{P}_2(\mathbb{R}^n)$ and if $\mu_1 = \mathscr{L}_X$, $\mu_2 = \mathcal{L}_Y$ are the corresponding distributions of random variables X and Y respectively, then

$$\mathbb{W}_2(\mu_1, \mu_2) \leqslant (\mathbb{E}|X - Y|^2)^{\frac{1}{2}},$$

where \mathbb{E} stands for the expectation with respect to \mathbb{P} .

2.2. **Derivatives for functions on** $\mathcal{P}_2(\mathbb{R}^d)$. In this subsection, we recall the definition of L-derivative for functions on $\mathcal{P}_2(\mathbb{R}^d)$. The definition was first introduced by Lions. Moreover, he used some abstract probability spaces to describe the L-derivatives. Here, for the convenience to understand the definition, we apply a straight way to state it (cf. [23]). Let I be the identify map on \mathbb{R}^n . For $\mu \in \mathcal{P}_2(\mathbb{R}^n)$ and $\phi \in L^2(\mathbb{R}^n, \mathscr{B}(\mathbb{R}^n), \mu; \mathbb{R}^n)$, $\mu(\phi) := \int_{\mathbb{R}^n} \phi(x) \mu(\mathrm{d}x)$. Moreover, by simple calculation, it holds that $\mu \circ (I + \phi)^{-1} \in$

Definition 2.1. (i) A function $h: \mathcal{P}_2(\mathbb{R}^n) \to \mathbb{R}$ is called L-differentiable at $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, if the functional

$$L^{2}(\mathbb{R}^{n}, \mathscr{B}(\mathbb{R}^{n}), \mu; \mathbb{R}^{n}) \ni \phi \to h(\mu \circ (I + \phi)^{-1})$$

is Fréchet differentiable at $0 \in L^2(\mathbb{R}^n, \mathscr{B}(\mathbb{R}^n), \mu; \mathbb{R}^n)$; that is, there exists a unique $\xi \in L^2(\mathbb{R}^n, \mathscr{B}(\mathbb{R}^n), \mu; \mathbb{R}^n)$ such that

$$\lim_{\mu(|\phi|^2)\to 0}\frac{h(\mu\circ (I+\phi)^{-1})-h(\mu)-\mu(\langle \xi,\phi\rangle)}{\sqrt{\mu(|\phi|^2)}}=0.$$

In the case, we denote $\partial_{\mu}h(\mu) = \xi$ and call it the L-derivative of h at μ . (ii) A function $h: \mathcal{P}_2(\mathbb{R}^n) \to \mathbb{R}$ is called L-differentiable on $\mathcal{P}_2(\mathbb{R}^n)$ if L-derivative $\partial_{\mu}h(\mu)$ exists for all $\mu \in \mathcal{P}_2(\mathbb{R}^n)$.

We say that a matrix-valued function $h(\mu) := (h_{ij}(\mu))$ differentiable at $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, if all its components are differentiable at μ , and set

$$\partial_{\mu}h(\mu) := (\partial_{\mu}h_{ij}(\mu)), \quad \|\partial_{\mu}h(\mu)\|_{L^{2}(\mu)}^{2} := \sum_{i,j} \int_{\mathbb{R}^{n}} |\partial_{\mu}h_{ij}(\mu)(\tilde{x})|^{2} \mu(\mathrm{d}\tilde{x}).$$

Moreover, we say $\partial_{\mu}h(\mu)(\tilde{x})$ differentiable at $\tilde{x} \in \mathbb{R}^n$, if all its components are differentiable at \tilde{x} , and set

$$\partial_{\tilde{x}}\partial_{\mu}h(\mu)(\tilde{x}) := (\partial_{\tilde{x}}\partial_{\mu}h_{ij}(\mu)(\tilde{x})), \quad \|\partial_{\tilde{x}}\partial_{\mu}h(\mu)\|_{L^{2}(\mu)}^{2} := \sum_{i,j} \int_{\mathbb{R}^{n}} \|\partial_{\tilde{x}}\partial_{\mu}h_{ij}(\mu)(\tilde{x})\|^{2}\mu(\mathrm{d}\tilde{x}).$$

Definition 2.2. The function $h(\mu)$ is said to be in $C^{(1,1)}(\mathcal{P}_2(\mathbb{R}^n))$, if $\mu \to h(\mu)$ is continuously L-differentiable, and for any $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, the mapping $\tilde{x} \to \partial_{\mu}h(\mu)(\tilde{x})$ is in $C(\mathbb{R}^n)$.

Definition 2.3. The function $h(x,\mu)$ is said to be in $C^{2,(1,1)}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$, if for any $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, the mapping $x \to h(x,\mu)$ is in $C^2(\mathbb{R}^n)$, and for any $x \in \mathbb{R}^n$, the mapping $\mu \to h(x,\mu)$ is in $C^{(1,1)}(\mathcal{P}_2(\mathbb{R}^n))$. If $h \in C^{2,(1,1)}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$, and all its derivatives are uniformly bounded, we say that $h \in C_b^{2,(1,1)}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$.

Definition 2.4. The function $h(x, \mu, y, \nu)$ is said to be in $C^{2,(1,1),2,(1,1)}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m))$, if for any $(y, \nu) \in \mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m)$, the mapping $(x, \mu) \to h(x, \mu, y, \nu)$ is in $C^{2,(1,1)}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$ and for any $(x, \mu) \in \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n)$, the mapping $(y, \nu) \to h(x, \mu, y, \nu)$ is in $C^{2,(1,1)}(\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m))$. If $h \in C^{2,(1,1),2,(1,1)}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m))$, and all its derivatives are uniformly bounded, we say that $h \in C_b^{2,(1,1),2,(1,1)}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^n))$.

2.3. **Assumptions.** In this subsection, we present all the assumptions used in the sequel: $(\mathbf{H}_{b_1,\sigma_1}^1)$ There exists a constant $L_{b_1,\sigma_1} > 0$ such that for $x_i \in \mathbb{R}^n$, $\mu_i \in \mathcal{P}_2(\mathbb{R}^n)$, $y_i \in \mathbb{R}^m$, $\nu_i \in \mathcal{P}_2(\mathbb{R}^m)$, i = 1, 2,

$$|b_1(x_1, \mu_1, y_1, \nu_1) - b_1(x_2, \mu_2, y_2, \nu_2)|^2 + ||\sigma_1(x_1, \mu_1) - \sigma_1(x_2, \mu_2)||^2$$

$$\leqslant L_{b_1, \sigma_1}(|x_1 - x_2|^2 + \mathbb{W}_2^2(\mu_1, \mu_2) + |y_1 - y_2|^2 + \mathbb{W}_2^2(\nu_1, \nu_2)).$$

 $(\mathbf{H}_{b_1}^2)$ $b_1 \in C_b^{2,(1,1),2,(1,1)}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m))$. In addition, for $y_i \in \mathbb{R}^m$, $\nu_i \in \mathcal{P}_2(\mathbb{R}^m)$, i = 1, 2,

$$\sup_{x \in \mathbb{R}^n, \mu \in \mathcal{P}_2(\mathbb{R}^n)} \|\partial_x b_1(x, \mu, y_1, \nu_1) - \partial_x b_1(x, \mu, y_2, \nu_2)\| \leqslant C(|y_1 - y_2| + \mathbb{W}_2(\nu_1, \nu_2)),$$

$$\sup_{x \in \mathbb{R}^n, \mu \in \mathcal{P}_2(\mathbb{R}^n)} \|\partial_{\mu} b_1(x, \mu, y_1, \nu_1) - \partial_{\mu} b_1(x, \mu, y_2, \nu_2)\|_{L^2(\mu)} \leqslant C(|y_1 - y_2| + \mathbb{W}_2(\nu_1, \nu_2)),$$

$$\sup_{x \in \mathbb{R}^n, \mu \in \mathcal{P}_2(\mathbb{R}^n)} \|\partial_y b_1(x, \mu, y_1, \nu_1) - \partial_y b_1(x, \mu, y_2, \nu_2)\| \leqslant C(|y_1 - y_2| + \mathbb{W}_2(\nu_1, \nu_2)),$$

$$\sup_{x \in \mathbb{R}^n, \mu \in \mathcal{P}_2(\mathbb{R}^n)} \|\partial_x^2 b_1(x, \mu, y_1, \nu_1) - \partial_x^2 b_1(x, \mu, y_2, \nu_2)\| \leqslant C(|y_1 - y_2| + \mathbb{W}_2(\nu_1, \nu_2)),$$

$$\sup_{x \in \mathbb{R}^n, \mu \in \mathcal{P}_2(\mathbb{R}^n)} \|\partial_{\tilde{x}} \partial_{\mu} b_1(x, \mu, y_1, \nu_1) - \partial_{\tilde{x}} \partial_{\mu} b_1(x, \mu, y_2, \nu_2)\|_{L^2(\mu)} \leqslant C(|y_1 - y_2| + \mathbb{W}_2(\nu_1, \nu_2)).$$

 $(\mathbf{H}_{b_2,\sigma_2}^1)$ There exists a constant $L_{b_2,\sigma_2} > 0$ such that for $y_i \in \mathbb{R}^m$, $\nu_i \in \mathcal{P}_2(\mathbb{R}^m)$, $i = 1, 2, \dots$

$$|b_2(y_1,\nu_1) - b_2(y_2,\nu_2)|^2 + ||\sigma_2(y_1,\nu_1) - \sigma_2(y_2,\nu_2)||^2 \leqslant L_{b_2,\sigma_2}(|y_1 - y_2|^2 + \mathbb{W}_2^2(\nu_1,\nu_2)).$$

 $(\mathbf{H}_{b_2,\sigma_2}^2)$ For some $p \geqslant 2$, there exist two constants $\beta_1 > 0, \beta_2 > 0$ satisfying $\beta_1 - \beta_2 > 6pL_{b_2,\sigma_2}$ such that for $y_i \in \mathbb{R}^m, \ \nu_i \in \mathcal{P}_2(\mathbb{R}^m), i = 1, 2,$

$$2\langle y_1 - y_2, b_2(y_1, \nu_1) - b_2(y_2, \nu_2) \rangle + (3p - 1) \|\sigma_2(y_1, \nu_1) - \sigma_2(y_2, \nu_2)\|^2$$

$$\leq -\beta_1 |y_1 - y_2|^2 + \beta_2 \mathbb{W}_2^2(\nu_1, \nu_2).$$

$$(\mathbf{H}_{b_2,\sigma_2}^3) \ b_2(y,\nu), \sigma_2(y,\nu) \in C_b^{2,(1,1)}(\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m)).$$

Remark 2.5. (i) $(\mathbf{H}_{b_1,\sigma_1}^1)$ yields that there exists a constant $\bar{L}_{b_1,\sigma_1} > 0$ such that for $x \in \mathbb{R}^n, \mu \in \mathcal{P}_2(\mathbb{R}^n), y \in \mathbb{R}^m, \nu \in \mathcal{P}_2(\mathbb{R}^m),$

$$|b_1(x,\mu,y,\nu)|^2 + ||\sigma_1(x,\mu)||^2 \le \bar{L}_{b_1,\sigma_1}(1+|x|^2+\mu(|\cdot|^2)+|y|^2+\nu(|\cdot|^2)).$$

(ii) $(\mathbf{H}_{b_2,\sigma_2}^1)$ implies there exists a constant $\bar{L}_{b_2,\sigma_2} > 0$ such that for $y \in \mathbb{R}^m$, $\nu \in \mathcal{P}_2(\mathbb{R}^m)$,

$$|b_2(y,\nu)|^2 + ||\sigma_2(y,\nu)||^2 \leqslant \bar{L}_{b_2,\sigma_2}(1+|y|^2+\nu(|\cdot|^2)).$$

(iii) By $(\mathbf{H}_{b_2,\sigma_2}^1)$ and $(\mathbf{H}_{b_2,\sigma_2}^2)$, we can obtain for $y \in \mathbb{R}^m$, $\nu \in \mathcal{P}_2(\mathbb{R}^m)$,

$$2\langle y, b_2(y, \nu) \rangle + (3p - 1) \|\sigma_2(y, \nu)\|^2 \leqslant -\alpha_1 |y|^2 + \alpha_2 \nu(|\cdot|^2) + C,$$

where $\alpha_1 := \beta_1 - 3pL_{b_2,\sigma_2}$, $\alpha_2 := \beta_2 + (3p-1)L_{b_2,\sigma_2}$, $\alpha_1 - \alpha_2 - L_{b_2,\sigma_2} > 0$, and C > 0 is a constant.

3. Main results

In this section, we provide main results of the paper.

3.1. The averaging principle for multiscale McKean-Vlasov SDEs. In this subsection, we state the averaging principle result for multiscale McKean-Vlasov SDEs.

Let us recall the system (1), i.e. for fixed T > 0,

$$\begin{cases} dX_t^{\varepsilon} = b_1(X_t^{\varepsilon}, \mathcal{L}_{X_t^{\varepsilon}}, Y_t^{\varepsilon, y_0, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_t^{\varepsilon, \xi}}) dt + \sigma_1(X_t^{\varepsilon}, \mathcal{L}_{X_t^{\varepsilon}}) dB_t, \\ X_0^{\varepsilon} = \varrho, \quad 0 \leqslant t \leqslant T, \\ dY_t^{\varepsilon, \xi} = \frac{1}{\varepsilon} b_2(Y_t^{\varepsilon, \xi}, \mathcal{L}_{Y_t^{\varepsilon, \xi}}) dt + \frac{1}{\sqrt{\varepsilon}} \sigma_2(Y_t^{\varepsilon, \xi}, \mathcal{L}_{Y_t^{\varepsilon, \xi}}) dW_t, \\ Y_0^{\varepsilon, \xi} = \xi, \quad 0 \leqslant t \leqslant T, \\ dY_t^{\varepsilon, y_0, \mathcal{L}_{\xi}} = \frac{1}{\varepsilon} b_2(Y_t^{\varepsilon, y_0, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_t^{\varepsilon, \xi}}) dt + \frac{1}{\sqrt{\varepsilon}} \sigma_2(Y_t^{\varepsilon, y_0, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_t^{\varepsilon, \xi}}) dW_t, \\ Y_0^{\varepsilon, y_0, \mathcal{L}_{\xi}} = y_0, \quad 0 \leqslant t \leqslant T, \end{cases}$$

where $\mathbb{E}|\varrho|^{3p} < \infty$ and $\mathbb{E}|\xi|^{3p} < \infty$. Under $(\mathbf{H}^1_{b_1,\sigma_1})$ and $(\mathbf{H}^1_{b_2,\sigma_2})$, it follows from [29, Theorem 2.1] that the system (1) has a unique strong solution $(X_{\cdot}^{\varepsilon}, Y_{\cdot}^{\varepsilon,\xi}, Y_{\cdot}^{\varepsilon,y_0,\mathcal{L}_{\xi}})$.

Consider the following system:

$$\begin{cases}
dY_t^{\xi} = b_2(Y_t^{\xi}, \mathcal{L}_{Y_t^{\xi}}) dt + \sigma_2(Y_t^{\xi}, \mathcal{L}_{Y_t^{\xi}}) dW_t, \\
Y_0^{\xi} = \xi, \quad 0 \leqslant t \leqslant T, \\
dY_t^{y_0, \mathcal{L}_{\xi}} = b_2(Y_t^{y_0, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_t^{\xi}}) dt + \sigma_2(Y_t^{y_0, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_t^{\xi}}) dW_t, \\
Y_0^{y_0, \mathcal{L}_{\xi}} = y_0, \quad 0 \leqslant t \leqslant T.
\end{cases} \tag{2}$$

Based on [29, Theorem 2.1], the above system admits a unique strong solution $(Y_{\cdot}^{\xi}, Y_{\cdot}^{y_0, \mathcal{L}_{\xi}})$ under $(\mathbf{H}_{b_2, \sigma_2}^1)$. Observing $(Y_{\cdot}^{y_0, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{\cdot}^{\xi}})$, [22, Theorem 4.11] ensures that a Markov process with the same distribution can be constructed on a new probability space. Furthermore, for any $\varphi \in \mathcal{B}_b(\mathbb{R}^m, \mathcal{P}_2(\mathbb{R}^m))$, define

$$\begin{split} P_t(y_0, \mathscr{L}_\xi; \cdot, \cdot) &:= \mathscr{L}_{Y_t^{y_0, \mathscr{L}_\xi}} \times \delta_{\mathscr{L}_{Y_t^\xi}}, \quad t \geqslant 0, \\ P_t\varphi(y_0, \mathscr{L}_\xi) &:= \int_{\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m)} \varphi(y, \nu) P_t(y_0, \mathscr{L}_\xi; \mathrm{d}y, \mathrm{d}\nu) = \mathbb{E}\varphi(Y_t^{y_0, \mathscr{L}_\xi}, \mathscr{L}_{Y_t^\xi}), \end{split}$$

 $\{P_t: t \geq 0\}$ is a transition semigroup on $\mathcal{B}_b(\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m))$. Under $(\mathbf{H}_{b_2,\sigma_2}^2)$, from [22, Theorem 4.12], it can be inferred that there exists a unique invariant probability measure $\eta \times \delta_{\eta}$ for this Markov process, where η is the unique invariant probability measure for the first equation of the system (2) (See [29, Theorem 3.1]). We then construct the averaging equation on $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0,T]}, \mathbb{P})$ as follows:

$$\begin{cases}
d\bar{X}_t = \bar{b}_1(\bar{X}_t, \mathcal{L}_{\bar{X}_t})dt + \sigma_1(\bar{X}_t, \mathcal{L}_{\bar{X}_t})dB_t, \\
\bar{X}_0 = \varrho,
\end{cases}$$
(3)

where $\bar{b}_1(x,\mu) = \int_{\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m)} b_1(x,\mu,y,\nu) \eta \times \delta_{\eta}(\mathrm{d}y,\mathrm{d}\nu).$

Now, it is the position to state the main result in this section.

Theorem 3.1. Supposed that $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{b_1}^2)$, $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^3)$ hold. Then there exists a positive constant $C_T > 0$ such that

$$\mathbb{E}\left(\sup_{0 \le t \le T} |X_t^{\varepsilon} - \bar{X}_t|^p\right) \le C_T \varepsilon^{\frac{p}{2}} (1 + \mathbb{E}|\varrho|^{3p} + |y_0|^{3p} + \mathbb{E}|\xi|^{3p}),$$

where \bar{X} is a solution of Eq.(3).

The proof of Theorem 3.1 is placed in Section 4.

3.2. The central limit theorem for multiscale McKean-Vlasov SDEs. In this subsection, we take p=4 and state the central limit theorem result for multiscale McKean-Vlasov SDEs.

First of all, we introduce some notations. Define the operator \mathcal{L} as follows: for every $\psi \in C_h^{2,(1,1)}(\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m))$

$$\mathcal{L}\psi(y,\nu) := \langle b_2(y,\nu), \partial_y \psi(y,\nu) \rangle + \frac{1}{2} Tr[\sigma_2 \sigma_2^*(y,\nu) \partial_y^2 \psi(y,\nu)]$$

$$+ \int_{\mathbb{R}^m} \left[\langle b_2(\tilde{y},\nu), \partial_\nu \psi(y,\nu)(\tilde{y}) \rangle + \frac{1}{2} Tr[\sigma_2 \sigma_2^*(\tilde{y},\nu) \partial_{\tilde{y}} \partial_\nu \psi(y,\nu)(\tilde{y})] \right] \nu(\mathrm{d}\tilde{y}),$$

and it is easy to see that \mathcal{L} is the infinitesimal generator of $(Y_t^{y,\nu}, \mathcal{L}_{Y_t^{\xi}})$, $\nu = \mathcal{L}_{\xi}$. Then consider the following Poisson equation:

$$-\mathcal{L}\Psi(x,\mu,y,\nu) = b_1(x,\mu,y,\nu) - \bar{b}_1(x,\mu). \tag{4}$$

Under the assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{b_1}^2)$ and $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^3)$, Eq.(4) admits a solution $\Psi(x,\mu,y,\nu)$ belongs to $C^{2,(1,1),2,(1,1)}(\mathbb{R}^n\times\mathcal{P}_2(\mathbb{R}^n)\times\mathbb{R}^m\times\mathcal{P}_2(\mathbb{R}^m),\mathbb{R}^n)$ (cf. Section 4.3). Finally, set

$$\begin{split} &\partial_y \Psi_{\sigma_2}(x,\mu,y,\nu) := \langle \partial_y \Psi(x,\mu,y,\nu), \sigma_2(y,\nu) \rangle, \\ &\overline{(\partial_y \Psi_{\sigma_2})(\partial_y \Psi_{\sigma_2})^*}(x,\mu) := \int_{\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m)} (\partial_y \Psi_{\sigma_2}(x,\mu,y,\nu)) (\partial_y \Psi_{\sigma_2}(x,\mu,y,\nu))^* \eta \times \delta_{\eta}(\mathrm{d}y,\mathrm{d}\nu), \\ &\Upsilon(x,\mu) := \left(\overline{(\partial_y \Psi_{\sigma_2})(\partial_y \Psi_{\sigma_2})^*}(x,\mu)\right)^{\frac{1}{2}}. \end{split}$$

The following central limit theorem is the main result in this section.

Theorem 3.2. Assume that $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{b_1}^2)$, $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^3)$ hold and $\sigma_1(\cdot,\cdot) \in C_b^{2,(1,1)}(\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m), \mathbb{R}^{n \times n})$. Then the process

$$U^{\varepsilon} := \frac{X^{\varepsilon} - \bar{X}}{\sqrt{\varepsilon}}$$

converges weakly to U in $C([0,T];\mathbb{R}^n)$ as $\varepsilon \to 0$, where the process U is the solution of the following equation

$$\begin{cases}
dU_{t} = \partial_{x}\bar{b}_{1}(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}})U_{t}dt + \mathbb{E}[\partial_{\mu}\bar{b}_{1}(u, \mathcal{L}_{\bar{X}_{t}})(\bar{X}_{t})U_{t}]|_{u=\bar{X}_{t}}dt + \partial_{x}\sigma_{1}(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}})U_{t}dB_{t}, \\
+ \mathbb{E}[\partial_{\mu}\sigma_{1}(u, \mathcal{L}_{\bar{X}_{t}})(\bar{X}_{t})U_{t}]|_{u=\bar{X}_{t}}dB_{t} + \Upsilon(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}})dV_{t}, \\
U_{0} = 0,
\end{cases} (5)$$

where V is a n-dimensional standard Brownian motion independent of B.

We postpone the proof of Theorem 3.2 to Section 5.

4. Proof of Theorem 3.1

In this section, we present the proof of Theorem 3.1. Our proof is divided into four parts. The first part (Subsection 4.1) and the second part (Subsection 4.2) providing estimates for systems (1) and (2). In the third part (Subsection 4.3), we analyze the regularity of the solution to the Poisson equation. Finally, in the fourth part (Subsection 4.4), we conclude the proof of Theorem 3.1 by integrating the results obtained in the preceding three subsections.

4.1. Some estimates for the system (1).

Lemma 4.1. Under assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{b_2,\sigma_2}^1)$ and $(\mathbf{H}_{b_2,\sigma_2}^2)$, for $p \geqslant 1$, there exist positive constants C and C_T independent of ε such that

$$\mathbb{E}\left(\sup_{0 \le t \le T} |X_t^{\varepsilon}|^{3p}\right) \le C_T(1 + \mathbb{E}|\varrho|^{3p} + |y_0|^{3p} + \mathbb{E}|\xi|^{3p}),\tag{6}$$

$$\sup_{0 \le t \le T} \mathbb{E}|Y_t^{\varepsilon,\xi}|^{3p} \le \mathbb{E}|\xi|^{3p} + C,\tag{7}$$

$$\sup_{0 \leqslant t \leqslant T} \mathbb{E} |Y_t^{\varepsilon, y_0, \mathcal{L}_{\xi}}|^{3p} \leqslant |y_0|^{3p} + C(1 + \mathbb{E}|\xi|^{3p}), \tag{8}$$

$$\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|Y_t^{\varepsilon,\xi}|^{3p}\right)\leqslant 2\mathbb{E}|\xi|^{3p}+\frac{C_T(1+\mathbb{E}|\xi|^{3p})}{\varepsilon},\tag{9}$$

$$\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|Y_t^{\varepsilon,y_0,\mathscr{L}_{\xi}}|^{3p}\right)\leqslant 2|y_0|^{3p}+\frac{C_T(1+\mathbb{E}|\xi|^{3p}+|y_0|^{3p})}{\varepsilon}.$$
 (10)

Proof. By the same deduction in [21, Lemma 4.1], we can derive the estimates (6)-(8). In the following, we will establish the estimates (9) and (10).

For $Y_t^{\varepsilon,\xi}$, applying the Itô formula to $|Y_t^{\varepsilon,\xi}|^{3p}$, we deduce that

$$|Y_{t}^{\varepsilon,\xi}|^{3p} = |\xi|^{3p} + \frac{3p}{\varepsilon} \int_{0}^{t} |Y_{s}^{\varepsilon,\xi}|^{3p-2} \langle Y_{s}^{\varepsilon,\xi}, b_{2}(Y_{s}^{\varepsilon,\xi}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) \rangle ds$$

$$+ \frac{3p}{\sqrt{\varepsilon}} \int_{0}^{t} |Y_{s}^{\varepsilon,\xi}|^{3p-2} \langle Y_{s}^{\varepsilon,\xi}, \sigma_{2}(Y_{s}^{\varepsilon,\xi}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) dW_{s} \rangle$$

$$+ \frac{3p(3p-2)}{2\varepsilon} \int_{0}^{t} |Y_{s}^{\varepsilon,\xi}|^{3p-4} ||\sigma_{2}(Y_{s}^{\varepsilon,\xi}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) Y_{s}^{\varepsilon,\xi}||^{2} ds$$

$$+ \frac{3p}{2\varepsilon} \int_{0}^{t} |Y_{s}^{\varepsilon,\xi}|^{3p-2} ||\sigma_{2}(Y_{s}^{\varepsilon,\xi}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}})||^{2} ds.$$

$$(11)$$

Note that (11) and (7) imply that

$$\begin{split} |Y^{\varepsilon,\xi}_t|^{3p} &\leqslant |\xi|^{3p} + \frac{3p}{2\varepsilon} \int_0^t |Y^{\varepsilon,\xi}_s|^{3p-2} \Big[2 \langle Y^{\varepsilon,\xi}_s, b_2(Y^{\varepsilon,\xi}_s, \mathcal{L}_{Y^{\varepsilon,\xi}_s}) \rangle \\ &\qquad \qquad + (3p-1) ||\sigma_2(Y^{\varepsilon,\xi}_s, \mathcal{L}_{Y^{\varepsilon,\xi}_s})||^2 \Big] \mathrm{d}s \\ &\qquad \qquad + \frac{3p}{\sqrt{\varepsilon}} \int_0^t |Y^{\varepsilon,\xi}_s|^{3p-2} \langle Y^{\varepsilon,\xi}_s, \sigma_2(Y^{\varepsilon,\xi}_s, \mathcal{L}_{Y^{\varepsilon,\xi}_s}) \mathrm{d}W_s \rangle \\ &\leqslant |\xi|^{3p} + \frac{3p}{2\varepsilon} \int_0^t |Y^{\varepsilon,\xi}_s|^{3p-2} \left[-\alpha_1 |Y^{\varepsilon,\xi}_s|^2 + \alpha_2 \mathbb{E} |Y^{\varepsilon,\xi}_s|^2 + C \right] \mathrm{d}s \\ &\qquad \qquad + \frac{3p}{\sqrt{\varepsilon}} \int_0^t |Y^{\varepsilon,\xi}_s|^{3p-2} \langle Y^{\varepsilon,\xi}_s, \sigma_2(Y^{\varepsilon,\xi}_s, \mathcal{L}_{Y^{\varepsilon,\xi}_s}) \mathrm{d}W_s \rangle \\ &\leqslant |\xi|^{3p} + \frac{3p}{2\varepsilon} \int_0^t \left[-\left(\alpha_1 - \frac{3p-2}{3p} \alpha_2 - L_{b_2,\sigma_2}\right) |Y^{\varepsilon,\xi}_s|^{3p} + \frac{2}{3p} \alpha_2 \mathbb{E} |Y^{\varepsilon,\xi}_s|^{3p} + C \right] \mathrm{d}s \\ &\qquad \qquad + \frac{3p}{\sqrt{\varepsilon}} \int_0^t |Y^{\varepsilon,\xi}_s|^{3p-2} \langle Y^{\varepsilon,\xi}_s, \sigma_2(Y^{\varepsilon,\xi}_s, \mathcal{L}_{Y^{\varepsilon,\xi}_s}) \mathrm{d}W_s \rangle \\ &\leqslant |\xi|^{3p} + \frac{C_T(1+\mathbb{E} |\xi|^{3p})}{\varepsilon} + \frac{3p}{\sqrt{\varepsilon}} \int_0^t |Y^{\varepsilon,\xi}_s|^{3p-2} \langle Y^{\varepsilon,\xi}_s, \sigma_2(Y^{\varepsilon,\xi}_s, \mathcal{L}_{Y^{\varepsilon,\xi}_s}) \mathrm{d}W_s \rangle \\ &\leqslant |\xi|^{3p} + \frac{C_T(1+\mathbb{E} |\xi|^{3p})}{\varepsilon} + \frac{3p}{\sqrt{\varepsilon}} \int_0^t |Y^{\varepsilon,\xi}_s|^{3p-2} \langle Y^{\varepsilon,\xi}_s, \sigma_2(Y^{\varepsilon,\xi}_s, \mathcal{L}_{Y^{\varepsilon,\xi}_s}) \mathrm{d}W_s \rangle. \end{split}$$

By the Burkholder-Davis-Gundy inequality, the Young inequality and Remark 2.5, we get

$$\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|Y_t^{\varepsilon,\xi}|^{3p}\right) \\
\leqslant \mathbb{E}|\xi|^{3p} + \frac{C_T(1+\mathbb{E}|\xi|^{3p})}{\varepsilon} + \frac{C}{\sqrt{\varepsilon}}\mathbb{E}\left[\int_0^T |Y_s^{\varepsilon,\xi}|^{6p-2} \|\sigma_2(Y_s^{\varepsilon,\xi},\mathcal{L}_{Y_s^{\varepsilon,\xi}})\|^2 \mathrm{d}s\right]^{\frac{1}{2}} \\
\leqslant \mathbb{E}|\xi|^{3p} + \frac{C_T(1+\mathbb{E}|\xi|^{3p})}{\varepsilon}$$

$$\begin{split} & + \frac{C}{\sqrt{\varepsilon}} \mathbb{E} \left[\sup_{0 \leqslant t \leqslant T} |Y_s^{\varepsilon,\xi}|^{3p} \int_0^T |Y_s^{\varepsilon,\xi}|^{3p-2} (1 + |Y_s^{\varepsilon,\xi}|^2 + \mathbb{E}|Y_s^{\varepsilon,\xi}|^2) \mathrm{d}s \right]^{\frac{1}{2}} \\ \leqslant & \mathbb{E} |\xi|^{3p} + \frac{C_T (1 + \mathbb{E}|\xi|^{3p})}{\varepsilon} + \frac{1}{2} \mathbb{E} \left(\sup_{0 \leqslant t \leqslant T} |Y_t^{\varepsilon,\xi}|^{3p} \right) + \frac{C}{\varepsilon} \int_0^T (1 + \mathbb{E}|Y_s^{\varepsilon,\xi}|^{3p}) \mathrm{d}s \\ \leqslant & \mathbb{E} |\xi|^{3p} + \frac{C_T (1 + \mathbb{E}|\xi|^{3p})}{\varepsilon} + \frac{1}{2} \mathbb{E} \left(\sup_{0 \leqslant t \leqslant T} |Y_t^{\varepsilon,\xi}|^{3p} \right), \end{split}$$

which implies that

$$\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|Y_t^{\varepsilon,\xi}|^{3p}\right)\leqslant 2\mathbb{E}|\xi|^{3p}+\frac{C_T(1+\mathbb{E}|\xi|^{3p})}{\varepsilon}.$$

Moreover, for $Y_t^{\varepsilon,y_0,\mathscr{L}_{\xi}}$, using a similar argument as for $Y_t^{\varepsilon,\xi}$, we can obtain the estimate (10). The proof is complete.

4.2. Some estimates for the system (2).

Lemma 4.2. Under assumptions $(\mathbf{H}_{b_2,\sigma_2}^1)$ and $(\mathbf{H}_{b_2,\sigma_2}^2)$, there exists a positive constant Cindependent of ε such that

$$\mathbb{E}|Y_t^{y_0,\mathcal{L}_{\xi}}|^2 \leqslant |y_0|^2 e^{-\alpha_1 t} + e^{-(\alpha_1 - \alpha_2)t} \mathbb{E}|\xi|^2 + C,$$

and for $p \ge 2$

$$\mathbb{E}|Y_t^{y_0,\mathcal{L}_{\xi}}|^{2p} \leqslant |y_0|^{2p} e^{-p(\alpha_1 - \alpha_2 - L_{b_2,\sigma_2})t} + C\mathbb{E}|\xi|^{2p} + C.$$

Proof. First of all, we estimate Y_t^{ξ} . Applying the Itô formula to $|Y_t^{\xi}|^{2p}$ and taking the expectation, we obtain

$$\mathbb{E}|Y_{t}^{\xi}|^{2p} = \mathbb{E}|\xi|^{2p} + 2p\mathbb{E}\int_{0}^{t}|Y_{s}^{\xi}|^{2p-2}\langle Y_{s}^{\xi}, b_{2}(Y_{s}^{\xi}, \mathcal{L}_{Y_{s}^{\xi}})\rangle ds
+2p(p-1)\mathbb{E}\int_{0}^{t}|Y_{s}^{\xi}|^{2p-4}\|\sigma_{2}(Y_{s}^{\xi}, \mathcal{L}_{Y_{s}^{\xi}})Y_{s}^{\xi}\|^{2}ds
+p\mathbb{E}\int_{0}^{t}|Y_{s}^{\xi}|^{2p-2}\|\sigma_{2}(Y_{s}^{\xi}, \mathcal{L}_{Y_{s}^{\xi}})\|^{2}ds.$$

By Remark 2.5, we have

$$\frac{\mathrm{d}}{\mathrm{d}t} \mathbb{E} |Y_{t}^{\xi}|^{2p} = 2p \mathbb{E} \Big[|Y_{t}^{\xi}|^{2p-2} \langle Y_{t}^{\xi}, b_{2}(Y_{t}^{\xi}, \mathcal{L}_{Y_{t}^{\xi}}) \rangle \Big] + 2p(p-1) \mathbb{E} \Big[|Y_{t}^{\xi}|^{2p-4} \| \sigma_{2}(Y_{t}^{\xi}, \mathcal{L}_{Y_{t}^{\xi}}) Y_{t}^{\xi} \|^{2} \Big] \\
+ p \mathbb{E} \Big[|Y_{t}^{\xi}|^{2p-2} \| \sigma_{2}(Y_{t}^{\xi}, \mathcal{L}_{Y_{t}^{\xi}}) \|^{2} \Big] \\
\leqslant p \mathbb{E} \Big[|Y_{t}^{\xi}|^{2p-2} \Big(2 \langle Y_{t}^{\xi}, b_{2}(Y_{t}^{\xi}, \mathcal{L}_{Y_{t}^{\xi}}) \rangle + (2p-1) \| \sigma_{2}(Y_{t}^{\xi}, \mathcal{L}_{Y_{t}^{\xi}}) \|^{2} \Big) \Big] \\
\leqslant p \mathbb{E} \Big[|Y_{t}^{\xi}|^{2p-2} \Big(-\alpha_{1} |Y_{t}^{\xi}|^{2} + \alpha_{2} \mathcal{L}_{Y_{t}^{\xi}} (|\cdot|^{2}) + C \Big) \Big] \\
\leqslant -p(\alpha_{1} - \alpha_{2} - L_{b_{2}, \sigma_{2}}) \mathbb{E} |Y_{t}^{\xi}|^{2p} + C.$$

The comparison theorem implies

$$\mathbb{E}|Y_t^{\xi}|^{2p} \leqslant \mathbb{E}|\xi|^{2p} e^{-p(\alpha_1 - \alpha_2 - L_{b_2, \sigma_2})t} + C. \tag{12}$$

Next, we deal with $Y_t^{y_0,\mathscr{L}_\xi}$. For any $\lambda>0$, the Itô formula yields that

$$\begin{split} & \mathbb{E}|Y_t^{y_0,\mathscr{L}_\xi}|^{2p}e^{\lambda t} \\ = & |y_0|^{2p} + \lambda \mathbb{E} \int_0^t e^{\lambda s}|Y_s^{y_0,\mathscr{L}_\xi}|^{2p}\mathrm{d}s + 2p\mathbb{E} \int_0^t e^{\lambda s}|Y_s^{y_0,\mathscr{L}_\xi}|^{2p-2}\langle Y_s^{y_0,\mathscr{L}_\xi}, b_2(Y_s^{y_0,\mathscr{L}_\xi},\mathscr{L}_{Y_s^\xi})\rangle \mathrm{d}s \\ & + 2p(p-1)\mathbb{E} \int_0^t e^{\lambda s}|Y_s^{y_0,\mathscr{L}_\xi}|^{2p-4} \|\sigma_2(Y_s^{y_0,\mathscr{L}_\xi},\mathscr{L}_{Y_s^\xi})Y_s^{y_0,\mathscr{L}_\xi}\|^2 \mathrm{d}s \\ & + p\mathbb{E} \int_0^t e^{\lambda s}|Y_s^{y_0,\mathscr{L}_\xi}|^{2p-2} \|\sigma_2(Y_s^{y_0,\mathscr{L}_\xi},\mathscr{L}_{Y_s^\xi})\|^2 \mathrm{d}s \\ \leqslant & |y_0|^{2p} + \lambda \mathbb{E} \int_0^t e^{\lambda s}|Y_s^{y_0,\mathscr{L}_\xi}|^{2p}\mathrm{d}s \\ & + p\mathbb{E} \int_0^t e^{\lambda s}|Y_s^{y_0,\mathscr{L}_\xi}|^{2p-2} \left[2\langle Y_s^{y_0,\mathscr{L}_\xi},b_2(Y_s^{y_0,\mathscr{L}_\xi},\mathscr{L}_{Y_s^\xi})\rangle + (2p-1)\|\sigma_2(Y_s^{y_0,\mathscr{L}_\xi},\mathscr{L}_{Y_s^\xi})\|^2\right]\mathrm{d}s \\ \leqslant & |y_0|^{2p} + \lambda \mathbb{E} \int_0^t e^{\lambda s}|Y_s^{y_0,\mathscr{L}_\xi}|^{2p}\mathrm{d}s \\ & + p \int_0^t e^{\lambda s} \left[-(\alpha_1 - \alpha_2 - L_{b_2,\sigma_2})\mathbb{E}|Y_s^{y_0,\mathscr{L}_\xi}|^{2p} + C\mathbb{E}|Y_s^\xi|^{2p} + C\right]\mathrm{d}s. \end{split}$$

Letting $\lambda = p(\alpha_1 - \alpha_2 - L_{b_2,\sigma_2})$, we have

$$\mathbb{E}|Y_t^{y_0,\mathcal{L}_{\xi}}|^{2p}e^{p(\alpha_1-\alpha_2-L_{b_2,\sigma_2})t} \leqslant |y_0|^{2p} + C \int_0^t e^{p(\alpha_1-\alpha_2-L_{b_2,\sigma_2})s} (\mathbb{E}|Y_s^{\xi}|^{2p} + 1) ds
\leqslant |y_0|^{2p} + C \int_0^t (\mathbb{E}|\xi|^{2p} + e^{p(\alpha_1-\alpha_2-L_{b_2,\sigma_2})s}) ds.$$

Then simple calculation implies that

$$\mathbb{E}|Y_t^{y_0,\mathscr{L}_{\xi}}|^{2p} \leqslant |y_0|^{2p} e^{-p(\alpha_1 - \alpha_2 - L_{b_2,\sigma_2})t} + C\mathbb{E}|\xi|^{2p} + C.$$

Finally, by similar arguments as above, we get

$$\mathbb{E}|Y_t^{\xi}|^2 \leqslant \mathbb{E}|\xi|^2 e^{-(\alpha_1 - \alpha_2)t} + C_{\xi}$$

and

$$\mathbb{E}|Y_t^{y_0,\mathcal{L}_{\xi}}|^2 \leqslant |y_0|^2 e^{-\alpha_1 t} + \mathbb{E}|\xi|^2 e^{-(\alpha_1 - \alpha_2)t} + C.$$

The proof is complete.

The following two results can be obtained through a deduction similar to that of [21, Lemma 4.5] and [21, Lemma 4.6].

Lemma 4.3. Suppose that assumptions $(\mathbf{H}^1_{b_2,\sigma_2})$ and $(\mathbf{H}^2_{b_2,\sigma_2})$ hold. Then it holds that for any $t \geq 0$, $y_i \in \mathbb{R}^m$, $\zeta_i \in L^2(\Omega, \mathscr{F}_0, \mathbb{P}; \mathbb{R}^m)$, i = 1, 2,

$$\mathbb{E}|Y_t^{y_1,\mathcal{L}_{\zeta_1}} - Y_t^{y_2,\mathcal{L}_{\zeta_2}}|^2 \leqslant |y_1 - y_2|^2 e^{-\beta_1 t} + \mathbb{E}|\zeta_1 - \zeta_2|^2 e^{-(\beta_1 - \beta_2)t}.$$

Lemma 4.4. Suppose that assumptions $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{b_2,\sigma_2}^1)$, $(\mathbf{H}_{b_2,\sigma_2}^2)$ hold. Then it holds that for any $t \geq 0$, $x \in \mathbb{R}^n$, $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, there exists a constant C > 0 such that

$$|\mathbb{E}b_1(x,\mu,Y_t^{y_0,\mathcal{L}_{\xi}},\mathcal{L}_{Y_t^{\xi}}) - \bar{b}_1(x,\mu)|^2 \leqslant Ce^{-(\beta_1-\beta_2)t}(1+|y_0|^2 + \mathcal{L}_{\xi}(|\cdot|^2)).$$

4.3. **Poisson equation.** In this subsection, we analyze the regularity of the solution to the Poisson equation.

Proposition 4.5. Suppose that $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{b_1}^2)$, $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^3)$ hold. Set

$$\Psi(x,\mu,y,\nu) := \int_0^\infty \left(\mathbb{E}b_1(x,\mu,Y_s^{y,\nu},\mathcal{L}_{Y_s^{\xi}}) - \bar{b}_1(x,\mu) \right) \mathrm{d}s, \quad \nu = \mathcal{L}_{\xi}.$$
 (13)

Then $\Psi(x,\mu,y,\nu)$ belongs to $C^{2,(1,1),2,(1,1)}(\mathbb{R}^n\times\mathcal{P}_2(\mathbb{R}^n)\times\mathbb{R}^m\times\mathcal{P}_2(\mathbb{R}^m),\mathbb{R}^n)$ and is the unique solution for Eq.(4). Moreover, it holds that for any $x \in \mathbb{R}^n$, $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, $y \in \mathbb{R}^m$, $\nu \in \mathcal{P}_2(\mathbb{R}^m),$

$$\sup_{x \in \mathbb{R}^{n}, \mu \in \mathcal{P}_{2}(\mathbb{R}^{n})} \max \left\{ |\Psi(x, \mu, y, \nu)|, \|\partial_{x}\Psi(x, \mu, y, \nu)\|, \|\partial_{\mu}\Psi(x, \mu, y, \nu)\|_{L^{2}(\mu)}, \right.$$

$$\left. \|\partial_{x}^{2}\Psi(x, \mu, y, \nu)\|, \|\partial_{\tilde{x}}\partial_{\mu}\Psi(x, \mu, y, \nu)(\cdot)\|_{L^{2}(\mu)} \right\}$$

$$\leqslant C(1 + |y|^{2} + \nu(|\cdot|^{2}))^{1/2}$$
(14)

and

$$\sup_{x \in \mathbb{R}^{n}, \mu \in \mathcal{P}_{2}(\mathbb{R}^{n}), y \in \mathbb{R}^{m}, \nu \in \mathcal{P}_{2}(\mathbb{R}^{m})} \max \left\{ \|\partial_{\nu} \Psi(x, \mu, y, \nu)\|_{L^{2}(\nu)}, \|\partial_{y} \Psi(x, \mu, y, \nu)\|, \\ \|\partial_{x} \partial_{y} \Psi(x, \mu, y, \nu)\|, \|\partial_{\mu} \partial_{y} \Psi(x, \mu, y, \nu)\|_{L^{2}(\mu)}, \\ \|\partial_{y}^{2} \Psi(x, \mu, y, \nu)\|, \|\partial_{\nu} \partial_{y} \Psi(x, \mu, y, \nu)(\cdot)\|_{L^{2}(\nu)} \right\} \leqslant C. \tag{15}$$

Proof. The proof is divided into three steps. First, we show that $\Psi(x,\mu,y,\nu)$ belongs to $C^{2,(1,1),2,(1,1)}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m), \mathbb{R}^n)$ and serves as the unique solution to Eq.(4). Next, in the second and third steps, we provide the verification for the regularity estimates (14) and (15) correspondingly.

Step 1. We prove $\Psi(x, \mu, y, \nu)$ belongs to $C^{2,(1,1),2,(1,1)}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m), \mathbb{R}^n)$ and is the unique solution for Eq.(4).

Since η is independent of x, we can take the derivative to $b(x,\mu)$ directly with respect to x, yielding

$$\partial_x^i \bar{b}_1(x,\mu) = \int_{\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m)} \partial_x^i b_1(x,\mu,y,\nu) \eta \times \delta_{\eta}(\mathrm{d}y,\mathrm{d}\nu), \quad i = 1, 2,$$

which implies that $\bar{b}_1(\cdot,\mu) \in C_b^2(\mathbb{R}^n)$. Similarly, it follows that $\bar{b}_1(x,\cdot) \in C_b^{(1,1)}(\mathcal{P}_2(\mathbb{R}^n))$, leading to the conclusion that $\bar{b}_1 \in C_b^{2,(1,1)}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n))$.

Under the assumptions $(\mathbf{H}_{b_1}^2)$ and $(\mathbf{H}_{b_2,\sigma_2}^3)$, and following a similar argument as in [3, Lemma 6.2], it can be concluded that $\Psi(x, \mu, y, \nu)$ belongs to $C^{2,(1,1),2,(1,1)}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathcal{P}_2(\mathbb{R}^n))$ $\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m), \mathbb{R}^n$). Then by acting the generator \mathcal{L} on $(Y_t^{y,\nu}, \mathscr{L}_{Y_t^{\xi}}), \nu = \mathscr{L}_{\xi}$, it holds that

$$\mathcal{L}\Psi(x,\mu,y,\nu) = \mathcal{L} \int_0^\infty \left(\mathbb{E}b_1(x,\mu,Y_s^{y,\nu},\mathcal{L}_{Y_s^{\xi}}) - \bar{b}_1(x,\mu) \right) ds$$

$$= \int_0^\infty (\mathcal{L}P_s) \left(b_1(x,\mu,y,\nu) - \bar{b}_1(x,\mu) \right) ds$$

$$= \int_0^\infty \frac{dP_s \left(b_1(x,\mu,y,\nu) - \bar{b}_1(x,\mu) \right)}{ds} ds$$

$$= P_s \Big(b_1(x, \mu, y, \nu) - \bar{b}_1(x, \mu) \Big) \Big|_0^{\infty}$$

$$= \lim_{s \to \infty} P_s \Big(b_1(x, \mu, y, \nu) \Big) - \bar{b}_1(x, \mu) - \Big(b_1(x, \mu, y, \nu) - \bar{b}_1(x, \mu) \Big)$$

$$= -\Big(b_1(x, \mu, y, \nu) - \bar{b}_1(x, \mu) \Big),$$

which yields that $\Psi(x, \mu, y, \nu)$ is a solution for Eq.(4). According to the Itô formula, the uniqueness can be achieved through the precise calculation.

Step 2. We establish the estimate (14).

$$\begin{aligned} |\Psi(x,\mu,y,\nu)| &\leqslant & \int_0^\infty \left| \mathbb{E}b_1(x,\mu,Y_s^{y,\nu},\mathcal{L}_{Y_s^{\xi}}) - \bar{b}_1(x,\mu) \right| \mathrm{d}s \\ &\leqslant & \int_0^\infty Ce^{\frac{-(\beta_1 - \beta_2)s}{2}} (1 + |y|^2 + \nu(|\cdot|^2))^{1/2} \mathrm{d}s \\ &\leqslant & C(1 + |y|^2 + \nu(|\cdot|^2))^{1/2}. \end{aligned}$$

Let $\check{b}_1(x,\mu,y,\nu,s) = \mathbb{E}b_1(x,\mu,Y_s^{y,\nu},\mathcal{L}_{Y_s^{\xi}}) - \bar{b}_1(x,\mu), \quad \nu = \mathcal{L}_{\xi}$. For any $s_0 > 0$, we define

$$\tilde{b}_{1,s_0}(x,\mu,y,\nu,s) := \hat{b}_1(x,\mu,y,\nu,s) - \hat{b}_1(x,\mu,y,\nu,s+s_0),$$

where $\hat{b}_1(x, \mu, y, \nu, s) := \mathbb{E}b_1(x, \mu, Y_s^{y,\nu}, \mathcal{L}_{Y_s^{\xi}})$. Hence,

$$\tilde{b}_{1,s_0}(x,\mu,y,\nu,s) = \mathbb{E}b_1(x,\mu,Y_s^{y,\nu},\mathcal{L}_{Y_s^{\xi}}) - \mathbb{E}b_1(x,\mu,Y_{s+s_0}^{y,\nu},\mathcal{L}_{Y_{s+s_0}^{\xi}})
= \mathbb{E}b_1(x,\mu,Y_s^{y,\nu},\mathcal{L}_{Y_s^{\xi}}) - \bar{b}_1(x,\mu)
- [\mathbb{E}b_1(x,\mu,Y_{s+s_0}^{y,\nu},\mathcal{L}_{Y_{s+s_0}^{\xi}}) - \bar{b}_1(x,\mu)].$$

Based on Lemma 4.4, we know that

$$\lim_{s_0 \to \infty} \tilde{b}_{1,s_0}(x, \mu, y, \nu, s) = \check{b}_1(x, \mu, y, \nu, s).$$

In order to prove

$$\|\partial_x \Psi(x, \mu, y, \nu)\| \le C(1 + |y|^2 + \nu(|\cdot|^2))^{1/2},$$

it suffices to show that for any $s_0 > 0$, $s \ge 0$, $x \in \mathbb{R}^n$, $\mu \in \mathcal{P}_2(\mathbb{R}^n)$, $y \in \mathbb{R}^m$, $\nu \in \mathcal{P}_2(\mathbb{R}^m)$

$$\|\partial_x \tilde{b}_{1,s_0}(x,\mu,y,\nu,s)\| \leqslant Ce^{-\frac{\beta_1-\beta_2}{2}s}(1+|y|^2+\nu(|\cdot|^2))^{1/2}.$$
 (16)

We prove (16) in the Appendix.

By similar arguments as above, we get

$$\|\partial_{\mu}\tilde{b}_{1,s_{0}}(x,\mu,y,\nu,s)\|_{L^{2}(\mu)} \leqslant Ce^{-\frac{\beta_{1}-\beta_{2}}{2}s}(1+|y|^{2}+\nu(|\cdot|^{2}))^{1/2},$$

$$\|\partial_{x}^{2}\tilde{b}_{1,s_{0}}(x,\mu,y,\nu,s)\| \leqslant Ce^{-\frac{\beta_{1}-\beta_{2}}{2}s}(1+|y|^{2}+\nu(|\cdot|^{2}))^{1/2},$$

$$\|\partial_{x}\partial_{\mu}\tilde{b}_{1,s_{0}}(x,\mu,y,\nu,s)\|_{L^{2}(\mu)} \leqslant Ce^{-\frac{\beta_{1}-\beta_{2}}{2}s}(1+|y|^{2}+\nu(|\cdot|^{2}))^{1/2},$$

which yields that

$$\|\partial_{\mu}\Psi(x,\mu,y,\nu)\|_{L^{2}(\mu)} \leq C(1+|y|^{2}+\nu(|\cdot|^{2}))^{1/2},$$

$$\|\partial_{x}^{2}\Psi(x,\mu,y,\nu)\| \leq C(1+|y|^{2}+\nu(|\cdot|^{2}))^{1/2},$$

$$\|\partial_{\tilde{x}}\partial_{\mu}\Psi(x,\mu,y,\nu)\|_{L^{2}(\mu)} \leq (1+|y|^{2}+\nu(|\cdot|^{2}))^{1/2}.$$

Step 3. We establish the estimate (15).

Relying on (13), $(\mathbf{H}_{b_1,\sigma_1}^1)$ and Lemma 4.3, we conclude that

$$|\Psi(x,\mu,y,\nu_{1}) - \Psi(x,\mu,y,\nu_{2})|$$

$$\leqslant C \int_{0}^{\infty} \mathbb{E}\left(|Y_{s}^{y,\nu_{1}} - Y_{s}^{y,\nu_{2}}| + \mathbb{W}_{2}(\mathcal{L}_{Y_{s}^{\xi_{1}}}, \mathcal{L}_{Y_{s}^{\xi_{2}}})\right) ds$$

$$\leqslant C(\mathbb{E}|\xi_{1} - \xi_{2}|^{2})^{\frac{1}{2}}, \tag{17}$$

for all ξ_1 , ξ_2 with $\mathcal{L}_{\xi_1} = \nu_1$, $\mathcal{L}_{\xi_2} = \nu_2$. Hence, taking at the right-hand side of (17) the infimum over all such ξ_1 , ξ_2 and considering the characterization of the L^2 -Wasserstein metric, we get

$$|\Psi(x, \mu, y, \nu_1) - \Psi(x, \mu, y, \nu_2)| \leq C \mathbb{W}_2(\nu_1, \nu_2),$$

which implies that

$$\sup_{x \in \mathbb{R}^n, \mu \in \mathcal{P}_2(\mathbb{R}^n), y \in \mathbb{R}^m, \nu \in \mathcal{P}_2(\mathbb{R}^m)} \|\partial_{\nu} \Psi(x, \mu, y, \nu)\|_{L^2(\nu)} \leqslant C.$$

Note that $\partial_y Y_s^{y,\nu}$ and $\partial_y^2 Y_s^{y,\nu}$ satisfy

$$\begin{cases}
d\partial_y Y_s^{y,\nu} = \partial_y b_2(Y_s^{y,\nu}, \mathcal{L}_{Y_s^{\xi}}) \partial_y Y_s^{y,\nu} ds + \partial_y \sigma_2(Y_s^{y,\nu}, \mathcal{L}_{Y_s^{\xi}}) \partial_y Y_s^{y,\nu} dW_s, \\
\partial_y Y_0^{y,\nu} = I,
\end{cases}$$

and

$$\begin{cases} d\partial_y^2 Y_s^{y,\nu} = [\partial_y^2 b_2(Y_s^{y,\nu}, \mathcal{L}_{Y_s^{\xi}}) \langle \partial_y Y_s^{y,\nu}, \partial_y Y_s^{y,\nu} \rangle + \partial_y b_2(Y_s^{y,\nu}, \mathcal{L}_{Y_s^{\xi}}) \partial_y^2 Y_s^{y,\nu}] ds \\ + [\partial_y^2 \sigma_2(Y_s^{y,\nu}, \mathcal{L}_{Y_s^{\xi}}) \langle \partial_y Y_s^{y,\nu}, \partial_y Y_s^{y,\nu} \rangle + \partial_y \sigma_2(Y_s^{y,\nu}, \mathcal{L}_{Y_s^{\xi}}) \partial_y^2 Y_s^{y,\nu}] dW_s, \\ \partial_y^2 Y_0^{y,\nu} = 0. \end{cases}$$

Under the assumption $(\mathbf{H}_{b_2,\sigma_2}^2)$, it holds that

$$\sup_{y \in \mathbb{R}^m, \nu \in \mathcal{P}_2(\mathbb{R}^m)} \mathbb{E} \|\partial_y Y_s^{y,\nu}\|^2 \leqslant C e^{-\gamma s}, \tag{18}$$

$$\sup_{y \in \mathbb{R}^m, \nu \in \mathcal{P}_2(\mathbb{R}^m)} \mathbb{E} \|\partial_y^2 Y_s^{y,\nu}\|^2 \leqslant C e^{-\gamma s}, \tag{19}$$

and

$$\sup_{y \in \mathbb{R}^m} \mathbb{E} \|\partial_y Y_s^{y,\nu_1} - \partial_y Y_s^{y,\nu_2}\|^2 \leqslant C e^{-\gamma s} \mathbb{E} |\xi_1 - \xi_2|^2, \tag{20}$$

where $\nu_1 = \mathcal{L}_{\xi_1}$, $\nu_2 = \mathcal{L}_{\xi_2}$. In view of (13), by $(\mathbf{H}_{b_1}^2)$ and (18), we deduce that

$$\|\partial_y \Psi(x,\mu,y,\nu)\| \leqslant \int_0^\infty \mathbb{E} \|\partial_y b_1(x,\mu,Y_s^{y,\nu},\mathcal{L}_{Y_s^{\xi}}) \cdot \partial_y Y_s^{y,\nu}\| ds \leqslant C \int_0^\infty \mathbb{E} \|\partial_y Y_s^{y,\nu}\| ds \leqslant C,$$

and

$$\|\partial_x \partial_y \Psi(x, \mu, y, \nu)\| \leqslant \int_0^\infty \mathbb{E} |\partial_x \partial_y b_1(x, \mu, Y_s^{y, \nu}, \mathcal{L}_{Y_s^{\xi}}) \cdot \partial_y Y_s^{y, \nu}| ds \leqslant C.$$

Similarly, we have

$$\|\partial_{\mu}\partial_{y}\Psi(x,\mu,y,\nu)\|_{L^{2}(\mu)} \leqslant C.$$

The term $\partial_y^2 \Psi(x, \mu, y, \nu)$ could be represented by

$$\partial_y^2 \Psi(x, \mu, y, \nu) = \int_0^\infty \mathbb{E}[\partial_y^2 b_1(x, \mu, Y_s^{y, \nu}, \mathcal{L}_{Y_s^{\xi}}) \cdot \langle \partial_y Y_s^{y, \nu}, \partial_y Y_s^{y, \nu} \rangle] ds$$

$$+ \int_0^\infty \mathbb{E}[\partial_y b_1(x,\mu,Y_s^{y,\nu},\mathscr{L}_{Y_s^{\xi}})\partial_y^2 Y_s^{y,\nu}] \mathrm{d}s.$$

Combining $(\mathbf{H}_{b_1}^2)$, (18) and (19), we have

$$\|\partial_y^2 \Psi(x,\mu,y,\nu)\| \leqslant C \int_0^\infty \mathbb{E} \|\partial_y Y_s^{y,\nu}\|^2 ds + C \int_0^\infty \mathbb{E} \|\partial_y^2 Y_s^{y,\nu}\| ds \leqslant C.$$

In the following we estimate $\partial_{\nu}\partial_{y}\Psi(x,\mu,y,\nu)$. Using $(\mathbf{H}_{b_{1}}^{2})$, (18), (20) and Lemma 4.3, we get

$$\begin{split} &\|\partial_{y}\Psi(x,\mu,y,\nu_{1})-\partial_{y}\Psi(x,\mu,y,\nu_{2})\| \\ \leqslant & \int_{0}^{\infty}\mathbb{E}\Big[\|\partial_{y}b_{1}(x,\mu,Y_{s}^{y,\nu_{1}},\mathcal{L}_{Y_{s}^{\xi_{1}}})-\partial_{y}b_{1}(x,\mu,Y_{s}^{y,\nu_{2}},\mathcal{L}_{Y_{s}^{\xi_{2}}})\|\cdot\|\partial_{y}Y_{s}^{y,\nu_{1}}\| \\ & +\|\partial_{y}b_{1}(x,\mu,Y_{s}^{y,\nu_{2}},\mathcal{L}_{Y_{s}^{\xi_{2}}})\|\cdot\|\partial_{y}Y_{s}^{y,\nu_{1}}-\partial_{y}Y_{s}^{y,\nu_{2}}\|\Big]\mathrm{d}s \\ \leqslant & C\int_{0}^{\infty}\Big[\Big(\mathbb{E}|Y_{s}^{y,\nu_{1}}-Y_{s}^{y,\nu_{2}}|^{2}\Big)^{\frac{1}{2}}+\Big(\mathbb{E}|Y_{s}^{\xi_{1}}-Y_{s}^{\xi_{2}}|^{2}\Big)^{\frac{1}{2}}\Big]\cdot\Big(\mathbb{E}\|\partial_{y}Y_{s}^{y,\nu_{1}}\|^{2}\Big)^{\frac{1}{2}}\mathrm{d}s \\ +& C\int_{0}^{\infty}\Big(\mathbb{E}\|\partial_{y}Y_{s}^{y,\nu_{1}}-\partial_{y}Y_{s}^{y,\nu_{2}}\|^{2}\Big)^{\frac{1}{2}}\mathrm{d}s \\ \leqslant & C(\mathbb{E}|\xi_{1}-\xi_{2}|^{2})^{\frac{1}{2}}, \end{split}$$

for all $\xi_1, \, \xi_2$ with $\mathcal{L}_{\xi_1} = \nu_1, \, \mathcal{L}_{\xi_2} = \nu_2$. Hence,

$$\|\partial_y \Psi(x,\mu,y,\nu_1) - \partial_y \Psi(x,\mu,y,\nu_2)\| \leqslant C \mathbb{W}_2(\nu_1,\nu_2),$$

which implies that

$$\sup_{x \in \mathbb{R}^n, \mu \in \mathcal{P}_2(\mathbb{R}^n), y \in \mathbb{R}^m, \nu \in \mathcal{P}_2(\mathbb{R}^m)} \|\partial_{\nu} \partial_y \Psi(x, \mu, y, \nu)\|_{L^2(\nu)} \leqslant C.$$

The proof is complete.

4.4. The proof of Theorem 3.1. First of all, we prove the Lipschitz continuity of the coefficient \bar{b}_1 .

By Lemma 4.4 and $(\mathbf{H}_{b_1,\sigma_1}^1)$, for any $x_i \in \mathbb{R}^n$, $\mu_i \in \mathcal{P}_2(\mathbb{R}^n)$, i = 1, 2, we have

$$\begin{aligned} &|\bar{b}_{1}(x_{1},\mu_{1}) - \bar{b}_{1}(x_{2},\mu_{2})|^{2} \\ &\leqslant & 3|\bar{b}_{1}(x_{1},\mu_{1}) - \mathbb{E}b_{1}(x_{1},\mu_{1},Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}})|^{2} \\ &+ 3|\mathbb{E}b_{1}(x_{2},\mu_{2},Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}}) - \bar{b}_{1}(x_{2},\mu_{2})|^{2} \\ &+ 3|\mathbb{E}b_{1}(x_{1},\mu_{1},Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}}) - \mathbb{E}b_{1}(x_{2},\mu_{2},Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}})|^{2} \\ &\leqslant & Ce^{-(\beta_{1}-\beta_{2})t}(1+|y_{0}|^{2}+||\mathcal{L}_{\xi}||^{2}) + C(|x_{1}-x_{2}|^{2}+\mathbb{W}_{2}^{2}(\mu_{1},\mu_{2})). \end{aligned}$$

Letting $t \to \infty$, we obtain

$$|\bar{b}_1(x_1, \mu_1) - \bar{b}_1(x_2, \mu_2)|^2 \leqslant C(|x_1 - x_2|^2 + \mathbb{W}_2^2(\mu_1, \mu_2)). \tag{21}$$

Combining the assumption $(\mathbf{H}_{b_1,\sigma_1}^1)$, we conclude that Eq.(3) admits a unique strong solution \bar{X} .

Note that

$$X_{t}^{\varepsilon} - \bar{X}_{t} = \int_{0}^{t} \left(b_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}}) - \bar{b}_{1}(\bar{X}_{s}, \mathcal{L}_{\bar{X}_{s}}) \right) ds$$

$$+ \int_{0}^{t} \left(\sigma_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}) - \sigma_{1}(\bar{X}_{s}, \mathcal{L}_{\bar{X}_{s}}) \right) dB_{s}$$

$$= \int_{0}^{t} \left(b_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}}) - \bar{b}_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}) \right) ds$$

$$+ \int_{0}^{t} \left(\bar{b}_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}) - \bar{b}_{1}(\bar{X}_{s}, \mathcal{L}_{\bar{X}_{s}}) \right) ds$$

$$+ \int_{0}^{t} \left(\sigma_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}) - \sigma_{1}(\bar{X}_{s}, \mathcal{L}_{\bar{X}_{s}}) \right) dB_{s}.$$

From the Hölder inequality, the Burkholder-Davis-Gundy inequality, $(\mathbf{H}_{b_1,\sigma_1}^1)$ and (21), it follows that for any $p \ge 2$,

$$\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}\left|X_{t}^{\varepsilon}-\bar{X}_{t}\right|^{p}\right)
\leqslant C\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}\left|\int_{0}^{t}\left(b_{1}(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}},Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{s}^{\varepsilon,\xi}})-\bar{b}_{1}(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}})\right)\mathrm{d}s\right|^{p}\right)
+CT^{p-1}\mathbb{E}\int_{0}^{T}\left|\bar{b}_{1}(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}})-\bar{b}_{1}(\bar{X}_{s},\mathcal{L}_{\bar{X}_{s}})\right|^{p}\mathrm{d}s
+C\mathbb{E}\left[\int_{0}^{T}\left\|\sigma_{1}(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}})-\sigma_{1}(\bar{X}_{s},\mathcal{L}_{\bar{X}_{s}})\right\|^{2}\mathrm{d}s\right]^{\frac{p}{2}}
\leqslant C\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}\left|\int_{0}^{t}\left(b_{1}(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}},Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{s}^{\varepsilon,\xi}})-\bar{b}_{1}(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}})\right)\mathrm{d}s\right|^{p}\right)
+C_{T}\int_{0}^{T}\mathbb{E}|X_{s}^{\varepsilon}-\bar{X}_{s}|^{p}\mathrm{d}s.$$

The Grownall inequality yields that

$$\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|X_{t}^{\varepsilon}-\bar{X}_{t}|^{p}\right)$$

$$\leqslant C_{T}\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}\left|\int_{0}^{t}\left(b_{1}(X_{s}^{\varepsilon},\mathscr{L}_{X_{s}^{\varepsilon}},Y_{s}^{\varepsilon,y_{0},\mathscr{L}_{\xi}},\mathscr{L}_{Y_{s}^{\varepsilon,\xi}})-\bar{b}_{1}(X_{s}^{\varepsilon},\mathscr{L}_{X_{s}^{\varepsilon}})\right)\mathrm{d}s\right|^{p}\right). \tag{22}$$

In the following, we aim to estimate the right-hand part of the inequality above. By Proposition 4.5, there exists a $\Psi(x, \mu, y, \nu)$ belonging to $C^{2,(1,1),2,(1,1)}(\mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m), \mathbb{R}^n)$ such that

$$-\mathcal{L}\Psi(x, \mu, y, \nu) = b_1(x, \mu, y, \nu) - \bar{b}_1(x, \mu).$$

Applying the Itô formula to $\Psi(X_t^{\varepsilon},\mathscr{L}_{X_t^{\varepsilon}},Y_t^{\varepsilon,y_0,\mathscr{L}_{\xi}},\mathscr{L}_{Y_t^{\varepsilon,\xi}})$, we have

$$\begin{split} &\Psi(X_t^{\varepsilon}, \mathscr{L}_{X_t^{\varepsilon}}, Y_t^{\varepsilon, y_0, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_t^{\varepsilon, \xi}}) = \Psi(\varrho, \mathscr{L}_{\varrho}, y_0, \mathscr{L}_{\xi}) \\ &+ \int_0^t \left\langle b_1(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}, Y_s^{\varepsilon, y_0, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_s^{\varepsilon, \xi}}), \partial_x \Psi(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}, Y_s^{\varepsilon, y_0, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_s^{\varepsilon, \xi}}) \right\rangle \mathrm{d}s \end{split}$$

$$+ \int_{0}^{t} \left\langle \partial_{x} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}), \sigma_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}) dB_{s} \right\rangle$$

$$+ \frac{1}{2} \int_{0}^{t} Tr \left[\sigma_{1} \sigma_{1}^{*}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}) \partial_{x}^{2} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) \right] ds$$

$$+ \tilde{\mathbb{E}} \int_{0}^{t} \left\langle b_{1}(\tilde{X}_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, \tilde{Y}_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}), \partial_{\mu} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) (\tilde{X}_{s}^{\varepsilon}) \right\rangle ds$$

$$+ \frac{1}{2} \tilde{\mathbb{E}} \int_{0}^{t} Tr \left[\sigma_{1} \sigma_{1}^{*}(\tilde{X}_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}) \partial_{x}^{*} \partial_{\mu} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) (\tilde{X}_{s}^{\varepsilon}) \right] ds$$

$$+ \frac{1}{\varepsilon} \int_{0}^{t} \left\langle b_{2}(Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}), \partial_{y} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) \right\rangle ds$$

$$+ \frac{1}{2\varepsilon} \int_{0}^{t} Tr \left[\sigma_{2} \sigma_{2}^{*}(Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) \partial_{y}^{2} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) \right) ds$$

$$+ \frac{1}{\sqrt{\varepsilon}} \int_{0}^{t} \left\langle \partial_{y} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon,\xi}}, Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) \partial_{y}^{2} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) \right] ds$$

$$+ \frac{1}{\varepsilon} \tilde{\mathbb{E}} \int_{0}^{t} \left\langle \partial_{y} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon,\xi}}), \partial_{\nu} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) dW_{s} \right\rangle$$

$$+ \frac{1}{\varepsilon} \tilde{\mathbb{E}} \int_{0}^{t} \left\langle b_{2}(\tilde{Y}_{s}^{\varepsilon,\xi}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}), \partial_{\nu} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) \tilde{Y}_{s}^{\varepsilon,\xi}) \right] ds$$

$$= \Psi(\varrho, \mathcal{L}_{\varrho}, y_{0}, \mathcal{L}_{\xi}) + N_{t}^{\varepsilon,1} + \frac{1}{\sqrt{\varepsilon}} N_{t}^{\varepsilon,2}$$

$$+ \tilde{\mathbb{E}} \int_{0}^{t} \left\langle b_{1}(\tilde{X}_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, \tilde{Y}_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) \partial_{\sigma} \partial_{\mu} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}}) \tilde{X}_{s}^{\varepsilon,\xi}) \right\rangle ds$$

$$+ \frac{1}{2} \tilde{\mathbb{E}} \int_{0}^{t} \left\langle b_{1}(\tilde{X}_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, \tilde{X}_{s}^{\varepsilon,y_{0},\mathcal{L}_$$

where the process $(\tilde{X}_s^{\varepsilon}, \tilde{Y}_s^{\varepsilon,y_0,\mathcal{L}_{\xi}}, \tilde{Y}_s^{\varepsilon,\xi})$ represents a copy of original process $(X_s^{\varepsilon}, Y_s^{\varepsilon,y_0,\mathcal{L}_{\xi}}, Y_s^{\varepsilon,\xi})$, constructed on a probability space $(\tilde{\Omega}, \tilde{\mathscr{F}}, \{\tilde{\mathscr{F}}_t\}_{t\in[0,T]}, \tilde{\mathbb{P}})$, which is an exact copy of the original probability space $(\Omega, \mathscr{F}, \{\mathscr{F}_t\}_{t\in[0,T]}, \mathbb{P})$,

$$N_{t}^{\varepsilon,1} := \int_{0}^{t} \left\langle \partial_{x} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}}), \sigma_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}) dB_{s} \right\rangle,$$

$$N_{t}^{\varepsilon,2} := \int_{0}^{t} \left\langle \partial_{y} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}}), \sigma_{2}(Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}}) dW_{s} \right\rangle,$$

and

$$\mathcal{L}_{1}\Psi(x,\mu,y,\nu) := \langle b_{1}(x,\mu,y,\nu), \partial_{x}\Psi(x,\mu,y,\nu) \rangle + \frac{1}{2}Tr[\sigma_{1}\sigma_{1}^{*}(x,\mu)\partial_{x}^{2}\Psi(x,\mu,y,\nu)].$$
₁₆

Then we have that

$$\begin{split} & \mathbb{E}\left(\sup_{0\leqslant t\leqslant T}\left|\int_{0}^{t}\left(b_{1}(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}},Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{s}^{\varepsilon,\xi}})-\bar{b}_{1}(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}})\right)\mathrm{d}s\right|^{p}\right)\\ & = & \mathbb{E}\left(\sup_{0\leqslant t\leqslant T}\left|\int_{0}^{t}\mathcal{L}\Psi(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}},Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{s}^{\varepsilon,\xi}})\mathrm{d}s\right|^{p}\right)\\ & \leqslant & C\varepsilon^{p}\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}\left|\Psi(X_{t}^{\varepsilon},\mathcal{L}_{X_{t}^{\varepsilon}},Y_{t}^{\varepsilon,y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{s}^{\varepsilon,\xi}})\right|^{p}\right)+C\varepsilon^{p}\mathbb{E}\left|\Psi(\varrho,\mathcal{L}_{\varrho},y_{0},\mathcal{L}_{\xi})\right|^{p}\\ & +C\varepsilon^{p}\mathbb{E}\tilde{\mathbb{E}}\left(\sup_{0\leqslant t\leqslant T}\left|\int_{0}^{t}\left\langle b_{1}(\tilde{X}_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}},\tilde{Y}_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{s}^{\varepsilon,\xi}}),\right.\\ & \left.\partial_{\mu}\Psi(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}},Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{s}^{\varepsilon,\xi}})(\tilde{X}_{s}^{\varepsilon})\right\rangle\mathrm{d}s\right|^{p}\right)\\ & +C\varepsilon^{p}\mathbb{E}\tilde{\mathbb{E}}\left(\sup_{0\leqslant t\leqslant T}\left|\int_{0}^{t}Tr\left[\sigma_{1}\sigma_{1}^{*}(\tilde{X}_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}})\right.\\ & \left.\partial_{\bar{x}}\partial_{\mu}\Psi(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}},Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{s}^{\varepsilon,\xi}})(\tilde{X}_{s}^{\varepsilon})\right]\mathrm{d}s\right|^{p}\right)\\ & +C\varepsilon^{p}\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}\left|\int_{0}^{t}\mathcal{L}_{1}\Psi(X_{s}^{\varepsilon},\mathcal{L}_{X_{s}^{\varepsilon}},Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{s}^{\varepsilon,\xi}})\mathrm{d}s\right|^{p}\right)\\ & +C\varepsilon^{p}\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}\left|N_{t}^{\varepsilon,1}\right|^{p}\right)+C\varepsilon^{\frac{p}{2}}\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}\left|N_{t}^{\varepsilon,2}\right|^{p}\right)\\ & = \sum_{i=1}^{7}\mathcal{J}_{i}. \end{split}$$

For \mathcal{J}_1 and \mathcal{J}_2 , by Proposition 4.5 and Lemma 4.1, we have

$$\mathcal{J}_{1} \leqslant C\varepsilon^{p} \left\{ 1 + \mathbb{E} \left(\sup_{0 \leqslant t \leqslant T} |Y_{t}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}|^{p} \right) + \mathbb{E} \left(\sup_{0 \leqslant t \leqslant T} |Y_{t}^{\varepsilon, \xi}|^{p} \right) \right\}
\leqslant C_{T}\varepsilon^{p-1} \left\{ 1 + |y_{0}|^{3p} + \mathbb{E}|\xi|^{3p} \right\},$$

and

$$\mathcal{J}_2 \leqslant C\varepsilon^p \{ 1 + |y_0|^p + \mathbb{E}|\xi|^p \}.$$

For \mathcal{J}_3 and \mathcal{J}_4 , based on the Hölder inequality, Remark 2.5 and Proposition 4.5, one can obtain that

$$\begin{split} \mathscr{J}_{3} &\leqslant C\varepsilon^{p}T^{p-1}\mathbb{E}\tilde{\mathbb{E}}\Big[\int_{0}^{T}|b_{1}(\tilde{X}_{s}^{\varepsilon},\mathscr{L}_{X_{s}^{\varepsilon}},\tilde{Y}_{s}^{\varepsilon,y_{0},\mathscr{L}_{\xi}},\mathscr{L}_{Y_{s}^{\varepsilon,\xi}})|^{p} \\ &\cdot \|\partial_{\mu}\Psi(X_{s}^{\varepsilon},\mathscr{L}_{X_{s}^{\varepsilon}},Y_{s}^{\varepsilon,y_{0},\mathscr{L}_{\xi}},\mathscr{L}_{Y_{s}^{\varepsilon,\xi}})\|_{L^{2}(\mu)}^{p}\mathrm{d}s\Big] \\ &\leqslant C\varepsilon^{p}T^{p}\left\{1+\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|X_{t}^{\varepsilon}|^{2p}\right)+\sup_{0\leqslant t\leqslant T}\mathbb{E}|Y_{t}^{\varepsilon,y_{0},\mathscr{L}_{\xi}}|^{2p}+\sup_{0\leqslant t\leqslant T}\mathbb{E}|Y_{t}^{\varepsilon,\xi}|^{2p}\right\}, \end{split}$$

and

$$\mathcal{J}_{4} \leqslant C\varepsilon^{p}T^{p-1}\mathbb{E}\tilde{\mathbb{E}}\int_{0}^{T} \|\sigma_{1}\sigma_{1}^{*}(\tilde{X}_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}})\|^{p} \\
\cdot \|\partial_{\tilde{x}}\partial_{\mu}\Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon,\xi}})\|_{L^{2}(\mu)}^{p} \mathrm{d}s \\
\leqslant C\varepsilon^{p}T^{p}\left\{1 + \mathbb{E}\left(\sup_{0 \leqslant t \leqslant T} |X_{t}^{\varepsilon}|^{3p}\right) + \sup_{0 \leqslant t \leqslant T} \mathbb{E}|Y_{t}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}|^{3p} + \sup_{0 \leqslant t \leqslant T} \mathbb{E}|Y_{t}^{\varepsilon,\xi}|^{3p}\right\}.$$

By similar arguments for \mathcal{J}_3 and \mathcal{J}_4 , we obtain

$$\mathscr{J}_5 \leqslant C\varepsilon^p T^p \left\{ 1 + \mathbb{E} \left(\sup_{0 \leqslant t \leqslant T} |X_t^{\varepsilon}|^{3p} \right) + \sup_{0 \leqslant t \leqslant T} \mathbb{E} |Y_t^{\varepsilon, y_0, \mathscr{L}_{\xi}}|^{3p} + \sup_{0 \leqslant t \leqslant T} \mathbb{E} |Y_t^{\varepsilon, \xi}|^{3p} \right\}.$$

For \mathcal{J}_6 and \mathcal{J}_7 , by the Hölder inequality, the Burkholder-Davis-Gundy inequality, Remark 2.5 and Proposition 4.5, we have

$$\mathcal{J}_{6} + \mathcal{J}_{7} \leqslant C\varepsilon^{p} \mathbb{E} \left[\int_{0}^{T} \|\partial_{x} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}}) \|^{2} \|\sigma_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}) \|^{2} ds \right]^{\frac{p}{2}} \\
+ C\varepsilon^{\frac{p}{2}} \mathbb{E} \left[\int_{0}^{T} \|\partial_{y} \Psi(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}}) \|^{2} \|\sigma_{2}(Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}}) \|^{2} ds \right]^{\frac{p}{2}} \\
\leqslant C\varepsilon^{p} T^{\frac{p}{2} - 1} \int_{0}^{T} \left(1 + \mathbb{E} |X_{s}^{\varepsilon}|^{2p} + \mathbb{E} |Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}|^{2p} + \mathbb{E} |Y_{s}^{\varepsilon, \xi}|^{2p} \right) ds \\
+ C\varepsilon^{\frac{p}{2}} T^{\frac{p}{2} - 1} \int_{0}^{T} \left(1 + \mathbb{E} |Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}|^{p} + \mathbb{E} |Y_{s}^{\varepsilon, \xi}|^{p} \right) ds \\
\leqslant C(\varepsilon^{p} + \varepsilon^{\frac{p}{2}}) T^{\frac{p}{2}} \left\{ 1 + \mathbb{E} \left(\sup_{0 \leqslant t \leqslant T} |X_{t}^{\varepsilon}|^{2p} \right) + \sup_{0 \leqslant t \leqslant T} \mathbb{E} |Y_{t}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}|^{2p} + \sup_{0 \leqslant t \leqslant T} \mathbb{E} |Y_{t}^{\varepsilon, \xi}|^{2p} \right\}. \tag{24}$$

Collecting the above deduction, by Lemma 4.1 we conclude that

$$\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}\left|\int_{0}^{t}\left(b_{1}(X_{s}^{\varepsilon},\mathscr{L}_{X_{s}^{\varepsilon}},Y_{s}^{\varepsilon,y_{0},\mathscr{L}_{\xi}},\mathscr{L}_{Y_{s}^{\varepsilon,\xi}})-\bar{b}_{1}(X_{s}^{\varepsilon},\mathscr{L}_{X_{s}^{\varepsilon}})\right)\mathrm{d}s\right|^{p}\right)$$

$$\leqslant C_{T}\varepsilon^{\frac{p}{2}}(1+\mathbb{E}|\varrho|^{3p}+|y_{0}|^{3p}+\mathbb{E}|\xi|^{3p}),$$
(25)

which together with (22) implies that

$$\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|X_t^{\varepsilon}-\bar{X}_t|^p\right)\leqslant C_T\varepsilon^{\frac{p}{2}}(1+\mathbb{E}|\varrho|^{3p}+|y_0|^{3p}+\mathbb{E}|\xi|^{3p}).$$

The proof is complete.

5. Proof of theorem 3.2

In this section, we give the proof of Theorem 3.2. The proof consists of three components. The first subsection (Subsection 5.1), we construct the auxiliary process ϑ^{ε} and demonstrate the difference $U^{\varepsilon}_{\cdot} - \vartheta^{\varepsilon}_{\cdot}$ converges to zero in the L^2 sense. In the subsequent part (Subsection 5.2), we verify the tightness of $\vartheta^{\varepsilon}_{\cdot}$ within $C([0,T],\mathbb{R}^n)$. In the final segment (Subsection 5.3), we establish the weak convergence of $\vartheta^{\varepsilon}_{\cdot}$ to the solution U_{\cdot} of equation (5) in $C([0,T];\mathbb{R}^n)$, subsequently confirming the validity of Theorem 3.2.

5.1. Construction of the auxiliary process. In this subsection, we construct an auxiliary process ϑ^{ε} and obtain that the difference $U^{\varepsilon}_{\cdot} - \vartheta^{\varepsilon}_{\cdot}$ converges to zero in the L^2 sense. First of all, by (1) and (3), the deviation process $\{U^{\varepsilon}_{t} := \frac{X^{\varepsilon}_{t} - \bar{X}_{t}}{\sqrt{\varepsilon}}\}_{t \geqslant 0}$ satisfies that

$$\begin{cases}
dU_t^{\varepsilon} = \frac{1}{\sqrt{\varepsilon}} [b_1(X_t^{\varepsilon}, \mathcal{L}_{X_t^{\varepsilon}}, Y_t^{\varepsilon, y_0, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_t^{\varepsilon, \xi}}) - \bar{b}_1(X_t^{\varepsilon}, \mathcal{L}_{X_t^{\varepsilon}})] dt \\
+ \frac{1}{\sqrt{\varepsilon}} [\bar{b}_1(X_t^{\varepsilon}, \mathcal{L}_{X_t^{\varepsilon}}) - \bar{b}_1(\bar{X}_t, \mathcal{L}_{\bar{X}_t})] dt + \frac{1}{\sqrt{\varepsilon}} [\sigma_1(X_t^{\varepsilon}, \mathcal{L}_{X_t^{\varepsilon}}) - \sigma_1(\bar{X}_t, \mathcal{L}_{\bar{X}_t})] dB_t, (26) \\
U_0^{\varepsilon} = 0.
\end{cases}$$

According to Theorem 3.1, there exists a constant C_T independent of ε such that

$$\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|U_t^{\varepsilon}|^4\right)\leqslant C_T(1+\mathbb{E}|\varrho|^{12}+|y_0|^{12}+\mathbb{E}|\xi|^{12}). \tag{27}$$

For any $\varepsilon \in (0,1)$, we consider the following auxiliary process $\vartheta_t^{\varepsilon}$:

$$\begin{cases}
d\vartheta_{t}^{\varepsilon} = \frac{1}{\sqrt{\varepsilon}} \left[b_{1}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}, Y_{t}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{t}^{\varepsilon, \xi}}) - \bar{b}_{1}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}) \right] dt \\
+ \partial_{x} \bar{b}_{1}(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}}) \vartheta_{t}^{\varepsilon} dt + \mathbb{E} \left[\partial_{\mu} \bar{b}_{1}(u, \mathcal{L}_{\bar{X}_{t}})(\bar{X}_{t}) \vartheta_{t}^{\varepsilon} \right] |_{u = \bar{X}_{t}} dt \\
+ \partial_{x} \sigma_{1}(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}}) \vartheta_{t}^{\varepsilon} dB_{t} + \mathbb{E} \left[\partial_{\mu} \sigma_{1}(u, \mathcal{L}_{\bar{X}_{t}})(\bar{X}_{t}) \vartheta_{t}^{\varepsilon} \right] |_{u = \bar{X}_{t}} dB_{t}, \\
\vartheta_{0}^{\varepsilon} = 0.
\end{cases} (28)$$

Proposition 5.1. Assume that $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{b_1}^2)$ and $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^3)$ hold, and $\sigma_1(\cdot,\cdot) \in C_b^{2,(1,1)}(\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m),\mathbb{R}^{n\times n})$. It holds that

$$\lim_{\varepsilon \to 0} \mathbb{E} \left(\sup_{0 \leqslant t \leqslant T} |U_t^{\varepsilon} - \vartheta_t^{\varepsilon}|^2 \right) = 0.$$

Proof. Set $\chi_t^{\varepsilon} := U_t^{\varepsilon} - \vartheta_t^{\varepsilon}$, we obtain

$$d\chi_{t}^{\varepsilon} = dU_{t}^{\varepsilon} - d\vartheta_{t}^{\varepsilon}$$

$$= \frac{1}{\sqrt{\varepsilon}} \Big[\bar{b}_{1}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}) - \bar{b}_{1}(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}}) - \partial_{x}\bar{b}_{1}(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}})\sqrt{\varepsilon}U_{t}^{\varepsilon}$$

$$-\mathbb{E}[\partial_{\mu}\bar{b}_{1}(u, \mathcal{L}_{\bar{X}_{t}})(\bar{X}_{t})\sqrt{\varepsilon}U_{t}^{\varepsilon}]|_{u=\bar{X}_{t}} \Big] dt$$

$$+\partial_{x}\bar{b}_{1}(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}})\chi_{t}^{\varepsilon}dt + \mathbb{E}[\partial_{\mu}\bar{b}_{1}(u, \mathcal{L}_{\bar{X}_{t}})(\bar{X}_{t})\chi_{t}^{\varepsilon}]|_{u=\bar{X}_{t}}dt$$

$$+\frac{1}{\sqrt{\varepsilon}} \Big[\sigma_{1}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}) - \sigma_{1}(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}}) - \partial_{x}\sigma_{1}(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}})\sqrt{\varepsilon}U_{t}^{\varepsilon}$$

$$-\mathbb{E}[\partial_{\mu}\sigma_{1}(u, \mathcal{L}_{\bar{X}_{t}})(\bar{X}_{t})\sqrt{\varepsilon}U_{t}^{\varepsilon}]|_{u=\bar{X}_{t}} \Big] dB_{t}$$

$$+\partial_{x}\sigma_{1}(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}})\chi_{t}^{\varepsilon}dB_{t} + \mathbb{E}[\partial_{\mu}\sigma_{1}(u, \mathcal{L}_{\bar{X}_{t}})(\bar{X}_{t})\chi_{t}^{\varepsilon}]|_{u=\bar{X}_{t}}dB_{t}.$$

By the Hölder inequality and the Burkholder-Davis-Gundy inequality, it holds that

$$\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|\chi_{t}^{\varepsilon}|^{2}\right)$$

$$\leqslant \frac{6T}{\varepsilon}\mathbb{E}\int_{0}^{T}\left|\bar{b}_{1}(X_{t}^{\varepsilon},\mathcal{L}_{X_{t}^{\varepsilon}})-\bar{b}_{1}(\bar{X}_{t},\mathcal{L}_{\bar{X}_{t}})-\partial_{x}\bar{b}_{1}(\bar{X}_{t},\mathcal{L}_{\bar{X}_{t}})\sqrt{\varepsilon}U_{t}^{\varepsilon}\right.$$

$$\left.-\mathbb{E}[\partial_{\mu}\bar{b}_{1}(u,\mathcal{L}_{\bar{X}_{t}})(\bar{X}_{t})\sqrt{\varepsilon}U_{t}^{\varepsilon}]|_{u=\bar{X}_{t}}\right|^{2}dt$$

$$+6T\mathbb{E}\int_{0}^{T}\left|\partial_{x}\bar{b}_{1}(\bar{X}_{t},\mathcal{L}_{\bar{X}_{t}})\chi_{t}^{\varepsilon}\right|^{2}dt+6T\mathbb{E}\int_{0}^{T}\left|\mathbb{E}[\partial_{\mu}\bar{b}_{1}(u,\mathcal{L}_{\bar{X}_{t}})(\bar{X}_{t})\chi_{t}^{\varepsilon}]|_{u=\bar{X}_{t}}\right|^{2}dt$$

$$+\frac{C}{\varepsilon}\mathbb{E}\int_{0}^{T} \left\|\sigma_{1}(X_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}) - \sigma_{1}(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}}) - \partial_{x}\sigma_{1}(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}})\sqrt{\varepsilon}U_{t}^{\varepsilon}\right\| -\mathbb{E}\left[\partial_{\mu}\sigma_{1}(u, \mathcal{L}_{\bar{X}_{t}})(\bar{X}_{t})\sqrt{\varepsilon}U_{t}^{\varepsilon}\right]_{u=\bar{X}_{t}}^{2} dt +C\mathbb{E}\int_{0}^{T} \left\|\partial_{x}\sigma_{1}(\bar{X}_{t}, \mathcal{L}_{\bar{X}_{t}})\chi_{t}^{\varepsilon}\right\|^{2} dt + C\mathbb{E}\int_{0}^{T} \left\|\mathbb{E}\left[\partial_{\mu}\sigma_{1}(u, \mathcal{L}_{\bar{X}_{t}})(\bar{X}_{t})\chi_{t}^{\varepsilon}\right]_{u=\bar{X}_{t}}^{2} dt \right\} =: \sum_{k=1}^{6} I_{k}^{\varepsilon}(T).$$

$$(29)$$

Next, we focus on proving the continuity and boundedness of the coefficients $\partial_x \bar{b}_1$ and $\partial_\mu \bar{b}_1$. According to Lemma 4.4, we have for any $x \in \mathbb{R}^n$, $\mu \in \mathcal{P}_2(\mathbb{R}^n)$

$$\bar{b}_1(x,\mu) = \lim_{t \to \infty} \mathbb{E}b_1(x,\mu,Y_t^{y_0,\mathcal{L}_\xi},\mathcal{L}_{Y_t^\xi}).$$

Then we have

$$\partial_x \bar{b}_1(x,\mu) = \lim_{t \to \infty} \mathbb{E} \left[\partial_x b_1(x,\mu, Y_t^{y_0, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_t^{\xi}}) \right]$$

and

$$\partial_{\mu}\bar{b}_{1}(x,\mu)(\tilde{x}) = \lim_{t \to \infty} \mathbb{E}\left[\partial_{\mu}b_{1}(x,\mu,Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}})(\tilde{x})\right].$$

For any sequence $\{(\mu_n, \tilde{x}_n)\}_{n\geqslant 1} \subset \mathcal{P}_2(\mathbb{R}^n) \times \mathbb{R}^n$ with $\mu_n \to \mu$ in \mathbb{W}_2 and $|\tilde{x}_n - \tilde{x}| \to 0$ as $n \to \infty$, it holds that

$$\sup_{t\geqslant 0} \mathbb{E}\|\partial_{\mu}b_{1}(x,\mu_{n},Y_{t}^{y_{0},\mathscr{L}_{\xi}},\mathscr{L}_{Y_{t}^{\xi}})(\tilde{x}_{n}) - \partial_{\mu}b_{1}(x,\mu,Y_{t}^{y_{0},\mathscr{L}_{\xi}},\mathscr{L}_{Y_{t}^{\xi}})(\tilde{x})\|$$

$$\leqslant \sup_{y\in\mathbb{R}^{m},\nu\in\mathcal{P}_{2}(\mathbb{R}^{m})}\|\partial_{\mu}b_{1}(x,\mu_{n},y,\nu)(\tilde{x}_{n}) - \partial_{\mu}b_{1}(x,\mu,y,\nu)(\tilde{x})\|.$$

By the uniform continuity of $\partial_{\mu}b_1$, we arrive at

$$\lim_{n \to \infty} \|\partial_{\mu} \bar{b}_1(x, \mu_n)(\tilde{x}_n) - \partial_{\mu} \bar{b}_1(x, \mu)(\tilde{x})\| = 0.$$

By similar argument, for any sequence $\{(x_n, \mu_n)\}_{n\geqslant 1} \subset \mathbb{R}^n \times \mathcal{P}_2(\mathbb{R}^n)$ satisfying $|x_n-x|\to 0$ and $\mu_n\to \mu$ in \mathbb{W}_2 as $n\to\infty$, we obtain

$$\lim_{n \to \infty} \|\partial_x \bar{b}_1(x_n, \mu_n) - \partial_x \bar{b}_1(x, \mu)\| = 0.$$

In addition, the uniform boundedness of $\partial_x b_1$ and $\partial_\mu b_1$ implies that

$$\sup_{x \in \mathbb{R}^n, \mu \in \mathcal{P}_2(\mathbb{R}^n)} \|\partial_x \bar{b}_1(x, \mu)\| < \infty,$$

and

$$\sup_{x \in \mathbb{R}^n, \mu \in \mathcal{P}_2(\mathbb{R}^n), \tilde{x} \in \mathbb{R}^n} \|\partial_{\mu} \bar{b}_1(x, \mu)(\tilde{x})\| < \infty.$$

In the following, we estimate $I_k^{\varepsilon}(T)$, $k=1,\cdots,6$ one by one. For $I_1^{\varepsilon}(T)$, according to the mean value theorem (See [16, Lemma 4.1]), there exists a constant $r \in (0,1)$ such that

$$I_{1}^{\varepsilon}(T) = 6T\mathbb{E} \int_{0}^{T} \left| \partial_{x} \bar{b}_{1}(\bar{X}_{t} + r\sqrt{\varepsilon}U_{t}^{\varepsilon}, \mathcal{L}_{X_{t}^{\varepsilon}}) U_{t}^{\varepsilon} \right. \\ \left. + \mathbb{E} \left[\partial_{\mu} \bar{b}_{1}(u, \mathcal{L}_{\bar{X}_{t} + r\sqrt{\varepsilon}U_{t}^{\varepsilon}}) (\bar{X}_{t} + r\sqrt{\varepsilon}U_{t}^{\varepsilon}) U_{t}^{\varepsilon} \right] \right|_{u = \bar{X}_{t}}$$

$$-\partial_{x}\bar{b}_{1}(\bar{X}_{t},\mathscr{L}_{\bar{X}_{t}})U_{t}^{\varepsilon} - \mathbb{E}[\partial_{\mu}\bar{b}_{1}(u,\mathscr{L}_{\bar{X}_{t}})(\bar{X}_{t})U_{t}^{\varepsilon}]|_{u=\bar{X}_{t}}\Big|^{2}dt$$

$$\leqslant C_{T}\left\{\int_{0}^{T}\left[\mathbb{E}\|\partial_{x}\bar{b}_{1}(\bar{X}_{t}+r\sqrt{\varepsilon}U_{t}^{\varepsilon},\mathscr{L}_{X_{t}^{\varepsilon}})-\partial_{x}\bar{b}_{1}(\bar{X}_{t},\mathscr{L}_{\bar{X}_{t}})\|^{4}\right]^{\frac{1}{2}}dt$$

$$+\mathbb{E}\int_{0}^{T}\mathbb{E}\Big[|\partial_{\mu}\bar{b}_{1}(u,\mathscr{L}_{\bar{X}_{t}+r\sqrt{\varepsilon}U_{t}^{\varepsilon}})(\bar{X}_{t}+r\sqrt{\varepsilon}U_{t}^{\varepsilon})-\partial_{\mu}\bar{b}_{1}(u,\mathscr{L}_{\bar{X}_{t}})(\bar{X}_{t})|^{2}\Big]\Big|_{u=\bar{X}_{t}}dt\right\}$$

$$\cdot\left[\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|U_{t}^{\varepsilon}|^{4}\right)\right]^{\frac{1}{2}}.$$

Note that $\partial_x \bar{b}_1(\cdot,\cdot)$ and $\partial_\mu \bar{b}_1(x,\cdot)(\cdot)$ are jointly continuous. Thus, by the boundedness of $\partial_x \bar{b}_1$ and $\partial_\mu \bar{b}_1$, and (27), it holds that

$$\lim_{\varepsilon \to 0} I_1^{\varepsilon}(T) = 0. \tag{30}$$

Using similar arguments as above, we obtain

$$\lim_{\varepsilon \to 0} I_4^{\varepsilon}(T) = 0. \tag{31}$$

From the boundedness of $\partial_x \bar{b}_1$, $\partial_\mu \bar{b}_1$, $\partial_x \sigma_1$ and $\partial_\mu \sigma_1$, it follows that

$$I_2^{\varepsilon}(T) + I_3^{\varepsilon}(T) + I_5^{\varepsilon}(T) + I_6^{\varepsilon}(T) \leqslant C_T \int_0^T \mathbb{E}\left(\sup_{0 \leqslant s \leqslant t} |\chi_s^{\varepsilon}|^2\right) dt.$$
 (32)

Finally, by (29), (32) and the Gronwall inequality, it immediately leads to

$$\mathbb{E}\left(\sup_{0 \le t \le T} |\chi_t^{\varepsilon}|^2\right) \le C_T(I_1^{\varepsilon}(T) + I_4^{\varepsilon}(T)),$$

which together with (30) and (31) yields that

$$\lim_{\varepsilon \to 0} \mathbb{E} \left(\sup_{0 \leqslant t \leqslant T} |\chi_t^\varepsilon|^2 \right) = 0.$$

The proof is complete.

5.2. **Tightness.** In this subsection, we intend to prove the tightness of the distribution of the solution ϑ^{ε} to Eq.(28) in $C([0,T],\mathbb{R}^n)$.

First of all, we write

$$\vartheta_t^\varepsilon = J_1^\varepsilon(t) + J_2^\varepsilon(t) + J_3^\varepsilon(t) + J_4^\varepsilon(t) + J_5^\varepsilon(t).$$

where

$$\begin{split} J_1^{\varepsilon}(t) &:= \frac{1}{\sqrt{\varepsilon}} \int_0^t [b_1(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}, Y_s^{\varepsilon, y_0, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_s^{\varepsilon, \xi}}) - \bar{b}_1(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}})] \mathrm{d}s, \\ J_2^{\varepsilon}(t) &:= \int_0^t \partial_x \bar{b}_1(\bar{X}_s, \mathscr{L}_{\bar{X}_s}) \vartheta_s^{\varepsilon} \mathrm{d}s, \\ J_3^{\varepsilon}(t) &:= \int_0^t \mathbb{E}[\partial_{\mu} \bar{b}_1(u, \mathscr{L}_{\bar{X}_s})(\bar{X}_s) \vartheta_s^{\varepsilon}]|_{u = \bar{X}_s} \mathrm{d}s, \\ J_4^{\varepsilon}(t) &:= \int_0^t \partial_x \sigma_1(\bar{X}_s, \mathscr{L}_{\bar{X}_s}) \vartheta_s^{\varepsilon} \mathrm{d}B_s, \end{split}$$

$$J_5^{\varepsilon}(t) := \int_0^t \mathbb{E}[\partial_{\mu}\sigma_1(u, \mathscr{L}_{\bar{X}_s})(\bar{X}_s)\vartheta_s^{\varepsilon}]|_{u=\bar{X}_s} dB_s.$$

Lemma 5.2. Suppose that $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{b_1}^2)$ and $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^3)$ hold, and $\sigma_1(\cdot,\cdot) \in C_b^{2,(1,1)}(\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m),\mathbb{R}^{n\times n})$. Then $\{\Pi^{\varepsilon}:=(\vartheta^{\varepsilon},J_1^{\varepsilon},J_2^{\varepsilon},J_3^{\varepsilon},J_4^{\varepsilon},J_5^{\varepsilon}),\varepsilon\in(0,1)\}$ is tight in $C([0,T];\mathbb{R}^{6n})$.

Proof. In view of the term $J_1^{\varepsilon}(t)$, it can be deduced from (23) that

$$\begin{split} J_{1}^{\varepsilon}(t) &= \frac{1}{\sqrt{\varepsilon}} \int_{0}^{t} \left[b_{1}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_{s}^{\varepsilon, \xi}}) - \bar{b}_{1}(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}) \right] \mathrm{d}s \\ &= -\frac{1}{\sqrt{\varepsilon}} \int_{0}^{t} \mathcal{L}\Psi(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_{s}^{\varepsilon, \xi}}) \mathrm{d}s \\ &= \sqrt{\varepsilon} \Big\{ - \Psi(X_{t}^{\varepsilon}, \mathscr{L}_{X_{t}^{\varepsilon}}, Y_{t}^{\varepsilon, y_{0}, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_{t}^{\varepsilon, \xi}}) + \Psi(\varrho, \mathscr{L}_{\varrho}, y_{0}, \mathscr{L}_{\xi}) \\ &+ \tilde{\mathbb{E}} \int_{0}^{t} \left\langle b_{1}(\tilde{X}_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}, \tilde{Y}_{s}^{\varepsilon, y_{0}, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_{s}^{\varepsilon, \xi}}), \partial_{\mu} \Psi(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_{s}^{\varepsilon, \xi}})(\tilde{X}_{s}^{\varepsilon}) \right\rangle \mathrm{d}s \\ &+ \frac{1}{2} \tilde{\mathbb{E}} \int_{0}^{t} Tr \left[\sigma_{1} \sigma_{1}^{*}(\tilde{X}_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}) \partial_{\tilde{x}} \partial_{\mu} \Psi(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_{s}^{\varepsilon, \xi}})(\tilde{X}_{s}^{\varepsilon}) \right] \mathrm{d}s \\ &+ \int_{0}^{t} \mathcal{L}_{1} \Psi(X_{s}^{\varepsilon}, \mathscr{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_{s}^{\varepsilon, \xi}}) \mathrm{d}s + N_{t}^{\varepsilon, 1} \Big\} + N_{t}^{\varepsilon, 2} \\ &=: J_{11}^{\varepsilon}(t) + N_{t}^{\varepsilon, 2}. \end{split}$$

From analogous deduction to that in Subsection 4.4, it follows that

$$\sup_{\varepsilon \in (0,1)} \mathbb{E} \left(\sup_{t \in [0,T]} |J_{11}^{\varepsilon}(t)|^4 \right) \leqslant C_T \varepsilon (1 + \mathbb{E}|\varrho|^{12} + |y_0|^{12} + \mathbb{E}|\xi|^{12}), \tag{33}$$

which implies that J_{11}^{ε} converges to 0 in probability in $C([0,T];\mathbb{R}^n)$. By Lemma 20.3 in [14, P.185], to establish the tightness of $\{\Pi^{\varepsilon}\}$, it is sufficient to verify the following conditions for i=2,3,4,5:

$$\sup_{\varepsilon \in (0,1)} \mathbb{E} \left(\sup_{t \in [0,T]} |J_i^{\varepsilon}(t)|^4 \right) < \infty, \tag{34}$$

and

$$\sup_{\varepsilon \in (0,1)} \mathbb{E}|J_i^{\varepsilon}(t_2) - J_i^{\varepsilon}(t_1)|^4 \leqslant C_T|t_2 - t_1|^2, \tag{35}$$

where the same conditions remain valid if $J_i^{\varepsilon}(t)$ is replaced by $N_t^{\varepsilon,2}$.

We divide the remaining proof into two steps.

Step 1. We aim to establish (34).

Firstly, for $N_t^{\varepsilon,2}$, we deduce from (24) and Lemma 4.1 that

$$\sup_{\varepsilon \in (0,1)} \mathbb{E} \left(\sup_{t \in [0,T]} |N_t^{\varepsilon,2}|^4 \right) \leqslant C_T (1 + |y_0|^{12} + \mathbb{E}|\xi|^{12}).$$

Secondly, in order to investigate the uniform estimates of $\{J_i^{\varepsilon}\}$, i=2,3,4,5, we need to prove that

$$\sup_{\varepsilon \in (0,1)} \mathbb{E} \left(\sup_{t \in [0,T]} |\vartheta_t^{\varepsilon}|^4 \right) \leqslant C_T (1 + \mathbb{E}|\varrho|^{12} + |y_0|^{12} + \mathbb{E}|\xi|^{12}). \tag{36}$$

In fact, using the uniform boundedness of $\partial_x \bar{b}_1$, $\partial_\mu \bar{b}_1$, $\partial_x \sigma_1$ and $\partial_\mu \sigma_1$, together with the Hölder inequality and the Burkholder-Davis-Gundy inequality, one can conclude that

$$\mathbb{E}\left(\sup_{t\in[0,T]}|\vartheta_{t}^{\varepsilon}|^{4}\right) \leqslant C\mathbb{E}\left(\sup_{t\in[0,T]}|J_{1}^{\varepsilon}(t)|^{4}\right) + C_{T}\int_{0}^{T}\mathbb{E}|\vartheta_{s}^{\varepsilon}|^{4}\mathrm{d}s
+C\mathbb{E}\left(\int_{0}^{T}|\partial_{x}\sigma_{1}(\bar{X}_{s},\mathscr{L}_{\bar{X}_{s}})\vartheta_{s}^{\varepsilon}|^{2}\mathrm{d}s\right)^{2}
+C\mathbb{E}\left(\int_{0}^{T}|\mathbb{E}[\partial_{\mu}\sigma_{1}(u,\mathscr{L}_{\bar{X}_{s}})(\bar{X}_{s})\vartheta_{s}^{\varepsilon}]|_{u=\bar{X}_{s}}|^{2}\mathrm{d}s\right)^{2}
\leqslant C_{T}(1+\mathbb{E}|\varrho|^{12}+|y_{0}|^{12}+\mathbb{E}|\xi|^{12})+C_{T}\int_{0}^{T}\mathbb{E}|\vartheta_{s}^{\varepsilon}|^{4}\mathrm{d}s.$$

As a result, (36) is derived by applying Gronwall's inequality. Consequently, it follows that $\{J_i^{\varepsilon}\}, i = 2, 3, 4, 5 \text{ satisfy (34)}.$

Step 2. We shall demonstrate (35).

For $N_t^{\varepsilon,2}$, by the Burkholder-Davis-Gundy inequality, the Hölder inequality, Proposition 4.5, Remark 2.5 and Lemma 4.1, we obtain

$$\mathbb{E}|N_{t_{2}}^{\varepsilon,2} - N_{t_{1}}^{\varepsilon,2}|^{4} \leqslant C\mathbb{E}\left(\int_{t_{1}}^{t_{2}} \|\sigma_{2}(Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{s}^{\varepsilon,\xi}})\|^{2} ds\right)^{2}$$

$$\leqslant C(t_{2} - t_{1}) \int_{t_{1}}^{t_{2}} (1 + \mathbb{E}|Y_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}|^{4} + \mathbb{E}|Y_{s}^{\varepsilon,\xi}|^{4}) ds$$

$$\leqslant C(1 + |y_{0}|^{12} + \mathbb{E}|\xi|^{12})|t_{2} - t_{1}|^{2}.$$

Similarly, by combining (36) with the uniform boundedness of $\partial_x \bar{b}_1$, $\partial_\mu \bar{b}_1$, $\partial_x \sigma_1$ and $\partial_\mu \sigma_1$, it holds that

$$\mathbb{E}|J_i^{\varepsilon}(t_2) - J_i^{\varepsilon}(t_1)|^4 \leqslant C_T(1 + \mathbb{E}|\varrho|^{12} + |y_0|^{12} + \mathbb{E}|\xi|^{12})|t_2 - t_1|^2, \quad i = 2, \dots, 5.$$

As a consequence, the terms $N_t^{\varepsilon,2}$ and $\{J_i^{\varepsilon}\}$ $i=2,\ldots,5$ fulfill (35). The proof is complete.

5.3. The proof of Theorem 3.2. In this subsection, we complete the proof of Theorem 3.2. The main objective is to identify the weak limit of $\vartheta_t^{\varepsilon}$ in $C([0,T];\mathbb{R}^n)$.

Proposition 5.3. Assume that $(\mathbf{H}_{b_1,\sigma_1}^1)$, $(\mathbf{H}_{b_1}^2)$ and $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^3)$ hold, and $\sigma_1(\cdot,\cdot) \in C_b^{2,(1,1)}(\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m),\mathbb{R}^{n\times n})$, then the solution ϑ^{ε} of equation (28) weakly converges to the solution U of equation (5) in $C([0,T];\mathbb{R}^n)$, as $\varepsilon \to 0$.

Proof. By lemma 5.2, it is sufficient to demonstrate that there exists a subsequence $\{\varepsilon_{n_k}\}_{k\geqslant 1}$ such that $\vartheta^{\varepsilon_{n_k}}$ converges weakly to U in $C([0,T];\mathbb{R}^n)$. Let $B^{\varepsilon_{n_k}}=B$, $W^{\varepsilon_{n_k}}=W$ and assume that $\{(\vartheta^{\varepsilon_{n_k}},J_1^{\varepsilon_{n_k}},J_2^{\varepsilon_{n_k}},J_3^{\varepsilon_{n_k}},J_4^{\varepsilon_{n_k}},J_5^{\varepsilon_{n_k}},B^{\varepsilon_{n_k}},W^{\varepsilon_{n_k}})\}_{k\geqslant 1}$ convergence weakly to

 $(\vartheta, J_1, J_2, J_3, J_4, J_5, B, W)$. Then by the Skorohod representation theorem (cf.[2, Theorem C.1]), there exists a new probability space denoted by $(\hat{\Omega}, \hat{\mathscr{F}}, \hat{\mathbb{P}})$ and a sequence

$$\{(\hat{\vartheta}^{\varepsilon_{n_k}},\hat{J}_1^{\varepsilon_{n_k}},\hat{J}_2^{\varepsilon_{n_k}},\hat{J}_3^{\varepsilon_{n_k}},\hat{J}_4^{\varepsilon_{n_k}},\hat{J}_5^{\varepsilon_{n_k}},\hat{X}^{\varepsilon_{n_k}},\hat{Y}^{\varepsilon_{n_k},y_0,\mathcal{L}_{\hat{\xi}}},\hat{Y}^{\varepsilon_{n_k},\hat{\xi}},\hat{\bar{X}},\hat{B}^{\varepsilon_{n_k}},\hat{W}^{\varepsilon_{n_k}})\}_{k\geqslant 1},$$

which has same law with $\{(\vartheta^{\varepsilon_{n_k}}, J_1^{\varepsilon_{n_k}}, J_2^{\varepsilon_{n_k}}, J_3^{\varepsilon_{n_k}}, J_4^{\varepsilon_{n_k}}, J_5^{\varepsilon_{n_k}}, X^{\varepsilon_{n_k}}, Y^{\varepsilon_{n_k}, y_0, \mathscr{L}_{\xi}}, Y^{\varepsilon_{n_k}, \xi}, \bar{X}, B^{\varepsilon_{n_k}}, W^{\varepsilon_{n_k}})\}_{k\geqslant 1}$ and the following properties hold

(i) $(\hat{\vartheta}^{\varepsilon_{n_k}}, \hat{J}_1^{\varepsilon_{n_k}}, \hat{J}_2^{\varepsilon_{n_k}}, \hat{J}_3^{\varepsilon_{n_k}}, \hat{J}_4^{\varepsilon_{n_k}}, \hat{J}_5^{\varepsilon_{n_k}}, \hat{B}^{\varepsilon_{n_k}}, \hat{W}^{\varepsilon_{n_k}})$ converges to $(\hat{\vartheta}, \hat{J}_1, \hat{J}_2, \hat{J}_3, \hat{J}_4, \hat{J}_5, \hat{B}, \hat{W})$ in $C([0, T]; \mathbb{R}^{7n+m})$, $\hat{\mathbb{P}}$ -a.s., as $k \to \infty$;

(ii) $(\hat{B}^{\varepsilon_{n_k}}(\hat{\omega}), \hat{W}^{\varepsilon_{n_k}}(\hat{\omega})) = (\hat{B}(\hat{\omega}), \hat{W}(\hat{\omega}))$ for all $\hat{\omega} \in \hat{\Omega}$.

The process $(\hat{X}_s^{\varepsilon_{n_k}}, \hat{Y}_s^{\varepsilon_{n_k}, y_0, \mathscr{L}_{\hat{\xi}}}, \hat{Y}_s^{\varepsilon_{n_k}, \hat{\xi}})$ is the solution of the system (1) where ε_{n_k} substitutes for ε , (\hat{B}, \hat{W}) replaces (B, W) and $(\hat{\varrho}, \hat{\xi})$ replaces (ϱ, ξ) . Similarly, \hat{X} is the solution of (3) with \hat{B} replacing B and $\hat{\varrho}$ replacing ϱ . Here, $\hat{\varrho}$ and $\hat{\xi}$ are random variables defined on $(\hat{\Omega}, \hat{\mathscr{F}}, \hat{\mathbb{P}})$, which have the same distributions as ϱ and ξ respectively.

According to Eq.(28) satisfied by $\{(\vartheta^{\varepsilon_{n_k}}, J_1^{\varepsilon_{n_k}}, J_2^{\varepsilon_{n_k}}, J_3^{\varepsilon_{n_k}}, J_4^{\varepsilon_{n_k}}, J_5^{\varepsilon_{n_k}}, B, W)\}_{k\geqslant 1}$, it can be deduced that

$$\hat{\vartheta}^{\varepsilon_{n_k}} = \hat{J}_1^{\varepsilon_{n_k}}(t) + \hat{J}_2^{\varepsilon_{n_k}}(t) + \hat{J}_3^{\varepsilon_{n_k}}(t) + \hat{J}_4^{\varepsilon_{n_k}}(t) + \hat{J}_5^{\varepsilon_{n_k}}(t),$$

where

$$\begin{split} \hat{J}_{1}^{\varepsilon_{n_{k}}}(t) &:= \frac{1}{\sqrt{\varepsilon_{n_{k}}}} \int_{0}^{t} [b_{1}(\hat{X}_{s}^{\varepsilon_{n_{k}}}, \mathcal{L}_{\hat{X}_{s}^{\varepsilon_{n_{k}}}}, \hat{Y}_{s}^{\varepsilon_{n_{k}}, y_{0}, \mathcal{L}_{\hat{\xi}}}, \mathcal{L}_{\hat{Y}_{s}^{\varepsilon_{n_{k}}, \hat{\xi}}}) - \bar{b}_{1}(\hat{X}_{s}^{\varepsilon_{n_{k}}}, \mathcal{L}_{\hat{X}_{s}^{\varepsilon_{n_{k}}}})] \mathrm{d}s, \\ \hat{J}_{2}^{\varepsilon_{n_{k}}}(t) &:= \int_{0}^{t} \partial_{x} \bar{b}_{1}(\hat{X}_{s}, \mathcal{L}_{\hat{X}_{s}}) \hat{\vartheta}_{s}^{\varepsilon_{n_{k}}} \mathrm{d}s, \\ \hat{J}_{3}^{\varepsilon_{n_{k}}}(t) &:= \int_{0}^{t} \hat{\mathbb{E}} [\partial_{\mu} \bar{b}_{1}(u, \mathcal{L}_{\hat{X}_{s}})(\hat{X}_{s}) \hat{\vartheta}_{s}^{\varepsilon_{n_{k}}}]|_{u=\hat{X}_{s}} \mathrm{d}s, \\ \hat{J}_{4}^{\varepsilon_{n_{k}}}(t) &:= \int_{0}^{t} \partial_{x} \sigma_{1}(\hat{X}_{s}, \mathcal{L}_{\hat{X}_{s}}) \hat{\vartheta}_{s}^{\varepsilon_{n_{k}}} \mathrm{d}\hat{B}_{s}, \\ \hat{J}_{5}^{\varepsilon_{n_{k}}}(t) &:= \int_{0}^{t} \hat{\mathbb{E}} [\partial_{\mu} \sigma_{1}(u, \mathcal{L}_{\hat{X}_{s}})(\hat{X}_{s}) \hat{\vartheta}_{s}^{\varepsilon_{n_{k}}}]|_{u=\hat{X}_{s}} \mathrm{d}\hat{B}_{s}, \end{split}$$

and $\hat{\mathbb{E}}$ is the expectation with respect to $\hat{\mathbb{P}}$.

We guess that the limit \hat{J}_i , corresponding to $\hat{J}_i^{\varepsilon_{n_k}}$ for i=2,3,4,5, take the following forms $\hat{\mathbb{P}}$ -a.s.:

$$\hat{J}_{2}(t) = \int_{0}^{t} \partial_{x} \bar{b}_{1}(\hat{\bar{X}}_{s}, \mathcal{L}_{\hat{X}_{s}}) \hat{\vartheta}_{s} ds,$$

$$\hat{J}_{3}(t) = \int_{0}^{t} \hat{\mathbb{E}}[\partial_{\mu} \bar{b}_{1}(u, \mathcal{L}_{\hat{X}_{s}})(\hat{\bar{X}}_{s}) \hat{\vartheta}_{s}]|_{u=\hat{X}_{s}} ds,$$

$$\hat{J}_{4}(t) = \int_{0}^{t} \partial_{x} \sigma_{1}(\hat{\bar{X}}_{s}, \mathcal{L}_{\hat{X}_{s}}) \hat{\vartheta}_{s} d\hat{B}_{s},$$

$$\hat{J}_{5}(t) = \int_{0}^{t} \hat{\mathbb{E}}[\partial_{\mu} \sigma_{1}(u, \mathcal{L}_{\hat{X}_{s}})(\hat{\bar{X}}_{s}) \hat{\vartheta}_{s}]|_{u=\hat{X}_{s}} d\hat{B}_{s}.$$

Indeed, from (36), it follows that

$$\sup_{k\geqslant 1} \hat{\mathbb{E}} \left(\sup_{t\in[0,T]} |\hat{\vartheta}_t^{\varepsilon_{n_k}}|^4 \right) \leqslant C(1+\hat{\mathbb{E}}|\hat{\varrho}|^{12}+|y_0|^{12}+\hat{\mathbb{E}}|\hat{\xi}|^{12}).$$

Since $\hat{\vartheta}^{\varepsilon_{n_k}} \to \hat{\vartheta}$ in $C([0,T];\mathbb{R}^n)$, $\hat{\mathbb{P}}$ -a.s, the Vitali convergence theorem infers that

$$\lim_{k \to \infty} \hat{\mathbb{E}} \left(\sup_{t \in [0,T]} |\hat{\vartheta}_t^{\varepsilon_{n_k}} - \hat{\vartheta}_t|^2 \right) = 0.$$
 (37)

As for $\hat{J}_{2}^{\varepsilon_{n_k}}(t)$, combining the uniform boundedness of $\partial_x \bar{b}_1(x,\mu)$ and (37), we conclude that

$$\lim_{k \to \infty} \hat{\mathbb{E}} \left(\sup_{t \in [0,T]} |\hat{J}_{2}^{\varepsilon_{n_{k}}}(t) - \hat{J}_{2}(t)|^{2} \right) = \lim_{k \to \infty} \hat{\mathbb{E}} \left(\sup_{t \in [0,T]} \left| \int_{0}^{t} \partial_{x} \bar{b}_{1} (\hat{\bar{X}}_{s}, \mathcal{L}_{\hat{X}_{s}}) (\hat{\vartheta}_{s}^{\varepsilon_{n_{k}}} - \hat{\vartheta}_{s}) ds \right|^{2} \right) \\
\leqslant C_{T} \lim_{k \to \infty} \hat{\mathbb{E}} \left(\sup_{t \in [0,T]} |\hat{\vartheta}_{t}^{\varepsilon_{n_{k}}} - \hat{\vartheta}_{t}|^{2} \right) = 0.$$

For $\hat{J}_3^{\varepsilon_{n_k}}(t)$, owing to the uniform boundedness of $\partial_{\mu}\bar{b}_1(x,\mu)(\tilde{x})$ and (37), it can be deduced that

$$\lim_{k \to \infty} \sup_{t \in [0,T]} |\hat{J}_3^{\varepsilon_{n_k}}(t) - \hat{J}_3(t)|^2 = \lim_{k \to \infty} \sup_{t \in [0,T]} \left| \int_0^t \hat{\mathbb{E}}[\partial_{\mu} \bar{b}_1(u, \mathcal{L}_{\hat{X}_s})(\hat{X}_s)(\hat{\mathcal{J}}_s^{\varepsilon_{n_k}} - \hat{\mathcal{J}}_s)]|_{u = \hat{X}_s} ds \right|^2$$

$$\leqslant C_T \lim_{k \to \infty} \hat{\mathbb{E}} \left(\sup_{t \in [0,T]} |\hat{\mathcal{J}}_t^{\varepsilon_{n_k}} - \hat{\mathcal{J}}_t|^2 \right) = 0.$$

For $\hat{J}_{4}^{\varepsilon_{n_k}}(t)$ and $\hat{J}_{5}^{\varepsilon_{n_k}}(t)$, by the uniform boundedness of $\partial_x \sigma_1(x,\mu)$, $\partial_\mu \sigma_1(x,\mu)(\tilde{x})$, (37) and the Burkholder-Davis-Gundy inequality, we obtain

$$\lim_{k \to \infty} \hat{\mathbb{E}} \left(\sup_{t \in [0,T]} |\hat{J}_{4}^{\varepsilon_{n_k}}(t) - \hat{J}_{4}(t)|^2 \right) \leqslant C \lim_{k \to \infty} \hat{\mathbb{E}} \left(\int_{0}^{T} \|\partial_x \sigma_1(\hat{\bar{X}}_s, \mathcal{L}_{\hat{\bar{X}}_s})(\hat{\vartheta}_s^{\varepsilon_{n_k}} - \hat{\vartheta}_s)\|^2 ds \right)$$

$$\leqslant C_T \lim_{k \to \infty} \hat{\mathbb{E}} \left(\sup_{t \in [0,T]} |\hat{\vartheta}_t^{\varepsilon_{n_k}} - \hat{\vartheta}_t|^2 \right) = 0,$$

and

$$\lim_{k \to \infty} \hat{\mathbb{E}} \left(\sup_{t \in [0,T]} |\hat{J}_{5}^{\varepsilon_{n_{k}}}(t) - \hat{J}_{5}(t)|^{2} \right)$$

$$\leqslant C \lim_{k \to \infty} \hat{\mathbb{E}} \left(\int_{0}^{T} \left| \hat{\mathbb{E}} [\partial_{\mu} \sigma_{1}(u, \mathcal{L}_{\hat{X}_{s}})(\hat{X}_{s})(\hat{\mathcal{J}}_{s}^{\varepsilon_{n_{k}}} - \hat{\mathcal{J}}_{s})] \right|_{u = \hat{X}_{s}} \right|^{2} ds \right)$$

$$\leqslant C_{T} \lim_{k \to \infty} \hat{\mathbb{E}} \left(\sup_{t \in [0,T]} |\hat{\mathcal{J}}_{t}^{\varepsilon_{n_{k}}} - \hat{\mathcal{J}}_{t}|^{2} \right) = 0.$$

Next, we focus on identifying the limiting process of $\hat{J}_1^{\varepsilon_{n_k}}$ in $C([0,T];\mathbb{R}^n)$. According to (33), J_{11}^{ε} converges to 0 in probability in $C([0,T];\mathbb{R}^n)$. As a result, \hat{J}_1 must coincide

with the limiting process \hat{N}_t^2 of $\hat{N}_t^{\varepsilon_{n_k},2}$ in $C([0,T];\mathbb{R}^n)$, where

$$\hat{N}_t^{\varepsilon_{n_k},2} = \int_0^t \partial_y \Psi_{\sigma_2}(\hat{X}_s^{\varepsilon_{n_k}}, \mathcal{L}_{\hat{X}_s^{\varepsilon_{n_k}}}, \hat{Y}_s^{\varepsilon_{n_k},y_0,\mathcal{L}_{\hat{\xi}}}, \mathcal{L}_{\hat{Y}_s^{\varepsilon_{n_k},\hat{\xi}}}) d\hat{W}_s$$

is a continuous martingale on the filtered probability space $(\hat{\Omega}, \hat{\mathscr{F}}, \{\hat{\mathscr{F}}_t\}_{t \geq 0}, \hat{\mathbb{P}})$ with quadratic variational process

$$\langle \hat{N}_{t}^{\varepsilon_{n_{k}},2} \rangle_{t} = \int_{0}^{t} (\partial_{y} \Psi_{\sigma_{2}}) (\partial_{y} \Psi_{\sigma_{2}})^{*} (\hat{X}_{s}^{\varepsilon_{n_{k}}}, \mathcal{L}_{\hat{X}_{s}^{\varepsilon_{n_{k}}}}, \hat{Y}_{s}^{\varepsilon_{n_{k}}, y_{0}, \mathcal{L}_{\hat{\xi}}}, \mathcal{L}_{\hat{Y}_{s}^{\varepsilon_{n_{k}}, \hat{\xi}}}) \mathrm{d}s, t \in [0, T],$$

and $\hat{\mathscr{F}}_t$ represents the natural filtration generated by $\{\hat{\varrho}, \hat{\xi}, \hat{B}_s, \hat{W}_s, s \leq t\}$.

Since $\hat{N}^{\varepsilon_{n_k},2}$ converges to \hat{N}^2 , $\hat{\mathbb{P}}$ -a.s., we obtain that \hat{N}^2 is a continuous martingale on $(\hat{\Omega},\hat{\mathscr{F}},\{\hat{\mathscr{F}}_t\}_{t\geqslant 0},\hat{\mathbb{P}})$. Note that the quadratic variational process of \hat{N}_t^2 is

$$\langle \hat{N}^2 \rangle_t = \int_0^t \overline{(\partial_y \Psi_{\sigma_2})(\partial_y \Psi_{\sigma_2})^*} (\hat{\bar{X}}_s, \mathcal{L}_{\hat{X}_s}) ds, \quad t \in [0, T].$$
 (38)

(See the detailed proof in the Appendix). Thus, by the martingale representation theorem (cf.[4, Theorem 8.2]), there exists a probability space $(\check{\Omega}, \check{\mathscr{F}}, \check{\mathbb{P}})$, a filtration $\{\check{\mathscr{F}}_t\}_{t\geq 0}$ and a *n*-dimensional standard Brownian motion \check{V} defined on $(\hat{\Omega} \times \check{\Omega}, \hat{\mathscr{F}} \times \check{\mathscr{F}}, \hat{\mathbb{P}} \times \check{\mathbb{P}})$, which is adapted to $\hat{\mathscr{F}} \times \check{\mathscr{F}}$, such that

$$\hat{N}_t^2 = \int_0^t \left(\overline{(\partial_y \Psi_{\sigma_2})(\partial_y \Psi_{\sigma_2})^*} \right)^{\frac{1}{2}} (\hat{\bar{X}}_s, \mathscr{L}_{\hat{\bar{X}}_s}) d\check{V}_s, \quad t \in [0, T],$$

where

$$\hat{N}_t^2(\hat{\omega}, \check{\omega}) = \hat{N}_t^2(\hat{\omega}), \hat{\bar{X}}_s(\hat{\omega}, \check{\omega}) = \hat{\bar{X}}_s(\hat{\omega}), (\hat{\omega}, \check{\omega}) \in \hat{\Omega} \times \check{\Omega}.$$

Observe that $\langle \hat{B}, \hat{N}^{\varepsilon_{n_k}, 2} \rangle_t = 0$ and $\hat{N}^{\varepsilon_{n_k}, 2}$ converges to \hat{N}^2 almost surely under $\hat{\mathbb{P}} \times \check{\mathbb{P}}$. Consequently, $\langle \hat{B}, \hat{N}^2 \rangle_t = 0$, which indicates that \check{V} is independent of \hat{B} .

Finally, based on the above discussion, $\hat{\vartheta}$ satisfies

$$\begin{split} \mathrm{d}\hat{\vartheta}_t &= \left. \partial_x \bar{b}_1(\hat{\bar{X}}_t, \mathscr{L}_{\hat{\bar{X}}_t}) \hat{\vartheta}_t \mathrm{d}t + \bar{\mathbb{E}}[\partial_\mu \bar{b}_1(u, \mathscr{L}_{\hat{\bar{X}}_t})(\hat{\bar{X}}_t) \hat{\vartheta}_t] \right|_{u = \hat{\bar{X}}_t} \mathrm{d}t + \partial_x \sigma_1(\hat{\bar{X}}_t, \mathscr{L}_{\hat{\bar{X}}_t}) \hat{\vartheta}_t \mathrm{d}\hat{B}_t, \\ &+ \bar{\mathbb{E}}[\partial_\mu \sigma_1(u, \mathscr{L}_{\hat{\bar{X}}_t})(\hat{\bar{X}}_t) \hat{\vartheta}_t] \right|_{u = \hat{\bar{X}}_t} \mathrm{d}\hat{B}_t + \left(\overline{(\partial_y \Psi_{\sigma_2})(\partial_y \Psi_{\sigma_2})^*} \right)^{\frac{1}{2}} (\hat{\bar{X}}_t, \mathscr{L}_{\hat{\bar{X}}_t}) \mathrm{d}\check{V}_t, \ \hat{\vartheta}_0 = 0, \end{split}$$

where $\bar{\mathbb{E}}$ is the expectation with respect to $\hat{\mathbb{P}} \times \check{\mathbb{P}}$ and

$$\hat{\vartheta}_t(\hat{\omega}, \check{\omega}) = \hat{\vartheta}_t(\hat{\omega}), \quad \hat{B}_t(\hat{\omega}, \check{\omega}) = \hat{B}_t(\hat{\omega}).$$

According to [29, Theorem 2.1], Eq.(5) has the weak uniqueness. Then $\hat{\vartheta}$ coincides with the solution U of Eq.(5) in the sense of distribution. The proof is complete.

Now, we are able to complete the proof of our main result.

Proof of Theorem 3.2: Combining Proposition 5.1 and 5.3, we immediate obtain that Theorem 3.2 holds. The proof is complete.

6. Appendix

In this section, we prove (16) and (38).

6.1. **Proof of** (16). Let $C([0,\infty);\mathbb{R}^m)$ be the collection of continuous functions from $[0,\infty)$ to \mathbb{R}^m with the uniform convergence topology. Set

$$\tilde{\Omega} := C([0, \infty); \mathbb{R}^m) \times C([0, \infty); \mathcal{P}_2(\mathbb{R}^m)),
\tilde{\mathscr{F}} := \mathscr{B}\Big(C([0, \infty); \mathbb{R}^m)\Big) \times \mathscr{B}\Big(C([0, \infty); \mathcal{P}_2(\mathbb{R}^m))\Big),
\tilde{\mathscr{F}}_t := \sigma(M_r, 0 \leqslant r \leqslant t), \quad t \geqslant 0.$$

By the similar deduction to that of [22, Theorem 4.11], there exists a unique probability measure $\tilde{\mathbb{P}}$ on $(\tilde{\Omega}, \tilde{\mathscr{F}})$ such that M is a Markov process with respect to $(\tilde{\mathscr{F}}_t)$ with the transition function $\{\mathscr{L}_{Y_t^{y,\nu}} \times \delta_{\mathscr{L}_{Y_t^{\varepsilon}}} : t \geqslant 0, (y,\nu) \in \mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m)\}$. Note that

$$\mathscr{L}_{M_t}^{\tilde{\mathbb{P}}} := \tilde{\mathbb{P}} \circ (M_t)^{-1} = \mathscr{L}_{Y_t^{y,\nu}} \times \delta_{\mathscr{L}_{Y_t^{\xi}}}.$$

Thus, it holds that

$$\tilde{b}_{1,s_{0}}(x,\mu,y,\nu,s) = \hat{b}_{1}(x,\mu,y,\nu,s) - \hat{b}_{1}(x,\mu,y,\nu,s+s_{0})
= \hat{b}_{1}(x,\mu,y,\nu,s) - \mathbb{E}b_{1}(x,\mu,Y_{s+s_{0}}^{y,\nu},\mathcal{L}_{Y_{s+s_{0}}^{\xi}})
= \hat{b}_{1}(x,\mu,y,\nu,s) - \tilde{\mathbb{E}}[\tilde{\mathbb{E}}[b_{1}(x,\mu,M_{s+s_{0}})|\tilde{\mathscr{F}}_{s_{0}}]]
= \hat{b}_{1}(x,\mu,y,\nu,s) - \tilde{\mathbb{E}}[\tilde{\mathbb{E}}^{M_{s_{0}}}[b_{1}(x,\mu,M_{s})]]
= \hat{b}_{1}(x,\mu,y,\nu,s) - \mathbb{E}\hat{b}_{1}(x,\mu,Y_{s_{0}}^{y,\nu},\mathcal{L}_{Y_{s_{0}}^{\xi}},s).$$
(39)

Then we obtain

$$\partial_x \tilde{b}_{1,s_0}(x,\mu,y,\nu,s) = \partial_x \hat{b}_1(x,\mu,y,\nu,s) - \partial_x \mathbb{E} \hat{b}_1(x,\mu,Y_{s_0}^{y,\nu},\mathcal{L}_{Y_{s_0}^{\xi}},s). \tag{40}$$

By $(\mathbf{H}_{b_1}^2)$ and Lemma 4.3, we get

$$\begin{aligned} &|\partial_{x}\hat{b}_{1}(x,\mu,y_{1},\nu_{1},s) - \partial_{x}\hat{b}_{1}(x,\mu,y_{2},\nu_{2},s)| \\ &\leqslant & \mathbb{E}\|\partial_{x}b_{1}(x,\mu,Y_{s}^{y_{1},\nu_{1}},\mathcal{L}_{Y_{s}^{\xi_{1}}}) - \partial_{x}b_{1}(x,\mu,Y_{s}^{y_{2},\nu_{2}},\mathcal{L}_{Y_{s}^{\xi_{2}}})\| \\ &\leqslant & C\mathbb{E}\Big(|Y_{s}^{y_{1},\nu_{1}} - Y_{s}^{y_{2},\nu_{2}}| + \mathbb{W}_{2}(\mathcal{L}_{Y_{s}^{\xi_{1}}},\mathcal{L}_{Y_{s}^{\xi_{2}}})\Big) \\ &\leqslant & C\Big[|y_{1} - y_{2}|e^{-\frac{\beta_{1}}{2}s} + (\mathbb{E}|\xi_{1} - \xi_{2}|^{2})^{\frac{1}{2}}e^{-\frac{\beta_{1} - \beta_{2}}{2}s}\Big]. \end{aligned}$$

$$\tag{41}$$

Combining (40), (41) and Lemma 4.2, we obtain

$$\|\partial_x \tilde{b}_{1,s_0}(x,\mu,y,\nu,s)\|$$

$$\leqslant C\mathbb{E}\Big[|y-Y_{s_0}^{y,\nu}|e^{-\frac{\beta_1}{2}s}+(\mathbb{E}|\xi-Y_{s_0}^{\xi}|^2)^{\frac{1}{2}}e^{-\frac{\beta_1-\beta_2}{2}s}\Big]$$

$$\leqslant Ce^{-\frac{\beta_1-\beta_2}{2}s}(1+|y|^2+\nu(|\cdot|^2))^{1/2}.$$

6.2. **Proof of** (38). Without loss of generality, we will use $(\Omega, \mathscr{F}, \{\mathscr{F}_t\}_{t\geqslant 0}, \mathbb{P})$ instead of $(\hat{\Omega}, \hat{\mathscr{F}}, \{\hat{\mathscr{F}}_t\}_{t\geqslant 0}, \hat{\mathbb{P}})$, and $(N^{\varepsilon,2}, N^2, X^{\varepsilon}, Y^{\varepsilon,y_0,\mathscr{L}_{\xi}}, Y^{\varepsilon,\xi}, \bar{X})$ instead of $(\hat{N}^{\varepsilon_{n_k},2}, \hat{N}^2, \hat{X}^{\varepsilon_{n_k}}, \hat{Y}^{\varepsilon_{n_k},2}, \hat{Y}^{\varepsilon_{n_k},2}, \hat{X}^{\varepsilon_{n_k},2}, \hat{X}^{\varepsilon_{n_k},2}, \hat{X}^{\varepsilon_{n_k},2})$ below.

We define the function

$$H(x, \mu, y, \nu) = [(\partial_y \Psi_{\sigma_2})(\partial_y \Psi_{\sigma_2})^*](x, \mu, y, \nu),$$

$$\bar{H}(x, \mu) = \int_{\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m)} H(x, \mu, y, \nu) \eta \times \delta_{\eta}(\mathrm{d}y, \mathrm{d}\nu).$$

Since $N^{\varepsilon,2}$ converges to N^2 , \mathbb{P} -a.s., we aim to prove

$$\langle N^2 \rangle_t = \int_0^t \overline{(\partial_y \Psi_{\sigma_2})(\partial_y \Psi_{\sigma_2})^*} (\bar{X}_s, \mathscr{L}_{\bar{X}_s}) ds, t \in [0, T].$$

We claim:

$$\lim_{\varepsilon \to 0} \mathbb{E} \left\| \int_0^t H(X_s^{\varepsilon}, \mathscr{L}_{X_s^{\varepsilon}}, Y_s^{\varepsilon, y_0, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_s^{\varepsilon, \xi}}) \mathrm{d}s - \int_0^t \bar{H}(\bar{X}_s, \mathscr{L}_{\bar{X}_s}) \mathrm{d}s \right\|^2 = 0. \tag{42}$$

Actually, if (42) holds, there exists a subsequence $\{\varepsilon_k\}_{k\geq 1}$ tending to 0 such that

$$\int_0^t H(X_s^{\varepsilon_k}, \mathscr{L}_{X_s^{\varepsilon_k}}, Y_s^{\varepsilon_k, y_0, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_s^{\varepsilon_k, \xi}}) \mathrm{d}s \to \int_0^t \bar{H}(\bar{X}_s, \mathscr{L}_{\bar{X}_s}) \mathrm{d}s, \quad \mathbb{P}\text{-a.s., as} \quad k \to \infty.$$

Moreover, note that $N_t^{\varepsilon_k,2} \otimes N_t^{\varepsilon_k,2} - \int_0^t H(X_s^{\varepsilon_k}, \mathscr{L}_{X_s^{\varepsilon_k}}, Y_s^{\varepsilon_k,y_0,\mathscr{L}_\xi}, \mathscr{L}_{Y_s^{\varepsilon_k,\xi}}) \mathrm{d}s$ is a matrix-valued martingale. Using the Vitali convergence theorem, we conclude that for any $0 \le s \le t \le t$ T

$$\begin{split} & \mathbb{E}\Big[\Big(N_t^2\otimes N_t^2 - \int_0^t \bar{H}(\bar{X}_r, \mathscr{L}_{\bar{X}_r})\mathrm{d}r\Big) - \Big(N_s^2\otimes N_s^2 - \int_0^s \bar{H}(\bar{X}_r, \mathscr{L}_{\bar{X}_r})\mathrm{d}r\Big)\Big|\mathscr{F}_s\Big] \\ &= \lim_{k\to\infty} \mathbb{E}\Big[\Big(N_t^{\varepsilon_k,2}\otimes N_t^{\varepsilon_k,2} - \int_0^t H(X_r^{\varepsilon_k}, \mathscr{L}_{X_r^{\varepsilon_k}}, Y_r^{\varepsilon_k,y_0,\mathscr{L}_\xi}, \mathscr{L}_{Y_r^{\varepsilon_k,\xi}})\mathrm{d}r\Big) \\ & - \Big(N_s^{\varepsilon_k,2}\otimes N_s^{\varepsilon_k,2} - \int_0^s H(X_r^{\varepsilon_k}, \mathscr{L}_{X_r^{\varepsilon_k}}, Y_r^{\varepsilon_k,y_0,\mathscr{L}_\xi}, \mathscr{L}_{Y_r^{\varepsilon_k,\xi}})\mathrm{d}r\Big)\Big|\mathscr{F}_s\Big] \\ &= 0, \quad \mathbb{P}\text{-a.s.}, \end{split}$$

which implies that $N_t^2 \otimes N_t^2 - \int_0^t \bar{H}(\bar{X}_r, \mathscr{L}_{\bar{X}_r}) dr$ is a matrix-valued martingale. Thus, we conclude that

$$\langle N^2 \rangle_t = \int_0^t \overline{(\partial_y \Psi_{\sigma_2})(\partial_y \Psi_{\sigma_2})^*} (\bar{X}_s, \mathscr{L}_{\bar{X}_s}) \mathrm{d}s, \quad t \in [0, T].$$

- 6.2.1. Proof of the claim (42). We divide the proof into three steps. The first step is to prove the locally Lipschitz continuity of $H(x, \mu, y, \nu)$. The second step is to demonstrate (42). The third step is to supplement the proof of estimate (56) required in the second step.
- **Step 1.** We prove that $H(x,\mu,y,\nu)$ is locally Lipschitz continuous, and $H(x,\mu)$ is Lipschitz continuous.
- (15) implies that $\partial_y \Psi(x, \mu, y, \nu)$ is bounded and Lipschitz continuous w.r.t. (x, μ, y, ν) , thus

$$\|\partial_{y}\Psi_{\sigma_{2}}(x_{1},\mu_{1},y_{1},\nu_{1}) - \partial_{y}\Psi_{\sigma_{2}}(x_{2},\mu_{2},y_{2},\nu_{2})\|$$

$$\leq \|\partial_{y}\Psi(x_{1},\mu_{1},y_{1},\nu_{1}) - \partial_{y}\Psi(x_{2},\mu_{2},y_{2},\nu_{2})\| \cdot \|\sigma_{2}(y_{1},\nu_{1})\|$$

$$+ \|\partial_{y}\Psi(x_{2},\mu_{2},y_{2},\nu_{2})\| \cdot \|\sigma_{2}(y_{1},\nu_{1}) - \sigma_{2}(y_{2},\nu_{2})\|$$

$$\leq C(|x_{1}-x_{2}| + \mathbb{W}_{2}(\mu_{1},\mu_{2}) + |y_{1}-y_{2}| + \mathbb{W}_{2}(\nu_{1},\nu_{2}))(1 + |y_{1}|^{2} + \nu_{1}(|\cdot|^{2}))^{1/2}, (43)$$

and

 $\|\partial_{\nu}\Psi_{\sigma_2}(x,\mu,y,\nu)\| \le \|\partial_{\nu}\Psi(x,\mu,y,\nu)\| \cdot \|\sigma_2(y,\nu)\| \le C(1+|y|^2+\nu(|\cdot|^2))^{1/2}.$ (44)Similarly, we have

$$\|\partial_y \Psi_{\sigma_2}(x_1, \mu_1, y_1, \nu_1) - \partial_y \Psi_{\sigma_2}(x_2, \mu_2, y_2, \nu_2)\|$$

$$\leq C(|x_1-x_2|+\mathbb{W}_2(\mu_1,\mu_2)+|y_1-y_2|+\mathbb{W}_2(\nu_1,\nu_2))(1+|y_2|^2+\nu_2(|\cdot|^2))^{1/2}.$$
 (45)

By (43)-(45), it holds that

$$||H(x_{1}, \mu_{1}, y_{1}, \nu_{1}) - H(x_{2}, \mu_{2}, y_{2}, \nu_{2})||$$

$$\leq ||\partial_{y}\Psi_{\sigma_{2}}(x_{1}, \mu_{1}, y_{1}, \nu_{1}) - \partial_{y}\Psi_{\sigma_{2}}(x_{2}, \mu_{2}, y_{2}, \nu_{2})|| \cdot ||\partial_{y}\Psi_{\sigma_{2}}(x_{1}, \mu_{1}, y_{1}, \nu_{1})||$$

$$+ ||\partial_{y}\Psi_{\sigma_{2}}(x_{2}, \mu_{2}, y_{2}, \nu_{2})|| \cdot ||\partial_{y}\Psi_{\sigma_{2}}(x_{1}, \mu_{1}, y_{1}, \nu_{1}) - \partial_{y}\Psi_{\sigma_{2}}(x_{2}, \mu_{2}, y_{2}, \nu_{2})||$$

$$\leq C(1 + |y_{1}|^{2} + \nu_{1}(|\cdot|^{2}) + |y_{2}|^{2} + \nu_{2}(|\cdot|^{2}))$$

$$\cdot (|x_{1} - x_{2}| + \mathbb{W}_{2}(\mu_{1}, \mu_{2}) + |y_{1} - y_{2}| + \mathbb{W}_{2}(\nu_{1}, \nu_{2})), \tag{46}$$

i.e. $H(x,\mu,y,\nu)$ is locally Lipschitz continuous. By uniform boundedness of $\partial_y \Psi(x,\mu,y,\nu)$ and Remark 2.5,we have

$$||H(x,\mu,y,\nu)|| = ||[(\partial_y \Psi_{\sigma_2})(\partial_y \Psi_{\sigma_2})^*](x,\mu,y,\nu)|| \le C(1+|y|^2+\nu(|\cdot|^2)). \tag{47}$$

Combining Lemma 4.2, Lemma 4.3 and (46), we obtain

$$\begin{split} & \left\| \mathbb{E}H(x,\mu,Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}}) - \bar{H}(x,\mu) \right\| \\ &= \left\| \mathbb{E}H(x,\mu,Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}}) - \int_{\mathbb{R}^{m}\times\mathcal{P}_{2}(\mathbb{R}^{m})} H(x,\mu,y,\nu)\eta \times \delta_{\eta}(\mathrm{d}y,\mathrm{d}\nu) \right\| \\ \mathcal{L}_{\xi}^{=\nu} & \left\| \mathbb{E}H(x,\mu,Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}}) - \int_{\mathbb{R}^{m}\times\mathcal{P}_{2}(\mathbb{R}^{m})} \mathbb{E}H(x,\mu,Y_{t}^{y,\nu},\mathcal{L}_{Y_{t}^{\zeta}})\eta \times \delta_{\eta}(\mathrm{d}y,\mathrm{d}\nu) \right\| \\ & \leqslant \int_{\mathbb{R}^{m}\times\mathcal{P}_{2}(\mathbb{R}^{m})} \mathbb{E} \left\| H(x,\mu,Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}}) - H(x,\mu,Y_{t}^{y,\nu},\mathcal{L}_{Y_{t}^{\zeta}}) \right\| \eta \times \delta_{\eta}(\mathrm{d}y,\mathrm{d}\nu) \\ & \stackrel{\mathcal{L}_{\xi}^{=\eta}}{=} \int_{\mathbb{R}^{m}} \mathbb{E} \left\| H(x,\mu,Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}}) - H(x,\mu,Y_{t}^{y,\eta},\mathcal{L}_{Y_{t}^{\zeta}}) \right\| \eta(\mathrm{d}y) \\ & \leqslant C \int_{\mathbb{R}^{m}} \left[1 + (\mathbb{E}|Y_{t}^{y_{0},\mathcal{L}_{\xi}}|^{4})^{\frac{1}{2}} + \mathbb{E}|Y_{t}^{\xi}|^{2} + (\mathbb{E}|Y_{t}^{y,\eta}|^{4})^{\frac{1}{2}} + \mathbb{E}|Y_{t}^{\zeta}|^{2} \right] \\ & \qquad \cdot \left[(\mathbb{E}|Y_{t}^{y_{0},\mathcal{L}_{\xi}} - Y_{t}^{y,\eta}|^{2})^{\frac{1}{2}} + (\mathbb{E}|Y_{t}^{\xi} - Y_{t}^{\zeta}|^{2})^{\frac{1}{2}} \right] \eta(\mathrm{d}y) \\ & \leqslant C \int_{\mathbb{R}^{m}} \left[1 + |y_{0}|^{2} + (\mathbb{E}|\xi|^{4})^{\frac{1}{2}} + |y|^{2} + (\mathbb{E}|\xi|^{4})^{\frac{1}{2}} \right] \\ & \qquad \cdot \left[|y_{0} - y|e^{-\frac{\beta_{1}}{2}t} + (\mathbb{E}|\xi - \tilde{\zeta}|^{2})^{\frac{1}{2}} e^{-\frac{\beta_{1}-\beta_{2}}{2}t} \right] \eta(\mathrm{d}y) \\ & \leqslant C e^{-\frac{\beta_{1}-\beta_{2}}{2}t} (1 + |y_{0}|^{4} + \mathbb{E}|\xi|^{4}). \end{split}$$

By integrating (46)-(48) and Lemma 4.2, we attain

$$\|\bar{H}(x_{1},\mu_{1}) - \bar{H}(x_{2},\mu_{2})\|$$

$$\leq \|\mathbb{E}H(x_{1},\mu_{1},Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}}) - \bar{H}(x_{1},\mu_{1})\| + \|\mathbb{E}H(x_{2},\mu_{2},Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}}) - \bar{H}(x_{2},\mu_{2})\|$$

$$+ \|\mathbb{E}H(x_{1},\mu_{1},Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}}) - \mathbb{E}H(x_{2},\mu_{2},Y_{t}^{y_{0},\mathcal{L}_{\xi}},\mathcal{L}_{Y_{t}^{\xi}})\|$$

$$\leq Ce^{-\frac{\beta_{1}-\beta_{2}}{2}t}(1+|y_{0}|^{4}+\mathbb{E}|\xi|^{4})$$

$$+C\mathbb{E}\Big[\Big(1+|y_{0}|^{2}e^{-\alpha_{1}t}+\mathbb{E}|\xi|^{2}e^{-(\alpha_{1}-\alpha_{2})t}\Big) \cdot (|x_{1}-x_{2}|+\mathbb{W}_{2}(\mu_{1},\mu_{2}))\Big],$$

and

$$\|\bar{H}(x,\mu)\| = \left\| \int_{\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m)} H(x,\mu,y,\nu) \eta \times \delta_{\eta}(\mathrm{d}y,\mathrm{d}\nu) \right\|$$

$$\stackrel{\mathcal{L}_{\zeta}=\nu}{=} \left\| \int_{\mathbb{R}^m \times \mathcal{P}_2(\mathbb{R}^m)} \mathbb{E} H(x,\mu,Y_t^{y,\nu},\mathcal{L}_{Y_t^{\zeta}}) \eta \times \delta_{\eta}(\mathrm{d}y,\mathrm{d}\nu) \right\|$$

$$\stackrel{\mathcal{L}_{\zeta}=\eta}{\leqslant} \int_{\mathbb{R}^m} \mathbb{E} \left\| H(x,\mu,Y_t^{y,\eta},\mathcal{L}_{Y_t^{\zeta}}) \right\| \eta(\mathrm{d}y)$$

$$\stackrel{\mathcal{L}_{\zeta}=\eta}{\leqslant} C \int_{\mathbb{R}^m} \left(1 + |y|^2 e^{-\alpha_1 t} + \mathbb{E} |\tilde{\zeta}|^2 e^{-(\alpha_1 - \alpha_2)t} \right) \eta(\mathrm{d}y)$$

$$\stackrel{\mathcal{L}_{\zeta}=\eta}{\leqslant} C + C e^{-(\alpha_1 - \alpha_2)t} (1 + \mathbb{E} |\xi|^2).$$

Letting $t \to \infty$, we obtain

$$\|\bar{H}(x_1, \mu_1) - \bar{H}(x_2, \mu_2)\| \leqslant C(|x_1 - x_2| + \mathbb{W}_2(\mu_1, \mu_2)). \tag{49}$$

and

$$\|\bar{H}(x,\mu)\| \leqslant C. \tag{50}$$

Step 2. We prove (42).

Note that

$$X_t^{\varepsilon} - X_{t(\Delta)}^{\varepsilon} = \int_{t(\Delta)}^t b_1(X_s^{\varepsilon}, \mathcal{L}_{X_s^{\varepsilon}}, Y_s^{\varepsilon, y_0, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_s^{\varepsilon, \xi}}) ds + \int_{t(\Delta)}^t \sigma_1(X_s^{\varepsilon}, \mathcal{L}_{X_s^{\varepsilon}}) dB_s,$$

where $t(\triangle) := \left[\frac{t}{\triangle}\right] \triangle$ and $\left[\frac{t}{\triangle}\right]$ denotes the integer part of $\frac{t}{\triangle}$. By Remark 2.5, Hölder inequality, Burkholder-Davis-Gundy inequality and Lemma 4.1, we have

$$\mathbb{E}|X_{t}^{\varepsilon} - X_{t(\Delta)}^{\varepsilon}|^{4}$$

$$\leqslant C\Delta^{3}\mathbb{E}\int_{t(\Delta)}^{t} \left|b_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}})\right|^{4} ds + C\mathbb{E}\left[\int_{t(\Delta)}^{t} \left\|\sigma_{1}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}})\right\|^{2} ds\right]^{2}$$

$$\leqslant C(\Delta^{3} + \Delta)\int_{t(\Delta)}^{t} \left[1 + \mathbb{E}|X_{s}^{\varepsilon}|^{4} + \mathbb{E}|Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}|^{4} + \mathbb{E}|Y_{s}^{\varepsilon, \xi}|^{4}\right] ds$$

$$\leqslant C_{T}(\Delta^{4} + \Delta^{2})(1 + \mathbb{E}|\varrho|^{12} + |y_{0}|^{12} + \mathbb{E}|\xi|^{12}).$$
(51)

Notice that

$$\int_{0}^{t} \left[H(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}}) - \bar{H}(\bar{X}_{s}, \mathcal{L}_{\bar{X}_{s}}) \right] ds$$

$$= \int_{0}^{t} \left[H(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}}) - H(X_{s(\Delta)}^{\varepsilon}, \mathcal{L}_{X_{s(\Delta)}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}}) \right] ds$$

$$+ \int_{0}^{t} \left[H(X_{s(\Delta)}^{\varepsilon}, \mathcal{L}_{X_{s(\Delta)}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}}) - \bar{H}(X_{s(\Delta)}^{\varepsilon}, \mathcal{L}_{X_{s(\Delta)}^{\varepsilon}}) \right] ds$$

$$+ \int_{0}^{t} \left[\bar{H}(X_{s(\Delta)}^{\varepsilon}, \mathcal{L}_{X_{s(\Delta)}^{\varepsilon}}) - \bar{H}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}) \right] ds$$

$$+ \int_{0}^{t} \left[\bar{H}(X_{s}^{\varepsilon}, \mathcal{L}_{X_{s}^{\varepsilon}}) - \bar{H}(\bar{X}_{s}, \mathcal{L}_{\bar{X}_{s}^{\varepsilon}}) \right] ds$$

$$=: \mathcal{O}_1(t) + \mathcal{O}_2(t) + \mathcal{O}_3(t) + \mathcal{O}_4(t). \tag{52}$$

For $\mathcal{O}_1(t)$, based on the Hölder inequality, (46), (51) and Lemma 4.1, one can obtain that

$$\mathbb{E}\|\mathscr{O}_{1}(t)\|^{2}$$

$$\leqslant C_{T}\mathbb{E}\int_{0}^{t} \left(1 + |Y_{s}^{\varepsilon,y_{0},\mathscr{L}_{\xi}}|^{2} + \|\mathscr{L}_{Y_{s}^{\varepsilon,\xi}}\|^{2}\right)^{2} \left(|X_{s}^{\varepsilon} - X_{s(\triangle)}^{\varepsilon}| + \mathbb{W}_{2}(\mathscr{L}_{X_{s}^{\varepsilon}}, \mathscr{L}_{X_{s(\triangle)}^{\varepsilon}})\right)^{2} ds$$

$$\leqslant C_{T}\int_{0}^{t} \left(1 + \mathbb{E}|Y_{s}^{\varepsilon,y_{0},\mathscr{L}_{\xi}}|^{8} + \mathbb{E}|Y_{s}^{\varepsilon,\xi}|^{8}\right)^{\frac{1}{2}} \cdot \left(\mathbb{E}|X_{s}^{\varepsilon} - X_{s(\triangle)}^{\varepsilon}|^{4}\right)^{\frac{1}{2}} ds$$

$$\leqslant C_{T}(\triangle^{2} + \triangle)(1 + \mathbb{E}|\varrho|^{12} + |y_{0}|^{12} + \mathbb{E}|\xi|^{12}).$$
(53)

With regard to $\mathcal{O}_3(t)$ and $\mathcal{O}_4(t)$, by the Hölder inequality, (49), (51) and Theorem 3.1, we get

$$\mathbb{E}\|\mathscr{O}_{3}(t)\|^{2} + \mathbb{E}\|\mathscr{O}_{4}(t)\|^{2} \leqslant C_{T} \int_{0}^{t} \left[\mathbb{E}|X_{s}^{\varepsilon} - X_{s(\Delta)}^{\varepsilon}|^{2} + \mathbb{E}|X_{s}^{\varepsilon} - \bar{X}_{s}|^{2}\right] ds$$

$$\leqslant C_{T}(\Delta^{2} + \Delta + \varepsilon)(1 + \mathbb{E}|\varrho|^{12} + |y_{0}|^{12} + \mathbb{E}|\xi|^{12}), \quad (54)$$

Now we focus on the term $\mathcal{O}_2(t)$. Notice that

$$\mathbb{E}\|\mathscr{O}_{2}(t)\|^{2} \\
= \mathbb{E}\left\|\sum_{k=0}^{[t/\Delta]-1} \int_{k\Delta}^{(k+1)\Delta} \left[H(X_{s(\Delta)}^{\varepsilon}, \mathscr{L}_{X_{s(\Delta)}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_{s}^{\varepsilon, \xi}}) - \bar{H}(X_{s(\Delta)}^{\varepsilon}, \mathscr{L}_{X_{s(\Delta)}^{\varepsilon}}) \right] ds \\
+ \int_{t(\Delta)}^{t} \left[H(X_{s(\Delta)}^{\varepsilon}, \mathscr{L}_{X_{s(\Delta)}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_{s}^{\varepsilon, \xi}}) - \bar{H}(X_{s(\Delta)}^{\varepsilon}, \mathscr{L}_{X_{s(\Delta)}^{\varepsilon}}) \right] ds \right\|^{2} \\
\leqslant \frac{C_{T}}{\Delta} \sum_{k=0}^{[t/\Delta]-1} \mathbb{E}\left\| \int_{k\Delta}^{(k+1)\Delta} \left[H(X_{s(\Delta)}^{\varepsilon}, \mathscr{L}_{X_{s(\Delta)}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_{s}^{\varepsilon, \xi}}) - \bar{H}(X_{s(\Delta)}^{\varepsilon}, \mathscr{L}_{X_{s(\Delta)}^{\varepsilon}}) \right] ds \right\|^{2} \\
+2\mathbb{E}\left\| \int_{t(\Delta)}^{t} \left[H(X_{s(\Delta)}^{\varepsilon}, \mathscr{L}_{X_{s(\Delta)}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathscr{L}_{\xi}}, \mathscr{L}_{Y_{s}^{\varepsilon, \xi}}) - \bar{H}(X_{s(\Delta)}^{\varepsilon}, \mathscr{L}_{X_{s(\Delta)}^{\varepsilon}}) \right] ds \right\|^{2} \\
=: \mathscr{U}_{1}(t) + \mathscr{U}_{2}(t). \tag{55}$$

For \mathcal{U}_1 , it holds that

$$\mathscr{U}_1(t) \leqslant C_T(1+|y_0|^{12}+\mathbb{E}|\xi|^{12})\frac{\varepsilon}{\Delta},\tag{56}$$

For \mathcal{U}_2 , combining the Hölder inequality, (47), (50) and Lemma 4.1, we obtain

$$\mathcal{U}_{2}(t) \leqslant C \triangle \mathbb{E} \int_{t(\triangle)}^{t} \left(\|H(X_{s(\triangle)}^{\varepsilon}, \mathcal{L}_{X_{s(\triangle)}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}})\|^{2} + \|\bar{H}(X_{s(\triangle)}^{\varepsilon}, \mathcal{L}_{X_{s(\triangle)}^{\varepsilon}})\|^{2} \right) ds$$

$$\leqslant C \triangle \mathbb{E} \int_{t(\triangle)}^{t} \left(1 + |Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}|^{4} + \|\mathcal{L}_{Y_{s}^{\varepsilon, \xi}}\|^{4} \right) ds$$

$$\leqslant C \triangle^{2} (1 + |y_{0}|^{12} + \mathbb{E}|\xi|^{12}). \tag{57}$$

Finally, (52)-(57) yield that

$$\mathbb{E} \left\| \int_0^t H(X_s^{\varepsilon}, \mathcal{L}_{X_s^{\varepsilon}}, Y_s^{\varepsilon, y_0, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_s^{\varepsilon, \xi}}) ds - \int_0^t \bar{H}(\bar{X}_s, \mathcal{L}_{\bar{X}_s}) ds \right\|^2$$

$$\leqslant C_T(\Delta^2 + \Delta + \varepsilon + \frac{\varepsilon}{\Delta}) (1 + \mathbb{E}|\varrho|^{12} + |y_0|^{12} + \mathbb{E}|\xi|^{12}).$$

Taking $\triangle = \varepsilon^{1/2}$, we infer that (42) holds.

Step 3. We prove (56).

Note that

$$\mathcal{U}_{1}(t)$$

$$\leq \frac{C_{T}}{\Delta} \sum_{k=0}^{[T/\Delta]-1} \mathbb{E} \left\| \int_{k\Delta}^{(k+1)\Delta} \left[H(X_{k\Delta}^{\varepsilon}, \mathcal{L}_{X_{k\Delta}^{\varepsilon}}, Y_{s}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s}^{\varepsilon, \xi}}) - \bar{H}(X_{k\Delta}^{\varepsilon}, \mathcal{L}_{X_{k\Delta}^{\varepsilon}}) \right] ds \right\|^{2}$$

$$\leq \frac{C_{T}}{\Delta^{2}} \sup_{0 \leq k \leq [T/\Delta]-1} \mathbb{E} \left\| \int_{0}^{\Delta} \left[H(X_{k\Delta}^{\varepsilon}, \mathcal{L}_{X_{k\Delta}^{\varepsilon}}, Y_{s+k\Delta}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s+k\Delta}^{\varepsilon, \xi}}) - \bar{H}(X_{k\Delta}^{\varepsilon}, \mathcal{L}_{X_{k\Delta}^{\varepsilon}}) \right] ds \right\|^{2} .$$

On one hand, for any $s \in [0, \triangle]$,

$$Y_{s+k\triangle}^{\varepsilon,y_0,\mathcal{L}_{\xi}} = Y_{k\triangle}^{\varepsilon,y_0,\mathcal{L}_{\xi}} + \frac{1}{\varepsilon} \int_{k\triangle}^{s+k\triangle} b_2(Y_r^{\varepsilon,y_0,\mathcal{L}_{\xi}}, \mathcal{L}_{Y_r^{\varepsilon,\xi}}) dr$$

$$+ \frac{1}{\sqrt{\varepsilon}} \int_{k\triangle}^{s+k\triangle} \sigma_2(Y_r^{\varepsilon,y_0,\mathcal{L}_{\xi}}, \mathcal{L}_{Y_r^{\varepsilon,\xi}}) dW_r$$

$$= Y_{k\triangle}^{\varepsilon,y_0,\mathcal{L}_{\xi}} + \frac{1}{\varepsilon} \int_0^s b_2(Y_{u+k\triangle}^{\varepsilon,y_0,\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{u+k\triangle}^{\varepsilon,\xi}}) du$$

$$+ \frac{1}{\sqrt{\varepsilon}} \int_0^s \sigma_2(Y_{u+k\triangle}^{\varepsilon,y_0,\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{u+k\triangle}^{\varepsilon,\xi}}) d\bar{W}_u,$$

where $\overline{W}_{\cdot} := W_{\cdot + k\triangle} - W_{k\triangle}$ is a m-dimensional standard Brownian motion. On the other hand, we construct a system as follows:

$$\begin{cases} d\check{Y}_{t}^{\varepsilon,\xi} = b_{2}(\check{Y}_{t}^{\varepsilon,\xi}, \mathcal{L}_{\check{Y}_{t}^{\varepsilon,\xi}})dt + \sigma_{2}(\check{Y}_{t}^{\varepsilon,\xi}, \mathcal{L}_{\check{Y}_{t}^{\varepsilon,\xi}})d\check{W}_{t}, \\ \check{Y}_{0}^{\varepsilon,\xi} = Y_{k\triangle}^{\varepsilon,\xi}, \quad 0 \leqslant t \leqslant \frac{\triangle}{\varepsilon}, \\ d\check{Y}_{t}^{\varepsilon,y_{0},\mathcal{L}_{\xi}} = b_{2}(\check{Y}_{t}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{\check{Y}_{t}^{\varepsilon,\xi}})dt + \sigma_{2}(\check{Y}_{t}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{\check{Y}_{t}^{\varepsilon,\xi}})d\check{W}_{t}, \\ \check{Y}_{0}^{\varepsilon,y_{0},\mathcal{L}_{\xi}} = Y_{k\triangle}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \quad 0 \leqslant t \leqslant \frac{\triangle}{\varepsilon}, \end{cases}$$

where \check{W} is a m-dimensional standard Brownian motion independent of B, W. For any $s \in [0, \Delta]$,

where $\check{\check{W}}:=\sqrt{\varepsilon}\check{W}_{\frac{\cdot}{\varepsilon}}$ is a m-dimensional standard Brownian motion.

Thus, $(Y_{s+k\triangle}^{\varepsilon,y_0,\mathscr{L}_{\xi}},\mathscr{L}_{Y_{s+k\triangle}^{\varepsilon,\xi}})$ and $(\check{Y}_{\frac{s}{\varepsilon}}^{\varepsilon,y_0,\mathscr{L}_{\xi}},\mathscr{L}_{\check{Y}_{\frac{s}{\varepsilon}}^{\varepsilon,\xi}})$ have the same distribution, which yields that

$$\mathbb{E} \left\| \int_{0}^{\triangle} \left[H(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, Y_{s+k\triangle}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s+k\triangle}^{\varepsilon,\xi}}) - \bar{H}(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \right] ds \right\|^{2} \\
= \mathbb{E} \left\| \int_{0}^{\triangle} \left[H(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, \check{Y}_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{\check{Y}_{s}^{\varepsilon,\xi}}) - \bar{H}(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \right] ds \right\|^{2} \\
= \varepsilon^{2} \mathbb{E} \left\| \int_{0}^{\triangle} \left[H(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, \check{Y}_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{\check{Y}_{s}^{\varepsilon,\xi}}) - \bar{H}(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \right] ds \right\|^{2} \\
= 2\varepsilon^{2} \int_{0}^{\triangle} \int_{r}^{\triangle} \mathbb{E} \left\langle H(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, \check{Y}_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{\check{Y}_{s}^{\varepsilon,\xi}}) - \bar{H}(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \right] ds dr \\
= 2\varepsilon^{2} \int_{0}^{\triangle} \int_{r}^{\triangle} \mathbb{E} \left[\mathbb{E} \left[\left\langle H(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, \check{Y}_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{\check{Y}_{s}^{\varepsilon,\xi}}) - \bar{H}(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \right\rangle ds dr \\
= 2\varepsilon^{2} \int_{0}^{\triangle} \int_{r}^{\triangle} \mathbb{E} \left[\mathbb{E} \left[\left\langle H(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, \check{Y}_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{\check{Y}_{s}^{\varepsilon,\xi}}) - \bar{H}(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \right\rangle ds dr \\
= 2\varepsilon^{2} \int_{0}^{\triangle} \int_{r}^{\triangle} \mathbb{E} \left[\mathbb{E} \left[\left\langle H(X_{k\triangle}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, \check{Y}_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{\check{Y}_{s}^{\varepsilon,\xi}}) - \bar{H}(X_{k\triangle}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \right\rangle ds dr \right] ds dr \\
= 2\varepsilon^{2} \int_{0}^{\triangle} \int_{r}^{\triangle} \mathbb{E} \left[\mathbb{E} \left[\left\langle H(X_{k\triangle}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, \check{Y}_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{\check{Y}_{s}^{\varepsilon,\xi}}) - \bar{H}(X_{k\triangle}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \right\rangle ds dr \right\} ds dr \right\} ds dr$$

where $\mathscr{G}_{k\triangle}$ is the σ -field generated by $X_{k\triangle}^{\varepsilon}$ and $\mathscr{F}_{0}^{\check{W}}$. In order to estimate the integrand in the right side of the above equality, we investigate the property of $(\check{Y}^{\varepsilon,y_0,\mathscr{L}_{\xi}},\mathscr{L}_{\check{Y}^{\varepsilon,\xi}})$. Let $C([0,\triangle/\varepsilon];\mathbb{R}^m)$ be the collection of continuous functions from $[0, \Delta/\varepsilon]$ to \mathbb{R}^m , endowed with the uniform convergence topology. Set

$$\tilde{\Omega} := C\left([0, \Delta/\varepsilon]; \mathbb{R}^m\right) \times C\left([0, \Delta/\varepsilon]; \mathcal{P}_2\left(\mathbb{R}^m\right)\right),
\tilde{\mathscr{F}} := \mathscr{B}\left(C\left([0, \Delta/\varepsilon]; \mathbb{R}^m\right)\right) \times \mathscr{B}\left(C\left([0, \Delta/\varepsilon]; \mathcal{P}_2\left(\mathbb{R}^m\right)\right)\right),
\tilde{\mathscr{F}}_t := \sigma\left(M_r, 0 \leqslant r \leqslant t\right), \quad 0 \leqslant t \leqslant \Delta/\varepsilon,$$

where M_i is the coordinate process. Following a similar reasoning to that in [22, Theorem 4.11], there exists a unique probability measure $\tilde{\mathbb{P}}$ on $(\tilde{\Omega}, \tilde{\mathscr{F}})$ such that M is a homogeneous Markov process with respect to $(\tilde{\mathscr{F}}_t)$, and its transition function is given by $\{\mathscr{L}_{\check{Y}_t^{\varepsilon,y_0,\mathscr{L}_{\xi}}} \times$ $\delta_{\mathscr{L}_{\hat{Y}^{\varepsilon,\xi}}}: 0 \leqslant t \leqslant \triangle/\varepsilon\}.$ Note that

$$\mathscr{L}_{M_t}^{\tilde{\mathbb{P}}} := \tilde{\mathbb{P}} \circ M_t^{-1} = \mathscr{L}_{\check{Y}_t^{\varepsilon,y_0},\mathscr{L}_{\xi}} \times \delta_{\mathscr{L}_{\check{Y}_t^{\varepsilon,\xi}}}.$$

Therefore, it follows that for $0 \le r \le s \le \Delta/\varepsilon$

$$\mathbb{E}\Big\langle H(x,\mathscr{L}_{X_{k\triangle}^{\varepsilon}},\check{Y}_{s}^{\varepsilon,y_{0},\mathscr{L}_{\xi}},\mathscr{L}_{\check{Y}_{s}^{\varepsilon,\xi}}) - \bar{H}(x,\mathscr{L}_{X_{k\triangle}^{\varepsilon}}),$$

$$H(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, \check{Y}_{r}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{\check{Y}_{r}^{\varepsilon,\xi}}) - \bar{H}(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \rangle$$

$$= \mathbb{E}\Big[\mathbb{E}\Big\langle H(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, \check{Y}_{s}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{\check{Y}_{s}^{\varepsilon,\xi}}) - \bar{H}(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}),$$

$$H(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, \check{Y}_{r}^{\varepsilon,y_{0},\mathcal{L}_{\xi}}, \mathcal{L}_{\check{Y}_{r}^{\varepsilon,\xi}}) - \bar{H}(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \Big\rangle \Big| \mathcal{F}_{r} \Big]$$

$$= \tilde{\mathbb{E}}\Big[\Big\langle \tilde{\mathbb{E}}\Big[H(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, M_{s}) - \bar{H}(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \Big| \tilde{\mathcal{F}}_{r} \Big],$$

$$H(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, M_{r}) - \bar{H}(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \Big\rangle \Big]$$

$$= \tilde{\mathbb{E}}\Big[\Big\langle \tilde{\mathbb{E}}^{M_{r}}\Big[H(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, M_{s-r}) - \bar{H}(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \Big],$$

$$H(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, M_{r}) - \bar{H}(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \Big\rangle \Big]$$

$$\leqslant \Big(\tilde{\mathbb{E}}\Big[\tilde{\mathbb{E}}^{M_{r}}\Big[H(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, M_{s-r}) - \bar{H}(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \Big]\Big|^{2}\Big)^{\frac{1}{2}}$$

$$\cdot \Big(\tilde{\mathbb{E}}\Big\|H(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, M_{r}) - \bar{H}(x, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \Big\|^{2}\Big)^{\frac{1}{2}},$$

$$(59)$$

where $\tilde{\mathbb{E}}$ is the expectation on $(\tilde{\Omega}, \tilde{\mathscr{F}}, {\{\tilde{\mathscr{F}}\}_{t\geq 0}, \tilde{\mathbb{P}}})$. Due to (48) and Lemma 4.1, we obtain that

$$\left(\tilde{\mathbb{E}}\Big|\tilde{\mathbb{E}}^{M_r}\left[H(x,\mathcal{L}_{X_{k\Delta}^{\varepsilon}},M_{s-r})-\bar{H}(x,\mathcal{L}_{X_{k\Delta}^{\varepsilon}})\right]\Big|^2\right)^{\frac{1}{2}}$$

$$\leqslant Ce^{-\frac{\beta_1-\beta_2}{2}(s-r)}\left(1+\mathbb{E}|Y_{k\Delta}^{\varepsilon,y_0,\mathcal{L}_{\xi}}|^8+\mathbb{E}|Y_{k\Delta}^{\varepsilon,\xi}|^8\right)^{\frac{1}{2}}$$

$$\leqslant Ce^{-\frac{\beta_1-\beta_2}{2}(s-r)}\left(1+|y_0|^{12}+\mathbb{E}|\xi|^{12}\right)^{\frac{1}{2}}.$$
(60)

By (47), (50) and Lemma 4.1, we have

$$\left(\tilde{\mathbb{E}} \left\| H(x, \mathcal{L}_{X_{k\Delta}^{\varepsilon}}, M_r) - \bar{H}(x, \mathcal{L}_{X_{k\Delta}^{\varepsilon}}) \right\|^2 \right)^{\frac{1}{2}} \\
\leqslant 2 \left(\tilde{\mathbb{E}} \left\| H(x, \mathcal{L}_{X_{k\Delta}^{\varepsilon}}, M_r) \right\|^2 + \left\| \bar{H}(x, \mathcal{L}_{X_{k\Delta}^{\varepsilon}}) \right\|^2 \right)^{\frac{1}{2}} \\
\leqslant C \left(1 + \mathbb{E} |Y_{k\Delta}^{\varepsilon, y_0, \mathcal{L}_{\xi}}|^4 + \mathbb{E} |Y_{k\Delta}^{\varepsilon, \xi}|^4 \right)^{\frac{1}{2}} \\
\leqslant C \left(1 + |y_0|^{12} + \mathbb{E} |\xi|^{12} \right)^{\frac{1}{2}},$$

which together with (58)-(60) yields that

$$\mathbb{E} \left\| \int_{0}^{\triangle} \left[H(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}, Y_{s+k\triangle}^{\varepsilon, y_{0}, \mathcal{L}_{\xi}}, \mathcal{L}_{Y_{s+k\triangle}^{\varepsilon, \xi}}) - \bar{H}(X_{k\triangle}^{\varepsilon}, \mathcal{L}_{X_{k\triangle}^{\varepsilon}}) \right] ds \right\|^{2} \\
\leqslant C \varepsilon^{2} \int_{0}^{\frac{\triangle}{\varepsilon}} \int_{r}^{\frac{\triangle}{\varepsilon}} e^{-\frac{\beta_{1} - \beta_{2}}{2}(s-r)} \left(1 + |y_{0}|^{12} + \mathbb{E}|\xi|^{12} \right) ds dr \\
\leqslant C \varepsilon \triangle \left(1 + |y_{0}|^{12} + \mathbb{E}|\xi|^{12} \right).$$

Therefore, the inequality (56) holds.

References

- [1] J. Bao, P. Ren and F-Y. Wang: Bismut formula for Lions derivative of distribution-path dependent SDEs, J. Differential Equations, 282(2021)285-329.
- [2] Z. Brzeźniak, E. Hausenblas and P.A. Razafimandimby: Stochastic reaction-diffusion equations driven by jump processes, *Potential Anal*, 49(2018)131-201.
- [3] R. Buckdahn, J. Li, S. Peng and C. Rainer: Mean-field stochastic differential equations and associated PDEs, *Ann. Probab.*, 45(2017)824-878.
- [4] G. Da Prato and J. Zabczyk: Stochastic equations in infinite dimensions, Cambridge University Press, Cambridge (1992).
- [5] X. Ding and H. Qiao: Euler-Maruyama approximations for stochastic McKean-Vlasov equations with non-Lipschitz coefficients, *J. Theoret. Probab.*, 34(2021)1408-1425.
- [6] X. Ding and H. Qiao: Stability for stochastic McKean-Vlasov equations with non-Lipschitz coefficients, SIAM J. Control Optim., 59(2021)887-905.
- [7] Z. Dong, X. Sun, H. Xiao and J. Zhai: Averaging principle for one dimensional stochastic Burgers equation, J. Differential Equations, 265(2018)4749-4797.
- [8] X. Fan, T. Yu and C. Yuan: Asymptotic behaviors for distribution dependent SDEs driven by fractional Brownian motions, *Stochastic Processes and their Applications*, 164(2023)383-415.
- [9] K. Fang, W. Liu, H. Qiao and F. Zhu: Asymptotic behaviors of small perturbation for multivalued McKean-Vlasov stochastic differential equations, Applied Mathematics and Optimization, 88(2023)22.
- [10] W. Hong, S. Li, W. Liu and X. Sun: Central limit type theorem and large deviation principle for multi-scale McKean-Vlasov SDEs, Probab. Theory Related Fields, 187(2023)133-201.
- [11] W. Hong, S. Li and W. Liu: Strong convergence rates in averaging principle for slow-fast McKean-Vlasov SPDEs, *J. Differential Equations*, 316(2022)94-135.
- [12] X. Huang and F.-Y. Wang: Distribution dependent SDEs with singular coefficients, Stochastic Process. Appl., 129(2019)4747-4770.
- [13] X. Huang, P. Ren and F.-Y. Wang: Distribution dependent stochastic differential equations, Front. Math. China, 16(2021)257-301.
- [14] Z. Huang: Basis of stochastic analysis (in Chinese), 2nd edn, Science Press, Beijing (2001).
- [15] R. Khasminskii: On the principle of averaging the Itô stochastic differential equations, Kibernetika, 4(1968)260-279.
- [16] C. Kumar and Neelima: On explicit Milstein-type scheme for McKean-Vlasov stochastic differential equations with super-linear drift coefficient, *Electron. J. Probab.*, 26(2021)1-32.
- [17] Y. Li and L. Xie: Functional law of large numbers and central limit theorem for slow-fast McKean-Vlasov equations, *Discrete Contin. Dyn. Syst. Ser. S*, 16(2023)846-877.
- [18] W. Liu, M. Röckner, X. Sun and Y. Xie: Averaging principle for slow-fast stochastic differential equations with time dependent locally Lipschitz coefficients, J. Differential Equations, 268(2020)2910-2948.
- [19] H.P. McKean: A class of Markov processes associated with nonlinear parabolic equations, Proc. Natl. Sci. U.S.A., 56(1966)1907-1911.
- [20] B. Pei, Y. Inahama and Y. Xu: Averaging principle for fast-slow system driven by mixed fractional Brownian rough path, J. Differential Equations, 301(2021)202-235.
- [21] H. Qiao and W. Wei: Strong approximation of nonlinear filtering for multiscale McKean-Vlasov stochastic systems, http://arxiv.org/abs/2206.05037.
- [22] P. Ren, M. Röckner and F.-Y. Wang: Linearization of nonlinear Fokker-Planck equations and applications, *J. Differential Equations*, 322(2022)1-37.
- [23] P. Ren and F.-Y. Wang: Space-Distribution PDEs for path independent additive functionals of McKean-Vlasov SDEs, Infin. Dimens. Anal. Quantum Probab. Relat. Top., 23(2020)2050018.
- [24] M. Röckner, X. Sun and Y. Xie: Strong convergence order for slow-fast McKean-Vlasov stochastic differential equations, *Ann. Inst. H. Poincaré Probab. Statist.*, 57(2021)547-576.
- [25] M. Röckner and X. Zhang: Well-posedness of distribution dependent SDEs with singular drifts, Bernoulli, 27(2021)1131-1158.

- [26] G. Shen, J. Xiang and J.-L. Wu: Averaging principle for distribution dependent stochastic differential equations driven by fractional Brownian motion and standard Brownian motion, *J. Differential Equations*, 321(2022)381-414.
- [27] G. Shen, J. Yin and J.-L. Wu: Stochastic averaging principle for two-time-scale SDEs with distribution dependent coefficients driven by fractional Brownian motion, Comm. Math. Statist (2023), https://doi.org/10.1007/s40304-023-00364-4.
- [28] Y. Suo and C. Yuan: Central limit theorem and moderate deviation principle for McKean-Vlasov SDEs, Acta Appl. Math., 175(2021)1-19.
- [29] F.-Y. Wang: Distribution dependent SDEs for Landau type equations, Stochastic Process. Appl., 128(2018)595-621.
- [30] F. Wu and G. Yin: Fast-slow-coupled stochastic functional differential equations, J. Differential Equations, 323(2022)1-37.
- [31] J. Xu, J. Liu and Y. Miao: Strong averaging principle for two-time-scale SDEs with non-Lipschitz coefficients, J. Math. Anal. Appl., 468(2018)116-140.
- [32] J. Xu and J. Liu: Stochastic averaging principle for two time-scale jump-diffusion SDEs under the non-Lipschitz coefficients, *Stochastics*, 93(2021)715-741.
- [33] J. Xu, J. Liu, J. Liu and Y. Miao: Strong averaging principle for two-time-scale stochastic McKean-Vlasov equations, *Appl. Math. Optim.*, 84(2021)837-867.